

A SIMPLIFIED PROOF OF PRIMES IN ALMOST ALL SHORT INTERVALS

RUNBO LI

ABSTRACT. We give a simplified proof of Jia's 1996 result which states that the interval $[n, n + n^{\frac{1}{20} + \varepsilon}]$ contains prime numbers for almost all n . Furthermore, we show that for almost all n , the shorter interval $[n, n + n^{\frac{1}{21} + \varepsilon}]$ still contains prime numbers. A delicate sieve decomposition and more accurate estimates for integrals are used to good effect.

CONTENTS

1. Introduction	1
2. An outline of the proof	3
3. Arithmetic Information I	3
4. Arithmetic Information II	22
5. Sieve error terms	23
6. Asymptotic formulas for type-II terms	23
7. The final decomposition	24
8. Further prospect	38
Acknowledgements	39
References	39
Appendix: Mathematica code for the theorem	40
Code for the asymptotic regions	40
Code for the numerical calculations	45
Code for the plot of regions	47

1. INTRODUCTION

One of the famous topics in number theory is to find prime numbers in short intervals. In 1937, Cramér [5] conjectured that every interval $[n, n + f(n)(\log n)^2]$ contains prime numbers for some $f(n) \rightarrow 1$ as $n \rightarrow \infty$. The Riemann Hypothesis implies that for all large n , the interval $[n, n + n^\theta]$ contains $\sim n^\theta (\log n)^{-1}$ prime numbers for every $\frac{1}{2} + \varepsilon \leq \theta \leq 1$. The first unconditional result of this asymptotic formula was proved by Hoheisel [17] in 1930 with $\theta = 1 - \frac{1}{33000}$. After the works of Hoheisel [17], Heilbronn [16], Chudakov [4], Ingham [19] and Montgomery [33], Huxley [18] proved in 1972 that the above asymptotic formula holds when $\theta > \frac{7}{12}$ by his zero density estimate. Very recently, Guth and Maynard [8] improved this to $\theta > \frac{17}{30}$ by a new zero density estimate.

In 1979, Iwaniec and Jutila [20] first introduced a sieve method into this problem. They established a lower bound with correct order (instead of an asymptotic formula) with $\theta = \frac{13}{23}$. After that breakthrough, many improvements were made and the value of θ was reduced successively to

$$\frac{5}{9} = 0.5556, \quad \frac{11}{20} = 0.5500, \quad \frac{17}{31} = 0.5484, \quad \frac{23}{42} = 0.5476,$$

$$\frac{1051}{1920} = 0.5474, \quad \frac{6}{11} = 0.5455 \quad \text{and} \quad \frac{7}{13} = 0.5385$$

2020 *Mathematics Subject Classification.* 11N05, 11N35, 11N36.

Key words and phrases. prime, sieve methods, Dirichlet polynomial.

by Iwaniec and Jutila [20], Heath–Brown and Iwaniec [15], Pintz [35] [36], Iwaniec and Pintz [21], Mozzochi [34] and Lou and Yao [30] [31] [32] respectively. In 1996, Baker and Harman [1] presented an alternative approach to this problem. They used the alternative sieve developed by Harman [10] [11] to reduce θ to 0.535. Finally, Baker, Harman and Pintz [2] further developed this sieve process and combined it with Watt’s theorem and showed $\theta = 0.525$.

However, if we only consider the prime numbers in ”almost all” intervals instead of ”all” intervals, the intervals will be much shorter than $n^{0.525}$. In 1943, under the Riemann Hypothesis, Selberg [37] showed that Cramér’s interval contains primes for almost all n if $f(n) \rightarrow \infty$ as $n \rightarrow \infty$. In the same paper, he also showed unconditionally that the interval $[n, n + n^{\frac{19}{77} + \varepsilon}]$ contains prime numbers for almost all n . In 1971, Montgomery [33] improved the exponent $\frac{19}{77}$ to $\frac{1}{5}$ with an asymptotic formula. The zero density estimate of Huxley [18] gives the exponent $\frac{1}{6}$ with an asymptotic formula, and the best asymptotic result now is also due to Guth and Maynard [8], where they proved the exponent $\frac{2}{15}$.

In 1982, Harman [9] used his alternative sieve method to showed that the interval $[n, n + n^{\frac{1}{10} + \varepsilon}]$ contains prime numbers for almost all n . His method can only provide a lower bound instead of an asymptotic formula. The exponent $\frac{1}{10}$ was reduced successively to

$$\begin{aligned} \frac{1}{12} = 0.0833, \quad \frac{1}{13} = 0.0769, \quad \frac{2}{27} = 0.0740, \quad \frac{1}{14} = 0.0714, \\ \frac{1}{15} = 0.0667, \quad \frac{1}{18} = 0.0556 \quad \text{and} \quad \frac{1}{20} = 0.0500 \end{aligned}$$

by Harman [10], Jia [23] [24], Li [28], Jia [22] (and Watt [39]), Li [29], Wong [40] (and Jia [26], Harman [[13], Chapter 9]) and Jia [25] respectively. As the strongest result above, Jia [25] used many powerful tools and arguments in his proof, both analytic and combinatorial, and these are extremely complicated.

In this paper, we obtain the following result.

Theorem 1.1. *The interval $[n, n + n^{\frac{1}{21} + \varepsilon}]$ contains prime numbers for almost all n . Specifically, suppose that B is a sufficiently large positive constant, ε is a sufficiently small positive constant and X is sufficiently large. Then for positive integers $n \in [X, 2X]$, except for $O(X(\log X)^{-B})$ values, the interval $[n, n + n^{\frac{1}{21} + \varepsilon}]$ contains $\gg n^{1/21 + \varepsilon}(\log n)^{-1}$ prime numbers.*

The proof of our Theorem 1.1 is much simpler than Jia’s proof. It is worth mentioning that Harman [[13], Chapter 9] gave a simplified proof with an exponent $\frac{1}{18}$, but this is still weaker than Jia’s result. In fact, Jia’s result requires some ”extra arithmetic information” which will be discussed in Section 4. Harman also used those information in his simplified proof. Without those extra information, Jia could also prove the exponent $\frac{1}{18}$ by the methods in [26].

Clearly our Theorem 1.1 has many interesting applications. The following application of our Theorem 1.1 is about prime values of the integer parts of real sequences, improving the previous result of Harman [12] by adding one more term on both of the sequences $[p^k \alpha]$ and $[(p\alpha)^k]$.

Theorem 1.2. *For almost all $\alpha > 0$, both of the following statements are true:*

- (1) *Infinitely often $p, [p\alpha], [p^2\alpha], \dots, [p^{21}\alpha]$ are all prime.*
- (2) *Infinitely often $p, [p\alpha], [(p\alpha)^2], \dots, [(p\alpha)^{21}]$ are all prime.*

Throughout this paper, we always suppose that B is a sufficiently large positive constant, ε is a sufficiently small positive constant, X is sufficiently large and $x \in [X, 2X]$. The letter p , with or without subscript, is reserved for prime numbers. Let c_0, c_1 and c_2 denote positive constants which may have different values at different places, and we write $m \sim M$ to mean that $c_1 M < m \leq c_2 M$. Let $\varepsilon_1 = \varepsilon^2$, $\delta = \varepsilon^{1/3}$ and $\eta = \frac{1}{2} X^{-\frac{20}{21} + \varepsilon}$. We use $M(s), N(s)$ and some other capital letters to denote the Dirichlet polynomials

$$M(s) = \sum_{m \sim M} a(m)m^{-s}, \quad N(s) = \sum_{n \sim N} b(n)n^{-s}$$

where $a(m), b(n)$ are complex numbers with $a(m) = O(1)$ and $b(n) = O(1)$. We define the boolean function as

$$\text{Boole}[\mathbf{X}] = \begin{cases} 1 & \text{if } \mathbf{X} \text{ is true,} \\ 0 & \text{if } \mathbf{X} \text{ is false.} \end{cases}$$

2. AN OUTLINE OF THE PROOF

Let $p_j = X^{t_j}$ in the following sections and put

$$\mathcal{A} = \{n : x < n \leq x + \eta x\}, \quad \mathcal{B} = \{n : x < n \leq 2x\}, \quad P(z) = \prod_{p < z} p, \quad S(\mathcal{A}, z) = \sum_{\substack{a \in \mathcal{A} \\ (a, P(z))=1}} 1.$$

Then we have

$$\pi(x + \eta x) - \pi(x) = S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right). \quad (1)$$

In order to prove Theorem 1.1, we only need to show that $S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right) > 0$. By Buchstab's identity, we have

$$\begin{aligned} S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right) &= S\left(\mathcal{A}, X^{\frac{13}{133}}\right) - \sum_{\frac{13}{133} \leq t_1 < \frac{1}{2}} S\left(\mathcal{A}_{p_1}, X^{\frac{13}{133}}\right) \\ &\quad + \sum_{\substack{\frac{13}{133} \leq t_1 < \frac{1}{2} \\ \frac{13}{133} \leq t_2 < \min(t_1, \frac{1}{2}(1-t_1))}} S(\mathcal{A}_{p_1 p_2}, p_2) \\ &= S_1 - S_2 + S_3. \end{aligned} \quad (2)$$

Our aim is to show that the sparser set \mathcal{A} contains the expected proportion of primes compared to the bigger set \mathcal{B} , which requires us to decompose $S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right)$ and prove asymptotic formulas (for almost $x \in [X, 2X]$ except for $O(X(\log X)^{-B})$ values) of the form

$$S(\mathcal{A}, z) = \eta(1 + o(1))S(\mathcal{B}, z) \quad (3)$$

for some parts of it, and drop the other positive parts. In Sections 3 and 4 we provide some arithmetic information of Dirichlet polynomials, and we shall use them to prove the asymptotic formulas for type-I terms $S(\mathcal{A}_{p_1 \dots p_n}, X^{c_0})$ in Section 5 and the asymptotic formulas for type-II terms $S(\mathcal{A}_{p_1 \dots p_n}, p_n)$ in Section 6. In Section 7 we will make further use of Buchstab's identity to decompose $S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right)$ and prove Theorem 1.1.

3. ARITHMETIC INFORMATION I

In the following two sections we provide some arithmetic information (i.e. mean value bounds for some Dirichlet polynomials) which will help us prove the asymptotic formulas for sieve functions. In this section we only use the classical mean value estimate and Halász method. In next section we will provide a crucial result which will help us give asymptotic formulas for even more terms.

Lemma 3.1. *Suppose that $MH = X$ where $M(s)$ is a Dirichlet polynomial, $H(s) = \sum_{h \sim H} \Lambda(h)h^{-s}$ and $X^\delta \ll H \ll X^{\frac{13}{133}}$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$, then for $T_0 \leq T \leq X$ we have*

$$\left(\min\left(\eta, \frac{1}{T}\right)\right)^2 \int_T^{2T} |M(b+it)H(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}.$$

Proof. The proof is similar to that of [[25], Lemma 1]. Let $s = b + it$ and by the zero-free region of the ζ function, for $|t| \leq 2X$ we have

$$H(s) = \frac{(c_2 H)^{1-s} - (c_1 H)^{1-s}}{1-s} + O\left((\log x)^{-2B/\varepsilon}\right). \quad (4)$$

So, for $T_0 \leq |t| \leq 2X$ we have $H(s) \ll (\log x)^{-B/\varepsilon}$. According to the discussion in , there are $O((\log X)^2)$ sets $S(V, W)$, where $S(V, W)$ is the set of $t_k (k = 1, \dots, K)$ with the property $|t_r - t_s| \geq 1$ ($r \neq s$). Moreover,

$$V \leq M^{\frac{1}{2}} |M(b + it_k)| < 2V, \quad W \leq H^{\frac{1}{2}} |H(b + it_k)| < 2W,$$

where $X^{-1} \leq M^{-\frac{1}{2}}V, X^{-1} \leq H^{-\frac{1}{2}}W$ and $V \ll M^{\frac{1}{2}}, W \ll H^{\frac{1}{2}}(\log x)^{-B/\varepsilon}$. Then we have

$$\int_T^{2T} |M(b+it)H(b+it)|^2 dt \ll V^2 W^2 x^{-1} (\log x)^2 |S(V, W)| + O(x^{-2\varepsilon_1}), \quad (5)$$

where $S(V, W)$ is one of sets with the above properties.

Assume $X^{\frac{1}{k+1}} \leq H < X^{\frac{1}{k}}$, where k is a positive integer, $k \geq 10$ and $k\delta \ll 1$. Applying the mean value estimate (see Lemma 7 of [23]) to $M(s)$ and $H^k(s)$, we have

$$\begin{aligned} |S(V, W)| &\ll V^{-2}(M+T)(\log x)^d, \\ |S(V, W)| &\ll W^{-2k}(H^k+T)(\log x)^d, \end{aligned}$$

where $d = c/\delta^2$. Applying the Halász method (see Lemma 7 of [23]) to $M(s)$ and $H^k(s)$, we have

$$\begin{aligned} |S(V, W)| &\ll (V^{-2}M + V^{-6}MT)(\log x)^d, \\ |S(V, W)| &\ll (W^{-2k}H^k + W^{-6k}H^kT)(\log x)^d. \end{aligned}$$

Thus,

$$V^2W^2|S(V, W)| \ll V^2W^2F(\log x)^d,$$

where

$$F = \min \{V^{-2}(M+T), V^{-2}M + V^{-6}MT, W^{-2k}(H^k+T), W^{-2k}H^k + W^{-6k}H^kT\}.$$

It will be proved that

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F \ll \eta^2 x(\log x)^{-B/\varepsilon}. \quad (6)$$

We consider four cases.

(a) $F \leq 2V^{-2}M, F \leq 2W^{-2k}H^k$. Then

$$\begin{aligned} V^2W^2F &\ll V^2W^2 \min \{V^{-2}M, W^{-2k}H^k\} \\ &\leq V^2W^2 (V^{-2}M)^{1-\frac{1}{2k}} (W^{-2k}H^k)^{\frac{1}{2k}} \\ &= V^{\frac{1}{k}}WM^{1-\frac{1}{2k}}H^{\frac{1}{2}} \\ &\ll x(\log x)^{-B/\varepsilon} \end{aligned}$$

and so

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F \ll \eta^2 x(\log x)^{-B/\varepsilon}.$$

(b) $F > 2V^{-2}M, F > 2W^{-2k}H^k$. Then

$$\begin{aligned} V^2W^2F &\ll V^2W^2 \min \{V^{-2}T, V^{-6}MT, W^{-2k}T, W^{-6k}H^kT\} \\ &\leq V^2W^2 (V^{-2})^{1-\frac{3}{2k}} (V^{-6}M)^{\frac{1}{2k}} (W^{-2k})^{\frac{1}{k}} T \\ &= M^{\frac{1}{2k}}T. \end{aligned}$$

Since $k \geq 10$, we have $H \geq X^{\frac{1}{k+1}} \geq X^{1-\frac{k}{11}} \geq X^{\frac{1}{22}}$, $M^{\frac{1}{2k}} \ll X^{\frac{1}{22}}$, and so

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F \ll \frac{\eta}{T} x^{\frac{1}{22}} T \ll \eta^2 x^{1-\varepsilon_1}.$$

(c) $F \leq 2V^{-2}M, F > 2W^{-2k}H^k$. Then

$$\begin{aligned} V^2W^2F &\ll V^2W^2 \min \{V^{-2}M, W^{-2k}T, W^{-6k}H^kT\} \\ &\leq V^2W^2 (V^{-2}M)^{1-\frac{1}{3k}} (W^{-6k}H^kT)^{\frac{1}{3k}} \\ &\ll MH^{\frac{1}{3}}T^{\frac{1}{3k}}, \end{aligned}$$

since $V \ll M^{\frac{1}{2}}$. As $H \geq X^{\frac{1}{k+1}} \geq X^{\frac{21}{44k}+\varepsilon}$ and $M \leq X^{1-\frac{21}{44k}}$, we have

$$\begin{aligned} \left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F &\ll \eta^{2-\frac{1}{3k}} \left(\frac{1}{T} \right)^{\frac{1}{3k}} (MH)^{\frac{1}{3}} M^{\frac{2}{3}} T^{\frac{1}{3k}} \\ &\ll x^{\frac{1}{3}-\frac{20}{21}(2-\frac{1}{3k})+\frac{2}{3}(1-\frac{21}{44k})} \\ &\ll \eta^2 x^{1-\varepsilon_1}. \end{aligned}$$

(d) $F > 2V^{-2}M, F \leq 2W^{-2k}H^k$. Then

$$\begin{aligned} V^2W^2F &\ll V^2W^2 \min \{V^{-2}T, V^{-6}MT, W^{-2k}H^k\} \\ &\leq V^2W^2 (V^{-2}T)^{1-\frac{3}{2k}} (V^{-6}MT)^{\frac{1}{2k}} (W^{-2k}H^k)^{\frac{1}{k}} \\ &= M^{\frac{1}{2k}}HT^{1-\frac{1}{k}}. \end{aligned}$$

If $k \geq 11$, then $H \leq X^{\frac{1}{k}} \leq X^{1-\frac{21(k-1)}{11(2k-1)}}$, $M \gg X^{\frac{21(k-1)}{11(2k-1)}}$, and so

$$\begin{aligned} \left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F &\ll \eta^{1+\frac{1}{k}} \left(\frac{1}{T} \right)^{1-\frac{1}{k}} (MH) M^{-\frac{2k-1}{2k}} T^{1-\frac{1}{k}} \\ &\ll x^{1-\frac{20}{21}(1+\frac{1}{k})-\frac{21(k-1)}{22k}} \\ &\ll \eta^2 x^{1-\varepsilon_1}. \end{aligned}$$

If $k = 10$, then $X^{\frac{1}{11}} \leq H \ll X^{\frac{13}{133}}$, $M \gg X^{\frac{120}{133}}$, and so

$$\begin{aligned} \left(\min \left(\eta, \frac{1}{T} \right) \right)^2 V^2W^2F &\ll \eta^{\frac{11}{10}} \left(\frac{1}{T} \right)^{\frac{9}{10}} (MH) M^{-\frac{19}{20}} T^{\frac{9}{10}} \\ &\ll x^{1-\frac{11}{10} \cdot \frac{20}{21} - \frac{19}{20} \cdot \frac{120}{133}} \\ &\ll \eta^2 x^{1-\varepsilon_1}. \end{aligned}$$

Combining the above cases, Lemma 3.1 is proved. \square

Lemma 3.2. *Suppose that $MHK = X$ where $M(s), H(s)$ and $K(s)$ are Dirichlet polynomials and $G(s) = M(s)H(s)K(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$. Assume further that for $T_0 \leq |t| \leq 2X$, $M(b+it) \ll (\log x)^{-B/\varepsilon}$ and $H(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that M and H satisfy one of the following 9 conditions:*

- (1) $MH \ll X^{\frac{47}{87}}$, $X^{\frac{40}{231}} \ll H$, $M^{29}H^{-1} \ll X^{\frac{209}{21}}$, $X^{\frac{40}{133}} \ll M$, $M^{-1}H^{29} \ll X^{\frac{43}{7}}$, $X^{\frac{40}{7}} \ll M^{12}H^{11}$;
- (2) $MH \ll X^{\frac{109}{189}}$, $M^{29}H^{19} \ll X^{\frac{103}{7}}$, $X^{\frac{160}{189}} \ll M^2H$, $M^2H^{11} \ll X^{\frac{20}{7}}$, $X^{\frac{200}{7}} \ll M^{58}H^{49}$;
- (3) $MH \ll X^{\frac{131}{231}}$, $X^{\frac{4}{21}} \ll H$, $M^6H \ll X^{\frac{46}{21}}$, $X^{\frac{40}{147}} \ll M$, $MH^8 \ll X^{\frac{16}{7}}$, $X^{\frac{20}{7}} \ll M^6H^5$;
- (4) $MH \ll X^{\frac{93}{133}}$, $X^{\frac{40}{231}} \ll H$, $M^{12}H \ll X^{\frac{44}{7}}$, $X^{\frac{40}{87}} \ll M$, $M^{-1}H^{19} \ll X^{\frac{79}{21}}$, $X^{\frac{160}{21}} \ll M^{12}H^{11}$;
- (5) $M^2H \ll X^{\frac{218}{189}}$, $M^{58}H^9 \ll X^{\frac{206}{7}}$, $X^{\frac{80}{189}} \ll M$, $MH^5 \ll X^{\frac{10}{7}}$, $X^{\frac{100}{7}} \ll M^{29}H^{10}$;
- (6) $MH \ll X^{\frac{107}{147}}$, $X^{\frac{4}{21}} \ll H$, $M^6H \ll X^{\frac{22}{7}}$, $X^{\frac{100}{231}} \ll M$, $M^{-1}H^7 \ll X^{\frac{9}{7}}$, $X^{\frac{80}{21}} \ll M^6H^5$;
- (7) $MH \ll X^{\frac{131}{231}}$, $M^{35}H^{23} \ll X^{\frac{125}{7}}$, $X^{\frac{200}{231}} \ll M^2H$, $M^2H^{13} \ll X^{\frac{64}{21}}$, $X^{\frac{240}{7}} \ll M^{70}H^{59}$;
- (8) $MH \ll X^{\frac{51}{91}}$, $M^{41}H^{27} \ll X^{21}$, $X^{\frac{80}{91}} \ll M^2H$, $M^2H^{15} \ll X^{\frac{68}{21}}$, $X^{40} \ll M^{82}H^{69}$;
- (9) $M^2H \ll X^{\frac{262}{231}}$, $M^{70}H^{11} \ll X^{\frac{250}{7}}$, $X^{\frac{100}{231}} \ll M$, $MH^6 \ll X^{\frac{32}{21}}$, $X^{\frac{120}{7}} \ll M^{35}H^{12}$.

Then for $T_0 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (7)$$

Proof. The proof is similar to that of [[25], Lemmas 5,7,9,10,12]. Using the method of Lemma 3.1, we only need to show that for $T = 1/\eta = 2X^{\frac{20}{21}-\varepsilon}$,

$$I = \int_T^{2T} |G(b+it)|^2 dt \ll (\log x)^{-10B}. \quad (8)$$

Assuming the condition (1). By applying the mean value estimate and Halász method to $M^3(s), H^5(s)$ and $K^2(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$\begin{aligned} F = \min \{ &V^{-6}(M^3+T), V^{-6}M^3+V^{-18}M^3T, W^{-10}(H^5+T), \\ &W^{-10}H^5+W^{-30}H^5T, U^{-4}(K^2+T), U^{-4}K^2+U^{-12}K^2T \}. \end{aligned}$$

We consider 8 cases:

(a) $F \leq 2V^{-6}M^3, F \leq 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-10}H^5, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{3}{8}} (W^{-10}H^5)^{\frac{1}{8}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= V^{\frac{1}{2}}M^{\frac{3}{4}}HK \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(b) $F \leq 2V^{-6}M^3, F \leq 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-10}H^5, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-4}T)^{\frac{9}{20}} (U^{-12}K^2T)^{\frac{1}{60}} \\ &= T^{\frac{7}{15}}MHK^{\frac{1}{30}} \\ &= T^{\frac{7}{15}}(MHK)^{\frac{1}{30}} \left(M^{\frac{29}{30}}H^{\frac{29}{30}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $MH \ll X^{\frac{47}{87}}$ is required.

(c) $F \leq 2V^{-6}M^3, F > 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-10}T, W^{-30}H^5T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-10}T)^{\frac{3}{20}} (W^{-30}H^5T)^{\frac{1}{60}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{1}{6}}MH^{\frac{1}{12}}K \\ &= T^{\frac{1}{6}}(MHK)H^{-\frac{11}{12}} \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{40}{231}} \ll H$ is required.

(d) $F \leq 2V^{-6}M^3, F > 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-10}T, W^{-30}H^5T, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-10}T)^{\frac{1}{5}} (U^{-4}T)^{\frac{9}{20}} (U^{-12}K^2T)^{\frac{1}{60}} \\ &= T^{\frac{2}{3}}MK^{\frac{1}{30}} \\ &= T^{\frac{2}{3}}(MHK)^{\frac{1}{30}} \left(M^{\frac{29}{30}}H^{-\frac{1}{30}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^{29}H^{-1} \ll X^{\frac{209}{21}}$ is required.

(e) $F > 2V^{-6}M^3, F \leq 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-10}H^5, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-6}T)^{\frac{17}{60}} (V^{-18}M^3T)^{\frac{1}{60}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{3}{10}}M^{\frac{1}{20}}HK \\ &= T^{\frac{3}{10}}(MHK)M^{-\frac{19}{20}} \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{40}{133}} \ll M$ is required.

(f) $F > 2V^{-6}M^3, F \leq 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-10}H^5, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-6}T)^{\frac{1}{3}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-4}T)^{\frac{9}{20}} (U^{-12}K^2T)^{\frac{1}{60}} \end{aligned}$$

$$\begin{aligned}
&= T^{\frac{4}{5}} H K^{\frac{1}{30}} \\
&= T^{\frac{4}{5}} (MHK)^{\frac{1}{30}} \left(M^{-\frac{1}{30}} H^{\frac{29}{30}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{-1}H^{29} \ll X^{\frac{43}{7}}$ is required.

(g) $F > 2V^{-6}M^3, F > 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-10}T, W^{-30}H^5T, U^{-4}K^2\} \\
&\leq U^2V^2W^2 (V^{-6}T)^{\frac{1}{3}} (W^{-10}T)^{\frac{3}{20}} (W^{-30}H^5T)^{\frac{1}{60}} (U^{-4}K^2)^{\frac{1}{2}} \\
&= T^{\frac{1}{2}} H^{\frac{1}{12}} K \\
&= T^{\frac{1}{2}} (MHK) \left(M^{-1}H^{-\frac{11}{12}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{40}{7}} \ll M^{12}H^{11}$ is required.

(h) $F > 2V^{-6}M^3, F > 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}, V^{-18}M^3, W^{-10}, W^{-30}H^5, U^{-4}, U^{-12}K^2\} T \\
&\leq U^2V^2W^2 (V^{-6})^{\frac{170}{60}} (V^{-18}M^3)^{\frac{1}{60}} (W^{-10})^{\frac{1}{5}} (U^{-4})^{\frac{1}{2}} T \\
&= TM^{\frac{1}{20}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M \ll X^{\frac{2459}{6699}}$ and $TM^{\frac{1}{20}} \ll X^{\frac{20}{21} + \frac{1}{20} \frac{2459}{6699}} \ll X^{\frac{43353}{44660}}$ are used. The former one can be obtained by using $MH \ll X^{\frac{47}{87}}$ and $X^{\frac{40}{231}} \ll H$.

Assuming the condition (2). By applying the mean value estimate and Halász method to $M^2(s)H(s), H^5(s)$ and $K^2(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$\begin{aligned}
F = \min \{ &V^{-4}W^{-2} (M^2H + T), V^{-4}W^{-2}M^2H + V^{-12}W^{-6}M^2HT, \\
&W^{-10} (H^5 + T), W^{-10}H^5 + W^{-30}H^5T, U^{-4} (K^2 + T), U^{-4}K^2 + U^{-12}K^2T \}.
\end{aligned}$$

We consider 8 cases:

(a) $F \leq 2V^{-4}W^{-2}M^2H, F \leq 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-10}H^5, U^{-4}K^2\} \\
&\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{3}{8}} (W^{-10}H^5)^{\frac{1}{8}} (U^{-4}K^2)^{\frac{1}{2}} \\
&= V^{\frac{1}{2}} M^{\frac{3}{4}} H K \\
&\ll x(\log x)^{-11B}.
\end{aligned}$$

(b) $F \leq 2V^{-4}W^{-2}M^2H, F \leq 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-10}H^5, U^{-4}T, U^{-12}K^2T\} \\
&\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}T)^{\frac{7}{20}} (U^{-12}K^2T)^{\frac{1}{20}} \\
&= T^{\frac{2}{5}} MHK^{\frac{1}{10}} \\
&= T^{\frac{2}{5}} (MHK)^{\frac{1}{10}} \left(M^{\frac{9}{10}} H^{\frac{9}{10}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH \ll X^{\frac{109}{189}}$ is required.

(c) $F \leq 2V^{-4}W^{-2}M^2H, F > 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-10}T, W^{-30}H^5T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(d) $F \leq 2V^{-4}W^{-2}M^2H, F > 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-10}T, W^{-30}H^5T, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (W^{-30}H^5T)^{\frac{1}{30}} (U^{-4}T)^{\frac{9}{20}} (U^{-12}K^2T)^{\frac{1}{10}} \\ &= T^{\frac{1}{2}}MH^{\frac{2}{3}}K^{\frac{1}{30}} \\ &= T^{\frac{1}{2}}(MHK)^{\frac{1}{30}} \left(M^{\frac{29}{30}}H^{\frac{19}{30}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^{29}H^{19} \ll X^{\frac{103}{189}}$ is required.

(e) $F > 2V^{-4}W^{-2}M^2H, F \leq 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-10}H^5, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{7}{20}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{20}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{2}{5}}M^{\frac{1}{10}}H^{\frac{11}{20}}K \\ &= T^{\frac{2}{5}}(MHK) \left(M^{-\frac{9}{10}}H^{-\frac{9}{20}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{160}{189}} \ll M^2H$ is required.

(f) $F > 2V^{-4}W^{-2}M^2H, F \leq 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-10}H^5, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{7}{20}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{20}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}T)^{\frac{1}{2}} \\ &= T^{\frac{9}{10}} \left(M^{\frac{1}{10}}H^{\frac{11}{20}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^2H^{11} \ll X^{\frac{20}{7}}$ is required.

(g) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-10}H^5, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-10}T, W^{-30}H^5T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{9}{20}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{60}} (W^{-30}H^5T)^{\frac{1}{30}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{1}{2}}M^{\frac{1}{30}}H^{\frac{11}{60}}K \\ &= T^{\frac{1}{2}}(MHK) \left(M^{-\frac{29}{30}}H^{-\frac{49}{60}}\right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{200}{7}} \ll M^{58}H^{49}$ is required.

(h) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-10}H^5, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}, V^{-12}W^{-6}M^2H, W^{-10}, W^{-30}H^5, U^{-4}, U^{-12}K^2\} T \\ &\leq U^2V^2W^2 (V^{-4}W^{-2})^{\frac{9}{20}} (V^{-12}W^{-6}M^2H)^{\frac{1}{60}} (W^{-30}H^5)^{\frac{1}{30}} (U^{-4})^{\frac{1}{2}} T \\ &= T \left(M^{\frac{1}{30}}H^{\frac{11}{60}}\right) \end{aligned}$$

$$\ll x^{1-\varepsilon_1},$$

where $M^2H^{11} \ll X^{\frac{20}{7}}$ is required.

Assuming the condition (3). By applying the mean value estimate and Halász method to $M^3(s), H^4(s)$ and $K^2(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$F = \min \{V^{-6}(M^3 + T), V^{-6}M^3 + V^{-18}M^3T, W^{-8}(H^4 + T), \\ W^{-8}H^4 + W^{-24}H^4T, U^{-4}(K^2 + T), U^{-4}K^2 + U^{-12}K^2T\}.$$

We consider 8 cases:

(a) $F \leq 2V^{-6}M^3, F \leq 2W^{-8}H^4, F \leq 2U^{-4}K^2$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-8}H^4, U^{-4}K^2\} \\ \leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{4}} (W^{-8}H^4)^{\frac{1}{4}} (U^{-4}K^2)^{\frac{1}{2}} \\ = V^{\frac{1}{2}}M^{\frac{3}{4}}HK \\ \ll x(\log x)^{-11B}.$$

(b) $F \leq 2V^{-6}M^3, F \leq 2W^{-8}H^4, F > 2U^{-4}K^2$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-8}H^4, U^{-4}T, U^{-12}K^2T\} \\ \leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-8}H^4)^{\frac{1}{4}} (U^{-4}T)^{\frac{3}{8}} (U^{-12}K^2T)^{\frac{1}{24}} \\ = T^{\frac{5}{12}}MHK^{\frac{1}{12}} \\ = T^{\frac{5}{12}}(MHK)^{\frac{1}{12}} \left(M^{\frac{11}{12}}H^{\frac{11}{12}}\right) \\ \ll x^{1-\varepsilon_1},$$

where $MH \ll X^{\frac{131}{231}}$ is required.

(c) $F \leq 2V^{-6}M^3, F > 2W^{-8}H^4, F \leq 2U^{-4}K^2$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-8}T, W^{-24}H^4T, U^{-4}K^2\} \\ \leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-8}T)^{\frac{1}{8}} (W^{-24}H^4T)^{\frac{1}{24}} (U^{-4}K^2)^{\frac{1}{2}} \\ = T^{\frac{1}{6}}MH^{\frac{1}{6}}K \\ = T^{\frac{1}{6}}(MHK)H^{-\frac{5}{6}} \\ \ll x^{1-\varepsilon_1},$$

where $X^{\frac{4}{21}} \ll H$ is required.

(d) $F \leq 2V^{-6}M^3, F > 2W^{-8}H^4, F > 2U^{-4}K^2$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-6}M^3, W^{-8}T, W^{-24}H^4T, U^{-4}T, U^{-12}K^2T\} \\ \leq U^2V^2W^2 (V^{-6}M^3)^{\frac{1}{3}} (W^{-8}T)^{\frac{1}{8}} (W^{-24}H^4T)^{\frac{1}{24}} (U^{-4}T)^{\frac{1}{2}} \\ = T^{\frac{2}{3}} \left(MH^{\frac{1}{6}}\right) \\ \ll x^{1-\varepsilon_1},$$

where $M^6H \ll X^{\frac{46}{21}}$ is required.

(e) $F > 2V^{-6}M^3, F \leq 2W^{-8}H^4, F \leq 2U^{-4}K^2$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-8}H^4, U^{-4}K^2\} \\ \leq U^2V^2W^2 (V^{-6}T)^{\frac{5}{24}} (V^{-18}M^3T)^{\frac{1}{24}} (W^{-8}H^4)^{\frac{1}{4}} (U^{-4}K^2)^{\frac{1}{2}} \\ = T^{\frac{1}{4}}M^{\frac{1}{8}}HK$$

$$\begin{aligned}
&= T^{\frac{1}{4}}(MHK)M^{-\frac{7}{8}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{40}{147}} \ll M$ is required.

(f) $F > 2V^{-6}M^3, F \leq 2W^{-8}H^4, F > 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-8}H^4, U^{-4}T, U^{-12}K^2T\} \\
&\leq U^2V^2W^2 (V^{-6}T)^{\frac{5}{24}} (V^{-18}M^3T)^{\frac{1}{24}} (W^{-8}H^4)^{\frac{1}{4}} (U^{-4}T)^{\frac{1}{2}} \\
&= T^{\frac{3}{4}} \left(M^{\frac{1}{8}} H \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH^8 \ll X^{\frac{16}{7}}$ is required.

(g) $F > 2V^{-6}M^3, F > 2W^{-8}H^4, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}T, V^{-18}M^3T, W^{-8}T, W^{-24}H^4T, U^{-4}K^2\} \\
&\leq U^2V^2W^2 (V^{-6}T)^{\frac{1}{3}} (W^{-8}T)^{\frac{1}{8}} (W^{-24}H^4T)^{\frac{1}{24}} (U^{-4}K^2)^{\frac{1}{2}} \\
&= T^{\frac{1}{2}} H^{\frac{1}{6}} K \\
&= T^{\frac{1}{2}}(MHK) \left(M^{-1}H^{-\frac{5}{6}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{20}{7}} \ll M^6H^5$ is required.

(h) $F > 2V^{-6}M^3, F > 2W^{-8}H^4, F > 2U^{-4}K^2$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-6}, V^{-18}M^3, W^{-8}, W^{-24}H^4, U^{-4}, U^{-12}K^2\} T \\
&\leq U^2V^2W^2 (V^{-6})^{\frac{5}{24}} (V^{-18}M^3)^{\frac{1}{24}} (W^{-8})^{\frac{1}{4}} (U^{-4})^{\frac{1}{2}} T \\
&= TM^{\frac{1}{8}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M \ll X^{\frac{29}{77}}$ and $TM^{\frac{1}{8}} \ll X^{\frac{20}{21} + \frac{1}{8} \cdot \frac{29}{77}} \ll X^{\frac{1847}{1848}}$ are used. The former one can be obtained by using $MH \ll X^{\frac{131}{231}}$ and $X^{\frac{4}{21}} \ll H$.

Assuming the condition (4). By applying the mean value estimate and Halász method to $M^2(s), H^5(s)$ and $K^3(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$\begin{aligned}
F &= \min \{V^{-4}(M^2 + T), V^{-4}M^2 + V^{-12}M^2T, W^{-10}(H^5 + T), \\
&\quad W^{-10}H^5 + W^{-30}H^5T, U^{-6}(K^3 + T), U^{-6}K^3 + U^{-18}K^3T\}.
\end{aligned}$$

We consider 8 cases:

(a) $F \leq 2V^{-4}M^2, F \leq 2W^{-10}H^5, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}H^5, U^{-6}K^3\} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-10}H^5)^{\frac{1}{6}} (U^{-6}K^3)^{\frac{1}{3}} \\
&= W^{\frac{1}{3}}MH^{\frac{5}{6}}K \\
&\ll x(\log x)^{-11B}.
\end{aligned}$$

(b) $F \leq 2V^{-4}M^2, F \leq 2W^{-10}H^5, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}H^5, U^{-6}T, U^{-18}K^3T\} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-6}T)^{\frac{17}{60}} (U^{-18}K^3T)^{\frac{1}{60}}
\end{aligned}$$

$$\begin{aligned}
&= T^{\frac{3}{10}} M H K^{\frac{1}{20}} \\
&= T^{\frac{3}{10}} (M H K)^{\frac{1}{20}} \left(M^{\frac{19}{20}} H^{\frac{19}{20}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M H \ll X^{\frac{93}{133}}$ is required.

(c) $F \leq 2V^{-4}M^2, F > 2W^{-10}H^5, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}T, W^{-30}H^5T, U^{-6}K^3\} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-10}T)^{\frac{3}{20}} (W^{-30}H^5T)^{\frac{1}{60}} (U^{-6}K^3)^{\frac{1}{3}} \\
&= T^{\frac{1}{6}} M H^{\frac{1}{12}} K \\
&= T^{\frac{1}{6}} (M H K) H^{-\frac{11}{12}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{40}{231}} \ll H$ is required.

(d) $F \leq 2V^{-4}M^2, F > 2W^{-10}H^5, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}T, W^{-30}H^5T, U^{-6}T, U^{-18}K^3T\} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-10}T)^{\frac{3}{20}} (W^{-30}H^5T)^{\frac{1}{60}} (U^{-6}T)^{\frac{1}{3}} \\
&= T^{\frac{1}{2}} \left(M H^{\frac{1}{12}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{12}H \ll X^{\frac{44}{7}}$ is required.

(e) $F > 2V^{-4}M^2, F \leq 2W^{-10}H^5, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}H^5, U^{-6}K^3\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{9}{20}} (V^{-12}M^2T)^{\frac{1}{60}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-6}K^3)^{\frac{1}{3}} \\
&= T^{\frac{7}{15}} M^{\frac{1}{30}} H K \\
&= T^{\frac{7}{15}} (M H K) M^{-\frac{29}{30}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{40}{87}} \ll M$ is required.

(f) $F > 2V^{-4}M^2, F \leq 2W^{-10}H^5, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}H^5, U^{-6}T, U^{-18}K^3T\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{1}{2}} (W^{-10}H^5)^{\frac{1}{5}} (U^{-6}T)^{\frac{17}{60}} (U^{-18}K^3T)^{\frac{1}{60}} \\
&= T^{\frac{4}{5}} H K^{\frac{1}{20}} \\
&= T^{\frac{4}{5}} (M H K)^{\frac{1}{20}} \left(M^{-\frac{1}{20}} H^{\frac{19}{20}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{-1}H^{19} \ll X^{\frac{79}{21}}$ is required.

(g) $F > 2V^{-4}M^2, F > 2W^{-10}H^5, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}T, W^{-30}H^5T, U^{-6}K^3\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{1}{2}} (W^{-10}T)^{\frac{3}{20}} (W^{-30}H^5T)^{\frac{1}{60}} (U^{-6}K^3)^{\frac{1}{3}} \\
&= T^{\frac{2}{3}} H^{\frac{1}{12}} K \\
&= T^{\frac{2}{3}} (M H K) \left(M^{-1} H^{-\frac{11}{12}} \right)
\end{aligned}$$

$$\ll x^{1-\varepsilon_1},$$

where $X^{\frac{160}{21}} \ll M^{12}H^{11}$ is required.

(h) $F > 2V^{-4}M^2, F > 2W^{-10}H^5, F > 2U^{-6}K^3$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}, V^{-12}M^2, W^{-10}, W^{-30}H^5, U^{-6}, U^{-18}K^3\} T \\ &\leq U^2V^2W^2 (V^{-4})^{\frac{1}{2}} (W^{-10})^{\frac{1}{5}} (U^{-6})^{\frac{17}{60}} (U^{-18}K^3)^{\frac{1}{60}} T \\ &= TK^{\frac{1}{20}} \\ &= T(MHK)^{\frac{1}{20}} \left(M^{-\frac{1}{20}} H^{-\frac{1}{20}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{1}{21}} \ll MH$ is required. This can be obtained by using $X^{\frac{40}{87}} \ll M$ and $X^{\frac{40}{231}} \ll H$.

Assuming the condition (5). By applying the mean value estimate and Halász method to $M^2(s), H^5(s)$ and $K^2(s)H(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$\begin{aligned} F = \min \{ &V^{-4}(M^2 + T), V^{-4}M^2 + V^{-12}M^2T, W^{-10}(H^5 + T), \\ &W^{-10}H^5 + W^{-30}H^5T, U^{-4}W^{-2}(K^2H + T), U^{-4}W^{-2}K^2H + U^{-12}W^{-6}K^2HT \}. \end{aligned}$$

We consider 8 cases:

(a) $F \leq 2V^{-4}M^2, F \leq 2W^{-10}H^5, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}H^5, U^{-4}W^{-2}K^2H\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(b) $F \leq 2V^{-4}M^2, F \leq 2W^{-10}H^5, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}H^5, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}W^{-2}T)^{\frac{7}{20}} (U^{-12}W^{-6}K^2HT)^{\frac{1}{20}} \\ &= T^{\frac{2}{5}}MH^{\frac{11}{20}}K^{\frac{1}{10}} \\ &= T^{\frac{2}{5}}(MHK)^{\frac{1}{10}} \left(M^{\frac{9}{10}}H^{\frac{9}{20}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^2H \ll X^{\frac{218}{189}}$ is required.

(c) $F \leq 2V^{-4}M^2, F > 2W^{-10}H^5, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}T, W^{-30}H^5T, U^{-4}W^{-2}K^2H\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(d) $F \leq 2V^{-4}M^2, F > 2W^{-10}H^5, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-10}T, W^{-30}H^5T, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-30}H^5T)^{\frac{1}{30}} (U^{-4}W^{-2}T)^{\frac{9}{20}} (U^{-12}W^{-6}K^2HT)^{\frac{1}{60}} \\ &= T^{\frac{1}{2}}MH^{\frac{11}{60}}K^{\frac{1}{30}} \end{aligned}$$

$$\begin{aligned}
&= T^{\frac{1}{2}}(MHK)^{\frac{1}{30}} \left(M^{\frac{29}{30}} H^{\frac{9}{60}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{58}H^9 \ll X^{\frac{206}{7}}$ is required.

(e) $F > 2V^{-4}M^2, F \leq 2W^{-10}H^5, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}H^5, U^{-4}W^{-2}K^2H\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{7}{20}} (V^{-12}M^2T)^{\frac{1}{20}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\
&= T^{\frac{2}{5}} M^{\frac{1}{10}} HK \\
&= T^{\frac{2}{5}}(MHK)M^{-\frac{9}{10}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{80}{189}} \ll M$ is required.

(f) $F > 2V^{-4}M^2, F \leq 2W^{-10}H^5, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}H^5, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{7}{20}} (V^{-12}M^2T)^{\frac{1}{20}} (W^{-10}H^5)^{\frac{1}{10}} (U^{-4}W^{-2}T)^{\frac{1}{2}} \\
&= T^{\frac{9}{10}} \left(M^{\frac{1}{10}} H^{\frac{1}{2}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH^5 \ll X^{\frac{10}{7}}$ is required.

(g) $F > 2V^{-4}M^2, F > 2W^{-10}H^5, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-10}T, W^{-30}H^5T, U^{-4}W^{-2}K^2H\} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{9}{20}} (V^{-12}M^2T)^{\frac{1}{60}} (W^{-30}H^5T)^{\frac{1}{30}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\
&= T^{\frac{1}{2}} M^{\frac{1}{30}} H^{\frac{2}{3}} K \\
&= T^{\frac{1}{2}}(MHK) \left(M^{-\frac{29}{30}} H^{-\frac{1}{3}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{100}{7}} \ll M^{29}H^{10}$ is required.

(h) $F > 2V^{-4}M^2, F > 2W^{-10}H^5, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}, V^{-12}M^2, W^{-10}, W^{-30}H^5, U^{-4}W^{-2}, U^{-12}W^{-6}K^2H\} T \\
&\leq U^2V^2W^2 (V^{-4})^{\frac{9}{20}} (V^{-12}M^2)^{\frac{1}{60}} (W^{-30}H^5)^{\frac{1}{30}} (U^{-4}W^{-2})^{\frac{1}{2}} T \\
&= T \left(M^{\frac{1}{30}} H^{\frac{1}{6}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH^5 \ll X^{\frac{10}{7}}$ is required.

Assuming the condition (6). By applying the mean value estimate and Halász method to $M^2(s), H^4(s)$ and $K^3(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$\begin{aligned}
F &= \min \{V^{-4}(M^2 + T), V^{-4}M^2 + V^{-12}M^2T, W^{-8}(H^4 + T), \\
&\quad W^{-8}H^4 + W^{-24}H^4T, U^{-6}(K^3 + T), U^{-6}K^3 + U^{-18}K^3T\}.
\end{aligned}$$

We consider 8 cases:

(a) $F \leq 2V^{-4}M^2, F \leq 2W^{-8}H^4, F \leq 2U^{-6}K^3$. Then

$$U^2V^2W^2F \ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-8}H^4, U^{-6}K^3\}$$

$$\begin{aligned}
&\leq U^2 V^2 W^2 (V^{-4} M^2)^{\frac{1}{2}} (W^{-8} H^4)^{\frac{1}{6}} (U^{-6} K^3)^{\frac{1}{3}} \\
&= W^{\frac{2}{3}} M H^{\frac{2}{3}} K \\
&\ll x(\log x)^{-11B}.
\end{aligned}$$

(b) $F \leq 2V^{-4}M^2, F \leq 2W^{-8}H^4, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{V^{-4} M^2, W^{-8} H^4, U^{-6} T, U^{-18} K^3 T\} \\
&\leq U^2 V^2 W^2 (V^{-4} M^2)^{\frac{1}{2}} (W^{-8} H^4)^{\frac{1}{4}} (U^{-6} T)^{\frac{5}{24}} (U^{-18} K^3 T)^{\frac{1}{24}} \\
&= T^{\frac{1}{4}} M H K^{\frac{1}{8}} \\
&= T^{\frac{1}{4}} (M H K)^{\frac{1}{8}} \left(M^{\frac{7}{8}} H^{\frac{7}{8}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH \ll X^{\frac{107}{147}}$ is required.

(c) $F \leq 2V^{-4}M^2, F > 2W^{-8}H^4, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{V^{-4} M^2, W^{-8} T, W^{-24} H^4 T, U^{-6} K^3\} \\
&\leq U^2 V^2 W^2 (V^{-4} M^2)^{\frac{1}{2}} (W^{-8} T)^{\frac{1}{8}} (W^{-24} H^4 T)^{\frac{1}{24}} (U^{-6} K^3)^{\frac{1}{3}} \\
&= T^{\frac{1}{6}} M H^{\frac{1}{6}} K \\
&= T^{\frac{1}{6}} (M H K) H^{-\frac{5}{6}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{4}{21}} \ll H$ is required.

(d) $F \leq 2V^{-4}M^2, F > 2W^{-8}H^4, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{V^{-4} M^2, W^{-8} T, W^{-24} H^4 T, U^{-6} T, U^{-18} K^3 T\} \\
&\leq U^2 V^2 W^2 (V^{-4} M^2)^{\frac{1}{2}} (W^{-8} T)^{\frac{1}{8}} (W^{-24} H^4 T)^{\frac{1}{24}} (U^{-6} T)^{\frac{1}{3}} \\
&= T^{\frac{1}{2}} \left(M H^{\frac{1}{6}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^6 H \ll X^{\frac{22}{7}}$ is required.

(e) $F > 2V^{-4}M^2, F \leq 2W^{-8}H^4, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{V^{-4} T, V^{-12} M^2 T, W^{-8} H^4, U^{-6} K^3\} \\
&\leq U^2 V^2 W^2 (V^{-4} T)^{\frac{3}{8}} (V^{-12} M^2 T)^{\frac{1}{24}} (W^{-8} H^4)^{\frac{1}{4}} (U^{-6} K^3)^{\frac{1}{3}} \\
&= T^{\frac{5}{12}} M^{\frac{1}{12}} H K \\
&= T^{\frac{5}{12}} (M H K) M^{-\frac{11}{12}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{100}{231}} \ll M$ is required.

(f) $F > 2V^{-4}M^2, F \leq 2W^{-8}H^4, F > 2U^{-6}K^3$. Then

$$\begin{aligned}
U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{V^{-4} T, V^{-12} M^2 T, W^{-8} H^4, U^{-6} T, U^{-18} K^3 T\} \\
&\leq U^2 V^2 W^2 (V^{-4} T)^{\frac{1}{2}} (W^{-8} H^4)^{\frac{1}{4}} (U^{-6} T)^{\frac{5}{24}} (U^{-18} K^3 T)^{\frac{1}{24}} \\
&= T^{\frac{3}{4}} H K^{\frac{1}{8}} \\
&= T^{\frac{3}{4}} (M H K)^{\frac{1}{8}} \left(M^{-\frac{1}{8}} H^{\frac{7}{8}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{-1}H^7 \ll X^{\frac{9}{7}}$ is required.

(g) $F > 2V^{-4}M^2, F > 2W^{-8}H^4, F \leq 2U^{-6}K^3$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}T, V^{-12}M^2T, W^{-8}T, W^{-24}H^4T, U^{-6}K^3\} \\ &\leq U^2V^2W^2 (V^{-4}T)^{\frac{1}{2}} (W^{-8}T)^{\frac{1}{8}} (W^{-24}H^4T)^{\frac{1}{24}} (U^{-6}K^3)^{\frac{1}{3}} \\ &= T^{\frac{2}{3}} H^{\frac{1}{6}} K \\ &= T^{\frac{2}{3}} (MHK) \left(M^{-1}H^{-\frac{5}{6}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{80}{21}} \ll M^6H^5$ is required.

(h) $F > 2V^{-4}M^2, F > 2W^{-8}H^4, F > 2U^{-6}K^3$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}, V^{-12}M^2, W^{-8}, W^{-24}H^4, U^{-6}, U^{-18}K^3\} T \\ &\leq U^2V^2W^2 (V^{-4})^{\frac{1}{2}} (W^{-8})^{\frac{1}{4}} (U^{-6})^{\frac{5}{24}} (U^{-18}K^3)^{\frac{1}{24}} T \\ &= TK^{\frac{1}{8}} \\ &= T(MHK)^{\frac{1}{8}} \left(M^{-\frac{1}{8}}H^{-\frac{1}{8}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{13}{21}} \ll MH$ is required. This can be obtained by using $X^{\frac{100}{231}} \ll M$ and $X^{\frac{4}{21}} \ll H$.

Assuming the condition (7). By applying the mean value estimate and Halász method to $M^2(s)H(s), H^6(s)$ and $K^2(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$F = \min \{V^{-4}W^{-2}(M^2H + T), V^{-4}W^{-2}M^2H + V^{-12}W^{-6}M^2HT, \\ W^{-12}(H^6 + T), W^{-12}H^6 + W^{-36}H^6T, U^{-4}(K^2 + T), U^{-4}K^2 + U^{-12}K^2T\}.$$

We consider 8 cases:

(a) $F \leq 2V^{-4}W^{-2}M^2H, F \leq 2W^{-12}H^6, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-12}H^6, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(b) $F \leq 2V^{-4}W^{-2}M^2H, F \leq 2W^{-12}H^6, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-12}H^6, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}T)^{\frac{3}{8}} (U^{-12}K^2T)^{\frac{1}{24}} \\ &= T^{\frac{5}{12}} MHK^{\frac{1}{12}} \\ &= T^{\frac{5}{12}} (MHK)^{\frac{1}{12}} \left(M^{\frac{11}{12}}H^{\frac{11}{12}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $MH \ll X^{\frac{131}{231}}$ is required.

(c) $F \leq 2V^{-4}W^{-2}M^2H, F > 2W^{-12}H^6, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-12}T, W^{-36}H^6T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \end{aligned}$$

$$\ll x(\log x)^{-11B}.$$

(d) $F \leq 2V^{-4}W^{-2}M^2H, F > 2W^{-12}H^6, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}M^2H, W^{-12}T, W^{-36}H^6T, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}M^2H)^{\frac{1}{2}} (W^{-36}H^6T)^{\frac{1}{36}} (U^{-4}T)^{\frac{11}{24}} (U^{-12}K^2T)^{\frac{1}{72}} \\ &= T^{\frac{1}{2}} MH^{\frac{2}{3}} K^{\frac{1}{36}} \\ &= T^{\frac{1}{2}} (MHK)^{\frac{1}{36}} \left(M^{\frac{35}{36}} H^{\frac{23}{36}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^{35}H^{23} \ll X^{\frac{125}{7}}$ is required.

(e) $F > 2V^{-4}W^{-2}M^2H, F \leq 2W^{-12}H^6, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-12}H^6, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{3}{8}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{24}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{5}{12}} M^{\frac{1}{12}} H^{\frac{13}{24}} K \\ &= T^{\frac{5}{12}} (MHK) \left(M^{-\frac{11}{12}} H^{-\frac{11}{24}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{200}{231}} \ll M^2H$ is required.

(f) $F > 2V^{-4}W^{-2}M^2H, F \leq 2W^{-12}H^6, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-12}H^6, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{3}{8}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{24}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}T)^{\frac{1}{2}} \\ &= T^{\frac{11}{12}} \left(M^{\frac{1}{12}} H^{\frac{13}{24}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^2H^{13} \ll X^{\frac{64}{21}}$ is required.

(g) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-12}H^6, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-12}T, W^{-36}H^6T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{11}{24}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{72}} (W^{-36}H^6T)^{\frac{1}{36}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{1}{2}} M^{\frac{1}{36}} H^{\frac{13}{72}} K \\ &= T^{\frac{1}{2}} (MHK) \left(M^{-\frac{35}{36}} H^{-\frac{59}{72}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{240}{7}} \ll M^7H^{59}$ is required.

(h) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-12}H^6, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}, V^{-12}W^{-6}M^2H, W^{-12}, W^{-36}H^6, U^{-4}, U^{-12}K^2\} T \\ &\leq U^2V^2W^2 (V^{-4}W^{-2})^{\frac{1}{2}} (W^{-12})^{\frac{1}{12}} (U^{-4})^{\frac{3}{8}} (U^{-12}K^2)^{\frac{1}{24}} T \\ &= TK^{\frac{1}{12}} \\ &= T(MHK)^{\frac{1}{12}} \left(M^{-\frac{1}{12}} H^{-\frac{1}{12}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{3}{7}} \ll MH$ is required. This can be obtained by using $X^{\frac{24}{49}} \ll MH^{\frac{59}{70}} \ll MH$.

Assuming the condition (8). By applying the mean value estimate and Halász method to $M^2(s)H(s)$, $H^7(s)$ and $K^2(s)$, we get

$$I \ll U^2 V^2 W^2 x^{-1} F(\log x)^c$$

where

$$F = \min \left\{ V^{-4} W^{-2} (M^2 H + T), V^{-4} W^{-2} M^2 H + V^{-12} W^{-6} M^2 H T, \right. \\ \left. W^{-14} (H^7 + T), W^{-14} H^7 + W^{-42} H^7 T, U^{-4} (K^2 + T) U^{-4} K^2 + U^{-12} K^2 T \right\}.$$

We consider 8 cases:

(a) $F \leq 2V^{-4}W^{-2}M^2H$, $F \leq 2W^{-14}H^7$, $F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{ V^{-4} W^{-2} M^2 H, W^{-14} H^7, U^{-4} K^2 \} \\ &\leq U^2 V^2 W^2 (V^{-4} W^{-2} M^2 H)^{\frac{1}{2}} (U^{-4} K^2)^{\frac{1}{2}} \\ &= W M H^{\frac{1}{2}} K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(b) $F \leq 2V^{-4}W^{-2}M^2H$, $F \leq 2W^{-14}H^7$, $F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{ V^{-4} W^{-2} M^2 H, W^{-14} H^7, U^{-4} T, U^{-12} K^2 T \} \\ &\leq U^2 V^2 W^2 (V^{-4} W^{-2} M^2 H)^{\frac{1}{2}} (W^{-14} H^7)^{\frac{1}{14}} (U^{-4} T)^{\frac{11}{28}} (U^{-12} K^2 T)^{\frac{1}{28}} \\ &= T^{\frac{3}{7}} M H K^{\frac{1}{14}} \\ &= T^{\frac{3}{7}} (M H K)^{\frac{1}{14}} \left(M^{\frac{13}{14}} H^{\frac{13}{14}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $MH \ll X^{\frac{51}{91}}$ is required.

(c) $F \leq 2V^{-4}W^{-2}M^2H$, $F > 2W^{-14}H^7$, $F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{ V^{-4} W^{-2} M^2 H, W^{-14} T, W^{-42} H^7 T, U^{-4} K^2 \} \\ &\leq U^2 V^2 W^2 (V^{-4} W^{-2} M^2 H)^{\frac{1}{2}} (U^{-4} K^2)^{\frac{1}{2}} \\ &= W M H^{\frac{1}{2}} K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(d) $F \leq 2V^{-4}W^{-2}M^2H$, $F > 2W^{-14}H^7$, $F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{ V^{-4} W^{-2} M^2 H, W^{-14} T, W^{-42} H^7 T, U^{-4} T, U^{-12} K^2 T \} \\ &\leq U^2 V^2 W^2 (V^{-4} W^{-2} M^2 H)^{\frac{1}{2}} (W^{-42} H^7 T)^{\frac{1}{42}} (U^{-4} T)^{\frac{13}{28}} (U^{-12} K^2 T)^{\frac{1}{84}} \\ &= T^{\frac{1}{2}} M H^{\frac{2}{3}} K^{\frac{1}{42}} \\ &= T^{\frac{1}{2}} (M H K)^{\frac{1}{42}} \left(M^{\frac{41}{42}} H^{\frac{9}{14}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^{41} H^{27} \ll X^{21}$ is required.

(e) $F > 2V^{-4}W^{-2}M^2H$, $F \leq 2W^{-14}H^7$, $F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2 V^2 W^2 F &\ll U^2 V^2 W^2 \min \{ V^{-4} W^{-2} T, V^{-12} W^{-6} M^2 H T, W^{-14} H^7, U^{-4} K^2 \} \\ &\leq U^2 V^2 W^2 (V^{-4} W^{-2} T)^{\frac{11}{28}} (V^{-12} W^{-6} M^2 H T)^{\frac{1}{28}} (W^{-14} H^7)^{\frac{1}{14}} (U^{-4} K^2)^{\frac{1}{2}} \\ &= T^{\frac{3}{7}} M^{\frac{1}{14}} H^{\frac{15}{28}} K \\ &= T^{\frac{3}{7}} (M H K) \left(M^{-\frac{13}{14}} H^{-\frac{13}{28}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{80}{91}} \ll M^2H$ is required.

(f) $F > 2V^{-4}W^{-2}M^2H, F \leq 2W^{-14}H^7, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-14}H^7, U^{-4}T, U^{-12}K^2T\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{11}{28}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{28}} (W^{-14}H^7)^{\frac{1}{14}} (U^{-4}T)^{\frac{1}{2}} \\ &= T^{\frac{13}{14}} \left(M^{\frac{1}{14}} H^{\frac{15}{28}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M^2H^{15} \ll X^{\frac{68}{21}}$ is required.

(g) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-14}H^7, F \leq 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}T, V^{-12}W^{-6}M^2HT, W^{-14}T, W^{-42}H^7T, U^{-4}K^2\} \\ &\leq U^2V^2W^2 (V^{-4}W^{-2}T)^{\frac{13}{88}} (V^{-12}W^{-6}M^2HT)^{\frac{1}{84}} (W^{-42}H^7T)^{\frac{1}{42}} (U^{-4}K^2)^{\frac{1}{2}} \\ &= T^{\frac{1}{2}} M^{\frac{1}{42}} H^{\frac{5}{28}} K \\ &= T^{\frac{1}{2}} (MHK) \left(M^{-\frac{41}{42}} H^{-\frac{23}{28}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{40} \ll M^{82}H^{69}$ is required.

(h) $F > 2V^{-4}W^{-2}M^2H, F > 2W^{-14}H^7, F > 2U^{-4}K^2$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}W^{-2}, V^{-12}W^{-6}M^2H, W^{-14}, W^{-42}H^7, U^{-4}, U^{-12}K^2\} T \\ &\leq U^2V^2W^2 (V^{-4}W^{-2})^{\frac{1}{2}} (W^{-14})^{\frac{1}{14}} (U^{-4})^{\frac{11}{28}} (U^{-12}K^2)^{\frac{1}{28}} T \\ &= TK^{\frac{1}{14}} \\ &= T(MHK)^{\frac{1}{14}} \left(M^{-\frac{1}{14}} H^{-\frac{1}{14}} \right) \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $X^{\frac{1}{3}} \ll MH$ is required. This can be obtained by using $X^{\frac{40}{91}} \ll MH^{\frac{1}{2}} \ll MH$.

Assuming the condition (9). By applying the mean value estimate and Halász method to $M^2(s), H^6(s)$ and $K^2(s)H(s)$, we get

$$I \ll U^2V^2W^2x^{-1}F(\log x)^c$$

where

$$F = \min \{V^{-4}(M^2 + T), V^{-4}M^2 + V^{-12}M^2T, W^{-12}(H^6 + T), W^{-12}H^6 + W^{-36}H^6T, \\ U^{-4}W^{-2}(K^2H + T), U^{-4}W^{-2}K^2H + U^{-12}W^{-6}K^2HT\}.$$

We consider 8 cases:

(a) $F \leq 2V^{-4}M^2, F \leq 2W^{-12}H^6, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-12}H^6, U^{-4}W^{-2}K^2H\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\ &= WMH^{\frac{1}{2}}K \\ &\ll x(\log x)^{-11B}. \end{aligned}$$

(b) $F \leq 2V^{-4}M^2, F \leq 2W^{-12}H^6, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min \{V^{-4}M^2, W^{-12}H^6, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT\} \\ &\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}W^{-2}T)^{\frac{3}{8}} (U^{-12}W^{-6}K^2HT)^{\frac{1}{24}} \\ &= T^{\frac{5}{12}} MH^{\frac{13}{24}} K^{\frac{1}{12}} \end{aligned}$$

$$\begin{aligned}
&= T^{\frac{5}{12}}(MHK)^{\frac{1}{12}} \left(M^{\frac{11}{12}} H^{\frac{11}{24}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^2H \ll X^{\frac{262}{231}}$ is required.

(c) $F \leq 2V^{-4}M^2, F > 2W^{-12}H^6, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{ V^{-4}M^2, W^{-12}T, W^{-36}H^6T, U^{-4}W^{-2}K^2H \} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\
&= WMH^{\frac{1}{2}}K \\
&\ll x(\log x)^{-11B}.
\end{aligned}$$

(d) $F \leq 2V^{-4}M^2, F > 2W^{-12}H^6, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{ V^{-4}M^2, W^{-12}T, W^{-36}H^6T, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT \} \\
&\leq U^2V^2W^2 (V^{-4}M^2)^{\frac{1}{2}} (W^{-36}H^6T)^{\frac{1}{36}} (U^{-4}W^{-2}T)^{\frac{1}{24}} (U^{-12}W^{-6}K^2HT)^{\frac{1}{72}} \\
&= T^{\frac{1}{2}}MH^{\frac{13}{72}}K^{\frac{1}{36}} \\
&= T^{\frac{1}{2}}(MHK)^{\frac{1}{36}} \left(M^{\frac{35}{36}} H^{\frac{11}{72}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $M^{70}H^{11} \ll X^{\frac{250}{7}}$ is required.

(e) $F > 2V^{-4}M^2, F \leq 2W^{-12}H^6, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{ V^{-4}T, V^{-12}M^2T, W^{-12}H^6, U^{-4}W^{-2}K^2H \} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{3}{8}} (V^{-12}M^2T)^{\frac{1}{24}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\
&= T^{\frac{5}{12}}M^{\frac{1}{12}}HK \\
&= T^{\frac{5}{12}}(MHK)M^{-\frac{11}{12}} \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{100}{231}} \ll M$ is required.

(f) $F > 2V^{-4}M^2, F \leq 2W^{-12}H^6, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{ V^{-4}T, V^{-12}M^2T, W^{-12}H^6, U^{-4}W^{-2}T, U^{-12}W^{-6}K^2HT \} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{3}{8}} (V^{-12}M^2T)^{\frac{1}{24}} (W^{-12}H^6)^{\frac{1}{12}} (U^{-4}W^{-2}T)^{\frac{1}{2}} \\
&= T^{\frac{11}{12}} \left(M^{\frac{1}{12}} H^{\frac{1}{2}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $MH^6 \ll X^{\frac{32}{21}}$ is required.

(g) $F > 2V^{-4}M^2, F > 2W^{-12}H^6, F \leq 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned}
U^2V^2W^2F &\ll U^2V^2W^2 \min \{ V^{-4}T, V^{-12}M^2T, W^{-12}T, W^{-36}H^6T, U^{-4}W^{-2}K^2H \} \\
&\leq U^2V^2W^2 (V^{-4}T)^{\frac{11}{24}} (V^{-12}M^2T)^{\frac{1}{72}} (W^{-36}H^6T)^{\frac{1}{36}} (U^{-4}W^{-2}K^2H)^{\frac{1}{2}} \\
&= T^{\frac{1}{2}}M^{\frac{1}{36}}H^{\frac{2}{3}}K \\
&= T^{\frac{1}{2}}(MHK) \left(M^{-\frac{35}{36}} H^{-\frac{1}{3}} \right) \\
&\ll x^{1-\varepsilon_1},
\end{aligned}$$

where $X^{\frac{120}{7}} \ll M^{35}H^{12}$ is required.

(h) $F > 2V^{-4}M^2, F > 2W^{-12}H^6, F > 2U^{-4}W^{-2}K^2H$. Then

$$\begin{aligned} U^2V^2W^2F &\ll U^2V^2W^2 \min\{V^{-4}, V^{-12}M^2, W^{-12}, W^{-36}H^6, U^{-4}W^{-2}, U^{-12}W^{-6}K^2H\}T \\ &\leq U^2V^2W^2 (V^{-4})^{\frac{3}{8}} (V^{-12}M^2)^{\frac{1}{24}} (W^{-12})^{\frac{1}{12}} (U^{-4}W^{-2})^{\frac{1}{2}} T \\ &= TM^{\frac{1}{12}} \\ &\ll x^{1-\varepsilon_1}, \end{aligned}$$

where $M \ll X^{\frac{4}{7}}$ is required. This can be obtained by using $M \ll MH^{\frac{11}{70}} \ll X^{\frac{25}{49}}$.

Finally, by combining all the cases above, Lemma 3.2 is proved. \square

Lemma 3.3. *Suppose that $NQRK = X$ where $N(s), Q(s), R(s)$ and $K(s)$ are Dirichlet polynomials and $G(s) = N(s)Q(s)R(s)K(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$. Assume further that for $T_0 \leq |t| \leq 2X$, $N(b+it)Q(b+it) \ll (\log x)^{-B/\varepsilon}$ and $R(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N, Q and R satisfy one of the following 8 conditions:*

- (1) $X^{\frac{4423}{10906}} \ll NQ \ll X^{\frac{113}{273}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-\frac{35}{23}} X^{\frac{125}{161}}$;
- (2) $X^{\frac{113}{273}} \ll NQ \ll X^{\frac{267}{637}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-1} X^{\frac{51}{91}}$;
- (3) $X^{\frac{267}{637}} \ll NQ \ll X^{\frac{2442}{5453}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-\frac{41}{27}} X^{\frac{21}{27}}$;
- (4) $X^{\frac{2124}{4655}} \ll NQ \ll X^{\frac{40}{87}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-\frac{1}{5}} X^{\frac{10}{35}}$.
- (5) $X^{\frac{40}{87}} \ll NQ \ll X^{\frac{10}{21}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{\frac{1}{2}} X^{\frac{9}{49}}$;
- (6) $X^{\frac{10}{21}} \ll NQ \ll X^{\frac{37}{77}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-1} X^{\frac{107}{147}}$;
- (7) $X^{\frac{37}{77}} \ll NQ \ll X^{\frac{61}{126}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-2} X^{\frac{262}{231}}$.
- (8) $X^{\frac{61}{126}} \ll NQ \ll X^{\frac{4607}{9310}}, X^{\frac{13}{133}} \ll R \ll Q \ll (NQ)^{-\frac{70}{11}} X^{\frac{250}{77}}$.

Then for $T_0 \leq T \leq X$, we have

$$\left(\min\left(\eta, \frac{1}{T}\right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (9)$$

Proof. The proof is similar to that of [[25], Lemma 13]. Let $M(s) = N(s)Q(s)$ and $H(s) = R(s)$. Then by Lemma 3.2, Lemma 3.3 is proved. \square

Lemma 3.4. *Suppose that $NQRK = X$ where $N(s), Q(s), R(s)$ and $K(s)$ are Dirichlet polynomials and $G(s) = N(s)Q(s)R(s)K(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$. Assume further that for $T_0 \leq |t| \leq 2X$, $N(b+it)R(b+it) \ll (\log x)^{-B/\varepsilon}$ and $Q(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N, Q and R satisfy one of the following 9 conditions:*

- (1) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{47}{87}}, X^{\frac{40}{231}} \ll Q, (NR)^{29}Q^{-1} \ll X^{\frac{209}{21}}, X^{\frac{40}{133}} \ll (NR), (NR)^{-1}Q^{29} \ll X^{\frac{43}{7}}, X^{\frac{40}{7}} \ll (NR)^{12}Q^{11}$;
- (2) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{109}{189}}, (NR)^{29}Q^{19} \ll X^{\frac{103}{7}}, X^{\frac{160}{189}} \ll (NR)^2Q, (NR)^2Q^{11} \ll X^{\frac{20}{7}}, X^{\frac{200}{7}} \ll (NR)^{58}Q^{49}$;
- (3) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{131}{231}}, X^{\frac{4}{21}} \ll Q, (NR)^6Q \ll X^{\frac{46}{21}}, X^{\frac{40}{147}} \ll (NR), (NR)Q^8 \ll X^{\frac{16}{7}}, X^{\frac{20}{7}} \ll (NR)^6Q^5$;
- (4) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{93}{133}}, X^{\frac{40}{231}} \ll Q, (NR)^{12}Q \ll X^{\frac{44}{7}}, X^{\frac{40}{87}} \ll (NR), (NR)^{-1}Q^{19} \ll X^{\frac{79}{21}}, X^{\frac{160}{21}} \ll (NR)^{12}Q^{11}$;
- (5) $X^{\frac{13}{133}} \ll R \ll Q, (NR)^2Q \ll X^{\frac{218}{189}}, (NR)^{58}Q^9 \ll X^{\frac{206}{7}}, X^{\frac{80}{189}} \ll (NR), (NR)Q^5 \ll X^{\frac{10}{7}}, X^{\frac{100}{7}} \ll (NR)^{29}Q^{10}$;
- (6) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{107}{147}}, X^{\frac{4}{21}} \ll Q, (NR)^6Q \ll X^{\frac{22}{7}}, X^{\frac{100}{231}} \ll (NR), (NR)^{-1}Q^7 \ll X^{\frac{9}{7}}, X^{\frac{80}{21}} \ll (NR)^6Q^5$;
- (7) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{131}{231}}, (NR)^{35}Q^{23} \ll X^{\frac{125}{7}}, X^{\frac{200}{231}} \ll (NR)^2Q, (NR)^2Q^{13} \ll X^{\frac{64}{21}}, X^{\frac{240}{7}} \ll (NR)^{70}Q^{59}$;
- (8) $X^{\frac{13}{133}} \ll R \ll Q, (NR)Q \ll X^{\frac{51}{91}}, (NR)^{41}Q^{27} \ll X^{21}, X^{\frac{80}{91}} \ll (NR)^2Q, (NR)^2Q^{15} \ll X^{\frac{68}{21}}, X^{40} \ll (NR)^{82}Q^{69}$;
- (9) $X^{\frac{13}{133}} \ll R \ll Q, (NR)^2Q \ll X^{\frac{262}{231}}, (NR)^{70}Q^{11} \ll X^{\frac{250}{7}}, X^{\frac{100}{231}} \ll (NR), (NR)Q^6 \ll X^{\frac{32}{21}}, X^{\frac{120}{7}} \ll (NR)^{35}Q^{12}$.

Then for $T_0 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (10)$$

Proof. Let $M(s) = N(s)R(s)$ and $H(s) = Q(s)$ and by Lemma 3.2, Lemma 3.4 is proved. \square

Lemma 3.5. *Suppose that $NQRK = X$ where $N(s), Q(s), R(s)$ and $K(s)$ are Dirichlet polynomials and $G(s) = N(s)Q(s)R(s)K(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$. Assume further that for $T_0 \leq |t| \leq 2X$, $Q(b+it)R(b+it) \ll (\log x)^{-B/\varepsilon}$ and $N(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N, Q and R satisfy one of the following 9 conditions:*

- (1) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{47}{87}}$, $X^{\frac{40}{231}} \ll N$, $(QR)^{29}N^{-1} \ll X^{\frac{209}{21}}$, $X^{\frac{40}{133}} \ll (QR)$, $(QR)^{-1}N^{29} \ll X^{\frac{43}{7}}$, $X^{\frac{40}{7}} \ll (QR)^{12}N^{11}$;
- (2) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{109}{189}}$, $(QR)^{29}N^{19} \ll X^{\frac{103}{7}}$, $X^{\frac{160}{189}} \ll (QR)^2N$, $(QR)^2N^{11} \ll X^{\frac{20}{7}}$, $X^{\frac{200}{7}} \ll (QR)^{58}N^{49}$;
- (3) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{131}{231}}$, $X^{\frac{4}{21}} \ll N$, $(QR)^6N \ll X^{\frac{46}{21}}$, $X^{\frac{40}{147}} \ll (QR)$, $(QR)N^8 \ll X^{\frac{16}{7}}$, $X^{\frac{20}{7}} \ll (QR)^6N^5$;
- (4) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{93}{133}}$, $X^{\frac{40}{231}} \ll N$, $(QR)^{12}N \ll X^{\frac{44}{7}}$, $X^{\frac{40}{87}} \ll (QR)$, $(QR)^{-1}N^{19} \ll X^{\frac{79}{21}}$, $X^{\frac{160}{21}} \ll (QR)^{12}N^{11}$;
- (5) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)^2N \ll X^{\frac{218}{189}}$, $(QR)^{58}N^9 \ll X^{\frac{206}{7}}$, $X^{\frac{80}{189}} \ll (QR)$, $(QR)N^5 \ll X^{\frac{10}{7}}$, $X^{\frac{100}{7}} \ll (QR)^{29}N^{10}$;
- (6) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{107}{147}}$, $X^{\frac{4}{21}} \ll N$, $(QR)^6N \ll X^{\frac{22}{7}}$, $X^{\frac{100}{231}} \ll (QR)$, $(QR)^{-1}N^7 \ll X^{\frac{9}{7}}$, $X^{\frac{80}{21}} \ll (QR)^6N^5$;
- (7) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{131}{231}}$, $(QR)^{35}N^{23} \ll X^{\frac{125}{7}}$, $X^{\frac{200}{231}} \ll (QR)^2N$, $(QR)^2N^{13} \ll X^{\frac{64}{21}}$, $X^{\frac{240}{7}} \ll (QR)^{70}N^{59}$;
- (8) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)N \ll X^{\frac{51}{91}}$, $(QR)^{41}N^{27} \ll X^{21}$, $X^{\frac{80}{91}} \ll (QR)^2N$, $(QR)^2N^{15} \ll X^{\frac{68}{21}}$, $X^{40} \ll (QR)^{82}N^{69}$;
- (9) $X^{\frac{13}{133}} \ll R \ll Q$, $(QR)^2N \ll X^{\frac{262}{231}}$, $(QR)^{70}N^{11} \ll X^{\frac{250}{7}}$, $X^{\frac{100}{231}} \ll (QR)$, $(QR)N^6 \ll X^{\frac{32}{21}}$, $X^{\frac{120}{7}} \ll (QR)^{35}N^{12}$.

Then for $T_0 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (11)$$

Proof. Let $M(s) = Q(s)R(s)$ and $H(s) = N(s)$ and by Lemma 3.2, Lemma 3.5 is proved. \square

Lemma 3.6. *Suppose that $NQRK = X$ where $N(s), Q(s), R(s)$ and $K(s)$ are Dirichlet polynomials and $G(s) = N(s)Q(s)R(s)K(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_0 = (\log X)^{B/\varepsilon}$. Assume further that for $T_0 \leq |t| \leq 2X$, $Q(b+it)R(b+it) \ll (\log x)^{-B/\varepsilon}$ and $N(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N, Q and R satisfy one of the following 9 conditions:*

- (1) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{47}{87}}$, $X^{\frac{40}{231}} \ll (QR)$, $N^{29}(QR)^{-1} \ll X^{\frac{209}{21}}$, $X^{\frac{40}{133}} \ll N$, $N^{-1}(QR)^{29} \ll X^{\frac{43}{7}}$, $X^{\frac{40}{7}} \ll N^{12}(QR)^{11}$;
- (2) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{109}{189}}$, $N^{29}(QR)^{19} \ll X^{\frac{103}{7}}$, $X^{\frac{160}{189}} \ll N^2(QR)$, $N^2(QR)^{11} \ll X^{\frac{20}{7}}$, $X^{\frac{200}{7}} \ll N^{58}(QR)^{49}$;
- (3) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{131}{231}}$, $X^{\frac{4}{21}} \ll (QR)$, $N^6(QR) \ll X^{\frac{46}{21}}$, $X^{\frac{40}{147}} \ll N$, $N(QR)^8 \ll X^{\frac{16}{7}}$, $X^{\frac{20}{7}} \ll N^6(QR)^5$;
- (4) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{93}{133}}$, $X^{\frac{40}{231}} \ll (QR)$, $N^{12}(QR) \ll X^{\frac{44}{7}}$, $X^{\frac{40}{87}} \ll N$, $N^{-1}(QR)^{19} \ll X^{\frac{79}{21}}$, $X^{\frac{160}{21}} \ll N^{12}(QR)^{11}$;
- (5) $X^{\frac{13}{133}} \ll R \ll Q$, $N^2(QR) \ll X^{\frac{218}{189}}$, $N^{58}(QR)^9 \ll X^{\frac{206}{7}}$, $X^{\frac{80}{189}} \ll N$, $N(QR)^5 \ll X^{\frac{10}{7}}$, $X^{\frac{100}{7}} \ll N^{29}(QR)^{10}$;
- (6) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{107}{147}}$, $X^{\frac{4}{21}} \ll (QR)$, $N^6(QR) \ll X^{\frac{22}{7}}$, $X^{\frac{100}{231}} \ll N$, $N^{-1}(QR)^7 \ll X^{\frac{9}{7}}$, $X^{\frac{80}{21}} \ll N^6(QR)^5$;
- (7) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{131}{231}}$, $N^{35}(QR)^{23} \ll X^{\frac{125}{7}}$, $X^{\frac{200}{231}} \ll N^2(QR)$, $N^2(QR)^{13} \ll X^{\frac{64}{21}}$, $X^{\frac{240}{7}} \ll N^{70}(QR)^{59}$;

(8) $X^{\frac{13}{133}} \ll R \ll Q$, $N(QR) \ll X^{\frac{51}{91}}$, $N^{41}(QR)^{27} \ll X^{21}$, $X^{\frac{80}{91}} \ll N^2(QR)$, $N^2(QR)^{15} \ll X^{\frac{68}{21}}$, $X^{40} \ll N^{82}(QR)^{69}$;

(9) $X^{\frac{13}{133}} \ll R \ll Q$, $N^2(QR) \ll X^{\frac{262}{231}}$, $N^{70}(QR)^{11} \ll X^{\frac{250}{7}}$, $X^{\frac{100}{231}} \ll N$, $N(QR)^6 \ll X^{\frac{32}{21}}$, $X^{\frac{120}{7}} \ll N^{35}(QR)^{12}$.

Then for $T_0 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (12)$$

Proof. Let $M(s) = N(s)$ and $H(s) = Q(s)R(s)$ and by Lemma 3.2, Lemma 3.6 is proved. \square

4. ARITHMETIC INFORMATION II

In [25], Jia used a corollary of a mean value bound of Deshouillers and Iwaniec [6], which makes an approximation to the sixth-power moment of the Riemann zeta-function. In [[13], Chapter 9] another mean value estimate of Watt [38] is used. Watt's bound is somewhere better than that of Deshouillers and Iwaniec, but we still use Jia's corollary for clarity. In fact, those two bounds have the same strength when the length of interval is $n^{\frac{1}{21}+\varepsilon}$ if we use them in a same way.

Lemma 4.1. ([25], Lemma 2]. Suppose that $N(s)$ is a Dirichlet polynomial and $L(s) = \sum_{l \sim L} l^{-s}$. Let $T \geq 1$, then

$$\int_T^{2T} \left| L \left(\frac{1}{2} + it \right) \right|^4 \left| N \left(\frac{1}{2} + it \right) \right|^2 dt \ll \left(T + N^2 T^{\frac{1}{2}} + N^{\frac{5}{4}} \left(T \min \left(L, \frac{T}{L} \right) \right)^{\frac{1}{2}} + NL^2 T^{-2} \right) T^{\varepsilon_1}.$$

Lemma 4.2. Suppose that $MHL = X$ where $M(s), H(s)$ are Dirichlet polynomials and $L(s) = \sum_{l \sim L} l^{-s}$. Let $b = 1 + \frac{1}{\log X}$, $T_1 = \sqrt{L}$. Moreover, suppose that M and H satisfy one of the following 2 conditions:

(1) $M^2 H \ll X^{2-\frac{20}{21}}$, $M^4 H^6 \ll X^{4-\frac{20}{21}}$, $M^6 H^7 \ll X^{6-\frac{40}{21}}$, $H^4 \ll X^{4-\frac{20}{7}}$, $M^{-2} H^3 \ll X^{6-\frac{40}{7}}$;

(2) $M \ll X^{\frac{11}{21}}$, $H \ll X^{\frac{5}{42}}$.

Then for $T_1 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |M(b+it)H(b+it)L(b+it)|^2 dt \ll \eta^2 x^{-2\varepsilon_1}. \quad (13)$$

Proof. The proof is similar to that of [[25], Lemmas 3,4] so we omit it here. \square

Lemma 4.3. Suppose that $NQRL = X$ where $N(s), Q(s)$ and $R(s)$ are Dirichlet polynomials, $L(s) = \sum_{l \sim L} l^{-s}$ and $G(s) = N(s)Q(s)R(s)L(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_1 = \sqrt{L}$. Assume further that for $T_1 \leq |t| \leq 2X$, $N(b+it)Q(b+it) \ll (\log x)^{-B/\varepsilon}$ and $R(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N , Q and R satisfy one of the following 7 conditions:

(1) $X^{\frac{13}{133}} \ll R \ll Q$, $(NQ)^2 R \ll X^{2-\frac{20}{21}}$, $(NQ)^4 R^6 \ll X^{4-\frac{20}{21}}$, $(NQ)^6 R^7 \ll X^{6-\frac{40}{21}}$, $R^4 \ll X^{4-\frac{20}{7}}$, $(NQ)^{-2} R^3 \ll X^{6-\frac{40}{7}}$;

(2) $X^{\frac{13}{133}} \ll R \ll Q$, $(NQ) \ll X^{\frac{11}{21}}$, $R \ll X^{\frac{5}{42}}$;

(3) $X^{\frac{13}{133}} \ll R \ll Q$, $X^{\frac{80}{189}} \ll (NQ) \ll X^{\frac{40}{87}}$, $R \ll (NQ)^{-\frac{1}{5}} X^{\frac{10}{35}}$;

(4) $X^{\frac{13}{133}} \ll R \ll Q$, $X^{\frac{40}{87}} \ll (NQ) \ll X^{\frac{10}{21}}$, $R \ll (NQ)^{\frac{1}{7}} X^{\frac{9}{49}}$;

(5) $X^{\frac{13}{133}} \ll R \ll Q$, $X^{\frac{10}{21}} \ll (NQ) \ll X^{\frac{37}{77}}$, $R \ll (NQ)^{-1} X^{\frac{107}{147}}$;

(6) $X^{\frac{13}{133}} \ll R \ll Q$, $X^{\frac{37}{77}} \ll (NQ) \ll X^{\frac{61}{126}}$, $R \ll (NQ)^{-2} X^{\frac{262}{231}}$;

(7) $X^{\frac{13}{133}} \ll R \ll Q$, $X^{\frac{61}{126}} \ll (NQ) \ll X^{\frac{289}{588}}$, $R \ll (NQ)^{-\frac{70}{11}} X^{\frac{250}{77}}$.

Then for $T_1 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (14)$$

Proof. The proof is similar to that of [[25], Lemmas 14,15]. Let $M(s) = N(s)Q(s)$ and $H(s) = R(s)$. Then by Lemma 4.2 and Lemma 3.2, Lemma 4.3 is proved. \square

Lemma 4.4. *Suppose that $NQRL = X$ where $N(s), Q(s)$ and $R(s)$ are Dirichlet polynomials, $L(s) = \sum_{l \sim L} l^{-s}$ and $G(s) = N(s)Q(s)R(s)L(s)$. Let $b = 1 + \frac{1}{\log X}$, $T_1 = \sqrt{L}$. Assume further that for $T_1 \leq |t| \leq 2X$, $Q(b+it)R(b+it) \ll (\log x)^{-B/\varepsilon}$ and $N(b+it) \ll (\log x)^{-B/\varepsilon}$. Moreover, suppose that N, Q and R satisfy one of the following 7 conditions:*

$$(1) X^{\frac{13}{133}} \ll R \ll Q, (QR)^2 N \ll X^{2-\frac{20}{21}}, (QR)^4 N^6 \ll X^{4-\frac{20}{21}}, (QR)^6 N^7 \ll X^{6-\frac{40}{21}}, N^4 \ll X^{4-\frac{20}{7}}, (QR)^{-2} N^3 \ll X^{6-\frac{40}{7}};$$

$$(2) X^{\frac{13}{133}} \ll R \ll Q, (QR) \ll X^{\frac{11}{21}}, N \ll X^{\frac{5}{42}};$$

$$(3) X^{\frac{13}{133}} \ll R \ll Q, X^{\frac{80}{189}} \ll (QR) \ll X^{\frac{40}{87}}, N \ll (QR)^{-\frac{1}{5}} X^{\frac{10}{35}};$$

$$(4) X^{\frac{13}{133}} \ll R \ll Q, X^{\frac{40}{87}} \ll (QR) \ll X^{\frac{10}{21}}, N \ll (QR)^{\frac{1}{7}} X^{\frac{9}{49}};$$

$$(5) X^{\frac{13}{133}} \ll R \ll Q, X^{\frac{10}{21}} \ll (QR) \ll X^{\frac{37}{77}}, N \ll (QR)^{-1} X^{\frac{107}{147}};$$

$$(6) X^{\frac{13}{133}} \ll R \ll Q, X^{\frac{37}{77}} \ll (QR) \ll X^{\frac{61}{126}}, N \ll (QR)^{-2} X^{\frac{262}{231}};$$

$$(7) X^{\frac{13}{133}} \ll R \ll Q, X^{\frac{61}{126}} \ll (QR) \ll X^{\frac{289}{538}}, N \ll (QR)^{-\frac{70}{11}} X^{\frac{250}{77}}.$$

Then for $T_1 \leq T \leq X$, we have

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B}. \quad (15)$$

Proof. Let $M(s) = Q(s)R(s)$ and $H(s) = N(s)$. Then by Lemma 4.2 and Lemma 3.2, Lemma 4.4 is proved. \square

5. SIEVE ERROR TERMS

For type-I terms we will use Iwaniec's linear sieve method and this requires us to control the errors occurred. The following two lemmas will help us deal with the error terms when we apply Iwaniec's linear sieve in Section 7.

Lemma 5.1. *Suppose that $M \ll X^{\frac{11}{21}}, H \ll X^{\frac{5}{42}}$ and $a(m) = O(1), b(h) = O(1)$. Then for real numbers $x \in [X, 2X]$, except for $O(X(\log X)^{-B})$ values, we have*

$$\sum_{\substack{m \sim M \\ h \sim H}} a(m)b(h) \left(\sum_{x < mhl \leq x + \eta x} 1 - \frac{\eta x}{mh} \right) \ll \eta x (\log x)^{-B}. \quad (16)$$

Proof. The proof is similar to that of [[25], Lemma 16] where the condition (2) of Lemma 4.2 is used. \square

Lemma 5.2. *Suppose that $a(n) = O(1), b(q) = O(1), c(r) = O(1)$ and N, Q, R satisfy one of the 14 conditions in Lemma 4.3 and Lemma 4.4. Let $b = 1 + \frac{1}{\log X}$, $T_1 = \sqrt{L}$ and $NQRL = X$. Assume further that for $T_1 \leq |t| \leq 2X$, $N(b+it)Q(b+it)R(b+it) \ll (\log x)^{-B/\varepsilon}$. Then for real numbers $x \in [X, 2X]$, except for $O(X(\log X)^{-B})$ values, we have*

$$\sum_{\substack{n \sim N \\ q \sim Q \\ r \sim R}} a(n)b(q)c(r) \left(\sum_{x < nqrl \leq x + \eta x} 1 - \frac{\eta x}{nqr} \right) \ll \eta x (\log x)^{-B}. \quad (17)$$

Proof. The proof is similar to that of [[25], Lemma 17]. \square

6. ASYMPTOTIC FORMULAS FOR TYPE-II TERMS

These lemmas will help us transform the functions $S(\mathcal{A}_{p_1 \dots p_n}, p_n)$ into $S(\mathcal{B}_{p_1 \dots p_n}, p_n)$ and estimate them.

Lemma 6.1. *Suppose that $X^{\frac{120}{133}} \ll M \ll X^{1-\delta}$ and that $0 \leq a(m) = O(1)$. If m has a prime factor $< X^\delta$, then $a(m) = 0$. Then for real numbers $x \in [X, 2X]$, except for $O(X(\log X)^{-B})$ values, we have*

$$\sum_{\substack{x < mp \leq x + \eta x \\ m \sim M}} a(m) = \eta \left(1 + O \left(\frac{1}{\log x} \right) \right) \sum_{\substack{x < mp \leq 2x \\ m \sim M}} a(m) + O(\eta x (\log x)^{-B}). \quad (18)$$

Proof. The proof is similar to that of [[25], Lemma 18] where Lemma 3.1 is used. \square

Lemma 6.2. *Suppose that $MHK = X$ where $M(s)$ and $H(s)$ are Dirichlet polynomials, $K(s) = \sum_{k \sim K} \Lambda(k)k^{-s}$. Suppose that $X^{\frac{13}{133}} \ll MH \ll X^{\frac{120}{133}}$ and that $0 \leq a(m) = O(1)$, $0 \leq b(h) = O(1)$. If m has a prime factor $< X^\delta$, then $a(m) = 0$. If h has a prime factor $< X^\delta$, then $b(h) = 0$. Let $G(s) = M(s)H(s)K(s)$, $b = 1 + \frac{1}{\log X}$ and $T_0 = (\log X)^{B/\varepsilon}$. If we have*

$$\left(\min \left(\eta, \frac{1}{T} \right) \right)^2 \int_T^{2T} |G(b+it)|^2 dt \ll \eta^2 (\log x)^{-10B} \quad (19)$$

for $T_0 \leq T \leq X$, then for real numbers $x \in [X, 2X]$, except for $O(X(\log X)^{-B})$ values, we have

$$\sum_{\substack{x < mhp \leq x + \eta x \\ m \sim M \\ h \sim H}} a(m)b(h) = \eta \left(1 + O \left(\frac{1}{\log x} \right) \right) \sum_{\substack{x < mhp \leq 2x \\ m \sim M \\ h \sim H}} a(m)b(h) + O(\eta x (\log x)^{-B}). \quad (20)$$

Proof. The proof is similar to that of [[24], Lemma 4]. □

Lemma 6.3. ([25], Lemma 21). *For sufficiently large x and z , we have*

$$S(\mathcal{B}, z) = \sum_{\substack{x < n \leq 2x \\ (n, P(z))=1}} 1 = (1 + o(1)) \frac{x}{\log z} \omega \left(\frac{\log x}{\log z} \right),$$

where $\omega(u)$ denote the Buchstab function determined by the following differential-difference equation

$$\begin{cases} \omega(u) = \frac{1}{u}, & 1 \leq u \leq 2, \\ (u\omega(u))' = \omega(u-1), & u \geq 2. \end{cases}$$

7. THE FINAL DECOMPOSITION

In this section, sets \mathcal{A} and \mathcal{B} are defined respectively. Let γ denote the Euler's constant, $F(s)$ and $f(s)$ are determined by the following differential-difference equation

$$\begin{cases} F(s) = \frac{2e^\gamma}{s}, & f(s) = 0, & 0 < s \leq 2, \\ (sF(s))' = f(s-1), & (sf(s))' = F(s-1), & s \geq 2, \end{cases}$$

and $\omega(u)$ denote the Buchstab function defined above. Moreover, we have the lower bound for $\omega(u)$:

$$\omega(u) \geq \omega_0(u) = \begin{cases} \frac{1}{u}, & 1 \leq u < 2, \\ \frac{1+\log(u-1)}{1+\log u}, & 2 \leq u < 3, \\ \frac{1+\log(u-1)}{u} + \frac{1}{u} \int_2^{u-1} \frac{\log(t-1)}{t} dt, & 3 \leq u < 4, \\ 0.5612, & u \geq 4. \end{cases}$$

We shall use $\omega_0(u)$ to give numerical lower bound for some sieve functions discussed below.

Beginning with (2), we shall use Iwaniec's linear sieve method to estimate S_1 and S_2 . By Buchstab's identity we have

$$S_1 = S \left(\mathcal{A}, X^{\frac{13}{133}} \right) = S(\mathcal{A}, X^\delta) - \sum_{\delta \leq t_1 < \frac{13}{133}} S(\mathcal{A}_{p_1}, p_1). \quad (21)$$

Let $z = X^\delta$ and $D = X^{\frac{11}{21}}$. Then by Iwaniec's linear sieve, Lemma 5.1 and arguments in [25] we have

$$\begin{aligned} S(\mathcal{A}, X^\delta) &\geq \frac{\eta x}{\log z} f \left(\frac{\log D}{\log z} \right) - O \left(\frac{\varepsilon \eta x}{\log x} \right) - \sum_{m \leq D} a(m) \left(|\mathcal{A}_m| - \frac{\eta x}{m} \right) \\ &= \frac{e^{-\gamma} \eta x}{\delta \log x} + O \left(\frac{\varepsilon \eta x}{\log x} \right); \end{aligned} \quad (22)$$

$$\begin{aligned} S(\mathcal{A}, X^\delta) &\leq \frac{\eta x}{\log z} F \left(\frac{\log D}{\log z} \right) + O \left(\frac{\varepsilon \eta x}{\log x} \right) + \sum_{m \leq D} a(m) \left(|\mathcal{A}_m| - \frac{\eta x}{m} \right) \\ &= \frac{e^{-\gamma} \eta x}{\delta \log x} + O \left(\frac{\varepsilon \eta x}{\log x} \right). \end{aligned} \quad (23)$$

So we have the asymptotic formula

$$S(\mathcal{A}, X^\delta) = \frac{e^{-\gamma}\eta x}{\delta \log x} + O\left(\frac{\varepsilon\eta x}{\log x}\right). \quad (24)$$

By Lemma 6.1 and Lemma 6.3 we have

$$\begin{aligned} \sum_{\delta \leq t_1 < \frac{13}{133}} S(\mathcal{A}_{p_1}, p_1) &= \eta \sum_{\delta \leq t_1 < \frac{13}{133}} S(\mathcal{B}_{p_1}, p_1) + O\left(\frac{\varepsilon\eta x}{\log x}\right) \\ &= \frac{\eta x}{\log x} \int_{\delta}^{\frac{13}{133}} \frac{1}{t^2} \omega\left(\frac{1-t}{t}\right) dt + O\left(\frac{\delta\eta x}{\log x}\right) \\ &= \frac{\eta x}{\log x} \int_{\frac{13}{133}}^{\frac{1}{\delta}} \omega(t-1) dt + O\left(\frac{\delta\eta x}{\log x}\right) \\ &= \frac{\eta x}{\log x} \left(\frac{1}{\delta} \omega\left(\frac{1}{\delta}\right) - \frac{133}{13} \omega\left(\frac{133}{13}\right) \right) + O\left(\frac{\delta\eta x}{\log x}\right) \\ &= \frac{e^{-\gamma}\eta x}{\delta \log x} - \frac{133}{13} \omega\left(\frac{133}{13}\right) \frac{\eta x}{\log x} + O\left(\frac{\delta\eta x}{\log x}\right). \end{aligned} \quad (25)$$

By (24) and (25) we have

$$\begin{aligned} S_1 &= S(\mathcal{A}, X^\delta) - \sum_{\delta \leq t_1 < \frac{13}{133}} S(\mathcal{A}_{p_1}, p_1) \\ &= \frac{133}{13} \omega\left(\frac{133}{13}\right) \frac{\eta x}{\log x} + O\left(\frac{\delta\eta x}{\log x}\right) \\ &\geq 5.744162 \frac{\eta x}{\log x}. \end{aligned} \quad (26)$$

Similarly, we have

$$\begin{aligned} S_2 &= \sum_{\frac{13}{133} \leq t_1 < \frac{1}{2}} S\left(\mathcal{A}_{p_1}, X^{\frac{13}{133}}\right) \\ &= \sum_{\frac{13}{133} \leq t_1 < \frac{1}{2}} S(\mathcal{A}_{p_1}, X^\delta) - \sum_{\delta \leq t_2 < \frac{13}{133} \leq t_1 < \frac{1}{2}} S(\mathcal{A}_{p_1 p_2}, p_2) \\ &= \frac{e^{-\gamma}\eta x}{\delta \log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \frac{1}{t} dt \\ &\quad - \left(\frac{e^{-\gamma}\eta x}{\delta \log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \frac{1}{t} dt - \frac{\eta x}{\log x} \frac{133}{13} \int_{\frac{13}{133}}^{\frac{1}{2}} \frac{1}{t} \omega\left(\frac{133}{13}(1-t)\right) dt \right) \\ &= \left(\frac{133}{13} \int_{\frac{13}{133}}^{\frac{1}{2}} \frac{1}{t} \omega\left(\frac{133}{13}(1-t)\right) dt \right) \frac{\eta x}{\log x} \\ &\leq 9.375923 \frac{\eta x}{\log x}. \end{aligned} \quad (27)$$

Before estimating S_3 , we first define the disjoint regions U_1 – U_5 as

$$\begin{aligned} U_1(t_1, t_2) &:= \{(t_1, t_2) \in U_{11} \cup U_{12} \cup \dots \cup U_{19}, \\ &\quad \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right)\}, \\ U_2(t_1, t_2) &:= \{(t_1, t_2) \in U_{21} \cup U_{22} \cup \dots \cup U_{28}, (t_1, t_2) \notin U_1, \\ &\quad \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right)\}, \\ U_3(t_1, t_2) &:= \{(t_1, t_2) \in U_{31} \cup U_{32} \cup \dots \cup U_{327}, (t_1, t_2) \notin U_1 \cup U_2, \end{aligned}$$

$$\begin{aligned}
& \left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min \left(t_1, \frac{1}{2}(1-t_1) \right) \right\}, \\
U_4(t_1, t_2) &:= \{(t_1, t_2) \in U_{41} \cup U_{42} \cup \dots \cup U_{47}, (t_1, t_2) \notin U_1 \cup U_2 \cup U_3, \\
& \left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min \left(t_1, \frac{1}{2}(1-t_1) \right) \right\}, \\
U_5(t_1, t_2) &:= \{(t_1, t_2) \in U_{51} \cup U_{52} \cup \dots \cup U_{57}, (t_1, t_2) \notin U_1 \cup U_2 \cup U_3 \cup U_4, \\
& \left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min \left(t_1, \frac{1}{2}(1-t_1) \right) \right\},
\end{aligned}$$

where

$$\begin{aligned}
U_{11}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{47}{87}, \frac{40}{231} \leq t_2, 29t_1 - t_2 \leq \frac{209}{21}, \frac{40}{133} \leq t_1, 29t_2 - t_1 \leq \frac{43}{7}, \frac{40}{7} \leq 12t_1 + 11t_2 \right\}, \\
U_{12}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{109}{189}, 29t_1 + 19t_2 \leq \frac{103}{7}, \frac{160}{189} \leq 2t_1 + t_2, 2t_1 + 11t_2 \leq \frac{20}{7}, \frac{200}{7} \leq 58t_1 + 49t_2 \right\}, \\
U_{13}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{131}{231}, \frac{4}{21} \leq t_2, 6t_1 + t_2 \leq \frac{46}{21}, \frac{40}{147} \leq t_1, t_1 + 8t_2 \leq \frac{16}{7}, \frac{20}{7} \leq 6t_1 + 5t_2 \right\}, \\
U_{14}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{93}{133}, \frac{40}{231} \leq t_2, 12t_1 + t_2 \leq \frac{44}{7}, \frac{40}{87} \leq t_1, 19t_2 - t_1 \leq \frac{79}{21}, \frac{160}{21} \leq 12t_1 + 11t_2 \right\}, \\
U_{15}(t_1, t_2) &:= \left\{ 2t_1 + t_2 \leq \frac{218}{189}, 58t_1 + 9t_2 \leq \frac{206}{7}, \frac{80}{189} \leq t_1, t_1 + 5t_2 \leq \frac{10}{7}, \frac{100}{7} \leq 29t_1 + 10t_2 \right\}, \\
U_{16}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{107}{147}, \frac{4}{21} \leq t_2, 6t_1 + t_2 \leq \frac{22}{7}, \frac{100}{231} \leq t_1, 7t_2 - t_1 \leq \frac{9}{7}, \frac{80}{21} \leq 6t_1 + 5t_2 \right\}, \\
U_{17}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{131}{231}, 35t_1 + 23t_2 \leq \frac{125}{7}, \frac{200}{231} \leq 2t_1 + t_2, 2t_1 + 13t_2 \leq \frac{64}{21}, \frac{240}{7} \leq 70t_1 + 59t_2 \right\}, \\
U_{18}(t_1, t_2) &:= \left\{ t_1 + t_2 \leq \frac{51}{91}, 41t_1 + 27t_2 \leq 21, \frac{80}{91} \leq 2t_1 + t_2, 2t_1 + 15t_2 \leq \frac{68}{21}, 40 \leq 82t_1 + 69t_2 \right\}, \\
U_{19}(t_1, t_2) &:= \left\{ 2t_1 + t_2 \leq \frac{262}{231}, 70t_1 + 11t_2 \leq \frac{250}{7}, \frac{100}{231} \leq t_1, t_1 + 6t_2 \leq \frac{32}{21}, \frac{120}{7} \leq 35t_1 + 12t_2 \right\}, \\
U_{21}(t_1, t_2) &:= \left\{ \frac{4423}{10906} \leq t_1 + t_2 \leq \frac{113}{273}, \frac{13}{133} \leq t_2 \leq -\frac{35}{23}(t_1 + t_2) + \frac{125}{161} \right\}, \\
U_{22}(t_1, t_2) &:= \left\{ \frac{113}{273} \leq t_1 + t_2 \leq \frac{263}{637}, \frac{13}{133} \leq t_2 \leq -(t_1 + t_2) + \frac{51}{91} \right\}, \\
U_{23}(t_1, t_2) &:= \left\{ \frac{263}{637} \leq t_1 + t_2 \leq \frac{2442}{5453}, \frac{13}{133} \leq t_2 \leq -\frac{41}{27}(t_1 + t_2) + \frac{21}{27} \right\}, \\
U_{24}(t_1, t_2) &:= \left\{ \frac{2124}{4655} \leq t_1 + t_2 \leq \frac{40}{87}, \frac{13}{133} \leq t_2 \leq -\frac{1}{5}(t_1 + t_2) + \frac{10}{35} \right\}, \\
U_{25}(t_1, t_2) &:= \left\{ \frac{40}{87} \leq t_1 + t_2 \leq \frac{10}{21}, \frac{13}{133} \leq t_2 \leq \frac{1}{7}(t_1 + t_2) + \frac{9}{49} \right\}, \\
U_{26}(t_1, t_2) &:= \left\{ \frac{10}{21} \leq t_1 + t_2 \leq \frac{37}{77}, \frac{13}{133} \leq t_2 \leq -(t_1 + t_2) + \frac{107}{147} \right\}, \\
U_{27}(t_1, t_2) &:= \left\{ \frac{37}{77} \leq t_1 + t_2 \leq \frac{61}{126}, \frac{13}{133} \leq t_2 \leq -2(t_1 + t_2) + \frac{262}{231} \right\}, \\
U_{28}(t_1, t_2) &:= \left\{ \frac{61}{126} \leq t_1 + t_2 \leq \frac{4607}{9310}, \frac{13}{133} \leq t_2 \leq -\frac{70}{11}(t_1 + t_2) + \frac{250}{77} \right\}, \\
U_{31}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{47}{87}, \frac{40}{231} \leq t_2, 29(t_1 + t_2) - t_2 \leq \frac{209}{21}, \frac{40}{133} \leq \left(t_1 + \frac{13}{133} \right) \right\},
\end{aligned}$$

$$\begin{aligned}
& 29t_2 - \left(t_1 + \frac{13}{133}\right) \leq \frac{43}{7}, \frac{40}{7} \leq 12 \left(t_1 + \frac{13}{133}\right) + 11t_2 \Big\}, \\
U_{32}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{109}{189}, 29(t_1 + t_2) + 19t_2 \leq \frac{103}{7}, \frac{160}{189} \leq 2 \left(t_1 + \frac{13}{133}\right) + t_2, \right. \\
& \left. 2(t_1 + t_2) + 11t_2 \leq \frac{20}{7}, \frac{200}{7} \leq 58 \left(t_1 + \frac{13}{133}\right) + 49t_2 \right\}, \\
U_{33}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{131}{231}, \frac{4}{21} \leq t_2, 6(t_1 + t_2) + t_2 \leq \frac{46}{21}, \frac{40}{147} \leq \left(t_1 + \frac{13}{133}\right), \right. \\
& \left. (t_1 + t_2) + 8t_2 \leq \frac{16}{7}, \frac{20}{7} \leq 6 \left(t_1 + \frac{13}{133}\right) + 5t_2 \right\}, \\
U_{34}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{93}{133}, \frac{40}{231} \leq t_2, 12(t_1 + t_2) + t_2 \leq \frac{44}{7}, \frac{40}{87} \leq \left(t_1 + \frac{13}{133}\right), \right. \\
& \left. 19t_2 - \left(t_1 + \frac{13}{133}\right) \leq \frac{79}{21}, \frac{160}{21} \leq 12 \left(t_1 + \frac{13}{133}\right) + 11t_2 \right\}, \\
U_{35}(t_1, t_2) &:= \left\{ 2(t_1 + t_2) + t_2 \leq \frac{218}{189}, 58(t_1 + t_2) + 9t_2 \leq \frac{206}{7}, \frac{80}{189} \leq \left(t_1 + \frac{13}{133}\right), \right. \\
& \left. (t_1 + t_2) + 5t_2 \leq \frac{10}{7}, \frac{100}{7} \leq 29 \left(t_1 + \frac{13}{133}\right) + 10t_2 \right\}, \\
U_{36}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{107}{147}, \frac{4}{21} \leq t_2, 6(t_1 + t_2) + t_2 \leq \frac{22}{7}, \frac{100}{231} \leq \left(t_1 + \frac{13}{133}\right), \right. \\
& \left. 7t_2 - \left(t_1 + \frac{13}{133}\right) \leq \frac{9}{7}, \frac{80}{21} \leq 6 \left(t_1 + \frac{13}{133}\right) + 5t_2 \right\}, \\
U_{37}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{131}{231}, 35(t_1 + t_2) + 23t_2 \leq \frac{125}{7}, \frac{200}{231} \leq 2 \left(t_1 + \frac{13}{133}\right) + t_2, \right. \\
& \left. 2(t_1 + t_2) + 13t_2 \leq \frac{64}{21}, \frac{240}{7} \leq 70 \left(t_1 + \frac{13}{133}\right) + 59t_2 \right\}, \\
U_{38}(t_1, t_2) &:= \left\{ (t_1 + t_2) + t_2 \leq \frac{51}{91}, 41(t_1 + t_2) + 27t_2 \leq 21, \frac{80}{91} \leq 2 \left(t_1 + \frac{13}{133}\right) + t_2, \right. \\
& \left. 2(t_1 + t_2) + 15t_2 \leq \frac{68}{21}, 40 \leq 82 \left(t_1 + \frac{13}{133}\right) + 69t_2 \right\}, \\
U_{39}(t_1, t_2) &:= \left\{ 2(t_1 + t_2) + t_2 \leq \frac{262}{231}, 70(t_1 + t_2) + 11t_2 \leq \frac{250}{7}, \frac{100}{231} \leq \left(t_1 + \frac{13}{133}\right), \right. \\
& \left. (t_1 + t_2) + 6t_2 \leq \frac{32}{21}, \frac{120}{7} \leq 35 \left(t_1 + \frac{13}{133}\right) + 12t_2 \right\}, \\
U_{310}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{47}{87}, \frac{40}{231} \leq t_1, 29(t_2 + t_2) - t_1 \leq \frac{209}{21}, \frac{40}{133} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
& \left. 29t_1 - \left(t_2 + \frac{13}{133}\right) \leq \frac{43}{7}, \frac{40}{7} \leq 12 \left(t_2 + \frac{13}{133}\right) + 11t_1 \right\}, \\
U_{311}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{109}{189}, 29(t_2 + t_2) + 19t_1 \leq \frac{103}{7}, \frac{160}{189} \leq 2 \left(t_2 + \frac{13}{133}\right) + t_1, \right. \\
& \left. 2(t_2 + t_2) + 11t_1 \leq \frac{20}{7}, \frac{200}{7} \leq 58 \left(t_2 + \frac{13}{133}\right) + 49t_1 \right\}, \\
U_{312}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{131}{231}, \frac{4}{21} \leq t_1, 6(t_2 + t_2) + t_1 \leq \frac{46}{21}, \frac{40}{147} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
& \left. (t_2 + t_2) + 8t_1 \leq \frac{16}{7}, \frac{20}{7} \leq 6 \left(t_2 + \frac{13}{133}\right) + 5t_1 \right\},
\end{aligned}$$

$$\begin{aligned}
U_{313}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{93}{133}, \frac{40}{231} \leq t_1, 12(t_2 + t_2) + t_1 \leq \frac{44}{7}, \frac{40}{87} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
&\quad \left. 19t_1 - \left(t_2 + \frac{13}{133}\right) \leq \frac{79}{21}, \frac{160}{21} \leq 12 \left(t_2 + \frac{13}{133}\right) + 11t_1 \right\}, \\
U_{314}(t_1, t_2) &:= \left\{ 2(t_2 + t_2) + t_1 \leq \frac{218}{189}, 58(t_2 + t_2) + 9t_1 \leq \frac{206}{7}, \frac{80}{189} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
&\quad \left. (t_2 + t_2) + 5t_1 \leq \frac{10}{7}, \frac{100}{7} \leq 29 \left(t_2 + \frac{13}{133}\right) + 10t_1 \right\}, \\
U_{315}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{107}{147}, \frac{4}{21} \leq t_1, 6(t_2 + t_2) + t_1 \leq \frac{22}{7}, \frac{100}{231} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
&\quad \left. 7t_1 - \left(t_2 + \frac{13}{133}\right) \leq \frac{9}{7}, \frac{80}{21} \leq 6 \left(t_2 + \frac{13}{133}\right) + 5t_1 \right\}, \\
U_{316}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{131}{231}, 35(t_2 + t_2) + 23t_1 \leq \frac{125}{7}, \frac{200}{231} \leq 2 \left(t_2 + \frac{13}{133}\right) + t_1, \right. \\
&\quad \left. 2(t_2 + t_2) + 13t_1 \leq \frac{64}{21}, \frac{240}{7} \leq 70 \left(t_2 + \frac{13}{133}\right) + 59t_1 \right\}, \\
U_{317}(t_1, t_2) &:= \left\{ (t_2 + t_2) + t_1 \leq \frac{51}{91}, 41(t_2 + t_2) + 27t_1 \leq 21, \frac{80}{91} \leq 2 \left(t_2 + \frac{13}{133}\right) + t_1, \right. \\
&\quad \left. 2(t_2 + t_2) + 15t_1 \leq \frac{68}{21}, 40 \leq 82 \left(t_2 + \frac{13}{133}\right) + 69t_1 \right\}, \\
U_{318}(t_1, t_2) &:= \left\{ 2(t_2 + t_2) + t_1 \leq \frac{262}{231}, 70(t_2 + t_2) + 11t_1 \leq \frac{250}{7}, \frac{100}{231} \leq \left(t_2 + \frac{13}{133}\right), \right. \\
&\quad \left. (t_2 + t_2) + 6t_1 \leq \frac{32}{21}, \frac{120}{7} \leq 35 \left(t_2 + \frac{13}{133}\right) + 12t_1 \right\}, \\
U_{319}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{47}{87}, \frac{40}{231} \leq \left(t_2 + \frac{13}{133}\right), 29t_1 - \left(t_2 + \frac{13}{133}\right) \leq \frac{209}{21}, \frac{40}{133} \leq t_1, \right. \\
&\quad \left. 29(t_2 + t_2) - t_1 \leq \frac{43}{7}, \frac{40}{7} \leq 12t_1 + 11 \left(t_2 + \frac{13}{133}\right) \right\}, \\
U_{320}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{109}{189}, 29t_1 + 19(t_2 + t_2) \leq \frac{103}{7}, \frac{160}{189} \leq 2t_1 + \left(t_2 + \frac{13}{133}\right), \right. \\
&\quad \left. 2t_1 + 11(t_2 + t_2) \leq \frac{20}{7}, \frac{200}{7} \leq 58t_1 + 49 \left(t_2 + \frac{13}{133}\right) \right\}, \\
U_{321}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{131}{231}, \frac{4}{21} \leq \left(t_2 + \frac{13}{133}\right), 6t_1 + (t_2 + t_2) \leq \frac{46}{21}, \frac{40}{147} \leq t_1, \right. \\
&\quad \left. t_1 + 8(t_2 + t_2) \leq \frac{16}{7}, \frac{20}{7} \leq 6t_1 + 5 \left(t_2 + \frac{13}{133}\right) \right\}, \\
U_{322}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{93}{133}, \frac{40}{231} \leq \left(t_2 + \frac{13}{133}\right), 12t_1 + (t_2 + t_2) \leq \frac{44}{7}, \frac{40}{87} \leq t_1, \right. \\
&\quad \left. 19(t_2 + t_2) - t_1 \leq \frac{79}{21}, \frac{160}{21} \leq 12t_1 + 11 \left(t_2 + \frac{13}{133}\right) \right\}, \\
U_{323}(t_1, t_2) &:= \left\{ 2t_1 + (t_2 + t_2) \leq \frac{218}{189}, 58t_1 + 9(t_2 + t_2) \leq \frac{206}{7}, \frac{80}{189} \leq t_1, \right. \\
&\quad \left. t_1 + 5(t_2 + t_2) \leq \frac{10}{7}, \frac{100}{7} \leq 29t_1 + 10 \left(t_2 + \frac{13}{133}\right) \right\}, \\
U_{324}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{107}{147}, \frac{4}{21} \leq \left(t_2 + \frac{13}{133}\right), 6t_1 + (t_2 + t_2) \leq \frac{22}{7}, \frac{100}{231} \leq t_1, \right.
\end{aligned}$$

$$\begin{aligned}
& \left. 7(t_2 + t_2) - t_1 \leq \frac{9}{7}, \frac{80}{21} \leq 6t_1 + 5 \left(t_2 + \frac{13}{133} \right) \right\}, \\
U_{325}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{131}{231}, 35t_1 + 23(t_2 + t_2) \leq \frac{125}{7}, \frac{200}{231} \leq 2t_1 + \left(t_2 + \frac{13}{133} \right), \right. \\
& \left. 2t_1 + 13(t_2 + t_2) \leq \frac{64}{21}, \frac{240}{7} \leq 70t_1 + 59 \left(t_2 + \frac{13}{133} \right) \right\}, \\
U_{326}(t_1, t_2) &:= \left\{ t_1 + (t_2 + t_2) \leq \frac{51}{91}, 41t_1 + 27(t_2 + t_2) \leq 21, \frac{80}{91} \leq 2t_1 + \left(t_2 + \frac{13}{133} \right), \right. \\
& \left. 2t_1 + 15(t_2 + t_2) \leq \frac{68}{21}, 40 \leq 82t_1 + 69 \left(t_2 + \frac{13}{133} \right) \right\}, \\
U_{327}(t_1, t_2) &:= \left\{ 2t_1 + (t_2 + t_2) \leq \frac{262}{231}, 70t_1 + 11(t_2 + t_2) \leq \frac{250}{7}, \frac{100}{231} \leq t_1, \right. \\
& \left. t_1 + 6(t_2 + t_2) \leq \frac{32}{21}, \frac{120}{7} \leq 35t_1 + 12 \left(t_2 + \frac{13}{133} \right) \right\}, \\
U_{41}(t_1, t_2) &:= \left\{ 2(t_1 + t_2) + t_2 \leq 2 - \frac{20}{21}, 4(t_1 + t_2) + 6t_2 \leq 4 - \frac{20}{21}, 6(t_1 + t_2) + 7t_2 \leq 6 - \frac{40}{21}, \right. \\
& \left. 4t_2 \leq 4 - \frac{20}{7}, 3t_2 - 2(t_1 + t_2) \leq 6 - \frac{40}{7} \right\}, \\
U_{42}(t_1, t_2) &:= \left\{ (t_1 + t_2) \leq \frac{11}{21}, t_2 \leq \frac{5}{42} \right\}, \\
U_{43}(t_1, t_2) &:= \left\{ \frac{80}{189} \leq (t_1 + t_2) \leq \frac{40}{87}, t_2 \leq -\frac{1}{5}(t_1 + t_2) + \frac{10}{35} \right\}, \\
U_{44}(t_1, t_2) &:= \left\{ \frac{40}{87} \leq (t_1 + t_2) \leq \frac{10}{21}, t_2 \leq \frac{1}{7}(t_1 + t_2) + \frac{9}{49} \right\}, \\
U_{45}(t_1, t_2) &:= \left\{ \frac{10}{21} \leq (t_1 + t_2) \leq \frac{37}{77}, t_2 \leq -(t_1 + t_2) + \frac{107}{147} \right\}, \\
U_{46}(t_1, t_2) &:= \left\{ \frac{37}{77} \leq (t_1 + t_2) \leq \frac{61}{126}, t_2 \leq -2(t_1 + t_2) + \frac{262}{231} \right\}, \\
U_{47}(t_1, t_2) &:= \left\{ \frac{61}{126} \leq (t_1 + t_2) \leq \frac{289}{588}, t_2 \leq -\frac{70}{11}(t_1 + t_2) + \frac{250}{77} \right\}, \\
U_{51}(t_1, t_2) &:= \left\{ 2(t_2 + t_2) + t_1 \leq 2 - \frac{20}{21}, 4(t_2 + t_2) + 6t_1 \leq 4 - \frac{20}{21}, 6(t_2 + t_2) + 7t_1 \leq 6 - \frac{40}{21}, \right. \\
& \left. 4t_1 \leq 4 - \frac{20}{7}, 3t_1 - 2 \left(t_2 + \frac{13}{133} \right) \leq 6 - \frac{40}{7} \right\}, \\
U_{52}(t_1, t_2) &:= \left\{ (t_2 + t_2) \leq \frac{11}{21}, t_1 \leq \frac{5}{42} \right\}, \\
U_{53}(t_1, t_2) &:= \left\{ \frac{80}{189} \leq \left(t_2 + \frac{13}{133} \right), (t_2 + t_2) \leq \frac{40}{87}, t_1 \leq -\frac{1}{5}(t_2 + t_2) + \frac{10}{35} \right\}, \\
U_{54}(t_1, t_2) &:= \left\{ \frac{40}{87} \leq \left(t_2 + \frac{13}{133} \right), (t_2 + t_2) \leq \frac{10}{21}, t_1 \leq \frac{1}{7} \left(t_2 + \frac{13}{133} \right) + \frac{9}{49} \right\}, \\
U_{55}(t_1, t_2) &:= \left\{ \frac{10}{21} \leq \left(t_2 + \frac{13}{133} \right), (t_2 + t_2) \leq \frac{37}{77}, t_1 \leq -(t_2 + t_2) + \frac{107}{147} \right\}, \\
U_{56}(t_1, t_2) &:= \left\{ \frac{37}{77} \leq \left(t_2 + \frac{13}{133} \right), (t_2 + t_2) \leq \frac{61}{126}, t_1 \leq -2(t_2 + t_2) + \frac{262}{231} \right\}, \\
U_{57}(t_1, t_2) &:= \left\{ \frac{61}{126} \leq \left(t_2 + \frac{13}{133} \right), (t_2 + t_2) \leq \frac{289}{588}, t_1 \leq -\frac{70}{11}(t_2 + t_2) + \frac{250}{77} \right\},
\end{aligned}$$

and let $U_0 = U_1 \cup U_2 \cup U_3 \cup U_4 \cup U_5$. The plot of the 5 regions can be found in Appendix. Then we have

$$\begin{aligned}
S_3 &= \sum_{(t_1, t_2) \in U_1} S(\mathcal{A}_{p_1 p_2}, p_2) + \sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&+ \sum_{(t_1, t_2) \in U_3} S(\mathcal{A}_{p_1 p_2}, p_2) + \sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&+ \sum_{(t_1, t_2) \in U_5} S(\mathcal{A}_{p_1 p_2}, p_2) + \sum_{(t_1, t_2) \notin U_0} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= S_{31} + S_{32} + S_{33} + \Sigma_4 + \Sigma_5 + \Sigma_0.
\end{aligned} \tag{29}$$

For S_{31} , by Lemma 3.2 and Lemma 6.2 we have

$$\begin{aligned}
S_{31} &= \sum_{(t_1, t_2) \in U_1} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= \eta \sum_{(t_1, t_2) \in U_1} S(\mathcal{B}_{p_1 p_2}, p_2) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&= \frac{\eta x}{\log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \text{Boole}[(t_1, t_2) \in U_1] \frac{\omega\left(\frac{1-t_1-t_2}{t_2}\right)}{t_1 t_2^2} dt_2 dt_1 \\
&\geq 0.806116 \frac{\eta x}{\log x}.
\end{aligned} \tag{30}$$

For S_{32} , we can apply Buchstab's identity again to get

$$\begin{aligned}
S_{32} &= \sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= \sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_2 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_2 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3).
\end{aligned} \tag{31}$$

By Lemma 5.1 and Iwaniec's linear sieve with $z = X^\delta$ and $D = \frac{X^{\frac{11}{21}}}{p_1 p_2}$, we have

$$\sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, X^\delta) = \eta \sum_{(t_1, t_2) \in U_2} S(\mathcal{B}_{p_1 p_2}, X^\delta) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{32}$$

By Lemma 6.1 we have

$$\sum_{\substack{(t_1, t_2) \in U_2 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) = \eta \sum_{\substack{(t_1, t_2) \in U_2 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{33}$$

By Lemma 3.3 and Lemma 6.2 with a small modification we have

$$\sum_{\substack{(t_1, t_2) \in U_2 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) = \eta \sum_{\substack{(t_1, t_2) \in U_2 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{34}$$

Combining together (31)–(34), we have

$$S_{32} = \sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, p_2)$$

$$\begin{aligned}
&= \sum_{(t_1, t_2) \in U_2} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_2 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_2 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&= \eta \left(\sum_{(t_1, t_2) \in U_2} S(\mathcal{B}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_2 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) \right. \\
&\quad \left. - \sum_{\substack{(t_1, t_2) \in U_2 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&= \eta \sum_{(t_1, t_2) \in U_2} S(\mathcal{B}_{p_1 p_2}, p_2) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&= \frac{\eta x}{\log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \text{Boole}[(t_1, t_2) \in U_2] \frac{\omega\left(\frac{1-t_1-t_2}{t_2}\right)}{t_1 t_2^2} dt_2 dt_1 \\
&\geq 0.496315 \frac{\eta x}{\log x}. \tag{35}
\end{aligned}$$

Similarly, by replacing Lemma 3.3 with Lemmas 3.4, 3.5 and 3.6 in above discussion, for S_{33} we have

$$\begin{aligned}
S_{33} &= \sum_{(t_1, t_2) \in U_3} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= \sum_{(t_1, t_2) \in U_3} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_3 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_3 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&= \eta \left(\sum_{(t_1, t_2) \in U_3} S(\mathcal{B}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_3 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) \right. \\
&\quad \left. - \sum_{\substack{(t_1, t_2) \in U_3 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&= \eta \sum_{(t_1, t_2) \in U_3} S(\mathcal{B}_{p_1 p_2}, p_2) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&= \frac{\eta x}{\log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \text{Boole}[(t_1, t_2) \in U_3] \frac{\omega\left(\frac{1-t_1-t_2}{t_2}\right)}{t_1 t_2^2} dt_2 dt_1 \\
&\geq 0.018318 \frac{\eta x}{\log x}. \tag{36}
\end{aligned}$$

We note that there is an extremely small region of U_3 with $t_1 + t_2 > \frac{11}{21}$. Fortunately we have $t_1 < \frac{1}{2}$ and $t_2 < \frac{5}{42}$ in this region. By this we can use Lemma 5.1 and Iwaniec's linear sieve with $z = X^\delta$ and $D = X^{\frac{1}{42}}$ to give asymptotic formulas.

For Σ_4 and Σ_5 , we can apply Buchstab's identity two more times to get

$$\begin{aligned}
\Sigma_4 &= \sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= \sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&= \left(\sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \right. \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, X^\delta) \\
&\quad \left. + \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \right) \\
&\quad + \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3))}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
&= S_{34} + \Sigma_{61} \tag{37}
\end{aligned}$$

and

$$\begin{aligned}
\Sigma_5 &= \sum_{(t_1, t_2) \in U_5} S(\mathcal{A}_{p_1 p_2}, p_2) \\
&= \sum_{(t_1, t_2) \in U_5} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\
&= \left(\sum_{(t_1, t_2) \in U_5} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \right) \\
&\quad - \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, X^\delta)
\end{aligned}$$

$$\begin{aligned}
& \left. \begin{aligned}
& + \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
& + \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3))}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4)
\end{aligned} \right) \\
& = S_{35} + \Sigma_{62}. \tag{38}
\end{aligned}$$

By Lemma 5.1 and Iwaniec's linear sieve with $z = X^\delta$ and $D = \frac{X^{\frac{11}{21}}}{p_1 p_2}$, we have

$$\sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, X^\delta) = \eta \sum_{(t_1, t_2) \in U_4} S(\mathcal{B}_{p_1 p_2}, X^\delta) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{39}$$

By shortening the length of the sum interval of t_1 in Σ_4 by 10^{-100} , using Iwaniec's linear sieve with $z = X^\delta$, $D = X^{10^{-100}}$ and by Lemma 5.2, we have

$$\sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, X^\delta) = \eta \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, X^\delta) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{40}$$

By Lemma 6.1 we have

$$\sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) = \eta \sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \tag{41}$$

and

$$\sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) = \eta \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3 p_4}, p_4) + O\left(\frac{\varepsilon \eta x}{\log x}\right). \tag{42}$$

Combining together (39)–(42) and by (37), we have

$$\begin{aligned}
S_{34} &= \sum_{(t_1, t_2) \in U_4} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, X^\delta) \\
&+ \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
&= \eta \left(\sum_{(t_1, t_2) \in U_4} S(\mathcal{B}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, X^\delta) \right)
\end{aligned}$$

$$\begin{aligned}
& + \left. \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3 p_4}, p_4) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
& = \eta \left(\sum_{(t_1, t_2) \in U_4} S(\mathcal{B}_{p_1 p_2}, X_{\frac{13}{133}}) - \sum_{\substack{(t_1, t_2) \in U_4 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, X_{\frac{13}{133}}) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
& \geq \frac{\eta x}{\log x} \left(\int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \text{Boole}[(t_1, t_2) \in U_4] \frac{\frac{133}{13} \omega\left(\frac{133}{13}(1-t_1-t_2)\right)}{t_1 t_2} dt_2 dt_1 \right. \\
& \quad \left. - \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{\frac{13}{133}}^{t_2} \text{Boole}[(t_1, t_2) \in U_4] \frac{\frac{133}{13} \omega\left(\frac{133}{13}(1-t_1-t_2-t_3)\right)}{t_1 t_2 t_3} dt_3 dt_2 dt_1 - 10^{-10} \right) \\
& \geq 2.212979 \frac{\eta x}{\log x} \tag{43}
\end{aligned}$$

where the total loss of shortening the sum interval of t_1 is less than 10^{-10} . Similarly, for S_{35} we have

$$\begin{aligned}
S_{35} & = \sum_{(t_1, t_2) \in U_5} S(\mathcal{A}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{A}_{p_1 p_2 p_3}, X^\delta) \\
& + \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
& = \eta \left(\sum_{(t_1, t_2) \in U_5} S(\mathcal{B}_{p_1 p_2}, X^\delta) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \delta \leq t_3 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, X^\delta) \right. \\
& \quad \left. + \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \delta \leq t_4 < \frac{13}{133}}} S(\mathcal{B}_{p_1 p_2 p_3 p_4}, p_4) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
& = \eta \left(\sum_{(t_1, t_2) \in U_5} S(\mathcal{B}_{p_1 p_2}, X_{\frac{13}{133}}) - \sum_{\substack{(t_1, t_2) \in U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2))}} S(\mathcal{B}_{p_1 p_2 p_3}, X_{\frac{13}{133}}) \right) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
& \geq \frac{\eta x}{\log x} \left(\int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \text{Boole}[(t_1, t_2) \in U_5] \frac{\frac{133}{13} \omega\left(\frac{133}{13}(1-t_1-t_2)\right)}{t_1 t_2} dt_2 dt_1 \right. \\
& \quad \left. - \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{\frac{13}{133}}^{t_2} \text{Boole}[(t_1, t_2) \in U_5] \frac{\frac{133}{13} \omega\left(\frac{133}{13}(1-t_1-t_2-t_3)\right)}{t_1 t_2 t_3} dt_3 dt_2 dt_1 - 10^{-10} \right) \\
& \geq 0.006401 \frac{\eta x}{\log x}. \tag{44}
\end{aligned}$$

Let $\Sigma_6 = \Sigma_{61} + \Sigma_{62}$ where Σ_{61} and Σ_{62} are defined in (37) and (38). We can give asymptotic formulas for part of Σ_6 if we can group t_1, t_2, t_3, t_4 into m, h satisfy Lemma 3.2. We define this part as S_{36} . By Lemma 6.2 with a small modification, we get a saving on S_{36} :

$$\begin{aligned}
S_{36} &= \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ can be partitioned into } (m, h) \in U_1}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
&= \eta \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ can be partitioned into } (m, h) \in U_1}} S(\mathcal{B}_{p_1 p_2 p_3 p_4}, p_4) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\
&\geq \frac{\eta x}{\log x} \left(\int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{\frac{13}{133}}^{\min(t_2, \frac{1-t_1-t_2}{2})} \int_{\frac{13}{133}}^{\min(t_3, \frac{1-t_1-t_2-t_3}{2})} \right. \\
&\quad \left. \text{Boole}[(t_1, t_2, t_3, t_4) \in U_6] \frac{\omega_0\left(\frac{1-t_1-t_2-t_3-t_4}{t_4}\right)}{t_1 t_2 t_3 t_4^2} dt_4 dt_3 dt_2 dt_1 \right) \\
&\geq 0.108021 \frac{\eta x}{\log x}, \tag{45}
\end{aligned}$$

where $U_6(t_1, t_2, t_3, t_4)$ is defined as

$$\begin{aligned}
U_6(t_1, t_2, t_3, t_4) &:= \left\{ (t_1, t_2) \in U_4 \cup U_5, \frac{13}{133} \leq t_3 < \min\left(t_2, \frac{1}{2}(1-t_1-t_2)\right), \right. \\
&\quad \frac{13}{133} \leq t_4 < \min\left(t_3, \frac{1}{2}(1-t_1-t_2-t_3)\right), \\
&\quad (t_1, t_2, t_3, t_4) \text{ can be partitioned into } (m, h) \in U_1, \\
&\quad \left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right) \right\}.
\end{aligned}$$

Let $\Sigma_7 = \Sigma_6 - S_{36}$. We can use Buchstab's identity in reverse to get asymptotic formulas for some almost-primes counted in Σ_7 . More specifically, when $t_4 < \frac{1}{2}(1-t_1-t_2-t_3-t_4)$, we have

$$\begin{aligned}
&\sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < \frac{1}{2}(1-t_1-t_2-t_3-t_4)}} S(\mathcal{A}_{p_1 p_2 p_3 p_4}, p_4) \\
&= \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < \frac{1}{2}(1-t_1-t_2-t_3-t_4)}} S\left(\mathcal{A}_{p_1 p_2 p_3 p_4}, \left(\frac{2X}{p_1 p_2 p_3 p_4}\right)^{\frac{1}{2}}\right) \\
&+ \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4)}} S(\mathcal{A}_{p_1 p_2 p_3 p_4 p_5}, p_5). \tag{46}
\end{aligned}$$

We can give asymptotic formulas for part of the last term if we can group t_1, t_2, t_3, t_4, t_5 into m, h satisfy Lemma 3.2. We define this part as S_{37} . More specifically,

$$S_{37} = \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4) \\ (t_1, t_2, t_3, t_4, t_5) \text{ can be partitioned into } (m, h) \in U_1}} S(\mathcal{A}_{p_1 p_2 p_3 p_4 p_5}, p_5). \quad (47)$$

In this way, by Lemma 6.2 with a small modification, we obtain a small saving on S_{37} :

$$\begin{aligned} S_{37} &\geq \frac{\eta x}{\log x} \left(\int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{\frac{13}{133}}^{\min(t_2, \frac{1-t_1-t_2}{2})} \int_{\frac{13}{133}}^{\min(t_3, \frac{1-t_1-t_2-t_3}{2})} \int_{t_4}^{\frac{1-t_1-t_2-t_3-t_4}{2}} \right. \\ &\quad \left. \text{Boole}[(t_1, t_2, t_3, t_4, t_5) \in U_7] \frac{\omega_0 \left(\frac{1-t_1-t_2-t_3-t_4-t_5}{t_5} \right)}{t_1 t_2 t_3 t_4 t_5^2} dt_5 dt_4 dt_3 dt_2 dt_1 \right) \\ &\geq 0.003994 \frac{\eta x}{\log x}, \end{aligned} \quad (48)$$

where $U_7(t_1, t_2, t_3, t_4, t_5)$ is defined as

$$\begin{aligned} U_7(t_1, t_2, t_3, t_4, t_5) := &\left\{ (t_1, t_2) \in U_4 \cup U_5, \frac{13}{133} \leq t_3 < \min\left(t_2, \frac{1}{2}(1-t_1-t_2)\right), \right. \\ &\frac{13}{133} \leq t_4 < \min\left(t_3, \frac{1}{2}(1-t_1-t_2-t_3)\right), \\ &(t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1, \\ &t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4), \\ &(t_1, t_2, t_3, t_4, t_5) \text{ can be partitioned into } (m, h) \in U_1, \\ &\left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right) \right\}. \end{aligned}$$

We can apply Buchstab's identity in reverse again to the remaining sums of the last term in (46). By Lemma 6.2 with a small modification, together with the above discussion, we have

$$\begin{aligned} S_{38} &= \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4) \\ (t_1, t_2, t_3, t_4, t_5) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_5 < t_6 < \frac{1}{2}(1-t_1-t_2-t_3-t_4-t_5) \\ (t_1, t_2, t_3, t_4, t_5, t_6) \text{ can be partitioned into } (m, h) \in U_1}} S(\mathcal{A}_{p_1 p_2 p_3 p_4 p_5 p_6}, p_6) \\ &= \eta \sum_{\substack{(t_1, t_2) \in U_4 \cup U_5 \\ \frac{13}{133} \leq t_3 < \min(t_2, \frac{1}{2}(1-t_1-t_2)) \\ \frac{13}{133} \leq t_4 < \min(t_3, \frac{1}{2}(1-t_1-t_2-t_3)) \\ (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4) \\ (t_1, t_2, t_3, t_4, t_5) \text{ cannot be partitioned into } (m, h) \in U_1 \\ t_5 < t_6 < \frac{1}{2}(1-t_1-t_2-t_3-t_4-t_5) \\ (t_1, t_2, t_3, t_4, t_5, t_6) \text{ can be partitioned into } (m, h) \in U_1}} S(\mathcal{B}_{p_1 p_2 p_3 p_4 p_5 p_6}, p_6) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\ &\geq \frac{\eta x}{\log x} \left(\int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{\frac{13}{133}}^{\min(t_2, \frac{1-t_1-t_2}{2})} \int_{\frac{13}{133}}^{\min(t_3, \frac{1-t_1-t_2-t_3}{2})} \int_{t_4}^{\frac{1-t_1-t_2-t_3-t_4}{2}} \int_{t_5}^{\frac{1-t_1-t_2-t_3-t_4-t_5}{2}} \right. \end{aligned}$$

$$\begin{aligned} & \text{Boole}[(t_1, t_2, t_3, t_4, t_5, t_6) \in U_8] \frac{\omega_0\left(\frac{1-t_1-t_2-t_3-t_4-t_5-t_6}{t_6}\right)}{t_1 t_2 t_3 t_4 t_5 t_6^2} dt_6 dt_5 dt_4 dt_3 dt_2 dt_1 \\ & \geq 0.000000000486547 \frac{\eta x}{\log x}, \end{aligned} \quad (49)$$

where $U_8(t_1, t_2, t_3, t_4, t_5, t_6)$ is defined as

$$\begin{aligned} U_8(t_1, t_2, t_3, t_4, t_5, t_6) := & \left\{ (t_1, t_2) \in U_4 \cup U_5, \frac{13}{133} \leq t_3 < \min\left(t_2, \frac{1}{2}(1-t_1-t_2)\right), \right. \\ & \frac{13}{133} \leq t_4 < \min\left(t_3, \frac{1}{2}(1-t_1-t_2-t_3)\right), \\ & (t_1, t_2, t_3, t_4) \text{ cannot be partitioned into } (m, h) \in U_1, \\ & t_4 < t_5 < \frac{1}{2}(1-t_1-t_2-t_3-t_4), \\ & (t_1, t_2, t_3, t_4, t_5) \text{ cannot be partitioned into } (m, h) \in U_1, \\ & t_5 < t_6 < \frac{1}{2}(1-t_1-t_2-t_3-t_4-t_5), \\ & (t_1, t_2, t_3, t_4, t_5, t_6) \text{ can be partitioned into } (m, h) \in U_1, \\ & \left. \frac{13}{133} \leq t_1 < \frac{1}{2}, \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right) \right\}. \end{aligned}$$

For the remaining Σ_0 , we can also use Buchstab's identity in reverse to get asymptotic formulas for some almost-primes counted. When $t_2 < \frac{1}{2}(1-t_1-t_2)$, we have

$$\sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < \frac{1}{2}(1-t_1-t_2)}} S(\mathcal{A}_{p_1 p_2}, p_2) = \sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < \frac{1}{2}(1-t_1-t_2)}} S\left(\mathcal{A}_{p_1 p_2}, \left(\frac{2X}{p_1 p_2}\right)^{\frac{1}{2}}\right) + \sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < t_3 < \frac{1}{2}(1-t_1-t_2)}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3). \quad (50)$$

Then, we can give asymptotic formulas for part of the last term if we can group t_1, t_2, t_3 into m, h satisfy Lemma 3.2. We define this part as S_{39} .

$$S_{39} = \sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < t_3 < \frac{1}{2}(1-t_1-t_2) \\ (t_1+t_2, t_3) \in U_1 \text{ or } (t_1+t_3, t_2) \in U_1 \text{ or } (t_2+t_3, t_1) \in U_1}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3). \quad (51)$$

Then by Lemma 6.2 with a small modification, we can get a small saving on S_{39} :

$$\begin{aligned} S_{39} &= \sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < t_3 < \frac{1}{2}(1-t_1-t_2) \\ (t_1+t_2, t_3) \in U_1 \text{ or } (t_1+t_3, t_2) \in U_1 \text{ or } (t_2+t_3, t_1) \in U_1}} S(\mathcal{A}_{p_1 p_2 p_3}, p_3) \\ &= \sum_{\substack{(t_1, t_2) \notin U_0 \\ t_2 < t_3 < \frac{1}{2}(1-t_1-t_2) \\ (t_1+t_2, t_3) \in U_1 \text{ or } (t_1+t_3, t_2) \in U_1 \text{ or } (t_2+t_3, t_1) \in U_1}} S(\mathcal{B}_{p_1 p_2 p_3}, p_3) + O\left(\frac{\varepsilon \eta x}{\log x}\right) \\ &= \frac{\eta x}{\log x} \int_{\frac{13}{133}}^{\frac{1}{2}} \int_{\frac{13}{133}}^{\min(t_1, \frac{1-t_1}{2})} \int_{t_2}^{\frac{1-t_1-t_2}{2}} \text{Boole}[(t_1, t_2, t_3) \in U_9] \frac{\omega\left(\frac{1-t_1-t_2-t_3}{t_3}\right)}{t_1 t_2 t_3^2} dt_3 dt_2 dt_1 \\ &\geq 0.020591 \frac{\eta x}{\log x}, \end{aligned} \quad (52)$$

where $U_9(t_1, t_2, t_3)$ is defined as

$$\begin{aligned} U_9(t_1, t_2, t_3) := & \left\{ t_2 < t_3 < \frac{1}{2}(1-t_1-t_2), (t_1, t_2) \notin U_0, \right. \\ & \left. (t_1+t_2, t_3) \in U_1 \text{ or } (t_1+t_3, t_2) \in U_1 \text{ or } (t_2+t_3, t_1) \in U_1, \right. \end{aligned}$$

$$\frac{13}{133} \leq t_1 < \frac{1}{2}, \quad \frac{13}{133} \leq t_2 < \min\left(t_1, \frac{1}{2}(1-t_1)\right)\}.$$

We remark that an extra use of Buchstab's identity in reverse on the remaining sums of the last term in (50) leads to no more saving.

Finally, by (1)–(2), (26), (28)–(30), (35)–(36), (43)–(45), (48)–(49) and (52) we have

$$\pi(x + \eta x) - \pi(x) = S\left(\mathcal{A}, (2X)^{\frac{1}{2}}\right) \geq 0.04 \frac{\eta x}{\log x},$$

and the proof of Theorem 1.1 is completed. Theorem 1.2 can be proved by replacing [[12], Lemma 4] by a variant of our Theorem 1.1. In this way, the ratio $\frac{20}{19}$ in [[12], Theorem 4] can be improved to $\frac{21}{20}$.

8. FURTHER PROSPECT

The main aim of the present paper is to give a simplified proof of Jia's theorem, so we don't consider further optimization. Here we point out some possible ways for improving the exponent $\frac{1}{21}$.

Just like in [25], we use only classical mean value estimate and Halász method to give arithmetic information in section 3. Very recently, Guth and Maynard [8] proved a new large value estimate for Dirichlet polynomials. One can employ this bound together with the Montgomery–Halász–Huxley bound to get more arithmetic information, and that would be of great benefit.

As we mentioned in Section 4, our proof uses Deshouillers–Iwaniec's theorem instead of Watt's theorem for the approximation of the sixth power moment of Riemann zeta function. In fact, the condition $M^{-2}H^3 \ll X^{6-\frac{40}{7}}$ in Lemma 4.2 can be removed if we follow the steps of the proof of [[2], Lemma 2] to get an analog of Lemma 4.1. However, removing this condition directly cannot provide us with more type-I information. One can try to use Watt's theorem in a more delicate way to obtain more type-I information and get an improvement of our exponent. Using [[7], Theorem 2] is another possible way to obtain more type-I information. For more type-II information, one can modify the methods used in [[2], Section 5] to get a similar result of [[2], Lemma 18] with $h = 2$.

Our asymptotic regions and sieve decompositions can be further optimized. For example, our definition of U_3 considers U_{31} – U_{327} separately, which means more arithmetic information may be obtainable by combining these subregions together (as we did on U_2). One can use the following code to give a unified definition of U_3 :

```
U3[t1_, t2_] := Resolve[ForAll[t3, SieveExponent <= t3 < t2, U1[t1 + t3, t2] || U1[t2 + t3, t1] || U1[t1, t2 + t3] || U1[t2, t1 + t3]]]
```

And calculating the integrals related to this region requires more powerful computers. Another example is that one can apply Buchstab's identity two more times on part of Σ_7 when $(t_1, t_2, t_3, t_4, t_4)$ can be grouped into (m, h) satisfies one of the conditions in Lemma 4.2 or the "extra Type-I region" (see Figure 3 in Appendix).

The present paper doesn't contain role-reversals in the sieve decompositions. In a role-reversal, one has the following formula for some region K satisfies $t_1 > 1 - t_1 - t_2$:

$$\begin{aligned} \sum_{(t_1, t_2) \in K} S(\mathcal{A}_{p_1 p_2}, p_2) &= \sum_{(t_1, t_2) \in K} S\left(\mathcal{A}_{h p_2}, \left(\frac{2X}{h p_2}\right)^{\frac{1}{2}}\right) \\ &= \sum_{(t_1, t_2) \in K} S\left(\mathcal{A}_{h p_2}, X^{\frac{13}{133}}\right) - \sum_{\substack{(t_1, t_2) \in K \\ \frac{13}{133} \leq t_3 < \frac{1}{2} t_1}} S(\mathcal{A}_{h p_2 p_3}, p_3), \end{aligned} \quad (53)$$

where $h = X^{1-t_1-t_2}$. Note that the left hand side of (53) counts $p_1 p_2 h$ while the right hand side counts $h p_2 p_1$. This device can also be used on triple sums $\sum S(\mathcal{A}_{p_1 p_2 p_3}, p_3)$ to produce positive quadruple sums. Harman used role-reversals on triple sums in his simplified proof of exponent $\frac{1}{18}$. We refer the readers to [27] for more applications of role-reversal. With this device, further decompositions are possible in more subregions.

The last thing is that one can use other sieve identities and inequalities other than Buchstab's identity. For example, Heath–Brown [14] used his own identity (see [[13], (2.5.6)]) in his proof of primes in all intervals

with $\theta = \frac{7}{12}$, and Brady [3] proved lots of sieve inequalities in his PhD thesis. Here we provide one of Brady's inequalities as the end of this section. For any $w \leq z$, we have

$$S(\mathcal{A}, z) \leq S(\mathcal{A}, w) - \frac{2}{3} \sum_{w \leq p_1 < z} S(\mathcal{A}_{p_1}, w) + \frac{1}{3} \sum_{w \leq p_2 < p_1 < z} S(\mathcal{A}_{p_1 p_2}, w).$$

ACKNOWLEDGEMENTS

The author would like to thank Professor Chaohua Jia and Professor Nigel Watt for some helpful discussions.

REFERENCES

- [1] R. C. Baker and G. Harman. The difference between consecutive primes. *Proc. London Math. Soc.*, 72(3):261–280, 1996.
- [2] R. C. Baker, G. Harman, and J. Pintz. The difference between consecutive primes, II. *Proc. London Math. Soc.*, 83(3):532–562, 2001.
- [3] Z. E. Brady. Sieves and iteration rules. *Ph.D. Thesis*, Stanford University, 2017.
- [4] N. G. Chudakov. On zeros of Dirichlet L-function. *Rec. Math. [Mat. Sbornik] N. S.*, 43(1):591–602, 1936.
- [5] H. Cramér. On the order of magnitude of the difference between consecutive prime numbers. *Acta Arith.*, 2:23–46, 1937.
- [6] J.-M. Deshouillers and H. Iwaniec. Power mean value of the Riemann zeta-function. *Mathematika*, 29:202–212, 1982.
- [7] J.-M. Deshouillers and H. Iwaniec. Power mean value of the Riemann zeta-function, II. *Acta Arith.*, 43(3):305–312, 1984.
- [8] L. Guth and J. Maynard. New large value estimates for Dirichlet polynomials. *arXiv e-prints*, page arXiv:2405.20552, May 2024.
- [9] G. Harman. Primes in short intervals. *Math. Z.*, 180:335–348, 1982.
- [10] G. Harman. On the distribution of αp modulo one. *J. London Math. Soc.*, 27(2):9–18, 1983.
- [11] G. Harman. On the distribution of αp modulo one II. *Proc. London Math. Soc.*, 72(3):241–260, 1996.
- [12] G. Harman. Metrical theorems on prime values of the integer parts of real sequences II. *J. London Math. Soc.*, 64(2):287–298, 2001.
- [13] G. Harman. *Prime-detecting Sieves*, volume 33 of *London Mathematical Society Monographs (New Series)*. Princeton University Press, Princeton, NJ, 2007.
- [14] D. R. Heath-Brown. Sieve identities and gaps between primes. *Astérisque.*, 94:61–65, 1982.
- [15] D. R. Heath-Brown and H. Iwaniec. On the difference between consecutive primes. *Invent. Math.*, 55:49–69, 1979.
- [16] H. Heilbronn. Über den Primzahlsatz von Herrn Hoheisel. *Math. Z.*, 36:394–423, 1933.
- [17] G. Hoheisel. Primzahlprobleme in der analysis. *Sitz. Preuss. Akad. Wiss.*, 2:1–13, 1930.
- [18] M. N. Huxley. On the difference between consecutive primes. *Invent. Math.*, 15:164–170, 1972.
- [19] A. E. Ingham. On the difference between consecutive primes. *Q. J. Math.*, 8:255–266, 1936.
- [20] H. Iwaniec and M. Jutila. Primes in short intervals. *Ark. Mat.*, 17:167–176, 1979.
- [21] H. Iwaniec and J. Pintz. Primes in short intervals. *Monatsh. Math.*, 98:115–143, 1984.
- [22] C. Jia. Difference between consecutive primes. *Sci. China Ser. A*, 38(10):1163–1186, 1995.
- [23] C. Jia. Goldbach numbers in short interval (I). *Sci. China Ser. A*, 38(4):385–406, 1995.
- [24] C. Jia. Goldbach numbers in short interval (II). *Sci. China Ser. A*, 38(5):513–523, 1995.
- [25] C. Jia. Almost all short intervals containing prime numbers. *Acta Arith.*, 76(1):21–84, 1996.
- [26] C. Jia. On the exceptional set of goldbach numbers in a short interval. *Acta Arith.*, 77(3):207–287, 1996.
- [27] C. Jia and M.-C. Liu. On the largest prime factor of integers. *Acta Arith.*, 95(1):17–48, 2000.
- [28] H. Li. Goldbach numbers in short intervals. *Sci. China Ser. A*, 38(6):641–652, 1995.
- [29] H. Li. Primes in short intervals. *Math. Proc. Camb. Phil. Soc.*, 122(2):193–205, 1997.
- [30] S. Lou and Q. Yao. A Chebyshev's type of prime number theorem in a short interval–II. *Hardy-Ramanujan J.*, 15:1–33, 1992.
- [31] S. Lou and Q. Yao. The number of primes in a short interval. *Hardy-Ramanujan J.*, 16:21–43, 1993.
- [32] S. Lou and Q. Yao. Estimate of sums of Dirichlet series. *Hardy-Ramanujan J.*, 17:1–31, 1994.
- [33] H. L. Montgomery. *Topics in Multiplicative Number Theory*. Lecture Notes in Math. 227. Springer, Berlin, 1971.
- [34] C. J. Mozzochi. On the difference between consecutive primes. *J. Number Theory*, 24:181–187, 1986.
- [35] J. Pintz. On primes in short intervals I. *Studia Sci. Math. Hungar.*, 16:395–414, 1981.
- [36] J. Pintz. On primes in short intervals II. *Studia Sci. Math. Hungar.*, 19:89–96, 1984.
- [37] A. Selberg. On the normal density of primes in short intervals, and the difference between consecutive primes. *Arch. Math. Naturvid.*, 47:87–105, 1943.
- [38] N. Watt. Kloosterman sums and a mean value for Dirichlet polynomials. *J. Number Theory*, 53:179–210, 1995.
- [39] N. Watt. Short intervals almost all containing primes. *Acta Arith.*, 72(2):131–167, 1995.
- [40] K. C. Wong. Contribution to analytic number theory. *Ph.D. Thesis*, Cardiff, 1996.

APPENDIX: MATHEMATICA CODE FOR THE THEOREM

Code for the asymptotic regions. The following Mathematica code shows the definition of the disjoint regions U_1-U_9 . In above calculations, we can give asymptotic formulas for all sums $S(\mathcal{A}_{p_1 p_2, p_2})$ where t_1 and t_2 lie in U_1, U_2 or U_3 . For the remaining sums which lie in U_4 or U_5 , we can give asymptotic formulas for part of these sums.

```
(*Our sieve exponent is 13/133*)
SieveExponent = 13/133;

(*The definition of the region U1*)
U11[t1_, t2_] := (t1 + t2 <= 47/87) && (40/231 <= t2) && (29*t1 - t2 <= 209/21) && (40/133
<= t1) && (29*t2 - t1 <= 43/7) && (40/7 <= 12*t1 + 11*t2)
U12[t1_, t2_] := (t1 + t2 <= 109/189) && (29*t1 + 19*t2 <= 103/7) && (160/189 <= 2*t1 + t2)
&& (2*t1 + 11*t2 <= 20/7) && (200/7 <= 58*t1 + 49*t2)
U13[t1_, t2_] := (t1 + t2 <= 131/231) && (4/21 <= t2) && (6*t1 + t2 <= 46/21) && (40/147 <=
t1) && (t1 + 8*t2 <= 16/7) && (20/7 <= 6*t1 + 5*t2)
U14[t1_, t2_] := (t1 + t2 <= 93/133) && (40/231 <= t2) && (12*t1 + t2 <= 44/7) && (40/87 <=
t1) && (19*t2 - t1 <= 79/21) && (160/21 <= 12*t1 + 11*t2)
U15[t1_, t2_] := (2*t1 + t2 <= 218/189) && (58*t1 + 9*t2 <= 206/7) && (80/189 <= t1) && (t1
+ 5*t2 <= 10/7) && (100/7 <= 29*t1 + 10*t2)
U16[t1_, t2_] := (t1 + t2 <= 107/147) && (4/21 <= t2) && (6*t1 + t2 <= 22/7) && (100/231 <=
t1) && (7*t2 - t1 <= 9/7) && (80/21 <= 6*t1 + 5*t2)
U17[t1_, t2_] := (t1 + t2 <= 131/231) && (35*t1 + 23*t2 <= 125/7) && (200/231 <= 2*t1 + t2)
&& (2*t1 + 13*t2 <= 64/21) && (240/7 <= 70*t1 + 59*t2)
U18[t1_, t2_] := (t1 + t2 <= 51/91) && (41*t1 + 27*t2 <= 21) && (80/91 <= 2*t1 + t2) && (2*
t1 + 15*t2 <= 68/21) && (40 <= 82*t1 + 69*t2)
U19[t1_, t2_] := (2*t1 + t2 <= 262/231) && (70*t1 + 11*t2 <= 250/7) && (100/231 <= t1) && (
t1 + 6*t2 <= 32/21) && (120/7 <= 35*t1 + 12*t2)
U1[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)])
&& (U11[t1, t2] || U12[t1, t2] || U13[t1, t2] || U14[t1, t2] || U15[t1, t2] || U16[t1,
t2] || U17[t1, t2] || U18[t1, t2] || U19[t1, t2])

(*The definition of the region U2*)
U21[t1_, t2_] := (4423/10906 <= t1 + t2 <= 113/273) && (SieveExponent <= t2 <= -(35/23) (t1
+ t2) + 125/161)
U22[t1_, t2_] := (113/273 <= t1 + t2 <= 267/637) && (SieveExponent <= t2 <= -(t1 + t2) +
51/91)
U23[t1_, t2_] := (267/637 <= t1 + t2 <= 2442/5453) && (SieveExponent <= t2 <= -(41/27) (t1 +
t2) + 21/27)
U24[t1_, t2_] := (2124/4655 <= t1 + t2 <= 40/87) && (SieveExponent <= t2 <= -(1/5) (t1 + t2)
+ 10/35)
U25[t1_, t2_] := (40/87 <= t1 + t2 <= 10/21) && (SieveExponent <= t2 <= 1/7 (t1 + t2) +
9/49)
U26[t1_, t2_] := (10/21 <= t1 + t2 <= 37/77) && (SieveExponent <= t2 <= -(t1 + t2) +
107/147)
U27[t1_, t2_] := (37/77 <= t1 + t2 <= 61/126) && (SieveExponent <= t2 <= -2 (t1 + t2) +
262/231)
U28[t1_, t2_] := (61/126 <= t1 + t2 <= 4607/9310) && (SieveExponent <= t2 <= -(70/11) (t1 +
t2) + 250/77)
U2[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)])
&& (U21[t1, t2] || U22[t1, t2] || U23[t1, t2] || U24[t1, t2] || U25[t1, t2] || U26[t1,
t2] || U27[t1, t2] || U28[t1, t2]) && (! U1[t1, t2])

(*The definition of the region U3*)
U31[t1_, t2_] := ((t1 + t2) + t2 <= 47/87) && (40/231 <= t2) && (29*(t1 + t2) - t2 <=
209/21) && (40/133 <= (t1 + SieveExponent)) && (29*t2 - (t1 + SieveExponent) <= 43/7) &&
(40/7 <= 12*(t1 + SieveExponent) + 11*t2)
U32[t1_, t2_] := ((t1 + t2) + t2 <= 109/189) && (29*(t1 + t2) + 19*t2 <= 103/7) && (160/189
<= 2*(t1 + SieveExponent) + t2) && (2*(t1 + t2) + 11*t2 <= 20/7) && (200/7 <= 58*(t1 +
SieveExponent) + 49*t2)
```



```

U33[t1_, t2_] := ((t1 + t2) + t2 <= 131/231) && (4/21 <= t2) && (6*(t1 + t2) + t2 <= 46/21)
&& (40/147 <= (t1 + SieveExponent)) && ((t1 + t2) + 8*t2 <= 16/7) && (20/7 <= 6*(t1 +
SieveExponent) + 5*t2)
U34[t1_, t2_] := ((t1 + t2) + t2 <= 93/133) && (40/231 <= t2) && (12*(t1 + t2) + t2 <= 44/7)
&& (40/87 <= (t1 + SieveExponent)) && (19*t2 - (t1 + SieveExponent) <= 79/21) &&
(160/21 <= 12*(t1 + SieveExponent) + 11*t2)
U35[t1_, t2_] := (2*(t1 + t2) + t2 <= 218/189) && (58*(t1 + t2) + 9*t2 <= 206/7) && (80/189
<= (t1 + SieveExponent)) && ((t1 + t2) + 5*t2 <= 10/7) && (100/7 <= 29*(t1 +
SieveExponent) + 10*t2)
U36[t1_, t2_] := ((t1 + t2) + t2 <= 107/147) && (4/21 <= t2) && (6*(t1 + t2) + t2 <= 22/7)
&& (100/231 <= (t1 + SieveExponent)) && (7*t2 - (t1 + SieveExponent) <= 9/7) && (80/21
<= 6*(t1 + SieveExponent) + 5*t2)
U37[t1_, t2_] := ((t1 + t2) + t2 <= 131/231) && (35*(t1 + t2) + 23*t2 <= 125/7) && (200/231
<= 2*(t1 + SieveExponent) + t2) && (2*(t1 + t2) + 13*t2 <= 64/21) && (240/7 <= 70*(t1 +
SieveExponent) + 59*t2)
U38[t1_, t2_] := ((t1 + t2) + t2 <= 51/91) && (41*(t1 + t2) + 27*t2 <= 21) && (80/91 <= 2*(
t1 + SieveExponent) + t2) && (2*(t1 + t2) + 15*t2 <= 68/21) && (40 <= 82*(t1 +
SieveExponent) + 69*t2)
U39[t1_, t2_] := (2*(t1 + t2) + t2 <= 262/231) && (70*(t1 + t2) + 11*t2 <= 250/7) &&
(100/231 <= (t1 + SieveExponent)) && ((t1 + t2) + 6*t2 <= 32/21) && (120/7 <= 35*(t1 +
SieveExponent) + 12*t2)
U310[t1_, t2_] := ((t2 + t2) + t1 <= 47/87) && (40/231 <= t1) && (29*(t2 + t2) - t1 <=
209/21) && (40/133 <= (t2 + SieveExponent)) && (29*t1 - (t2 + SieveExponent) <= 43/7) &&
(40/7 <= 12*(t2 + SieveExponent) + 11*t1)
U311[t1_, t2_] := ((t2 + t2) + t1 <= 109/189) && (29*(t2 + t2) + 19*t1 <= 103/7) && (160/189
<= 2*(t2 + SieveExponent) + t1) && (2*(t2 + t2) + 11*t1 <= 20/7) && (200/7 <= 58*(t2 +
SieveExponent) + 49*t1)
U312[t1_, t2_] := ((t2 + t2) + t1 <= 131/231) && (4/21 <= t1) && (6*(t2 + t2) + t1 <= 46/21)
&& (40/147 <= (t2 + SieveExponent)) && ((t2 + t2) + 8*t1 <= 16/7) && (20/7 <= 6*(t2 +
SieveExponent) + 5*t1)
U313[t1_, t2_] := ((t2 + t2) + t1 <= 93/133) && (40/231 <= t1) && (12*(t2 + t2) + t1 <=
44/7) && (40/87 <= (t2 + SieveExponent)) && (19*t1 - (t2 + SieveExponent) <= 79/21) &&
(160/21 <= 12*(t2 + SieveExponent) + 11*t1)
U314[t1_, t2_] := (2*(t2 + t2) + t1 <= 218/189) && (58*(t2 + t2) + 9*t1 <= 206/7) && (80/189
<= (t2 + SieveExponent)) && ((t2 + t2) + 5*t1 <= 10/7) && (100/7 <= 29*(t2 +
SieveExponent) + 10*t1)
U315[t1_, t2_] := ((t2 + t2) + t1 <= 107/147) && (4/21 <= t1) && (6*(t2 + t2) + t1 <= 22/7)
&& (100/231 <= (t2 + SieveExponent)) && (7*t1 - (t2 + SieveExponent) <= 9/7) && (80/21
<= 6*(t2 + SieveExponent) + 5*t1)
U316[t1_, t2_] := ((t2 + t2) + t1 <= 131/231) && (35*(t2 + t2) + 23*t1 <= 125/7) && (200/231
<= 2*(t2 + SieveExponent) + t1) && (2*(t2 + t2) + 13*t1 <= 64/21) && (240/7 <= 70*(t2 +
SieveExponent) + 59*t1)
U317[t1_, t2_] := ((t2 + t2) + t1 <= 51/91) && (41*(t2 + t2) + 27*t1 <= 21) && (80/91 <= 2*(
t2 + SieveExponent) + t1) && (2*(t2 + t2) + 15*t1 <= 68/21) && (40 <= 82*(t2 +
SieveExponent) + 69*t1)
U318[t1_, t2_] := (2*(t2 + t2) + t1 <= 262/231) && (70*(t2 + t2) + 11*t1 <= 250/7) &&
(100/231 <= (t2 + SieveExponent)) && ((t2 + t2) + 6*t1 <= 32/21) && (120/7 <= 35*(t2 +
SieveExponent) + 12*t1)
U319[t1_, t2_] := (t1 + (t2 + t2) <= 47/87) && (40/231 <= (t2 + SieveExponent)) && (29*t1 -
(t2 + SieveExponent) <= 209/21) && (40/133 <= t1) && (29*(t2 + t2) - t1 <= 43/7) &&
(40/7 <= 12*t1 + 11*(t2 + SieveExponent))
U320[t1_, t2_] := (t1 + (t2 + t2) <= 109/189) && (29*t1 + 19*(t2 + t2) <= 103/7) && (160/189
<= 2*t1 + (t2 + SieveExponent)) && (2*t1 + 11*(t2 + t2) <= 20/7) && (200/7 <= 58*t1 +
49*(t2 + SieveExponent))
U321[t1_, t2_] := (t1 + (t2 + t2) <= 131/231) && (4/21 <= (t2 + SieveExponent)) && (6*t1 + (
t2 + t2) <= 46/21) && (40/147 <= t1) && (t1 + 8*(t2 + t2) <= 16/7) && (20/7 <= 6*t1 +
5*(t2 + SieveExponent))
U322[t1_, t2_] := (t1 + (t2 + t2) <= 93/133) && (40/231 <= (t2 + SieveExponent)) && (12*t1 +
(t2 + t2) <= 44/7) && (40/87 <= t1) && (19*(t2 + t2) - t1 <= 79/21) && (160/21 <= 12*t1
+ 11*(t2 + SieveExponent))
U323[t1_, t2_] := (2*t1 + (t2 + t2) <= 218/189) && (58*t1 + 9*(t2 + t2) <= 206/7) && (80/189
<= t1) && (t1 + 5*(t2 + t2) <= 10/7) && (100/7 <= 29*t1 + 10*(t2 + SieveExponent))

```

```

U324[t1_, t2_] := (t1 + (t2 + t2) <= 107/147) && (4/21 <= (t2 + SieveExponent)) && (6*t1 + (
t2 + t2) <= 22/7) && (100/231 <= t1) && (7*(t2 + t2) - t1 <= 9/7) && (80/21 <= 6*t1 +
5*(t2 + SieveExponent))
U325[t1_, t2_] := (t1 + (t2 + t2) <= 131/231) && (35*t1 + 23*(t2 + t2) <= 125/7) && (200/231
<= 2*t1 + (t2 + SieveExponent)) && (2*t1 + 13*(t2 + t2) <= 64/21) && (240/7 <= 70*t1 +
59*(t2 + SieveExponent))
U326[t1_, t2_] := (t1 + (t2 + t2) <= 51/91) && (41*t1 + 27*(t2 + t2) <= 21) && (80/91 <= 2*
t1 + (t2 + SieveExponent)) && (2*t1 + 15*(t2 + t2) <= 68/21) && (40 <= 82*t1 + 69*(t2 +
SieveExponent))
U327[t1_, t2_] := (2*t1 + (t2 + t2) <= 262/231) && (70*t1 + 11*(t2 + t2) <= 250/7) &&
(100/231 <= t1) && (t1 + 6*(t2 + t2) <= 32/21) && (120/7 <= 35*t1 + 12*(t2 +
SieveExponent))
U3[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)])
&& (U31[t1, t2] || U32[t1, t2] || U33[t1, t2] || U34[t1, t2] || U35[t1, t2] || U36[t1,
t2] || U37[t1, t2] || U38[t1, t2] || U39[t1, t2] || U310[t1, t2] || U311[t1, t2] || U312
[t1, t2] || U313[t1, t2] || U314[t1, t2] || U315[t1, t2] || U316[t1, t2] || U317[t1, t2]
|| U318[t1, t2] || U319[t1, t2] || U320[t1, t2] || U321[t1, t2] || U322[t1, t2] || U323
[t1, t2] || U324[t1, t2] || U325[t1, t2] || U326[t1, t2] || U327[t1, t2]) && (! U1[t1,
t2])

(*The definition of the region U4*)
U41[t1_, t2_] := (2*(t1 + t2) + t2 <= 2 - 20/21) && (4*(t1 + t2) + 6*t2 <= 4 - 20/21) &&
(6*(t1 + t2) + 7*t2 <= 6 - 40/21) && (4*t2 <= 4 - 20/7) && (3*t2 - 2*(t1 + t2) <= 6 -
40/7)
U42[t1_, t2_] := ((t1 + t2) <= 11/21) && (t2 <= 5/42)
U43[t1_, t2_] := (80/189 <= (t1 + t2) <= 40/87) && (5/42 <= t2 <= -(1/5) (t1 + t2) + 10/35)
U44[t1_, t2_] := (40/87 <= (t1 + t2) <= 10/21) && (5/42 <= t2 <= 1/7 (t1 + t2) + 9/49)
U45[t1_, t2_] := (10/21 <= (t1 + t2) <= 37/77) && (5/42 <= t2 <= -(t1 + t2) + 107/147)
U46[t1_, t2_] := (37/77 <= (t1 + t2) <= 61/126) && (5/42 <= t2 <= -2 (t1 + t2) + 262/231)
U47[t1_, t2_] := (61/126 <= (t1 + t2) <= 289/588) && (5/42 <= t2 <= -(70/11) (t1 + t2) +
250/77)
U4[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)])
&& (U41[t1, t2] || U42[t1, t2] || U43[t1, t2] || U44[t1, t2] || U45[t1, t2] || U46[t1,
t2] || U47[t1, t2]) && (! U1[t1, t2]) && (! U2[t1, t2]) && (! U3[t1, t2])

(*The definition of the region U5*)
U51[t1_, t2_] := (2*(t2 + t2) + t1 <= 2 - 20/21) && (4*(t2 + t2) + 6*t1 <= 4 - 20/21) &&
(6*(t2 + t2) + 7*t1 <= 6 - 40/21) && (4*t1 <= 4 - 60/21) && (3*t1 - 2*(t2 +
SieveExponent) <= 6 - 120/21)
U52[t1_, t2_] := ((t2 + t2) <= 11/21) && (t1 <= 5/42)
U53[t1_, t2_] := (80/189 <= (t2 + SieveExponent)) && ((t2 + t2) <= 40/87) && (5/42 <= t1 <=
-(1/5) (t2 + t2) + 10/35)
U54[t1_, t2_] := (40/87 <= (t2 + SieveExponent)) && ((t2 + t2) <= 10/21) && (5/42 <= t1 <=
1/7 (t2 + SieveExponent) + 9/49)
U55[t1_, t2_] := (10/21 <= (t2 + SieveExponent)) && ((t2 + t2) <= 37/77) && (5/42 <= t1 <=
-(t2 + t2) + 107/147)
U56[t1_, t2_] := (37/77 <= (t2 + SieveExponent)) && ((t2 + t2) <= 61/126) && (5/42 <= t1 <=
-2 (t2 + t2) + 262/231)
U57[t1_, t2_] := (61/126 <= (t2 + SieveExponent)) && ((t2 + t2) <= 289/588) && (5/42 <= t1
<= -(70/11) (t2 + t2) + 250/77)
U5[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)])
&& (U51[t1, t2] || U52[t1, t2] || U53[t1, t2] || U54[t1, t2] || U55[t1, t2] || U56[t1,
t2] || U57[t1, t2]) && (! U1[t1, t2]) && (! U2[t1, t2]) && (! U3[t1, t2]) && (! U4[t1,
t2])

(*The definition of the region U6*)
U6[t1_, t2_, t3_, t4_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1
- t1)]) && (U4[t1, t2] || U5[t1, t2]) && (SieveExponent <= t3 < Min[t2, 1/2 (1 - t1 -
t2)]) && (SieveExponent <= t4 < Min[t3, 1/2 (1 - t1 - t2 - t3)]) && (U1[t1 + t2 + t3, t4
] || U1[t1 + t2 + t4, t3] || U1[t1 + t3 + t4, t2] || U1[t2 + t3 + t4, t1] || U1[t1 + t2,
t3 + t4] || U1[t1 + t3, t2 + t4] || U1[t1 + t4, t2 + t3] || U1[t2 + t3, t1 + t4] || U1[
t2 + t4, t1 + t3] || U1[t3 + t4, t1 + t2] || U1[t1, t2 + t3 + t4] || U1[t2, t1 + t3 + t4
] || U1[t3, t1 + t2 + t4] || U1[t4, t1 + t2 + t3])

```

```
(*The definition of the region U7*)
U7[t1_, t2_, t3_, t4_, t5_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1,
1/2 (1 - t1)]) && (U4[t1, t2] || U5[t1, t2]) && (SieveExponent <= t3 < Min[t2, 1/2 (1 -
t1 - t2)]) && (SieveExponent <= t4 < Min[t3, 1/2 (1 - t1 - t2 - t3)]) && (t4 < 1/2 (1 -
t1 - t2 - t3 - t4)) && (t4 < t5 < 1/2 (1 - t1 - t2 - t3 - t4)) && (! (U1[t1 + t2 + t3,
t4] || U1[t1 + t2 + t4, t3] || U1[t1 + t3 + t4, t2] || U1[t2 + t3 + t4, t1] || U1[t1 +
t2, t3 + t4] || U1[t1 + t3, t2 + t4] || U1[t1 + t4, t2 + t3] || U1[t2 + t3, t1 + t4] ||
U1[t2 + t4, t1 + t3] || U1[t3 + t4, t1 + t2] || U1[t1, t2 + t3 + t4] || U1[t2, t1 + t3 +
t4] || U1[t3, t1 + t2 + t4] || U1[t4, t1 + t2 + t3])) && (U1[t1, t2 + t3 + t4 + t5] ||
U1[t2, t1 + t3 + t4 + t5] || U1[t3, t1 + t2 + t4 + t5] || U1[t4, t1 + t2 + t3 + t5] ||
U1[t5, t1 + t2 + t3 + t4] || U1[t1 + t2, t3 + t4 + t5] || U1[t1 + t3, t2 + t4 + t5] ||
U1[t1 + t4, t2 + t3 + t5] || U1[t1 + t5, t2 + t3 + t4] || U1[t2 + t3, t1 + t4 + t5] ||
U1[t2 + t4, t1 + t3 + t5] || U1[t2 + t5, t1 + t3 + t4] || U1[t3 + t4, t1 + t2 + t5] ||
U1[t3 + t5, t1 + t2 + t4] || U1[t4 + t5, t1 + t2 + t3] || U1[t1 + t2 + t3, t4 + t5] ||
U1[t1 + t2 + t4, t3 + t5] || U1[t1 + t2 + t5, t3 + t4] || U1[t1 + t3 + t4, t2 + t5] ||
U1[t1 + t3 + t5, t2 + t4] || U1[t1 + t4 + t5, t2 + t3] || U1[t2 + t3 + t4, t1 + t5] ||
U1[t2 + t3 + t5, t1 + t4] || U1[t2 + t4 + t5, t1 + t3] || U1[t3 + t4 + t5, t1 + t2] ||
U1[t1 + t2 + t3 + t4, t5] || U1[t1 + t2 + t3 + t5, t4] || U1[t1 + t2 + t4 + t5, t3] ||
U1[t1 + t3 + t4 + t5, t2] || U1[t2 + t3 + t4 + t5, t1])
```

```
(*The definition of the region U8*)
U8[t1_, t2_, t3_, t4_, t5_, t6_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[
t1, 1/2 (1 - t1)]) && (U4[t1, t2] || U5[t1, t2]) && (SieveExponent <= t3 < Min[t2, 1/2
(1 - t1 - t2)]) && (SieveExponent <= t4 < Min[t3, 1/2 (1 - t1 - t2 - t3)]) && (t4 < 1/2
(1 - t1 - t2 - t3 - t4)) && (t4 < t5 < 1/2 (1 - t1 - t2 - t3 - t4)) && (t5 < 1/2 (1 - t1
- t2 - t3 - t4 - t5)) && (t5 < t6 < 1/2 (1 - t1 - t2 - t3 - t4 - t5)) && (! (U1[t1 + t2
+ t3, t4] || U1[t1 + t2 + t4, t3] || U1[t1 + t3 + t4, t2] || U1[t2 + t3 + t4, t1] || U1
[t1 + t2, t3 + t4] || U1[t1 + t3, t2 + t4] || U1[t1 + t4, t2 + t3] || U1[t2 + t3, t1 +
t4] || U1[t2 + t4, t1 + t3] || U1[t3 + t4, t1 + t2] || U1[t1, t2 + t3 + t4] || U1[t2, t1
+ t3 + t4] || U1[t3, t1 + t2 + t4] || U1[t4, t1 + t2 + t3])) && (! (U1[t1, t2 + t3 + t4
+ t5] || U1[t2, t1 + t3 + t4 + t5] || U1[t3, t1 + t2 + t4 + t5] || U1[t4, t1 + t2 + t3
+ t5] || U1[t5, t1 + t2 + t3 + t4] || U1[t1 + t2, t3 + t4 + t5] || U1[t1 + t3, t2 + t4 +
t5] || U1[t1 + t4, t2 + t3 + t5] || U1[t1 + t5, t2 + t3 + t4] || U1[t2 + t3, t1 + t4 +
t5] || U1[t2 + t4, t1 + t3 + t5] || U1[t2 + t5, t1 + t3 + t4] || U1[t3 + t4, t1 + t2 +
t5] || U1[t3 + t5, t1 + t2 + t4] || U1[t4 + t5, t1 + t2 + t3] || U1[t1 + t2 + t3, t4 +
t5] || U1[t1 + t2 + t4, t3 + t5] || U1[t1 + t2 + t5, t3 + t4] || U1[t1 + t3 + t4, t2 +
t5] || U1[t1 + t3 + t5, t2 + t4] || U1[t1 + t4 + t5, t2 + t3] || U1[t2 + t3 + t4, t1 +
t5] || U1[t2 + t3 + t5, t1 + t4] || U1[t2 + t4 + t5, t1 + t3] || U1[t3 + t4 + t5, t1 +
t2] || U1[t1 + t2 + t3 + t4, t5] || U1[t1 + t2 + t3 + t5, t4] || U1[t1 + t2 + t4 + t5,
t3] || U1[t1 + t3 + t4 + t5, t2] || U1[t2 + t3 + t4 + t5, t1]))
```

```

&& (U1[t1, t2 + t3 + t4 + t5 + t6] || U1[t2, t1 + t3 + t4 + t5 + t6] || U1[t3, t1 + t2 + t4
+ t5 + t6] || U1[t4, t1 + t2 + t3 + t5 + t6] || U1[t5, t1 + t2 + t3 + t4 + t6] || U1[t6,
t1 + t2 + t3 + t4 + t5] || U1[t1 + t2, t3 + t4 + t5 + t6] || U1[t1 + t3, t2 + t4 + t5 +
t6] || U1[t1 + t4, t2 + t3 + t5 + t6] || U1[t1 + t5, t2 + t3 + t4 + t6] || U1[t1 + t6,
t2 + t3 + t4 + t5] || U1[t2 + t3, t1 + t4 + t5 + t6] || U1[t2 + t4, t1 + t3 + t5 + t6]
|| U1[t2 + t5, t1 + t3 + t4 + t6] || U1[t2 + t6, t1 + t3 + t4 + t5] || U1[t3 + t4, t1 +
t2 + t5 + t6] || U1[t3 + t5, t1 + t2 + t4 + t6] || U1[t3 + t6, t1 + t2 + t4 + t5] || U1[
t4 + t5, t1 + t2 + t3 + t6] || U1[t4 + t6, t1 + t2 + t3 + t5] || U1[t5 + t6, t1 + t2 +
t3 + t4] || U1[t1 + t2 + t3, t4 + t5 + t6] || U1[t1 + t2 + t4, t3 + t5 + t6] || U1[t1 +
t2 + t5, t3 + t4 + t6] || U1[t1 + t2 + t6, t3 + t4 + t5] || U1[t1 + t3 + t4, t2 + t5 +
t6] || U1[t1 + t3 + t5, t2 + t4 + t6] || U1[t1 + t3 + t6, t2 + t4 + t5] || U1[t1 + t4 +
t5, t2 + t3 + t6] || U1[t1 + t4 + t6, t2 + t3 + t5] || U1[t1 + t5 + t6, t2 + t3 + t4] ||
U1[t2 + t3 + t4, t1 + t5 + t6] || U1[t2 + t3 + t5, t1 + t4 + t6] || U1[t2 + t3 + t6, t1
+ t4 + t5] || U1[t2 + t4 + t5, t1 + t3 + t6] || U1[t2 + t4 + t6, t1 + t3 + t5] || U1[t2
+ t5 + t6, t1 + t3 + t4] || U1[t3 + t4 + t5, t1 + t2 + t6] || U1[t3 + t4 + t6, t1 + t2
+ t5] || U1[t3 + t5 + t6, t1 + t2 + t4] || U1[t4 + t5 + t6, t1 + t2 + t3] || U1[t1 + t2
+ t3 + t4, t5 + t6] || U1[t1 + t2 + t3 + t5, t4 + t6] || U1[t1 + t2 + t3 + t6, t4 + t5]
|| U1[t1 + t2 + t4 + t5, t3 + t6] || U1[t1 + t2 + t4 + t6, t3 + t5] || U1[t1 + t2 + t5 +
t6, t3 + t4] || U1[t1 + t3 + t4 + t5, t2 + t6] || U1[t1 + t3 + t4 + t6, t2 + t5] || U1[
t1 + t3 + t5 + t6, t2 + t4] || U1[t1 + t4 + t5 + t6, t2 + t3] || U1[t2 + t3 + t4 + t5,
t1 + t6] || U1[t2 + t3 + t4 + t6, t1 + t5] || U1[t2 + t3 + t5 + t6, t1 + t4] || U1[t2 +
t4 + t5 + t6, t1 + t3] || U1[t3 + t4 + t5 + t6, t1 + t2] || U1[t1 + t2 + t3 + t4 + t5,
t6] || U1[t1 + t2 + t3 + t4 + t6, t5] || U1[t1 + t2 + t3 + t5 + t6, t4] || U1[t1 + t2 +
t4 + t5 + t6, t3] || U1[t1 + t3 + t4 + t5 + t6, t2] || U1[t2 + t3 + t4 + t5 + t6, t1])

(*The definition of the region U9*)
U9[t1_, t2_, t3_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1
)]) && (t2 < 1/2 (1 - t1 - t2)) && (t2 < t3 < 1/2 (1 - t1 - t2)) && (! U1[t1, t2]) && (!
U2[t1, t2]) && (! U3[t1, t2]) && (! U4[t1, t2]) && (! U5[t1, t2]) && (U1[t1 + t2, t3]
|| U1[t1 + t3, t2] || U1[t2 + t3, t1])

```

Code for the numerical calculations. The following Mathematica code shows all the numerical calculations performed on Mathematica 14.

```
(*The linear sieve functions F[u] and f[u], in Galway's SieveFunctions package*)
FF[u_] := UpperSieveFunc[1, u]
ff[u_] := LowerSieveFunc[1, u]

(*The Buchstab-Selberg function \[Omega][u]*)
\[Omega][u_] := (FF[u] + ff[u])/(2 Exp[EulerGamma])

(*A numerical lower bound for \[Omega][u] which will be used in calculating the value of
sums related to regions U6, U7 and U8*)
\[Omega]0[u_] := Piecewise[{{1/u, 1 <= u < 2}, {(1 + Log[u - 1])/u, 2 <= u < 3}, {(1 + Log[u
- 1])/u + (\[Pi]^2/12 + Log[-2 + u] Log[-1 + u] + PolyLog[2, 2 - u])/u, 3 <= u < 4},
{0.5612, u >= 4}}];

(*Numerical values of S1 and S2*)
S1 = 133/13 \[Omega][133/13]
S2 = 133/13 NIntegrate[\[Omega][133/13 (1 - t)]/t, {t, 13/13, 1/2}]
(*S1 = 5.744162508397124*)
(*S2 = 9.37592280506135*)

(*Numerical value of S31*)
S31 = NIntegrate[Boole[U1[t1, t2]]*\[Omega][(1 - t1 - t2)/t2]/(t1 t2^2), {t1, SieveExponent,
1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, Method -> {"GlobalAdaptive", "
MaxErrorIncreases" -> 30000}, MinRecursion -> 9]
(*S31 = 0.8061844833009656 with error less than 0.00006778442811040491*)

(*Numerical value of S32*)
S32 = NIntegrate[Boole[U2[t1, t2]]*\[Omega][(1 - t1 - t2)/t2]/(t1 t2^2), {t1, SieveExponent,
1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, Method -> {"GlobalAdaptive", "
MaxErrorIncreases" -> 30000}, MinRecursion -> 9]
(*S32 = 0.4964270854951454 with error less than 0.00011163981682258737*)

(*Numerical value of S33*)
S33 = NIntegrate[Boole[U3[t1, t2]]*\[Omega][(1 - t1 - t2)/t2]/(t1 t2^2), {t1, SieveExponent,
1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, Method -> {"GlobalAdaptive", "
MaxErrorIncreases" -> 1000}, MinRecursion -> 9]
(*S33 = 0.018359721032820922 with error less than 0.000041504121500542376*)

(*Numerical value of S34*)
S34 = NIntegrate[Boole[U4[t1, t2]]*(133/13 \[Omega][133/13 (1 - t1 - t2)]/(t1 t2), {t1,
SieveExponent, 1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, Method -> {"
GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9] - NIntegrate[Boole[U4[
t1, t2]]*(133/13 \[Omega][133/13 (1 - t1 - t2 - t3)]/(t1 t2 t3), {t1, SieveExponent
, 1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, {t3, SieveExponent, t2}, Method -> {
"GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9]
(*S34 = 2.213806035987798 with error less than 0.000826872988914825*)

(*Numerical value of S35*)
S35 = NIntegrate[Boole[U5[t1, t2]]*(133/13 \[Omega][133/13 (1 - t1 - t2)]/(t1 t2), {t1,
SieveExponent, 1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, Method -> {"
GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9] - NIntegrate[Boole[U5[
t1, t2]]*(133/13 \[Omega][133/13 (1 - t1 - t2 - t3)]/(t1 t2 t3), {t1, SieveExponent
, 1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, {t3, SieveExponent, t2}, Method -> {
"GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9]
(*S35 = 0.006463811030973154 with error less than 0.00006266570111378758*)

(*Numerical value of S36, using ParallelSum with 80 Wolfram kernels*)
SubIntervals = Table[{SieveExponent + (i - 1)*(1/2 - SieveExponent)/80, SieveExponent + i
*(1/2 - SieveExponent)/80}, {i, 1, 80}];
```

```

S36 = Total[ParallelSum[NIntegrate[Boole[U6[t1, t2, t3, t4]]*\[Omega]0[(1 - t1 - t2 - t3 -
t4)/t4]/(t1 t2 t3 t4^2), {t1, interval[[1]], interval[[2]]}, {t2, SieveExponent, Min[t1,
1/2 (1 - t1)]}, {t3, SieveExponent, Min[t2, 1/2 (1 - t1 - t2)]}, {t4, SieveExponent,
Min[t3, 1/2 (1 - t1 - t2 - t3)}], Method -> {"GlobalAdaptive", "MaxErrorIncreases" ->
30000}, MinRecursion -> 9], {interval, SubIntervals}]]
(*S36 = 0.1091869726983783 with total error less than 0.001165500896567007*)

(*Numerical value of S37, using ParallelSum with 80 Wolfram kernels*)
SubIntervals = Table[{SieveExponent + (i - 1)*(1/2 - SieveExponent)/80, SieveExponent + i
*(1/2 - SieveExponent)/80}, {i, 1, 80}];

S37 = Total[ParallelSum[NIntegrate[Boole[U7[t1, t2, t3, t4, t5]]*\[Omega]0[(1 - t1 - t2 - t3
- t4 - t5)/t5]/(t1 t2 t3 t4 t5^2), {t1, interval[[1]], interval[[2]]}, {t2,
SieveExponent, Min[t1, 1/2 (1 - t1)]}, {t3, SieveExponent, Min[t2, 1/2 (1 - t1 - t2)]},
{t4, SieveExponent, Min[t3, 1/2 (1 - t1 - t2 - t3)]}, {t5, t4, 1/2 (1 - t1 - t2 - t3 -
t4)}], Method -> {"GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9], {
interval, SubIntervals}]]
(*S37 = 0.004073696897902602 with total error less than 0.00007900973483599141*)

(*Numerical value of S38, using ParallelSum with 80 Wolfram kernels*)
SubIntervals = Table[{SieveExponent + (i - 1)*(1/2 - SieveExponent)/80, SieveExponent + i
*(1/2 - SieveExponent)/80}, {i, 1, 80}];

S38 = Total[ParallelSum[NIntegrate[Boole[U8[t1, t2, t3, t4, t5, t6]]*\[Omega]0[(1 - t1 - t2
- t3 - t4 - t5 - t6)/t6]/(t1 t2 t3 t4 t5 t6^2), {t1, interval[[1]], interval[[2]]}, {t2,
SieveExponent, Min[t1, 1/2 (1 - t1)]}, {t3, SieveExponent, Min[t2, 1/2 (1 - t1 - t2)]},
{t4, SieveExponent, Min[t3, 1/2 (1 - t1 - t2 - t3)]}, {t5, t4, 1/2 (1 - t1 - t2 - t3 -
t4)}, {t6, t5, 1/2 (1 - t1 - t2 - t3 - t4 - t5)}], Method -> {"GlobalAdaptive", "
MaxErrorIncreases" -> 30000}, MinRecursion -> 9], {interval, SubIntervals}]]
(*S38 = 0.0000000013638108439152523 with total error less than 0.00000000877263349687567*)

(*Numerical value of S39*)
S39 = NIntegrate[Boole[U9[t1, t2, t3]]*\[Omega]0[(1 - t1 - t2 - t3)/t3]/(t1 t2 t3^2), {t1,
SieveExponent, 1/2}, {t2, SieveExponent, Min[t1, 1/2 (1 - t1)]}, {t3, t2, 1/2 (1 - t1 -
t2)}, Method -> {"GlobalAdaptive", "MaxErrorIncreases" -> 30000}, MinRecursion -> 9]
(*S39 = 0.02094925331280443 with error less than 0.00035737496886306376*)

```

Code for the plot of regions. The following Mathematica code gives the plot of regions U_1-U_5 , the part of U_1 which produces U_2 , the extra two dimensional Type-I information region produced by U_1 and the three-dimensional plot of U_9 .

```
RegionPlot[{{SieveExponent <= t1 < 1/2 &&
SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)]}, {U1[t1, t2]}, {U2[t1, t2]}, {U3[t1, t2]},
{U4[t1, t2]}, {U5[t1, t2]}}, {t1, SieveExponent, 1/2}, {t2, SieveExponent, 1/2},
PlotLegends -> {"TotalRegion", "U1", "U2", "U3", "U4", "U5"}, MaxRecursion -> 13]
```




Figure 1: Plot of regions U_1-U_5

```

U2V21[t1_, t2_] := (4423/10906 <= t1 <= 113/273) && (SieveExponent <= t2 <= -(35/23) t1 +
125/161)
U2V22[t1_, t2_] := (113/273 <= t1 <= 267/637) && (SieveExponent <= t2 <= -t1 + 51/91)
U2V23[t1_, t2_] := (267/637 <= t1 <= 2442/5453) && (SieveExponent <= t2 <= -(41/27) t1 +
21/27)
U2V24[t1_, t2_] := (2124/4655 <= t1 <= 40/87) && (SieveExponent <= t2 <= -(1/5) t1 + 10/35)
U2V25[t1_, t2_] := (40/87 <= t1 <= 10/21) && (SieveExponent <= t2 <= 1/7 t1 + 9/49)
U2V26[t1_, t2_] := (10/21 <= t1 <= 37/77) && (SieveExponent <= t2 <= -t1 + 107/147)
U2V27[t1_, t2_] := (37/77 <= t1 <= 61/126) && (SieveExponent <= t2 <= -2 t1 + 262/231)
U2V28[t1_, t2_] := (61/126 <= t1 <= 4607/9310) && (SieveExponent <= t2 <= -(70/11) t1 +
250/77)
U2in2Variables[t1_, t2_] := U2V21[t1, t2] || U2V22[t1, t2] || U2V23[t1, t2] || U2V24[t1, t2]
|| U2V25[t1, t2] || U2V26[t1, t2] || U2V27[t1, t2] || U2V28[t1, t2]

RegionPlot[{{SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)]}, {U1[
t1, t2]}, {U2in2Variables[t1, t2]}}, {t1, SieveExponent, 1/2}, {t2, SieveExponent, 1/2},
PlotLegends -> {"TotalRegion", "U1", "U2 in 2 variables"}, MaxRecursion -> 7]

```




Figure 2: Plot of the part of region U_1 which produces U_2


```

DTypeI1[t1_, t2_] := (2*t1 + t2 <= 2 - 20/21) && (4*t1 + 6*t2 <= 4 - 20/21) && (6*t1 + 7*t2
<= 6 - 40/21) && (4*t2 <= 4 - 20/7) && (3*t2 - 2*t1 <= 6 - 40/7)
DTypeI2[t1_, t2_] := (t1 <= 11/21) && (t2 <= 5/42)
ExtraTypeI1[t1_, t2_] := (80/189 <= t1 <= 40/87) && (5/42 <= t2 <= -(1/5) t1 + 10/35)
ExtraTypeI2[t1_, t2_] := (40/87 <= t1 <= 10/21) && (5/42 <= t2 <= 1/7 t1 + 9/49)
ExtraTypeI3[t1_, t2_] := (10/21 <= t1 <= 37/77) && (5/42 <= t2 <= -t1 + 107/147)
ExtraTypeI4[t1_, t2_] := (37/77 <= t1 <= 61/126) && (5/42 <= t2 <= -2 t1 + 262/231)
ExtraTypeI5[t1_, t2_] := (61/126 <= t1 <= 289/588) && (5/42 <= t2 <= -(70/11) t1 + 250/77)
DTypeI[t1_, t2_] := (SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1
)]) && (DTypeI1[t1, t2] || DTypeI2[t1, t2])
ExtraTypeI[t1_, t2_] := (ExtraTypeI1[t1, t2] || ExtraTypeI2[t1, t2] || ExtraTypeI3[t1, t2]
|| ExtraTypeI4[t1, t2] || ExtraTypeI5[t1, t2]) && (! DTypeI[t1, t2])

RegionPlot[{{SieveExponent <= t1 < 1/2 && SieveExponent <= t2 < Min[t1, 1/2 (1 - t1)]}, {U1[
t1, t2]}, {DTypeI[t1, t2]}, {ExtraTypeI[t1, t2]}}, {t1, SieveExponent, 1/2}, {t2,
SieveExponent, 1/2}, PlotLegends -> {"TotalRegion", "U1", "Deshouillers-Iwaniec Type I",
"Extra Type I"}, MaxRecursion -> 7]

```




Figure 3: Plot of the 2 dimensional Type-I information

```

RegionPlot3D[{{U9[t1, t2, t3]}}, {t1, 0.2, 0.4}, {t2, SieveExponent, 0.3}, {t3,
SieveExponent, 0.3}, PlotLegends -> {"U9"}]

RegionPlot3D[{{U9[t1, t2, t3]}, {U1[t1, t2]}, {U2[t1, t2]}, {U3[t1, t2]}, {U4[t1, t2]}, {U5[
t1, t2]}}, {t1, 0.2, 0.4}, {t2, SieveExponent, 0.3}, {t3, SieveExponent, 0.3},
PlotLegends -> {"U9", "U1", "U2", "U3", "U4", "U5"}]

```




Figure 4: Three-dimensional plot of region U_9 (Default view, Top view, Front view, With other regions)

THE HIGH SCHOOL AFFILIATED TO RENMIN UNIVERSITY OF CHINA INTERNATIONAL CURRICULUM CENTER, BEIJING, CHINA
Email address: runbo.li.carey@gmail.com