

BILINEAR ESTIMATE FOR SCHRÖDINGER EQUATION ON $\mathbb{R} \times \mathbb{T}$

YANGKENDI DENG, BONING DI, CHENJIE FAN, AND ZEHUA ZHAO

ABSTRACT. We continue our study of bilinear estimates on waveguide $\mathbb{R} \times \mathbb{T}$ started in [DFY⁺24, Den23]. The main point of the current article is, comparing to previous work [Den23], that we obtain estimates beyond the semiclassical time regime. Our estimate is sharp in the sense that one can construct examples which saturate this estimate.

CONTENTS

1. Introduction	1
1.1. Statement of main results	1
1.2. Background, motivations and previous works	3
1.3. More results and some preliminary reductions	4
1.4. Structure of the article	5
1.5. Notations	5
Acknowledgment	7
2. An overview	7
2.1. A recall of Tao's bilinear estimates on Euclidean space	7
2.2. Overview of our proof	8
3. Proof of main results	9
3.1. Local estimate: integral version	9
3.2. Wave packet decomposition and semiclassical regime	10
3.3. Local estimate: tube version	12
3.4. Periodic property and the bilinear estimate	12
Appendix A: Waveguide version epsilon-removal lemma	13
Appendix B: Some examples for explaining the sharpness	17
References	18

1. INTRODUCTION

1.1. **Statement of main results.** In this paper, we consider (linear) Schrödinger equation on waveguide $\mathbb{R} \times \mathbb{T}$ as follows,

$$(1) \quad \begin{cases} iu_t - \Delta u = 0, \\ u(x, 0) = f(x). \end{cases}$$

We use $e^{it\Delta_{\mathbb{R} \times \mathbb{T}}}$ to denote the linear (Schrödinger) propagator. When there is no confusion, we short it as $e^{it\Delta}$ for convenience.

Note that the physical space here is $\mathbb{R} \times \mathbb{T}$, thus the corresponding (Fourier) frequency space is $\mathbb{R} \times \mathbb{Z}$. We use Q to denote cubes on $\mathbb{R} \times \mathbb{Z}$, and P_Q to the associated Littlewood-Paley projections¹.

The main result of this paper reads,

Theorem 1.1. *Consider equation (1). Let $p \in (\frac{5}{3}, 2)$. Let Q_1, Q_2 be two cubes of length $R \geq 1$ on $\mathbb{R} \times \mathbb{Z}$, and we assume they are R separated, i.e. their centers are of distance $\sim R$. Let T_p be defined as $R^{\frac{4p-8}{3-p}}$. Then for all $\varepsilon > 0$, one has for all $f, g \in L^2(\mathbb{R} \times \mathbb{T})$,*

$$(2) \quad \|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, T_p R^{-\varepsilon}])} \lesssim_\varepsilon R^{\frac{2p-4}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Remark 1.2. Under the same assumption of Theorem 1.1, it follows directly from (2) that

$$(3) \quad \|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, T_p])} \lesssim_\varepsilon R^{\frac{2p-4}{p} + \varepsilon} \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

And indeed, for (3), we can also cover the endpoint $p = \frac{5}{3}$.

But it should be noted that (3) cannot imply (2) and it, in some senses, loses derivative and is not very useful when one studies L^2 -critical nonlinear Schrödinger equations (NLS) problems².

One may observe T_p in Theorem 1.1 equals R^{-1} when $p = \frac{5}{3}$, and goes to 1 as $p \rightarrow 2$. The two endpoints are both special. When $p = \frac{5}{3}$ and $T_p = \frac{1}{R}$, the estimate falls in the so-called self similar regime, and estimate (2) is expected to follow from the Euclidean case, [Tao03] (since within this time scale the Schrödinger wave does not quite see the periodic effect of \mathbb{T}), and the associated estimate has been studied in [Den23], but in the spirit of [TVV98], that is

$$(4) \quad \|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^2_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, R^{-1}])} \lesssim R^{\frac{2q-4}{q}} \|\widehat{f}\|_{L^q(\mathbb{R} \times \mathbb{Z})} \|\widehat{g}\|_{L^q(\mathbb{R} \times \mathbb{Z})}, \quad q = \frac{12}{7}.$$

Up to ε -loss, the above estimate can be deduced by interpolating (3) for the $p = \frac{5}{3}$ and an $L^1 \times L^1 \rightarrow L^\infty$ estimate. It should be noted that (4) does not suffer from the ε -loss, and is more useful within the semiclassical time scale $T \lesssim \frac{1}{R}$.

When $p = 2$, Theorem 1.1 follows directly from the Strichartz estimate on waveguide $\mathbb{R} \times \mathbb{T}$, [TT01] (see also [BCP21]).

The **main point** of our estimate (2) is that p can be smaller than 2, and T_p can be pushed to 1 as p approaches 2, and in particular, $T_p \gg \frac{1}{R}$ for all $p > \frac{5}{3}$. We also note that estimate (2) is **sharp** in the sense that one can construct examples which saturate this estimate. See Appendix B for these examples.

¹See Notations 1.5 for more details.

²We refer to [Dod12] for the notion of L^2 -criticality where the long time behavior for L^2 -critical NLS is under studied.

1.2. Background, motivations and previous works. To explain the background explicitly, one needs to consider (1) also on \mathbb{R}^2 (Euclidean space), and \mathbb{T}^2 (periodic space, i.e. 2D tori). In this subsection, we will use $e^{it\Delta_{\mathbb{R} \times \mathbb{T}}}$, $e^{it\Delta_{\mathbb{R}^2}}$, $e^{it\Delta_{\mathbb{T}^2}}$ to denote linear Schrödinger operators on different manifolds to avoid unnecessary confusions.

Strichartz estimates for those linear models play a fundamental role in the study of corresponding nonlinear problems. In \mathbb{R}^2 , see, [KT98, Caz03], it reads as

$$(5) \quad \|e^{it\Delta_{\mathbb{R}^2}} f\|_{L_{x,t}^4(\mathbb{R}^2 \times \mathbb{R})} \lesssim \|f\|_{L^2(\mathbb{R}^2)}.$$

The parallel estimate on \mathbb{T}^2 (rational or irrational), see [Bou93, BD15], reads as

$$(6) \quad \|e^{it\Delta_{\mathbb{T}^2}} P_N f\|_{L_{x,t}^4(\mathbb{T}^2 \times [0,1])} \lesssim_\varepsilon N^\varepsilon \|f\|_{L^2(\mathbb{T}^2)}.$$

Estimate (6) is weaker than (5), but optimal³. Note that it only holds local in time and loses ε derivative. And in particular, it makes the study of 2D cubic (mass-critical) NLS ($iu_t + \Delta u = +|u|^2 u$) at L^2 regularity on torus out of reach at the moment. Meanwhile, global well-posedness (GWP) and scattering of (defocusing) mass-critical NLS on Euclidean space is one landmark result in dispersive PDEs, see [Dod12, Dod15, Dod16a].

The parallel estimate on $\mathbb{R} \times \mathbb{T}$ can still only hold local in time, but with no loss of derivative, [TT01],

$$(7) \quad \|e^{it\Delta_{\mathbb{R} \times \mathbb{T}}} f\|_{L_{x,t}^4(\mathbb{R} \times \mathbb{T} \times [0,1])} \lesssim \|f\|_{L^2(\mathbb{R} \times \mathbb{T})}.$$

In some sense, this makes $\mathbb{R} \times \mathbb{T}$ an interesting manifold to study mass-critical NLS, and the associated Strichartz estimate which does **not lose** derivative.

Barron [Bar21] proves global-in-time Strichartz estimates for Schrödinger equations on product spaces $\mathbb{R}^n \times \mathbb{T}^m$ with an additional ε -derivative loss via the ℓ^2 -decoupling method. This loss is significant at the critical endpoint of the Strichartz estimates⁴. These Strichartz estimates are then applied to prove small-data-scattering at the scaling critical regularity for certain nonlinear NLS models. Moreover, Barron-Christ-Pausader's recent work [BCP21] establishes global-in-time⁵ Strichartz estimates for Schrödinger equations on product spaces $\mathbb{R} \times \mathbb{T}$, overcoming derivative losses.

It is also natural to study the bilinear version of estimates (5),(6),(7).

One direction is to consider $\|e^{it\Delta} P_N f e^{it\Delta} P_M g\|_{L_{x,t}^2}$, i.e. one exploits extra smallness when f, g are localized at different (dyadic) frequencies N, M . It indicates direct applications for low regularity GWP results of NLS, we refer to [Bou98, CGSY23, DFY⁺24, DSPST07, FSWW18, ZZ21]⁶.

Another direction is to study bilinear restriction estimates as in [Tao03]. And in \mathbb{R}^2 , it reads as

$$(8) \quad \|e^{it\Delta_{\mathbb{R}^2}} P_{Q'_1} f e^{it\Delta_{\mathbb{R}^2}} P_{Q'_2} g\|_{L_{x,t}^p(\mathbb{R}^2 \times \mathbb{R})} \lesssim \|f\|_{L_x^2} \|g\|_{L_x^2}, \quad \frac{5}{3} < p < 2.$$

³See Appendix in [TT01].

⁴Interestingly, Barron shows that this ε -loss can be removed for exponents away from the critical endpoint. This removal enhances the strength of the estimates for those specific exponents.

⁵Compared with [TT01], the estimate in [BCP21] is global-in-time.

⁶We also refer to the Introduction in [DFY⁺24] for a brief introduction for the topic “NLS on waveguide manifolds” (see also the references therein), which has been intensively studied in recent decades. We do not emphasize this point since we focus on the bilinear estimates in this paper.

Here Q'_1, Q'_2 are two cubes of scale 1, and they are separated by distance 1.

Estimate (8) is of great importance in the study of modern harmonic analysis, see [Tao01, Tao03, TVV98, Wol01]. And it is directly related to the concentration compactness theory in dispersive PDEs, see [Bou98, BG99, KM06, KM06, MV98]. And one may refer to [BV07] for a systematic approach to transfer estimate (8) into a profile decomposition.

By the scaling symmetry of \mathbb{R}^2 , estimate (8) is equivalent to

$$(9) \quad \|e^{it\Delta_{\mathbb{R}^2}} P_{Q_1} f e^{it\Delta_{\mathbb{R}^2}} P_{Q_2} g\|_{L^p_{x,t}(\mathbb{R}^2 \times \mathbb{R})} \lesssim R^{\frac{2p-4}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}.$$

Here Q_1, Q_2 are two cubes of scale $R \geq 1$, and they are separated by distance R .

The goal of the current article is to study the analogous estimate of (9) in waveguide $\mathbb{R} \times \mathbb{T}$, and hopefully it can be helpful to further applied to implement the scheme of [IPS12, IP12] to transfer the (large data) GWP results in the Euclidean setting [Dod16b]. for mass-critical NLS to the waveguide $\mathbb{R} \times \mathbb{T}$ case.

1.3. More results and some preliminary reductions. Theorem 1.1 estimates bilinear interactions of Schrödinger wave localizing at frequency $|\xi| \sim R$ up to time scale $T_p := R^{\frac{4p-8}{3-p}}$. One may wonder what can be said for time scale $T_p \leq T \leq 1$. Via interpolation Theorem 1.1 with (the rather straightforward) bound

$$(10) \quad \|e^{it\Delta} f e^{it\Delta} g\|_{L^1_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim T \|f\|_{L^2_x} \|g\|_{L^2_x},$$

one obtains the following corollary,

Corollary 1.3. *Let Q_1, Q_2 be two cubes of length $R \geq 1$ on $\mathbb{R} \times \mathbb{Z}$, and we assume that they are R separated. Then for $p \in (\frac{5}{3}, 2]$, $T \in [T_p, 1]$, and all $f, g \in L^2(\mathbb{R} \times \mathbb{T})$, there holds*

$$\|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p(\mathbb{R} \times \mathbb{T} \times [0, TR^{-\varepsilon}])} \lesssim_{\varepsilon} T^{\frac{3-p}{2p}} \|f\|_{L^2_x} \|g\|_{L^2_x}, \quad \forall \varepsilon > 0.$$

For $p \in (\frac{5}{3}, 2]$, $T \in [R^{-2}, T_p]$, there holds

$$\|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p(\mathbb{R} \times \mathbb{T} \times [0, TR^{-\varepsilon}])} \lesssim_{\varepsilon} R^{\frac{2p-4}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}, \quad \forall \varepsilon > 0.$$

For $p \in [1, \frac{5}{3}]$, $T \in [R^{-1}, 1]$, there holds

$$\|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim_{\varepsilon} T^{\frac{3-p}{2p}-\varepsilon} \|f\|_{L^2_x} \|g\|_{L^2_x}, \quad \forall \varepsilon > 0.$$

For $p \in [1, \frac{5}{3}]$, $T \in [R^{-2}, R^{-1}]$, there holds

$$\|e^{it\Delta} P_{Q_1} f e^{it\Delta} P_{Q_2} g\|_{L^p(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim_{\varepsilon} T^{\frac{5p-3}{2p}-\varepsilon} R^{\frac{1-p}{p}} \|f\|_{L^2_x} \|g\|_{L^2_x}, \quad \forall \varepsilon > 0.$$

Remark 1.4. The above estimate is sharp up to ε -loss, see Appendix B. Our methods can also be applied to get similar results for general waveguide $\mathbb{R}^m \times \mathbb{T}^n$ ($m, n \geq 1$), with some suitable modifications.

Remark 1.5. As mentioned in [Tao03], it is expected to apply Corollary 1.3 to obtain a **pointwise convergence** result for the Schrödinger equation on $\mathbb{R} \times \mathbb{T}$ in a standard way. There has been intensive studies for the pointwise convergence in Euclidean situation, and we refer to [Bou16, DGL17, DZ19] for some classical results. We thank Prof. Jiqiang Zheng for helpful communications on this point.

We reformulate (2) in the formal of extension operators. The study of free Schrödinger solution is directly related to the restriction problem in harmonic analysis, see for instance [Gut23] and the reference therein. This is because the space-time Fourier transform of the linear Schrödinger equation is supported in the paraboloid. And that is why estimate (8) are usually called bilinear restriction estimates in the literature. Let extension operator \tilde{E} be defined as

$$(11) \quad (\tilde{E}h)(x, t) := \int_{\mathbb{R}^2} h(\xi) e^{ix\xi + it|\xi|^2} d\xi.$$

Here $x \in \mathbb{R}^2, \xi \in \mathbb{R}^2$. Note that one has $e^{-it\Delta_{\mathbb{R}^2}} f \equiv \tilde{E}\hat{f}$.

Let Q be a cube in \mathbb{R}^2 , one naturally define $\tilde{E}_Q h$ as $\tilde{E}(\chi_Q h)$.

Let Q'_1, Q'_2 be as in (8), then (8) is equivalent to

$$(12) \quad \|E_{Q'_1} h_1 E_{Q'_2} h_2\|_{L^p_{x,t}} \lesssim \|h_1\|_{L^2_\xi} \|h_2\|_{L^2_\xi}.$$

And one may also reformulate (9) in a similar fashion.

We now set up the extension operator in the setting of waveguide, i.e. we define the extension operator E as

$$(13) \quad Eh(x, t) := \int_{\mathbb{R} \times \mathbb{Z}} e^{ix\xi + it|\xi|^2} h(\xi) d\xi \equiv \sum_{\xi_2 \in \mathbb{Z}} \int_{\mathbb{R}} e^{i(x_1\xi_1 + x_2\xi_2) + it(\xi_1^2 + \xi_2^2)} h(\xi_1, \xi_2) d\xi_1,$$

where $x \in \mathbb{R} \times \mathbb{T}, t \in \mathbb{R}$ and $h \in L^2(\mathbb{R} \times \mathbb{Z})$. Strictly speaking, we should use $e^{2\pi ix\xi}$ rather than $e^{ix\xi}$ in (13), we neglect this issue for notation simplicity.

And one defines $E_Q h$ as $E(\chi_Q h)$.

Now, let Q_1, Q_2 be two cubes of scale R , and separated by distance $\sim R$ as in Theorem 1.1, the main estimate (2) can be **rewritten** as

$$(14) \quad \|E_{Q_1} h_1 E_{Q_2} h_2\|_{L^p_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, T_p R^{-\varepsilon}])} \lesssim R^{\frac{2p-4}{p}} \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z})},$$

where $h_1, h_2 \in L^2(\mathbb{R} \times \mathbb{Z})$. We now focus on estimate (14) in the rest of the article.

1.4. Structure of the article. The rest of the article is structured as follows. We end this section with some notations which will be used later. In Section 2, we recall some classical Euclidean bilinear estimates of Tao [Tao03], and then state the outline of our bilinear estimate on waveguide manifold. In Section 3, we present the detailed proof of our main result Theorem 1.1. Finally, we show a waveguide version epsilon-removal lemma in Appendix A, and then show the aforementioned sharpness examples in Appendix B.

1.5. Notations. We write $A \lesssim B$ to say that there is a constant C such that $A \leq CB$. We use $A \sim B$ when $A \lesssim B \lesssim A$. Particularly, we write $A \lesssim_u B$ to express that $A \leq C(u)B$ for some constant $C(u)$ depending on u .

Then we give some more preliminaries in the setting of waveguide manifold. Throughout this paper, we regularly refer to the spacetime norms

$$(15) \quad \|u\|_{L_x^q L_t^p(\mathbb{R}^m \times \mathbb{T}^n \times I)} := \left(\int_I \left(\int_{\mathbb{R}^m \times \mathbb{T}^n} |u(t, x)|^q dx \right)^{\frac{p}{q}} dt \right)^{\frac{1}{p}}.$$

Moreover, we turn to the Fourier transformation and Littlewood-Paley theory. We define the Fourier transform on $\mathbb{R}^m \times \mathbb{T}^n$ as follows:

$$(16) \quad \widehat{f}(\xi) := \int_{\mathbb{R}^m \times \mathbb{T}^n} f(x) e^{-ix\xi} dx,$$

where $\xi = (\xi_1, \xi_2, \dots, \xi_d) \in \mathbb{R}^m \times \mathbb{Z}^n$ and $d = m + n$. We also note the Fourier inversion formula

$$(17) \quad f(x) = c \sum_{(\xi_{m+1}, \dots, \xi_d) \in \mathbb{Z}^n} \int_{(\xi_1, \dots, \xi_m) \in \mathbb{R}^m} \widehat{f}(\xi) e^{ix\xi} d\xi_1 \dots d\xi_m.$$

Moreover, we define the Schrödinger propagator $e^{it\Delta}$ by

$$(18) \quad \widehat{e^{it\Delta} f}(\xi) := e^{it|\xi|^2} \widehat{f}(\xi).$$

We are now ready to define the Littlewood-Paley projections. First, we fix $\eta_1 : \mathbb{R} \rightarrow [0, 1]$, a smooth even function satisfying

$$(19) \quad \eta_1(\xi) = \begin{cases} 1, & |\xi| \leq 1, \\ 0, & |\xi| \geq 2, \end{cases}$$

and $N = 2^j$ a dyadic integer. Let $\eta^d = \mathbb{R}^d \rightarrow [0, 1]$, $\eta^d(\xi) = \eta_1(\xi_1)\eta_1(\xi_2)\eta_1(\xi_3)\dots\eta_1(\xi_d)$. We define the Littlewood-Paley projectors $P_{\leq N}$ and P_N by

$$(20) \quad \widehat{P_{\leq N} f}(\xi) := \eta^d\left(\frac{\xi}{N}\right) \widehat{f}(\xi), \quad \xi \in \mathbb{R}^m \times \mathbb{Z}^n,$$

and

$$(21) \quad P_N f := P_{\leq N} f - P_{\leq \frac{N}{2}} f.$$

Littlewood-Paley projection P_Q can be also defined in a natural way where Q is a cube on $\mathbb{R} \times \mathbb{Z}$.

Let the rescaled torus be defined as $\mathbb{T}_R := R\mathbb{T}$ for $R \geq 1$. Correspondingly $\mathbb{Z}_{1/R} := R^{-1}\mathbb{Z}$. We define

$$\int_{\mathbb{R} \times \mathbb{Z}_{1/R}} h(\xi) d\xi := R^{-1} \sum_{\xi_2 \in \mathbb{Z}_{1/R}} \int_{\mathbb{R}} h(\xi_1, \xi_2) d\xi_1,$$

and the rescaled extension operator E^R as follows

$$E^R h(x, t) := \int_{\mathbb{R} \times \mathbb{Z}_{1/R}} e^{ix\xi + it|\xi|^2} h(\xi) d\xi.$$

For a subset $Q' \subset \mathbb{R} \times \mathbb{Z}_{1/R}$, we introduce the operator $E_{Q'}^R$ as

$$E_{Q'}^R h(x, t) := \int_{Q'} e^{ix\xi + it|\xi|^2} h(\xi) d\xi.$$

Based on these notations, notice that

$$E_Q h\left(\frac{x}{R}, \frac{t}{R^2}\right) = R E_{Q'}^R h_R(x, t), \quad h_R(\xi) := R h(R\xi), \quad Q' := R^{-1}Q.$$

For a finite set S , we use the notation $\#S$ to denote the number of its elements.

Acknowledgment. B. Di was supported by the Postdoctoral Fellowship Program of CPSF under Grant Number GZB20230812. C. Fan was partially supported by the National Key R&D Program of China, 2021YFA1000800, CAS Project for Young Scientists in Basic Research, Grant No. YSBR-031, and NSFC grant no. 12288201. Z. Zhao was supported by the NSF grant of China (No. 12101046, 12271032) and the Beijing Institute of Technology Research Fund Program for Young Scholars.

2. AN OVERVIEW

2.1. A recall of Tao's bilinear estimates on Euclidean space. In this subsection, due to the aforementioned connections between free (linear) Schrödinger equations and Fourier restriction operators, we recall some of Tao's classical bilinear restriction estimates on Euclidean space ([Tao03]). There are mainly three steps summarized as follows.

Step 1: As shown in the literature [Tao03, TVV98, Wol01], to establish some desired bilinear restriction estimates, one typical method is to establish a local estimate (which may have ε -loss but have endpoint index) and then apply an ε -removal lemma. More precisely, to establish the desired global-type bilinear restriction estimate (8), it suffices to show that the following local-type bilinear restriction estimate

$$(22) \quad \left\| e^{it\Delta_{\mathbb{R}^2}} P_{Q'_1} f e^{it\Delta_{\mathbb{R}^2}} P_{Q'_2} g \right\|_{L_{x,t}^{\frac{5}{3}}(B_r(x_0, t_0))} \lesssim C_\alpha r^\alpha \|f\|_{L_x^2} \|g\|_{L_x^2}$$

holds for all spacetime balls $B_r(x_0, t_0)$ of radius r and all $\alpha > 0$. Note that this kind of reductions unavoidably lead to some non-endpoint index (non-sharp) phenomena.

Step 2: Next, based on this local-type reduction in the first step, one need to further investigate the local-type bilinear estimate (22). Here one typical tool is called *wave packet decomposition*, which can be viewed as a quantity version of the uncertainty principle. Roughly speaking, wave packet decomposition states that the solution $e^{it\Delta_{\mathbb{R}^2}} f$ can be decomposed as a summation of wave packets ϕ_T , which have good localization in both physical space and frequency space. Formally, all these wave packets ϕ_T can be viewed as characteristic functions of tubes T , which have base radii of r and lengths of r^2 . More precisely, there exists a wave packet decomposition

$$e^{it\Delta_{\mathbb{R}^2}} P_{Q'_1} f(x) = \sum_{T_1 \in \mathcal{T}'_1} c_{T_1} \phi_{T_1}(x, t),$$

where \mathcal{T}'_1 is a collection of tubes that is associated to the cube Q'_1 , and the coefficients c_{T_1} obey the following ℓ^2 bound $\|c_{T_1}\|_{\ell^2} \lesssim \|f\|_{L_x^2}$. Similar decomposition also holds for the item $e^{it\Delta_{\mathbb{R}^2}} P_{Q'_2} g$. Thus by putting the wave packet decomposition into the desired estimate (22), one can reduce this desired estimate to a tube-type (wave packet) estimate. Note that these kind of reductions can transfer the integral-type problem into a tube-type (wave packet) problem.

Step 3: Moreover, based on this *tube-type reduction* in the second step, one need to further investigate the associated *tube-type estimate*. To do so, under certain transverse condition, Tao has established an incidence estimate of tubes, which is his main conclusion in [Tao03]. More precisely, to show the desired local bilinear restriction estimate (22), Tao established an incidence estimate on wave packets, which claims that the following estimate

$$(23) \quad \left\| \sum_{T_1 \in \mathcal{T}'_1} \sum_{T_2 \in \mathcal{T}'_2} \phi_{T_1} \phi_{T_2} \right\|_{L^{\frac{5}{3}}_{x,t}([0,r]^3)} \lesssim C_\alpha r^\alpha \left(\#\mathcal{T}'_{1,[0,r]^3} \right)^{1/2} \left(\#\mathcal{T}'_{2,[0,r]^3} \right)^{1/2}$$

holds for all radii $r > 0$ and all parameters $\alpha > 0$. Here the collections \mathcal{T}'_1 and \mathcal{T}'_2 satisfy certain transverse condition, due to the geometric property of paraboloid; and the notation $\mathcal{T}'_{i,[0,r]^3}$ are defined as

$$\mathcal{T}'_{i,[0,r]^3} := \{T : T \in \mathcal{T}'_i, T \cap [0,r]^3 \neq \emptyset\}.$$

We point out that this incidence estimate is highly nontrivial, and is proved by induction on scales with the help of certain combinatorial argument. In this article, we will use this incidence-type estimate as a *blackbox*.

We **emphasize** that the aforementioned wave packet decomposition is very important for us, since it helps us connect Schrödinger wave on \mathbb{R}^2 and on $\mathbb{R} \times \mathbb{T}$, through incidence-type estimate. This intuition will be achieved later in the proof of our main result.

2.2. Overview of our proof. Here we show the outline of our proof, which basically follows the aforementioned Euclidean steps. However, some special phenomena have appeared due to our waveguide $\mathbb{R} \times \mathbb{T}$ setting. Thus there need some necessary adjustments to handle these phenomena.

First, we establish a waveguide version ε -removal lemma, which will lead to the ε -loss in our main estimate (2). In summary, applying the local-type reduction in the waveguide setting, we not only lose the endpoint Lebesgue space index but also lose the $R^{-\varepsilon}$ time regime. That is why we have ε -loss in our main Theorem 1.1. To state the waveguide version ε -removal argument more precisely, we naturally introduce the cubes B_r in rescaled-waveguide $\mathbb{R} \times \mathbb{T}_R \times \mathbb{R}$ as follows

$$B_r := [0,r] \times [0,r] \times [0,r] \quad \text{for } r \leq R, \quad B_r := [0,r] \times [0,R] \times [0,r] \quad \text{for } r > R.$$

Then, by the waveguide version ε -removal lemma and some rescaled arguments, we can reduce our desired global-type estimate (14) to the following local-type bilinear estimate

$$\left\| E_{Q'_1}^R h_1 E_{Q'_2}^R h_2 \right\|_{L^p_{x,t}(B_r)} \lesssim C_\alpha r^\alpha \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})}, \quad \forall r \in [1, R^{\frac{2p-2}{3-p}}].$$

Note that the waveguide cube B_r is different from the classical Euclidean cubes when $r > R$. This fact prevents us from directly applying Tao's classical incidence estimate. We will deal with this difference later, by making use of certain periodic property.

Second, we establish a waveguide version wave packet decomposition, which has some periodic properties due to the waveguide $\mathbb{R} \times \mathbb{T}$ setting. Roughly speaking, the waveguide version wave packet decomposition satisfies all the properties of Euclidean version wave packet decomposition,

and further satisfies one more periodic property in the second spatial direction due to the half-periodic manifold $\mathbb{R} \times \mathbb{T}$. More precisely, there exists a decomposition

$$E_{Q'_1}^R h_1 = \sum_{T_1} \tilde{c}_{T_1} \psi_{T_1},$$

where T_1 ranges over a collection that is associated to the cube Q'_1 , and the coefficients \tilde{c}_{T_1} obey the ℓ^2 bound $\|\tilde{c}_{T_1}\|_{\ell^2} \lesssim \|h_1\|_{L^2_\xi}$. Moreover, for each fixed T_1 , the function ψ_{T_1} can be viewed as a characteristic function of the following periodic-type tube

$$\bigcup_{m \in \mathbb{Z}} [T_1 + (0, mR, 0)],$$

where each translated tube $T_1 + (0, mR, 0)$ has base radial r and length r^2 . Similar decomposition also holds for the item $E_{Q'_2}^R h_2$. Hence, we can reduce our desired local-type bilinear estimate to a tube-type (wave packet) estimate, and the localized waveguide spacetime cube B_r . As mentioned above, to apply Tao's classical tube-type bilinear estimate, it remains to transfer the waveguide cube B_r to the standard Euclidean cube $[0, r]^3$.

Finally, we use the aforementioned periodic property to extend the data, and transfer our B_r -type wave packet estimate to the standard $[0, r]^3$ -type wave packet estimate. This periodic argument allows us to apply Tao's incidence estimate in our situation, and we hence complete the proof.

3. PROOF OF MAIN RESULTS

3.1. Local estimate: integral version. In this subsection, we do the local-type reduction. Recall that we are going to show the desired bilinear estimate (14). By rescaling, it suffices to show

$$\left\| E_{Q'_1}^R h_1 E_{Q'_2}^R h_2 \right\|_{L^p(\mathbb{R} \times \mathbb{T}_R \times [0, T_p R^{2-\varepsilon}])} \lesssim \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})}.$$

It is natural to introduce the waveguide cubes B_r in $\mathbb{R} \times \mathbb{T}_R \times \mathbb{R}$ as follows

$$B_r := [0, r] \times [0, r] \times [0, r] \quad \text{for } r \leq R, \quad B_r := [0, r] \times [0, R] \times [0, r] \quad \text{for } r > R.$$

Then by the waveguide version ε -removal lemma, which will be presented in the Appendix A, it suffices to show the following local estimate.

Proposition 3.1. *Given $p \in [\frac{5}{3}, 2]$ and $R \gg 1$. For the dyadic cubes*

$$Q'_1 \in R^{-1} Q_R, \quad Q'_2 \in R^{-1} Q_R, \quad \text{dist}(Q'_1, Q'_2) \gg 1,$$

the following bilinear estimate

$$(24) \quad \left\| E_{Q'_1}^R h_1 E_{Q'_2}^R h_2 \right\|_{L^p_{x,t}(B_r)} \lesssim_\alpha r^\alpha \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})},$$

holds for all dyadic numbers $r \in [1, R^{\frac{2p-2}{3-p}}]$.

3.2. Wave packet decomposition and semiclassical regime. Similar to the Euclidean case, we need a waveguide version wave packet decomposition to investigate the local estimate (24). In this subsection, we are going to establish the desired waveguide version wave packet decomposition, which can also help us directly handle the semiclassical regime, see Remark 3.3 below. The classical Euclidean version wave packet decomposition can be seen in [Tao03, Lemma 4.1], see also [Dem20, Section 2].

For a base set $Q'_i \subset \mathbb{R} \times \mathbb{Z}_{1/R}$ and the associated cap

$$S_i := \{(\xi, |\xi|^2) : \xi \in Q'_i\}, \quad S_i \subset \mathbb{R} \times \mathbb{Z}_{1/R} \times \mathbb{R},$$

denote a S_i -tube (with scale r) to be any set of the form

$$T := \{(x, t) : \frac{r}{2} \leq t \leq r, |x - (x_T + tv_T)| \leq r^{\frac{1}{2}}\},$$

with

$$x_T \in r^{\frac{1}{2}}\mathbb{Z}^2, \quad v_T \in r^{-\frac{1}{2}}\mathbb{Z}^2 \cap Q'_i, \quad (1, v_T) \perp S_i.$$

For convenience, we further denote the single-period \tilde{S}_i -tube (with scale r) to be any set of the form

$$T := \{(x, t) : \frac{r}{2} \leq t \leq r, |x - (x_T + tv_T)| \leq r^{\frac{1}{2}}\},$$

with

$$x_T \in r^{\frac{1}{2}}\mathbb{Z}^2 \cap (\mathbb{R} \times [0, R]), \quad v_T \in r^{-\frac{1}{2}}\mathbb{Z}^2 \cap Q'_i, \quad (1, v_T) \perp S_i.$$

Lemma 3.2 (Waveguide wave packet decomposition at scale r). *Let $1 \leq \sqrt{r} \leq R$ and the sets $Q'_i \subset \mathbb{R} \times \mathbb{Z}_{1/R}$. For each $h_i \in L^2(Q'_i)$, there exists a decomposition*

$$(25) \quad E_{Q'_i}^R h_i(x, t) = \sum_{T_i} c_{T_i} \phi_{T_i}(x, t)$$

where T_i ranges over all S_i -tubes. This decomposition satisfies the following properties: for each fixed tube T_i and time $t \in [\frac{r}{2}, r]$, the function $\phi_{T_i}(t)$ has Fourier support in the set

$$\{\xi \in \mathbb{R}^2 : \xi = v_T + O(r^{-\frac{1}{2}})\}$$

and obeys the pointwise decay estimates

$$(26) \quad |\phi_{T_i}(x, t)| \lesssim r^{-\frac{1}{2}} \left(1 + \frac{|x - (x_T + tv_T)|}{r^{\frac{1}{2}}}\right)^{-N}, \quad \forall N > 0;$$

then for any collection \mathcal{T}_i of S_i -tubes, there holds the following probability estimate

$$(27) \quad \left\| \sum_{T_i \in \mathcal{T}_i} \phi_{T_i}(x, t) \right\|_{L^2}^2 \lesssim \#\mathcal{T}_i;$$

and the complex-valued coefficients c_{T_i} obey the periodic property

$$(28) \quad |c_{T_i}| = |c_{T_{i,m}}|, \quad T_{i,m} := T_i + (0, mR, 0) \quad \text{with } m \in \mathbb{Z};$$

moreover, by periodization, if we rewrite this decomposition (25) as follows

$$(29) \quad E_{Q_i}^R h_i(x, t) = \sum_{T_i} \tilde{c}_{T_i} \psi_{T_i}(x, t), \quad \psi_{T_i}(x, t) := e^{ixv_T + it|v_T|^2} \sum_{m \in \mathbb{Z}} \phi_{T_i, m}(x, t),$$

where T_i ranges over all \tilde{S}_i -tubes, then the single-period coefficients \tilde{c}_{T_i} obey the ℓ^2 bound

$$(30) \quad \|\tilde{c}_{T_i}\|_{\ell^2} \lesssim \|h_i\|_{L^2}.$$

Remark 3.3. By the waveguide wave packet decomposition Lemma 3.2, for the desired local estimate Proposition 3.1, the case $r \lesssim R$ can be viewed as Euclidean case. The key observation is that there are only $O(1)$ single-period \tilde{S}_i -tubes in the decomposition (25), and all the Euclidean properties similarly hold on waveguide.

Proof of Lemma 3.2. We imitate the proof for classical Euclidean version wave packet decomposition, which can be seen in [Dem20, Section 2]. Consider a smooth function $\Upsilon : [-4, 4]^2 \rightarrow [0, \infty)$ such that

$$\sum_{\eta \in \mathbb{Z}^2} \Upsilon(\xi - \eta) = 1, \quad \forall \xi \in \mathbb{R}^2.$$

Then for each tube T with the position $x_T \in r^{\frac{1}{2}}\mathbb{Z}^2$ and velocity $v_T \in r^{-\frac{1}{2}}\mathbb{Z}^2$, we define the associated function

$$\psi_T(x, t) := r^{-\frac{1}{2}} e^{ixv_T + it|v_T|^2} \int_{\mathbb{R} \times \mathbb{Z}_{\sqrt{r}/R}} e^{\frac{i(x-x_T+tv_T)\xi}{\sqrt{r}} + \frac{it|\xi|^2}{r}} \Upsilon(\xi) d\xi,$$

and the associated coefficient

$$\tilde{c}_T := r^{-\frac{1}{2}} e^{-ix_T v_T} \int_{\mathbb{R} \times \mathbb{Z}_{1/R}} e^{-ix_T \xi} h(\xi) \Upsilon(r^{\frac{1}{2}}(\xi - v_T)) d\xi.$$

Note that $\psi_T(x_1, x_2, t) = \psi_T(x_1, x_2 + mR, t)$ up to the constant 2π in the phase function. Denote the direction $e_2 := (0, 1, 0)$. By the Poisson summation formula, one can obtain

$$\psi_T(x, t) = e^{ixv_T + it|v_T|^2} \sum_m \phi_{T_m}(x, t), \quad \phi_{T_m} := r^{-\frac{1}{2}} \int_{\mathbb{R}^2} e^{\frac{i(x-x_T+tv_T-mRe_2)\xi}{\sqrt{r}} + \frac{it|\xi|^2}{r}} \Upsilon(\xi) d\xi.$$

Then by the partition of unit and Fourier series theory, we obtain the desired wave packet decomposition (25) and (29). Applying the non-stationary phase estimate and the $\mathbb{R} \times \mathbb{T}_R$ version of Parseval identity, one can check all these desired properties. Hence the proof is finished. \square

Notice the following implication

$$r \leq R^{\frac{2p-2}{3-p}} \quad \Rightarrow \quad r^{\frac{5-3p}{2p}} (rR^{-1})^{\frac{1}{p'}} \leq 1.$$

Thus to show Proposition 3.1, recalling the Remark 3.3, it suffices to show the following lemma.

Lemma 3.4. *Given $p \in [\frac{5}{3}, 2]$ and $R \gg 1$. For the dyadic cubes*

$$Q_1 \in R^{-1}\mathcal{Q}_R, \quad Q_2 \in R^{-1}\mathcal{Q}_R, \quad \text{dist}(Q_1, Q_2) \gg 1,$$

we have the following bilinear estimate

$$(31) \quad \|E_{Q_1}^R h_1 E_{Q_2}^R h_2\|_{L_{x,t}^p(B_r)} \lesssim_\alpha r^\alpha r^{\frac{5-3p}{2p}} (rR^{-1})^{\frac{1}{p'}} \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z}_{1/R})}, \quad \forall r \in [R, R^{\frac{2p-2}{3-p}}].$$

3.3. Local estimate: tube version. In this subsection, we reduce the desired local bilinear integral estimate (31) to a tube-type incidence estimate. Note that the waveguide version wave packet decomposition satisfies the periodic property (29) and single-period bound (30). Therefore, we only need to consider the periodic collections \mathcal{T}_i of S_i -tubes, which means

$$T \in \mathcal{T}_i \quad \Rightarrow \quad T + (0, mR, 0) \in \mathcal{T}_i, \quad \forall m \in \mathbb{Z}.$$

For a subset $B \subset \mathbb{R}^3$, we introduce the intersection notation

$$\mathcal{T}_{i,B} := \{T : T \in \mathcal{T}_i, T \cap B \neq \emptyset\}.$$

Applying the waveguide version wave packet decomposition Lemma 3.2 and some dyadic pigeonholing arguments as shown in [Tao03, page 1369-1370], to establish the desired bilinear estimate (31), it suffices to establish the following single-period incidence estimate

$$(32) \quad \left\| \sum_{T_1 \in \mathcal{T}_1} \sum_{T_2 \in \mathcal{T}_2} \phi_{T_1} \phi_{T_2} \right\|_{L^p(B_r)} \lesssim r^{\frac{5-3p}{2p}} (rR^{-1})^{\frac{1}{p'}} (\#\mathcal{T}_{1,B_r})^{1/2} (\#\mathcal{T}_{2,B_r})^{1/2}, \quad \forall r \in [R, R^{\frac{2p-2}{3-p}}]$$

Here the notation $A \lesssim B$ means that $A \leq C_\alpha r^\alpha B$ holds for all $\alpha > 0$.

At the end of this subsection, we point out that our waveguide problem has been reduced to an Euclidean problem now. This is because that the wave packets ϕ_{T_i} can be formally viewed as characteristic function of Euclidean tubes, and the waveguide cubes B_r are indeed Euclidean cuboids (all these objects are in the Euclidean situation now).

3.4. Periodic property and the bilinear estimate. As mentioned above, by applying the waveguide version wave packet decomposition, one can reduce a waveguide problem to an associated Euclidean problem. Therefore, based on this reduction, one can apply the classical Euclidean-type bilinear restriction estimate to establish an associated waveguide-type bilinear estimate.

However, compared with the Euclidean case, one can see that the waveguide cubes B_r appeared in (32) is different from the standard cubes in \mathbb{R}^3 . To overcome this problem, we make use of the aforementioned periodic property of \mathcal{T}_i , so that we can extend the integral range B_r to the range $[0, r]^3$. Indeed, by the periodic property of \mathcal{T}_i , to show the desired single-period incidence estimate (32), it suffices to establish the following multiple-period incidence estimate

$$(33) \quad \left\| \sum_{T_1 \in \mathcal{T}_1} \sum_{T_2 \in \mathcal{T}_2} \phi_{T_1} \phi_{T_2} \right\|_{L^p([0,r]^3)} \lesssim r^{\frac{5-3p}{2p}} (\#\mathcal{T}_{1,[0,r]^3})^{1/2} (\#\mathcal{T}_{2,[0,r]^3})^{1/2}, \quad \forall r \in [R, R^{\frac{2p-2}{3-p}}].$$

Next we will explain that this desired estimate essentially follows from Tao's bilinear estimate. In his paper [Tao03], Tao mainly states his result for the crucial index $p = \frac{5}{3}$, which indeed comes from interpolation with the L^1 -estimate and L^2 -estimate. Here we restate Tao's bilinear restriction result for general index $p \in [1, 2]$. In fact, Tao's bilinear estimate on wave packets can be stated as follows

$$(34) \quad \left\| \sum_{T_1 \in \mathcal{T}'_1} \sum_{T_2 \in \mathcal{T}'_2} \phi_{T_1} \phi_{T_2} \right\|_{L^p([0,r]^3)} \lesssim r^{\frac{5-3p}{2p}} (\#\mathcal{T}'_{1,[0,r]^3})^{1/2} (\#\mathcal{T}'_{2,[0,r]^3})^{1/2}, \quad \forall r \in [R, R^{\frac{2p-2}{3-p}}].$$

Here \mathcal{T}'_i is an arbitrary collection of S_i -tubes (may be not periodic), and the notation $\mathcal{T}'_{i,[0,r]^3}$ is parallelly defined as

$$\mathcal{T}'_{i,[0,r]^3} := \{T : T \in \mathcal{T}'_i, T \cap [0, r]^3 \neq \emptyset\}.$$

Finally, it can be seen that Tao's bilinear estimate on wave packets (34) directly gives our desired estimate (33), and thus we complete the proof.

APPENDIX A: WAVEGUIDE VERSION EPSILON-REMOVAL LEMMA

Here we show a sketch proof of the ε -removal lemma in our waveguide setting. To begin with, we introduce the rescaled waveguide cubes B_r^R in $\mathbb{R} \times [0, 1] \times [0, T]$ as follows

$$B_r^R := [0, rR^{-1}] \times [0, rR^{-1}] \times [0, rR^{-2}] \quad \text{for } r \leq R, \quad B_r^R := [0, rR^{-1}] \times [0, 1] \times [0, rR^{-2}] \quad \text{for } r > R.$$

We consider $r \in [1, R^2T]$ to be dyadic numbers. Then our ε -removal lemma can be stated as follows.

Lemma 3.5. *For $p_0 \in [\frac{5}{3}, 2)$ and $R^{-2} \lesssim T \leq 1$, if there holds the local bilinear estimate*

$$(35) \quad \|E_{Q_1^*} h_1 E_{Q_2^*} h_2\|_{L_{x,t}^{p_0}(B_r^R)} \lesssim_\alpha r^\alpha R^{2-\frac{4}{p_0}} \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z})}, \quad \forall r \in [1, R^2T], \quad \forall \alpha > 0,$$

then there holds the following global estimate

$$\|E_{Q_1} h_1 E_{Q_2} h_2\|_{L_{x,t}^p(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim R^{2-\frac{4}{p}} \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z})}, \quad \forall p \in (p_0, 2].$$

Proof of Lemma 3.5. The proof mainly relies on two facts: the Fourier transform of waveguide-paraboloid measure has some decay, and the L^2 - L^4 Strichartz estimate on the waveguide $\mathbb{R} \times \mathbb{T}$ (see [TT01]). Here we adapt some ideas from [TV00, Lemma 2.4].

Since the estimate

$$\|E_{Q_1^*} h_1 E_{Q_2^*} h_2\|_{L_{x,t}^\infty(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim R^2 \|h_1\|_{L^2(\mathbb{R} \times \mathbb{Z})} \|h_2\|_{L^2(\mathbb{R} \times \mathbb{Z})}$$

holds, it suffices to show the following weak type estimate

$$|B| \lesssim \lambda^{-p} R^{-4}, \quad B := \{(x, t) \in \mathbb{R} \times \mathbb{T} \times [0, T] : |E_{Q_1} h_1 E_{Q_2} h_2| > \lambda R^2\}$$

holds for all $0 < \lambda \lesssim 1$ and $\|h_1\|_{L^2} = \|h_2\|_{L^2} = 1$. Moreover, it suffices to consider the case $|B| \gtrsim R^{-4}$. By following the arguments in [TV00, Proof of Lemma 2.4], for fixed (λ, h_1, h_2) which means fixed B , it suffices to show the following bilinear estimate

$$\|\chi_B E_{Q_1} g_1 E_{Q_2} g_2\|_{L_{x,t}^1} \lesssim R^{2-\frac{4}{p}} |B|^{1-\frac{1}{p}} \|g_1\|_{L^2} \|g_2\|_{L^2}, \quad \forall g_1 \in L^2, \quad \forall g_2 \in L^2.$$

Similarly, by applying some duality arguments, for fixed g_2 with $\|g_2\|_{L^2} \sim 1$, it suffices to show

$$(36) \quad \left\langle \widetilde{F} * E\chi_{Q_1}, \widetilde{F} \right\rangle_{L_{x,t}^2} \lesssim R^{4-\frac{8}{p}} |B|^{2-\frac{2}{p}}, \quad \|F\|_{L^\infty} \lesssim 1, \quad \widetilde{F} := \chi_B \overline{E_{Q_2} g_2} F.$$

Next we divide the time interval into three regions

$$I_1 := \{t : |t| \gg T\}, \quad I_2 := \{t : |t| \gg M\}, \quad I_3 := \{t : |t| \lesssim M, |t| \lesssim T\}.$$

Here the constant M will be determined later. Choose a non-negative Schwartz function ϕ satisfying

$$\phi : \mathbb{R} \rightarrow \mathbb{R}, \quad \text{supp} \phi \subset [-1, 1], \quad \phi \equiv 1 \quad \text{on} \quad \left[-\frac{1}{2}, \frac{1}{2}\right], \quad \widehat{\phi} \gtrsim 1 \quad \text{on} \quad [-1, 1].$$

The fact that $\text{supp}(\widetilde{F}) \subset \mathbb{R} \times \mathbb{T} \times [0, T]$ implies that the first region I_1 can be computed as follows

$$\left\langle \widetilde{F} * \left(E\chi_{Q_1} \left(1 - \phi \left(\frac{t}{2^{10}T} \right) \right) \right), \widetilde{F} \right\rangle_{L^2_{x,t}} = 0.$$

Then we consider the second region I_2 , where we need two facts: the scale-invariant Strichartz estimate and the Fourier transform of surface measure has some decay. First, the Strichartz estimate on $\mathbb{R} \times \mathbb{T}$ and Hölder's inequality implies that

$$\|\widetilde{F}\|_{L^1} \lesssim \|F\|_{L^\infty} \|\chi_B\|_{L^{\frac{4}{3}}} \|\chi_{\mathbb{R} \times \mathbb{T} \times [0, T]} E_{Q_2} g_2\|_{L^4} \lesssim |B|^{\frac{3}{4}}.$$

Second, the Van der Corput lemma on the first variable ξ_1 can imply

$$\|E\chi_{Q_1} \left(1 - \phi \left(\frac{t}{M} \right) \right)\|_{L^\infty} \lesssim M^{-\frac{1}{2}} R.$$

Note that our decay in $\mathbb{R} \times \mathbb{T}$ is $M^{-\frac{1}{2}}$ rather than M^{-1} in the Euclidean case. However, as we will see later, this decay is enough to establish the desired epsilon-removal consequence. Combining these aforementioned two estimates, we conclude

$$\left\langle \widetilde{F} * \left(E\chi_{Q_1} \left(1 - \phi \left(\frac{t}{M} \right) \right) \right), \widetilde{F} \right\rangle_{L^2_{x,t}} \lesssim |B|^{\frac{3}{2}} M^{-\frac{1}{2}} R.$$

Recalling the desired estimate (36), we can choose $M := |B|^{4/p-1} R^{16/p-6}$ and then consider the third region I_3 . Here we introduce the notation

$$r_0 := \min \left\{ 2^{10} T R^2, (|B| R^4)^{4/p-1} \right\},$$

it remains to show the following

$$(37) \quad \left\langle \widetilde{F} * \left(E1_{Q_1} \phi \left(\frac{t}{r_0 R^{-2}} \right) \right), \widetilde{F} \right\rangle_{L^2} \lesssim R^{4-\frac{8}{p}} |B|^{2-\frac{2}{p}}.$$

Next we are going to use the assumption (35) with $r \sim r_0$ to prove this result. Notice that the index p in the right hand side of (37) is strict larger than p_0 . Hence, we can take a suitable α in (35) to show the desired estimate (37). The details are shown as follows.

Let \mathcal{F} denote the space-time Fourier transform for (x, t) . Define

$$\Gamma_{i,\gamma} = \{(\xi, \tau) : \xi \in Q_i, |r_0 R^{-2}(\tau - |\xi|^2)| \sim \gamma\}, \quad i = 1, 2.$$

For $(\xi, \tau) \in \Gamma_{1,\gamma}$,

$$|\mathcal{F} \left(E1_{Q_1} \phi \left(\frac{t}{r_0 R^{-2}} \right) \right) (\xi, \tau)| = r_0 R^{-2} 1_{Q_1}(\xi) |\widehat{\phi}(r_0 R^{-2}(\tau - |\xi|^2))| \lesssim_N r_0 R^{-2} \gamma^{-N}.$$

So

$$\begin{aligned} \left\langle \widetilde{F} * \left(E1_{Q_1} \phi \left(\frac{t}{r_0 R^{-2}} \right) \right), \widetilde{F} \right\rangle_{L^2} &\lesssim \int |\mathcal{F}(\widetilde{F})|^2 |\mathcal{F} \left(E1_{Q_1} \phi \left(\frac{t}{r_0 R^{-2}} \right) \right)| \\ &\lesssim_N \sum_{\gamma_1} \int_{\Gamma_{1,\gamma_1}} |\mathcal{F}(\widetilde{F})|^2 r_0 R^{-2} \gamma_1^{-N}, \end{aligned}$$

it suffices to show that

$$\left(\int_{\Gamma_{1,\gamma_1}} |\mathcal{F}(\widetilde{F})|^2 \right)^{\frac{1}{2}} \lesssim \gamma_1^N r_0^{-1/2} R^{3-\frac{4}{p}} |B|^{1-\frac{1}{p}}$$

for all $\gamma_1 \geq 1$ and some $N \geq 1$. From the definition of \widetilde{F} , this will follow if we show

$$\|1_B E_{Q_2} g_2 F\|_{L^2(\Gamma_{1,\gamma_1})} \lesssim \gamma_1^N r_0^{-1/2} R^{3-\frac{4}{p}} |B|^{1-\frac{1}{p}} \|F\|_{L^\infty} \|g_2\|_{L^2},$$

for all F, g_2 . By duality, it suffices to prove

$$\|1_B \mathcal{F}(G_1) E_{Q_2} g_2\|_{L^1} \lesssim \gamma_1^N r_0^{-1/2} R^{3-\frac{4}{p}} |B|^{1-\frac{1}{p}} \|G_1\|_{L^2_{\xi,\tau}} \|g_2\|_{L^2_{\xi,\tau}}$$

for all g_2 and all G_1 supported on Γ_{1,γ_1} .

We repeat the similar argument on g_2 when we fix G_1 , so it suffices to show that

$$\|1_B \mathcal{F}(G_1) \mathcal{F}(G_2)\|_{L^1} \lesssim (\gamma_1 \gamma_2)^N r_0^{-1} R^{4-\frac{4}{p}} |B|^{1-\frac{1}{p}} \|G_1\|_{L^2_{\xi,\tau}} \|G_2\|_{L^2_{\xi,\tau}}$$

for all $\gamma_i \geq 1$, G_i supported on Γ_{i,γ_i} , $i = 1, 2$ and some $N \geq 1$. By the Hölder inequality and $B \subset A$, it suffices to show that

$$\|1_B \mathcal{F}(G_1) \mathcal{F}(G_2)\|_{L^{p_0}} \lesssim (\gamma_1 \gamma_2)^N r_0^{-1} R^{4-\frac{4}{p}} |B|^{\frac{1}{p_0}-\frac{1}{p}} \|G_1\|_{L^2_{\xi,\tau}} \|G_2\|_{L^2_{\xi,\tau}}.$$

Choose $r \in [1, TR^2]$ such that $r \sim r_0$. We break the proof into two cases.

Case 1: there holds $r < R$. Choose a function $\psi_r : \mathbb{Z} \rightarrow \mathbb{R}$ satisfying

$$\text{supp } \psi_r \subset [-r^{-1}R, r^{-1}R] \cap \mathbb{Z}, \quad \psi_r \equiv 1 \text{ on } \left[-\frac{r^{-1}R}{2}, \frac{r^{-1}R}{2}\right], \quad \widehat{\psi_r} \gtrsim 1 \text{ on } [-rR^{-1}, rR^{-1}].$$

So

$$\text{supp} \left(\mathcal{F}^{-1} \left(\widehat{\phi} \left(\frac{x_1 - \widetilde{x}_1}{rR^{-1}} \right) \widehat{\psi_r}(x_2 - \widetilde{x}_2) \widehat{\phi} \left(\frac{t - \widetilde{t}}{rR^{-2}} \right) \mathcal{F}(G_i) \right) \right) \subset N_{\gamma_i r^{-2} R^2}(S_i^*), \quad i = 1, 2,$$

for any $(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t}) \in \mathbb{R} \times \mathbb{T} \times \mathbb{R}$. Here we have used the notations

$$(38) \quad S_i^* := \{(\xi, |\xi|^2) : \xi \in Q_i^*\}, \quad i = 1, 2,$$

and

$$(39) \quad N_\gamma(S_i^*) := \{(\xi, \tau) \in (\mathbb{R} \times \mathbb{Z}) \times \mathbb{R} : \xi \in Q_i^*, |\tau - |\xi|^2| \lesssim \gamma\}, \quad i = 1, 2.$$

By the Cauchy-Schwarz inequality, the Lemma 3.6 stated below and the fact $B \subset \mathbb{R} \times \mathbb{T} \times [0, T]$, we conclude

$$\begin{aligned} \|1_B \mathcal{F}(G_1) \mathcal{F}(G_2)\|_{L^{p_0}} &\leq \sum_{(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t})} \|\mathcal{F}(G_1) \mathcal{F}(G_2)\|_{L^{p_0}(B_r^R+(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t}))} \\ &\lesssim \sum_{(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t})} \left\| \left(\widehat{\phi} \left(\frac{x_1 - \widetilde{x}_1}{rR^{-1}} \right) \widehat{\psi_r}(x_2 - \widetilde{x}_2) \widehat{\phi} \left(\frac{t - \widetilde{t}}{rR^{-2}} \right) \right)^2 \mathcal{F}(G_1) \mathcal{F}(G_2) \right\|_{L^{p_0}(B_r^R+(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t}))} \\ &\lesssim_\alpha \sum_{(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t})} R^{2-\frac{4}{p_0}} r^\alpha \prod_{i=1}^2 \left((\gamma_i r^{-2} R^2)^{\frac{1}{2}} \|\widehat{\phi} \left(\frac{x_1 - \widetilde{x}_1}{rR^{-1}} \right) \widehat{\psi_r}(x_2 - \widetilde{x}_2) \widehat{\phi} \left(\frac{t - \widetilde{t}}{rR^{-2}} \right) \mathcal{F}(G_i)\|_{L^2} \right) \end{aligned}$$

$$\lesssim R^{2-\frac{4}{p_0}} r^\alpha (\gamma_1 \gamma_2)^{\frac{1}{2}} r^{-1} R^2 \|G_1\|_{L^2} \|G_2\|_{L^2},$$

where the $(\widetilde{x}_1, \widetilde{x}_2, \widetilde{t})$ is in

$$\left\{ (k_1 r R^{-1}, k_2 r R^{-1}, k_3 r R^{-2}) : k_1, k_2, k_3 \in \mathbb{Z}, 0 \leq k_2 \lesssim \frac{1}{r R^{-1}}, 0 \leq k_3 \lesssim \frac{T}{r R^{-2}} \right\}.$$

Recalling the assumption $|B| \gtrsim R^{-4}$, we will be done if

$$r^\alpha \sim r_0^\alpha \lesssim (|B|R^4)^{\frac{2}{p_0} - \frac{2}{p}}.$$

From the definition of r_0 , we only need to choose $\alpha < (\frac{2}{p} - \frac{1}{2})^{-1} (\frac{1}{p_0} - \frac{1}{p})$.

Case 2: There holds $r \geq R$. Note that

$$\text{supp} \left(\mathcal{F}^{-1} \left(\widehat{\phi} \left(\frac{x_1 - \widetilde{x}_1}{r R^{-1}} \right) \widehat{\phi} \left(\frac{t - \widetilde{t}}{r R^{-2}} \right) \mathcal{F}(G_i) \right) \right) \subset N_{\gamma_i r^{-2} R^2}(S_i^*), \quad i = 1, 2,$$

and the rest of the discussion is similar. \square

Recalling the notations in (38) and (39), it remains to prove the following thick-bilinear restriction estimate.

Lemma 3.6. *Given $\gamma_1, \gamma_2 > 0$. Suppose the local bilinear estimate (35) holds, then we have*

$$\left\| \mathcal{F}^{-1}(F_1) \mathcal{F}^{-1}(F_2) \right\|_{L^{p_0}(B_r^R)} \lesssim_\alpha R^{2-\frac{4}{p_0}} r^\alpha \gamma_1^{\frac{1}{2}} \gamma_2^{\frac{1}{2}} \|F_1\|_{L^2(N_{\gamma_1}(S_1^*))} \|F_2\|_{L^2(N_{\gamma_2}(S_2^*))}$$

holds for all F_i supported on $N_{\gamma_i}(S_i^*)$, $i = 1, 2$, where \mathcal{F}^{-1} is the space-time inverse Fourier transform.

Proof. Direct computations can give that

$$\begin{aligned} \mathcal{F}^{-1}(F_i)(x, t) &= \int_{N_{\gamma_i}(S_i^*)} e^{ix\xi + it\tau} F_i(\xi, \tau) d\xi d\tau \\ &= \int_{-\gamma_i}^{\gamma_i} \int_{Q_i^*} e^{ix\xi + it|\xi|^2} e^{it\tau_i} F_i(\xi, |\xi|^2 + \tau_i) d\xi d\tau_i \\ &= \int_{-\gamma_i}^{\gamma_i} e^{it\tau_i} (E_{Q_i^*} h_{i, \tau_i})(x, t) d\tau_i, \end{aligned}$$

where $h_{i, \tau_i}(\xi) = F_i(\xi, |\xi|^2 + \tau_i)$, $i = 1, 2$. Then by the Cauchy-Schwarz inequality, the Minkowski inequality and the assumption (35), we have

$$\begin{aligned} \left\| \mathcal{F}^{-1}(F_1) \mathcal{F}^{-1}(F_2) \right\|_{L^{p_0}(B_r^R)} &= \left\| \int_{-\gamma_1}^{\gamma_1} \int_{-\gamma_2}^{\gamma_2} e^{it(\tau_1 + \tau_2)} E_{Q_1^*} h_{1, \tau_1} E_{Q_2^*} h_{2, \tau_2} d\tau_2 d\tau_1 \right\|_{L^{p_0}(B_r^R)} \\ &\leq \int_{-\gamma_1}^{\gamma_1} \int_{-\gamma_2}^{\gamma_2} \|E_{Q_1^*} h_{1, \tau_1} E_{Q_2^*} h_{2, \tau_2}\|_{L^{p_0}(B_r^R)} d\tau_2 d\tau_1 \\ &\lesssim_\alpha R^{2-\frac{4}{p_0}} r^\alpha \int_{-\gamma_1}^{\gamma_1} \int_{-\gamma_2}^{\gamma_2} \|h_{1, \tau_1}\|_{L^2(Q_1^*)} \|h_{2, \tau_2}\|_{L^2(Q_2^*)} d\tau_2 d\tau_1 \\ &\lesssim_\alpha R^{2-\frac{4}{p_0}} r^\alpha \gamma_1^{\frac{1}{2}} \gamma_2^{\frac{1}{2}} \|F_1\|_{L^2(N_{\gamma_1}(S_1^*))} \|F_2\|_{L^2(N_{\gamma_2}(S_2^*))}. \end{aligned}$$

This completes the proof. \square

Remark 3.7. For higher dimensional waveguide $\mathbb{R}^m \times \mathbb{T}^n$ with $m \geq 1$, the corresponding epsilon-removal result still holds by applying some noncritical scale-invariant Strichartz estimates, see for instance [Bar21, KV16].

APPENDIX B: SOME EXAMPLES FOR EXPLAINING THE SHARPNESS

Here we give some examples to show that our conclusion Corollary 1.3 is sharp up to the ε -loss. These three examples are shown as follows.

Let $C(R, T, p)$ be the sharp constant that

$$\|E_{Q_1} h_1 E_{Q_2} h_2\|_{L^p_{x,t}(\mathbb{R} \times \mathbb{T} \times [0, T])} \lesssim C(R, T, p) \|h_1\|_{L^2} \|h_2\|_{L^2}, \quad \forall h_1, h_2 \in L^2(\mathbb{R} \times \mathbb{Z}).$$

We only consider the situation $T \geq R^{-2}$.

First, we choose the following

$$Q_1 = [0, R] \times \{0, 1, \dots, R\}, \quad Q_2 = [0, R] \times \{100R, 100R + 1, \dots, 101R\}, \quad h_1 = \chi_{Q_1}, \quad h_2 = \chi_{Q_2},$$

then

$$|E_{Q_1} h_1(x, t)| \gtrsim R^2, \quad |E_{Q_2} h_2(x, t)| \gtrsim R^2, \quad \forall (x, t) \in \{|x| \lesssim R^{-1}, |t| \lesssim R^{-2}\}.$$

These implies the constant $C(R, T, p)$ satisfies

$$C(R, T, p) \gtrsim R^{\frac{2p-4}{p}}.$$

Second, we choose the following

$$Q_1 = [0, T^{-1/2}] \times \{0\}, \quad Q_2 = [0, T^{-1/2}] \times \{100R\}, \quad h_1 = \chi_{Q_1}, \quad h_2 = \chi_{Q_2},$$

then

$$|E_{Q_1} h_1(x, t)| \gtrsim T^{-1/2}, \quad |E_{Q_2} h_2(x, t)| \gtrsim T^{-1/2}, \quad \forall (x, t) \in \{|x_1| \lesssim T^{1/2}, x_2 \in \mathbb{T}, |t| \lesssim T\}.$$

These implies the constant $C(R, T, p)$ satisfies

$$C(R, T, p) \gtrsim T^{\frac{3-p}{2p}}.$$

Third, we choose the following

$$Q_1 = [0, a] \times ([0, a^2 R^{-1}] \cap \mathbb{Z}), \quad Q_2 = [0, a] \times ([100R, 100R + a^2 R^{-1}] \cap \mathbb{Z}), \quad h_1 = \chi_{Q_1}, \quad h_2 = \chi_{Q_2},$$

where $a = \max\{R^{1/2}, T^{-1/2}\}$, then

$$|E_{Q_1} h_1(x, t)| \gtrsim a^3 R^{-1}, \quad |E_{Q_2} h_2(x, t)| \gtrsim a^3 R^{-1}, \quad \forall (x, t) \in S_a,$$

where the set S_a is defined as

$$S_a := \{|x_1| \lesssim a^{-1}, |x_2| \lesssim a^{-2} R, |x_2 + 200Rt| \lesssim a^{-2} R, |t| \lesssim a^{-2}\}.$$

These implies

$$C(R, T, p) \gtrsim (R + T^{-1})^{\frac{3p-5}{2p}} R^{\frac{1-p}{p}}.$$

In summary, we have proved the sharpness of our Corollary 1.3 up to the ε -loss.

REFERENCES

- [Bar21] Alex Barron. On global-in-time Strichartz estimates for the semiperiodic Schrödinger equation. *Anal. PDE*, 14(4):1125–1152, 2021.
- [BCP21] Alex Barron, Michael Christ, and Benoit Pausader. Global endpoint Strichartz estimates for Schrödinger equations on the cylinder $\mathbb{R} \times \mathbb{T}$. *Nonlinear Analysis*, 206:112172, 2021.
- [BD15] Jean Bourgain and Ciprian Demeter. The proof of the l^2 decoupling conjecture. *Ann. of Math. (2)*, 182(1):351–389, 2015.
- [BG99] Hajer Bahouri and Patrick Gérard. High frequency approximation of solutions to critical nonlinear wave equations. *American Journal of Mathematics*, 121(1):131–175, 1999.
- [Bou93] Jean Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations. *Geom. Funct. Anal.*, 3(2):107–156, 1993.
- [Bou98] Jean Bourgain. Refinements of Strichartz inequality and applications to 2D-NLS with critical nonlinearity. *Internat. Math. Res. Notices*, 5:253–283, 1998.
- [Bou16] Jean Bourgain. A note on the Schrödinger maximal function. *J. Anal. Math.*, 130:393–396, 2016.
- [BV07] Pascal Bégout and Ana Vargas. Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation. *Trans. Amer. Math. Soc.*, 359(11):5257–5282, 2007.
- [Caz03] Thierry Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- [CGSY23] Xianghong Chen, Zihua Guo, Minxing Shen, and Lixin Yan. On smoothing estimates for Schrödinger equations on product spaces $\mathbb{T}^m \times \mathbb{R}^n$. [arXiv:2301.05450](https://arxiv.org/abs/2301.05450), 2023.
- [Dem20] Ciprian Demeter. *Fourier restriction, decoupling, and applications*, volume 184 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2020.
- [Den23] Yangkendi Deng. On a bilinear restriction estimate for Schrödinger equations on 2D waveguide. [arXiv:2311.18541](https://arxiv.org/abs/2311.18541), 2023.
- [DFY⁺24] Yangkendi Deng, Chenjie Fan, Kailong Yang, Zehua Zhao, and Jiqiang Zheng. On bilinear Strichartz estimates on waveguides with applications. [arXiv:2402.02916](https://arxiv.org/abs/2402.02916), 2024.
- [DGL17] Xiumin Du, Larry Guth, and Xiaochun Li. A sharp Schrödinger maximal estimate in \mathbb{R}^2 . *Ann. of Math. (2)*, 186(2):607–640, 2017.
- [Dod12] Benjamin Dodson. Global well-posedness and scattering for the defocusing, mass-critical nonlinear Schrödinger equation when $d \geq 3$. *Journal of the American Mathematical Society*, 25(2):429–463, 2012.
- [Dod15] Benjamin Dodson. Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state. *Advances in Mathematics*, 285:1589–1618, 2015.
- [Dod16a] Benjamin Dodson. Global well-posedness and scattering for the defocusing, L^2 -critical, nonlinear Schrödinger equation when $d = 1$. *American Journal of Mathematics*, 138(2):531–569, 2016.
- [Dod16b] Benjamin Dodson. Global well-posedness and scattering for the defocusing, L^2 -critical, nonlinear Schrödinger equation when $d = 2$. *Duke Math. J.*, 165(18):3435–3516, 2016.
- [DSPST07] Daniela De Silva, Nataša Pavlović, Gigliola Staffilani, and Nikolaos Tzirakis. Global well-posedness for a periodic nonlinear Schrödinger equation in 1D and 2D. *Discrete Contin. Dyn. Syst.*, 19(1):37–65, 2007.
- [DZ19] Xiumin Du and Ruixiang Zhang. Sharp L^2 estimates of the Schrödinger maximal function in higher dimensions. *Ann. of Math. (2)*, 189(3):837–861, 2019.
- [FSWW18] Chenjie Fan, Gigliola Staffilani, Hong Wang, and Bobby Wilson. On a bilinear Strichartz estimate on irrational tori. *Anal. PDE*, 11(4):919–944, 2018.
- [Gut23] Larry Guth. Decoupling estimates in Fourier analysis. In *ICM—International Congress of Mathematicians. Vol. 2. Plenary lectures*, pages 1054–1089. EMS Press, Berlin, [2023] ©2023.
- [IP12] Alexandru D. Ionescu and Benoit Pausader. The energy-critical defocusing NLS on \mathbb{T}^3 . *Duke Mathematical Journal*, 2012.
- [IPS12] Alexandru D. Ionescu, Benoit Pausader, and Gigliola Staffilani. On the global well-posedness of energy-critical Schrödinger equations in curved spaces. *Anal. PDE*, 5(4):705–746, 2012.

- [KM06] Carlos E Kenig and Frank Merle. Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. *Inventiones mathematicae*, 166(3):645–675, 2006.
- [KT98] Markus Keel and Terence Tao. Endpoint Strichartz estimates. *Amer. J. Math.*, 120(5):955–980, 1998.
- [KV16] Rowan Killip and Monica Visan. Scale invariant Strichartz estimates on tori and applications. *Math. Res. Lett.*, 23(2):445–472, 2016.
- [MV98] Frank Merle and Luis Vega. Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D. *International Mathematics Research Notices*, 1998(8):399–425, 1998.
- [Tao01] Terence Tao. Endpoint bilinear restriction theorems for the cone, and some sharp null form estimates. *Math. Z.*, 238(2):215–268, 2001.
- [Tao03] Terence Tao. A sharp bilinear restrictions estimate for paraboloids. *Geom. Funct. Anal.*, 13(6):1359–1384, 2003.
- [TT01] Hideo Takaoka and Nikolay Tzvetkov. On 2D nonlinear Schrödinger equations with data on $\mathbb{R} \times \mathbb{T}$. *J. Funct. Anal.*, 182(2):427–442, 2001.
- [TV00] Terence Tao and Ana Vargas. A bilinear approach to cone multipliers. I. Restriction estimates. *Geom. Funct. Anal.*, 10(1):185–215, 2000.
- [TVV98] Terence Tao, Ana Vargas, and Luis Vega. A bilinear approach to the restriction and Keakeya conjectures. *J. Amer. Math. Soc.*, 11(4):967–1000, 1998.
- [Wol01] Thomas H. Wolff. A sharp bilinear cone restriction estimate. *Ann. of Math. (2)*, 153(3):661–698, 2001.
- [ZZ21] Zehua Zhao and Jiqiang Zheng. Long time dynamics for defocusing cubic nonlinear Schrödinger equations on three dimensional product space. *SIAM J. Math. Anal.*, 53(3):3644–3660, 2021.

YANGKENDI DENG

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES,
BEIJING, 100190, PEOPLE’S REPUBLIC OF CHINA
E-mail address: dengyangkendi@amss.ac.cn

BONING DI

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES,
BEIJING, 100190, PEOPLE’S REPUBLIC OF CHINA
E-mail address: diboning@amss.ac.cn

CHENJIE FAN

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES,
BEIJING, 100190, PEOPLE’S REPUBLIC OF CHINA
HUA LOO-KENG KEY LABORATORY OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES,
BEIJING, 100190, PEOPLE’S REPUBLIC OF CHINA
E-mail address: fancj@amss.ac.cn

ZEHUA ZHAO

DEPARTMENT OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY,
BEIJING, PEOPLE’S REPUBLIC OF CHINA
KEY LABORATORY OF ALGEBRAIC LIE THEORY AND ANALYSIS OF MINISTRY OF EDUCATION,
BEIJING, PEOPLE’S REPUBLIC OF CHINA
E-mail address: zzh@bit.edu.cn