

AN INVERSE HYPERBOLIC OBSTACLE PROBLEM

MOURAD CHOULLI AND HIROSHI TAKASE

ABSTRACT. We establish Hölder stability of an inverse hyperbolic obstacle problem. Mainly, we study the problem of reconstructing an unknown function defined on the boundary of the obstacle from two measurements taken on the boundary of a domain surrounding the obstacle.

1. INTRODUCTION AND MAIN RESULT

Let $n \geq 2$ be an integer. Throughout this text, we use the Einstein summation convention for quantities with indices. If in any term appears twice, as both an upper and lower index, that term is assumed to be summed from 1 to n .

Let $(g_{k\ell}) \in W^{2,\infty}(\mathbb{R}^n; \mathbb{R}^{n \times n})$ be a symmetric matrix-valued function satisfying

$$g_{k\ell}(x)\xi^k\xi^\ell \geq \kappa|\xi|^2 \quad x, \xi \in \mathbb{R}^n,$$

where $\kappa > 0$ is a constant. Note that $(g^{k\ell})$ the matrix inverse to g is uniformly positive definite as well. Recall that the Laplace-Beltrami operator associated with the metric tensor $g = g_{k\ell}dx^k \otimes dx^\ell$ is given by

$$\Delta_g u := \left(1/\sqrt{|g|}\right) \partial_k \left(\sqrt{|g|}g^{k\ell} \partial_\ell u\right),$$

where $|g| = \det(g)$.

For convenience, we recall the following usual notations,

$$\langle X, Y \rangle = g_{k\ell}X^kY^\ell, \quad X = X^k \frac{\partial}{\partial x^k}, \quad Y = Y^k \frac{\partial}{\partial x^k},$$

$$\nabla_g w = g^{k\ell} \partial_k w \frac{\partial}{\partial x^\ell},$$

$$|\nabla_g w|_g^2 = \langle \nabla_g w, \nabla_g w \rangle = g^{k\ell} \partial_k w \partial_\ell w.$$

Also, recall that the Christoffel symbols $\Gamma_{k\ell}^m$, $1 \leq k, \ell, m \leq n$, are given as follows

$$\Gamma_{k\ell}^m := (1/2)g^{jm}(\partial_k g_{j\ell} + \partial_\ell g_{jk} - \partial_j g_{k\ell}),$$

and the Hessian matrix $\nabla_g^2 w$ is defined by

$$(\nabla_g^2 w)_{k\ell} := \partial_k \partial_\ell w - \Gamma_{k\ell}^m \partial_m w.$$

If dx is the Lebesgue measure on \mathbb{R}^n , set $dV_g = \sqrt{|g|}dx$. Let O be an open subset of \mathbb{R}^n . We assume hereinafter that $L^2(O)$, $H^1(O)$ and $H^2(O)$ are endowed

Date: July 9, 2024.

2020 Mathematics Subject Classification. 35R30, 35L20, 58J45.

Key words and phrases. hyperbolic obstacle problem, inverse problem, Hölder stability, Carleman estimate.

respectively with the norms

$$\begin{aligned}\|w\|_2 &:= \left(\int_O |w|^2 dV_g \right)^{1/2}, \\ \|w\|_{1,2} &:= \left(\int_O [|w|^2 + |\nabla_g w|_g^2] dV_g \right)^{1/2}, \\ \|w\|_{2,2} &:= \left(\int_O [|w|^2 + |\nabla_g w|_g^2 + |\nabla_g^2 w|_g^2] dV_g \right)^{1/2}.\end{aligned}$$

Here

$$|\nabla_g^2 w|_g^2 := g^{k_1 \ell_1} g^{k_2 \ell_2} (\nabla_g^2 w)_{k_1 k_2} (\nabla_g^2 w)_{\ell_1 \ell_2}.$$

In the rest of this text, $\phi \in C^3(\mathbb{R}^n)$ satisfies the following properties

$$\inf_{\{\phi > 0\}} |\nabla_g \phi|_g > 0, \quad \inf_{\{\phi > 0\}} \inf_{|\xi|=1} \nabla_g^2 \phi(\xi, \xi) > 0$$

and $\{\phi < 0\} \subset \mathbb{R}^n$ is bounded. Let us give an example of such a function.

Example 1.1. We denote the unit ball of \mathbb{R}^n by \mathbb{B} and set $\mathbb{B}_r = r\mathbb{B}$, $r > 0$. For fixed $r_0 > 0$, let $\phi = |x|^2 - r_0^2$. Furthermore, assume that $g = \mathbf{I}$ in $\{|y| \geq r_1\}$, where $r_1 > r_0$ and \mathbf{I} is the identity matrix of \mathbb{R}^n . Then, we have $|\nabla_g \phi(x)|_g = 2|x| \geq 2r_1$ if $x \in \{|y| > r_1\}$ and, since g^{-1} is non singular, we have

$$|\nabla_g \phi(x)|_g > 0, \quad x \in \overline{\mathbb{B}_{r_1}} \setminus \mathbb{B}_{r_0}.$$

On the other hand, as $\nabla_g^2 \phi = 2\mathbf{I}$ in $\{|y| > r_1\}$, using again a compactness argument to show that $\nabla_g^2 \phi \geq \mathbf{I}$ provided that g has a sufficiently small norm in $W^{1,\infty}(\mathbb{R}^n, \mathbb{R}^{n \times n})$.

Before stating our main result, we introduce new definitions. Let $T > 0$, $B := \{\phi < 0\}$, $\Gamma := \{\phi = 0\}$ ($= \partial B$), $U := \mathbb{R}^n \setminus \overline{B}$ and $\Sigma := \Gamma \times (0, T)$.

Since we will consider an IBVP with nonhomogenous boundary condition, we introduce the space

$$\mathcal{Y} := \{(f, u_0, u_1) = (H|_\Sigma, H(\cdot, 0), \partial_t H(\cdot, 0)); H \in \mathcal{Z}\},$$

where

$$\mathcal{Z} := W^{3,1}((0, T); L^2(U)) \cap W^{2,1}((0, T); H^1(U)) \cap W^{1,1}((0, T); H^2(U)).$$

We equip \mathcal{Z} with its natural norm denoted by $\|\cdot\|_{\mathcal{Z}}$, and we observe that \mathcal{Z} is continuously embedded in \mathcal{X} , where

$$\mathcal{X} := C^2([0, T]; L^2(U)) \cap C^1([0, T]; H^1(U)) \cap C([0, T]; H^2(U))$$

with its natural norm that we denote by $\|\cdot\|_{\mathcal{X}}$. Define

$$\overbrace{(f, u_0, u_1)} := \{H \in \mathcal{Z}; (H|_\Sigma, H(\cdot, 0), \partial_t H(\cdot, 0)) = (f, u_0, u_1)\}.$$

We endow \mathcal{Y} with the quotient norm

$$\|(f, u_0, u_1)\|_{\mathcal{Y}} := \inf\{\|H\|_{\mathcal{Z}}; H \in \overbrace{(f, u_0, u_1)}\}.$$

Consider the nonhomogeneous IBVP

$$(1.1) \quad \begin{cases} (\partial_t^2 - \Delta_g)u = 0 & \text{in } U \times (0, T), \\ u|_\Sigma = f, \\ (u(\cdot, 0), \partial_t u(\cdot, 0)) = (u_0, u_1). \end{cases}$$

We prove in Appendix A that, for each $(f, u_0, u_1) \in \mathcal{Y}$, (1.1) admits a unique solution $u = u(f, u_0, u_1) \in \mathcal{X}$ and

$$(1.2) \quad \|u\|_{\mathcal{X}} \leq \mathbf{c} \|(f, u_0, u_1)\|_{\mathcal{Y}},$$

where $\mathbf{c} = \mathbf{c}(g, \phi) > 0$ is a constant.

Let $\Omega \ni B$ be an arbitrary fixed bounded domain with boundary $\Gamma_0 := \partial\Omega$. We aim to establish Hölder stability of the determination of the unknown function f from the measurements $u(f, 0, 0)|_{\Sigma_0}$ and $\partial_{\nu_g} u(f, 0, 0)|_{\Sigma_0}$, where $\Sigma_0 := \Gamma_0 \times (0, T)$. This problem can be seen as the determination of an unknown source emanating from the obstacle from two measurements performed far from the obstacle.

The same type of inverse problem was considered by Choulli-Takase [4] for elliptic and parabolic equations. See for example Isakov [5] and Bourgeois-Ponomarev-Dardé [1] for an inverse obstacle problem, which determines interior regions from finite-time Cauchy data. Recently, the first author obtained stability inequality for an inverse source problem for an obstacle problem for the wave equation. We refer to Choulli [3]. Other references have been mentioned in this work.

Henceforth, we assume that Ω is $C^{0,1}$ and we set $D := \Omega \setminus \overline{B}$ and $\Lambda := \partial D$. Define

$$\mathcal{F}_0 := \left\{ f = \mathbf{a} \otimes \mathbf{b}; \mathbf{a} \in H^{3/2}(\Gamma), \mathbf{b} \in W^{3,1}((0, T)), \mathbf{b}(0) = \mathbf{b}'(0) = 0 \right\}.$$

It is worth observing that

$$\{(f, 0, 0); f \in \mathcal{F}_0\} \subset \mathcal{Y}.$$

The following additional notations will be used in the sequel

$$\begin{aligned} \nu_g &= (\nu_g)^k \frac{\partial}{\partial x^k}, \quad (\nu_g)^k = \frac{g^{k\ell} \nu_\ell}{\sqrt{g^{ij} \nu_i \nu_j}}, \\ \partial_{\nu_g} w &= \langle \nu_g, \nabla_g w \rangle. \end{aligned}$$

Also, define $\nabla_{\tau_g} w$ by

$$\nabla_{\tau_g} w = \nabla_g w - (\partial_{\nu_g} w) \nu_g.$$

We check that

$$|\nabla_g w|_g^2 = |\nabla_{\tau_g} w|_g^2 + |\partial_{\nu_g} w|^2.$$

Let $\alpha > 0$ and $\beta > 0$ be fixed and define the admissible set of the unknown function f by

$$\mathcal{F} := \left\{ f = \mathbf{a} \otimes \mathbf{b} \in C^1(\Sigma) \cap \mathcal{F}_0; |\mathbf{a}| \geq \alpha \text{ and } |\nabla_{\tau_g} \mathbf{a}|_g \leq \beta \right\}.$$

Next, set

$$\mathcal{D} := \|u\|_{H^1(\Sigma)} + \|\partial_{\nu_g} u\|_{L^2(\Sigma)}$$

and

$$\mathcal{D}_0 := \|u\|_{H^1(\Sigma_0)} + \|\partial_{\nu_g} u\|_{L^2(\Sigma_0)},$$

where ν denotes the outer unit normal to Λ . Here $L^2(\Sigma)$ and $H^1(\Sigma)$ are endowed respectively with the norms

$$\begin{aligned} \|w\|_{L^2(\Sigma)} &:= \left(\int_{\Sigma} |w|^2 dS_g dt \right)^{1/2}, \\ \|w\|_{H^1(\Sigma)} &:= \left(\int_{\Sigma} [|w|^2 + |\partial_t w|^2 + |\nabla_{\tau_g} w|_g^2] dS_g dt \right)^{1/2}, \end{aligned}$$

and similarly for $L^2(\Sigma_0)$ and $H^1(\Sigma_0)$. Here $dS_g := \sqrt{|g|}dS$, where dS denotes the surface measure on Λ .

Finally, define $\mathbf{m} := \max_{\overline{D}} \phi$, $\delta := \min_{\overline{D}} |\nabla_g \phi|_g$ and $T^* := \mathbf{m}/\delta$.

Theorem 1.2. *Assume that $T > T^*$. Let $\tau \in (0, T - T^*)$ and $\zeta = (g, \phi, \tau, \beta/\alpha)$. Then there exist $C = C(\zeta) > 0$ and $\theta = \theta(\zeta) \in (0, 1)$ such that for any $f \in \mathcal{F}$ we have*

$$\|f\|_{L^2((0,\tau);H^1(\Gamma))} \leq C (\mathcal{D}_0 + \mathcal{D}^{1-\theta} \mathcal{D}_0^\theta).$$

If \mathbf{b} is known and satisfies

$$\|\mathbf{b}\|_{L^2((0,\tau))} > 0,$$

where $\tau \in (0, T - T^*)$ is as in Theorem 1.2, we derive the following result.

Corollary 1.3. *Assume that $T > T^*$. Let $\tau \in (0, T - T^*)$ and $\zeta = (g, \phi, \tau, \beta/\alpha)$. Then there exist $C = C(\zeta) > 0$ and $\theta = \theta(\zeta) \in (0, 1)$ such that for any $f = \mathbf{a} \otimes \mathbf{b} \in \mathcal{F}$ we have*

$$\|\mathbf{a}\|_{H^1(\Gamma)} \leq C \|\mathbf{b}\|_{L^2((0,\tau))}^{-1} (\mathcal{D}_0 + \mathcal{D}^{1-\theta} \mathcal{D}_0^\theta).$$

To our knowledge, there are no other results in the literature comparable to those established in the present work.

2. PROOF OF THEOREM 1.2

Let $Q := D \times (0, T)$, $\Upsilon := \Lambda \times (0, T)$, $0 < \gamma < \delta$ and

$$\varphi(x, t) := \phi(x) - \gamma t$$

The proof of Theorem 1.2 is based on the following Carleman inequality.

Proposition 2.1. *There exist $s_* = s_*(g, \phi, \gamma) > 0$ and $C = C(g, \phi, \gamma) > 0$ such that for any $s \geq s_*$ and $u \in H^2(Q)$ satisfying $u = \partial_t u = 0$ on $D \times \{0\}$ we have*

$$\begin{aligned} & C \left(\int_Q e^{2s\varphi} (s|\nabla_g u|^2 + s^3|u|^2) dV_g dt + \int_\Sigma e^{2s\varphi} s^3|u|^2 dS_g dt \right) \\ & \leq \int_Q e^{2s\varphi} |(\partial_t^2 - \Delta_g)u|^2 dV_g dt + \int_\Sigma e^{2s\varphi} s|\nabla_{\tau_g} u|_g^2 dS_g dt \\ & \quad + \int_{\Sigma_0} e^{2s\varphi} (s|\partial_t u|^2 + s|\nabla_g u|_g^2 + s^3|u|^2) dS_g dt \\ & \quad + \int_{D \times \{T\}} e^{2s\varphi} (s|\partial_t u|^2 + s|\nabla_g u|_g^2 + s^3|u|^2) dV_g. \end{aligned}$$

The proof of this proposition is quite technical and will be given in Appendix B. We also need an energy inequality. To this end, set for any $u \in H^2(Q)$

$$\mathcal{E}(u)(t) := \int_{D \times \{t\}} (|\partial_t u|^2 + |\nabla_g u|_g^2 + |u|^2) dV_g.$$

We check that $\mathcal{E}(u) \in W^{1,1}((0, T))$.

Proposition 2.2. *For any $t \in [0, T]$ and $u \in H^2(Q)$ satisfying $(\partial_t^2 - \Delta_g)u = 0$ in Q and $u = \partial_t u = 0$ on $D \times \{0\}$ we have*

$$\mathcal{E}(u)(t) \leq e^t \left(\|u\|_{H^1(\Upsilon)}^2 + \|\partial_{\nu_g} u\|_{L^2(\Upsilon)}^2 \right).$$

Proof. Without loss of generality, we assume that u is real valued. Multiplying $2\partial_t u$ to the equation $(\partial_t^2 - \Delta_g)u = 0$ and integrating it over D yield

$$\begin{aligned} 0 &= \frac{d}{dt} \int_D |\partial_t u|^2 dV_g - \int_D 2\Delta_g u \partial_t u dV_g \\ &= \frac{d}{dt} \int_D (|\partial_t u|^2 + |\nabla_g u|_g^2) dV_g - \int_\Lambda 2\partial_{\nu_g} u \partial_t u dS_g. \end{aligned}$$

On the other hand, we have trivially

$$\frac{d}{dt} \int_D |u|^2 dV_g \leq \int_D (|\partial_t u|^2 + |u|^2) dV_g \leq \mathcal{E}(u)(t).$$

In consequence, we have

$$\mathcal{E}'(t) \leq \mathcal{E}(t) + 2 \int_\Lambda |\partial_{\nu_g} u| |\partial_t u| dV_g.$$

Grönwall's inequality gives

$$\begin{aligned} \mathcal{E}(t) &\leq 2e^t \int_{\Lambda \times (0,t)} |\partial_{\nu_g} u| |\partial_t u| dS_g dt \\ &\leq e^t \left(\|u\|_{H^1(\Upsilon)}^2 + \|\partial_{\nu_g} u\|_{L^2(\Upsilon)}^2 \right). \end{aligned}$$

The proof is then complete. \square

Proof of Theorem 1.2. Let $\tau \in (0, T - T^*)$ and choose $\gamma \in (0, \delta)$ in such a way that $\mathfrak{m} - \gamma(T - \tau) < 0$ and define

$$\mu := -\mathfrak{m} + \gamma(T - \tau) > 0.$$

Let $f = \mathbf{a} \otimes \mathbf{b} \in \mathcal{F}$ and $u = u(f, 0, 0)$. Then we have

$$|\nabla_{\tau_g} u|_g = |\nabla_{\tau_g} \mathbf{a}|_g |\mathbf{b}| \leq (\beta/\alpha) |f| \quad \text{on } \Sigma.$$

Henceforth, $C = C(\zeta) > 0$ and $c = c(\zeta)$ denote a generic constants. By applying Proposition 2.1 to u yields

$$\begin{aligned} &C \int_\Sigma e^{2s\varphi} s^3 |f|^2 dS_g dt \\ &\leq \int_{\Sigma_0} e^{2s\varphi} (s|\partial_t u|^2 + s|\nabla_g u|_g^2 + s^3|u|^2) dS_g dt + \int_\Sigma e^{2s\varphi} s |\nabla_{\tau_g} f|_g^2 dS_g dt \\ &\quad + \int_{D \times \{T\}} e^{2s\varphi} (s|\partial_t u|^2 + s|\nabla_g u|_g^2 + s^3|u|^2) dV_g \\ &\leq s^3 e^{cs} \mathcal{D}_0^2 + (\beta/\alpha)^2 \int_\Sigma e^{2s\varphi} s |f|^2 dS_g dt + s^3 e^{2(\mathfrak{m}-\gamma T)s} \mathcal{E}(u)(T), \quad s \geq s_*, \end{aligned}$$

where s_* is as in Proposition 2.1. Upon modifying s_* , we may and do assume that $Cs^3 - (\beta/\alpha)^2 s > Cs^3/2$. In this case, we have

$$C \int_\Sigma e^{2s\varphi} |f|^2 dS_g dt \leq e^{cs} \mathcal{D}_0^2 + e^{2(\mathfrak{m}-\gamma T)s} \mathcal{E}(u)(T), \quad s \geq s_*.$$

Since

$$\int_\Sigma e^{2s\varphi} |f|^2 dS_g dt \geq \int_{\Gamma \times (0,\tau)} e^{2s\varphi} |f|^2 dS_g dt \geq e^{-2\gamma\tau s} \|f\|_{L^2(\Gamma \times (0,\tau))}^2,$$

we obtain

$$C \|f\|_{L^2(\Gamma \times (0,\tau))}^2 \leq e^{cs} \mathcal{D}_0^2 + e^{-2\mu s} \mathcal{E}(u)(T).$$

By applying Proposition 2.2 to the second term of the right-hand side yields

$$\mathcal{E}(u)(T) \leq e^T (\mathcal{D}_0^2 + \mathcal{D}^2).$$

Whence

$$C\|f\|_{L^2(\Gamma \times (0, \tau))} \leq e^{cs}\mathcal{D}_0 + e^{-\mu s}\mathcal{D}.$$

By replacing C by Ce^{cs^*} , we see that the above inequality holds for all $s > 0$. When $\mathcal{D}_0 \geq \mathcal{D}$, $\|f\|_{L^2(\Gamma \times (0, \tau))} \leq e^{cs}\mathcal{D}_0$ holds. When $\mathcal{D}_0 < \mathcal{D}$, we choose $s > 0$ so that

$$e^{cs}\mathcal{D}_0 = e^{-\mu s}\mathcal{D},$$

that is,

$$s = \frac{1}{c + \mu} \log \frac{\mathcal{D}}{\mathcal{D}_0}.$$

Thus, we get

$$\|f\|_{L^2(\Gamma \times (0, \tau))} \leq C\mathcal{D}^{1-\theta}\mathcal{D}_0^\theta$$

for

$$\theta := \frac{\mu}{c + \mu} \in (0, 1).$$

As

$$\begin{aligned} \|f\|_{L^2(\Gamma \times (0, \tau))} &= \|\mathbf{a}\|_{L^2(\Gamma)} \|\mathbf{b}\|_{L^2((0, \tau))}, \\ \|\nabla_{\tau_g} f\|_{L^2(\Gamma \times (0, \tau))} &\leq (\beta/\alpha) \|\mathbf{a}\|_{L^2(\Gamma)} \|\mathbf{b}\|_{L^2((0, \tau))}, \end{aligned}$$

the expected inequality follows. \square

APPENDIX A. ANALYSIS OF THE IBVP

Let $A : L^2(U) \rightarrow L^2(U)$ be the unbounded operator given by

$$Au = \Delta_g u \quad \text{and} \quad D(A) = H_0^1(U) \cap H^2(U).$$

Let $u \in D(A)$. By Green formula, we have

$$\int_U \Delta_g u \bar{u} dV_g = - \int_U |\nabla u|^2 dV_g \leq 0,$$

that is, A is dissipative. Let $\lambda > 0$ and $u_0 \in L^2(U)$. Since

$$B(u, v) = \lambda \int_U u \bar{v} dV_g + \int_U \langle \nabla_g u, \nabla_g \bar{v} \rangle dV_g, \quad u, v \in H_0^1(U)$$

is an equivalent scalar product on $H_0^1(U)$, Riesz representation theorem shows that there exists a unique $u \in H_0^1(U)$ such that

$$\lambda \int_U u \bar{v} dV_g + \int_U \langle \nabla_g u, \nabla_g \bar{v} \rangle dV_g = \int_U u_0 \bar{v} dV_g, \quad v \in H_0^1(U).$$

Hence, $(\lambda - \Delta_g)u = u_0$ in the distributional sense and then $u \in D(A)$ according to the elliptic regularity (e.g [2, Theorem 9.25] and Remark 24 just after [2, Theorem 9.26]).

We endow the Hilbert space $\mathcal{H} = H_0^1(U) \times L^2(U)$ with the scalar product

$$\langle (v_1, w_1) | (v_2, w_2) \rangle = \int_U [\langle \nabla_g v_1, \nabla_g \bar{v}_2 \rangle + w_1 \bar{w}_2] dV_g, \quad (v_j, w_j) \in \mathcal{H}, \quad j = 1, 2.$$

Define the unbounded operator $\mathcal{A} : \mathcal{H} \rightarrow \mathcal{H}$ as follows

$$\mathcal{A}(u, v) = (v, Au) \quad \text{and} \quad D(\mathcal{A}) = \mathcal{H}_0 := (H_0^1(U) \cap H^2(U)) \times H_0^1(U).$$

Let $(u, v) \in D(\mathcal{A})$. As

$$\begin{aligned} \langle \mathcal{A}(u, v) | (u, v) \rangle &= \int_U \langle \nabla_g \bar{u}, \nabla_g v \rangle dV_g + \int_U \Delta_g u \bar{v} dV_g \\ &= \int_U \langle \nabla_g \bar{u}, \nabla_g v \rangle dV_g - \int_U \langle \nabla_g u, \nabla_g \bar{v} \rangle dV_g, \end{aligned}$$

we obtain

$$\Re \langle \mathcal{A}(u, v) | (u, v) \rangle = 0,$$

that is, \mathcal{A} is dissipative.

Let $\lambda > 0$ and $(u_0, u_1) \in \mathcal{H}$ and consider the equation

$$\lambda(u, v) - \mathcal{A}(u, v) = (u_0, u_1).$$

This equation is equivalent to the following system

$$(A.1) \quad (\lambda^2 - A)u = u_1 + \lambda u_0, \quad v = \lambda u - u_0.$$

We have seen above that A is m -dissipative. Hence, (A.1) admits unique solution $(u, v) \in \mathcal{D}(\mathcal{A})$. In other words, we proved that \mathcal{A} is m -dissipative and therefore, according to Lumer-Phillips theorem, \mathcal{A} generates a contraction semigroup (e.g. [7, Theorem 3.8.4]).

Let \mathcal{X}_0 be the subspace of \mathcal{X} given by

$$\mathcal{X}_0 = C^2([0, T]; L^2(U)) \cap C^1([0, T]; H_0^1(U)) \cap C([0, T]; H_0^1(U) \cap H^2(U)).$$

As \mathcal{A} is the generator of a contraction semigroup, for each $(u_0, u_1) \in \mathcal{H}_0$ and $F \in W^{1,1}((0, T); L^2(U))$ the IBVP

$$(A.2) \quad \begin{cases} (\partial_t^2 - \Delta_g)v = F & \text{in } U \times (0, T), \\ v|_{\Sigma} = 0, \\ (v(\cdot, 0), \partial_t v(\cdot, 0)) = (u_0, u_1) \end{cases}$$

has a unique solution $v = v(u_0, u_1, F) \in \mathcal{X}_0$. Furthermore, we have

$$(A.3) \quad \|v\|_{\mathcal{X}} \leq \mathbf{c} (\|(u_0, u_1)\|_{\mathcal{H}_0} + \|F\|_{W^{1,1}((0, T); L^2(U))}), \quad t \geq 0,$$

where $\mathbf{c} = \mathbf{c}(g, \phi) > 0$ is a constant.

Next, consider the IBVP (1.1). Let $(f, u_0, u_1) \in \mathcal{Z}$ and $H \in \overbrace{(f, u_0, u_1)}$. If $F := -(\partial_t^2 - \Delta_g)H \in W^{1,1}((0, T); L^2(U))$ then we check that $u := u(f, u_0, u_1) := H + v(0, 0, F) \in \mathcal{X}$ is the unique solution of (1.1). Furthermore, as $H \in \overbrace{(f, u_0, u_1)}$ was taken arbitrary, (A.3) implies (1.2).

Let us remark that, according to perturbation theory of strongly continuous semigroups (e.g., [6]), the result for the IBVP (1.1) still holds when Δ_g is replaced by $\Delta_g + pu + q\partial_t u$, where $p, q \in L^\infty(U)$.

APPENDIX B. PROOF OF PROPOSITION 2.1

Without loss of generality, we limit our proof to real-valued function. Also, for simplicity, we set $\square := (\partial_t^2 - \Delta_g)$ and $w' := \partial_t w$. Let $u \in H^2(Q)$, $z := e^{s\varphi}u$, and

$$P_s z := e^{s\varphi} \square (e^{-s\varphi} z),$$

A direct calculation yields $P_s z = P_s^+ z + P_s^- z$, where

$$\begin{cases} P_s^+ z := \square z + s^2(|\varphi'|^2 - |\nabla_g \varphi|_g^2)z, \\ P_s^- z := -2s\varphi'z' + 2s\langle \nabla_g \varphi, \nabla_g z \rangle - s\square\varphi z. \end{cases}$$

Hereinafter, the integrals on Q are with respect to the measure $dV_g dt$ and those on Υ are with respect to the surface measure $dS_g dt$.

Set

$$(P_s^+ z, P_s^- z)_g =: \sum_{k=1}^9 I_k,$$

where

$$\begin{aligned} I_1 &:= -2 \int_Q s\varphi'z'z'', \\ I_2 &:= 2 \int_Q s\langle \nabla_g \varphi, \nabla_g z \rangle z'', \\ I_3 &:= - \int_Q s\square\varphi z z'', \\ I_4 &:= -2 \int_Q s\varphi'z'\Delta_g z, \\ I_5 &:= -2 \int_Q s\Delta_g z \langle \nabla_g \varphi, \nabla_g z \rangle, \\ I_6 &:= \int_Q s\Delta_g z \square\varphi z, \\ I_7 &:= -2 \int_Q s^3\varphi'(|\varphi'|^2 - |\nabla_g \varphi|_g^2)z z', \\ I_8 &:= 2 \int_Q s^3(|\varphi'|^2 - |\nabla_g \varphi|_g^2)\langle \nabla_g \varphi, \nabla_g z \rangle z, \\ I_9 &:= - \int_Q s^3\square\varphi(|\varphi'|^2 - |\nabla_g \varphi|_g^2)|z|^2. \end{aligned}$$

Integrations by parts yield

$$\begin{aligned} I_1 &= - \int_Q s\varphi'\partial_t(|z'|^2) = \int_Q s\varphi''|z'|^2 - \int_D [s\varphi'|z'|^2]_0^T, \\ I_2 &= - \int_Q s\langle \nabla_g \varphi, \nabla_g |z'|^2 \rangle + 2 \int_D [sz'\langle \nabla_g \varphi, \nabla_g z \rangle]_0^T \\ &= \int_Q s\Delta_g \varphi |z'|^2 - \int_\Upsilon s\partial_{\nu_g} \varphi |z'|^2 + 2 \int_D [sz'\langle \nabla_g \varphi, \nabla_g z \rangle]_0^T, \\ I_3 &= \int_Q s\square\varphi |z'|^2 + \int_Q s\partial_t \square\varphi z \partial_t z - \int_D [s\square\varphi z z']_0^T \\ &= \int_Q s\square\varphi |z'|^2 - (1/2) \int_Q s\partial_t^2 \square\varphi |z|^2 - \int_D [s\square\varphi z z']_0^T \\ &\quad + (1/2) \int_D [s\partial_t \square\varphi |z|^2]_0^T, \end{aligned}$$

$$\begin{aligned}
I_4 &= - \int_Q s\varphi' \partial_t (|\nabla_g z|_g^2) + 2 \int_{\Upsilon} s\varphi' \partial_{\nu_g} z z' \\
&= \int_Q s\varphi'' |\nabla_g z|_g^2 + 2 \int_{\Upsilon} s\varphi' \partial_{\nu_g} z z' - \int_D [s\varphi' |\nabla_g z|_g^2]_0^T,
\end{aligned}$$

$$\begin{aligned}
I_5 &= 2 \int_Q s \langle \nabla_g z, \nabla_g \langle \nabla_g \varphi, \nabla_g z \rangle \rangle - 2 \int_{\Upsilon} s \partial_{\nu_g} z \langle \nabla_g \varphi, \nabla_g z \rangle \\
&= 2 \int_Q s \nabla_g^2 \varphi (\nabla_g z, \nabla_g z) + \int_Q s \langle \nabla_g \varphi, \nabla_g (|\nabla_g z|_g^2) \rangle \\
&\quad - 2 \int_{\Upsilon} s \partial_{\nu_g} z \langle \nabla_g \varphi, \nabla_g z \rangle \\
&= 2 \int_Q s \nabla_g^2 \varphi (\nabla_g z, \nabla_g z) - \int_Q s \Delta_g \varphi |\nabla_g z|_g^2 \\
&\quad - 2 \int_{\Upsilon} s \partial_{\nu_g} z \langle \nabla_g \varphi, \nabla_g z \rangle + \int_{\Upsilon} s \partial_{\nu_g} \varphi |\nabla_g z|_g^2,
\end{aligned}$$

$$I_6 = - \int_Q s \square \varphi |\nabla_g z|_g^2 - \int_Q s \langle \nabla_g \square \varphi, \nabla_g z \rangle z + \int_{\Upsilon} s \square \varphi \partial_{\nu_g} z z,$$

$$\begin{aligned}
I_7 &= - \int_Q s^3 \varphi' (|\varphi'|^2 - |\nabla_g \varphi|_g^2) \partial_t (|z|^2) \\
&= \int_Q s^3 \partial_t (\varphi' (|\varphi'|^2 - |\nabla_g \varphi|_g^2)) |z|^2 - \int_D [s^3 \varphi' (|\varphi'|^2 - |\nabla_g \varphi|_g^2) |z|^2]_0^T,
\end{aligned}$$

$$\begin{aligned}
I_8 &= \int_Q s^3 (|\varphi'|^2 - |\nabla_g \varphi|_g^2) \langle \nabla_g \varphi, \nabla_g (|z|^2) \rangle \\
&= - \int_Q s^3 \operatorname{div}((|\varphi'|^2 - |\nabla_g \varphi|_g^2) \nabla_g \varphi) |z|^2 + \int_{\Upsilon} s^3 (|\varphi'|^2 - |\nabla_g \varphi|_g^2) \partial_{\nu_g} \varphi |z|^2.
\end{aligned}$$

Adding all of these together yields

$$\begin{aligned}
&(P_s^+ z, P_s^- z)_g + \mathcal{B} = \\
&\int_Q 2s [\varphi'' |z'|^2 + \nabla_g^2 \varphi (\nabla_g z, \nabla_g z)] - \int_Q s [\langle \nabla_g \square \varphi, \nabla_g z \rangle + (1/2) \partial_t^2 \square \varphi z] z \\
&+ \int_Q [s^3 \varphi' \partial_t (|\varphi'|^2 - |\nabla_g \varphi|_g^2) - s^3 \langle \nabla_g (|\varphi'|^2 - |\nabla_g \varphi|_g^2), \nabla_g \varphi \rangle] |z|^2
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{B} := & \int_{\Upsilon} [s^3(|\nabla_g \varphi|_g^2 - |\varphi'|^2) \partial_{\nu_g} \varphi] |z|^2 \\
& + \int_{\Upsilon} [-2s\varphi' z' + 2s\langle \nabla_g \varphi, \nabla_g z \rangle - s\Box \varphi z] \partial_{\nu_g} z \\
& + \int_{\Upsilon} [s\partial_{\nu_g} \varphi |z'|^2 - s\partial_{\nu_g} \varphi |\nabla_g z|_g^2] \\
& + \int_{D \times \{T\}} [(-s/2)\partial_t \Box \varphi - s^3\varphi'(|\nabla_g \varphi|_g^2 - |\varphi'|^2)] |z|^2 \\
& + \int_{D \times \{T\}} [-2s\langle \nabla_g \varphi, \nabla_g z \rangle + s\Box \varphi z] z' \\
& + \int_{D \times \{T\}} s\varphi' [|z'|^2 + |\nabla_g z|_g^2].
\end{aligned}$$

Henceforth, $C = C(g, \phi, \gamma) > 0$ and $s_* = s_*(g, \phi, \gamma) > 0$ denote generic constants. Then we have

$$\varphi'' |z'|^2 + \nabla_g^2 \varphi (\nabla_g z, \nabla_g z) = \nabla_g^2 \phi (\nabla_g z, \nabla_g z) \geq C |\nabla_g z|_g^2,$$

and

$$\begin{aligned}
& \varphi' \partial_t (|\varphi'|^2 - |\nabla_g \varphi|_g^2) - \langle \nabla_g (|\varphi'|^2 - |\nabla_g \varphi|_g^2), \nabla_g \varphi \rangle \\
& = 2\nabla_g^2 \phi (\nabla_g \phi, \nabla_g \phi) \geq C |\nabla_g \phi|_g^2
\end{aligned}$$

In consequence, we get

$$C \int_Q (s |\nabla_g z|_g^2 + s^3 |z|^2) \leq (P_s^+ z, P_s^- z)_g + \int_Q s [\langle \nabla_g \Box \varphi, \nabla_g z \rangle + (1/2) \partial_t^2 \Box \varphi z] z + \mathcal{B}.$$

Set

$$\begin{aligned}
\mathcal{B}_{\Sigma_0} := & \int_{\Sigma_0} [s^3(|\nabla_g \varphi|^2 - |\varphi'|^2) \partial_{\nu_g} \varphi] |z|^2 \\
& + \int_{\Sigma_0} [-2s\varphi' z' + 2s\langle \nabla_g \varphi, \nabla_g z \rangle - s\Box \varphi z] \partial_{\nu_g} z \\
& + \int_{\Sigma_0} [s\partial_{\nu_g} \varphi |z'|^2 - s\partial_{\nu_g} \varphi |\nabla_g z|_g^2], \\
\mathcal{B}_{\Sigma} := & \int_{\Sigma} [s^3(|\nabla_g \varphi|^2 - |\varphi'|^2) \partial_{\nu_g} \varphi] |z|^2 \\
& + \int_{\Sigma} [-2s\varphi' z' + 2s\langle \nabla_g \varphi, \nabla_g z \rangle - s\Box \varphi z] \partial_{\nu_g} z \\
& + \int_{\Sigma} [s\partial_{\nu_g} \varphi |z'|^2 - s\partial_{\nu_g} \varphi |\nabla_g z|_g^2],
\end{aligned}$$

and

$$\begin{aligned} \mathcal{B}_T &:= \int_{D \times \{T\}} [(-s/2)\partial_t \square \varphi - s^3 \varphi' (|\nabla_g \varphi|^2 - |\varphi'|^2)] |z|^2 \\ &\quad + \int_{D \times \{T\}} [-2s \langle \nabla_g \varphi, \nabla_g z \rangle + s \square \varphi z] z' \\ &\quad + \int_{D \times \{T\}} s \varphi' [|z'|^2 + |\nabla_g z|^2] \end{aligned}$$

so that

$$\mathcal{B} = \mathcal{B}_{\Sigma_0} + \mathcal{B}_\Sigma + \mathcal{B}_T.$$

We check that

$$\mathcal{B}_{\Sigma_0} \leq C \int_{\Sigma_0} [s|z'|^2 + s|\nabla_g z|_g^2 + s^3|z|^2],$$

and

$$\mathcal{B}_T \leq C \int_{D \times \{T\}} [s|z'|^2 + s|\nabla_g z|_g^2 + s^3|z|^2].$$

On the other hand, we have

$$|\nabla_g z|_g^2 = |\nabla_{\tau_g} z|_g^2 + |\partial_{\nu_g} z|^2, \quad \langle \nabla_g \varphi, \nabla_g z \rangle = -|\nabla_g \phi|_g \partial_{\nu_g} z \quad \text{on } \Sigma$$

and $\nu_g = -\frac{\nabla_g \phi}{|\nabla_g \phi|_g}$ on Σ . Whence

$$\begin{aligned} \mathcal{B}_\Sigma &= \int_\Sigma [-s^3 (|\nabla_g \phi|_g^2 - \gamma^2) |\nabla_g \phi|_g] |z|^2 \\ &\quad + \int_\Sigma [2\gamma s z' - 2s |\nabla_g \phi|_g \partial_{\nu_g} z - s \square \varphi z] \partial_{\nu_g} z \\ &\quad + \int_\Sigma [-s |\nabla_g \phi|_g |z'|^2 + s |\nabla_g \phi|_g (|\nabla_{\tau_g} z|_g^2 + |\partial_{\nu_g} z|^2)] \\ &= \int_\Sigma [-s |\nabla_g \phi|_g |z'|^2 + 2\gamma s \partial_{\nu_g} z z' - s |\nabla_g \phi|_g |\partial_{\nu_g} z|^2 + s |\nabla_g \phi|_g |\nabla_{\tau_g} z|^2 \\ &\quad - s \square \varphi \partial_{\nu_g} z z - s^3 (|\nabla_g \phi|_g^2 - \gamma^2) |\nabla_g \phi|_g |z|^2] \\ &\leq \int_\Sigma s [-\delta |z'|^2 + (2\gamma z' - \square \varphi z) \partial_{\nu_g} z - \delta |\partial_{\nu_g} z|^2 + |\nabla_g \phi|_g |\nabla_{\tau_g} z|^2 \\ &\quad - s^2 (\delta^2 - \gamma^2) \delta |z|^2]. \end{aligned}$$

By $0 < \gamma < \delta$, we note that

$$\begin{aligned} &-\delta |z'|^2 + (2\gamma z' - \square \varphi z) \partial_{\nu_g} z - \delta |\partial_{\nu_g} z|^2 \\ &= -\delta \left| z' - \frac{\gamma}{\delta} \partial_{\nu_g} z \right|^2 - \frac{\delta^2 - \gamma^2}{\delta} |\partial_{\nu_g} z|^2 - \square \varphi z \partial_{\nu_g} z \\ &\leq -\frac{\delta^2 - \gamma^2}{\delta} \left| \partial_{\nu_g} z + \frac{\delta \square \varphi z}{2(\delta^2 - \gamma^2)} \right|^2 + C|z|^2 \\ &\leq C|z|^2. \end{aligned}$$

Therefore, we obtain

$$\mathcal{B}_\Sigma \leq \int_\Sigma s [C |\nabla_{\tau_g} z|^2 + C |z|^2 - s^2 (\delta^2 - \gamma^2) \delta |z|^2]$$

and then

$$\mathcal{B}_\Sigma + (1/2) \int_\Sigma (\delta^2 - \gamma^2) \delta s^3 |z|^2 \leq C \int_\Sigma s |\nabla_{\tau_g} z|^2, \quad s \geq s_*.$$

Combining the estimates above, we get

$$\begin{aligned} & C \left(\int_Q (s |\nabla_g z|_g^2 + s^3 |z|^2) + \int_\Sigma s^3 |z|^2 \right) \\ & \leq (P_s^+ z, P_s^- z)_g + \int_Q s [\langle \nabla_g \square \varphi, \nabla_g z \rangle + (1/2) \partial_t^2 \square \varphi z] z \\ & \quad + \mathcal{B}_{\Sigma_0} + \mathcal{B}_T + \int_\Sigma s |\nabla_{\tau_g} z|_g^2 \\ & \leq \|P_s z\|_g^2 + \int_Q (|\nabla_g z|_g^2 + s^2 |z|^2) + \int_{\Sigma_0} (s |z'|^2 + s |\nabla_g z|_g^2 + s^3 |z|^2) \\ & \quad + \int_\Sigma s |\nabla_{\tau_g} z|_g^2 + \int_{D \times \{T\}} (s |z'|^2 + s |\nabla_g z|_g^2 + s^3 |z|^2), \quad s \geq s_*. \end{aligned}$$

As the second term in the right-hand side can be absorbed by the left-hand side, we find

$$\begin{aligned} & C \left(\int_Q (s |\nabla_g z|_g^2 + s^3 |z|^2) + \int_\Sigma s^3 |z|^2 \right) \\ & \leq \|P_s z\|_g^2 + \int_{\Sigma_0} (s |z'|^2 + s |\nabla_g z|_g^2 + s^3 |z|^2) \\ & \quad + \int_\Sigma s |\nabla_{\tau_g} z|_g^2 + \int_{D \times \{T\}} (s |z'|^2 + s |\nabla_g z|_g^2 + s^3 |z|^2), \quad s \geq s_*. \end{aligned}$$

Since $u = e^{-s\varphi} z$ and $\nabla_{\tau_g} u = \nabla_{\tau_g} z$ holds by $\nabla_{\tau_g} \phi = 0$, we end up getting

$$\begin{aligned} & C \left(\int_Q e^{2s\varphi} (s |\nabla_g u|_g^2 + s^3 |u|^2) + \int_\Sigma e^{2s\varphi} s^3 |u|^2 \right) \\ & \leq \int_Q e^{2s\varphi} |\square u|^2 + \int_{\Sigma_0} e^{2s\varphi} (s |u'|^2 + s |\nabla_g u|_g^2 + s^3 |u|^2) \\ & \quad + \int_\Sigma e^{2s\varphi} s |\nabla_{\tau_g} u|_g^2 + \int_{D \times \{T\}} e^{2s\varphi} (s |u'|^2 + s |\nabla_g u|_g^2 + s^3 |u|^2), \quad s \geq s_*. \end{aligned}$$

The proof is then complete.

ACKNOWLEDGEMENT

This work was supported by JSPS KAKENHI Grant Number JP23KK0049.

DECLARATIONS

Conflict of interest. The authors declare that they have no conflict of interest.

Data availability. Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

REFERENCES

- [1] L. Bourgeois, D. Ponomarev, and J. Dardé. An inverse obstacle problem for the wave equation in a finite time domain. *Inverse Problems and Imaging*, 19:377–400, 2019.
- [2] H. Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer, 2011.
- [3] M. Choulli. Stable determination of the initial data in an ibvp for the wave equation outside a non-trapping obstacle. *Communications on Applied Mathematics and Computation*, 2024.
- [4] M. Choulli and H. Takase. Lipschitz stability for an elliptic inverse problem with two measurements. *arXiv:2404.13901*, 4 2024.
- [5] V. Isakov. Inverse obstacle problems. *Inverse Problems*, 25:18pp, 2009.
- [6] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer-Verlag, New York, 1983.
- [7] M. Tucsnak and G. Weiss. *Observation and Control for Operator Semigroups*. Birkhäuser Verlag, 2009.

UNIVERSITÉ DE LORRAINE, 34 COURS LÉOPOLD, 54052 NANCY CEDEX, FRANCE

Email address: `mourad.choulli@univ-lorraine.fr`

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU, FUKUOKA 819-0395, JAPAN

Email address: `htakase@imi.kyushu-u.ac.jp`