

# PROOF OF ZAMOLODCHIKOV CONJECTURE FOR SEMI-CLASSICAL CONFORMAL BLOCKS ON TORUS

HARINI DESIRAJU, PROMIT GHOSAL, AND ANDREI PROKHOROV

ABSTRACT. In 1986, Zamolodchikov conjectured an exponential structure for the semi-classical limit of conformal blocks on a sphere. This paper provides a rigorous proof of the analog of Zamolodchikov conjecture for Liouville conformal blocks on a one-punctured torus, using their probabilistic construction and show the existence of a positive radius of convergence of the semi-classical limit. As a consequence, we obtain a closed form expression for the solution of the Lamé equation, and show a relation between its accessory parameter and the classical action of the non-autonomous elliptic Calogero-Moser model evaluated at specific values of the solution.

## CONTENTS

1. Introduction	1
2. Probabilistic conformal blocks	6
3. Derivative gaussian multiplicative chaos measure	9
4. Semi-classical limits for deformed conformal blocks	23
5. Semi-classical limit of undeformed conformal blocks	34
6. Asymptotics of semi-classical conformal block as $\tau \rightarrow i\infty$	37
Appendix A. Useful notation and elliptic functions	41
Appendix B. Non-autonomous elliptic Calogero-Moser model	42
References	47

## 1. INTRODUCTION

Conformal field theory (CFT), a subclass of quantum field theory provides a framework for constructing random functions on Riemannian manifolds, which transform covariantly under conformal (i.e., angle-preserving) mappings. Since the groundbreaking work by Belavin, Polyakov, and Zamolodchikov (BPZ) in 1984 [BPZ84], two-dimensional (2D) CFT has evolved into a major area of theoretical physics. It has significant applications in 2D statistical physics, string theory and has profound implications for mathematics. See [Rib14] for a review and references therein.

An important subclass of conformal field theories (CFTs) is known as Liouville CFTs (LCFTs). These CFTs are characterized by a parameter called the central charge, denoted as  $c$ , which for LCFTs lies in the range  $c \in [25, \infty)$ . LCFT originated from Polyakov's work on two-dimensional quantum gravity and bosonic string theory in 1981 [Pol81]. It was rigorously constructed using the path integral formalism of quantum field theory, initially on the sphere in [Dav+16] and subsequently on other Riemann surfaces in [DRV16; HRV18; GRV19]. We refer the reader to [RV] for a review.

The paper by Belavin, Polyakov, and Zamolodchikov in 1984 introduced a schematic approach known as the conformal bootstrap. This program aims to exactly solve the correlation functions of a given two-dimensional CFT by expressing them in terms of its three-point sphere correlation functions and specific power series called conformal blocks. These conformal blocks are entirely determined by the Virasoro algebra, which encodes the infinitesimal local conformal symmetries. The conformal blocks depend on the particular CFT solely through the central charge.

Recently a construction of Liouville conformal blocks using rigorous path integral formulation based on probabilistic ideas was introduced by [Gho+24]. The construction is via the Gaussian multiplicative chaos (GMC) measure, a random measure defined by exponentiating the Gaussian free field (see e.g. [DRV16; Ber17; Aru20]). This paper explores the semi-classical limit of the

probabilistic representation of Liouville conformal blocks, focusing on the case of torus with a single marked point. For a given in the upper half-plane, we consider the flat torus  $\mathbb{T}_\tau$  characterized by the modular parameter  $\tau$ , and the nome  $q := e^{i\pi\tau}$ . The 1-point toric correlation function of LCFT, rigorously constructed in [DRV16], has the form  $\langle e^{\alpha\Phi(0)} \rangle_\tau$ , where  $\langle \dots \rangle_\tau$  is the average over the random field  $\phi$  for LCFT on  $\mathbb{T}_\tau$  and  $\alpha$  is called the vertex insertion weight. Conformal bootstrap formula for torus which is recently shown in [Gui+21, Theorem 9] expresses  $\langle e^{\alpha\Phi(0)} \rangle_\tau$  via one point toric conformal block  $\mathcal{G}_{\gamma,P}^\alpha(q)$ ,

$$\langle e^{\alpha\Phi(0)} \rangle_\tau = \int_{\mathbb{R}} C_\gamma(\alpha, Q - iP, Q + iP) |\mathcal{G}_{\gamma,P}^\alpha(q)|^2 dP,$$

where the parameter  $\gamma$  dictates the central charge through the relation

$$c = 1 + 6Q^2 \in (25, \infty), \quad Q := \frac{\gamma}{2} + \frac{2}{\gamma} \in (2, \infty).$$

Here  $C_\gamma(\alpha_1, \alpha_2, \alpha_3)$  is the Liouville 3-point correlation function. The exact expression of  $C_\gamma$  as first proposed in [DO94; ZZ96] and later proved in [KRV20]. It is worthwhile to note that the conformal bootstrap for sphere is recently proven in [KRV20] and [Gui+21] showed it for other geometry including torus.

In a recent work, [Gho+24, Theorem 1.1] introduced a probabilistic description of  $\mathcal{G}_{\gamma,P}^\alpha(q)$  using GMC measure. This probabilistic definition of  $\mathcal{G}_{\gamma,P}^\alpha(q)$  is shown to match with the q-power series formula of the same (defined via Virasoro algebra in [BPZ84]) through rigorously proving the Zamolodchikov recursion (as proposed in [Zam84; Zam87a; HJS10]) for the GMC expression of  $\mathcal{G}_{\gamma,P}^\alpha(q)$ . In fact, [Gho+24] showed that  $\mathcal{G}_{\gamma,P}^\alpha(q)$  is defined for  $\alpha \in (-\frac{4}{\gamma}, Q)$ ,  $P \in \mathbb{R}$  and is analytic as a function of  $q \in \mathbb{C}$  in a ball  $B_r(0)$  of radius  $r$  greater than  $\frac{1}{2}$  when  $\alpha \in [0, Q)$ . We refer the reader to [Gho+24, Appendix A] and [Gho+, Section 1.1] for a summary of the algebraic approach to conformal blocks on one-point torus.

Our objective is to understand the behavior and properties of these Liouville conformal blocks  $\mathcal{G}_{\gamma,P}^\alpha(q)$  in the semi-classical limit  $\gamma \rightarrow 0$ , shedding light on the underlying probabilistic structures associated with conformal field theories on tori. Such a proof for all other geometries remains an open question. Starting from the probabilistic definition of conformal blocks on the four point sphere, one can easily adapt our methods to study their semi-classical limits.

**Theorem 1.1.** *The semi-classical limit of the conformal block  $\mathcal{G}_{\gamma,P}^\alpha(q)$  exists and has the form*

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma,P}^\alpha(q) = \phi(\alpha_0, P_0, q) + \tilde{f}(\alpha_0, P_0); \quad \alpha = \frac{\alpha_0}{\gamma}, \quad P = \frac{P_0}{\gamma}, \quad (1.1)$$

for  $q = e^{i\pi\tau} \in B_r(0)$  for some  $r > 0$  small,  $\alpha_0 \in (-4, 2)$  and  $P_0 \in \mathbb{R}$ . Here,  $\phi(\cdot)$  is an analytic function in  $q$  and  $\tilde{f}(\cdot)$  is a normalization constant such that

$$\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma, P_0/\gamma}^{\alpha_0/\gamma}(q) = 0.$$

Moreover,  $\phi(\cdot)$  determines the accessory parameter of the Lamé equation that can be written explicitly as

$$\left( \partial_z^2 - \frac{\alpha_0}{4} \left( \frac{\alpha_0}{4} - 1 \right) \wp(z) + \frac{\pi i}{2} \partial_\tau \phi(\alpha_0, P_0, q) \right) \tilde{\Gamma}(z; \alpha_0, P_0, q) = 0, \quad (1.2)$$

where  $\wp(\cdot)$  is the Weierstrass  $\wp$ -function (see (A.6)) and the function  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is given by

$$\tilde{\Gamma}(z; \alpha_0, P_0, q) := e^{\frac{P_0 z \pi}{2}} \exp \left( - \frac{\alpha_0 \int_0^1 \log \left( \frac{\theta_1(z+x)}{\theta_1(z)} \right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{4 \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} \right). \quad (1.3)$$

**Remark 1.1.** *The limit in (1.1) can be shown for  $q \in B_r(0)$  for some  $r > \frac{1}{2}$  when  $\alpha_0 \in [0, 2)$ . Furthermore, since  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  has an explicit expression in (1.3), one can use Lamé equation to write explicit form of the accessory parameter  $\pi i \partial_\tau \phi(\alpha_0, P_0, q)$  (see Corollary 4.40). The analytic properties with respect to  $\alpha_0, P_0, q$  of the semi-classical limit are accessible through this explicit form.*

**1.1. Zamolodchikov conjecture.** The main result of this paper rigorously proves the Zamolodchikov conjecture for the case of conformal blocks on a one-point torus. A long standing conjecture formulated by Zamolodchikov in 1986 [Zam86, (2.30)-(2.32)] postulates that, in the semi-classical limit ( $c \rightarrow \infty$ ), conformal blocks on a four-point sphere have an exponential structure. Furthermore, it is known that, assuming such an exponential structure, the semi-classical limit of the sphere 4-point conformal block is described by the accessory parameter of the Heun's equation.

The evidence for such a structure for the case of the four point sphere comes from the analysis of the power series representation of conformal blocks [Zam87b], from the oscillator formalism of Virasoro algebra [BDK20], and from the combinatorial expression in terms of the Nekrasov-Okounkov functions describing the partition functions of supersymmetric gauge theories given by the Alday-Gaiotto-Tachikawa (AGT) correspondence [NPS18; NS10].

The semi-classical limit of conformal blocks carries significant importance in mathematical physics as it plays a crucial role in a variety of areas from black hole physics [AGH22; CC22] and supersymmetric gauge theories [NS10] (see also the introduction in [Pia+22]) to Teichmüller theory (see [BCP17; Tak94] and references therein).

Of particular interest to this paper is the connection between semi-classical conformal blocks and the Heun equation, a second-order differential equation featuring a free parameter known as the *accessory parameter*. This parameter is crucial in the semi-classical analysis as it is characterized by conformal blocks on the sphere [Zam86]. The study of the Heun equation is interesting on its own. See [RA95] for a bibliographical account and [Hor18] for a review on applications to physics.

In the recent years, interest has grown in understanding the relationships between conformal blocks, accessory parameters, and Painlevé equations [BGG22; LN21; PP19; LN18]. Specifically, the semi-classical limit of conformal blocks is related to the Painlevé VI equation through its Hamiltonian at special values of the solution [Lit+14; KT13]. In turn the Hamiltonian of the Painlevé VI equation is given by  $c = 1$  conformal blocks on sphere, and hence the aforementioned relation provides a non-trivial link between the  $c \rightarrow \infty$  and  $c = 1$  conformal blocks.

More recently, a relation between  $c \rightarrow \infty$  conformal blocks and the connection problem for the degenerations of the Heun equation was conjectured in [Bon+23] and is dubbed 'the Trieste formula' [LN22], with the case of the Heun equation (without degenerations) verified in [LN22]. So, the proof of Zamolodchikov conjecture paves the way for a rigorous treatment of the aforementioned problems. Indeed most of above claims could be easily adapted to the one-point torus case. In particular, the role of the Heun equation is now played by the Lamé equation and furthermore, its accessory parameter, under the assumption that the Zamolodchikov conjecture holds, is known to be related to the semi-classical conformal block on the torus [Pia14] and to the elliptic form of the Painlevé VI equation with special value of the parameters [BGG22]. In this context, we also want to highlight a similar study of the semiclassical limit of Liouville CFT on a sphere [LRV], where they prove the Takhtajan-Zograf theorem relating the classical Liouville action at special points and accessory parameters of a second order Fuchsian equation of holomorphic functions on the unit disk [TZ03].

In Theorem 1.1, we showed that, as the central charge  $c$  goes to  $\infty$  or equivalently  $\gamma \rightarrow 0$ , the torus one point Liouville conformal block behaves as  $\mathcal{G}_{\gamma,P}^\alpha(q) \sim \exp(\phi(\alpha_0, P_0, q)/\gamma^2)$  where  $\phi(\alpha_0, P_0, q)$  is the semi-classical conformal block. Moreover, (1.2) showed that  $\phi(\alpha_0, P_0, q)$  is related to the accessory parameter of Lamé equation. As a result, we provide a rigorous proof of the analogue of Zamolodchikov conjecture for one-point torus conformal blocks. Consequently, we prove that  $\phi(\dots)$  is also related to the Hamiltonian of the non-autonomous elliptic Calogero-Moser (NAECM) model (see Theorem 4.2) for special values of the solution [BGG22], which in turn is related to  $c = 1$  conformal blocks on a one-point torus [DDG23]. Such a proof for all other geometries remains an open question.

**1.2. Ideas of the proofs.** Our main goal in this paper is to prove the existence and estimate the radius of convergence for, the semi-classical limit ( $\gamma \rightarrow 0$ ) of the conformal block  $\mathcal{G}_{\gamma,P}^\alpha(q)$ . Henceforth, we refer to this as the *undeformed* conformal block. The proofs of these statements involve several steps. One of the main challenges is to show that the conformal block has exponential structure in the semi-classical limit, i.e., as  $\gamma \rightarrow 0$ , the torus 1-point conformal block  $\mathcal{G}_{\gamma,P}^\alpha$  behaves

as  $\exp\left(-\frac{1}{\gamma^2}(\phi(\alpha_0, P_0, q) + \tilde{f}(\alpha_0, P_0))\right)$ . For showing this, we use the probabilistic description of  $\mathcal{G}_{\gamma, P}^\alpha$  given as

$$\mathcal{G}_{\gamma, P}^\alpha \sim \mathbb{E} \left[ \left( \text{IGMC}_{\gamma, P}^\alpha(q) \right)^{-\frac{\alpha}{\gamma}} \right], \quad \text{IGMC}_{\gamma, P}^\alpha(q) := \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma P x} e^{\gamma:Y(x; q):} dx \quad (1.4)$$

where  $:Y(x; q):$  is a renormalized log-correlated field on  $[0, 1]$ . See Section 2.1 for its definition. The above representation of the conformal block was introduced by [Gho+24] and expressed in terms of the Gaussian multiplicative chaos measure constructed using the log-correlated field  $Y(x; q)$  defined for  $x \in [0, 1]$ . See Definition 2.2 for further discussion. With the choice of scaling  $P = \frac{P_0}{\gamma}$  and  $\alpha = \frac{\alpha_0}{\gamma}$  for  $P_0 \in \mathbb{R}$  and  $\alpha_0 \in \mathbb{R}_{>0}$ , the first difficulty is to show that  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$  does not blow up to infinity. We overcome this difficulty by showing that the (deep) tail probabilities  $\gamma^{-1} \log(\text{IGMC}_{\gamma, P}^\alpha(q))$  behave as the tails of a mean zero and finite variance Gaussian distribution. With replacing  $\gamma^{-1} \log(\text{IGMC}_{\gamma, P}^\alpha(q))$  in (1.4) by a mean zero and finite variance Gaussian random variable  $X$  yields

$$\mathcal{G}_{\gamma, P}^\alpha \sim \mathbb{E} \left[ \exp\left(-\frac{\alpha_0}{\gamma} X\right) \right] \sim \exp\left(\frac{C\alpha_0}{\gamma^2}\right)$$

for some constant  $C \in \mathbb{R}$ . This shows the tightness of the sequence  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$  as  $\gamma$  approaches 0. We studied the tail probabilities of  $\gamma^{-1} \log(\text{IGMC}_{\gamma, P}^\alpha(q))$  using the derivative GMC measure. For small  $\gamma$ , mean value theorem suggests  $\gamma^{-1} \log(\text{IGMC}_{\gamma, P}^\alpha(q))$  behaves as  $\text{DGMC}_{\gamma', P}^\alpha(q)$  for some  $\gamma' \in (0, \gamma)$  where, for the renormalized log-correlated field  $:Y(x; q):$ ,

$$\text{DGMC}_{\gamma', P}^\alpha(q) := \frac{1}{\int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} \int_0^1 :Y(x; q): \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\gamma:Y(x; q):} dx.$$

In Theorem 3.1, we show that the tail probabilities of  $\text{DGMC}_{\gamma', P}^\alpha(q)$  are like the Gaussian distribution. The proof of Theorem 3.1 is one of the most technical part of our paper and is built on ideas from several previous works in the study of GMC measure including the local decomposition of log-correlated fields as sum of \*-scale invariant field and a continuous field [JSW18] and the tools based on derivative GMC measure which are required to study the path integral of quantum Mabuchi K-energy [lacoïn2022path].

Once the tightness of  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$  is shown, it remains to show that all subsequential limits of  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$  are the same, and they are all uniquely characterized by the function  $\phi(\cdot)$  described in Theorem 1.1 that is related to the accessory parameter of the Lamé equation. To this end, we use the deformation of the conformal block  $\mathcal{G}_{\gamma, P}^\alpha$  which we will later denote as  $\psi_{\chi, P}^\alpha(z, \tau)$  (see Definition 2.3) for  $\chi \in \{\frac{\gamma}{2}, \frac{2}{\gamma}\}$ ,  $z \in (0, 1)$  and  $\alpha, P, \tau$  are same as before. Deformed conformal block is uniquely determined as the solution of a differential equation named the Belavin-Polyakov-Zamolodchikov equation,

$$\left( \partial_{zz} - l_\chi(l_\chi + 1)\wp(z) + 2i\pi\chi^2\partial_\tau \right) \psi_{\chi, P}^\alpha(z, \tau) = 0, \quad \ell_\chi = \frac{\chi^2}{2} - \frac{\alpha\chi}{2}.$$

For  $\chi = \frac{\gamma}{2}$ , we show in Theorem 4.2 that the deformed conformal block  $\psi_{\chi, P}^\alpha(z, \tau)$  factorizes approximately as  $\tilde{\Gamma}(z; \alpha_0, P_0, q) \times \mathcal{G}_{\gamma, P}^\alpha$  as  $\gamma \rightarrow 0$  where  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is same as in (1.3). Similar decomposition of deformed conformal block was previously observed in [Lit+14; Pia14]. Substituting this into the BPZ equation and letting  $\gamma \rightarrow 0$  with the scaling  $\alpha = \frac{\alpha_0}{\gamma}$  and  $P = \frac{P_0}{\gamma}$  gives us the Lamé equation as shown in (1.2). Since  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is explicit, the accessory parameter of (1.2) which could be  $\tau$ -derivative of any subsequential limit of  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$  is now uniquely identified which in turn shows the existence of the limit of the sequence  $\gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha$ .

On the other hand, for  $\chi = \frac{2}{\gamma}$ , the BPZ equation degenerates to the Hamilton-Jacobi equation as  $\gamma \rightarrow 0$ . Using this observation and by showing the tightness of  $\gamma^2 \log \psi_{\chi, P}^\alpha$  using the tails of the associated derivative GMC measure, we demonstrate in Theorem 4.1 that the semi-classical limit of  $\psi_{\chi, P}^\alpha$  solves the Hamilton-Jacobi equation for the non-autonomous elliptic Calogero Moser model (NAECM). See Remark 4.1 for a discussion. This connection between the deformed conformal

block and Calogero-Moser model is used later in Theorem 4.2 to show that the accessory parameter of the Lamé equation (1.2) is in fact related to the Hamiltonian of the NAECM model.

**1.3. Outline of the paper.** This paper is organized as follows. In Section 2, we review the probabilistic construction of the *undeformed* conformal block  $\mathcal{G}_{\gamma,P}^\alpha(q)$ , and the *deformed* conformal block which satisfies the PDE known as the Belavin-Polyakov-Zamolodchikov (BPZ) equation.

In Section 3, we introduce the main tool to analyse the semi-classical limit of the conformal blocks, namely, the DGMC measure, which is the  $\gamma$  derivative of the GMC measure. The main result of Section 3 is Theorem 3.1 which proves the tail estimates of the DGMC measure.

In Section 4, we formulate the semi-classical limit of the *deformed* conformal block. In Subsection 4.1 we study the semi-classical limit of the deformed conformal block for  $\chi = \frac{2}{\gamma}$ . The main result of this Subsection is Theorem 4.1 which establishes an explicit description of such semi-classical limit. In Subsection 4.2 we analyse the semi-classical limit of the deformed conformal block for  $\chi = \frac{\gamma}{2}$ . The main result of this Subsection is Theorem 4.2 which gives explicit description of that semi-classical limit in terms of the solution of Lamé equation. As a consequence of this theorem, we obtain explicit expressions for the solution and accessory parameter of the Lamé equation.

We prove Theorem 1.1 in Section 5. We further prove that the radius of convergence of the semi-classical conformal block is positive in Theorem 5.2.

Finally, in Section 6, we study the asymptotic behaviour of the semi-classical limit of the conformal blocks for  $\tau \rightarrow \mathbf{i}\infty$ . The key component of the analysis is the expression of the normalization constant of the undeformed conformal block in terms of double Gamma functions. Results in this section are used to prove results in Section 4.

**1.4. Outlook.** Our paper presents two key takeaways. Firstly, the methods we introduced in proving the Zamolodchikov conjecture offer a road map for studying semi-classical conformal blocks on other Riemann surfaces and for the analysis of conformal blocks with different central charges. Secondly, the explicit formulas for the conformal blocks and the Lamé equation obtained here provide profound insights into the analytic and geometric aspects of conformal field theories (CFTs) and the associated Heun-type equations. Let us now describe some open questions of immediate interest.

- A natural question that arises is the generalization of our techniques to other geometries such as the four-point sphere and higher genus surfaces. The key to achieving this would be to develop a systematic approach to defining probabilistic conformal blocks on Riemann surfaces. The formulation of probabilistic conformal blocks on a four point sphere is expected to have a slightly varied construction [Gho+], and the irregular conformal blocks can then be constructed through appropriate degeneration limits. The proof of the Zamolodchikov conjecture in the aforementioned cases is expected to follow a similar formalism to that presented here.
- Semi-classical limit of Liouville CFT corresponds to taking the central charge  $c$  to  $\infty$ . It is natural to ask what happens when  $c \rightarrow -\infty$ . This question is related to the semi-classical limit of CFTs with  $c \leq 1$  (i.e.,  $\gamma$  purely imaginary). Recently, in a series of works [Wan21; Wan19a; Wan19b], Yilin Wang has investigated the large deviation problem of Schramm-Loewner evolution (SLE $_{\kappa}$ ) as  $\kappa \rightarrow 0_+$  and has shown that the large deviation rate function is equal to the Loewner energy function. Since  $\kappa \rightarrow 0_+$  corresponds to taking  $c \rightarrow -\infty$  and SLE is related to the correlation function of certain CFT, the results of [Wan21; Wan19a; Wan19b] describes semi-classical limit for  $c \rightarrow -\infty$ . In a more recent work [PW24], Peltola and Wang characterized minimizers of the Loewner potential using tools from real algebraic geometry which provides unique insight about the CFT when the central charges diminishes to  $-\infty$ .

Besides the above questions, the explicit formulation of the semi-classical conformal block in terms of the DGMC measure presented in this paper merits an investigation of its own. On the other hand, one can obtain a series representation of the semi-classical conformal block through the Zamolodchikov recursion relations and a combinatorial relation coming from AGT correspondence (see [FL10]). It is important to study the interrelation between these explicit formulations. An important aspect of the explicit representation is the dependence of the semi-classical conformal

block on the monodromy data of the NAECM model and the Lamé equation, which would lead to interesting outcomes. We list a few a few of them.

- The Trieste formula [Bon+23, (4.1.16) - (4.1.17)], [LN22, eq. 2.20b] connects the connection formulae of the Heun's equation with the semi-classical limit of the fusion kernel of the 4-point sphere conformal blocks. Our description of the semi-classical conformal block may enable the proof of an analogue to the Trieste formula connecting the semi-classical limit of the modular kernel (see [Gho+, (5.3)]) of one-point torus conformal block with the connection formulae of the Lamé equation. A probabilistic description of the four-point conformal block and its degenerations could prove the Trieste formula in full generality.
- In [LN21], the authors found a series representation of the accessory parameter of the Heun's equation where the coefficients satisfies some recurrence relation. This is reminiscent of the Zamolodchikov recurrence relation satisfied by the coefficients of the conformal blocks. Following a similar procedure as in [LN21], one can also derive a series representation of the accessory parameter of Lamé equation once the monodromy data associated to  $\alpha_0, P_0$  is determined. Furthermore, the accessory parameter is also explicitly given in terms of the derivative GMC measure in (4.24), and in terms of integrals (4.40). Such formulae would allow to study its asymptotic properties [Nov86; XXZ21].
- Through the AGT correspondence, the semi-classical conformal block on a one-point torus determines the Nekrasov-Shatashvili (NS) limit of the corresponding  $\mathcal{N} = 2^*$  gauge theory, or equivalently, the effective twisted superpotential [Tes11]. The explicit form of the classical conformal block presented in this paper would enable a rigorous proof of the Nekrasov-Rosly-Shatashvili conjecture that relates the effective twisted superpotential of the gauge theory to the so-called Yang-Yang function of the associated quantum Hitchin system [NS09; NRS11; HK18].
- The conformal blocks for arbitrary central charge solve certain bilinear relations called the Nakajima-Yoshioka blow-up relations [NY03]. Such relations for the one-point torus have been written in [Gu+20; BGG22] and it will be interesting to see if such relations could be proved using probabilistic conformal blocks. An interesting property of the relations is that the blow-up relation connects the semi-classical conformal blocks with the conformal block at central charge  $c = 1$  [NS10], in turn relating Painlevé equations to Heun equations for the sphere case and the NAECM model to Lamé equation (see also [BGG22]) for the torus case. We show the latter relation in Theorem 4.2 based on probabilistic ideas. Showing the full blow-up relation is beyond the scope of this paper and we leave this for a future work.

**Acknowledgements.** We thank Alba Grassi, Aleksandra Korzhenkova, Oleg Lisovyi, Alex Moll, Eveliina Peltola, Vincent Vargas for informative discussions. Part of this work was done during the authors' residence at the SLMath (formerly, Mathematical Sciences Research Institute) in Berkeley, California, during the program 'Universality and Integrability in Random Matrix Theory and Interacting Particle Systems'. They acknowledge the support of the National Science Foundation Grant No. DMS-1928930. H.D. also acknowledges the support of the Okinawa Institute of Science and Technology (OIST), where she completed part of this work during her stay for the Theoretical Sciences Visiting Program (TSVP) "Exact Asymptotics: From Fluid Dynamics to Quantum Geometry."

The work of H.D. is partly supported by Australian Research Council Discovery Project #DP200100210, INI-Simons fellowship, and SMRI postdoctoral fellowship. The work of P.G. is partially supported by the NSF grant DMS-2346685. A.P. is supported NSF MSPRF grant DMS-2103354, and RSF grant 22-11-00070.

## 2. PROBABILISTIC CONFORMAL BLOCKS

In this section we recall the construction of the conformal block in [Gho+24]. We begin by defining Gaussian Multiplicative Chaos (GMC) measures in terms of log-correlated fields, which will then be used to define the *probabilistic* one point conformal block on the torus. We then introduce the notion of a *deformed* conformal block which solves the PDE known as the Belavin-Polyakov-Zamolodchikov (BPZ) equation.

**2.1. Log-correlated fields and gaussian multiplicative chaos (GMC) measures.** For  $x \in [0, 1]$ , the random fields  $Y_N(x; q)$ ,  $Y_N(x)$ , with  $N \in \mathbb{N}$  are defined as

$$\tilde{Y}_N(x) = \sum_{n=1}^N \sqrt{\frac{2}{n}} \left( \tilde{a}_n \cos(2\pi n x) + \tilde{b}_n \sin(2\pi n x) \right), \quad (2.1)$$

$$\tilde{Y}_N(x; q) = \tilde{Y}_N(x) + 2 \underbrace{\sum_{n,m=1}^{\infty} \frac{q^{nm}}{\sqrt{n}} \left( \tilde{a}_{n,m} \cos(2\pi n x) + \tilde{b}_{n,m} \sin(2\pi n x) \right)}_{=: F(x; q)}, \quad (2.2)$$

where  $\{\tilde{a}_n\}_{n \geq 1}$ ,  $\{\tilde{b}_n\}_{n \geq 1}$ ,  $\{\tilde{a}_{n,m}\}_{n,m \geq 1}$ ,  $\{\tilde{b}_{n,m}\}_{n,m \geq 1}$  are sequences of i.i.d standard real Gaussian random variables, the modular parameter  $\tau \in \mathbb{H}$ , and  $q = e^{i\pi\tau}$ . In what follows we use the notation

$$Y(x) := \lim_{N \rightarrow \infty} \tilde{Y}_N(x), \quad Y(x; q) := \lim_{N \rightarrow \infty} \tilde{Y}_N(x; q). \quad (2.3)$$

Indeed, the above limit exists. The second limit exist in the sense of weak convergence in the Sobolev space  $H^{-s}([0, 1])$  and the second limit exists due to the first limit and the almost convergence of infinite series of  $F(x; q)$  due to the term  $q^{nm}$ . The resulting random fields  $Y(\cdot; q)$  and  $Y(\cdot)$  are log-correlated, and the covariance kernels of (2.3) for  $\tau \in \mathbf{iR}_{>0}$  are given as follows.

- The field  $Y(x)$  is log-correlated on  $[0, 1]$ :

$$\mathbb{E}[Y(x)Y(y)] = -2 \log |e^{2i\pi x} - e^{2i\pi y}|, \quad (2.4)$$

which is the usual expression  $\log |x - y|$  in cylindrical coordinates obtained by the transformation  $z \rightarrow e^{2\pi i z}$ .

- The  $q$ -dependent field  $Y(x; q)$  is log correlated on a doubly-periodic lattice

$$\mathbb{E}[Y(x; q)Y(y; q)] = -2 \log |\theta_1(x - y)| + 2 \log |q^{1/6} \eta(q)|. \quad (2.5)$$

Note that for  $q \rightarrow 0$  and  $(x - y) \rightarrow 0$ , the above correlation functions (2.4), (2.5) coincide up to a trivial factor.

Using [Gho+24, Lemma 2.5], one finds that  $\mathbb{E}[F(x; q)^2] = -4 \log |q^{-1/12} \eta(q)|$ . As a result, we get

$$\mathbb{E}[\tilde{Y}_N(x; q)^2] = \mathbb{E}[\tilde{Y}_N(x)^2] - 4 \log |q^{-1/12} \eta(q)|. \quad (2.6)$$

With the above log-correlated fields, we now define the following random measures.

**Definition 2.1** (GMC measure). *For  $\gamma \in (0, 2)$  and  $\tau \in \mathbf{iR}_{>0}$ , the GMC measures  $e^{\frac{\gamma}{2}Y(x)} dx$  and  $e^{\frac{\gamma}{2}Y(x; q)} dx$  are defined to be the weak limits of measures in probability*

$$\begin{aligned} e^{\frac{\gamma}{2}Y(x)} dx &:= \lim_{N \rightarrow \infty} e^{\frac{\gamma}{2}\tilde{Y}_N(x) - \frac{\gamma^2}{8}\mathbb{E}[\tilde{Y}_N(x)^2]} dx, \\ e^{\frac{\gamma}{2}Y(x; q)} dx &:= \lim_{N \rightarrow \infty} e^{\frac{\gamma}{2}\tilde{Y}_N(x; q) - \frac{\gamma^2}{8}\mathbb{E}[\tilde{Y}_N(x; q)^2]} dx. \end{aligned} \quad (2.7)$$

For a test function  $f(x)$ , these measures are interpreted as

$$\int_0^1 f(x) e^{\frac{\gamma}{2}Y(x; q)} dx = \lim_{N \rightarrow \infty} \int_0^1 f(x) e^{\frac{\gamma}{2}\tilde{Y}_N(x; q) - \frac{\gamma^2}{8}\mathbb{E}[\tilde{Y}_N(x; q)^2]} dx, \quad (2.8)$$

and a similar expression holds for the  $q$ -independent field  $Y(x)$ .

The existence of the limits above can be shown in the following way. Note that the identity

$$\mathbb{E}[e^{c(\tilde{Y}_k - \tilde{Y}_{k-1})}] = e^{\frac{c^2}{2}},$$

implies that the sequence in (2.8) is a martingale. Since it is positive, almost sure convergence is guaranteed by the martingale convergence theorem, which in turn proves the existence of the limit in (2.8).

**2.2. One-point (undeformed) conformal block.** We now recall the definition of the one-point conformal block in [Gho+24].

**Definition 2.2.** For  $\alpha \in (-\frac{4}{\gamma}, Q)$ , and  $P \in \mathbb{R}$ , we define the probabilistic 1-point toric conformal block as

$$\mathcal{G}_{\gamma,P}^\alpha(q) := \frac{1}{Z_{\gamma,P}^\alpha} \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x;q)} dx \right)^{-\frac{\alpha}{\gamma}} \right], \quad (2.9)$$

with the normalization

$$Z_{\gamma,P}^\alpha(q) := q^{\frac{1}{12}(\frac{\alpha\gamma}{2} + \frac{\alpha^2}{2} - 1)} \eta(q)^{\alpha^2 + 1 - \frac{\alpha\gamma}{2}} \mathbb{E} \left[ \left( \int_0^1 [-2 \sin(\pi x)]^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right]. \quad (2.10)$$

Note that the normalization constant  $Z_{\gamma,P}^\alpha$  ensures that

$$\lim_{q \rightarrow 0} \mathcal{G}_{\gamma,P}^\alpha(q) = 1, \quad \lim_{P \rightarrow \infty} \mathcal{G}_{\gamma,P}^\alpha(q) = q^{\frac{1}{12}} \eta(q)^{-1}.$$

A key consequence of the present construction is that the analytic properties of the conformal block (2.9) become accessible, and we recall the following result.

**Theorem 2.1** ([Gho+24, Theorem 1.1]). For  $\gamma \in (0, 2)$ ,  $\alpha \in (-\frac{4}{\gamma}, Q)$ , and  $P \in \mathbb{R}$ , the conformal block in (2.9) is analytic in a complex neighborhood of  $q = 0$ . Moreover, for  $\alpha \in [0, Q)$ , the analytic extension of (2.9) exists in a neighborhood of  $q = 0$  with radius greater than  $\frac{1}{2}$ .

Furthermore, (2.9) can be represented a series with the coefficients satisfying the so-called Zamolodchikov recursion [Gho+24, Proposition 2.11], and as a combinatorial series in terms of Nekrasov-Okounkov functions [Gho+24, Section 2.3] due to the Alday-Gaiotto-Tachikawa (AGT) correspondence [Neg16].

**Remark 2.1.** For  $-\frac{\alpha}{\gamma} = N < \frac{4}{\gamma^2}$  with  $N \in \mathbb{N}$ , (2.9) can be expressed as a Dotsenko-Fadeev integral [MMS10; Fat+09]

$$\begin{aligned} \mathcal{G}_{\gamma,P}^\alpha(q) &= \frac{1}{Z_{\gamma,P}^\alpha} \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\gamma\pi Px} e^{\frac{\gamma}{2}Y(x;q)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ &= \frac{1}{C} \int_{[0,1]^N} \prod_{1 \leq i < j \leq N} |\theta_1(x_i - x_j)|^{-\frac{\gamma^2}{4}} \prod_{i=1}^N \theta_1(x_i)^{\frac{N\gamma^2}{2}} e^{\pi\gamma Px_i} \prod_{i=1}^N dx_i, \end{aligned}$$

with

$$C = \frac{e^{\frac{N\gamma P\pi}{2}}}{\left(\Gamma\left(1 - \frac{\gamma^2}{8}\right)\right)^N} \left(\frac{\theta_1'(0)}{2}\right)^{\frac{N(N+1)\gamma^2}{8}} \prod_{k=1}^N \frac{\Gamma\left(1 - \frac{k\gamma^2}{8}\right) \Gamma\left(1 + \frac{(2N+1-k)\gamma^2}{8}\right)}{\Gamma\left(1 + \frac{k\gamma^2}{8} + \frac{i\gamma P}{2}\right) \Gamma\left(1 + \frac{k\gamma^2}{8} - \frac{i\gamma P}{2}\right)}. \quad (2.11)$$

Let us highlight the following result that will prove useful to study the  $q \rightarrow 0$  asymptotics of the semi-classical conformal blocks.

**Proposition 2.1.** The partition function  $Z$  in (2.9) can be expressed in terms of double Gamma functions as follows [Gho+24, (2.15), Proposition 6.4]

$$Z_{\gamma,P}^\alpha = q^{\frac{1}{12}(\frac{\alpha\gamma}{2} + \frac{\alpha^2}{2} - 1)} \eta(q)^{\alpha^2 + 1 - \frac{\alpha\gamma}{2}} \mathcal{A}_{\gamma,P}(\alpha), \quad (2.12)$$

where the  $q$ -independent factor is

$$\begin{aligned} \mathcal{A}_{\gamma,P}(\alpha) &= e^{\frac{i\pi\alpha^2}{2}} \left(\frac{\gamma}{2}\right)^{\frac{\gamma\alpha}{4}} e^{-\frac{\pi\alpha P}{2}} \Gamma\left(1 - \frac{\gamma^2}{4}\right)^{\frac{\alpha}{\gamma}} \\ &\times \frac{\Gamma_{\frac{\gamma}{2}}\left(Q - \frac{\alpha}{2}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma} + \frac{\alpha}{2}\right) \Gamma_{\frac{\gamma}{2}}\left(Q - \frac{\alpha}{2} - iP\right) \Gamma_{\frac{\gamma}{2}}\left(Q - \frac{\alpha}{2} + iP\right)}{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(Q - iP\right) \Gamma_{\frac{\gamma}{2}}\left(Q + iP\right) \Gamma_{\frac{\gamma}{2}}\left(Q - \alpha\right)}, \end{aligned} \quad (2.13)$$

where the function  $\Gamma_{\frac{\gamma}{2}}(\cdot)$  is the ratio of double Gamma functions

$$\Gamma_{\frac{\gamma}{2}}(x) = \frac{\Gamma_2(x|\gamma, \gamma^{-1})}{\Gamma_2(Q/2|\gamma, \gamma^{-1})},$$

and  $Q$  is given by (A.1)

**2.3.  $z$ -deformed conformal block.** We now define the probabilistic equivalent of inserting degenerate vertex operator  $\mathcal{V}$  inside conformal block as

$$\mathcal{V}_{\chi, P}^{\alpha}(z, q) := \int_0^1 \theta_1(z+x)^{\frac{\gamma\chi}{2}} \theta_1(z)^{-\frac{\gamma\chi}{2}} \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x; q)} dx \quad \text{for } z \in \mathfrak{B}, \quad \chi \in \left\{ \frac{2}{\gamma}, \frac{\gamma}{2} \right\}$$

with  $\mathfrak{B} := \{z : 0 < \Im(z) < \frac{3}{4}\Im(\tau)\}$ . The deformed conformal block is then in terms of  $\mathcal{V}$  as follows.

**Definition 2.3.** For  $x \in (0, 1)$ ,  $q \in (0, 1)$ ,  $\alpha \in \left(-\frac{4}{\gamma} + \chi, Q\right)$ ,  $P \in \mathbb{R}$ , the two point toric conformal block with one degenerate insertion, known as the  $z$ -deformed probabilistic conformal block is defined as

$$\psi_{\chi, P}^{\alpha}(z, \tau) := \mathcal{W}(q) e^{\chi P z \pi} \mathbb{E} \left[ \left( \mathcal{V}_{\chi, P}^{\alpha}(z, q) \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right], \quad (2.14)$$

where

$$\mathcal{W}(q) := q^{\frac{P^2}{2} + \frac{\gamma l_{\chi}}{12\chi} - \frac{1}{6} \frac{l_{\chi}^2}{\chi^2}} \theta_1'(0) - \frac{2l_{\chi}^2}{3\chi^2} + \frac{l_{\chi}}{3} + \frac{4l_{\chi}}{3\gamma\chi}, \quad l_{\chi} = \frac{\chi^2}{2} - \frac{\alpha\chi}{2}, \quad (2.15)$$

and  $\theta_1'(0)$  given by (A.5).

**Theorem 2.2** ([Gho+24, Proposition 3.2]). The deformed conformal block (2.14) is convergent in the domain  $D_{\chi}^{\alpha}$  where

$$D_{\chi}^{\alpha} := \{(q, z) : |q| < r_{\alpha-\chi} \text{ and } z \in \mathfrak{B}\}, \quad \mathfrak{B}_q := \{z : 0 < \Im(z) < \frac{3}{4}\Im(\tau)\}$$

Here  $r_{\alpha} > 0$  for  $\alpha \in \left(-\frac{4}{\gamma}, Q\right)$ .

**Remark 2.2.**

1. The choice of the pre-factor  $\mathcal{W}(q)$  in (2.14) is dictated by the BPZ equation which will be satisfied by  $\psi_{\chi, P}^{\alpha}(z, \tau)$  (see Theorem 2.3). The expression for  $\mathcal{W}(q)$  will play a major role in fixing the monodromy of the probabilistic conformal block, and this will be expanded upon in a future work.
2. The deformed and the undeformed conformal blocks are related as

$$\lim_{z \rightarrow 0} \left( \theta_1(z)^{l_{\chi}} \psi_{\chi, P}^{\alpha}(z, \tau) \right) = \mathcal{W}(q) Z_{\gamma, P}^{\alpha-\chi}(q) \mathcal{G}_{\gamma, P}^{\alpha-\chi}(q). \quad (2.16)$$

We now state the BPZ equation for  $\psi_{\chi, P}^{\alpha}(z, \tau)$ .

**Theorem 2.3** ([Gho+24, Theorem 3.5]). The  $z$ -deformed toric conformal block satisfies the following BPZ equation

$$\left( \partial_{zz} - l_{\chi}(l_{\chi} + 1)\wp(z) + 2i\pi\chi^2\partial_{\tau} \right) \psi_{\chi, P}^{\alpha}(z, \tau) = 0 \quad \text{for } (q, z) \in D_{\chi}^{\alpha}, \quad (2.17)$$

where  $\alpha \in \left(-\frac{4}{\gamma} + \chi, Q\right)$ ,  $\chi \in \left\{ \frac{\gamma}{2}, \frac{2}{\gamma} \right\}$ , and  $\wp(z)$  is Weierstrass  $\wp$ -function.

### 3. DERIVATIVE GAUSSIAN MULTIPLICATIVE CHAOS MEASURE

The semi-classical limit of the undeformed conformal block is the  $\gamma \rightarrow 0$  limit of  $\mathcal{G}_{\gamma, P}^{\alpha}(q)$ . From (2.9), it turns out to be the case that the convergence of the semi-classical limit is controlled by the convergence of the  $\gamma$  derivative of the GMC measure (2.8), known as the derivative GMC (DGMC) measure.

**Definition 3.1** (DGMC measure). For a bounded test function  $f(x, \tau) \in L_\infty([0, 1])$  for every  $\tau \in \mathbb{H}$  the DGMC measure is defined as

$$(\text{DM}_{\gamma, N} f)(\tau) := \int_0^1 \left( \frac{1}{2} Y_N(x; q) - \frac{\gamma}{4} \mathbb{E}[Y_N^2(x; q)] \right) e^{\frac{\gamma}{2} Y_N(x; q) - \frac{\gamma^2}{8} \mathbb{E}[Y_N^2(x; q)]} f(x, \tau) \frac{dx}{\int_0^1 f(x, \tau) dx}. \quad (3.1)$$

The  $N \rightarrow \infty$  limit of the measure above is treated in the same way as (2.7):

$$(\text{DM}_\gamma f)(\tau) = \lim_{N \rightarrow \infty} (\text{DM}_{\gamma, N} f)(\tau). \quad (3.2)$$

**Theorem 3.1.** For a bounded function  $f$ , there exist  $c_1 = c_1(\tau) > 0$ ,  $c_2 = c_2(\tau) > 0$ ,  $c_3 > 0$ ,  $c_4 > 0$  such that for all  $v > 0$ ,

$$\mathbb{P}[(\text{DM}_{\gamma, N} f)(\tau) \leq -v] \leq c_1(\tau) e^{-c_2(\tau)v^2}, \quad \mathbb{P}[(\text{DM}_{\gamma, N} f)(\mathbf{i}\infty) \leq -v] \leq c_3 e^{-c_4 v^2}. \quad (3.3)$$

**Remark 3.1.** It is worthwhile to note that [LRV22, Theorem 4.4] proved similar result for the DGMC when the underlying field of the GMC satisfy the so called smooth white noise decomposition property (see Assumption 4.1 of [LRV22]). However the log-correlated fields  $Y_\infty$  and  $Y_\tau$  does not satisfy all those assumptions. Motivated by their results, we provide a self-contained proof of Theorem 3.1 by utilizing the infinite series representation of the fields  $Y_\infty$  and  $Y_\tau$  (as shown in (2.2) and (2.1)).

The remainder of this section is dedicated to proving Theorem 3.1. We restrict to  $Y_N(x)$  and prove the second inequality in (3.3). Because of (2.6), and the  $N$  independence of  $F(x; q)$ , the bound of  $(\text{DM}_{\gamma, N} f)(\tau)$  will coincide with the bounds of  $(\text{DM}_{\gamma, N} f)(\mathbf{i}\infty)$ , a fact that can be verified by repeating the statements in the proof below for  $Y_N(x; q)$ .

We begin the proof of Theorem 3.1 by introducing the following shorthand notation and proving some important properties of the DGMC measure. These properties will later be used to show the tail of  $(\text{DM}_{\gamma, N} f)(\mathbf{i}\infty)$ .

By Theorem B of [JSW18], a log-correlated field  $Y$  can be locally decomposed as  $Y = \mathcal{L} + \mathcal{R}$  where the local component  $\mathcal{L}$  is almost  $*$ -scale invariant field and the global component  $\mathcal{R}$  is a regular Gaussian field with Hölder continuous realizations. Moreover,  $\mathcal{L}$  and  $\mathcal{R}$  are independent.

Let  $\{P_1, \dots, P_m\}$  be a partition of the interval  $[0, 1]$  such that  $Y$  could be expressed as  $Y(x) = \mathcal{L}^{(\ell)} + \mathcal{R}^{(\ell)}$  for some  $*$ -scale invariant field  $\mathcal{L}^{(\ell)}$  and a continuous field  $\mathcal{R}^{(\ell)}$ . In the following discussion, we focus on bounding the tail probabilities of  $(\text{DM}_{\gamma, N} f)(\tau)$  for any bounded function supported on one of the partitioning intervals  $\{P_1, \dots, P_m\}$ . For notational convenience, we denote the  $*$ -scale invariant field  $\mathcal{L}^{(\ell)}$  and  $\mathcal{R}^{(\ell)}$  for the interval  $P_\ell$  as  $\mathcal{L}$  and  $\mathcal{R}$ . However, the tail probability of  $(\text{DM}_{\gamma, N} f)(\tau)$  for any other bounded function  $f$  could be found by applying the union bound.

Let  $\mathcal{K}_\mathcal{L}$  be the covariance kernel of the field  $\mathcal{L}$ . Since  $\mathcal{L}$  is a  $*$ -scale invariant field,  $\mathcal{K}_\mathcal{L}$  is translation invariant and it could be written as

$$\mathcal{K}_\mathcal{L}(x, y) =: \mathcal{K}_\mathcal{L}(x - y) = \int_0^\infty K(e^u(x - y)) du, \quad \forall x \neq y \in [0, 1]$$

where  $k$  is a covariance in  $\mathbb{R}$ ,  $K(0) = 1$  and  $|K(x)| \lesssim |x|^{-a}$  for some  $a < 1$ . In particular, we can choose  $a \in (0, 1)$ . This choice of  $a$  determines the number of intervals in the partition  $\{P_1, \dots, P_m\}$  of  $[0, 1]$ . Furthermore, by Proposition 4.1 of [JSW18], there exists sequence of Gaussian fields  $\{\mathcal{L}_n\}_{k \geq 1}$  and sigma algebras  $\mathfrak{F}_1 \subset \mathfrak{F}_2 \subset \mathfrak{F}_3 \subset \dots$  such that  $\mathcal{L}_k$  is measurable w.r.t.  $\mathfrak{F}_k$  and,

$$\mathbb{E}[\mathcal{L}_{n_1}(x) \mathcal{L}_{n_2}(y)] = \int_0^{n_1 \wedge n_2} K(e^u(x - y))(1 - e^{-\delta u}) du, \quad (3.4)$$

where  $n_1 \wedge n_2$  indicates  $\min\{n_1, n_2\}$ . We define  $Y_k := \mathcal{L}_k + \mathcal{R}$  and further introduce the following notations:

$$\Lambda_k(x) := Y_k(x) - Y_{k-1}(x), \quad \Omega_k(x) := \frac{1}{2} \mathbb{E}[Y_k^2(x)], \quad (3.5)$$

By the independence between  $\Lambda_k$  and  $Y_{k-1}$ , we get

$$e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{8}\mathbb{E}[Y_k^2(x)]} \stackrel{(3.5)}{=} e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{8}\mathbb{E}[Y_{k-1}^2(x)]} e^{\frac{\gamma}{2}\Lambda_k(x) - \frac{\gamma^2}{8}\mathbb{E}[\Lambda_k(x)^2]}.$$

To begin with, we show that  $(\mathcal{D}_k, \mathfrak{F}_k)_{k \geq 1}$  is a martingale, where

$$\mathcal{D}_k := \int_0^1 W_k^0(x) f(x) dx, \quad \Omega_k(x) = \frac{1}{2} \mathbb{E}[(\mathcal{L}_k(x) + \mathcal{R}(x))^2], \quad (3.6)$$

and,

$$W_k^0(x) := (Y_k(x) - \gamma \Omega_k(x)) e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{4}\Omega_k(x)}. \quad (3.7)$$

The test function  $f(x)$  is  $\gamma$  independent, and  $Y_k(x) = \mathcal{L}_k(x) + \mathcal{R}(x)$  for  $k \in \mathbb{R}$  and  $x \in [0, 1]$ .

**Proposition 3.1.** *The following statements are true*

- (1)  $(\mathcal{D}_k, \mathfrak{F}_k)_{k \geq 1}$  is a martingale and  $\mathcal{D}_k$  is bounded below uniformly in  $k$ .
- (2)  $\mathcal{D}_k$  converges almost surely to  $\mathcal{D}_\infty$ .

*Proof.*

(1): To prove this statement, it suffices to show that the conditional expectation

$$\mathbb{E}[\mathcal{D}_k | \mathfrak{F}_{k-1}] = \mathcal{D}_{k-1}.$$

We define  $\Lambda_k(x) := Y_k(x) - Y_{k-1}(x)$  for all  $x \in [0, 1]$ . Indeed, we have

$$\begin{aligned} \mathbb{E}[\mathcal{D}_k | \mathfrak{F}_{k-1}] &= \int_0^1 (Y_{k-1}(x) - \gamma \Omega_{k-1}(x)) \mathbb{E}[e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{4}\Omega_k(x)} | \mathfrak{F}_{k-1}] f(x) dx \\ &\quad + \int_0^1 \mathbb{E}\left[\left(\Lambda_k(x) - \frac{\gamma}{2}\mathbb{E}[\Lambda_k^2(x)]\right) e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{4}\Omega_k(x)} | \mathfrak{F}_{k-1}\right] f(x) dx \\ &= \int_0^1 (Y_{k-1}(x) - \gamma \Omega_{k-1}(x)) e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)} f(x) dx = \mathcal{D}_{k-1}. \end{aligned}$$

To obtain the last step, we use the identities below that follow from (3.5):

$$\begin{aligned} \mathbb{E}[e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{4}\Omega_k(x)} | \mathfrak{F}_{k-1}] &= \mathbb{E}[e^{\frac{\gamma}{2}(Y_{k-1}(x) + \Lambda_k(x)) - \frac{\gamma^2}{4}(\Omega_{k-1} + \mathbb{E}[\Lambda_k^2(x)])} | \mathfrak{F}_{k-1}] \\ &= e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)}, \end{aligned}$$

and

$$\int_0^1 \mathbb{E}\left[\left(\Lambda_k(x) - \frac{\gamma}{2}\mathbb{E}[\Lambda_k^2(x)]\right) e^{\frac{\gamma}{2}Y_k(x) - \frac{\gamma^2}{4}\Omega_k(x)} | \mathfrak{F}_{k-1}\right] f(x) dx = 0.$$

This proves the claim that  $(\mathcal{D}_k, \mathfrak{F}_k)_{k \geq 1}$  is a martingale.

(2): We show that  $\{\mathcal{D}_k\}_{k \geq 1}$  is a Cauchy sequence in  $L_2$ . Thus, by Doob's martingale convergence theorem,  $\mathcal{D}_k$  almost surely converges to  $\mathcal{D}_\infty$ .

Let us compute the following expression for  $k > n$ :

$$\begin{aligned} &\mathbb{E}[(\mathcal{D}_k - \mathcal{D}_n)^2] \\ &= \mathbb{E}[\mathcal{D}_k^2] + \mathbb{E}[\mathcal{D}_n^2] - 2\mathbb{E}[\mathcal{D}_k \mathcal{D}_n] = \mathbb{E}[\mathcal{D}_k^2] + \mathbb{E}[\mathcal{D}_n^2] - 2\mathbb{E}[\mathcal{D}_n \mathbb{E}[\mathcal{D}_k | \mathfrak{F}_n]] \\ &= \mathbb{E}[\mathcal{D}_k^2] - \mathbb{E}[\mathcal{D}_n^2] \\ &\stackrel{(3.6)}{=} \int_0^1 \int_0^1 \mathbb{E}[W_k^0(x) W_k^0(y)] f(x) f(y) dx dy - \int_0^1 \int_0^1 \mathbb{E}[W_n^0(x) W_n^0(y)] f(x) f(y) dx dy, \quad (3.8) \end{aligned}$$

where we use the tower property to obtain the last line. Note that, from the decomposition  $Y_k = \mathcal{L}_k + \mathcal{R}$ , and since  $K(0) = 1$ ,

$$\Omega_k(x) \stackrel{(3.4)}{=} \frac{1}{2} \mathbb{E}[\mathcal{R}(x)^2] + \frac{1}{2} \int_0^k (1 - e^{-\delta u}) du. \quad (3.9)$$

In what follows, we denote the bound of  $\Omega_k(x)$  by  $\Omega_k$ . Let us now expand the second term in (3.8):

$$\begin{aligned} & \mathbb{E} [W_n^0(x)W_n^0(y)] \\ & \stackrel{(3.7)}{=} \mathbb{E} \left[ (Y_n(x) - \gamma \Omega_n(x)) (Y_n(y) - \gamma \Omega_n(y)) e^{\frac{\gamma}{2}Y_n(x) - \frac{\gamma^2}{4}\Omega_n(x)} e^{\frac{\gamma}{2}Y_n(y) - \frac{\gamma^2}{4}\Omega_n(y)} \right] \\ & = e^{-\frac{\gamma^2}{4}(\Omega_n(x) + \Omega_n(y))} \left( \mathbb{E} \left[ Y_n(x)Y_n(y) e^{\frac{\gamma}{2}(Y_n(x) + Y_n(y))} \right] + \gamma^2 \Omega_n(x)\Omega_n(y) \mathbb{E} \left[ e^{\frac{\gamma}{2}(Y_n(x) + Y_n(y))} \right] \right. \\ & \quad \left. - \gamma \Omega_n(y) \mathbb{E} \left[ Y_n(x) e^{\frac{\gamma}{2}(Y_n(x) + Y_n(y))} \right] - \gamma \Omega_n(x) \mathbb{E} \left[ Y_n(y) e^{\frac{\gamma}{2}(Y_n(x) + Y_n(y))} \right] \right). \end{aligned}$$

Similarly, the first term in (3.8) can be written as

$$\begin{aligned} & \mathbb{E} [W_k^0(x)W_k^0(y)] \\ & \stackrel{(3.7)}{=} \mathbb{E} \left[ (Y_n(x) + \sum_{\ell=n+1}^k \Lambda_\ell(x) - \gamma \Omega_k) (Y_n(y) + \sum_{\ell=n+1}^k \Lambda_\ell(y) - \gamma \Omega_k) \right. \\ & \quad \left. \times e^{\frac{\gamma}{2}(Y_n(y) + \sum_{\ell=n+1}^k \Lambda_\ell(y)) - \frac{\gamma^2}{4}\Omega_k(y)} e^{\frac{\gamma}{2}(Y_n(x) + \sum_{\ell=n+1}^k \Lambda_\ell(x)) - \frac{\gamma^2}{4}\Omega_k(x)} \right] \\ & = \mathbb{E} [W_n^0(x)W_n^0(y)] (1 + \Xi(x, y)) \\ & + \mathbb{E} \left[ \left( \sum_{\ell=n+1}^k \Lambda_\ell(x) - \gamma(\Omega_k - \Omega_n) \right) \left( \sum_{\ell=n+1}^k \Lambda_\ell(y) - \gamma(\Omega_k - \Omega_n) \right) e^{\frac{\gamma}{2} \sum_{\ell=n+1}^k (\Lambda_\ell(y) + \Lambda_\ell(x)) - \frac{\gamma^2}{2}(\Omega_k - \Omega_n)} \right] \\ & + \mathbb{E} \left[ (Y_n(x) - \gamma \Omega_n(x)) \left( \sum_{\ell=n+1}^k \Lambda_\ell(y) - \gamma(\Omega_k - \Omega_n) \right) e^{\frac{\gamma}{2}(Y_n(x) + Y_n(y) + \sum_{\ell=n+1}^k \Lambda_\ell(y)) - \frac{\gamma^2}{4}\Omega_k(x)} \right] \\ & + \mathbb{E} \left[ (Y_n(y) - \gamma \Omega_n(y)) \left( \sum_{\ell=n+1}^k \Lambda_\ell(x) - \gamma(\Omega_k - \Omega_n) \right) e^{\frac{\gamma}{2}(Y_n(y) + Y_n(x) + \sum_{\ell=n+1}^k \Lambda_\ell(x)) - \frac{\gamma^2}{4}\Omega_k(y)} \right]. \quad (3.10) \end{aligned}$$

By the tower property of the conditional expectation, the last two terms of the expression above equal to 0. The second term in the right hand side of the above expression can be analysed in the following way. By Gaussian integration by parts, we get

$$\begin{aligned} & \mathbb{E} \left[ \left( \sum_{\ell=n+1}^k \Lambda_\ell(x) - \gamma(\Omega_k - \Omega_n) \right) \left( \sum_{\ell=n+1}^k \Lambda_\ell(y) - \gamma(\Omega_k - \Omega_n) \right) e^{\frac{\gamma}{2} \sum_{\ell=n+1}^k (\Lambda_\ell(y) + \Lambda_\ell(x)) - \frac{\gamma^2}{2}(\Omega_k - \Omega_n)} \right] \\ & = e^{\frac{\gamma^2}{2} \sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)]} \left[ \left( \gamma \mathbb{E} \left[ \sum_{\ell=n+1}^n (\Lambda_\ell(x) + \Lambda_\ell(y)) \Lambda_\ell(x) \right] - \gamma(\Omega_k - \Omega_n) \right) \right. \\ & \quad \left. \times \left( \gamma \mathbb{E} \left[ \sum_{\ell=n+1}^n (\Lambda_\ell(x) + \Lambda_\ell(y)) \Lambda_\ell(y) \right] - \gamma(\Omega_k - \Omega_n) \right) + \mathbb{E} \left[ \sum_{\ell=n+1}^n \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \right] \right] \\ & = e^{\frac{\gamma^2}{2} \sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)]} \left[ \gamma^2 \left( \sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \right)^2 + \sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \right]. \end{aligned}$$

For any  $x \neq y \in [0, 1]$ ,  $\sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \rightarrow 0$  as  $n, k \rightarrow \infty$  because  $\sum_{\ell=1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)]$  converges to  $\mathbb{E}[\mathcal{L}(x)\mathcal{L}(y)]$  as  $k \rightarrow \infty$ . This implies that the last line of the above display converges to 0 as  $k, n \rightarrow \infty$  and  $x \neq y$ . We further notice that

$$\sum_{\ell=n+1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \leq \sum_{\ell=1}^k \mathbb{E}[\Lambda_\ell(x)\Lambda_\ell(y)] \leq -2 \log |\sin(2\pi(x - y))|.$$

Thus, the last line of the above display can be uniformly bounded by  $C|\sin(2\pi(x - y))|^{-\gamma^2 + \delta}$  for some  $\delta > 0$ . As a result, the second term of (3.10) is uniformly integrable and hence, integral of that term we obtain in (3.8) converges to 0. Using this fact into the right hand side of (3.8)

shows that  $\mathbb{E}[(\mathcal{D}_k - \mathcal{D}_n)^2] \rightarrow 0$  as  $k, n \rightarrow \infty$ . Since  $\{(\mathcal{D}_k, \mathfrak{F}_k)\}_k$  is a martingale and a  $L_2$  Cauchy sequence,  $\mathcal{D}_k$  almost surely converges to  $\mathcal{D}_\infty$  by Doob's martingale convergence theorem.  $\square$

We recall that the quadratic variation  $\langle \cdot \rangle_n$  of the martingale  $\mathcal{D}$  is defined as

$$\langle \mathcal{D} \rangle_n := \sum_{k=1}^n \mathbb{E} \left[ \left( \mathcal{D}_k - \mathcal{D}_{k-1} \right)^2 \middle| \mathfrak{F}_{k-1} \right].$$

**Proposition 3.2.** *Let  $\langle \cdot \rangle_n$  be the quadratic variation of  $\mathcal{D}$ . Then we have*

$$\begin{aligned} \langle \mathcal{D} \rangle_n &= \sum_{k=1}^n \int_0^1 \int_0^1 \int_{k-1}^k K(e^u(x-y))(1 - e^{-\delta u}) du \left( 1 + \frac{\gamma}{2} (Y_{k-1} - \gamma \Omega_{k-1}) \right) \\ &\quad \times \left( 1 + \frac{\gamma}{2} (Y_{k-1}(y) - \gamma \Omega_{k-1}(y)) \right) e^{\frac{\gamma}{2}(Y_{k-1}(x) + Y_{k-1}(y)) - \frac{\gamma^2}{4} \mathbb{E}[Y_{k-1}^2(x)]} f(x)f(y) dx dy + \mathcal{E}_n, \end{aligned}$$

where the error term  $\mathcal{E}_n > 0$  for all  $n$  positive and  $\sup_n \mathcal{E}_n < \infty$ .

*Proof.* The proof of this proposition follows along the same lines as the proof of Proposition 3.1, with the conditional expectation instead. We begin with the following statement

$$\mathbb{E} \left[ \left( \mathcal{D}_k - \mathcal{D}_{k-1} \right)^2 \middle| \mathfrak{F}_{k-1} \right] = \mathbb{E} \left[ \mathcal{D}_k^2 \middle| \mathfrak{F}_{k-1} \right] - \mathcal{D}_{k-1}^2. \quad (3.11)$$

Let us now analyze the first term in the expression above in a similar manner to (3.10). We first compute the conditional expectation and then consider expansion around  $\gamma = 0$ :

$$\begin{aligned} &\mathbb{E} \left[ W_k^0(x) W_k^0(y) \middle| \mathfrak{F}_{k-1} \right] \\ &= \mathbb{E} \left[ \left( Y_{k-1}(x) - \gamma \Omega_{k-1}(x) + \Lambda_k(x) - \gamma (\Omega_k(x) - \Omega_{k-1}(x)) \right) \right. \\ &\quad \left. \left( Y_{k-1}(y) - \gamma \Omega_{k-1}(y) + \Lambda_k(y) - \gamma (\Omega_k(y) - \Omega_{k-1}(y)) \right) \right. \\ &\quad \left. \times e^{\frac{\gamma}{2} Y_{k-1}(x) - \frac{\gamma^2}{4} \Omega_{k-1}(x)} e^{\frac{\gamma}{2} Y_{k-1}(y) - \frac{\gamma^2}{4} \Omega_{k-1}(y)} e^{\frac{\gamma}{2} \Lambda_k(x) - \frac{\gamma^2}{4} (\Omega_k(x) - \Omega_{k-1}(x))} e^{\frac{\gamma}{2} \Lambda_k(y) - \frac{\gamma^2}{4} (\Omega_k(y) - \Omega_{k-1}(y))} \middle| \mathfrak{F}_{k-1} \right] \\ &= \left( Y_{k-1}(x) - \gamma \Omega_{k-1}(x) \right) \left( Y_{k-1}(y) - \gamma \Omega_{k-1}(y) \right) e^{\frac{\gamma}{2} Y_{k-1}(x) - \frac{\gamma^2}{4} \Omega_{k-1}(x)} e^{\frac{\gamma}{2} Y_{k-1}(y) - \frac{\gamma^2}{4} \Omega_{k-1}(y)} \\ &\quad \times e^{-\frac{\gamma^2}{4} (\Omega_k(x) + \Omega_k(y) - \Omega_{k-1}(x) - \Omega_{k-1}(y))} \mathbb{E} \left[ e^{\frac{\gamma}{2} (\Lambda_k(x) + \Lambda_k(y))} \middle| \mathfrak{F}_{k-1} \right] \\ &\quad + \left( Y_{k-1}(x) - \gamma \Omega_{k-1}(x) \right) e^{\frac{\gamma}{2} Y_{k-1}(x) - \frac{\gamma^2}{4} \Omega_{k-1}(x)} e^{\frac{\gamma}{2} Y_{k-1}(y) - \frac{\gamma^2}{4} \Omega_{k-1}(y)} e^{-\frac{\gamma^2}{4} (\Omega_k(x) + \Omega_k(y) - \Omega_{k-1}(x) - \Omega_{k-1}(y))} \\ &\quad \times \mathbb{E} \left[ \left( \Lambda_k(y) - \gamma (\Omega_k(y) - \Omega_{k-1}(y)) \right) e^{\frac{\gamma}{2} (\Lambda_k(x) + \Lambda_k(y))} \middle| \mathfrak{F}_{k-1} \right] \\ &\quad + \left( Y_{k-1}(y) - \gamma \Omega_{k-1}(y) \right) e^{\frac{\gamma}{2} Y_{k-1}(x) - \frac{\gamma^2}{4} \Omega_{k-1}(x)} e^{\frac{\gamma}{2} Y_{k-1}(y) - \frac{\gamma^2}{4} \Omega_{k-1}(y)} e^{-\frac{\gamma^2}{4} (\Omega_k(x) + \Omega_k(y) - \Omega_{k-1}(x) - \Omega_{k-1}(y))} \\ &\quad \times \mathbb{E} \left[ \left( \Lambda_k(x) - \gamma (\Omega_k(x) - \Omega_{k-1}(x)) \right) e^{\frac{\gamma}{2} (\Lambda_k(x) + \Lambda_k(y))} \middle| \mathfrak{F}_{k-1} \right] \\ &\quad + e^{\frac{\gamma}{2} Y_{k-1}(x) - \frac{\gamma^2}{4} \Omega_{k-1}(x)} e^{\frac{\gamma}{2} Y_{k-1}(y) - \frac{\gamma^2}{4} \Omega_{k-1}(y)} e^{-\frac{\gamma^2}{4} (\Omega_k(x) + \Omega_k(y) - \Omega_{k-1}(x) - \Omega_{k-1}(y))} \\ &\quad \times \mathbb{E} \left[ \left( \Lambda_k(x) - \gamma (\Omega_k(x) - \Omega_{k-1}(x)) \right) \left( \Lambda_k(y) - \gamma (\Omega_k(y) - \Omega_{k-1}(y)) \right) e^{\frac{\gamma}{2} (\Lambda_k(x) + \Lambda_k(y))} \middle| \mathfrak{F}_{k-1} \right] \\ &= \text{(I)} + \text{(II)} + \text{(III)} + \text{(IV)} \end{aligned} \quad (3.12)$$

We first provide simplified expression of **(I)**, **(II)**, **(III)** and **(IV)** which will be followed by explanations.

$$\begin{aligned} \text{(I)} &= W_{k-1}^0(x)W_{k-1}^0(y) \\ &+ \frac{\gamma^2}{4}(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] + O(\gamma))(Y_{k-1}(x) - \gamma\Omega_{k-1}(x))(Y_{k-1}(y) - \gamma\Omega_{k-1}(y)) \\ &\quad \times e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)}e^{\frac{\gamma}{2}Y_{k-1}(y) - \frac{\gamma^2}{4}\Omega_{k-1}(y)}, \end{aligned} \quad (3.13)$$

$$\text{(II)} = \frac{\gamma}{2}(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] + O(\gamma))(Y_{k-1}(x) - \gamma\Omega_{k-1}(x))e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)}e^{\frac{\gamma}{2}Y_{k-1}(y) - \frac{\gamma^2}{4}\Omega_{k-1}(y)}, \quad (3.14)$$

$$\text{(III)} = \frac{\gamma}{2}(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] + O(\gamma))(Y_{k-1}(y) - \gamma\Omega_{k-1}(y))e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)}e^{\frac{\gamma}{2}Y_{k-1}(y) - \frac{\gamma^2}{4}\Omega_{k-1}(y)}, \quad (3.15)$$

$$\text{(IV)} = \left(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] + \frac{\gamma^2}{2}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] + O(\gamma^3)\right)e^{\frac{\gamma}{2}Y_{k-1}(x) - \frac{\gamma^2}{4}\Omega_{k-1}(x)}e^{\frac{\gamma}{2}Y_{k-1}(y) - \frac{\gamma^2}{4}\Omega_{k-1}(y)}, \quad (3.16)$$

while ignoring the subleading terms. The expressions above are obtained by first noticing that

$$\mathbb{E}\left[e^{\frac{\gamma}{2}(\Lambda_k(x)+\Lambda_k(y))}\middle|\mathfrak{F}_{k-1}\right] = \mathbb{E}\left[e^{\frac{\gamma}{2}(\Lambda_k(x)+\Lambda_k(y))}\right] = e^{\frac{\gamma^2}{4}(\Omega_k(x)-\Omega_{k-1}(x))+\frac{\gamma^2}{4}(\Omega_k(y)-\Omega_{k-1}(y))}e^{\frac{\gamma}{4}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]}.$$

Taylor expanding the term  $e^{\frac{\gamma^2}{2}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]}$  w.r.t  $\gamma$  yields the equality (3.13). To see the second equality (3.14), we take the derivative w.r.t.  $\gamma$  of both sides of the following identity

$$e^{-\frac{\gamma^2}{4}(\Omega_k(x)-\Omega_{k-1}(x))-\frac{\gamma^2}{4}(\Omega_k(y)-\Omega_{k-1}(y))}\mathbb{E}\left[e^{\frac{\gamma}{2}(\Lambda_k(x)+\Lambda_k(y))}\right] = e^{\frac{\gamma^2}{4}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]},$$

and symmetrize w.r.t.  $x$  and  $y$ . As a result, we get the following equation

$$\mathbb{E}\left[(\Lambda_k(x) - \gamma(\Omega_k(x) - \Omega_{k-1}(x)))e^{\frac{\gamma}{2}(\Lambda_k(x)+\Lambda_k(y))}\right] = \frac{\gamma}{2}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]e^{\frac{\gamma^2}{4}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]}.$$

Plugging the above into the expression of **(II)** in (3.12) and Taylor expanding the term  $e^{\frac{\gamma^2}{4}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]}$  w.r.t.  $\gamma$ , we obtain the second identity (3.14). The third equality (3.15) follows by a similar argument. The fourth equality (3.16) is obtained by noticing that

$$\begin{aligned} &\mathbb{E}\left[(\Lambda_k(x) - \gamma(\Omega_k(x) - \Omega_{k-1}(x)))(\Lambda_k(y) - \gamma(\Omega_k(y) - \Omega_{k-1}(y)))e^{\frac{\gamma}{2}(\Lambda_k(x)+\Lambda_k(y))}\middle|\mathfrak{F}_{k-1}\right] \\ &= e^{\frac{\gamma^2}{4}(\Omega_k(x)+\Omega_k(y)-\Omega_{k-1}(x)-\Omega_{k-1}(y))}e^{\frac{\gamma^2}{4}\mathbb{E}[\Lambda_k(x)\Lambda_k(y)]}\left(\gamma^2(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)])^2 + \mathbb{E}[\Lambda_k(x)\Lambda_k(y)]\right), \end{aligned}$$

and

$$\mathbb{E}[\Lambda_k(x)\Lambda_k(y)] = \int_{k-1}^k K(e^u(x-y))(1 - e^{-\delta u})du.$$

Substituting the expressions of **(I)**, **(II)**, **(III)**, **(IV)** in (3.11) and rearranging the terms gives

$$\begin{aligned} &\mathbb{E}\left[\left(\mathcal{D}_k - \mathcal{D}_{k-1}\right)^2\middle|\mathfrak{F}_{k-1}\right] \\ &= \int_0^1 \int_0^1 \int_{k-1}^k K(e^u(x-y))(1 - e^{-\delta u})\left(1 + \frac{\gamma}{2}(Y_{k-1}(x) - \gamma\Omega_{k-1}(x))\right) \\ &\quad \times \left(1 + \frac{\gamma}{2}(Y_{k-1}(y) - \gamma\Omega_{k-1}(y))\right)e^{\frac{\gamma}{2}(Y_{k-1}(x)+Y_{k-1}(y))-\frac{\gamma^2}{2}(\Omega_{k-1}(x)+\Omega_{k-1}(y))}f(x)f(y)dudxdy \\ &\quad + \gamma^2\text{Err}_k, \end{aligned} \quad (3.17)$$

where  $\text{Err}_k$  is the residual sub-leading term. In fact,  $\text{Err}_k$  can also be written as integral w.r.t.  $x$  and  $y$  where the leading term of the integrands is  $(\mathbb{E}[\Lambda_k(x)\Lambda_k(y)])^2$ . Since the integrands of  $\text{Err}_k$  are absolutely summable, for  $\mathcal{E}_n := \sum_{k=1}^n \text{Err}_k$ , we have  $\sup_n \mathcal{E}_n < \infty$ . This completes the proof.  $\square$

Now we start the proof of Theorem 3.1.

*Proof of Theorem 3.1.* We see from the above expression (3.17) that most of the contribution to  $\langle \mathcal{D} \rangle_n$  is given by high values of  $Y_k(x)$ . However, these high values should not contribute much to the negative tail of  $\langle \mathcal{D} \rangle_n$  since they tend to yield high positive values of the integrand. Hence our idea is to compare  $\mathcal{D}_k$  with a martingale obtained by replacing  $Y_k(x)$  by an alternative martingale  $W_k(x)$  which does not sum the variation of  $\mathcal{D}_n$  when the value of  $Y_k(x)$  is too large. Our definition of  $\tilde{\mathcal{D}}_k$  has to be carefully chosen so that it compares well with  $\mathcal{D}_k$ .

To analyse the quadratic variation, we begin by defining the following quantities: the variance of  $Y_n$  for all  $n \in \mathbb{N}$ , and the sequence of stopping times

$$\begin{aligned} T_k^x &:= \inf_n \left\{ \Omega_n > R_{k-1}^x : Y_n(x) \geq (\gamma + \eta)\Omega_n + A \right\}, \\ R_k^x &:= \inf \left\{ \Omega_n > T_k^x : Y_n(x) \leq \gamma \Omega_n \right\}, \end{aligned} \quad (3.18)$$

where  $\Omega_n$  is the bound of  $\Omega_n(x)$ , and  $R_0^x = 0, A > \gamma + 2$ . By introducing the notation

$$\mathcal{R}_x = \cup_{k=1}^{\infty} [R_{k-1}^x, T_k^x], \quad : \gamma Y_k(x) := \frac{\gamma}{2} Y_k(x) - \frac{\gamma^2}{4} \Omega_k,$$

we can define

$$W_k(x) := \sum_{j=1}^k (W_j^0(x) - W_{j-1}^0(x)) \mathbf{1}_{\Omega(j) \in \mathcal{R}_x},$$

where  $W_j^0(x)$  is given by (3.7), and its integral is denoted by

$$\tilde{\mathcal{D}}_k := \int_0^1 W_k(x) f(x) dx. \quad (3.19)$$

The function  $W_k(x)$  has alternative representation in terms of stopping times

$$W_k(x) = \begin{cases} (Y_k(x) - \gamma \Omega_k) e^{:\gamma Y_k(x):} + \sum_{\ell=1}^n (A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x} + \xi_k; & \Omega_k \in [R_n^x, T_{n+1}^x], \\ \sum_{\ell=1}^n (A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x} + \xi_k; & \Omega_k \in [T_n^x, R_n^x], \end{cases} \quad (3.20)$$

where  $\xi_k$  is the sum of the residual between  $W_\ell$  and  $(A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x}$  for  $\ell = 1, \dots, n$ . The residual corresponding to the  $\ell$ -th difference is bounded by  $\max\{0, \Lambda_\ell - \eta T_\ell^x - A\} e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x}$ . As a result, we have

$$\xi_k \leq \sum_{\ell=1}^n \max\{0, \Lambda_\ell - \eta T_\ell^x - A\} e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x}.$$

Repeating the calculations leading to Proposition 3.2, we find that the quadratic variation of the function  $\tilde{\mathcal{D}}_k$  in (3.19) is

$$\begin{aligned} &\langle \tilde{\mathcal{D}} \rangle_{k'} \\ &= \sum_{k=1}^{k'} \int_0^1 \int_0^1 \int_{k-1}^k K(e^u(x-y))(1 - e^{-\delta u}) \left(1 + \frac{\gamma}{2} (Y_{k-1}(x) - \gamma \Omega_{k-1})\right) \left(1 + \frac{\gamma}{2} (Y_{k-1}(y) - \gamma \Omega_{k-1})\right) \\ &\quad \times \mathbf{1}_{k \in \mathcal{R}_x \cap \mathcal{R}_y} e^{:Y_{k-1}(x) + :Y_{k-1}(y):} \mathcal{G}_k(x) \mathcal{G}_k(y) du dx dy + \tilde{\mathcal{E}}_{k'}, \end{aligned} \quad (3.21)$$

for some  $\mathcal{G}_k(x)$  which can be computed starting from (3.19) and repeating the computations in the proof of Proposition 3.2. We also note from Proposition 3.3 that  $\tilde{\mathcal{D}}_k$  is a martingale which is bounded in  $L^2$ , and by the Doob's martingale convergence theorem, there exists an almost sure limit of  $\tilde{\mathcal{D}}_k$  we call  $\tilde{\mathcal{D}}_\infty$ .

Furthermore, we know that  $\mathcal{D}_k$  in (3.6) also converges almost surely due to Proposition 3.1, and let that limit be equal to  $\mathcal{D}_\infty$ . From the definitions (3.6), (3.19), and noting that the residual term  $\xi_n \geq 0$  for all  $n \in \mathbb{Z}$ , it then follows that

$$\mathcal{D}_\infty \geq \tilde{\mathcal{D}}_\infty - Q, \quad (3.22)$$

where

$$Q := \sum_{\ell=1}^{\infty} \int_0^1 (A + \eta T_{\ell}^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_{\ell}^x} f(x) dx. \quad (3.23)$$

Note that, the definitions (3.1), (3.6) imply that  $\mathcal{D}_{\infty}$  is proportional to  $\text{DM}_N f(\mathbf{i}\infty)$ .

In Propositions 3.3 and 3.4, we present sub-Gaussian tail bounds for  $\tilde{\mathcal{D}}_{\infty}$  and  $Q$ , respectively. By using these bounds in the inequality (3.22), we obtain the following result:

$$\mathbb{P}[(\text{DM}_{\gamma, N} f)(\mathbf{i}\infty) \leq -v] \leq c_3 e^{-c_4 v^2},$$

where the constant  $c_3 = \int_0^1 f(x) dx$  comes from the normalization in (3.1). Notice that

$$\begin{aligned} \mathbb{P}\left(\tilde{\mathcal{D}}_{\infty} - Q \leq -s\right) &\leq \mathbb{P}\left(\tilde{\mathcal{D}}_{\infty} \leq -\frac{s}{2}\right) + \mathbb{P}\left(Q \geq \frac{s}{2}\right) \leq e^{-\lambda \frac{s}{2}} \mathbb{E}[e^{\lambda \tilde{\mathcal{D}}_{\infty}}] + e^{-\lambda \frac{s}{2}} \mathbb{E}[e^{\lambda Q}] \leq e^{-\lambda s/2 + c\lambda^2} \\ &\leq e^{-\frac{s^2}{16c}}. \end{aligned}$$

The first inequality in the above display follows by union bound, The second inequality is obtained by applying Markov inequality. To obtain the third inequality, we bound  $\mathbb{E}[e^{\lambda \tilde{\mathcal{D}}_{\infty}}]$  by  $e^{c\lambda^2}$  using Proposition 3.3 and bound  $\mathbb{E}[e^{\lambda Q}]$  by  $e^{c\lambda^2}$  using Proposition 3.4. Here  $c$  is a positive constant independent of  $\lambda$ . The last inequality of the above display follows by substituting  $\lambda = \frac{s}{4c}$ .

If we repeat the above calculations with  $Y_N(x; q)$  instead, the main difference is that the constant  $c$  in the sub-Gaussian tails of Propositions 3.3 and 3.4 will depend on  $\tau$ . We denote this  $\tau$ -dependent constant as  $c(\tau)$ . Additionally, we define the corresponding normalization constant as  $c_1(\tau) = \int_0^1 f(x, \tau) dx$ . Consequently, we obtain the following result:

$$\mathbb{P}[(\text{DM}_{\gamma, N} f)(\tau) \leq -v] \leq c_1(\tau) e^{-c_2(\tau) v^2},$$

proving the theorem.  $\square$

The following reasoning leads to the proof of convergence. If a process  $M_k$  is a martingale and  $\mathbb{E}[e^{sM_k}] < \infty$  for any  $s \in \mathbb{R}$ , then  $e^{-sM_k - \frac{s^2}{2} \langle M_k \rangle_k}$  is also a martingale, with  $\langle \cdot \rangle_k$  representing the quadratic variation. Furthermore, if  $M_0 = 0$ , then

$$\mathbb{E}[e^{-sM_k - \frac{s^2}{2} \langle M_k \rangle}] = 1 \quad (3.24)$$

for all  $k$ , leading to the precise bound. We will therefore, aim to show that  $\text{DM}_{\gamma, N} f(\tau)$  is a martingale and obtain the bound through the quadratic variation.

The convergence of  $\mathcal{D}_{\infty}$  will be proved through the convergence of the expectation of the exponential of the terms  $\tilde{\mathcal{D}}_{\infty}$ ,  $Q$  in Propositions 3.3, 3.4 respectively.

**Proposition 3.3.** *The following statements hold true:*

- (1) *The quadratic variation  $\langle \tilde{\mathcal{D}} \rangle_{k'}$  in (3.21) is  $L^2$  bounded and  $\mathcal{D}_{k'}$  converges to  $\mathcal{D}_{\infty}$  almost surely.*
- (2) *For any  $s \in \mathbb{R}_{<0}$  and  $c > 0$ , we can establish the following upper bound on the expectation of the exponential of  $\tilde{\mathcal{D}}_{\infty}$ , which represents the limit of  $\tilde{\mathcal{D}}_k$  defined in (3.19):*

$$\mathbb{E}[e^{s\tilde{\mathcal{D}}_{\infty}}] \leq e^{\frac{cs^2}{2}}.$$

*Proof.*

(1): To prove the first part of this proposition, it suffices to show that  $\langle \tilde{\mathcal{D}} \rangle_{k'}$  is bounded and  $\mathbb{E}[e^{s\tilde{\mathcal{D}}_{\infty}}] < \infty$ . We show these as follows. To examine the bound on the quadratic variation (3.21), it is enough to focus on  $k \in [R_k, T_{k+1}]$ . By referring to (3.18), we can derive the following expression:

$$Y_{k-1} \leq A + (\gamma + \eta) \Omega_{k-1}, \quad (3.25)$$

and consequently,

$$\begin{aligned} \max_{(3.25)} \left(1 + \frac{\gamma}{2} (Y_{k-1}(x) - \Omega_{k-1})\right) e^{\frac{\gamma}{2} Y_{k-1} - \frac{\gamma^2}{4} \Omega_{k-1}} \\ = \left(1 + \frac{\gamma}{2} (A + \eta \Omega_{k-1})\right) e^{\frac{\gamma A}{2} + \frac{\gamma}{2} (\eta + \frac{\gamma}{2}) \Omega_{k-1}}. \end{aligned}$$

Substituting the above expression in (3.21) we get

$$\begin{aligned} \langle \tilde{\mathcal{D}} \rangle_{k'} \leq & \sum_{k=1}^{k'} \int \int \int_{k-1}^k K(e^u(x-y))(1-e^{-\delta u}) \left(1 + \frac{\gamma}{2}(A + \eta\Omega_{k-1})\right)^2 \\ & \times e^{\gamma A + \gamma(\eta + \frac{\gamma}{2})\Omega_{k-1}} \mathbf{1}_{k \in \mathcal{R}_x \cap \mathcal{R}_y} \mathcal{G}_k(x) \mathcal{G}_k(y) du dx dy, \end{aligned}$$

We now claim that  $\sup_{k'} \langle \tilde{\mathcal{D}} \rangle_{k'} < \infty$  for all small  $\gamma > 0$ . Note that

$$\Omega_k \stackrel{(3.9)}{=} \frac{k}{2} - \frac{1 - e^{-\delta k}}{2\delta} + \frac{1}{2} \mathbb{E}[\mathcal{R}^2(x)]. \quad (3.26)$$

As a result  $\Omega_k$  grows linearly with  $k$ . Substituting this bound into the integrand of the above inequality yields

$$(A + \eta\Omega_{k-1})^2 e^{\gamma A + \gamma(\eta + \frac{\gamma}{2})\Omega_{k-1}} \lesssim (\eta k)^2 e^{\gamma(\eta + \frac{\gamma}{2})k}.$$

Combining this bound with the inequality  $K(e^u(x-y)) \leq e^{-au}|x-y|^{-a}$  for some  $a \in (0, 1)$ , we get

$$\langle \tilde{\mathcal{D}} \rangle_{k'} \lesssim \int \int \sum_{k=1}^{k'} (\eta k)^2 e^{-(a-\gamma(\eta + \frac{\gamma}{2})k)} |x-y|^{-a} \mathbf{1}_{k \in \mathcal{R}_x \cap \mathcal{R}_y} \mathcal{G}_k(x) \mathcal{G}_k(y) du dx dy + \tilde{\mathcal{E}}_{k'}.$$

Note that  $k^2 e^{-ak} e^{\gamma(\eta + \frac{\gamma}{2})k}$  is absolutely summable for  $\gamma$  small and  $|x-y|^{-a}$  is integrable. Therefore,  $\langle \tilde{\mathcal{D}} \rangle_{k'}$  is bounded as  $k'$  tends to  $\infty$ .

(2): Now we proceed to show  $\mathbb{E}[e^{s\tilde{\mathcal{D}}_\infty}] < \infty$  for any  $s < 0$ . For all  $x$ , the function  $W_k(x) - \xi_k(x)$  in (3.20) has the following upper bound

$$W_k(x) - \xi_k(x) \geq -\left(c + |Y_k(x) - \gamma\Omega_k|\right) e^{A + \eta\Omega_k} + \sum_{m=1}^k (A + \eta T_m^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})T_m^x}.$$

Since  $Y_k(x)$  is a Gaussian distribution,  $\mathbb{E}[e^{s|Y_k(x)|}] < \infty$  for any  $s \in \mathbb{R}$ , and therefore,

$$\mathbb{E}\left[\exp\left(s\tilde{\mathcal{D}}_k - s \int_0^1 \xi_k(x) f(x) dx\right)\right] < \infty$$

for any  $s < 0$ . As a result,  $\mathbb{E}[e^{s\tilde{\mathcal{D}}_k}] < \infty$  for  $s < 0$  since  $\xi_k(x) > 0$  for all  $x \in [0, 1]$  and any integer  $k$ . Moreover,  $\tilde{\mathcal{D}}_k$  is a martingale and its quadratic variation is bounded. Since  $\mathbb{E}[e^{s\tilde{\mathcal{D}}_k}] < \infty$ , We get  $\mathbb{E}[e^{s\tilde{\mathcal{D}}_k}] = e^{s^2 \langle \tilde{\mathcal{D}} \rangle_k / 2}$  by Wald's identity. The identity (3.24) then suggests that,

$$\mathbb{E}[e^{s\tilde{\mathcal{D}}_k}] \leq e^{cs^2/2}.$$

And since the bound is independent of  $k$ , the above inequality hold for  $\tilde{\mathcal{D}}_\infty$ .  $\square$

The final ingredient needed for the proof of Theorem 3.1 is to bound the exponential expectation of  $Q$  in (3.22).

**Proposition 3.4.** *For any  $s \in \mathbb{R}$  and  $c > 0$ , the expectation of the exponential of  $Q$  defined in (3.23) has the following upper bound*

$$\mathbb{E}[e^{sQ}] \leq e^{\frac{cs^2}{2}}. \quad (3.27)$$

*Proof.* We prove this by using tools from martingale theory. Recall the definition of  $Q$  from (3.23). Recall that sigma algebras  $\mathfrak{F}_1 \subset \mathfrak{F}_2 \subset \mathfrak{F}_3 \subset \dots$  such that  $\mathcal{L}_k$  is measurable with respect to  $\mathfrak{F}_k$

$$Q_k = \mathbb{E}[Q | \mathfrak{F}_k]. \quad (3.28)$$

We now aim to prove the convergence of  $Q_k$ , for which it is sufficient to prove the boundedness of the integrand of  $Q$  in (3.23).

Consider the variable

$$Q^x := \sum_{\ell=1}^{\infty} (A + \eta T_{\ell}^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_{\ell}^x}. \quad (3.29)$$

For  $T_{\ell}^x \in \mathbb{Z}$ , the above expression satisfies the inequality

$$Q^x \leq \sum_{m=1}^{\infty} \sum_{\ell=1}^{\infty} (A + \eta T_{\ell}^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_{\ell}^x} \mathbf{1}(T_{\ell}^x \in [m-1, m]) =: \sum_{m=1}^{\infty} Q^{x,m}. \quad (3.30)$$

The problem at hand then simplifies to proving the convergence of  $Q^{x,m}$ . To compute its bound, we begin by noting that

$$\begin{aligned} \mathbb{E}[Q^{x,m}] &= (A + \eta m) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) m} \sum_{\ell=1}^{\infty} \mathbb{P}\left(\left\{T_{\ell}^x \in [m-1, m]\right\}\right) \\ &= (A + \eta m) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) m} \mathbb{E}\left[\#\left\{1 \leq \ell \leq \infty : T_{\ell}^x \in [m-1, m]\right\}\right]. \end{aligned}$$

The remainder of the proof can be divided into several steps, with each step involving the demonstration of a lemma. The inequality (3.31) in Lemma 3.1 implies that

$$\mathbb{E}[Q^{x,m}] \stackrel{(3.31)}{\leq} (A + \eta m) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) m} \frac{c}{\sqrt{m-1}} e^{-\frac{(\gamma+\eta)^2(m-1)}{2}},$$

and therefore due to (3.30),

$$\begin{aligned} \mathbb{E}[Q^x] &\stackrel{(3.30)}{\leq} \sum_{m=1}^{\infty} (A + \eta m) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) m} \frac{c}{\max\{\sqrt{(m-1)}, 1\}} e^{-\frac{(\gamma+\eta)^2(m-1)}{2}} \\ &\lesssim \sum_{m=1}^{\infty} \frac{1}{\max\{\sqrt{m-1}, 1\}} (A + \eta m) e^{-\frac{\eta^2 m}{2}}. \end{aligned}$$

Having the above expression available, the only remaining task to obtain (3.27) is to evaluate the quadratic variation of  $Q^x$ , which can be obtained through Lemma 3.2. The quadratic variation of  $Q_k$  is bounded due to Lemma 3.2 proving this proposition.  $\square$

**Lemma 3.1.** *For an arbitrary constant  $c$ , the following inequality holds for stopping time  $T_{\ell}^x$  defined in (3.18)*

$$\mathbb{E}\left[\#\left\{1 \leq \ell \leq \infty : T_{\ell}^x \in [m-1, m]\right\}\right] \leq \frac{c}{\max\{\sqrt{(m-1)}, 1\}} e^{-\frac{(\gamma+\eta)^2(m-1)}{2}}. \quad (3.31)$$

*Proof.* The definition of the stopping times (3.18) implies that, for  $T_x^{\ell} \in [m-1, m]$ ,

$$\Omega_n \in [m-1, m], \quad Y_n(x) \geq (\gamma + \eta)\Omega_n + A.$$

Since  $\Omega_n (= \int_0^n K(0)(1 - e^{-\delta u}) du + c_2)$  (see (3.26)) grows linearly in  $n$ , we have  $c_1(m-1) \leq n \leq c_1 m$  for some constant  $c_1 > 0$  because  $\Omega_n \in [m-1, m]$ . Under the assumption  $A > \gamma + \eta$  and recognizing that  $Y_{\ell}(x)$  is the sum of independent Gaussian random variable  $\sum_{t=1}^{\ell} (\mathcal{L}_t - \mathcal{L}_{t-1}) + \mathcal{R}(x)$ , we can derive the following expression utilizing the maximal Bernstein's inequality:

$$\begin{aligned} \mathbb{P}\left(\left\{\exists \ell \geq 1 : T_{\ell}^x \in [m-1, m]\right\}\right) &\leq \mathbb{P}\left(\sup_{1 \leq \ell \leq c_1(m-1)} Y_{\ell}(x) \geq (\gamma + \eta)\Omega_{\ell} + A\right) \\ &\leq \frac{c}{\max\{\sqrt{\Omega_{c_1(m-1)}}, 1\}} \exp\left(-(\gamma + \eta)^2 \Omega_{c_1(m-1)}\right). \quad (3.32) \end{aligned}$$

The inequality (3.31) is then obtained as follows

$$\begin{aligned}
 \mathbb{E}\left[\#\left\{1 \leq \ell \leq \infty : T_\ell^x \in [m-1, m]\right\}\right] &= \sum_{k=1}^{\infty} \mathbb{P}\left(\#\{\ell : T_\ell^x \in [m-1, m]\} \geq k\right) \\
 &\stackrel{(3.33)}{\leq} \sum_{k=1}^{\infty} \mathbb{P}\left(\#\{\ell : T_\ell^x \in [m-1, m]\} \geq 1\right) \frac{1}{2^k} \\
 &\stackrel{(3.32)}{\leq} \frac{c}{\max\{\sqrt{\Omega_{c_1(m-1)}}, 1\}} \exp\left(-(\gamma + \eta)^2 \Omega_{c_1(m-1)}\right) \\
 &\stackrel{(3.26)}{\leq} \frac{c}{\max\{\sqrt{(m-1)}, 1\}} \exp\left(-(\gamma + \eta)^2 \frac{(m-1)}{2}\right).
 \end{aligned}$$

The following inequality was used to obtain the result above

$$\mathbb{P}\left(\#\{\ell : T_\ell^x \in [m-1, m]\} \geq k\right) \leq 2^{-k} \mathbb{P}\left(\exists \ell \geq 1 : T_\ell^x \in [m-1, m]\right). \quad (3.33)$$

The proof of the above expression is as follows.

We begin by noting that the inequality (3.33) can be rewritten as

$$\mathbb{P}\left(\#\{\ell : T_\ell^x \in [m-1, m]\} \geq k \mid \#\{\ell : T_\ell^x \in [m-1, m]\} \geq k-1\right) \leq \frac{1}{2}.$$

The increments of  $Y_\ell(x)$  are independent Gaussian random variables, i.e.,

$$Y_{\ell+1}(x) - Y_\ell(x) \sim N\left(0, \int_\ell^{\ell+1} (1 - e^{-\delta u}) du\right).$$

Consequently,  $Y_\ell(x)$  is regarded as the evaluation of a Brownian motion  $B_t$  at time  $\int_0^\ell (1 - e^{-\delta u}) du$  and  $B_0 := \mathcal{R}(x)$ . Recall the definition of  $T_\ell^x$  in (3.18). The fact that the increments of Brownian motion  $B_t$  are stationary then implies that, for  $A \geq \gamma + 2$ ,

$$\mathbb{P}\left(\#\{\ell : T_\ell^x \in [m-1, m]\} \geq k \mid \#\{\ell : T_\ell^x \in [m-1, m]\} \geq k-1\right) \leq \mathbb{P}\left(\inf_{t \in [0, 1]} B_t \leq \gamma - A\right) \leq \frac{1}{2}.$$

This completes the proof of the lemma. □

**Lemma 3.2.** *The quadratic variation of  $Q_k$  defined in (3.28) can be expressed as*

$$\langle Q_k \rangle_k = \sum_{\ell=1}^k \int_0^1 \int_0^1 \mathbb{E}\left[\mathfrak{A}_\ell^x \mathfrak{A}_\ell^y \mid \mathfrak{F}_\ell\right] f(x) f(y) dx dy, \quad (3.34)$$

for some random function  $\mathfrak{A}_\ell^x$  (defined in (3.42)) such that

$$|\mathfrak{A}_\ell^x| \leq C \left(1 + \Omega_\ell\right) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_\ell}. \quad (3.35)$$

The constant  $C > 0$  is independent of  $x \in [0, 1]$  and the integer  $\ell$ .

*Proof.* Recall that  $Q_k$  in (3.28) is a martingale. Taking into account the definition of the integrand of  $Q$  given by (3.29):

$$Q^x = \sum_{\ell=1}^{\infty} (A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})T_\ell^x},$$

we define the martingale

$$Q_\ell^x := \mathbb{E}[Q^x \mid \mathfrak{F}_\ell], \quad \ell \in \mathbb{N}.$$

With these notations, the quadratic variation of  $Q_k$  is given by

$$\begin{aligned} \langle Q_k \rangle_k &= \sum_{\ell=1}^k \mathbb{E} \left[ (Q_{\ell+1} - Q_\ell)^2 \middle| \mathfrak{F}_\ell \right] \\ &= \sum_{\ell=1}^k \int \int \mathbb{E} \left[ (Q_{\ell+1}^x - Q_\ell^x)(Q_{\ell+1}^y - Q_\ell^y) \middle| \mathfrak{F}_\ell \right] f(x)f(y) dx dy. \end{aligned} \quad (3.36)$$

In what follows, we rewrite  $Q_{\ell+1}^x - Q_\ell^x$  in terms a new set of stopping time and use that expression to find almost sure bound on  $Q_{\ell+1}^x - Q_\ell^x$ .

Let  $k'' < k'$  represent the number of distinct values of  $T_\ell^x$  between 0 and  $\Omega_{k'}$ . We divide the discussion into two cases:

- (1) The first case arises when  $\Omega_{k'} \in [T_{k''}^x, R_{k''}^x]$ .
- (2) The second cases occurs when  $\Omega_{k'} \in [R_{k''}^x, T_{k''+1}^x]$ .

In the first case, we write

$$Q_{k'}^x = \sum_{\ell=1}^{k''} (A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x} + \mathbb{E} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widehat{T}_\ell^x < +\infty\}} (A + \eta \widehat{T}_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) (\widehat{T}_\ell^x + \Omega_{k'})} \right],$$

where the set of stopping times  $\widehat{T}_\ell^x$  are defined in the following way. Starting with  $\widehat{T}_0^x := 0$ , for  $\ell \geq 1$ ,

$$\widehat{R}_\ell^x := \inf \left\{ \Omega_{s+k'} \geq \widehat{T}_{\ell-1}^x : H_{s+k'} \leq \eta \Omega_\ell \right\}, \quad (3.37)$$

$$\widehat{T}_\ell^x := \inf \left\{ \Omega_{s+k'} \geq \widehat{R}_\ell^x : H_{s+k'} \geq (\eta + \gamma) \Omega_\ell + A \right\}, \quad (3.38)$$

where

$$H_s(x) := Y_{k'}(x) + \sum_{n=k'+1}^s (\mathcal{L}_n(x) - \mathcal{L}_{n-1}(x)), \quad s > k'. \quad (3.39)$$

Similarly, in the second case, we can express the value of  $Q_{k'}^x$  as

$$Q_{k'}^x = \sum_{\ell=1}^{k''} (A + \eta T_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_\ell^x} + \mathbb{E} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widetilde{T}_\ell^x < +\infty\}} (A + \eta \widetilde{T}_\ell^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) (\widetilde{T}_\ell^x + \Omega_{k'})} \right].$$

Here we define  $\widetilde{R}_0^x := 0$ , and for  $\ell \geq 1$ , the stopping times are defined as

$$\widetilde{T}_\ell^x := \inf \left\{ \Omega_{s+k'} \geq \widetilde{R}_{\ell-1}^x : H_{s+k'} \geq (\eta + \gamma) \Omega_\ell + A \right\}, \quad (3.40)$$

$$\widetilde{R}_\ell^x := \inf \left\{ \Omega_{s+k'} \geq \widetilde{T}_\ell^x : H_{s+k'} \leq \gamma \Omega_\ell \right\}, \quad (3.41)$$

with the definition (3.39) of  $H_s$ .

With the above notation, the integrand of the quadratic variation (3.36) has the expression

$$\begin{aligned} (Q_{k'+1}^x - Q_{k'}^x) &=: \mathfrak{A}_{k'}^x \\ &= (A + \eta T_{k''+1}^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) T_{k''+1}^x} \mathbf{1}_{(k''_{k'+1} \neq k''_{k'})} \\ &\quad + \mathbb{E}_{Y_{k'+1}} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widehat{T}_\ell^x < +\infty\}} (A + \eta \widehat{T}_\ell^x + \eta \Omega_{k'+1}) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) (\widehat{T}_\ell^x + \Omega_{k'+1})} \right] \\ &\quad - \mathbb{E}_{Y_{k'}} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widehat{T}_\ell^x < +\infty\}} (A + \eta \widehat{T}_\ell^x + \eta \Omega_{k'}) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2}) (\widehat{T}_\ell^x + \Omega_{k'})} \right], \end{aligned} \quad (3.42)$$

and substituting the above in (3.36), we obtain (3.34). We now proceed to complete showing (3.35). We seek to show uniform bound on  $\mathfrak{A}_{k'}^x$ . To this end, we introduce the following two

events:

$$\mathbf{1}_{\text{Good}}^{(1)} := \mathbf{1}\left(Y_{k'}(x) \geq \gamma \Omega_{k'}\right), \quad \mathbf{1}_{\text{Good}}^{(2)} := \mathbf{1}\left(Y_{k'}(x) \leq (\gamma + \eta)\Omega_{k'}\right). \quad (3.43)$$

We now complete proof of (3.35) using the following two claims whose proofs are deferred to the end of this section. The Claims 3.1, 3.2 therefore imply that  $\mathfrak{A}_{k'}^x$  is uniformly bounded.  $\square$

**Claim 3.1.** *There exists  $C_1 = C_1(A, \gamma, \eta) > 0$  such that*

$$\left|\mathfrak{A}_{k'}^x \mathbf{1}_{\text{Good}}^{(1)}\right| \leq C_1 \left(1 + \Omega_{k'}\right) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}.$$

*Proof.* Let us rewrite (3.42) as

$$\mathfrak{A}_{k'}^x = I_1 + I_2 - I_3$$

where

$$I_1 := (A + \eta T_{k''_{k'+1}}^x) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})T_{k''_{k'+1}}^x} \mathbf{1}(k''_{k'+1} \neq k''_{k'}), \quad (3.44)$$

$$I_2 := \mathbb{E}_{Y_{k'+1}} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widehat{T}_\ell^x(k'+1) < +\infty\}} \left( A + \eta \widehat{T}_\ell^x + \eta \Omega_{k'+1} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widehat{T}_\ell^x + \Omega_{k'+1})} \right], \quad (3.45)$$

$$I_3 := \mathbb{E}_{Y_{k'}} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}_{\{\widehat{T}_\ell^x(k') < +\infty\}} \left( A + \eta \widehat{T}_\ell^x + \eta \Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widehat{T}_\ell^x + \Omega_{k'})} \right]. \quad (3.46)$$

Our aim in the following analysis is to illustrate that

$$|I_1| \mathbf{1}_{\text{Good}}^{(1)} \leq C(1 + \Omega_{k'}) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}, \quad |I_2 - I_3| \mathbf{1}_{\text{Good}}^{(1)} \leq C(1 + \Omega_{k'}) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}. \quad (3.47)$$

To prove the first inequality in (3.47), we begin by recalling the expressions (3.43), (3.44). When the conditions

$$k''_{k'+1} \neq k''_{k'}, \quad \mathbf{1}_{\text{Good}}^{(1)} = \mathbf{1}\left(Y_{k'}(x) \geq \gamma \Omega_{k'}\right),$$

hold simultaneously, the definitions (3.37), (3.38) imply that

$$\Omega_{k'+1} = R_{k''}^x, \quad \Omega_{k'+1} \geq T_{k''_{k'+1}}^x.$$

Resultantly,

$$T_{k''_{k'+1}}^x \leq \Omega_{k'+1} \leq (1 + o(1)) \Omega_{k'}. \quad (3.48)$$

Substituting the above bound in the expression for  $I_1$  (3.44), we obtain the first inequality in (3.47).

Continuing with the proof of the second inequality, we divide the discussion into two cases. First, we consider the case when  $\Omega_{k'}, \Omega_{k'+1}$  lie in the same interval,  $\Omega_{k'}, \Omega_{k'+1} \in [T_{k''}^x, R_{k''}^x]$ . In this scenario, from (3.37), (3.38), observe that

$$\{\widehat{T}_\ell^x(k'), \widehat{R}_\ell^x(k')\} = \left\{ \widehat{T}_\ell^x(k'+1) + (\Omega_{k'+1} - \Omega_{k'}), \widehat{R}_\ell^x(k'+1) + (\Omega_{k'+1} - \Omega_{k'}) \right\},$$

and as a result  $I_2, I_3$  in (3.45), (3.46) coincide with each other. The second case is when  $\Omega_{k'}, \Omega_{k'+1}$  belong to subsequent intervals,

$$\Omega_{k'} \in [T_{k''}^x, R_{k''}^x], \quad \Omega_{k'+1} \in [R_{k''}^x, T_{k''_{k'+1}}^x].$$

In this case, we observe from the expressions for  $I_2, I_3$  in (3.45), (3.46) that

$$I_2 - I_3 = -\mathbb{E}_{Y_{k'}} \left[ \mathbf{1}_{\{\widehat{T}_1^x < +\infty\}} \left( A + \eta \widehat{T}_1^x + \eta \Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widehat{T}_1^x + \Omega_{k'})} \right]. \quad (3.49)$$

We may further rewrite the right hand side of the above equation as

$$\text{r.h.s. of (3.49)} = -\mathbb{E}_{Y_{k'}} \left[ \sum_{\ell=1}^{\infty} \mathbf{1}(\widehat{T}_1^x \in [\ell-1, \ell]) \left( A + \eta \widehat{T}_1^x + \eta \Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widehat{T}_1^x + \Omega_{k'})} \right].$$

Starting from the expression above, we obtain the following inequality

$$\begin{aligned} |I_2 - I_3| &\leq \sum_{\ell=1}^{\infty} (A + \eta\ell + \eta\Omega_{k'}) \mathbb{E} \left[ \mathbf{1}(\widehat{T}_1^x \in [\ell-1, \ell]) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widehat{T}_1^x + \Omega_{k'})} \right] \\ &\leq \sum_{\ell=1}^{\infty} (A + \eta\ell + \eta\Omega_{k'}) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\ell + \Omega_{k'})} \mathbb{E} \left[ \#(\widehat{T}_1^x \in [\ell-1, \ell]) \right]. \end{aligned} \quad (3.50)$$

By employing a similar argument as in Lemma 3.1, we arrive at the following result:

$$\mathbb{E} \left[ \#(\widehat{T}_1^x \in [\ell-1, \ell]) \right] \leq \frac{C}{\max\{\sqrt{\ell-1}, 1\}} e^{-\frac{(\gamma+\eta)^2}{2}(\ell-1)}.$$

Finally, plugging the above inequality into the right hand side (3.50) shows that

$$\begin{aligned} |I_2 - I_3| &\leq \sum_{\ell=1}^{\infty} (A + \eta\ell + \eta\Omega_{k'}) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\ell + \Omega_{k'})} \frac{C}{\max\{\sqrt{\ell-1}, 1\}} e^{-\frac{(\gamma+\eta)^2}{2}(\ell-1)} \\ &\leq C \left( 1 + \eta\Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}, \end{aligned} \quad (3.51)$$

which implies the second inequality in (3.47). This completes the proof of Claim 3.1.  $\square$

**Claim 3.2.** *There exists  $C_2 = C_2(A, \gamma, \eta) > 0$  such that*

$$\left| \mathfrak{A}_{k'}^x \mathbf{1}_{\text{Good}}^{(2)} \right| \leq C_2 \left( 1 + \Omega_{k'} \right) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}.$$

*Proof of Claim 3.2.* Let us recall the notations of  $I_1, I_2, I_3$  from the proof of Claim 3.1. Similar to the previous result, we demonstrate that

$$|I_1| \mathbf{1}_{\text{Good}}^{(2)} \leq C(1 + \Omega_{k'}) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}, \quad |I_2 - I_3| \mathbf{1}_{\text{Good}}^{(2)} \leq C \left( 1 + \Omega_{k'} \right) e^{(\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}. \quad (3.52)$$

The first inequality can be obtained by noting that the conditions  $k''_{k'+1} \neq k''_{k'}$  and  $\mathbf{1}_{\text{Good}}^{(2)} = 1$ , imply that  $\Omega_{k'+1} = T_{k''_{k'+1}}^x$ . Consequently, following a similar approach to (3.48), we have

$$T_{k''_{k'+1}}^x \leq (1 + o(1))\Omega_{k'}$$

which yields the first inequality in the above display by a direct substitution of the above inequality in  $I_1$  (3.44).

To prove the second inequality, we repeat the arguments used in the proof of Claim 3.1. Once again we consider the following two cases. In the first case,  $\Omega_{k'}, \Omega_{k'+1} \in \left[ R_{k''_{k'}}^x, T_{k''_{k'+1}}^x \right]$ , and from (3.41), (3.40), we have

$$\left\{ \widetilde{T}_\ell^x(k'), \widetilde{R}_\ell^x(k') \right\} = \left\{ \widetilde{T}_\ell^x(k'+1) + (\Omega_{k'+1} - \Omega_{k'}), \widetilde{R}_\ell^x(k'+1) + (\Omega_{k'+1} - \Omega_{k'}) \right\},$$

which shows that  $I_2 - I_3 = 0$ . In the second case where

$$\Omega_{k'} \in \left[ R_{k''_{k'}}^x, T_{k''_{k'+1}}^x \right], \quad \text{and} \quad \Omega_{k'} \in \left[ T_{k''_{k'+1}}^x, R_{k''_{k'+1}}^x \right],$$

we obtain the expression below from (3.45), (3.46):

$$I_2 - I_3 = -\mathbb{E}_{Y_{k'}} \left[ \mathbf{1}_{\{\widetilde{T}_1^x < +\infty\}} \left( A + \eta \widetilde{T}_1^x + \eta \Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})(\widetilde{T}_1^x + \Omega_{k'})} \right].$$

Now, following the similar argument as in (3.51), we arrive at

$$|I_2 - I_3| \leq C \left( 1 + \eta\Omega_{k'} \right) e^{\gamma A + (\gamma\eta + \frac{\gamma^2}{2})\Omega_{k'}}.$$

Combining the bounds on  $|I_2 - I_3|$  in the above two cases yields the second inequality in (3.52). This completes the proof of Claim 3.2.  $\square$

## 4. SEMI-CLASSICAL LIMITS FOR DEFORMED CONFORMAL BLOCKS

In this section, we carry out the analysis for the semi-classical limit of deformed conformal blocks (2.3) in two cases, for  $\chi = \frac{2}{\gamma}$  and  $\chi = \frac{2}{\gamma}$ . It is notable that the structure of the conformal blocks in each of these cases is different, and in the limit  $\gamma \rightarrow 0$ , the BPZ equation satisfied by the deformed conformal block reduces to the following equations:

- For  $\chi = \frac{2}{\gamma}$ , it reduces to the Hamilton-Jacobi equation of the non-autonomous elliptic Calogero-Moser model, (see Theorem 4.1).
- For  $\chi = \frac{2}{\gamma}$ , it reduces to the Lamé equation, (see Theorem 4.2).

As a result of these two reductions, the *accessory parameter* of the Lamé equation [BGG22] and the action from the Hamilton-Jacobi equation can be related by using the semi-classical analogue of the connection (see (2.16)) between the undeformed conformal block and deformed conformal blocks for different values of  $\chi$ .

The above results, while interesting on their own, are vital to prove the existence of the semi-classical limit of the undeformed conformal block, which we study in Section 5. Furthermore, the relation between the semi-classical limits of the deformed and undeformed conformal blocks as revealed in Theorem 4.2 and relation (4.38) lets us prove the structure of the conformal block (1.1) in Theorem 1.1.

Before proceeding to the discussion of main results in this section, we introduce few notations which will be used throughout the rest of the paper. In the remainder of the paper when we write  $\alpha$  and  $P$  we mean  $\frac{\alpha_0}{\gamma}$  and  $\frac{P_0}{\gamma}$ . We also define the DGMC measures (3.1)

$$\text{DM}_{\gamma,N}\Theta(\tau) := \text{DM}_{\gamma,N}\theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x}, \quad \text{DM}_{\gamma,N}\Theta(\mathbf{i}\infty) = \text{DM}_{\gamma,N}(\sin(\pi x))^{-\frac{\alpha_0}{2}} e^{\pi P_0 x}, \quad (4.1)$$

that arise in the semi-classical limit of the undeformed conformal block (2.9), and the measure

$$\text{DM}_{\gamma,N}\tilde{\Theta}(z, \tau; \chi) := \text{DM}_{\gamma,N} \frac{\theta_1(z+x)}{\theta_1(z)} \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x}, \quad (4.2)$$

will be useful in obtaining the semi-classical limit of the deformed conformal block (2.3) for  $\chi = \frac{2}{\gamma}$ . We recall the notation (3.2) from the previous section:

$$\text{DM}_\bullet = \lim_{N \rightarrow \infty} \text{DM}_{\bullet,N}.$$

which means  $\text{DM}_\gamma\Theta(\tau)$ ,  $\text{DM}_\gamma\Theta(\mathbf{i}\infty)$  and  $\text{DM}_\gamma\tilde{\Theta}(z, \tau; \chi)$  are the large  $N$  limits of  $\text{DM}_{\gamma,N}\Theta(\tau)$ ,  $\text{DM}_{\gamma,N}\Theta(\mathbf{i}\infty)$  and  $\text{DM}_{\gamma,N}\tilde{\Theta}(z, \tau; \chi)$ .

**4.1. Semi-classical limit of the deformed conformal blocks for  $\chi = \frac{2}{\gamma}$  and the Hamilton-Jacobi equation.** In this section, we study the semi-classical limit of the deformed block  $\psi_\chi^\alpha(z, q)$  when the deformation parameter  $\chi$  is equal to  $\frac{2}{\gamma}$ . We show that in this case, the semi-classical limit is given by the solution of a Hamilton-Jacobi equation with the potential given by the Weierstrass  $\wp$ -function as described in physics literature [Lit+14; BGG22]. Our proof heavily relies on the BPZ equation satisfied by the  $\psi_\chi^\alpha(z, q)$  and the tail probabilities of the derivative GMC which we found in Section 3. In what follows, we first state the main result (Theorem 4.1) of this subsection. Before proceeding to the proof, we state and prove two main tools in Proposition 4.1 and Proposition 4.2. These two propositions claim and prove the boundedness and the tightness of the sequence of the deformed conformal block  $\psi_{2/\gamma}^\alpha(z, q)$  as  $\gamma \rightarrow 0$ . The proof of Theorem 4.1 will be detailed in Section 4.1.1.

**Theorem 4.1.** *For  $\chi = \frac{2}{\gamma}$ ,  $\alpha_0 \in (-2, 2)$ ,  $P_0 \in \mathbb{R}$ ,  $q \in (0, 1)$ , the semi-classical limit of deformed conformal blocks exists and assumes the form*

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q) = \tilde{\phi}(z, q), \quad (4.3)$$

where

$$\begin{aligned} \tilde{\phi}(z, q) := & \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12} \log(-2\pi) + \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12} \log q + 2P_0 z \\ & - (\alpha_0 - 2) \left( \log \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) \\ & + \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - 2}{\gamma} \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma} \right) \right) \right]. \end{aligned} \quad (4.4)$$

Furthermore,  $\tilde{\phi}(z, q)$  satisfies the following the partial differential equation which arises as the semi-classical limit of the BPZ equation (2.17):

$$(\partial_z \tilde{\phi}(z, \tau))^2 - (2 - \alpha_0)^2 \wp(z) + 2\pi i \partial_\tau \tilde{\phi}(z, \tau) = 0, \quad (4.5)$$

which is the Hamilton-Jacobi equation for the non-autonomous Calogero Moser model (see below) with  $\tilde{\phi}(z, \tau)$  describing the action.

**Remark 4.1.** The non-autonomous elliptic Calogero-Moser model is defined by the equations of motion [Man98]

$$v(\tau) = 2\pi i \frac{du(\tau)}{d\tau}, \quad 2\pi i \frac{dv(\tau)}{d\tau} = m^2 \wp'(2u(\tau)), \quad (4.6)$$

of the Hamiltonian

$$H(\tau) = v(\tau)^2 - m^2 \wp(2u(\tau)|\tau). \quad (4.7)$$

The solution  $u(\tau)$  depends on  $\tau$ ,  $m$ , and the initial conditions or equivalently, on the monodromy data through the Riemann-Hilbert correspondence [Bon+20; DDG23]. The equation (4.5) is then the Hamilton-Jacobi equation of the system above with the identification

$$z = 2u(\tau), \quad m^2 = (2 - \alpha_0)^2.$$

Given initial condition at  $\tau \rightarrow i\infty$  the solution of the Hamilton-Jacobi equation is unique and coincides with the classical action [Arn13] corresponding to the Hamiltonian above (see [BGG22], [Bus80, p. 258])

$$\tilde{\phi}(2u(\tau), \tau) = \int_{i\infty}^{\tau} (v(\tau)^2 + m^2 \wp(2u(\tau)|\tau)) \frac{d\tau}{2\pi i}. \quad (4.8)$$

**Proposition 4.1.** For  $\chi = \frac{2}{\gamma}$ ,  $q \in (0, 1)$  and with the parameters  $\alpha_0 \in (-2, 2)$ ,  $P_0 \in \mathbb{R}$ , the sequence of the semi-classical limit of the deformed conformal block (2.14) is bounded

$$\limsup_{\gamma \rightarrow 0} \gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z; q) < \infty, \quad \liminf_{\gamma \rightarrow 0} \gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z; q) > -\infty.$$

*Proof.* Recall the expression of the deformed conformal block (4.26) reads:

$$\psi_\chi^{\alpha_0/\gamma}(z, \tau) = \mathcal{W}(q) e^{\chi P_0 z \pi / \gamma} \mathbb{E} \left[ \mathcal{V}_{\chi, P}^{\alpha_0/\gamma}(z, q)^{-\frac{\alpha_0}{\gamma^2} + \frac{\chi}{\gamma}} \right], \quad (4.9)$$

where

$$\mathcal{V}_{\chi, P}^{\alpha_0/\gamma}(z, q) := \int_0^1 \theta_1(z+x)^{\frac{\gamma\chi}{2}} \theta_1(z)^{-\frac{\gamma\chi}{2}} \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx.$$

We analyse each term of the above expression. The parameter  $l_\chi$  for  $\chi = 2/\gamma$  simplifies as

$$l_\chi = \frac{\chi^2}{2} - \frac{\alpha\chi}{2} = \frac{(2 - \alpha_0)}{\gamma^2}.$$

The pre-factor of (4.9) defined in (2.15) then assumes the form

$$\mathcal{W}(q) e^{\chi P z \pi} = (-2\pi)^{\frac{1}{24}(2-\alpha_0) + \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12\gamma^2}} q^{\frac{1}{24}(2-\alpha_0) + \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12\gamma^2}} e^{\frac{2P_0 z \pi}{\gamma^2}} + \mathcal{O}(q^{\gamma^2}). \quad (4.10)$$

Note that the above expression consists of terms that are dependent on both  $z$  and  $\gamma$ .

We now consider the term  $\mathcal{V}_{\chi,P}^\alpha(z, q)$  and focus on obtaining uniform bounds of the expression  $\gamma^2 \log \mathbb{E} \left[ \mathcal{V}_{\chi,P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2} \right]$  as functions of  $z$  and  $q$  as they vary on a compact set. To this end, we note that

$$\begin{aligned} \mathbb{E} \left[ \mathcal{V}_{\chi,P}^\alpha(z, q)^{\frac{(2-\alpha_0)}{\gamma^2}} \right] &= \left( \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\left(\frac{\alpha_0-2}{\gamma^2}\right)} \\ &\quad \times \mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma^2} \left( \log \mathcal{V}_{\chi,P}^\alpha(z, q) - \log \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) \right) \right]. \end{aligned} \quad (4.11)$$

The last term in the expression above can be simplified as

$$\begin{aligned} &\mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma^2} \left( \log \mathcal{V}_{\chi,P}^\alpha(z, q) - \log \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) \right) \right] \\ &\stackrel{(4.2)}{=} \mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma^2} \int_0^\gamma \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma} \right) d\gamma' \right) \right]. \end{aligned} \quad (4.12)$$

By the Jensen's inequality, the right hand side of the above display is bounded by

$$\frac{1}{\gamma} \int_0^\gamma \mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma} \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma'} \right) \right) \right] d\gamma'. \quad (4.13)$$

We first now investigate the growth of the above expression when  $\alpha_0 \geq 2$  and then, show the growth in the case when  $\alpha_0 < 2$ . The first case is for theoretical interests since the actual range of  $\alpha_0$  for any fixed  $\gamma > 0$  is in between  $(-2, 2 + \frac{\gamma^2}{2})$ . The boundedness of the above expression follows from the property of  $z$ -deformed DGMC provided by Lemma 4.1:

$$\mathbb{P} \left( \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma} \right) \leq -v \right) \leq \tilde{c}_1 e^{-\tilde{c}_2 v^2}.$$

for any  $v > 0$ . We then obtain the inequality

$$\begin{aligned} \mathbb{E} \left[ e^{-\frac{\alpha_0-2}{\gamma} \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma} \right)} \right] &\leq 1 + \int_0^\infty e^{\frac{(\alpha_0-2)}{\gamma} v} v \mathbb{P} \left( \text{DM}_{\gamma'} \tilde{\Theta} \left( z, \tau; \frac{2}{\gamma} \right) \leq -v \right) dv \\ &\leq 1 + \tilde{c}_1 e^{\frac{(\alpha_0-2)^2}{2\tilde{c}_2\gamma^2}} \end{aligned}$$

for any  $\alpha_0 \geq 2$ . Substituting this above bound into (4.13) shows that  $\limsup_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, \tau) < \infty$  when  $\alpha_0 \geq 2$ .

Now we proceed to prove the same claim when  $\alpha_0 < 2$ . For any positive integer  $N$ , we can write  $\mathbb{E} [\mathcal{V}_{\chi,P}^\alpha(z, q)^N]$  as a Selberg-type integral

$$\mathbb{E} [\mathcal{V}_{\chi,P}^\alpha(z, q)^N] = \int_{[0,1]^N} \dots \int_0^1 \prod_{i < j}^N |\theta_1(x_i - x_j)|^{-\frac{\gamma^2}{2}} \prod_{i=1}^N \theta_1(z + x_i)^{\frac{\gamma x}{2}} \theta_1(z)^{-\frac{\gamma x}{2}} \theta_1(x_i)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x_i} dx_i.$$

Notice that  $\theta(z) = \mathbf{p}(z; \tau) \sin(\pi z)$  where  $c_1 \leq \mathbf{p}(z; \tau) \leq c_2$  as  $z$  varies in  $(0, 1)$ . As a result, we can prove that the right hand side of the above integral is bounded by above and below by some constant multiple of

$$\int_{[0,1]^N} \prod_{i < j}^N |\sin(\pi(x_i - x_j))|^{-\frac{\gamma^2}{2}} \prod_{i=1}^N \sin(x_i)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x_i} dx_i.$$

The above integral has an explicit expression as stated in (2.11) of Remark 2.1. As  $N$  grows, the ratios of gamma function in (2.11) grows as  $\exp(cN^2\gamma^2)$ . When  $\alpha_0 < 2$ ,  $\mathbb{E} [\mathcal{V}_{\chi,P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}]$  can be bounded by  $\mathbb{E} [\mathcal{V}_{\chi,P}^\alpha(z, q)^N]$  for some integer  $N = \lfloor (2-\alpha_0)/\gamma^2 \rfloor$ . Now the large  $N$  asymptotics of the above integral (obtained through the asymptotics of the ratios of gamma function in (2.11)) shows that  $\mathbb{E} [\mathcal{V}_{\chi,P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}]$  is bounded by  $\exp(c/\gamma^2)$  for some constant  $c = c(\alpha_0) > 0$  when  $\alpha_0 < 2$ . As a result, we get  $\limsup_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, \tau) < \infty$  as  $\gamma \rightarrow 0$ .

It remains now to show that  $\liminf_{\gamma \rightarrow 0} \gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau) > -\infty$ . For any  $\alpha_0 \in (-2, 2)$ , notice that  $h(x) = x^{(2-\alpha_0)/\gamma^2}$  is a convex function of  $x$  for all small  $\gamma$ . Hence, by Jensen's inequality, we have

$$\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}] \geq \left(\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)]\right)^{(2-\alpha_0)/\gamma^2}.$$

With the explicit form of  $\mathcal{V}_{\chi, P}^\alpha(z, q)$ ,

$$\left(\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)]\right)^{(2-\alpha_0)/\gamma^2} = \left(\int_0^1 \theta_1(z+x)\theta_1(z)^{-1}\theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} \frac{2-\alpha_0}{\gamma^2}\right),$$

we see that  $\liminf_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}] > -\infty$  due to the above inequality. Therefore, along with the expression (4.10), we have  $\liminf_{\gamma \rightarrow 0} \gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau) > -\infty$  for the deformed conformal block (4.9). This completes the proof of this proposition.  $\square$

**Proposition 4.2.** *As  $\gamma \downarrow 0$ ,  $\{\gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)\}_{\gamma > 0}$  is a tight sequence of functions in  $C^2([-M, M] \times [0, q_0])$  for any  $M > 0$  and  $q_0 \in (0, 1)$ . The parameters  $\alpha_0 \in (-2, 2)$ ,  $P_0 \in \mathbb{R}$ .*

*Proof.* Let us start by rewriting the deformed conformal block (2.14) as (see also eq. (3.5) in [Gho+24])

$$\psi_\chi^\alpha(z, q) = q^{\left(\frac{P^2}{2} + \frac{1}{6\chi^2} l_\chi(l_\chi+1)\right)} \widehat{\psi}_\chi^\alpha(z, q). \quad (4.14)$$

To prove this proposition, it is then sufficient to show that the limits  $\gamma \rightarrow 0$  and  $q \rightarrow 0$  commute for  $\widehat{\psi}_\chi^\alpha(z, q)$ . The key to proving the commutativity is to express the function  $\widehat{\psi}_\chi^\alpha(z, q)$  such that the integrand inside the expectation is  $q$  independent. Such a structure can be obtained via Girsanov's theorem (see [Gho+24, Theorem C.5]) that lets us re-express the conformal block in terms of the  $q$ -independent measure  $e^{\frac{\tau}{2}Y(x)}$ , resulting in the following expression for the function  $\widehat{\psi}_\chi^\alpha(z, q)$  in the equation above:

$$\begin{aligned} \widehat{\psi}_\chi^\alpha(z, q) &= C(q) e^{\chi P z \pi} \theta_1(z)^{-l_\chi} C_1(q) \left(-ie^{-i\pi z} q^{1/6} \eta(q)\right)^{l_\chi} \\ &\times \mathbb{E} \left[ e^{\frac{\alpha}{2} F_\tau(0)} \mathcal{Q}(q) e^{\chi \mathcal{Y}(z, q)} e^{\chi \mathcal{X}(z, q) - \frac{\chi^2}{2} \mathbb{E}[\mathcal{X}(z, q)^2]} \left( \int_0^1 dx e^{\frac{\tau}{2} Y(x)} e^{\pi \gamma P x} (2 \sin(\pi x))^{-\frac{\alpha \gamma}{2}} \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right], \end{aligned} \quad (4.15)$$

where

$$\begin{aligned} C(q) &:= q^{\frac{\gamma l_\chi}{12\chi} - \frac{1}{6} \frac{l_\chi^2}{\chi^2} - \frac{1}{6\chi^2} l_\chi(l_\chi+1)} \theta_1'(0) - \frac{2l_\chi^2}{3\chi^2} + \frac{l_\chi}{3} + \frac{4l_\chi}{3\gamma\chi} e^{-\frac{1}{2} i\pi \alpha \gamma \left(-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}\right)}, \\ C_1(q) &:= \left(q^{1/6} \eta(q)\right)^{\alpha(\alpha-\chi)/2} e^{\left(\frac{\alpha\gamma}{8} - \frac{\gamma\chi}{8} - \frac{\alpha^2}{8}\right) \mathbb{E}[F_\tau(0)^2]}, \\ \mathcal{Q}(q) &:= \exp \left[ \sqrt{2} \sum_{m, n=1}^{\infty} q^{nm} (\alpha_{n, m} \alpha_n + \beta_{n, m} \beta_n) - \sum_{n=1}^{\infty} \left( \sum_{m=1}^{\infty} q^{nm} \alpha_{m, n} \right)^2 - \sum_{n=1}^{\infty} \left( \sum_{m=1}^{\infty} q^{nm} \beta_{m, n} \right)^2 \right], \\ F_\tau(0) &:= 2 \sum_{n, m \geq 1} \frac{q^{nm}}{\sqrt{n}} \alpha_{n, m} \\ \mathcal{X}(z, q) &:= -\sqrt{2} \sum_{n, m=1}^{\infty} \frac{q^{(2m-1)n}}{\sqrt{n}} \left( \cos \left( 2\pi \left( z - \frac{\tau}{2} \right) n \right) \alpha_n - \sin \left( 2\pi \left( z - \frac{\tau}{2} \right) n \right) \beta_n \right), \\ \mathcal{Y}(z, q) &:= 2 \sum_{m, n, k \geq 1} \alpha_{n, m} \frac{q^{(2k-1+m)n}}{\sqrt{n}} \cos \left( 2\pi \left( z - \frac{\tau}{2} \right) n \right) - 2 \sum_{m, n, k \geq 1} \beta_{n, m} \frac{q^{(2k-1+m)n}}{\sqrt{n}} \sin \left( 2\pi \left( z - \frac{\tau}{2} \right) n \right), \end{aligned} \quad (4.16)$$

and for convenience we define the  $\gamma$ -independent entity

$$\tilde{\mathcal{X}}(z, q) := \frac{\alpha_0}{2} F_\tau(0) + 2(\mathcal{X}(z, q) + \mathcal{Y}(z, q)). \quad (4.17)$$

With all the above definitions, the equation (4.15) simplifies as

$$\begin{aligned} \widehat{\psi}_\chi^\alpha(z, q) &= C(q) e^{\chi P z \pi} \theta_1(z)^{-l_\chi} C_1(q) \left( -\mathbf{i} e^{-i\pi z} q^{1/6} \eta(q) \right)^{\frac{\chi}{2}(\chi - \alpha)} e^{-\frac{\chi^2}{2} \mathbb{E}[\mathcal{X}(z, q)^2]} \\ &\times \mathbb{E} \left[ e^{\frac{1}{\gamma} \tilde{\mathcal{X}}(z, q)} \mathcal{Q}(q) \left( \int_0^1 dx e^{\frac{\gamma}{2} Y(x)} e^{\pi \gamma P x} (2 \sin(\pi x))^{-\frac{\alpha \gamma}{2}} \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right]. \end{aligned} \quad (4.18)$$

Restricting to  $\chi = \frac{2}{\gamma}$ , we observe that the term

$$C(q) e^{\chi P z \pi} \theta_1(z)^{-l_\chi} C_1(q) \left( -\mathbf{i} e^{-i\pi z} q^{1/6} \eta(q) \right)^{\frac{\chi}{2}(\chi - \alpha)}$$

is independent of  $q$ . Therefore, the first term in (4.18) has the following limit

$$\begin{aligned} &\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left[ C(q) e^{\chi P z \pi} \theta_1(z)^{-l_\chi} C_1(q) \left( -\mathbf{i} e^{-i\pi z} q^{1/6} \eta(q) \right)^{\frac{\chi}{2}(\chi - \alpha)} e^{-\frac{\chi^2}{2} \mathbb{E}[\mathcal{X}(z, q)^2]} \right] \\ &= -2 \lim_{q \rightarrow 0} \mathbb{E} [\mathcal{X}(z, q)^2]. \end{aligned} \quad (4.19)$$

As for the second term of (4.18), the facts that  $\lim_{q \rightarrow 0} \mathcal{Q}(q) = 1$ , and the term containing the integral is  $q$ -independent allow us to factorize the terms inside the expectation as follows:

$$\begin{aligned} &\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ e^{\frac{1}{\gamma} \tilde{\mathcal{X}}(z, q)} \mathcal{Q}(q) \left( \int_0^1 dx e^{\frac{\gamma}{2} Y(x)} e^{\pi \gamma P x} (2 \sin(\pi x))^{-\frac{\alpha \gamma}{2}} \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right] \\ &= \lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ e^{\frac{1}{\gamma} \tilde{\mathcal{X}}(z, q)} \right] \\ &+ \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( \int_0^1 dx e^{\frac{\gamma}{2} Y(x)} e^{\pi \gamma P x} (2 \sin(\pi x))^{-\frac{\alpha \gamma}{2}} \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right]. \end{aligned} \quad (4.20)$$

The second term in the expression above is a constant

$$C_2(\alpha_0, P_0) := \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( \int_0^1 dx e^{\frac{\gamma}{2} Y(x)} e^{\pi \gamma P x} (2 \sin(\pi x))^{-\frac{\alpha \gamma}{2}} \right)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right],$$

while the first term of the expression (4.20) simplifies as follows due to the identity  $\mathbb{E}[e^{\frac{1}{\gamma} \tilde{\mathcal{X}}(z, q)}] = e^{\frac{1}{2\gamma^2} \mathbb{E}[(\tilde{\mathcal{X}}(z, q))^2]}$ .

$$\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E}[e^{\frac{1}{\gamma} \tilde{\mathcal{X}}(z, q)}] = \lim_{q \rightarrow 0} \frac{1}{2} \mathbb{E}[(\tilde{\mathcal{X}}(z, q))^2]. \quad (4.21)$$

With (4.19), (4.20), (4.21), the limit of the regularized  $z$ -deformed conformal block (4.18) is therefore

$$\begin{aligned} \lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \widehat{\psi}_{2/\gamma}^{\alpha_0/\gamma} &= -2 \lim_{q \rightarrow 0} \mathbb{E} [\mathcal{X}(z, q)^2] + \lim_{q \rightarrow 0} \frac{1}{2} \mathbb{E}[(\tilde{\mathcal{X}}(z, q))^2] + C_2(\alpha_0, P_0) \\ &\stackrel{(4.17)}{=} \lim_{q \rightarrow 0} \frac{1}{2} \mathbb{E} \left[ \frac{\alpha_0^2}{4} F_\tau(0)^2 + 2(2\mathcal{Y} + \alpha_0 F_\tau(0)) (\mathcal{X} + \mathcal{Y}) + 4\mathcal{X}\mathcal{Y} \right] + C_2(\alpha_0, P_0), \end{aligned}$$

which is an analytic function in  $q$  in some open ball  $|q| < r_0$  for all  $z \in (0, 1)$ . This shows that the above convergence is uniform on the ball  $|q| < r_0 - \epsilon$  for any  $\epsilon > 0$ . The limits  $\gamma \rightarrow 0$ ,  $q \rightarrow 0$  are therefore commutative, proving this proposition.  $\square$

The following lemmas which are simple corollary of the results from Section 3 played important role in the proofs of Proposition 4.1 and Proposition 4.2.

**Lemma 4.1.** *There exist  $\tilde{c}_1 = \tilde{c}_1(z, \tau)$ ,  $\tilde{c}_2 = \tilde{c}_2(\tau)$  such that for all  $v > 0$ , the DGMC measure (4.2) satisfies the following inequality*

$$\mathbb{P} \left[ \mathbf{DM}_{\gamma, N} \tilde{\Theta}(z, \tau; \chi) \leq -v \right] \leq \tilde{c}_1 e^{-\tilde{c}_2 v^2}. \quad (4.22)$$

*Proof.* Note that (4.2) is (3.1) with  $f(z, x, \tau) = \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x}$ , and  $f(\cdot, z, \tau) \in L_1([0, 1])$ . Moreover, from the proof of Theorem 3.1, we can observe that the exponent in (3.3) arises from the explicit form of  $Y(x; q)$  whereas the proportionality constant is given by the normalization factor  $\int_0^1 f(z, x, \tau)$ , resulting in (4.22).  $\square$

**Lemma 4.2.** *The semi-classical limits of the following  $z$  and  $\tau$  derivatives of the conformal block are bounded*

$$\begin{aligned} \limsup_{\gamma \rightarrow 0} \gamma^2 \partial_z \log(\psi_\chi^{\alpha/\gamma}(z, q)) < \infty, & \quad \liminf_{\gamma \rightarrow 0} \gamma^2 \partial_z \log(\psi_\chi^{\alpha/\gamma}(z, q)) > -\infty \\ \limsup_{\gamma \rightarrow 0} \gamma^2 \partial_\tau \log(\psi_\chi^{\alpha/\gamma}(z, q)) < \infty, & \quad \liminf_{\gamma \rightarrow 0} \gamma^2 \partial_\tau \log(\psi_\chi^{\alpha/\gamma}(z, q)) > -\infty \end{aligned}$$

*Proof.* Using the form of the conformal block (4.14),(4.15) where the  $z$  and  $\tau$  dependence is separated from the GMC measure, we can prove the boundedness of the  $z$  and  $\tau$  derivatives  $\partial_z \log(\psi_\chi^{\alpha/\gamma}(z, q))$ ,  $\partial_\tau \log(\psi_\chi^{\alpha/\gamma}(z, q))$  in the following way. Repeating the arguments of the proof of Proposition 4.1, we observe that the expressions for the  $z$ ,  $\tau$  derivative above is determined by the DGMC measure (3.1) with appropriate test functions. The proof of this lemma then follows from the tail of the measure given by the second expression in (3.3) with suitable constant functions.  $\square$

4.1.1. *Proof of Theorem 4.1.* The explicit form of the semi-classical conformal block can be obtained in the following manner. We begin with the following term in the conformal block (4.9):

$$\begin{aligned} \mathbb{E} \left[ \mathcal{V}_{\gamma, P}^\alpha(z, q)^{-\frac{(2-\alpha_0)}{\gamma^2}} \right] &\stackrel{(4.11),(4.12)}{=} \left( \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\left(\frac{\alpha_0-2}{\gamma^2}\right)} \\ &\quad \mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma} \mathbf{DM}_{\gamma'} \tilde{\Theta}(z, \tau; \frac{2}{\gamma}) \right) \right]. \end{aligned}$$

Along with the expression (4.10), (4.9) reads

$$\begin{aligned} \psi_{2/\gamma}^\alpha(z, \tau) &= (-2\pi)^{\frac{1}{24}(2-\alpha_0) + \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12\gamma^2}} q^{\frac{1}{24}(2-\alpha_0) + \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12\gamma^2}} e^{\frac{2P_0 z \pi}{\gamma^2}} \\ &\quad \times \left( \int_0^1 \frac{\theta_1(z+x)}{\theta_1(z)} \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\left(\frac{\alpha_0-2}{\gamma^2}\right)} \mathbb{E} \left[ \exp \left( -\frac{\alpha_0-2}{\gamma} \mathbf{DM}_{\gamma'} \tilde{\Theta}(z, \tau; \frac{2}{\gamma}) \right) \right] + \mathcal{O}(\gamma^2). \end{aligned}$$

Proposition 4.2 implies that  $\gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$ , is a tight sequence of functions in  $C^2([-M, M] \times [0, q_0])$  for any  $M > 0$  and  $q_0 \in (0, 1)$ . As a result, any subsequence of  $\gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$  has a further converging subsequence. Now we show that all subsequential limits of  $\gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$  are same. To this end, let us fix a subsequential limit

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q) = \tilde{\phi}(z, q). \quad (4.23)$$

Furthermore, due to Lemma 4.2,  $\gamma^2 \partial_z \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$ ,  $\gamma^2 \partial_\tau \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$  are also tight sequences of functions in  $C^2([-M, M] \times [0, q_0])$  for any  $M > 0$  and  $q_0 \in (0, 1)$ . Consequently, with (4.23), the BPZ equation (2.17) in the limit  $\gamma \rightarrow 0$  reads

$$(\partial_z \tilde{\phi}(z, q))^2 - (2 - \alpha_0)^2 \wp(z) + 2\pi i \partial_\tau \tilde{\phi}(z, q) = 0.$$

Note that the above equation is the Hamilton-Jacobi equation corresponding to the non-autonomous elliptic Calogero-Moser model, see Remark 4.1.

From all of the above, any sub-sequential limit of  $\gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$  will solve (4.5). Note that the commutativity of limits  $\gamma \rightarrow 0$  and  $q \rightarrow 0$  uniquely determines the initial condition of Hamilton-Jacobi equation (4.5) through Proposition 6.3. Then, by the uniqueness of the solution of the Hamilton-Jacobi equation (see Remark 4.1) and the fact that all sub-sequential limits of  $\tilde{\phi}(z, q)$  satisfy (4.5), we obtain that all the sub-sequential limits of  $\tilde{\phi}(z, q)$  are same. As a result,  $\lim_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z; q)$  exists and coincides with (4.4).

**4.2. Semi-classical limit of the deformed conformal blocks for  $\chi = \frac{\gamma}{2}$  and the Lamé equation.** As in the previous section, we study the semi-classical limit of the deformed conformal block for the deformation parameter  $\chi = \frac{\gamma}{2}$ . We show that the semi-classical limit of the deformed conformal block solves Lamé equation as noted in [Piq14]. Furthermore, we derive relation between the accessory parameter of the Lamé equation and the Hamiltonian mentioned in (4.7) (see also [BGG22]). We first state this semi-classical limit in the following result (Theorem 4.2). We prove this result after stating and proving Proposition 4.3 which shows the boundedness of the sequence of deformed conformal block  $\psi_{\gamma/2}^\alpha(z, q)$  as  $\gamma \rightarrow 0$ .

**Theorem 4.2.** *Consider  $\chi = \frac{\gamma}{2}$ ,  $q \in (0, 1)$ , and the constants  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ . Then, the following limit exists*

$$\lim_{\gamma \rightarrow 0} \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) e^{-\phi(q)/\gamma^2} =: \tilde{\Gamma}(z; \alpha_0, P_0, q),$$

where  $\phi(q)$  is given by

$$\begin{aligned} \phi(q) := & -\frac{\alpha_0(\alpha_0 + 4)}{6} \log(-2\pi) + \frac{6P_0^2 - \alpha_0(\alpha_0 + 2)}{12} \log q - \alpha_0 \left( \log \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) \\ & + \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - 2\gamma^2}{\gamma} \text{DM}_{\gamma'} \Theta(\tau) \right) \right]. \end{aligned} \quad (4.24)$$

Moreover,  $\phi(q)$  has the following properties:

(1)  $\partial_\tau \phi(q)$  is assuming the role of the accessory parameter of the Lamé equation

$$\left( \partial_z^2 - \frac{\alpha_0}{4} \left( \frac{\alpha_0}{4} - 1 \right) \wp(z) + \frac{\pi \mathbf{i}}{2} \partial_\tau \phi(q) \right) \tilde{\Gamma}(z; \alpha_0, P_0, q) = 0.$$

where the solution of the above equation  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is given by

$$\tilde{\Gamma}(z; \alpha_0, P_0, q) = e^{\frac{P_0 z \pi}{2}} \exp \left( -\frac{\alpha_0 \int_0^1 \log \left( \frac{\theta_1(z+x)}{\theta_1(z)} \right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{4 \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} \right).$$

(2) Furthermore,  $\partial_\tau \phi(q)$  is also described by the Hamiltonian in (4.7) as:

$$2\pi \mathbf{i} \partial_{\tau_*} \phi(q_*) = H(\tau_*), \quad q_* = e^{i\pi \tau_*} \quad (4.25)$$

for all  $\tau_*$  such that  $u(\tau_*) = 0$  (see Remark 4.1 for the definition of  $u(\tau)$ ).

Let us now detail the propositions needed to prove the above theorem.

**Proposition 4.3.** *For  $\chi = \frac{\gamma}{2}$ ,  $q \in (0, 1)$ , and the constants  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ , the sequence of the deformed conformal blocks (2.14) is bounded*

$$\limsup_{\gamma \rightarrow 0} \gamma^2 \log \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) < \infty, \quad \liminf_{\gamma \rightarrow 0} \gamma^2 \log \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) > -\infty.$$

*Proof.* We begin by recalling the expression for the deformed conformal block (2.14) with the scaling  $\alpha = \alpha_0/\gamma$ ,  $P = P_0/\gamma$ :

$$\psi_\chi^{\alpha_0/\gamma}(z, \tau) = \mathcal{W}(q) e^{\chi \frac{P_0}{\gamma} z \pi} \mathbb{E} \left[ \mathcal{V}_{\chi, P_0/\gamma}^{\alpha_0/\gamma}(z, q)^{-\frac{\alpha_0}{\gamma^2} + \frac{\chi}{\gamma}} \right], \quad (4.26)$$

where

$$\mathcal{V}_{\chi, P_0/\gamma}^{\alpha_0/\gamma}(z, q) := \int_0^1 \theta_1(z+x)^{\frac{\gamma \chi}{2}} \theta_1(z)^{-\frac{\gamma \chi}{2}} \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx. \quad (4.27)$$

Let us examine each of the terms above separately, beginning with the parameter  $l_\chi$  for  $\chi = \gamma/2$ :

$$l_\chi = \frac{\chi^2}{2} - \frac{\alpha\chi}{2} = \frac{\gamma^2}{8} - \frac{\alpha_0}{4}.$$

The pre-factor in (4.26) that is defined by (2.15), simplifies as follows in the limit  $\gamma \rightarrow 0$ :

$$\begin{aligned} \mathcal{W}(q) e^{\chi \frac{P_0}{\gamma} z \pi} &\stackrel{(2.15)}{=} q^{\frac{P_0^2}{2\gamma^2} + \frac{\gamma l_\chi}{12\chi} - \frac{1}{6} \frac{l_\chi^2}{\chi^2}} \theta_1'(0) - \frac{2l_\chi^2}{3\chi^2} + \frac{l_\chi}{3} + \frac{4l_\chi}{3\gamma\chi} e^{P_0 z \pi/2} \\ &\stackrel{(A.5)}{=} (-2\pi)^{-\frac{\alpha_0(\alpha_0+4)}{6\gamma^2} + \frac{1}{12}(\alpha_0+4)} q^{\frac{1}{48}(\alpha_0+4) + \frac{6P_0^2 - \alpha_0(\alpha_0+2)}{12\gamma^2}} e^{\frac{P_0 z \pi}{2}} + \mathcal{O}(q^{\gamma^2}). \end{aligned} \quad (4.28)$$

It is worth noting that the term above can be divided into two components: one dependent on  $z$  and the other dependent on  $\gamma$ . Such a structure will be important in the subsequent analysis.

Next we observe that

$$\theta_1(z+x)\gamma^{2/4}\theta_1(z)^{-\gamma^2/4} = e^{\gamma^2/4 \log(\theta_1(z+x)/\theta_1(z))} = 1 + \frac{\gamma^2}{4} \log\left(\frac{\theta_1(z+x)}{\theta_1(z)}\right) + \mathcal{O}(\gamma^3).$$

Substituting the above expression in (4.27), we obtain the following:

$$\begin{aligned} \mathcal{V}_{\chi, P_0/\gamma}^{\alpha_0/\gamma}(z, q) &= \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx \\ &\quad + \frac{\gamma^2}{4} \int_0^1 \log\left(\frac{\theta_1(z+x)}{\theta_1(z)}\right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx + \mathcal{R}_0^\gamma(z, q), \end{aligned} \quad (4.29)$$

where  $|\mathcal{R}_0^\gamma(z, q)| \lesssim \gamma^3 \mathcal{V}_{\chi, P}^\alpha(z, q)$ . With the above expansion, we can now write

$$\log \mathcal{V}_{\chi, P}^\alpha(z, q) = \log\left(\int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx\right) + \log(1 + \tilde{\mathcal{R}}_0^\gamma(z, q)),$$

such that

$$\tilde{\mathcal{R}}_0^\gamma(z, q) = \frac{\frac{\gamma^2}{4} \int_0^1 \log\left(\frac{\theta_1(z+x)}{\theta_1(z)}\right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx + \mathcal{R}_0^\gamma(z, q)}{\int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx}.$$

As  $\gamma \rightarrow 0$ , we obtain that

$$\log\left(1 + \tilde{\mathcal{R}}_0^\gamma(z, q)\right) = \frac{\gamma^2 \int_0^1 \log\left(\frac{\theta_1(z+x)}{\theta_1(z)}\right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{\int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} + \mathcal{O}(\gamma^3). \quad (4.30)$$

With the expressions (4.29) and (4.30), the term with the expectation in (4.26) reads

$$\begin{aligned} &\mathbb{E}\left[\mathcal{V}_{\chi, P}^\alpha(z, q)^{\frac{\gamma^2/2 - \alpha_0}{\gamma^2}}\right] \\ &= \left(\int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx\right)^{-\frac{\alpha_0}{\gamma^2} + \frac{1}{2}} \\ &\quad \times \mathbb{E}\left[\exp\left(-\frac{\alpha_0 - \gamma^2/2}{\gamma^2} (\log \mathcal{V}_{\chi, P}^\alpha(z, q)) + \frac{\alpha_0 - \gamma^2/2}{\gamma^2} \left(\log \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx\right)\right)\right]. \end{aligned} \quad (4.31)$$

The second term in the above expression can be further simplified as

$$\begin{aligned} &\mathbb{E}\left[\exp\left(-\frac{\alpha_0 - \gamma^2/2}{\gamma^2} (\log \mathcal{V}_{\chi, P}^\alpha(z, q)) + \frac{\alpha_0 - \gamma^2/2}{\gamma^2} \left(\log \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx\right)\right)\right] \\ &= \mathbb{E}\left[\exp\left(-\frac{\alpha_0 - \gamma^2/2}{\gamma^2} \left(\log \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx\right) \right. \right. \\ &\quad \left. \left. + \frac{\alpha_0 - \gamma^2/2}{\gamma^2} \left(\log \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx\right)\right)\right] \\ &\quad \times \exp\left(-\frac{\alpha_0 \int_0^1 \log\left(\frac{\theta_1(z+x)}{\theta_1(z)}\right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{4 \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} + \mathcal{O}(\gamma^2)\right), \end{aligned} \quad (4.32)$$

where the leading term is  $z$ -independent, and the sub-leading term is  $\gamma$ -independent. We first show that the above expression is bounded when  $\alpha_0 \in (-4, 0)$ . For this, we use the integer moment formulas of  $\mathcal{V}_{\chi, P}^\alpha(z, q)$ . For any positive integer  $N$ , we can write  $\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^N]$  as a Selberg-type integral

$$\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^N] = \int_{[0, 1]^N} \dots \int_0^1 \prod_{i < j}^N |\theta_1(x_i - x_j)|^{-\frac{\gamma^2}{2}} \prod_{i=1}^N \theta_1(z + x_i)^{\frac{\gamma x}{2}} \theta_1(z)^{-\frac{\gamma x}{2}} \theta_1(x_i)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x_i} dx_i.$$

Notice that  $\theta(z) = \mathfrak{p}(z; \tau) \sin(\pi z)$  where  $c_1 \leq \mathfrak{p}(z; \tau) \leq c_2$  as  $z$  varies in  $(0, 1)$ . As a result, we can upper and lower bound the right hand side of the above integral by some constant multiple of

$$\int_{[0, 1]^N} \prod_{i < j}^N |\sin(\pi(x_i - x_j))|^{-\frac{\gamma^2}{2}} \prod_{i=1}^N \sin(x_i)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x_i} dx_i.$$

The above integral has an explicit expression as stated in (2.11) of Remark 2.1. As  $N$  grows, the ratios of gamma function in (2.11) grows as  $\exp(cN^2\gamma^2)$ . When  $\alpha_0 < 2$ ,  $\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}]$  can be bounded by  $\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^N]$  for some integer  $N = \lfloor -\alpha_0/\gamma^2 \rfloor$ . Now the large  $N$  asymptotics of the above integral (obtained through the asymptotics of the ratios of gamma function in (2.11)) shows that  $\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)^{(2-\alpha_0)/\gamma^2}]$  bounded by  $\exp(c/\gamma^2)$  for some constant  $c = c(\alpha_0) > 0$ .

Now it remains to show similar bound when  $\alpha_0 \in [0, 2)$ . By the mean value theorem, there exists a  $\gamma' \in (0, \gamma)$  such that

$$\begin{aligned} & \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - \gamma^2/2}{\gamma^2} \left( \log \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx \right) + \frac{\alpha_0}{\gamma^2} \left( \log \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} \right) \right) \right] \\ & \stackrel{(4.1)}{=} \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - \gamma^2/2}{\gamma^2} \int_0^\gamma \text{DM}_{\gamma'} \Theta(\tau) d\gamma' \right) \right]. \end{aligned} \quad (4.33)$$

By Jensen's inequality, we have

$$\mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - \gamma^2/2}{\gamma^2} \int_0^\gamma \text{DM}_{\gamma'} \Theta(\tau) d\gamma' \right) \right] \leq \frac{1}{\gamma} \int_0^\gamma \mathbb{E} \left[ \exp \left( -\frac{\alpha_0}{\gamma} \text{DM}_{\gamma'} \Theta(\tau) \right) \right] d\gamma'$$

Due to the above relation, proving the boundedness of  $\gamma^2 \log \mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{\frac{2\gamma^2 - \alpha_0}{\gamma^2}} \right]$  as  $\gamma \rightarrow 0$  is equivalent to showing that, for some constants  $c > 0, \tilde{c} > 0$ ,

$$\mathbb{E} \left[ \exp \left( -\frac{\alpha_0}{\gamma} \text{DM}_{\gamma'} \Theta(\tau) \right) \right] < \tilde{c} e^{c \frac{\alpha_0}{\gamma^2}}.$$

We show the above when  $\alpha_0 \in [0, 2)$ . Indeed, from Lemma 4.3, we obtain that

$$\mathbb{E} \left[ e^{-\frac{\alpha_0}{\gamma} \text{DM}_{\gamma'} \Theta(\tau)} \right] \leq 1 + \frac{\alpha_0}{\gamma} \int_0^\infty e^{\frac{\alpha_0}{\gamma} v} \mathbb{P}(\text{DM}_{\gamma'} \Theta(\tau) \leq -v) dv \leq 1 + c_1 e^{\frac{\alpha_0^2}{2c_2 \gamma^2}} \quad (4.34)$$

for  $c_1 = c_1(\tau)$ ,  $c_2 = c_2(\tau)$ , and  $c_1 > 0$ ,  $c_2 > 0$ . The above computation shows that

$$\limsup_{\gamma \rightarrow 0} \gamma^2 \log \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) < \infty.$$

It remains to show  $\liminf_{\gamma \rightarrow 0} \gamma^2 \log \log \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) > -\infty$ . Since  $h(x) = x^{-(\alpha_0 - \gamma^2/2)/\gamma^2}$  is a convex function for small  $\gamma > 0$ , we bound

$$\mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{\frac{\gamma^2/2 - \alpha_0}{\gamma^2}} \right] \geq (\mathbb{E}[\mathcal{V}_{\chi, P}^\alpha(z, q)])^{-\frac{(\alpha_0 - \frac{\gamma^2}{2})}{\gamma^2}} = \left( \int_0^1 \theta_1(z+x)^{\frac{\gamma^2}{4}} \theta_1(z)^{-\frac{\gamma^2}{4}} \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\frac{(\alpha_0 - \frac{\gamma^2}{2})}{\gamma^2}}$$

via Jensen's inequality. From the the above display, it follows that

$$\liminf_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{\frac{\gamma^2/2 - \alpha_0}{\gamma^2}} \right] > -\infty.$$

This implies that  $\liminf_{\gamma \rightarrow 0} \gamma^2 \log \log \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) > -\infty$ .  $\square$

**Lemma 4.3.** *There exist some constants  $c_1 = c_1(\tau), c_2 = c_2(\tau), c_3, c_4 > 0$  such that the DGMC measures (4.1) satisfy the following inequalities for  $v > 0$*

$$\mathbb{P} [\text{DM}_{\gamma, N} \Theta(\tau) \leq -v] \leq c_1 e^{-c_2 v^2}, \quad \mathbb{P} [\text{DM}_{\gamma, N} \Theta(\mathbf{i}\infty) \leq -v] \leq c_3 e^{-c_4 v^2}. \quad (4.35)$$

*Proof.* Note that (4.1) is simply (3.1) with the choice of the test function  $f(x, \tau) = \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x}$ . Since  $f(\cdot, \tau) \in L_1([0, 1])$  for all  $\tau \in \mathbb{H}$ , the statement of this Lemma follows from Theorem 3.1.  $\square$

#### 4.2.1. Proof of Theorem 4.2.

*Proving (1):* We begin by obtaining the explicit expression for the conformal block. Gathering the expressions (4.31), (4.32), (4.33), the following term in (4.26) can be written as

$$\begin{aligned} \mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{\frac{2\gamma^2 - \alpha_0}{\gamma^2}} \right] &= \left( \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\frac{\alpha_0}{\gamma^2} + \frac{1}{2}} \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - 2\gamma^2}{\gamma} \text{DM}_{\gamma'} \Theta(\tau) \right) \right] \\ &\quad \times \exp \left( -\frac{\alpha_0 \int_0^1 \log \left( \frac{\theta_1(z+x)}{\theta_1(z)} \right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{4 \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} + O(\gamma^2) \right). \end{aligned}$$

Substituting the above equation and (4.28) in (4.26), we have

$$\begin{aligned} \psi_\chi^{\alpha_0/\gamma}(z, \tau) &= (-2\pi)^{-\frac{\alpha_0(\alpha_0+4)}{6\gamma^2} + \frac{1}{12}(\alpha_0+4)} q^{\frac{1}{48}(\alpha_0+4) + \frac{6P_0^2 - \alpha_0(\alpha_0+2)}{12\gamma^2}} e^{\frac{P_0 z \pi}{2}} \left( \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\frac{\alpha_0}{\gamma^2} + \frac{1}{2}} \\ &\quad \times \mathbb{E} \left[ \exp \left( -\frac{\alpha_0 - 2\gamma^2}{\gamma} \text{DM}_{\gamma'} \Theta(\tau) \right) \right] \exp \left( \frac{\int_0^1 \log \left( \frac{\theta_1(z+x)}{\theta_1(z)} \right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{\int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} \right) + O(\gamma^2). \end{aligned} \quad (4.36)$$

By Theorem 3.1, we know that  $\gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau)$  is tight as  $\gamma \rightarrow 0$ . As a result,  $\gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau)$  has subsequential limit. We now show that all subsequential limits of  $\gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau)$  are same. Suppose  $\phi(q)$  is a subsequential limit of  $\gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau)$ , i.e., there exists a subsequence  $\gamma_n$  converging to 0 such that  $\lim_{\gamma_n \rightarrow 0} \gamma_n^2 \log \psi_{\chi_n}^{\alpha_0/\gamma_n}(z, \tau) = \phi(q)$ . As a result, we get the following expression up to a  $z$ -independent constant  $c_0(q)$ :

$$\begin{aligned} \lim_{\gamma \rightarrow 0} \psi_{\gamma/2}^\alpha e^{-\phi(q)/\gamma^2} &=: c_0(q) \tilde{\Gamma}(z; \alpha_0, P_0, q) \\ &= c_0(q) e^{\frac{P_0 z \pi}{2}} \exp \left( -\frac{\alpha_0 \int_0^1 \log \left( \frac{\theta_1(z+x)}{\theta_1(z)} \right) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx}{4 \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx} \right). \end{aligned} \quad (4.37)$$

Substituting the above expression for the solution of the BPZ equation (2.17), it is straightforward to see that it reduces to the Lamé equation in the semi-classical limit:

$$\left( \partial_z^2 - \frac{\alpha_0}{4} \left( \frac{\alpha_0}{4} - 1 \right) \wp(z) + \frac{\pi \mathbf{i}}{2} \partial_\tau \phi(q) \right) \tilde{\Gamma}(z; \alpha_0, P_0, q) = 0,$$

where  $\pi \mathbf{i} \partial_\tau \phi(q)/2$  is the accessory parameter, and  $\alpha_0(4 + \alpha_0) \geq 0$  since  $\alpha_0 \in (0, 2)$ . Note that  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is explicit in (4.37). From [Ere+22; Inc40], it follows that  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  uniquely determine the accessory parameter  $\frac{\pi \mathbf{i}}{2} \partial_\tau \phi(q)$ . This in turn shows that all the subsequential limit of  $\gamma^2 \log \psi_\chi^{\alpha_0/\gamma}(z, \tau)$  are same and uniquely determined by  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$ .

*Proving (2):* Recall that the semi-classical limits of the deformed and undeformed conformal blocks in (2.9), (2.14) are related through (2.16):

$$\begin{aligned} \lim_{z \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(\chi-\alpha)} \psi_\chi^\alpha(z, \tau) \right) &= \lim_{z \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(\chi-\alpha)} \mathcal{W}(q) \mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right] \right) \\ &= \lim_{\gamma \rightarrow 0} \lim_{z \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(\chi-\alpha)} \mathcal{W}(q) \mathbb{E} \left[ \mathcal{V}_{\chi, P}^\alpha(z, q)^{-\frac{\alpha}{\gamma} + \frac{\chi}{\gamma}} \right] \right) \\ &= \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha(q) + \lim_{\gamma \rightarrow 0} \gamma^2 \log Z. \end{aligned} \quad (4.38)$$

To obtain the last line we use the fact that the limits  $z \rightarrow 0$  and  $\gamma \rightarrow 0$  for the above expression are commutative as can be seen by substituting the alternative description of the conformal block (4.15) in the expression above and repeating the arguments of the proof of Proposition 4.2.

Note that the RHS of the expression (4.38) is independent of the choice of  $\chi$ . Using this fact, we can compare its LHS for the two choices of  $\chi$ . The relation (4.38) for  $\chi = \frac{2}{\gamma}$  reads:

$$\lim_{z \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(\chi-\alpha)} \psi_{2/\gamma}^\alpha(z, \tau) \right) \stackrel{(4.23)}{=} \lim_{z \rightarrow 0} \tilde{\phi}(z, \tau) + (2 - \alpha_0) \lim_{z \rightarrow 0} \log \theta_1(z).$$

Note that the term  $\lim_{z \rightarrow 0} \log \theta_1(z)$  cancels the singularity at  $z \rightarrow 0$  in (4.4). The RHS of the above expression is therefore analytic in  $z$ .

Similarly, for  $\chi = \frac{2}{\gamma}$ , we have (4.37), and the expression (4.38) reads:

$$\lim_{z \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(\chi-\alpha)} \psi_{\gamma/2}^\alpha(z, \tau) \right) = \phi(\tau). \quad (4.39)$$

As (4.38) is independent of the choice of  $\chi$ , a direct comparison of the above limits gives us that

$$\lim_{z \rightarrow 0} \tilde{\phi}(z, \tau) + (2 - \alpha_0) \lim_{z \rightarrow 0} \log \theta_1(z) = \phi(q).$$

Furthermore,  $\tilde{\phi}(\cdot)$  is associated to the action of the non-autonomous elliptic Calogero-Moser model (see Remark 4.1) with  $z = 2u(\tau)$ . Near its zero,  $u(\tau)$  has the following behaviour (B.17):

$$u(\tau) = e^{\mp i\pi/4} \sqrt{\frac{m}{2\pi}} (\tau - \tau_\star)^{1/2} \left( 1 \pm \frac{H_\star}{4\pi i m} (\tau - \tau_\star) \right) + \mathcal{O} \left( (\tau - \tau_\star)^{5/2} \right), \quad u(\tau_\star) = 0.$$

Using the above expression, we can obtain the asymptotic behaviour of  $v(\tau)$  through (4.6). Then a direct substitution of  $u(\tau), v(\tau)$  in the expression (4.8) give the following behaviour of the action

$$\tilde{\phi}(2u(\tau), \tau) = \pm \int \frac{m}{2(\tau - \tau_\star)} d\tau + \mathcal{O}(1).$$

The logarithmic singularity in the equation above is then cancelled by the asymptotic behaviour of the term  $(2 - \alpha_0) \lim_{u(\tau) \rightarrow 0} \log \theta_1(2u(\tau))$  near  $u(\tau) \rightarrow 0$  with  $m = \mp(2 - \alpha_0)$ . Therefore,

$$2\pi i \partial_\tau \phi(\tau) \Big|_{\tau=\tau_\star} = \partial_{\tau_\star} \left( \lim_{u(\tau) \rightarrow 0} \left( \tilde{\phi}(2u(\tau), \tau) + (2 - \alpha_0) \log \theta_1(2u(\tau)) \right) \right) = H(\tau_\star).$$

**Corollary 4.1.** *The accessory parameter has the explicit form*

$$\frac{\pi i}{2} \partial_\tau \phi(q, \alpha_0, P_0) = \left( -\frac{\partial_z^2 \tilde{\Gamma}(z; \alpha_0, P_0, q)}{\tilde{\Gamma}(z; \alpha_0, P_0, q)} + \frac{\alpha_0}{4} \left( \frac{\alpha_0}{4} - 1 \right) \wp(z) \right), \quad (4.40)$$

where  $\tilde{\Gamma}(z; \alpha_0, P_0, q)$  is given by (1.3).

**Remark 4.2.** *It is noteworthy that the accessory parameter has three equivalent descriptions in terms of the Hamiltonian (4.25), in terms of the DGMC measure that one can obtain through  $\tau$  derivative of the expression (4.24), and in terms of integrals of theta functions as (4.40).*

## 5. SEMI-CLASSICAL LIMIT OF UNDEFORMED CONFORMAL BLOCKS

The goal of this section is to prove our main result Theorem 1.1, which we present towards the end of this section. We begin the section with Theorems 5.1 and 5.2 that establish the existence and analyticity of the semi-classical limit of the undeformed conformal block as a function of the modular parameter  $q$  respectively. We then use these theorems along with the results from the previous section to complete the proof of Theorem 1.1.

**Theorem 5.1.** *The semi-classical limit  $\gamma \rightarrow 0$  of the one-point conformal block on the torus (2.9) exists for  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ . Explicitly, the semi classical limit of the conformal block is described by the DGMC measures as*

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma, P}^\alpha(q) = -\alpha_0 \mathfrak{Z}(\alpha_0, P_0, q) + \lim_{\gamma \rightarrow 0} \mathcal{M}(P_0, \alpha_0, q), \quad (5.1)$$

where the function

$$\mathcal{M}(P_0, \alpha_0, q) := \gamma^2 \left( \log \mathbb{E} \left[ e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)} \right] - \log \mathbb{E} \left[ e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\infty)} \right] \right),$$

with  $\text{DM}_\gamma \Theta(\tau)$ ,  $\text{DM}_\gamma \Theta(\mathbf{i}\infty)$  defined in (4.1), and

$$\begin{aligned} \mathfrak{Z}(\alpha_0, P_0, q) = \log \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) - \log \left( \int_0^1 (-2 \sin(\pi x))^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right) \\ + \alpha_0 \log(q^{\frac{1}{12}} \eta(q)^{-1}). \end{aligned}$$

**Theorem 5.2** (Radius of convergence). *There exists  $r_0 > 0$  which is independent of  $\alpha_0, P_0$  such that the semi-classical limit of the torus conformal block  $\mathcal{G}_{\gamma, P}^\alpha(q)$  is well defined and analytic as a function of  $q$  for all  $|q| < r_0$ .*

We now proceed to prove the above two theorems.

**5.1. Proofs of Theorems 5.1 and 5.2.** The main approach to proving Theorem 5.1 is as follows: to establish boundedness, we express the semi-classical limit of the conformal block  $\mathcal{G}_{\gamma, P}^\alpha(q)$  in (2.9) in terms of the DGMC measures (4.1), and use their property (4.35). The uniqueness of the conformal block then follows from the relation (4.38) which implies the uniqueness of the undeformed conformal block based on the uniqueness of the deformed conformal block established in Theorems 4.3 and 4.1.

*Proof of Theorem 5.1.* Recall the definition (2.9):

$$\mathcal{G}_{\gamma, P}^\alpha(q) := \frac{1}{Z} \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2} Y(x; q)} dx \right)^{-\frac{\alpha}{\gamma}} \right].$$

We can rewrite it as

$$\mathcal{G}_{\gamma, P}^\alpha(q) = g(q) \mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}} \right] / \mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(0))^{-\frac{\alpha}{\gamma}} \right], \quad (5.2)$$

where

$$\mathcal{F}_{\gamma, P}^{(\alpha)}(q) := q^{\frac{\alpha\gamma}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2} Y(x; q)} dx, \quad g(q) = q^{\frac{1}{12}(-\frac{\alpha\gamma}{2} + \alpha^2 + 1)} \eta(q)^{-\alpha^2 - 1 + \frac{\alpha\gamma}{2}}. \quad (5.3)$$

With the above expression, the semi-classical limit in (5.1) can be estimated through the asymptotics of  $\mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}} \right]$  and  $\mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(0))^{-\frac{\alpha}{\gamma}} \right]$ . Let us now evaluate each of the aforementioned terms. We will start by simplifying the following expression with the scaling  $\alpha = \frac{\alpha_0}{\gamma}$ ,  $P = \frac{P_0}{\gamma}$ :

$$\begin{aligned} \mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}} \right] &= \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right)^{-\frac{\alpha_0}{\gamma^2}} \\ &\quad \times \mathbb{E} \left[ \exp \left( -\frac{\alpha_0}{\gamma^2} \left[ \log \mathcal{F}_{\gamma, P_0/\gamma}^{(\alpha_0/\gamma)}(q) - \log q^{\frac{\alpha_0}{8}} \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx \right] \right) \right]. \end{aligned}$$

Our goal is to show that  $\mathbb{E} \left[ (\mathcal{F}_{\gamma, P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}} \right]$  grows as  $\exp(c/\gamma^2)$  for  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$  and  $q \in B_r(0)$  for some small  $r > 0$ . We first show the case when  $\alpha_0 \in (-4, 0)$ . In this case it suffices

to show that  $\mathbb{E}\left[(\mathcal{F}_{\gamma,P}^{(\alpha)}(q))^N\right]$  grows as  $\exp(c/\gamma^2)$  for any integer  $N = O(1/\gamma^2)$ . However such moment bounds is very similar to one showed in the proof of Proposition 4.3 for bounding (4.31).

Now we proceed to show the bound  $\mathbb{E}\left[(\mathcal{F}_{\gamma,P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}}\right]$  when  $\alpha_0 \in (0, 2)$ . By the mean value theorem, for any small  $\gamma > 0$ , there exists a  $\gamma' \in (0, \gamma)$  such that

$$\begin{aligned} & \mathbb{E}\left[\exp\left(-\frac{\alpha_0}{\gamma^2}\left[\log \mathcal{F}_{\gamma,P_0/\gamma}^{(\alpha_0/\gamma)}(q) - \log q^{\frac{\alpha_0}{8}} \int_0^1 \theta(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} dx\right]\right)\right] \\ & \stackrel{(4.1)}{=} \mathbb{E}\left[\exp\left(-\frac{\alpha_0}{\gamma^2} \int_0^\gamma \text{DM}_{\gamma'} \Theta(\tau) d\tau\right)\right]. \end{aligned}$$

Due to the above relation, boundedness of  $\gamma^2 \log \mathbb{E}\left[(\mathcal{F}_{\gamma,P_0/\gamma}^{(\alpha_0/\gamma)}(q))^{-\alpha_0/\gamma^2}\right]$  as  $\gamma \rightarrow 0$  is implied by the relation (4.34) for the DGMC measure. Similarly, boundedness of the second term in (5.2)  $\gamma^2 \log \mathbb{E}\left[(\mathcal{F}_{\gamma,P_0/\gamma}^{(\alpha_0/\gamma)}(0))^{-\alpha_0/\gamma^2}\right]$  follows from the relation below which is obtained using the second inequality in (4.35):

$$\mathbb{E}\left[\exp\left(-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\mathbf{i}\infty)\right)\right] \leq 1 + \tilde{c}_0 e^{\frac{\alpha_0^2}{2c_0\gamma^2}}, \quad (5.4)$$

for some constants  $\tilde{c}_0 > 0$ ,  $c_0 > 0$ . The inequalities (4.34) and (5.4) imply the upper bounds to the numerators and denominator of (5.2). The lower bounds could be easily derived by applying Jensen's inequality,

$$\mathbb{E}\left[(\mathcal{F}_{\gamma,P}^{(\alpha)}(q))^{-\frac{\alpha}{\gamma}}\right] \geq \left(\mathbb{E}[\mathcal{F}_{\gamma,P}^{(\alpha)}(q)]\right)^{-\frac{\alpha}{\gamma}}, \quad \mathbb{E}\left[(\mathcal{F}_{\gamma,P}^{(\alpha)}(0))^{-\frac{\alpha}{\gamma}}\right] \geq \left(\mathbb{E}[\mathcal{F}_{\gamma,P}^{(\alpha)}(0)]\right)^{-\frac{\alpha}{\gamma}}.$$

From the above bounds, it follows that

$$\gamma^2 \left| \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right] - \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\mathbf{i}\infty)}\right] \right| \leq C\alpha_0^2$$

for some positive constant  $C = C(\tau)$ ,  $C > 0$ . This above result shows the tightness of the term  $\gamma^2 \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right]$  as a sequence in  $\gamma$ . In order to prove that limit of this sequence exists, it suffices to that all subsequential limits of this sequence are same. For this, we proceed as in the proof of statement 1 of Theorem 4.2 using the ideas of the deformed conformal block. Recall  $\psi_{\gamma/2}^{\alpha_0/\gamma}(z, \tau)$  from (4.36), and  $\tilde{\Gamma}(z, \tau)$  from (4.37). Since  $\tilde{\Gamma}(z, \tau)$  is explicit and solves Lamé equation, the solution theory of Lamé equation (see [Ere+22; Inc40]) guarantees that the accessory parameter of the Lamé equation is uniquely identified. Moreover, by (4.36), it is clear that the accessory parameter of the Lamé equation associated to  $\tilde{\Gamma}(z, \tau)$  is given by the derivative of the subsequential limit of  $\gamma^2 \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right]$  upto some explicit constant. As a result, all the subsequential limits of  $\gamma^2 \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right]$  are same and hence,  $\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right]$  exists.

Therefore, the semi-classical limit of the conformal block (5.2) is given as

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma,P}^\alpha(q) = \lim_{\gamma \rightarrow 0} \gamma^2 \log g(q) + \lim_{\gamma \rightarrow 0} \gamma^2 \left( \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\tau)}\right] - \log \mathbb{E}\left[e^{-\frac{\alpha_0}{\gamma} \text{DM}_\gamma \Theta(\mathbf{i}\infty)}\right] \right)$$

The explicit expression for the conformal block (5.1) is then obtained by substituting the value of  $g(q)$  in (5.3) in the expression above.

Finally, the Theorems 4.1 and 4.3 prove the uniqueness of the deformed conformal block, which in turn imply the uniqueness of the undeformed conformal block due to (4.38).  $\square$

Let us now show that the semi-classical conformal block has a positive radius of convergence as stated in Theorem 5.2. The main element of the proof is the expression of conformal block that can be obtained using the Girsanov theorem [Gho+24, Theorem C.5] such that the GMC measure is made  $q$ -independent. Such an expression enables analytic control over the  $q$ -dependent part and helps us obtain the final result.

*Proof of Theorem 5.2.* Recall that

$$\mathcal{G}_{\gamma,P}^\alpha(q) = \frac{1}{Z} \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x;q)} dx \right)^{-\frac{\alpha}{\gamma}} \right]$$

where

$$Z := q^{\frac{1}{12}(\frac{\alpha\gamma}{2} + \frac{\alpha^2}{2} - 1)} \eta(q)^{\alpha^2 + 1 - \frac{\alpha\gamma}{2}} \mathbb{E} \left[ \left( \int_0^1 [-2 \sin(\pi x)]^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y_\infty(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right].$$

Note that  $\lim_{\gamma \rightarrow \infty} \gamma^2 \log Z$  is an analytic function of  $q$  for all  $|q| < 1$ . It then suffices to show that the expression

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x;q)} dx \right)^{-\frac{\alpha}{\gamma}} \right]$$

is well defined and analytic function for all  $q$  in  $|q| < r_0$  for some  $r_0$ . By Girsanov's theorem we obtain that

$$\begin{aligned} & \mathbb{E} \left[ \left( \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x;q)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ &= C_1(q) \mathbb{E} \left[ e^{\frac{\alpha}{2}F_\tau(0)} \mathcal{Q}(q) \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right], \end{aligned}$$

where

$$\mathcal{Q}(q) := \exp \left( \sqrt{2} \sum_{m,n=1}^{\infty} q^{2nm} (\alpha_{n,m} \alpha_n + \beta_{n,m} \beta_n) - \sum_{n=1}^{\infty} \left( \sum_{m=1}^{\infty} q^{nm} \alpha_{n,m} \right)^2 - \sum_{n=1}^{\infty} \left( \sum_{m=1}^{\infty} q^{nm} \beta_{n,m} \right)^2 \right).$$

The function above can be written as  $\mathcal{Q}(q) =: 1 + q^2 \Theta(q)$  where  $\mathbb{E}[\Theta(q)^{p_2}] < \infty$  for any  $p_2 > 1$ . Suppose we choose  $p_1, p_2, p_3 \in (1, \infty)$  such that  $\frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} = 1$  and

$$\mathbb{E} \left[ \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha p_3}{\gamma}} \right] < \infty.$$

Then, by Hölder's inequality, we have

$$\begin{aligned} & \mathbb{E} \left[ \left| e^{\frac{\alpha}{2}F_\tau(0)} \Theta(q) \right| \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ & \leq \mathbb{E} \left[ \left| e^{\frac{\alpha}{2}F_\tau(0)} \right|^{p_1} \right]^{\frac{1}{p_1}} \mathbb{E} \left[ \left| \Theta(q) \right|^{p_2} \right]^{\frac{1}{p_2}} \mathbb{E} \left[ \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha p_3}{\gamma}} \right]^{\frac{1}{p_3}}. \end{aligned}$$

Now we choose a sufficiently small  $r_0$  such that for all  $|q| < r_0$  we obtain the inequality

$$\begin{aligned} & \mathbb{E} \left[ e^{\frac{\alpha}{2}F_\tau(0)} \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ & \geq 2q \mathbb{E} \left[ \left| e^{\frac{\alpha}{2}F_\tau(0)} \right|^{p_1} \right]^{\frac{1}{p_1}} \mathbb{E} \left[ \left| \Theta(q) \right|^{p_2} \right]^{\frac{1}{p_2}} \mathbb{E} \left[ \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha p_3}{\gamma}} \right]^{\frac{1}{p_3}}. \end{aligned}$$

As a result, for all  $|q| < r_0$ , we have

$$\begin{aligned} & \mathbb{E} \left[ e^{\frac{\alpha}{2}F_\tau(0)} \mathcal{Q}(q) \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ & \geq \frac{1}{2} \mathbb{E} \left[ e^{\frac{\alpha}{2}F_\tau(0)} \left( \int_0^1 (-2 \sin(\pi x))^{\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2}Y(x)} dx \right)^{-\frac{\alpha}{\gamma}} \right]. \end{aligned}$$

The above display shows that  $\log \mathcal{G}_{\gamma,P}^\alpha(q)$  is well defined for all  $|q| < r_0$ . Furthermore, repeating the computations in the proof of Proposition (4.2) for the present case, it is immediate to see that  $\gamma^2 \log \mathcal{G}_{\gamma,P}^\alpha(q)$  uniformly converges as  $\gamma \rightarrow 0$  for all  $|q| < r_0$ . Therefore, by Morera's theorem, the semi-classical limit of the undeformed conformal block  $\lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{G}_{\gamma,P}^\alpha(q)$  is analytic for all  $|q| < r_0$ . In other words, it can be expressed as a convergent power series in  $q$  for all  $|q| < r_0$ .  $\square$

**5.2. Proof of Theorem 1.1.** Theorem 5.1 proves existence of the semi-classical undeformed conformal block. Combining the expressions (4.38) and (4.39) proves (1.1), with (1.2) being the Lamé equation and the accessory parameter given by the solution as (4.40). This completes the proof of Theorem 1.1.

## 6. ASYMPTOTICS OF SEMI-CLASSICAL CONFORMAL BLOCK AS $\tau \rightarrow \mathbf{i}\infty$

In this section, we study the asymptotic behaviour of the deformed conformal blocks  $\psi_\chi^\alpha(z, q)$  (for  $\chi = \frac{\gamma}{2}$  and  $\chi = \frac{2}{\gamma}$ ) and the partition function  $Z_{\gamma, P}^\alpha(q)$  (see Definition 2.2) by setting  $q \rightarrow 0$  in the semi-classical limit  $\gamma \rightarrow 0$ . These results played a major role in Section 4 to prove the semi-classical limit of the deformed conformal blocks. In order to find the desired asymptotics of this section, a key role will be played by the representation of the partition function  $Z_{\gamma, P}^\alpha$  in terms of double Gamma functions (see Proposition 2.1).

We find the semi-classical asymptotics of the partition function  $Z_{\gamma, P}^\alpha(q)$  in Theorem 6.1 around  $q = 0$ . This is used in Theorem 6.2 and Theorem 6.3 to show the semi-classical asymptotics of the deformed conformal blocks  $\psi_{\gamma/2}^\alpha(z, q)$  and  $\psi_{2/\gamma}^\alpha(z, q)$  by first setting  $q = 0$ . In what follows, we presents the ingredients of finding the asymptotics of Theorem 6.1 in few lemmas. We start by noting that the function  $\mathcal{A}_{\gamma, P}(\alpha)$ , which is related to the partition function  $Z_{\gamma, P}^\alpha(q)$  as shown in (2.12), can be expressed in the following way by comparing (2.10) and (2.13):

$$\begin{aligned} \mathcal{A}_{\gamma, P}(\alpha) &= \lim_{q \rightarrow 0} \mathbb{E} \left[ \left( q^{\frac{\alpha\gamma}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha\gamma}{2}} e^{\pi\gamma Px} e^{\frac{\gamma}{2} Y(x; q)} dx \right)^{-\frac{\alpha}{\gamma}} \right] \\ &= e^{\frac{\mathbf{i}\pi\alpha^2}{2}} \left( \frac{\gamma}{2} \right)^{\frac{\gamma\alpha}{2}} e^{-\frac{\pi\alpha P}{2}} \Gamma \left( 1 - \frac{\gamma^2}{4} \right)^{\frac{\alpha}{\gamma}} \frac{\Gamma_{\frac{\gamma}{2}}(Q - \frac{\alpha}{2}) \Gamma_{\frac{\gamma}{2}}(\frac{2}{\gamma} + \frac{\alpha}{2}) \Gamma_{\frac{\gamma}{2}}(Q - \frac{\alpha}{2} - \mathbf{i}P) \Gamma_{\frac{\gamma}{2}}(Q - \frac{\alpha}{2} + \mathbf{i}P)}{\Gamma_{\frac{\gamma}{2}}(\frac{2}{\gamma}) \Gamma_{\frac{\gamma}{2}}(Q - \mathbf{i}P) \Gamma_{\frac{\gamma}{2}}(Q + \mathbf{i}P) \Gamma_{\frac{\gamma}{2}}(Q - \alpha)}. \end{aligned} \quad (6.1)$$

We first find the asymptotics of  $\mathcal{A}_{\gamma, P}(\alpha)$  in Proposition 6.1. To this end, We first observe the following asymptotics for the double Gamma functions.

**Lemma 6.1.** *For any  $\xi \in \mathbb{C}$  such that  $\arg(\xi) < \pi$ , we have*

$$\begin{aligned} \lim_{\gamma \rightarrow 0} \left[ \gamma^2 \log \Gamma_{\frac{\gamma}{2}} \left( \frac{2\xi}{\gamma} \right) - \gamma^2 \log \frac{\gamma}{2} \int_0^{\operatorname{Re}(\xi)} (\xi - x) dx - \gamma^2 \log \Gamma_{\frac{\gamma}{2}} \left( \frac{2\mathbf{i}\operatorname{Im}(\xi)}{\gamma} \right) \right] \\ = \begin{cases} -4\operatorname{Re}(\xi) \log \sqrt{2\pi} + \int_{-\infty}^{\operatorname{Re}(\xi)} \log \Gamma(\xi - x) dx & \text{if } \operatorname{Re}(\xi) > 0, \\ 4\operatorname{Re}(\xi) \log \sqrt{2\pi} + \int_{\operatorname{Re}(\xi)}^0 \log \Gamma(\xi + x) dx & \text{if } \operatorname{Re}(\xi) < 0. \end{cases} \end{aligned} \quad (6.2)$$

Here  $\log \Gamma(\cdot)$  is taken w.r.t. the principle branch of logarithm.

*Proof.* We prove the above statement for  $\operatorname{Re}(\xi) > 0$ . The remaining case follows from a similar argument. From the identities of the double Gamma function, we obtain the following shift relation for the function  $\Gamma_{\frac{\gamma}{2}}(\cdot)$  for any  $\operatorname{Re}(\xi) > 0$  [Pon04]:

$$\Gamma_{\frac{\gamma}{2}}(z + \chi) = \sqrt{2\pi} \frac{\chi^{\chi z - \frac{1}{2}}}{\Gamma(\chi z)} \Gamma_{\frac{\gamma}{2}}(z), \quad \chi = \left\{ \frac{2}{\gamma}, \frac{\gamma}{2} \right\}, \quad (6.3)$$

which can also be expressed as

$$\Gamma_{\frac{\gamma}{2}} \left( \frac{2\xi}{\gamma} - (n+1) \frac{\gamma}{2} \right) = \frac{1}{\sqrt{2\pi}} \left( \frac{\gamma}{2} \right)^{-\xi + (n+1) \frac{\gamma^2}{4} + \frac{1}{2}} \Gamma(\xi - (n+1) \frac{\gamma^2}{4}) \Gamma_{\frac{\gamma}{2}} \left( \frac{2\xi}{\gamma} - n \frac{\gamma}{2} \right).$$

Applying the above relation for all  $n$  such that  $n \frac{\gamma^2}{4} \leq \operatorname{Re}(\xi)$ , we get

$$\Gamma_{\frac{\gamma}{2}} \left( \frac{2\mathbf{i}\operatorname{Im}(\xi)}{\gamma} \right) = \prod_{n=1}^{\lfloor \frac{4\operatorname{Re}(\xi)}{\gamma^2} \rfloor} \frac{1}{\sqrt{2\pi}} \left( \frac{\gamma}{2} \right)^{-\xi + (n+1) \frac{\gamma^2}{4} + \frac{1}{2}} \Gamma(\xi - (n+1) \frac{\gamma^2}{4}) \times \Gamma_{\frac{\gamma}{2}} \left( \frac{2\xi}{\gamma} \right).$$

Taking logarithms on both sides of the above equation, multiplying by  $\gamma^2$  and taking the limit  $\gamma \rightarrow 0$ , we arrive at (6.2).  $\square$

In the next result, we utilize this asymptotics to find semi-classical limit of  $\mathcal{A}_{\gamma,P}(\alpha)$ .

**Proposition 6.1.** *Consider the scaling  $\alpha = \frac{\alpha_0}{\gamma}$ ,  $P = \frac{P_0}{\gamma}$  with  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ . As  $\gamma \rightarrow 0$ , the semi-classical limit of  $\mathcal{A}_{\gamma,P}(\alpha)$  has the following expression:*

$$\begin{aligned} \mathcal{B}(P_0, \alpha_0) &:= \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{A}_{\gamma,P}(\alpha) \\ &= \frac{i\pi\alpha_0(\alpha_0 + iP_0)}{2} + 4\alpha_0\sqrt{2\pi} - \int_{\operatorname{Re}(1-\frac{\alpha_0}{4})}^{\operatorname{Re}(1)+\frac{\alpha_0}{4}} \log \Gamma\left(1-x-\frac{\alpha_0}{4}\right) dx \\ &\quad - \int_{\operatorname{Re}(1+\frac{\alpha_0}{4})}^{\operatorname{Re}(1+\frac{\alpha_0}{2})+\frac{\alpha_0}{4}} \log \Gamma\left(1-x+\frac{\alpha_0}{4}\right) dx - \int_{\operatorname{Re}(1-\frac{iP_0}{2}-\frac{\alpha_0}{4})}^{\operatorname{Re}(1-\frac{iP_0}{2})+\frac{\alpha_0}{4}} \log \Gamma\left(1-x-\frac{iP_0}{2}-\frac{\alpha_0}{4}\right) dx \\ &\quad - \int_{\operatorname{Re}(1+\frac{iP_0}{2}-\frac{\alpha_0}{4})}^{\operatorname{Re}(1+\frac{iP_0}{2})+\frac{\alpha_0}{4}} \log \Gamma\left(1-x+\frac{iP_0}{2}-\frac{\alpha_0}{4}\right) dx. \end{aligned} \quad (6.4)$$

*Proof.* With the scaling  $\alpha = \alpha_0/\gamma$ ,  $P = P_0/\gamma$ , and using the shift relations of the double gamma function as noted in (6.3) with  $\chi = \frac{\gamma}{2}$ , the expression (6.1) simplifies as

$$\begin{aligned} &\mathcal{A}_{\gamma,P_0}(\alpha_0) \\ &= e^{\frac{i\pi\alpha_0(\alpha_0+iP_0)}{2\gamma^2}} \Gamma\left(1-\frac{\gamma^2}{4}\right)^{\frac{\alpha_0}{\gamma^2}} \frac{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{iP_0}{\gamma}-\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{iP_0}{\gamma}-\frac{\alpha_0}{2\gamma}\right)}{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{\alpha_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{iP_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{iP_0}{\gamma}\right)} \\ &\quad \times \frac{\Gamma\left(1-\frac{\alpha_0}{2}\right) \Gamma\left(1-\frac{iP_0}{2}\right) \Gamma\left(1+\frac{iP_0}{2}\right)}{\Gamma\left(1-\frac{\alpha_0}{4}\right) \Gamma\left(1-\frac{\alpha_0}{4}-\frac{iP_0}{2}\right) \Gamma\left(1-\frac{\alpha_0}{4}+\frac{iP_0}{2}\right)}. \end{aligned}$$

Taking logarithm of both sides of the above equation, multiplying both sides of the equation with  $\gamma$  and letting  $\gamma \rightarrow 0$  reads

$$\begin{aligned} &\lim_{\gamma \rightarrow 0} \gamma^2 \lim_{\gamma \rightarrow 0} \mathcal{A}_{\gamma,P_0}(\alpha_0) \\ &= \frac{i\pi\alpha_0(\alpha_0 + iP_0)}{2} + \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \frac{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{iP_0}{\gamma}-\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{iP_0}{\gamma}-\frac{\alpha_0}{2\gamma}\right)}{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{\alpha_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}-\frac{iP_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}+\frac{iP_0}{\gamma}\right)} \right). \end{aligned} \quad (6.5)$$

Observe that the last term in the expression above can be written as a product of ratios of the form  $\frac{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}(\xi-\alpha_0/4)\right)}{\Gamma_{\frac{\gamma}{2}}\left(\frac{2\xi}{\gamma}\right)}$ . It is therefore sufficient to study the term

$$\begin{aligned} &\lim_{\gamma \rightarrow 0} \gamma^2 \log \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}(\xi-\alpha_0/4)\right) - \lim_{\gamma \rightarrow 0} \gamma^2 \log \Gamma_{\frac{\gamma}{2}}\left(\frac{2\xi}{\gamma}\right) \\ &\stackrel{(6.2)}{=} \alpha_0\sqrt{2\pi} + \int_{-\infty}^{\operatorname{Re}(\xi-\frac{\alpha_0}{4})} \log \Gamma\left(\xi-\frac{\alpha_0}{4}-x\right) dx - \int_{-\infty}^{\operatorname{Re}(\xi)} \log \Gamma(\xi-x) dx \\ &= \alpha_0\sqrt{2\pi} - \int_{\operatorname{Re}(\xi-\frac{\alpha_0}{4})}^{\operatorname{Re}(\xi)+\frac{\alpha_0}{4}} \log \Gamma\left(\xi-x-\frac{\alpha_0}{4}\right) dx. \end{aligned}$$

Simplifying the expression (6.5) using the identity above proves the statement of this proposition.  $\square$

**Theorem 6.1.** *For any  $|q| < r$  for some very small  $r > 0$ , partition function  $Z_{\gamma,P}^\alpha(q)$  show the following asymptotics as  $\gamma$  approaches 0,*

$$\lim_{\gamma \rightarrow 0} \gamma^2 \log Z_{\gamma,P}^\alpha(q) = -\alpha_0^2 \sum_{k=1}^{\infty} \log(1-q^{2k}) + \mathcal{B}(P_0, \alpha_0),$$

where  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ , with  $\mathcal{B}(P_0, \alpha_0)$  defined in Proposition 6.1.

*Proof.* Recall from (2.12) that

$$Z_{\gamma,P}^{\alpha} = q^{\frac{1}{12}(\frac{\alpha\gamma}{2} + \frac{\alpha^2}{2} - 1)} \eta(q)^{\alpha^2 + 1 - \frac{\alpha\gamma}{2}} \mathcal{A}_{\gamma,P}(\alpha).$$

Taking logarithm on both sides and multiplying by  $\gamma^2$  yields

$$\gamma^2 \log Z_{\gamma,P}^{\alpha} = \gamma^2 \left( \frac{\alpha_0^2}{24\gamma^2} + \frac{1}{12} \left( \frac{\alpha_0}{2} - 1 \right) \right) \log q + \gamma^2 \left( \frac{\alpha_0^2}{\gamma^2} + \left( \frac{\alpha_0}{2} - 1 \right) \right) \log \eta(q) + \gamma^2 \log \mathcal{A}_{\gamma,P}(\alpha)$$

where we have used the scaling rule  $\alpha = \alpha_0/\gamma$ . Letting  $\gamma \rightarrow 0$  both sides yields the desired conclusion.  $\square$

**Theorem 6.2.** *As  $q \rightarrow 0$ , the semi-classical deformed conformal block  $\lim_{\gamma \rightarrow 0} \gamma^2 \log \psi_{\gamma}^{\alpha}$  in (4.37) behaves as*

$$\begin{aligned} & \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\alpha}{4}(\chi - \alpha)} \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) \right) \\ &= -\frac{\alpha_0(\alpha_0 + 4)}{6} \log(-2\pi) + \left( \frac{P_0^2}{2} + \frac{\alpha_0(\alpha_0 + 2)}{24} \right) \log q + \mathcal{B}(P_0, \alpha_0), \end{aligned} \quad (6.6)$$

where  $\alpha_0 \in (-4, 2)$ ,  $P_0 \in \mathbb{R}$ , and  $\mathcal{B}(P_0, \alpha_0)$  is given by (6.4).

*Proof.* By using the explicit form of (4.37), we can obtain the following expression the following expression

$$\begin{aligned} \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\gamma}{4} \left( \frac{\gamma}{2} - \frac{\alpha_0}{\gamma} \right)} \psi_{\gamma/2}^{\alpha_0/\gamma}(z, q) \right) &= -\frac{\alpha_0(\alpha_0 + 4)}{6} \log(-2\pi) + \left( \frac{P_0^2}{2} + \frac{\alpha_0(\alpha_0 + 2)}{24} \right) \log q + \\ &+ \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx \right)^{-\frac{\alpha_0}{\gamma^2}} \right]. \end{aligned} \quad (6.7)$$

for the semiclassical limit from (2.14) and (4.28). In obtaining the final expression, we note that the  $\gamma \rightarrow 0$  and  $q \rightarrow 0$  limit of the following

$$\mathcal{T} := \gamma^2 \log \mathbb{E} \left[ \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x; q)} dx \right)^{-\frac{\alpha_0}{\gamma^2}} \right]$$

commute. To see this, using Girsanov's theorem, we write the above expression as (see [Gho+24, Eq. 2.225])

$$\mathcal{T} = (\alpha_0^2 - \alpha_0\gamma/2) \log \left( q^{-1/12} \eta(q) \right) + \gamma^2 \log \mathbb{E} \left[ e^{\frac{\alpha_0}{2\gamma} F_{\tau}(0)} \mathcal{Q}(q) \left( \int_0^1 (2 \sin \pi x)^{-\alpha_0/2} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y(x)} dx \right)^{-\frac{\alpha_0}{\gamma^2}} \right],$$

where  $F_{\tau}(0)$  and  $\mathcal{Q}(q)$  are defined as in (4.16). Using the above decomposition expression analogous and following the arguments after (4.20) in the proof of Proposition 4.2, one can prove the commutativity of the limits. As a consequence, we get

$$\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \mathcal{T} = \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathcal{A}_{\gamma,P}(\alpha) = \mathcal{B}(P_0, \alpha_0).$$

Taking  $q \rightarrow 0$  on both sides of (6.7) and substituting  $\mathcal{B}(P_0, \alpha_0)$  in place of  $\lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \mathcal{T}$  gives us (6.6).  $\square$

**Theorem 6.3.** *Consider  $z = a\tau + \mathbf{b}$  for some  $\mathbf{a}, \mathbf{b} \in (0, 1)$ ,  $\mathbf{a} \neq \mathbb{Z} + \frac{1}{2}$ . As  $\tau \rightarrow \mathbf{i}\infty$ , the semi-classical deformed conformal block  $\lim_{\gamma \rightarrow 0} \gamma^2 \log \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q)$  in (4.3) has the following behaviour*

with  $\alpha_0 \in (-2, 2)$ ,  $P_0 \in \mathbb{R}$ :

$$\begin{aligned} & \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(x-\alpha)} \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q) \right) \\ &= C(\tau) + 2\pi P_0 + \int_{\operatorname{Re}(\frac{2-3\alpha_0}{4})}^{\operatorname{Re}(-\frac{\alpha_0}{2}) + \frac{\alpha_0-2}{4}} \log \Gamma \left( -x + \frac{2-3\alpha_0}{4} \right) dx \\ &+ \int_{\operatorname{Re}(\frac{2+\alpha_0}{4})}^{\operatorname{Re}(\frac{\alpha_0}{2}) + \frac{\alpha_0-2}{4}} \log \Gamma \left( -x + \frac{2+\alpha_0}{4} \right) dx - \int_{\operatorname{Re}(\frac{iP_0}{2} + \frac{2-\alpha_0}{4}) + \frac{\alpha_0-2}{4}}^{\operatorname{Re}(\frac{iP_0}{2} + \frac{2-\alpha_0}{2})} \log \Gamma \left( \frac{iP_0}{2} - x + \frac{2-\alpha_0}{2} \right) dx \\ &- \int_{\operatorname{Re}(-\frac{iP_0}{2})}^{\operatorname{Re}(-\frac{iP_0}{2} - \frac{2-\alpha_0}{4}) + \frac{\alpha_0-2}{4}} \log \Gamma \left( -\frac{iP_0}{2} - x \right) dx, \end{aligned}$$

where

$$\begin{aligned} \mathcal{C}(\tau) := & \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12} \log(-2\pi) + i\pi(2 - \alpha_0) + \left( \frac{\alpha_0(\alpha_0 - 10)}{24} + \frac{P_0^2}{2} + \frac{1}{6} \right) \log q \\ & - 2\pi i(2 - \alpha_0)(a\tau + b) - 4P_0(a\tau + b). \end{aligned} \quad (6.8)$$

*Proof.* Combining (2.14) with (4.10), we obtain the following expression for  $\chi = \frac{2}{\gamma}$ :

$$\begin{aligned} & \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(x-\alpha)} \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q) \right) \\ &= \frac{-\alpha_0^2 + \alpha_0 + 6P_0^2 + 2}{12} \log(-2\pi) + \left( \frac{\alpha_0^2 - 10\alpha_0 + 12P_0^2 + 16}{24} \right) \log q + 2P_0 z \\ &+ \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( q^{\frac{\alpha_0-2}{8}} \int_0^1 \theta_1(z+x) \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi P_0 x} e^{\frac{\gamma}{2} Y_\tau(x)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right]. \end{aligned} \quad (6.9)$$

We identify  $z$  with the solution of the non-autonomous elliptic Calogero-Moser model  $u(\tau)$  by letting  $z = 2u(\tau)$ . As  $\tau \rightarrow i\infty$ , the variable  $u(\tau) = a\tau + b$ , with  $\mathbf{a}, \mathbf{b}$  parameterization the A, B-cycle monodromies of the linear system associated to the non-autonomous elliptic Calogero-Moser model (see Appendix C). From (A.2), we then obtain the following behaviour of the theta function in the same limit:

$$\begin{aligned} \theta_1(z) \equiv \theta_1(2u(\tau)) &= -2q^{1/4} \sin(2\pi u(\tau)) \prod_{k=1}^{\infty} (1 - q^{2k})(1 - 2\cos(4\pi u(\tau))q^{2k} + q^{4k}) \\ &= iq^{1/4} e^{-2i\pi(a\tau+b)} (1 + \mathcal{O}(q^2)), \end{aligned}$$

and resultantly, as  $\tau \rightarrow i\infty$ ,

$$\theta_1(z+x) = iq^{1/4} e^{-2i\pi(a\tau+b)-i\pi x} (1 + \mathcal{O}(q^2)).$$

Plugging the above expressions into the right hand side of (6.9) shows that (see also [Gho+24, Lemma 5.3])

$$\begin{aligned} & \lim_{\tau \rightarrow i\infty} \left[ \lim_{\gamma \rightarrow 0} \gamma^2 \log \left( \theta_1(z)^{\frac{\chi}{2}(x-\alpha)} \psi_{2/\gamma}^{\alpha_0/\gamma}(z, q) \right) - \mathcal{C}(\tau) \right] \\ &= \lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi x(P_0-i)} e^{\frac{\gamma}{2} Y(x;q)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right], \end{aligned} \quad (6.10)$$

where  $C(\tau)$  is defined in (6.8). Due to Proposition 4.2, the limits  $\lim_{q \rightarrow 0}$  and  $\lim_{\gamma \rightarrow 0}$  commute. So, we first begin with the  $q \rightarrow 0$  limit

$$\mathbb{E} \left[ \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi x(P_0-i)} e^{\frac{\gamma}{2} Y(x;q)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right] = \mathbb{E} \left[ \left( \int_0^1 (\sin \pi x)^{-\frac{\alpha_0}{2}} e^{\pi x(P_0-i)} e^{\frac{\gamma}{2} Y(x;q)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right].$$

We now note the following identity [Ang+24]

$$\begin{aligned} & \mathbb{E} \left[ \left( \int_0^1 (\sin \pi x)^{-\frac{\alpha_0}{2}} e^{\pi x(P_0 - i)} e^{\frac{\gamma}{2} Y(x; q)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right] \\ &= e^{2\pi P_0/\gamma^2} \frac{\Gamma\left(\frac{\alpha_0}{4} - \frac{\gamma^2}{4}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(-\frac{\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{iP_0}{\gamma} + \frac{2-\alpha_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(-\frac{iP_0}{\gamma}\right)}{\Gamma\left(1 - \frac{\gamma^2}{4}\right) \frac{2}{\gamma} \left(\frac{2-\alpha_0}{2\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{1}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{1-\alpha_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(\frac{iP_0}{\gamma} + \frac{2-\alpha_0}{\gamma}\right) \Gamma_{\frac{\gamma}{2}}\left(-\frac{iP_0}{\gamma} - \frac{2-\alpha_0}{\gamma}\right)} \end{aligned} \quad (6.11)$$

To simplify the above equation, we first note that the expression above can be written in terms of ratios of the form

$$\frac{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\left(\xi + \frac{2-\alpha_0}{4}\right)\right)}{\Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\xi\right)}.$$

We then use the identity

$$\begin{aligned} & \lim_{\gamma \rightarrow 0} \gamma^2 \log \Gamma_{\frac{\gamma}{2}}\left(\frac{2}{\gamma}\left(\xi + \frac{2-\alpha_0}{4}\right)\right) - \lim_{\gamma \rightarrow 0} \gamma^2 \log \Gamma_{\frac{\gamma}{2}}\left(\frac{2\xi}{\gamma}\right) \\ & \stackrel{(6.2)}{=} (\alpha_0 - 2) \sqrt{2\pi} + \int_{-\infty}^{\operatorname{Re}(\xi + \frac{2-\alpha_0}{4})} \log \Gamma\left(\xi + \frac{2-\alpha_0}{4} - x\right) dx - \int_{-\infty}^{\operatorname{Re}(\xi)} \log \Gamma(\xi - x) dx \\ & = (\alpha_0 - 2) \sqrt{2\pi} - \int_{\operatorname{Re}(\xi + \frac{2-\alpha_0}{4})}^{\operatorname{Re}(\xi) + \frac{\alpha_0 - 2}{4}} \log \Gamma\left(\xi - x + \frac{2-\alpha_0}{4}\right) dx. \end{aligned}$$

Then, we repeat the procedure used in the proof of Proposition 6.1. One can obtain similar expressions to the one above for the remaining terms in (6.11). Substituting them in (6.10), we obtain the expression

$$\begin{aligned} & \lim_{q \rightarrow 0} \lim_{\gamma \rightarrow 0} \gamma^2 \log \mathbb{E} \left[ \left( q^{\frac{\alpha_0}{8}} \int_0^1 \theta_1(x)^{-\frac{\alpha_0}{2}} e^{\pi x(P_0 - i)} e^{\frac{\gamma}{2} Y(x)} dx \right)^{\frac{(2-\alpha_0)}{\gamma^2}} \right] \\ & \stackrel{(6.1), (6.4)}{=} 2\pi P_0 + \int_{\operatorname{Re}(\frac{2-3\alpha_0}{4})}^{\operatorname{Re}(-\frac{\alpha_0}{2}) + \frac{\alpha_0 - 2}{4}} \log \Gamma\left(-x + \frac{2-3\alpha_0}{4}\right) dx \\ & + \int_{\operatorname{Re}(\frac{2+\alpha_0}{4})}^{\operatorname{Re}(\frac{\alpha_0}{2}) + \frac{\alpha_0 - 2}{4}} \log \Gamma\left(-x + \frac{2+\alpha_0}{4}\right) dx - \int_{\operatorname{Re}(\frac{iP_0}{2} + \frac{2-\alpha_0}{2})}^{\operatorname{Re}(\frac{iP_0}{2} + \frac{2-\alpha_0}{4}) + \frac{\alpha_0 - 2}{4}} \log \Gamma\left(\frac{iP_0}{2} - x + \frac{2-\alpha_0}{2}\right) dx \\ & - \int_{\operatorname{Re}(-\frac{iP_0}{2})}^{\operatorname{Re}(-\frac{iP_0}{2} - \frac{2-\alpha_0}{4}) + \frac{\alpha_0 - 2}{4}} \log \Gamma\left(-\frac{iP_0}{2} - x\right) dx, \end{aligned}$$

completing the proof of this Theorem. □

## APPENDIX A. USEFUL NOTATION AND ELLIPTIC FUNCTIONS

The parameter  $\gamma \in (0, 2)$  defines the central charge of the CFT as

$$c = 1 + 6Q^2, \quad Q := \frac{\gamma}{2} + \frac{2}{\gamma}. \quad (\text{A.1})$$

We use two other parameters

$$l_\chi = \frac{\chi^2}{2} - \frac{\alpha\chi}{2}, \quad \chi = \left\{ \frac{2}{\gamma}, \frac{\gamma}{2} \right\}.$$

The modular parameter  $\tau \in \mathbb{H}$  and the nome is defined as  $q := e^{i\pi\tau}$ . The elliptic theta functions  $\theta_i(z) = \theta_i(z|\tau)$  are defined as [DLMF, 20.5(i)]

$$\theta_1(z) = -2q^{1/4} \sin(\pi z) \prod_{k=1}^{\infty} (1 - q^{2k})(1 - 2\cos(2\pi z)q^{2k} + q^{4k}), \quad (\text{A.2})$$

$$\theta_2(z) = 2q^{1/4} \cos(\pi z) \prod_{k=1}^{\infty} (1 - q^{2k})(1 + 2\cos(2\pi z)q^{2k} + q^{4k}) \quad (\text{A.3})$$

$$\theta_3(z) = \prod_{k=1}^{\infty} (1 - q^{2k})(1 + 2\cos(2\pi z)q^{2k-1} + q^{4k}) \quad (\text{A.4})$$

$$\theta_1'(0) = -2\pi q^{1/4} \prod_{k=1}^{\infty} (1 - q^{2k})^3 = -2\pi\eta(q)^3. \quad (\text{A.5})$$

They have the following periodicity properties

$$\theta_1(z+1) = -\theta_1(z), \quad \theta_1(z+\tau) = -q^{-1}e^{-2\pi iz}\theta_1(z).$$

The Dedekind  $\eta$  function is defined as

$$\eta(q) := q^{\frac{1}{12}} \prod_{k=1}^{\infty} (1 - q^{2k}),$$

and has the following relation to the theta function

$$\eta(\tau) = \left( \frac{\theta_1'(\tau)}{2\pi} \right)^{1/3}.$$

The Weierstrass cubic is

$$(\wp'(z))^2 = 4\wp^3(z) - g_2\wp(z) - g_3.$$

The Weierstrass  $\wp$ -function solves the cubic above and is given by [DLMF, 23.2(ii)]

$$\wp(z) = \frac{1}{z^2} + \sum_{w \in \mathbb{L} \setminus \{0\}} \left( \frac{1}{(z-w)^2} - \frac{1}{w^2} \right), \quad (\text{A.6})$$

where  $\mathbb{L}$  denotes the lattice. The above function has a double pole at  $z = 0$ , and is doubly periodic

$$\wp(z+1) = \wp(z), \quad \wp(z+\tau) = \wp(z).$$

## APPENDIX B. NON-AUTONOMOUS ELLIPTIC CALOGERO-MOSER MODEL

**B.1. Hamiltonian and the action.** The 2-particle nonautonomous elliptic Calogero-Moser system is represented by the Hamiltonian<sup>1</sup>

$$H(\tau) = v(\tau)^2 - m^2\wp(2u(\tau)|\tau), \quad (\text{B.1})$$

and the equations of motion read

$$v(\tau) = 2\pi\mathbf{i} \frac{dv(\tau)}{d\tau}, \quad (2\pi\mathbf{i}) \frac{dv(\tau)}{d\tau} = m^2\wp'(2u|\tau). \quad (\text{B.2})$$

The above equations of motion give the elliptic form of the Painlevé VI equation for special values of the parameters (see [Des21, Chapter 1], [Tak01]).

The equation (B.2) is *integrable* in the sense that it has an associated *Lax pair*

$$\begin{aligned} \partial_z Y(z, \tau) &:= L(z, \tau)Y(z, \tau) = \begin{pmatrix} v(\tau) & mx(2u(\tau), z) \\ mx(-2u(\tau), z) & -v(\tau) \end{pmatrix} Y(z, \tau), \\ 2\pi\mathbf{i}\partial_\tau Y(z, \tau) &:= M(z, \tau)Y(z, \tau) = m \begin{pmatrix} 0 & y(2u(\tau), z) \\ y(-2u(\tau), z) & 0 \end{pmatrix} Y(z, \tau), \end{aligned} \quad (\text{B.3})$$

<sup>1</sup>Note that the Hamiltonian below differs by the one in [DDG23] by a normalization factor of  $2m^2\eta_1(\tau)$ .

and (B.2) is given by the consistency condition

$$2\pi\mathbf{i}\partial_z\partial_\tau Y(z, \tau) = 2\pi\mathbf{i}\partial_z\partial_\tau Y(z, \tau) \Rightarrow \partial_\tau L(z, \tau) - \partial_z M(z, \tau) + [L(z, \tau), M(z, \tau)] = 0.$$

In the above expressions, the functions  $x(\xi, z)$ ,  $y(\xi, z)$  (B.3) are defined as

$$x(\xi, z) := \frac{\theta_1(z - \xi|\tau)\theta_1'(0|\tau)}{\theta_1(z|\tau)\theta_1(\xi|\tau)}, \quad y(\xi, z) := \partial_\xi x(\xi, z).$$

Furthermore, the Hamiltonian (B.1) is given by the A-cycle contour integral

$$H(\tau) = \oint_A dz \frac{1}{2} \operatorname{tr} L^2(z, \tau) = v(\tau)^2 - m^2 \wp(2u(\tau)).$$

The functions  $(u(\tau), v(\tau))$  are the canonical coordinates of the system and they describe the position and momentum of a particle in an elliptic potential respectively. Therefore, the Lagrangian of the above system is given by

$$\mathcal{L} = v(\tau)^2 + m^2 \wp(2u(\tau)),$$

and the action functional is defined as the integral of the Lagrangian

$$S(\tau) := \int^\tau (v(\tau')^2 + m^2 \wp(2u(\tau'))) \frac{d\tau'}{2\pi\mathbf{i}}.$$

It is a known fact from classical mechanics that the action solves the Hamilton-Jacobi equation

$$2\pi\mathbf{i}\partial_\tau S = (\partial_{u(\tau)} S)^2 - m^2 \wp(2u(\tau)) S.$$

**B.2. Monodromy data.** The monodromy data of the linear system (B.3) is obtained by observing that  $L, M$  are defined on a torus with a simple pole at  $z = 0$  and satisfy the relations

$$L(z + 1, \tau) = L(z, \tau), \quad L(z + \tau, \tau) = e^{2\pi\mathbf{i}u(\tau)\sigma_3} L(z, \tau) e^{-2\pi\mathbf{i}u(\tau)\sigma_3}.$$

Subsequently, the solution  $Y(z, \tau)$  (B.3) has the following monodromy properties around the A, B cycles of the torus and around the puncture  $z = 0$ :

$$Y(z + 1, \tau) = M_A Y(z, \tau), \quad Y(z + \tau, \tau) = M_B Y(z, \tau) e^{2\pi\mathbf{i}u(\tau)\sigma_3},$$

$$Y(e^{2\pi\mathbf{i}} z, \tau) = M_0 Y(z, \tau),$$

with the constraint

$$M_0 = M_A^{-1} M_B^{-1} M_A M_B.$$

Choosing the matrix  $M_A$  to be diagonal, one obtains the following explicit expressions for the monodromy matrices

$$M_A = e^{2\pi\mathbf{i}a\sigma_3}, \quad M_0 \sim e^{2\pi\mathbf{i}m\sigma_3}, \quad M_B = \begin{pmatrix} \frac{\sin \pi(2(a-m))}{\sin 2\pi a} e^{-\mathbf{i}\nu/2} & \frac{\sin \pi m}{\sin 2\pi a} \\ -\frac{\sin \pi m}{\sin 2\pi a} & \frac{\sin \pi(2(a+m))}{\sin 2\pi a} e^{\mathbf{i}\nu/2} \end{pmatrix}, \quad (\text{B.4})$$

where  $\sim$  denotes the conjugacy class,  $\sigma_3$  is the Pauli sigma matrix, and  $m$  is the parameter of the equation (B.2).

**B.3. Asymptotic behaviour of  $u(\tau)$ .** For the sake of completeness, in this section we detail the asymptotics of the solution of the NAECM model  $u(\tau)$  for  $\tau \rightarrow \mathbf{i}\infty$ , and  $\tau \rightarrow \tau_\star$  that were also computed in [Bon+20; BGG22].

**Proposition B.1.** *In the limit  $\tau \rightarrow \mathbf{i}\infty$ , the solution of the equation (B.2)  $u(\tau)$  behaves as*

$$u(\tau) = a\tau + \mathbf{b} + \frac{m^2}{8\pi\mathbf{i}a^2} e^{4\pi\mathbf{i}(a\tau+\mathbf{b})} + \mathcal{O}(q e^{4\pi\mathbf{i}(a\tau+\mathbf{b})}),$$

where  $q = e^{\mathbf{i}\pi\tau}$ ,  $\mathbf{a}, m$  parameterize the monodromies around the A-cycle and the puncture respectively (B.4), and the monodromy around the B-cycle is related to  $\mathbf{b}$  as

$$e^{2\pi\mathbf{i}\mathbf{b}} := e^{\mathbf{i}\nu/2} \frac{\Gamma(2\mathbf{a} - m)\Gamma(1 - 2\mathbf{a})}{\Gamma(2\mathbf{a})\Gamma(1 - m - 2\mathbf{a})}. \quad (\text{B.5})$$

*Proof.* Our starting point is the following identity satisfied by the function  $u(\tau)$  that was shown in [Bon+20, equation (3.56)] (see also [Des22]):

$$\frac{\theta_3(2u|2\tau)}{\theta_2(2u|2\tau)} = \frac{Z_0^D(\tau)}{Z_{1/2}^D(\tau)}, \quad (\text{B.6})$$

where  $Z_{0,1/2}^D$  are dual Nekrasov-Okounkov functions with integer and half integer shifts and are defined as

$$Z_{\epsilon/2}^D := \sum_{n \in \mathbb{Z} + \frac{\epsilon}{2}} C_n e^{in\nu} q^{2(a+n)^2} \mathcal{B}(a+n, m, q).$$

The coefficients

$$C_n = \frac{G(1-m+2(a+n))G(1-m-2(a+n))}{G(1+2(a+n))G(1-2(a+n))} \times \frac{G(1+2a)G(1-2a)}{G(1-m+2a)G(1-m-2a)},$$

and the combinatorial term  $\mathcal{B}(\cdot)$  is given by

$$\mathcal{B}(a, m, q) = \prod_{n=1}^{\infty} (1-q^{2n})^{1-2m^2} \sum_{Y_+, Y_-} q^{2(|Y_+|+|Y_-|)} \prod_{\epsilon, \epsilon' = \pm} \frac{N_{Y_\epsilon, Y_{\epsilon'}}(m + (\epsilon - \epsilon')a)}{N_{Y_\epsilon, Y_{\epsilon'}}((\epsilon - \epsilon')a)},$$

where

$$N_{\lambda, \mu}(x) = \prod_{s \in \lambda} (x + a_\lambda(s) + l_\mu(s) + 1) \prod_{t \in \mu} (x - a_\mu(t) - l_\lambda(t) - 1),$$

with  $a_Y(\square)$  and  $l_Y(\square)$  denoting the arm and leg length of the box  $\square$  in the Young diagram  $Y$  respectively. In the limit  $\tau \rightarrow i\infty$ , the RHS of (B.6) reads

$$\frac{Z_0^D(\tau)}{Z_{1/2}^D(\tau)} = q^{-1/2} \left[ \frac{C_0}{C_{-1/2} e^{-i\nu/2} q^{-2a} + C_{1/2} e^{i\nu/2} q^{2a}} + \mathcal{O}(q) \right]. \quad (\text{B.7})$$

The coefficients  $C_0, C_{\pm 1/2}$  can be simplified using the relation  $G(z+1) = \Gamma(z)G(z)$ :

$$\begin{aligned} C_0 &= \frac{G(1-m+2a)G(1-m-2a)}{G(1+2a)G(1-2a)} \times \frac{G(1+2a)G(1-2a)}{G(1-m+2a)G(1-m-2a)} = 1, \\ C_{-1/2} &= \frac{G(1-m+2(a-1/2))G(1-m-2(a-1/2))}{G(1+2(a-1/2))G(1-2(a-1/2))} \times \frac{G(1+2a)G(1-2a)}{G(1-m+2a)G(1-m-2a)} \\ &= \frac{\Gamma(2a)\Gamma(1-m-2a)}{\Gamma(2a-m)\Gamma(1-2a)}, \end{aligned} \quad (\text{B.8})$$

$$\begin{aligned} C_{1/2} &= \frac{G(1-m+2(a+1/2))G(1-m-2(a+1/2))}{G(1+2(a+1/2))G(1-2(a+1/2))} \times \frac{G(1+2a)G(1-2a)}{G(1-m+2a)G(1-m-2a)} \\ &= \left(1 - \frac{m^2}{4a^2}\right) \frac{\Gamma(2a-m)\Gamma(1-2a)}{\Gamma(1-m-2a)\Gamma(2a)}. \end{aligned} \quad (\text{B.9})$$

Substituting (B.8), (B.9), (B.5) in (B.7) we get that

$$\begin{aligned} \frac{Z_0^D(\tau)}{Z_{1/2}^D(\tau)} &= q^{-1/2} \left[ \frac{1}{e^{-2\pi i\beta} q^{-2a} + e^{2\pi i\beta} q^{2a}} + \mathcal{O}(q) \right] \\ &= \frac{q^{-1/2}}{2 \cos(2\pi(a\tau + b))} (1 + \mathcal{O}(\cos(4\pi(a\tau + b)))). \end{aligned} \quad (\text{B.10})$$

Now for the LHS of (B.6), substituting the series expansions of the theta functions (A.3), (A.4), we see that

$$\frac{\theta_3(2u|2\tau)}{\theta_2(2u|2\tau)} = \frac{q^{-1/2}}{2 \cos(2\pi u)} (1 + \mathcal{O}(q^2)). \quad (\text{B.11})$$

Comparing the RHS (B.10) and the LHS (B.11),

$$\begin{aligned} 2 \cos(2u) &= e^{-2\pi i b} q^{-2a} + e^{2\pi i b} q^{2a} - \frac{m^2}{4a^2} e^{2\pi i b} q^{2a} \\ &= 2 \cos(2\pi(a\tau + b)) - \frac{m^2}{4a^2} e^{2\pi i(a\tau + b)}. \end{aligned}$$

We can therefore make the following ansatz for  $u(\tau)$  in the limit  $\tau \rightarrow i\infty$ :

$$u(\tau) = a\tau + b + \frac{1}{2\pi i} \sum_{n=0}^{\infty} \sum_{k=-n}^{\infty} c_{n,k} q^n e^{4\pi i k(a\tau + b)} = a\tau + b + \frac{X}{2\pi i}.$$

The first derivative

$$\frac{u(\tau)}{d\tau} = a + \frac{1}{2\pi i} \sum_{n=0}^{\infty} \sum_{k=-n}^{\infty} c_{n,k} (2\pi i) (n + 2k) q^n e^{4\pi i k(a\tau + b)},$$

and the second derivative

$$\frac{d^2 u(\tau)}{d\tau^2} = (2\pi i) \sum_{n=0}^{\infty} \sum_{k=-n}^{\infty} c_{n,k} (n + 2ka)^2 q^n e^{4\pi i k(a\tau + b)}. \quad (\text{B.12})$$

Let us now look at our equation

$$(2\pi i)^2 \frac{d^2 u(\tau)}{d\tau^2} = m^2 \wp'(2u|\tau). \quad (\text{B.13})$$

The Weierstrass  $\wp$  function has the following expression in terms of theta functions

$$\wp(z|\tau) = -\partial_z^2 \log \theta_1(z|\tau) - 2\eta_1(\tau),$$

and we now analyze the term  $\partial_z^2 \log \theta_1(z|\tau)$  in the limit  $q \rightarrow 0$ . Starting from the series representation of the theta function

$$\theta_1(z|\tau) = -i \sum_{n \in \mathbb{Z}} (-1)^n q^{(n+\frac{1}{2})^2} e^{2\pi i z(n+\frac{1}{2})},$$

the logarithmic derivative

$$\begin{aligned} \partial_z \log \theta_1(z|\tau) - \pi \cot \pi z &= 4\pi \sum_{k=1}^{\infty} q^k \sum_{n=0}^{\infty} q^{kn} \sin(2\pi k z) \\ &= -(2\pi i) \left( \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} q^{kn} e^{i2\pi k z} - \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} q^{kn} e^{-i2\pi k z} \right). \end{aligned}$$

The above computation implies that

$$\partial_z^2 \log \theta_1(z|\tau) + \pi^2 \csc^2 \pi z = -(2\pi i)^2 \left( \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k q^{kn} e^{i2\pi k z} + \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k q^{kn} e^{-i2\pi k z} \right). \quad (\text{B.14})$$

Moreover,  $\csc^2(z)$  has the following series representation:

$$\pi^2 \csc^2 \pi z = \frac{\pi^2}{\sin^2 \pi z} = (2\pi i)^2 e^{2i\pi z} \sum_{k=1}^{\infty} k e^{2i\pi(k-1)z} = (2\pi i)^2 \sum_{k=1}^{\infty} k e^{2i\pi k z}. \quad (\text{B.15})$$

Combining (B.15) and (B.14) we get

$$\begin{aligned} \partial_z^2 \log \theta_1(z|\tau) &= -(2\pi i)^2 \left( \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k q^{kn} e^{i2\pi k z} + \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k q^{kn} e^{-i2\pi k z} + \sum_{k=1}^{\infty} k e^{2i\pi k z} \right) \\ &= -(2\pi i)^2 \left( \sum_{k=1}^{\infty} \sum_{n=0}^{\infty} k q^{kn} e^{i2\pi k z} + \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k q^{kn} e^{-i2\pi k z} \right). \end{aligned}$$

Therefore,

$$\wp'(z|\tau) = -\partial_z^3 \log \theta_1(z|\tau) = (2\pi\mathbf{i})^3 \left( \sum_{k=1}^{\infty} \sum_{n=0}^{\infty} k^2 q^{kn} e^{i2\pi kz} - \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k^2 q^{kn} e^{-i2\pi kz} \right). \quad (\text{B.16})$$

Substituting (B.12), (B.16) in (B.13) we get

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{k=-n}^{\infty} c_{n,k} (n+2k\mathbf{a})^2 q^n e^{4\pi\mathbf{i}k(\mathbf{a}\tau+\mathbf{b})} &= m^2 \sum_{k=1}^{\infty} \sum_{n=0}^{\infty} k^2 q^{kn} e^{4\pi\mathbf{i}k(\mathbf{a}\tau+\mathbf{b})} e^{kX} \\ &\quad - m^2 \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} k^2 q^{kn} e^{-4\pi\mathbf{i}k(\mathbf{a}\tau+\mathbf{b})} e^{-kX}. \end{aligned}$$

Computing the first few values of  $c_{n,k}$ :

$$\begin{aligned} c_{0,0} &= 0, \\ c_{0,1} &= \frac{m^2}{4\mathbf{a}^2}, \\ c_{1,-1} &= -\frac{m^2}{(1-2\mathbf{a})^2} \end{aligned}$$

Therefore,

$$u(\tau) = \mathbf{a}\tau + \mathbf{b} + \frac{m^2}{8\pi\mathbf{i}\mathbf{a}^2} e^{4\pi\mathbf{i}(\mathbf{a}\tau+\mathbf{b})} + \mathcal{O}(qe^{4\pi\mathbf{i}(\mathbf{a}\tau+\mathbf{b})}).$$

□

**Proposition B.2.** *Near its zero  $u(\tau_*) = 0$ , the solution of the equation (B.2)  $u(\tau)$  behaves as*

$$u(\tau) = e^{\mp\mathbf{i}\pi/4} \sqrt{\frac{m}{2\pi}} (\tau - \tau_*)^{1/2} \left( 1 \pm \frac{H_* + 2m^2\eta_1(\tau)}{4\pi\mathbf{i}m} (\tau - \tau_*) \right) + \mathcal{O}\left((\tau - \tau_*)^{5/2}\right). \quad (\text{B.17})$$

*Proof.* The equation (B.2) near  $u(\tau) = 0$  reads as

$$(2\pi\mathbf{i})^2 \frac{d^2 u(\tau)}{d\tau^2} = -\frac{m^2}{4u(\tau)^3} \quad (\text{B.18})$$

due to the following behaviour of the Weierstrass  $\wp$  function at zero

$$\lim_{z \rightarrow 0} \wp(z) = \frac{1}{z^2}.$$

The solution  $u(\tau)$  therefore has the following behaviour

$$u(\tau) = c_1 \sqrt{\tau - \tau_*} + c_2 (\tau - \tau_*)^{3/2} + \mathcal{O}\left((\tau - \tau_*)^{5/2}\right) = c_1 (\tau - \tau_*)^{1/2} \left( 1 + \frac{c_2}{c_1} (\tau - \tau_*) \right) + \mathcal{O}\left((\tau - \tau_*)^{5/2}\right), \quad (\text{B.19})$$

and the second derivative of the above function

$$\frac{d^2 u(\tau)}{d\tau^2} = -\frac{c_1}{4} (\tau - \tau_*)^{-3/2} \left( 1 - \frac{3c_2}{c_1} (\tau - \tau_*) \right). \quad (\text{B.20})$$

Substituting (B.19), (B.20) in (B.18), we obtain that

$$c_1 = e^{\mp\mathbf{i}\pi/4} \frac{\sqrt{m}}{\sqrt{2\pi}}. \quad (\text{B.21})$$

The coefficient of the sub-leading term  $c_2$  can be obtained by studying the Hamiltonian in the same limit:

$$\begin{aligned}
\lim_{\tau \rightarrow \tau_*} H &= \lim_{\tau \rightarrow \tau_*} \left( (2\pi\mathbf{i})^2 \left( \frac{du(\tau)}{d\tau} \right)^2 - \frac{m^2}{4u(\tau)^2} \right) \\
&\stackrel{\text{(B.19)}}{=} (2\pi\mathbf{i})^2 \frac{c_1^2}{4} (\tau - \tau_*)^{-1} \left( 1 + \frac{6c_2}{c_1} (\tau - \tau_*) \right) - m^2 \frac{1}{4c_1^2} (\tau - \tau_*)^{-1} \left( 1 - 2\frac{c_2}{c_1} (\tau - \tau_*) \right) + \mathcal{O}(\tau - \tau_*) \\
&\stackrel{\text{(B.21)}}{=} \frac{c_2}{2c_1} \left( 3(2\pi\mathbf{i})^2 \frac{m}{(2\pi)} e^{\mp i\pi/2} + \frac{m^2(2\pi)}{m} e^{\pm i\pi/2} \right) + \mathcal{O}(\tau - \tau_*) \\
&= \pm \frac{c_2}{c_1} (4\pi\mathbf{i}m) + \mathcal{O}(\tau - \tau_*) \\
&=: H_* + \mathcal{O}(\tau - \tau_*). \\
\Rightarrow \frac{c_2}{c_1} &= \pm \frac{H_*}{4\pi\mathbf{i}m}.
\end{aligned}$$

With the constants  $c_1, c_2$  above,

$$\lim_{\tau \rightarrow \tau_*} u(\tau) = e^{\mp i\pi/4} \sqrt{\frac{m}{2\pi}} (\tau - \tau_*)^{1/2} \left( 1 \pm \frac{H_* + 2m^2\eta_1(\tau)}{4\pi\mathbf{i}m} (\tau - \tau_*) \right) + \mathcal{O}\left((\tau - \tau_*)^{5/2}\right).$$

□

## REFERENCES

- [AGH22] G. Aminov, A. Grassi, and Y. Hatsuda. “Black hole quasinormal modes and Seiberg–Witten theory”. In: *Annales Henri Poincaré*. Vol. 23. 6. Springer. 2022, pp. 1951–1977. arXiv: [2006.06111](#).
- [Ang+24] M. Ang, G. Remy, X. Sun, and T. Zhu. *Derivation of all structure constants for boundary Liouville CFT*. 2024. arXiv: [2305.18266](#) [[math.PR](#)].
- [Arn13] V. I. Arnold. *Mathematical methods of classical mechanics*. Vol. 60. Springer Science & Business Media, 2013.
- [Aru20] J. Aru. “Gaussian multiplicative chaos through the lens of the 2D Gaussian free field”. In: *Markov Process. Related Fields* 26.1 (2020), pp. 17–56. ISSN: 1024-2953. MR: [4237161](#). arXiv: [1709.04355](#) [[math.PR](#)].
- [BCP17] J. Barragán Amado, B. Carneiro da Cunha, and E. Pallante. “On the Kerr-AdS/CFT correspondence”. In: *J. High Energy Phys.* 8 (2017), 094, front matter+25. ISSN: 1126-6708,1029-8479. DOI: [10.1007/JHEP08\(2017\)094](#). MR: [3697388](#). arXiv: [1702.01016](#) [[hep-th](#)].
- [BDK20] M. Beşken, S. Datta, and P. Kraus. “Semi-classical Virasoro blocks: proof of exponentiation”. In: *J. High Energy Phys.* 1 (2020), pp. 109, 15. ISSN: 1126-6708,1029-8479. DOI: [10.1007/jhep01\(2020\)109](#). MR: [4088210](#). arXiv: [1910.04169](#) [[hep-th](#)].
- [Ber17] N. Berestycki. “An elementary approach to Gaussian multiplicative chaos”. In: *Electron. Commun. Probab.* 22 (2017), Paper No. 27, 12. DOI: [10.1214/17-ECP58](#). MR: [3652040](#). arXiv: [1506.09113](#) [[math.PR](#)].
- [BG22] M. Bershtein, P. Gavrylenko, and A. Grassi. “Quantum spectral problems and isomonodromic deformations”. In: *Comm. Math. Phys.* 393.1 (2022), pp. 347–418. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-022-04369-y](#). MR: [4440693](#). arXiv: [2105.00985](#) [[math-ph](#)].
- [Bon+20] G. Bonelli, F. Del Monte, P. Gavrylenko, and A. Tanzini. “ $\mathcal{N} = 2^*$  gauge theory, free fermions on the torus and Painlevé VI”. In: *Comm. Math. Phys.* 377.2 (2020), pp. 1381–1419. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-020-03743-y](#). MR: [4115020](#). arXiv: [1901.10497v2](#) [[hep-th](#)].
- [Bon+23] G. Bonelli, C. Iossa, D. P. Lichtig, and A. Tanzini. “Irregular Liouville correlators and connection formulae for Heun functions”. In: *Comm. Math. Phys.* 397.2 (2023), pp. 635–727. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-022-04497-5](#). MR: [4538706](#). arXiv: [2201.04491](#) [[hep-th](#)].

- [BPZ84] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov. “Infinite conformal symmetry in two-dimensional quantum field theory”. In: *Nuclear Physics B* 241.2 (1984), pp. 333–380.
- [Bus80] V. S. Buslaev. *Variational calculus*. Izd. Leningrad. Univ., 1980.
- [CC22] B. C. da Cunha and J. P. Cavalcante. “Expansions for semiclassical conformal blocks”. In: (2022). arXiv: [2211.03551](https://arxiv.org/abs/2211.03551) [[hep-th](#)].
- [Dav+16] F. David, A. Kupiainen, R. Rhodes, and V. Vargas. “Liouville quantum gravity on the Riemann sphere”. In: *Comm. Math. Phys.* 342.3 (2016), pp. 869–907. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-016-2572-4](https://doi.org/10.1007/s00220-016-2572-4). MR: [3465434](#). arXiv: [1410.7318](https://arxiv.org/abs/1410.7318) [[math.PR](#)].
- [DDG23] F. Del Monte, H. Desiraju, and P. Gavrylenko. “Isomonodromic tau functions on a torus as Fredholm determinants, and charged partitions”. In: *Comm. Math. Phys.* 398.3 (2023), pp. 1029–1084. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-022-04458-y](https://doi.org/10.1007/s00220-022-04458-y). MR: [4561797](#). arXiv: [2011.06292](https://arxiv.org/abs/2011.06292) [[math-ph](#)].
- [Des21] H. Desiraju. “Painlevé tau-functions and Fredholm determinants”. PhD thesis. SISSA, 2021. URL: <https://hdl.handle.net/20.500.11767/118727>.
- [Des22] H. Desiraju. “Painlevé/CFT correspondence on a torus”. In: *J. Math. Phys.* 63.8 (2022), Paper No. 081102, 16. ISSN: 0022-2488,1089-7658. DOI: [10.1063/5.0089867](https://doi.org/10.1063/5.0089867). MR: [4462569](#). arXiv: [2305.04240](https://arxiv.org/abs/2305.04240) [[math-ph](#)].
- [DLMF] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.1.10 of 2023-06-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [DO94] H. Dorn and H.-J. Otto. “Two-and three-point functions in Liouville theory”. In: *Nuclear Physics B* 429.2 (1994), pp. 375–388.
- [DRV16] F. David, R. Rhodes, and V. Vargas. “Liouville quantum gravity on complex tori”. In: *J. Math. Phys.* 57.2 (2016), pp. 022302, 25. DOI: [10.1063/1.4938107](https://doi.org/10.1063/1.4938107). MR: [3450564](#). arXiv: [1504.00625](https://arxiv.org/abs/1504.00625) [[math.PR](#)].
- [Ere+22] A. Eremenko, A. Gabrielov, G. Mondello, and D. Panov. “Moduli spaces for Lamé functions and Abelian differentials of the second kind”. In: *Commun. Contemp. Math.* 24.2 (2022), Paper No. 2150028, 68. DOI: [10.1142/S0219199721500280](https://doi.org/10.1142/S0219199721500280). MR: [4380114](#). arXiv: [2006.16837](https://arxiv.org/abs/2006.16837) [[math.CV](#)].
- [Fat+09] V. A. Fateev, A. V. Litvinov, A. Neveu, and E. Onofri. “A differential equation for a four-point correlation function in Liouville field theory and elliptic four-point conformal blocks”. In: *J. Phys. A* 42.30 (2009), pp. 304011, 29. ISSN: 1751-8113,1751-8121. DOI: [10.1088/1751-8113/42/30/304011](https://doi.org/10.1088/1751-8113/42/30/304011). MR: [2521330](#). arXiv: [0902.1331v3](https://arxiv.org/abs/0902.1331v3) [[hep-th](#)].
- [FL10] V. A. Fateev and A. V. Litvinov. “On AGT conjecture”. In: *J. High Energy Phys.* 2 (2010), pp. 014, 17. ISSN: 1126-6708,1029-8479. DOI: [10.1007/JHEP02\(2010\)014](https://doi.org/10.1007/JHEP02(2010)014). MR: [2672761](#). arXiv: [0912.0504v2](https://arxiv.org/abs/0912.0504v2) [[hep-th](#)].
- [Gho+] P. Ghosal, G. Remy, X. Sun, and Y. Sun. “Analyticity and Symmetry of Virasoro Conformal blocks via Liouville CFT”. In: (). URL: [https://drive.google.com/file/d/1CKpOICEbezvoma16ABVekEv5xhJ0-0\\_Q/view?pli=1](https://drive.google.com/file/d/1CKpOICEbezvoma16ABVekEv5xhJ0-0_Q/view?pli=1).
- [Gho+24] P. Ghosal, G. Remy, X. Sun, and Y. Sun. “Probabilistic conformal blocks for Liouville CFT on the torus”. In: *Duke Math. J.* 173.6 (2024), pp. 1085–1175. DOI: [10.1215/00127094-2023-0031](https://doi.org/10.1215/00127094-2023-0031). MR: [4748792](#). arXiv: [2003.03802](https://arxiv.org/abs/2003.03802) [[math.PR](#)].
- [GRV19] C. Guillarmou, R. Rhodes, and V. Vargas. “Polyakov’s formulation of  $2d$  bosonic string theory”. In: *Publ. Math. Inst. Hautes Études Sci.* 130 (2019), pp. 111–185. ISSN: 0073-8301,1618-1913. DOI: [10.1007/s10240-019-00109-6](https://doi.org/10.1007/s10240-019-00109-6). MR: [4028515](#). arXiv: [1607.08467](https://arxiv.org/abs/1607.08467) [[math-ph](#)].
- [Gu+20] J. Gu, B. Haghighat, A. Klemm, K. Sun, and X. Wang. “Elliptic blowup equations for  $6d$  SCFTs. Part III. E-strings, M-strings and chains”. In: *J. High Energy Phys.* 7 (2020), pp. 135, 55. ISSN: 1126-6708,1029-8479. DOI: [10.1007/jhep07\(2020\)135](https://doi.org/10.1007/jhep07(2020)135). MR: [4137961](#). arXiv: [1911.11724](https://arxiv.org/abs/1911.11724) [[hep-th](#)].

- [Gui+21] C. Guillarmou, A. Kupiainen, R. Rhodes, and V. Vargas. *Segal’s axioms and bootstrap for Liouville Theory*. 2021. arXiv: [2112.14859](#) [[math.PR](#)].
- [HJS10] L. Hadasz, Z. Jaskólski, and P. Suchanek. “Recursive representation of the torus 1-point conformal block”. In: *Journal of High Energy Physics* 2010.1 (2010), pp. 1–13.
- [HK18] L. Hollands and O. Kidwai. “Higher length-twist coordinates, generalized Heun’s opers, and twisted superpotentials”. In: *Adv. Theor. Math. Phys.* 22.7 (2018), pp. 1713–1822. ISSN: 1095-0761,1095-0753. DOI: [10.4310/ATMP.2018.v22.n7.a2](#). MR: [3976899](#). arXiv: [1710.04438](#) [[hep-th](#)].
- [Hor18] M. Hortaçsu. “Heun functions and some of their applications in physics”. In: *Adv. High Energy Phys.* (2018), Art. ID 8621573, 14. DOI: [10.1155/2018/8621573](#). MR: [3834316](#). arXiv: [1101.0471v11](#) [[math-ph](#)].
- [HRV18] Y. Huang, R. Rhodes, and V. Vargas. “Liouville quantum gravity on the unit disk”. In: *Ann. Inst. Henri Poincaré Probab. Stat.* 54.3 (2018). DOI: [10.1214/17-AIHP852](#). MR: [3825895](#). arXiv: [1502.04343](#) [[math.PR](#)].
- [Inc40] E. L. Ince. “Further investigations into the periodic Lamé functions”. In: *Proc. Roy. Soc. Edinburgh* 60 (1940), pp. 83–99. ISSN: 0370-1646. MR: [2400](#).
- [JSW18] J. Junnila, E. Saksman, and C. Webb. “Decompositions of log-correlated fields with applications”. In: *arXiv e-prints* (Aug. 2018). DOI: [10.48550/arXiv.1808.06838](#). arXiv: [1808.06838](#) [[math.PR](#)].
- [KRV20] A. Kupiainen, R. Rhodes, and V. Vargas. “Integrability of Liouville theory: proof of the DOZZ formula”. In: *Annals of Mathematics* 191.1 (2020), pp. 81–166.
- [KT13] A.-K. Kashani-Poor and J. Troost. “Transformations of spherical blocks”. In: *J. High Energy Phys.* 10 (2013). DOI: [10.1007/JHEP10%282013%29009](#). MR: [3110708](#). arXiv: [1305.7408](#) [[hep-th](#)].
- [Lit+14] A. Litvinov, S. Lukyanov, N. Nekrasov, and A. Zamolodchikov. “Classical conformal blocks and Painlevé VI”. In: *J. High Energy Phys.* 7 (2014), 144, front matter+19. ISSN: 1126-6708,1029-8479. DOI: [10.1007/JHEP07\(2014\)144](#). MR: [3250114](#). arXiv: [1309.4700](#) [[hep-th](#)].
- [LN18] M. Lencsés and F. Novaes. “Classical conformal blocks and accessory parameters from isomonodromic deformations”. In: *J. High Energy Phys.* 4 (2018), 096, front matter+36. ISSN: 1126-6708,1029-8479. DOI: [10.1007/jhep04\(2018\)096](#). MR: [3804362](#). arXiv: [1709.03476](#) [[hep-th](#)].
- [LN21] O. Lisovyy and A. Naidiuk. “Accessory parameters in confluent Heun equations and classical irregular conformal blocks”. In: *Lett. Math. Phys.* 111.6 (2021), Paper No. 137, 28. ISSN: 0377-9017,1573-0530. DOI: [10.1007/s11005-021-01400-6](#). MR: [4338702](#). arXiv: [2101.05715](#) [[math-ph](#)].
- [LN22] O. Lisovyy and A. Naidiuk. “Perturbative connection formulas for Heun equations”. In: *J. Phys. A* 55.43 (2022), Paper No. 434005, 22. ISSN: 1751-8113,1751-8121. DOI: [10.1088/1751-8121/ac9ba7](#). MR: [4506508](#). arXiv: [2208.01604](#) [[math-ph](#)].
- [LRV] H. Lacoïn, R. Rhodes, and V. Vargas. “The semiclassical limit of Liouville conformal field theory”. In: (). arXiv: [1903.08883](#).
- [LRV22] H. Lacoïn, R. Rhodes, and V. Vargas. “Path integral for quantum Mabuchi K-energy”. In: *Duke Math. J.* 171.3 (2022), pp. 483–545. DOI: [10.1215/00127094-2021-0007](#). MR: [4383250](#). arXiv: [1807.01758](#) [[math-ph](#)].
- [Man98] Y. I. Manin. “Sixth Painlevé equation, universal elliptic curve, and mirror of  $\mathbf{P}^2$ ”. In: *Geometry of differential equations*. Vol. 186. Amer. Math. Soc. Transl. Ser. 2. Amer. Math. Soc., Providence, RI, 1998, pp. 131–151. DOI: [10.1090/trans2/186/04](#). MR: [1732409](#). arXiv: [alg-geom/9605010](#) [[math.AG](#)].
- [MMS10] A. Mironov, A. Morozov, and S. Shakirov. “Conformal blocks as Dotsenko-Fateev integral discriminants”. In: *Internat. J. Modern Phys. A* 25.16 (2010), pp. 3173–3207. ISSN: 0217-751X,1793-656X. DOI: [10.1142/S0217751X10049141](#). MR: [2659902](#). arXiv: [1001.0563](#) [[hep-th](#)].

- [Neg16] A. Neguț. “Exts and the AGT relations”. In: *Lett. Math. Phys.* 106.9 (2016), pp. 1265–1316. ISSN: 0377-9017,1573-0530. DOI: [10.1007/s11005-016-0865-3](https://doi.org/10.1007/s11005-016-0865-3). MR: [3533570](https://arxiv.org/abs/1510.05482). arXiv: [1510.05482](https://arxiv.org/abs/1510.05482) [hep-th].
- [Nov86] V. Y. Novokshënov. “Moving poles of solutions of the Painlevé equation of the third type and their connection with Mathieu functions”. In: *Funktsional. Anal. i Prilozhen.* 20.2 (1986), pp. 38–49, 96. ISSN: 0374-1990. DOI: [10.1007/BF01077265](https://doi.org/10.1007/BF01077265). MR: [847137](https://arxiv.org/abs/847137).
- [NPS18] N. Nekrasov, V. Pestun, and S. Shatashvili. “Quantum geometry and quiver gauge theories”. In: *Comm. Math. Phys.* 357.2 (2018), pp. 519–567. ISSN: 0010-3616,1432-0916. DOI: [10.1007/s00220-017-3071-y](https://doi.org/10.1007/s00220-017-3071-y). MR: [3767702](https://arxiv.org/abs/1312.6689). arXiv: [1312.6689](https://arxiv.org/abs/1312.6689) [hep-th].
- [NRS11] N. Nekrasov, A. Rosly, and S. Shatashvili. “Darboux coordinates, Yang-Yang functional, and gauge theory”. In: *Nuclear Phys. B Proc. Suppl.* 216 (2011), pp. 69–93. ISSN: 0920-5632. DOI: [10.1016/j.nuclphysbps.2011.04.150](https://doi.org/10.1016/j.nuclphysbps.2011.04.150). MR: [2851597](https://arxiv.org/abs/2851597). arXiv: [1103.3919](https://arxiv.org/abs/1103.3919) [hep-th].
- [NS09] N. A. Nekrasov and S. L. Shatashvili. “Supersymmetric vacua and Bethe ansatz”. In: *Nuclear Phys. B Proc. Suppl.* 192/193 (2009), pp. 91–112. ISSN: 0920-5632. DOI: [10.1016/j.nuclphysbps.2009.07.047](https://doi.org/10.1016/j.nuclphysbps.2009.07.047). MR: [2570974](https://arxiv.org/abs/2570974). arXiv: [0901.4744](https://arxiv.org/abs/0901.4744) [hep-th].
- [NS10] N. A. Nekrasov and S. L. Shatashvili. “Quantization of integrable systems and four dimensional gauge theories”. In: *XVIIth International Congress on Mathematical Physics*. World Sci. Publ., Hackensack, NJ, 2010, pp. 265–289. ISBN: 978-981-4304-62-7. DOI: [10.1142/9789814304634\\_0015](https://doi.org/10.1142/9789814304634_0015). MR: [2730782](https://arxiv.org/abs/2730782). arXiv: [0908.4052](https://arxiv.org/abs/0908.4052) [hep-th].
- [NY03] H. Nakajima and K. Yoshioka. “Lectures on instanton counting”. In: *Algebraic structures and moduli spaces*. Vol. 38. CRM Proc. Lecture Notes. Amer. Math. Soc., Providence, RI, 2003, pp. 31–101. ISBN: 0-8218-3568-8. DOI: [10.1090/crmp/038/02](https://doi.org/10.1090/crmp/038/02). MR: [2095899](https://arxiv.org/abs/2095899). arXiv: [math/0311058](https://arxiv.org/abs/math/0311058) [math.AG].
- [Piğ+22] M. Piğtek, R. G. Nazmitdinov, A. Puente, and A. R. Pietrykowski. “Classical conformal blocks, Coulomb gas integrals and Richardson-Gaudin models”. In: *J. High Energy Phys.* 4 (2022), Paper No. 098, 48. DOI: [10.1007/JHEP04%282022%29098](https://doi.org/10.1007/JHEP04%282022%29098). MR: [4429642](https://arxiv.org/abs/4429642). arXiv: [2110.15009](https://arxiv.org/abs/2110.15009) [hep-th].
- [Piğ14] M. Piğtek. “Classical torus conformal block,  $\mathcal{N} = 2^*$  twisted superpotential and the accessory parameter of Lamé equation”. In: *Journal of High Energy Physics* 2014.3 (2014), pp. 1–39. DOI: [10.1007/JHEP03\(2014\)124](https://doi.org/10.1007/JHEP03(2014)124). arXiv: [1309.7672v3](https://arxiv.org/abs/1309.7672v3) [hep-th].
- [Pol81] A. M. Polyakov. “Quantum geometry of bosonic strings”. In: *Physics Letters B* 103.3 (1981), pp. 207–210. DOI: [10.1016/0370-2693\(81\)90743-7](https://doi.org/10.1016/0370-2693(81)90743-7).
- [Pon04] B. Ponsot. “Recent progress in Liouville field theory”. In: *Proceedings of 6th International Workshop on Conformal Field Theory and Integrable Models*. Vol. 19. 2004, pp. 311–335. DOI: [10.1142/S0217751X0402049X](https://doi.org/10.1142/S0217751X0402049X). MR: [2087118](https://arxiv.org/abs/2087118). arXiv: [hep-th/0301193](https://arxiv.org/abs/hep-th/0301193).
- [PP19] M. Piğtek and A. R. Pietrykowski. “Solving Heun’s equation using conformal blocks”. In: *Nuclear Phys. B* 938 (2019), pp. 543–570. DOI: [10.1016/j.nuclphysb.2018.11.021](https://doi.org/10.1016/j.nuclphysb.2018.11.021). MR: [3883275](https://arxiv.org/abs/3883275). arXiv: [1708.06135](https://arxiv.org/abs/1708.06135) [hep-th].
- [PW24] E. Peltola and Y. Wang. “Large deviations of multichordal  $\text{SLE}_{0+}$ , real rational functions, and zeta-regularized determinants of Laplacians”. In: *J. Eur. Math. Soc. (JEMS)* 26.2 (2024), pp. 469–535. ISSN: 1435-9855,1435-9863. DOI: [10.4171/jems/1274](https://doi.org/10.4171/jems/1274). MR: [4705656](https://arxiv.org/abs/4705656). arXiv: [2006.08574](https://arxiv.org/abs/2006.08574) [math-ph].
- [RA95] A. Ronveaux and F. M. Arscott. *Heun’s differential equations*. Clarendon Press, 1995.
- [Rib14] S. Ribault. *Conformal field theory on the plane*. 2014. arXiv: [1406.4290](https://arxiv.org/abs/1406.4290) [hep-th].
- [RV] R. Rhodes and V. Vargas. “Lecture notes on Liouville theory and the DOZZ formula”. In: *Topics in statistical mechanics*. Vol. 59. Panor. Synthèses. Soc. Math. France, Paris, pp. 185–229. ISBN: 978-2-85629-970-8. MR: [4619314](https://arxiv.org/abs/4619314). arXiv: [1712.00829](https://arxiv.org/abs/1712.00829) [math.PR].
- [Tak01] K. Takasaki. “Painlevé-Calogero correspondence revisited”. In: *J. Math. Phys.* 42.3 (2001), pp. 1443–1473. ISSN: 0022-2488,1089-7658. DOI: [10.1063/1.1348025](https://doi.org/10.1063/1.1348025). MR: [1814699](https://arxiv.org/abs/1814699). arXiv: [math/0004118](https://arxiv.org/abs/math/0004118) [math-qa].
- [Tak94] L. A. Takhtajan. “Topics in the quantum geometry of Riemann surfaces: two-dimensional quantum gravity”. In: *Quantum groups and their applications in physics (Varenna,*

- 1994). Vol. 127. Proc. Internat. School Phys. Enrico Fermi. IOS, Amsterdam, 1994, pp. 541–579. ISBN: 90-5199-247-5. MR: [1415866](#). arXiv: [hep-th/9409088](#).
- [Tes11] J. Teschner. “Quantization of the Hitchin moduli spaces, Liouville theory and the geometric Langlands correspondence I”. In: *Adv. Theor. Math. Phys* 15 (2011), pp. 471–564. arXiv: [1005.2846](#).
- [TZ03] L. Takhtajan and P. Zograf. “Hyperbolic 2-spheres with conical singularities, accessory parameters and Kähler metrics on  $\mathcal{M}_{0,n}$ ”. In: *Transactions of the American mathematical society* 355.5 (2003), pp. 1857–1867. arXiv: [math/0112170](#).
- [Wan19a] Y. Wang. “Equivalent descriptions of the Loewner energy”. In: *Invent. Math.* 218.2 (2019), pp. 573–621. ISSN: 0020-9910,1432-1297. DOI: [10.1007/s00222-019-00887-0](#). MR: [4011706](#). arXiv: [1802.01999](#) [[math.CV](#)].
- [Wan19b] Y. Wang. “The energy of a deterministic Loewner chain: reversibility and interpretation via SLE $_{0+}$ ”. In: *J. Eur. Math. Soc. (JEMS)* 21.7 (2019), pp. 1915–1941. ISSN: 1435-9855,1435-9863. DOI: [10.4171/JEMS/876](#). MR: [3959854](#). arXiv: [1601.05297](#) [[math.CV](#)].
- [Wan21] Y. Wang. “A note on Loewner energy, conformal restriction and Werner’s measure on self-avoiding loops”. In: *Ann. Inst. Fourier (Grenoble)* 71.4 (2021), pp. 1791–1805. DOI: [10.5802/aif.3427](#). MR: [4398248](#). arXiv: [1810.04578](#) [[math.CV](#)].
- [XXZ21] J. Xia, S.-X. Xu, and Y.-Q. Zhao. “Isomonodromy sets of accessory parameters for Heun class equations”. In: *Stud. Appl. Math.* 146.4 (2021). DOI: [10.1111/sapm.12370](#). MR: [4300140](#). arXiv: [2101.02864](#) [[math.CA](#)].
- [Zam84] A. B. Zamolodchikov. “Conformal symmetry in two dimensions: an explicit recurrence formula for the conformal partial wave amplitude”. In: *Communications in mathematical physics* 96 (1984), pp. 419–422.
- [Zam86] A. B. Zamolodchikov. “Two-dimensional conformal symmetry and critical four-spin correlation functions in the Ashkin-Teller model”. In: *Soviet Journal of Experimental and Theoretical Physics* 63.5 (1986), pp. 1061–1066. MR: [869395](#).
- [Zam87a] A. B. Zamolodchikov. “Conformal symmetry in two-dimensional space. On a recurrent representation of the conformal block”. In: *Teor. Mat. Fiz.;(USSR)* 73.1 (1987).
- [Zam87b] A. B. Zamolodchikov. “Conformal symmetry in two-dimensional space: on a recurrent representation of the conformal block”. In: *Teoret. Mat. Fiz.* 73.1 (1987), pp. 103–110. ISSN: 0564-6162. DOI: [10.1007/bf01022967](#). MR: [939798](#).
- [ZZ96] A. Zamolodchikov and A. Zamolodchikov. “Conformal bootstrap in Liouville field theory”. In: *Nuclear Physics B* 477.2 (1996), pp. 577–605.

(Harini Desiraju) SYDNEY MATHEMATICAL RESEARCH INSTITUTE (SMRI), SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, CAMPERDOWN, NSW 2006, AUSTRALIA.

Email address: [harini.desiraju@sydney.edu.au](mailto:harini.desiraju@sydney.edu.au)

(Promit Ghosal) UNIVERSITY OF CHICAGO, DEPARTMENT OF STATISTICS, 5747 S ELLIS AVE, CHICAGO, IL 60637, USA

Email address: [promit@uchicago.edu](mailto:promit@uchicago.edu)

(Andrei Prokhorov) UNIVERSITY OF MICHIGAN, 500 S STATE ST, ANN ARBOR, MI 48109, USA, AND ST. PETERSBURG UNIVERSITY, 7/9 UNIVERSITetskAYA NAB. 199034 ST. PETERSBURG, RUSSIA.

Email address: [andreip@umich.edu](mailto:andreip@umich.edu)