

L^p ESTIMATES FOR THE LAPLACIAN VIA BLOW-UP

JAN LEWENSTEIN-SANPERA AND XAVIER ROS-OTON

ABSTRACT. In this note we provide a new proof of the $W^{2,p}$ Calderón-Zygmund regularity estimates for the Laplacian, i.e., $\Delta u = f$ and its parabolic counterpart $\partial_t u - \Delta u = f$. Our proof is an adaptation of a contradiction and compactness argument that so far had been only used to prove estimates in Hölder spaces. This new approach is simpler than previous ones, and avoids the use of any interpolation theorem.

1. INTRODUCTION

The main concern for this article is the following classical theorem:

Theorem 1.1 ([CZ52]). *Let $1 < p < \infty$, and $u \in H^1(B_1)$ be any weak solution to:*

$$\Delta u = f \quad \text{in } B_1,$$

with $f \in L^p(B_1)$. Then u is in $W^{2,p}$ inside B_1 and the following estimate holds:

$$\int_{B_{1/2}} |D^2 u|^p \leq C \left(\int_{B_1} |u|^p + \int_{B_1} |f|^p \right). \quad (1.1)$$

The constant C depends only on n and p .

The classical proof of this result, which can be found in [GT98], is as follows:

- (i) Prove the result for $p = 2$ by an easy integration by parts argument.
- (ii) Using the Calderón-Zygmund decomposition, and the maximal function of Hardy-Littlewood, prove a *weak* L^1 estimate corresponding to the case $p = 1$.
- (iii) Prove the Marcinkiewicz interpolation theorem, and use it to deduce the result for all $1 < p < 2$.
- (iv) By a duality argument, deduce the result for all $2 < p < \infty$.

This result and its proof is a landmark in elliptic PDE, and connects this subject to harmonic analysis.

Up to date, some other approaches have been introduced to find new proofs of this result. In particular, an alternative proof is presented in [Le13], in which (ii)-(iii)-(iv) are replaced by:

- (ii') Prove an L^∞ -to-BMO estimate, corresponding to the case $p = \infty$.
- (iii') Prove the Stampacchia interpolation theorem, by using the one due to Marcinkiewicz as well as the sharp maximal function of Fefferman-Stein, and use it to deduce the result for all $2 < p < \infty$.
- (iv') By a duality argument, deduce the result for all $1 < p < 2$.

On the other hand, a completely different (and more geometric) approach was developed in [Ca89] to treat fully nonlinear equations for $p > n$, and later in [Wa13] for the Laplacian for all $1 < p < \infty$. The proof in [Wa13] uses only the maximal function, energy estimates, and Vitali covering lemma, to

2020 *Mathematics Subject Classification.* 35B65, 35J05, 35K05.

Key words and phrases. Calderón-Zygmund estimates, elliptic equations, parabolic equations.

establish a decay estimate for the superlevel sets of $\mathcal{M}|D^2u|^2$ in terms of those of $\mathcal{M}|f|^2$ (where \mathcal{M} is the maximal function).

Our goal in this paper is to give a new proof of this result, which we believe is simpler and easier to follow than the one in [Wa13, Ca89]. Our proof uses only the sharp maximal function, combined with a quite elementary contradiction and compactness argument in the spirit of those in [Si97, Se15]; see also [FR22].

In the last decade there have been many regularity estimates that have been established by a contradiction and compactness estimate in the spirit of those in [Si97, Se15]. However, this type of argument had been always used to prove estimates in Hölder spaces, and the present paper is the first one to establish a $W^{2,p}$ estimate via a blow-up argument. Notice that L^p regularity is quite different than $C^{0,\alpha}$, which is pointwise. To solve this issue we use the sharp maximal function, and establish the pointwise inequality (2.4) below. The idea to establish such a pointwise inequality comes from [LZ24], where a simple proof of the L^∞ to BMO estimate is given, with a different argument.

1.1. Acknowledgements. X. R. was supported by the European Research Council under the Grant Agreements No. 801867 (EllipticPDE) and No. 101123223 (SSNSD), by the AEI project PID2021-125021NA-I00 (Spain), by the grant RED2022-134784-T funded by AEI/10.13039/501100011033, by AGAUR Grant 2021 SGR 00087 (Catalunya), and by the Spanish State Research Agency through the María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M).

1.2. Organization of the paper. The paper is organized as follows: In section 2 the L^p regularity estimate for the Laplacian is proved, while in section 3 we treat parabolic equations.

2. ELLIPTIC EQUATIONS

In this section we focus on the regularity of the weak solutions to Poisson equation

$$\Delta u = f \quad \text{in } B_1. \quad (2.1)$$

2.1. Sharp maximal function. We provide a very brief introduction to the tools needed in order to prove Theorem 1.1. Further information can be found in [LL01, Ev10, GM12].

We define the 2-sharp maximal function (see, e.g. [ST89]) for a locally integrable function w as

$$\mathcal{M}_2^\# w(x) = \sup_{r>0} \int_{B_r(x)} |w - \bar{w}_{B_r(x)}|^2,$$

where $\bar{w}_E := \int_E w$.

Then, we have the following estimate for any $1 < p < \infty$

$$c\|w\|_{L^p(B_r)} \leq \|\mathcal{M}_2^\# w\|_{L^p(B_r)}^{1/2} + \|w\|_{L^1(B_r)} \leq C\|w\|_{L^p(B_r)}, \quad (2.2)$$

where c, C depend only on p, n and r . Indeed, the first inequality follows from a classical theorem of Fefferman and Stein on the sharp maximal function [GM12] and Hölder's inequality, while the second from the trivial bound $\mathcal{M}_2^\# w \leq \mathcal{M}w$ and the strong L^p - L^p inequality for the maximal function [GT98].

When applied to a Hessian matrix, we still denote $\mathcal{M}_2^\# D^2w(x) = \sup_{r>0} \int_{B_r(x)} |D^2w - \overline{D^2w}_{B_r(x)}|^2$.

2.2. Regularity estimates. We want to establish the estimate (1.1) for $p \neq 2$. Indeed, recall that in the simplest case $p = 2$, i.e.,

$$\|D^2u\|_{L^2(B_{1/2})} \leq C (\|u\|_{L^2(B_1)} + \|f\|_{L^2(B_1)}), \quad (2.3)$$

it follows easily from the identity

$$\int_{\mathbb{R}^n} |D^2v|^2 = \int_{\mathbb{R}^n} |\Delta v|^2 \quad \forall v \in C_c^\infty(\mathbb{R}^n);$$

see e.g. [FR22, Remark 2.13].

The estimate (1.1) $2 < p < \infty$ will follow from the following:

Proposition 2.1. *Let $u, f \in C^\infty(B_1)$, with $\Delta u = f$ in B_1 . Then,*

$$\mathcal{M}_2^\# D^2u(x) \leq C (\|u\|_{L^2(B_1)}^2 + \|f\|_{L^2(B_1)}^2 + \mathcal{M}_2^\# f(x)), \quad (2.4)$$

for any $x \in B_{1/2}$, where C depends only on n .

To prove this result, we need the following elementary Lemma.

Lemma 2.2. *Let $\{u_k\}_k \subset W^{2,2}(B_1)$ be a sequence of functions satisfying $\|D^2u_k\|_{L^2(B_1)} \leq C_0$ and*

$$\sup_k \sup_{r>0} \int_{B_r} |D^2u_k(x) - \overline{D^2u_k}_{B_r}|^2 \rightarrow \infty.$$

Then, for any $0 < \delta < 1$ there exists $r_m \rightarrow 0$ and a subsequence k_m such that:

$$(1 - \delta) \int_{B_\rho} |D^2u_k(x) - \overline{D^2u_k}_{B_\rho}|^2 \leq \int_{B_{r_m}} |D^2u_{k_m}(x) - \overline{D^2u_{k_m}}_{B_{r_m}}|^2 \quad \forall k \in \mathbb{N}, \rho \geq r_m.$$

Proof. The quantity:

$$\Theta(r) = \sup_k \sup_{\rho \geq r} \int_{B_\rho} |D^2u_k(x) - \overline{D^2u_k}_{B_\rho}|^2.$$

satisfies that $\theta(r) \rightarrow \infty$ as $r \rightarrow 0$. Hence, for any $\varepsilon > 0$ there exists a $r_\varepsilon \geq \varepsilon$ such that:

$$(1 - \delta)\Theta(\varepsilon) \leq \int_{B_{r_\varepsilon}} |D^2u_k(x) - \overline{D^2u_k}_{B_{r_\varepsilon}}|^2.$$

Take a sequence $\varepsilon_m \rightarrow 0$. Since $\|D^2u_k\|_{L^2(B_1)}$ is bounded, then for $\int_{B_{r_\varepsilon}} |D^2u_k(x) - \overline{D^2u_k}_{B_{r_\varepsilon}}|^2 \rightarrow \infty$ we need that $r_m \rightarrow 0$ as well. Moreover, by monotonicity of $\theta(r)$, we have that $\theta(r_\varepsilon) \leq \theta(\varepsilon)$ with:

$$(1 - \delta)\Theta(r_\varepsilon) \leq \int_{B_{r_\varepsilon}} |D^2u_k(x) - \overline{D^2u_k}_{B_{r_\varepsilon}}|^2,$$

and the lemma follows. \square

We next give the:

Proof of Proposition 2.1. It suffices to establish the result for $x = 0$. Let us show the result by contradiction through a blow-up method. Suppose that the estimate (2.4) does not hold. Then, there exist functions $u_k, f_k \in C^\infty(B_1)$, such that $\Delta u_k = f_k$ in B_1 , and

$$\mathcal{M}_2^\# D^2u_k(0) > k (\|u_k\|_{L^2(B_1)}^2 + \|f_k\|_{L^2(B_1)}^2 + \mathcal{M}_2^\# f_k(0))$$

for all $k \in \mathbb{N}$. Moreover, dividing u_k by a constant if necessary, we may assume

$$\|u_k\|_{L^2(B_1)}^2 + \|f_k\|_{L^2(B_1)}^2 + \mathcal{M}_2^\# f_k(0) \leq 1,$$

It then follows that

$$\sup_k \sup_{r>0} \int_{B_r} |D^2 u_k(x) - \overline{D^2 u_k}_{B_r}|^2 \rightarrow \infty.$$

Let $r_m \rightarrow 0$ and k_m be given by Lemma 2.2 (applied with $\delta = \frac{1}{2}$) and define new functions $v_m, g_m \in C^\infty(B_{1/2r_m})$ as

$$v_m(x) = \frac{u_{k_m}(r_m x) - p_m(x)}{r_m^2 \Theta^{1/2}(r_m)}, \quad g_m(x) = \frac{f_{k_m}(r_m x) - \overline{f_{k_m}}_{B_{r_m}}}{\Theta^{1/2}(r_m)},$$

where

$$\int_{B_\rho} |D^2 u_k(x) - \overline{D^2 u_k}_{B_\rho}|^2 \leq 2 \int_{B_{r_m}} |D^2 u_{k_m}(x) - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2 =: \Theta(r_m) \quad \forall k \in \mathbb{N}, \rho \geq r_m. \quad (2.5)$$

Notice also that it follows from the previous inequality that $\Theta(r_m) \rightarrow \infty$.

Here, $p_m(x)$ denotes a quadratic polynomial such that $\overline{v_m}_{B_1} = \overline{\nabla v_m}_{B_1} = \overline{D^2 v_m}_{B_1} = 0$. Notice that $\Delta v_m = g_m$ in $B_{1/2r_m}$.

Let us show that v_m are bounded in the L^2 norm in B_R with $R > 1$:

$$\begin{aligned} \|D^2 v_m\|_{L^2(B_R)}^2 &= \frac{\int_{B_R} |D^2 u_{k_m}(r_m x) - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2}{\Theta(r_m)} = \frac{R^n \int_{B_{Rr_m}} |D^2 u_{k_m}(y) - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2}{\Theta(r_m)} \\ &\leq \frac{2R^n \left[\int_{B_{Rr_m}} |D^2 u_{k_m}(y) - \overline{D^2 u_{k_m}}_{B_{Rr_m}}|^2 + |\overline{D^2 u_{k_m}}_{B_{Rr_m}} - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2 \right]}{\Theta(r_m)} \leq 2R^n(1 + R^n). \end{aligned}$$

where we used (2.5) and

$$\begin{aligned} |\overline{D^2 u_{k_m}}_{B_{Rr_m}} - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2 &\leq \left(\int_{B_{r_m}} |D^2 u_{k_m}(x) - \overline{D^2 u_{k_m}}_{B_{Rr_m}}| \right)^2 \\ &\leq \int_{B_{r_m}} |D^2 u_{k_m}(x) - \overline{D^2 u_{k_m}}_{B_{Rr_m}}|^2 \\ &\leq R^n \int_{B_{Rr_m}} |D^2 u_{k_m}(x) - \overline{D^2 u_{k_m}}_{B_{Rr_m}}|^2 \leq R^n \Theta(r_m) \end{aligned}$$

where we used (2.5) again. Moreover, using the same arguments as above, one gets

$$\|g_m\|_{L^2(B_R)}^2 \leq \frac{R^n(1 + R^n) \mathcal{M}_\#^2 f_{k_m}(0)}{\Theta(r_m)}.$$

Since $\mathcal{M}_\#^2 f_{k_m}(0) \leq 1$ by hypothesis, then $\|g_m\|_{L^2(B_R)}^2 \rightarrow 0$ as $m \rightarrow \infty$. Note as well that

$$\|D^2 v_m\|_{L^2(B_1)}^2 = \frac{\int_{B_1} |D^2 u_{k_m}(r_m x) - \overline{D^2 u_{k_m}}_{B_{r_m}}|^2}{\Theta(r_m)} = \frac{1}{2},$$

by definition.

Using the estimate (2.3) (the case $p = 2$), we deduce $\frac{1}{2} \leq \|D^2 v_m\|_{L^2(B_1)} \leq C(\|v_m\|_{L^2(B_2)} + \|g_m\|_{L^2(B_2)})$ and therefore, for m large enough,

$$\|v_m\|_{L^2(B_2)} \geq c_0 > 0.$$

Finally, for any fixed $R > 1$ we have:

$$\begin{aligned} \int_{B_R} |D^2 v_m(x) - \overline{D^2 v_m}_{B_R}|^2 &\leq \frac{\int_{B_R} |D^2 u_{k_m}(r_m x) - \overline{D^2 u_{k_m}(r_m x)}_{B_R}|^2}{\Theta(r_m)} \\ &\leq \frac{\int_{B_{Rr_m}} |D^2 u_{k_m}(y) - \overline{D^2 u_{k_m}}_{B_{Rr_m}}|^2}{\Theta(r_m)} \leq 1. \end{aligned}$$

Since $D^2 v_m$ is bounded in L^2 norm on compact sets, then by Poincaré inequality v_m is bounded in $W^{2,2}$ and hence there exists a weakly convergent series in $W^{2,2}$, such that $v_m \rightarrow v$ and $\nabla v_m \rightarrow \nabla v$ strongly in the L^2 norm, while $D^2 v_m \rightarrow D^2 v$ weakly in L^2 norm, for every compact set. Moreover, since $g_m \rightarrow 0$ strongly in L^2 , we have that

$$\Delta v = 0 \quad \text{in } \mathbb{R}^n.$$

Making use of the definition of weak convergence, and lower semicontinuity of the $W^{2,2}$ norm, we find that v satisfies the following conditions:

$$\overline{v}_{B_1} = \overline{\nabla v}_{B_1} = \overline{D^2 v}_{B_1} = 0 \tag{2.6}$$

$$\|v\|_{L^2(B_2)} \geq c_0 > 0 \tag{2.7}$$

$$\|D^2 v\|_{L^2(B_R)}^2 \leq R^n(1 + R^n) \text{ for any } R > 1. \tag{2.8}$$

$$\int_{B_R} |D^2 v(x) - \overline{D^2 v}_{B_R}|^2 \leq 1 \text{ for any } R > 1. \tag{2.9}$$

Using the mean value property, for any $x \in B_{R/2}$ and any $R > 1$ we have

$$|D^2 v(x)| \leq \int_{B_{R/2}(x)} |D^2 v| \leq \frac{C}{R^{n/2}} \|D^2 v\|_{L^2(B_R)} \leq CR^{3n/2}.$$

Hence, $D^2 v$ is a harmonic with polynomial growth, and the Liouville theorem implies that it is a polynomial. However, the first condition (2.6) and again the mean value property

$$\int_{B_R} D^2 v(x) = \int_{B_1} D^2 v(x) = 0,$$

meaning that the last condition (2.9) yields $\int_{B_R} |D^2 v|^2 \leq 1$ for any $R > 1$. The last two conditions yield that the polynomial $D^2 v$ must be identically zero, which means that v is an affine function. Again making use of conditions (2.6) we reach that $v \equiv 0$, which contradicts (2.7). This means that the estimate (2.4) must hold. \square

Finally, we provide the:

Proof of Theorem 1.1. We split the proof into two steps.

Step 1. We first prove the result for $2 < p < \infty$.

Let u be a solution in B_1 from $\Delta u = f$, with $f \in L^p(B_1)$. Let $\eta \in C_c^\infty(B_1)$ be any smooth function with $\eta \geq 0$ and $\int_{B_1} \eta = 1$ and $\eta_\varepsilon = \varepsilon^{-n} \eta(x/\varepsilon)$.

Then, the function $u_\varepsilon(x) = u * \eta_\varepsilon(x)$ is C^∞ and satisfies

$$\Delta u_\varepsilon = f * \eta_\varepsilon =: f_\varepsilon \quad \text{in } B_{1-\varepsilon}.$$

Since u_ε is C^∞ we can use the estimates (2.2), (2.3), and Proposition 2.1 to get:

$$\|D^2 u_\varepsilon\|_{L^p(B_{1/2})} \leq C(\|\mathcal{M}_2^\# D^2 u_\varepsilon\|_{L^p(B_1)}^{1/2} + \|D^2 u_\varepsilon\|_{L^1(B_{1/2})})$$

$$\leq C(\|u_\varepsilon\|_{L^2(B_1)} + \|f_\varepsilon\|_{L^2(B_1)} + \|f_\varepsilon\|_{L^p(B_1)}).$$

Thanks to Young's convolution inequality for L^p norms we have $\|u_\varepsilon\|_{L^2(B_1)} \leq \|u\|_{L^2(B_1)}$ and $\|f_\varepsilon\|_{L^p(B_1)} \leq \|f\|_{L^p(B_1)}$. The result then follows by letting $\varepsilon \rightarrow 0$ and using the lower semicontinuity of the $W^{2,p}$ norm.

Step 2. The result for $1 < p < 2$ then follows from Step 1 and a standard duality argument, which we sketch next. First, it suffices prove the estimate

$$\|D^2u\|_{L^p(\mathbb{R}^n)} \leq C\|f\|_{L^p(\mathbb{R}^n)}, \quad 1 < p < 2,$$

for global solutions $u \in C_c^\infty(\mathbb{R}^n)$ of $\Delta u = f$ in \mathbb{R}^n . Once we have this, the desired estimate follows easily by combining this with the interior regularity of harmonic functions.

Recall that the L^p -norm of a function f can be characterized by

$$\|f\|_{L^p(\Omega)} = \sup_{g \in L^{p'}(\Omega)} \left\{ \int_{\Omega} fg : \|g\|_{L^{p'}(\Omega)} = 1 \right\}.$$

Given u as above, consider a test function $g \in C_c^\infty(\mathbb{R}^n)$ with $\Delta v = g$ and integrate by parts to get

$$\int_{\mathbb{R}^n} D^2u g = \int_{\mathbb{R}^n} f D^2v \leq \|f\|_{L^p(\mathbb{R}^n)} \|D^2v\|_{L^{p'}(\mathbb{R}^n)} \leq C\|f\|_{L^p(\mathbb{R}^n)} \|g\|_{L^{p'}(\mathbb{R}^n)},$$

where we used the result in Step 1, for $p' > 2$

Taking the supremum one reaches $\|D^2u\|_{L^p(\mathbb{R}^n)} \leq C\|f\|_{L^p(\mathbb{R}^n)}$, as desired. \square

Remark 2.3. Once we have the $W^{2,p}$ estimates for the Laplacian, one can prove by the method of “freezing coefficients” that the same result holds for general operators in non-divergence form with continuous coefficients, i.e.,

$$\operatorname{tr}(A(x)D^2u(x)) = \sum_{i,j=1}^n a_{ij}(x)\partial_{ij}u(x) = f(x) \quad \text{in } B_1,$$

where $A(x) = (a_{ij}(x))_{i,j}$ is uniformly elliptic and continuous; see e.g. [GT98, FR22].

3. PARABOLIC EQUATIONS

Our method can also be used to give a new proof for the following parabolic $W^{2,p}$ estimate:

Theorem 3.1. *Let $1 < p < \infty$ and u be any weak solution of*

$$\partial_t u - \Delta u = f \text{ in } Q_1,$$

with $f \in L^p(Q_1)$. Then D^2u and u_t are in L^p inside Q_1 and the following estimate holds:

$$\int_{Q_{1/2}} |D^2u|^p + |\partial_t u|^p \leq C \left(\int_{Q_1} |u|^p + \int_{Q_1} |f|^p \right), \quad (3.1)$$

where C depends only on n .

Here, $Q_r(x_o, t_o) = B_r(x_o) \times (t_o - \frac{r^2}{2}, t_o + \frac{r^2}{2})$ represents the parabolic cube. As before, the case $p = 2$ follows from an easy integration by parts argument [Ev10], and we will give a new proof for $p \neq 2$.

Parabolic sharp maximal function. We define the parabolic 2-sharp maximal function as

$$\mathcal{M}_{2,\text{par}}^\# w(x, t) = \sup_{r>0} \fint_{Q_r(x,t)} |w - \overline{w}_{Q_r(x,t)}|^2,$$

for which the analogous bounds to (2.2) in Q_r hold; see [GM12].

Regularity estimates. In order to prove Theorem 3.1 we follow the steps as in the elliptic case. First, we will show an equivalent result for the non-homogeneous heat equation:

Proposition 3.2. *Let $u, f \in C^\infty(Q_1)$, with $u_t - \Delta u = f$. Then,*

$$\mathcal{M}_{2,\text{par}}^\# \partial_t u(x, t) + \mathcal{M}_{2,\text{par}}^\# D^2 u(x, t) \leq C(\|u\|_{L^2(Q_1)}^2 + \|f\|_{L^2(Q_1)}^2 + \mathcal{M}_{2,\text{par}}^\# f(x, t)), \quad (3.2)$$

for any $(x, t) \in Q_{1/2}$, where C depends only on n .

Proof. Following the same steps and calculations as in the elliptic case, we will see through contradiction that the result holds. Assume as before that $(x_\circ, t_\circ) = (0, 0)$, and suppose that the estimate (3.2) does not hold. Then, there exist a set of functions $u_k, f_k \in C^\infty(Q_1)$, such that $\partial_t u_k - \Delta u_k = f_k$ in Q_1 and:

$$\mathcal{M}_{2,\text{par}}^\# \partial_t u_k(0, 0) + \mathcal{M}_{2,\text{par}}^\# D^2 u_k(x_0, t_0) > k(\|u_k\|_{L^2(B_1)}^2 + \|f_k\|_{L^2(B_1)}^2 + \mathcal{M}_{2,\text{par}}^\# f_k(0, 0))$$

Moreover we can take $\|u_k\|_{L^2(B_1)}^2 + \|f_k\|_{L^2(B_1)}^2 + \mathcal{M}_{2,\text{par}}^\# f_k(0, 0) \leq 1$. Rewrite the inequality as:

$$\sup_k \sup_{r>0} \fint_{Q_r} \left[|D^2 u_k(x) - \overline{D^2 u_k}_{Q_r}|^2 + |\partial_t u_k(x) - \overline{\partial_t u_k}_{Q_r}|^2 \right] \rightarrow \infty,$$

and exactly as in the elliptic case, there exist sequences $r_m \rightarrow 0$ and k_m such that the new functions

$$v_m(x) = \frac{u_{k_m}(r_m x, r_m^2 t) - p_m(x, t)}{r_m^2 \Theta^{1/2}(r_m)}, \quad g_m(x) = \frac{f_{k_m}(r_m x, r_m^2 t) - \overline{f_{k_m}}_{Q_{r_m}}}{\Theta^{1/2}(r_m)}.$$

satisfy

$$\fint_{Q_\rho} |D^2 u_k - \overline{D^2 u_k}_{Q_\rho}|^2 + |\partial_t u_k - \overline{\partial_t u_k}_{Q_\rho}|^2 \leq 2 \fint_{Q_{r_m}} |D^2 u_{k_m} - \overline{D^2 u_{k_m}}_{Q_{r_m}}|^2 + |\partial_t u_{k_m} - \overline{\partial_t u_{k_m}}_{Q_{r_m}}|^2 =: \Theta(r_m)$$

for all $k \in \mathbb{N}$, $\rho \geq r_m$.

Here, $p_m(x, t)$ denotes a quadratic polynomial in x and linear in t such that $\overline{v_m}_{Q_1} = \overline{\nabla v_m}_{Q_1} = \overline{D^2 v_m}_{Q_1} = \overline{\partial_t v_m}_{Q_1} = 0$. Notice once more that $\partial_t v_m - \Delta v_m = g_m$ in $B_{1/2r_m}$.

Through the same calculations as in the elliptic case, we find the following bounds:

$$\|D^2 v_m\|_{L^2(Q_R)}^2 + \|\partial_t v_m\|_{L^2(Q_R)}^2 \leq 2R^{2+n}(1 + R^{2+n}),$$

while for our g_m we have

$$\|g_m\|_{L^2(Q_R)}^2 \leq \frac{R^{2+n}(1 + R^{2+n})\mathcal{M}_{2,\text{par}}^\# f_{k_m}(0, 0)}{\Theta(r_m)} \rightarrow 0,$$

for any fixed $R > 1$. Note as well that

$$\|D^2 v_m\|_{L^2(Q_1)}^2 + \|\partial_t v_m\|_{L^2(Q_1)}^2 \geq \frac{1}{2},$$

which leads to $\frac{1}{2} \leq \|D^2 v_m\|_{L^2(Q_1)} + \|\partial_t v_m\|_{L^2(Q_1)} \leq C(\|v_m\|_{L^2(Q_2)} + \|g_m\|_{L^2(Q_2)})$ thanks to the case $p = 2$. For a large enough m , it means that $\|v_m\|_{L^2(Q_2)} \geq c_0 > 0$. Finally, following the same steps, for any fixed $R > 1$ we have:

$$\int_{Q_R} \left[|D^2 v_m(x, t) - \overline{D^2 v_m}_{Q_R}|^2 + |\partial_t v_m(x, t) - \overline{\partial_t v_m}_{Q_R}|^2 \right] \leq 1.$$

Since $D^2 v_m, \partial_t v_m$ are bounded in the L^2 norm, then there exists a weakly convergent sequence, such that $v_m \rightarrow v$ strongly, and $\nabla v_m \rightarrow \nabla v$ and $\partial_t v_m \rightarrow \partial_t v$ weakly in L^2 norm because of the compact Sobolev embeddings. We see as well that $D^2 v_m \rightarrow D^2 v$ weakly in the L^2 norm and since $g_m \rightarrow 0$ strongly in L^2 , we reach that $\partial_t v - \Delta v = 0$ in $\mathbb{R}^n \times \mathbb{R}$.

Taking limits we find that

$$\overline{v}_{Q_1} = \overline{\nabla v}_{Q_1} = \overline{D^2 v}_{Q_1} = \overline{\partial_t v}_{Q_1} = 0. \quad (3.3)$$

$$0 < \|v\|_{L^2(Q_2)}. \quad (3.4)$$

$$\|D^2 v\|_{L^2(Q_R)}^2 + \|\partial_t v\|_{L^2(Q_R)}^2 \leq R^{2+n}(1 + R^{2+n}) \text{ for any } R > 1. \quad (3.5)$$

$$\int_{Q_R} \left[|D^2 v(x, t) - \overline{D^2 v}_{Q_R}|^2 + |\partial_t v(x, t) - \overline{\partial_t v}_{Q_R}|^2 \right] \leq 1 \text{ for any } R > 1. \quad (3.6)$$

By interior regularity for the heat equation, we can bound

$$\max_{Q_{R/2}} |D_x^k D_t^l D^2 v(x, t)| \leq \frac{C_{kl}}{R^{k+2l+n+2}} \|D^2 v\|_{L^1(Q_R)},$$

which with condition (3.5) means that $D^2 v$ and $\partial_t v$ have to be polynomials of degree N at most. By Lemma 3.3, this would mean that:

$$\int_{Q_R} \left[|D^2 v(x, t) - \overline{D^2 v}_{Q_R}|^2 + |\partial_t v(x, t) - \overline{\partial_t v}_{Q_R}|^2 \right] \geq cR^{2N} > 0 \quad \text{for any } R > 1.$$

which fulfils condition (3.6) only if $N = 0$. This means that $D^2 v$ and $\partial_t v$ are constants and that v is a quadratic polynomial in space and linear in time. By condition (3.3) we reach that $v \equiv 0$ which is a contradiction, and therefore the estimate (3.1) must hold. \square

We used this elementary result.

Lemma 3.3. *Let $p(x, t)$ be a polynomial of parabolic degree N . Then, large enough $R > 1$ we have*

$$\int_{Q_R} |p - c_R|^2 \geq cR^{2N},$$

for some constant $c > 0$.

Proof. Write $p = p_0 + \dots + p_N$, where each p_i is a parabolically homogeneous polynomial of degree $i = 0, \dots, N$. By triangle inequality,

$$\frac{1}{2} |p_N - c_R|^2 - |p - p_N|^2 \leq |p - c_R|^2 \leq 2|p - p_N|^2 + 2|p_N - c_R|^2.$$

Notice that $p - p_N$ is of degree $N - 1$ and therefore $\int_{Q_R} |p - p_N|^2 \leq CR^{2N-2}$. On the other hand, simple computations lead to

$$\int_{Q_R} |p_N(x, t) - c_R|^2 dx = \int_{Q_1} |p_N(Ry, R^2s) - c_R|^2 dy = R^{2N} \int_{Q_1} \left| p_N(y, s) - \frac{c_R}{R^N} \right|^2 \geq cR^{2N} > 0$$

for any constant c_R .

Hence, for large enough $R > 1$

$$\int_{Q_R} |p - c_R|^2 \geq cR^{2N},$$

and the Lemma follows. □

Finally, we sketch the:

Proof of Theorem 3.1. The proof is essentially the same as in the elliptic case, the only nontrivial difference is in the duality argument. Namely, once we have the result for $p > 2$, we fix $1 < p < 2$ and proceed as follows.

Consider $\Omega = \mathbb{R}^n \times (0, 1)$ and let $u \in C_c^\infty(\mathbb{R}^n \times [0, 1])$ be such that $\partial_t u - \Delta u = f$ in Ω , with $u(x, 0) = 0$. For any $g \in C_c^\infty(\Omega)$ let v be such that $\partial_t v - \Delta v = g$ with $v(x, 0) = 0$. Consider the same problem but backwards in time, i.e., $\tilde{g} = g(x, 1 - t)$ as well as $\tilde{v} = v(x, 1 - t)$ which satisfies $-\partial_t \tilde{v} - \Delta \tilde{v} = \tilde{g}$ and $\tilde{v}(x, 1) = 0$.

Then, integrating by parts we find

$$\begin{aligned} \int_{\Omega} \partial_t u \tilde{g} &= - \int_{\Omega} \partial_t u \partial_t \tilde{v} - \int_{\Omega} \partial_t u \Delta \tilde{v} = - \int_{\Omega} (\partial_t u - \Delta u) \partial_t \tilde{v} + \left[\int_{\mathbb{R}^n} u \Delta \tilde{v} \right]_0^1 = - \int_{\Omega} f \partial_t \tilde{v} \leq \\ &\leq \|f\|_{L^p(\Omega)} \|\partial_t \tilde{v}\|_{L^{p'}(\Omega)} = C \|f\|_{L^p(\Omega)} \|\tilde{g}\|_{L^{p'}(\Omega)}. \end{aligned}$$

Finally, take the supremum on the left hand side to reach the desired estimate for $\partial_t u$ is in $L^p(\Omega)$. By definition of f , it means that Δu is bounded in $L^p(\Omega)$ as well, and therefore that $D^2 u$ is in $L^p(\Omega)$. This, combined with interior regularity for the heat equation, yields the result for $1 < p < 2$. □

REFERENCES

[Ca89] L. Caffarelli, *Interior a priori estimates for solutions of fully nonlinear equations*, Ann. of Math. **130** (1989), 189-213.

[CZ52] A. Calderón, A. Zygmund, *On the existence of certain singular integrals*, Acta Math. **88**, (1952), 85-139.

[Ev10] L. Evans, *Partial Differential Equations, 2nd ed.*, Graduate studies in mathematics 19, AMS, 2010

[FR22] X. Fernández-Real, X. Ros-Oton, *Regularity Theory For Elliptic PDE*, EMS Press, Zurich, 2022.

[GM12] M. Giaquinta, L. Martinazzi, *An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs. 2nd ed.*, Edizioni della Normale Pisa, Pisa, 2012.

[GT98] D. Gilbarg, N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics 224, Springer, 1998.

[Le13] G. Leoni, *Partial Differential Equations I*, Carnegie Mellon University, 2013, Lecture Notes available at the webpage of the author.

[LZ24] Y. Lian, K. Zhang, *A note on the BMO and Calderón-Zygmund estimate*, Collect. Math. **75**, (2024), 1-8.

[LL01] E. Lieb, M. Loss, *Analysis, 2nd ed.*, Graduate studies in mathematics 14, AMS, 2001.

[Sel15] J. Serra, *Regularity for fully nonlinear nonlocal parabolic equations with rough kernels*, Calc. Var. Partial Differential Equations **54**, (2015), 615-629.

[Si97] L. Simon, *Schauder estimates by scaling*, Calc. Var. Partial Differential Equations **5** (1997), 391-407.

[ST89] J. Strömberg, A. Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Mathematics, Springer, 1989.

[Wa13] L. Wang, *A Geometric Approach to the Calderón-Zygmund Estimates*, Acta Math. Sin. **19**, (2003), 381-396.

ICREA, PG. LLUÍS COMPANYS 23, 08010 BARCELONA, SPAIN & UNIVERSITAT DE BARCELONA, DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, GRAN VIA DE LES CORTS CATALANES 585, 08007 BARCELONA, SPAIN & CENTRE DE RECERCA MATEMÀTICA, BARCELONA, SPAIN

Email address: `xros@icrea.cat`