

# Some Novel Results on $(\alpha, \beta)$ -Ricci-Yamabe Soliton and its Spacetime

<sup>1</sup> Pankaj Pandey and <sup>2</sup> Kamakshi Sharma

<sup>1,2</sup>Department of Mathematics (SCEPS)

Lovely Professional University, Phagwara, Punjab-144411, India

<sup>1</sup> Email: pankaj.anvarat@gmail.com

<sup>2</sup> Email: kamakshisharma82@gmail.com

**Abstract:** This article aims to investigate the characteristics of  $(\alpha, \beta)$ -Ricci-Yamabe Soliton (briefly:  $(\alpha, \beta) - (RYS)_n$ ) and its spacetime. The inclusion of killing vector field and the Lorentzian metrics make the Ricci-Yamabe soliton richer and interesting. We study the cosmological and dust fluid model on  $(RYS)_4$  equipped with Lorentzian para Sasakian  $(LPS)_4$  spacetime. The cases of  $\eta$ -parallel Ricci tensor and the Poisson structure have been studied on  $(RYS)_n$  equipped with  $(LPS)_n$  manifold. Gradient  $(RYS)_n$  equipped with  $(LPS)_n$  manifold also reveal. Finally, we establish an example of four-dimensional LP-Sasakian manifold  $(LPS)_4$  that satisfy  $(\alpha, \beta) - (RYS)_4$  and some results.

**AMS Mathematics Subject Classification (2010):** 53C25.

**Keywords:** Ricci-Yamabe soliton; Gradient Ricci-Yamabe soliton; LP Sasakian manifold; Poisson manifold; Partial differential equation; Ricci flow; Heat Equation; Cosmological model; Perfect fluid; Killing vector field; Parallel Ricci tensor.

## 1 Introduction

Hamilton [20] revealed the concept of Ricci flow (resp. Yamabe flow) in the last quarter of twenties in order to discuss new striking results in Riemannian geometry. The idea of Ricci soliton recognized as a generalization of an Einstein metric and governing as the solution of partial differential equation representing Ricci flow which is isomorphically equivalent to heat equation. In case of Ricci flow, the Riemannian metric is proportional to a  $(0, 2)$  type tensor  $(\frac{1}{2}Lg + Ric)$ . In this case the proportionality constant is called the soliton constant. The Ricci flow in terms of soliton constant  $\lambda$  (say) is governed by the equation [20]

$$(1.1) \quad -\frac{1}{2}(L_v g)(\zeta_1, \zeta_2) = S(\zeta_1, \zeta_2) + \lambda g(\zeta_1, \zeta_2),$$

which is equivalent to

$$(1.2) \quad \frac{\partial g(t)}{\partial t} = -2S(t), \quad g_0 = g(0).$$

The Yamabe flow is discribed by an evolving equation [17, 20]

$$(1.3) \quad \frac{\partial g(t)}{\partial t} = -r(t)g(t), \quad g_0 = g(0),$$

equivalently

$$(1.4) \quad \frac{1}{2}(L_v g)(\zeta_1, \zeta_2) = (r - \lambda)g(\zeta_1, \zeta_2).$$

In (1.1) and (1.4), the Lie derivative  $L$  is taken along the complete vector field  $v$  and is denoted by  $L_v g$ . The Ricci denotes  $S$  and the  $r$  is scalar tensor. The vector fields  $\zeta_1, \zeta_2$  belong to the set of an algebra of tangent vectors denoted by  $\chi(M)$ . The nature of the soliton can be expressed in terms of soliton constant  $\lambda$  and is said to explore for  $\lambda$  positive. We say the case is of compacting soliton or is of constant soliton, if  $\lambda < 0$  or  $\lambda = 0$ . In general the Ricci and the Yamabe solitons are distinct for higher dimension but coincide in case of dimension 2. The reason behind this distinction is that the Yamabe soliton preserves the metric conformality in nature where as Ricci soliton denied it.

Guler and Crasmareanu [19] (2019) investigated another geometric flow restricted by Ricci-Yamabe map. Authors pronounced with a special call:  $(\alpha, \beta)$  type Ricci-Yamabe flow (briefly denoted  $(\alpha, \beta) - (RYF)_n$ ). Particularly, it is noted that the  $(\alpha, \beta) - (RYF)_n$  is nothing but the  $\alpha$ -Ricci soliton for  $\beta = 0$  and it turns into  $\beta$ -Yamabe soliton, in case of vanishing  $\alpha$ . The above said flow, in the Riemannian manifold, is defined by

$$(1.5) \quad \frac{\partial}{\partial t}g(t) = \beta r(t)g(t) - 2\alpha S(t), \quad g_0 = g(0),$$

where  $S$  being Ricci tensor,  $r$ : the scalar curvature, and  $\lambda, \alpha, \beta \in R$ . As a restriction to solve the  $(\alpha, \beta) - (RYF)_n$ , the solution of nonlinear partial differential equation (1.5) notified as  $(\alpha, \beta) - (RYS)_n$ .

If we denote  $(M^n, g)$  by an  $n$ -dimensional  $(\alpha, \beta) - (RYS)_n$ , on  $(M^n, g)$  having structure  $(g, V, \lambda, \alpha, \beta)$  that gratify the equation

$$(1.6) \quad (L_V g)(\zeta_1, \zeta_2) = -2\alpha S(\zeta_1, \zeta_2) - (2\lambda - \beta r)g(\zeta_1, \zeta_2).$$

The aforementioned soliton is known as the  $(\alpha, \beta)$  type gradient Ricci-Yamabe soliton (briefly  $(\alpha, \beta) - (GRYS)_n$ ) if  $V$  becomes a gradient of  $f$  (if exist such a smooth function) and (1.6) reveals

$$(1.7) \quad \nabla^2 f + \alpha S = (\lambda - \frac{1}{2}\beta r)g,$$

where  $\nabla^2 f$  is defined as Hessian of the smooth function  $f$  and denoted in general by  $Hess(f)$ . The nature of the  $(\alpha, \beta) - (RYS)_n$  is quite similar and depends on the soliton constant  $\lambda$ . Accordingly, the respective soliton is called steady for  $\lambda = 0$  otherwise natured as shrinking for being  $\lambda$  negative. The positiveness of  $\lambda$  explored as expanding.

An  $(\alpha, \beta) - (RYS)_n$  turns into almost  $(\alpha, \beta) - (RYS)_n$  (briefly  $(\alpha, \beta) - (ARYS)_n$ ) for  $(\lambda, \alpha, \beta)$  being smooth function. The  $(\alpha, \beta) - (RYS)_n$  simplifying as Ricci soliton in special case  $(\alpha = 1, \beta = 0)$  and it turns into Yamabe soliton in other special case

$(\alpha = 0, \beta = 1)$ . Also, for  $(\alpha = 1, \beta = -1)$  and  $(\alpha = 1, \beta = -2\rho)$ , it pointed into Einstein and  $\rho$ -Einstein soliton. Some times, we call it proper for the values  $\alpha \neq 0, 1$ . It is important to keep in mind that the Ricci Yamabe soliton corresponds to a specific class of generalized Ricci solitons [16] if scalar curvature  $r$  is constant in (1.6). Therefore, studying Ricci Yamabe soliton with non-constant scalar curvature rather than the constant scalar curvature is more appropriate and natural.

The Lorentzian manifold, which is among the most significant sub classes of pseudo-Riemannian manifold, has a significant impact on the advancement of the theory of relativity and cosmology. Ahsan and Ali [1] looked into the symmetries of soliton spacetime. The geometrical characteristics discussed by Blaga [3] in fluid spacetime under assumptions of Einstein/Ricci soliton. In [3] author examined the geometrical importance of  $(PFST)_n$  conformal Ricci solitons. Chaturvedi et.al. studied the Kähler spacetime manifold under consideration of Bochner flatness. They determined that the energy-momentum tensor of a perfect fluid Lorentzian Kähler spacetime manifold exhibits hybrid characteristics. They also analyzed the behaviour of dust fluid Lorentzian Kähler spacetime manifold where the Bochner curvature vanishes (see also authored by Pokhriyal and Chaubey [37]).

The LP-Kenmotsu manifold was investigated by Haseeb and Prasad in [23]. Haseeb et.al. [26] proposed the study on  $(\alpha, \beta) - (RYS)_n$  and  $(\alpha, \beta) - (GRYS)_n$  conditionally related to the  $\zeta$ -conformally flat  $m$ -dimensional LP Kenmotsu manifold. They proved that the scalar tensor of  $(LPK)_m$  manifold admitting Ricci Yamabe soliton satisfies the Poisson equation. Sardar and Sarkar [39] described Kenmotsu 3-manifold equipped with  $(\alpha, \beta) - (RYS)_n$  and  $(\alpha, \beta) - (GRYS)_n$  metric that satisfy  $\zeta$ -parallel Ricci tensor. Recently, Pal and Chaudhary [31] revealed the idea of Poisson fluid flow to investigate almost soliton like Ricci soliton (briefly:  $(ASLRS)_n$ ) by exploring the statistical structure.

De and De [13] (2020) studied almost Ricci soliton and gradient almost Ricci soliton on para-Sasakian manifolds. Kundu [28] (2021) explored para-Kenmotsu metric as an  $\eta$ -Ricci soliton. In [11] (2021), De discussed Sasakian 3-manifolds admitting a gradient Ricci-Yamabe soliton. Sarkar and Bhakta [40] obtained some interesting results on certain solitons on generalized  $(\kappa, \mu)$  contact metric manifolds. D. Kar and P. Majhi [27] (2019) concentrated on Beta-almost Ricci solitons on almost Co-Kähler manifolds. Vishwas, Das, Baishya and Bakshi [43] studied  $\eta$ -Ricci solitons on Kenmotsu manifolds admitting general connection. T. Mandal [?] (2022) given certain results on three-dimensional  $f$ -Kenmotsu manifolds with conformal Ricci solitons. Lee, Kim and Choi [29] classified the warped product spaces with gradient Ricci solitons. We propose the papers ([2, 4, 6, 7, 10, 17, 18, 22, 23, 24, 25, 30, 32, 33, 34, 35, 36]) for deep understanding and various interesting results on respective solitons and various examples.

Motivated by the above research we study  $(\alpha, \beta) - (RYS)_n$  and  $(\alpha, \beta) - (GRYS)_n$  on  $(LPS)_n$ . The preliminaries discussed in section 2. After that we study  $(\alpha, \beta) - (RYS)_n$  in section 3. In section 4 we study  $(\alpha, \beta) - (RYS)_n$  in perfect fluid flow. Further in section 5 and 6, we deal with the cosmological models. Section 7 has been explored with  $\eta$ -parallel Ricci tensor and in section 8, we study Poisson manifold admitting  $(\alpha, \beta) - (RYS)_n$ . The gradient case  $(\alpha, \beta) - (GRYS)_n$  has been studied in section 9. The last section includes an example of  $(\alpha, \beta) - (RYS)_n$  admitting  $(LPS)_n$ .

## 2 Preliminaries

Let  $M^n$  (Riemannian manifold) denotes the inclusion of the triplet data  $(\phi, \xi, \eta)$  then we say  $(M^n, \phi, \xi, \eta)$  is an almost contact metric structure that satisfy

$$(2.1) \quad \begin{aligned} \phi\xi &= 0, \\ \eta(\phi\zeta_1) &= 0, \\ \eta(\xi) &= -1, \\ \phi^2\zeta_1 &= \zeta_1 + \eta(\zeta_1)\xi. \end{aligned}$$

Here,  $\phi$  denotes a tensor field ((1, 1)-type) and  $\eta$  we call 1-form on  $M^n$  defined by  $\eta(\zeta_1) = g(\zeta_1, \xi)$ . Some additional following properties also hold on almost contact metric manifold.

$$(2.2) \quad g(\phi\zeta_1, \phi\zeta_2) = g(\zeta_1, \zeta_2) - \eta(\zeta_1)\eta(\zeta_2)$$

equivalently

$$(2.3) \quad \begin{aligned} g(\zeta_1, \phi\zeta_2) &= -g(\phi\zeta_1, \zeta_2) \\ g(\zeta_1, \xi) &= \eta(\zeta_1), \end{aligned}$$

for all the vector fields  $\zeta_1, \zeta_2 \in \chi(M)$ .

The structure  $(M^n, \phi, \xi, \eta, g)$  poured to  $K$ -contact, for being killing to characteristic vector field  $\xi$ . For Lorentzian para Sasakian  $(LPS)_n$  manifold with killing vector field, we have

$$(2.4) \quad \nabla_{\zeta_1}\xi = \phi\zeta_1,$$

$$(2.5) \quad (L_\xi g)(\zeta_1, \zeta_2) = 0,$$

$$(2.6) \quad \phi\xi = 0, \quad \eta(\phi\zeta_1) = 0, \quad \text{rank } \phi = n - 1.$$

Also

$$(2.7) \quad (\nabla_{\zeta_1}\eta)(\zeta_2) = \nabla_{\zeta_1}\eta(\zeta_2) - \eta(\nabla_{\zeta_1}\zeta_2) = g(\zeta_2, \nabla_{\zeta_1}\zeta_2).$$

In view of (2.4) and (2.7) we have

$$(2.8) \quad (\nabla_{\zeta_1}\eta)(\zeta_2) = g(\zeta_2, \phi\zeta_1).$$

Let structure  $(\phi, \xi, \eta, g)$  on  $(LPS)_n$   $M^n$  ( $n$  being dimension) then

$$(2.9) \quad g(R(\zeta_1, \zeta_2)\zeta_3, \xi) = \eta(R(\zeta_1, \zeta_2)\zeta_3) = g(\zeta_2, \zeta_3)\eta(\zeta_1) - g(\zeta_1, \zeta_3)\eta(\zeta_2),$$

$$(2.10) \quad R(\xi, \zeta_1)\zeta_2 = g(\zeta_1, \zeta_2)\xi - \eta(\zeta_2)\zeta_1,$$

$$(2.11) \quad R(\zeta_1, \zeta_2)\xi = \eta(\zeta_2)\zeta_1 - \eta(\zeta_1)\zeta_2,$$

$$(2.12) \quad R(\xi, \zeta_1)\xi = \zeta_1 + \eta(\zeta_1)\xi,$$

$$(2.13) \quad S(\zeta_1, \xi) = (n - 1)\eta(\zeta_1),$$

$$(2.14) \quad S(\phi\zeta_1, \phi\zeta_2) = S(\zeta_1, \zeta_2) + (n - 1)\eta(\zeta_1)\eta(\zeta_2),$$

for the vector fields  $\zeta_1, \zeta_2, \zeta_3$ , where  $R$  and  $S$  denote the tensor of Riemann and Ricci.

### 3 $(\alpha, \beta) - (RYS)_n$ on $(M^n, \phi, \xi, \eta, g)$

**Lemma 3.1.** *Let  $M^n$  be an Lorentzian para Sasakian  $(LPS)_n$  manifold. If the  $(\alpha, \beta) - (RYS)_n$  admits an  $(LPS)_n$  structure then it is Einstein manifold and its Ricci curvature is given by equation (3.1).*

*Proof.* Taking  $V = \xi$  in (1.6) and using (2.5), we get

$$(3.1) \quad S(\zeta_1, \zeta_2) = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) g(\zeta_1, \zeta_2).$$

Hence proved.  $\square$

**Lemma 3.2.** *Let  $M^n$  be an  $(LPS)_n$  manifold. If the  $(\alpha, \beta) - (RYS)_n$  admits an  $(LPS)_n$  structure then (3.2), (3.3), (3.4), (3.5) and (3.6) hold good.*

*Proof.* From (3.1), it is very obvious to obtain the following results:

$$(3.2) \quad S(\zeta_1, \xi) = S(\xi, \zeta_1) = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) \eta(\zeta_1),$$

$$(3.3) \quad S(\xi, \xi) = \left( \frac{2\lambda - \beta r}{2\alpha} \right),$$

$$(3.4) \quad Q\zeta_1 = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) \zeta_1,$$

$$(3.5) \quad Q\xi = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) \xi,$$

$$(3.6) \quad SCA_{(\alpha, \beta) - RYS} = n \left( \frac{\beta r - 2\lambda}{2\alpha} \right).$$

$SCA_{(\alpha, \beta) - RYS}$  denotes the scalar curvature of  $(\alpha, \beta) - (RYS)_n$  admitting  $(LPS)_n$ .  $\square$

**Theorem 3.1.** *Let  $(\alpha, \beta) - (RYS)_n$  admits an  $(LPS)_n$  manifold having a non-constant scalar then  $\alpha$  and  $\beta$  are related by  $\alpha = \frac{n\beta}{2}$ .*

*Proof.* The derivative (covariant) of (3.4) yields

$$(3.7) \quad (\nabla_K Q\zeta_1) = \frac{\beta}{2\alpha} (Kr)\zeta_1.$$

Applying the metric both side, above equation reduces to

$$(3.8) \quad g((\nabla_K Q)(\zeta_1, \zeta_2)) = \frac{\beta}{2\alpha} (Kr)g(\zeta_1, \zeta_2).$$

Arranging  $\zeta_1 = \zeta_2 = e_i$  in (3.8) and contracting, we get

$$(3.9) \quad g((\nabla_K Q)e_i, e_i) = n \frac{\beta}{2\alpha}(Kr),$$

which implies

$$(3.10) \quad (Kr) \left( \frac{n\beta}{2\alpha} - 1 \right) = 0.$$

For being non-constant scalar curvature  $r$ , we have the result.  $\square$

**Theorem 3.2.** *Let  $(\alpha, \beta)$ – $(RYS)_n$  admits an  $(LPS)_n$  manifold having a non-constant scalar curvature then soliton function is given by*

$$\lambda = \frac{\beta}{2}(r - n(n-1)).$$

*Proof.* Using (2.5) in (1.6), we get

$$(3.11) \quad 2\alpha S(\zeta_1, \zeta_2) = (\beta r - 2\lambda)g(\zeta_1, \zeta_2).$$

Substituting  $\zeta_2 = \xi$  in (3.11) and then applying (2.13), result obtain

$$(3.12) \quad 2\alpha(n-1)\eta(\zeta_1) = (\beta r - 2\lambda)\eta(\zeta_1).$$

Since,  $\eta(\zeta_1)$  is non-vanishing, the above equation produces

$$(3.13) \quad \lambda = \frac{\beta r}{2} - (n-1)\alpha.$$

With the help of theorem (3.1), equation (3.13) yields

$$(3.14) \quad \lambda = \frac{\beta}{2}(r - n(n-1)),$$

which completes the proof.  $\square$

## 4 $(\alpha, \beta) - (RYS)_4$ Admitting $(LPS)_4$ Space-time

In this section, we consider the  $(LPS)_4$  spacetime to study. The Einstein's field equation (*briefly: EFE*) defines the following relation

$$(4.1) \quad S(\zeta_1, \zeta_2) - \frac{r}{2}g(\zeta_1, \zeta_2) + \mu g(\zeta_1, \zeta_2) = \tau T(\zeta_1, \zeta_2).$$

Here,  $\mu$  denotes the cosmological term and  $\zeta_1, \zeta_2$ , the vector fields.  $\tau$  being the gravitational constant and  $T$  deals for  $(0, 2)$  type energy momentum tensor (energy tensor). The below equation reveals the information regarding perfect fluid and govern by

$$(4.2) \quad T(\zeta_1, \zeta_2) = (\rho - p)A(\zeta_1)A(\zeta_2) + pg(\zeta_1, \zeta_2).$$

Here, energy tensor and energy density function denote by  $T$ , whereas  $p$  indicates for fluid isotropic pressure. Also, we consider here a non-zero 1-form by  $A$  that define as  $g(\zeta_1, V) = A(\zeta_1)$ ;  $V$  being the flow vector field.

In our discussion, we consider  $\xi$  as the flow vector field and (4.2) changes in a new form as

$$(4.3) \quad T(\zeta_1, \zeta_2) = (\rho - p)\eta(\zeta_1)\eta(\zeta_2) + pg(\zeta_1, \zeta_2).$$

Substituting the value of Ricci tensor from (3.1) in (4.1), we have

$$(4.4) \quad T(\zeta_1, \zeta_2) = \frac{1}{\tau} \left[ \mu - \left( \frac{\beta r - 2\lambda}{2\alpha} \right) \right] g(\zeta_1, \zeta_2).$$

Using theorem (3.1), the energy tensor takes the form

$$(4.5) \quad T(\zeta_1, \zeta_2) = \frac{1}{4\tau} \left[ 4\mu - r + \frac{2\lambda}{\beta} \right] g(\zeta_1, \zeta_2).$$

The conclusion is below:

**Theorem 4.1.** *Let  $(\alpha, \beta) - (RYS)_4$  admits an  $(LPS)_4$  space-time with non-constant scalar curvature then energy tensor is an Einstein like tensor and given by*

$$T(\zeta_1, \zeta_2) = \frac{1}{4\tau} \left[ 4\mu - r + \frac{2\lambda}{\beta} \right] g(\zeta_1, \zeta_2).$$

**Theorem 4.2.** *Let  $(\alpha, \beta) - (RYS)_4$  admits an  $(LPS)_4$  space-time with non-constant scalar curvature then soliton function  $\lambda$  is given by*

$$\lambda = \frac{\beta}{2} \{ 4\tau(2p - \rho) + r - 4\mu \}.$$

*Proof.* Equating (4.3) and (4.5), we obtain

$$(4.6) \quad \frac{1}{4\tau} \left[ 4\mu - r + \frac{2\lambda}{\beta} \right] g(\zeta_1, \zeta_2) = (\rho - p)\eta(\zeta_1)\eta(\zeta_2) + pg(\zeta_1, \zeta_2).$$

Arranging  $\zeta_1 = \zeta_2 = \xi$ , the above equation (4.6) yields the soliton function

$$(4.7) \quad \lambda = \frac{\beta}{2} \{ 4\tau(2p - \rho) + r - 4\mu \}.$$

Hence the proof has been over. □

**Theorem 4.3.** *Let  $(\alpha, \beta)$  steady- $(RYS)_n$  admits an  $(LPS)_n$  space-time then non-constant scalar curvature is four times of cosmological scalar function  $\mu$ , that is:*

$$r = 4\mu.$$

*Proof.* For the steady soliton we know that  $\lambda$  vanishes. Therefore, using (4.2) we can calculate the value of cosmological constant  $\mu$  as

$$(4.8) \quad \mu = \frac{\beta r - 2\lambda}{2\alpha}.$$

Applying theorem (3.1), completes the proof. □

## 5 Cosmological Model on $(\alpha, \beta) - (RYS)_4$ Admitting $(LPS)_4$ Space-time

This section, contains the results on  $(LPS)_4$  space-time with considering a killing vector field. Here, we consider the (EFE), that does not consist cosmological term, given by

$$(5.1) \quad S(\zeta_1, \zeta_2) - \frac{r}{2}g(\zeta_1, \zeta_2) = \tau T(\zeta_1, \zeta_2).$$

Using equation (3.1), (4.3) and (5.1), we obtain

$$(5.2) \quad \left[ \frac{r(\beta - \alpha) - 2\lambda}{2\alpha} - \tau p \right] g(\zeta_1, \zeta_2) = \kappa(\rho + p)A(\zeta_1)A(\zeta_2).$$

Contracting  $\zeta_1$  and  $\zeta_2$  in (5.2), we get

$$(5.3) \quad r = \frac{\alpha\tau(-\rho + 3p) + 2\lambda}{2(\beta - \alpha)}.$$

Substituting (5.3) in (3.1), we get

$$(5.4) \quad S(\zeta_1, \zeta_2) = \left[ \frac{\beta\alpha\tau(-\rho + 3p) - 2\lambda(\beta - 2\alpha)}{4\alpha(\beta - \alpha)} \right] g(\zeta_1, \zeta_2).$$

As we know the Ricci operator  $Q$  is defined as

$$(5.5) \quad g(Q\zeta_1, \zeta_2) = S(\zeta_1, \zeta_2) \quad \text{and} \quad S(Q\zeta_1, \zeta_2) = S^2(\zeta_1, \zeta_2).$$

Thus, we have

$$(5.6) \quad A(Q\zeta_1) = g(Q\zeta_1, \xi) = S(\zeta_1, \xi).$$

Then from equation (5.4) and (5.5), we have

$$(5.7) \quad S(Q\zeta_1, \zeta_2) = S^2(\zeta_1, \zeta_2) = \left[ \frac{\beta\alpha\tau(-\rho + 3p) - 2\lambda(\beta - 2\alpha)}{4\alpha(\beta - \alpha)} \right]^2 g(\zeta_1, \zeta_2).$$

Contraction over  $\zeta_1$  and  $\zeta_2$  in (5.7), we have

$$(5.8) \quad \|Q\|^2 = \left[ \frac{\beta\alpha\tau(-\rho + 3p) - 2\lambda(\beta - 2\alpha)}{2\alpha(\beta - \alpha)} \right]^2.$$

Using theorem (3.1), we get

$$(5.9) \quad \|Q\|^2 = \frac{1}{16} \left[ \tau(\rho - 3p) - \frac{3\lambda}{\beta} \right]^2.$$

**Theorem 5.1.** *If the perfect fluid space-time on  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$  satisfy the (EFE) not consisting the cosmological term, the length of Ricci operator is given by*

$$\|Q\|^2 = \frac{1}{16} \left[ \tau(\rho - 3p) - \frac{3\lambda}{\beta} \right]^2.$$

Now, if we put  $\rho = 3p$  (the condition of perfect fluid to be radiation fluid) in equation (5.9), then we have

**Theorem 5.2.** *If the perfect fluid is radiation fluid (i.e  $\rho = 3p$ ) in  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$ , then length of Ricci operator is given by*

$$\|Q\|^2 = \left[ \frac{3\lambda}{4\beta} \right]^2.$$

## 6 Dust Fluid on $(\alpha, \beta) - (RYS)_4$ Admitting $(LPS)_4$ Space-time

The dust fluid and the energy tensor can be modelled by

$$(6.1) \quad T(\zeta_1, \zeta_2) = \rho A(\zeta_1)A(\zeta_2).$$

Using equation (5.1) and (6.1), the expression becomes

$$(6.2) \quad S(\zeta_1, \zeta_2) - \frac{r}{2}g(\zeta_1, \zeta_2) = \tau\rho A(\zeta_1)A(\zeta_2).$$

On contracting over  $\zeta_1$  and  $\zeta_2$  in above equation, we get

$$(6.3) \quad r = \tau\rho.$$

By using equation (6.3) in equation (3.1), the Ricci tensor of perfect fluid  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$  spacetime becomes

$$(6.4) \quad S(\zeta_1, \zeta_2) = \left[ \frac{\beta\tau\rho - 2\lambda}{2\alpha} \right] g(\zeta_1, \zeta_2).$$

Thus, by virtue of equation (5.7) and (6.4), we obtain

$$(6.5) \quad S(Q\zeta_1, \zeta_2) = \left[ \frac{\beta\tau\rho - 2\lambda}{2\alpha} \right]^2 g(\zeta_1, \zeta_2).$$

Contracting  $\zeta_1$  and  $\zeta_2$  in above equation, we have

$$(6.6) \quad \|Q\|^2 = \left[ \frac{\beta\tau\rho - 2\lambda}{2\alpha} \right]^2.$$

Using theorem (3.1), we have

$$(6.7) \quad \|Q\|^2 = \frac{1}{16} \left[ \tau\rho - \frac{2\lambda}{\beta} \right]^2.$$

**Theorem 6.1.** *If the dust cosmological model on  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$  space-time satisfy the (EFE) not consisting the cosmological term, the length of Ricci operator is given as:*

$$\|Q\|^2 = \frac{1}{16} \left[ \tau\rho - \frac{2\lambda}{\beta} \right]^2.$$

Now, using equation (3.1) and (6.2), we get

$$(6.8) \quad \left[ \frac{\beta r - 2\lambda - \alpha r}{2\alpha} \right] g(\zeta_1, \zeta_2) = \tau \rho A(\zeta_1) A(\zeta_2).$$

Contracting over  $\zeta_1$  and  $\zeta_2$  in equation (6.8), we get

$$(6.9) \quad r = \frac{-\alpha \rho \tau + 4\lambda}{2(\beta - \alpha)}.$$

By multiplying  $A(\zeta_3)$  in equation (6.8) and then taking contraction over  $\zeta_2$  and  $\zeta_3$ , we have

$$(6.10) \quad r = \frac{2(\alpha \rho \tau + \lambda)}{\beta - \alpha}.$$

From equation (6.9) and (6.10), we obtain

$$(6.11) \quad \alpha \rho \tau = 0.$$

As we know  $\tau$  is gravitational constant, therefore  $\tau \neq 0$ . Thus, we have either  $\alpha = 0$  or  $\rho = 0$ .

If  $\rho = 0$ , then from equation (6.1) and (6.11), we obtain

$$(6.12) \quad T(\zeta_1, \zeta_2) = 0.$$

Therefore, we can state the following theorem:

**Theorem 6.2.** *If the dust cosmological model on  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$  space-time satisfy the (EFE) not consisting the cosmological term the it is vacuum.*

## 7 $\eta$ -PRT on $(\alpha, \beta) - (RYS)_n$ Admitting $(LPS)_n$

This section examines  $\eta$ -parallel Ricci tensor (*briefly:  $\eta$ -PRT*) on  $(\alpha, \beta) - (RYS)_n$  admitting  $(LPS)_n$ . If an  $(LPS)_n$  consist  $\eta$ -PRT then

$$(7.1) \quad g((\nabla_K Q)\zeta_1, \zeta_2) = 0.$$

for all smooth vector fields  $\zeta_1, \zeta_2$ .

The following is the very general expansion of Ricci operator

$$(7.2) \quad (\nabla_K Q)\zeta_1 = \nabla_K Q\zeta_1 - Q(\nabla_K \zeta_1).$$

Using (3.4) in (7.2), yields

$$(7.3) \quad (\nabla_K Q)\zeta_1 = \frac{\beta}{2\alpha}(Kr)\zeta_1.$$

Using (7.3) in (7.1), we obtain

$$(7.4) \quad g((\nabla_K Q)\zeta_1, \zeta_2) = \frac{\beta}{2\alpha}(Kr)g(\zeta_1, \zeta_2).$$

In view of (3.10), we have  $Kr = 0$  only if  $(\frac{n\beta}{2\alpha} - 1) \neq 0$ .

**Theorem 7.1.** *There exists an  $(\alpha, \beta) - (RYS)_n$  admitting  $(LPS)_n$  with an  $\eta$ -parallel Ricci tensor only if  $(\frac{n\beta}{2\alpha} - 1) \neq 0$ .*

## 8 Poisson $(\alpha, \beta) - (RYS)_n$ Admitting $(LPS)_n$ Manifold

This section provides a definition of the statistical Poisson manifold and highlights several significant findings related to the statistical structure on the manifold.

**Definition 8.1.** The Poisson manifold consist the codazzi structure and explored by

$$(8.1) \quad (\nabla_{\zeta_3} S)(\zeta_1, \zeta_2) = (\nabla_{\zeta_2} S)(\zeta_3, \zeta_1).$$

Taking covariant derivative of (3.1) and using (2.4), we get

$$(8.2) \quad (\nabla_{\zeta_3} S)(\zeta_1, \zeta_2) = \frac{\beta}{2\alpha}(\zeta_3 r)g(\zeta_1, \zeta_2).$$

Also, we can calculate,

$$(8.3) \quad (\nabla_{\zeta_2} S)(\zeta_3, \zeta_1) = \frac{\beta}{2\alpha}(\zeta_2 r)g(\zeta_3, \zeta_1).$$

In view of (8.1), (8.2) and (8.3), we have

$$(8.4) \quad \frac{\beta}{2\alpha}(\zeta_3 r)g(\zeta_1, \zeta_2) = \frac{\beta}{2\alpha}(\zeta_2 r)g(\zeta_3, \zeta_1).$$

Put  $\zeta_1 = \xi$  in (8.4), we obtain

$$(8.5) \quad \frac{\beta}{2\alpha}[(\zeta_3 r)\eta(\zeta_2) - (\zeta_2 r)\eta(\zeta_3)] = 0.$$

Put  $\zeta_3 = \xi$  in (8.5) and using lemma (3.1), we have

$$(8.6) \quad \frac{\beta}{2\alpha}(\zeta_2 r) = 0.$$

So, we have either  $\frac{\beta}{2\alpha} = 0$  or  $(\zeta_2 r) = 0$ .

As defined earlier  $\alpha$  and  $\beta$  are non-vanishing constants, therefore  $\zeta_2 r = 0$  which is possible only if  $(\frac{n\beta}{2\alpha} - 1) \neq 0$ . Thus, we have

**Theorem 8.2.** *An  $(\alpha, \beta) - (RYS)_n$  admitting  $(LPS)_n$  is the statistical Poisson manifold only if  $(\frac{n\beta}{2\alpha} - 1) \neq 0$  and the scalar curvature is constant.*

## 9 Gradient $(\alpha, \beta) - (RYS)_n$ Admitting $(LPS)_n$ Manifold

Suppose that  $(\alpha, \beta) - (GRYS)_n$  admits  $(LPS)_n$  manifold then (1.7) turns to

$$(9.1) \quad \nabla_{\zeta_1} Df = (\lambda - \frac{1}{2}\beta r)\zeta_1 - \alpha Q\zeta_1.$$

The derivative (covariantly) of (9.1) along  $\zeta_2$ , produces

$$(9.2) \quad \nabla_{\zeta_2} \nabla_{\zeta_1} Df = (\lambda - \frac{1}{2}\beta r)\nabla_{\zeta_2} \zeta_1 - \frac{\beta}{2}(\zeta_2 r)\zeta_1 - \alpha \nabla_{\zeta_2} Q\zeta_1.$$

Interchanging  $\zeta_1$  and  $\zeta_2$  in (9.2) refer

$$(9.3) \quad \nabla_{\zeta_1} \nabla_{\zeta_2} Df = \left(\lambda - \frac{1}{2}\beta r\right) \nabla_{\zeta_1} \zeta_2 - \frac{\beta}{2} (\zeta_1 r) \zeta_2 - \alpha \nabla_{\zeta_1} Q \zeta_2.$$

It is quite simple to arrange the above information to form the following

$$(9.4) \quad R(\zeta_1, \zeta_2) Df = \frac{\beta}{2} [(\zeta_2 r) \zeta_1 - (\zeta_1 r) \zeta_2] - \alpha [(\nabla_{\zeta_1} Q) \zeta_2 - (\nabla_{\zeta_2} Q) \zeta_1].$$

On the other hand, equation (3.4) results as

$$(9.5) \quad (\nabla_{\zeta_1} Q) \zeta_2 - (\nabla_{\zeta_2} Q) \zeta_1 = \frac{\beta}{2\alpha} [(\zeta_1 r) \zeta_2 - (\zeta_2 r) \zeta_1].$$

Using (9.5) in (9.4), we get

$$(9.6) \quad R(\zeta_1, \zeta_2) Df = \frac{\beta}{2} [(\zeta_2 r) \zeta_1 - (\zeta_1 r) \zeta_2] - \alpha \left[ \frac{\beta}{2\alpha} [(\zeta_1 r) \zeta_2 - (\zeta_2 r) \zeta_1] \right],$$

which can be simplified as

$$(9.7) \quad R(\zeta_1, \zeta_2) Df = \beta [(\zeta_2 r) \zeta_1 - (\zeta_1 r) \zeta_2].$$

Contracting (9.7) and using lemma (3.1) we get

$$(9.8) \quad S(\zeta_2, Df) = n\beta(\zeta_2 r).$$

Alteration of  $\zeta_1$  by  $Df$  in (3.1), yields

$$(9.9) \quad S(\zeta_2, Df) = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) (\zeta_2 f).$$

In view of (9.8) and (9.9), we have

$$(9.10) \quad n\beta(\zeta_2 r) = \left( \frac{\beta r - 2\lambda}{2\alpha} \right) (\zeta_2 f).$$

Since  $\zeta_2 r = 0$  is possible only if  $\left( \frac{n\beta}{2\alpha} - 1 \right) \neq 0$ . Hence, we have

$$(9.11) \quad \left( \frac{\beta r - 2\lambda}{2\alpha} \right) (\xi f) = 0.$$

As  $(\xi f) \neq 0$ , the soliton function reduce to

$$(9.12) \quad \lambda = \frac{\beta r}{2}.$$

**Theorem 9.1.** *In a  $(\alpha, \beta)$ -(GRYS) $_n$  admitting (LPS) $_n$  manifold the soliton function is given by.*

$$\lambda = \frac{\beta r}{2}.$$

*provided  $\left( \frac{n\beta}{2\alpha} - 1 \right) \neq 0$ .*

The metric inner product on (9.8) with  $\xi$  results

$$(9.13) \quad \eta(\zeta_2)(\zeta_1 f) - \eta(\zeta_1)(\zeta_2 f) = \frac{\beta}{2}[(\zeta_2 r)\eta(\zeta_1) - (\zeta_1 r)\eta(\zeta_2)].$$

Put  $\zeta_1 = \xi$  in (9.13) and using lemma (3.1), we have

$$(9.14) \quad \eta(\zeta_2)(\xi f) - \eta(\xi)(\zeta_2 f) = \frac{\beta}{2}[(\zeta_2 r)\eta(\xi) - (\xi r)\eta(\zeta_2)].$$

Simplifying, above equation reduces to

$$(9.15) \quad (\zeta_2 r) = -\frac{2}{\beta}(\zeta_2 f).$$

Using (9.15) in (9.10) and then applying theorem (3.1), we receive

$$(9.16) \quad \lambda = \frac{\beta(r + 2n^2)}{2}.$$

Thus, we can state

**Theorem 9.2.** *The soliton function of  $(\alpha, \beta)$ -(GRYS) $_n$  admitting (LPS) $_n$  Manifold, having non-constant scalar curvature, is given by*

$$\lambda = \frac{\beta(r + 2n^2)}{2}.$$

## 10 Example

**Example 10.1.** Let's have a look at a differentiable manifold in four dimensions,  $M^4 = (\zeta_1, \zeta_2, \zeta_3, \zeta_4) \in \mathfrak{R} : (\zeta_1, \zeta_2, \zeta_3, \zeta_4) \neq 0$ , where  $(\zeta_1, \zeta_2, \zeta_3, \zeta_4)$  is the standard coordinate in four-dimensional real space. Considering that each point on M has a collection of linearly independent vector fields, let's call them  $(f_1, f_2, f_3, f_4)$ , and is defined by

$$(10.1) \quad f_1 = f^{\zeta_1 - a\zeta_4} \frac{\partial}{\partial \zeta_1}, \quad f_2 = f^{\zeta_2 - a\zeta_4} \frac{\partial}{\partial \zeta_2}, \quad f_3 = f^{\zeta_3 - a\zeta_4} \frac{\partial}{\partial \zeta_3}, \quad f_4 = \frac{\partial}{\partial \zeta_4}$$

where  $a$  is non-zero constant and  $\{\frac{\partial}{\partial \zeta_1}, \frac{\partial}{\partial \zeta_2}, \frac{\partial}{\partial \zeta_3}, \frac{\partial}{\partial \zeta_4}\}$  denotes the standard basis of  $M^4$ . Let the metric (Lorentzian) and the 1-form  $\eta$  components on  $M^4$  are concreted by

$$(10.2) \quad g_{cd} = g(f_c, f_d) = \begin{cases} -1, & \text{if } c = d = 4 \\ 0, & \text{if } c \neq d \\ 1, & \text{if } c = d \neq 4 \end{cases}$$

and

$$(10.3) \quad \eta(\zeta_1) = g(\zeta_1, f_4).$$

for any  $\zeta_1 \in \Gamma(TM)$  on  $M^4$ .

If  $\phi(f_1) = f_1, \phi(f_2) = f_2, \phi(f_3) = f_3, \phi(f_4) = 0$ , are the tensor field then the following relationships may be easily verified by  $g$ 's and  $\phi$ 's linearity properties:

$$(10.4) \quad \eta(f_4) = -1, \quad \phi^2\zeta_1 = \zeta_1 + \eta(X)f_4, \quad g(\phi\zeta_1, \phi\zeta_2) = g(\zeta_1, \zeta_2) - \eta(\zeta_1)\eta(\zeta_2).$$

The Lie bracket's (non-vanishing) components are determined as follows:

$$(10.5) \quad [f_1, f_4] = af_1, \quad [f_2, f_4] = af_2, \quad [f_3, f_4] = af_3.$$

The Koszul's formula yields for  $f_4 = \xi$

$$(10.6) \quad \begin{aligned} \nabla_{f_1}f_1 &= af_4, & \nabla_{f_1}f_2 &= 0, & \nabla_{f_1}f_3 &= 0, & \nabla_{f_1}f_4 &= af_1, \\ \nabla_{f_2}f_1 &= 0, & \nabla_{f_2}f_2 &= af_4, & \nabla_{f_2}f_3 &= 0, & \nabla_{f_2}f_4 &= af_2, \\ \nabla_{f_3}f_1 &= 0, & \nabla_{f_3}f_2 &= 0, & \nabla_{f_3}f_3 &= af_4, & \nabla_{f_3}f_4 &= af_3, \\ \nabla_{f_4}f_1 &= 0, & \nabla_{f_4}f_2 &= 0, & \nabla_{f_4}f_3 &= 0, & \nabla_{f_4}f_4 &= 0. \end{aligned}$$

The curvature components  $R$ , Ricci components  $S$  and the scalar tensor  $r$  are obtained respectively

$$(10.7) \quad \begin{aligned} R(f_1, f_2)f_1 &= -a^2f_2, & R(f_1, f_3)f_1 &= -a^2f_3, & R(f_1, f_4)f_1 &= -a^2f_4, \\ R(f_1, f_2)f_2 &= a^2f_1, & R(f_2, f_3)f_2 &= -a^2f_3, & R(f_2, f_4)f_2 &= -a^2f_4, \\ R(f_1, f_3)f_3 &= a^2f_1, & R(f_2, f_3)f_3 &= a^2f_2, & R(f_3, f_4)f_3 &= -a^2f_4, \\ R(f_1, f_4)f_4 &= -a^2f_1, & R(f_2, f_4)f_4 &= -a^2f_2, & R(f_3, f_4)f_4 &= -a^2f_3. \end{aligned}$$

$$(10.8) \quad S(f_1, f_1) = S(f_2, f_2) = S(f_3, f_3) = 3a^2, \quad S(f_4, f_4) = -3a^2,$$

$$(10.9) \quad r = 6a^2.$$

If we consider  $\lambda = 3a^2(\beta - \alpha)$  and the values from the equation (10.8) and (10.9), equation (3.1) satisfies.

Hence, it is an example of four-dimensional  $(\alpha, \beta) - (RYS)_4$  admitting  $(LPS)_4$ .

Using theorem (3.1) in  $\lambda = 3a^2(\beta - \alpha)$ , we get

$$(10.10) \quad \lambda = -3\beta a^2.$$

Equating (10.10) and (3.14), yield

$$(10.11) \quad a = \pm 1.$$

Hence, for  $a = \pm 1$ , the defined metric in (10.1) and the values obtained in (10.8) and (10.9) satisfy the theorem (3.1) and theorem (3.2).

## Conclusion

The Lorentzian metric and its generalizations play very important role in the study of cosmology. The perfect fluid, that depends on the lorentzian metric, is an example

to represents the universe. Here, we study the soliton metric on Lorentzian manifold and discuss some fruitful investigation. Specially, the cosmological models have been discussed and the length of Ricci operator has been obtained and proved that spacetime is vaccum. The Poisson structure is an interesting topic to investigate in cosmology and mathematical physics. The soliton function has been discussed under the two different conditions, whether the scalar curvature is constant or not.

## Acknowledgement

There does not any financial support provide to conduct this research. All the authors made equal contribution to complete the manuscript.

## References

- [1] M. ALI and Z. AHSAN: *Ricci solitons and symmetries of spacetime manifold of general relativity*, Journal of Advanced Research on Classical and Modern Geometries, **1(2)**, (2014), 75-84.
- [2] A.M. BLAGA: *Some geometrical aspects of Einstein, Ricci and Yamabe solitons*, J. Geom. Symmetry Phys., (**52**), (2019), 17–26.
- [3] A.M. BLAGA: *Solitons and geometrical structures in a perfect fluid spacetime*, Rocky Mountain J. Math., **50(1)**, (2020), 41-53. DOI: doi.org/10.1216/rmj.2020.50.41
- [4] R. BURING, D. LIPPER and A.V. KISELEV: *The hidden symmetry of Kontsevich's graph flows on the spaces of Nambu-determinant Poisson brackets*, Open Communications in Nonlinear Mathematical Physics (OCNMP), **2**, (2022), 186–215. DOI: doi.org/10.46298/ocnmp.8844
- [5] B.B. CHATURVEDI, P. BHAGAT, M.N.I. KHAN, T. RANA and R.K. MISHRA: *Proposed theorems on a Lorentzian Kähler space-time manifold admitting Bochner curvature tensor*, Results in Nonlinear Analysis, **6(4)**, (2023), 140-148.
- [6] B.B. CHATURVEDI, K.B. KAUSHIK, P. BHAGAT and M.N.I. KHAN: *Characterization of solitons in a pseudo-quasi-conformally flat and pseudo  $W_8$  flat Lorentzian Kahler space-time manifolds*, AIMS Mathematics, **9(7)**, (2024), 19515–19528. DOI: doi.org/10.3934/math.2024951
- [7] B.B. CHATURVEDI and P. PANDEY: *Some Examples on Weak Symmetries*, General Mathematics Notes, **29(1)**, (2015), 61-66.
- [8] R.S. CHAUDHARY and B. PAL: *On gradient Ricci soliton space-time warped product*, Afrika Matematika, **34(3)**, (2023), 34-37.
- [9] S.K. CHAUBEY and U.C. DE: *Three-dimensional Riemannian manifolds and Ricci solitons*, Quaestiones Mathematicae, **45**, (2022), 667-825.
- [10] S. CHIDANANDA and V. VENKATESHA: *Yamabe soliton and Riemann soliton on Lorentzian para-Sasakian manifold*, Commun. Korean Math. Soc., **37**, (2022), 213-228.

- [11] D. DE: *Sasakian 3-manifolds admitting a gradient Ricci-Yamabe soliton*, Korean J. Math., **29(3)**, (2021), 547-554. DOI: doi.org/10.11568/kjm.2021.29.3.547
- [12] U.C. DE, S.K. CHAUBEY and S. SHENAWY: *Perfect fluid spacetimes and Yamabe solitons*, J. Math. Phys., **62**, (2021). DOI: doi.org/10.1063/5.0033967.
- [13] K. DE, and U.C. DE: *A note on almost Ricci soliton and gradient almost Ricci soliton on para-Sasakian manifolds*, Korean J. Math., **28(4)**, (2020) 739-751. DOI: doi.org/10.11568/kjm.2020.28.4.739
- [14] K. DE, U.C. DE, A.A. SYIED, N.B. TURKI and S. ALSAEED: *Perfect fluid spacetimes and gradient solitons*, Journal of Nonlinear Mathematical Physics, **29**, (2022), 843-848.
- [15] U.C. DE, C.A. MANTICA and Y. SUH: *Perfect fluid spacetimes and gradient solitons*, Filomat, **36(3)**, (2022), 829-842.
- [16] U.C. DE, A. SARDAR and K. DE: *Ricci-Yamabe solitons and 3-dimensional Riemannian manifolds*, Turkish Journal of Mathematics, 46(3), (2022), 1078-1088.
- [17] S. DESHMUKH and B.Y. CHEN: *A note on Yamabe solitons*, Balk. J. Geom. Its Appl., **23**, (2018), 37-43.
- [18] D. GANGULY, S. DEY, A. ALI and A. BHATTACHARYYA: *Conformal Ricci soliton and quasi-Yamabe soliton on generalized Sasakian space form*, J. Geom. Phys., **169**, (2021), (104339).
- [19] S. GULER and M. CRASMAREANU: *Ricci-Yamabe maps for Riemannian flows and their volume variation and volume entropy*, Turkish Journal of Mathematics, **46(5)**, (2019), 2631-2641.
- [20] R.S. HAMILTON: *Three-manifolds with positive Ricci curvature*, Journal of Differential Geometry, **17**, (1982), 255-306.
- [21] A. HASEEB and S.K. CHAUBEY: *Lorentzian para Sasakian manifolds and \*Ricci solitons*, Kragujevac Journal of Mathematics, **48 (2)**, (2024), 167-179.
- [22] A. HASEEB, S.K. CHAUBEY and M.A. KHAN: *Riemannian 3-manifolds and Ricci-Yamabe solitons* Int. J. Geom. Methods Mod. Phys., **20**, (2023), 2350015. DOI: doi.org/10.1142/S0219887823500159.
- [23] A. HASEEB and R. PRASAD: *Certain results on Lorentzian para-Kenmotsu manifolds*, Bol. Soc. Parana. Mat., **39**, (2021), 201-220.
- [24] A. HASEEB, R. PRASAD, S.K. CHAUBEY and A.T. VANLI: *A note on \*-conformal and gradient \*-conformal  $\eta$ -Ricci solitons in  $\alpha$ -cosymplectic manifolds*, Honam Math. J. **44**, (2022), 231-243.
- [25] A. HASEEB, R. PRASAD and F. MOFARREH: *Sasakian manifolds admitting  $*\eta$ -Ricci-Yamabe solitons*, Adv. Math. Phys., **2022**, (2022), 5718736. DOI: doi.org/10.1155/2022/5718736

- [26] A. HASEEB, M. BILAL, S.K. CHAUBEY and A.A.H. AHMADINI: *zeta-Conformally flat LP-Kenmotsu manifolds and Ricci–Yamabe solitons*, Mathematics, **11(1)**, 212, (2023). DOI: doi.org/10.3390/math11010212
- [27] D. KAR, and P. MAJHI: *Beta-almost Ricci solitons on almost CoKahler manifolds*, Korean J. Math., **27(3)**, (2019), 691-705. DOI: doi.org/10.11568/kjm.2019.27.3.691
- [28] S. KUNDU: *Para-Kenmotsu metric as a  $\eta$ -Ricci soliton*, Korean J. Math., **29(2)**, (2021), 445-453. DOI: doi.org/10.11568/kjm.2021.29.2.445
- [29] S.D. LEE, B.H. KIM, and J.H. CHOI; *On a classification of warped product spaces with gradient Ricci solitons*, Korean J. Math. **24(4)**, (2016), 627-636. DOI: doi.org/10.11568/kjm.2016.24.4.627
- [30] Y. LI, A. HASEEB and M. ALI: *LP-Kenmotsu manifolds admitting  $\eta$ -Ricci solitons and spacetime*, J. Math., **2022**, (2022), 6605127. DOI: doi.org/10.1155/2022/6605127
- [31] B. PAL and R.S. CHAUDHARY: *An introduction to gradient solitons on statistical Poisson manifold and perfect fluid Poisson manifold*, Journal of Geometry and Physics, **195**, (2024), 105039.
- [32] P. PANDEY: *On weakly cyclic generalized Z-symmetric manifolds*, National Academy Science Letters, **43(04)**, (2020), 347-350.
- [33] P. PANDEY: *Study of Kahler Structure on 4-dimensional Space-time*, Differential Geometry-Dynamical System, **24**, (2022), 157-163.
- [34] P. PANDEY and B.B. CHATURVEDI: *On a Lorentzian complex space form*, National Academy Science Letters, **43(04)**, (2020), 351-353.
- [35] P. PANDEY and K. SHARMA: *Ricci Soliton Admitting Generalized Z-tensor on Sasakian 3-Manifold*, AIP: Conference Proceedings, **3087-060001**, (2024). DOI: doi.org/10.1063/5.0199390.
- [36] P. PANDEY and K. SHARMA: *Some Results on Ricci Soliton on Contact Metric Manifolds*, Facta Universitatis: Mathematics and Informatics, **38(3)**, (2023), 473-486.
- [37] G.P. POKHARIYAL and S.K. CHAUBEY: *Study of P-curvature tensor in the space-time of general relativity*, Honam Mathematical J., **45(2)**, (2023), 316–324. DOI: doi.org/10.5831/HMJ.2023.45.2.316
- [38] A. SARDAR and U.C. DE: *Characterizations of GRW spacetimes admitting Ricci-Yamabe solitons*, International Journal of Geometric Methods in Modern Physics, **2440002**, (2023). DOI: doi.org/10.1142/S0219887824400024
- [39] A. SARDAR and A. SARKAR: *Ricci-Yamabe solitons and gradient Ricci-Yamabe solitons on Kenmotsu 3-manifolds*, Kyungpook Math. J., **61**, (2021), 813-822.
- [40] A. SARKAR, and P. BHAKTA: *Certain solitons on generalized  $(\kappa, \mu)$  contact metric manifolds*, Korean J. Math., **28(4)**, (2020) 847-863. DOI: doi.org/10.11568/kjm.2020.28.4.847

- [41] M. D. SIDDIQI: *Ricci  $\rho$ -soliton and geometrical structure in a dust fluid and viscous fluid spacetime*, Bulg. J. Phys. **46**, (2019), 163-173.
- [42] M.D. SIDDIQI and U.C. DE: *Relativistic perfect fluid spacetimes and Ricci-Yamabe solitons*, Letters in Mathematical Physics, **112(1)**, (2022). DOI: 10.1007/s110 05-021-01493-z
- [43] A. VISHWAS, A. DAS, K.K. BAISHYA and M.R. BAKSHI:  *$\eta$ -Ricci solitons on Kenmotsu manifolds admitting general connection*, Korean J. Math., **28(4)**, (2020), 803-817. DOI: <https://doi.org/10.11568/kjm.2020.28.4.803>
- [44] S.K. YADAV, S.K. CHAUBEY and S.K HUI: *On the perfect fluid Lorentzian para-Sasakian spacetimes*, Bulg. J. Phys., **46**, (2019), 1-15.
- [45] Y.J. SUH and S.K. CHAUBEY: *Ricci solitons on general relativistic spacetimes*, Physica Scripta, **98(6)**, (2023), 065207. DOI: 10.1088/1402-4896/accf41
- [46] P. ZHANG, Y. LI, S. ROY, S. DEY and A. BHATTACHARYA: *Geometrical structure in a perfect fluid spacetime with conformal Ricci-Yamabe soliton*, 14(3), 594, (2022). DOI: [doi.org/10.3390/sym14030594](https://doi.org/10.3390/sym14030594)