

# All You Need is a Paraboloid: Quadratic Cuts for Non-Convex MINLP

ADRIAN GÖß\*, ROBERT BURLACU, ALEXANDER MARTIN

JULY 9, 2024

**ABSTRACT.** It is only half the job to find a good solution for a mathematical optimization problem, as one needs to verify its quality by specifying a dual bound. When it comes to mixed-integer nonlinear programming (MINLP), strong prerequisites such as constraint qualifications appear suitable, but may be difficult to verify computationally. In practice, solvers apply local refinement or convexification strategies to retrieve tight dual bounds. However, these concepts require appropriate big-M formulations, generate new sub-problems, or struggle to represent non-convex characteristics in terms of high accuracy, all of which can lead to long running times.

As an alternative, we aim to leverage recent advances in mixed-integer quadratically-constrained programming (MIQCP) and propose a global approximation of constraint functions by paraboloids, i.e., univariate quadratic terms. The approximation is retrieved as a solution to a mixed-integer linear programming (MIP) problem. Further, for each nonlinear constraint function, we solve such MIPs and determine small numbers of paraboloids approximating it from either side. A replacement of the nonlinearities with the corresponding quadratic functions leads to a quadratically-constrained relaxation of the original problem. Solving the MIQCP relaxation then leads to a dual bound whose tightness depends on the approximation guarantee of the paraboloids. In summary, this approach enables solvers that are explicitly tailored for quadratic constraints to solve MINLPs to global optimality.

## ACKNOWLEDGMENTS

We thank the Deutsche Forschungsgemeinschaft for their support within project A05 in the “Sonderforschungsbereich/Transregio 154 Mathematical Modelling, Simulation and Optimization using the Example of Gas Networks”.

## 1. INTRODUCTION

We consider the problem

$$\min c(\mathbf{x}) \tag{1a}$$

$$f_j(\mathbf{x}) \leq 0, \quad j \in J, \tag{1b}$$

$$\mathbf{x} \in \Omega, \tag{1c}$$

where  $\Omega = [\underline{\mathbf{x}}, \bar{\mathbf{x}}] \cap (\mathbb{Z}^p \times \mathbb{R}^{n-p}) \neq \emptyset$  for  $p, n \in \mathbb{N}$  with  $0 \leq p \leq n$  and  $|J| < \infty$ . We assume the involved functions  $c, f_j : \mathbb{R}^n \mapsto \mathbb{R}$  to be continuous on the domain  $[\underline{\mathbf{x}}, \bar{\mathbf{x}}]$ ,

---

\*CORRESPONDING AUTHOR

(A. Göß, R. Burlacu, A. Martin) UNIVERSITY OF TECHNOLOGY NUREMBERG (UTN),  
ANALYTICS & OPTIMIZATION LAB, ULMENSTR. 52I, 90443 NUREMBERG, GERMANY

*E-mail address:* {adrian.goess, robert.burlacu, alexander.martin}@utn.de.

*2010 Mathematics Subject Classification.* 90C11, 90C20, 90C26, 90C30 .

*Key words and phrases.* mixed-integer nonlinear programming, quadratic programming, non-convex programming, cutting scheme .

$j \in J$ . In general, we consider at least one involved function to be nonlinear and thus call (1) a mixed-integer nonlinear program (MINLP). Please note that we can assume linearity of the objective by introduction of a separate variable  $z$ , which is then minimized subject to  $c(\mathbf{x}) \leq z$  and the remaining constraints. This does not increase the theoretical character of the resulting program. For clear notation, we assume the existence of integer variables throughout the paper, i.e.,  $p > 0$ , if not stated otherwise and refer to (1) when writing about MINLPs.

The solution methods for a MINLP can differ greatly depending on the properties of the involved functions. See [17] for an overview over global solution strategies (mainly on continuous variables) and [5] for a survey on the most common components of practical approaches for MINLPs.

If all involved functions are convex, the feasible set of the continuous relaxation of the MINLP is convex and therefore (1) is typically called a *convex* MINLP. Such problems are usually tackled with cutting methods, e.g., outer approximation [12] or extended cutting-plane methods [28], and decomposition schemes, e.g., generalized Benders decomposition [15].

However, the situation changes drastically if at least one of the functions is not convex, which coins the term *non-convex* MINLP. Note here that equality constraints  $f_j(\mathbf{x}) = 0$  can be modeled via  $f_j(\mathbf{x}) \leq 0$  and  $-f_j(\mathbf{x}) \leq 0$ , but directly enforce non-convexity for  $f_j$  nonlinear (even if convex). Non-convex MINLPs are typically solved by some form of convex relaxation of the (non-convex) nonlinearities in combination with a refinement mechanism. Useful convexification techniques are for instance proposed by [26]. As such may lack a sufficiently tight representation of the original feasible region, the latter can be split to create new subproblems with tighter convex relaxations. This divide and conquer approach is performed in spatial-branch-and-bound (compare, for instance, [23]) by reformulation or in  $\alpha$ -branch-and-bound by  $\alpha$ -convexification [2]. In the last decade, the approximation of the non-convex functions by piecewise linear functions and re-modeling as a mixed-integer linear program (MIP) has established itself as a viable alternative, especially for real-world applications with considerable combinatorics [9, 14]. See [27] for a comprehensive overview on theoretical properties of the various MIP-based models for piecewise linear functions.

A natural generalization of the piecewise linear approach is the piecewise approximation by polynomials of a certain degree. Such an approximation is performed in an optimization setting using quadratic functions in [8]. In certain algebraic domains, collections of more general such polynomials are typically referred to as *splines*, and a distinction is made between approximations with fixed and variable knots, i.e., discretization points of an interval. There is a vast amount of literature starting from the 1960s on splines. We refer the interested reader to [25] for an introduction to the topic and to [22] for a recap of the first approximately 30 years.

From an optimization point of view, there is no clear benefit in the piecewise approximation by higher-order polynomials instead of linear functions, as this involves the introduction of both non-convex constraints and integer variables. However, if we can perform a one-sided spline approximation, i.e., an underestimation on the entire domain of the function, this can serve as a globally valid relaxation, which does not require piecewise modeling. In [7], the authors prove existence and uniqueness for a best one-sided approximation with maximal degree. A computational procedure to compute such an approximation with one polynomial is presented in [18].

Surprisingly enough, we did not find any (recent) literature that treats the one-sided approximation by a spline with variable knots. That is, the required approximations are either not one-sided (see again [25] or [1]) and thus do not yield a globally valid relaxation of the MINLP or deal with the best approximation by a

single polynomial (see [11]). Furthermore, to the best of our knowledge, there is no approach for solving MINLP problems that globally approximates non-convex functions on their respective domains to an arbitrary accuracy without introducing additional variables, i.e. increasing the dimension of the problem, or branching on continuous variables.

We start by presenting a MIP model in Section 2 for the one-sided approximation by univariate quadratic functions and prove its correctness for this purpose. In Section 3 we formally introduce the resulting method to approximate non-convex constraint functions by a small number of such quadratic functions. Both sections are complemented by computational showcases in Section 4 before we close the article with concluding remarks and further research directions in Section 5.

**1.1. Contribution.** Motivated by recent developments in non-convex mixed-integer quadratically constrained programming (MIQCP) (see, for instance, [3, 4, 29]), we propose a novel approach that solves non-convex MINLPs by reducing them to MIQCPs. The main part is based on the construction of non-convex parabolic relaxations with arbitrary tightness of rather general non-linear Lipschitz continuous functions. In a sense, these parabolic relaxations can be considered as non-convex cuts for MINLPs, since we construct them to be globally valid on the domain of the function.

More particularly, our contribution is twofold: On the one hand, we present a MIP approach to globally approximate multi-dimensional Lipschitz functions from one side, where the anti-derivatives to these may or may not be known. That is, the latter case even includes black-box functions. The correctness of this approach is shown by theoretical means and a practical version is showcased on one-dimensional constraint functions. We conduct these computations for frequently occurring functions in the MINLPLib [10] on certain domains to provide a lookup table that can be used a priori for the following part. Due to its character each approximation gives a global under-/overestimator when taking the pointwise maximum/minimum, respectively.

On the other hand, we introduce a novel framework to solve non-convex MINLPs on bounded domains. Our approach computes a (in some sense) minimal number of parabolic functions for each non-convex constraint and then replaces the original constraints with their global one-sided parabolic approximations. This can be interpreted as a parabolic relaxation and reduces the MINLP to a MIQCP. We stress the integration of the lookup table mentioned above and demonstrate the potential of our method with preliminary numerical results for some instances from the MINLPLib.

**1.2. Notation.** A vector  $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$  is denoted with bold and upright letters, whereas components  $x_i$  are not bold and italic. A comparison between two vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , e.g.,  $\mathbf{x} \leq \mathbf{y}$ , is always interpreted component-wise. During the theoretical section, we make use of the 1-norm of a vector  $\mathbf{x}$  which is defined as  $\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i|$ . For a domain  $\mathcal{D} \subseteq \mathbb{R}^n$ , we define the set of continuous functions on  $\mathcal{D}$  as  $\mathcal{C}^0(\mathcal{D})$ . Further,  $\mathcal{P}(\mathcal{D})$  denotes the power set of  $\mathcal{D}$ . Lastly, for a positive integer  $m \in \mathbb{N}$ , we state the set of indexes up to  $m$  as  $[m] := \{1, \dots, m\}$ .

Note that we present our theoretical parts for constraint functions on a multi-dimensional domain. We aim for their approximation by quadratic functions without bivariate terms and refer to such as *paraboloids*. Hence, the global one-sided approximation by paraboloids is called a *parabolic approximation*. Later on, when we substitute constraint functions with mentioned approximations, we name such a *parabolic relaxation*, stressing the character of this procedure.

## 2. PARABOLIC APPROXIMATION – MIP APPROACH

In order to construct a relaxation to problem (1), we aim to approximate its constraint functions from one or either side – if necessary – and replace it by the approximation. Hereby, “from one side” means that the approximation serves as a global underestimator (or overestimator) on the entire domain. In this work, this is supposed to be realized as a maximum (or minimum) of a set of paraboloids, offering the advantage to be incorporated into the optimization problems by simple inequalities with respect to all such paraboloids. Given an arbitrary but fixed number of paraboloids  $K \in \mathbb{N}$ , we introduce a mixed-integer programming (MIP) formulation whose optimal solution is guaranteed to return the desired approximation if its objective is zero.

In mathematical terms, let  $f : \mathcal{D} \mapsto \mathbb{R}$  be a Lipschitz continuous function with respect to  $\|\cdot\|_1$  and some  $L > 0$ , where  $\mathcal{D} = [\mathbf{a}, \mathbf{b}]$  a non-empty, fulldimensional box defined by the vectors  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ . That is, for all  $\mathbf{x}, \mathbf{y} \in \mathcal{D}$ , it holds true that

$$|f(\mathbf{x}) - f(\mathbf{y})| \leq L\|\mathbf{x} - \mathbf{y}\|_1.$$

Here,  $f$  represents the constraint function, for which we want to find a global underestimator by paraboloids. The case for a global overestimator can be treated analogously.

Now, given an approximation guarantee  $\varepsilon > 0$  and respective  $K \in \mathbb{N}$ , we aim to determine  $K$  paraboloids of degree 2, i.e.,  $p^l(\mathbf{x}) = \sum_{i=1}^n \alpha_i^l x_i^2 + \sum_{i=1}^n \beta_i^l x_i + \gamma^l$  for  $l \in [K]$ , such that

$$\max_{l \in [K]} p^l(\mathbf{x}) \geq f(\mathbf{x}) - \varepsilon, \quad (2)$$

and

$$\max_{l \in [K]} p^l(\mathbf{x}) \leq f(\mathbf{x}), \quad (3)$$

for all  $\mathbf{x} \in \mathcal{D}$ . We note that the authors of [19] use the minimum of such paraboloids to compute a non-convex underestimator of a non-convex objective function.

In the following, we state a MIP formulation in order to determine the coefficients for these paraboloids. Afterwards, we show, on the one hand, that a solution to the MIP formulation with objective zero exists for sufficiently high number of paraboloids and, on the other hand that a solution with objective zero inherits the desired approximation conditions (2) and (3).

**2.1. MIP formulation.** In addition to the statements from the introduction of this section, we assume to know a function  $\mu_f : \mathcal{P}(\mathcal{D}) \mapsto \mathbb{R}_+$  which measures the Lebesgue integral of  $f$  on any connected sub-domain of  $\mathcal{D}$ . In the one-dimensional case, such a function is typically the anti-derivative of  $f$ .

Besides an approximation guarantee  $\varepsilon > 0$  and a number of paraboloids  $K \in \mathbb{N}$ , we need to define parameters  $\delta \in (0, \varepsilon)$  and  $\nu \in (0, \delta/\varepsilon)$ . Informally their choice defines how the parabolic approximation is centered inside the  $\varepsilon$ -tube between  $f$  and  $f - \varepsilon$ . This influence becomes clearer throughout this section. Lastly, in order to avoid “spiking” paraboloids, we choose a constant  $C \geq L$  which bounds the maximal absolute slope of each paraboloid as shown below.

With these parameters at hand, we define the grids of  $\mathcal{D}$

$$\mathcal{G}_\varepsilon := \times_{i=1}^n \left\{ a_i + k\Delta t_i \mid k = 0, 1, \dots, \frac{b_i - a_i}{\Delta t_i} \right\},$$

and

$$\mathcal{G} := \times_{i=1}^n \left\{ a_i + k\Delta d_i \mid k = 0, 1, \dots, \frac{b_i - a_i}{\Delta d_i} \right\},$$

for achieving the conditions (2) and (3), respectively. Without loss of generality, we assume the discretization widths  $\Delta t_i$  and  $\Delta d_i$  to be chosen such that  $(b_i - a_i)/\Delta t_i \in \mathbb{N}$  for all  $i \in [n]$ . In addition, they need to fulfill  $\Delta t_i, \Delta d_i \geq 0$  for all  $i \in [n]$  as well as

$$\sum_{i=1}^n \Delta t_i \leq \frac{n+1}{n} \frac{\varepsilon - \delta}{3L}, \quad (4)$$

and

$$\Delta d_i \leq \frac{2\nu\varepsilon}{(\sqrt{3}-1)n(C+L)}, \quad \text{for all } i \in [n], \quad (5)$$

respectively. Because the right-hand side of above inequalities is strictly positive, such a choice is always possible.

For clear notation, we summarize the widths as  $\Delta \mathbf{t} = (\Delta t_1, \dots, \Delta t_n)^\top$  and  $\Delta \mathbf{d} = (\Delta d_1, \dots, \Delta d_n)^\top$ . Note that this allows to rewrite  $\mathcal{D}$  as

$$\mathcal{D} = \bigcup_{\mathbf{t} \in \mathcal{G}_\varepsilon \cap [\mathbf{a}, \mathbf{b}]} [\mathbf{t}, \mathbf{t} + \Delta \mathbf{t}] = \bigcup_{\mathbf{d} \in \mathcal{G} \cap [\mathbf{a}, \mathbf{b}]} [\mathbf{d}, \mathbf{d} + \Delta \mathbf{d}], \quad (6)$$

where in each union the arguments at most intersect on their boundaries. We abbreviate  $\mathcal{B}(\mathbf{d}) = [\mathbf{d}, \mathbf{d} + \Delta \mathbf{d}]$ .

In order to make the following model statement even more readable, we denote for a  $\mathbf{t}$  all neighboring points in  $\mathcal{G}_\varepsilon$  which differ in each coordinate by exactly  $\Delta t_i$  in absolute terms as

$$\mathcal{N}(\mathbf{t}) = \left\{ \mathbf{t} + \sum_{i=1}^n u_i \Delta t_i \mathbf{e}_i \in \mathcal{G}_\varepsilon \mid \mathbf{u} \in \{-1, 1\}^n \right\},$$

where  $\mathbf{e}_i$  denotes the  $i$ th unit vector.

Lastly, we apply the so-called *big-M method* in our model and, thus, determine  $M_1, M_2 > 0$  sufficiently large.

In terms of variables in the model, we define  $\alpha_i^l, \beta_i^l, \gamma^l$  for  $i \in [n]$  and  $l \in [K]$ , specifying the  $l$ th paraboloid as  $p^l(\mathbf{x}) = \sum_{i=1}^n \alpha_i^l x_i^2 + \sum_{i=1}^n \beta_i^l x_i + \gamma^l$ . Further, we introduce binary variables  $s_{\mathbf{t}}^l \in \{0, 1\}$  for all  $l \in [K]$  and all grid points  $\mathbf{t} \in \mathcal{G}$ . These indicate whether paraboloid  $l$  “contains” the approximation point  $(\mathbf{t}, f(\mathbf{t}) - \delta)$  in the sense that  $p^l(\mathbf{t}) \geq f(\mathbf{t}) - \delta$ . As formally shown below, this ensures property (2). Condition (3), however, is controlled by continuous variables  $v_{\mathbf{d}}^l \geq 0$  for  $l \in [K]$  and  $\mathbf{d} \in \mathcal{G}$  which track violations of the integral between  $f$  and a paraboloid  $p^l$ .

After the introduction of all necessary parameters and variables, we can finally formulate the MIP. We conduct this implicitly and comment on it afterwards:

$$\min \sum_{l \in [K]} \sum_{\mathbf{d} \in \mathcal{G} \cap [\mathbf{a}, \mathbf{b}]} v_{\mathbf{d}}^l \quad (7a)$$

$$\text{s.t.} \quad p^l(\mathbf{t}) \geq f(\mathbf{t}) - \delta - M_1(1 - s_{\mathbf{t}}^l), \quad l \in [K], \mathbf{t} \in \mathcal{G}_\varepsilon, \quad (7b)$$

$$\sum_{l \in [K]} s_{\mathbf{t}}^l \geq 1, \quad l \in [K], \mathbf{t} \in \mathcal{G}_\varepsilon, \quad (7c)$$

$$\left| \frac{d}{dx_i} p^l(\mathbf{t}') \right| \leq 2L + M_2(1 - s_{\mathbf{t}}^l), \quad l \in [K], i \in [n], \mathbf{t} \in \mathcal{G}_\varepsilon, \mathbf{t}' \in \mathcal{N}(\mathbf{t}), \quad (7d)$$

$$p^l(\mathbf{d}) \leq f(\mathbf{d}) - \nu\varepsilon, \quad l \in [K], \mathbf{d} \in \mathcal{G}, \quad (7e)$$

$$v_{\mathbf{d}}^l \geq \int_{\mathcal{B}(\mathbf{d})} p^l(\mathbf{x}) - (f(\mathbf{x}) - \nu\varepsilon) d\mathbf{x}, \quad l \in [K], \mathbf{d} \in \mathcal{G}, \quad (7f)$$

$$\left| \frac{d}{dx_i} p^l(\mathbf{a}) \right| \leq C, \quad l \in [K], i \in [n], \quad (7g)$$

$$\left| \frac{d}{dx_i} p^l(\mathbf{b}) \right| \leq C, \quad l \in [K], i \in [n], \quad (7h)$$

$$\alpha_i^l, \beta_i^l, \gamma_i^l \in \mathbb{R}, \quad l \in [K], i \in [n], \quad (7i)$$

$$s_{\mathbf{t}}^l \in \{0, 1\}, \quad l \in [K], \mathbf{t} \in \mathcal{G}_\varepsilon, \quad (7j)$$

$$v_{\mathbf{d}}^l \geq 0, \quad l \in [K], \mathbf{d} \in \mathcal{G}. \quad (7k)$$

Note that  $\frac{d}{dx_i} p^l(\mathbf{x}) = 2\alpha_i^l x_i + \beta_i^l$  for  $l \in [K]$  which is a linear constraint in the variables  $\alpha_i^l$  and  $\beta_i^l$ . Further, for a function  $h(\mathbf{x}) : \mathcal{D} \mapsto \mathbb{R}$ , we can re-write  $|h(\mathbf{x})| \leq C$  as  $h(\mathbf{x}) \leq C$  and  $h(\mathbf{x}) \geq -C$ . This allows to rewrite constraints (7d), (7g), and (7h) as linear inequalities. In addition, the argument of the integral in (7f) can also be evaluated by the help of  $\mu_f$  such that it reduces to another set of linear inequalities. This concludes the formulation of the MIP to find a parabolic approximation for given parameters.

**2.2. Proof of correctness.** We have to show two aspects regarding model (7): the existence of a solution for appropriate choice of parameters and the validity of conditions (2) and (3) for such a solution. The following theorem provides the first one.

**Theorem 2.1** (Existence of solution). *Let  $0 < \Delta t \leq \min\{\frac{n+1}{n^2} \frac{\varepsilon - \delta}{3L}, \frac{2\delta}{nL}\}$ . If we choose  $\Delta t_i = \Delta t$  for all  $i \in [n]$  and set  $C = 2L\|\mathbf{b} - \mathbf{a}\|_\infty / \Delta t$  as well as  $\nu = \delta / (2\varepsilon)$ , problem (7) has an optimal solution with objective value zero and*

$$K = |\mathcal{G}_\varepsilon| = \prod_{i=1}^n \left\lceil \frac{b_i - a_i}{\Delta t} \right\rceil.$$

*Proof.* First, see that

$$\sum_{i=1}^n \Delta t_i = n\Delta t \leq \frac{n+1}{n} \frac{\varepsilon - \delta}{3L},$$

thus, the choice of  $\Delta t_i$  is in accordance with condition (4).

We want to prove the statement in a constructive way and want to define for every  $\mathbf{t} \in \mathcal{G}$  an explicit paraboloid  $p_{\mathbf{t}}$  such that these paraboloids fulfill the constraints of problem (7).

In particular, consider an arbitrary but fixed  $\mathbf{t} \in \mathcal{G}$ . Then, the vertices of  $[\mathbf{t} - \Delta \mathbf{t}, \mathbf{t} + \Delta \mathbf{t}]$  are exactly the points in  $\mathcal{N}(\mathbf{t})$  which are further implicitly given by all  $\mathbf{u} \in \{-1, 1\}^n$ , i.e.,  $\mathbf{v}_{\mathbf{u}} := \mathbf{t} + \Delta t \sum_{i=1}^n u_i \mathbf{e}_i$ . Note that  $u_i$  represents the  $i$ th entry of  $\mathbf{u}$  and  $\mathbf{e}_i$  is the  $i$ th unit vector.

Now, we define  $p_{\mathbf{t}} = \sum_{i=1}^n \alpha_i x_i^2 + \beta_i x_i + \gamma$  such that

$$p_{\mathbf{t}}(\mathbf{t}) = f(\mathbf{t}) - \delta, \quad (8)$$

and for all  $\mathbf{u} \in \{-1, 1\}^n$  and  $i \in [n]$ ,

$$\frac{d}{dx_i} p(\mathbf{v}_{\mathbf{u}}) = -u_i 2L. \quad (9)$$

The latter can be re-written as

$$2\alpha_i(t_i + \Delta t) + \beta_i = -2L \quad \wedge \quad 2\alpha_i(t_i - \Delta t) + \beta_i = 2L, \quad (10)$$

for all  $i \in [n]$ . Solving this system of  $2n$  equations, we receive  $\alpha_i = -L/\Delta t$  and  $\beta_i = 2Lt_i/\Delta t$ . As  $\gamma$  is the only degree of freedom left, we calculate it by solving (8). Therefore, such a  $p_{\mathbf{t}}$  exists and is uniquely determined.

With (8), we can set  $s_{\mathbf{t}} = 1$  for the current paraboloid and, thus, with (9), our construction satisfies (7b)-(7d) for the current  $\mathbf{t}$ .

In order to prove conformity with constraints (7e) and (7f), we abstract the situation and assume  $\mathbf{t} = \mathbf{0}$ . From (8) then follows  $\gamma = f(\mathbf{0}) - \delta$  and (9) (or

rather (10)) gives  $\alpha_i = -L/\Delta t$  and  $\beta_i = 0$ . If we can show  $f(\mathbf{x}) - \nu\varepsilon - p(\mathbf{x}) \geq 0$  for all  $\mathbf{x} \in [-\Delta\mathbf{t}, \Delta\mathbf{t}]$ , this transfers to the original box  $[\mathbf{t} - \Delta\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  which directly gives (7e). Even further, it implies the validity of  $v_{\mathbf{d}} = 0$  in (7f) for any  $\mathbf{d} \in \mathcal{G} \cap [\mathbf{t} - \Delta\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$ , as the integral's argument is shown to be non-positive.

From the Lipschitz continuity of  $f$ , we have  $f(\mathbf{x}) \geq f(\mathbf{0}) - L\|\mathbf{x}\|_1 =: \Lambda(\mathbf{x})$ . Note that for the (sub)gradient, it is  $\nabla\Lambda(\mathbf{x}) = -L\text{sgn}(\mathbf{x})$ , where  $\text{sgn}(\mathbf{x})$  is the component-wise sign function. Now, defining the auxiliary function  $g(\mathbf{x}) := \Lambda(\mathbf{x}) - \nu\varepsilon - p(\mathbf{x})$ , we want to show  $\min_{\mathbf{x}} g(\mathbf{x}) \geq 0$ . Suitable candidates for this minimum, have to fulfill  $\nabla g(\mathbf{x}) = \mathbf{0}$ , which gives

$$\mathbf{0} = \nabla g(\mathbf{x}) = -L\text{sgn}(\mathbf{x}) + (2L/\Delta t)\mathbf{x}.$$

So, a vector  $\mathbf{x}'$  solving this equation must fulfill  $x'_i = \pm\Delta t/2$ . From above, we note that  $p(\mathbf{x}') = \sum_{i=1}^n (-L/\Delta t)(\Delta t/2)^2 + f(\mathbf{0}) - \delta = -nL\Delta t/4 + f(\mathbf{0}) - \delta$ . The respective value of  $g$  is

$$\begin{aligned} g(\mathbf{x}') &= f(\mathbf{0}) - L\|\mathbf{x}'\|_1 - \nu\varepsilon - p(\mathbf{x}') \\ &= f(\mathbf{0}) - Ln\Delta t/2 - \nu\varepsilon + nL\Delta t/4 - f(\mathbf{0}) + \delta \\ &= \delta/2 - nL\Delta t/4 \geq \delta/2 - nL\delta/(2nL) = 0, \end{aligned}$$

using the choice  $\nu = \delta/(2\varepsilon)$  and  $\Delta t \leq (2\delta)/(nL)$ . Evaluating  $g$  at  $\mathbf{0}$ , we get  $g(\mathbf{0}) = f(\mathbf{0}) - \delta/2 - p(\mathbf{0}) = \delta/2 > 0$  and it follows  $\min_{\mathbf{x}} g(\mathbf{x}) \geq 0$ . So, we conclude

$$f(\mathbf{x}) - \nu\varepsilon - p(\mathbf{x}) \geq \Lambda(\mathbf{x}) - \nu\varepsilon - p(\mathbf{x}) = g(\mathbf{x}) \geq \min_{\mathbf{x}} g(\mathbf{x}) \geq 0.$$

In particular, (7e) is true for  $p$ , as well as (7f) for  $v_{\mathbf{d}} = 0$ ,  $\mathbf{d} \in \mathcal{G}$ .

Lastly, we prove the feasibility of  $p$  for constraints (7g) and (7h). From above, we have the explicit representation of  $\alpha_i$  and  $\beta_i$ ,  $i \in [n]$ . Hence, we directly derive

$$\begin{aligned} \left| \frac{d}{dx_i} p(\mathbf{a}) \right| &= |2\alpha_i a_i + \beta_i| = |-2La_i/\Delta t + 2Lt_i/\Delta t| = \frac{2L}{\Delta t} |t_i - a_i| \\ &= \frac{2L}{\Delta t} (t_i - a_i) \leq \frac{2L}{\Delta t} (b_i - a_i) \leq \frac{2L}{\Delta t} \|\mathbf{b} - \mathbf{a}\|_{\infty} = C. \end{aligned}$$

This shows that  $p$  satisfies (7g). The case for (7h) follows analogously. This concludes the proof.  $\square$

It remains to show the validity of a solution with zero objective value for the conditions (2) and (3). Before going into detail, we generally investigate the influence of the bounded partial derivatives in (7d), (7g), (7h) on the Lipschitz continuity for the paraboloids.

**Lemma 2.2.** *Let  $p(\mathbf{x}) = \sum_{i=1}^n \alpha_i x_i^2 + \sum_{i=1}^n \beta_i x_i + \gamma$  be a paraboloid of degree two and let  $\mathcal{D}' = [\mathbf{a}', \mathbf{b}']$ . Then, for all  $i \in [n]$ , if it holds true that*

$$\left| \frac{d}{dx_i} p(\mathbf{a}') \right| \leq C \quad \wedge \quad \left| \frac{d}{dx_i} p(\mathbf{b}') \right| \leq C,$$

for some  $C > 0$ , it follows that

$$\forall \mathbf{x} \in \mathcal{D}' : \left| \frac{d}{dx_i} p(\mathbf{x}) \right| \leq C.$$

In particular,  $p$  is Lipschitz continuous on  $\mathcal{D}'$  with Lipschitz constant  $nC$  with respect to  $\|\cdot\|_{\infty}$  and with Lipschitz constant  $C$  with respect to  $\|\cdot\|_1$ .

*Proof.* A detailed proof of this statement can be found in Appendix A.1  $\square$

Due to this lemma, from now on, all  $p^l$  which result from a solution of model (7), in particular, (7g) and (7h), can be treated as Lipschitz continuous with constant  $C$  with respect to  $\|\cdot\|_1$ . Additionally, inequalities (7d) ensure that a paraboloid  $p^l$  with  $s_{\mathbf{t}}^l = 1$  is Lipschitz continuous on the neighborhood  $\mathcal{N}(\mathbf{t})$  of  $\mathbf{t}$  with constant  $2L$  with respect to  $\|\cdot\|_1$ . This will come beneficial for the proofs below.

As mentioned above, we now turn to showing the validity of a solution to (7) with objective zero to properties (2) and (3). For either one, we start by abstracting the situation and show a corresponding lemma first. Afterwards, the validity of the respective property is concluded and shown in a theorem.

For property (2), inequalities eqs. (7b) to (7d) in the model are supposed to do the trick. The following lemma demonstrates in an abstract setting, how the value of a function  $g$  with a Lipschitz property and certain inequality conditions can be bounded from below. Later,  $g$  is replaced by the difference of  $f$  and a respective paraboloid.

**Lemma 2.3.** *Let  $g \in \mathcal{C}^0(\mathcal{D}')$  be Lipschitz continuous with  $L_g > 0$  with respect to  $\|\cdot\|_1$ , where  $\mathcal{D}' = [\mathbf{a}', \mathbf{b}'] \subseteq \mathcal{D}$ . Further, let  $g(\mathbf{v}) \geq 0$  for all vertices  $\mathbf{v}$  of  $\mathcal{D}'$ . Then it holds true that*

$$\min_{\mathbf{x} \in \mathcal{D}'} g(\mathbf{x}) \geq -\frac{L_g n}{n+1} \|\mathbf{b}' - \mathbf{a}'\|_1.$$

*Proof.* A detailed proof of this statement can be found in Appendix A.2.  $\square$

In other terms, the lemma states that the minimum of a Lipschitz function which is non-negative at the vertices of a box is bounded by a term depending on the Lipschitz constant and the “size” of the box (in the respective norm for the Lipschitz continuity). We note that this result analogously holds when we consider non-positivity of  $g$  at the vertices and an upper bound to the maximum of it is claimed, where the right-hand side has a positive sign. This could be used to ensure property (3) too, but we included the variables  $v_{\mathbf{d}}^l$  in (7) and made advantage of the objective in order to get a stronger bound (compare Lemma 2.5).

However, by means of Lemma 2.3, we can show the validity of (2) for a solution of eq. (7).

**Theorem 2.4** (Validity of (2)). *Let  $(\alpha_i^l, \beta_i^l, \gamma^l)$ ,  $s_{\mathbf{t}}^l$ , for  $\mathbf{t} \in \mathcal{G}_\varepsilon$ ,  $l \in [K]$ ,  $i \in [n]$ , be a solution for model (7), in particular for (7b) to (7d). Then, (2) holds true for all  $\mathbf{x} \in \mathcal{D}$ .*

*Proof.* Let  $\mathbf{t} \in \mathcal{G}_\varepsilon \cap [\mathbf{a}, \mathbf{b}]$ . As stated in (6), the union of  $[\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  over all such  $\mathbf{t}$  gives  $\mathcal{D}$ . Hence, we focus on the fixed box  $[\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  for the moment.

Observe that all vertices of  $[\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  are again vectors in  $\mathcal{G}_\varepsilon$ . Therefore, (7b) and (7c) ensure that for all such vertices  $\mathbf{v}$ , there exists  $l_{\mathbf{v}} \in [K]$  such that  $p^{l_{\mathbf{v}}}(\mathbf{v}) \geq f(\mathbf{v}) - \delta$ . We collect these indices in

$$I_{\mathbf{t}} = \{l \in [K] \mid \exists \mathbf{v} \text{ vertex of } [\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}] : p^l(\mathbf{v}) \geq f(\mathbf{v}) - \delta\}.$$

Now, the inequalities (7d) with Lemma 2.2 ensure that for all  $l \in I_{\mathbf{t}}$ , the paraboloid  $p^l$  is Lipschitz continuous on  $[\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  with constant  $2L > 0$  with respect to  $\|\cdot\|_1$ . Indeed, this Lipschitz property transfers to  $\max_{l \in I_{\mathbf{t}}} p^l(\mathbf{x})$ . In order to show this, consider  $\mathbf{x}, \mathbf{y} \in [\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$  and set  $p_{\mathbf{x}}(\mathbf{x}) := \max_{l \in I_{\mathbf{t}}} p^l(\mathbf{x})$ , as well as  $p_{\mathbf{y}}(\mathbf{y}) := \max_{l \in I_{\mathbf{t}}} p^l(\mathbf{y})$ . If  $p_{\mathbf{x}}(\mathbf{x}) \geq p_{\mathbf{y}}(\mathbf{y})$ , we can derive

$$\begin{aligned} \left| \max_{l \in I_{\mathbf{t}}} p^l(\mathbf{x}) - \max_{l \in I_{\mathbf{t}}} p^l(\mathbf{y}) \right| &= |p_{\mathbf{x}}(\mathbf{x}) - p_{\mathbf{y}}(\mathbf{y})| = p_{\mathbf{x}}(\mathbf{x}) - p_{\mathbf{y}}(\mathbf{y}) \\ &\leq p_{\mathbf{x}}(\mathbf{x}) - p_{\mathbf{x}}(\mathbf{y}) \leq |p_{\mathbf{x}}(\mathbf{x}) - p_{\mathbf{x}}(\mathbf{y})| \leq C \|\mathbf{x} - \mathbf{y}\|_1, \end{aligned}$$

where we used the fact  $p_{\mathbf{y}}(\mathbf{y}) \geq p^l(\mathbf{y})$  for all  $l \in I_{\mathbf{t}}$ . The case  $p_{\mathbf{x}}(\mathbf{x}) \leq p_{\mathbf{y}}(\mathbf{y})$  follows analogously. Hence, setting  $g(\mathbf{x}) := \max_{l \in I_{\mathbf{t}}} p^l(\mathbf{x}) - (f(\mathbf{x}) - \delta)$  and using



the triangle inequality, we can show that  $g$  is Lipschitz continuous with constant  $L_g = L + 2L = 3L$  with respect to  $\|\cdot\|_1$ . From above, we additionally have  $g(\mathbf{v}) \geq 0$  for all vertices  $\mathbf{v}$  of  $[\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$ .

This allows to apply Lemma 2.3 and we receive

$$\min_{\mathbf{x}' \in [\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]} g(\mathbf{x}') \geq -\frac{3Ln}{n+1} \|\Delta\mathbf{t}\|_1.$$

From (4), it must hold  $\|\Delta\mathbf{t}\|_1 = \sum_{i=1}^n \Delta t_i \leq \frac{n+1}{n} \frac{\varepsilon - \delta}{3L}$ , which then gives that for all  $\mathbf{x} \in [\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]$ , it is

$$\max_{l \in [K]} p^l(\mathbf{x}) - (f(\mathbf{x}) - \delta) \geq g(\mathbf{x}) \geq \min_{\mathbf{x}' \in [\mathbf{t}, \mathbf{t} + \Delta\mathbf{t}]} g(\mathbf{x}') \geq \delta - \varepsilon,$$

which is equivalent to

$$\max_{l \in [K]} p^l(\mathbf{x}) \geq f(\mathbf{x}) - \varepsilon.$$

As this holds for all such  $\mathbf{t}$ , by the introducing comment of this proof it also holds for the entire  $\mathcal{D}$  which shows the claim.  $\square$

It remains to show the validity of eq. (3) for a solution of (7). We follow an analogous structure and abstract the situation first, before presenting the final proof in a clear manner. As mentioned earlier, the following lemma gives in general a stronger bound than an analogous result of Lemma 2.3, but requires an additional condition on the integral of the function in question.

**Lemma 2.5.** *Let  $g \in \mathcal{C}^0(\mathcal{D}')$  be Lipschitz continuous with  $L_g > 0$  with respect to  $\|\cdot\|_1$ , where  $\mathcal{D}' = [\mathbf{a}', \mathbf{b}'] \subseteq \mathcal{D}$  full dimensional. Further, let  $g(\mathbf{v}) \leq 0$  for all vertices  $\mathbf{v}$  of  $\mathcal{D}'$  and  $\int_{\mathcal{D}'} g(\mathbf{x}) d\mathbf{x} \leq 0$ . Then it holds true that*

$$\max_{\mathbf{x} \in \mathcal{D}'} g(\mathbf{x}) \leq \frac{\sqrt{3}-1}{2} \Delta_{\max} n L_g,$$

where  $\Delta_{\max} = \max_{i \in [n]} b'_i - a'_i$ .

*Proof.* A detailed proof of this statement can be found in Appendix A.3.  $\square$

Considering  $\|\mathbf{b}' - \mathbf{a}'\|_1 \approx n \Delta_{\max}$ , we now formally note that this bound is tighter than the one obtained in Lemma 2.3 (in absolute terms). For this, compare  $0.366 \approx (\sqrt{3}-1)/2 < n/(n+1)$  for all  $n \in \mathbb{N}$ .

By leveraging this lemma and the choices made, we can finally show that property (3) holds true.

**Theorem 2.6** (Validity of (3)). *Let  $(\alpha_i^l, \beta_i^l, \gamma^l), v_{\mathbf{d}}^l$ , for  $\mathbf{d} \in \mathcal{G}$ ,  $l \in [K]$ ,  $i \in [n]$ , be a solution for model (7), in particular for (7e) and (7f), with objective value (7a) of zero. Then, (3) holds true for all  $\mathbf{x} \in \mathcal{D}$ .*

*Proof.* Let  $\mathbf{d} \in \mathcal{G} \cap [\mathbf{a}, \mathbf{b}]$ . As stated in (6), the union of  $[\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]$  over all such  $\mathbf{d}$  gives  $\mathcal{D}$ . Hence, we focus on the fixed box  $[\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]$  for the moment.

Observe that all vertices of this box are again vectors in  $\mathcal{G}$ . Therefore, (7e) ensure that for all  $l \in [K]$ , it is true that  $p^l(\mathbf{v}) \leq f(\mathbf{v}) - \nu\varepsilon$  for all such vertices  $\mathbf{v}$ .

Now, we fix  $l \in [K]$  and investigate  $g(\mathbf{x}) := p^l(\mathbf{x}) - (f(\mathbf{x}) - \nu\varepsilon)$ . The upper considerations give  $g(\mathbf{v}) \leq 0$  for all vertices  $\mathbf{v}$  of  $[\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]$ . In addition, the triangle inequality gives that  $g$  is Lipschitz continuous with constant  $L_g = C + L$  with respect to  $\|\cdot\|_1$ . With (7f) and the assumed objective value of zero it follows that  $0 \geq \int_{[\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]} g(\mathbf{x}) d\mathbf{x}$ . Therefore, we can apply Lemma 2.5 and receive

$$p^l(\mathbf{x}) - (f(\mathbf{x}) - \nu\varepsilon) = g(\mathbf{x}) \leq \max_{\mathbf{x}' \in [\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]} g(\mathbf{x}') \leq \frac{\sqrt{3}-1}{2} n(C+L) \max_{i \in [n]} \Delta d_i,$$

for all  $\mathbf{x} \in [\mathbf{d}, \mathbf{d} + \Delta\mathbf{d}]$ . From the choice  $\Delta d_i \leq (2\nu\varepsilon)/((\sqrt{3}-1)n(C+L))$ , we conclude

$$p^l(\mathbf{x}) \leq f(\mathbf{x}).$$

As  $l \in [K]$  was chosen arbitrarily and the statement holds for all such  $\mathbf{d}$ , the introducing comment implies that this statement holds for all  $\mathbf{x} \in \mathcal{D}$ . This shows the claim.  $\square$

We have now established a model which is suitable to find an approximation with the desired properties. In the following section, this approach is embedded in a binary search to determine a minimal number of paraboloids in order to incorporate the result as a relaxation into the original problem.

### 3. THE GBM METHOD: REDUCING MINLP TO MIQCP

Endowed with the approximate optimization problem from Section 2 that delivers a parabolic approximation, we are now ready to state our framework, which is capable of turning an MIQCP solver into a rather general MINLP solver. In an attempt to summarize the following framework in one expression, we came up with “global one-sided best approximations and relaxation by a minimal number of paraboloids”. We propose to abbreviate it with the first letters of “global”, “best”, and “minimal”. This leads to the GBM method which is incidentally also an acronym of the authors’ surnames.

In more detail, the method describes a two-part algorithmic framework that first determines a small set of paraboloids for each nonlinear, non-quadratic function in (1) to approximate the latter. Second, it replaces the constraints that involve these non-quadratic functions by means of inequalities with respect to the computed paraboloids. The resulting problem constitutes a relaxation to the original one and is finally solved to optimality in order to receive a dual bound, as well as a potential starting point for a refined solving process. In general, feasible points to MINLP problems are considered feasible with respect to some predefined accuracy  $\varepsilon > 0$ . That is, from a practical perspective if the parabolic approximation is tight enough, i.e., with respect to  $\varepsilon$ , the found incumbent might even serve as a global optimal solution for the original problem.

For a detailed breakdown of the GBM method, we start with one theoretical and one practical algorithm to compute a small number of paraboloids that approximate a given Lipschitz continuous function. Afterwards, we describe the algorithmic framework to find approximative optimal solutions and address its benefits, as well as limits.

**3.1. The Search for a Small  $K$ .** For the first part, i.e., determining a *small* set of paraboloids for approximation, we start by setting the prerequisites. We assume to have a function  $f$  with Lipschitz constant  $L$  with respect to  $\|\cdot\|_1$ , which is to approximate. Further, let  $\varepsilon > 0$  be given and consider a choice of  $\delta, \nu, C, \Delta t, \Delta d > 0$  that fits the requirements of (4), (5), as well as Theorem 2.1. The latter gives the existence of a solution to Problem (7) with  $\bar{K} = |\mathcal{G}_\varepsilon|$ , which meets the conditions (2) and (3), i.e., the necessary approximation accuracy.

As there might exist a number of paraboloids  $K^* < \bar{K}$ , we propose a binary search in the discrete interval  $[1, \bar{K}] =: [\underline{K}, \bar{K}]$  to find it. That is, omitting a potentially necessary rounding for the moment, we start by solving (7) with  $K = \bar{K}/2$ . If an optimal solution with objective zero is returned, we set  $\bar{K} \leftarrow K$ , otherwise,  $\underline{K} \leftarrow K + 1$ , and restart the search with respect to the new interval. This procedure is terminated when  $\bar{K} \leq \underline{K}$  and  $K^* \leftarrow \bar{K}$  is returned as the minimal number of paraboloids. A formalization can be found in Algorithm 1.

---

**Algorithm 1** Number of Paraboloids – Binary Search

---

**Input:** an upper bound of paraboloids  $\bar{K}$ **Output:** the minimal number of paraboloids  $K^*$  satisfying (2) and (3)

```

1: Set  $\underline{K} \leftarrow 1$ 
2: while  $\bar{K} > \underline{K}$  do
3:   Set  $K \leftarrow \lfloor (\bar{K} + \underline{K})/2 \rfloor$ 
4:   Solve (7) for  $K \rightarrow$  “infeasible” or objective  $c^K$ 
5:   if “infeasible” or  $c^K > 0$  then
6:     Set  $\underline{K} \leftarrow K + 1$ 
7:   else
8:     Set  $\bar{K} \leftarrow K$ 
9:   end if
10: end while
11: return  $K^* \leftarrow \bar{K}$ 

```

---

We note that the minimality is considered with respect to our MIP formulation. In particular, there might exist an approximation with fewer paraboloids that satisfies the conditions (2) and (3), but is infeasible for (7), e.g., as two of the paraboloids do not overlap at a predefined discretization point. In addition, dependent on the particular form of  $f$ , an analytical solution may be computable by means of analytical and numerical methods. However, we aim for a computational procedure and, thus, do not want to depend on case-by-case analysis.

Regarding the running time of Algorithm 1, we note that the while loop is executed  $\mathcal{O}(\log_2(\bar{K}))$  times. Taking the size of  $\bar{K} = |\mathcal{G}_\varepsilon|$  into account, this is  $\mathcal{O}(n \log_2(n/\varepsilon))$  loops. Further, there exist  $\bar{K} \cdot |\mathcal{G}_\varepsilon| = |\mathcal{G}_\varepsilon|^2$  binary variables in (7) which is in  $\mathcal{O}((n/\varepsilon)^{2n})$ . Considering that (7) can be solved in polynomial time if we fix an assignment of the binary variables, we get  $\mathcal{O}^*((n/\varepsilon)^{2n} \log_2(n/\varepsilon))$  as a rough estimate of the running time, where the asterisk denotes an omission of polynomial terms. Note that we could have refined some bounds, e.g., by excluding certain binary assignments via (7c), but only want to highlight the order of magnitude by this theoretical framework.

Such a running time appears tremendous, which motivates certain adjustments in the practical implementation of above methodology. First, we relax (7) by neglecting constraints (7d). Second,  $\Delta t_i$  and  $\Delta d_i$  are not chosen as claimed in (4) and (5), but according to fixed numbers of discretization points.

These measures lead to significant lower solution times, as the number of inequalities and (binary) variables is decreased, but a solution may lack the desired properties (2) and (3). In order to overcome this issue, we check for these properties by solving the unconstrained NLP problems  $\min_{\mathbf{x}}(f(\mathbf{x}) - \varepsilon - \max_l p^l(\mathbf{x}))$  and  $\min_{\mathbf{x}}(\max_l p^l(\mathbf{x}) - f(\mathbf{x}))$  on the domain  $\mathbf{x} \in \mathcal{D}$  for a given solution  $(p^l)_l$ . This is achieved by rearrangement to  $K$  inequalities or by solving  $K$  simpler problems, respectively. If the respective objective is negative, the solution fulfills the desired property and we terminate. Otherwise, the number of paraboloids and/or the number of discretization points are increased and another loop is executed. We note that we “shift” a solution  $(p^l)_l$  by decreasing their constant terms for the objective value of the first NLP to solve, if this value is negative. We summarize the practical implementation in Algorithm 2.

**3.2. Parabolic Relaxation.** In the previous subsection, we established one theoretical and one practical method to compute a one-sided parabolic approximation of a function  $f$  that satisfies an approximation guarantee  $\varepsilon > 0$ . We note once again

**Algorithm 2** Number of Paraboloids – Practical Implementation

---

**Input:** an upper bound of paraboloids  $\bar{K}$ , start numbers of discretization points  $T_0, D_0 \in \mathbb{N}$  and paraboloids  $K_0 < \bar{K}$

**Output:** a small number of paraboloids  $K^*$  satisfying (2) and (3)

- 1: Set  $T \leftarrow T_0, D \leftarrow D_0, K \leftarrow K_0$
- 2: **while**  $K < \bar{K}$  **do**
- 3:   Set  $T$ -bound  $\leftarrow$  **false**,  $D$ -bound  $\leftarrow$  **false**
- 4:   Compute  $\Delta t_i, \Delta d_i$  according to  $T, D$  for  $i \in [n]$
- 5:   Solve (7) without (7d), but with  $\Delta t_i, \Delta d_i, i \in [n] \rightarrow$  “infeasible” or solution  $(p^l)_{l \in [K]}$  with objective  $c^K$
- 6:   **if** “infeasible” **or**  $c^K > 0$  **then**
- 7:     Increment  $K$
- 8:   **else**
- 9:     Check (3) by minimizing  $c^l := p^l - f, l \in [K]$
- 10:    **if**  $\exists l \in [K] : c^l > 0$  **then**
- 11:     Increase  $T$
- 12:    **else**
- 13:     Set  $T$ -bound  $\leftarrow$  **true**
- 14:    **end if**
- 15:    Shift each  $p^l$  by  $-c^l, l \in [K]$
- 16:    Check (2) by minimizing  $c_\varepsilon := y - (f - \varepsilon)$  s.t.  $y \geq p^l, l \in [K]$
- 17:    **if**  $c_\varepsilon < 0$  **then**
- 18:     Increase  $D$
- 19:    **else**
- 20:     Set  $D$ -bound  $\leftarrow$  **true**
- 21:    **end if**
- 22:   **end if**
- 23:   **if**  $T$ -bound **and**  $D$ -bound **then**
- 24:     Exit loop
- 25:   **end if**
- 26: **end while**
- 27: **return**  $K^* \leftarrow K$

---

that these methods are analogously valid when approximating from “the other side”, as one can consider  $-f$  and flip the coefficients’ signs of the resulting paraboloids.

Now, in a direct approach, we can compute the approximations for every function occurring in the constraints of (1) and then replace them. This reduces the stated MINLP to a MIQCP. In particular, for every  $j \in J$ , we compute a small number  $K^j \in \mathbb{N}$  of paraboloids  $(p_j^l)_{l \in [K^j]}$  that approximate  $f_j$  from one side with respect to  $\varepsilon > 0$ . That is, it holds

$$f(\mathbf{x}) - \varepsilon \leq \max_{l \in [K^j]} p_j^l(\mathbf{x}) \leq f(\mathbf{x}), \quad \text{for all } \mathbf{x} \in \Omega.$$

Then, we replace each  $f_j$  in (1) with its approximation, leading to

$$\begin{aligned} & \min c(\mathbf{x}) \\ & \text{s.t. } \max_{l \in [K^j]} p_j^l(\mathbf{x}) \leq 0, \quad j \in J, \\ & \mathbf{x} \in \Omega, \end{aligned}$$

or, equivalently,

$$\min c(\mathbf{x}) \tag{11a}$$

$$\text{s.t. } p_j^l(\mathbf{x}) \leq 0, \quad j \in J, l \in [K^j], \tag{11b}$$

$$\mathbf{x} \in \Omega. \tag{11c}$$

Without loss of generality, we can assume that  $c(\mathbf{x})$  is a linear function, as otherwise we can reformulate it as a (possibly non-convex) constraint and approximate it too. Hence, the original MINLP problem was entirely reformulated in terms of a MIQCP problem, which allows solvers tailored for the latter (e.g., GloMIQO [20], Gurobi [16], SCIP [6]) to be applied for a more general problem class. If an exact solution to (11) can be obtained in such a way, it directly serves as an  $\varepsilon$ -approximate solution for (1) and, thus, solves the original problem (with respect to the accuracy).

Naturally, the reduction to the more specific problem class of MIQCP problems is not without cost. Some computation time must be invested in finding the paraboloids in the first place, which, if exceeding the solution time for the original problem, appears useless on first sight. However, similar to approaches in the field of artificial intelligence, once computed, we can reuse the approximations in every optimization problem that contains such constraints on smaller or equal bounds. This motivates the creation of some sort of lookup table a-prior that can be consulted when solving a bounded MINLP with “common” constraint functions.

Although the theoretical running time of determining a small number of approximating paraboloids is exponential in the dimension, its computation allows for a direct replacement of the multi-dimensional constraints without introduction of further variables. Nevertheless, if available, one can try to leverage factorability, as explained in the following.


In particular, we note that most practical problems of type (1) involve only factorable functions, which allows for a representation as an expression tree, see, for instance, [24] and we call the problem *factorable* MINLP. That is, each constraint function comprises of nested elementary operations (such as ‘+’, ‘.’ or one-dimensional, univariate functions), variables and constants. This can be displayed in terms of an acyclic graph where every node is such an operation, variable, or constant. By introduction of auxiliary variables a factorable MINLP can be transformed to only contain one-dimensional functions, as well as linear and bivariate terms. For an explanatory introduction of this procedure, see [21].

The MINLPLib [10] constitutes a test set of various MINLPs, which are in main parts available in the OSIL format [13] that mimics the expression tree in XML format. In Figure 1, we display the number of occurrences of one dimensional functions for all instances that are available in OSIL format (1594 out of 1601, on 23rd May 2024). Note that we excluded the square function, as this is considered to be kept anyways. Further, we just counted the occurrence of ‘power’ without distinguishing whether the exponent or base is a variable/constant.

As we can see, the exponential, the cosine, the sine, and the square root function construct most problems in this test set. Hence, parabolic approximation to these functions can be used repeatedly when tackling a subset of these problems. Naturally, the domain of their arguments varies dependent on the particular instance. However, an approximation for a wider domain also shows the desired behavior for a narrower one. Therefore, we take up the idea from above and propose to approximate these one dimensional functions with respect to some (e.g., the most common) domains and different approximation guarantees a-priori. The result would serve as a lookup table when tackling problems from the MINLPLib.

We note that a concatenation of approximations, i.e., sets of paraboloids, of two overlapping domains might not be a one-sided approximation for the united domain. As we compute each of the paraboloids with respect to the original domain, one paraboloid can even intersect the function to approximate at a bound and leads to a violation of property (3) right outside the domain.

FIGURE 1. Cumulative occurrences of one dimensional function types in MINLPLib instances in OSIL format



**Remark 3.1.** For the special case of the sine (and, thus, cosine) function, we state an enhanced procedure which allows for concatenation. At first, one can find a parabolic approximation to the sine function from  $-\pi/2$  to  $3\pi/2$  and hereby restrict the sign of the quadratic coefficients to be non-positive. Then, we can observe that the approximation fulfills (3) on  $\mathbb{R}$  entirely, as the paraboloids intersect somewhere below  $-1$  at the bounds and decrease outside of the domain. At a second step, an approximation for an arbitrary (finite) domain can be computed by simply shifting the paraboloids appropriately and concatenating the solutions due to the periodicity of the sine. We even propose an approach by integer programming for unbounded domains for sine functions, where the integer variable keeps track of the period as an offset with respect to  $[-\pi/2, 3\pi/2]$  and shifts the paraboloids accordingly. Naturally, both ideas work analogously for the cosine function.

Now, we assume to have a factorable MINLP at hand. Further, suppose its reformulation in one dimensional functions and bivariate terms as described above. In order to find appropriate approximations for the functions, we propose bound propagation, as this may have crucial impact for tightening their domains. Then, a one dimensional function  $f$  in an inequality is replaced by its approximations in the analogous form, whereas an equality constraint is reformulated in terms of both,  $\leq$ - and  $\geq$ -constraints. Again, this reduces the character of a MINLP to the one of a MIQCP.

If we tackle this reformulated problem with the GBM method, we potentially decrease the computation time for the parabolic approximation, as the dimension for the functions to approximate is reduced to one. However, the dimensionality is in some sense shifted to the problem itself due to the lifting in the reformulation. In addition, the approximation guarantee for a one-dimensional component of an original constraint  $f_j$  might not propagate directly. The latter can worsen the overall accuracy and requires thorough analysis of the expression tree's depth. Nevertheless, we think of these aspects as accountable and propose a combination with established methods such as spatial branch-and-bound for future research.

**Remark 3.2.** *At last, we want to emphasize that with the introduction of the first paraboloid for a one-sided approximation of a nonlinear function, the corresponding relaxation of the MINLP becomes a MIQCP as it introduces the required quadratic functions. When solving this MIQCP, state-of-the-art solvers typically lift the problem to a higher dimension using expression trees, in which quadratic functions are reduced to simple squares. Each additional paraboloid can therefore be considered as an additional linear constraint in the extended space. This allows us to control the complexity of our approach, since tighter paraboloid relaxations can be realized by adding linear constraints. From this point of view, it is natural to refer to the one-sided approximations as non-convex cuts for MINLPs.*

#### 4. COMPUTATIONAL RESULTS

In order to supplement the above theoretical statements, we carry out computational showcases in this section. We assume factorizability of the problem at hand and thus a reformulation to one-dimensional constraint functions, for instance with the help of expression trees, as shown in the previous section. Under this assumption, we first provide one-sided parabolic approximations for common functions in MINLPs with respect to fixed domains and several accuracy values. Second, we employ parts of these findings to apply the GBM-method to specific instances of the MINLPLib [10] for the first time.

All computational experiments have been conducted on a machine with Apple M2 Pro chip and 32 GB of RAM. Regarding software, macOS 14.4.1 with Darwin 23.4.0 kernel was used. All implementations are based on Python 3.11.7. For solving the respective optimization problems, we leveraged GAMS in version 46.4.0, which includes Gurobi 11.0.1 [16] and SCIP 8.1 [6]. If not stated otherwise, default settings are used. All code is available at <https://github.com/adriangoess/gbm-method> under a MIT license.

**4.1. Practical Parabolic Approximation.** As displayed in Figure 1, the four function types exponential, cosine, sine and square root are the most common ones in the well-known test set MINLPLib [10]. Since exponential growth is presumably hard to represent by quadratic functions, i.e., paraboloids, we take the considerably tight domain  $[-2, 2]$  into account. As the results presented below indicate, this provides a challenging problem already. Regarding the trigonometric functions, we note that the cosine is equivalent to the sine function up to a shift. Hence, we omit it and consider only the latter on one entire period  $[-\pi/2, 3\pi/2]$ . Since we can reformulate the square root into a quadratic function by squaring, we will not approximate this function. In addition to the four most common one-dimensional function types from above, we also check for higher order polynomial approximations, in particular  $x^3$ . For a comparison to the exponential function, we consider the same domain. Regarding accuracy, we test exponentially decaying, i.e.,  $\varepsilon \in \{10^0, 10^{-1}, 10^{-2}\}$ , in a standardized way. At the end of this section, we address the handling of higher accuracies and the necessary computing resources as well as the tailored choice of parameters.

Since we suggested the practical Algorithm 2 for finding a small number of paraboloids, we need to specify a choice for  $T_0$ ,  $D_0$ ,  $K_0$ . This is determined by the function to be approximated (especially its Lipschitz constant), the domain width, and the approximation accuracy. We give the following formula as a rule of thumb:

$$T_0 = D_0 = \lceil L|D|/(10\varepsilon) \rceil.$$

In particular, we included the respective values as seemed natural to us, i.e., the steeper the function, larger the domain, or the smaller the approximation accuracy, the more discretization points are required. To keep the initial solution times low,

we added a division by 10. For  $K_0$ , we initialized a value of 1 for the largest  $\varepsilon$ . Any smaller  $\varepsilon$  was given the “minimum” number of paraboloids found for the next larger one as a starting value.

The resulting MIPs are solved using Gurobi from within the GAMS framework. We enforced a time limit of 3600 s for each MIP in the search loop and increased the number of paraboloids if no solution was found, i.e., counting it as an infeasible run. Please note that we approximated the mentioned functions *exp* and *sin* from below and above. For  $x^3$ , an approximation from below suffices, as it is symmetric around the origin and, therefore,  $-p(-x)$  provides the corresponding approximation from above, where  $p$  is an approximating paraboloid from below. For the implementation, an approximation from above is performed by applying the algorithm for the underestimating case to the negative of the original function and swapping the sign of the parabolic parameters at the end. In Table 1, we present the number of paraboloids determined by this procedure depending on accuracy, function type, and approximation direction.

		$\varepsilon$		
		$10^0$	$10^{-1}$	$10^{-2}$
sin	below	1	3	12
	above	1	5	46
exp	below	2	4	15
	above	2	5	20
$x^3$	either	3	10	79

TABLE 1. “Minimal” number of paraboloids to approximate the functions from either side with different accuracy.

Interestingly, the numbers of paraboloids appear to be in the same order of magnitude as the accuracy  $\varepsilon$  yet reversed, i.e., they increase exponentially. We notice that approximating the sine from above with  $\varepsilon = 10^{-2}$  requires significantly more paraboloids in comparison to the approximation from below and the exponential function from both sides. We attribute this to the special shape of the sine function on the chosen domain. Specifically, it is symmetric and requires an approximation on both the “left” and the “right” side. See Figure 2 for a graphical representation of this result. For a visualization of the other results see Figure 5, Figure 6, and Figure 7 in Appendix B.2.

As far as  $x^3$  is concerned, we need a significantly higher number of paraboloids compared to the other two functions. We strongly suspect that the comparatively high maximum slope of 12 on the chosen interval and the additional concatenation of a concave and a convex part are the main reasons for this. Since paraboloids are either convex or concave, the latter property increases the difficulty of being approximated by them.

For each setting, which consists of a function  $\sin/\exp/x^3$ , the approximation direction, and an accuracy  $\varepsilon$ , every iteration of the search loop is divided into three computationally relevant steps: Modeling the optimization problem, solving it, and evaluating a solution for the required approximation properties. The modeling and evaluation step took a maximum of 15 s and 41 s, respectively. We obtained similar values for the solution step with  $\varepsilon \in \{10^0, 10^{-1}\}$  except for  $x^3$  with  $\varepsilon = 10^{-1}$ . However, for  $\varepsilon = 10^{-2}$ , the time spent solving the MIP problems was about 13 000 s for the sine, 18 000 s to 20 000 s for the exp function and 33 000 s for  $x^3$ ; see Table 5 in Appendix B.1 for details. In other words, for the low accuracy cases and for the modeling and evaluation steps, the numbers appear negligible compared to those for






FIGURE 2. Approximation of  $\sin(x)$  for  $x \in [-\pi/2, 3\pi/2]$  from above by 46 paraboloids each with  $\varepsilon = 10^{-2}$ .

solving the MIP problems up to high accuracies. This results from exceeding the time limit of one hour several times, which is interpreted as an infeasible solution, see above.

However, we emphasize once more the one-time necessity of this procedure. With the result available in the form of a lookup table, we can reuse it for any instance that includes the approximated function up to the specified domain (or even more, compare Remark 3.1). Hence, when faced with a large number of similar problem instances, approximating the involved constraint functions once in advance seems to be a fair trade.

As the reader may notice, we do not report results for smaller values of  $\varepsilon$ . In fact, we conducted runs with time limits of 3600s and 14400s for the resulting MIPs in Algorithm 2, including smaller values compared to the rule of thumb (1) for the number of discretization points. However, after up to six hours, we received memory errors for all but one setting. This shows its current limitations, and we propose improvements in this regard as a future research direction, e.g., by identifying valid cuts for significant speedups in the solution process.

**4.2. Testing the GBM Method.** In a first showcase of the GBM method, we use the repeated occurrence of identical functions and variable bounds. For this purpose, we choose the `lnts[m]` instances of the MINLPLib for  $m \in \{50, 100, 200, 400\}$ , as they contain sine and cosine functions on the domain  $[-\pi/2, \pi/2]$ . In particular, the number  $m$  represents the number of occurrences of the two functions, while there are  $5m + 6$  continuous variables and  $4m$  constraints.

In order to assess the quality of our relaxations, we use the optimality gap, which is most commonly used in the literature, as a measure. For the sake of completeness, we define it in the following as an absolute and relative number. Hence, let  $c^*$  be the optimal or best known value of Problem (1) and  $\bar{d}$  be a given lower (i.e. dual) bound for  $c^*$ . Naturally,  $\bar{d} \leq c^*$  and we define the corresponding *absolute optimality gap* as

$$\text{absgap} = c^* - \bar{d}.$$

Its relative analog is then computed with respect to  $c^* \neq 0$  as

$$\text{relgap} = (c^* - \bar{d})/c^* = \text{absgap}/c^*.$$

The best values for `absgap` for the `lnts` instances detected so far in the MINLPLib range from about 0.05 to 0.1 and correspond to `relgap` values of roughly 8.7% to 18.9%.

In terms of one-sided parabolic approximations, we run the practical implementation of Algorithm 2 once more, as the relevant domain is significantly smaller than in




FIGURE 3. Approximation of  $\sin(x)$  for  $x \in [-\pi/2, \pi/2]$  from below and above by eight paraboloids each with  $\varepsilon = 10^{-2}$ .




FIGURE 4. Approximation of  $\cos(x)$  for  $x \in [-\pi/2, \pi/2]$  from below and above by three and five paraboloids, respectively, with  $\varepsilon = 10^{-2}$ .

	lnts50		lnts100		lnts200		lnts400	
	orig	para	orig	para	orig	para	orig	para
gurobi	14 400	14 400	14 400	14 400	14 400	14 400	14 400	14 400
scip	14 400	10	14 400	26	14 400	155	14 400	13 531

TABLE 2. Comparison of running times in seconds.

the computational study of Section 4.1. Hereby, we directly consider  $\varepsilon = 10^{-2}$ , as the practical problem complexity is not supposed to increase significantly, compare Remark 3.2. This accuracy results in a considerably small number of paraboloids (i.e., three, five, and eight) from both sides. We illustrated the approximations in Figure 3 for the sine and in Figure 4 for the cosine function.

For the incorporation of these approximations, we followed the procedure described in Section 3.2. To this end, we reformulated each instance in terms of one-dimensional functions and linear/quadratic parts. This was complemented by an expression tree based bound propagation of the natural bounds of the respective arguments. As all sine and cosine functions are involved in equality constraints, we replaced them by a variable and bounded it by the paraboloids from either side. Regarding the instances, we distinguish the relaxed MIQCP instance from above by the short notation *para* compared to the original problem *orig*.

We solved all four instances in both, original and relaxed form with the commercial solver Gurobi and the non-commercial solver SCIP. For the limits, we set a maximal time limit of 14 400s and a relative optimality gap of 0.001%. A summary of the respective solution times can be found in Table 2.

	lnts50		lnts100		lnts200		lnts400	
	orig	para	orig	para	orig	para	orig	para
gurobi	0.0001	0.0023	0.0098	0.0031	0.0063	0.0033	0.0064	0.1046
scip	0.0154	0.0019	0.0398	0.0019	0.0474	0.0019	0.1036	0.0019

TABLE 3. Comparison of absgap.

	lnts50		lnts100		lnts200		lnts400	
	orig	para	orig	para	orig	para	orig	para
gurobi	0.01	0.42	1.77	0.57	1.13	0.59	1.15	18.86
scip	2.78	0.34	7.17	0.34	8.54	0.34	18.69	0.34

TABLE 4. Comparison of relgap in %.

We observe that Gurobi fails to solve either form of each instance within the specified time limit. SCIP, however, hits the time limit only for the original problem formulation, but succeeds in solving the relaxed MIQCP problems to optimality. For lnts50, lnts100, and lnts200, this is even achieved in a mere fraction of the time limit. This demonstrates that the problem resulting from the GBM method is significantly easier to tackle for SCIP. As the problems are solved with respect to a tight relative gap, we can even consider their primal values as dual values for the original problem.

As introduced above, we now use the optimality gap in order to measure the quality of the solutions received by the *para* formulation. Therefore, we use the best known primal value contained in the MINLPLib for either problem formulation. In fact, these values are as least as good as the ones found during the solution of the original problems (*orig*), hence we always consider the best known values. For the dual bounds, we used the computed ones in either formulation *orig* or *para*. Table 3 and Table 4 show the corresponding comparison of the gaps.

We observe ambiguous results for Gurobi. On the one hand, the solution to the relaxations for the two medium sized instances return a better dual bound in comparison to the original instances within the time limit. On the other hand, Gurobi achieves by far better values on the smallest and the largest instances.

The situation differs for the runs with SCIP. Here, the parabolic approximation is solved to optimality within the time limit and achieves an approximate relative optimality gap of 0.34% throughout all instances. We account this effect to the similar structure shown by all tested instances. The respective values on the original instances exceed this gap by factor 8 to 54 or, in terms of differences, by 2.4% to 18.3%. With increasing number of constraints, the solution time for the reformulated problems increases, though, so does the gap differences achieved over 14 400 s on the original problem. Except for the smallest instance the achieved bounds even provide a better result than Gurobi does on the original instances. We cannot emphasize enough that our method improves the best known relative optimality gaps for lnts50, lnts100, and lnts200 from 8.7%–9.8% to 0.34% in just a few minutes.

In conclusion, the results indicate a remarkable boost in the computation of dual bounds for nonlinear programming problems. We believe to have opened a door for novel approaches in the world of MINLPs that utilize some sort of non-convex cuts in the form of paraboloids. The next natural step is to embed the GBM method into a general MINLP solver and determine its practical relevance through extensive computational studies. Hence, we dedicate this to future research, as it is not in the scope of the current article.

## 5. CONCLUSION &amp; FUTURE WORK

We introduced a MIP approach to compute a global one-sided approximation of multi-dimensional Lipschitz functions by means of paraboloids and showed its correctness. This was leveraged in a novel method, which we call GBM, to determine a small amount of such paraboloids for each occurring constraint function and replacing the respective constraint with its corresponding parabolic approximation. We proposed to store the resulting paraboloids in a lookup table and to make use of them repeatedly, reducing the original MINLP to a MIQCP. Here, depending on the approximation accuracy chosen and the definition of an  $\varepsilon$ -optimal solution, the MIQCP serves as a relaxation or even an equivalent problem statement.

The computational results regarding the approximations demonstrate the functionality of the algorithm on frequently appearing one-dimensional functions in the MINLPLib. They also show the limits of the current approach when tackling high accuracy, steep functions in the sense of a large Lipschitz constant, and/or wide domains. Here, we point out that a program is referred to as a MINLP if only one involved function is nonlinear, regardless of its structure. Thus, in our point of view, there exists a certain lack of gradation within the area of MINLP. One way to account for this could be the ability to approximate the involved functions well enough by polynomials of a certain degree. The current article could hence serve as a first contribution in this direction.

Turning to the showcase for the GBM method, our MIQCP relaxations for three out of four instances are solved in up to three minutes using SCIP. Even for the moderate approximation accuracy used, their solution values reduced the reported optimality gap in the MINLPLib tremendously, indicating the potential for a remarkable boost with GBM.

There is certainly room for improvement that we would dedicate to future research directions and elaborate shortly in the following. To start, we propose the development of cuts for the MIP to compute the one-sided parabolic approximations. This allows to use our approach at runtime when a constraint function is not available in the lookup table or when dealing with black-box functions. In order to at least avoid the first case, one could establish a large lookup table with a variety of function, domains, and accuracies. Supposing a sufficiently large lookup table, the idea of a stand-alone MINLP solver based on our approach appears very promising. One could iteratively make use of higher accuracies, therefore minimize the quadratic constraints and adaptively provide dual bounds. Even further, local search techniques can be used to determine feasible solutions from the dual ones.

With respect to the use of expression trees, our approach is theoretically capable of solving MINLP without factorization and lifting. Therefore, we expect a huge impact on instances for which large variable bounds disfavor the use of expression trees, as feasibility tolerance propagates drastically here. Hence, a comparison of solution accuracy and time when using expression trees or not constitutes one more interesting research direction.

## REFERENCES

- [1] J. H. Ahlberg, E. N. Nilson, and J. L. Walsh. *The Theory of Splines and Their Applications: Mathematics in Science and Engineering: A Series of Monographs and Textbooks*. Vol. 38. Elsevier, 2016.
- [2] I. P. Androulakis, C. D. Maranas, and C. A. Floudas. “ $\alpha$ BB: A global optimization method for general constrained nonconvex problems.” In: *Journal of Global Optimization* 7.4 (Dec. 1, 1995), pp. 337–363. DOI: [10.1007/BF01099647](https://doi.org/10.1007/BF01099647). URL: <https://doi.org/10.1007/BF01099647>.

- [3] B. Beach, R. Burlacu, A. Bärman, L. Hager, and R. Hildebrand. “Enhancements of discretization approaches for non-convex mixed-integer quadratically constrained quadratic programming: Part I.” In: *Computational Optimization and Applications* 87.3 (Apr. 1, 2024), pp. 835–891. DOI: [10.1007/s10589-023-00543-7](https://doi.org/10.1007/s10589-023-00543-7). URL: <https://doi.org/10.1007/s10589-023-00543-7>.
- [4] B. Beach, R. Hildebrand, and J. Huchette. “Compact mixed-integer programming formulations in quadratic optimization.” In: *Journal of Global Optimization* 84.4 (2022), pp. 869–912.
- [5] P. Belotti, C. Kirches, S. Leyffer, J. Linderoth, J. Luedtke, and A. Mahajan. “Mixed-integer nonlinear optimization.” In: *Acta Numerica* 22 (2013), pp. 1–131. DOI: [10.1017/S0962492913000032](https://doi.org/10.1017/S0962492913000032).
- [6] K. Bestuzheva, M. Besançon, W.-K. Chen, A. Chmiela, T. Donkiewicz, J. van Doornmalen, L. Eifler, O. Gaul, G. Gamrath, A. Gleixner, L. Gottwald, C. Graczyk, K. Halbig, A. Hoen, C. Hojny, R. van der Hulst, T. Koch, M. Lübbecke, S. J. Maher, F. Matter, E. Mühmer, B. Müller, M. E. Pfetsch, D. Rehfeldt, S. Schlein, F. Schlösser, F. Serrano, Y. Shinano, B. Sofranac, M. Turner, S. Vigerske, F. Wegscheider, P. Wellner, D. Weninger, and J. Witzig. “Enabling Research through the SCIP Optimization Suite 8.0.” In: 49.2 (2023). DOI: [10.1145/3585516](https://doi.org/10.1145/3585516). URL: <https://doi.org/10.1145/3585516>.
- [7] R. Bojanic and R. DeVore. “On polynomials of best one-sided approximation.” In: *Enseign. Math* 12 (1966), pp. 139–164.
- [8] L. Buchheim Christoph and Trieu. “Quadratic Outer Approximation for Convex Integer Programming with Box Constraints.” In: *Experimental Algorithms*. Ed. by C. Bonifaci Vincenzo and Demetrescu and A. Marchetti-Spaccamela. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013, pp. 224–235.
- [9] R. Burlacu. “Adaptive mixed-integer refinements for solving nonlinear problems with discrete decisions.” 2020.
- [10] M. R. Bussieck, A. S. Drud, and A. Meeraus. “MINLPLib—a collection of test models for mixed-integer nonlinear programming.” In: *INFORMS Journal on Computing* 15.1 (2003), pp. 114–119.
- [11] M. V. Deikalova and A. Y. Torgashova. “Best One-Sided Approximation in the Mean of the Characteristic Function of an Interval by Algebraic Polynomials.” In: *Proceedings of the Steklov Institute of Mathematics* 308.1 (Apr. 1, 2020), pp. 68–82. DOI: [10.1134/S0081543820020066](https://doi.org/10.1134/S0081543820020066). URL: <https://doi.org/10.1134/S0081543820020066>.
- [12] M. A. Duran and I. E. Grossmann. “An outer-approximation algorithm for a class of mixed-integer nonlinear programs.” In: *Mathematical programming* 36.3 (1986), pp. 307–339.
- [13] R. Fourer, J. Ma, and K. Martin. “OSiL: An instance language for optimization.” In: *Computational Optimization and Applications* 45.1 (2010), pp. 181–203.
- [14] B. Geißler, A. Martin, A. Morsi, and L. Schewe. “Using Piecewise Linear Functions for Solving MINLPs.” In: *Mixed Integer Nonlinear Programming*. Ed. by J. Lee and S. Leyffer. New York, NY: Springer New York, 2012, pp. 287–314.
- [15] A. M. Geoffrion. “Generalized benders decomposition.” In: *Journal of optimization theory and applications* 10.4 (1972), pp. 237–260.
- [16] Gurobi Optimization, LLC. *Gurobi Optimizer Reference Manual, Version 11.0*. 2023. URL: <https://www.gurobi.com>.
- [17] R. Horst and H. Tuy. *Global optimization: Deterministic approaches*. Third. Springer Science & Business Media, 1996.

- [18] J. T. Lewis. “Computation of best one-sided  $L_1$  approximation.” In: *Mathematics of Computation* 24.111 (1970), pp. 529–536.
- [19] O. L. Mangasarian, J. B. Rosen, and M. E. Thompson. “Nonconvex Piecewise-Quadratic Underestimation for Global Minimization.” In: *Journal of Global Optimization* 34.4 (2006), pp. 475–488.
- [20] R. Misener and C. A. Floudas. “GloMIQO: Global mixed-integer quadratic optimizer.” In: *Journal of Global Optimization* 57.1 (Sept. 1, 2013), pp. 3–50. DOI: [10.1007/s10898-012-9874-7](https://doi.org/10.1007/s10898-012-9874-7). URL: <https://doi.org/10.1007/s10898-012-9874-7>.
- [21] A. Morsi. “Solving MINLPs on Loosely-coupled Networks with Applications in Water and Gas Network Optimization.” 2013.
- [22] G. Nürnberger. *Approximation by spline functions*. Vol. 1. Springer, 1989.
- [23] H. S. Ryoo and N. V. Sahinidis. “A branch-and-reduce approach to global optimization.” In: *Journal of Global Optimization* 8.2 (Mar. 1, 1996), pp. 107–138. DOI: [10.1007/BF00138689](https://doi.org/10.1007/BF00138689). URL: <https://doi.org/10.1007/BF00138689>.
- [24] H. Schichl and A. Neumaier. “Interval Analysis on Directed Acyclic Graphs for Global Optimization.” In: *Journal of Global Optimization* 33.4 (Dec. 1, 2005), pp. 541–562. DOI: [10.1007/s10898-005-0937-x](https://doi.org/10.1007/s10898-005-0937-x). URL: <https://doi.org/10.1007/s10898-005-0937-x>.
- [25] L. Schumaker. *Spline functions: basic theory*. Cambridge university press, 2007.
- [26] M. Tawarmalani and N. V. Sahinidis. *Convexification and global optimization in continuous and mixed-integer nonlinear programming: theory, algorithms, software, and applications*. Vol. 65. Springer Science & Business Media, 2013.
- [27] J. P. Vielma. “Mixed Integer Linear Programming Formulation Techniques.” In: *SIAM Review* 57.1 (2015), pp. 3–57. DOI: [10.1137/130915303](https://doi.org/10.1137/130915303). eprint: <https://doi.org/10.1137/130915303>. URL: <https://doi.org/10.1137/130915303>.
- [28] T. Westerlund and F. Pettersson. “An extended cutting plane method for solving convex MINLP problems.” In: *Computers & chemical engineering* 19 (1995), pp. 131–136.
- [29] S. Wiese. *A computational practicability study of MIQCQP reformulations*. 2021. URL: <https://docs.mosek.com/whitepapers/miqcqp.pdf> (visited on 06/17/2024).

## APPENDIX A. PROOFS FOR LEMMATA IN SECTION 2

## A.1. Proof of Lemma 2.2.

*Proof.* Let  $i \in [n]$  be arbitrary but fixed. Observe that  $\frac{d}{dx_i}p(\mathbf{x}) = 2\alpha_i x_i + \beta_i$  which is linear in  $x_i$ . As  $\mathbf{x} \in \mathcal{D}'$  implies  $a'_i \leq x_i \leq b'_i$ , the first claim about the general bound of the derivative on  $\mathcal{D}'$  follows directly.

Now, let  $\mathbf{x}, \mathbf{y} \in \mathcal{D}'$ , and  $g(\lambda) := p((1-\lambda)\mathbf{x} + \lambda\mathbf{y})$ . By the chain rule, it follows  $g'(\lambda) = \frac{d}{d\lambda}g(\lambda) = \nabla p((1-\lambda)\mathbf{x} + \lambda\mathbf{y})^\top(\mathbf{y} - \mathbf{x})$ . Using the mean value theorem, it exists  $\hat{\lambda} \in [0, 1]$  such that

$$g(1) - g(0) = g'(\hat{\lambda})(1 - 0) = g'(\hat{\lambda}),$$

and it follows

$$|p(\mathbf{x}) - p(\mathbf{y})| = |g(1) - g(0)| = |g'(\hat{\lambda})| = |\nabla p((1-\hat{\lambda})\mathbf{x} + \hat{\lambda}\mathbf{y})^\top(\mathbf{y} - \mathbf{x})|.$$

In order to show the Lipschitz continuity of  $p$ , we now have to find a bound for the upper. This is achieved by an application of Hölder's inequality, in particular,

$$|p(\mathbf{x}) - p(\mathbf{y})| \leq \|\nabla p((1-\hat{\lambda})\mathbf{x} + \hat{\lambda}\mathbf{y})\|_1 \|\mathbf{y} - \mathbf{x}\|_\infty \leq nC \|\mathbf{y} - \mathbf{x}\|_\infty,$$

and

$$|p(\mathbf{x}) - p(\mathbf{y})| \leq \|\nabla p((1-\hat{\lambda})\mathbf{x} + \hat{\lambda}\mathbf{y})\|_\infty \|\mathbf{y} - \mathbf{x}\|_1 \leq C \|\mathbf{y} - \mathbf{x}\|_1,$$

respectively, where we used the bounded partial derivative in each last step. This shows the second claim.  $\square$

## A.2. Proof of Lemma 2.3.

*Proof.* Let  $g^* := g(\mathbf{x}^*) := \min_{\mathbf{x} \in \mathcal{D}'} g(\mathbf{x})$ . Note that  $g^*$  is finite, as  $g$  is (Lipschitz) continuous and  $\mathcal{D}'$  is compact. Analogously, the point  $\mathbf{x}^* \in \mathcal{D}'$  exists.

Now, by Caratheodory's theorem, there further exist vertices  $\mathbf{v}^1, \dots, \mathbf{v}^{n+1}$  of  $\mathcal{D}'$  such that

$$\mathbf{x}^* = \sum_{j=1}^{n+1} \lambda_j \mathbf{v}^j \quad \wedge \quad \sum_{j=1}^{n+1} \lambda_j = 1 \quad \wedge \quad \lambda_j \geq 0, \text{ for all } j \in [n+1].$$

Next, let  $k \in [n+1]$ . Then, we can re-write the first equality from above to

$$\mathbf{x}^* - \mathbf{v}^k = \sum_{\substack{j=1, \\ j \neq k}}^{n+1} \lambda_j \mathbf{v}^j + (\lambda_k - 1)\mathbf{v}^k = \sum_{\substack{j=1, \\ j \neq k}}^{n+1} \lambda_j \mathbf{v}^j - \sum_{\substack{j=1, \\ j \neq k}}^{n+1} \lambda_j \mathbf{v}^k = \sum_{j=1}^{n+1} \lambda_j (\mathbf{v}^j - \mathbf{v}^k),$$

where we used that the sum of  $\lambda_j$  is one in the second and  $\mathbf{v}^k - \mathbf{v}^k = \mathbf{0}$  in the last step.

By leveraging  $g(\mathbf{v}^k) \geq 0$  from the assumption, the minimality of  $g(\mathbf{x}^*)$ , and the Lipschitz continuity of  $g$  (in this order), we derive

$$\begin{aligned} -g(\mathbf{x}^*) &= 0 - g(\mathbf{x}^*) \leq g(\mathbf{v}^k) - g(\mathbf{x}^*) = |g(\mathbf{v}^k) - g(\mathbf{x}^*)| \leq L_g \|\mathbf{v}^k - \mathbf{x}^*\|_1 \\ &= L_g \left\| \sum_{j=1, j \neq k}^{n+1} \lambda_j (\mathbf{v}^j - \mathbf{v}^k) \right\|_1 = L_g \sum_{i=1}^n \left| \sum_{j=1}^{n+1} \lambda_j (v_i^j - v_i^k) \right|. \end{aligned}$$

Note that the  $i$ th entry of any vertex  $\mathbf{v}$  is either  $a'_i$  or  $b'_i$ . This implies that the inner bracket is non-negative if  $v_i^k = a'_i$  and non-positive if  $v_i^k = b'_i$ . Having this in mind

and rearranging the sums, we can further rewrite the right-hand side of the upper inequality as

$$\begin{aligned}
L_g \sum_{i \in [n]} \left| \sum_{j=1}^{n+1} \lambda_j (v_i^j - v_i^k) \right| &= L_g \left[ \sum_{\substack{i \in [n], j=1 \\ v_i^k = a'_i}}^{n+1} \lambda_j (v_i^j - v_i^k) + \sum_{\substack{i \in [n], j=1 \\ v_i^k = b'_i}}^{n+1} \lambda_j (v_i^k - v_i^j) \right] \\
&= L_g \left[ \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{i \in [n], \\ v_i^k = a'_i}} (v_i^j - v_i^k) + \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{i \in [n], \\ v_i^k = b'_i}} (v_i^k - v_i^j) \right] \\
&= L_g \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{i \in [n], \\ v_i^k \neq v_i^j}} (b'_i - a'_i),
\end{aligned}$$

where  $\lambda_j \geq 0$  allowed to omit the absolute value in the first equality and we took into account that the inner brackets are  $b'_i - a'_i$  if and only if  $v_i^k \neq v_i^j$ . In conclusion, we receive

$$-g(\mathbf{x}^*) \leq L_g \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{i \in [n], \\ v_i^k \neq v_i^j}} (b'_i - a'_i).$$

Summing this over  $k$  gives

$$\begin{aligned}
-(n+1)g(\mathbf{x}^*) &\leq L_g \sum_{k=1}^{n+1} \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{i \in [n], \\ v_i^k \neq v_i^j}} (b'_i - a'_i) = L_g \sum_{j=1}^{n+1} \lambda_j \sum_{k=1}^{n+1} \sum_{\substack{i \in [n], \\ v_i^k \neq v_i^j}} (b'_i - a'_i) \\
&= L_g \sum_{j=1}^{n+1} \lambda_j \sum_{\substack{k=1, i \in [n], \\ k \neq j, v_i^k \neq v_i^j}}^{n+1} (b'_i - a'_i) \leq L_g \sum_{j=1}^{n+1} \lambda_j n \sum_{i=1}^n (b'_i - a'_i) \\
&= L_g n \sum_{j=1}^{n+1} \lambda_j \|\mathbf{b}' - \mathbf{a}'\|_1 = L_g n \|\mathbf{b}' - \mathbf{a}'\|_1.
\end{aligned}$$

A division by  $-(n+1)$  then gives the desired bound and finishes the proof.  $\square$

### A.3. Proof of Lemma 2.5.

*Proof.* Let  $g^* := g(\mathbf{x}^*) := \max_{\mathbf{x} \in \mathcal{D}'} g(\mathbf{x})$ . Again, note that  $g^*$  is finite and such a point  $\mathbf{x}^*$  exists due to the (Lipschitz) continuity of  $g$  and the compactness of  $\mathcal{D}'$ .

As  $g$  is Lipschitz, we have

$$g^* - g(\mathbf{x}) = |g^* - g(\mathbf{x})| \leq L_g \|\mathbf{x} - \mathbf{x}^*\|_1,$$

which is equivalent to

$$g(\mathbf{x}) \geq g^* - L_g \|\mathbf{x} - \mathbf{x}^*\|_1 =: \Lambda(\mathbf{x}), \quad (12)$$

for all  $\mathbf{x} \in \mathcal{D}'$ . Now, with the assumption and by writing  $V := \text{vol}(\mathcal{D}')$ , we derive

$$\begin{aligned}
0 &\geq \int_{\mathcal{D}'} g(\mathbf{x}) d\mathbf{x} \geq \int_{\mathcal{D}'} \Lambda(\mathbf{x}) d\mathbf{x} = \int_{\mathcal{D}'} g^* - L_g \|\mathbf{x} - \mathbf{x}^*\|_1 d\mathbf{x} \\
&= V g^* - L_g \int_{\mathcal{D}'} \|\mathbf{x} - \mathbf{x}^*\|_1 d\mathbf{x} = V g^* - L_g \int_{[\mathbf{a}' - \mathbf{x}^*, \mathbf{b}' - \mathbf{x}^*]} \|\mathbf{x}\|_1 d\mathbf{x}
\end{aligned}$$



$$\begin{aligned}
&= Vg^* - \\
&L_g \left[ \frac{1}{2} \text{vol}([\mathbf{a}' - \mathbf{x}^*, \mathbf{b}' - \mathbf{x}^*]) \sum_{i=1}^n \left( b'_i - x_i^* - (a'_i - x_i^*) + 2 \frac{(b'_i - x_i^*)(a'_i - x_i^*)}{b'_i - x_i^* - (a'_i - x_i^*)} \right) \right] \\
&= Vg^* - L_g \frac{1}{2} \text{vol}([\mathbf{a}', \mathbf{b}']) \left[ \sum_{i=1}^n (b'_i - a'_i) + 2 \frac{(b'_i - x_i^*)(a'_i - x_i^*)}{b'_i - a'_i} \right] \\
&= Vg^* - \frac{L_g V}{2} \left[ \|\mathbf{b}' - \mathbf{a}'\|_1 + 2 \sum_{i=1}^n \frac{(b'_i - x_i^*)(a'_i - x_i^*)}{b'_i - a'_i} \right].
\end{aligned}$$

A rearrangement gives

$$g^* \leq \frac{L_g}{2} \|\mathbf{b}' - \mathbf{a}'\|_1 + L_g \sum_{i=1}^n \frac{(b'_i - x_i^*)(a'_i - x_i^*)}{b'_i - a'_i}, \quad (13)$$

where it remains to investigate the sum. We note that  $0 < b'_i - a'_i \leq \Delta_{\max}$  for all  $i \in [n]$  by definition. In addition, each vertex  $\mathbf{v}$  of  $\mathcal{D}'$  is either  $a'_i$  or  $b'_i$  in its  $i$ th entry. Hence, there exists a vertex  $\mathbf{v}$  of  $\mathcal{D}'$  which attains the minimum in the inequality

$$(b'_i - x_i^*)(x_i^* - a'_i) \geq \min\{(b'_i - x_i^*), (x_i^* - a'_i)\}^2 = (v_i - x_i^*)^2,$$

for all  $i \in [n]$ . Using this  $\mathbf{v}$ , we can bound the sum in (13) as

$$\sum_{i=1}^n \frac{(b'_i - x_i^*)(a'_i - x_i^*)}{b'_i - a'_i} \leq -\frac{1}{\Delta_{\max}} \sum_{i=1}^n (v_i - x_i^*)^2 = -\frac{1}{\Delta_{\max}} \|\mathbf{v} - \mathbf{x}^*\|_2^2.$$

As (12) is especially true for  $\mathbf{v}$ , the assumption to be non-positive at vertices gives  $0 \geq g(\mathbf{v}) \geq g^* - L_g \|\mathbf{v} - \mathbf{x}^*\|_1$  which is equivalent to  $\|\mathbf{v} - \mathbf{x}^*\|_1 \geq g^*/L_g$ . Combined with the well-known estimate  $\sqrt{n}\|\mathbf{x}\|_2 \geq \|\mathbf{x}\|_1$ , which is equivalent to  $\|\mathbf{x}\|_2^2 \geq \frac{1}{n}\|\mathbf{x}\|_1^2$ , we receive

$$-\frac{1}{\Delta_{\max}} \|\mathbf{v} - \mathbf{x}^*\|_2^2 \leq -\frac{1}{\Delta_{\max} n} \|\mathbf{v} - \mathbf{x}^*\|_1^2 \leq -\frac{(g^*)^2}{\Delta_{\max} n L_g^2}.$$

In summary, (13) breaks down to

$$g^* \leq \frac{L_g}{2} \|\mathbf{b}' - \mathbf{a}'\|_1 - \frac{(g^*)^2}{\Delta_{\max} n L_g},$$

which can be rearranged to

$$\frac{1}{2}(g^*)^2 + \frac{1}{2}\Delta_{\max} n L_g g^* - \frac{1}{4}\Delta_{\max} L_g^2 n \|\mathbf{b}' - \mathbf{a}'\|_1 \leq 0.$$

Solving this quadratic (in)equality for the positive solution leads to

$$\begin{aligned}
g^* &\leq -\frac{1}{2}\Delta_{\max} n L_g + \sqrt{\frac{1}{4}\Delta_{\max}^2 n^2 L_g^2 + \frac{1}{2}\Delta_{\max} L_g^2 n \|\mathbf{b}' - \mathbf{a}'\|_1} \\
&\leq -\frac{1}{2}\Delta_{\max} n L_g + \sqrt{\frac{1}{4}\Delta_{\max}^2 n^2 L_g^2 + \frac{1}{2}\Delta_{\max}^2 n^2 L_g^2} \\
&= -\frac{1}{2}\Delta_{\max} n L_g + \sqrt{\frac{3}{4}\Delta_{\max}^2 n^2 L_g^2} = -\frac{1}{2}\Delta_{\max} n L_g + \frac{\sqrt{3}}{2}\Delta_{\max} n L_g \\
&= \frac{\sqrt{3}-1}{2}\Delta_{\max} n L_g,
\end{aligned}$$

where we used in the second inequality that  $\|\mathbf{b}' - \mathbf{a}'\|_1 \leq n\Delta_{\max}$ . This gives the desired bound.  $\square$

## APPENDIX B. PARABOLIC APPROXIMATIONS – COMPUTATION TIMES &amp; PLOTS

## B.1. Running Times for Computing Parabolic Approximations.

		$\varepsilon$								
		$10^0$	$10^{-1}$	$10^{-2}$	$10^0$	$10^{-1}$	$10^{-2}$	$10^0$	$10^{-1}$	$10^{-2}$
sin	below	1	1	2	1	1	13 015	1	1	4
	above	1	1	15	1	2	13 291	1	3	41
exp	below	1	1	1	1	8	18 652	1	3	6
	above	1	1	2	1	6	20 550	1	4	3
$x^3$	either	1	1	12	1	6 384	33 321	1	4	11

TABLE 5. Computation times for modeling, solving, evaluation in seconds with values below 1 rounded up.

## B.2. Plots for the Parabolic Approximations.




FIGURE 5. Approximation of  $\sin(x)$  for  $x \in [-\pi/2, 3\pi/2]$  from below and above with  $\varepsilon = 10^0, 10^{-1}, 10^{-2}$  and  $\varepsilon = 10^0, 10^{-1}$  by 1, 3, 12 and 1, 5 paraboloids, respectively.




FIGURE 6. Approximation of  $\exp(x)$  for  $x \in [-2, 2]$  from below and above with  $\varepsilon = 10^0, 10^{-1}, 10^{-2}$  by 2, 4, 15 and 2, 5, 20 parabolooids, respectively.




FIGURE 7. Approximation of  $x^3$  for  $x \in [-2, 2]$  from below with  $\varepsilon = 10^0, 10^{-1}, 10^{-2}$  by 3, 10, 79 paraboloids, respectively.