

Explanation of the excesses in associated di-photon production at 152 GeV in 2HDM

Sumit Banik^{1,2,*} and Andreas Crivellin^{1,2,†}

¹*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

²*Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

Statistically significant excesses exist at around 152 GeV in associated di-photon production ($\gamma\gamma + X$) in the sidebands of SM Higgs analyses of ATLAS (using the full run-2 dataset). They are most pronounced in the single- τ , missing-transverse-energy, four-jet and $\geq 1\ell + \geq 1b$ -jet channels ($\approx 3\sigma$) and can be explained by the Drell-Yan production of new Higgs bosons, i.e. $pp \rightarrow W^* \rightarrow H^\pm H^0$. We first examine the excesses in a simplified model approach, considering that H^\pm decays to $\tau\nu$, WZ or tb . Both the $\tau\nu$ and tb decay modes individually lead to a significance of $\lesssim 4\sigma$ while for WZ one can obtain at most 3.5σ . This is because the decays of WZ lead to multiple leptons contributing to the two-lepton channel which does not show an excess at 152 GeV. Next, we consider two-Higgs-doublet models where the charged Higgs does not decay to WZ at tree-level, finding a significance of $\gtrsim 4\sigma$ for a branching ratio of the new neutral Higgs to photons of $\approx 2\%$. Even though this branching fraction is quite sizable, it can be obtained in composite models or via the Lagrangian term $\lambda_6 H_1^\dagger H_1 H_2^\dagger H_1 + \text{h.c.}$ breaking the commonly imposed Z_2 symmetry.

I. INTRODUCTION

Uncovering new (fundamental) particles is one of the primary goals of particle physics. Because the Standard Model (SM) is complete since the Higgs discovery in 2012 [1–3], this would take us into uncharted territory and, hopefully, closer to the ultimate theory describing Nature at the subatomic level. The scalar sector of the SM is particularly promising to contain beyond-the-SM physics because no symmetry or underlying principle enforces its minimality. Furthermore, Higgs bosons naturally couple proportionally to the masses of the SM particles, resulting in LHC constraints which are relatively weak [4, 5] since valence quarks are nearly massless.

Statistically significant indications for a new Higgs with a mass of 152 GeV have emerged [6, 7]¹. They are most pronounced in the sidebands of the di-photon search for the SM Higgs in associated production channels ($\gamma\gamma + X$). More specifically, Ref. [16] studied 22 different signal regions such as missing transverse energy ($X = \text{MET}$), leptons ($X = \ell$) etc., and Ref. [17] contains in addition the single- τ channel. The most significant excesses at ≈ 152 GeV are in the $\text{MET} > 100$ GeV, τ , 4-jet and ℓb (at least one b jet and at least one light lepton) channels with significances of $\approx 3\sigma$. They are consistent with the Drell-Yan production of an $SU(2)_L$ triplet with $Y = 0$ [18], resulting in a combined significance of $\approx 4\sigma$ [19].

However, while a reasonable agreement was obtained for the preferred signal strength among the different signal regions, i.e. a χ^2 of ≈ 24 for 8 channels, which cor-




FIG. 1. Feynman diagram showing the Drell-Yan production process $pp \rightarrow W^* \rightarrow H^\pm H$ with $H \rightarrow \gamma\gamma$ and $H^\pm \rightarrow \tau\nu$, $H^\pm \rightarrow WZ$.

responds to a χ^2 of 3 per degree of freedom (DoF), for the “actual” theory describing the data one expects a χ^2 of 1 per DoF. This motivates searching for a model in which the signal strengths from the different channels agree better, yielding a (even) higher combined significance. Since there is an excess in the single- τ and single-lepton channels but not in di- τ and di-lepton signal regions, this suggests that Drell-Yan $pp \rightarrow W^* \rightarrow HH^\pm$ is the main production mechanism.² However, because in the triplet model the relative signal strengths in the single-light-lepton and MET channels are too big compared to the ℓb and τ channels, one could aim at avoiding the decay of the charged Higgs to WZ . This is naturally achieved within a two-Higgs-doublet model (2HDM) where the charged component does not couple at tree-level to gauge bosons.

In the next section, we will perform a simplified model analysis considering the decay channels $H^\pm \rightarrow \tau\nu$, $H^\pm \rightarrow tb$ and $H^\pm \rightarrow WZ$, separately. We will then study the 2HDM where only the first two channels are

* sumit.banik@psi.ch

† andreas.crivellin@cern.ch

¹ There are also indirect (i.e. non-resonant) hints for new Higgses at the electroweak scale, encoded in the LHC multi-lepton anomalies [8–11], including differential WW [12] and top-quark distributions [13, 14]. They are compatible with the 152 GeV excess and could be related to the 95 GeV excess [15].

² Here H labels any neutral Higgs. In the context of the 2HDM, it could also be the CP -odd component A .




FIG. 2. $\Delta\chi^2 = \chi_{\text{NP}}^2 - \chi_{\text{SM}}^2$ as a function of $\text{BR}(H \rightarrow \gamma\gamma)$ for $m_{H^\pm} = 150$ GeV, normalized to the production cross-section of a doublet Higgs of 92fb. Note that the dependence on the charged Higgs mass is quite weak; it mostly leads to a rescaling of the preferred value for $\text{BR}(H \rightarrow \gamma\gamma)$ due to the varying Drell-Yan production cross-section.

relevant and perform a combined fit assuming that the charged Higgs couplings to fermions are proportional to the fermion masses (i.e. type-I like).

II. SIMPLIFIED MODEL ANALYSIS

As motivated in the introduction, we consider the Drell-Yan production $pp \rightarrow W^* \rightarrow HH^\pm$ with $m_H = 152$ GeV (see Fig. 1). This is the most important production mechanism if the new Higgs transforms non-trivially under $SU(2)_L$, has a small vacuum expectation value, no direct (or tiny) Yukawa couplings and small mixing with the SM Higgs. Furthermore, one can search for this process in associated di-photon production ($\gamma\gamma + X$). This additional requirement significantly reduces the SM background and enhances the new physics sensitivity.

The dominant decay modes of Higgs boson are generally expected to be the heaviest SM particles, albeit suppression by kinematic factors. More specifically, for the charged Higgs we have the decay modes $H^\pm \rightarrow tb$ (including that the top quark can be off-shell) and $H^\pm \rightarrow \tau\nu$ (for electroweak scale masses) if it originates from an $SU(2)_L$

doublet as well as WZ for the triplet case. For these cases the relevant di-photon associated production channels (X) are light leptons ($X = 1\ell$ and $\geq 2\ell$), taus (τ), leptonic top decays ($t_{\text{lep}} = 1\ell + 1b$), ℓb ($\geq 1\ell + \geq 1b$), missing energy ($\text{MET} > 100\text{GeV}$ and $\text{MET} > 200\text{GeV}$) as well as four jets.³ They are shown in Fig. 5 in the appendix.

We simulated the processes $pp \rightarrow W^* \rightarrow (H^\pm \rightarrow \tau\nu, tb, WZ)(H \rightarrow \gamma\gamma)$ using MadGraph5aMC@NLO [20]. The parton showering was performed by Pythia8.3 [21] and the ATLAS detector [16] was modeled with Delphes [22]. The UFO model file at next-to-leading order (NLO) [23] was built using FeynRules [24]. We normalize the production cross section $\sigma(pp \rightarrow W^* \rightarrow HH^\pm)$ to the one

³ The signal region targeting hadronic top decays is not included because ATLAS uses a boosted decision tree (BDT) which targets top-pair production via a tight cut on the BDT score of 0.9. Our signal with a bottom quark and an off-shell top quark is quite different resulting in a very small efficiency. We also used the single lepton category from Ref. [17], rather than from Ref. [16], since the bottom-quark jet veto leads to a nearly uncorrelated data set w.r.t. the ℓb category of Ref. [16].

of an $SU(2)_L$ doublet, with $m_{H^\pm} = 150$ GeV, of 92 fb, obtained by rescaling the cross section calculated with `MadGraph5aMC@NLO` by the NNLL and NLO QCD correction factor of Refs. [25, 26].

For fitting the signal and extracting the significance for the 152 GeV Higgs, we follow the analysis of Ref. [19]. We subtract from the measured ATLAS data the prediction from the SM Higgs signal events as well as the events due to the 152 GeV Higgs and describe the remaining continuous background by the function

$$\left(1 - \frac{m}{\sqrt{s}}\right)^b + (m/\sqrt{s})^{a_0+a_1 \log(m/s)}. \quad (1)$$

The free parameters a_0, a_1 and b are fitted, $\sqrt{s} = 13$ TeV is the LHC run-2 center-of-mass energy, and m the invariant mass of the di-photon pair. We then perform a simultaneous log-likelihood ratio (\mathcal{L}_R) fit using Poisson statistics and convert it to a χ^2 function via $\Delta\chi^2 = -2\ln(\mathcal{L}_R)$.

Starting with the assumption $\text{BR}(H^\pm \rightarrow tb) = 100\%$, one can see from Fig. 2 that the dominant associated production channels are ℓb , t_{lep} , 1ℓ and MET. Since there is no indication for a signal in t_{lep} at 152 GeV, the best-fit value $\text{BR}(H \rightarrow \gamma\gamma) \approx 3\%$ leads to a significance of around 3.8σ . Also, note that $H^\pm \rightarrow tb$ only gives a small effect in the τ channel and thus cannot explain the corresponding excess.

For the case of $\text{BR}(H^\pm \rightarrow \tau\nu) = 100\%$, the 1ℓ and MET channels, receiving contributions from leptonic tau decays, prefer a smaller BR to photons than the single- τ channel. This means that the 1ℓ and MET channels must display an under-fluctuation and/or that the τ channel contains an over-fluctuation. However, the tensions are within the expected range for statistical fluctuations and are included in the combined significance of around 3.8σ at $\text{BR}(H \rightarrow \gamma\gamma) \approx 1\%$. As expected, the decay mode $H^\pm \rightarrow \tau\nu$ cannot address the excesses in the ℓb and $4j$ signal regions.

Finally, the WZ channel results in dominant effects in 1ℓ , 2ℓ (where no excess is seen) and MET channels with a combined significance of 3.5σ at $\text{BR}(H \rightarrow \gamma\gamma) \approx 2.5\%$. However, it has only a small contribution in the single- τ , $4j$ and ℓb channels.

Let us now look at possible combinations of the charged Higgs decay channels. As we already studied the case with all three charged Higgs decay channels in Ref. [19] for the triplet with $Y = 0$, finding a significance of $\approx 4\sigma$, let us therefore focus on combining only $H^\pm \rightarrow \tau\nu$ and $H^\pm \rightarrow t^*b$. They are complementary in their effects in the ℓb and τ signal regions and the dominant modes in 2HDMs with natural flavour conservation [27, 28], MFV [29, 30] or alignment [31, 32]. Therefore, we consider the $\text{BR}(H^\pm \rightarrow tb)$ – $\text{BR}(H^0 \rightarrow \gamma\gamma)$ plane in Fig. 3. This means we assume that tb and $\tau\nu$ are the dominant decay modes of the charged Higgs such that $\text{BR}(H^\pm \rightarrow tb) + \text{BR}(H^\pm \rightarrow \tau\nu) \approx 1$. Note that $H^\pm \rightarrow cs$ does not lead to a relevant effect in any signal region, such that its presence can be included by rescaling $\text{BR}(H^0 \rightarrow \gamma\gamma)$. For $m_{H^\pm} = 130$ GeV, $m_{H^\pm} = 150$ GeV

and $m_{H^\pm} = 170$ GeV, one can see that with an increasing mass, the $-\Delta\chi^2$ maximum slightly increases. The reason for this is the relative efficiency in the ℓb channel (where an excess is seen) compared to the t_{lep} channel (where no indication for an excess at 152 GeV is seen) is enhanced. Furthermore, with an increasing charged Higgs mass a higher branching ratio to photons is preferred, mainly to compensate for the decreasing Drell-Yan production cross-section. The blue horizontal bands, whose widths are due to the sizable uncertainty of the decay width to off-shell top quarks [33], show the one-dimensional case in which the charged Higgs couplings are proportional to the masses of the involved fermions, as in the case of the type-I 2HDM. In this setup, a combined significance of $> 4\sigma$ can be obtained for $m_{H^\pm} = 150$ GeV.

III. 2HDM PHENOMENOLOGY

So far, we have studied a simplified model, considering $pp \rightarrow W^* \rightarrow HH^\pm$ with $\text{BR}(H \rightarrow \gamma\gamma)$ being a free parameter. Let us now examine the excesses at 152 GeV within 2HDMs [34, 35] where, after spontaneous symmetry breaking, an additional CP -odd Higgs boson (A) emerges (for more details and our conventions see e.g. Ref. [36]). To study the effect of $pp \rightarrow Z^* \rightarrow HA$, which we disregarded so far, we consider the benchmark point $m_{H^\pm} = 130$ GeV and $m_A = 200$ GeV with small Yukawa couplings of A to fermions such that $\text{BR}(A \rightarrow WH^\pm) \approx 100\%$. The resulting impact can be seen by comparing Fig. 4 (left) to Fig. 3 (left). The $-\Delta\chi^2$ slightly decreases since $pp \rightarrow Z^* \rightarrow HA$ with $A \rightarrow W(H^\pm \rightarrow tb)$ has a higher relative efficiency in ℓb vs t_{lep} signal region than $pp \rightarrow Z^* \rightarrow H(H^\pm \rightarrow tb)$.

A branching ratio of H to photons at the percent level is needed to explain the excesses. This cannot be achieved in plain Z_2 -symmetric 2HDM with natural flavour conservation, since decays to gauge bosons and fermions are correlated to photon decays, because it is induced by loops involving them. To circumvent this, one could add the effective operator

$$F^{\mu\nu} F_{\mu\nu} \phi_1^\dagger \phi_2 + \text{h.c.}, \quad (2)$$

which dominantly contributes to the decays of the new Higgs bosons for $\tan\beta = v_2/v_1 \gg 1$. This corresponding Wilson coefficient is generated in composite Higgs models which can naturally lead to large decay widths of the new Higgses to photon [37–40] due to the new (excited) states in the loop. In this case, fermion loops are in general dominant and one predicts $\Gamma(A \rightarrow \gamma\gamma) = \frac{9}{4}\Gamma(H \rightarrow \gamma\gamma)$ for $m_A = m_H$.

Alternatively, one can give up on the usually employed Z_2 symmetry and include in the Lagrangian

$$\mathcal{L} \in -\lambda_6 H_1^\dagger H_1 H_2^\dagger H_2 + \text{h.c.}, \quad (3)$$

This term leads to a modification of the width of $H \rightarrow \gamma\gamma$




FIG. 3. Contours of $-\Delta\chi^2 = \chi_{\text{NP}}^2 - \chi_{\text{SM}}^2$ for $m_H = 152$ GeV and $m_{H^\pm} = 130$ GeV, 150 GeV and 170 GeV assuming that $\tau\nu$ and tb are the dominant decay modes of the charged Higgs. Because the decay to cs , which can be numerically sizable, has a very small impact on all signal regions, its effect can be reabsorbed into a rescaling of $\text{BR}(H \rightarrow \gamma\gamma)$.

via the charged Higgs loop with the amplitude

$$\mathcal{M}(H \rightarrow \gamma\gamma) \propto \frac{\alpha_{\text{em}} g_2 v^2 \lambda_6}{2m_{H^\pm}^2} \beta_H^0 \left(\frac{4m_{H^\pm}^2}{m_H^2} \right), \quad (4)$$

$$\beta_H^0(x) = -x[1 - xf(x)], \quad f(x) = (\sin^{-1}(1/\sqrt{x}))^2$$

for $x \geq 1$ [41], in the limit of $\tan\beta \rightarrow \infty$, but not to the SM Higgs.

We illustrate this in Fig. 4 (right) where the preferred regions in the α - λ_6 plane for $\tan\beta = 20$ and $m_{12}^2 = 1100\text{GeV}^2$ in the type-I 2HDM are shown.⁴ One can see that it is possible to obtain the preferred size of the branching ratio to photons for order-one values of λ_6 . The resulting $\Delta\chi^2$ minimum is ≈ -23 (corresponding to 4.4σ for our 2-dimensional hypothesis), including the contribution to χ^2 from the constraints of the inclusive $\gamma\gamma$ channel [6].

IV. CONCLUSIONS AND OUTLOOK

We study possible simplified-model explanations of the excesses at 152 GeV in associated di-photon production via the Drell-Yan process $pp \rightarrow W^* \rightarrow HH^\pm$ for the three decay modes $H^\pm \rightarrow \tau\nu$, $H^\pm \rightarrow tb$ and $H^\pm \rightarrow WZ$. The $H^\pm \rightarrow \tau\nu$ decay channel predicts interesting correlations between the single- τ , single-lepton and MET

signal regions. Furthermore, its signatures in the signal regions which show excesses, are complementary to the ones of the $H^\pm \rightarrow tb$ decay, which can address the lb channel. On the other hand, the $H^\pm \rightarrow WZ$ decay in general leads to too many leptons because no excess in the two-lepton signal region is observed.

Therefore, we considered in a second step 2HDMs where the decay $H^\pm \rightarrow WZ$ is not allowed at tree-level. We find that for $m_{H^\pm} \approx 150$ GeV, assuming it has type-I-like BRs, one can obtain a combined significance of 4.3σ , which even slightly increases if $pp \rightarrow Z^* \rightarrow HA$ with $A \rightarrow WH^\pm$ is included. Because a large $\text{BR}(H \rightarrow \gamma\gamma)$ at the percent level is required, this suggests supplementing the 2HDM with higher dimensional operators which could be realized within a composite setup. Alternatively, the Lagrangian term $\lambda_6 H_1^\dagger H_1 H_2^\dagger H_1$ can lead to sizable BR in the alignment limit ($\alpha \rightarrow 0$, $\beta \rightarrow \pi/2$) and, including the inclusive di-photon channel, result in a significance of 4.4σ . While we considered $m_H = 152$ GeV, also A could be the 152 GeV boson. In this case, an imaginary part of λ_6 would be needed to modify $\text{BR}(A \rightarrow \gamma\gamma)$, giving at the same time rise to interesting correlated effects in electric dipole moments.

ACKNOWLEDGMENTS

We thank Sezen Sekmen for useful discussions and Saiyad Ashanujjaman, Guglielmo Coloretti, Siddharth P. Maharathy and Bruce Mellado for agreeing to reuse some results of Ref. [19]. This work is supported by a professorship grant from the Swiss National Science Foundation (No. PP00P21.76884).

⁴ Note that without the commonly used Z_2 symmetry one cannot impose natural flavour conservation, but flavour-changing neutral currents can be avoided via MFV [29, 30] or alignment [31, 32]. Furthermore, in the aligned 2HDM one can effectively recover the type-I Yukawa structure in the limit of large couplings ξ_f (see Refs. [31, 32]). We checked with ScannerS [42, 43], based on HDECAY [44, 45] and EVADE [46], that vacuum stability and perturbativity that $A \rightarrow WH$ is, in fact, the dominant decay mode for our benchmark point.




FIG. 4. Left: Same as Fig. 3 but including $pp \rightarrow Z^* \rightarrow AH$ for $m_A = 200$ GeV and $\text{BR}(A \rightarrow WH^\pm \approx 100\%)$. Right: Preferred 1σ and 2σ regions in the α - λ_6 plane, assuming λ_6 to be real for $\tan(\beta) = 20$. The minimal $\Delta\chi^2$ is ≈ -23 corresponding to 4.4σ for this two-dimensional hypothesis.

Appendix A: Fit to the individual signal regions

We show the eight relevant signal regions of the ATLAS analyses [16, 17] in Fig. 5 with the individual best fits for a 152 GeV Higgs signals [19].

REFERENCES

- [1] ATLAS Collaboration, G. Aad et al., “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett. B* **716** (2012) 1–29, [arXiv:1207.7214 \[hep-ex\]](#).
- [2] CMS Collaboration, S. Chatrchyan et al., “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC,” *Phys. Lett. B* **716** (2012) 30–61, [arXiv:1207.7235 \[hep-ex\]](#).
- [3] CDF, D0 Collaboration, T. Aaltonen et al., “Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson searches at the Tevatron,” *Phys. Rev. Lett.* **109** (2012) 071804, [arXiv:1207.6436 \[hep-ex\]](#).
- [4] ATLAS Collaboration, G. Aad et al., “ATLAS searches for additional scalars and exotic Higgs boson decays with the LHC Run 2 dataset,” [arXiv:2405.04914 \[hep-ex\]](#).
- [5] CMS Collaboration, A. Hayrapetyan et al., “Searches for Higgs Boson Production through Decays of Heavy Resonances,” [arXiv:2403.16926 \[hep-ex\]](#).
- [6] A. Crivellin, Y. Fang, O. Fischer, S. Bhattacharya, M. Kumar, E. Malwa, B. Mellado, N. Rapheeha, X. Ruan, and Q. Sha, “Accumulating evidence for the associated production of a new Higgs boson at the LHC,” *Phys. Rev. D* **108** no. 11, (2023) 115031, [arXiv:2109.02650 \[hep-ph\]](#).
- [7] S. Bhattacharya, G. Coloretti, A. Crivellin, S.-E. Dahbi, Y. Fang, M. Kumar, and B. Mellado, “Growing Excesses of New Scalars at the Electroweak Scale,” [arXiv:2306.17209 \[hep-ph\]](#).
- [8] S. von Buddenbrock, N. Chakrabarty, A. S. Cornell, D. Kar, M. Kumar, T. Mandal, B. Mellado, B. Mukhopadhyaya, R. G. Reed, and X. Ruan, “Phenomenological signatures of additional scalar bosons at the LHC,” *Eur. Phys. J. C* **76** no. 10, (2016) 580, [arXiv:1606.01674 \[hep-ph\]](#).
- [9] S. von Buddenbrock, A. S. Cornell, A. Fadol, M. Kumar, B. Mellado, and X. Ruan, “Multi-lepton signatures of additional scalar bosons beyond the Standard Model at the LHC,” *J. Phys. G* **45** no. 11, (2018) 115003, [arXiv:1711.07874 \[hep-ph\]](#).
- [10] S. Buddenbrock, A. S. Cornell, Y. Fang, A. Fadol Mohammed, M. Kumar, B. Mellado, and K. G. Tomiwa, “The emergence of multi-lepton anomalies at the LHC and their compatibility with new physics at the EW scale,” *JHEP* **10** (2019) 157, [arXiv:1901.05300 \[hep-ph\]](#).
- [11] O. Fischer et al., “Unveiling hidden physics at the LHC,” *Eur. Phys. J. C* **82** no. 8, (2022) 665, [arXiv:2109.06065 \[hep-ph\]](#).
- [12] G. Coloretti, A. Crivellin, S. Bhattacharya, and B. Mellado, “Searching for low-mass resonances decaying into W bosons,” *Phys. Rev. D* **108** no. 3, (2023) 035026, [arXiv:2302.07276 \[hep-ph\]](#).
- [13] ATLAS Collaboration, G. Aad et al., “Inclusive and differential cross-sections for dilepton $t\bar{t}$ production measured in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *JHEP* **07** (2023) 141, [arXiv:2303.15340 \[hep-ex\]](#).




FIG. 5. Di-photon invariant mass distributions for eight relevant signal regions from Ref. [19]. The data (black) is shown together with the continuum background (blue) given in the ATLAS analyses and the total number events (red). The latter contains the refitted background (not shown for brevity), the predicted SM Higgs signals at 125 GeV (magenta) and the individual best fits for a 152 GeV signal (green).

- [14] S. Banik, G. Coloretti, A. Crivellin, and B. Mellado, “Uncovering New Higgses in the LHC Analyses of Differential $t\bar{t}$ Cross Sections,” [arXiv:2308.07953 \[hep-ph\]](#).
- [15] G. Coloretti, A. Crivellin, and B. Mellado, “Combined Explanation of LHC Multi-Lepton, Di-Photon and Top-Quark Excesses,” [arXiv:2312.17314 \[hep-ph\]](#).
- [16] **ATLAS** Collaboration, G. Aad et al., “Model-independent search for the presence of new physics in events including $H \rightarrow \gamma\gamma$ with $\sqrt{s} = 13$ TeV pp data recorded by the ATLAS detector at the LHC,” *JHEP* **07** (2023) 176, [arXiv:2301.10486 \[hep-ex\]](#).
- [17] **ATLAS** Collaboration, “Search for non-resonant Higgs boson pair production in final states with leptons, taus and photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,”.
- [18] S. Ashanujjaman, S. Banik, G. Coloretti, A. Crivellin, S. P. Maharathy, and B. Mellado, “Explaining the $\gamma\gamma + X$ Excesses at ≈ 151.5 GeV via the Drell-Yan Production of a Higgs Triplet,” [arXiv:2402.00101 \[hep-ph\]](#).
- [19] A. Crivellin, S. Ashanujjaman, S. Banik, G. Coloretti, S. P. Maharathy, and B. Mellado, “Growing Evidence for a Higgs Triplet,” [arXiv:2404.14492 \[hep-ph\]](#).
- [20] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP* **07** (2014) 079, [arXiv:1405.0301](#)

- [hep-ph].
- [21] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, “An introduction to PYTHIA 8.2” *Comput. Phys. Commun.* **191** (2015) 159–177, [arXiv:1410.3012](#) [hep-ph].
- [22] **DELPHES 3** Collaboration, J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi, “DELPHES 3, A modular framework for fast simulation of a generic collider experiment,” *JHEP* **02** (2014) 057, [arXiv:1307.6346](#) [hep-ex].
- [23] Y. Wu, “Ufo model file for cp conserving 2hdm,” Aug., 2023. <https://doi.org/10.5281/zenodo.8207058>.
- [24] C. Degrande, “Automatic evaluation of UV and R2 terms for beyond the Standard Model Lagrangians: a proof-of-principle,” *Comput. Phys. Commun.* **197** (2015) 239–262, [arXiv:1406.3030](#) [hep-ph].
- [25] R. Ruiz, “QCD Corrections to Pair Production of Type III Seesaw Leptons at Hadron Colliders,” *JHEP* **12** (2015) 165, [arXiv:1509.05416](#) [hep-ph].
- [26] A. A. H., B. Fuks, H.-S. Shao, and Y. Simon, “Precision predictions for exotic lepton production at the Large Hadron Collider,” *Phys. Rev. D* **107** no. 7, (2023) 075011, [arXiv:2301.03640](#) [hep-ph].
- [27] S. L. Glashow and S. Weinberg, “Natural Conservation Laws for Neutral Currents,” *Phys. Rev. D* **15** (1977) 1958.
- [28] E. A. Paschos, “Diagonal Neutral Currents,” *Phys. Rev. D* **15** (1977) 1966.
- [29] G. D’Ambrosio, G. F. Giudice, G. Isidori, and A. Strumia, “Minimal flavor violation: An Effective field theory approach,” *Nucl. Phys. B* **645** (2002) 155–187, [arXiv:hep-ph/0207036](#).
- [30] F. J. Botella, G. C. Branco, and M. N. Rebelo, “Minimal Flavour Violation and Multi-Higgs Models,” *Phys. Lett. B* **687** (2010) 194–200, [arXiv:0911.1753](#) [hep-ph].
- [31] A. Pich and P. Tuzon, “Yukawa Alignment in the Two-Higgs-Doublet Model,” *Phys. Rev. D* **80** (2009) 091702, [arXiv:0908.1554](#) [hep-ph].
- [32] P. Tuzon and A. Pich, “The Aligned two-Higgs Doublet model,” *Acta Phys. Polon. Supp.* **3** (2010) 215–220, [arXiv:1001.0293](#) [hep-ph].
- [33] **LHC Higgs Cross Section Working Group** Collaboration, D. de Florian et al., “Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector,” [arXiv:1610.07922](#) [hep-ph].
- [34] T. D. Lee, “A Theory of Spontaneous T Violation,” *Phys. Rev. D* **8** (1973) 1226–1239.
- [35] H. E. Haber and G. L. Kane, “The Search for Supersymmetry: Probing Physics Beyond the Standard Model,” *Phys. Rept.* **117** (1985) 75–263.
- [36] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, “Theory and phenomenology of two-Higgs-doublet models,” *Phys. Rept.* **516** (2012) 1–102, [arXiv:1106.0034](#) [hep-ph].
- [37] G. Cacciapaglia, A. Deandrea, A. M. Iyer, and K. Sridhar, “Tera-Z stage at future colliders and light composite axionlike particles,” *Phys. Rev. D* **105** no. 1, (2022) 015020, [arXiv:2104.11064](#) [hep-ph].
- [38] R. Benbrik et al., “Signatures of vector-like top partners decaying into new neutral scalar or pseudoscalar bosons,” *JHEP* **05** (2020) 028, [arXiv:1907.05929](#) [hep-ph].
- [39] G. Cacciapaglia, G. Ferretti, T. Flacke, and H. Serôdio, “Light scalars in composite Higgs models,” *Front. in Phys.* **7** (2019) 22, [arXiv:1902.06890](#) [hep-ph].
- [40] A. Belyaev, G. Cacciapaglia, H. Cai, G. Ferretti, T. Flacke, A. Parolini, and H. Serodio, “Di-boson signatures as Standard Candles for Partial Compositeness,” *JHEP* **01** (2017) 094, [arXiv:1610.06591](#) [hep-ph]. [Erratum: *JHEP* **12**, 088 (2017)].
- [41] M. Spira, “Higgs Boson Production and Decay at Hadron Colliders,” *Prog. Part. Nucl. Phys.* **95** (2017) 98–159, [arXiv:1612.07651](#) [hep-ph].
- [42] R. Coimbra, M. O. P. Sampaio, and R. Santos, “ScannerS: Constraining the phase diagram of a complex scalar singlet at the LHC,” *Eur. Phys. J. C* **73** (2013) 2428, [arXiv:1301.2599](#) [hep-ph].
- [43] M. Mühlleitner, M. O. P. Sampaio, R. Santos, and J. Wittbrodt, “ScannerS: parameter scans in extended scalar sectors,” *Eur. Phys. J. C* **82** no. 3, (2022) 198, [arXiv:2007.02985](#) [hep-ph].
- [44] A. Djouadi, J. Kalinowski, and M. Spira, “HDECAY: A Program for Higgs boson decays in the standard model and its supersymmetric extension,” *Comput. Phys. Commun.* **108** (1998) 56–74, [arXiv:hep-ph/9704448](#).
- [45] **HDECAY** Collaboration, A. Djouadi, J. Kalinowski, M. Muehlleitner, and M. Spira, “HDECAY: Twenty++ years after,” *Comput. Phys. Commun.* **238** (2019) 214–231, [arXiv:1801.09506](#) [hep-ph].
- [46] W. G. Hollik, G. Weiglein, and J. Wittbrodt, “Impact of Vacuum Stability Constraints on the Phenomenology of Supersymmetric Models,” *JHEP* **03** (2019) 109, [arXiv:1812.04644](#) [hep-ph].