

# D-instanton Induced Effective Action and its Gauge Invariance

Ashoke Sen

*International Centre for Theoretical Sciences - TIFR  
Bengaluru - 560089, India*

E-mail: ashoke.sen@icts.res.in

## Abstract

The effect of D-instantons on closed string scattering amplitudes may be encoded into an effective action obtained by integrating out the (transient) open string fields. In order that this respects the gauge invariance of the theory, the sum of the perturbative closed string field theory action and the D-instanton induced effective action must satisfy the quantum Batalin-Vilkovisky master equation. In a previous paper this was proved for the effective action induced by up to two instantons. We generalize the proof for arbitrary number of D-instantons.

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Review and the statement of the problem</b>	<b>3</b>
<b>3</b>	<b>Multi-instanton contribution to the effective action and its gauge invariance</b>	<b>8</b>
<b>4</b>	<b>Non-identical D-instantons</b>	<b>10</b>

## 1 Introduction

The action of closed string field theory satisfies the quantum Batalin-Vilkovisky (BV) master equation [1, 2] encoding the gauge invariance of the theory [3, 4]. However besides the perturbative contribution to the action, closed string field theory also receives non-perturbative contributions to the action from D-instantons. A systematic algorithm for computing D-instanton contribution to the closed string field theory action was described in [5], but the explicit construction of the action was carried out only up to two instanton contribution. It was also checked that the action, so constructed, satisfies the BV master equation up to corrections involving three instanton terms.

Our goal in this paper will be to construct the action of closed string field theory after taking into account the effect of arbitrary number of instantons. We also show that the action so constructed satisfies the quantum BV master equation to all orders in the instanton expansion. Our main result, describing the instanton induced effective action due to arbitrary number of identical D-instantons, is given in (3.1). Its generalization to arbitrary numbers of non-identical D-instantons, is given in (4.3).

One immediate application of our results is the proof of gauge invariance of the closed string amplitudes even after the inclusion of D-instanton effects. Since for computing D-instanton contribution to closed string amplitudes we need to modify the world-sheet rules, separating out the zero mode integrals and doing these integrals at the end, the usual perturbative proof of gauge invariance showing the decoupling of BRST trivial states no longer holds. However the fact that the closed string effective action satisfies the quantum BV master equation immediately implies the gauge invariance of on-shell closed string amplitudes.

## 2 Review and the statement of the problem

In this section we shall review the relevant results of [5] using a somewhat different notation and paying special attention to the zero mode integrals.

Let us consider closed strings interacting with some fixed number of D-branes. Even though our eventual goal is to study D-instantons, let us first consider the case where the D-branes extend along more than two non-compact space-time directions so that they can be regarded as fixed background. The observables in this system are scattering amplitudes involving external open and closed strings and these amplitudes can be reproduced by a gauge invariant field theory of open and closed strings [6]. The interaction vertices of this field theory are expressed as integrals over appropriate subspaces of the moduli spaces of punctured, *connected* Riemann surfaces, with the integrand given in terms of correlation functions of off-shell open and closed strings inserted at the punctures, together with some additional insertions of ghost and / or picture changing operators. The subspaces of the moduli space on which we carry out the integration avoid all degenerate Riemann surfaces and hence the integrals do not suffer from any divergences. More details have been reviewed in [4].

There is a standard procedure, also reviewed in [4], by which we can integrate out a set of degrees of freedom of the theory and write down an expression for the effective action involving the rest of the degrees of freedom [7–9]. In particular, if we are interested in amplitudes involving only the external closed strings, we can integrate out the open strings and write down an effective action for closed string fields only. The interaction vertices of the resulting effective closed string field theory are still given by integrals over appropriate subspaces of the moduli spaces of punctured connected Riemann surfaces. The difference is that now the integrand has correlation functions of external closed strings only and the integration region now includes degenerate Riemann surfaces involving open string degeneration but still avoids degenerate Riemann surfaces involving closed string degeneration. Since the integrals may get divergent contributions from the region of the moduli space where there is open string degeneration, the definition of the effective action given above is somewhat formal. For D-instantons that will be of interest to us, the solution to this problem has been well understood and reviewed in [4]. We shall discuss some consequences of this when we turn to D-instantons.

Let  $S_{(r)}$  denote the total closed string field theory effective action for  $r$  identical D-branes.<sup>1</sup>  $S_{(r)}$  includes the contribution from *connected* Riemann surfaces without boundaries as well as

---

<sup>1</sup>Here identical means D-branes that differ from each other at most by their open string moduli *e.g.* the position of the D-brane in transverse directions. They have the same tension.

*connected* Riemann surfaces with boundaries lying on any of the  $r$  D-branes.  $S_{(0)}$  will denote the closed string field theory action without any D-brane in the background. Then  $S_{(r)}$  will satisfy the quantum BV master equation [4]

$$\frac{1}{2} \{S_{(r)}, S_{(r)}\} + \Delta S_{(r)} = 0, \quad \text{for } r \geq 0, \quad (2.1)$$

where for a pair of functions  $F$  and  $G$  of closed string fields  $\{\psi^\alpha\}$  and their anti-fields  $\{\psi_\alpha^*\}$ , we have

$$\{F, G\} = \frac{\partial_R F}{\partial \psi^\alpha} \frac{\partial_L G}{\partial \psi_\alpha^*} - \frac{\partial_R F}{\partial \psi_\alpha^*} \frac{\partial_L G}{\partial \psi^\alpha}, \quad \Delta F = (-1)^{\psi_\alpha} \frac{\partial_L}{\partial \psi^\alpha} \left( \frac{\partial_L F}{\partial \psi_\alpha^*} \right), \quad (2.2)$$

where  $\partial_L$  and  $\partial_R$  denotes left and right derivatives respectively and  $(-1)^{\psi_\alpha}$  takes value 1 ( $-1$ ) for Grassmann even (odd)  $\psi_\alpha$ . For even functions  $F$ , we have the useful property,

$$\Delta e^F = \frac{1}{2} \{F, F\} e^F + \Delta F e^F. \quad (2.3)$$

Using this, (2.1) may be written as

$$\Delta e^{S_{(r)}} = 0. \quad (2.4)$$

We shall see later that for D-instantons the definition of  $S_{(r)}$  given above requires some modification, so it will be useful to give a formal definition of  $S_{(r)}$  that is valid in all cases. Let  $\tilde{S}^{(r)}$  denote the full BV master action of open-closed string field theory in the presence of  $r$  D-branes before we integrate out any fields. Then  $e^{S_{(r)}}$  is defined as the integral of  $e^{\tilde{S}^{(r)}}$  over half of the open string fields, the other half being fixed by ‘gauge choice’.  $e^{\tilde{S}^{(r)}}$  satisfies the BV master equation  $(\Delta + \Delta_{open})e^{\tilde{S}^{(r)}} = 0$  [4,6]. Integrating this over half of the open string fields, we see that the term involving  $\Delta_{open}e^{\tilde{S}^{(r)}}$  integrates to zero since it is a total derivative, and we are led to (2.4). (2.1) then follows as a consequence of (2.3) after defining  $S_{(r)}$  as  $\ln e^{S_{(r)}}$ . If we construct  $S_{(r)}$  using perturbation theory, then by the standard rules  $S_{(r)}$  will be given by the sum of connected Feynman diagrams, with the fields that are integrated out appearing in the propagators. In string theory in the presence of D-branes, the sum over connected Feynman diagrams translates to sum over connected world-sheets with boundaries, leading to the description of  $S_{(r)}$  given earlier.

We shall now introduce some new definitions. We shall denote by  $\widehat{S}_i$  the contribution to the closed string effective action from the sum of all *connected* world-sheets with all boundaries on the  $i$ -th D-brane.  $\widehat{S}_{ij}$  will denote the contribution from the sum of all connected world-sheets with each boundary on either the  $i$ -th D-brane or the  $j$ -th D-brane, at least one boundary on the  $i$ -th D-brane and at least one boundary on the  $j$ -th D-brane. Similar definitions may be

given for  $\widehat{S}_{i_1 \dots i_k}$  when  $k$  or larger number of D-branes are present. It follows from the definition that  $\widehat{S}_{i_1 \dots i_k}$  is symmetric under the exchange of the indices. Then we have, for example,

$$S_{(1)} = S_{(0)} + \widehat{S}_1, \quad S_{(2)} = S_{(0)} + \widehat{S}_1 + \widehat{S}_2 + \widehat{S}_{12}, \quad S_{(3)} = S_{(0)} + \widehat{S}_1 + \widehat{S}_2 + \widehat{S}_3 + \widehat{S}_{12} + \widehat{S}_{23} + \widehat{S}_{13} + \widehat{S}_{123}, \quad (2.5)$$

etc.

Let us now turn to the case of D-instantons. As usual, we shall use a general definition of D-instantons – D-branes that satisfy Dirichlet boundary condition for all *non-compact* space-time directions. In this case the construction of the closed string effective action requires some modification. There are several reasons for this that we enumerate below.

1. Unlike in the case of extended D-branes where the contribution to the closed string effective action comes from a sum over connected world-sheets with boundaries, for computing D-instanton contribution to the closed string effective action, we have to include contribution from *disconnected* world-sheets [10]. This is related to the infrared divergences associated with open string degeneration that was alluded to earlier. The problem comes from integration over the open string zero modes, including the D-instanton position. The presence of these zero modes makes the building blocks  $\widehat{S}_{i_1 \dots i_k}$  ill defined. The remedy is to remove the contribution of these zero modes from the internal open string propagators (and include a contribution from an out of Siegel gauge mode propagator [11]), leading to modified rules for world-sheet computation. This gives finite  $\widehat{S}_{i_1 \dots i_k}$ . We shall denote this by  $\widehat{S}'_{i_1 \dots i_k}$  to indicate that the rules for world-sheet computation have been modified. However since the construction of the closed string effective action requires integrating out all the open string modes, the zero mode integrals must be performed at the end. As a result, disconnected world-sheets that end on a common D-instanton will have their zero mode integrals performed together and will contribute to the closed string effective action. We shall call them *target space connected contributions*. Since the overall energy momentum conservation is implemented only after integration over the zero modes, a pair of disconnected world-sheets that are target space connected will have one overall energy-momentum conserving delta function instead of separate energy-momentum conservation associated with individual world-sheets. For example the product of  $\widehat{S}'_{12}$  and  $\widehat{S}'_{23}$  is target space connected since both end on the D-instanton 2 at one of their boundaries. Hence its contribution is included in the closed string effective action  $S$  after performing integration over the zero modes of instantons 1 and 2. But not all disconnected world-sheets are allowed, *e.g.* the product of  $\widehat{S}'_{12}$  and  $\widehat{S}'_{34}$  is not target space connected and

does not contribute to the four instanton effective action.<sup>2</sup> However if we multiply this by  $\widehat{S}'_{13}$  then the world-sheets are connected in the target space and this product becomes an allowed four instanton contribution to the effective action after we integrate over the zero modes of the instantons 1,2,3 and 4.

2. Since instantons have finite action, they do not represent fixed backgrounds but correspond to saddle points of the path integral that contribute to the closed string amplitudes in the vacuum. Hence we must sum over the contributions from different number of instantons. An  $r$  instanton contribution to the effective action is multiplied by a factor of  $\mathcal{N}^r$  where  $\mathcal{N} = e^{-\mathcal{T}}$ ,  $\mathcal{T}$  being the D-instanton tension.<sup>3</sup>
3. Since the path integral over the open string fields on the D-instanton includes integration over the D-instanton positions and other moduli, after the integration different D-instantons become indistinguishable. This gives rise to an extra symmetry factor  $1/r!$  for an  $r$ -instanton contribution.

With these rules the contribution to the closed string effective action up to two instanton sector can be written as [4,7]

$$S \equiv S_{(0)} + \mathcal{N} \int_z \left( e^{\widehat{S}'_1} - 1 \right) + \frac{1}{2} \mathcal{N}^2 \int_z \left( e^{\widehat{S}'_{12} + \widehat{S}'_1 + \widehat{S}'_2} - 1 - e^{\widehat{S}'_1 + \widehat{S}'_2} + 1 \right) + \mathcal{O}(\mathcal{N}^3), \quad (2.6)$$

where  $\int_z$  signs on the right hand side of (2.6) denote integration over the zero modes of the instantons. In order to determine what zero modes we should integrate over, we should expand the exponentials in power series in  $\widehat{S}'_{i_1 \dots i_k}$  and in each term carry out the integration over the zero modes of the instantons that appear in the product. For example for  $\widehat{S}'_{12}$  we integrate over the zero modes of instantons 1 and 2 while for  $\widehat{S}'_{12} \widehat{S}'_{23}$  we integrate over the zero modes of the instantons 1, 2 and 3. In (2.6)  $S_{(0)}$  is the perturbative contribution to the effective action and the term proportional to  $\mathcal{N}^r$  represents the  $r$  instanton contribution to the effective action.  $e^{\widehat{S}'_1} - 1$ , expanded in a power series in  $\widehat{S}'_1$ , generates connected and disconnected world-sheets where each connected component has at least one boundary and all the boundaries end on a single D-instanton. All of these diagrams are connected in the target space and represent one instanton contribution to the effective action. This explains why they are multiplied by  $\mathcal{N}$ . To see how the coefficient of  $\mathcal{N}^2$  arises, we note that the  $e^{\widehat{S}'_{12} + \widehat{S}'_1 + \widehat{S}'_2} - 1$  factor, expanded in a

<sup>2</sup>Instead, it should be viewed as a contribution to the square of the two instanton effective action.

<sup>3</sup>In [4,7] we included also the exponential of the annulus amplitude in the definition of  $\mathcal{N}$  but here we shall include this in the action.

power series in  $\widehat{S}'_{12}$ ,  $\widehat{S}'_1$  and  $\widehat{S}'_2$ , generates connected and disconnected world-sheets, where each connected component has at least one boundary lying on a D-instanton. As long as we have terms with at least one factor of  $\widehat{S}'_{12}$ , the diagram is connected in the target space and should be included in the closed string effective action. However the terms without any factor of  $\widehat{S}'_{12}$  are either disconnected in the target space or has all its boundaries lying on one instanton. These contributions are not genuine two instanton contribution to the effective action and must be removed. The subtraction term  $e^{\widehat{S}'_1 + \widehat{S}'_2} - 1$  precisely achieves this.

We shall now express (2.6) in terms of the  $S_{(r)}$ 's whose general definition was given below (2.4). For D-instantons we cannot use (2.5) since the contribution from the open string zero modes are removed from the definition of  $\widehat{S}'_{i_1 \dots i_k}$ . Instead, we can express  $S_{(r)}$  via,

$$e^{S_{(1)}} = \int_z e^{S_{(0)}} e^{\widehat{S}'_1}, \quad e^{S_{(2)}} = \int_z e^{S_{(0)}} e^{\widehat{S}'_1 + \widehat{S}'_2 + \widehat{S}'_{12}}, \quad e^{S_{(3)}} = \int_z e^{S_{(0)}} e^{\widehat{S}'_1 + \widehat{S}'_2 + \widehat{S}'_3 + \widehat{S}'_{12} + \widehat{S}'_{23} + \widehat{S}'_{13} + \widehat{S}'_{123}} \quad (2.7)$$

etc. Due to the argument given below (2.4),  $S_{(r)}$  satisfies the BV master equation (2.4), (2.1). Using (2.7), we can express (2.6) as

$$S = S_{(0)} + \mathcal{N} (e^{S_{(1)} - S_{(0)}} - 1) + \frac{1}{2} \mathcal{N}^2 (e^{S_{(2)} - S_{(0)}} - e^{2(S_{(1)} - S_{(0)})}) + \mathcal{O}(\mathcal{N}^3), \quad (2.8)$$

where we used the fact that after carrying out the zero mode integrals,  $e^{\widehat{S}'_1}$  and  $e^{\widehat{S}'_2}$  become identical to  $e^{S_{(1)} - S_{(0)}}$ .

For consistency,  $S$  given in (2.8) should satisfy the BV master equation. To prove this, we use (2.1) with  $r = 0, 1, 2$  and (2.3) to get [4, 7]

$$\frac{1}{2} \{S, S\} + \Delta S = \mathcal{O}(\mathcal{N}^3), \quad (2.9)$$

as required.

We are now ready to state the problem. This can be divided into two parts:

1. We need to find the  $r$  instanton contribution to  $S$ , i.e. find the coefficient of the  $\mathcal{N}^r$  term in (2.8). This requires us to generalize (2.6) or (2.8) for larger number of instantons. The subtraction terms now take more complicated form, *e.g.* for three instantons we need to keep terms that contain at least one factor of  $\widehat{S}'_{123}$ , but we need to also keep terms that contain  $\widehat{S}'_{12} \widehat{S}'_{23}$  or  $\widehat{S}'_{12} \widehat{S}'_{13}$  or  $\widehat{S}'_{13} \widehat{S}'_{23}$  as a factor.
2. Once we have found  $S$ , we need to prove that the left hand side of (2.9) vanishes exactly.

### 3 Multi-instanton contribution to the effective action and its gauge invariance

In this section we shall answer the questions posed at the end of the last section. We claim that the full action takes the form

$$S = S_{(0)} + \ln \left( 1 + \sum_{r=1}^{\infty} \frac{1}{r!} \mathcal{N}^r e^{S_{(r)} - S_{(0)}} \right) - \mathcal{N}. \quad (3.1)$$

The explicit computation of  $e^{S_{(r)}}$  will involve sum of all connected and disconnected world-sheets in the presence of  $r$  D-instantons with the modified world-sheet rules, followed by integration over the zero modes of these instantons.

Before giving a proof of (3.1), let us first verify that its expansion to order  $\mathcal{N}^2$  agrees with (2.8). By expanding (3.1) to order  $\mathcal{N}^2$ , we get

$$S = S_{(0)} + \mathcal{N}(e^{S_{(1)} - S_{(0)}} - 1) + \frac{1}{2!} \mathcal{N}^2 (e^{S_{(2)} - S_{(0)}} - e^{2(S_{(1)} - S_{(0)})}) + \mathcal{O}(\mathcal{N}^3), \quad (3.2)$$

in agreement with (2.8).

Let us now turn to the proof of (3.1). For this we rewrite this equation as

$$e^S = e^{-\mathcal{N}} e^{S_{(0)}} \left[ 1 + \sum_{r=1}^{\infty} \frac{1}{r!} \mathcal{N}^r e^{S_{(r)} - S_{(0)}} \right]. \quad (3.3)$$

On the left hand side, since  $S$  is supposed to be the effective action, the expansion of  $e^S$  should generate all possible combination of interaction vertices that arise in a Feynman diagram – with the understanding that the actual Feynman diagrams are constructed by joining the vertices to each other and to the external lines by propagators. Therefore this should correspond to all possible combinations of connected and disconnected world-sheets with external closed strings.<sup>4</sup> Thus checking (3.3) amounts to checking that the right hand side generates all possible world-sheets, including disconnected world-sheets that are not even connected in the target space. First consider the case of world-sheets without boundaries. These come from terms that contain only factors of  $S_{(0)}$  but no factors of  $S_{(r)} - S_{(0)}$ . Picking the 1 term in the expansion of  $e^{S_{(r)} - S_{(0)}}$  in (3.3), we see that the coefficient of the  $e^{S_{(0)}}$  term is

$$e^{-\mathcal{N}} \left[ 1 + \sum_{r=1}^{\infty} \frac{1}{r!} \mathcal{N}^r \right] = 1. \quad (3.4)$$

---

<sup>4</sup>Here by ‘all possible’ world-sheets we mean only those Riemann surfaces that appear in the construction of the interaction vertices of closed string effective field theory. Other Riemann surfaces that are constructed from these by sewing with closed string propagators are not included.



Therefore the right hand side of (3.3) has the contribution of connected and disconnected world-sheets, each without any boundary, with the correct coefficient.

Next we focus on the  $s$  instanton contribution from the right hand side of (3.3). Here by  $s$ -instanton contribution we mean the contribution from those connected or disconnected Riemann surfaces for which each of the boundaries lies on one of the  $s$  instantons and each of the  $s$  instantons has at least one boundary lying on it. One source of such terms is the expansion of  $e^{S_{(s)}-S_{(0)}} - 1$ . However the expansion of  $e^{S_{(r)}-S_{(0)}} - 1$  for any  $r \geq s$  also contains  $s$  instanton contribution, since in the product of the  $S_{(r)} - S_{(0)}$  factors, we have a contribution where all the boundaries lie on  $s$  of the  $r$  instantons and each of the  $s$  instantons has at least one boundary lying on it. Since  $s$  instantons can be chosen from  $r$  instantons in  $\binom{r}{s}$  ways, the net coefficient of the  $s$  instanton contribution to the amplitude will be given by

$$e^{-\mathcal{N}} \sum_{r \geq s} \frac{1}{r!} \mathcal{N}^r \binom{r}{s} = e^{-\mathcal{N}} \sum_{r \geq s} \frac{1}{s!(r-s)!} \mathcal{N}^r = \frac{1}{s!} \mathcal{N}^s. \quad (3.5)$$

This is precisely the desired contribution, since we need a factor of  $\mathcal{N}^s$  to accompany the  $s$  instanton contribution and the  $1/s!$  term reflects the fact that we have  $s$  identical instantons.

This establishes that the right hand side of (3.3) gives the perturbative and instanton contribution to  $e^S$  with the correct coefficient.

It is instructive to study the coefficient of the  $\mathcal{N}^3/3!$  term in (3.1). It is given by

$$\left[ e^{S_{(3)}-S_{(0)}} - 3 e^{S_{(2)}-S_{(0)}+S_{(1)}-S_{(0)}} + 2 e^{3(S_{(1)}-S_{(0)})} \right]. \quad (3.6)$$

With the definition of  $\widehat{S}'_{i_1 \dots i_k}$  given in section 2, (3.6) may be expressed as:

$$\int_z \left[ e^{\widehat{S}'_{123}+\widehat{S}'_{12}+\widehat{S}'_{23}+\widehat{S}'_{13}+\widehat{S}'_1+\widehat{S}'_2+\widehat{S}'_3} - e^{\widehat{S}'_{12}+\widehat{S}'_1+\widehat{S}'_2+\widehat{S}'_3} - e^{\widehat{S}'_{13}+\widehat{S}'_1+\widehat{S}'_3+\widehat{S}'_2} - e^{\widehat{S}'_{23}+\widehat{S}'_2+\widehat{S}'_3+\widehat{S}'_1} + 2 e^{\widehat{S}'_1+\widehat{S}'_2+\widehat{S}'_3} \right]. \quad (3.7)$$

Expanding this in a power series in  $\widehat{S}'_i$ ,  $\widehat{S}'_{ij}$  for  $1 \leq i, j \leq 3$  and  $\widehat{S}'_{123}$ , we can verify that each term in the expansion has one of the following four factors:  $\widehat{S}'_{123}$ ,  $\widehat{S}'_{12}\widehat{S}'_{23}$ ,  $\widehat{S}'_{12}\widehat{S}'_{13}$  or  $\widehat{S}'_{13}\widehat{S}'_{23}$ . This shows that each world-sheet diagram contributing to (3.6) is connected in the target space as desired.

We now turn to the second part of the problem, namely to show that  $S$  defined in (3.1) satisfies the BV master equation

$$\frac{1}{2}\{S, S\} + \Delta S = 0. \quad (3.8)$$

For this we express (3.3) as

$$e^S = e^{-\mathcal{N}} \left[ e^{S(0)} + \sum_{r=1}^{\infty} \frac{1}{r!} \mathcal{N}^r e^{S(r)} \right]. \quad (3.9)$$

It now follows from (2.4) that we have

$$\Delta e^S = 0. \quad (3.10)$$

This leads to (3.8) as a consequence of (2.3).

## 4 Non-identical D-instantons

The generalization of these results to non-identical D-instantons is straightforward. Let us suppose that the theory contains  $k$  different kinds of instantons of tensions  $\mathcal{T}_1, \dots, \mathcal{T}_k$  and let  $\mathcal{N}_i = e^{-\mathcal{T}_i}$  for  $1 \leq i \leq k$ . Let us denote by  $e^{S(r_1, r_2, \dots, r_k)}$  the sum over contributions from all connected and disconnected world-sheets in the presence of  $r_1$  instantons of the first kind,  $r_2$  instantons of the second kind etc., followed by integration over the D-instanton positions and other zero modes.  $S_{(r_1, r_2, \dots, r_k)}$  is then defined to be the logarithm of  $e^{S(r_1, r_2, \dots, r_k)}$ . Then  $S_{(r_1, r_2, \dots, r_k)}$  satisfies the BV master equation

$$\frac{1}{2} \{S_{(r_1, r_2, \dots, r_k)}, S_{(r_1, r_2, \dots, r_k)}\} + \Delta S_{(r_1, r_2, \dots, r_k)} = 0 \quad \Leftrightarrow \quad \Delta e^{S_{(r_1, r_2, \dots, r_k)}} = 0. \quad (4.1)$$

Our goal is to construct the effective action for closed strings after taking into account the effect of the weight factors  $\prod_{i=1}^k \mathcal{N}_i^{r_i}$  and including the contribution from the disconnected world-sheets that are target space connected. By following the same logic that established the validity of (3.3), we arrive at

$$e^S = e^{-\mathcal{N}_1 - \mathcal{N}_2 - \dots - \mathcal{N}_k} \sum_{r_1, \dots, r_k=0}^{\infty} \frac{1}{r_1! r_2! \dots r_k!} \mathcal{N}_1^{r_1} \mathcal{N}_2^{r_2} \dots \mathcal{N}_k^{r_k} e^{S_{(r_1, r_2, \dots, r_k)}}. \quad (4.2)$$

This can also be written as

$$S = S_{(\vec{0})} + \ln \left[ 1 + \sum_{\substack{r_1, \dots, r_k=0 \\ (r_1, \dots, r_k) \neq \vec{0}}}^{\infty} \frac{1}{r_1! r_2! \dots r_k!} \mathcal{N}_1^{r_1} \mathcal{N}_2^{r_2} \dots \mathcal{N}_k^{r_k} e^{S_{(r_1, r_2, \dots, r_k)} - S_{(\vec{0})}} \right] - \mathcal{N}_1 - \mathcal{N}_2 - \dots - \mathcal{N}_k. \quad (4.3)$$

Here  $S_{(\vec{0})}$  denotes the perturbative contribution to the effective action.

We now see that as a consequence of (4.1) and (4.2), we have

$$\Delta e^S = 0 \quad \Leftrightarrow \quad \frac{1}{2}\{S, S\} + \Delta S = 0. \quad (4.4)$$

This establishes that the full closed string effective action induced by multiple D-instantons of different kinds satisfy the BV master equation.

**Acknowledgement:** We thank Barton Zwiebach for useful discussions and comments and suggestions on an earlier draft of the manuscript that helped improve the presentation. This work was supported by the ICTS-Infosys Madhava Chair Professorship, the J. C. Bose fellowship of the Department of Science and Technology, India and the Department of Atomic Energy, Government of India, under project no. RTI4001.

## References

- [1] I. A. Batalin and G. A. Vilkovisky, “Gauge Algebra and Quantization,” *Phys. Lett.* **102B**, 27 (1981). doi:10.1016/0370-2693(81)90205-7
- [2] I. A. Batalin and G. A. Vilkovisky, “Quantization of Gauge Theories with Linearly Dependent Generators,” *Phys. Rev. D* **28**, 2567 (1983) Erratum: [*Phys. Rev. D* **30**, 508 (1984)]. doi:10.1103/PhysRevD.28.2567, 10.1103/PhysRevD.30.508
- [3] B. Zwiebach, “Closed string field theory: Quantum action and the B-V master equation,” *Nucl. Phys. B* **390**, 33-152 (1993) doi:10.1016/0550-3213(93)90388-6 [arXiv:hep-th/9206084 [hep-th]].
- [4] A. Sen and B. Zwiebach, “String Field Theory: A Review,” [arXiv:2405.19421 [hep-th]].
- [5] A. Sen, “Cutkosky rules and unitarity (violation) in D-instanton amplitudes,” *JHEP* **07** (2021), 205 doi:10.1007/JHEP07(2021)205 [arXiv:2012.00041 [hep-th]].
- [6] B. Zwiebach, “Oriented open - closed string theory revisited,” *Annals Phys.* **267** (1998), 193-248 doi:10.1006/aphy.1998.5803 [arXiv:hep-th/9705241 [hep-th]].
- [7] A. Sen, “Wilsonian Effective Action of Superstring Theory,” *JHEP* **01**, 108 (2017) doi:10.1007/JHEP01(2017)108 [arXiv:1609.00459 [hep-th]].

- [8] H. Kajiura, “Homotopy algebra morphism and geometry of classical string field theory,” Nucl. Phys. B **630** (2002), 361-432 doi:10.1016/S0550-3213(02)00174-8 [arXiv:hep-th/0112228 [hep-th]].
- [9] H. Kajiura, “Noncommutative homotopy algebras associated with open strings,” Rev. Math. Phys. **19**, 1-99 (2007) doi:10.1142/S0129055X07002912 [arXiv:math/0306332 [math.QA]].
- [10] J. Polchinski, “Combinatorics of boundaries in string theory,” Phys. Rev. D **50** (1994), R6041-R6045 doi:10.1103/PhysRevD.50.R6041 [arXiv:hep-th/9407031 [hep-th]].
- [11] A. Sen, “D-instantons, string field theory and two dimensional string theory,” JHEP **11** (2021), 061 doi:10.1007/JHEP11(2021)061 [arXiv:2012.11624 [hep-th]].