

# THE SPECTRUM OF THE VLADIMIROV SUB-LAPLACIAN ON THE COMPACT ENGEL GROUP

J.P. VELASQUEZ-RODRIGUEZ

ABSTRACT. Let  $p > 3$  be a prime number. In this note, we calculate explicitly the unitary dual and the matrix coefficients of the Engel group over the  $p$ -adic integers  $\mathcal{B}_4(\mathbb{Z}_p)$ . We use this information to calculate explicitly the spectrum of the Vladimirov sub-Laplacian, and show how it defines a globally hypoelliptic operator on  $\mathcal{B}_4$ .

## CONTENTS

1. Introduction	1
2. Preliminaries	6
2.1. The field of $p$ -adic numbers $\mathbb{Q}_p$	6
2.2. The Engel group over $\mathbb{Z}_p$	9
2.3. Directional VT operators	10
3. Representation theory of the Engel group	12
3.1. Level 1:	12
3.2. Level 2:	14
3.3. General case and Fourier series	17
4. The Vladimirov Sub-Laplacian on the Engel group	19
5. Acknowledgements	21
References	22

## 1. INTRODUCTION

When considering a partial differential operator  $L$ , a fundamental question in PDE theory is: *If  $Lu = f$  and  $f$  is smooth, does this imply that  $u$  is also smooth?* This question, which plays a central role in the theory of pseudo-differential operators on smooth manifolds, introduces the concept of *hypoellipticity*, see for instance [2] and related references. For example, in 1967, Hörmander demonstrated that, under the assumption that the system of smooth vector fields  $X_1, \dots, X_\kappa$  spans the entire tangent space at every point, a condition now known as *Hörmander's condition*, the

---

*Date:* July 10, 2024.

*2020 Mathematics Subject Classification.* Primary; 22E35, 58J4 ; Secondary: 20G05, 35R03, 42A16.

*Key words and phrases.* Pseudo-differential operators,  $p$ -adic Lie groups, representation theory, compact groups, Vladimirov-Taibleson operator.

The author is supported by the FWO Odysseus 1 Grant G.0H94.18N: Analysis and Partial Differential Equations, and the Methusalem programme of the Ghent University Special Research Fund (BOF) (Grant number 01M01021).

operator

$$L := \sum_{i=1}^k X_i^2 + X_0 + c \quad (1)$$

is hypoelliptic [6]. Such operators are commonly referred to as *Hörmander's operators*. Special cases include the *Hörmander's* sum of squares when  $X_0 = 0$  and  $c = 0$ , and important examples of Hörmander's operators include sub-Laplacians on Lie groups [2].

This paper addresses the issue of global hypoellipticity on compact stratified groups over non-archimedean local fields. Specifically, we introduce the concept of *directional Vladimirov–Taibleson operators* and discuss associated polynomials. These operators provide a framework for understanding differentiability on functions defined over profinite groups, with directional VT operators bearing similarities to directional derivatives.

**Definition 1.1.** Let  $\mathbb{K}$  be a non-archimedean local field with ring of integers  $\mathcal{O}_{\mathbb{K}}$ , prime ideal  $\mathfrak{p} = \mathfrak{p}\mathcal{O}_{\mathbb{K}}$ , and residue field  $\mathbb{F}_q = \mathcal{O}_{\mathbb{K}}/\mathfrak{p}\mathcal{O}_{\mathbb{K}}$ . Let  $\mathfrak{g} = \text{span}_{\mathcal{O}_{\mathbb{K}}}\{X_1, \dots, X_d\}$  be a nilpotent  $\mathcal{O}_{\mathbb{K}}$ -Lie algebra, and let  $G$  be the exponential image of  $\mathfrak{g}$ , so that  $G$  is a compact nilpotent  $\mathbb{K}$ -Lie group. We will use the symbol  $\partial_X^\alpha$  to denote the directional Vladimirov–Taibleson operator in the direction of  $X \in \mathfrak{g}$ , or directional VT operator for short, which we define as

$$\partial_X^\alpha f(x) := \frac{1 - q^\alpha}{1 - q^{-(\alpha+1)}} \int_{\mathcal{O}_{\mathbb{K}}} \frac{f(x \cdot \exp(tX)^{-1}) - f(x)}{|t|_{\mathbb{K}}^{\alpha+1}} dt, \quad \alpha > 0, \quad f \in \mathcal{D}(G).$$

Here  $\mathcal{D}(G)$  denotes the space of smooth functions on  $G$ , i.e., the collection of locally constant functions with a fixed index of local constancy.

**Remark 1.2.** Notice how the above definition is independent of any coordinate system we choose for the group as a manifold. Even though we are using the exponential map in the definition, this does not mean we are using the exponential system of coordinates. First,  $x \in G$  is not written in coordinates, and second, we know there is a one to one correspondence between elements of the Lie algebra  $\mathfrak{g}$  and one-parameter subgroups of  $G$ . Here a one-parameter subgroup of  $G$  is an analytic homomorphism  $\gamma_X : \mathcal{O}_{\mathbb{K}} \rightarrow G$ . Using this fact, given any  $X \in \mathfrak{g}$ , we can define the directional VT operator in the direction of  $X \in \mathfrak{g}$  via the formula

$$\partial_X^\alpha f(x) := \frac{1 - q^\alpha}{1 - q^{-(\alpha+1)}} \int_{\mathcal{O}_{\mathbb{K}}} \frac{f(x \cdot \gamma_X(t)^{-1}) - f(x)}{|t|_{\mathbb{K}}^{\alpha+1}} dt.$$

Since  $G$  is compact, we can identify it with a matrix group, where all one parameter subgroups have the form  $\gamma_X(t) = e^{tX}$ , which justifies our initial definition. More generally, we can write any analytic vector field on  $G$  as

$$X(x) := \sum_{j=1}^d c_j(x) X_j,$$

where the coefficient functions  $c_j$  are analytic. For this vector field, the directional VT operator is defined in a similar way as

$$\partial_X^\alpha f(x) := \frac{1 - q^\alpha}{1 - q^{-(\alpha+1)}} \int_{\mathcal{O}_{\mathbb{K}}} \frac{f(x \cdot \exp(tX(x))^{-1}) - f(x)}{|t|_{\mathbb{K}}^{\alpha+1}} dt.$$

However, in this work we wont consider the case where the coefficients are not constant, because such operators are not necessarily invariant.

By using Definition 2.5, we can link an specific Vladimirov-type operator to each direction  $X \in \mathfrak{g}$  and subsequently study the resulting operators. While this association does not preserve the Lie algebra structure, as seen in Lie groups over the real numbers, the resulting operators are nonetheless interesting and share similarities with partial differential operators on Lie groups. We want to study polynomials in these operators and, specifically, a distinguished operator called here the *Vladimirov sub-Laplacian*. For this operator, which we define in principle for compact stratified groups, we would like to pose the following conjecture:

**Conjecture.** Let  $\mathbb{K}$  be a non-archimedean local field with ring of integers  $\mathcal{O}_{\mathbb{K}}$ , prime ideal  $\mathfrak{p} = \mathfrak{p}\mathcal{O}_{\mathbb{K}}$ , and residue field  $\mathbb{F}_q = \mathcal{O}_{\mathbb{K}}/\mathfrak{p}\mathcal{O}_{\mathbb{K}}$ . Let  $\mathfrak{g} = \text{span}_{\mathcal{O}_{\mathbb{K}}} \{X_1, \dots, X_d\}$  be a stratified  $\mathcal{O}_{\mathbb{K}}$ -Lie algebra, and let  $G$  be the exponential image of  $\mathfrak{g}$ , so that  $G$  is a compact stratified  $\mathbb{K}$ -Lie group. Let  $X_1, \dots, X_{\kappa}$ ,  $1 \leq \kappa \leq d$ , be a basis for the first stratum of  $\mathfrak{g}$ , so that  $X_1, \dots, X_{\kappa}$  generates  $\mathfrak{g}$ , and let  $G$  be the exponential image of  $\mathfrak{g}$ . Then the *Vladimirov sub-Laplacian of order  $\alpha > 0$*  is a hypoelliptic operator on  $G$ , and it is invertible in the space of mean-zero functions. Here the Vladimirov sub-Laplacian is the pseudo-differential operator  $\mathcal{L}_{sub}^{\alpha}$ , defined on the space of smooth functions  $D(G)$  via the formula

$$\mathcal{L}_{sub}^{\alpha} f(x) := \sum_{k=1}^{\kappa} \partial_{X_k}^{\alpha} f(x),$$

where  $X_1, \dots, X_{\kappa}$  generates  $\mathfrak{g}$ .

Now, let  $p > 3$  be a prime number. In this note, we want to address the particular case of what we call here *the Engel group over the  $p$ -adic integers*, which we will denote by  $\mathcal{B}_4(\mathbb{Z}_p)$ , or simply  $\mathcal{B}_4$ . It can be defined in simple terms as  $\mathbb{Z}_p^4$  endowed with the non-commutative operation

$$x \star y := (x_1 + y_1, x_2 + y_2, x_3 + y_3 + x_1 y_2, x_4 + y_4 + x_1 y_3 + \frac{1}{2} x_1^2 y_2).$$

As an analytic manifold over  $\mathbb{Q}_p$ , we can identify its tangent space at the identity with  $\mathfrak{b}_4$ , the 4-dimensional Engel algebra, that is, the  $\mathbb{Z}_p$ -Lie algebra generated by  $X_1, \dots, X_4$ , with the commutation relations

$$[X_1, X_2] = X_3, \quad [X_1, X_3] = X_4, \quad \mathfrak{b}_4 = \bigoplus_{j=1}^3 \mathcal{V}_j,$$

where

$$\mathcal{V}_1 = \text{Span}_{\mathbb{Z}_p} \{X_1, X_2\}, \quad \mathcal{V}_2 = \text{Span}_{\mathbb{Z}_p} \{X_3\}, \quad \mathcal{V}_3 = \text{Span}_{\mathbb{Z}_p} \{X_4\}.$$

This algebra is stratified because the pair  $\{X_1, X_2\}$  generates the whole algebra, which is actually Hörmanders condition for a Lie group, so its associated *Vladimirov sub-Laplacian* is going to be:

$$\mathcal{L}_{sub}^{\alpha} f(x) := (\partial_{X_1}^{\alpha} + \partial_{X_2}^{\alpha}) f(x).$$

**Remark 1.3.** To the knowledge of the author, an operator like the Vladimirov sub-Laplacian has not appeared before in the mathematical literature. The closest thing is probably the study of the *Vladimirov-Laplacian* by Bendikov, Grigoryan, Pittet, and Woess [1]. There the authors study the operators

$$\mathcal{L}^{\alpha} f(x) := \sum_{k=1}^d \partial_{X_k}^{\alpha} f(x),$$

as the generators of a Dirichlet form associated with a certain jump kernel. See [1, Section 5] for more details.

So, drawing an analogy between the real case and the non-archimedean case, we can think on the Vladimirov sub-Laplacian as an analog of the sub-Laplacian or Hörmander's sum of squares for the non-archimedean case. Consequently, it is reasonable to expect the global hypoellipticity for these operators, and the main goal of this paper is to demonstrate that this is indeed the case for the 4-dimensional Engel group over the  $p$ -adic integers. See for instance [3, Example 2.2.7] for the definition of the Engel algebra and the Engel group over the reals.

**Remark 1.4.** *In this paper, we will identify each equivalence class  $\lambda$  in  $\widehat{\mathbb{Z}}_p \cong \mathbb{Q}_p/\mathbb{Z}_p$ , with its associated representative in the complete system of representatives*

$$\{1\} \cup \left\{ \sum_{k=1}^{\infty} \lambda_k p^{-k} : \text{only finitely many } \lambda_k \text{ are non-zero.} \right\}$$

The convention introduced above is important because it is used in the description of the unitary dual of the Heisenberg group over the  $p$ -adic integers, exposed in detail in [8]. We recall it here because we will use it to simplify the description of the unitary dual of  $\mathcal{B}_4$ .

**Theorem 1.5.** *Let  $\mathbb{H}_1(\mathbb{Z}_p)$ , or simply  $\mathbb{H}_1$  for short, be the 3-dimensional Heisenberg group over the  $p$ -adic integers. Let us denote by  $\widehat{\mathbb{H}}_1$  the unitary dual of  $\mathbb{H}_1$ , i.e., the collection of all equivalence classes of unitary irreducible representations of  $\mathbb{H}_1$ . Then we can identify  $\widehat{\mathbb{H}}_1$  with the following subset of  $\widehat{\mathbb{Z}}_p^3 \cong \mathbb{Q}_p^3/\mathbb{Z}_p^3$ :*

$$\widehat{\mathbb{H}}_1 = \{(\xi_1, \xi_2, \xi_3) \in \widehat{\mathbb{Z}}_p^3 : (\xi_1, \xi_2) \in \mathbb{Q}_p^2/|\xi_3|_p^{-1}\mathbb{Z}_p^2\}.$$

Moreover, each non-trivial representation  $[\pi_{(\xi_1, \xi_2, \xi_3)}] \in \widehat{\mathbb{H}}_1$  can be realized in the finite dimensional sub-space  $\mathcal{H}_{\xi_3}$  of  $L^2(\mathbb{Z}_p)$  defined as

$$\mathcal{H}_{\xi_3} := \text{span}_{\mathbb{C}}\{\varphi_h : h \in \mathbb{Z}_p/|\xi_3|_p\mathbb{Z}_p\}, \quad \varphi_h(u) := |\xi_3|_p^{1/2} \mathbb{1}_{h+|\xi_3|_p\mathbb{Z}_p}(u), \quad \dim(\mathcal{H}_{\xi_3}) = |\xi_3|_p,$$

where the representation acts on functions  $\varphi \in \mathcal{H}_{\xi_3}$  according to the formula

$$\pi_{(\xi_1, \xi_2, \xi_3)}(x)\varphi(u) := e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_3(x_3 + ux_2)\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_3}.$$

With this explicit realization, and by choosing the basis  $\{\varphi_h : h \in \mathbb{Z}_p/|\xi_3|_p\mathbb{Z}_p\}$  for each representation space, the associated matrix coefficients are given by

$$(\pi_{(\xi_1, \xi_2, \xi_3)})_{hh'} = e^{2\pi i\{\xi_3(x_3 + h'x_2) + (x_1\xi_1 + x_2\xi_2)\}_p} \mathbb{1}_{h-h'+|\xi_3|_p\mathbb{Z}_p}(x_1).$$

If we denote by  $\mathcal{Z}(\mathcal{B}_4)$  the center of  $\mathcal{B}_4(\mathbb{Z}_p)$ , then we can say there are two different kind of representations: those which are trivial on the center and the one that are not. If an unitary irreducible representation of  $\mathcal{B}_4(\mathbb{Z}_p)$  is trivial on the center, then it must be one of the representations of  $\mathcal{B}_4(\mathbb{Z}_p)/\mathcal{Z}(\mathcal{B}_4) \cong \mathbb{H}_1(\mathbb{Z}_p)$ , and these were calculated in [8]. Because of this, part of the work is already done and here we only need to focus on those representations which act non-trivially on the center.

As the main results of this paper, we will provide a complete description of the unitary dual, the matrix coefficients of the representations, and the spectrum of the Vladimirov sub-Laplacian on  $\mathcal{B}_4(\mathbb{Z}_p)$ . To be more precise, our goal in this work is to establish the following two theorems:

**Theorem 1.6.** Let  $\mathcal{B}_4(\mathbb{Z}_p)$ , or simply  $\mathcal{B}_4$  for short, be the 4-dimensional Engel group over the  $p$ -adic integers. Let us denote by  $\widehat{\mathcal{B}}_4$  the unitary dual of  $\mathcal{B}_4$ , i.e., the collection of all unitary irreducible representations of  $\mathcal{B}_4$ . Then we can identify  $\widehat{\mathcal{B}}_4$  with the following subset of  $\widehat{\mathbb{Z}}_p^4 \cong \mathbb{Q}_p^4/\mathbb{Z}_p^4$ :

$$\widehat{\mathcal{B}}_4 := \{\xi \in \widehat{\mathbb{Z}}_p^4 : 1 \leq |\xi_4|_p < |\xi_3|_p \wedge (\xi_1, \xi_2, \xi_3) \in \widehat{\mathbb{H}}_1, \text{ or } |\xi_3|_p = 1 \wedge \xi_1 \in \mathbb{Q}_p/|\xi_3|_p^{-1}\mathbb{Z}_p\}.$$

Moreover, each non-trivial representation  $[\pi_\xi]$  is equivalent to one of the representation  $[\pi_\xi] \in \widehat{\mathcal{B}}_4$  which can be realized in the following finite dimensional sub-space  $\mathcal{H}_\xi$  of  $L^2(\mathbb{Z}_p)$ :

$$\mathcal{H}_\xi := \text{span}_{\mathbb{C}}\{d_\xi^{1/2}\mathbb{1}_{h+d_\xi\mathbb{Z}_p} : h \in \mathbb{Z}_p/d_\xi\mathbb{Z}_p\},$$

where  $d_\xi := \dim_{\mathbb{C}}(\mathcal{H}_\xi) = \max\{|\xi_3|_p, |\xi_4|_p\}$ , and the representation acts on functions  $\varphi \in \mathcal{H}_\xi$  according to the formula

$$\pi_\xi(x)\varphi(u) := e^{2\pi i\{\xi_1x_1 + \xi_2x_2 + \xi_3(x_3 + ux_2) + \xi_4(x_4 + ux_3 + \frac{u^2}{2}x_2)\}_p} \varphi(u + x_1).$$

With this explicit realization of  $[\pi_\xi] \in \widehat{\mathcal{B}}_4$ , and with the natural choice of basis for each representation space, the associated matrix coefficients  $(\pi_\xi)_{hh'}$  are given by

$$(\pi_\xi(x))_{hh'} = e^{2\pi i\{\xi \cdot x\}_p} \mathbb{1}_{h-h' + \|\xi\|_p\mathbb{Z}_p}(x_1) \left( \|\xi\|_p \int_{h'+\|\xi\|_p\mathbb{Z}_p} e^{2\pi i\{(\xi_3x_2 + \xi_4x_3)u + \xi_4x_2\frac{u^2}{2}\}_p} du \right)$$

Sometimes we will use the notation

$$\mathcal{V}_\xi := \text{span}_{\mathbb{C}}\{(\pi_\xi)_{hh'} : h, h' \in \mathbb{Z}_p/\|\xi\|_p\mathbb{Z}_p\},$$

With the description of the unitary dual given in Theorem 1.6, we can prove the following spectral theorem for the Vladimirov sub-Laplacian on  $\mathcal{B}_4$ :

**Theorem 1.7.** The Vladimirov sub-Laplacian associated to the basis  $\{X_1, X_2\}$  of the first stratum of  $\mathfrak{b}_4$

$$\mathcal{L}_{sub}^\alpha := \partial_{X_1}^\alpha + \partial_{X_2}^\alpha,$$

defines a left-invariant, self-adjoint, sub-elliptic operator on  $\mathcal{B}_4$ . The spectrum of this operator is purely punctual, and its associated eigenfunctions form an orthonormal basis of  $L^2(\mathcal{B}_4)$ . Furthermore, the symbol of  $\mathcal{L}_{sub}^\alpha$  acts on each representation space as a  $p$ -adic Schrödinger type operator, and the space  $L^2(\mathcal{B}_4)$  can be written as the direct sum

$$L^2(\mathcal{B}_4) = \overline{\bigoplus_{[\xi] \in \widehat{\mathcal{B}}_4} \bigoplus_{h' \in \mathbb{Z}_p/\|\xi\|_p\mathbb{Z}_p} \mathcal{V}_\xi^{h'}}}, \quad \mathcal{V}_\xi = \bigoplus_{h' \in \mathbb{Z}_p/\|\xi\|_p\mathbb{Z}_p} \mathcal{V}_\xi^{h'}$$

where each finite-dimensional sub-space

$$\mathcal{V}_\xi^{h'} := \text{span}_{\mathbb{C}}\{(\pi_\xi)_{hh'} : h \in \mathbb{Z}_p/\|\xi\|_p\mathbb{Z}_p\},$$

is an invariant sub-space where  $\mathcal{L}_{sub}^\alpha$  acts like the Schrödinger-type operator

$$\partial_{x_1}^\alpha + Q(\xi, h) - \frac{1 - p^{-1}}{1 - p^{-(\alpha+1)}}.$$

Consequently, the spectrum of  $\mathcal{L}_{sub}^\alpha$  restricted to  $\mathcal{V}_\xi^{h'}$  is purely punctual, and it is given by

$$\begin{cases} |\xi_1|_p^\alpha + |\xi_2|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } |\xi_3|_p = |\xi_4|_p = 1, \\ |\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_3 h'|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } 1 = |\xi_4|_p < |\xi_3|_p, \\ (|\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_3 h' + \xi_4 \frac{(h')^2}{2}|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}})e_{\xi, h', \tau}(x) & \text{if } 1 < |\xi_4|_p < |\xi_3|_p, \\ (|\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_4 \frac{(h')^2}{2}|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}})e_{\xi, h', \tau}(x) & \text{if } 1 = |\xi_3|_p < |\xi_4|_p, \end{cases} \quad h \in \mathbb{Z}_p/|\xi_4|_p\mathbb{Z}_p,$$

where  $\tau \in \mathbb{Z}_p/\|(\xi_3, \xi_4)\|_p\mathbb{Z}_p$ , and the corresponding eigenfunctions are given by

$$e_{\xi, h', \tau}(x) := e^{2\pi i\{\xi x + \tau x_1\}_p} \left( d_\xi \int_{h' + d_\xi \mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right),$$

where

$$\xi \in \widehat{B}_4, h' \in \mathbb{Z}_p/\|(\xi_3, \xi_4)\|_p\mathbb{Z}_p, 1 \leq |\tau|_p \leq \|(\xi_3, \xi_4)\|_p.$$

## 2. PRELIMINARIES

**2.1. The field of  $p$ -adic numbers  $\mathbb{Q}_p$ .** Throughout this article  $p > 2$  will denote a fixed prime number. The field of  $p$ -adic numbers, usually denoted by  $\mathbb{Q}_p$ , can be defined as the completion of the field of rational numbers  $\mathbb{Q}$  with respect to the  $p$ -adic norm  $|\cdot|_p$ , defined as

$$|u|_p := \begin{cases} 0 & \text{if } u = 0, \\ p^{-\gamma} & \text{if } u = p^\gamma \frac{a}{b}, \end{cases}$$

where  $a$  and  $b$  are integers coprime with  $p$ . The integer  $\gamma := \text{ord}(u)$ , with  $\text{ord}(0) := +\infty$ , is called the  $p$ -adic order of  $u$ . The unit ball of  $\mathbb{Q}_p^d$  with the  $p$ -adic norm

$$\|u\|_p := \max_{1 \leq j \leq d} |u_j|_p,$$

is called the group of  $p$ -adic integers, and it will be denoted by  $\mathbb{Z}_p^d$ . Any  $p$ -adic number  $u \neq 0$  has a unique expansion of the form

$$u = p^{\text{ord}(u)} \sum_{j=0}^{\infty} u_j p^j,$$

where  $u_j \in \{0, 1, \dots, p-1\}$  and  $u_0 \neq 0$ . With this expansion we define the fractional part of  $u \in \mathbb{Q}_p$ , denoted by  $\{u\}_p$ , as the rational number

$$\{u\}_p := \begin{cases} 0 & \text{if } u = 0 \text{ or } \text{ord}(u) \geq 0, \\ p^{\text{ord}(u)} \sum_{j=0}^{-\text{ord}(u)-1} u_j p^j, & \text{if } \text{ord}(u) < 0. \end{cases}$$

$\mathbb{Z}_p^d$  is compact, totally disconnected, i.e. profinite, and abelian. Its dual group in the sense of Pontryagin, the collection of characters of  $\mathbb{Z}_p^d$  will be denoted by  $\widehat{\mathbb{Z}}_p^d$ . The dual group of the  $p$ -adic integers is known to be the Prüfer group  $\mathbb{Z}(p^\infty)$ , the unique  $p$ -group in which every element has  $p$  different  $p$ -th roots. The Prüfer group may be identified with the quotient group  $\mathbb{Q}_p/\mathbb{Z}_p$ . In this way the characteres of the group  $\mathbb{Z}_p^d$  may be written as

$$\chi_p(\tau u) := e^{2\pi i\{\tau \cdot u\}_p}, \quad u \in \mathbb{Z}_p^d, \quad \tau \in \widehat{\mathbb{Z}}_p^d \cong \mathbb{Q}_p^d/\mathbb{Z}_p^d.$$

**Remark 2.1.** Along this work, it will be convenient at several points to identify the elements of  $\widehat{\mathbb{Z}}_p$ , which are equivalence classes  $[\xi] \in \mathbb{Q}_p/\mathbb{Z}_p$ , with the representative of the class  $\tau = \sum_{k=1}^{\infty} \tau_k p^{-k}$ , where only finitely many  $\tau_k \in \mathbb{F}_p$  are non-zero.

By the Peter-Weyl theorem the elements of  $\widehat{\mathbb{Z}}_p^d$  constitute an orthonormal basis for the Hilbert space  $L^2(\mathbb{Z}_p^d)$ , which provide us a Fourier analysis for suitable functions defined on  $\mathbb{Z}_p$  in such a way that the formula

$$\varphi(u) = \sum_{\tau \in \widehat{\mathbb{Z}}_p^d} \widehat{\varphi}(\tau) \chi_p(\tau u),$$

holds almost everywhere in  $\mathbb{Z}_p$ . Here  $\mathcal{F}_{\mathbb{Z}_p^d}[\varphi]$  denotes the Fourier transform of  $f$ , in turn defined as

$$\mathcal{F}_{\mathbb{Z}_p^d}[\varphi](\tau) := \int_{\mathbb{Z}_p^d} \varphi(u) \overline{\chi_p(\tau u)} du,$$

where  $du$  is the normalised Haar measure on  $\mathbb{Z}_p^d$ .

Another important tool from  $p$ -adic analysis that we are going to be using, is the calculation of the  $p$ -adic Gaussian integral on the disk, see [9, p.p. 65].

**Lemma 2.2.** Let  $p \neq 2$  be a prime number, and take any  $\gamma \in \mathbb{Z}$ . Then, for any  $a, b \in \mathbb{Q}_p$ , we have

$$\int_{p^\gamma \mathbb{Z}_p} e^{2\pi i \{au^2 + bu\}_p} du = \begin{cases} p^{-\gamma} \mathbb{1}_{p^{-\gamma} \mathbb{Z}_p}(b), & \text{if } |a|_p \leq p^{2\gamma}, \\ \lambda_p(a) |a|_p^{-1/2} e^{2\pi i \{-\frac{b^2}{4a}\}_p} \mathbb{1}_{p^\gamma \mathbb{Z}_p}(b/a) & \text{if } |a|_p > p^{2\gamma}. \end{cases}$$

Here  $\lambda_p : \mathbb{Q}_p \rightarrow \mathbb{C}$  is the function defined as

$$\lambda_p(a) := \begin{cases} 1, & \text{if } \text{ord}(u) \text{ is even,} \\ \left(\frac{a_0}{p}\right), & \text{if } \text{ord}(u) \text{ is odd and } p \equiv 1 \pmod{4}, \\ i \left(\frac{a_0}{p}\right), & \text{if } \text{ord}(u) \text{ is odd and } p \equiv 3 \pmod{4}, \end{cases}$$

where  $\left(\frac{a_0}{p}\right)$  denotes the Legendre symbol of  $a_0$ . Here, to simplify our notation, we are going to define

$$\Lambda(a, b) := \lambda_p(a) |a|_p^{-1/2} e^{2\pi i \{-\frac{b^2}{4a}\}_p},$$

so that  $|\Lambda(a, b)| = |a|_p^{-1/2}$ , and in this way

$$\int_{p^\gamma \mathbb{Z}_p} e^{2\pi i \{au^2 + bu\}_p} du = p^{-\gamma} \mathbb{1}_{p^{-2\gamma} \mathbb{Z}_p}(a) \cdot \mathbb{1}_{p^{-\gamma} \mathbb{Z}_p}(b) + \mathbb{1}_{\mathbb{Q}_p \setminus p^{-2\gamma} \mathbb{Z}_p}(a) \cdot \Lambda(a, b) \cdot \mathbb{1}_{p^\gamma \mathbb{Z}_p}(b/a).$$

With this tool we can prove the following auxiliary lemma:

**Lemma 2.3.** Let  $(x_2, x_3) \in \mathbb{Z}_p^2$ , and take  $\xi_3, \xi_4 \in \widehat{\mathbb{Z}}_p$ . Then

$$\int_{\mathbb{Z}_p^2} \left| \int_{\mathbb{Z}_p} e^{2\pi i \{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right|^2 dx_2 dx_3 = \max\{|\xi_3|_p, |\xi_4|_p\}^{-1} = \|(\xi_3, \xi_4)\|_p^{-1}.$$

*Proof.* Let us define the auxiliary function

$$f(x_2, x_3) := \int_{\mathbb{Z}_p} e^{2\pi i \{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du.$$

Then, as a function on  $\mathbb{Z}_p^2$ , we can compute its Fourier transform as

$$\begin{aligned}\mathcal{F}_{\mathbb{Z}_p^2}[f](\alpha, \beta) &= \int_{\mathbb{Z}_p^2} \int_{\mathbb{Z}_p} e^{2\pi i \{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} - \alpha x_2 - \beta x_3\}_p} dudx_2 dx_3 \\ &= \int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p^2} e^{2\pi i \{(\xi_3 u + \xi_4 \frac{u^2}{2} - \alpha)x_2 - (\xi_4 u - \beta)x_3\}_p} dx_2 dx_3 du \\ &= \int_{\mathbb{Z}_p} \mathbb{1}_{\mathbb{Z}_p}(\xi_3 u + \xi_4 \frac{u^2}{2} - \alpha) \mathbb{1}_{\mathbb{Z}_p}(\xi_4 u - \beta) du.\end{aligned}$$

Notice how  $|\xi_4 u - \beta|_p \leq 1$  implies that  $\xi_4 u = \beta$  in  $\mathbb{Q}_p/\mathbb{Z}_p$ . In this way, we have the following equality modulo  $\mathbb{Z}_p$ :

$$\xi_3 u + \xi_4 \frac{u^2}{2} = \frac{u}{2}(2\xi_3 + \xi_4 u) = \frac{u}{2}(2\xi_3 + \beta) \in \mathbb{Q}_p/\mathbb{Z}_p,$$

therefore

$$\begin{aligned}\mathcal{F}_{\mathbb{Z}_p^2}[f](\alpha, \beta) &= \int_{\mathbb{Z}_p} \mathbb{1}_{\mathbb{Z}_p}(u(\xi_3 + \beta/2) - \alpha) \mathbb{1}_{\mathbb{Z}_p}(\xi_4 u - \beta) du \\ &= \int_{\mathbb{Z}_p} \mathbb{1}_{\frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p}(u) \mathbb{1}_{\frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p}(u) du \\ &= \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right|.\end{aligned}$$

Here we have to notice how, since we are integrating over  $u \in \mathbb{Z}_p$ , in order for the above integral to be different from zero we need

$$\left| \frac{\alpha}{\xi_3 + \beta/2} \right|_p \leq 1, \text{ and } \left| \frac{\beta}{\xi_4} \right|_p \leq 1,$$

so that only finitely many terms  $\mathcal{F}_{\mathbb{Z}_p^2}[f](\alpha, \beta)$  can be non-zero. Also

$$\mathcal{F}_{\mathbb{Z}_p^2}[f](\alpha, \beta) = \begin{cases} 0, & \text{if the balls are disjoint,} \\ \max\{|\xi_4|_p, |\xi_3 + \beta/2|_p\}^{-1}, & \text{if the balls are not disjoint.} \end{cases}$$

Finally, we use this Fourier transform to calculate the  $L^2$ -norm of  $f$  in two different cases. When  $|\xi_3|_p > |\xi_4|_p$ , we have  $|\xi_3 + \beta/2|_p = |\xi_3|_p$ , so that

$$\begin{aligned}\|f\|_{L^2(\mathbb{Z}_p^2)}^2 &= \sum_{(\alpha, \beta) \in \widehat{\mathbb{Z}}_p^2} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right|^2 \\ &= \sum_{(\alpha, \beta) \in \widehat{\mathbb{Z}}_p^2} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right|^2 \\ &= \sum_{\beta \in \widehat{\mathbb{Z}}_p} |\xi_3|_p^{-1} \sum_{\alpha \in \widehat{\mathbb{Z}}_p} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right| \\ &= \sum_{\beta \in \widehat{\mathbb{Z}}_p} |\xi_3|_p^{-1} \left| \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right| = |\xi_3|_p^{-1}.\end{aligned}$$



When  $|\xi_3|_p \leq |\xi_4|_p$ , we get  $|\xi_3 + \beta/2|_p \leq |\xi_4|_p$ , so we conclude in a similar way

$$\begin{aligned} \|f\|_{L^2(\mathbb{Z}_p^2)}^2 &= \sum_{(\alpha, \beta) \in \widehat{\mathbb{Z}}_p^2} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right|^2 \\ &= \sum_{\alpha \in \widehat{\mathbb{Z}}_p} \sum_{\beta \in \widehat{\mathbb{Z}}_p} |\xi_4|_p^{-1} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p \cap \frac{\beta}{\xi_4} + |\xi_4|_p \mathbb{Z}_p \right| \\ &= \sum_{\alpha \in \widehat{\mathbb{Z}}_p} |\xi_4|_p^{-1} \left| \frac{\alpha}{\xi_3 + \beta/2} + |\xi_3 + \beta/2|_p \mathbb{Z}_p \right| = |\xi_4|_p^{-1}. \end{aligned}$$

This concludes the proof.  $\square$

**2.2. The Engel group over  $\mathbb{Z}_p$ .** Let  $p > 2$  be a prime number. Let us denote by  $\mathfrak{b}_4$  the  $\mathbb{Z}_p$ -Lie algebra generated by  $X_1, \dots, X_4$ , with the commutation relations

$$[X_1, X_2] = X_3, \quad [X_1, X_3] = X_4, \quad \mathfrak{b}_4 = \bigoplus_{j=1}^3 \mathcal{V}_j,$$

where

$$\mathcal{V}_1 = \text{Span}_{\mathbb{Z}_p} \{X_1, X_2\}, \quad \mathcal{V}_2 = \text{Span}_{\mathbb{Z}_p} \{X_3\}, \quad \mathcal{V}_3 = \text{Span}_{\mathbb{Z}_p} \{X_4\}.$$

We call the  $\mathbb{Z}_p$ -Lie algebra  $\mathfrak{b}_4$  the 4-dimensional *Engel algebra*, and its exponential image, which we denote here by  $B_4$ , is called the *Engel group* over the  $p$ -adic integers. Let us consider the realization of  $\mathfrak{b}_4$  as the matrix algebra

$$\mathfrak{b}_4(\mathbb{Z}_p) = \left\{ \begin{bmatrix} 0 & x_1 & 0 & x_4 - \frac{1}{2}x_1(x_3 - \frac{1}{2}x_1x_2) - \frac{1}{6}x_1^2x_2 \\ 0 & 0 & x_1 & x_3 - \frac{1}{2}x_1x_2 \\ 0 & 0 & 0 & x_2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \in \mathcal{M}_4(\mathbb{Z}_p) : x \in \mathbb{Z}_p^4 \right\}.$$

With this realization, and using the usual matrix exponential map, we can think on  $B(\mathbb{Z}_p)$  as the matrix group

$$B_4(\mathbb{Z}_p) = \left\{ \begin{bmatrix} 1 & x_1 & \frac{1}{2}x_1^2 & x_4 \\ 0 & 1 & x_1 & x_3 \\ 0 & 0 & 1 & x_2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \in \mathcal{M}_4(\mathbb{Z}_p) : x \in \mathbb{Z}_p^4 \right\},$$

which analytically isomorphic to  $\mathbb{Z}_p^4$  with the operation

$$x \star y := (x_1 + y_1, x_2 + y_2, x_3 + y_3 + x_1y_2, x_4 + y_4 + x_1y_3 + \frac{1}{2}x_1^2y_2).$$

The exponential map transforms sub-ideals of the Lie algebra  $\mathfrak{b}_4$  to subgroups of  $B_4 \cong (\mathfrak{b}_4, *)$ , which can be endowed with the sequence of subgroups  $J_n := (\mathfrak{b}_4(p^n \mathbb{Z}_p), *)$ , where

$$\mathfrak{b}_4(p^n \mathbb{Z}_p) = p^n \mathbb{Z}_p X_1 + p^n \mathbb{Z}_p X_2 + p^n \mathbb{Z}_p X_3 + p^n \mathbb{Z}_p X_4,$$

so  $B_4$  is a compact Vilenkin group, together with the sequence of compact open subgroups

$$G_n := B_4(p^n \mathbb{Z}_p) = \mathbf{exp}(p^n \mathfrak{b}_4), \quad n \in \mathbb{N}_0.$$

Notice how the sequence  $\mathcal{G} = \{G_n\}_{n \in \mathbb{N}_0}$  forms a basis of neighbourhoods at the identity, so the group is metrizable, and we can endow it with the natural ultrametric

$$|xy^{-1}|_{\mathcal{G}} := \begin{cases} 0 & \text{if } x = y, \\ |G_n| = p^{-4n} & \text{if } x \star y^{-1} \in G_n \setminus G_{n+1}. \end{cases}$$

Nevertheless, instead of this ultrametric we will use the  $p$ -adic norm

$$\|x\|_p := \max_{1 \leq j \leq 4} \|x_j\|_p.$$

Notice that how  $\|x\|_p^4 = |x|_{\mathcal{G}}$ , for any  $x \in B_4$

**2.3. Directional VT operators.** One important idea from the theory of differential and pseudo-differential operators on Lie groups, is the correspondence between directional derivatives and elements of the Lie algebra. However, in the  $p$ -adic case, there are plenty of non-trivial locally constant functions, due to the fact that  $p$ -adic numbers are totally disconnected. This means that the usual notion of derivative does not apply, and therefore we need to find an alternative kind of operators to talk about differentiability on these groups. A first approach to this problem can be the *Vladimirov-Taibleson operator* [4, 5], which we define for general compact  $\mathbb{K}$ -Lie groups as follows:

**Definition 2.4.** Let  $\mathbb{K}$  be a non-archimedean local field with ring of integers  $\mathcal{O}_{\mathbb{K}}$ , prime ideal  $\mathfrak{p} = \mathfrak{p}\mathcal{O}_{\mathbb{K}}$  and residue field  $\mathbb{F}_q \cong \mathcal{O}_{\mathbb{K}}/\mathfrak{p}\mathcal{O}_{\mathbb{K}}$ . Let  $G \leq GL_m(\mathcal{O}_{\mathbb{K}})$  be a compact  $d$ -dimensional  $\mathbb{K}$ -Lie group. We define the *Vladimirov-Taibleson operator* on  $G$  via the formula

$$D^\alpha f(x) := \frac{1 - q^\alpha}{1 - q^{-(\alpha+d)}} \int_G \frac{f(xy^{-1}) - f(x)}{\|y\|_{\mathbb{K}}^{\alpha+d}} dy,$$

where

$$\|y\|_{\mathbb{K}} := \begin{cases} 1, & \text{if } y \in G \setminus GL_m(\mathfrak{p}\mathcal{O}_{\mathbb{K}}), \\ q^{-n}, & \text{if } y \in GL_m(\mathfrak{p}^n\mathcal{O}_{\mathbb{K}}) \setminus GL_m(\mathfrak{p}^{n+1}\mathcal{O}_{\mathbb{K}}). \end{cases}$$

Here  $dy$  denotes the unique normalized Haar measure on  $G$ . Sometimes it will be convenient to consider the operator

$$\mathbb{D}^\alpha f(x) := \frac{1 - q^{-d}}{1 - q^{-(\alpha+d)}} f(x) + \frac{1 - q^\alpha}{1 - q^{-(\alpha+d)}} \int_G \frac{f(xy^{-1}) - f(x)}{\|y\|_{\mathbb{K}}^{\alpha+d}} dy,$$

The Vladimirov-Taibleson operator can be considered as a fractional Laplacian for functions on totally disconnected spaces, and it provides a first notion of differentiability. However, for functions of several variables it is natural to consider the differentiability of the function in each variable, or in a certain given direction. For that reason, we introduce the following definition:

**Definition 2.5.** Let  $\mathfrak{g}$  be the  $\mathcal{O}_{\mathbb{K}}$ -Lie module associated to  $G$ , and assume that  $\mathfrak{g}$  is nilpotent. Let  $\alpha > 0$ . Given a  $V \in \mathfrak{g}$ , we define the *directional VT operator of order  $\alpha$  in the direction of  $V$*  as the linear invariant operator  $\partial_V^\alpha$  acting on smooth functions via the formula

$$\partial_V^\alpha f(x) := \frac{1 - q^\alpha}{1 - q^{-(\alpha+1)}} \int_{\mathcal{O}_{\mathbb{K}}} \frac{f(x \cdot \exp(tV)^{-1}) - f(x)}{|t|_{\mathbb{K}}^{\alpha+1}} dt, \quad f \in \mathcal{D}(G).$$

**Remark 2.6.** *Directional VT operators are interesting because they associate a certain pseudo-differential operator to each element of the Lie algebra. Nonetheless, it is important to remark how this association does not follow the same patten as in the locally connected case, where the correspondence between vectors and operators preserves the Lie algebra structure.*

Just to give an example, in the particular case when  $G = \mathfrak{g} = \mathbb{Z}_p^d$ , these operators take the form

$$\partial_V^\alpha f(x) := \frac{1 - p^\alpha}{1 - p^{-(\alpha+1)}} \int_{\mathbb{Z}_p^d} \frac{f(x - tV) - f(x)}{|t|_p^{\alpha+1}} dt,$$

and one can easily compute its associated symbol:  $\sigma_{\partial_V^\alpha}(\xi) = 0$ , if  $V \cdot \xi = 0$ , and

$$\sigma_{\partial_V^\alpha}(\xi) = |V \cdot \xi|_p^\alpha - \frac{1 - p^{-1}}{1 - p^{-(\alpha+1)}}, \quad \xi \in \widehat{\mathbb{Z}}_p^d,$$

in other case. If we define  $\partial_{x_i}^\alpha := \partial_{e_i}^\alpha$ , where  $e_i$ ,  $1 \leq i \leq d$  are the canonical vectors of  $\mathbb{Q}_p^d$ , then

$$\sigma_{\partial_{x_i}^\alpha}(\xi) = \begin{cases} 0, & \text{if } |\xi_i|_p = 1, \\ |\xi_i|_p^\alpha - \frac{1 - p^{-1}}{1 - p^{-(\alpha+1)}} & \text{if } |\xi_i|_p > 1, \end{cases}$$

which resembles the symbol of the usual partial derivatives on  $\mathbb{R}^d$ , justifying that way our choice of notation. However, we want to be emphatic about the fact that these directional VT operators are not derivatives, but rather some special kind of pseudo-differential operators which we will study with the help of the Fourier analysis on compact groups.

Before proceeding to the next section, let us introduce some notation.

**Definition 2.7.**

- The symbol  $Rep(\mathcal{B}_4)$  will denote the collection of all unitary finite-dimensional representations of  $\mathcal{B}_4$ . We will denote by  $\widehat{\mathcal{B}}_4$  the *unitary dual* of  $\mathcal{B}_4$ .
- Let  $K$  be a normal sub-group of  $\mathcal{B}_4$ . We denote by  $K^\perp$  the *annihilator* of  $K$ , here defined as

$$K^\perp := \{[\pi] \in Rep(\mathcal{B}_4) : \pi|_K = I_{d_\pi}\}.$$

Also, we will use the notation

$$B_{\widehat{\mathcal{B}}_4}(n) := \widehat{\mathcal{B}}_4 \cap \mathcal{B}_4(p^n \mathbb{Z}_p)^\perp,$$

and  $\widehat{\mathcal{B}}_4(n) := B_{\widehat{\mathcal{B}}_4}(n) \setminus B_{\widehat{\mathcal{B}}_4}(n-1)$ .

- We say that a function  $f : \mathcal{B}_4 \rightarrow \mathbb{C}$  is a *smooth function*, if  $f$  is a locally constant function with a fixed index of local constancy, i.e., there is an  $n_f \in \mathbb{N}_0$ , which we always choose to be the minimum possible, such that

$$f(x \star y) = f(x), \quad \text{for all } y \in \mathcal{B}_4(p^n \mathbb{Z}_p).$$

We will denote by  $D(\mathcal{B}_4)$  the collection of all smooth functions on  $\mathcal{B}_4$ , and  $D_n(\mathcal{B}_4)$  will denote the collection of smooth functions with index of local constancy equal to  $n \in \mathbb{N}_0$ .

### 3. REPRESENTATION THEORY OF THE ENGEL GROUP

Given a representation  $[\pi] \in \widehat{\mathcal{B}}_4$  we can say that there are two possibilities: the representation is trivial on the center, or it is not. When the representation is trivial on the center, since  $\mathcal{B}_4/\mathcal{Z}(\mathcal{B}_4) \cong \mathbb{H}_1(\mathbb{Z}_p)$ , the representation must descend to an unitary irreducible representation of  $\mathbb{H}_1(\mathbb{Z}_p)$ . These representations were completely described in [8], and they have the form

$$\pi_{(\xi_1, \xi_2, \xi_3)}(x)\varphi(u) := e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_3(x_3 + ux_2)\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_3},$$

where

$$\mathcal{H}_{\xi_3} := \text{Span}_{\mathbb{C}}\{\varphi_h : h \in \mathbb{Z}_p/|\xi_3|_p\mathbb{Z}_p\}, \quad \varphi_h(u) := |\xi_3|^{1/2} \mathbb{1}_{h+p^{o(\xi_3)}\mathbb{Z}_p}(u), \quad \dim(\mathcal{H}_{\xi_3}) = |\xi_3|_p.$$

See [8] for the full calculation of  $\widehat{\mathbb{H}}_d(\mathbb{Z}_p)$ . This shorten our calculations here by a lot, and actually, the ideas introduced in [8] are very useful to deduce how to obtain the unitary dual for the more complicated case of  $\mathcal{B}_4$ .

**3.1. Level 1:** We will use the fact that any representation of  $\mathcal{B}_4$  reduces to the representation of some finite group, obtained as the quotient between  $\mathcal{B}_4$  and one of it compact open subgroups. See again the notation introduced in Definition 2.7.

First, as it is obvious,  $\mathcal{B}_{\widehat{\mathcal{B}}_4}(0)$ , the collection of unitary irreducible representations that are trivial on  $G_0 = \mathcal{B}_4(\mathbb{Z}_p)$ , contains only the identity representation. To calculate  $\mathcal{B}_{\widehat{\mathcal{B}}_4}(1)$ , the collection of unitary irreducible representations that are trivial on  $G_1 = \mathcal{B}_4(p\mathbb{Z}_p)$ , we have to find those representations which descend to an element of the unitary dual of  $G/G_1 \cong \mathcal{B}_4(\mathbb{Z}_p/p\mathbb{Z}_p) \cong \mathcal{B}_4(\mathbb{F}_p)$ . The usual representation theory of nilpotent groups applied to  $\mathcal{B}_4(\mathbb{F}_p)$  provides us with three different kind of representations, which correspond to the three different kind of co-adjoint orbits obtained via the Kirilov orbit method:

- We have  $p^2$  one dimensional representations given by the  $p$ -adic characters

$$\pi_{(\xi_1, \xi_2, 1, 1)}(x) = \chi_{\xi_1, \xi_2}(x) = e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2\}_p}, \quad \|(\xi_1, \xi_2)\|_p \leq p.$$

Clearly it holds  $\int_{\mathcal{B}_4} |\chi_{\xi_1, \xi_2}(x)|^2 dx = 1$ .

- There are  $p - 1$  non-commutative representations which are trivial on the center of  $\mathcal{B}_4(\mathbb{F}_p)$ , and therefore descend to a representation of  $\mathbb{H}_1(\mathbb{F}_p)$ . These have the form

$$\pi_{(1, 1, \xi_3, 1)}(x) = \pi_{\xi_3}(x)\varphi(u) := e^{2\pi i\{\xi_3(x_3 + ux_2)\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_3},$$

where  $|\xi_3|_p = p$ , and

$$\mathcal{H}_{\xi_3} := \text{span}_{\mathbb{C}}\{\varphi_h : h \in \mathbb{Z}_p/p\mathbb{Z}_p\}, \quad \varphi_h(u) := p^{1/2} \mathbb{1}_{h+p\mathbb{Z}_p}(u), \quad \dim(\mathcal{H}_{\xi_3}) = p.$$

The characters associated to these representations have the form

$$\chi_{\pi_{\xi_3}}(x) = \sum_{h \in \mathbb{Z}_p/p\mathbb{Z}_p} e^{2\pi i\{\xi_3(x_3 + hx_2)\}_p} \mathbb{1}_{p\mathbb{Z}_p}(x_1) = p e^{2\pi i\{\xi_3 x_3\}_p} \mathbb{1}_{p\mathbb{Z}_p}(x_1) \mathbb{1}_{p\mathbb{Z}_p}(x_2),$$

thus the irreducibility of  $[\pi_{\xi_3}]$  is proved by the condition

$$\int_{\mathcal{B}_4} |\chi_{\pi_{\xi_3}}(x)|^2 dx = p^2 \int_{\mathcal{B}_4} |\mathbb{1}_{p\mathbb{Z}_p}(x_1) \mathbb{1}_{p\mathbb{Z}_p}(x_2)|^2 dx = 1.$$

- There are  $p(p-1)$  representations of the form

$$\pi_{(1, \xi_2, 1, \xi_4)}(x) = \pi_{(\xi_2, \xi_4)}(x) := e^{2\pi i \{\xi_4(x_4 + ux_3 + \frac{1}{2}u^2x_2) + \xi_2x_2\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_4},$$

where  $1 \leq |\xi_2|_p \leq p$  and  $|\xi_4|_p = p$ . The representation space has to be

$$\mathcal{H}_{\xi_4} := \text{span}_{\mathbb{C}} \{ \varphi_h : h \in \mathbb{Z}_p/p\mathbb{Z}_p \}, \quad \varphi_h(u) := p^{1/2} \mathbb{1}_{h+p\mathbb{Z}_p}(u), \quad \dim(\mathcal{H}_{\xi_4}) = p.$$

By using this basis we can compute

$$\chi_{\pi_{(\xi_2, \xi_4)}}(x) = pe^{2\pi i \{\xi_2x_2 + \xi_4x_4\}_p} \mathbb{1}_{p\mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{\xi_4x_3u + \frac{1}{2}\xi_4x_2u^2\}_p} du,$$

where the above integral is a  $p$ -adic Gaussian integral. Using Lemma 2.2, see the calculations in [9, p.p. 65], we obtain

$$\begin{aligned} & \chi_{\pi_{(\xi_4, \xi_2)}}(x) \\ &= pe^{2\pi i \{\xi_2x_2 + \xi_4x_4\}_p} \mathbb{1}_{p\mathbb{Z}_p}(x_1) \times \begin{cases} \mathbb{1}_{\mathbb{Z}_p}(\xi_4x_3), & \text{if } |\xi_4x_2|_p \leq 1, \\ \lambda_p\left(\frac{\xi_4x_2}{2}\right) |\xi_4x_2|_p^{-1/2} e^{2\pi i \left\{-\frac{(\xi_4x_3)^2}{2\xi_4x_2}\right\}_p} \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right), & \text{if } |\xi_4x_2|_p > 1. \end{cases} \end{aligned}$$

See again Lemma 2.2. Now, considering the fact that  $|\xi_4|_p = p$ , the irreducibility of the representation follows from the calculation

$$\begin{aligned} \int_{B_4} |\chi_{\pi_{(\xi_4, \xi_2)}}(x)|^2 dx &= \int_{\mathbb{Z}_p^3} \int_{p\mathbb{Z}_p} p^2 \mathbb{1}_{p\mathbb{Z}_p}(x_1) \mathbb{1}_{p\mathbb{Z}_p}(x_3) dx_2(dx_1 dx_3 dx_4) \\ &+ \int_{\mathbb{Z}_p^3} \int_{\mathbb{Z}_p \setminus \mathbb{Z}_p} p^2 |\mathbb{1}_{p\mathbb{Z}_p}(x_1) \lambda_p\left(\frac{\xi_4x_2}{2}\right) |\xi_4x_2|_p^{-1/2} \mathbb{1}_{\mathbb{Z}_p}(x_3/x_2)|^2 dx_2(dx_1 dx_3 dx_4). \end{aligned}$$

In one hand

$$\int_{\mathbb{Z}_p^3} \int_{p\mathbb{Z}_p} p^2 \mathbb{1}_{p\mathbb{Z}_p}(x_1) \mathbb{1}_{p\mathbb{Z}_p}(x_3) dx_2(dx_1 dx_3 dx_4) = 1/p.$$

On the other

$$\begin{aligned} & \int_{\mathbb{Z}_p^3} \int_{\mathbb{Z}_p \setminus \mathbb{Z}_p} p^2 |\mathbb{1}_{p\mathbb{Z}_p}(x_1) \lambda_p\left(\frac{\xi_4x_2}{2}\right) |\xi_4x_2|_p^{-1/2} \mathbb{1}_{\mathbb{Z}_p}(x_3/x_2)|^2 dx_2(dx_1 dx_3 dx_4) = \\ & \int_{\mathbb{Z}_p^3} \int_{\mathbb{Z}_p \setminus p\mathbb{Z}_p} p^2 |\mathbb{1}_{p\mathbb{Z}_p}(x_1) p^{-1/2} |x_2|_p^{-1/2} \mathbb{1}_{p\mathbb{Z}_p}(x_3)|^2 dx_2(dx_1 dx_3 dx_4) \\ &= \int_{\mathbb{Z}_p^3} \int_{\mathbb{Z}_p \setminus \mathbb{Z}_p} p^2 |\mathbb{1}_{p\mathbb{Z}_p}(x_1) \mathbb{1}_{p\mathbb{Z}_p}(x_3)|^2 dx_2(dx_1 dx_3 dx_4) = 1 - 1/p. \end{aligned}$$

Notice how these representations are indexed by the sets

$$\begin{aligned} & B_{\hat{\beta}_4}(1) = \\ & \{ \|\xi\| \leq p : 1 \leq |\xi_4|_p < |\xi_3|_p \wedge (\xi_1, \xi_2, \xi_3) \in \hat{\mathbb{H}}_1, \text{ or } |\xi_3|_p = 1, p = |\xi_4|_p \wedge \xi_1 \in \mathbb{Q}_p/p^{-1}\mathbb{Z}_p \}, \end{aligned}$$

and we can check how these are all the elements in  $B_{\widehat{B}_4}(1) = \widehat{B}_4 \cap B_4(p^1\mathbb{Z}_p)^\perp$  since

$$\begin{aligned} & \sum_{\|(\xi_1, \xi_2)\|_p \leq p} \dim(\chi_{\xi_1, \xi_2})^2 + \sum_{|\xi_3|_p = p} \dim(\pi_{\xi_3})^2 + \sum_{|\xi_2|_p \leq |\xi_4|_p = p} \dim(\pi_{(\xi_4, \xi_2)})^2 \\ &= \sum_{\|(\xi_1, \xi_2)\|_p \leq p} 1^2 + \sum_{|\xi_3|_p = p} p^2 + \sum_{|\xi_2|_p \leq |\xi_4|_p = p} p^2 = p^4 = |G/G_1|. \end{aligned}$$

Now, for the next level, we will have to start mixing these three different kind of representations.

**3.2. Level 2:** We want to describe the elements of  $B_{\widehat{B}_4}(2) = \widehat{B}_4 \cap B_4(p^2\mathbb{Z}_p)^\perp$ . Using once again how  $B_4/\mathcal{Z}(B_4) \cong \mathbb{H}_1(\mathbb{Z}_p)$ , and the description of the unitary dual given of  $\mathbb{H}_d(\mathbb{Z}_p)$  given in [8], we will obtain the new representations in  $\widehat{B}_4(2)$  by considering three cases.

The first case is when an element of  $B_{\widehat{B}_4}(2)$  acts trivially on the center, and it must descend to a representation of  $\mathbb{H}_1(\mathbb{Z}_p)$ , so it has the form

$$\pi_\xi(x)\varphi(u) = \pi_{(\xi_1, \xi_2, \xi_3, 1)}(x)\varphi(u) := e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_3(x_3 + ux_2)\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_3},$$

where the representation space is

$$\mathcal{H}_\xi = \mathcal{H}_{\xi_3} := \text{Span}_{\mathbb{C}}\{\varphi_h : h \in \mathbb{Z}_p/|\xi_3|_p\mathbb{Z}_p\}, \quad \varphi_h(u) := |\xi_3|_p^{1/2} \mathbb{1}_{h+|\xi_3|_p\mathbb{Z}_p}(u), \quad \|\xi\|_p \leq p^2,$$

the associated character is

$$\chi_{\pi_{(\xi_1, \xi_2, \xi_3, 1)}}(x) = |\xi_3|_p e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_3 x_3\}_p} \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_1) \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_2),$$

and we clearly have  $\dim(\mathcal{H}_{\xi_3}) = |\xi_3|_p$ . For these representations it holds that

$$\sum_{\xi} d_\xi^2 = p^6 = |\mathbb{H}_1(p^2\mathbb{Z}_p)|.$$

On the other hand, if a representation is in  $B_{\widehat{B}_4}(2)$ , and it is not trivial on the center of  $B_4(\mathbb{Z}_p)$ , then we need to examine two remaining cases.

The second case we need to examine is when  $|\xi_4|_p = p^2$  and  $1 \leq |\xi_2|_p \leq p^2$ . Then we can define

$$\pi_{(\xi_2, \xi_4)}(x) := e^{2\pi i\{\xi_4(x_4 + ux_3 + \frac{1}{2}u^2 x_2) + \xi_2 x_2\}_p} \varphi(u + x_1), \quad \varphi \in \mathcal{H}_{\xi_4},$$

where  $1 \leq |\xi_2|_p \leq p^2$ , and

$$\mathcal{H}_{\xi_4} := \text{span}_{\mathbb{C}}\{\varphi_h : h \in \mathbb{Z}_p/|\xi_4|_p\mathbb{Z}_p\}, \quad \varphi_h(u) := |\xi_4|_p^{1/2} \mathbb{1}_{h+|\xi_4|_p\mathbb{Z}_p}(u), \quad \dim(\mathcal{H}_{\xi_4}) = |\xi_4|_p.$$

These representations are unitary and irreducible, since their associated character is given by

$$\begin{aligned} \chi_{\pi_{(\xi_4, \xi_2)}}(x) &= |\xi_4|_p e^{2\pi i\{\xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_1) \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_3) \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_2) \\ &+ |\xi_4|_p e^{2\pi i\{\xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{|\xi_4|_p\mathbb{Z}_p}(x_1) \cdot \Lambda\left(\frac{\xi_4 x_2}{2}, \xi_4 x_3\right) \cdot \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right) \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p}(\xi_4 x_2), \end{aligned}$$

which defines an  $L^2$ -normalized function. Here  $\Lambda(a, b)$  is defined as in Lemma 2.2. See [9, p.p. 65] for the calculations with  $p$ -adic Gaussian integrals.

Finally, for the third and last case, we have to examine mixed representations, which occur when  $|\xi_4|_p = p$ . These have the form

$$\pi_\xi(x)\varphi(u) := e^{2\pi i\{\xi \cdot x + u(\xi_3 x_2 + \xi_4 x_3) + \frac{u^2}{2} \xi_4 x_2\}_p} \varphi(u + x_1),$$

where  $1 \leq |\xi_3|_p \leq 2$ , and they can be realized in the sub-spaces  $\mathcal{H}_\xi$  of  $L^2(\mathbb{Z}_p)$  defined as

$$\mathcal{H}_\xi := \text{span}_{\mathbb{C}} \{ \varphi_h = \|\xi\|_p^{-1/2} \mathbb{1}_{h + \|\xi\|_p \mathbb{Z}_p}(u) : h \in \mathbb{Z}_p / \|\xi\|_p \mathbb{Z}_p \}.$$

Let us define here  $d_\xi := \max\{|\xi_3|_p, |\xi_4|_p\}$ . With this realization and the natural choice of basis, the associated matrix coefficients of the representations are given by

$$\begin{aligned} (\pi_\xi(x))_{hh'} &= (\pi_\xi(x)\varphi_h, \varphi_{h'})_{L^2(\mathbb{Z}_p)} \\ &= d_\xi \int_{\mathbb{Z}_p} e^{2\pi i \{ \xi \cdot x + u(\xi_3 x_2 + \xi_4 x_3) + \frac{u^2}{2} \xi_4 x_2 \}_p} \mathbb{1}_h(u + x_1) \mathbb{1}_{h'}(u) du \\ &= d_\xi e^{2\pi i \{ \xi \cdot x \}_p} \mathbb{1}_{h-h'}(x_1) \int_{h'+d_\xi \mathbb{Z}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} \}_p} du \\ &= e^{2\pi i \{ \xi \cdot x \}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)(h') + \xi_4 x_2 \frac{(h')^2}{2} \}_p} \mathbb{1}_{h-h'+d_\xi \mathbb{Z}_p}(x_1) \\ &\quad \times \left( d_\xi \int_{d_\xi \mathbb{Z}_p^d} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2 + 2uh'}{2} \}_p} \right). \end{aligned}$$

From here it is clear how

$$\begin{aligned} \chi_{\pi_\xi}(x) &= \sum_{h \in \mathbb{Z}_p / d_\xi \mathbb{Z}_p} (\pi_\xi(x))_{hh} \\ &= \sum_{h \in \mathbb{Z}_p / d_\xi \mathbb{Z}_p} d_\xi e^{2\pi i \{ \xi \cdot x + u(\xi_3 x_2 + \xi_4 x_3) + \frac{u^2}{2} \xi_4 x_2 \}_p} \mathbb{1}_{h-h'}(x_1) \int_{h'+d_\xi \mathbb{Z}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} \}_p} du \\ &= d_\xi e^{2\pi i \{ \xi \cdot x \}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} \}_p} du, \end{aligned}$$

where one can write explicitly

$$\begin{aligned} \int_{\mathbb{Z}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} \}_p} &= \mathbb{1}_{\mathbb{Z}_p} \left( \frac{\xi_4 x_2}{2} \right) \cdot \mathbb{1}_{\mathbb{Z}_p}(\xi_3 x_2 + \xi_4 x_3) \\ &\quad + \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p} \left( \frac{\xi_4 x_2}{2} \right) \cdot \Lambda \left( \frac{\xi_4 x_2}{2}, \xi_3 x_2 + \xi_4 x_3 \right) \cdot \mathbb{1}_{\mathbb{Z}_p} \left( (\xi_3 x_2 + \xi_4 x_3) / \left( \frac{\xi_4 x_2}{2} \right) \right). \end{aligned}$$

Let us analyze these characters:

**Case  $|\xi_3|_p \leq |\xi_4|_p$ :** In this case  $d_\xi = |\xi_4|_p$ . Since  $p > 2$ , we have the equality  $\mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) = \mathbb{1}_{\mathbb{Z}_p} \left( \frac{\xi_4 x_2}{2} \right)$ . Next

$$\begin{aligned} \mathbb{1}_{\mathbb{Z}_p} \left( \frac{\xi_4 x_2}{2} \right) \cdot \mathbb{1}_{\mathbb{Z}_p}(\xi_3 x_2 + \xi_4 x_3) &= \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{\mathbb{Z}_p}(\xi_3 x_2 + \xi_4 x_3) \\ &= \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{\mathbb{Z}_p}(\xi_4 x_3) \\ &= \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_3), \end{aligned}$$

where the equality before the last one holds because  $\xi_3 x_2 \in \mathbb{Z}_p$  when  $|\xi_3|_p \leq |\xi_4|_p$ . For the remaining part of the function, we have

$$(\xi_3 x_2 + \xi_4 x_3) / \left( \frac{\xi_4 x_2}{2} \right) = 2 \left( \frac{\xi_3}{\xi_4} + \frac{x_3}{x_2} \right),$$

so that actually

$$\mathbb{1}_{\mathbb{Z}_p}\left(2\left(\frac{\xi_3}{\xi_4} + \frac{x_3}{x_2}\right)\right) = \mathbb{1}_{\mathbb{Z}_p}\left(\frac{\xi_3}{\xi_4} + \frac{x_3}{x_2}\right) = \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right).$$

By using all these we can rewrite

$$\begin{aligned} \int_{\mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} &= \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_3) \\ &+ \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p}\left(\frac{\xi_4 x_2}{2}\right) \cdot e^{-2\pi i\{\frac{\xi_3^2}{2\xi_4} x_2 + \xi_3 x_3\}_p} h\left(\frac{\xi_4 x_2}{2}, \xi_4 x_3\right) \cdot \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right), \end{aligned}$$

and therefore

$$\begin{aligned} \chi_{\pi_\xi}(x) &= \\ &|\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_3 x_3 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_3) \\ &+ |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_3 x_3 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p}\left(\frac{\xi_4 x_2}{2}\right) \cdot e^{-2\pi i\{\frac{\xi_3^2}{2\xi_4} x_2 + \xi_3 x_3\}_p} h\left(\frac{\xi_4 x_2}{2}, \xi_4 x_3\right) \cdot \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right), \\ &= |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_3) \\ &+ |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + (\xi_2 - \frac{\xi_3^2}{2\xi_4})x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p}\left(\frac{\xi_4 x_2}{2}\right) \cdot h\left(\frac{\xi_4 x_2}{2}, \xi_4 x_3\right) \cdot \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right). \end{aligned}$$

So, in the end the variable  $\xi_3$  disappears, and the characters are given by

$$\begin{aligned} \chi_{\pi_\xi}(x) &= |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_2) \cdot \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_3) \\ &+ |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \mathbb{1}_{\mathbb{Q}_p \setminus \mathbb{Z}_p}\left(\frac{\xi_4 x_2}{2}\right) \cdot h\left(\frac{\xi_4 x_2}{2}, \xi_4 x_3\right) \cdot \mathbb{1}_{\mathbb{Z}_p}\left(\frac{x_3}{x_2}\right) \\ &= |\xi_4|_p e^{2\pi i\{\xi_1 x_1 + \xi_2 x_2 + \xi_4 x_4\}_p} \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i\{\xi_4 x_3 u + \xi_4 x_2 \frac{u^2}{2}\}_p} du. \end{aligned}$$

Counting all the different functions among these, we can see how we have  $(p-1)p^3$  different representations, all with dimension  $d_\xi = |\xi_4|_p = p^2$ .

**Case  $p^2 = |\xi_3|_p > |\xi_4|_p$ :** In this case  $d_\xi = |\xi_3|_p$  and  $|\xi_1|_p = |\xi_2|_p = 1$ , so that  $\xi_1$  and  $\xi_2$  are not appearing. The associated character is

$$\begin{aligned} \chi_{\pi_\xi}(x) &= |\xi_3|_p e^{2\pi i\{\xi_3 x_3 + \xi_4 x_4\}_p} \mathbb{1}_{|\xi_3|_p \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du, \\ &= |\xi_3|_p e^{2\pi i\{\xi_3 x_3 + \xi_4 x_4\}_p} \mathbb{1}_{|\xi_3|_p \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du. \end{aligned}$$

Counting all the different functions among these, we can see how we have  $(p-1)^2 p$  different representations, all with dimension  $d_\xi = |\xi_3|_p$ .

Summing up, all the obtained representations can be indexed by the set

$$\begin{aligned} &\mathbf{B}_{\widehat{B}_4}(2) \\ &= \{\|\xi\| \leq p^2 : 1 \leq |\xi_4|_p < |\xi_3|_p, (\xi_1, \xi_2, \xi_3) \in \widehat{\mathbb{H}}_1, \text{ or } |\xi_3|_p = 1, \xi_1 \in \mathbb{Q}_p / |\xi_4|_p^{-1} \mathbb{Z}_p, \xi_2 \in \widehat{\mathbb{Z}}_p\} \end{aligned}$$



To convince ourselves that these representations are indeed irreducible we just apply Lemma 2.3 to see that

$$\int_{\mathcal{B}_4} |\chi_{\pi_\xi}(x)|^2 dx = d_\xi^2 \int_{\mathbb{Z}_p^4} \left| \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right|^2 dx = 1.$$

Notice how these are indeed all the desired representations since

$$\begin{aligned} \sum_{\xi \in \mathcal{B}_{\hat{\mathcal{B}}_4}(2)} d_\xi^2 &= \sum_{\|\xi\|_p \leq p^2, (\xi_1, \xi_2, \xi_3) \in \hat{\mathbb{H}}_1} d_\xi^2 + \sum_{|\xi_4|_p = p \wedge |\xi_3|_p = 1} d_\xi^2 + \sum_{|\xi_4|_p = p \wedge |\xi_3|_p = p^2} d_\xi^2 + \sum_{|\xi_4|_p = p^2} d_\xi^2 \\ &= p^6 + \sum_{|\xi_4|_p = p \wedge |\xi_3|_p = 1} p^2 + \sum_{|\xi_4|_p = p \wedge |\xi_3|_p = p^2} (p^2)^2 + \sum_{|\xi_4|_p = p^2} (p^2)^2 \\ &= p^6 + (p-1)p^3(p^2) + (p-1)^2 p(p^2)^2 + p^3(p-1)(p^2)^2 \\ &= p^6 + (p-1)p^5 + (p-1)^2 p^7 + (p-1)p^7 \\ &= p^6 + (p-1)p^5 + (p-1)^2 p^5 + (p-1)p^7 \\ &= p^8 = (p^2)^4 = |\mathcal{B}_4(\mathbb{F}_{p^2})| = |G/G_2|. \end{aligned}$$

**3.3. General case and Fourier series.** We can use all the ideas collected so far to deduce the general case. First, we can see how the representations we are looking for must have the form

$$\pi_\xi(x)\varphi(u) := e^{2\pi i \{ \xi_1 x_1 + \xi_2 x_2 + \xi_3(x_3 + ux_2) + \xi_4(x_4 + ux_3 + \frac{u^2}{2}x_2) \}_p} \varphi(u + x_1) = e^{2\pi i \{ \xi \cdot x + u(\xi_3 x_2 + \xi_4 x_3) + \frac{u^2}{2} \xi_4 x_2 \}_p} \varphi(u + x_1),$$

where we are taking the function  $\varphi$  to be an element of the following sub-space of  $L^2(\mathbb{Z}_p)$  :

$$\mathcal{H}_\xi := \text{span}_{\mathbb{C}} \{ d_\xi^{1/2} \mathbb{1}_{h + d_\xi \mathbb{Z}_p} : h \in \mathbb{Z}_p / d_\xi \mathbb{Z}_p \}, \quad d_\xi := \dim_{\mathbb{C}}(\mathcal{H}_\xi) = \max\{ |\xi_3|_p, |\xi_4|_p \}.$$

One can easily check how these are all unitary representations of  $\mathcal{B}_4$ , for any  $\xi \in \hat{\mathbb{Z}}_p^4$ , and their associated characters are going to have the form

$$\chi_{\pi_\xi}(x) = d_\xi e^{2\pi i \{ \xi \cdot x \}_p} \mathbb{1}_{d_\xi \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{ (\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2} \}_p} du.$$

Let us use these to count all the different non-equivalent representations in  $\mathcal{B}_{\hat{\mathcal{B}}_4}(n) := \hat{\mathcal{B}}_4 \cap \mathcal{B}_4(p^n \mathbb{Z}_p)^\perp$ :

- If  $|\xi_4|_p = 1$ , then the representation is a Heisenberg representation, and for these it holds that

$$\sum_{\{\|\xi\|_p \leq p^n : |\xi_4|_p = 1\}} d_\xi^2 = p^{3n} = |\mathbb{H}_1(\mathbb{Z}_p)/\mathbb{H}_1(p^n \mathbb{Z}_p)| = |\mathbb{H}_1(\mathbb{F}_{p^n})|.$$

- We saw before how in the case  $|\xi_3|_p \leq |\xi_4|_p$  the character reduces to

$$\chi_{\pi_\xi}(x) = |\xi_4|_p e^{2\pi i \{ \xi_1 x_1 + \xi_2 x_2 + \xi_4 x_4 \}_p} \mathbb{1}_{|\xi_4|_p \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{ \xi_4 x_3 u + \xi_4 x_2 \frac{u^2}{2} \}_p} du.$$

So they all have dimension  $|\xi_4|_p$ , and we can index all the non-equivalent classes among these with the set

$$\{\|\xi\|_p \leq p^n : 1 < |\xi_4|_p \leq p^n, |\xi_3|_p = 1, \xi_1 \in \mathbb{Q}_p / |\xi_4|_p^{-1} \mathbb{Z}_p\}.$$

- If  $1 < |\xi_4|_p < |\xi_3|_p$ , the character is going to be

$$\chi_{\pi_\xi}(x) = |\xi_3|_p e^{2\pi i \{ \xi_1 x_1 \xi_2 x_2 + \xi_3 x_3 + \xi_4 x_4 \}_p} \mathbb{1}_{|\xi_3|_p \mathbb{Z}_p}(x_1) \int_{\mathbb{Z}_p} e^{2\pi i \{ \xi_4 x_3 u + \xi_4 x_2 \frac{u^2}{2} \}_p} du.$$

We can index all the non-equivalent classes among these with the set

$$\{ \|\xi\|_p \leq p^n : 1 < |\xi_4|_p < |\xi_3|_p, (\xi_1, \xi_2, \xi_3) \in \widehat{\mathbb{H}}_1 \}.$$

In total, we can index our representations with the set

$$\widehat{B}_4 := \{ \xi \in \widehat{\mathbb{Z}}_p^4 : 1 \leq |\xi_4|_p < |\xi_3|_p \wedge (\xi_1, \xi_2, \xi_3) \in \widehat{\mathbb{H}}_1, \text{ or } |\xi_3|_p = 1 \wedge \xi_1 \in \mathbb{Q}_p / |\xi_3|_p^{-1} \mathbb{Z}_p \}.$$

These are all the desired representations since

$$\begin{aligned} \sum_{\{\xi \in B_4 : \|\xi\|_p \leq p^n\}} d_\xi^2 &= \sum_{\{\|\xi\|_p \leq p^n : |\xi_4|_p = 1\}} d_\xi^2 + \sum_{\{\|\xi\|_p \leq p^n : |\xi_4|_p > 1, |\xi_3|_p = 1\}} d_\xi^2 + \sum_{\{\|\xi\|_p \leq p^n : 1 < |\xi_4|_p < |\xi_3|_p\}} d_\xi^2 \\ &= p^{3n} + \sum_{1 < |\xi_4|_p \leq p^n} \sum_{|\xi_4|_p < |\xi_3|_p \leq p^n} \sum_{(\xi_1, \xi_2) \in \mathbb{Q}_p^2 / |\xi_3|_p^{-1} \mathbb{Z}_p^2} |\xi_3|_p^2 \\ &\quad + \sum_{1 < |\xi_4|_p \leq p^n} \sum_{|\xi_2|_p \leq p^n} \sum_{\xi_1 \in \mathbb{Q}_p / |\xi_4|_p^{-1} \mathbb{Z}_p} |\xi_4|_p^2 \\ &= p^{3n} + \sum_{1 < |\xi_4|_p \leq p^n} \sum_{|\xi_4|_p < |\xi_3|_p \leq p^n} |\xi_3|_p^2 (p^{2n} |\xi_3|_p^{-2}) + \sum_{1 < |\xi_4|_p \leq p^n} |\xi_4|_p^2 (p^n) (p^n |\xi_4|_p^{-1}) \\ &= p^{3n} + p^{2n} \left( \sum_{1 < |\xi_4|_p \leq p^n} \sum_{|\xi_4|_p < |\xi_3|_p \leq p^n} 1 + \sum_{1 < |\xi_4|_p \leq p^n} |\xi_4|_p \right) \\ &= p^{3n} + p^{2n} \left( \sum_{1 < |\xi_4|_p \leq p^n} (p^n - |\xi_4|_p) + \sum_{1 < |\xi_4|_p \leq p^n} |\xi_4|_p \right) \\ &= p^{3n} + p^{3n} \left( \sum_{1 < |\xi_4|_p \leq p^n} 1 \right) \\ &= p^{3n} + p^{3n} (p^n - 1) = p^{4n} = |B_4 / B_4(p^n \mathbb{Z}_p)|. \end{aligned}$$

Finally to conclude this section, notice how from the analysis done so far, and the Fourier analysis in general compact groups, we obtain that

$$f(x) = \sum_{\xi \in \widehat{B}_4} \|(\xi_3, \xi_4)\|_p \text{Tr}[\pi_\xi(x) \widehat{f}(\xi)], \quad f \in L^2(B_4),$$

where

$$\widehat{f}(\xi) \varphi = F_{B_4}[f](\xi) \varphi := \int_{B_4} f(x) \pi_\xi^*(x) dx, \quad f \in L^2(B_4).$$

In this way, the action of any densely defined left-invariant operator on  $\mathcal{D}(B_4)$  can be expressed as

$$Tf(x) = \sum_{\xi \in \widehat{B}_4} \|(\xi_3, \xi_4)\|_p \text{Tr}[\pi_\xi(x) \sigma_T(\xi) \widehat{f}(\xi)],$$

where the symbol  $\sigma_T$  of the operator  $T$  is the mapping

$$\sigma_T : \widehat{\mathcal{B}}_4 \rightarrow \bigcup_{\xi \in \widehat{\mathcal{B}}_4} \mathcal{L}(\mathcal{H}_\xi), \text{ defined as } \sigma_T(\xi) := \pi_\xi^*(x)T\pi_\xi(x) \in \mathcal{L}(\mathcal{H}_\xi).$$

We will use all this information in the next section, where we apply it to study an important example of a left invariant operator on  $\mathcal{B}_4$  which we call *the Vladimirov sub-Laplacian*.

#### 4. THE VLADIMIROV SUB-LAPLACIAN ON THE ENGEL GROUP

We already calculated

$$\begin{aligned} (\pi_\xi(x))_{hh'} &= e^{2\pi i\{\xi \cdot x\}_p} \mathbb{1}_{h-h'+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p}(x_1) \left( \|(\xi_3, \xi_4)\|_p \int_{h'+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right) \\ &= e^{2\pi i\{\xi \cdot x\}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)(h') + \xi_4 x_2 \frac{(h')^2}{2}\}_p} \mathbb{1}_{h-h'+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p}(x_1) \\ &\quad \times \left( \|(\xi_3, \xi_4)\|_p \int_{\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2 + 2uh'}{2}\}_p} du \right). \end{aligned}$$

In order to prove Theorem 1.7 we want to use this calculate the associated symbol of  $\mathcal{L}_{sub}^\alpha$ , and its respective invariant sub-spaces. Let us start by computing the symbols of the directional VT operators  $\partial_{X_1}^\alpha, \partial_{X_2}^\alpha, \alpha > 0$ , by using what we got for the matrix coefficients. For  $\partial_{X_2}^\alpha$  we have that its associated symbol  $\sigma_{\partial_{X_2}^\alpha}(\xi) = \partial_{X_2}^\alpha \pi_\xi|_{x=0}$  is the diagonal matrix

$$\begin{aligned} &\int_{\mathbb{Z}_p} \frac{\pi_\xi(0, x_2, 0, 0) - I_{d_\xi}}{|x_2|_p^{\alpha+1}} dx_2 \\ &= \text{Diag} \left( \int_{\mathbb{Z}_p} \frac{1}{|x_2|_p^{\alpha+1}} \left( d_\xi \int_{h+d_\xi \mathbb{Z}_p} e^{2\pi i\{\xi_2 x_2 + \xi_3 x_2 u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right) - 1 \right) dx_2 : h \in \mathbb{Z}_p / d_\xi \mathbb{Z}_p \\ &= \text{Diag} \left( d_\xi \int_{h+d_\xi \mathbb{Z}_p} \int_{\mathbb{Z}_p} \frac{e^{2\pi i\{\xi_2 x_2 + \xi_3 x_2 u + \xi_4 x_2 \frac{u^2}{2}\}_p} - 1}{|x_2|_p^{\alpha+1}} dx_2 du : h \in \mathbb{Z}_p / d_\xi \mathbb{Z}_p \right) \\ &= \text{Diag} \left( \|(\xi_3, \xi_4)\|_p \int_{h+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p} |\xi_2 + \xi_3 u + \xi_4 \frac{u^2}{2}|_p^\alpha du : h \in \mathbb{Z}_p / \|(\xi_3, \xi_4)\|_p \mathbb{Z}_p \right) \\ &= \text{Diag} \left( |\xi_2 + \xi_3 h + \xi_4 \frac{h^2}{2}|_p^\alpha : h \in \mathbb{Z}_p / \|(\xi_3, \xi_4)\|_p \mathbb{Z}_p \right). \end{aligned}$$

Now we consider the different cases:

$$\sigma_{\partial_{X_2}^\alpha}(\xi)_{hh} = \begin{cases} |\xi_2|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } |\xi_3|_p = |\xi_4|_p = 1, \\ |\xi_2 + \xi_3 h|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } 1 = |\xi_4|_p < |\xi_3|_p, h \in \mathbb{Z}_p / |\xi_3|_p \mathbb{Z}_p, \\ |\xi_2 + \xi_3 h + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } 1 < |\xi_4|_p < |\xi_3|_p, h \in \mathbb{Z}_p / |\xi_3|_p \mathbb{Z}_p, \\ |\xi_2 + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } 1 = |\xi_3|_p < |\xi_4|_p, h \in \mathbb{Z}_p / |\xi_4|_p \mathbb{Z}_p. \end{cases}$$

For  $\partial_{X_1}^\alpha$ , let us introduce some notation: let  $\varphi_\xi : \mathbb{Z}_p \rightarrow \mathbb{C}^{d_\xi \times d_\xi}$  be the matrix function with entries

$$(\varphi_\xi)_{hh'}(u) := e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p}(u), \quad u \in \mathbb{Z}_p.$$

The associated associated symbol  $\sigma_{\partial_{X_1}^\alpha}(\xi) = \partial_{X_1}^\alpha \pi_\xi|_{x=0}$  is a Toeplitz matrix which can be written as

$$\sigma_{\partial_{X_1}^\alpha}(\xi)_{hh'} = \begin{cases} |\xi_2|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } |\xi_3|_p = |\xi_4|_p = 1, \\ \partial_u^\alpha(e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+|\xi_3|_p \mathbb{Z}_p})|_{u=0} & \text{if } |\xi_4|_p < |\xi_3|_p, h \in \mathbb{Z}_p/|\xi_3|_p \mathbb{Z}_p, \\ \partial_u^\alpha(e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+|\xi_4|_p \mathbb{Z}_p})|_{u=0} & \text{if } 1 = |\xi_3|_p < |\xi_4|_p, h \in \mathbb{Z}_p/|\xi_4|_p \mathbb{Z}_p, \end{cases}$$

where  $\partial_u^\alpha$  is the VT operator

$$\partial_u^\alpha \psi(u) := \int_{\mathbb{Z}_p} \frac{\psi(u-t) - \psi(u)}{|t|_p^{\alpha+1}} dt, \quad \psi \in \mathcal{D}(\mathbb{Z}_p).$$

Summing up, the symbol of the Vladimirov Sub-Laplacian  $\mathcal{L}_{sub}^\alpha$  is:

$$\sigma_{\mathcal{L}_{sub}^\alpha}(\xi) = \begin{cases} |\xi_1|_p^\alpha + |\xi_2|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } |\xi_3|_p = |\xi_4|_p = 1, \\ (\partial_u^\alpha + |\xi_2 + \xi_3 h|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})\psi_\xi(u)|_{u=0}, & \text{if } 1 = |\xi_4|_p < |\xi_3|_p, \\ (\partial_u^\alpha + |\xi_2 + \xi_3 h + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})\psi_\xi(u)|_{u=0}, & \text{if } 1 < |\xi_4|_p < |\xi_3|_p, \\ (\partial_u^\alpha + |\xi_2 + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})\psi_\xi(u)|_{u=0}, & \text{if } 1 = |\xi_3|_p < |\xi_4|_p, \end{cases}$$

or component-wise  $\sigma_{\mathcal{L}_{sub}^\alpha}(\xi)_{hh'} =$

$$\begin{cases} |\xi_1|_p^\alpha + |\xi_2|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}} & \text{if } |\xi_3|_p = |\xi_4|_p = 1, \\ (\partial_u^\alpha + |\xi_2 + \xi_3 h|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})(e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+|\xi_3|_p \mathbb{Z}_p})(u)|_{u=0}, & \text{if } 1 = |\xi_4|_p < |\xi_3|_p, \\ (\partial_u^\alpha + |\xi_2 + \xi_3 h + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})(e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+|\xi_3|_p \mathbb{Z}_p})(u)|_{u=0}, & \text{if } 1 < |\xi_4|_p < |\xi_3|_p, \\ (\partial_u^\alpha + |\xi_2 + \xi_4 \frac{h^2}{2}|_p^\alpha - \frac{1-p^{-1}}{1-p^{-(\alpha+1)}})(e^{2\pi i\{\xi_1 u\}_p} \mathbb{1}_{h-h'+|\xi_4|_p \mathbb{Z}_p})(u)|_{u=0}, & \text{if } 1 = |\xi_3|_p < |\xi_4|_p, \end{cases}$$

Let us decompose now each space  $\mathcal{V}_\xi := \text{span}_{\mathbb{C}}\{(\pi_\xi(x))_{hh'} : h, h' \in \mathbb{Z}_p/d_\xi \mathbb{Z}_p\}$  in the direct sum

$$\mathcal{V}_\xi = \bigoplus_{h' \in \mathcal{I}_\xi} \mathcal{V}_\xi^{h'},$$

where

$$\mathcal{V}_\xi^{h'} := \text{span}_{\mathbb{C}}\{(\pi_\xi)_{hh'} : h \in \mathbb{Z}_p/\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p\}.$$

Functions in this space can actually be described as

$$\mathcal{V}_\xi^{h'} = \left\{ e^{2\pi i\{\xi \cdot x\}_p} \varphi(x_1) \left( \|(\xi_3, \xi_4)\|_p \int_{h'+\|(\xi_3, \xi_4)\|_p \mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right) : \varphi \in \mathcal{D}_{d_\xi}(\mathbb{Z}_p) \right\},$$

where we are using the notation

$$\mathcal{D}_{d_\xi}(\mathbb{Z}_p^d) := \{ \varphi \in \mathcal{D}(\mathbb{Z}_p^d) : \varphi(u+v) = \varphi(u), |v|_p \leq d_\xi^{-1} \}, \quad \dim_{\mathbb{C}}(\mathcal{D}_n(\mathbb{Z}_p)) = \|(\xi_3, \xi_4)\|_p.$$

For this space, we have the alternative choice of basis

$$e_{\xi, h', \tau}(x) := e^{2\pi i\{\xi x + \tau x_1\}_p} \left( d_\xi \int_{h'+d_\xi \mathbb{Z}_p} e^{2\pi i\{(\xi_3 x_2 + \xi_4 x_3)u + \xi_4 x_2 \frac{u^2}{2}\}_p} du \right), \quad 1 \leq \tau \leq \|(\xi_3, \xi_4)\|_p,$$

and for these functions we can calculate  $(\partial_{X_1}^\alpha + \partial_{X_2}^\alpha)e_{\xi, h', \tau}(x) =$

$$\begin{cases} (|\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_3 h'|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}})e_{\xi, h', \tau}(x) & \text{if } 1 = |\xi_4|_p < |\xi_3|_p, \\ (|\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_3 h' + \xi_4 \frac{(h')^2}{2}|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}})e_{\xi, h', \tau}(x) & \text{if } 1 < |\xi_4|_p < |\xi_3|_p, \\ (|\xi_1 + \tau|_p^\alpha + |\xi_2 + \xi_4 \frac{(h')^2}{2}|_p^\alpha - 2\frac{1-p^{-1}}{1-p^{-(\alpha+1)}})e_{\xi, h', \tau}(x) & \text{if } 1 = |\xi_3|_p < |\xi_4|_p. \end{cases}$$

So, the Vladimirov sub-Laplacian acts on each finite-dimensional sub-space  $\mathcal{V}_\xi^{h'}$  like the Schrödinger-type operator  $\partial_{x_1}^\alpha + Q(\xi, h)$ , or more precisely, in this special sub-spaces it is just a translation of the directional Vladimirov-Taibleson operator in the direction of  $x_1$ . Moreover, the functions  $e_{\xi, h', \tau}(x)$  form a complete system of eigenfunctions in  $V_\xi^{h'}$ , proving in this way how

$$\|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi^{h'}}\|_{op} = \max_{\tau, h'} \lambda_{\xi, h', \tau}(\mathcal{L}_{sub}^\alpha) \lesssim \|\xi\|_p^\alpha.$$

In the same way

$$\|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi^{h'}}\|_{inf} = \min_{\tau, h'} \lambda_{\xi, h', \tau}(\mathcal{L}_{sub}^\alpha) \gtrsim |\xi_1|_p^\alpha + |\xi_2|_p^\alpha.$$

This implies that in each representation space we have the estimates

$$|\xi_1|_p^\alpha + |\xi_2|_p^\alpha \lesssim \|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi}\|_{inf} \leq \|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi}\|_{op} \lesssim \|\xi\|_p^\alpha,$$

but we know that

$$\|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi}\|_{inf} = \|\sigma_{\mathcal{L}_{sub}^\alpha}(\xi)\|_{inf}, \quad \|\mathcal{L}_{sub}^\alpha|_{\mathcal{V}_\xi}\|_{op} = \|\sigma_{\mathcal{L}_{sub}^\alpha}(\xi)\|_{op},$$

proving how  $\mathcal{L}_{sub}^\alpha$  provides an example of an operator in the class we define next:

**Definition 4.1.** Let  $\sigma$  be an invariant symbol, and let  $T_\sigma$  be its associated pseudo-differential operator. We say that  $\sigma$  is an *elliptic symbol of order*  $m \in \mathbb{R}$ , if there are  $C_1, C_2 > 0$  and  $m \in \mathbb{R}$  such that

$$C_1 \|\xi\|_p^m \leq \|\sigma(\xi)\|_{inf} \leq \|\sigma(\xi)\|_{op} \leq C_2 \|\xi\|_p^m,$$

for  $\|\xi\|_p$  large enough. Similarly, if there is an  $0 < \delta \leq m$  such that

$$C_1 \|\xi\|_p^{m-\delta} \leq \|\sigma(\xi)\|_{inf} \leq \|\sigma(\xi)\|_{op} \leq C_2 \|\xi\|_p^m,$$

for  $\|\xi\|_p$  large enough, we say that  $\sigma$  is a *sub-elliptic symbol*. We say that an invariant operator  $T_\sigma \in \mathcal{L}(H_2^{s+m}(\mathcal{B}_4), H_2^s(\mathcal{B}_4))$  is elliptic or sub-elliptic, if its associated symbol is elliptic or sub-elliptic, respectively.

The collection of the results proved so far constitute the statement of Theorem 1.7, whose proof is now completed.

## 5. ACKNOWLEDGEMENTS

The author wants to thank Lea Schmitt for her unconditional support during the time of the development of the ideas that led to this work.

## REFERENCES

- [1] BENDIKOV, A., GRIGORYAN, A., PITTET, C., AND WOESS, W. Isotropic Markov semigroups on ultra-metric spaces. *Russian Mathematical Surveys* 69, 4 (Aug. 2014), 589–680.
- [2] BRAMANTI, M. *An Invitation to Hypoelliptic Operators and Hörmander’s Vector Fields*. Springer International Publishing, 2014.
- [3] CORWIN, L., AND GREENLEAF, F. P. *Cambridge studies in advanced mathematics: Volume 1 representations of nilpotent Lie groups and their applications: Series number 18: Basic theory and examples part 1*. Cambridge University Press, Cambridge, England, June 2004.
- [4] DRAGOVICH, B., KHRENNIKOV, A., KOZYREV, S., AND VOLOVICH, I. On the occasion of the centenary of the birth of V. S. Vladimirov (1923–2012). *p-Adic Numbers, Ultrametric Analysis and Applications* 15, 3 (Sept. 2023), 243–244.
- [5] DRAGOVICH, B., KHRENNIKOV, A., KOZYREV, S., VOLOVICH, I., AND ZELENOV, E. p-adic mathematical physics: the first 30 years. *p-Adic Numbers, Ultrametric Analysis and Applications* 9, 2 (Apr. 2017), 87–121.
- [6] HÖRMANDER, L. Hypoelliptic second order differential equations. *Acta Mathematica* 119, 0 (1967), 147–171.
- [7] RUZHANSKY, M., AND TURUNEN, V. *Pseudo-Differential Operators and Symmetries*. Birkhäuser Basel, 2010.
- [8] VELASQUEZ-RODRIGUEZ, J. The spectrum of the Vladimirov sub-Laplacian on the compact Heisenberg group. *Researchgate* (01 2024).
- [9] VLADIMIROV, V., VOLOVICH, I., AND ZELENOV, E. *P-adic analysis and mathematical physics*. Series on Soviet and East European mathematics, V.1. World Scientific, 1994.

J.P. Velasquez-Rodriguez, DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS, GHENT UNIVERSITY, BELGIUM, AND DEPARTAMENTO DE MATEMATICAS, UNIVERSIDAD DEL VALLE, CALI-COLOMBIA

*E-mail address:* [juanpablo.velasquezrodriguez@ugent.be](mailto:juanpablo.velasquezrodriguez@ugent.be) / [velasquez.juan@correounivalle.edu.co](mailto:velasquez.juan@correounivalle.edu.co)