

**CORRECTIONS TO A PAPER OF ALLARD AND ALMGREN
ON THE UNIQUENESS OF TANGENT CONES**

WILLIAM K. ALLARD

CONTENTS

1. Introduction.	2
2. Preliminaries.	2
2.1. Some linear algebra.	2
2.2. A useful notational convention.	3
3. Submanifolds of Euclidean spaces.	4
3.1.	4
3.2. Smooth functions on M .	4
3.3. More on differentiation.	6
3.4. Diffeomorphisms.	6
3.5. \mathcal{L}_m , \mathcal{P}_m and \mathcal{W}_m .	7
3.6. The second fundamental form of M .	7
4. Vector bundles.	8
4.1. The mean curvature normal of M .	10
5. Reach.	11
5.1.	13
5.2. θ and ι .	15
6. η and \mathbf{o} .	15
7. f .	16
7.1. γ .	17
7.2. F .	18
8. Families of submanifolds of \mathbb{S}^{n-1} .	18
8.1. ϵ_1 .	19
8.2. \underline{M} .	22
8.3. $\underline{\tau}$, $\underline{\nu}$, $\underline{\gamma}$, \underline{B} , \underline{A} .	23
8.4. \underline{N} , $\underline{\Xi}$, $\underline{\xi}$, $\underline{\sigma}$.	23
8.5. \underline{f} , $\underline{\nu}$, $\underline{\rho}$, \underline{V} , \underline{g} .	24
9. F_s^t .	24
9.1. ϵ_2 and \mathcal{G} .	26
9.2. \mathcal{L}_m , \mathcal{W}_m and ϵ_3 .	28
10. The family of interest.	29
10.1. The symmetric quadratic functionals P, Q, R, S .	30
10.2. Φ ; Ψ ; \mathbf{L} ; and E_λ , $\lambda \in \mathbb{R}$.	30
10.3. The Hölder spaces $\mathcal{C}^{k,\alpha}(\gamma)$.	30
10.4. Construction of the family of interest.	31

Date: July 10, 2024.

Supported in part by Los Alamos National Laboratory.

10.5. Symmetric bilinear functions.	35
11. L_s^t and ϵ_4 .	35
12. ϵ_5 .	39
12.1. Π_s .	40
12.2. ϵ_6 .	41
13. Proof of a statement similar to 5.4(4) of [AA81].	42
13.1. $\zeta_{Z,s}$.	42
13.2. Finding a zero of $\zeta_{Z,s}$.	44
References	45

1. INTRODUCTION.

The paper referred to in the title is [AA81].

Several months ago Francesco Maggi emailed me saying that the inequalities 5.3(4),(5) of [AA81] were wrong. In fact, as he pointed out, their incorrectness is immediately apparent if one takes $Z = 0$ there. Maggi and his coauthor wanted to use these inequalities in their paper [MN]. They were able to obtain a version of these inequalities which suffice for the carrying out the work in [MN].

I started writing this paper in order to provide a version of these inequalities as needed in [AA81]. In thinking about this material I began to realize there were other problems with the paper. As a result I ended up *completely rewriting 5.1-5.4 on pages 243-248 of [AA81]*; this rewrite is the contents of this paper. In addition to many annoying misprints many of the needed definitions and proofs in 5.1-5.4 are incomplete or absent. For example, it is not said where z in 5.1(2) comes from; this omission completely surprised me since I remember doing a lot of work to come up with z when [AA81] was being written. Also, much of the necessary material about the reach of a submanifold as in [FE2] is not provided in [AA81].

The table of contents can serve as an index. In particular one sees there where the constants ϵ_1 through ϵ_6 are introduced.

This material is extremely technical. One way to navigate this paper would be to start looking at Proposition 9.5 and work backwards.

2. PRELIMINARIES.

If X is a set we let

$$i_X = \{(x, x) : x \in X\};$$

thus $i_X : X \rightarrow X$ and $i_X(x) = x$ for $x \in X$; we call i_X the **identity map of X** ; if $X \subset Y$ the i_X is also called the **inclusion map of X into Y** .

If (X, ρ) and (Y, σ) are metric spaces and $f : X \rightarrow Y$ we let

$$\mathbf{Lip} f = \sup \left\{ \frac{\sigma(f(x_1), f(x_2))}{\rho(x, y)} : x_i \in X, i = 1, 2, \text{ and } x_1 \neq x_2 \right\}.$$

2.1. **Some linear algebra.** be the vector space of linear maps from V to W . We let

$$\mathbf{End}(V) = \mathbf{Lin}(V, V).$$

If V_1, \dots, V_N and W are vector spaces we let

$$\mathbf{MultiLin}(V_1, \dots, V_N, W)$$

be the vector space of multilinear maps from $V_1 \times \cdots \times V_N$ into W .

If V and W are vector spaces we let

$$\mathbb{L}\text{in}(V, W) = \mathbb{M}\text{ulti}\mathbb{L}\text{in}(V, W).$$

If V is a vector space we let

$$\mathbb{E}\text{nd}(V) = \mathbb{L}\text{in}(V, V).$$

A **Euclidean space** is a finite dimensional inner product space. If E is a Euclidean space and $l \in \mathbb{N}$ we let

$$\mathbb{G}_l(E) \text{ and } \mathbb{O}_l(E)$$

be, respectively, the family of l dimensional linear subspaces of E and the family of orthogonal projections of E onto a member of $\mathbb{G}_l(E)$.

If E_i , $i = 1, \dots, m$ and F are Euclidean spaces and $\mu : E_1 \times \cdots \times E_m \rightarrow F$ is multilinear we let

$$\|\mu\| = \sup\{|\mu(v_1, \dots, v_m)| : v_i \in E_i \text{ and } |v_i| \leq 1, i = 1, \dots, m\};$$

the vector space of multilinear maps from $E_1 \times \cdots \times E_m$ into F has a natural Euclidean structure and the norm with respect to this structure of μ as above will be larger than $\|\cdot\|$ unless the E_i are one dimensional. For example, if i is the identity map of the Euclidean space E then $\|i\| = 1$ but the Euclidean norm of i is the square root of the dimension of E .

Proposition 2.1. *Suppose $S \in \mathbb{G}_m(\mathbb{R}^n)$, $L \in \mathbb{L}\text{in}(S, \mathbb{R}^n)$, $\ker L = \{0\}$. and $T = \text{rng } L$. and $\dim T = m$.*

There are unique A and O such that $A \in \mathbb{E}\text{nd}(T)$, $O \in \mathbb{L}\text{in}(S, T)$, $A^ = A$, $A(v) \bullet v > 0$ for $v \in T$, $A^2 = L \circ L^*$, O is an isometry and $L = A \circ O$.*

Proof. $L \circ L^*$ is a positive definite and carries T isomorphically onto T . Let A be the unique positive definite square root of $L \circ L^*$ and let $O = A^{-1} \circ L$. \square

Suppose S and T are finite dimensional inner product spaces. For $L, M \in \mathbb{L}\text{in}(S, T)$

$$L \bullet M = \text{trace } L^* \circ M.$$

This is an inner product on $\mathbb{L}\text{in}(S, T)$. If u_1, \dots, u_m is an orthonormal basis for S we have

$$L \bullet M = \sum_{i=1}^m L(u_i) \bullet M(u_i).$$

It follows that

$$|L \bullet M| \leq m \|L\| \|M\|;$$

note that if L is an isometry then $\|L\| = 1$ and $|L| = \sqrt{m}$.

2.2. A useful notational convention. Suppose T , X and Y are nonempty sets and $W \subset T \times X$.

If $h : T \times X \rightarrow Y$ and $t \in \mathcal{J}$ we let

$$h_t = \{(x, h(t, x)) : (t, x) \in W\}.$$

Thus

$$h_t : \{x : (t, x) \in W\} \rightarrow Y \quad \text{for } t \in \mathcal{J};$$

of course $h_t = \emptyset$ if $W \cap (\{t\} \times X) = \emptyset$.

3. SUBMANIFOLDS OF EUCLIDEAN SPACES.

Definition 3.1. Suppose E is a Euclidean space, $m \in \mathbb{N}^+$ and $m \leq \dim E$. We let

$$\mathcal{M}_m(E)$$

be the family of nonempty subsets M of E such that for each $a \in M$ there are U, V, ψ, ϕ such that V is an open subset of E ; U is an open subset of \mathbb{R}^n , $\psi : V \rightarrow U$, ψ is smooth, $\phi : U \rightarrow V$, ϕ is smooth, $a \in V$ and

$$\psi(\phi(u)) = u \quad \text{for } u \in U.$$

Remark 3.1. This is a slight variation of the definition in [FE, 3.1.19(4)]. Note that $\phi = (\psi|_M)^{-1}$

A member of $\mathcal{M}_m(E)$ is called a **smooth m dimensional submanifold of E** .

Evidently, $M \in \mathcal{M}_{\dim E}(E)$ if and only if M is an open subset of E . Also, if $L \subset M$ then $L \in \mathcal{M}_m(E)$ if and only if L is nonempty and is open relative to M .

It is elementary that

$$\mathbf{Tan}M, \phi(u) = \mathbf{rng} D\phi(u) \in \mathbb{G}_m(E) \quad \text{for } u \in U.$$

It follows that m is determined by M ; we call it the **dimension** of M . We let

$$\tau : M \rightarrow \mathbb{G}_m(E) \quad \text{and} \quad \nu : M \rightarrow \mathbb{G}_{\dim E - m}(E)$$

be the **tangent map** and the **normal map** of M , respectively; by definition, $\tau(x)$ is orthogonal projection of E onto $\mathbf{Tan}M, x$ and $\nu(x) = \tau(x)^\perp$ for $x \in M$.

We now fix $m, n \in \mathbb{N}^+$ with $m \leq n$.

3.1. For the remainder of this Section we fix $M \in \mathcal{M}_m(\mathbb{R}^n)$ and a normed vector space Y .

3.2. **Smooth functions on M .** A function $f : M \rightarrow Y$ is **smooth** if $f \circ \phi$ is smooth if whenever a, U, V, ϕ, ψ are as in Definition 3.1.

$$\mathcal{E}(M, Y)$$

be the vector space of smooth maps from M into Y .

Proposition 3.1. Suppose Y is a Euclidean space, $f : M \rightarrow Y$, f is smooth and $a \in M$. There is and only one

$$Df : M \rightarrow \mathbb{L}\text{in}(\mathbb{R}^n, Y)$$

such that if $a \in M$ then

$$Df(a)(D\phi(\psi(a))(v)) = D(f \circ \phi)(\psi(a))(v) \quad \text{for } v \in \mathbb{R}^n$$

and

$$Df(a) \circ \nu(a) = 0.$$

Moreover, if N is a smooth submanifold of some Euclidean space and $F : N \rightarrow M$ is smooth then $f \circ F$ is smooth and

$$D(f \circ F)(b) = Df(F(b)) \circ Df(b) \quad \text{for } b \in N.$$

Proof. Suppose a, U, V, ϕ, ψ are as in Definition 3.1. Then $(f \circ F)|_{F^{-1}[V]} = (f \circ \phi) \circ (\psi \circ F)|_{F^{-1}[V]}$. \square

If $f \in \mathcal{E}(M, Y)$, $a \in M$ and $v \in E$ we let

$$D_v f(a) = Df(a)(v).$$

If $k \in \mathbb{N}^+$ and $v = (v_1, \dots, v_k) \in E^k$ we let

$$D_v f(a) = D_{v_1} \cdots D_{v_k} f(a).$$

Proposition 3.2. *Suppose $f \in \mathcal{E}(M, Y)$, $a \in M$, W is an open neighborhood of a in \mathbb{R}^n , $F : W \rightarrow Y$ and $F|_M = f|_M$. Then $Df(a)(v) = DF(a)(v)$ whenever $a \in M \cap W$ and $v \in \mathbf{Tan}M, a$.*

Proof. If U, V, ϕ, ψ are as in Definition 3.1 then $F \circ \phi$ equals $f \circ \phi$ in an open neighborhood of $\psi(a)$. \square

We leave the straightforward proof of the following Proposition to the reader.

Proposition 3.3. *Suppose E is a Euclidean space, $m \in \mathbb{N}^+$ and $m \leq \dim E$, M is a nonempty subset of E and for each $a \in M$ there are U, V, ψ, ϕ such that V is an open subset of E ; U is an open subset of some m dimensional submanifold of some Euclidean space, $\psi : V \rightarrow U$, ψ is smooth, $\phi : U \rightarrow V$, ϕ is smooth, $a \in V$ and*

$$\psi(\phi(u)) = u \quad \text{for } u \in U.$$

Then $M \in \mathcal{M}_m(E)$.

If i_M is the inclusion map of M in \mathbb{R}^n then i_M is smooth and

$$Di_M(x) = \tau(x) \quad \text{for } x \in M.$$

It follows that τ is smooth and since $\nu(x) = i_{\mathbb{R}^n} - \tau(x)$ for $x \in M$ we find that ν is smooth.

Definition 3.2. *For each $k \in \mathbb{N}$ we define*

$$D^k f(a) \in \text{MultiLin}(\underbrace{E, \dots, E}_{k \text{ times}}, Y)$$

by letting $D^0 f = f$, $D^1 f = Df$ and, if $k \geq 2$, requiring that

$$D^k f(a)(D\phi(\psi(a))(v_1), \dots, D\phi(\psi(a))(v_k)) = D^k(f \circ \phi)(\psi(a))(v_1, \dots, v_k)$$

whenever a, U, V, ϕ, ψ are as in Definition 3.1 and $v_1, \dots, v_k \in \mathbb{R}^m$ and that

$$D^k f(a)(w_1, \dots, w_k) = 0 \quad \text{whenever } w_1, \dots, w_k \in E \text{ and } w_i \in \mathbf{Nor}M, a \text{ for some } i = 1, \dots, k.$$

As is well known $D^k f(a)$ is symmetric. We have

$$D^k f(a)(w_1, \dots, w_k) = D_{w_1} \cdots D_{w_k} f(a) \quad \text{whenever } w_1, \dots, w_k \in E.$$

Definition 3.3. *Suppose Y is a Euclidean space and $f : M \rightarrow Y$ is smooth. For each $k \in \mathbb{N}$ we let*

$$\chi^k(f) = \sup\{\|D^k f(x)\| : x \in M\}$$

and we let

$$\chi^{[k]}(f) = \max\{\chi^j(f) : j \in \mathbb{N} \text{ and } j \leq k\}.$$

Any of these numbers may be ∞ .

Remark 3.2. *We will not use χ as on page 229 of [AA81].*

3.3. More on differentiation. Suppose $N \in \mathbb{N}^+$ and, for each $i = 1, \dots, N$, E_i is a Euclidean space. For each $i \in \{1, \dots, N\}$ we let

$$\iota_i : E_i \rightarrow E_1 \times \cdots \times E_N$$

be such that for each $u \in E_i$ and $j \in \{1, \dots, N\}$

$$(\iota_i)_j(u) = \begin{cases} u & \text{if } j = i, \\ 0 & \text{if } j \neq i. \end{cases}$$

For each $j \in \mathbb{N}^+$ we let

$$\iota_i^j E_i^j \rightarrow (E_1 \times \cdots \times E_N)^j$$

be such that

$$\iota_i^j(u_1, \dots, u_j) = (\iota_i(u_1), \dots, \iota_i(u_j)) \quad \text{for } (u_1, \dots, u_j) \in E_i^j.$$

Suppose Y is a normed vector space and $f : M_1 \times \cdots \times M_N \rightarrow Y$ is smooth and $(k_1, \dots, k_N) \in \mathbb{N}^N$ and $K > 0$ where K is the number of positive k_i , $i = 1, \dots, N$.

Let l_j , $j = 1, \dots, K$, be such that

$$l_1 < \cdots < l_K \quad \text{and} \quad \{l_j : j = 1, \dots, K\} = \{i \in \{1, \dots, N\} : k_i > 0\}.$$

We let

$$D^{k_1, \dots, k_N} f : M_1 \times \cdots \times M_N \rightarrow \text{MultiLin}(E_{l_1}^{k_{l_1}}, \dots, E_{l_K}^{k_{l_K}}, Y)$$

be such that

$$D^{k_1, \dots, k_N} f(x_1, \dots, x_N)(U_1, \dots, U_K) = D^{\sum_{i=1}^K k_i} f(x_1, \dots, x_N)(\iota_{l_1}^{k_{l_1}}(U_1), \dots, \iota_{l_K}^{k_{l_K}}(U_K))$$

for $(U_1, \dots, U_K) \in E_{l_1}^{k_{l_1}}, \dots, E_{l_K}^{k_{l_K}}$. We also let

$$\chi^{k_1, \dots, k_N}(f) = \sup\{\|D^{k_1, \dots, k_N} f(x_1, \dots, x_N)\| : (x_1, \dots, x_N) \in M_1 \times \cdots \times M_N\}.$$

Example 3.1. *Supposed $N = 3$, $K = 2$ and $(k_1, k_2, k_3) = (3, 0, 2)$ and*

$$f : E_1 \times E_2 \times E_3 \rightarrow Y.$$

Then $(l_1, l_2) = (1, 3)$ and, if $U_1 \in E_1^3$ and $U_2 \in E_3^2$ then

$$D^{3,0,2} f(x_1, x_2, x_3)(U_1, U_2) = D^5 f(x_2, x_2, x_3)((U_1)_1, (U_1)_2, (U_1)_3, (U_2)_1, (U_2)_2).$$

for $(x_1, x_2, x_3) \in E_1 \times E_2 \times E_3$.

e

3.4. Diffeomorphisms.

Definition 3.4. *Suppose*

$$f : X \rightarrow Y.$$

We say f carries X diffeomorphically onto Y if

- (i) *there are $m \in \mathbb{N}^+$ and Euclidean spaces E and F such that $X \in \mathcal{M}_m(E)$ and $Y \in \mathcal{M}_m(F)$;*
- (ii) *f is smooth and univalent;*
- (iii) *$\text{rng } f = Y$ and f^{-1} is smooth.*

Proposition 3.4. *Suppose*

- (i) *$m \in \mathbb{N}^+$ and E and F are Euclidean spaces;*
- (ii) *$X \in \mathcal{M}_m(E)$, $Y \in \mathcal{M}_m(F)$ and*

$$f : X \rightarrow Y;$$

- (iii) f is smooth and univalent;
- (iv) $\mathbf{rank} Df(a) = m$ for each $a \in M$;
- (v) X is compact and X and Y are homeomorphic.

Then $\mathbf{rng} f = Y$ and f carries X diffeomorphically onto Y .

Remark 3.3. *This Proposition need not hold if X is not compact.*

Proof. Let \mathcal{C} be the family of connected components of Y . \mathcal{C} is finite since Y is compact.

Suppose $a \in X$. Since f is smooth and $\mathbf{rank} Df(a) = m$ is elementary that there are G and H such that G is a relatively open subset of X , $a \in G$, H is an open subset of F and $f[G] = H \cap \mathbf{rng} f$. Let $d = \mathbf{dist}(a, Y \sim H)$. Then $d > 0$ since $Y \sim H$ is compact and $a \notin Y \sim H$. So there is an open subset W of H such that $a \in W$ and $W \cap Y = W \cap f[G]$. Thus $\mathbf{rng} f$ is open relative to Y . So there is a subfamily \mathcal{B} of \mathcal{C} such that $\mathbf{rng} f = \bigcup \mathcal{B}$. Since $\mathbf{rng} f$ is homeomorphic to X and X is homeomorphic to Y we find that $\mathcal{B} = \mathcal{C}$ so $\mathbf{rng} f = Y$. That f^{-1} is smooth follows from the Inverse Function Theorem. So f carries X diffeomorphically onto Y \square

3.5. $\mathcal{L}_m, \mathcal{P}_m$ and \mathcal{W}_m . Let $\mathcal{L}_m = \{L \in \mathbb{E}\mathbf{nd}(\mathbb{R}^n) : \mathbf{rank} L = m\}$; as is well known, $\mathcal{L}_m \in \mathcal{M}_{m(n-m)}(\mathbb{E}\mathbf{nd}(\mathbb{R}^n))$.

Let

$$\mathcal{P}_m : \mathcal{L}_m \rightarrow \mathbb{O}_m(\mathbb{R}^n)$$

be such that $\mathcal{P}_m(L)$ is orthogonal projection of \mathbb{R}^n onto $\mathbf{rng} L$; as is well known, \mathcal{P}_m is smooth and for each $R \in \mathbb{R}^+$ there is constant $C_{\mathcal{P}_m, R}$ such that

$$(1) \quad \|\mathcal{P}_m(L) - \mathcal{P}_m(M)\| \leq C_{\mathcal{P}_m, R} \|L - M\|.$$

whenever $L, M \in \mathcal{L}_m$ and $\max\{\|L\|, \|M\|\} \leq R$.

Let

$$\mathcal{W}_m : \mathcal{L}_m \rightarrow \mathbb{R}^+$$

be such that $\mathcal{W}_m(L) = |\bigwedge_m L|$ for $L \in \mathcal{L}_m$. It is elementary that \mathcal{W}_m is smooth and that

$$|\mathcal{W}_m(L) - 1| \leq (2^m - 1) \|L - \mathcal{P}_m(L)\|$$

3.6. **The second fundamental form of M .** We define

$$B : M \rightarrow \odot^2(\mathbb{R}^n, \mathbb{R}^n)$$

by requiring that, for any $a \in M$,

$$B(a)(D\phi(\psi(a)(s)), D\phi(\psi(a)(t))) = \nu(a)(D^2\phi(\psi(a))(s, t))$$

where U, V, ψ, ψ are as in Definition 3.1 and $s, t \in \mathbb{R}^m$ and that

$$B(a)(u, v) \bullet w = 0 \quad \text{if either } u \in \mathbf{Nor}M, a \text{ or } v \in \mathbf{Nor}M, a \text{ or } w \in \mathbf{Tan}M, a.$$

One calls B the **second fundamental form of M** .

Proposition 3.5. *Suppose $a \in M$, $u, v \in \mathbf{Tan}M, a$. Then*

$$D\tau(a)(u)(v) = B(a)(u, v) = -D\nu(a)(u)(v).$$

Proof. Let U, V, ϕ, ψ are as in Definition 3.1, let $b = \psi(a)$, note that $\phi(b) = a$ and let $s, t \in \mathbb{R}^m$ be such that $u = D\phi(b)(s)$ and $v = D\phi(b)(t)$. Then

$$(2) \quad \nu(\phi(x))(D\phi(x)(t)) = 0 \quad \text{for } x \in U.$$

We differentiate (2) at $x = b$ in the direction s and find that

$$\begin{aligned} 0 &= D\nu(a)(D\phi(b)(s))(D\phi(b)(t)) + \nu(a)(D(D\phi)(b)(s, t)) \\ &= D\nu(a)(D\phi(b)(s))(D\phi(b)(t)) + \nu(a)(D^2\phi(b)(s, t)) \\ &= D\nu(a)(u)(v) + B(a)(u, v). \end{aligned}$$

□

We define

$$A : M \rightarrow \text{Lin}(\mathbb{R}^n, \text{Lin}(\mathbb{R}^n))$$

by requiring that, for any $a \in M$,

$$A(a)(u)(v) \bullet w = u \bullet B(a)(v, w) \quad \text{whenever } u, v, w \in \mathbb{R}^n.$$

It follows that

$$A(a)(u)(v) \bullet w = A(a)(u)(w) \bullet v \quad \text{if } u, v, w \in \mathbb{R}^n.$$

We have

$$\|B\| = \sup\{|B(a)(u, v) \bullet w| : a \in M \text{ and } u, v, w \in \mathbb{S}^{n-1}\};$$

It is elementary that $\|B\| < \infty$ if M is compact and that $\|B\| = 0$ if and only if for each point $a \in M$ there are an open neighborhood V of a and $S \in \mathbb{G}_m(\mathbb{R}^n)$ such that $M \cap V = (a + S) \cap V$. We let

$$R = \frac{1}{\|B\|} \in [0, \infty];$$

it is elementary that $0 < R < \infty$ if M is compact.

4. VECTOR BUNDLES.

Theorem 4.1. $\mathbb{O}_m(\mathbb{R}^n) \in \mathcal{M}_{m(n-m)}(\text{End}(\mathbb{R}^n))$.

Proof. Let

$$\mathcal{P}(l) = l \circ l - l \quad \text{for } l \in \mathbb{S}\text{ym}(\mathbb{R}^n).$$

Suppose $p \in \mathbb{O}_m(\mathbb{R}^n)$. Suppose $l \in \ker D\mathcal{P}(p)$. We have

$$0 = D\mathcal{P}(p)(l) = l \circ p + l \circ l - l = l \circ p - p^\perp \circ l;$$

this implies that $p \circ l \circ p = 0$ and $p^\perp \circ l \circ p^\perp = 0$ so $l = p \circ l \circ p^\perp + p^\perp \circ l \circ p$ and since $l^* = l$ we have $p^\perp \circ l \circ p = (p \circ l \circ p^\perp)^*$. □

Suppose l, N are integers and $1 \leq l \leq N$. Suppose

$$\gamma : M \rightarrow \mathbb{O}_l(\mathbb{R}^N)$$

and γ is smooth. We let

$$\mathbf{E}(\gamma) = \{(x, y) \in M \times \mathbb{R}^N : y \in \text{rng } \gamma(x)\}$$

and for $r \in \mathbb{R}^+$ we let

$$\mathbf{E}(\gamma, r) = \{(x, y) \in \mathbf{E}(\gamma) : |y| < r\}.$$

We let

$$\mathbf{Tan}M = \mathbf{E}(\tau) \quad \text{and we let} \quad \mathbf{Nor}M = \mathbf{E}(\nu).$$

Proposition 4.1. *Suppose $a \in M$ and $u \in \mathbf{Tan}M, a$. Then*

$$D\gamma(a)(u) = (D\gamma(a)(u))^*$$

and

$$D\gamma(a)(u) = \gamma(a) \circ D\gamma(a)(u) \circ \gamma(a)^\perp + \gamma(a)^\perp \circ D\gamma(a)(u) \circ \gamma(a).$$

Proof. The first equation follows from $\gamma(x)^* = \gamma(x)$ for all $x \in M$.

Since $\gamma(x) \circ \gamma(x) = \gamma(x)$ for $x \in M$ we have

$$D\gamma(a)(u) \circ \gamma(a) + \gamma(a) \circ D\gamma(a)(u) = D\gamma(a)(u)$$

for $u \in \mathbb{R}^n$. Premultiplying and postmultiplying this equation by $\gamma(a)$ we infer that $\gamma(a) \circ D\gamma(a)(u) \circ \gamma(a) = 0$. Premultiplying and postmultiplying this equation by $\gamma(a)^\perp$ we infer that $\gamma(a)^\perp \circ D\gamma(a)(u) \circ \gamma(a)^\perp = 0$. \square

Theorem 4.2. $\mathbf{E}(\gamma) \in \mathcal{M}_{m+l}(\mathbb{R}^n \times \mathbb{R}^N)$. Moreover, for any $(x, y) \in \mathbf{E}(\gamma)$ we have

(3)

$$\begin{aligned} \mathbf{TanE}(\gamma), (x, y) &= \{(u, v) \in \mathbf{Tan}M, x \times \mathbb{R}^N : \gamma(x)^\perp(v) = D\gamma(x)(u)(y)\} \\ &= \{(u, D\gamma(x)(u)(y)) : u \in \mathbf{Tan}M, x\} \oplus \{(0, v) : v \in \mathbf{rng} \gamma(x)\} \end{aligned}$$

Proof. We have

$$\mathbf{Tan}M \times \mathbb{R}^N, (x, y) = \mathbf{Tan}M, x \oplus \mathbb{R}^N \quad \text{for } (x, y) \in M \times \mathbb{R}^N.$$

Let $F : M \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ be such that

$$F(x, y) = \gamma(x)^\perp(y) \quad \text{for } (x, y) \in M \times \mathbb{R}^N.$$

Then $\mathbf{E}(\gamma) = \{(x, y) \in M \times \mathbb{R}^N : F(x, y) = 0\}$.

Suppose $(x, y) \in \mathbf{E}(\gamma)$. We have

$$DF(x, y)(u, 0) = -D\gamma(x)(u)(y) \in \mathbf{rng} \gamma(x)^\perp \quad \text{for } u \in \mathbf{Tan}M, x$$

as well as

$$DF(x, y)(0, v) = \gamma(x)^\perp(v) \quad \text{for } v \in \mathbb{R}^N;$$

it follows that

$$\mathbf{rng} DF(x, y) = \mathbf{rng} \gamma(x)^\perp; \quad \mathbf{dim} \mathbf{rng} DF(x, y) = N - l;$$

and

$$\mathbf{ker} DF(x, y) = \{(u, v) \in \mathbf{Tan}M, x \times \mathbb{R}^N : \gamma(x)^\perp(v) = D\gamma(x)(u)(y)\}.$$

The Theorem now follows from the Implicit Function Theorem. \square

Corollary 4.1. *Suppose $Z \in \mathcal{E}(\gamma)$, $x \in M$ and $u \in \mathbf{Tan}M, x$. Then*

$$D\gamma(x)(u)(Z(x)) = \gamma^\perp(x)(DZ(x)(u)).$$

Proof. $(u, DZ(x)(u)) \in \mathbf{TanE}(\gamma), (x, Z(x))$. \square

If $Z \in \mathcal{E}(\gamma)$ we let

$$D_\gamma Z : M \rightarrow \mathbb{L}\text{in}(\mathbb{R}^n, \mathbb{R}^l)$$

be such that $D_\gamma Z(x) = \gamma(x) \circ DZ(x)$ for $x \in \mathbb{R}^n$. Evidently, $\mathbf{rng} D_\gamma Z(x) \subset \mathbf{rng} \gamma(x)$ and $\mathbf{Nor}M, x \subset \mathbf{ker} D_\gamma Z(x)$. The preceding Corollary implies that

$$(4) \quad DZ(x) = D_\gamma Z(x) + D\gamma(x)(u)(Z(x)) \quad \text{for } x \in M.$$

Corollary 4.2. *If $(x, y) \in \mathbf{Tan}M$ then*

$$\mathbf{TanTan}M, (x, y) = \{(u, B(x)(u, y) + v) : (u, v) \in \mathbf{Tan}M, x \times \mathbf{Tan}M, x\}$$

and if $(x, y) \in \mathbf{Nor}M$ then

$$\mathbf{TanNor}M, (x, y) = \{(u, -A(x)(y)(u) + v) : u \in \mathbf{Tan}M, x \times \mathbf{Nor}M, x\}.$$

Theorem 4.3 (Weingarten Equations). *Suppose $a \in M$ and $u \in \mathbf{Tan}M, a$. Then*

$$(5) \quad \tau(a)(DN(a)(u)) = -A(a)(N(a))(u) \quad \text{if } N \in \mathcal{E}(\nu)$$

and

$$(6) \quad \nu(a)(DT(a)(u)) = B(a)(T(a), u) \quad \text{if } T \in \mathcal{E}(\tau).$$

Proof. Suppose $N \in \mathcal{E}(\nu)$. Then $\tau(x)(N(x)) = 0$ for $x \in M$; differentiating at $x = a$ we find that

$$\tau(a)(DN(a)(u)) = -D\tau(a)(u)(N(a)) = -A(a)(N(a))(u).$$

Suppose $T \in \mathcal{E}(\tau)$. Then $\nu(x)(T(x)) = 0$ for $x \in M$. Differentiating at $x = a$ we find that

$$\nu(a)(DT(a)(u)) = -D\nu(a)(T(a))(u) = D\tau(a)(T(a))(u) = B(a)(T(a), u).$$

□

4.1. The mean curvature normal of M . We let

$$H \in \mathcal{E}(\nu)$$

be such that

$$H(x) \bullet u = \mathbf{trace} A(x)(u) \quad \text{for } x \in M \text{ and } u \in \mathbf{Nor}M, x.$$

H is called the **mean curvature normal of M** .

Proposition 4.2. *Suppose*

(i) *I is an open interval, $0 \in I$,*

$$h : I \times M \rightarrow \mathbb{R}^n$$

and h is smooth;

(ii) *for each $t \in I$, $M_t = h_t[M] \in \mathcal{M}_m(\mathbb{R}^n)$ and h_t carries M diffeomorphically onto M_t ;*

(iii) *for each $t \in I$, $X_t : M_t \rightarrow \mathbb{R}^n$ is such that*

$$X_t(y) = \frac{d}{d\epsilon} h_{t+\epsilon}(h_t^{-1}(y))|_{\epsilon=0}$$

for $y \in M_t$.

Then X_t is smooth and

$$\frac{d}{dt} \mathcal{H}^m(h_t[M]) = \int X_t \bullet H_t d\|M_t\|$$

for each $t \in I$.

For the proof see the literature.

5. REACH.

Suppose $M \in \mathcal{M}_m(\mathbb{R}^n)$.

Definition 5.1. We let

$$\Xi : \mathbf{Nor}M \rightarrow \mathbb{R}^n$$

be such that $\Xi(x, y) = x + y$ for $(x, y) \in \mathbf{Nor}M$ and we let

$$\mathbf{r} = \sup(\{0\} \cup \{r \in (0, \infty) : \Xi|_{\mathbf{E}(\nu, r)} \text{ is univalent.}\})$$

Proposition 5.1. Ξ is smooth.

Proof. Ξ is the restriction to $\mathbf{E}(\nu)$ of the smooth map

$$\mathbb{R}^n \times \mathbb{R}^n \ni (x, y) \mapsto x + y \in \mathbb{R}^n.$$

□

Proposition 5.2. Suppose $0 < r < \infty$. The following statements are equivalent:

- (i) $r \leq \mathbf{r}$;
- (ii) $M \cap \mathbf{U}(a + v, |v|) = \{a\}$ whenever $(a, v) \in \mathbf{Nor}M$ and $0 < |v| < r$;
- (iii) $|\nu(a)(x - a)| \leq \frac{|x - a|^2}{2r}$ whenever $a, x \in M$.

Proof. Suppose (i) holds, $a \in M$ and $u \in \mathbf{Nor}M, a \cap \mathbb{S}^{n-1}$. Let $I = \{s \in (0, r) : a + su \notin M\}$. Suppose I were not empty. Then $0 \leq \inf I < r$. We have $\inf I > 0$ since $u \notin \mathbf{Tan}M, a$. Let $t = \inf I/2$. Then $\Xi(a, tu) = a + tu = \Xi(a + 2tu, -tu)$ which is impossible. Suppose $s \in I$ and $x \in M$ is such that $\mathbf{dist}(a + su, M) = |(a + su) - x|$. Then $(a + su) - x \in \mathbf{Nor}M, x$ so $\Xi(x, a + su) = \Xi(a, su)$ so, as $s \leq r \leq \mathbf{r}$ we have $M \cap \mathbf{U}(a + su, s) = \{a\}$ which implies that $\mathbf{U}(a + ru, r) \cap M = \emptyset$. Thus (ii) holds.

It is a simple matter to verify that (ii) implies (i) and that (ii) and (iii) are equivalent. □

Proposition 5.3. $\mathbf{r} \leq \frac{1}{\|B\|}$.

Proof. Suppose $a \in M$. Suppose U, V, ϕ, ψ are as in Definition 3.1 and $u \in \mathbb{R}^m$. We have

$$B(a)(D\phi(0)(u), D\phi(0)(u)) = D^2\phi(0)(u, u)$$

as well as, for t sufficiently small,

$$\phi(tu) = \phi(0) + tD\phi(0)(u) + \frac{t^2}{2}D^2\phi(0)(u, u) + \rho(t)$$

where $\rho(t)/t^2 \rightarrow 0$ as $t \rightarrow 0$. So

$$\begin{aligned} \frac{\nu(a)(\phi(tu) - \phi(0))}{t^2} &= \frac{1}{2}\nu(a)(D^2\phi(0)(u, u)) + \frac{\nu(a)(\rho(t))}{t^2} \\ &= \frac{1}{2}(B(a)(D\phi(0)(u), D\phi(0)(u)) + \frac{\nu(a)(\rho(t))}{t^2}). \end{aligned}$$

But

$$\frac{|\nu(a)(\phi(tu) - \phi(0))|}{t^2} \leq \frac{|\phi(tu) - \phi(0)|^2}{2r t^2} \rightarrow \frac{|D\phi(0)(u)|^2}{2r} \text{ as } t \rightarrow 0.$$

Thus $|B(a)(D\phi(0)(u), D\phi(0)(u))| \leq \frac{|D\phi(0)(u)|^2}{r}$.

□

Theorem 5.1. *Suppose $(a, b) \in \mathbf{Nor}M$ and $(u, v) \in \mathbf{Tan}M, a \times \mathbf{Nor}M, a$. Then*

$$(7) \quad D\Xi(a, b)(u - A(a)(b)(u), v) = u - A(a)(b)(u) + v$$

and

$$(8) \quad |D\Xi(a, b)(u - A(a)(b)(u), v)|^2 = |u - A(a)(b)(u)|^2 + |v|^2.$$

In particular, if $|b| < 1/\|B\|$, then

$$(9) \quad |D\Xi(a, b)(u - A(a)(b)(u), v)| \geq |u - A(a)(b)(u)| \geq 1 - \|B\||v| > 0$$

and $D\Xi(a, b)$ carries $\mathbf{TanNor}M, (a, b)$ isomorphically onto \mathbb{R}^n .

Proof. $(u - A(a)(b)(u), v) \in \mathbf{TanNor}M, a, (a, b)$ by Corollary 4.2.

Let $F(x, y) = x + y$ for $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$. Since $\Xi = F|_{\mathbf{Nor}M}$, (7) follows from Corollary 4.2. (8) holds since $u - A(a)(b)(u)(v) \in \mathbf{Tan}M, a$ and $v \in \mathbf{Nor}M, a$. If $|b| < 1/\|B\|$ then $|u - A(a)(b)(u)| > 1 - \|B\||v|$ and (9) follows from (8). (9) implies $D\Xi(a, b)|_{\mathbf{TanNor}M, (a, b)}$ is univalent. This and the fact that $\mathbf{Nor}M \in \mathcal{M}_n(\mathbb{R}^n \times \mathbb{R}^n)$ implies the final conclusion. \square

Definition 5.2. *For $r \in (0, \infty]$ we set*

$$N(r) = \{z \in \mathbb{R}^n : \mathbf{dist}(z, M) < r\}.$$

Obviously, $N(r)$ is an open subset of \mathbb{R}^n .

Theorem 5.2. *Suppose $\mathbf{r} > 0$. Then Ξ carries $\mathbf{E}(\nu, \mathbf{r})$ diffeomorphically onto $N(\mathbf{r})$.*

Proof. $\Xi|_{\mathbf{E}(\nu, \mathbf{r})}$ is univalent by the definition of \mathbf{r} . If $(a, b) \in \mathbf{E}(\nu, \mathbf{r})$ then $D\Xi(a, b)$ carries $\mathbf{TanNor}M, (a, b)$ isomorphically onto $\mathbf{Tan}N(\mathbf{r}), a + b = \mathbb{R}^n$ by the preceding Theorem. \square

Proposition 5.4. $\mathbf{r} = \infty$ if and only if M is a subset of some member of $\mathbb{G}_m(\mathbb{R}^n)$.

Proof. Suppose $\mathbf{r} = \infty$ and $a \in M$. It follows from (ii) of the preceding Proposition that $M \subset \mathbf{Tan}M, a$. \square

Proposition 5.5. *Suppose $M \subset \mathbb{S}^{n-1}$. If $a \in M$ then $A(a)(u) = u$ for $u \in \mathbf{Tan}M, a$; in particular, $\|B\| \geq 1$ and $\mathbf{r} \leq 1$.*

Proof. Let $X(x) = x$ for $x \in M$. Then $X \in \mathcal{E}(\nu)$ and, if $u \in \mathbf{Tan}M, a$, $DX(a)(u) = u$ so so $A(a)(u) = u$ by a (5); also, if $|u| = 1$, then $\|B\| \geq \|A(a)(u)\| = 1$. \square

Lemma 5.1. *Suppose*

- (i) $Y_i, i = 1, 2$, is a Euclidean space;
- (ii) $N_i \in \mathcal{M}_m(Y_i)$ for $i = 1, 2$;
- (iii) K is a compact subset of N_1 ; $F : N_1 \rightarrow N_2$ is smooth; $F|_K$ is univalent and, for each $x \in K$, $DF(x)$ carries $\mathbf{Tan}N_{1,x}$ isomorphically onto $\mathbf{Tan}N_{2,F(x)}$;
- (iv) $U_\epsilon, 0 < \epsilon < \infty$, is an open subset of N_1 such that
 - (a) if $0 < \epsilon_1 < \epsilon_2 < \infty$ then $U_{\epsilon_1} \subset U_{\epsilon_2}$;
 - (b) if V is an open subset of N_1 and $K \subset V$ then $U_\epsilon \subset V$ for some $\epsilon > 0$;
 - (c) $K = \bigcap_{\epsilon > 0} U_\epsilon$.

Then

- (v) there is $\epsilon > 0$ such that $DF(x)$ carries $\mathbf{Tan}N_{1,x}$ isomorphically onto $\mathbf{Tan}N_{2,F(x)}$ for $x \in U_\epsilon$ and such that $F|_{U_\epsilon}$ is univalent;

- (vi) if ϵ is as in (v) then $F[U_\epsilon]$ is an open neighborhood in N_2 of $F[K]$ and F carries U_ϵ diffeomorphically onto $F[U_\epsilon]$.

Remark 5.1. What follows is a paraphrase of the argument on page 116 of [MS].

Proof. Let V_1 be the set of $x \in N_1$ such that $DF(x)$ carries $\mathbf{Tan}N_1, x$ isomorphically onto $\mathbf{Tan}N_2, F(x)$. Then V_1 is an open subset of N_1 so there is by (iii)(b) $\epsilon > 0$ such that $U_\epsilon \subset V_1$.

Suppose (v) is false. Let N_* be a positive integer such that the closure of K of U_{1/N_*} in N_1 is compact. For each integer N such that $N \geq N_*$ there are $a_N, b_N \in U_{1/N}$ such that $a_N \neq b_N$ and $F(a_N) = F(b_N)$. Since K is compact there are a_* and b_* in K which are accumulation points of $\{a_N : N = 1, 2, \dots\}$ and $\{b_N : N = 1, 2, \dots\}$, respectively. Then $F(a_*) = F(b_*)$ which implies $\lim_{N \rightarrow \infty} |a_N - b_N| = 0$. Thus F would not be univalent in any open neighborhood of a_* . But as $DF(a_*)$ carries $\mathbf{Tan}N_1, a_*$ isomorphically onto $\mathbf{Tan}N_2, F(a_*)$, F must be univalent in some open neighborhood of a_* .

(vi) is an elementary consequence of (v) and the smoothness of F . \square

Corollary 5.1. Suppose $M \in \mathcal{M}_m(\mathbb{R}^n)$ and M is compact. Then $\mathbf{r} > 0$.

Proof. Let

$$Y_1 = \mathbb{R}^n \times \mathbb{R}^n; \quad Y_2 = \mathbb{R}^n; \quad N_1 = \mathbf{Nor}M; \quad N_2 = \mathbb{R}^n;$$

let $F = \Xi$ and, for $\epsilon > 0$, let $U_\epsilon = \mathbf{E}(\gamma, \epsilon)$; and let $K = \{(x, 0) : x \in M\} \subset N_1$. Then (i)-(iv) of Lemma 5.1 hold. So if ϵ is as in Lemma 5.1 then $\mathbf{reach}M \geq \epsilon$. \square

5.1. Suppose M is compact. Since Ξ carries $\mathbf{E}(\nu, \mathbf{r})$ diffeomorphically onto $N(\mathbf{r})$ we may define smooth functions

$$\xi : N(\mathbf{r}) \rightarrow \mathbb{R}^n \quad \text{and} \quad \sigma : N(\mathbf{r}) \rightarrow \mathbb{R}^n$$

by requiring that

$$(\xi, \sigma) = (\Xi|_{\mathbf{E}(\nu, \mathbf{r})})^{-1} : N(\mathbf{r}) \rightarrow \mathbf{E}(\nu, \mathbf{r}).$$

Note that

$$(10) \quad z = \Xi(\xi(z), \sigma(z)) = \xi(z) + \sigma(z) \quad \text{for } z \in N(\mathbf{r}).$$

Theorem 5.3. Suppose $b \in N(\mathbf{r})$, $a = \xi(b)$ and $c = \sigma(b)$. Then

- (i) $a \in M$, $c \in \mathbf{Nor}M, a$, $b = a + c$ and $|c| = \mathbf{dist}(b, M)$;
- (ii) $D\xi(b)(u - A(a)(c)(u)) = u$ for $u \in \mathbf{Tan}M, a$ and $\ker D\xi(b) = \mathbf{Nor}M, a$;
- (iii) $D\sigma(b)(u - A(a)(c)(u)) = -A(a)(c)(u)$ for $u \in \mathbf{Tan}M, a$ and $D\sigma(b)(v) = v$ for $v \in \mathbf{Nor}M, b$;
- (iv) $\|D\xi(b) - \tau(a)\| \leq \frac{\|B\||c|}{1 - \|B\||c|} \leq \frac{\mathbf{r}|c|}{\mathbf{r} - |c|}$.

Remark 5.2. Keep in mind that

$$\mathbf{Tan}M, a \ni u \mapsto u - A(a)(c)(u) \in \mathbf{Tan}M, a$$

is a linear isomorphism since $\|B\||c| < 1$.

Proof. We have $\xi \circ \Xi(x, y) = x$ and $\sigma \circ \Xi(x, y) = y$ for $(x, y) \in \mathbf{E}(\nu, \mathbf{r})$. (i)-(iii) now follow from (7) and the Chain Rule.

Let $l \in \mathbb{L}\text{End}(\mathbf{Tan}M, a)$ and $L \in \mathbb{L}\text{in}(\mathbf{Tan}M, a, \mathbb{R}^n)$ be such that $l(u) = A(a)(c)(u)$ and $L(u) = u - l(u)$ for $u \in \mathbf{Tan}M, a$. Then $\|l\| \leq \|B\||c| < 1$ and $\|L\| \geq 1 - \|l\|$

so L is invertible and $\|L^{-1}\| \leq (1 - \|l\|)^{-1}$. Let i be the identity map of $\mathbf{Tan}M, a$. (ii) implies that $D\xi(b) \circ L = i$ so $D\xi_b|_{\mathbf{Tan}M, a} = L^{-1}$ which implies that

$$\|D\xi(b)|_{\mathbf{Tan}M, a} - i\| = \|L^{-1} - i\| \leq \|L^{-1}\| \|i - L\| \leq \frac{\|l\|}{1 - \|l\|} \leq \frac{\|B\| \|c\|}{1 - \|B\| \|c\|}.$$

(iv) follows since (ii) and (iii) imply that $\mathbf{Nor}M, a \subset \ker D\xi(b)$. \square

Theorem 5.4. *Suppose $z \in N(\mathbf{r})$. Then*

$$(11) \quad D\xi(\xi(z)) = \tau(\xi(z))$$

and

$$(12) \quad \|D\xi(z) - \tau(\xi(z))\| \leq \chi^2(\xi) \mathbf{dist}(z, M).$$

Proof. (11) follows from (ii) of Theorem 5.3. Let $C(\epsilon) = (1 - \epsilon)z + \epsilon\xi(z)$ for $\epsilon \in (0, 1)$. Since $\mathbf{U}(z, \mathbf{dist}z, M) \subset N(\mathbf{r}) = \mathbf{dmn}\xi$ we find that $\mathbf{rng}C \subset \mathbf{dmn}\xi$ so that

$$D\xi(z) - \tau(\xi(z)) = D\xi(z) - D\xi(\xi(z)) = \int_0^1 D(D\xi)(C(\epsilon))(C'(\epsilon)) d\epsilon$$

so, as $C'(\epsilon) = z - \xi(z)$ for $\epsilon \in (0, 1)$, (12) holds. \square

Theorem 5.5. *Suppose f is a smooth function on M with values in a normed space Y . Then*

$$\mathbf{Lip} f \leq \left(1 + \frac{1}{\mathbf{r}}\right) \chi^{[1]}(f).$$

Proof. Suppose $0 < \lambda < 1$ and $|a - b| \leq 2\lambda\mathbf{r}$. Let $C(t) = (1 - t)a + tb$ for $t \in (0, 1)$. Then $\mathbf{dist}(C(t), M) < \lambda\mathbf{r}$ for $t \in (0, 1)$ so $\mathbf{rng}C \subset \mathbf{dmn}\xi$ and, by Theorem 5.3(ii),

$$\|D\xi(C(t))\| \leq \frac{1}{1 - \|B\| \mathbf{dist}(C(t), M)} \leq \frac{1}{1 - \lambda}$$

so

$$\frac{d}{dt}(f \circ \xi \circ C)(t) \leq \chi^1(f) \|D\xi(C(t))\| \|C'(t)\| \leq \frac{\chi^1(f)}{1 - \lambda} |a - b|.$$

Integrating this inequality from 0 to 1 we infer that

$$|f(a) - f(b)| \leq \frac{\chi^1(f)}{1 - \lambda} |a - b|.$$

On the other hand, if $|a - b| > 2\lambda\mathbf{r}$ then

$$|f(a) - f(b)| \leq \mathbf{diam} \mathbf{rng} f \leq 2\chi^0(f) < \frac{\chi^0(f)}{\lambda\mathbf{r}} |a - b|.$$

Thus

$$|f(a) - f(b)| \leq \chi^{[1]} \max \left\{ \frac{1}{1 - \lambda}, \frac{1}{\lambda\mathbf{r}} \right\} |a - b|$$

so we should set $\lambda = 1/(1 + \mathbf{r})$. \square

5.2. θ and ι . Let

$$\theta : \mathbb{S}^{n-1} \rightarrow \mathbb{O}_{n-1}(\mathbb{R}^n) \quad \text{and} \quad \iota : \mathbb{S}^{n-1} \rightarrow \mathbb{O}_1(\mathbb{R}^n)$$

be the tangent map of \mathbb{S}^{n-1} and normal map of \mathbb{S}^{n-1} , respectively.

Corollary 5.2. *Suppose $(x, v) \in \mathbf{E}(\theta)$. Then*

$$D\iota(x)(u) = \mathbf{TanE}(\theta), (x, v) = \{(u, w) \in \mathbf{TanS}^{n-1}, x \times \mathbb{R}^n : w \bullet x = -u \bullet v\}.$$

6. η AND \mathbf{o} .

We define

$$\eta : \mathbb{R}^n \sim \{0\} \rightarrow \mathbb{S}^{n-1} \quad \text{and} \quad \mathbf{o} : \mathbb{R}^n \sim \{0\} \rightarrow \mathbb{G}_{n-1}(\mathbb{R}^n)$$

at $x \in \mathbb{R}^n \sim \{0\}$ by letting $\eta(x) = |x|^{-1}x$ and letting $\mathbf{o}(x)$ be orthogonal projection of \mathbb{R}^n onto $\mathbf{TanS}^{n-1}, x = \{w \in \mathbb{R}^n : w \bullet x = 0\}$.

Proposition 6.1. *Suppose $x, y \in \mathbb{S}^{n-1} \sim \{0\}$. Then*

$$(13) \quad |\eta(x) - \eta(y)| \leq \frac{|x - y|}{\min\{|x|, |y|\}}.$$

Proof. Let $s, t \in \mathbb{R}^+$ and $u, v \in \mathbb{S}^{n-1}$ be such that $x = su$ and $y = tv$. Since both sides of (13) are symmetric in x and y we may suppose that $0 < s = |x| \leq |y| = t$. Then (13) amounts to

$$|u - v| \leq \frac{|su - tv|}{s} = |u - \lambda v|$$

where we have set $\lambda = t/s \geq 1$. We have

$$|u - \lambda v| - |u - v| = 1 - 2\lambda u \bullet v + \lambda^2 - (1 - 2u \bullet v + 1) = (\lambda - 1)(\lambda + 1 - 2u \bullet v) \geq 0.$$

□

Proposition 6.2. *For $x \in \mathbb{R}^n \sim \{0\}$ we have*

$$D\eta(x) = \frac{\mathbf{o}(x)}{|x|};$$

in particular,

$$D\eta(x)(w) = |x|^{-3}(|x|^2 w - (w \bullet x)x) \quad \text{for } w \in \mathbb{R}^n.$$

Proof. Suppose $w \in \mathbb{R}^n$. Then

$$D\eta(x)(w) = \frac{d}{d\epsilon} \frac{x + \epsilon w}{|x + \epsilon w|} \Big|_{\epsilon=0} = \frac{w}{|x|} - \frac{1}{|x|^2} \left(w \bullet \frac{x}{|x|} \right) x = |x|^{-3}(|x|^2 w - (w \bullet x)x).$$

□

Proposition 6.3. *Suppose $x \in \mathbb{S}^{n-1}$ and $v \in \mathbb{R}^n$ and $v \bullet x = 0$. Then*

$$|\eta(x + v) - x| \leq |v|.$$

Proof. We have

$$|x - \eta(x + v)|^2 = 2 \left(1 - x \bullet \left(\frac{x + v}{|x + v|} \right) \right) = 2 \left(1 - \frac{1}{\sqrt{1 + |v|^2}} \right) \leq |v|^2.$$

□

7. f .

We let

$$f : \mathbf{E}(\nu, \mathbf{r}) \rightarrow \mathbb{R}^n$$

be such that

$$f(x, y) = (1 + y \bullet x) \boldsymbol{\eta}(x + y) \quad \text{for } (x, y) \in \mathbf{E}(\nu, \mathbf{r}).$$

We let

$$v : N(\mathbf{r}) \rightarrow \mathbb{R}$$

be such that $v(z) = z \bullet \xi(z)$ for $z \in N(\mathbf{r})$.

Remark 7.1. *On page 221 of [AA81] f there is based on the exponential map of \mathbb{S}^{n-1} applied to $\Xi|\mathbf{E}(\gamma)$. The f here is as in [FN] which I believe is easier to work with. The modifications to required to accomodate this f in [AA81] are straightforward.*

Proposition 7.1. $v > 0$.

Proof. Suppose $z \in N(\mathbf{r})$ and $x = \xi(z)$. Then

$$z \bullet \xi(z) = (x + (z - x)) \bullet x = 1 + (z - x) \bullet x \geq 1 - |z - x| > 1 - \mathbf{r} \geq 0.$$

□

We define the smooth function

$$\rho : N(\mathbf{r}) \rightarrow \mathbb{R}^n$$

by setting

$$\rho(z) = \frac{z}{v(z)} - \xi(z) \quad \text{for } z \in N(\mathbf{r}).$$

Theorem 7.1. f is a diffeomorphism with range an open subset V of $N(\mathbf{r})$ and with inverse $(\xi, \rho)|V$. In particular,

(14)

$$D\xi(f(a, b))(Df(a, b)(u - A(a)(b)(u), v)) = DP(a, b)(u - A(a)(b)(u), v) = u - A(a)(b)(u)$$

and

$$(15) \quad D\rho(f(a, b))(Df(a, b)(u - A(a)(b)(u), v)) = DQ(a, b)(u - A(a)(b)(u), v) = v$$

for any $(a, b) \in \mathbf{E}(\nu, \mathbf{r})$ and $(u, v) \in \mathbf{Tan}M, a \times \mathbf{Nor}M, a$ and where

$$P(a, b) = a \quad \text{and} \quad Q(a, b) = b \quad \text{for } (a, b) \in \mathbf{E}(\nu, \mathbf{r}).$$

Proof. Suppose $(x, y) \in \mathbf{E}(\nu, \mathbf{r})$ and $z = f(x, y)$. Then

- (i) $1 + y \bullet x > 1 - \mathbf{r} \geq 0$ and $|z| = 1 + y \bullet x$;
- (ii) $z - x \in \mathbf{Nor}M, x$;
- (iii) $z \in \mathbf{dmn} \xi$ and $x = \xi(z) = x$;
- (iv) $y = \rho(z)$.

$y \bullet x \leq |y| < \mathbf{r}$ and $|z| = 1 + y \bullet x$ so (i) holds.

$\{x, y\} \subset \mathbf{Nor}M, x$ so (ii) holds.

Let $\alpha(r) = (1+r)^2 - 2\frac{(1+r)^2}{1-r} + 1$ for $r \in [0, 1)$. Then $(1-r)(r^2 - \alpha(r)) = 2(2|y|^2 + |y| + 1) > 0$ so $\alpha(|y|) < |y|^2 < \mathbf{r}^2$. We infer from $1 + y \bullet x \leq 1 + |y|$ and $|x - y| > 1 - |y|$ that

$$\begin{aligned} |z - x|^2 &= \left| (1 + y \bullet x)^2 - 2(1 + y \bullet x) \frac{x + y}{|x + y|} \bullet x + 1 \right|^2 \\ &= (1 + |y|)^2 - 2\frac{(1 + |y|)^2}{1 - |y|} + 1 \\ &= \alpha(|y|) \\ &< \mathbf{r}^2 \end{aligned}$$

so that $\xi(z) = x$ and (iii) holds.

(i) implies that

$$z = |z| \frac{x + y}{|x + y|};$$

Taking the inner product with $\xi(z) = x$ we find that

$$z \bullet \xi(z) = \frac{1 + y \bullet x}{|x + y|} = \frac{|z|}{|x + y|}$$

so that $|x + y| = \frac{|z|}{z \bullet \xi(z)}$ and

$$z = |z| \frac{z \bullet \xi(z)}{|z|} (x + y) = (z \bullet \xi(z))(\xi(z) + y)$$

so (iv) holds.

We have shown that $(\xi, \rho) \circ f(x, y) = (x, y)$ for $(x, y) \in \mathbf{E}(\nu, \mathbf{r})$. Thus f is univalent and, by the Chain Rule, $Df(x, y)$ carries $\mathbf{TanE}(\nu, \mathbf{r})$, (x, y) isomorphically onto \mathbb{R}^n so f is an open mapping. The Theorem follows. \square

7.1. γ . Let

$$\gamma(x) = \nu(x) - \iota(x) \quad \text{for } x \in M;$$

evidently, $\gamma : M \rightarrow \mathbb{O}_{n-m-1}(\mathbb{R}^n)$, γ is smooth and $\mathbf{rng} \gamma(x) = \mathbf{Nor} M, x \cap \mathbf{TanS}^{n-1}, x$ for any $x \in M$.

Proposition 7.2. *Suppose $a \in M$, $u \in \mathbf{Tan} M, a$ and $y \in \mathbf{Nor} M, a$. Then*

$$(16) \quad D\gamma(a)(u)(y) = -A(a)(y)(u) - \begin{cases} u & \text{if } y = a, \\ 0 & \text{if } y \in \mathbf{rng} \gamma(a) \end{cases}$$

and

$$(17) \quad DW(a) = D_\gamma W(a) - A(a)(W(a)) \quad \text{for } W \in \mathcal{E}(\gamma).$$

Moreover,

$$(18) \quad \mathbf{TanE}(\gamma), (a, v) = \{(u, A(a)(a)(u)) : u \in \mathbf{Tan} M, a\} \oplus \{(0, w) : w \in \mathbf{rng} \gamma(a)\}.$$

Proof. We have

$$\gamma(x)(y) = \nu(x)(y) - (y \bullet x)x \quad \text{for } x \in M \text{ and } y \in \mathbb{R}^n.$$

For $u \in \mathbf{Tan} M, a$ we have

$$D\gamma(a)(u)(y) = -A(a)(y)(u) - (y \bullet u)a - (y \bullet a)u.$$

Since

$$(y \bullet u)a + (y \bullet a)u = \begin{cases} u & \text{if } y = a, \\ 0 & \text{if } y \in \mathbf{rng} \gamma(a) \end{cases}$$

(4) implies (16). (18) now follows from (3). \square

The following Theorem is a Corollary of Theorem 7.1.

Theorem 7.2. *f carries $\mathbf{E}(\gamma, \mathbf{r})$ diffeomorphically onto $V \cap \mathbb{S}^{n-1}$.*

Proposition 7.3. *Suppose $x \in M$. Then*

$$D\rho(x)(w) = \gamma(x)(w) \quad \text{whenever } w \in \mathbf{Nor}M, x \cap \mathbf{Tan}\mathbb{S}^{n-1}, x.$$

Proof. We have $D\xi(x) = \tau(x)$.

Suppose $w \in \mathbf{Nor}M, x \cap \mathbf{Tan}\mathbb{S}^{n-1}, x$. Then

$$Dv(x)(w) = w \bullet x + x \bullet D\xi(x)(w) = 0.$$

so, as $v(x) = 1$,

$$D\rho(x)(w) = w - D\xi(x)(w) = \nu(x)(w) = \gamma(x)(w).$$

\square

7.2. *F.* Whenever $W \in \mathcal{E}(\gamma, \mathbf{r})$ we define

$$F(W) : M \rightarrow \mathbb{S}^{n-1}$$

by letting $F(W) = f \circ (i_M, W)$; it follows from Theorem 7.1 that $F(W)$ carries M diffeomorphically onto its range; in particular,

$$(19) \quad \mathbf{rng} F(W) \in \mathcal{M}_m(\mathbb{R}^n), \quad \xi \circ F = i_M \quad \text{and} \quad \rho \circ F(W) = W.$$

8. FAMILIES OF SUBMANIFOLDS OF \mathbb{S}^{n-1} .

We suppose throughout this section that \mathcal{J} is a Euclidean space. For each $\epsilon \in \mathbb{R}^+$ and $s \in \mathcal{J}$ we let

$$\mathcal{J}(\epsilon) = \{t \in \mathcal{J} : |t| < \epsilon\} \quad \text{and we let} \quad \mathcal{J}(s, \epsilon) = \{t \in \mathcal{J} : |s - t| < \epsilon\}.$$

We suppose

- (i) \mathcal{S} is a open subset of \mathcal{J} and $0 \in \mathcal{S}$;
- (II) $M \in \mathcal{M}_m(\mathbb{R}^n)$, $M \subset \mathbb{S}^{n-1}$ and M is compact;
- (III) $\mathbf{z} : \mathcal{S} \times M \rightarrow \mathbb{S}^{n-1}$ and \mathbf{z} is smooth;
- (IV) \mathbf{z}_s is a diffeomorphism for each $s \in \mathcal{S}$;
- (IV) $\mathbf{z}_0(p) = p$ for $p \in M$.

Let

\mathbf{r}

be the reach of M .

Let

$$M_s = \mathbf{z}_s[M] \quad \text{for each } s \in \mathcal{S};$$

it follows that $M_s \in \mathcal{M}_m(\mathbb{R}^n)$, M_s is compact and $M_s \subset \mathbb{S}^{n-1}$.

We let

$$T : \mathcal{S} \times M \rightarrow \mathcal{L}_m$$

be such that $T(s, p) = \mathcal{P}_m(D\mathbf{z}_s(p))$ for $(s, p) \in \mathcal{S} \times M$.

For each $s \in \mathcal{S}$ we let

$$\tau_s, \nu_s, B_s, A_s \quad \text{be as in Section 3;}$$

$$\begin{aligned} \mathbf{r}_s, \Xi_s, N_s(r), r \in \mathbb{R}^+, \xi_s, \sigma_s & \text{ be as in Section 5;} \\ \nu_s, \rho_s, f_s, V_s, F_s & \text{ be as in Section 7} \end{aligned}$$

with M replaced by M_s .

We have

$$T(s, p) = \tau_s(\mathbf{z}_s(p)) \quad \text{for } (s, p) \in \mathbf{S} \times M.$$

8.1. ϵ_1 . We fix

$$\epsilon_1 \in \mathbb{R}^+$$

such that the closure of $\mathcal{J}(\epsilon_1)$ is a compact subset of \mathcal{S} .

For each $(j, k) \in \mathbb{N} \times \mathbb{N}$ we let

$$(20) \quad C_{j,k} = \chi^{j,k}(\mathbf{z}(\mathcal{J}(\epsilon_1) \times M)) \quad \text{for any } (j, k) \in \mathbb{N}.$$

and note that $C_{j,k} < \infty$.

Theorem 8.1. *There is a constant C_T such that*

$$(21) \quad \|T(s, p) - T(t, q)\| \leq C_T(\epsilon)(|s - t| + |p - q|) \quad \text{whenever } s, t \in \mathcal{J}(\epsilon_1) \text{ and } p, q \in M$$

Proof. Suppose $s, t \in \mathcal{J}(\epsilon_1)$ and $p, q \in M$. By (20) we have

$$T(s, p) - T(t, p) = \int_0^1 \frac{d}{d\lambda} T((1 - \lambda)t + \lambda s, p) d\lambda = \int_0^1 DT((1 - \lambda)t + \lambda s, p)(s - t, 0) d\lambda$$

and, by Theorem 5.5,

$$\|T(t, p) - T(t, q)\| \leq \left(1 + \frac{1}{\mathbf{r}}\right) \chi^{[1]}(T_t) |p - q|.$$

□

Let

$$\underline{\mathbf{r}} = \inf\{\mathbf{r}_s : s \in \mathcal{J}(\epsilon_1)\} \quad \text{and let} \quad \underline{\mathbf{b}} = \sup\{\|B_s\| : s \in \mathcal{J}(\epsilon_1)\}.$$

Theorem 8.2. *We have*

$$0 < \underline{\mathbf{r}} \leq 1 \quad \text{and} \quad \underline{\mathbf{b}} \leq \frac{1}{\underline{\mathbf{r}}}.$$

Proof. Let \mathcal{H} be the closure of $\mathcal{J}(\epsilon_1)$. Let

$$Y_1 = (\mathcal{J} \oplus \mathbb{R}^n) \oplus \mathbb{R}^n, \quad Y_2 = \mathcal{J} \oplus \mathbb{R}^n, \quad N_1 = \mathbf{E}(T), \quad N_2 = \mathcal{J} \oplus \mathbb{R}^n,$$

let $F : N_1 \rightarrow N_2$ be such that $F((s, p), v) = \mathbf{z}(s, p) + v$ for $((s, p), v) \in \mathbf{E}(T)$. Let $K = \{((s, p), 0) : (s, p) \in \mathcal{H} \times M\}$ and for each $\epsilon \in \mathbb{R}^+$ let

$$U_\epsilon = \{((s, p), v) \in \mathbf{E}(T) : \mathbf{dist}(s, K) < \epsilon \text{ and } |v| < \epsilon\}.$$

Keeping in mind Theorem 4.2 we find that (i)-(iv) of Lemma 5.1 hold. Let ϵ_* be as in (v) of Lemma 5.1. Then $\epsilon_* \leq \mathbf{r}_s$ for any $s \in \mathcal{J}(\epsilon_1)$.

$\underline{\mathbf{b}} \leq 1/\underline{\mathbf{r}}$ since $\|B_s\| \mathbf{r}_s \leq 1$ for $s \in \mathcal{J}$. □

For each $(s, t) \in \mathcal{J}(\epsilon_1)$ we let

$$\mathbf{c}_{s,t} : [0, 1] \rightarrow \mathcal{J}(\epsilon_1)$$

be such that

$$\mathbf{c}_{s,t}(\lambda) = t + \lambda(s - t) \text{ for } \lambda \in [0, 1].$$

Proposition 8.1. *Suppose $s, t \in \mathcal{J}(\epsilon_1)$. Then*

$$|\mathbf{z}_s(p) - \mathbf{z}_t(p)| \leq C_{1,0}|s - t| \quad \text{for any } p \in M.$$

Moreover,

$$M_t \subset N_s(r) \quad \text{if } C_{1,0}|s - t| < r \leq \underline{\mathbf{r}}.$$

Proof. For any $p \in M$ we have

$$|\mathbf{z}_s(p) - \mathbf{z}_t(p)| \left| \int_0^1 \frac{d}{d\lambda} \mathbf{z}(\mathbf{c}_{s,t}(\lambda), p) d\lambda \right| = \left| \int_0^1 D\mathbf{z}(\mathbf{c}_{s,t}(\lambda), p)(s - t, 0) d\lambda \right| \leq C_{1,0}|s - t|.$$

Suppose $z \in M_t$. Let $p \in M$ be such that $\mathbf{z}_t(p) = z$. Then

$$\mathbf{dist}(z, M_s) \leq |\mathbf{z}_t(p) - \mathbf{z}_s(p)| \leq C_{1,0}|s - t|.$$

□

Let

$$\delta_1 \in \mathbb{R}^+$$

be such that $C_{1,0}\delta_1 < \underline{\mathbf{r}}$.

Corollary 8.1. *If $s, t \in \mathcal{J}(\epsilon_1)$ and $|s - t| < \delta_1$ then*

$$M_t \subset N_s(\underline{\mathbf{r}}).$$

We define

$$\mu : \mathcal{S} \times M \rightarrow \mathbb{R}^+$$

by letting

$$\mu(s, p) = \sup\{\lambda \in [0, \infty) : |D\mathbf{z}_s(p)(u)| \geq \lambda|u| \text{ for } u \in \mathbb{S}^{n-1} \cap \mathbf{Tan}M, p\}.$$

We let

$$\langle \mu \rangle = \inf\{\mu(s, p) : (s, p) \in \mathcal{J}(\epsilon_1) \times M\}$$

We leave as an elementary exercise for the reader to prove the following Proposition.

Proposition 8.2. *μ is positive and continuous and $\langle \mu \rangle$ is positive.*

Proposition 8.3. *There is a constant*

$$C_\mu$$

such that if $s, t \in \mathcal{J}(\epsilon_1)$ and $|s - t| < \delta_1$ then $M_t \subset N_s(\underline{\mathbf{r}})$ and

$$\|T(t, p) - \tau_s(\xi_s(\mathbf{z}_t(p)))\| \leq C_\mu|s - t| \quad \text{for } p \in M.$$

Proof. Suppose $s, t \in \mathcal{J}(\epsilon_1) \cap \mathcal{J}(\delta_1)$. From Corollary 8.1 we infer that

$$M_t \subset N_s(\underline{\mathbf{r}}) \subset \mathbf{dmn} \xi_s.$$

Suppose $p \in M$. Then

$$\|D\mathbf{z}_s^{-1}(p)\| \leq \frac{1}{\mu(s, p)} \leq \frac{1}{\langle \mu \rangle}.$$

Let $q \in M$ be such that $\xi_s(\mathbf{z}_t(p)) = \mathbf{z}_s(q)$. If

$$c = \left(1 + \frac{1}{\underline{\mathbf{r}}}\right) \left(\frac{1}{\langle \mu \rangle}\right)$$

we infer from Theorem 5.5 that

$$\begin{aligned}
|p - q| &= |\mathbf{z}_s^{-1}(\mathbf{z}_s(p)) - \mathbf{z}_s^{-1}(\mathbf{z}_s(q))| \\
&\leq \text{Lip}(\mathbf{z}_s^{-1}) |\mathbf{z}_s(p) - \mathbf{z}_s(q)| \\
&\leq \left(1 + \frac{1}{\mathbf{r}_s}\right) \left(\frac{1}{\underline{\mu}}\right) |\mathbf{z}_s(p) - \mathbf{z}_s(q)| \\
&\leq c |\mathbf{z}_s(p) - \mathbf{z}_s(q)| \\
&\leq c (|\mathbf{z}_s(p) - \mathbf{z}_t(p)| + |\mathbf{z}_t(p) - \xi_s(\mathbf{z}_t(p))|) \\
&\leq 2c |\mathbf{z}_s(p) - \mathbf{z}_t(p)| \\
&\leq 2cC_{1,0} |s - t|.
\end{aligned}$$

Thus

$$\begin{aligned}
\|T(t, p) - \tau_s(\xi_s(\mathbf{z}_t(p)))\| &= \|T(t, p) - T(s, q)\| \\
&\leq \|T(s, p) - T(t, p)\| + \|T(t, p) - T(t, q)\| \\
&\leq C_{1,0} \chi^1(T, \mathcal{J}(\epsilon_1) \times M) |s - t| + C_T |p - q| \\
&\leq C_\mu |s - t|
\end{aligned}$$

where

$$C_\mu = \chi^1(T, \mathcal{J}(\epsilon_1) \times M) |s - t| + C_T (2cC_{1,0}).$$

□

Theorem 8.3. *Suppose $s \in \mathcal{J}(\epsilon_1)$. There is $\delta \in (0, \epsilon_1 - |s|]$ such that if $t \in \mathcal{J}(s, \delta)$ then $t \in \mathcal{J}(\epsilon_1)$, $M_t \subset N_s(\underline{\mathbf{x}})$ and*

$$\Lambda_t \text{ carries } M_t \text{ diffeomorphically onto } M_s$$

where

$$\Lambda : \mathcal{J}(s, \delta) \times M \rightarrow M$$

is such that

$$\Lambda(t, q) = \mathbf{z}_s^{-1} \circ \xi_s \circ \mathbf{z}_t(q) \quad \text{for } (t, q) \in \mathcal{J}(s, \delta) \times M.$$

Proof. That $s, t \in \mathcal{J}(\epsilon_1)$ and $|s - t| < \delta_1$. Then $M_t \subset N_s(\underline{\mathbf{x}})$ by Corollary 8.1.

For each $(t, x) \in \mathcal{J}(s, \delta_1) \times M_t$ we let $\mathcal{N}(t, x) = \{z \in M_t : \xi_s(z) = x\}$ and we let $N(t, x)$ equal the cardinality of $\mathcal{N}(t, x)$.

We choose $\delta_2 \in (0, \delta_1]$ such that $C_\mu \delta_2 < 1$ where C_μ is as in Proposition 8.3. Suppose $p \in M$. By Proposition 8.3 we have

$$\|T(t, p) - \tau_s(\xi_s(\mathbf{z}_t(p)))\| < 1 \quad \text{if } t \in \mathcal{J}(s, \delta_2).$$

It follows from (ii) of Theorem 5.3 that $\mathbf{Tan}M_t, \mathbf{z}_t(p)$ meets $\ker D\xi_s(\mathbf{z}_t(p))$ transversely so that

$$D\xi_s(\mathbf{z}_t(p)) \text{ carries } \mathbf{Tan}M_t, \mathbf{z}_t(p) \text{ isomorphically onto } \mathbf{Tan}M_s, \xi_s(\mathbf{z}_t(p)).$$

This implies that there is an open neighborhood of $\mathbf{z}_t(p)$ such that \mathbf{z}_s carries $U \cap M_t$ diffeomorphically onto an open neighborhood of $\mathbf{z}_s(\mathbf{z}_t(p))$ in M_s . Since M_t is compact we find that if $x \in \mathbf{rng} \xi_s|_{M_t}$ then $N_t(x)$ is finite and there are an open neighborhood U of x in M_s and a map \mathcal{U} with domain $\mathcal{N}_t(x)$ such that $\mathcal{U}(z)$ is an open neighborhood of z in M_t which ξ_s carries diffeomorphically onto U . It follows that $\mathcal{N}(t, x)$ is finite and that N_t is locally constant.

We choose $\delta_3 \in (0, \delta_2]$ as follows. Let \mathcal{C} be the family of connected components of M_s ; since M_s is compact \mathcal{C} is finite. For $x \in M_s$ we let $C_x \in \mathcal{C}$ be such that $x \in C_x$. If \mathcal{C} has one member we let $\delta_2 = \delta_1$. Otherwise we let

$$d = \inf\{|x - y| : x, y \in M_s \text{ and } C_x \neq C_y\}.$$

We choose $\delta_3 \in (0, \delta_2]$ such that $C_{1,0}\delta_2 < d$. Suppose $|s - t| < \delta_3$ and $x \in M_s$. There is $z \in M_t$ such that $|x - z| = \mathbf{dist}(x, M_t) \leq C_{1,0}|s - t| < d$. Hence $|\xi_s(z) - x| = |\mathbf{dist} z, M_s| \leq |x - z| < d$ so $z \in C_x$ and $N_t(\xi_s(z)) \geq 1$. Since N_t is locally constant we must have $N_t(x) \geq 1$. Thus $N_t(x) \geq 1$ for all $x \in M_s$.

Let $\mathbf{z}_s^t = \mathbf{z}_t \circ \mathbf{z}_s^{-1}$ for $t \in \mathcal{J}(s, \delta_2)$.

$$\begin{aligned} \int N_t d\|M_s\| &= \int \mathcal{W}_m(D\xi_s(z) \circ \tau_t(z)) d\|M_t\|z \\ &= \int \mathcal{W}_m(D\xi_s(\mathbf{z}_s^t(x))) \mathcal{W}_m(D\mathbf{z}_s^t(x)) d\|M_s\|x \quad \rightarrow \quad \|M_s\|(\mathbb{R}^n) \end{aligned}$$

since

$$\limsup_{t \rightarrow s} \{|\mathcal{W}_m(D\xi_s(\mathbf{z}_s^t(x))) \mathcal{W}_m(D\mathbf{z}_s^t(x)) - 1| : x \in M_s\} = 0.$$

Thus there is $\delta \in (0, \delta_3]$ such that

$$N_t(x) = 1 \quad \text{if } t \in \mathcal{J}(s, \delta) \text{ and } x \in M_s.$$

It follows that Λ_t carries M_s diffeomorphically onto M_t for such t . □

8.2. M . We let

$$\underline{M} = \{(s, z) : s \in (0, \epsilon_1) \text{ and } z \in M_s\}.$$

Theorem 8.4. $\underline{M} \in \mathcal{M}_{j+m}(\mathcal{J} \oplus \mathbb{R}^n)$.

Proof. Suppose $s \in \mathcal{J}(\epsilon_1)$. Let $\delta = \epsilon_1 - |s|$. Let

$$\Psi : \mathcal{J}(s, \delta) \times N_s(\mathbf{r}) \rightarrow \mathcal{J}(s, \delta) \times N_s(\mathbf{r})$$

be such that $\Psi(t, z) = (t, \mathbf{z}_s^{-1}(\xi_s(z)))$ for $(t, z) \in \mathcal{J}(s, \delta) \times N_s(\mathbf{r})$ and note that Ψ is smooth.

Let δ and Λ be as in Theorem 8.3; note that Λ is smooth and let

$$\Gamma : \mathcal{J}(s, \delta) \times M \rightarrow \mathcal{J}(s, \delta) \times M$$

be such that $\Gamma_t = \Lambda_t^{-1}$ for $t \in \mathcal{J}(s, \delta)$ and note that Γ is smooth.

Let

$$\Phi : \mathcal{J}(s, \delta) \times M \rightarrow \mathcal{J}(s, \delta) \times N_s(\mathbf{r})$$

be such that

$$\Phi(t, p) = (t, \mathbf{z}_t(\Lambda_t^{-1}(p))) \quad \text{for } (t, p) \in \mathcal{J}(s, \delta) \times M.$$

Note that Φ is smooth.

If $(t, p) \in \mathcal{J}(s, \delta) \times M$ then

$$\begin{aligned} \Psi(\Phi(t, p)) &= \Psi(t, \mathbf{z}_t(\Lambda_t^{-1}(p))) \\ &= (t, \mathbf{z}_t(\mathbf{z}_s^{-1}(\xi_s(\Lambda_t^{-1}(p)))) \\ &= (t, p). \end{aligned}$$

The Theorem now follows from Proposition 3.3. □

8.3. $\underline{\tau}, \underline{\nu}, \underline{\gamma}, \underline{B}, \underline{A}$. We define smooth maps

$$\underline{\tau} : \underline{M} \rightarrow \mathbb{O}_m(\mathbb{R}^n); \quad \underline{\nu} : \underline{M} \rightarrow \mathbb{O}_{n-m}(\mathbb{R}^n); \quad \underline{\gamma} : \underline{M} \rightarrow \mathbb{O}_{n-m-1}(\mathbb{R}^n)$$

by requiring that

$$\underline{\tau}(s, x)(u) = Di_{\underline{M}}(x, s)(u, 0) \quad \text{for } (s, x) \in \underline{M} \text{ and } u \in \mathbb{R}^n$$

and letting

$$\underline{\nu}(s, x) = i_{\mathbb{R}^n} - \underline{\tau}(x, s); \quad \underline{\gamma}(s, x) = \underline{\nu}(s, x) - \iota(x)$$

for $(s, x) \in \underline{M}$ and $u \in \mathbb{R}^n$. It follows that

$$\underline{\tau}(s, x) = \tau_s(x); \quad \underline{\nu}(s, x) = \nu_s(x); \quad \underline{\gamma}(s, x) = \gamma_s(x)$$

for $(x, s) \in \underline{M}$.

We define the smooth maps

$$\underline{B} : \underline{M} \rightarrow \text{MultiLin}(\mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n) \quad \text{and} \quad \underline{A} : \underline{M} \rightarrow \text{Lin}(\mathbb{R}^n, \text{End}(\mathbb{R}^n))$$

by letting

$$\underline{B}(s, x)(u, v) = \underline{\nu}(s, x)(D\underline{\tau}(s, x)(0, u)(0, v)) \quad \text{whenever } (s, x) \in \underline{M} \text{ and } u, v \in \mathbb{R}^n$$

and requiring that

$$\underline{A}(s, x)(u)(v) \bullet w = u \bullet \underline{B}(s, x)(v, w) \quad \text{whenever } (s, x) \in \underline{M} \text{ and } u, v, w \in \mathbb{R}^n.$$

It follows that

$$\underline{B}(s, x)(u, v) = B_s(x)(u, v) \quad \text{and} \quad \underline{A}(s, x)(u) = A_s(x)(u)$$

whenever $(s, x) \in \underline{M}$ and $u, v \in \mathbb{R}^n$.

8.4. $\underline{N}, \underline{\Xi}, \underline{\xi}, \underline{\sigma}$. For each $r \in (0, \mathbf{r}]$ we let

$$\underline{N}(r) = \{(s, z) \in \mathcal{J}(\epsilon_1) \times \mathbb{R}^n : \mathbf{dist}(z, M_s) < r\}.$$

We let

$$\underline{\Xi} : \mathbf{E}(\underline{\nu}, \mathbf{r}) \rightarrow \underline{N}(\mathbf{r})$$

be such that

$$\underline{\Xi}((s, x), v) = (s, \Xi_s(x, v)) \quad \text{for } ((s, x), v) \in \mathbf{E}(\underline{\nu}, \mathbf{r}).$$

It follows from Theorem 5.2 that $\underline{\Xi}$ carries $\mathbf{E}(\underline{\nu}, \mathbf{r})$ diffeomorphically onto $\underline{N}(\mathbf{r})$; in particular, $\underline{N}(r)$ is an open subset of $\mathcal{J} \times \mathbb{R}^n$ whenever $0 < r \leq \mathbf{r}$. Since $\underline{\Xi}$ carries $\mathbf{E}(\underline{\nu}, \mathbf{r})$ diffeomorphically onto $\underline{N}(\mathbf{r})$ we may define smooth functions

$$\underline{\xi} : \underline{N}(\mathbf{r}) \rightarrow \mathbb{R}^n \quad \text{and} \quad \underline{\sigma} : \underline{N}(\mathbf{r}) \rightarrow \mathbb{R}^n$$

by requiring that

$$(\underline{\xi}, \underline{\sigma}) = (\underline{\Xi}|_{\mathbf{E}(\underline{\nu}, \mathbf{r})})^{-1} : \underline{N}(\mathbf{r}) \rightarrow \mathbf{E}(\underline{\nu}, \mathbf{r}).$$

Note that

$$(22) \quad (s, z) = \underline{\Xi}(\underline{\xi}(s, z), \underline{\sigma}(s, z)) = (s, \underline{\xi}(s, z) + \underline{\sigma}(s, z)) \quad \text{for } (s, z) \in \underline{N}(\mathbf{r}).$$

8.5. \underline{f} , \underline{v} , $\underline{\rho}$, \underline{V} , \underline{g} . We define the smooth maps

$$\underline{f} : \mathbf{E}(\underline{v}, \mathbf{r}) \rightarrow \mathcal{J} \times \mathbb{R}^n; \quad \underline{v} : \underline{M}(\mathbf{r}) \rightarrow \mathbb{R}^+; \quad \underline{\rho} : \underline{M}(\mathbf{r}) \rightarrow \mathbb{R}^n$$

by letting

$$\underline{f}((t, z), v) = (t, f_t(x, v)); \quad \underline{v}((t, z), v) = (t, v_t(x, v)); \quad \underline{\rho}((t, z), v) = (t, \rho_t(x, v))$$

for $((t, x), v) \in \mathbf{E}(\underline{v})$. We also set

$$\underline{V} = \mathbf{rng} \underline{f}.$$

Theorem 8.5. \underline{V} is an open subset of $\mathcal{J} \times \mathbb{R}^n$ and \underline{f} carries $\mathbf{E}(\underline{v}, \mathbf{r})$ diffeomorphically onto \underline{V} with inverse $(\underline{\xi}, \underline{\rho})$.

Moreover, \underline{f} carries $\mathbf{E}(\underline{\gamma}, \mathbf{r})$ diffeomorphically onto $\underline{V} \cap \mathbb{S}^{n-1}$.

Proof. This follows from the results of Section 7. \square

For $r \in (0, \mathbf{r})$ we let

$$\underline{Q}(r) = \{(s, t, z) \in \mathcal{J}(\epsilon_1) \times \mathcal{J}(\epsilon_1) \times \mathbb{R}^n : (s, z) \in \underline{N}(r) \text{ and } (t, z) \in \underline{N}(r)\}.$$

Since $\underline{N}(r)$ is an open subset of $\mathcal{J} \times \mathbb{R}^n$ it follows that $\underline{Q}(r)$ is an open subset of $\mathcal{J} \times \mathcal{J} \times \mathbb{R}^n$.

Lemma 8.1. Suppose $s, t \in \mathcal{J}(\epsilon_1)$, $q, r \in \mathbb{R}^+$, $q + r \leq \mathbf{r}$ and $(s, t, z) \in \underline{Q}(r)$. If $\chi^1(\mathbf{z})|s - t| \leq q$ then

$$\{(\mathbf{c}_{s,t}(\lambda), z) : \lambda \in [0, 1]\} \subset \underline{Q}(q + r)$$

Proof. Since $(s, z) \in \underline{N}(r)$, $\mathbf{dist}(z, M_s) < r \leq \mathbf{r} \leq \mathbf{r}_s$ so $z \in \mathbf{dmn} \xi_s$ which implies that there is $w \in M_s$ such that $\mathbf{dist}(z, M_s) = |z - w|$. Suppose $\lambda \in [0, 1]$ and $u = (1 - \lambda)s + \lambda t$. Then

$\mathbf{dist}(z, M_u) \leq |z - \mathbf{z}_u(p)| \leq |z - w| + |\mathbf{z}_s(p) - \mathbf{z}_u(p)| < r + \chi^1(\mathbf{z})\lambda|t - s| \leq r + q$
so $z \in \underline{N}(q + r)$. By the same argument with s replaced by t we find that $z \in \underline{N}(q + r)$. \square

9. F_s^t .

Suppose $s \in \mathcal{S}$ and $W \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$. Let

$$F_s(W) : M_s \rightarrow V_s$$

be such that

$$F_s(W) = f_s \circ (i_{M_s}, W).$$

Then $\mathbf{rng} F_s(W)$ is a smooth compact submanifold of \mathbb{S}^{n-1} , $F_s(W)$ carries M_s diffeomorphically onto $\mathbf{rng} F_s(W)$,

$$\xi_s \circ F_s(W) = i_{M_s} \quad \text{and} \quad \rho_s \circ F_s(W) = 0.$$

Proposition 9.1. Suppose $t \in \mathcal{S}$, $W_i \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$, $i = 1, 2$ and $\mathbf{rng} F_s(W_1) = \mathbf{rng} F_s(W_2)$. Then $W_1 = W_2$.

Proof. Suppose $x_1 \in M_s$. There is $x_2 \in M_s$ such that $F_s(W_2)(x_2) = F_s(W_1)(x_1)$. Thus

$$f_s(x_1, W_1(x_1)) = F_s(W_1)(x_1) = F_s(W_2)(x_2) = f_s(x_2, W_2(x_2)).$$

Applying g_s we find that $(x_1, W_1(x_1)) = (x_2, W_2(x_2))$. \square

Proposition 9.2. *Suppose $s \in \mathcal{S}$, $X \in \mathcal{M}_m(\mathbb{R}^n)$ and $X \subset V_s$. There is $W \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$ such that $\mathbf{rng} F_s(W) = X$ if and only if $\xi_s|X$ carries X diffeomorphically onto M_s .*

Proof. Suppose $W \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$ and $\mathbf{rng} F_s(W) = X$. Then $F_s(W) = f_s \circ (i_{M_s}, W)$ so

$$(\xi_s \circ F_s(W), \rho_s \circ F_s(W)) = g_s \circ F_s(W) = g_s \circ f_s \circ (i_{M_s}, W) = (i_{M_s}, W)$$

so $\xi_s \circ F_s(W) = i_{M_s}$. It follows that $\xi_s|X$ carries X diffeomorphically onto M_s .

Suppose $\xi_s|X$ carries X diffeomorphically onto M_s . Let $W \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$ be such that $W \circ (\xi_s|X) = \rho_s|X \in \mathcal{E}(M_s, \gamma_s)$. Then

$$F_s(W) \circ (\xi_s|X) = f_s \circ (i_{M_s}, W) \circ (\xi_s|X) = f_s \circ (\xi_s|X, \rho_s|X) = i_X$$

so $\mathbf{rng} F_s(W) = X$. □

Definition 9.1. *For $s, t \in \mathcal{S}$ we let*

$$F_s^t = \{(W, Z) \in \mathcal{E}(\gamma_s, \mathbf{r}_s) \times \mathcal{E}(\gamma_t, \mathbf{r}_t) : \mathbf{rng} F_s(W) = \mathbf{rng} F_t(Z)\}.$$

Proposition 9.3. *Suppose $s, t \in \mathcal{S}$, $W \in \mathcal{E}(\gamma_s, \mathbf{r}_s)$ and $Z \in \mathcal{E}(\gamma_t, \mathbf{r}_t)$. The following statements are equivalent.*

- (I) $(W, Z) \in F_s^t$;
- (II) $\xi_t \circ F_s(W)$ carries M_s diffeomorphically onto M_t and

$$(23) \quad Z \circ \xi_t \circ F_s(W) = \rho_t \circ F_s(W).$$

Proof. Suppose (I) holds. Let $g = F_t(Z)^{-1} \circ F_s(W)$. Then g carries M_s diffeomorphically onto M_t and $F_t(Z) \circ g = F_s(W)$ so for any $y \in M_t$ we have

$$(i_{M_t}, Z) \circ g = (\xi_t, \rho_t) \circ F_t(Z) \circ g = (\xi_t, \rho_t) \circ F_s(W)$$

so $g = \xi_t \circ F_s(W)$ and $Z \circ g = \rho_t \circ F_s(W)$ so (II) holds.

Suppose (II) holds. Let $X = \mathbf{rng} F_s(W)$. (23) implies that $Z \circ (\xi_t|X) = \rho_t|X$. So

$$F_t(Z) \circ (\xi_t|X) = f_t \circ (i_{M_t}, Z) \circ (\xi_t|X) = f_t \circ (\xi_t|X, \rho_t|X) = f_t \circ g_t \circ i_X = i_X$$

so $\mathbf{rng} F_t(Z) = \mathbf{rng} i_X = X = \mathbf{rng} F_s(W)$. □

Corollary 9.1. *If $s, t \in \mathcal{S}$ then $0 \in \mathbf{dmn} F_s^t$ if and only if ξ_t carries M_s diffeomorphically onto M_t .*

Proposition 9.4. *Suppose $s \in \mathcal{S}$ and $W \in \mathcal{E}(\gamma_s)$. Then*

$$\mathbf{Lip} W \leq \left(1 + \frac{8}{\mathbf{r}_s}\right) \chi^{[1]}(W) \quad \text{and} \quad \mathbf{Lip} F_s(W) \leq 1 + \left(1 + \frac{8}{\mathbf{r}_s}\right) \chi^{[1]}(W)$$

Proof. The first inequality follows from Proposition 5.5.

Suppose $x_i \in M_s$, $i = 1, 2$. Using Proposition 6.1 we find that

$$\begin{aligned} |F_s(W)(x_1) - F_s(W)(x_2)| &= |\boldsymbol{\eta}(x_1 + W(x_2)) - \boldsymbol{\eta}(x_2 + W(x_2))| \\ &\leq |x_1 - x_2| + |W(x_1) - W(x_2)| \end{aligned}$$

so the second inequality holds. □

Proposition 9.5. *Suppose E is a Euclidean space, $h : \underline{N}(\mathbf{r}) \rightarrow E$, h is smooth and*

$$\square h : \underline{Q}(\mathbf{r}/2) \rightarrow \mathbb{L}\text{in}(\mathcal{J}, E)$$

is such that

$$\square h(s, t, z)(u) = \int_0^1 D^{1,0}h(\mathbf{c}_{s,t}(\lambda), z)(u) d\lambda \quad \text{for } (s, t, z) \in \underline{Q}(\mathbf{r}/2) \text{ and } u \in \mathcal{J}.$$

Then $\square h$ is smooth and

$$(24) \quad h(s, z) - h(t, z) = \square h(s, t, z)(s - t) \quad \text{for } (s, t, z) \in \underline{Q}(\mathbf{r}/2).$$

If $(s, t, z_i) \in \underline{Q}(\mathbf{r}/2)$ for $i = 1, 2$ then

$$(25) \quad \|\square h(s, t, z_1) - \square h(s, t, z_2)\| \leq C_{1,\square}(h)|z_1 - z_2|$$

where we have set

$$C_{1,\square}(h) = \max \left\{ \chi^2(h), 8 \frac{\chi^0(h)}{\mathbf{r}} \right\}.$$

Proof. Since $\underline{Q}(\mathbf{r})$ is an open subset of $\mathcal{J}(\epsilon_1) \times \mathcal{J}(\epsilon_1) \times \mathbb{R}^n$ it is elementary that $\square h$ is smooth. (24) follows since

$$h(s, z) - h(t, z) = \int_0^1 \frac{d}{d\lambda} h(\mathbf{c}_{s,t}(\lambda), z) d\lambda = \int D^{1,0}h(\mathbf{c}_{s,t}(\lambda), z)(s - t) d\lambda$$

whenever $(s, t, z) \in \underline{Q}(\mathbf{r}/2)$.

Suppose $(s, t, z_i) \in \underline{Q}(\mathbf{r}/2)$ for $i = 1, 2$. If $|z_2 - z_1| < \mathbf{r}/4$ then Corollary 8.1 implies that $(s, t, w) \in \underline{Q}(\mathbf{r}/2)$ if $w \in \mathbf{U}(z_1, \mathbf{r}/4)$ so that

$$\|\square h(s, t, z_1) - \square h(s, t, z_2)\| \leq \chi^1(\square h)|z_1 - z_2| \leq \chi^2(h)|z_1 - z_2|.$$

On the other hand, if $|z_1 - z_2| \geq \mathbf{r}/4$ then

$$\|\square h(s, t, z_1) - \square h(s, t, z_2)\| \leq 2\chi^1(h) \leq 8 \frac{\chi^1(h)}{\mathbf{r}}|z_1 - z_2|.$$

So (25) holds. □

9.1. ϵ_2 and \mathcal{G} . Let

$$\epsilon_2 \in (0, \epsilon_1]$$

be such that

$$C_{1,\square}(\underline{\xi})(2\epsilon_2) < \frac{1}{2}.$$

Let

$$\mathcal{G} = \left\{ (s, W) \in \mathcal{J}(\epsilon_1) \times \mathcal{E}(\gamma_s) : \left(1 + \frac{8}{\mathbf{r}}\right) \chi^{[1]}(W) < 1 \right\}.$$

Proposition 9.6. *Suppose $(s, W) \in \mathcal{G}$. Then*

$$\chi^0(W) < \frac{\mathbf{r}}{8}, \quad \mathbf{Lip} W < 1 \quad \text{and} \quad \mathbf{Lip} F_s(W) < 2.$$

Proof. $\chi^0(W) \leq \chi^{[1]}(W) < \mathbf{r}/8$. The second and third inequalities follow directly from Proposition 9.4. □

Proposition 9.7. *Suppose $s, t \in \mathcal{J}(\epsilon_2)$ and $(s, W) \in \mathcal{G}$. Then*

$$(26) \quad \{(s, t, F_s(W)(x)) : x \in M_s\} \subset \underline{Q}(\mathbf{r}/2);$$

in particular, $\text{rng } F_s(W) \subset \text{dmn } \xi_t$.

Proof. Suppose $x \in M_s$. There is $p \in M$ such that $\mathbf{z}_s(p) = x$. By Proposition 6.3 we have

$$\mathbf{dist}(F_s(W)(x), M_s) \leq |\boldsymbol{\eta}(x + W(x)) - x| \leq \chi^0(W) \leq \frac{\mathbf{r}}{4}$$

and

$$\mathbf{dist}(F_s(W)(x), M_t) \leq |\boldsymbol{\eta}(x + W(x)) - x| + |\mathbf{z}(s, p) - \mathbf{z}(t, p)| \leq \chi^0(W) + C_{1,0}|s - t| \leq \frac{\mathbf{r}}{2}$$

so (26) holds and, since $s, t \in \mathcal{J}(\epsilon_2)$, $\mathbf{rng} F_s(W) \subset \mathbf{dmn} \xi_t$ since $C_{1,\square}(\underline{\xi})(2\epsilon_2) < 1/2$. \square

For $(s, W) \in \mathcal{G}$ and $(t, x) \in \underline{Q}(\mathbf{r}/2)$ we let

$$A_s^t(W)(x) = \square \underline{\xi}(s, t, F_s(W)(x))(s - t)$$

and we let

$$B_s^t(W)(x) = \square \underline{\rho}(s, t, F_s(W)(x))(s - t).$$

Theorem 9.1. *Suppose $s, t \in \mathcal{J}(\epsilon_2)$ and $(s, W) \in \mathcal{G}$.*

$$(27) \quad \xi_t(F_s(W)(x)) = x - A_s^t(W)(x);$$

$$(28) \quad \rho_t(F_s(W)(x)) = W(x) - B_s^t(W)(x);$$

$$(29) \quad \begin{aligned} & \frac{1}{2}|x_1 - x_2| \\ & \leq (1 - 2C_{1,\square}(\underline{\xi})|s - t|)|x_1 - x_2| \\ & \leq |\xi_t(F_s(W)(x_1)) - \xi_t(F_s(W)(x_2))| \\ & \leq (1 + 2C_{1,\square}(\underline{\xi})|s - t|)|x_1 - x_2| \\ & \leq \frac{3}{2}|x_1 - x_2| \end{aligned}$$

if $x_i \in M_s$, $i = 1, 2$; and

$$(30) \quad W \in \mathbf{dmn} F_s^t.$$

Proof. (27) and (28) follow from (24) since $\xi_s(F_s(W)(x)) = x$ and $\rho_s(F_s(W)(x)) = W(x)$ for $x \in M_s$.

Suppose $x_i \in M_s$, $i = 1, 2$. From (27) we infer that

$$\begin{aligned} & \underline{\xi}(t, F_s(W)(x_1)) - \underline{\xi}(t, F_s(W)(x_2)) \\ & = x_1 - x_2 - (\square \underline{\xi}(s, t, F_s(W)(x_1))(s - t) - \square \underline{\xi}(s, t, F_s(W)(x_2))(s - t)) \end{aligned}$$

and, by Proposition 9.5 and Proposition 9.4,

$$\begin{aligned} & |\square \underline{\xi}(s, t, F_s(W)(x_1))(s - t) - \square \underline{\xi}(s, t, F_s(W)(x_2))(s - t)| \\ & \leq C_{1,\square}(\underline{\xi})|F_s(W)(x_2) - F_s(W)(x_1)||s - t| \\ & \leq C_{1,\square}(\underline{\xi})2|x_1 - x_2||s - t| \\ & \leq \frac{1}{2} \end{aligned}$$

It now follows from Proposition 3.4 that $W \in \mathbf{dmn} F_s^t$. \square

Theorem 9.2. *Suppose $s, t \in \mathcal{J}(\epsilon_2)$ and $(s, W) \in \mathcal{G}$. Then*

$$(31) \quad D(\xi_t \circ F_s(W))(x) = \tau_s(x) + \square(D^{0,1}\underline{\xi})(t, s, F_s(W)(x))(t-s) \circ DF_s(W)(x);$$

$$(32) \quad D(\rho_t \circ F_s(W))(x) = DW(x) + \square(D^{0,1}\underline{\rho})(t, s, F_s(W)(x))(t-s) \circ DF_s(W)(x);$$

$$(33) \quad \chi^0(F_s^t(W)) \leq \chi^0(D^{1,0}\underline{\rho})|s-t| + \chi^0(W)$$

and

$$(34) \quad \chi^0(DF_s^t(W)) \leq \chi^0(D^{1,1}\underline{\rho})|s-t| + \chi^0(DW).$$

Proof. Suppose $x \in M_s$. Using (24) we find that

$$\begin{aligned} D(\xi_t \circ F_s(W))(x) - D(\xi_s \circ F_s(W))(x) &= (D^{0,1}\underline{\xi}(t, F_s(W)(x)) - D^{0,1}\underline{\xi}(s, F_s(W)(x))) \circ DF_s(W)(x) \\ &= (\square(D^{0,1}\underline{\xi})(t, s, F_s(W))(t-s)) \circ DF_s(W)(x). \end{aligned}$$

Moreover, $\xi_s \circ F_s(W) = i_{M_s}$ so $D(\xi_s \circ F_s(W))(x) = \tau_s(x)$ so (31) holds. With regard to (32) we have $\rho_s \circ F_s(W) = W$ so a similar analysis yields (32).

By (23) we have

$$F_s^t(W) \circ \xi_t \circ F_s(W) = \rho_t \circ F_s(W).$$

(33) follows from (28). The Chain Rule implies that for any $x \in M_s$ we have

$$DF_s^t(W)((\xi_t \circ F_s(W)(x))) \circ D(\xi_t \circ F_s(W))(x) = D(\rho_t \circ F_s(W))(x);$$

(34) now follows from (31) and (32). \square

9.2. $\mathcal{L}_m, \mathcal{W}_m$ and ϵ_3 . Let $\mathcal{L}_m = \{L \in \mathbb{E}nd(\mathbb{R}^n) : \mathbf{rank} L = m\}$ and let

$$\mathcal{W}_m : \mathcal{L}_m \rightarrow \mathbb{R}^+$$

be such that $\mathcal{W}_m(L) = |\bigwedge_m L|$ for $L \in \mathcal{L}_m$. It is elementary that if $L \in \mathcal{L}_m$ and p is orthogonal projection of \mathbb{R}^n onto the kernel of L then

$$|\mathcal{W}_m(L) - 1| \leq (2^m - 1)\|L - p\|.$$

Suppose $(s, W) \in \mathcal{G}$ and $(s, t, x) \in \underline{Q}(\mathbf{r}/2)$. Let

$$\mathbf{c}_s^t(W)(x) = \mathcal{W}_m(D(\xi_t \circ F_s(W))(x)) - 1.$$

Since $D(\xi_t \circ F_s(W))(x) \in \mathcal{L}_m$ we infer from (9.4) and (32) that

$$\begin{aligned} |\mathbf{c}_s^t(W)(x)| &\leq (2^m - 1)\|D(\xi_t \circ F_s(W))(x) - D(\xi_s \circ F_s(W))(x)\| \\ (35) \quad &\leq (2^m - 1)\|D\xi_t(F_s(W)(x)) - D\xi_s(F_s(W)(x))\| \|D(F_s(W))(x)\| \\ &\leq C_{\mathcal{W}_m}|s-t| \end{aligned}$$

where we have set

$$C_{\mathcal{W}_m} = (2^m - 1)2\chi^{1,1}(\underline{\xi}).$$

Let

$$\epsilon_3 \in (0, \epsilon_2]$$

be such that $C_{\mathcal{W}_m}(2\epsilon_3) < 1$. Suppose $s, t \in \mathcal{J}(\epsilon_3)$. Then $-|\mathbf{c}_s^t(W)| \leq \mathbf{c}_s^t(W) \leq |\mathbf{c}_s^t(W)|$ so

$$(36) \quad 0 \leq 1 - C_{\mathcal{W}_m}|s-t| \leq 1 - |\mathbf{c}_s^t(W)| \leq 1 - \mathbf{c}_s^t(W) \leq 1 + |\mathbf{c}_s^t(W)| \leq 1 + C_{\mathcal{W}_m}|s-t|.$$

Theorem 9.3. *Suppose $s, t \in \mathcal{J}(\epsilon_3)$ and $(s, W) \in \mathcal{G}$. Then*

$$\{(s, t, F_s(W)(x)) : x \in M_s\} \subset \underline{Q}(\underline{\mathbf{r}}/2)$$

and, for any $\epsilon \in \mathbb{R}^+$,

$$|P_t(F_s^t(W), F_s^t(W)) - P_s(W, W)| \leq (\epsilon + C_{F,1}|s-t|)P_s(W, W) + \left(1 + \frac{1}{\epsilon}\right) C_{F,2}|s-t|^2$$

where

$$C_{F,1} = C_{\mathcal{W}_m} \quad \text{and} \quad C_{F,2} = \chi^0(\square\rho)^2 \|M_s\|(\mathbb{R}^n)(1 + C_{\mathcal{W}_m})(2\epsilon_3).$$

Proof. Keeping in mind (36) we have that

$$\begin{aligned} P_t(F_s^t(W), F_s^t(W)) &= \int |F_s^t(W)|^2 d\|M_t\| \\ &= \int |F_s^t(W) \circ \xi_t \circ F_s(W)|^2 (1 + \mathbf{c}_s^t(W)) d\|M\|_s \\ &= \int |\rho_t \circ F_s(W)|^2 (1 + \mathbf{c}_s^t(W)) d\|M\|_s \\ &= \int |W - B_s^t(W)|^2 (1 + \mathbf{c}_s^t(W)) d\|M\|_s \\ &= \int (|W|^2 - 2W \bullet B_s^t(W) + |B_s^t(W)|^2) (1 + \mathbf{c}_s^t(W)) d\|M\|_s \end{aligned}$$

so, using (35), we find that

$$\begin{aligned} &|P_t(F_s^t(W), F_s^t(W)) - P_s(W, W)| \\ &= \left| \int \mathbf{c}_s^t(W) |W|^2 + (-2W \bullet B_s^t(W) + |B_s^t(W)|^2) (1 + \mathbf{c}_s^t(W)) d\|M\|_s \right| \\ &\leq \int (C_{\mathcal{W}_m}|s-t| + \epsilon) |W|^2 \\ &\quad + \left(1 + \frac{1}{\epsilon}\right) |B_s^t(W)|^2 (1 + C_{\mathcal{W}_m}|s-t|) d\|M_s\|. \end{aligned}$$

Since $|B_s^t(W)|^2 \leq (\chi^0(\square\rho))^2 |s-t|^2$ our estimate holds. \square

10. THE FAMILY OF INTEREST.

Suppose $M \in \mathcal{M}_m(\mathbb{R}^n)$, $M \subset \mathbb{S}^{n-1}$ and

$$(37) \quad H(x) = -x \quad \text{for } x \in M$$

where H is the mean curvature normal of M . As is well known and is implied by Proposition 4.2 (37) is equivalent to the statement that

$$(38) \quad \frac{d}{d\epsilon} \mathcal{H}^m(h_t[M]) \Big|_{\epsilon=0} = 0$$

whenever I is an open interval I containing 0, $h : I \times M \rightarrow \mathbb{S}^{n-1}$, h is smooth and $h_0(x) = x$ for $x \in M$.

Let γ be as in 7.1 and let \mathbf{r} be the reach of M .

10.1. **The symmetric quadratic functionals P, Q, R, S .** We define symmetric quadratic functionals

$$P, Q, R, S : \mathcal{E}(\gamma) \times \mathcal{E}(\gamma) \rightarrow \mathbb{R}$$

by letting

$$\begin{aligned} P(X, Y) &= \int X \bullet Y d\|M\|; \\ Q(X, Y) &= \int D_\gamma X \bullet D_\gamma Y d\|M\|; \\ R(X, Y) &= \int A(x)(X(x)) \bullet A(x)(Y(x)) d\|M\|_x; \\ S(X, Y) &= Q(X, Y) - mP(X, Y) - R(X, Y) \end{aligned}$$

for $X, Y \in \mathcal{E}(\gamma)$.

10.2. **$\Phi; \Psi; \mathbf{L}$; and E_λ , $\lambda \in \mathbb{R}$.** For $Z \in \mathcal{E}(\gamma, \mathbf{r})$ we let $F(Z)$ be as in (7.2) and we let

$$\Phi(Z) = \mathcal{H}^m(\mathbf{rng} F(Z)).$$

We define

$$\Psi : \mathcal{C}^{2,\alpha}(\gamma, \mathbf{r}) \rightarrow \mathcal{C}^{0,\alpha}(\gamma)$$

by requiring that

$$P(\Psi(Z), W) = \frac{d}{d\epsilon} \Phi(Z + \epsilon W)|_{\epsilon=0}.$$

(In fact, the right hand side of this formula is an integral over M of linear function of DW and the preceding formula follows by an integration by parts in the integral to obtain an integral over M of a linear function of W .) Let

$$\mathbf{L} \in \mathbb{L}\text{in}(\mathcal{E}(\gamma), \mathcal{E}(\gamma))$$

be the linearization at 0 of Ψ ; that is,

$$\mathbf{L}(Z) = \frac{d}{d\eta} \Psi(\eta Z)|_{\eta=0} \quad \text{for } Z \in \mathcal{E}(\gamma).$$

As is well known,

$$(39) \quad P(\mathbf{L}(Z), W) = P(Z, \mathbf{L}(W)) = S(Z, W) \quad \text{whenever } Z, W \in \mathcal{E}(\gamma).$$

For each $\lambda \in \mathbb{R}$ we let

$$E_\lambda = \{Z \in \mathcal{E}(\gamma) : \mathbf{L}(Z) = \lambda Z\};$$

so $Z \in E_\lambda$ if and only if

$$S(Z, W) = \lambda P(Z, W) \quad \text{for } W \in \mathcal{E}(\gamma).$$

10.3. **The Hölder spaces $\mathcal{C}^{k,\alpha}(\gamma)$.** Let us fix $\alpha \in (0, 1)$. For each $k \in \mathbb{N}$ we let $\mathcal{C}^{k,\alpha}(\gamma)$ be the vector space of maps $Z : M \rightarrow \mathbb{R}^n$ such that $\gamma(x)(Z(x)) = Z(x)$ for each $x \in M$; $D^j Z$ exists and is continuous for each $j \in \mathbb{N}$ such that $j \leq k$; and $D^k Z$ is Hölder continuous with exponent α . $\mathcal{C}^{k,\alpha}(\gamma)$ has a natural Banach space structure. Clearly, P and R have unique continuous extensions, still denoted by P and R , to $\mathcal{C}^{0,\alpha}(\gamma)$ and Q and S have natural continuous extensions, still denoted by Q and S , to $\mathcal{C}^{1,\alpha}(\gamma)$. \mathbf{L} has a unique extension, still denoted by \mathbf{L} to a continuous map from $\mathcal{C}^{2,\alpha}(\gamma)$ to $\mathcal{C}^{0,\alpha}(\gamma)$. Moreover, (39) holds when $Z, W \in \mathcal{C}^{2,\alpha}$.

Φ has a unique continuous extension to $\{Z \in \mathcal{C}^{1,\alpha}(\gamma) : |Z| < \mathbf{r}\}$ and Ψ has a unique extension to a continuous map from $\{Z \in \mathcal{C}^{2,\alpha}(\gamma)\}$ to $\mathcal{C}^{0,\alpha}(\gamma)$. These extensions are real analytic.

Let

$$\mathcal{J} = \ker \mathbf{L} \text{ and let } \mathbf{j} = \dim \mathcal{J}.$$

The members of \mathcal{J} are called **Jacobi fields relative to** \mathbb{S}^{n-1} . *The inner product on \mathcal{J} is the restriction to $\mathcal{J} \times \mathcal{J}$ of P .* So if $s, t \in \mathcal{J}$ then

$$s \bullet t = \int s(x) \bullet t(x) d\|M\|.$$

By standard elliptic theory,

\mathbf{j} is finite;

for each $\epsilon \in \mathbb{R}^+$ there is a constant C such that

$$\sum_{|\lambda| \leq N} \dim E_\lambda \leq (1 + N)^{m/2+\epsilon} \quad \text{whenever } N \in \mathbb{R}^+;$$

in particular, $\dim \text{rng } E_\lambda$ is finite for each $\lambda \in \mathbb{R}$ and $\{\lambda \in \mathbb{R} : E_\lambda \neq 0\}$ is discrete; for each $s \in \mathbb{R}$ and each $Z \in \mathcal{E}(\gamma)$,

$$\Sigma_s(Z) = P(E_0(Z), E_0(Z)) + \sum_{\lambda \in \mathbb{R} \setminus \{0\}} |\lambda|^2 P(E_\lambda(Z), E_\lambda(Z)) < \infty$$

and $\{\Sigma_s : s \in \mathbb{R}\}$ is a family of seminorms which define the strong topology on $\mathcal{E}(\gamma)$.

10.4. Construction of the family of interest. For $k = 0, 2$ let \mathcal{H}_k be the P orthogonal complement of \mathcal{J} in $\mathcal{C}^{k,\alpha}(\gamma)$; since \mathbf{j} is finite \mathcal{H}_k is closed; let Π_k be the P orthogonal projection of $\mathcal{C}^{k,\alpha}(\gamma)$ onto \mathcal{J} and let Π_k^\perp be the complementary projection; thus $Z = \Pi_k(Z) + \Pi_k^\perp(Z)$ for $Z \in \mathcal{C}^{k,\alpha}(\gamma)$. By standard elliptic theory,

$$\mathbf{L}|_{\mathcal{H}_2} \text{ is Banach space isomorphism from } \mathcal{H}_2 \text{ onto } \mathcal{H}_0.$$

Since $\Psi(0) = 0$ and since Ψ is real analytic it follows from the Implicit Function Theorem that there are open neighborhoods \mathcal{S} of 0 in \mathcal{J} and of 0 in \mathcal{V} of 0 in $\text{rng } \Pi_0^\perp \tilde{\Gamma}$ and a real analytic function $\Upsilon : \mathcal{S} \rightarrow \text{rng } \Pi_0^\perp$ such that $\{s + \Upsilon(s) : s \in \mathcal{S}\} \subset \mathcal{V}$ and such that if $s \in \mathcal{S}$, $W \in \mathcal{J}^\perp$ and $s + W \in \mathcal{V}$ then

$$\Pi_0^\perp(\Psi(s + W)) = 0 \Leftrightarrow W = \Upsilon(s).$$

We let $\Omega : \mathcal{S} \rightarrow \mathcal{J}$ be such that

$$\Omega(s) = \Pi_0(\Psi(s + \Upsilon(s))) \quad \text{for } s \in \mathcal{S}$$

and we let

$$\mathcal{A} = \{s \in \mathcal{S} : \Omega(s) = 0\}.$$

Note that Ω is a real analytic function so \mathcal{A} is a real analytic set. Evidently,

$$\{Z \in \mathcal{V} : \Psi(Z) = 0\} = \{s + \Upsilon(s) : s \in \mathcal{S} \text{ and } \Omega(s) = 0\}.$$

Since $\Psi(0) = 0$ we have

$$\Upsilon(0) = 0 \quad \text{and} \quad \Omega(0) = 0.$$

We now assume that

$$(40) \quad \mathbf{Tan} \mathcal{A}, 0 = \mathcal{J};$$

owing to the real analyticity of \mathcal{A} this implies that

Ω vanishes near 0 in \mathcal{S} .

That $\mathbf{Tan}\mathcal{A}, 0 = \mathcal{J}$ is clearly implied by the assumption about Jacobi fields made in (1) of the Introduction of [AA81]. So we have

$$\{Z \in \mathcal{V} : \Psi(Z) = 0\} = \{s + \Upsilon(s) : s \in \mathcal{S}\}.$$

We define

$$\mathbf{z} : \mathcal{S} \times M \rightarrow \mathbb{S}^{n-1}$$

by letting

$$\mathbf{z}(s, x) = F(s + \Upsilon(s))(x) = f(x, (s + \Upsilon(s))(x)) \in \mathbb{S}^{n-1} \quad \text{for } (s, x) \in \mathcal{S} \times M.$$

Letting

$$M_s = \mathbf{rng} F(s + \Upsilon(s))$$

we find that

M_s is Φ stationary.

It follows from Proposition 4.2 that

$$(41) \quad \mathcal{H}^m(M_s) = \mathcal{H}^m(M) \quad \text{for } s \text{ in the connected component of } 0 \text{ in } \mathcal{S}.$$

Remark 10.1. *In fact (41) holds without assuming (40) owing to the fact that the real analyticity of \mathcal{A} implies that \mathcal{A} is locally rectifiably arcwise connected.*

Let

$$\mathcal{J}_s$$

be the Jacobi fields on M_s relative to \mathbb{S}^{n-1} .

For $s \in \mathcal{S}$ we let

$$\gamma_s : M_s \rightarrow \mathbb{O}_m(\mathbb{R}^n)$$

be such that $\gamma_s(y)$ equals orthogonal projection of \mathbb{R}^n onto $\mathbf{Nor}M_s, y \cap \mathbf{Tan}\mathbb{S}^{n-1}, y$. For each $s \in \mathcal{S}$ let

$$\mathbf{J}_s \in \mathbb{L}\mathbf{in}(\mathcal{J}, \mathcal{E}(\gamma_s))$$

be such that

$$\mathbf{J}_s(t)(\mathbf{z}(s, x)) = \gamma_s(\mathbf{z}(s, x)) \left(\left. \frac{d}{d\epsilon} \mathbf{z}(s + \epsilon t, x) \right|_{\epsilon=0} \right) \quad \text{for } t \in \mathcal{J}.$$

Since $M_{s+\epsilon t}$ is stationary for sufficiently small ϵ we find that

$$\mathbf{rng} \mathbf{J}_s \subset \mathcal{J}_s.$$

Theorem 10.1. *$D\Upsilon(0) = 0$ and $\mathbf{J}_0(t) = t$ for $t \in \mathcal{J}$. Moreover,*

$$(42) \quad s \in \mathcal{S} \Rightarrow \mathbf{ker} \mathbf{J}_s = \{0\}.$$

Proof. Suppose $s \in \mathcal{S}$, $t \in \mathcal{J}$ and $W = s + \Upsilon(s) \in \mathcal{E}(\gamma_s)$. Let $X = t + D\Upsilon(s)(t) \in \mathcal{E}(\gamma_s)$. Suppose $x \in M$ and $y = \mathbf{z}(s, x)$. Using Proposition 6.2) we find that

$$\begin{aligned} \mathbf{J}_s(t)(y) &= \gamma_s(y) \left(\frac{d}{d\epsilon} (F(s + \epsilon t))(x) \Big|_{\epsilon=0} \right) \\ &= \gamma_s(y) \left(\frac{d}{d\epsilon} \boldsymbol{\eta}(x + (s + \epsilon t + \Upsilon(s + \epsilon t))(x)) \Big|_{\epsilon=0} \right) \\ &= \gamma_s(y) D\boldsymbol{\eta}(x + W(x))(X(x)) \\ &= \gamma_s(y) \left(1 + |W(x)|^{-3/2} \left((1 + |W(x)|^2)X(x) - (X(x) \bullet W(x))(x + W(x)) \right) \right) \\ &= (1 + |W(x)|)^{-3/2} \left((1 + |W(x)|^2)X(x) - (X(x) \bullet W(x))(W(x)) \right). \end{aligned}$$

In particular, if $s = 0$ then $W = 0$ so that

$$\mathbf{J}_0(t) = X = t + D\Upsilon(0)(t).$$

Since $\mathbf{J}_0(t) \in \mathcal{J}$ and $D\Upsilon(0)(t) \in \mathcal{H}_0$ we find that $D\Upsilon(0)(t) = 0$ and $\mathbf{J}_0(t) = t$.

Now suppose $t \in \ker \mathbf{J}_s$. Then

$$X(x) = \frac{(X(x) \bullet W(x))W(x)}{1 + |W(x)|^2}$$

which implies that

$$|X(x)| \leq |X(x)| \frac{|W(x)|^2}{1 + |W(x)|^2}$$

which in turn implies that $X(x) = 0$ which implies that $t(x) = 0$. □

Corollary 10.1. *Suppose $s, t \in \mathcal{J}$. Then*

$$s \bullet t = P_0(\mathbf{J}_0(s), \mathbf{J}_0(t)).$$

Proof. We have

$$P_0(\mathbf{J}_0(s), \mathbf{J}_0(t)) = \int \mathbf{J}_0(s)(x) \bullet \mathbf{J}_0(t)(x) d\|M_0\|_x = \int s(x) \bullet t(x) d\|M\| = s \bullet t. \quad \square$$

Theorem 10.2. *Suppose $t \in \mathcal{J}(\epsilon_1)$, $u \in \mathcal{J}$ and $x \in M_t$. Then*

$$\mathbf{J}_t(u)(x) = -D\rho(t, x)(u, 0).$$

Remark 10.2. *This is 5.4(7) of [AA81].*

Proof. Let $x = \mathbf{z}(t, p) \in M_t$. Let $I = \{\epsilon \in \mathbb{R} : t + \epsilon u \in \mathcal{J}\}$ and note that I is an open neighborhood of 0 in \mathcal{S} . For $\epsilon \in I$ let

$$g(\epsilon) = \underline{f}(t + \epsilon u, (\mathbf{z}(t + \epsilon u, p), 0)) = \boldsymbol{\eta}(\mathbf{z}(t + \epsilon u, p)).$$

We have $g(0) = \mathbf{z}(t, p) = x$ and, by Proposition 6.2,

$$g'(0) = \mathbf{o}(\mathbf{z}(t, p))(D\mathbf{z}(t, p)(u, 0)) = D\mathbf{z}(t, p)(u, 0)$$

since $|\mathbf{z}(t, p)| = 1$ for $(t, p) \in \mathcal{S} \times M$. Since $g(\epsilon) \in M_{t+\epsilon u}$ we have

$$0 = \underline{\rho}(t + \epsilon u, g(\epsilon)) \quad \text{for } \epsilon \in I;$$

Differentiating at $\epsilon = 0$ we find using Proposition 7.3 that

$$\begin{aligned}
0 &= \frac{d}{d\epsilon} \underline{\rho}(t + \epsilon u, g(\epsilon)) \Big|_{\epsilon=0} \\
&= D\underline{\rho}(t, \mathbf{z}(t, p))(u, g'(0)) \\
&= D\underline{\rho}(t, \mathbf{z}(t, p))(u, D\mathbf{z}(t, p)(u, 0)) \\
&= D\underline{\rho}(t, \mathbf{z}(t, p))(u, 0) + \gamma_t(\mathbf{z}(t, p))(D\mathbf{z}(t, p)(u, 0)).
\end{aligned}$$

□

Lemma 10.1. *There is a constant $C_{\mathbf{J},0}$ such that*

$$|\mathbf{J}_s(u)(x) - \mathbf{J}_t(u)(\xi_t(x))| \leq C |s - t| |u|$$

Suppose $s, t \in \mathcal{J}(\epsilon_1)$, $u \in \mathcal{J}$ and $x \in M_s$. Then

Proof.

$$\begin{aligned}
&\mathbf{J}_s(u)(x) - \mathbf{J}_t(u)(\xi_t(x)) \\
&= (D\underline{\rho}(s, x) - D\underline{\rho}(t, \xi_t(x)))(u, 0) \\
&= (D\underline{\rho}(s, \xi_s(x)) - D\underline{\rho}(t, \xi_t(x)))(u, 0) \\
&= (D\underline{\rho}(s, \xi_s(x)) - D\underline{\rho}(s, \xi_t(x)))(u, 0) + (D\underline{\rho}(s, \xi_t(x)) - D\underline{\rho}(t, \xi_t(x)))(u, 0).
\end{aligned}$$

Now apply Proposition 9.5. □

Corollary 10.2. *There is a constant $C_{\mathbf{J}}$ such that*

$$|P_s(\mathbf{J}_s(u), \mathbf{J}_s(v)) - P_t(\mathbf{J}_t(u), \mathbf{J}_t(v))| \leq C_{\mathbf{J}} |s - t| |u| |v| \quad \text{for } s, t \in \mathcal{J}(\epsilon_1) \text{ and } u, v \in \mathcal{J}.$$

Proof. Suppose $s, t \in \mathcal{J}(\epsilon_1)$ and $u, v \in \mathcal{J}$. For each $x \in M_s$ let

$$\begin{aligned}
\Gamma_1(x) &= \mathbf{J}_s(u)(x); \\
\Gamma_2(x) &= \mathbf{J}_s(v)(x); \\
\Gamma_3(x) &= \mathbf{J}_t(u)(\xi_t(x)); \\
\Gamma_4(x) &= \mathbf{J}_t(v)(\xi_t(x)); \\
\Gamma_5(x) &= \mathbf{J}_t(u)(\xi_t(x)) \bullet \mathbf{J}_t(v)(\xi_t(x)) \mathbf{c}_s^t(0)(x).
\end{aligned}$$

Then

$$\begin{aligned}
&|P_s(\mathbf{J}_s(u), \mathbf{J}_s(v)) - P_t(\mathbf{J}_t(u), \mathbf{J}_t(v))| \\
&= \int \Gamma_1 \bullet \Gamma_2 - \Gamma_3 \bullet \Gamma_4 - \Gamma_5 \, d\|M_s\| \\
&= \int (\Gamma_1 - \Gamma_2) \bullet \Gamma_3 + \Gamma_3 \bullet (\Gamma_2 - \Gamma_4) - \Gamma_5 \, d\|M_s\|.
\end{aligned}$$

Now apply the preceding Lemma and (35). □

The following Corollary follows by letting $t = 0$ above.

Corollary 10.3. *If $u \in \mathcal{J}$ and $s \in \mathcal{J}(\epsilon_1)$ then*

$$P_s(\mathbf{J}_s(u), \mathbf{J}_s(u)) \leq (1 + C_{\mathbf{J}}) |s| |u|.$$

10.5. Symmetric bilinear functions. Suppose V is an inner product space, \mathbf{B} is the vector space of real value symmetric bilinear functions on $V \times V$. We say $T \in \mathbf{B}$ is **nonnegative** if $T(v, v) \geq 0$ for $v \in V$. and \mathbf{B}^+ is the set of those $T \in \mathbf{B}$ such that $T(v, v) \geq 0$ for $v \in V$. Then

$$(43) \quad T(v, w) = \frac{1}{2}(T(v+w, v+w) - T(v, v) - T(w, w))$$

and, if T is nonnegative,

$$(44) \quad |T(v, w)| \leq \sqrt{T(v, v)T(w, w)} \leq \frac{1}{2}(T(v, v) + T(w, w))$$

for $v, w \in V$.

Proposition 10.1. *Suppose $T_i \in \mathbf{B}$, $i = 1, 2$, $U \in \mathbf{B}$, U is nonnegative and*

$$|T_1(v, v) - T_2(v, v)| \leq U(v, v) \quad \text{for } v \in V.$$

Then

$$|T_1(v, w) - T_2(v, w)| \leq \frac{3}{2}(U(u, u) + U(v, v)) \quad \text{for } u, v \in V.$$

Proof. Suppose $v, w \in V$. Then

$$\begin{aligned} & |T_1(v, w) - T_2(v, w)| \\ &= \frac{1}{2} |(T_1(v+w, v+w) - T_1(v, v) - T_1(w, w)) - (T_2(v+w, v+w) - T_2(v, v) - T_2(w, w))| \\ &= \frac{1}{2} |(T_1(v+w, v+w) - T_2(v+w, v+w)) - (T_1(v) - T_2(v)) - (T_1(w) - T_2(w))| \\ &\leq \frac{1}{2} (U(v+w, v+w) + U(v, v) + U(w, w)) \\ &= U(v, v) + U(w, w) + U(v, w) \\ &\leq \frac{3}{2} (U(v, v) + U(w, w)), \end{aligned}$$

□

11. L_s^t AND ϵ_4 .

Suppose $s, t \in \mathcal{J}(\epsilon_2)$. We define a linear map

$$L_s^t : \mathcal{E}(\gamma_s) \rightarrow \mathcal{E}(\gamma_t)$$

as follows. Keeping in mind that Proposition 9.1 implies that ξ_t carries M_s diffeomorphically onto M_t we require that

$$(45) \quad L_s^t(W)(\xi_t(x)) = \gamma_t(\xi_t(x))(W(x)) \quad \text{for } W \in \mathcal{E}(\gamma_s) \text{ and } x \in M_s.$$

Remark 11.1. *This is not the L_s^t of Part Two of the proof of Lemma 5.4 in [AA81]. There we see*

$$L_s^t(W) = \left. \frac{d}{d\epsilon} F_s^t(\epsilon W) \right|_{\epsilon=0}.$$

This has the pleasant property that $L_s^u = L_t^u \circ L_s^t$ which L_s^t above does not have but it is harder to work with.

We let

$$h_L : \underline{N}(\mathbf{r}) \rightarrow \mathbb{E}\text{nd}(\mathbb{R}^n)$$

be such that $h_L(s, z) = \gamma_s(\xi_s(z))$ for $(s, z) \in \underline{N}(\mathbf{r})$; note that h_L is smooth. We infer from Proposition 9.5 that

$$\gamma_s(\xi_s(z)) - \gamma_t(\xi_t(z)) = \square h_L(s, t, z)(s - t) \quad \text{for } (s, t, z) \in \underline{Q}(\mathbf{r}/2).$$

If $W \in \mathcal{E}(\gamma_s)$ and $x \in M_s$ we have

$$(46) \quad L_s^t(W)(\xi_t(x)) = W(x) - \square h_L(s, t, \xi_t(x))(s - t)(W(x))$$

whenever $x \in M_s$ since $\gamma_s(\xi_s(x))(W(x)) = \gamma_s(x)(W(x)) = W(x)$. Let

$$C_L = \sup\{|\square h_L(s, t, z)(u)| : (s, t, z) \in \underline{Q}(\mathbf{r}), u \in \mathcal{J} \text{ and } |u| \leq 1\}.$$

Then $C_L < \infty$ and

$$(47) \quad \|\gamma_s(\xi_s(z)) - \gamma_t(\xi_t(z))\| \leq C_L |s - t| \quad \text{whenever } z \in \underline{Q}(\mathbf{r}/2)$$

and

$$(48) \quad (1 - C_L |s - t|)|W(x)| \leq |L_s^t(W)(\xi_t(x))| \leq (1 + C_L |s - t|)|W(x)|.$$

Let

$$\epsilon_4 \in (0, \min\{1, \epsilon_3\}]$$

be such that $2C_L(2\epsilon_4) < 1$.

The following Theorem is an immediate consequence of (48).

Theorem 11.1. *Suppose $s, t \in \mathcal{J}(\epsilon_4)$. Then L_s^t carries $\mathcal{E}(\gamma_s)$ isomorphically onto $\mathcal{E}(\gamma_t)Su$*

Theorem 11.2. *and there is a constant $C_{L,P}$ such that*

$$|P_t(L_s^t(X), L_s^t(Y)) - P_s(X, Y)| \leq C_{L,P} |s - t| P_s(X, Y).$$

Proof. Suppose $W \in \mathcal{E}(\gamma_s)$ and \mathbf{c}_s^t is as in (9.2). Using 46 we find that

$$\begin{aligned} & \int |L_s^t(W)|^2 - |W|^2 d||M_t|| \\ &= \int |L_s^t(W) \circ \xi_t|^2 \mathcal{W}_m(D(\xi_t|M_s)) - |W|^2 d||M_s|| \\ &= \int |L_s^t(W) \circ \xi_t|^2 (1 + \mathbf{c}_s^t(0)) d||M_s|| \\ &= \int |W(x) - \square h_L(s, t, \xi_t(x))(s - t)|^2 (1 + \mathbf{c}_s^t(0)(x)) - |W(x)|^2 d||M_s|| \end{aligned}$$

Now use (36) and Proposition 10.1 to complete the proof. \square

Proposition 11.1. *Suppose $s \in \mathcal{J}(\epsilon_1)$ and $W \in \mathcal{E}(\gamma_s)$. Then*

$$|R_s(W, W)| \leq m_{\mathbf{r}}^2 P_s(W, W).$$

Proof. Suppose u_1, \dots, u_n is an orthonormal basis for \mathbb{R}^n . For any $x \in M_s$ we have

$$|A_s(x)(W(x))|^2 = \sum_{i=1}^m |A_s(x)(W(x))(u_i)|^2 \leq m \|B_s(x)\|^2 \|W(x)\|^2 \leq m_{\mathbf{b}}^2 |W(x)|^2.$$

\square

Theorem 11.3. *Suppose $s, t \in \mathcal{J}(\epsilon_4)$. There is a constant $C_{L,R}$ such that*

$$|R_t(L_s^t(X), L_s^t(Y)) - R_s(X, Y)| \leq C_{L,R}|s - t|P_s(X, Y) \quad \text{for } X, Y \in \mathcal{E}(\gamma_s).$$

Proof. We let

$$h_A(s, z) = \underline{A}(s, \underline{\xi}(s, z))(\underline{\gamma}(s, \underline{\xi}(s, z))) \quad \text{for } (s, z) \in \underline{Q}(\mathbf{r}).$$

Suppose $W \in \mathcal{E}(\gamma_s)$. Since $(s, t, x) \in \underline{Q}(\mathbf{r}/2)$ we have

$$A_s(x)(W(x)) = A_s(\xi_s(x))(\gamma_s(\xi_s(x)))(W(x)) = h_A(s, x)(W(x))$$

as well as

$$\begin{aligned} A_t(\xi_t(x))(L_s^t(W)(\xi_t(x))) &= A_t(\xi_t(x))(\gamma_t(\xi_t(x)))(W(x)) \\ &= \underline{A}(t, \underline{\xi}(t, x))(\underline{\gamma}(t, \underline{\xi}(t, x)))(W(x)) \\ &= h_A(t, x)(W(x)) \end{aligned}$$

so

$$\begin{aligned} R_t(L_s^t(W), L_s^t(W)) - R_s(W, W) &= \int |A_t(y)(L_s^t(W)(y))|^2 d\|M_t\|y - \int |A_s(x)(W(x))|^2 d\|M_s\|x \\ &= \int |A_t(\xi_t(x))(L_s^t(W)(\xi_t(x)))|^2 (1 + \mathbf{c}_s^t(0)(x)) d\|M_s\|x \\ &= \int h_A(t, x)(W(x))(1 + \mathbf{c}_s^t(0)(x)) - h_A(s, x)(W(x)) d\|M_s\| \\ &= \int (h_A(t, x) - h_A(s, x))(W(x)) + h_A(t, x)(W(x))\mathbf{c}_s^t(0)(x) d\|M_s\|x. \end{aligned}$$

The existence of the constant $C_{L,R}$ now follows from Proposition 9.5, (35) and Proposition 10.1. \square

Theorem 11.4. *Suppose $s, t \in \mathcal{J}(\epsilon_4)$. There is a constant $C_{L,Q}$ such that for $X, Y \in \mathcal{E}(\gamma_s, \mathbf{r})$ we have*

$$|Q_t(L_s^t(X), L_s^t(Y)) - Q_s(X, Y)| \leq C_{L,Q}|s - t|((P_t + Q_t)(X, X) + (P_t + Q_t)(Y, Y))$$

Proof. For $u \in \mathcal{J}(\epsilon_4)$ and $U, V \in \mathcal{E}(\gamma_u, \mathbf{r})$ we set

$$T_u(U, V) = \int DU \bullet DV d\|M_u\|$$

and observe that

$$\begin{aligned} T_u(U, V) &= \int (D_\gamma U(y) + A_u(y)(U(y))) \bullet (D_\gamma V(y) + A_u(y)(V(y))) d\|M_u\|y \\ &= \int D_\gamma U(y) \bullet D_\gamma V(y) + A_u(y)(U(y)) \bullet A_u(y)(V(y)) d\|M_u\|y \\ &= Q_u(U, V) + R_u(U, V). \end{aligned}$$

Suppose $W \in \mathcal{E}(\gamma_s, \mathbf{r})$. We calculate

$$\begin{aligned} Q_t(L_s^t(W), L_s^t(W)) &= \int DL_s^t(W)(y) \bullet DL_s^t(W)(y) d\|M_t\|y \\ &= \int DL_s^t(W)(\xi_t(x)) \bullet DL_s^t(W)(\xi_t(x))(1 + \mathbf{c}_s^t(x)) d\|M_s\|x. \end{aligned}$$

Now by (ii) of Theorem 5.3 and (46) we have for any $W \in \mathcal{E}(\gamma_s, \mathbf{r})$ that

$$\begin{aligned} (49) \quad DL_s^t(W)(\xi_t(x))(u) &= DL_s^t(W)(\xi_t(x)) \circ \tau_t(\xi_t(x))(u) \\ &= DL_s^t(W)(\xi_t(x)) \circ D\xi_t(\xi_t(x))(u) \\ &= D(L_s^t(W) \circ \xi_t)(\xi_t(x))(u) \\ &= DW(x)(u) - D^{0,0,1}\square h_L(s, t, \xi_t(x))(u)(s-t)(W(x)) \\ &\quad - \square h_L(s, t, \xi_t(x))(s-t)(DW(x)(u)) \\ &= \sum_{i=0}^2 \Gamma_i(s, t, x)(DW(x), s-t, W(x))(u) \end{aligned}$$

where

$$\Gamma_i \in \mathcal{E}(\underline{Q}(\mathbf{r}/2), \text{MultiLin}(\text{End}(\mathbb{R}^n) \times \mathcal{J} \times \mathbb{R}^n, \text{End}(\mathbb{R}^n))) \quad i = 0, 1, 2,$$

are such that

$$\begin{aligned} \Gamma_0(s, t, x)(l, v, w)(u) &= l(u); \\ \Gamma_1(s, t, x)(l, v, w)(u) &= -D^{0,0,1}\square h_L(s, t, \xi_t(x))(u)(v)(w); \\ \Gamma_2(s, t, x)(l, v, w)(u) &= -\square h_L(s, t, \xi_t(x))(v)(l(u)) \end{aligned}$$

For $i, j = 0, 1, 2$ we let

$$Z_{i,j}(W, W) \in \mathcal{E}(\underline{Q}(\mathbf{r}/2))$$

be such that

$$Z_{i,j}(W, W)(s, t, x) = \Gamma_i(s, t, x)(DW(x), s-t, W(x)) \bullet \Gamma_j(s, t, x)(DW(x), s-t, W(x))$$

for $i, j = 0, 1, 2$ and $(s, t, x) \in \underline{Q}(\mathbf{r}/2)$. Then (49) implies that

$$\begin{aligned} T_t(L_s^t(W), L_s^t(W)) &= \int DL_s^t(W)(\xi_t(x)) \bullet DL_s^t(W)(\xi_t(x))(1 + \mathbf{c}_s^t(0)) d\|M_s\|x \\ &= \int \sum_{i,j=0}^2 Z_{i,j}(W, W)(s, t, x)(1 + \mathbf{c}_s^t(0)(x)) d\|M_s\|x. \end{aligned}$$

We have

$$|Z_{0,0}(W, W) - |DW|^2| = |\mathbf{c}_s^t||DW|^2 \leq C_{\mathcal{W}_m}|s-t||DW|^2.$$

If $i, j \in \{0, 1, 2\}$ and not both i and j are 0 there is a constant c_1 such that

$$|Z_{i,j}(W, W)(s, t, x)(1 + \mathbf{c}_s^t(x))| \leq c_1|s-t|(|W|^2 + |DW|^2).$$

It follows that

$$|T_t(L_s^t(W), L_s^t(W)) - T_s(W, W)| \leq (C_{\mathcal{W}_m} + 5c_1)C_{\mathcal{W}_m}|s-t|(P_s(W, W) + T_s(W, W)).$$

Since $Q_s = T_s - R_s$ the Theorem now follows from Theorem 11.3 and Proposition 10.1. \square

Theorem 11.5. *Suppose $s, t \in \mathcal{J}(\epsilon_4)$. There is a constant $C_{L,S}$ such that*

$$|S_t(L_s^t(X), L_s^t(Y)) - S_s(X, Y)| \leq C_{L,S}|s-t|((P_s+Q_s)(X, X)+(P_s+Q_s)(Y, Y)) \quad \text{for } X, Y \in \mathcal{E}(\gamma_s).$$

Proof. Suppose $X, Y \in \mathcal{E}(\gamma_s)$. Then

$$\begin{aligned} & |S_t(L_s^t(X), L_s^t(Y)) - S_s(X, Y)| \\ &= |(Q_t - R_t - mP_t)(L_s^t(X), L_s^t(Y)) - (Q_s - R_s - mP_s)(X, Y)| \\ &\leq |Q_t(L_s^t(X), L_s^t(Y)) - Q_s(X, Y)| \\ &\quad + |R_t(L_s^t(X), L_s^t(Y)) - R_s(X, Y)| \\ &\quad + m|P_t(L_s^t(X), L_s^t(Y)) - P_s(X, Y)| \end{aligned}$$

Now apply Theorems 11.2, 11.3 and 11.4. \square

1 b

12. ϵ_5 .

Let

$$\epsilon_5 \in (0, \epsilon_4]$$

be such that

$$(50) \quad 2C_{L,Q\epsilon_5} < \frac{1}{2}.$$

Remark 12.1. *The following Lemma is a replacement for Part Four of Lemma 5.4 of [AA81] which is false; to see this let λ there equal zero. Here is a corrected version for which the inequalities derived in the proof of Part Four of Lemma 5.4 of [AA81] do not suffice to prove the replacement.*

Lemma 12.1. *There is a constant $C_{L,*}$ such that if $s, t \in \mathcal{J}(\epsilon_5)$, $X \in \mathcal{E}(\gamma_s)$, $X^* = X - E_{s,0}(X)$ and*

$$S_t(L_s^t(X), Y) = \lambda P_t(L_s^t(X), Y) \quad \text{for } Y \in \mathcal{E}(\gamma_t)$$

then

$$\kappa_s P_s(X^*, X^*) \leq (|\lambda|(1 + C_{L,*}|s-t|) + C_{L,*}|s-t|)P_s(X, X).$$

Proof. We have

$$\kappa_s P_s(X^*, X^*) \leq |S_s(X, X^*)| \leq \Gamma_1 + \Gamma_2$$

where

$$\Gamma_1 = |S_t(L_s^t(X), L_s^t(X^*))| \quad \text{and} \quad \Gamma_2 = |S_t(L_s^t(X), L_s^t(X^*)) - S_s(X, X^*)|.$$

$$\begin{aligned} \Gamma_1 &= |\lambda| |P_t(L_s^t(X), L_s^t(X^*))| \\ &\leq |\lambda| (|P_s(X, X^*) + C_{L,P}|s-t|(P_s(X, X) + P_s(X^*, X^*))) \\ &\leq |\lambda| (1 + 2C_{L,P}|s-t|)P_s(X, X) \end{aligned}$$

and

$$\begin{aligned} \Gamma_2 &\leq C_{L,S}|s-t|((P_s+Q_s)(X, X) + (P_s+Q_s)(X^*, X^*)) \\ &\leq C_{L,S}|s-t|(2P_s(X, X) + Q_s(X, X) + Q_s(X^*, X^*)). \end{aligned}$$

We estimate

$$\begin{aligned}
Q_s(X, X) &\leq Q_t(L_s^t(X), L_s^t(X)) + C_{L,Q}|s-t|(P_s(X, X) + Q_s(X, X)) \\
&= (S_t + mP_t + R_t)(L_s^t(X), L_s^t(X) + C_{L,Q}|s-t|(P_s(X, X) + Q_s(X, X))) \\
&= \lambda P_t(L_s^t(X), L_s^t(X)) + (mP_t + R_t)(L_s^t(X), L_s^t(X)) \\
&\quad + C_{L,Q}|s-t|(P_s(X, X) + Q_s(X, X)) \\
&\leq ((|\lambda| + m)(1 + C_{L,P}|s-t|) + (1 + C_{L,R}|s-t|))P_s(X, X) \\
&\quad + C_{L,Q}|s-t|(P_s(X, X) + Q_s(X, X))
\end{aligned}$$

so, as $1 - C_{L,Q}|s-t| \geq 1 - C_{L,Q}(2\epsilon_5) \geq 1/2$ we have

$$Q_s(X, X) \leq 2((|\lambda| + m)(1 + C_{L,P}|s-t|) + (1 + C_{L,R}|s-t|))P_s(X, X).$$

Since $S_s(X_0, Y) = 0$ for any $Y \in \mathcal{E}(\gamma_s)$ we estimate

$$\begin{aligned}
Q_s(X^*, X^*) &= Q_s(X, X) - 2Q_s(X, X_0) + Q_s(X_0, X_0) \\
&= Q_s(X, X) + mP_s(X, X_0) + R_s(X, X_0) + mP_s(X_0, X_0) + R_s(X_0, X_0) \\
&\leq Q_s(X, X) + (2m + 2m^2\mathbf{b}^2)P_s(X, X).
\end{aligned}$$

The existence of $C_{L,*}$ follows. \square

12.1. Π_s . For each $s \in \mathcal{J}(\epsilon_5)$ let

$$\Pi_s$$

be Pi_s orthogonal projection of $\mathcal{E}(\gamma_s)$ onto $\mathbf{rng} \mathbf{J}_s$.

Lemma 12.2. *There is a constant C_Π such that*

$$|P_t(\Pi_t(L_s^t(W), \Pi_t(L_s^t(W))) - P_s(\Pi_s(W), \Pi_s(W)))| = C_\Pi|s-t|P_s(W, W)$$

whenever $s \in \mathcal{J}(\epsilon_5)$ and $W \in \mathcal{E}(\gamma_s)$.

Proof. Let $A_s = \mathbf{J}_s^* \circ \mathbf{J}_s$. Then $\mathbf{J}_s \circ A_s^{-1/2}$ is an isometry from \mathcal{J} to $\mathbf{rng} \mathbf{J}_s$. Indeed, if $u, v \in \mathcal{J}$ we have

$$\begin{aligned}
(51) \quad &P_s(\mathbf{J}_s \circ A_s^{-1/2}(u), \bullet \mathbf{J}_s \circ A_s^{-1/2}(v)) \\
&= \mathbf{J}_s^* \circ \mathbf{J}_s \circ A_s^{-1/2}(u) \bullet v \\
&= A_s \circ A_s^{-1/2}(u) \bullet A_s^{-1/2}(v) \\
&= u \bullet v.
\end{aligned}$$

Suppose $W \in \mathcal{E}(\gamma_s)$. If u_1, \dots, u_j is an orthonormal basis for \mathcal{J} and $U_{s,i} = \mathbf{J}_s^* \circ A_s^{-1/2}(u_i) \in \mathcal{E}(\gamma_s)$ then

$$\Pi_s(W) = \sum_{i=1}^j P_s(W, U_{s,i})U_{s,i}$$

which implies that

$$\Pi_t(L_s^t(W), L_s^t(W)) = \sum_{i=1}^j P_t(L_s^t(W), U_{t,i})^2.$$

Thus

$$\begin{aligned}
P_t(L_s^t(W), U_{t,i}) &= \int L_s^t(W) \bullet U_{t,i} d\|M_t\| \\
&= \int L_s^t(W)(\xi_t(x)) \bullet U_{t,i}(\xi_t(x))(1 + \mathbf{c}_s^t(x)) d\|M_s\|x \\
&= \int \gamma_t(\xi_t(x))(W(x)) \bullet U_{t,i}(\xi_t(X))(1 + \mathbf{c}_s^t(x)) d\|M_s\|x
\end{aligned}$$

The Lemma now follows from the estimates occurring in the proof of Theorem 11.3. \square

12.2. ϵ_6 .

Remark 12.2. *The following Lemma is, essentially, Part Six of Lemma 5.4 of [AA81].*

Lemma 12.3. *Suppose $s, t \in \mathcal{J}(\epsilon_5)$, $\Pi_s = E_{s,0}$, $\lambda \in \mathbb{R}$ and $\Pi_t \neq E_{t,\lambda}$. Then*

$$\kappa_s(1 - C_{L,P}|s - t|) \leq |\lambda|(1 + C_{L,*}|s - t|) + C_{L,*}|s - t|.$$

Proof. Choose Y such that $Y \in \mathbf{rng} E_{t,\lambda} \sim \{0\}$ and $\Pi_t(Y) = 0$; if $\lambda = 0$ this is possible since $\mathbf{rng} \Pi_t \subset E_{t,0}$ and if $\lambda \neq 0$ this is possible since $\mathbf{rng} \Pi_t \subset E_{t,0}$ which does not meet $E_{t,\lambda}$. Let $X \in \mathcal{E}(\gamma_s) \sim \{0\}$ be such that $Y = L_s^t(X)$; this is possible since L_s^t carries $\mathcal{E}(\gamma_s)$ isomorphically onto $\mathcal{E}(\gamma_t)$. Rescaling Y if necessary may assume that $P_s(X, X) = 1$.

We have

$$\begin{aligned}
P_s(\Pi_s(X), \Pi_s(X)) &= |P_t(Y, Y) - P_s(\Pi_s(X), \Pi_s(X))| \\
&= |P_t(L_s^t(X), L_s^t(X)) - P_s(\Pi_s(X), \Pi_s(X))| \\
&\leq C_{L,P}|s - t|P_s(X, X) \\
&= C_{L,P}|s - t|
\end{aligned}$$

so that, as $\mathbf{rng} \Pi_s \subset \mathbf{rng} E_{s,0}$,

$$\begin{aligned}
\kappa_s(1 - C_{L,P}|s - t|) &\leq \kappa_s(1 - P_s(\Pi_s(X), \Pi_s(X))) \\
&\leq \kappa_s(P_s(X, X) - P_s(E_{s,0}(X), E_{s,0}(X)));
\end{aligned}$$

now apply Lemma 12.1. \square

Let

$$\epsilon_6 \in (0, \epsilon_5]$$

be such that $\kappa_0 > (C_{L,P} + C_{L,*})\epsilon_6$.

Corollary 12.1. $\mathbf{rng} \mathbf{J}_t = \mathbf{rng} E_{t,0}$ for $t \in \mathcal{J}(\epsilon_6)$.

Proof. Suppose $t \in \mathcal{J}(\epsilon_6)$ and $\Pi_t \neq E_{t,0}$. Applying Lemma 12.3 with $\lambda = 0$ we would have $\kappa_0(1 - C_{L,P}|t|) \leq C_{L,*}|t|$ which would imply $\kappa_0 \leq (C_{L,S}(1 + C)|t| < (C_{L,P} + C_{L,*})\epsilon_6$. \square

Corollary 12.2. *There is a constant C_κ such that*

$$s, t \in \mathcal{J}(\epsilon_6) \Rightarrow |\kappa_s - \kappa_t| \leq C_\kappa|s - t|.$$

Proof. For $u \in \mathcal{J}(\epsilon_6)$ let $\Lambda_u = \{\lambda \in \mathbb{R} \sim \{0\} : E_{t,\lambda} \neq 0\}$.

Suppose $u \in \mathcal{J}(\epsilon_6)$ and $\lambda \in \Lambda_u$. Since $\Pi_0 = E_{0,0}$ and $\Pi_u = E_{u,0} \neq E_{u,\lambda}$ Lemma 12.3 implies

$$\kappa_u(1 - C_{L,P}|u|) \leq |\lambda|(1 + C_{L,*}|u|) + C_{L,*}|u|.$$

Since $\kappa_0 = \inf \Lambda_0$ we have

$$\kappa_u(1 - C_{L,P}|u|) \leq \kappa_0(1 + C_{L,*}|u|) + C_{L,*}|u|.$$

Now suppose $s, t \in \mathcal{J}(\epsilon_6)$, $\kappa_t \geq \kappa_s$ and $\lambda \in \Lambda_s$. Since $\Pi_s = E_{s,\lambda}$, Lemma 12.3 implies that

$$\kappa_t(1 - C_{L,P}|s - t|) \leq |\lambda|(1 + C_{L,*}|s - t|) + C_{L,*}|s - t|.$$

Since $\kappa_s = \inf \Lambda_s$ we find that

$$\kappa_t(1 - C_{L,P}|s - t|) \leq \kappa_s(1 + C_{L,*}|s - t|) + C_{L,*}|s - t|;$$

subtracting $\kappa_s(1 - C_{L,P}|s - t|)$ from both sides we obtain

$$\begin{aligned} & (\kappa_t - \kappa_s)(1 - C_{L,P}|s - t|) \\ & \leq \kappa_s(C_{L,*} + C_{L,P})|s - t| + C_{L,*}|s - t| \\ & \leq \frac{\kappa_0(1 + C_{L,*}|s|) + C_{L,*}|s|}{1 - C_{L,P}|s|} (C_{L,*} + C_{L,P})|s - t| + C_{L,*}|s - t| \end{aligned}$$

so the Corollary holds. \square

13. PROOF OF A STATEMENT SIMILAR TO 5.4(4) OF [AA81].

Theorem 13.1. *There is $\delta \in \mathbb{R}^+$ such that if $(s, Z) \in \mathcal{G}$ and $|s| < \epsilon_6/2$ and $\chi^{[1]}(Z) < \delta$ there is $t \in \mathcal{J}(s, \epsilon_6/2)$ such that*

$$|s - t|^2 \leq 2(1 + C_{\mathbf{J}}|s|)P_s(E_{s,0}(Z), E_{s,0}(Z))$$

and

$$E_{t,0}(F_s^t(Z)) = 0.$$

13.1. $\zeta_{Z,s}$. For $(s, Z) \in \mathcal{G}$ with $|s| < \epsilon_6/2$ we define

$$\zeta_{Z,s} : \mathcal{J}(\epsilon_6) \rightarrow \mathcal{J}$$

by requiring that

$$\zeta_{Z,s}(t) \bullet v = \int F_s^t(Z) \bullet \mathbf{J}_t(v) d\|M_t\| \quad \text{for } t \in \mathcal{J}(\epsilon_6) \text{ and } v \in \mathcal{J}.$$

Since $F_s^s(Z) = Z$ we infer from Corollary 10.3 that

$$\begin{aligned} |\zeta_{Z,s}(s) \bullet v| & \leq \chi^0(Z) \int |\mathbf{J}_s(v)| d\|M_s\| \\ & \leq \chi^0(Z) P_s(\mathbf{J}_s(v), \mathbf{J}_s(v))^{1/2} \\ & \leq \chi^0(Z) (1 + C_{\mathbf{J}}|s|)^{1/2} |v| \end{aligned}$$

so

$$(52) \quad |\zeta_{Z,s}(s)| \leq \frac{\epsilon_6}{4} \quad \text{if} \quad \chi^0(Z) \left(1 + C_{\mathbf{J}} \frac{\epsilon_6}{2}\right)^{1/2} < \frac{\epsilon_6}{4}.$$

Lemma 13.1. *There is a constant $C_{1,\zeta}$ such that*

$$\|D\zeta_{Z,s}(t) - D\zeta_{0,s}(t)\| \leq C_{1,\zeta} \chi^{[1]}(Z).$$

Proof. We have

$$\begin{aligned}\zeta_{Z,s}(t) \bullet v &= \int F_s^t(Z)(\xi_t(F_s(Z)(x))) \bullet \mathbf{J}_t(v)(\xi_t(F_s(Z)(x))) \mathcal{W}_m(D(\xi_t \circ F_s(Z))(x)) d\|M_s\| \\ &= \int \rho_t(F_s(Z)(x)) \bullet \mathbf{J}_t(v)(\xi_t(F_s(Z)(x))) \mathcal{W}_m(D(\xi_t \circ F_s(Z)) d\|M_s\| x \\ &= \int A_s(Z)(t, x) B_s(Z)(t, x)(v) C_s(Z)(t, x) d\|M_s\| x\end{aligned}$$

where

$$\begin{aligned}A_s(Z)(t, x) &= \rho_t(F_s(Z)(x)) = \underline{\rho}(t, F_s(Z)(x)); \\ B_s(Z)(t, x)(v) &= \mathbf{J}_t(v)(\xi_t(F_s(Z)(x))) = -D^{1,0} \underline{\rho}(t, \underline{\xi}(t, F_s(Z)(x)))(v); \\ C_s(Z)(t, x) &= \mathcal{W}_m(D(\xi_t \circ F_s(Z))(x)) = \mathcal{W}_m(D^{0,1} \underline{\xi}(t, F_s(Z)(x)) \circ DF_s(Z)(x));\end{aligned}$$

Consequently,

$$\begin{aligned}D^{1,0}A(Z)(t, x)(u) &= D^{1,0} \underline{\rho}(t, F_s(Z)(x))(u); \\ D^{1,0}B(Z)(t, x)(u)(v) &= -D(D^{1,0} \underline{\rho})(t, \underline{\xi}(t, F_s(Z)(x)))(u, D^{1,0} \underline{\xi}(t, F_s(Z)(x))(u))(v); \\ D^{1,0}C(W)(t, x)(u) &= D\mathcal{W}_m(D^{0,1} \underline{\xi}(t, F_s(Z)(x)) \circ DF_s(Z)(x)) \\ &\quad (D^{1,0}(D^{0,1} \underline{\xi})(t, F_s(Z)(x)(u) \circ DF_s(Z)(x))).\end{aligned}$$

In particular,

$$\begin{aligned}D^{1,0}A(0)(t, x)(u) &= D^{1,0} \underline{\rho}(t, x)(u); \\ D^{1,0}B(0)(t, x)(u)(v) &= -D(D^{1,0} \underline{\rho})(t, \underline{\xi}(t, x))(u, D^{1,0} \underline{\xi}(t, x)(u))(v); \\ D^{1,0}C(W)(t, x)(u) &= D\mathcal{W}_m(D^{0,1} \underline{\xi}(t, x) \circ \tau_s(x)) \\ &\quad (D^{1,0}(D^{0,1} \underline{\xi})(t, x)(u) \circ \tau_x(x)).\end{aligned}$$

Lemma 13.2. *There are constants c_A, c_B, c_C such that*

$$\begin{aligned}\|D^{1,0}A(Z)(t, x) - D^{1,0}A(0)(t, x)\| &\leq c_A \chi^{[1]}(Z); \\ \|D^{1,0}B(Z)(t, x) - D^{1,0}B(0)(t, x)\| &\leq c_B \chi^{[1]}(Z); \\ \|D^{1,0}C(Z)(t, x) - D^{1,0}C(0)(t, x)\| &\leq c_C \chi^{[1]}(Z).\end{aligned}$$

Proof. Observe that $F_s(Z)(x) = f_s(x, Z(x))$ and $F_s(0)(x) = f_s(x, 0)$. This implies that $DF_s(Z)(x) = Df_s(x, Z(x)) \circ (\tau_s(x), DZ(x))$ and $DF_s(0)(x) = Df_s(x, 0) \circ (\tau_s(x), 0)$. □

□

Lemma 13.3. *There is a constant $C_{2,\zeta}$ such that*

$$\|D\zeta_{0,s}(t) - D\zeta_{0,s}(s)\| \leq C_{2,\zeta} |s - t| \quad \text{for } s \in \mathcal{J}(\epsilon_6) \text{ and } t \in \mathcal{J}(\epsilon_6).$$

Proof. We have

$$\zeta_{0,s}(t) \bullet v = \int \rho_t(x) \bullet D^{1,0} \underline{\rho}(t, x)(v) \mathcal{W}_m(D\xi_t(x) \circ \tau_s(x)) d\|M_s\|.$$

It follows that

$$\begin{aligned} D\zeta_{0,s}(t)(u) \bullet v &= \int D^{1,0}\underline{\rho}(t,x)(u) \bullet D^{1,0}\underline{\rho}(t,x)(v) \mathcal{W}_m(D\xi_t(x) \circ \tau_s(x)) \\ &\quad + \rho_t(x) \bullet D^{2,0}\underline{\rho}(t,x)(u \odot v) \mathcal{W}_m(D\xi_t(x) \circ \tau_s(x)) \\ &\quad + \rho_t(x) \bullet D^{1,0}\underline{\rho}(t,x)(v) D\mathcal{W}_m(D\xi_t(x) \circ \tau_s(x))(D^{1,0}\underline{\xi}(t,x)(u) \circ \tau(x)) d\|M_s\|x \end{aligned}$$

Since $\rho_s(x) = 0$ and $D\xi_s(x) \circ \tau_s(x) = \tau_s(x)$ for $x \in M_s$ we have

$$(53) \quad D\zeta_{0,s}(s)(u) \bullet v = \int D^{1,0}\underline{\rho}(s,x)(u) \bullet D^{1,0}\underline{\rho}(s,x)(v) d\|M_s\|x$$

If $(t,x) \in \underline{N}(\mathbf{r})$ then $\underline{\rho}(t,x) = -\square\rho(s,t,x)(s-t)$ and

$$D^{1,0}\underline{\rho}(t,x) = D^{1,0}\underline{\rho}(s,x) - \square(D^{1,0}\underline{\rho})(s,t,x)(s-t).$$

□

Lemma 13.4. *We have*

$$\|D\zeta_{0,s}(s) - i_{\mathcal{J}}\| \leq C_{\mathbf{J}}|s| \quad \text{for } s \in \mathcal{J}(\epsilon_1).$$

Proof. Suppose $s \in \mathcal{J}(\epsilon_6)$ and $u, v \in \mathcal{J}$. Then

$$\begin{aligned} D\zeta_{0,s}(s)(u) \bullet v - u \bullet v &= \int D^{1,0}\underline{\rho}(s,x)(u) \bullet D^{1,0}\underline{\rho}(s,x)(v) d\|M_s\|x - \int \mathbf{J}_0(u) \bullet \mathbf{J}_0(v) d\|M_0\| \\ &= \int \mathbf{J}_s(u) \bullet \mathbf{J}_s(v) d\|M_s\| - \int \mathbf{J}_0(u) \bullet \mathbf{J}_0(v) d\|M_0\| \\ &= P_s(\mathbf{J}_s(u), \mathbf{J}_s(v)) - P_0(\mathbf{J}_0(u), \mathbf{J}_0(v)); \end{aligned}$$

Now apply Corollary 10.2. □

13.2. Finding a zero of $\zeta_{Z,s}$. Let $B = \{u \in \mathcal{J} : |u - s| \leq \zeta_6/2\} \subset \mathcal{J}(\epsilon_6)$; let $g : B \rightarrow \mathcal{J}$ be such that $g(u) = \zeta_{Z,s}(u)$ for $u \in B$; and let

It follows from Lemmas (13.1), (13.3) 13.4 that

$$\|D\zeta_{Z,s}(t) - i_{\mathcal{J}}\| \leq \|D\zeta_{Z,s}(t) - D\zeta_{0,s}(s)\| + \|D\zeta_{0,s}(s) - i_{\mathcal{J}}\| \leq C_{1,\zeta}\chi^{[1]}(Z) + C_{2,\zeta}|s-t| + C_{\mathbf{J}}|s|.$$

Suppose (52); then

$$|g(s)| < \frac{\epsilon_6}{4}.$$

Suppose

$$(54) \quad C_{1,\zeta}\chi^{[1]}(Z) + C_{2,\zeta}|s-t| + C_{\mathbf{J}}|s| \leq \frac{1}{2}.$$

$p : B \rightarrow \mathcal{J}$ be such that $p(u) = (g(u) - g(s)) - (u - s)$ for $u \in B$. Then $p(s) = 0$ and $\mathbf{Lip} p \leq 1/2$. Let $C : B \rightarrow \mathcal{J}$ be such that $C(u) = s - g(s) - p(u)$ for $u \in B$. Then $\mathbf{Lip} C = \mathbf{Lip} p \leq 1/2$.

Suppose $u \in B$. Then

$|C(u) - s| = |g(s) + p(u)| = |g(s) + p(u) - p(s)| \leq |g(s)| + \mathbf{Lip} p|u - s| \leq \epsilon_6 - |s|$
so $C[B] \subset B$. Since B is closed and $\mathbf{Lip} C \leq 1/2 < 1$, C has a unique fixed point $t \in B$. Since $t = C(t) = s - g(s) - p(t) = s - g(s) - (p(t) - p(s))$ we have

$$|s - t| = |g(s) + (p(t) - p(s))| \leq |g(s)| + \frac{|s - t|}{2}$$

so

$$|s - t| \leq 2|g(s)|.$$

Suppose $u \in \mathcal{J}$ and $|u| = 1$. Then

$$g(s) \bullet u = \int Z \bullet \mathbf{J}_s(u) d\|M_s\| = P_s(Z, \mathbf{J}_s(u)) = P_s(E_{s,0}(Z), \mathbf{J}_s(u))$$

so by Corollary 10.2 we have

$$\begin{aligned} |g(s) \bullet u|^2 &\leq P_s(E_{s,0}(Z), E_{s,0}(Z))P_s(\mathbf{J}_s(u), \mathbf{J}_s(u)) \\ &\leq P_s(E_{s,0}(Z), E_{s,0}(Z))(P_0(\mathbf{J}_0(u), \mathbf{J}_0(u)) + C_{\mathbf{J}}|s|) \\ &\leq P_s(E_{s,0}(Z), E_{s,0}(Z))(1 + C_{\mathbf{J}}|s|) \end{aligned}$$

so that

$$|s - t|^2 \leq 2|g(s)| \leq P_s(E_{s,0}(Z), E_{s,0}(Z))(1 + C_{\mathbf{J}}|s|).$$

REFERENCES

- [AW1] W. K. Allard. On the first variation of a varifold. *Ann. Math.*, 95:417–491, 1972.
- [AA81] William K. Allard and Frederick J. Almgren, Jr. On the radial behavior of minimal surfaces and the uniqueness of their tangent cones. *Ann. of Math. (2)*, 113(2):215–265, 1981.
- [FE1] H. Federer: *Geometric Measure Theory*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Band 153, Springer Verlag, (1969)
- [FE2] H. Federer: *Curvature measures*, Trans. Amer. Math. Soc., 95:3 (1959), pp. 418-491.
- [MN] Francesco Maggi and Michael Novack, Isoperimetric Residues and a Mesoscale Flatness Criterion for Hypersurfaces with Bounded Mean Curvature, preprint.
- [MS] John W. Milnor and James D. Stasheff, *Characteristic Classes*, Annals of Mathematics Studies, Number 76, Princeton University Press.