Sinkhorn algorithms and linear programming solvers for optimal partial transport problems

Yikun Bai

June 2024

Abstract

In this note, we generalize the classical optimal partial transport (OPT) problem by modifying the mass destruction/creation term to function-based terms, introducing what we term "generalized optimal partial transport" problems. We then discuss the dual formulation of these problems and the associated Sinkhorn solver. Finally, we explore how these new OPT problems relate to classical optimal transport (OT) problems and introduce a linear programming solver tailored for these generalized scenarios.

1 Notation and assumptions

- $\mathbb{R} = (-\infty, \infty), \mathbb{R}_+ = [0, \infty), \mathbb{R}_{++} = (0, \infty).$
- Ω : A non-empty convex open subset of \mathbb{R}^d for some $d \in \mathbb{N}$.
- $c: \Omega^2 \to \mathbb{R}_+$: A lower semi-continuous function, referred to as the "cost function" in this note. Given a topological set $E, f: E \to \mathbb{R}$ is said to be lower-semi continuous at $x \in E$ if the following holds:

Pick $x_n \to x$ (where the convergence is determined by the topology of E), then

$$\liminf_{n} f(x_n) \ge f(x).$$

- $\mathcal{P}(\Omega)$: The set of all probability measures defined on Ω .
- $\mathcal{M}_+(\Omega)$: The set of all positive Radon measures defined on Ω .
- $\mathcal{M}(\Omega)$: The set of all Radon measures defined on Ω .
- $C_0(\Omega)$: set of all C_0 functions. In particular, it is defined as

 $C_0(\Omega) := \{ f : \Omega \to \mathbb{R}, f \text{ is continuous}, \exists \text{compact set} K \subset \Omega, s.t.f(x) = 0, \forall x \notin K \}.$

- γ : A element in $\mathcal{P}(\Omega^2)$ or $\mathcal{M}_+(\Omega^2)$.
- μ, ν : elements in $\mathcal{P}(\Omega)$ or $\mathcal{M}_+(\Omega)$.
- $\pi_1, \pi_2: \Omega^2 \to \Omega$: Canonical projections defined on Ω^2 . Specifically, $\pi_1((x, y)) = x$ and $\pi_2((x, y)) = y$.
- $f_{\#\mu}$: The push-forward of measure μ by a function $f: \Omega \to \Omega$. Specifically, $f_{\#\mu}(A) = \mu(f^{-1}(A))$ for any Borel set A.
- γ_1, γ_2 : $\gamma_1 = (\pi_1)_{\#} \mu$ is the first marginal of γ . γ_2 is defined similarly.
- $\mu = \nu$: Indicates that two measures are equal. For each Borel set A, $\mu(A) = \nu(A)$, or equivalently, for each test function $\phi \in C_0$, we have

$$\int \phi d\mu = \int \phi d\nu.$$

• $\mu \leq \nu$: Indicates that μ is dominated by ν . Specifically, for each Borel set A, $\mu(A) \leq \nu(A)$. Equivaletly, for each test function $\phi \in C_0(\Omega)$, $\phi \geq 0$, we have

$$\int \phi d\mu \leq \int \phi d\nu.$$

- $\Gamma(\mu,\nu) := \{\gamma \in \mathcal{P}(\Omega^2) : \gamma_1 = \mu, \gamma_2 = \nu\}.$
- $\Gamma_{\leq}(\mu,\nu) := \{\gamma \in \mathcal{M}_{+}(\Omega^{2}) : \gamma_{1} \leq \mu, \gamma_{2} \leq \nu\}.$
- $\Gamma^{\eta}_{<}(\mu,\nu) := \{\gamma \in \mathcal{M}_{+}(\Omega^{2}) : \gamma_{1} \leq \mu, \gamma_{2} \leq \nu, |\gamma| = \eta\}, \text{ where } \eta \in [0, \min\{|\mu|, |\nu|\}].$
- $\operatorname{Supp}(\mu)$, $\operatorname{Supp}(\nu)$: The support sets of μ and ν .
- \mathcal{L} : The Lebesgue measure defined on Ω .
- Continuous setting: In this case, we suppose $\mu = f d\mathcal{L}$ and $\nu = g d\mathcal{L}$, where $f, g \in L_1(\Omega)$.
- Discrete setting: In this case, we suppose $\mu = \sum_{i=1}^{n} p_i \delta_{x_i}$ and $\nu = \sum_{j=1}^{m} q_j \delta_{y_j}$, where $n, m \in \mathbb{N}$ and $p_i, q_j > 0$ for all i, j. We denote $X = \{x_i : i \in [1:n]\}$ and $Y = \{y_j : j \in [1:m]\}$.
- dx, dy: dx is a reference measure in Ω such that μ ≪ dx, and similarly, dy is a reference measure in Ω such that ν ≪ dy. Both are set to be probability measures.
 Typically, we assume dµ/dx, dν/dy are L₁ functions.
- $ight cally, we assume <math>a\mu/ax, a\nu/ay$ are D_1 functions.
- $dxdy = dx \otimes dy$: The product measure of dx and dy, defined as the reference measure on Ω^2 .
- $\frac{d\mu}{dx}, \frac{d\nu}{dy}$: Density functions of μ and ν with respect to dx and dy, respectively.
- Inner product and integration: In this note, we use the following identity:

$$\int f d\mu = \mu(f) = \langle f, \mu \rangle,$$

where f is a measurable function defined on Ω .

• $\|\cdot\|_{TV}$: Total variation norm defined by:

$$\|\mu\|_{TV} = \int d|\mu|,$$

where $|\mu| = \mu_+ + \mu_-$, and μ_+, μ_- are the unique measure decomposition of the signed measure $\mu \in \mathcal{M}(\Omega)$.

- $\mu = \frac{d\mu}{d\nu}\nu + \mu^{\perp}$: The Lebesgue decomposition of μ . By classical measure theory, this decomposition is unique.
- $\mu \ll \nu$: Indicates that μ is absolutely continuous with respect to ν .
- $\|\mu\|, \|\nu\|$: Total masses of μ and ν , respectively. $\|p\|, \|q\|$ are defined similarly.
- $KL(\mu \parallel \nu)$: The Kullback-Leibler (KL) divergence between μ and ν . See (3).
- $\|\mu \nu\|_{TV} = \|\mu \nu\|_{TV}$, $P\|\mu \nu\|_{TV} = \|\mu \nu\|_{PTV}$: Total variation and partial total variation between μ and ν . See (5) and (7).
- f, D_f : f is an entropy function, and D_f is the corresponding f-divergence.
- $f'_{\infty} = \lim_{x \to \infty} \frac{f(x)}{x} \in \mathbb{R} \cup \{\infty\}$: The growth rate of f at infinity.
- ι_A , where $A \in \Omega$: indicator function such that $\iota_A(x) = 0$ if $x \in A$ and $\iota_A(x) = \infty$ elsewhere.

• (E, E^*) : E is a convex set, and E^* is its dual space. The dual pairing is defined by:

 $(x, x^*) \mapsto x^*(x) = \langle x, x^* \rangle \in \mathbb{R}.$

- $\lambda > 0$: A constant used in the OPT problem (18).
- $\lambda_1, \lambda_2 : \Omega \to \mathbb{R}_+$: Positive bounded functions.
- $\eta \in [0, \min\{|\mu|, |\nu|\}]$: A constant used in the MOPT problem (21).
- $\epsilon > 0$: A constant used in the entropic OPT problem (24).
- $K = e^{-c/\epsilon}$
- $\mu|_A$, where $A \subset \Omega$: The restriction of the measure μ on set A. Specifically,

$$\mu|_A(B) = \mu(B \cap A)$$

for all Borel sets $B \subset \Omega$.

• $\phi|_A$, where $A \subset \Omega$: The restriction of the function ϕ on A. Specifically,

$$\phi|_A(x) = \begin{cases} \phi(x) & \text{if } x \in A \\ 0 & \text{elsewhere} \end{cases}$$

Note, the above definition can be equivalently redefined as

 $\Phi \mid_A : A \to \mathbb{R}$

with $\Phi \mid_A (x) = \Phi(x), \forall x \in A$. In this note, for conveneince, we to not distinguish these two formulations.

- $x \odot y, \frac{x}{y}$ where x, y are vectors or matrices with same shape: Point-wise multiplication/division.
- 1_n : a $n \times 1$ vector where entry is 1. Similarly, we can define $1_m, 1_{n \times m}, 0_n, 0_m \dots$

2 Outlines

In section 3, we introduce the foundational concepts of f-divergence, unbalanced optimal transport, and various optimal partial transport problems. Section 4 presents the formulations of new optimal partial transport problems, their entropic versions, and the corresponding dual forms. Section 4.4.2 discusses the Sinkhorn algorithms tailored for these new OPT problems. In section 4.4.3, we explore the relationship between the generalized OPT problems and classical OT problems, including discussions on the linear programming solver adapted for generalized OPT problems. Section 5 is dedicated to examining both the Sinkhorn and linear programming solvers for mass-constrained OPT problems.

3 Background

3.1 Entropy Function and *f*-Divergence

An entropy function $f : \mathbb{R} \to \mathbb{R} \cup \{\infty\}$ is lower semi-continuous, convex, and its domain $D(f) := \{x : f(x) < \infty\}$ is contained within $[0, \infty)$ and intersects $(0, \infty)$. The growth rate of f at infinity is defined as:

$$f'_{\infty} = \lim_{x \to \infty} \frac{f(x)}{x} \in \mathbb{R} \cup \{\infty\}.$$

Given measures $\mu, \nu \in \mathcal{M}(\Omega)$ with the Lebesgue decomposition $\mu = \frac{d\mu}{d\nu}\nu + \mu^{\perp}$, the *f*-divergence is defined by:

$$D_f(\mu \parallel \nu) := \int_{\Omega} f\left(\frac{d\mu}{d\nu}\right) d\nu + f'_{\infty} |\mu^{\perp}|, \qquad (1)$$

assuming the convention $0 \cdot \infty = 0$.

Example 1. We consider the following examples of entropy functions:

• Kullback-Leibler (KL) Divergence:

$$f(s) = f_{KL}(s) := \begin{cases} s \ln(s) - s + 1 & \text{if } s > 0\\ 1 & \text{if } s = 0 \\ \infty & \text{if } s < 0 \end{cases}$$
(2)

Here, $(f_{KL})'_{\infty} = \infty$, and the KL divergence is:

$$KL(\mu \parallel \nu) := \begin{cases} \int \ln\left(\frac{d\mu}{d\nu}\right) \frac{d\mu}{d\nu} d\nu - \|\mu\| + \|\nu\| & \text{if } \mu \ll \nu\\ \infty & \text{otherwise} \end{cases}, \tag{3}$$

with $\ln(0) \cdot 0 = 0$ assumed.

• Total Variation:

$$f(s) = f_{TV}(s) := \begin{cases} |s-1| & \text{if } s \ge 0\\ \infty & \text{if } s < 0 \end{cases},$$
(4)

where $(f_{TV})'_{\infty} = 1$, and the total variation divergence is:

$$\|\mu - \nu\|_{TV} := \int_{\Omega} d|\mu - \nu|,$$
 (5)

• Partial Total Variation:

$$f(s) = f_{PTV}(s) := \begin{cases} 1 - s & \text{if } s \in [0, 1] \\ \infty & \text{if } s < 0 \text{ or } s > 1 \end{cases},$$
(6)

with $(f_{PTV})'_{\infty} = \infty$. The corresponding divergence is termed "partial total variation divergence":

$$\|\mu - \nu\|_{PTV} := \begin{cases} \|\nu - \mu\|_{TV} = \|\nu\| - \|\mu\| & \text{if } \mu \le \nu \\ \infty & \text{otherwise} \end{cases},$$
(7)

• Equality Constraint Divergence:

$$f(s) = \iota_{\{1\}}(s) := \begin{cases} 0 & \text{if } s = 1\\ \infty & \text{otherwise} \end{cases},$$
(8)

where $(\iota_{\{1\}})'_{\infty} = \infty$. The associated divergence is termed equality-constraint divergence:

$$\iota_{=}(\mu \parallel \nu) := \begin{cases} 0 & if \ \mu = \nu \\ \infty & otherwise \end{cases},$$
(9)

• Non-constraint Divergence Function:

$$f(s) = \mathbf{0}(s) := 0,\tag{10}$$

with $(\mathbf{0})'_{\infty} = 0$, the associate divergence, termed "D₀", is:

$$D_{\mathbf{0}}(\mu \parallel \nu) = 0, \tag{11}$$

Note, **0** does not meet classical entropy function conditions as it does not ensure $D(\mathbf{0}) \subset [0, \infty)$.

• Partial-Transportation Constraint Divergence Function:

$$f(s) = \iota_{[0,1]}(s) = \begin{cases} 0 & \text{if } s \in [0,1] \\ \infty & \text{if } s < 0 \text{ or } s > 1 \end{cases},$$
(12)

where $(\iota_{[0,1]})_{\infty}=\infty.$ The corresponding divergence, termed " ι_{\leq} ", is:

$$\iota_{\leq}(\mu \parallel \nu) = \begin{cases} 0 & \text{if } \mu \leq \nu \\ \infty & \text{otherwise} \end{cases}.$$
(13)

Remark 1. It is straightforward to verify, $f_{PTV} = f_{TV} + \iota_{[0,1]}$, and thus,

$$PTV(\mu \parallel \nu) = \|\mu - \nu\|_{TV} + \iota_{\leq}(\mu \parallel \nu).$$

3.2 Optimal Transport and Optimal Partial Transport

Optimal transport (OT) problems, as studied in [13, 12], seek the most cost-efficient way to transfer mass between two probability measures. Specifically, given $\mu, \nu \in \mathcal{P}(\Omega)$, the OT problem is defined as:

$$OT(\mu,\nu) := \min_{\gamma \in \Gamma(\mu,\nu)} \int_{\Omega^2} c(x,y) d\gamma(x,y) = \min_{\gamma \in \Gamma(\mu,\nu)} \langle c,\gamma \rangle$$
(14)

Classical OT theory ensures the existence of a minimizer, as c is lower semi-continuous, allowing the substitution of inf with min.

In a discrete setting, where $\mu = \sum_{i=1}^{n} p_i \delta_{x_i}$ and $\nu = \sum_{j=1}^{m} q_j \delta_{y_j}$, which are discrete probability measures, the problem becomes:

$$OT(\mu,\nu) := \min_{\gamma \in \Gamma(p,q)} \langle c, \gamma \rangle \tag{15}$$

Here, $c \in \mathbb{R}^{n \times m}$ with $c_{i,j} = c(x_i, y_j)$ for all i, j, and $\Gamma(p, q) = \{\gamma \in \mathbb{R}^{n \times m}_+ : \gamma \mathbf{1}_m = p, \gamma^\top \mathbf{1}_n = q\}.$

The problem (14), (15) is referred to as the **balanced optimal transport problem** because it requires $\|\mu\| = \|\nu\|$. In an unbalanced setting, where $\mu, \nu \in \mathcal{M}_+(\Omega)$ may not be equal, the problem (14) can be generalized as follows:

$$UOT(\mu,\nu; D_{f_1}, D_{f_2}) := \inf_{\gamma \in \mathcal{M}_+(\Omega^2)} \langle c, \gamma \rangle + D_{f_1}(\gamma_1 \parallel \mu) + D_{f_2}(\gamma_2 \parallel \nu),$$
(16)

where D_{f_1}, D_{f_2} are f-divergences defined in (1).

=

Remark 2. When $D_{f_1}, D_{f_2} = \iota_=$, the above formulation reverts to the balanced OT (14) when $\|\mu\| = \|\nu\|$, and to ∞ otherwise. Thus, (14) can be redefined as:

$$OT(\mu,\nu) := \inf_{\gamma \in \mathcal{M}_+(\Omega^2)} \langle c, \gamma \rangle + \iota_=(\gamma_1 \parallel \mu) + \iota_=(\gamma_2 \parallel \nu) = \begin{cases} \min_{\gamma \in \Gamma(\mu,\nu)} \langle c, \gamma \rangle & \text{if } \|\mu\| = \|\nu\| \\ \infty & \text{otherwise} \end{cases}.$$
(17)

A specific class of unbalanced optimal transport, termed **optimal partial transport** and discussed in [5, 10, 8, 9], is formulated by setting $D_{f_1}, D_{f_2} = \lambda T V$:

$$OPT(\mu,\nu;\lambda) := \inf_{\gamma \in \mathcal{M}_{+}(\Omega^{2})} \langle c,\gamma \rangle + \lambda \left(TV(\gamma_{1} \parallel \mu) + TV(\gamma_{2} \parallel \nu) \right)$$
(18)

$$= \inf_{\gamma \in \gamma_{\leq}(\mu,\nu)} \langle c, \gamma \rangle + \lambda(\|\mu - \gamma_{1}\| + \|\nu - \gamma_{2}\|)$$
(19)

$$= \inf_{\gamma \in \mathcal{M}_{+}(\Omega^{2})} \langle c, \gamma \rangle + \lambda (PTV(\gamma_{1} \parallel \mu) + PTV(\gamma_{2} \parallel \nu))$$
(20)

where

$$\Gamma_{\leq}(\mu,\nu) = \{\gamma \in \mathcal{M}_{+}(\Omega^{2}) : \gamma_{1} \leq \mu, \gamma_{2} \leq \nu\};$$

(19) has been validated by [10], and (20) directly follows from the definition of PTV (see (7)).

Additionally, an equivalent formulation of (19), termed "mass-constraint optimal partial transport", is proposed in this paper:

$$MOPT_{\eta}(\mu,\nu) := \inf_{\gamma \in \Gamma_{\leq}^{n}(\mu,\nu)} \langle c, \gamma \rangle$$
(21)

where $\eta \in [0, \min\{|\mu|, |\nu|\}],\$

$$\inf_{\gamma \in \Gamma_{\leq}^{\eta}} (\mu, \nu) = \{ \gamma \in \mathcal{M}_{+}(\Omega^{2}) : \gamma_{1} \leq \mu, \gamma_{2} \leq \nu, \|\gamma\| = \eta \}$$

Optimal partial transport problems (18), (21) admit a minimizer.

4 A Generalized Version of the OPT Problem

In this section, we assume $\lambda_1, \lambda_2 : \Omega \to \mathbb{R}_+$ are measurable, bounded functions. We introduce the generalized OPT problem defined as:

$$GOPT(\mu,\nu;\lambda_1,\lambda_2) := \inf_{\gamma \in \mathcal{M}_+(\Omega^2)} \langle c,\gamma \rangle + \begin{cases} TV(\lambda_1\gamma_1 \parallel \lambda_1\mu) \\ \text{or} \\ \overline{PTV}(\lambda_1\gamma_1 \parallel \lambda_1\mu) \end{cases} + \begin{cases} TV(\lambda_2\gamma_2 \parallel \lambda_2\nu) \\ \text{or} \\ \overline{PTV}(\lambda_2\gamma_2 \parallel \lambda_2\nu) \end{cases}$$
(22)

where $\overline{PTV}(\lambda_1\gamma_1 \parallel \lambda_1\mu) = PTV(\lambda_1\gamma_1 \parallel \lambda_1\mu) + \iota_{\leq}(\gamma_1 \parallel \mu)$, and $\overline{PTV}(\lambda_2\gamma_2 \parallel \lambda_2\nu)$ is defined similarly. We refer to $TV(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ (or similarly for \overline{PTV}) as the "first penalty term" and $TV(\lambda_2\gamma_2 \parallel \lambda_2\nu)$ (and

We refer to $TV(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ (or similarly for PTV) as the "first penalty term" and $TV(\lambda_2\gamma_2 \parallel \lambda_2\nu)$ (and similarly for \overline{PTV}) as the "second penalty term". Intuitively, $\lambda_1(x)$ models the penalty of mass destruction/creation on the source measure μ at point x, $\lambda_2(y)$ models the penalty of mass destruction/creation on ν at point y.

Remark 3. When $\lambda_1 = \lambda_2 \equiv \lambda$, this formulation reduces to the classical OPT problem as defined in (18).

Remark 4. Unlike the classical OPT problem (18) and (19), in this formulation, when selecting $TV(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ and $TV(\lambda_2\gamma_2 \parallel \lambda_2\nu)$, we are not restricted to searching within $\Gamma_{\leq}(\mu,\nu)$ in $\mathcal{M}_{+}(\Omega^2)$. For instance, consider $\mu = \delta_0$ and $\nu = \delta_0 + \delta_1$ with $\lambda_1 \equiv 0$ and $\lambda_2 \equiv 100$. The optimal transportation plan, denoted by γ , would be $\delta_{(0,0)} + \delta_{(1,1)}$, yielding a zero cost in (22). However, $\gamma_1 = \delta_0 + \delta_1 \nleq \delta_0 = \mu$.

In the rest of this note, we will use
$$\begin{cases} A \\ B \end{cases}$$
 to denote the $\begin{cases} A \\ or \\ B \end{cases}$ for simplicity.

4.1 Entropic Regularization

Given $\gamma \in \mathcal{M}_+(\Omega^2)$, if $\gamma \ll dxdy$, the negative entropy term is defined as:

$$-H(\gamma) := \int_{\Omega^2} \left(\ln\left(\frac{d\gamma}{dxdy}\right) - 1 \right) \frac{d\gamma}{dxdy} dxdy = KL(\gamma \parallel dxdy) - 1.$$
(23)

If $\gamma \ll dxdy$, we emphasize the difference by still using $\frac{d\gamma}{dxdy}dxdy$ in the definition to indicate that $\frac{d\gamma}{dxdy}dxdy \neq d\gamma$ in this case.

With a chosen $\epsilon > 0$, we define the entropic (generalized) optimal transport problems based on (22):

$$EGOPT(\mu,\nu;\lambda_{1},\lambda_{2},\epsilon) := \inf_{\substack{\gamma \in \mathcal{M}_{+}(\Omega^{2})\\\gamma \ll dxdy}} \langle c,\gamma \rangle - \epsilon H(\gamma) + \begin{cases} TV(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu)\\PTV(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu) \end{cases} + \begin{cases} TV(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu)\\PTV(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu) \end{cases}$$
$$= \inf_{\substack{\gamma \in \mathcal{M}_{+}(\Omega^{2})\\\gamma \ll dxdy}} -\epsilon KL(\gamma \parallel K) + \begin{cases} TV(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu)\\PTV(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu) \end{cases} + \begin{cases} TV(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu)\\PTV(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu) \end{cases} + \text{constant} \end{cases}$$

where in (24), K is defined as $K := e^{-c/\epsilon}$. The equation follows from the fact:

$$KL(\gamma|Kdxdy) = \int \left(\ln\left(\frac{d\gamma}{Kdxdy}\right) - 1 \right) \frac{d\gamma}{dxdy} dxdy + \|Kdxdy\|$$

$$= \int \left(\ln\left(\frac{d\gamma}{dxdy}\right) - 1 \right) \frac{d\gamma}{dxdy} dxdy + \int \frac{c(x,y)}{\epsilon} d\gamma + \|Kdxdy\|$$

$$= \frac{1}{\epsilon} \left(-H(\gamma) + \langle c, \gamma \rangle \right) + \|Kdxdy\|.$$
(25)

As ||Kdxdy|| is a non-negative constant, independent of γ , it is disregarded in further discussions.

Remark 5. Rigorously speaking, whether in the general entropic unbalanced OT case or in the entropic GOPT, setting $\epsilon = 0$ does not guarantee equivalence between the entropic and the original versions since the optimal solution in the original version might not satisfy $\gamma \ll dxdy$. The convergence of the minimizer of EGOPT (24) to a minimizer in GOPT (22) under general conditions remains an open question.

However, if a minimizer in (22) is absolutely continuous with respect to dxdy and Ω is finite, as $\epsilon \to 0$, the minimizer of EGOPT converges to this minimizer of GOPT, as shown in [7, Proposition 3.1].

Dual formulation of entropic (generalized) OPT problems 4.2

In this section, we introduce the dual formulation of the entropic (generalized) OPT problem defined in (24), which equal the original problems (up to a constant):

$$Dual - EGOPT(\mu, \nu; \lambda_1, \lambda_2) = \sup_{\substack{\phi \in L^{\infty}(\Omega, dx) \\ \psi \in L^{\infty}(\Omega, dy)}} -\epsilon \langle (e^{\frac{\phi \oplus \psi}{\epsilon}} - 1)K, dxdy \rangle + \begin{cases} \langle \min(\phi, \lambda_1), \mu \rangle + \iota_{\phi \in \mathcal{S}(\lambda_1, dx)} \\ \langle \min(\phi, \lambda_1), \mu \rangle \end{cases} + \begin{cases} \langle \min(\psi, \lambda_2), \nu \rangle + \iota_{\psi \in \mathcal{S}(\lambda_2, dy)} \\ \langle \min(\psi, \lambda_2), \nu \rangle \end{cases} \end{cases}$$

$$(26)$$

where

$$\mathcal{S}(\lambda_1, dx) := \{\phi : \phi \ge -\lambda_1, dx - a.s.; \phi = 0, dx \mid_{\lambda_1 = 0} -a.s.\}$$

and $\mathcal{S}(\lambda_2, dy)$ is defined similarly.

• • •

The rest of this section is the derivation of this formulations (in particular, see Remark 6) and related basic concepts. The main idea is based on [7, Theorem 3.1] and related concepts.

We first setup some notations and settings. Since $\gamma \ll dxdy$ in EGOPT problem. We have $\gamma_1 \ll dx$ and $\gamma_2 \ll dy$. Thus we can write $\gamma_1 = \frac{\gamma_1}{dx}dx$, $\gamma_2 = \frac{d\gamma_2}{dy}dy$ and can use $r = \frac{\gamma_1}{dx}$ to represent γ_1 , where $r \in L_1(\Omega, dx)$ and similar to γ_2 .

4.2.1 Background: convex function, conjugate, dual formulation.

Let (E, E^*) be a pair of convex Hausdorff topologically paired vector spaces, so that the elements of each space can be identified with a continuous linear functional on the other. We refer to [11] for details. The topological pairing is the bi-linear form:

$$(E, E^*) \ni (s, s') \mapsto \langle s, s' \rangle \in \mathbb{R}$$

$$(27)$$

Example 2. $(L_1(\Omega, dx), L_{\infty}(\Omega, dx))$ is such a topologically paired vector space. In particular, we have: $(L_1(\Omega, dx))^* = L_\infty(\Omega, dx)$ and $(L_\infty(\Omega, dx))^* \supset L_1(\Omega, dx)$. Additionally, the topological pairing is defined by:

$$\langle f,g\rangle = \int_{\Omega} fgdx$$

Example 3. $(\mathbb{R}^d, \mathbb{R}^d)$ is another instance of topologically paired vector spaces. Specifically, $(\mathbb{R}^d)^* = \mathbb{R}^d$. The topological pairing can be described by:

$$\langle s, s' \rangle = \sum_{i=1}^{d} s_i s'_i$$

A function $f: E \to \overline{\mathbb{R}} = [-\infty, \infty]$ is termed a convex function if it satisfies:

$$f((1-\alpha)x + \alpha y) \le (1-\alpha)f(x) + \alpha f(y)$$
(28)

for all $x, y \in E$ and $\alpha \in [0, 1]$. The domain of f, denoted D(f), is:

$$D(f) = \{ x \in E \mid f(x) < \infty \}.$$

f is a proper function if $D(f) \cap \{f > -\infty\} \neq \emptyset$.

The convex conjugate of a function $f: E \to [-\infty, \infty]$, denoted as $f^*: E^* \to [-\infty, \infty]$, is defined as:

$$f^*(s') = \sup_{x \in E} \langle x, s' \rangle - f(x) \tag{29}$$

Example 4. The entropy functions defined in previous examples have the following convex conjugates:

$$f_{KL}^*(s') = \exp(s') - 1 \tag{30}$$

$$f_{TV}^{*}(s') = \begin{cases} \max(s', -1) & \text{if } s' \le 1\\ \infty & \text{if } s' > 1 \end{cases}$$
(31)

$$f_{PTV}^*(s') = \max(s', -1)$$
(32)

$$\iota_{\{0\}}^*(s') = s' \tag{33}$$

$$\mathbf{0}^*(s') = \iota_{\{0\}} \tag{34}$$

$$\iota_{[0,1]}^*(s') = \begin{cases} 0 & \text{if } s' \le 0\\ s' & \text{if } s' \ge 0 \end{cases}$$
(35)

Example 5. Let $E = L_1(X, dx)$ and $E^* = L_{\infty}(X, dx)$. Consider the mapping

$$E \ni r \mapsto D_f(rdx \parallel \mu) := \langle f(\frac{r}{d\mu/dx}), \mu \rangle + f'_{\infty} \parallel (rdx)^{\perp} \parallel \in \mathbb{R} \cup \{\infty\},$$

where f is an entropy function.

According to [7, Theorem A.3], this mapping is a proper convex, weakly lower-semi continuous function in $L_1(X, dx)$. Its conjugate is given by

$$E^* \ni \phi \mapsto D_f^*(\phi dx \parallel \mu) := \langle f^*(\phi), \mu \rangle + \langle \iota_{\leq f'_{\infty}} \phi, dx \rangle.$$
(36)

Thus, we have the following:

• Let $E = L_1(\Omega^2, dxdy)$ and $E^* = L_{\infty}(\Omega^2, dxdy)$, we have:

$$KL^*(\phi dxdy \parallel K dxdy) = \langle (e^{\frac{\varphi}{\epsilon}} - 1)K, dxdy \rangle$$
(37)

• Let $E = L_1(\Omega, dx)$ and $E^* = L_{\infty}(\Omega, dx)$. Consider the mapping:

$$r \mapsto TV(\lambda_1 r dx \parallel \lambda_1 \mu). \tag{38}$$

First, it is proper convex since

$$r' \mapsto TV(r'dx \parallel \lambda_1 \mu)$$

is a proper convex function, and

$$E_1 \ni r \mapsto r' = \lambda_1 r \in E_1$$

is a linear operator.

By Lemma 1, its conjugate, denoted as $TV^*(\lambda_1\phi dx \parallel \lambda_1\mu)$, is given by:

$$TV^*(\phi\lambda_1 dx \parallel \lambda_1 \mu) = \begin{cases} \langle \max(\phi, -\lambda_1), \mu \rangle & \text{if } \phi \leq \lambda_1, dx \text{-} a.s.; \phi = 0, dx \mid_{\lambda_1 = 0} \text{-} a.s. \\ \infty & elsewhere \end{cases}$$
(39)

Similarly, replace TV with PTV, we have:

$$PTV^{*}(\phi\lambda_{1}dx \parallel \lambda_{1}\mu) = \begin{cases} \langle \max(\phi, -\lambda_{1}), \mu \rangle & \text{if } \phi = 0, dx \mid_{\lambda_{1}=0} \\ \infty & \text{elsewhere} \end{cases}$$
(40)

In addition, by Lemma 3, we have:

$$\overline{PTV}^{*}(\phi\lambda_{1}dx \parallel \lambda_{1}\mu) = \langle \max(\phi, -\lambda_{1}), \mu \rangle$$
(41)

• Let $E = L_1(\Omega, dx)$ and $E^* = L_{\infty}(\Omega, dx)$. We have:

$$\iota_{=}^{*}(\phi dx \parallel \mu) = \langle \phi, \mu \rangle \tag{42}$$

$$D_{\mathbf{0}}^{*}(\phi dx \parallel \mu) = \begin{cases} 0 & \text{if } \phi \leq 0, dx \text{-} a.s. \\ \infty & \text{elsewhere} \end{cases}$$
(43)

$$\iota_{<}^{*}(\phi dx \parallel \mu) = \langle \max(\phi, 0), \mu \rangle \tag{44}$$

Lemma 1. We claim (39) holds.

Proof. From (36), we have for each $\phi' \in L_{\infty}(\Omega, dx)$:

$$TV^*(\phi'dx \parallel \lambda_1\mu) := \max_r \left(\langle \lambda_1 r \phi', dx \rangle - TV(\lambda_1 r dx \parallel \lambda_1\mu) \right) = \langle \max(\phi', -1), \lambda_1\mu \rangle + \langle \iota_{\geq 1}(\phi'), dx \rangle$$
(45)

Let $\Omega_1 = \{x : \phi > 0\}$ and $\Omega_2 = \{x : \phi = 0\}$. For each function $r \in E$, let $r \mid_{\Omega_1}, r \mid_{\Omega_2}$ denote the restriction of r on Ω_1 and Ω_2 respectively, and the restriction of measure is defined similarly. We have:

$$TV^*(\lambda_1 \phi dx \parallel \lambda_1 \mu) = \max_r \left(\langle r\phi, dx \rangle - TV(\lambda_1 r dx \parallel \lambda_1 \mu) \right)$$

=
$$\max_r \left(\langle (r \mid_{\Omega_1} + r \mid_{\Omega_2})\phi, dx \rangle - TV(\lambda_1 r \mid_{\Omega_1} dx \parallel \lambda_1 \mu) \right)$$

=
$$\max_r \left(\langle r \mid_{\Omega_1} \phi, dx \rangle - TV(\lambda_1 r \mid_{\Omega_1} dx \parallel \lambda_1 \mu) \right) + \max_{r \mid_{\Omega_2}} \langle r \mid_{\Omega_2} \phi, dx \rangle$$

=
$$\underbrace{TV^* \left(\frac{\phi \mid_{\Omega_1}}{\lambda_1 \mid_{\Omega_1}} dx \parallel \lambda_1 \mu \right)}_{B_1} + \underbrace{\max_{r \mid_{\Omega_2}} \langle r \mid_{\Omega_2} \phi, dx \rangle}_{B_2}$$

where we use the convention $\frac{0}{0} = 1$ for the notation $\frac{\phi|_{\Omega_1}}{\lambda_1|_{\Omega_1}}$. It is straightforward to verify $B_1, B_2 \ge -\infty$. Case 1: If $dx \mid_{\Omega_2} (\{\phi \neq 0\}) > 0$, then $B_2 = \infty$. Case 2: If $dx \mid_{\Omega_1} (\{\phi > \lambda_1\}) > 0$, then we have:

$$B_1 \ge \langle \iota_{\ge 1} \frac{\phi \mid_{\Omega_1}}{\lambda_1 \mid_{\Omega_1}}, dx \mid_{\Omega_1} \rangle - \|\lambda_1 \mu\| = \infty.$$

Case 3: If $\phi \leq \lambda_1$, dx-a.s., and $\phi = 0$, dx $|_{\lambda_1=0}$ -a.s., then:

$$B_1 = \langle \max(\phi, -\lambda_1), \mu \rangle, B_2 = 0 \tag{46}$$

Thus, we complete the proof.

Next, we introduce the sub-differential operator of a convex function f:

$$\partial f(x) = \{ y \in E^* : \langle x' - x, y \rangle \le f(x') - f(x), \forall x' \in E \}$$

$$\tag{47}$$

Lemma 2. Given topological pairs (E_1, E_1^*) and (E_2, E_2^*) , it is a well-known result that $(E_1 \times E_2, E_1^* \times E_2^*)$ defines a topological pair with the pairing

$$(E_1 \times E_2, E_1^* \times E_2^*) \ni ((x_1, x_2), (x_1^*, x_2^*)) \mapsto \langle x_1, x_1^* \rangle + \langle x_2, x_2^* \rangle.$$

If $E_1 \ni x_1 \mapsto f_1(x_1)$ and $E_2 \ni x_2 \mapsto f_2(x_2)$ are two functions, then $(f_1 \oplus f_2)^* = f_1^* \oplus f_2^*$. In addition, if f_1 and f_2 are convex functions, then $f_1 \oplus f_2$ is convex.

The proof directly follows from the definition of convexity and conjugate functions.

Lemma 3. We claim that the function $r \mapsto \overline{D}(\lambda_1 r dx \parallel \lambda_1 \mu)$ is a proper, lower-semi continuous, convex function. In addition, \overline{PTV}^* is given by (41).

Proof. We can split the space Ω into $\Omega_1 = \{x : \lambda_1 > 0\}$ and $\Omega_2 = \{x : \lambda_1 = 0\}$. Then,

$$PTV(\lambda_1 r dx \parallel \lambda_1 \mu) = PTV(\lambda_1 r dx \parallel \lambda_1 \mu) + \iota_{\leq}(r dx \parallel \mu)$$
$$= \underbrace{PTV(\lambda_1 r \mid_{\Omega_1} dx \parallel \lambda_1 \mu \mid_{\Omega_1})}_{A_1} + \underbrace{\iota_{\leq}(r \mid_{\Omega_2} dx \parallel \mu \mid_{\Omega_2})}_{A_2}$$

The mappings $r \mid_{\Omega_1} \to A_1$ and $r \mid_{\Omega_2} \to A_2$ are proper, convex, weakly lower semi-continuous functions. Thus, $\overline{PTV}(\lambda_1 r dx \parallel \lambda_1 \mu)$ is proper, convex, and weakly lower semi-continuous.

In addition, its conjugate is given by

$$\overline{PTV}^*(\phi\lambda_1 dx \parallel \lambda_1 \mu) = PTV^*(\lambda_1 \phi \mid_{\Omega_1} \parallel \lambda_1 dx \mid_{\Omega_1}) + \iota_{\leq}^*(\phi \mid_{\Omega_2} \parallel \mu \mid_{\Omega_2})$$
$$= \int_{\Omega_1} \max(\phi, -\lambda_1) d\mu + \int_{\Omega_2} \max(\phi, 0) d\mu$$
$$= \langle \max(\phi, -\lambda_1), \mu \rangle$$

and this completes the proof.

Given topological pairs (E_1, E_1^*) and (E_2, E_2^*) , suppose $A : E_1 \to E_2$ is a continuous linear operator. Its dual operator $A^* : E_2^* \to E_1^*$ is a continuous linear operator such that

$$\langle As_1, s_2^* \rangle = \langle s_1, A^* s_2^* \rangle, \forall s_1 \in E_1, s_2^* \in E_2^*.$$

Example 6. Consider $(E_1 = L_1(\Omega^2, dxdy), E_1^* = L_{\infty}(\Omega^2, dxdy))$ and $(E_2 = L_1(\Omega, dx) \times L_1(\Omega, dy), E_2^* = L_{\infty}(\Omega, dx) \times L_{\infty}(\Omega, dy))$. For (E_2, E_2^*) , the topological pairing is defined by:

$$\left(\left(\frac{d\gamma_1}{dx}, \frac{d\gamma_2}{dy}\right), (\phi, \psi)\right) := \left\langle \frac{d\gamma_1}{dx}, \phi dx \right\rangle + \left\langle \frac{d\gamma_2}{dy}, \psi dy \right\rangle.$$
(48)

The following is a linear continuous mapping:

$$E_1 \ni \frac{d\gamma}{dxdy} \to A\left(\frac{d\gamma}{dxdy}\right) = \left(\frac{\gamma_1}{dx}, \frac{\gamma_2}{dy}\right) \in E_2,$$

then its dual operator A^* is defined by:

$$E_2^* \ni (\phi, \psi) \to A^*(\phi, \psi) = \phi \oplus \psi \in E_1^*.$$

Next, we introduce the following celebrated theorem in convex analysis:

Theorem 1 (Rockafellar Theorem, [Theorem 3, [11], [Theorem A.1, [7]). Given topological pairs (E_1, E_1^*) and (E_2, E_2^*) , a linear continuous mapping $A : E_1 \to E_2$ with its dual operator $A^* : E_2^* \to E_1$. Suppose $F : E_1 \to [-\infty, \infty]$ and $G : E_2 \to [-\infty, \infty]$ are proper, lower semi-continuous functions. In addition, there exists $x \in Dom(F)$ such that G is continuous at Ax.

Then the following strong duality holds:

$$\inf_{x_1 \in E_1} F(x_1) + G(Ax_1) = \sup_{x_2^* \in E_2^*} -F^*(A^*x_2^*) - G^*(-x_2^*).$$
(49)

Moreover, the infimum is attained by some x_1 . If $x_2^* \in E_2^*$ is a maximizer, there exists a minimizer $x_1 \in E_1$ such that $Ax_1 \in \partial G^*(-x_2^*)$ and $A^*x_2^* \in \partial F(x_1)$.

Remark 6. Now we discuss the derivation of (26). Apply $(E_1, E_1^*), (E_2, E_2^*)$, operator A, defined in Example 6, set F, G as

$$E_{1} \ni \frac{d\gamma}{dxdy} \mapsto F\left(\frac{d\gamma}{dxdy}\right) := KL(\gamma \parallel Kdxdy)$$

$$E_{2} \ni \left(\frac{d\gamma_{1}}{dx}, \frac{d\gamma_{2}}{dy}\right) \mapsto G\left(\frac{d\gamma_{1}}{dx}, \frac{d\gamma_{2}}{dy}\right) := \begin{cases} TV(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu) \\ \overline{PTV}(\lambda_{1}\gamma_{1} \parallel \lambda_{1}\mu) \end{cases} + \begin{cases} TV(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu) \\ \overline{PTV}(\lambda_{2}\gamma_{2} \parallel \lambda_{2}\nu) \end{cases}$$
(50)

Then, by lemma 2 and the convex conjugate functions (39), (41), we have the convex conjugate functions F^*, G^* as:

$$E_{1}^{*} \ni \Phi \mapsto F^{*}(\Phi) = \langle (e^{-\Phi/\epsilon} - 1)K, dxdy \rangle$$

$$E_{2}^{*} \ni (\phi, \psi) \mapsto G^{*}(\phi, \psi) = \begin{cases} TV^{*}(\lambda_{1}\phi \parallel \lambda_{1}\mu) \\ PTV^{*}(\lambda_{1}\phi \parallel \lambda_{1}\mu) \end{cases} + \begin{cases} TV^{*}(\lambda_{2}\psi \parallel \lambda_{2}\nu) \\ PTV^{*}(\lambda_{2}\psi \parallel \lambda_{2}\nu) \end{cases}$$

$$= \langle \max(\phi, -\lambda_{1}), \mu \rangle + \langle \max(\psi, -\lambda_{2}), \nu \rangle + \begin{cases} \iota_{\phi \in \mathcal{S}^{-}(-\lambda_{1}, dx)} \\ 0 \end{cases} + \begin{cases} \iota_{\psi \in \mathcal{S}^{-}(-\lambda_{2}, dy)} \\ 0 \end{cases}$$
(51)

where

$$\mathcal{S}^{-}(-\lambda_{1}, dx) := \{ \phi \ge -\lambda_{1}, dx - a.s.; \phi = 0, dx \mid_{\lambda_{1}=0} -a.s. \},\$$

and $\mathcal{S}^{-}(-\lambda_2, dy)$ is defined similarly.

4.2.2 Alternative dual optimization

By [7, Theorem 3.1] (or Theorem 1), suppose (ϕ, ψ) is a maximizer for the dual problem (26), then the corresponding minimizer can be obtained by $\gamma = e^{\frac{\phi}{\epsilon}} K e^{\frac{\psi}{\epsilon}} dx dy$.

It remains to find optimal dual variations ϕ, ψ in the dual problem (26). Given ψ , we will find the optimal ϕ . First, we set $u = e^{\phi/\epsilon}, v = e^{\psi/\epsilon}, Kv = \langle K(x, \cdot)v(\cdot), dy \rangle, K^{\top}u = \langle K(\cdot, y)u(\cdot), dx \rangle$.

We choose the $TV(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ term, then optimizing ϕ is equivalent to

$$\max_{\phi \ge -\lambda_1, dx - \text{a.s.}} -\epsilon \langle e^{\frac{\phi}{\epsilon}} K, dx dy \rangle + \langle \min(\phi, \lambda_1), \mu \rangle.$$

By the first-order derivative method, combined with the fact that $\phi \ge -\lambda_1, dx$ – a.s. and $\phi = 0, dx \mid_{\lambda_1=0}$ –a.s., we obtain the optimal ϕ (or u) given by:

$$u(x) = \exp(\phi/\epsilon) = \underbrace{\frac{\operatorname{clip}\left(\frac{d\mu}{dx}, [e^{-\lambda_1/\epsilon}, e^{\lambda_1/\epsilon}]Kv\right)(x)}{Kv(x)}}_{Kv(x)} = \operatorname{clip}\left(\frac{d\mu/dx}{Kv(x)}, [e^{-\lambda_1/\epsilon}, e^{\lambda_1/\epsilon}]\right),$$
(52)

where $\operatorname{clip}(a, [a_1, a_2]) = \begin{cases} a & \text{if } a \in [a_1, a_2] \\ a_1 & \text{if } a < a_1 \\ a_2 & \text{if } a > a_2 \end{cases}$, and $\operatorname{prox}_{TV}^{KL}$ is called the **proximal operator** for the KL divergence and the functional $\frac{1}{\epsilon}TV(\lambda_1 \cdot \parallel \lambda_1 \mu)$:

$$\operatorname{prox}_{F_1/\epsilon}^{KL}(z) := \arg\min_s F_1(s) + \epsilon KL(s|z).$$

In addition, the operator

$$Kv \mapsto \operatorname{proxdiv}_{F_1/\epsilon}^{KL}(Kv) := \frac{\operatorname{prox}_{F_1/\epsilon}^{KL}(Kv)}{Kv}$$

is called the proximal-divide operator.

Similarly, when we choose the $\overline{PTV}(\gamma_1 \parallel \mu)$ term, ϕ (or $u = e^{\phi/\epsilon}$) is updated by the corresponding proximal-divide operator:

$$u = \operatorname{proxdiv}_{PTV(\lambda_1 \cdot \|\lambda_1\mu)/\epsilon}^{KL}(Kv) = \min\left(\frac{d\mu/dx}{Kv}, e^{\lambda_1/\epsilon}\right).$$
(53)

Similarly, for fixed ϕ (or u), the optimal ψ (or v) is given by

$$v = \exp(\psi/\epsilon) = \operatorname{proxdiv}_{TV(\lambda_2 \cdot |\lambda_2 \nu)/\epsilon}^{KL}(K^{\top}u) = \operatorname{clip}\left(\frac{\nu/dy}{K^{\top}u}, [e^{-\lambda_2/\epsilon}, e^{\lambda_2/\epsilon}]\right), \quad \text{for } TV(\lambda_2\gamma_2 \parallel \lambda_2\nu) \quad (54)$$

$$v = \exp(\psi/\epsilon) = \operatorname{proxdiv}_{\overline{PTV}(\lambda_2 \cdot | \lambda_2 \nu)/\epsilon}^{KL}(K^{\top}u) = \min\left(\frac{\nu/dy}{K^{\top}u}, e^{\lambda_2/\epsilon}\right), \quad \text{for } \overline{PTV}(\lambda_2\gamma_2 \parallel \lambda_2\nu).$$
(55)

4.3 Relation between GOPT and classical OT problem

It has been studied that the classical OPT problem (18), (19) is equivalent to a classical balanced OT problem. By this relation, one can apply the linear programming solver in OT for the OPT problem. In this section, we extend this relation to GOPT (22).

In particular, we choose $\overline{PTV}(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ and $\overline{PTV}(\lambda_2\gamma_2 \parallel \lambda_2\nu)$ in GOPT, and we obtain:

$$GOPT(\mu,\nu) = \inf_{\gamma \in \Gamma_{\leq}(\mu,\nu)} \langle c - (\lambda_1 \oplus \lambda_2), \gamma \rangle + \underbrace{\langle \lambda_1, \mu \rangle + \langle \lambda_2, \nu \rangle}_{\text{constant}}.$$
(56)

Introduce an auxiliary point $\hat{\infty}$ and let $\hat{\Omega} = \Omega \cup \{\hat{\infty}\}$. We define $\hat{c} : \hat{\Omega}^2 \to \mathbb{R}$ with

$$\hat{c}(x,y) := \begin{cases} c(x,y) - (\lambda_1(x) + \lambda_2(y)) & \text{if } (x,y) \in \Omega^2\\ 0 & \text{elsewhere} \end{cases}$$
(57)

In addition, define

$$\hat{\mu} = \mu + |\nu| \delta_{\hat{\infty}}$$
$$\hat{\nu} = \nu + |\mu| \delta_{\hat{\infty}}$$

Thus, $\|\hat{\mu}\| = \|\hat{\nu}\|$. By [5, 2], the following mapping:

$$\Gamma_{\leq}(\mu,\nu) \ni \gamma \mapsto \hat{\gamma} := \gamma + (\mu - \gamma_1) \otimes \delta_{\hat{\infty}} + \delta_{\hat{\infty}} \otimes (\nu - \gamma_2) + |\gamma| \delta_{(\hat{\infty},\hat{\infty})} \in \Gamma(\hat{\mu},\hat{\nu})$$
(58)

is a well-defined bijection.

We claim the following:

Lemma 4. Consider the balanced OT problem:

$$\inf_{\hat{\gamma}\in\Gamma(\hat{\mu},\hat{\nu})}\int_{\hat{\Omega}^2}\hat{c}d\hat{\gamma}$$
(59)

Choose $\gamma \in \Gamma_{\leq}(\mu,\nu)$ and find $\hat{\gamma}$ by (58). We have that $\hat{\gamma}$ is a minimizer of (59) if and only if γ is a minimizer for (56).

Proof. We have:

$$\int_{\hat{\Omega}^2} \hat{c} d\hat{\gamma} = \int_{\Omega^2} (c - (\lambda_1 \oplus \lambda_2)) d\gamma$$

Since the mapping (58) is a bijection, we have that $\hat{\gamma}$ is a minimizer for the OT problem (59) if and only if γ is a minimizer for the OPT problem (56).

In addition, it is straightforward to verify the following:

Lemma 5. In problem (56), choose $\epsilon' > 0$, then for each optimal $\gamma \in \Gamma_{\leq}(\mu, \nu)$, we have

$$\gamma(\{c - (\lambda_1 \oplus \lambda_2) \ge \epsilon'\}) = 0.$$

Proof. Pick an optimal $\gamma \in \Gamma_{<}(\mu, \nu)$ and let $\mathcal{A} = \{c - (\lambda_1 \oplus \lambda_2) \geq \epsilon'\}$. Suppose $\gamma(\mathcal{A}) > 0$. We have $\gamma \mid_{\mathcal{A}} \leq \gamma$, thus $\gamma \mid_{\mathcal{A}^c} \in \Gamma_{\leq}(\mu, \nu)$. In addition,

$$\langle c - (\lambda_1 \oplus \lambda_2), \gamma \rangle - \langle c - (\lambda_1 \oplus \lambda_2), \gamma \mid_{\mathcal{A}^c} \rangle = \langle c - (\lambda_1 \oplus \lambda_2), \gamma \mid_{\mathcal{A}} \rangle \ge \epsilon' \|\gamma \mid_{\mathcal{A}} \| > 0$$

This is a contradiction since γ is optimal.

Lemma 6. In problem (56), suppose $|\nu| \leq |\mu|$. If for all $(x, y) \in \Omega^2$, $c - (\lambda_1 \oplus \lambda_2) < -\epsilon_1$ for some $\epsilon_1 > 0$, then for each optimal transportation plan γ , we have $\gamma_2 = \nu$.

Proof. Choose an optimal γ and suppose $|\gamma| < |\nu|$. Then by [Lemma E.1 [1]], there exists a $\gamma' \in \Gamma(\mu, \nu)$ such that $\gamma'_2 = \nu, \gamma \leq \gamma'$, and $\gamma'_1 \leq \mu$. Then we have:

$$\langle c - (\lambda_1 \oplus \lambda_2), \gamma' \rangle - \langle c - (\lambda_1 \oplus \lambda_2), \gamma \rangle = \langle c - (\lambda_1 \oplus \lambda_2), \gamma' - \gamma \rangle \le -\epsilon_1 \|\gamma' - \gamma\| < 0$$

This is a contradiction to the fact that γ is an optimal transportation plan.

Algorithms for GOPT problems 4.4

4.4.1 OPT problems in a discrete setting

Suppose $\mu = \sum_{i=1}^{n} p_i \delta_{x_i}$ and $\nu = \sum_{j=1}^{m} q_j \delta_{y_j}$, where $p > 0_n$ and $q > 0_m$ for all i, j. We set $dx = \frac{1}{n} \sum_{i=1}^{n} \delta_{x_i}$ and $dy = \frac{1}{m} \sum_{j=1}^{m} \delta_{y_j}$. Without loss of generality, we can redefine dx and dy as $dx = \sum_{i=1}^{n} \delta_{x_i}$ and $dy = \sum_{j=1}^{m} \delta_{y_j}$ since the (positive) scaling of reference measures will not change the OPT problems. In additionance dx = 0, dx = 0, dx = 0, dx = 0, dx = 0. addition, we set $\lambda_1 \in \mathbb{R}^n_+$ and $\lambda_2 \in \mathbb{R}^m_+$.

Finally, choose $\epsilon > 0$. We adopt $c \in \mathbb{R}^{n \times m}$ as the cost function (matrix) in OPT problems and define $K = e^{-c/\epsilon}$. The negative entropy is defined by $-H(\gamma) := \sum_{i,j} (\ln(\gamma_{ij}) - 1)\gamma_{ij}$. Then the generalized OPT (22), the entropic version (24), and the corresponding dual forms (26), can

be written as follows:

$$GOPT(\mu,\nu;\lambda_{1},\lambda_{2}) = \min_{\gamma \in \mathbb{R}^{n \times m}_{+}} \sum_{i,j} c_{ij}\gamma_{ij} + \begin{cases} \sum_{i} (\lambda_{1})_{i} |(p_{i} - (\gamma \mathbf{1}_{m})_{i})| \\ \sum_{i} (\lambda_{1})_{i} (p_{i} - (\gamma \mathbf{1}_{m})_{i}) + \iota_{\{\gamma \mathbf{1}_{m} \leq p\}} \end{cases} \\ + \begin{cases} \sum_{j} (\lambda_{2})_{j} |(q_{j} - (\gamma^{\top} \mathbf{1}_{n})_{j})| \\ \sum_{j} (\lambda_{2})_{j} (q_{j} - (\gamma^{\top} \mathbf{1}_{n})_{j}) + \iota_{\{\gamma^{\top} \mathbf{1}_{n} \leq q\}} \end{cases}$$
(60)

$$EGOPT(\mu,\nu;\lambda_{1},\lambda_{2},\epsilon) = \min_{\gamma \in \mathbb{R}^{n \times m}_{+}} \sum_{i,j} c_{ij}\gamma_{ij} + \epsilon \sum_{i,j} (\ln(\gamma_{ij}) - 1)\gamma_{ij} + \begin{cases} \sum_{i} (\lambda_{1})_{i} |(p_{i} - (\gamma \mathbf{1}_{m})_{i})| \\ \sum_{i} (\lambda_{1})_{i} (p_{i} - (\gamma \mathbf{1}_{m})_{i}) + \iota_{\{\gamma \mathbf{1}_{m} \leq p\}} \end{cases} + \begin{cases} \sum_{j} (\lambda_{2})_{j} |(q_{j} - (\gamma^{\top} \mathbf{1}_{n})_{j})| \\ \sum_{j} (\lambda_{2})_{j} (q_{j} - (\gamma^{\top} \mathbf{1}_{n})_{j}) + \iota_{\{\gamma^{\top} \mathbf{1}_{n} \leq q\}} \end{cases}$$
(61)

$$Dual - EGOPT(\mu, \nu; \lambda_1, \lambda_2, \epsilon) = \max_{\substack{\phi \in \mathbb{R}^n \\ \psi \in \mathbb{R}^m}} -\epsilon \sum_{i,j} (e^{\frac{\phi_i + \psi_j}{\epsilon}} - 1) K_{ij} + \begin{cases} \sum_i \min((\lambda_1)_i, \phi_i) p_i \\ \sum_i \min((\lambda_1)_i, \phi_i) p_i + \iota_{\{\phi \ge -\lambda_1\}} \end{cases} \\ + \begin{cases} \sum_j \min((\lambda_2)_j, \psi_j) q_j \\ \sum_j \min((\lambda_2)_j, \psi_j) q_j + \iota_{\{\psi \ge -\lambda_2\}} \end{cases}$$
(62)

4.4.2 Sinkhorn algorithm

The Sinkhorn algorithm aims to solve (62). In particular, set $u = e^{\phi/\epsilon} \in \mathbb{R}^n$ and $v = e^{\psi/\epsilon} \in \mathbb{R}^m$. From the fact $dx \ll \mu$ in the discrete setting, the alternative optimization steps (52) (or (53)), (54) (or (55)) become:

$$u \leftarrow \operatorname{proxdiv}_{TV(\lambda_1 \cdot | \lambda_1 p)/\epsilon}^{KL}(Kv) = \operatorname{clip}\left(\frac{p}{Kv}, [e^{-\lambda_1/\epsilon}, e^{\lambda_1/\epsilon}]\right) \qquad \text{for } TV(\lambda_1 \gamma_1 \parallel \lambda_1 \mu) \tag{63}$$

$$u \leftarrow \operatorname{proxdiv}_{\overline{PTV}(\lambda_1 \cdot |\lambda_1 p)/\epsilon}^{KL}(Kv) = \min\left(\frac{p}{Kv}, e^{\lambda_1/\epsilon}\right) \qquad \text{for } \overline{PTV}(\lambda_1 \gamma_1 \parallel \lambda_1 \mu) \qquad (64)$$

$$v \leftarrow \operatorname{proxdiv}_{TV(\lambda_2 \cdot | \lambda_2 q)/\epsilon}^{KL}(K^{\top}u) = \operatorname{clip}\left(\frac{q}{K^{\top}u}, [e^{-\lambda_2/\epsilon}, e^{\lambda_2/\epsilon}]\right) \qquad \text{for } TV(\lambda_2\gamma_2 \parallel \lambda_2\nu) \tag{65}$$

$$v \leftarrow \operatorname{proxdiv}_{\overline{PTV}(\lambda_2 \cdot | \lambda_2 q)/\epsilon}^{KL}(K^{\top} u) = \min\left(\frac{q}{K^{\top} u}, e^{\lambda_2/\epsilon}\right) \qquad \text{for } \overline{PTV}(\lambda_2 \gamma_2 \parallel \lambda_2 \nu) \qquad (66)$$

where "proxdiv" denotes the "proximal-divide operator".

In summary, the Sinkhorn algorithm is presented in Algorithm 1. When solving the (entropic) GOPT problem, if we choose $TV(\lambda_1\gamma_1 \parallel \lambda_2\mu)$ as the first penalty term, we apply (63) as $\operatorname{proxdiv}_{F_1}$; if we choose $\overline{PTV}(\lambda_1\gamma_1 \parallel \lambda_2\mu)$ as the first penalty term, we apply (64) as $\operatorname{proxdiv}_{F_1}$. Similarly, we set $\operatorname{proxdiv}_{F_2}$ accordingly.

Algorithm 1: opt-Sinkhorn

Input: c, ϵ , proxdiv_{F1}, proxdiv_{F2} Output: γ 1 Initialize $v = 1_m, K = e^{-c/\epsilon}$ 2 for l = 1, 2, ... do 3 $\begin{bmatrix} u = \operatorname{proxdiv}_{F_1}^{KL}(Kv) \text{ by (63) or (64)} \\ v = \operatorname{proxdiv}_{F_2}^{KL}(K^{\top}u) \text{ by (65) or (66)} \\ \text{If } (u, v) \text{ converge, break} \end{bmatrix}$ 6 $\gamma \leftarrow (u_i K_{ij} v_j)_{ij}$

4.4.3 Linear programming

When we select PTV as the first and second penalty terms, based on the relation between GOPT and OT as discussed in Lemma (4), we proceed as follows: Let $\hat{c} \in \mathbb{R}^{(n+1)\times(m+1)}$ with

$$\hat{c}_{i,j} = \begin{cases} c_{i,j} - (\lambda_1)_i - (\lambda_2)_j & \text{if } i \in [1:n], j \in [1:m] \\ 0 & \text{elsewhere} \end{cases}$$
(67)

Similarly, set $\hat{p} \in \mathbb{R}^{n+1}, \hat{q} \in \mathbb{R}^{m+1}$ with

$$\hat{p}_i = \begin{cases} p_i & \text{if } i \in [1:n] \\ \sum_j q_j & \text{if } i = n+1 \end{cases}$$

$$\tag{68}$$

$$\hat{q}_{j} = \begin{cases} q_{j} & \text{if } j \in [1:m] \\ \sum_{i} p_{i} & \text{if } j = m+1 \end{cases}$$
(69)

The linear programming solver is given in Algorithm 2.

4.4.4 Extreme case

Suppose $|\mu| \ge |\nu|$. We consider the following special optimal partial transport problem:

$$SOPT(\mu,\nu) := \inf_{\substack{\gamma \in \Gamma_{\leq}(\mu,\nu)\\\gamma_2 = \nu}} \langle c, \gamma \rangle$$
(70)

$$= \inf_{\gamma \in \mathcal{M}_{+}(\Omega^{2})} \langle c, \gamma \rangle + \iota_{\leq}(\gamma_{1}|\mu) + \iota_{=}(\gamma_{2}|\nu)$$
(71)

Algorithm 2: opt-lp Input: $c, p, q, \lambda_1, \lambda_2$ Output: γ 1 Compute $\hat{c}, \hat{p}, \hat{q}$ using (67), (68), (69) 2 Solve the OT problem $\hat{\gamma}^* = \arg \min_{\hat{\gamma} \in \Gamma(\hat{p}, \hat{q})} \langle \hat{c}, \hat{\gamma} \rangle$ 3 Set $\gamma \leftarrow \hat{\gamma}^* [1:n, 1:m]$

Remark 7. In the discrete case, we can apply the linear programming method (2) to solve it. In particular, we set $\lambda_1 \equiv 0$, $\lambda_2 \equiv \max(c) + 1$ in the PGOPT problem (56). By Lemma 6, the optimal solution satisfies $\gamma_2 = \nu$ and $\gamma_1 \leq \mu$.

The entropic version of the above Special-OPT (up to a constant) is:

$$ESOPT(\mu,\nu) := \inf_{\substack{\gamma \in \mathcal{M}_{+}(\Omega^{2})\\\gamma \ll dxdy}} \epsilon KL(\gamma \parallel Kdxdy) + \iota_{\leq}(\gamma_{1}\mid\mu) + \iota_{=}(\gamma_{2}\mid\nu)$$
(72)

By the convex-conjugate of $\iota_{=}$ and ι_{\leq} (see (42), (44)), we can derive the equivalent dual form:

$$Dual - SOPT(\mu, \nu) = \sup_{\substack{\phi \in L_{\infty}(\Omega, dx)\\\psi \in L_{\infty}(\Omega, dy)}} -\epsilon \langle e^{\phi \oplus \psi} K, dx dy \rangle + \langle \min(\phi, 0), \mu \rangle + \langle \psi, \nu \rangle$$
(73)

Remark 8. The problem (70) and its entropic version (72) can be obtained by setting λ_1, λ_2 to extreme values in GOPT problems (22),(24).

Indeed, for each $D \subset \Omega$, and let $D^c = \Omega \setminus \{D\}$, it is clear that as $\lambda_1 \to \infty$ on D, and λ_1 is fixed on D^c , we have:

$$TV(\lambda_1 \cdot \parallel \lambda_1 \mu) \to \iota_{=}(\cdot \mid_D \parallel \mu \mid_D) + TV(\lambda_1 \cdot \mid_{D^c} \parallel \lambda_1 \mu \mid_{D^c})$$
(74)

$$\overline{PTV}(\lambda_1 \cdot \parallel \lambda_1 \mu) \to \iota_{=}(\cdot \mid_D \parallel \mu \mid_D) + \overline{PTV}(\lambda_1 \cdot \mid_{D^c} \parallel \lambda_1 \mu \mid_{D^c})$$
(75)

Similarly, by setting $\lambda_1 = 0$ on D, we obtain:

$$TV(\lambda_1 \cdot \| \lambda_1 \mu) = TV(\lambda_1 \cdot |_{D^c} \| \lambda_1 \mu|_{D^c})$$
(76)

$$\overline{PTV}(\lambda_1 \cdot \parallel \lambda_1 \mu) = \iota_{\leq}(\cdot, \mu) + \overline{PTV}(\lambda_1 \cdot \mid_{D^c} \parallel \lambda_1 \mu \mid_{D^c})$$
(77)

We choose $\overline{PTV}(\lambda_1\gamma_1 \parallel \lambda_1\mu)$ as the first penalty term and $\overline{PTV}(\lambda_2\gamma_2 \parallel \lambda_2\nu)$ as the second penalty term. Setting $D = \Omega$, and let $\lambda_1 \equiv 0, \lambda_2 \to \infty$. Then we have (26) becomes (73).

It is straightforward to verify that the Sinkhorn iteration of (73) in step is

$$\begin{cases} u = e^{\phi} := \operatorname{proxdiv}_{F_1}(Kv) = \min\left(\frac{d\mu/dx}{Kv}, 1_n\right) = \min\left(\frac{p}{Kv}, 1_n\right) \\ v = e^{\psi} := \operatorname{proxdiv}_{F_2}(K^{\top}u) = \frac{d\nu/dy}{Kv} = \frac{q}{K^{\top}u} \end{cases}$$
(78)

5 Related Work: Mass-Constrained Partial Transport Problem

The mass-constrained optimal partial transport (MOPT) problem is defined by (21). In this section, we introduce the discrete version and related computations.

First, suppose $\mu = \sum_{i=1}^{n} p_i \delta_{x_i}$ and $\nu = \sum_{j=1}^{m} q_j \delta_{y_j}$ with $\eta \in [0, \min(\|p\|, \|q\|)]$, where $\|p\| = \sum_{i=1}^{n} p_i$ and $\|q\|$ is defined similarly. Then the discrete version of (21) is defined by

$$MOPT(\mu,\nu) := \min_{\gamma \in \Gamma_{\leq}^{\eta}(p,q)} \sum_{i,j} c_{ij} \gamma_{ij}$$
(79)

where $\Gamma^{\eta}_{\leq}(p,q) := \{ \gamma \in \mathbb{R}^{n \times m}_{+} : \gamma \mathbf{1}_n \leq q, \gamma^{\top} \mathbf{1}_m \leq p, \sum_{ij} \gamma = \eta \}.$

By [6], it can be solved by the following: Let $\hat{c} \in \mathbb{R}^{(n+1)\times(m+1)}, \hat{p} \in \mathbb{R}^{n+1}_+, \hat{q} \in \mathbb{R}^{m+1}_+$ as

$$\hat{c}_{ij} = \begin{cases}
c_{i,j} & \text{if } i \in [1:n], j \in [1:m] \\
\max(c) + 2\alpha + \beta & \text{if } i = n+1, j = m+1 \\
\alpha & \text{elsewhere}
\end{cases}$$
(80)

$$\hat{p}_{i} = \begin{cases} p_{i} & \text{if } i \in [1:n] \\ \|q\| - \eta & \text{if } i = n+1 \end{cases} \quad \hat{q}_{j} = \begin{cases} q_{j} & \text{if } j \in [1:m] \\ \|p\| - \eta & \text{if } j = m+1 \end{cases}$$
(81)

where $\alpha \geq 0, \beta > 0$.

Then the OT problem

$$\min_{\hat{\gamma} \in \Gamma(\hat{p}, \hat{q})} \sum_{i, j} \hat{c}_{ij} \hat{\gamma}_{i, j}$$

is equivalent to the MOPT problem (21). $\hat{\gamma}$ is optimal for the above OT problem if and only if $\gamma = \hat{\gamma}[1 : n, 1 : m]$ is optimal for the MOPT problem (21).

Thus, applying $\hat{c}, \hat{p}, \hat{q}$ in Algorithm (2) returns the solution for the MOPT problem (79).

The entropic regularization version of problem (79) (up to a constant) is defined as

$$EMOPT(\mu,\nu) := \min_{\gamma \in \Gamma_{\leq}^{\eta}(p,q)} KL(\gamma \parallel K)$$
(82)

where $K = e^{-c/\epsilon}$ and $\epsilon > 0$.

Note,

$$\gamma \in \Gamma^{\eta}_{\leq}(p,q) = \bigcap_{i=1}^{3} \mathcal{C}_{i}$$

where $C_1 = \{\gamma \in \mathbb{R}^{n \times m}_+ : \gamma \mathbb{1}_m \leq q\}$, $C_2 = \{\gamma \in \mathbb{R}^{n \times m}_+ : \gamma^\top \mathbb{1}_n \leq p\}$, and $C_3 = \{\gamma \in \mathbb{R}^{n \times m}_+ : \sum \gamma = \eta\}$. C_1 and C_2 are convex, but not affine sets. C_3 is an affine set.

The entropic MOPT problem (82) can be solved by the following so-called Dykstra's algorithm [3]:

Algorithm 3: mopt-Dykstra
Input: p, q, η, c
Output: γ
1 Initialization
2 for <i>i</i> =1,2,3 do
$3 \begin{bmatrix} \xi^i \leftarrow 1_{n \times m} \end{bmatrix}$
$4 \gamma^{(0)} \leftarrow K \frac{m}{\ K\ }$
5 Main loop
6 $k = 0$ for $l = 0, 1, 2,$ do
7 for $i=1,2,3$ do
\mathbf{s} $k \leftarrow k+1$
9 $\gamma^{(k)} \leftarrow \operatorname{Proj}_{\mathcal{C}_i}^{KL}(\gamma^{(k-1)} \odot \xi^i)$
10 $\xi^i \leftarrow \xi^i \odot \frac{\gamma^{(k-1)}}{\gamma^{(k)}}$
11 Break if $\gamma^{(k)}$ converges

where for i = 1, 2, 3,

$$\operatorname{Proj}_{\mathcal{C}_{i}}^{KL}(\gamma) := \min_{\gamma^{i} \in \mathcal{C}_{i}} KL(\gamma^{i} \parallel \gamma)$$
(83)

is called the "**Bregman projection**" (which can be regarded the equivalent primal form of the proximaldivide operator discussed in (52)) onto the convex set C_i . By [4, Proposition 5], these projection operators are given by:

$$\operatorname{Proj}_{\mathcal{C}^{1}}(\gamma) = \operatorname{diag}\left(\min\left(\frac{p}{\gamma 1_{m}}, 1_{n}\right)\right)\gamma\tag{84}$$

$$\operatorname{Proj}_{\mathcal{C}^2}(\gamma) = \gamma \operatorname{diag}\left(\min\left(\frac{q}{\gamma^{\top} \mathbf{1}_n}, \mathbf{1}_m\right)\right)$$
(85)

$$\operatorname{Proj}_{\mathcal{C}^{3}}(\gamma) = \gamma \frac{\eta}{\|\gamma\|}$$
(86)

6 Acknowledgement

The author thanks Dr. Soheil Kolouri (soheil.kolouri@vanderbilt.edu) for helpful discussion.

References

- Yikun Bai, Rocio Diaz Martin, Hengrong Du, Ashkan Shahbazi, and Soheil Kolouri. Efficient solvers for partial gromov-wasserstein. arXiv preprint arXiv:2402.03664, 2024.
- [2] Yikun Bai, Bernard Schmitzer, Mathew Thorpe, and Soheil Kolouri. Sliced optimal partial transport. arXiv preprint arXiv:2212.08049, 2023.
- [3] Heinz H Bauschke and Adrian S Lewis. Dykstras algorithm with bregman projections: A convergence proof. Optimization, 48(4):409–427, 2000.
- [4] Jean-David Benamou, Guillaume Carlier, Marco Cuturi, Luca Nenna, and Gabriel Peyré. Iterative bregman projections for regularized transportation problems. SIAM Journal on Scientific Computing, 37(2):A1111-A1138, 2015.
- [5] Luis A Caffarelli and Robert J McCann. Free boundaries in optimal transport and monge-ampere obstacle problems. Annals of mathematics, pages 673–730, 2010.
- [6] Laetitia Chapel, Mokhtar Z Alaya, and Gilles Gasso. Partial optimal tranport with applications on positive-unlabeled learning. Advances in Neural Information Processing Systems, 33:2903–2913, 2020.
- [7] Lenaic Chizat, Gabriel Peyré, Bernhard Schmitzer, and François-Xavier Vialard. Scaling algorithms for unbalanced optimal transport problems. *Mathematics of Computation*, 87(314):2563–2609, 2018.
- [8] Alessio Figalli. The optimal partial transport problem. Archive for rational mechanics and analysis, 195(2):533-560, 2010.
- [9] Alessio Figalli and Nicola Gigli. A new transportation distance between non-negative measures, with applications to gradients flows with dirichlet boundary conditions. *Journal de mathématiques pures et* appliquées, 94(2):107–130, 2010.
- [10] Benedetto Piccoli and Francesco Rossi. Generalized wasserstein distance and its application to transport equations with source. Archive for Rational Mechanics and Analysis, 211(1):335–358, 2014.
- [11] Ralph Rockafellar. Duality and stability in extremum problems involving convex functions. Pacific Journal of Mathematics, 21(1):167–187, 1967.
- [12] Cedric Villani. Optimal transport: old and new. Springer, 2009.
- [13] Cédric Villani. Topics in optimal transportation, volume 58. American Mathematical Soc., 2021.