

HARMONICS AND GRADED EHRHART THEORY

VICTOR REINER AND BRENDON RHOADES

ABSTRACT. The Ehrhart polynomial and Ehrhart series count lattice points in integer dilations of a lattice polytope. We introduce and study a q -deformation of the Ehrhart series, based on the notions of harmonic spaces and Macaulay's inverse systems for coordinate rings of finite point configurations. We conjecture that this q -Ehrhart series is a rational function, and introduce and study a bigraded algebra whose Hilbert series matches the q -Ehrhart series. Defining this algebra requires a new result on Macaulay inverse systems for Minkowski sums of point configurations.

1. INTRODUCTION

A *lattice polytope* P in \mathbb{R}^n is the convex hull of a finite set of points in \mathbb{Z}^n . With their deep connections to commutative algebra and toric geometry, lattice polytopes are a central object in algebraic combinatorics. Our goal is to introduce in a canonical way an extra q -parameter into two important algebro-combinatorial objects associated to a lattice polytope P : the Ehrhart polynomials/series and the affine semigroup ring associated to P .

1.1. Classical Ehrhart theory and a q -analogous conjecture. The classical story to be generalized is one of lattice point enumeration. Given a d -dimensional lattice polytope $P \subset \mathbb{R}^n$ with boundary $\text{Bd}(P)$ and (*relative*) *interior* $\text{int}(P) = P \setminus \text{Bd}(P)$, for $m \geq 0$, one has its m -fold *dilate* $mP := \{mp : p \in P\} \subseteq \mathbb{R}^n$, along with its interior $\text{int}(mP)$. Define these integer point enumerators $i_P(m)$, $\bar{i}_P(m)$, and their generating functions $E_P(t)$, $\bar{E}_P(t)$ in $\mathbb{Z}[[t]]$:

$$(1) \quad i_P(m) := \#(\mathbb{Z}^n \cap mP) \text{ for } m \geq 0, \quad E_P(t) := \sum_{m=0}^{\infty} i_P(m)t^m,$$

$$(2) \quad \bar{i}_P(m) := \#(\mathbb{Z}^n \cap \text{int}(mP)) \text{ for } m \geq 1, \quad \bar{E}_P(t) := \sum_{m=1}^{\infty} \bar{i}_P(m)t^m.$$

We summarize here some of the results from the original work of Ehrhart [12], as well as Macdonald [33] and Stanley [41, 43]; see also expositions and surveys in Braun [7], Beck and Robins [4], Beck and Sanyal [5], Stanley [45, Ch. 1] [47, §4.6].

Classical Ehrhart Theorem. *Let P be a d -dimensional lattice polytope.*

- (i) *Both $i_P(m)$ and $\bar{i}_P(m)$ are polynomial functions of m of degree d . Both generating functions $E_P(t)$ and $\bar{E}_P(t)$ are rational functions which can be written over the denominator $(1-t)^{d+1}$, and whose numerator polynomials lie in $\mathbb{Z}[t]$, with numerator of degree at most d in the case of $E_P(t)$ and degree exactly $d+1$ in the case of $\bar{E}_P(t)$.*
- (ii) *The integer coefficients $\{h_i^*\}_{i=0}^d$ defined uniquely by*

$$E_P(t) = \frac{\sum_{i=0}^d h_i^* t^i}{(1-t)^{d+1}}$$

1

are nonnegative, with $h_0^* = 1$ and $\sum_{i=0}^d h_i^* = v$ the normalized d -volume $v := \text{vol}_d(P)$.
 (iii) For P a lattice d -simplex with vertices $\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(d+1)}$, one can interpret the $\{h_i^*\}_{i=0}^d$ as

$$h_i^* = \#\{(x_0, \mathbf{z}) \in \Pi \cap \mathbb{Z}^{n+1} : x_0 = i\},$$

where $\Pi \subseteq \mathbb{R}^{n+1} = \mathbb{R}^1 \times \mathbb{R}^n$ is this semi-open parallelepiped spanned by $\{(1, \mathbf{v}^{(j)})\}_{j=1, \dots, d+1}$:

$$(3) \quad \Pi := \left\{ \sum_{j=1}^{d+1} c_j \cdot (1, \mathbf{v}^{(j)}) : 0 \leq c_i < 1 \right\}.$$

(iv) (Ehrhart-Macdonald reciprocity) The series $E_P(t), \bar{E}_P(t)$ determine each other uniquely by $\bar{E}_P(t) = (-1)^{d+1} E_P(t^{-1})$.


Our goal is to introduce a natural q -parameter into this picture¹. Let \mathbf{k} be a field and let $\mathcal{Z} \subseteq \mathbf{k}^n$ be a finite locus of points in affine n -space over \mathbf{k} . The *point-orbit method* (also known as the *orbit harmonics method*) is a machine which produces from \mathcal{Z} a graded quotient $R(\mathcal{Z})$ of the polynomial ring $S := \mathbf{k}[x_1, \dots, x_n]$. Let $\mathbf{I}(\mathcal{Z}) \subseteq S$ be the vanishing ideal of \mathcal{Z} given by

$$(4) \quad \mathbf{I}(\mathcal{Z}) := \{f(\mathbf{x}) \in S : f(\mathbf{z}) = 0 \text{ for all } \mathbf{z} \in \mathcal{Z}\}.$$

Since \mathcal{Z} is finite, by Lagrange interpolation one has an identification of the space $\mathbf{k}[\mathcal{Z}]$ of functions $\mathcal{Z} \rightarrow \mathbf{k}$ and the quotient ring $S/\mathbf{I}(\mathcal{Z})$. We let

$$(5) \quad R(\mathcal{Z}) := S/\text{gr } \mathbf{I}(\mathcal{Z})$$

where $\text{gr } \mathbf{I}(\mathcal{Z}) \subseteq S$ is the *associated graded ideal* of $\mathbf{I}(\mathcal{Z})$, generated by the top degree homogeneous components $\tau(f)$ of nonzero polynomials $f \in \mathbf{I}(\mathcal{Z})$. The ideal $\text{gr } \mathbf{I}(\mathcal{Z})$ is homogeneous, so the quotient ring $R(\mathcal{Z})$ is graded. Geometrically, the construction $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ is a flat deformation of the reduced subscheme $\mathcal{Z} \subseteq \mathbf{k}^n$ deforming linearly to a zero dimensional subscheme of degree $|\mathcal{Z}|$ supported at the origin, as illustrated by the picture below.



The construction $\mathcal{Z} \rightsquigarrow R(\mathcal{Z})$ is a canonical way to put a graded structure on the finite locus \mathcal{Z} . One has an isomorphism of \mathbf{k} -vector spaces

$$(6) \quad \mathbf{k}[\mathcal{Z}] = S/\mathbf{I}(\mathcal{Z}) \cong R(\mathcal{Z}).$$

In particular, introducing a grading variable q , one finds that the *Hilbert series*

$$\text{Hilb}(R(\mathcal{Z}), q) := \sum_{d=0}^{\infty} \dim_{\mathbf{k}} R(\mathcal{Z})_d \cdot q^d$$

is a polynomial in q with nonnegative integer coefficients giving a geometrically motivated and canonical q -analogue of the cardinality of \mathcal{Z} , that is,

$$(7) \quad [\text{Hilb}(R(\mathcal{Z}), q)]_{q=1} = \dim_{\mathbf{k}} R(\mathcal{Z}) = \dim_{\mathbf{k}} \mathbf{k}[\mathcal{Z}] = \#\mathcal{Z}.$$

Furthermore, when \mathcal{Z} is stable under the action of a finite subgroup $G \subseteq GL_n(\mathbf{k})$ (as suggested by the previous picture where $G \cong \mathfrak{S}_3$ and the transpositions in \mathfrak{S}_3 act by reflections in the three lines), the vector space isomorphism (6) is also a *Brauer isomorphism* of $\mathbf{k}G$ -modules, that is, $\mathbf{k}[\mathcal{Z}]$

¹Note that Chapoton [8] introduced a *different*, less canonical q -analogue, discussed in Remark 3.16 below.

and $R(\mathcal{Z})$ share the same composition multiplicities for all simple $\mathbf{k}G$ -modules. In particular, when $|G|$ is invertible in \mathbf{k}^\times , such as when \mathbf{k} characteristic zero, then (6) is a $\mathbf{k}G$ -module isomorphism.

However, the deformation $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ does not have perfect memory: since $R(\mathcal{Z})$ is not reduced for $|\mathcal{Z}| > 1$, the rings $S/\mathbf{I}(\mathcal{Z})$ and $R(\mathcal{Z})$ are almost never isomorphic. Despite this amnesia, we will exhibit an unexpected multiplicative structure on direct sums of point-orbit rings (or rather their associated *harmonic spaces*) coming from dilates of a lattice polytope.

Introduced by Kostant [29] in his study of coinvariant rings of finite reflection groups, the rings $R(\mathcal{Z})$ have proven to be an enormously fruitful source of graded quotients of S . The rings $R(\mathcal{Z})$ present the cohomology of Springer fibers [18], have ties to Coxeter-Catalan theory [2], give generalized coinvariant rings with ties to Macdonald-theoretic delta operators [25, 22], give graded modules over (0-)Hecke algebras [27, 28], prove cyclic sieving results [36], and encode the Viennot shadow line construction of the Schensted correspondence [38, 31]. The rings $R(\mathcal{Z})$ have also been used to understand [37] Donaldson-Thomas invariants of symmetric quivers when \mathcal{Z} consists of lattice points in certain zonotopes. In all of these examples, it is crucial to choose a point locus $\mathcal{Z} \subseteq \mathbf{k}^n$ with strategic organization.

Here we will take $\mathbf{k} = \mathbb{R}$ and consider the case where $\mathcal{Z} = P \cap \mathbb{Z}^n$ consists of the integer points of an arbitrary lattice polytope $P \subseteq \mathbb{R}^n$, together with its dilates. Define two q -Ehrhart series for P , analogous to those in (1), (2)

$$(8) \quad E_P(t, q) := \sum_{m \geq 0} i_P(m; q) \cdot t^m, \quad \text{where } i_P(m; q) := \text{Hilb}(R(\mathbb{Z}^n \cap mP), q)$$

$$(9) \quad \bar{E}_P(t, q) := \sum_{m \geq 1} \bar{i}_P(m; q) \cdot t^m, \quad \text{where } \bar{i}_P(m; q) := \text{Hilb}(R(\mathbb{Z}^n \cap \text{int}(mP)), q).$$

Thus $E_P(t, q), \bar{E}_P(t, q)$ lie in $\mathbb{Z}[q][[t]]$, and (7) implies they specialize to $E_P(t), \bar{E}_P(t)$ when $q \rightarrow 1$. The following is our main conjecture, on the form of $E_P(t, q), \bar{E}_P(t, q)$.

Conjecture 1.1. *Let P be a d -dimensional lattice polytope in \mathbb{R}^n , having V vertices. Then both of the series (8),(9) lie in $\mathbb{Q}(t, q)$, and are expressible as rational functions*

$$E_P(t, q) = \frac{N_P(t, q)}{D_P(t, q)} \quad \text{and} \quad \bar{E}_P(t, q) = \frac{\bar{N}_P(t, q)}{D_P(t, q)},$$

over the same denominator of the form $D(t, q) = \prod_{i=1}^\nu (1 - q^{a_i} t^{b_i})$. Furthermore, there exists such an expression with all of these properties:

- (i) the number of denominator factors ν lies in the range $\dim(P) + 1 \leq \nu \leq V$,
- (ii) the numerators $N_P(t, q), \bar{N}_P(t, q)$ lie in $\mathbb{Z}[t, q]$,
- (iii) $N_P(t, q), \bar{N}_P(t, q)$, have t -degrees at most $b-1$ and exactly b , respectively, where $b := \sum_{i=1}^\nu b_i$,
- (iv) the denominator factors $1 - q^{a_i} t^{b_i}$ all have $b_i \leq \#(P \cap \mathbb{Z}^n) - 1$,
- (v) for P a simplex, $N_P(t, q), \bar{N}_P(t, q)$ have nonnegative coefficients as polynomials in t, q .

And lastly one has a q -analogue of Ehrhart-Macdonald reciprocity:

- (vi) the two series $E_P(t, q), \bar{E}_P(t, q)$ determine each other via

$$q^d \cdot \bar{E}_P(t, q) = (-1)^{d+1} E_P(t^{-1}, q^{-1}).$$

Example. Consider a 1-dimensional lattice polytope $P \subset \mathbb{R}^1$ of volume $v \geq 1$, that is, a line segment $P = [a, a + v]$ for some integer a . Thus for $m \geq 0$, one has $mP = [ma, m(a + v)]$ and

$$\mathbb{Z}^1 \cap mP = \{ma, ma + 1, ma + 2, \dots, m(a + v)\}.$$

Therefore one can calculate

$$\begin{aligned}
\mathbf{I}(\mathbb{Z}^1 \cap mP) &= (x - ma)(x - ma - 1) \cdots (x - ma - mv) \subset S = \mathbb{R}[x], \\
\text{gr } \mathbf{I}(\mathbb{Z}^1 \cap mP) &= (x^{mv+1}), \text{ so that } R(\mathbb{Z}^1 \cap mP) = \mathbb{R}[x]/(x^{mv+1}), \\
\text{Hilb}(R(\mathbb{Z}^1 \cap mP); q) &= 1 + q + q^2 + \cdots + q^{mv} = (1 - q^{mv+1})/(1 - q), \\
E_P(t, q) &= \sum_{m \geq 0} t^m \cdot (1 - q^{mv+1})/(1 - q) = \frac{1}{1 - q} \sum_{m \geq 0} t^m (1 - q^{mv+1}) \\
(10) \quad &= \frac{1}{1 - q} \left(\frac{1}{1 - t} - \frac{q}{1 - tq^v} \right) = \frac{1 + t(q + q^2 + \cdots + q^{v-1})}{(1 - t)(1 - tq^v)} = \frac{1 + t \cdot q[v - 1]_q}{(1 - t)(1 - tq^v)},
\end{aligned}$$

where we are using here a standard q -analogue of the positive integer m

$$[m]_q := 1 + q + q^2 + \cdots + q^{m-1}.$$

Note that $i(P, m) = mv + 1$, and hence, as expected, one has

$$E_P(t) = \sum_{m=0}^{\infty} (mv + 1)t^m = \frac{vt}{(1-t)^2} + \frac{1}{1-t} = \frac{1 + (v-1)t}{(1-t)^2} = [E_P(t, q)]_{q=1}.$$

Since mP has interior lattice points

$$\mathbb{Z}^1 \cap \text{int}(mP) = \{ma + 1, ma + 2, \dots, m(a + v) - 1\},$$

a similar calculation shows that

$$\begin{aligned}
\bar{E}_P(t, q) &= \sum_{m \geq 1} t^m \cdot (1 - q^{mv-1})/(1 - q) \\
(11) \quad &= \frac{1}{1 - q} \left(\frac{t}{1 - t} - \frac{q^{v-1}t}{1 - tq^v} \right) = \frac{t(1 + q + q^2 \cdots + q^{v-2}) + t^2 q^{v-1}}{(1 - t)(1 - tq^v)} = \frac{t[v - 1]_q + t^2 q^{v-1}}{(1 - t)(1 - tq^v)},
\end{aligned}$$

with $[\bar{E}_P(t, q)]_{q=1} = \frac{(v-1)t + t^2}{(1-t)^2} = \bar{E}_P(t)$, as expected. Furthermore, the line segment P is a lattice simplex, and one can check that (10), (11) are consistent with all parts (i)-(vi) of Conjecture 1.1.

1.2. The affine semigroup ring and the harmonic algebra. For lattice d -polytopes $P \subset \mathbb{R}^n$, work of Stanley (see, e.g., [45]) explains the Classical Ehrhart Theorem by reinterpreting $E_P(t)$ as the *Hilbert series* of a certain commutative graded algebra associated to P , reviewed here.

One embeds P in \mathbb{R}^{n+1} as $\{1\} \times P \subseteq \mathbb{R}^1 \times \mathbb{R}^n = \mathbb{R}^{n+1}$. The *cone* over P is

$$\text{cone}(P) := \mathbb{R}_{\geq 0} \cdot (\{1\} \times P).$$

It is an affine polyhedral $(d + 1)$ -dimensional cone in \mathbb{R}^{n+1} , and its lattice points $\mathbb{Z}^{n+1} \cap \text{cone}(P)$ form a semigroup under addition. The *affine semigroup ring* of P over a field \mathbf{k} is

$$(12) \quad A_P := \mathbf{k}[\mathbb{Z}^{n+1} \cap \text{cone}(P)].$$

The first coordinate x_0 on $\mathbb{R}^1 \times \mathbb{R}^n = \mathbb{R}^{n+1}$ endows A_P with the structure of a graded algebra, having Krull dimension $d + 1$. Inside A_P , one has the (homogeneous) *interior ideal* \bar{A}_P , the \mathbf{k} -span of all monomials corresponding to the interior lattice points $\text{int}(\text{cone}(P)) \cap \mathbb{Z}^{n+1}$. Then one has

$$\begin{aligned}
E_P(t) &= \text{Hilb}(A_P, t), \\
\bar{E}_P(t) &= \text{Hilb}(\bar{A}_P, t).
\end{aligned}$$

Although A_P is not always generated in degree one, there is always a tower of integral ring extensions

$$\mathbf{k}[\Theta] := \mathbf{k}[\theta_1, \dots, \theta_{d+1}] \subset B_P \subset A_P$$

where B_P is the subalgebra of A_P generated by the monomials corresponding to the vertices of P , and $\Theta := (\theta_1, \dots, \theta_{d+1})$ is any choice of a linear (degree one) system of parameters for B_P ; their existence is guaranteed by Noether's Normalization Lemma [35]. Most of the assertions (i),(ii),(iii),(iv) from the Classical Ehrhart Theorem are then explained by these four facts, respectively:

- (i) A_P is a finitely-generated $\mathbf{k}[\Theta]$ -module, and hence so is the ideal \overline{A}_P .
- (ii) A_P is even a *free* $\mathbf{k}[\Theta]$ -module, as Hochster [26] showed A_P is a *Cohen-Macaulay* ring.
- (iii) When P is a simplex, so that $\mathbf{k}[\Theta] = B_P$, then A_P has a $\mathbf{k}[\Theta]$ -basis given by monomials corresponding to the lattice points in the semi-open parallelepiped (3).
- (iv) Work of Danilov [11] and Stanley [44] independently showed that \overline{A}_P is isomorphic to the *canonical module* Ω_{A_P} of the Cohen-Macaulay ring A_P .


This might inspire one to approach Conjecture 1.1 by introducing a bigraded deformation of A_P . Since $A_P = \bigoplus_{m \geq 0} (A_P)_m$ where $(A_P)_m$ comes from the $x_0 = m$ slice of $\text{cone}(P)$

$$(13) \quad \text{cone}(P) \cap (\{m\} \times \mathbb{R}^n) = \{m\} \times mP,$$

one might look for such a bigraded object by defining a graded multiplication on the direct sum

$$(14) \quad \bigoplus_{m \geq 0} R(mP \cap \mathbb{Z}^n).$$

That is, one would like a way to ‘multiply’ elements $f \in R(mP \cap \mathbb{Z}^n)$ and $f' \in R(m'P \cap \mathbb{Z}^n)$ to produce a new element $f \star f' \in R((m+m')P \cap \mathbb{Z}^n)$ in a fashion which respects the polynomial degrees of f and f' . Geometrically, this corresponds to deforming the lattice points in $mP \times \{m\} \subseteq \text{cone}(P)$ linearly onto the coning axis and defining a multiplication between fat point loci. The relevant picture is shown below when $P = [-1, 1] \times [-1, 1]$ is a square in \mathbb{R}^2 .



The naïve approach of looking at the ring $R(\text{cone}(P) \cap \mathbb{Z}^{n+1})$ has two problems:

- the transformation $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ deforms a locus $\mathcal{Z} \subseteq \mathbf{k}^{n+1}$ to the origin, but we want to deform to the coning axis, and
- if P is full-dimensional, then $\text{cone}(P) \cap \mathbb{Z}^{n+1}$ is Zariski-dense and $\mathbf{I}(\text{cone}(P) \cap \mathbb{Z}^{n+1}) = 0$. Consequently, the deformation $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ for $\mathcal{Z} = P \cap \mathbb{Z}^{n+1}$ remembers only the affine span of P , not P itself.

We achieve the task of defining a multiplication on the direct sum (14) in Section 5 by replacing each summand $R(mP \cap \mathbb{Z}^n)$ with its *harmonic space* (or *Macaulay inverse system*). The resulting bigraded ring \mathcal{H}_P is the *harmonic algebra* attached to P . The bigraded Hilbert series of \mathcal{H}_P equals the q -Ehrhart series $E_P(t, q)$. Although not isomorphic to A_P in general, we conjecture that \mathcal{H}_P enjoys analogous algebraic properties (finite generation, Cohen-Macaulayness, identification of the canonical module) that would explain much of Conjecture 1.1.

In spite of all of these properties remaining conjectural, we are able to show that the q -Ehrhart series $E_P(t, q)$ and harmonic algebra \mathcal{H}_P behave in a predictable fashion when performing three well-studied operations on lattice polytopes P :

- *dilation* by an positive integer factor d , sending P to dP ,
- *Cartesian product*, sending $P \subset \mathbb{R}^n$ and $Q \subset \mathbb{R}^m$ to $P \times Q \subset \mathbb{R}^{n+m}$,
- *free join*, sending P, Q to $P * Q \subseteq \mathbb{R}^{1+n+m}$ defined by

$$(15) \quad P * Q := \{(t, t\mathbf{p}, (1-t)\mathbf{q}) : 0 \leq t \leq 1, \mathbf{p} \in P, \mathbf{q} \in Q\}.$$

The next result is proven in Section 6, and relates the above operations to the constructions of *Veronese subalgebras*, *Segre products* and *graded tensor products* for harmonic algebras $\mathcal{H}_P, \mathcal{H}_Q$.

Theorem 1.2. *Let P, Q be lattice polytopes.*

- (i) *For positive integers d , the dilation dP has $E_{dP}(t, q)$ given by*

$$E_{dP}(t, q) = \sum_{m=0}^{\infty} i_P(dm; q)t^m.$$

- (ii) *The Cartesian product $P \times Q$ has $E_{P \times Q}(t, q)$ given by the Hadamard product of series*

$$E_{P \times Q}(t, q) = \sum_{m=0}^{\infty} i_P(m; q) \cdot i_Q(m; q) \cdot t^m.$$

- (iii) *The free join $P * Q$ has $E_{P * Q}(t, q)$ given by*

$$E_{P * Q}(t, q) = \frac{1-t}{1-qt} \cdot E_P(t, q) \cdot E_Q(t, q).$$

1.3. Harmonic spaces and Minkowski addition. The fact that the harmonic algebra \mathcal{H}_P is closed under multiplication is not obvious, and rests upon a surprising new property of harmonic spaces for arbitrary finite loci $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$. Define their *Minkowski sum* to be the finite point locus

$$(16) \quad \mathcal{Z} + \mathcal{Z}' := \{\mathbf{z} + \mathbf{z}' : \mathbf{z} \in \mathcal{Z}, \mathbf{z}' \in \mathcal{Z}'\}.$$

The point-orbit rings $R(\mathcal{Z}), R(\mathcal{Z}')$, and $R(\mathcal{Z} + \mathcal{Z}')$ are graded quotients of $S = \mathbf{k}[x_1, \dots, x_n]$. When \mathbf{k} has characteristic zero, the partial differentiation action of S on itself gives rise to a \mathbf{k} -linear perfect pairing $\langle -, - \rangle$ on each homogeneous component S_d of S , and we may replace $R(\mathcal{Z}), R(\mathcal{Z}'), R(\mathcal{Z} + \mathcal{Z}')$ by their *harmonic spaces*

$$(17) \quad V_{\mathcal{Z}} := \text{gr } \mathbf{I}(\mathcal{Z})^{\perp}, \quad V_{\mathcal{Z}'} := \text{gr } \mathbf{I}(\mathcal{Z}')^{\perp}, \quad V_{\mathcal{Z} + \mathcal{Z}'} := \text{gr } \mathbf{I}(\mathcal{Z} + \mathcal{Z}')^{\perp}.$$

If $I \subseteq S$ is a homogeneous ideal, the harmonic space $I^{\perp} \subseteq S$ is a graded subspace with the same Hilbert series as S/I . Since elements of I^{\perp} are honest polynomials f , whereas elements of S/I are cosets $f + I$, working in I^{\perp} avoids coset-related issues which arise in proving, e.g., linear independence results. On the other hand, unlike the graded ring S/I , the subspace I^{\perp} has the defect of not being closed under multiplication. Nevertheless, in Section 4 we prove the following.

Theorem 1.3. *For any pair of finite point loci $\mathcal{Z}, \mathcal{Z}'$ in \mathbf{k}^n over any field \mathbf{k} , one has*

$$V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z} + \mathcal{Z}'}$$

This containment $V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z} + \mathcal{Z}'}$ may be interpreted as saying that the rings $R(\mathcal{Z})$ ‘remember’ the structure of Minkowski sums via multiplication of their harmonic spaces. As the deformation $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ does not respect ring structure, we find Theorem 1.3 a bit unexpected².

Note Theorem 1.3 is stated for finite point loci inside \mathbf{k}^n where \mathbf{k} is *any* field, not just $\mathbf{k} = \mathbb{R}$ or a field of characteristic zero. This requires defining harmonic spaces $V_{\mathcal{Z}}$ over an arbitrary field \mathbf{k} , which occurs already in the theory of *Macaulay’s inverse systems* over all fields \mathbf{k} discussed, e.g., in Geramita [20], and reviewed in Section 4 below. These more general harmonic spaces $V_{\mathcal{Z}}$ are defined not inside a polynomial ring over \mathbf{k} , but rather in the *divided power algebra* over \mathbf{k} . When \mathbf{k} has characteristic zero, these two rings are the same, and the definitions of $V_{\mathcal{Z}}$ coincide.

The remainder of the paper is structured as follows.

Section 2 reviews commutative algebra of associated graded ideals and rings, Gröbner bases, harmonic spaces and Macaulay’s inverse systems (first in characteristic zero, and then over all fields). It then briefly reviews some aspects of groups acting on rings and representation theory.

Section 3 defines the q -Ehrhart series $E_P(t, q)$, reviews Conjecture 1.1, and then examines several families of examples. It also incorporates symmetries of P in an equivariant q -Ehrhart series $E_P^G(t, q)$, and computes some highly symmetric examples, such as simplices and cross-polytopes.

Section 4 develops the proof of Theorem 1.3 on Minkowski sums, working over arbitrary fields.

Section 5 then uses Theorem 1.3 to define the harmonic algebra \mathcal{H}_P , and states Conjecture 5.5, explaining its connection to Conjecture 1.1.

Section 6 examines the behavior of $E_P(t, q)$ and \mathcal{H}_P under the lattice polytope operations of dilation, Cartesian product, and joins. In particular, it proves Theorem 1.2.

ACKNOWLEDGEMENTS

The authors thank Christian Haase for pointing them to the work of Balletti [3]. Authors partially supported by NSF grants DMS-2246846 and DMS-1745638, respectively.

2. BACKGROUND

2.1. Commutative algebra. Let \mathbf{k} be a field, let $n \geq 0$, and let $\mathbf{x} = (x_1, \dots, x_n)$ be a list of n variables. We write $S := \mathbf{k}[\mathbf{x}] = \mathbf{k}[x_1, \dots, x_n]$ for the polynomial ring in x_1, \dots, x_n over \mathbf{k} with its standard grading induced by $\deg(x_i) = 1$ for all i .

We will consider various graded \mathbf{k} -subspaces and quotients of S , as well as other rings. If $V = \bigoplus_{i \geq 0} V_i$ is a graded \mathbf{k} -vector space with each piece V_i finite-dimensional and q is variable, the *Hilbert series* of V is the formal power series

$$(18) \quad \text{Hilb}(V, q) := \sum_{i \geq 0} \dim_{\mathbf{k}}(V_i) \cdot q^i.$$

More generally, if $V = \bigoplus_{i, j \geq 0} V_{i, j}$ is a bigraded vector space, the *bigraded Hilbert series* is

$$(19) \quad \text{Hilb}(V, t, q) := \sum_{i, j \geq 0} \dim_{\mathbf{k}}(V_{i, j}) \cdot t^i q^j.$$

²It is reminiscent of another unexpected fact, about standard monomials for $\mathbf{I}(\mathcal{Z}), \mathbf{I}(\mathcal{Z}'), \mathbf{I}(\mathcal{Z} + \mathcal{Z}')$ with respect to a chosen monomial ordering on $S = \mathbb{R}[\mathbf{x}]$, observed by F. Gundlach [23]; see Remark 2.8 below.

Given $f \in S$ a nonzero polynomial, write $\tau(f) \in S$ for the top degree homogeneous component of f . That is, if $f = f_d + \cdots + f_1 + f_0$ with f_i homogeneous of degree i and $f_d \neq 0$, we have $\tau(f) = f_d$. If $I \subseteq S$ is an ideal, the *associated graded ideal* $\text{gr } I \subseteq S$ is given by

$$(20) \quad \text{gr } I := (\tau(f) : f \in I \setminus \{0\}) \subseteq S.$$

The ideal $\text{gr } I \subseteq S$ is homogeneous by construction, so that $S/\text{gr } I$ is a graded ring. In fact, we wish to explain why it is isomorphic to the *associated graded ring*

$$(21) \quad \text{gr}_{\mathcal{F}}(S/I) = \bigoplus_{d=0}^{\infty} F_d/F_{d-1} = F_0 \oplus F_1/F_0 \oplus F_2/F_1 \oplus \cdots$$

for the ascending filtration $\mathcal{F} = \{F_d\}_{d=0,1,2,\dots}$ on S/I

$$(22) \quad (\mathbf{k} =) F_0 \subseteq F_1 \subseteq F_2 \subseteq \cdots \subseteq S/I$$

where F_d is the image of the polynomials $S_{\leq d} := S_0 \oplus S_1 \oplus \cdots \oplus S_d$ of degree at most d under the surjection $S \twoheadrightarrow S/I$. Note that this filtration satisfies $F_i \cdot F_j \subseteq F_{i+j}$, so that the graded multiplication $F_i/F_{i-1} \times F_j/F_{j-1} \rightarrow F_{i+j}/F_{i+j-1}$ in $\text{gr}_{\mathcal{F}}(S/I)$ is well-defined.

Proposition 2.1. *Define a \mathbf{k} -algebra map $\varphi : S = \mathbf{k}[\mathbf{x}] \rightarrow \text{gr}_{\mathcal{F}}(S/I)$ sending $x_i \mapsto \bar{x}_i$ in F_1/F_0 .*

(i) *The map φ is surjective, with kernel $\text{gr } I$, inducing an \mathbb{N} -graded \mathbf{k} -algebra isomorphism*

$$S/\text{gr } I \cong \text{gr}_{\mathcal{F}}(S/I).$$

(ii) *Consequently, any homogeneous polynomials $\{f_j\}_{j \in J}$ whose images $\{f_j + \text{gr } I\}_{j \in J}$ give a \mathbf{k} -basis of $S/\text{gr } I$ will also have their images $\{f_j + I\}_{j \in J}$ giving a \mathbf{k} -basis of S/I .*

(iii) *In particular, whenever S/I is Artinian, that is, $\dim_{\mathbf{k}} S/I$ is finite, one can view $\text{Hilb}(S/\text{gr } I, q)$ as a q -analogue of $\dim_{\mathbf{k}}(S/I)$ in this sense:*

$$[\text{Hilb}(S/\text{gr } I, q)]_{q=1} = \dim_{\mathbf{k}}(S/\text{gr } I) = \dim_{\mathbf{k}}(S/I).$$

Proof. For (i), the surjectivity of φ holds because S is generated by x_1, \dots, x_n . Hence one has $S_{\leq d} = \text{span}_{\mathbf{k}}\{x_{i_1} \cdots x_{i_\ell}\}_{\ell \leq d}$, and therefore $F_d = \text{span}_{\mathbf{k}}\{\bar{x}_{i_1} \cdots \bar{x}_{i_\ell}\}_{\ell \leq d}$.

To show $\ker(\varphi) \supseteq \text{gr } I$, we check $\tau(f) \in \ker(\varphi)$ for $f \in I$. If $f = \sum_{i=0}^d f_i$ and $f_d \neq 0$, then

$$\varphi(\tau(f)) = \overline{f_d(\mathbf{x})} \equiv - \sum_{i=0}^{d-1} \overline{f_i(\mathbf{x})} \pmod{I},$$

so that $\varphi(\tau(f)) = 0$ in F_d/F_{d-1} .

To prove $\ker(\varphi) \subseteq \text{gr } I$, it suffices to show every *homogeneous* f in $\ker(\varphi)$ lies in $\text{gr } I$. If $\deg(f) = d$, then $f \in \ker(\varphi)$ implies $\bar{f} \in F_{d-1} \pmod{I}$, say $f = f' + f''$ with $f' \in S_{\leq d-1}$, $f'' \in I$. But then $f'' = f - f'$ has $\tau(f'') = f$, so $f \in \text{gr } I$.

For (ii), it suffices to check that for each $m \geq 0$, the set $\{f_j + I : j \in J, \deg(f_j) \leq m\}$ is a \mathbf{k} -basis for $F_m = \text{im}(S_{\leq m} \hookrightarrow S \twoheadrightarrow S/I)$. However, this follows by induction on m , since our hypotheses imply that $\{f_j + F_{m-1} : j \in J, \deg(f_j) = m\}$ is a \mathbf{k} -basis for F_m/F_{m-1} .

Assertion (iii) then follows immediately from (ii). \square

We have another consequence in the case $\dim_{\mathbf{k}} S/I = d < \infty$: the quotient S/I will be determined by the intersection $I_{\leq d-1} := I \cap S_{\leq d-1}$ with the first summand in the \mathbf{k} -vector space direct sum decomposition

$$(23) \quad S = S_{\leq d-1} \oplus S_{\geq d} \quad \text{where } S_{\geq d} := \bigoplus_{m=d}^{\infty} S_m = (x_1, \dots, x_n)^d.$$

Lemma 2.2. *Assume the ideal $I \subseteq S = \mathbf{k}[\mathbf{x}]$ has $d := \dim_{\mathbf{k}} S/I$ finite.*

- (i) *The inclusion $S_{\leq m}/I_{\leq m} \hookrightarrow S/I$ is an isomorphism for all $m \geq d-1$.*
- (ii) *The graded ring $S/\text{gr } I$ vanishes in degrees $m \geq d$, that is, $(S/\text{gr } I)_{\geq d} = 0$.*

Proof. Recall that $S/\text{gr } I \cong \text{gr}_{\mathcal{F}}(S/I) = \bigoplus_{m=0}^{\infty} F_m/F_{m-1}$ for the filtration $\mathcal{F} = \{F_m\}_{m=0,1,2,\dots}$ on S/I from (22), in which F_m is the image of the composite $S_{\leq m} \hookrightarrow S \twoheadrightarrow S/I$. The composite has kernel $I_{\leq m} := I \cap S_{\leq m}$, so $S_{\leq m}/I_{\leq m} \cong F_m$, and hence

$$(24) \quad \dim_{\mathbf{k}} S_{\leq m}/I_{\leq m} = \dim_{\mathbf{k}} F_m = \dim_{\mathbf{k}}(S/\text{gr } I)_{\leq m}.$$

Note that since $S/\text{gr } I$ is a graded \mathbf{k} -algebra generated in degree one, its nonzero graded components form an initial segment of degrees. Since $\dim_{\mathbf{k}} S/\text{gr } I = \dim_{\mathbf{k}} S/I = d$, one concludes that $\dim_{\mathbf{k}}(S/\text{gr } I)_{\leq m} = d$ for all $m \geq d-1$, and consequently, $\dim_{\mathbf{k}}(S/\text{gr } I)_m = 0$ for all $m \geq d$, proving assertion (ii). For assertion (i), note (24) also shows that $\dim_{\mathbf{k}} S_{\leq m}/I_{\leq m} = d$ for all $m \geq d-1$, and therefore the inclusion $S_{\leq m}/I_{\leq m} \hookrightarrow S/I$ must be an isomorphism. \square

If $I \subseteq S$ is an ideal with generating set $I = (f_1, \dots, f_s)$, we have $(\tau(f_1), \dots, \tau(f_s)) \subseteq \text{gr } I$, but this containment is strict in general. A finite generating set for $\text{gr } I$ may be computed using graded term orderings and Gröbner theory as follows; see Cox, Little, O’Shea [10] for more background.

Definition 2.3. A total order \preceq on the monomials of S is a *term order* if $1 \preceq m$ for all monomials m , and whenever $m_1 \preceq m_2$ one also has $m_1 m_3 \preceq m_2 m_3$ for all monomials m_1, m_2, m_3 .

For \prec a term order and $f \in S \setminus \{0\}$, write $\text{in}_{\prec}(f)$ for the \prec -largest monomial appearing in f .

Example 2.4. The *lexicographical* term order is defined by $x_1^{a_1} \cdots x_n^{a_n} <_{\text{lex}} x_1^{b_1} \cdots x_n^{b_n}$ if there exists i such that $a_1 = b_1, \dots, a_{i-1} = b_{i-1}$, and $a_i < b_i$. The *graded lex* term order is defined by

$$x^a <_{\text{grlex}} x^b \text{ if } \deg x^a < \deg x^b \text{ or } (\deg x^a = \deg x^b \text{ and } x^a <_{\text{lex}} x^b)$$

where $a = (a_1, \dots, a_n), b = (b_1, \dots, b_n) \in \mathbb{Z}_{\geq 0}^n$.

A term order \preceq is *graded* if $m \prec m'$ whenever $\deg m < \deg m'$. Equivalently, \preceq is graded if and only if one has for all $f \in S \setminus \{0\}$ that

$$(25) \quad \text{in}_{\prec}(f) = \text{in}_{\prec}(\tau(f)).$$

Example 2.5. Lexicographic order $<_{\text{lex}}$ is *not* graded for $n \geq 2$, but $<_{\text{grlex}}$ is always graded.

Let $I \subseteq S$ be an ideal and let \prec be a term order. The *initial ideal* of I is the monomial ideal

$$(26) \quad \text{in}_{\prec}(I) := (\text{in}_{\prec}(f) : f \in I \setminus \{0\}) \subseteq S.$$

Definition 2.6. A finite subset $G \subseteq I$ is a *Gröbner basis* of I if $\text{in}_{\prec}(I) = (\text{in}_{\prec}(g) : g \in G)$. Equivalently, for every f in I there exists some g in G with $\text{in}_{\prec}(g)$ dividing $\text{in}_{\prec}(f)$.

One can show that a Gröbner basis G for I always generates I as an ideal. We also have the following useful \mathbf{k} -basis for S/I . Say that monomial m in S is a *standard monomial* of I (with respect to \prec) if $m \notin \text{in}_{\prec}(I)$. Equivalently, this means that $\text{in}_{\prec}(g)$ does not divide m for all $g \in G$, where G is a Gröbner basis of I with respect to \prec . Then the set

$$(27) \quad \{m + I : m \text{ a standard monomial of } I\}$$

is a \mathbf{k} -basis of the quotient ring S/I . It is uniquely determined by the term order \prec , and called the *standard monomial basis* of S/I . The following can then be proven easily using (25).

Proposition 2.7. *Fix a graded term order \prec on S . Then for any ideal $I \subset S$, a Gröbner basis G for I (with respect to \prec) gives rise to a Gröbner basis with respect to \prec*

$$\tau(G) := (\tau(g) : g \in G)$$

for the homogeneous ideal $\text{gr } I$. Consequently, $\tau(G)$ also generates $\text{gr } I$ as an ideal:

$$\text{gr } I = (\tau(g) : g \in G)$$

Furthermore, $I, \text{gr } I$ share the same set of standard monomials \mathcal{B} with respect to \prec , which descend to \mathbf{k} -bases $\{m + I\}_{m \in \mathcal{B}}$ and $\{m + \text{gr } I\}_{m \in \mathcal{B}}$ for S/I and $S/\text{gr } I$, respectively.

Remark 2.8. When $I = \mathbf{I}(\mathcal{Z})$ is the vanishing ideal in $S = \mathbf{k}[\mathbf{x}]$ for a finite point set $\mathcal{Z} \subseteq \mathbf{k}^n$, Gundlach [23, Lem. 4] gives a very interesting alternate characterization of the \prec -standard monomials for $\mathbf{I}(\mathcal{Z})$. Let $\mathbf{k}[\mathcal{Z}]$ denote the \mathbf{k} -vector space of all functions $f : \mathcal{Z} \rightarrow \mathbf{k}$, with pointwise addition and \mathbf{k} -scaling. Endow $\mathbf{k}^{\mathcal{Z}}$ with a nondegenerate \mathbf{k} -bilinear form $(-, -) : \mathbf{k}[\mathcal{Z}] \times \mathbf{k}[\mathcal{Z}] \rightarrow \mathbf{k}$ given by

$$(f_1, f_2) := \sum_{\mathbf{z} \in \mathcal{Z}} f_1(\mathbf{z})f_2(\mathbf{z}).$$

Let U^\perp denote perp with respect to $(-, -)$ for \mathbf{k} -subspaces $U \subseteq \mathbf{k}^{\mathcal{Z}}$. Nondegeneracy of $(-, -)$ implies $U_1 \subsetneq U_2 \Leftrightarrow U_1^\perp \supsetneq U_2^\perp$. By multivariate Lagrange interpolation, the map $S \rightarrow \mathbf{k}[\mathcal{Z}]$ restricting polynomials $f(\mathbf{x})$ to functions \bar{f} on \mathcal{Z} is surjective. Since its kernel is $\mathbf{I}(\mathcal{Z})$, it gives a \mathbf{k} -vector space isomorphism $S/\mathbf{I}(\mathcal{Z}) \rightarrow \mathbf{k}[\mathcal{Z}]$. This implies, that for any $f \in \mathbf{k}[\mathcal{Z}] \setminus \{0\}$, there must exist some monomials m in S for which $(f, \bar{m}) \neq 0$. Consequently, *having fixed the monomial order \prec* , for each $f \in \mathbf{k}[\mathcal{Z}] \setminus \{0\}$, there will be a \prec -smallest such monomial associated to f , since \prec is a well-ordering:

$$\text{sm}_\prec(f) := \min_{\prec} \{ \text{monomials } m \in S : (f, \bar{m}) \neq 0 \in \mathbf{k} \}.$$

Proposition 2.9. [23, Lem.4] *For any finite subset $\mathcal{Z} \subset \mathbf{k}^n$, and for any choice of monomial order \prec on $S = \mathbf{k}[\mathbf{x}]$, one has this equality of sets:*

$$\{ \prec\text{-standard monomials for } \mathbf{I}(\mathcal{Z}) \} = \{ \text{sm}_\prec(f) : f \in \mathbf{k}[\mathcal{Z}] \setminus \{0\} \}.$$

Proof. For any monomial m in S one has the following:

$$\begin{aligned} m \text{ is } \prec\text{-standard for } \mathbf{I}(\mathcal{Z}) &\Leftrightarrow \nexists g \in \mathbf{I}(\mathcal{Z}) \text{ with } \text{in}_\prec(g) \text{ dividing } m \\ &\Leftrightarrow \nexists \hat{g} \in \mathbf{I}(\mathcal{Z}) \text{ with } \text{in}_\prec(\hat{g}) = m \\ &\Leftrightarrow \nexists \hat{g} \in \mathbf{I}(\mathcal{Z}) \text{ of the form } \hat{g} = m + \sum_{m' : m' \prec m} c_{m'} m' \text{ with } c_{m'} \in \mathbf{k} \\ &\Leftrightarrow \bar{m} \notin \text{span}_{\mathbf{k}} \{ \bar{m}' : m' \prec m \} \text{ inside } S/\mathbf{I}(\mathcal{Z}) = \mathbf{k}[\mathcal{Z}]. \\ &\Leftrightarrow \text{span}_{\mathbf{k}} \{ \bar{m}' : m' \prec m \} \subsetneq \text{span}_{\mathbf{k}} (\{ \bar{m} \} \cup \{ \bar{m}' : m' \prec m \}) \\ &\Leftrightarrow \text{span}_{\mathbf{k}} \{ \bar{m}' : m' \prec m \}^\perp \supsetneq \text{span}_{\mathbf{k}} (\{ \bar{m} \} \cup \{ \bar{m}' : m' \prec m \})^\perp \\ &\Leftrightarrow \exists f \in \text{span}_{\mathbf{k}} \{ \bar{m}' : m' \prec m \}^\perp \setminus \text{span}_{\mathbf{k}} (\{ \bar{m} \} \cup \{ \bar{m}' : m' \prec m \})^\perp \\ &\Leftrightarrow \exists f \in \mathbf{k}[\mathcal{Z}] \setminus \{0\} \text{ with } \text{sm}(f) = m. \quad \square \end{aligned}$$

2.2. Homogeneous harmonics in characteristic zero. A good reference for much of this material is Geramita [20, §2]. Let \mathbf{k} be a field of characteristic zero. We wish to set up two polynomial rings over \mathbf{k} , one of which acts on the other by partial derivatives. Let \mathbf{k}^n and its \mathbf{k} -dual $(\mathbf{k}^n)^*$ have dual ordered \mathbf{k} -bases (y_1, \dots, y_n) and (x_1, \dots, x_n) with respect to the usual \mathbf{k} -bilinear pairing of functionals and vectors

$$(28) \quad \langle -, - \rangle : (\mathbf{k}^n)^* \times \mathbf{k}^n \rightarrow \mathbf{k}$$

so that $\langle x_i, y_j \rangle = \delta_{ij}$. If one considers the polynomial algebras

$$\begin{aligned} S &:= \mathbf{k}[\mathbf{x}] = \mathbf{k}[x_1, \dots, x_n], \\ \mathbb{D} &:= \mathbf{k}[\mathbf{y}] = \mathbf{k}[y_1, \dots, y_n] \end{aligned}$$

then one can extend this to a \mathbb{D} -valued pairing

$$(29) \quad \odot : S \times \mathbb{D} \rightarrow \mathbb{D}$$

by requiring that each x_i act on \mathbb{D} as a derivation. That is, x_i acts as $\frac{\partial}{\partial y_i} : \mathbb{D} \rightarrow \mathbb{D}$, and for polynomials $f(\mathbf{x}), g(\mathbf{y})$, one has

$$f(\mathbf{x}) \odot g(\mathbf{y}) := f\left(\frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_n}\right) g(\mathbf{y}).$$

In this way, one obtains an S -module structure on \mathbb{D} . This S -module structure on \mathbb{D} is degree-lowering for the usual gradings on S, \mathbb{D} in which $\deg(x_i) = \deg(y_j) = 1$, in the sense that it restricts to a map $\odot : S_m \times \mathbb{D}_{m'} \rightarrow \mathbb{D}_{m'-m}$. This lets one extend the pairing $\langle -, - \rangle$ from (28) to a \mathbf{k} -bilinear pairing $\langle -, - \rangle : S \times \mathbb{D} \rightarrow \mathbf{k}$ defined by

$$(30) \quad \langle f(\mathbf{x}), g(\mathbf{y}) \rangle := \text{the constant term of } f \odot g$$

Employing an exponential notation for monomials $\mathbf{x}^{\mathbf{a}} := x_1^{a_1} \cdots x_n^{a_n}$ in S , where $\mathbf{a} = (a_1, \dots, a_n)$ lies in $\{0, 1, 2, \dots\}^n$, and similarly for monomials $\mathbf{y}^{\mathbf{b}}$ in \mathbb{D} , one can check that

$$\mathbf{x}^{\mathbf{a}} \odot \mathbf{y}^{\mathbf{b}} = \begin{cases} \prod_{i=1}^n \frac{b_i!}{(b_i - a_i)!} \cdot \mathbf{y}^{\mathbf{b} - \mathbf{a}} & \text{if } a_i \leq b_i \text{ for } i = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

One sees that $\langle -, - \rangle$ pairs orthonormally the \mathbf{k} -dual bases $\{\mathbf{x}^{\mathbf{a}}\}$ and $\left\{\frac{\mathbf{y}^{\mathbf{a}}}{a_1! \cdots a_n!}\right\}$, and hence restricts to a perfect \mathbf{k} -linear pairing on the (finite-dimensional!) spaces

$$\begin{aligned} \langle -, - \rangle : S_m \times \mathbb{D}_m &\longrightarrow \mathbf{k} \\ \langle -, - \rangle : S_{\leq m} \times \mathbb{D}_{\leq m} &\longrightarrow \mathbf{k}. \end{aligned}$$

Note that S_m and $\mathbb{D}_{m'}$ are perpendicular with respect to the pairing $\langle -, - \rangle$ whenever $m \neq m'$.

Definition 2.10. For any homogeneous ideal $I \subseteq S = \mathbf{k}[\mathbf{x}]$, the *harmonic space* (or *Macaulay inverse system*) $I^\perp \subseteq \mathbb{D} = \mathbf{k}[\mathbf{y}]$ is the graded vector space

$$(31) \quad I^\perp := \{g(\mathbf{y}) \in \mathbb{D} : \langle f(\mathbf{x}), g(\mathbf{y}) \rangle = 0 \text{ for all } f \in I\}$$

$$(32) \quad = \{g(\mathbf{y}) \in \mathbb{D} : f(\mathbf{x}) \odot g(\mathbf{y}) = 0 \text{ for all } f \in I\}.$$

The equality of the two sets on the right in (31), (32) is justified as follows. If $f \odot g = 0$ then $\langle f, g \rangle = 0$, showing the set from (32) is contained in the set from (31). For the reverse inclusion, note that if $f(\mathbf{x}) \odot g(\mathbf{y}) \neq 0$ for some $f(\mathbf{x})$ in I , say $f(\mathbf{x}) \odot g(\mathbf{y}) = \sum_{\mathbf{a}} c_{\mathbf{a}} \mathbf{y}^{\mathbf{a}}$ with some $c_{\mathbf{a}} \neq 0$, then the ideal I contains $\mathbf{x}^{\mathbf{a}} f(\mathbf{x})$ with $\langle \mathbf{x}^{\mathbf{a}} f(\mathbf{x}), g(\mathbf{y}) \rangle = c_{\mathbf{a}} \prod_i a_i! \neq 0$.

Note that in each degree m , the perfect pairing $\langle -, - \rangle : S_m \times \mathbb{D}_m \rightarrow \mathbf{k}$ gives a \mathbf{k} -vector space isomorphism $\mathbb{D}_m \rightarrow S_m^*$ sending $g(\mathbf{y}) \mapsto \langle -, g \rangle$. This induces a \mathbf{k} -vector space isomorphism

$$(33) \quad I_m^\perp \rightarrow (S_m/I_m)^*,$$

showing that $\dim_{\mathbf{k}} I^\perp = \dim_{\mathbf{k}} (S_m/I_m)^*$. Hence as graded \mathbf{k} -vector spaces one has

$$(34) \quad \text{Hilb}(I^\perp, q) = \text{Hilb}(S/I, q).$$

2.3. Homogeneous harmonics for all fields: divided powers. In order to define harmonic spaces over arbitrary fields, as needed in Theorem 1.3, we will need to replace $\mathbb{D} = \mathbf{k}[\mathbf{y}]$ with a *divided power algebra* over a field \mathbf{k} . We therefore review divided power algebras here; a reader who is content with seeing Theorem 1.3 stated and/or proven only in characteristic zero can mostly skip this section. Useful references for this material are Eisenbud [13, A2.4] and Geramita [20, §9].

Definition 2.11. Let \mathbf{k} be any field. For $n = 0, 1, 2, \dots$, let $\mathbf{y} = (y_1, \dots, y_n)$ be a list of n variables, thought of as the \mathbf{k} -basis for \mathbf{k}^n . Then the *divided power algebra* of rank n over \mathbf{k} is defined as a free R -module $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ with “monomial” \mathbf{k} -basis given by the symbols $\mathbf{y}^{(\mathbf{a})} := y_1^{(a_1)} \cdots y_n^{(a_n)}$ for $\mathbf{a} = (a_1, \dots, a_n) \in \{0, 1, 2, \dots\}^n$. One can then define a multiplication on $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ which is \mathbf{k} -bilinear and determined on monomials by the rule

$$(35) \quad \mathbf{y}^{(\mathbf{a})} \cdot \mathbf{y}^{(\mathbf{b})} = (y_1^{(a_1)} \cdots y_n^{(a_n)}) \cdot (y_1^{(b_1)} \cdots y_n^{(b_n)}) := \binom{a_1 + b_1}{a_1, b_1} \cdots \binom{a_n + b_n}{a_n, b_n} y_1^{(a_1 + b_1)} \cdots y_n^{(a_n + b_n)}$$

where the binomial coefficients $\binom{a_i + b_i}{a_i, b_i}$ are regarded as elements of \mathbf{k} in the natural way; some will vanish when \mathbf{k} has positive characteristic. This makes $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ an associative, commutative \mathbf{k} -algebra with unit $1 = y_1^{(0)} \cdots y_n^{(0)}$. It has a grading $\mathbb{D}_{\mathbf{k}}(\mathbf{y}) = \bigoplus_{m \geq 0} \mathbb{D}_{\mathbf{k}}(\mathbf{y})_m$ where $\mathbb{D}_{\mathbf{k}}(\mathbf{y})_m$ has \mathbf{k} -basis $\{\mathbf{y}^{(\mathbf{a})} : a_1 + \cdots + a_n = m\}$.

Roughly speaking, the symbol $y_i^{(d)} \in \mathbb{D}_{\mathbf{k}}(\mathbf{y})$ plays the role of “ $y_i^d/d!$ ”, even when $d! = 0$ in \mathbf{k} . When \mathbf{k} has characteristic zero, the \mathbf{k} -vector space isomorphism $\mathbb{D}_{\mathbf{k}}(\mathbf{y}) \cong \mathbf{k}[\mathbf{y}]$ given by

$$\begin{aligned} \mathbb{D}_{\mathbf{k}}(\mathbf{y}) &\longrightarrow \mathbf{k}[\mathbf{y}] \\ \mathbf{y}^{(\mathbf{a})} = y_1^{(a_1)} y_2^{(a_2)} \cdots y_n^{(a_n)} &\longmapsto \frac{y_1^{a_1} y_2^{a_2} \cdots y_n^{a_n}}{a_1! a_2! \cdots a_n!} \end{aligned}$$

is a ring isomorphism. In fact, with conventions $y_i^{(0)} := 1$ and $y_i^{(1)} := y_i$, the set map $y_i \mapsto y_i^{(d)}$ for $i = 1, 2, \dots, n$ extends to what is called a *system of divided powers* on $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$: a collection of maps for $d = 0, 1, 2, \dots$

$$\begin{aligned} \mathbb{D}_{\mathbf{k}}(\mathbf{y})_m &\longrightarrow \mathbb{D}_{\mathbf{k}}(\mathbf{y})_{dm} \\ a &\longmapsto a^{(d)} \end{aligned}$$

satisfying these axioms (modeled on properties of the maps $a \mapsto \frac{a^d}{d!}$ that exist whenever $\mathbf{k} \supseteq \mathbb{Q}$):

$$(36) \quad a^{(0)} = 1, a^{(1)} = a$$

$$(37) \quad a^{(d)} a^{(e)} = \binom{d+e}{d, e} a^{(d+e)}$$

$$(38) \quad \left(a^{(d)}\right)^{(e)} = \frac{1}{e!} \binom{de}{d, d, \dots, d} a^{(de)}$$

$$(39) \quad (ab)^{(d)} = d! \cdot a^{(d)}b^{(d)}$$

$$(40) \quad (a+b)^{(d)} = \sum_{\substack{(d_1, d_2): \\ d_1+d_2=d}} a^{(d_1)}b^{(d_2)}$$

For example, the reader might wish to check that iterating (37) implies $a^d = (a^{(1)})^d = d! \cdot a^{(d)}$, so that one has no choice but to define $a^{(d)} = a^d/d!$ whenever $d! \in \mathbf{k}^\times$. Similarly, if one iterates (40), which Eisenbud [13, A2.4] calls the “beginner’s binomial theorem”, one obtains the “beginner’s multinomial theorem”:

$$(41) \quad (a_1 + \cdots + a_m)^{(d)} = \sum_{\substack{(d_1, \dots, d_m): \\ d_1 + \cdots + d_m = d}} a_1^{(d_1)} \cdots a_m^{(d_m)}.$$

It is also not hard to check that one has a graded \mathbf{k} -algebra isomorphism

$$(42) \quad \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \cong \mathbb{D}_{\mathbf{k}}(y_1) \otimes_{\mathbf{k}} \cdots \otimes_{\mathbf{k}} \mathbb{D}_{\mathbf{k}}(y_n).$$

For the sake of defining harmonics and inverse systems, let $S := \mathbf{k}[\mathbf{x}] = \mathbf{k}[x_1, \dots, x_n]$ as before. The algebra $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ attains a unique S -module structure $\odot : S \times \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \rightarrow \mathbb{D}_{\mathbf{k}}(\mathbf{y})$ by having x_i act on $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ as a derivation, extending the rule

$$(43) \quad x_i \odot y_j^{(k)} = \begin{cases} y_i^{(k-1)} & \text{if } i = j \text{ and } k \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

This gives rise to a \mathbf{k} -bilinear pairing $\langle -, - \rangle : S \times \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \rightarrow R$ given by

$$(44) \quad \langle f, g \rangle := \text{the constant term of } f \odot g$$

under which

$$(45) \quad \langle \mathbf{x}^{\mathbf{a}}, \mathbf{y}^{(\mathbf{b})} \rangle = \langle x_1^{a_1} \cdots x_n^{a_n}, y_1^{(b_1)} \cdots y_n^{(b_n)} \rangle = \begin{cases} 1 & \text{if } a_i = b_i \text{ for all } i, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, the \mathbf{k} -bilinear pairing $\langle -, - \rangle : S \times \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \rightarrow \mathbf{k}$ generalizes the one from (83) when $\mathbf{k} \supseteq \mathbb{Q}$. It again leads to perfect \mathbf{k} -bilinear pairings on these finite-dimensional spaces:

$$\begin{aligned} \langle -, - \rangle : S_m \times \mathbb{D}_{\mathbf{k}}(\mathbf{y})_m &\longrightarrow \mathbf{k} \\ \langle -, - \rangle : S_{\leq m} \times \mathbb{D}_{\mathbf{k}}(\mathbf{y})_{\leq m} &\longrightarrow \mathbf{k}. \end{aligned}$$

This leads to the following generalization of Definition 2.10.

Definition 2.12. For \mathbf{k} any field and any homogeneous ideal $I \subseteq S = \mathbf{k}[\mathbf{x}]$, the *harmonic space* (or *Macaulay inverse system*) $I^\perp \subseteq \mathbb{D}_{\mathbf{k}}(\mathbf{y})$ is the graded \mathbf{k} -vector space

$$\begin{aligned} I^\perp &:= \{g(\mathbf{y}) \in \mathbb{D}_{\mathbf{k}}(\mathbf{y}) : \langle f(\mathbf{x}), g(\mathbf{y}) \rangle = 0 \text{ for all } f \in I\} \\ &= \{g(\mathbf{y}) \in \mathbb{D}_{\mathbf{k}}(\mathbf{y}) : f(\mathbf{x}) \odot g(\mathbf{y}) = 0 \text{ for all } f \in I\}. \end{aligned}$$

Note that, as before, the perfect pairing $\langle -, - \rangle : S_m \times \mathbb{D}_{\mathbf{k}}(\mathbf{y})_m \rightarrow \mathbf{k}$ gives a \mathbf{k} -linear isomorphism $\mathbb{D}_{\mathbf{k}}(\mathbf{y})_m \rightarrow S_m^*$ sending $g(\mathbf{y}) \mapsto \langle -, g \rangle$. As in (33), this induces a \mathbf{k} -linear isomorphism

$$(46) \quad I_m^\perp \rightarrow (S_m/I_m)^*,$$

showing $\dim_{\mathbf{k}} I^\perp = \dim_{\mathbf{k}} (S_m/I_m)^*$. Hence one has this generalization of (34):

$$(47) \quad \text{Hilb}(I^\perp, q) = \text{Hilb}(S/I, q).$$

2.4. Symmetry. The natural action of $GL_n(\mathbf{k})$ on \mathbf{k}^n and a given basis y_1, \dots, y_n induces a left-action on the polynomials $\mathbb{D} = \mathbf{k}[y_1, \dots, y_n]$ by linear substitutions. The contragredient action on $(\mathbf{k}^n)^*$ precomposes functionals with h^{-1} , that is, sending $h : f \mapsto f \circ h^{-1}$, thereby acting on x_1, \dots, x_n , as well as on the polynomials $S = \mathbf{k}[x_1, \dots, x_n]$. Explicitly, if h in $GL_n(\mathbf{k})$ acts in the basis y_1, \dots, y_n via the matrix A in $\mathbf{k}^{n \times n}$, then

$$\begin{aligned} h : g(\mathbf{y}) &\mapsto g(A\mathbf{y}), \\ h : f(\mathbf{x}) &\mapsto f((A^{-1})^t \mathbf{x}). \end{aligned}$$

Note that the pairing $\langle -, - \rangle : (\mathbf{k}^n)^* \times \mathbf{k}^n \rightarrow \mathbf{k}$ between functionals and vectors in (28) satisfies a certain *invariance* with respect to these $GL_n(\mathbf{k})$ -actions: for any linear functional f in $(\mathbf{k}^n)^*$, vector y in \mathbf{k}^n , and h in $GL_n(\mathbf{k})$, one has

$$\langle h(f), h(y) \rangle = (f \circ h^{-1})(h(y)) = f(y) = \langle f, y \rangle.$$

Consequently, the pairings \odot and $\langle -, - \rangle$ are similarly invariant with respect to the $GL_n(\mathbf{k})$ -action:

$$(48) \quad h(f(\mathbf{x})) \odot h(g(\mathbf{y})) = f(\mathbf{x}) \odot g(\mathbf{y}),$$

$$(49) \quad \langle h(f(\mathbf{x})), h(g(\mathbf{y})) \rangle = \langle f(\mathbf{x}), g(\mathbf{y}) \rangle.$$

Remark 2.13. For arbitrary fields \mathbf{k} when one replaces the polynomial algebra $\mathbb{D} = \mathbf{k}[\mathbf{y}]$ with the divided power algebra $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$, it is still true the $GL_n(\mathbf{k})$ -action on $\mathbf{k}^n = \mathbb{D}_{\mathbf{k}}(\mathbf{y})_1$ extends to an action via graded \mathbf{k} -algebra automorphisms on all of $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$. This fact is more apparent when one constructs multiplication in $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ as the (*graded*) *dual* of the coalgebra structure $\Delta : S \rightarrow S \otimes S$ on $S = \mathbf{k}[\mathbf{x}]$ in which $\Delta(x_i) = 1 \otimes x_i + x_i \otimes 1$, that is, each $x \in S_1 = (\mathbf{k}^n)^*$ is *primitive*. See Akin, Buchsbaum and Weyman [1, §I.4], Eisenbud [13, A2.4], Geramita [20, §9] for more on this alternate construction of $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$.

A consequence of the $GL_n(\mathbf{k})$ -invariance (49) is that the isomorphisms (33), (46) become $GL_n(\mathbf{k})$ -equivariant. This shows that for each m , one has isomorphisms $I_m^\perp \cong (S/I)^*$ as $GL_n(\mathbf{k})$ -representations. The same holds for the action of any subgroup $G \subseteq GL_n(\mathbf{k})$ on S, \mathbb{D} by restriction.

2.5. Representation theory. We will be interested in polytopes and point loci in \mathbb{R}^n with symmetry, and wish to keep track of the representations of their symmetry groups on the various \mathbf{k} -vector spaces that we construct. We review one way to do such bookkeeping, using the language of $\mathbf{k}[G]$ -modules and representation rings.

Definition 2.14. For a field \mathbf{k} and finite group G , define its *representation ring* $\text{Rep}_{\mathbf{k}}(G)$ as follows.

- As a \mathbb{Z} -module, $\text{Rep}_{\mathbf{k}}(G)$ is the quotient of free \mathbb{Z} -module with \mathbb{Z} -basis elements $[V]$ indexed by all isomorphism classes finite-dimensional $\mathbf{k}[G]$ -modules V , in which one mods out by the submodule \mathbb{Z} -spanned by all relations

$$(50) \quad [V \oplus V'] - ([V] + [V']).$$

- As a \mathbb{Z} -algebra, its multiplication is induced by the rule

$$[V] \cdot [V'] := [V \otimes V'].$$

- The operation $V \mapsto V^*$ of taking the contragredient $\mathbf{k}[G]$ -module leads to a \mathbb{Z} -automorphism and involution on $(-)^* : \text{Rep}_{\mathbf{k}}(G) \rightarrow \text{Rep}_{\mathbf{k}}(G)$

$$[V]^* := [V^*].$$

Whenever $\#G$ lies in \mathbf{k}^\times , Maschke's Theorem asserts that $\mathbf{k}[G]$ -modules are completely reducible, which shows that $\text{Rep}_{\mathbf{k}}(G)$ is a free \mathbb{Z} -module on the \mathbb{Z} -basis $[V_1], [V_2], \dots, [V_N]$ where V_1, \dots, V_N are the non-isomorphic *simple/irreducible* $\mathbf{k}[G]$ -modules.

For any field \mathbf{k} , the map sending $[V]$ to its *character* $\text{ch}(V) : G \rightarrow \mathbf{k}$ defined by

$$\text{ch}(V)(g) := \text{trace}(g : V \rightarrow V),$$

becomes an algebra map from $\text{Rep}_{\mathbf{k}}(G)$ into the *ring of class functions* $\{f : G \rightarrow \mathbf{k}\}$, that is, functions which are constant on G -conjugacy classes. The ring of class functions is given pointwise addition, multiplication, and the involution $f \mapsto f^*$ defined by $f^*(g) = f(g^{-1})$. Whenever \mathbf{k} has characteristic zero, this algebra map is injective, and in particular, two $\mathbf{k}[G]$ -modules V, V' are isomorphic (that is, $[V] = [V']$) if and only if they have the same character $\text{ch}(V) = \text{ch}(V')$.

More generally, for graded $\mathbf{k}[G]$ -modules $V = \bigoplus_{m \geq 0} V_m$, with each V_m a finite-dimensional $\mathbf{k}[G]$ -module, we will track the representation with a power series in $\text{Rep}_{\mathbf{k}}(G)[[q]]$:

$$(51) \quad [V]_q := \sum_{m \geq 0} [V_m] \cdot q^m.$$

In particular, when $I \subseteq S$ is a homogeneous ideal which is stable under the action of a finite subgroup G of $GL_n(\mathbf{k})$, both the quotient S/I and the harmonic space I^\perp inherit the structure of graded $\mathbf{k}[G]$ -modules. Then (46) implies a graded $\mathbf{k}[G]$ -module isomorphism $I^\perp \cong (S/I)^*$, and hence $[I^\perp]_q = [S/I]_q^*$ in $\text{Rep}_{\mathbf{k}}(G)[[q]]$.

On the other hand, we will also consider potentially *inhomogeneous* ideals $I \subseteq S$ that are stable under the action of a finite subgroup $G \subseteq GL_n(\mathbf{k})$, e.g., $I = \mathbf{I}(\mathcal{Z})$ where $\mathcal{Z} \subseteq \mathbf{k}^n$ is a G -stable locus.

Proposition 2.15. *Assume $\#G$ lies in \mathbf{k}^\times . For any ideal $I \subseteq S = \mathbf{k}[\mathbf{x}]$ which is stable under a finite subgroup G of $GL_n(\mathbf{k})$, if $\dim_{\mathbf{k}} S/I$ finite, then one has a $\mathbf{k}[G]$ -module isomorphism $S/I \cong S/\text{gr } I$.*

Proof. Recall (21) gave an isomorphism $S/\text{gr } I \cong \text{gr}_{\mathcal{F}}(S/I) = \bigoplus_{d=0}^M F^d/F^{d-1}$, where the sum on the right is finite here due to the finiteness assumption on $\dim_{\mathbf{k}} S/I$. This isomorphism is easily seen to be G -equivariant. Since the action of G preserves degree, the filtration \mathcal{F} of S/I is G -stable, and its filtration factors match the $\mathbf{k}[G]$ -module structures on the graded components of $S/\text{gr } I$. Complete reducibility of $\mathbf{k}[G]$ -modules then shows $S/I \cong S/\text{gr } I$. \square

This proof also shows, for any field \mathbf{k} , the $\mathbf{k}[G]$ -modules $S/\text{gr } I$ and S/I are Brauer-isomorphic.

3. THE q -EHRHART SERIES AND CONJECTURE 1.1

Throughout this section, we take $\mathbf{k} = \mathbb{R}$ and consider finite point loci $\mathcal{Z} \subset \mathbb{R}^n$. The locus \mathcal{Z} has two associated ideals inside $S = \mathbb{R}[\mathbf{x}] = \mathbb{R}[x_1, \dots, x_n]$: the (inhomogeneous) *vanishing ideal*

$$\mathbf{I}(\mathcal{Z}) = \{f(\mathbf{x}) \in S : f(\mathbf{z}) = 0 \text{ for all } \mathbf{z} \in \mathcal{Z}\}$$

and its *associated graded ideal*

$$\text{gr } \mathbf{I}(\mathcal{Z}) := (\tau(f) : f \in \mathbf{I}(\mathcal{Z})).$$

Within the polynomial ring $\mathbb{D} = \mathbb{R}[\mathbf{y}] = \mathbb{R}[y_1, \dots, y_n]$ in the dual variables \mathbf{y} , the *harmonic space* $(\text{gr } \mathbf{I}(\mathcal{Z}))^\perp$ of $\text{gr } \mathbf{I}(\mathcal{Z})$ will play a crucial role in our work. To reduce notational clutter, we write

$$(52) \quad V_{\mathcal{Z}} := (\text{gr } \mathbf{I}(\mathcal{Z}))^\perp$$

for this harmonic space. This notation emphasizes the role of $V_{\mathcal{Z}}$ as a graded subspace of \mathbb{D} which is almost never closed under multiplication. On the other hand, (34) shows that, as a graded

vector space, it has the same Hilbert series (actually a polynomial here) as the quotient ring $R(\mathcal{Z}) := S/\text{gr } \mathbf{I}(\mathcal{Z})$ that was defined in (5)

$$\text{Hilb}(V_{\mathcal{Z}}, q) = \text{Hilb}(R(\mathcal{Z}), q).$$

Note Proposition 2.1(iii) shows this Hilbert series is a q -analogue of the cardinality $\#\mathcal{Z}$, that is,

$$[\text{Hilb}(V_{\mathcal{Z}}, q)]_{q=1} = [\text{Hilb}(R(\mathcal{Z}), q)] = \dim_{\mathbf{k}}(S/\mathbf{I}(\mathcal{Z})) = \#\mathcal{Z}.$$

Furthermore, when G is a finite subgroup of $GL_n(\mathbb{R})$ that preserves \mathcal{Z} setwise, it acts via ring automorphisms and (graded) $\mathbb{R}[G]$ -modules on all of the objects under consideration:

$$S, \mathbb{D}, \mathbf{I}(\mathcal{Z}), \text{gr } \mathbf{I}(\mathcal{Z}), S/\mathbf{I}(\mathcal{Z}), R(\mathcal{Z}), V_{\mathcal{Z}}.$$

Because finite-dimensional $\mathbb{R}[G]$ -modules V are all self-contragredient ($V^* \cong V$), one can check that (33) implies graded $\mathbb{R}[G]$ -module isomorphisms

$$(53) \quad V_{\mathcal{Z}} \cong R(\mathcal{Z})$$

and then Proposition 2.15 implies a further ungraded $\mathbf{k}[G]$ -module isomorphisms

$$(54) \quad V_{\mathcal{Z}} \cong R(\mathcal{Z}) \cong S/\mathbf{I}(\mathcal{Z}),$$

which are all three isomorphic to the $\mathbb{R}[G]$ -permutation module on the points \mathcal{Z} .

3.1. Definition of q -Ehrhart series and the conjecture. As in the introduction, a *lattice polytope* $P \subset \mathbb{R}^n$ is the convex hull of a finite set of points in the lattice \mathbb{Z}^n . For each integer $m = 0, 1, 2, \dots$ one obtains a finite point locus $\mathbb{Z}^n \cap mP$. For each $m \geq 1$, one has the interior point locus $\mathbb{Z}^n \cap \text{int}(mP)$, where $\text{int}(P) = P \setminus \text{Bd}(P)$ is the (relative) interior where one removes the union $\text{Bd}(P)$ of all boundary faces of P .

Definition 3.1. For a lattice polytope $P \subset \mathbb{R}^n$, define two q -Ehrhart series in $\mathbb{Z}[q][[t]]$:

$$(55) \quad E_P(t, q) := \sum_{m=0}^{\infty} i_P(m; q) \cdot t^m,$$

where $i_P(m; q) := \text{Hilb}(V_{\mathbb{Z}^n \cap mP}, q) = \text{Hilb}(R(\mathbb{Z}^n \cap mP), q),$

$$(56) \quad \bar{E}_P(t, q) := \sum_{m=1}^{\infty} \bar{i}_P(m; q) \cdot t^m$$

where $\bar{i}_P(m; q) = \text{Hilb}(V_{\mathbb{Z}^n \cap \text{int}(mP)}, q) = \text{Hilb}(R(\mathbb{Z}^n \cap \text{int}(mP)), q).$

Note the series $E_P(t, q), \bar{E}_P(t, q)$ reduce to the classical Ehrhart series $E_P(t), \bar{E}_P(t)$ at $q \rightarrow 1$.

Example 3.2. Recall the Example in the Introduction looked at the general 1-dimensional lattice polytope $P = [a, a + v] \subset \mathbb{R}^1$ with $a, v \in \mathbb{Z}$ and volume $v \geq 1$, finding that

$$(57) \quad E_P(t, q) = \frac{1 + tq[v-1]_q}{(1-t)(1-tq^v)},$$

$$(58) \quad \bar{E}_P(t, q) = \frac{t[v-1]_q + t^2q^{v-1}}{(1-t)(1-tq^v)},$$

where $[m]_q := 1 + q + q^2 + \dots + q^{m-1}$.

Note that Example 3.2 shows that both $E_P(t, q), \bar{E}_P(t, q)$ for lattice polytopes $P \subset \mathbb{R}^1$ depend only upon a single parameter, which one could take either to be the volume v , or the number of lattice points $\#\mathbb{Z}^1 \cap P = v + 1$, or the h^* -vector entry $h_1^* = v - 1$. This illustrates a certain affine-lattice invariance that one might expect, similar to classical Ehrhart theory. Recall that the group of *affine transformations* $\text{Aff}(\mathbb{R}^n)$ of \mathbb{R}^n is a semidirect product $\text{Aff}(\mathbb{R}^n) = GL_n(\mathbb{R}) \ltimes \mathbb{R}^n$, where $GL_n(\mathbb{R})$ is the subgroup fixing the origin, and \mathbb{R}^n is the translation subgroup. This restricts to a semidirect product decomposition $\text{Aff}(\mathbb{Z}^n) = GL_n(\mathbb{Z}) \ltimes \mathbb{Z}^n$.

Proposition 3.3. *For any $g \in \text{Aff}(\mathbb{Z}^n)$, one has $E_{gP}(t, q) = E_P(t, q)$ and $\bar{E}_{gP}(t, q) = \bar{E}_P(t, q)$.*

Proof. We give the argument for $E_{gP}(t, q)$; the argument for $\bar{E}_{gP}(t, q)$ is similar.

Note that for each $m = 0, 1, 2, \dots$, the point locus $\mathbb{Z}^n \cap m \cdot gP$ is an affine transformation of the locus $\mathbb{Z}^n \cap mP$, namely by an affine transformation h whose translation vector $m \cdot v$ is scaled by m from the translation vector v of g . It therefore suffices to check for finite point loci $\mathcal{Z} \subseteq \mathbb{R}^n$ and for any $h \in \text{Aff}(\mathbb{R}^n)$, that one has

$$(59) \quad \text{Hilb}(R(h\mathcal{Z}), q) = \text{Hilb}(R(\mathcal{Z}), q).$$

Note that $h \in \text{Aff}(\mathbb{R}^n)$ acting on \mathbf{x} variables by $h\mathbf{x} = h_0\mathbf{x} + v$ for some $h_0 \in GL_n(\mathbb{R}), v \in \mathbb{R}^n$ has

$$\tau(h(f))(\mathbf{x}) = \tau(f(h_0\mathbf{x} + v)) = \tau(f)(h_0\mathbf{x}).$$

Therefore $\text{gr } \mathbf{I}(h\mathcal{Z}) = h_0(\text{gr } \mathbf{I}(\mathcal{Z}))$. Since $h_0 \in GL_n(\mathbb{R})$ acts via a graded \mathbb{R} -algebra automorphism on $S = \mathbb{R}[\mathbf{x}]$, this means it induces a graded \mathbb{R} -algebra isomorphism $R(\mathcal{Z}) \cong R(h\mathcal{Z})$ implying (59):

$$R(\mathcal{Z}) = S/\text{gr } \mathbf{I}(\mathcal{Z}) \xrightarrow{h_0} S/h_0(\text{gr } \mathbf{I}(\mathcal{Z})) = S/\text{gr } \mathbf{I}(h\mathcal{Z}) = R(h\mathcal{Z}). \quad \square$$

Before discussing more examples, recall the main conjecture from the Introduction.

Conjecture 1.1. *Let P be a d -dimensional lattice polytope in \mathbb{R}^n , having V vertices. Then both of the series (8),(9) lie in $\mathbb{Q}(t, q)$, and are expressible as rational functions*

$$E_P(t, q) = \frac{N_P(t, q)}{D_P(t, q)} \quad \text{and} \quad \bar{E}_P(t, q) = \frac{\bar{N}_P(t, q)}{D_P(t, q)},$$

over the same denominator of the form $D(t, q) = \prod_{i=1}^{\nu} (1 - q^{a_i} t^{b_i})$. Furthermore, there exists such an expression with all of these properties:

- (i) the number of denominator factors ν lies in the range $\dim(P) + 1 \leq \nu \leq V$,
- (ii) the numerators $N_P(t, q), \bar{N}_P(t, q)$ lie in $\mathbb{Z}[t, q]$,
- (iii) $N_P(t, q), \bar{N}_P(t, q)$, have t -degrees at most $b-1$ and exactly b , respectively, where $b := \sum_{i=1}^{\nu} b_i$,
- (iv) the denominator factors $1 - q^{a_i} t^{b_i}$ all have $b_i \leq \#(P \cap \mathbb{Z}^n) - 1$,
- (v) for P a simplex, $N_P(t, q), \bar{N}_P(t, q)$ have nonnegative coefficients as polynomials in t, q .

And lastly one has a q -analogue of Ehrhart-Macdonald reciprocity:

- (vi) the two series $E_P(t, q), \bar{E}_P(t, q)$ determine each other via

$$q^d \bar{E}_P(t, q) = (-1)^{d+1} E_P(t^{-1}, q^{-1}).$$

3.2. Examples: lattice polygons. Having computed $E_P(t, q)$ for all one-dimensional lattice polytopes (line segments) in the Introduction and Example 3.2, one might wish to see data for lattice *polygons*. Proposition 3.3 allows one to consider them only up to the action of $\text{Aff}(\mathbb{Z}^2)$. If one bounds the normalized volume of a d -dimensional lattice polytope P in \mathbb{R}^d , a well-known result of Lagarias and Ziegler [30] shows that there are only finitely many such P up to the action of $\text{Aff}(\mathbb{Z}^d)$. Work of Balletti [3] gives an algorithm to list them, including an online database at














normalized area of P	vertices of P	$h_P^*(t) = 1 + h_1^*t + h_2^*t^2$	$E_P(t, q)$	Aff(\mathbb{Z}^2)-equivalent to antiblocking?
1	$(0, 0), (1, 0), (0, 1)$ 	1	$\frac{1}{(1-t)(1-qt)^2}$	Yes
2	$(0, 0), (1, 0), (1, 2)$ 	$1 + t$	$\frac{1+qt}{(1-t)(1-qt)(1-q^2t)}$	Yes
2	$(0, 0), (1, 0), (0, 1), (1, 1)$ 	$1 + t$	$\frac{1+qt}{(1-t)(1-qt)(1-q^2t)}$	Yes
3	$(0, 0), (1, 0), (1, 3)$ 	$1 + 2t$	$\frac{1+qt+q^2t}{(1-t)(1-qt)(1-q^2t)}$	Yes
3	$(0, 0), (1, 0), (2, 3)$ 	$1 + t + t^2$	$\frac{(1+qt)(1+qt+q^2t^2)}{(1-t)(1-q^2t)(1-q^3t^2)}$	No
3	$(0, 0), (1, 0), (0, 1), (-2, 1)$ 	$1 + 2t$	$\frac{1+qt-q^2t^2-q^3t^2}{(1-t)(1-qt)(1-q^2t)^2}$	Yes
4	$(0, 0), (1, 0), (1, 4)$ 	$1 + 3t$	$\frac{1+t(q+q^2+q^3)}{(1-t)(1-qt)(1-q^4t)}$	Yes
4	$(0, 0), (1, 0), (3, 4)$ 	$(1 + t)^2$	$\frac{(1+qt)^2}{(1-t)(1-q^2t)^2}$	No
4	$(0, 0), (2, 0), (0, 2)$ 	$1 + 3t$	$\frac{1+2qt+q^2t}{(1-t)(1-q^2t)^2}$	Yes
4	$(0, 0), (1, 0), (0, 1), (-3, 1)$ 	$1 + 3t$	$\frac{1+qt+q^2t-q^2t^2-q^3t^2-q^4t^2}{(1-t)(1-qt)(1-q^2t)(1-q^3t)}$	Yes
4	$(0, 0), (1, 0), (0, 2), (1, 2)$ 	$1 + 3t$	$\frac{1+qt+q^2t-q^2t^2-q^3t^2-q^4t^2}{(1-t)(1-qt)(1-q^2t)(1-q^3t)}$	Yes
4	$(0, 0), (2, 0), (0, 1), (1, -1)$ 	$(1 + t)^2$	$\frac{(1+qt)^2}{(1-t)(1-q^2t)^2}$	No
4	$(0, 0), (1, 0), (1, 2), (2, 2)$ 	$(1 + t)^2$	$\frac{(1+qt)^2}{(1-t)(1-q^2t)^2}$	No

FIGURE 1. $E_P(t, q)$ for lattice polygons P of area at most 4, up to Aff(\mathbb{Z}^2).

vertices of P	$h_P^*(t) =$ $1 + h_1^*t + h_2^*t^2$	$E_P(t, q)$	Aff(\mathbb{Z}^2)-equivalent to antiblocking?
$(0, 0), (1, 0), (1, 5)$	$1 + 4t$	$\frac{1+t(q+q^2+q^3+q^4)}{(1-t)(1-qt)(1-q^3t)}$	Yes
$(0, 0), (1, 0), (2, 5)$	$1 + 2t + 2t^2$	$\frac{(1+qt)(1+t(q+q^2)+t^2(q^3+q^4))}{(1-t)(1-q^2t)(1-q^5t^2)}$	No
$(0, 0), (1, 0), (0, 1), (-4, 1)$	$1 + 4t$	$\frac{1+qt+q^2t+q^3t-q^2t^2-q^3t^2-q^4t^2-q^5t^2}{(1-t)(1-qt)(1-q^2t)(1-q^4t)}$	Yes
$(0, 0), (2, 0), (0, 1), (-3, 1)$	$1 + 4t$	$\frac{1+qt+2q^2t-q^2t^2-q^3t^2-q^4t^2-q^5t^2}{(1-t)(1-qt)(1-q^3t)^2}$	Yes
$(0, 0), (1, 0), (2, 3), (2, 1)$	$1 + 3t + t^2$	$\frac{1+2qt+2q^2t+2q^3t^2+2q^4t^2+q^5t^3}{(1-t)(1-q^2t)(1-q^5t^2)}$	No
$(0, 0), (1, 0), (1, 2), (2, 2), (0, -1)$	$1 + 3t + t^2$	$\frac{1+2qt+2q^2t+2q^3t^2+2q^4t^2+q^5t^3}{(1-t)(1-q^2t)(1-q^5t^2)}$	No

FIGURE 2. $E_P(t, q)$ for lattice polygons of area 5, up to Aff(\mathbb{Z}^2).

<https://github.com/gabrieleballetti/small-lattice-polytopes> listing equivalence classes of lattice polytopes up to dimension 6 of relatively small normalized volumes. In particular, it includes lattice triangles up to normalized area 1000.

Using this data, Figures 1 and 2 present the q -Ehrhart series $E_P(t, q)$ of Aff(\mathbb{Z}^2)-equivalence classes of lattice polygons P of normalized volume at most 5. These were first guessed using `Macaulay2` to compute $i_P(m; q) := \text{Hilb}(R(\mathbb{Z}^2 \cap mP), q)$ for $m = 0, 1, 2, \dots$ up to some reasonably large values of m . The guesses were then proven correct via some extra computation in `Macaulay2` that uses our results on the *harmonic algebras* \mathcal{H}_P in Section 5; see Remark 5.8 below for an explanation. Note the last column of the tables, indicating whether the polygon P is Aff(\mathbb{Z}^2)-equivalent to one in the tamer subclass of *antiblocking* lattice polytopes, discussed in Section 3.3.

The data in Figures 1 and 2 is consistent with Conjecture 1.1. However, we close this subsection with remarks on some cautionary features.

Remark 3.4. The specialization $[E_P(t, q)]_{q=1} = E_P(t)$ sometimes has interesting numerator and denominator cancellations when $q \rightarrow 1$.

Remark 3.5. Note that, for fixed d , the classical Ehrhart series $E_P(t) \in \mathbb{Q}(t)$ for a lattice d -polytope can be expressed as an affine-linear function of the d real parameters (h_1^*, \dots, h_d^*) :

$$E_P(t) = \frac{1}{(1-t)^{d+1}} + h_1^* \cdot \frac{t}{(1-t)^{d+1}} + \dots + h_d^* \cdot \frac{t^d}{(1-t)^{d+1}}.$$

One has no such affine-linear formula for $E_P(t, q)$, already at $d = 1$, since the formulas (57)

$$E_P(t, q) = \frac{1 + tq[v-1]_q}{(1-t)(1-tq^v)}$$


give affine-linearly independent functions of $h_1^* = v - 1$ where $v = \text{vol}_1(P)$. On the other hand this formula does express $E_P(t, q)$ for $d = 1$ as a function (which is *not* affine-linear) of the one real parameter h_1^* . When $d = 2$, there can be no such function of the two parameters (h_1^*, h_2^*) , since one can have two lattice polygons P, P' with the same classical series $E_P(t) = E_{P'}(t)$ but different q -Ehrhart series $E_P(t, q) \neq E_{P'}(t, q)$; this happens for two area 3 lattice polygons in Figure 1.

3.3. Examples: antiblocking polytopes. For certain polytopes P introduced by Fulkerson [16, 17] in the context of combinatorial optimization, both q -Ehrhart series $E_P(t, q), \bar{E}_P(t, q)$ have a simpler description as classical weighted lattice point enumerators, avoiding harmonic spaces. This will allow us to verify Conjecture 1.1 for such polytopes.

In this section, abbreviate the nonnegative reals by $\mathbb{R}_{\geq 0} := \{z \in \mathbb{R} : z \geq 0\}$, the nonnegative orthant by $\mathbb{R}_{\geq 0}^n$, and the nonnegative integers $\mathbb{N} := \{0, 1, 2, \dots\}$. Define the componentwise partial order on $\mathbb{R}_{\geq 0}^n$ via $\mathbf{z} \leq \mathbf{z}'$ if $z_i \leq z'_i$ for all i .

Definition 3.6. Say that a convex polytope P is *antiblocking* (or *of antiblocking type*) if $P \subset \mathbb{R}_{\geq 0}^n$, and P forms a (lower) order ideal in the componentwise order on $\mathbb{R}_{\geq 0}^n$, that is, whenever $\mathbf{0} \leq \mathbf{z} \leq \mathbf{z}'$ with $\mathbf{z}' \in P$ then also $\mathbf{z} \in P$.

Here is an example of an antiblocking polytope P inside the orthant $\mathbb{R}_{\geq 0}^2$.



Intersecting antiblocking polytopes with \mathbb{Z}^n gives rise to point loci \mathcal{Z} with a restrictive property that vastly simplifies the ring $R(\mathcal{Z})$ and harmonic space $V_{\mathcal{Z}}$.

Definition 3.7. Call a subset $\mathcal{Z} \subset \mathbb{R}^n$ *shifted* if $\mathcal{Z} \subset \mathbb{N}^n$ and \mathcal{Z} forms a (lower) order ideal in the componentwise order on \mathbb{N}^n , that is, whenever $\mathbf{0} \leq \mathbf{z} \leq \mathbf{z}'$ and $\mathbf{z}' \in \mathcal{Z}$ then also $\mathbf{z} \in \mathcal{Z}$. Equivalently, $\mathcal{Z} \subset \mathbb{N}^n$ is shifted if and only if the \mathbf{k} -vector space $I := \text{span}_{\mathbf{k}}\{\mathbf{x}^{\mathbf{a}} : \mathbf{a} \in \mathbb{N}^n \setminus \mathcal{Z}\}$ forms a *monomial ideal* inside $S := \mathbf{k}[\mathbf{x}]$.

Lemma 3.8. For any finite shifted point locus $\mathcal{Z} \subset \mathbb{N}^n$,

- (i) the ideal $\text{gr } \mathbf{I}(\mathcal{Z}) \subset S = \mathbb{R}[\mathbf{x}]$ is monomial, with \mathbb{R} -basis $\{\mathbf{x}^{\mathbf{a}} : \mathbf{a} \in \mathbb{N}^n \setminus \mathcal{Z}\}$,
- (ii) the quotient ring $R(\mathcal{Z}) = S/\text{gr } \mathbf{I}(\mathcal{Z})$ of $S = \mathbb{R}[\mathbf{x}]$ has \mathbb{R} -basis $\{\bar{\mathbf{x}}^{\mathbf{z}} : \mathbf{z} \in \mathcal{Z}\}$,
- (iii) the harmonic space $V_{\mathcal{Z}} \subseteq S$ has \mathbb{R} -basis $\{\mathbf{y}^{\mathbf{z}} : \mathbf{z} \in \mathcal{Z}\}$ inside $\mathbb{D} = \mathbb{R}[\mathbf{y}]$.

Proof. Since \mathcal{Z} is shifted, the \mathbb{R} -vector space $I := \text{span}_{\mathbb{R}}\{\mathbf{x}^{\mathbf{a}} : \mathbf{a} \in \mathbb{N}^n \setminus \mathcal{Z}\}$ is a monomial ideal. Note that all assertions in the lemma would follow from showing (i), that is, $I = \text{gr } \mathbf{I}(\mathcal{Z})$.

To see this, we first show the inclusion $I \subseteq \text{gr } \mathbf{I}(\mathcal{Z})$. Since \mathcal{Z} is shifted, we claim that for any $\mathbf{a} \in \mathbb{N}^n \setminus \mathcal{Z}$, the polynomial

$$f_{\mathbf{a}}(\mathbf{x}) := \prod_{i=1}^n x_i(x_i - 1) \cdots (x_i - (a_i - 1))$$

lies in the vanishing ideal $\mathbf{I}(\mathcal{Z})$ for \mathcal{Z} . To see this claim, assume not, so there exists some point $\mathbf{z} \in \mathcal{Z}$ having $f(\mathbf{z}) \neq 0$. But then for each i one must have $z_i \in \mathbb{N} \setminus \{0, 1, 2, \dots, a_i - 1\}$, implying $\mathbf{z} \geq \mathbf{a}$, and contradicting $\mathbf{a} \notin \mathcal{Z}$. Hence $\text{gr } \mathbf{I}(\mathcal{Z})$ contains $\tau(f_{\mathbf{a}}) = \prod_{i=1}^n x_i^{a_i} = \mathbf{x}^{\mathbf{a}}$, showing $I \subseteq \text{gr } \mathbf{I}(\mathcal{Z})$.

To show the opposite inclusion $I \supseteq \text{gr } \mathbf{I}(\mathcal{Z})$, note that the surjection $S/I \rightarrow S/\text{gr } \mathbf{I}(\mathcal{Z})$ must be an isomorphism by dimension-counting, since $\dim_{\mathbb{R}} S/I = \#\mathcal{Z} = \dim_{\mathbb{R}} S/\text{gr } \mathbf{I}(\mathcal{Z})$. \square

If one defines the following q -weight enumerator for finite point loci $\mathcal{Z} \subset \mathbb{N}^n$

$$(60) \quad w_{\mathcal{Z}}(q) := \sum_{\mathbf{z} \in \mathcal{Z}} q^{|\mathbf{z}|} \quad \text{where } |\mathbf{z}| := z_1 + \cdots + z_n,$$

then the next corollary is immediate from Lemma 3.8

Corollary 3.9. *For finite shifted point loci $\mathcal{Z} \subset \mathbb{N}^n$, one has*

$$\text{Hilb}(R(\mathcal{Z}), q) = \text{Hilb}(V_{\mathcal{Z}}, q) = w_{\mathcal{Z}}(q).$$

We next explain how antiblocking polytopes P give rise to shifted point loci. For $0 \leq d \leq n$, let

$$(\mathbf{1}_d, \mathbf{0}_{n-d}) := \underbrace{(1, 1, \dots, 1)}_{d \text{ times}}, \underbrace{(0, 0, \dots, 0)}_{n-d \text{ times}}.$$

Proposition 3.10. *Let P be an antiblocking d -polytope in \mathbb{R}^n .*

- (i) *The finite point locus $\mathbb{Z}^n \cap P \subset \mathbb{N}^n$ is shifted, and the same holds for $\mathbb{Z}^n \cap mP$ for all $m \in \mathbb{N}$.*
- (ii) *There exists a re-indexing of the coordinates in \mathbb{R}^n such that the translated set*

$$\mathbb{Z}^n \cap \text{int}(P) - (\mathbf{1}_d, \mathbf{0}_{n-d}) := \{\mathbf{z} - (\mathbf{1}_d, \mathbf{0}_{n-d}) : \mathbf{z} \in \mathbb{Z}^n \cap \text{int}(P)\}$$

is shifted, and the same holds for the set $\mathbb{Z}^n \cap \text{int}(mP) - (\mathbf{1}_d, \mathbf{0}_{n-d})$ for all $m \in \mathbb{N}$.

Proof. First note that whenever P is antiblocking, then so is mP . Thus one only needs to check the first part of both assertions (i),(ii) for P .

Assertion (i) for P is straightforward from the definitions of antiblocking and shiftedness.

To prove assertion (ii) for P , consider the *support sets* $\text{supp}(\mathbf{z}) := \{i = 1, 2, \dots, n : z_i \neq 0\}$ for $\mathbf{z} \in P$. Since P is convex and lies in $\mathbb{R}_{\geq 0}$, whenever $\mathbf{z}, \mathbf{z}' \in P$, the points \mathbf{z}'' in the interior of the line segment between them have $\text{supp}(\mathbf{z}'') = \text{supp}(\mathbf{z}) \cup \text{supp}(\mathbf{z}')$. Consequently, there exists $\mathbf{z} \in P$ with $\text{supp}(\mathbf{z}) = \bigcup_{\mathbf{z}' \in P} \text{supp}(\mathbf{z}')$. Reindexing coordinates so that $\text{supp}(\mathbf{z}) = \{1, 2, \dots, e\}$, one concludes that P contains the e -dimensional parallelepiped of vectors componentwise between $\mathbf{0}$ and \mathbf{z} , and also P is contained in the e -dimensional coordinate subspace $\mathbb{R}^e \times \{\mathbf{0}_{n-d}\}$. Since P is d -dimensional, this forces $e = d$, and the last $n - d$ coordinates vanish on all points of P .

Therefore in the rest of the proof that $\mathbb{Z}^n \cap \text{int}(P) - (\mathbf{1}_d, \mathbf{0}_{n-d})$ is shifted, it suffices to project away the last $n - d$ coordinates which all vanish on P , and assume $d = n$. As a full d -dimensional antiblocking polytope $P \subset \mathbb{R}^d$, starting with any inequality description

$$P = \{\mathbf{z} \in \mathbb{R}^d : \mathbf{z} \geq \mathbf{0}, A\mathbf{z} \leq \mathbf{b}\}$$

where A, \mathbf{b} have all entries in $\mathbb{R}_{>0}$, one can describe $\text{int}(P)$ as

$$\text{int}(P) = \{\mathbf{z} \in \mathbb{R}^d : \mathbf{z} > \mathbf{0}, A\mathbf{z} < \mathbf{b}\}.$$

Then the points of $\mathbb{Z}^d \cap \text{int}(P)$ have this description:

$$\mathbb{Z}^d \cap \text{int}(P) = \{\mathbf{z} \in \mathbb{Z}^d : \mathbf{z} \geq \mathbf{1}_d, A\mathbf{z} < \mathbf{b}\}.$$

Hence the points of $\mathbb{Z}^d \cap \text{int}(P) - \mathbf{1}_d$ have this description, which defines a shifted subset of \mathbb{N}^n :

$$\mathbb{Z}^d \cap \text{int}(P) - \mathbf{1}_d = \{\mathbf{z} \in \mathbb{Z}^d : \mathbf{z} \geq \mathbf{0}_d, A\mathbf{z} < \mathbf{b} - A\mathbf{1}_d\}. \quad \square$$

Proposition 3.10 lets us re-cast the q -Ehrhart series for an antiblocking polytope in terms of the following simpler and more classical weight enumerators, involving $w_{\mathcal{Z}}(q)$ from (60).

Definition 3.11. For any lattice polytope $P \subset \mathbb{R}^m$, define two q -weight enumerator series

$$W_P(t, q) := \sum_{m=0}^{\infty} w_{\mathbb{Z}^m \cap mP}(q) \cdot t^m, \quad \text{and} \quad \overline{W}_P(t, q) := \sum_{m=0}^{\infty} w_{\mathbb{Z}^m \cap \text{int}(mP)}(q) \cdot t^m.$$

Combining Proposition 3.10, Corollary 3.9, and Proposition 3.3 yields this consequence.

Corollary 3.12. For any d -dimensional lattice polytope $P \subset \mathbb{R}^n$ which is antiblocking, one has

$$(61) \quad E_P(t, q) = W_P(t, q),$$

$$(62) \quad q^d \cdot \overline{E}_P(t, q) = \overline{W}_P(t, q).$$

Classical results now apply to the weight enumerators $W_P(t, q), \overline{W}_P(t, q)$, showing that they have rational expressions with predictable properties, for *any* lattice polytopes P .

Proposition 3.13. Let $P \subset \mathbb{R}^n$ be any d -dimensional lattice polytope, with vertices $\mathbf{v}^{(1)}, \mathbf{v}^{(2)}, \dots, \mathbf{v}^{(\nu)}$.

(i) Both $W_P(t, q), \overline{W}_P(t, q)$ have rational expressions with the same denominator

$$W_P(t, q) = \frac{N_P(t, q)}{\prod_{i=1}^{\nu} (1 - tq^{|\mathbf{v}^{(i)}|})} \quad \text{and} \quad \overline{W}_P(t, q) = \frac{\overline{N}_P(t, q)}{\prod_{i=1}^{\nu} (1 - tq^{|\mathbf{v}^{(i)}|})}$$

and numerators $N_P(t, q), \overline{N}_P(t, q) \in \mathbb{Z}[t, q]$.

(ii) For a lattice d -simplex P with vertices $\{\mathbf{v}^{(i)}\}_{i=1,2,\dots,d+1}$, one has these numerators in $\mathbb{N}[t, q]$

$$N_P(t, q) = \sum_{\mathbf{z} \in \mathbb{Z}^{n+1} \cap \Pi} t^{z_{n+1}} q^{z_1 + \dots + z_n} \quad \text{and} \quad \overline{N}_P(t, q) = \sum_{\mathbf{z} \in \mathbb{Z}^{n+1} \cap \Pi^{\text{opp}}} t^{z_{n+1}} q^{z_1 + \dots + z_n},$$

where Π is the semi-open parallelepiped from (3), and Π^{opp} its ‘‘opposite’’, defined by

$$\Pi := \left\{ \sum_{j=1}^{d+1} c_j \cdot (1, \mathbf{v}^{(j)}) : 0 \leq c_i < 1 \right\} \quad \text{and} \quad \Pi^{\text{opp}} := \left\{ \sum_{j=1}^{d+1} c_j \cdot (1, \mathbf{v}^{(j)}) : 0 < c_i \leq 1 \right\}.$$

(iii) The two series $W_P(t, q), \overline{W}_P(t, q)$ determine each other via

$$\overline{W}_P(t, q) = (-1)^{d+1} W_P(t^{-1}, q^{-1}).$$

Proof. These are all bivariate specializations of results of Stanley on multivariable lattice point enumerators for rational polyhedral cones, implicit in [40], and more explicit in [47, §4.5]. One must apply them to $\text{cone}(P) \subset \mathbb{R}^{n+1}$ which is defined to be the $(d+1)$ -dimensional cone nonnegatively spanned by the vectors $\{(1, \mathbf{v}^{(j)})\}_{j=1,\dots,\nu}$. In variable set $\mathbf{x} = (x_0, x_1, \dots, x_n)$, Stanley considers

$$E_{\text{cone}(P)}(\mathbf{x}) := \sum_{\mathbf{z} \in \text{cone}(P)} \mathbf{x}^{\mathbf{z}} \quad \text{and} \quad \overline{E}_{\text{cone}(P)}(\mathbf{x}) := \sum_{\mathbf{z} \in \text{int}(\text{cone}(P))} \mathbf{x}^{\mathbf{z}}.$$

He proves [47, Thm. 4.5.11] that one has rational expressions of the form

$$(63) \quad E_{\text{cone}(P)}(\mathbf{x}) = \frac{N_P(\mathbf{x})}{\prod_{i=1}^{\nu} (1 - \mathbf{x}^{\mathbf{v}^{(i)}})} \quad \text{and} \quad \bar{E}_{\text{cone}(P)}(\mathbf{x}) = \frac{\bar{N}_P(\mathbf{x})}{\prod_{i=1}^{\nu} (1 - \mathbf{x}^{\mathbf{v}^{(i)}})}$$

for some polynomials $N_P(\mathbf{x}), \bar{N}_P(\mathbf{x})$ in $\mathbb{Z}[\mathbf{x}]$. He also proves [47, Cor. 4.5.8] that a d -simplex P has

$$(64) \quad N_P(\mathbf{x}) = \sum_{\mathbf{z} \in \mathbb{Z}^{n+1} \cap \Pi} \mathbf{x}^{\mathbf{z}} \quad \text{and} \quad \bar{N}_P(\mathbf{x}) = \sum_{\mathbf{z} \in \mathbb{Z}^{n+1} \cap \Pi^{\text{opp}}} \mathbf{x}^{\mathbf{z}}.$$

And he shows in [47, Thm. 4.5.14] (often called *Stanley's Reciprocity Theorem*) that

$$(65) \quad E_P(x_0^{-1}, x_1^{-1}, \dots, x_n^{-1}) = (-1)^{d+1} \bar{E}_P(x_0, x_1, \dots, x_n).$$

Assertions (i),(ii),(iii) are the specializations at $x_0 = t, x_1 = \dots = x_n = q$ of (63),(64),(65). \square

Corollary 3.14. *Conjecture 1.1 holds for all lattice polytopes P which are antiblocking.*

Proof. We claim that Proposition 3.13 implies that Conjecture 1.1 holds for antiblocking polytopes when one specializes all assertions (i)-(vi) in the conjecture, taking $\nu = V$ and with the denominator factors of the more explicit form $1 - q^{a_i} t^{b_i} = 1 - tq^{|\mathbf{v}^{(i)}|}$. Therefore all $b_i = 1$, and $b := \sum_{i=1}^{\nu} b_i = \nu$.

Most assertions of the conjecture are seen to follow easily. The trickiest is assertion (iii) on the t -degrees of $N(t, q), \bar{N}(t, q)$. But since $[E_P(t, q)]_{q=1} = E_P(t)$ satisfies assertion (i) of the Classical Ehrhart Theorem from the Introduction, this implies $E_P(t, q)$ satisfies Conjecture 1.1(iii). \square

Remark 3.15. Not all lattice polytopes P are equivalent via $\text{Aff}(\mathbb{Z}^d)$ to antiblocking polytopes. In particular, sometimes $\text{gr } \mathbf{I}(\mathbb{Z}^d \cap P)$ is not a monomial ideal in $\mathbb{R}[\mathbf{x}]$, and $V_{\mathbb{Z}^d \cap P} \subset \mathbb{R}[\mathbf{y}]$ has no monomial \mathbb{R} -basis. For example, the lattice triangle P shown here with vertices $\{(0, 0), (2, 1), (1, 2)\}$



is equivalent to the second triangle of area 3 in Figure 1, with vertices $\{(0, 0), (1, 0), (2, 3)\}$. It has

$$E_P(t) = \frac{1 + t + t^2}{(1 - t)^3},$$

$$E_P(t, q) = \frac{(1 + qt)(1 + qt + q^2 t^2)}{(1 - t)(1 - q^2 t)(1 - q^3 t^2)}.$$

Via hand calculation or using `Macaulay2`, one can check that

$$(66) \quad \text{gr } \mathbf{I}(\mathbb{Z}^2 \cap P) = (x_1^2 - x_2^2, 2x_1x_2 - x_2^2, x_2^3) \subseteq \mathbb{R}[x_1, x_2]$$

and this ideal cannot be generated by monomials. Again hand calculation or `Macaulay2` shows that

$$V_{\mathbb{Z}} = \text{span}_{\mathbb{R}}\{1, y_1, y_2, y_1^2 + y_1y_2 + y_2^2\}$$

which has no \mathbb{R} -basis of monomials.

Remark 3.16. The q -weight enumerators in Definition 3.11 bear some resemblance to work of Chapoton [8]. He introduced q -analogues of $E_P(t)$ that first require a choice a \mathbb{Z} -linear functional $\lambda : \mathbb{Z} \rightarrow \mathbb{Z}$ which is nonconstant along edges of P and has $\lambda(\mathbf{z}^{(i)}) \geq 0$ for all vertices of P . Having made such a choice, he defined

$$W_{\lambda}(P, q) := \sum_{\mathbf{z} \in \mathbb{Z}^n \cap P} q^{\lambda(\mathbf{z})} \quad \text{and} \quad \text{Ehr}_{P, \lambda}(t, q) := \sum_{m \geq 0} t^m W_{\lambda}(mP, q).$$

Therefore Corollary 3.12 shows that if a lattice polytope P is antiblocking, and happens to have that the functional $\lambda(\mathbf{z}) := |\mathbf{z}| = z_1 + \cdots + z_n$ is nonconstant along its edges, then $\text{Ehr}_{P,\lambda}(t, q) = E_P(t, q)$.

3.4. Definition of the equivariant q -Ehrhart series. We next incorporate symmetries of subgroups of $GL_n(\mathbb{Z})$ acting on a lattice polytope. Such groups are often called *crystallographic groups*.

Definition 3.17. Let P be a lattice polytope in \mathbb{R}^n which is stable under the action of a finite crystallographic group $G \subset GL_n(\mathbb{Z})$. Define the *equivariant q -Ehrhart series*

$$E_P^G(t, q) := \sum_{m=0}^{\infty} [V_{\mathbb{Z}^n \cap mP}]_q \cdot t^m \quad \left(= \sum_{m=0}^{\infty} [R(\mathbb{Z}^n \cap mP)]_q \cdot t^m \right)$$

where recall $[U]_q = \sum_{d \geq 0} [U_d] q^d$ is the class in $\text{Rep}_{\mathbb{Z}}(G)[[q]]$ of any graded $\mathbf{k}[G]$ -module $U = \bigoplus_{d \geq 0} U_d$. Hence $E_P^G(t, q)$ lies in the power series ring $\text{Rep}_{\mathbb{Z}}(G)[[t, q]]$, and even lies in its subring $\text{Rep}_{\mathbb{Z}}(G)[q][[t]]$.

When $q \rightarrow 1$, this series $E_P^G(t, q)$ specializes to Stapledon's *equivariant Ehrhart series* [48] involving the ungraded characters of $V_{mP \cap \mathbb{Z}^n}$, or equivalently the permutation $\mathbb{R}[G]$ -modules $\mathbb{Z}^n \cap mP$.

Example 3.18. We return to Example 3.2, where P is a lattice polytope $P \subset \mathbb{R}^1$, but now assume it is stable under the action of a nontrivial crystallographic group G . This implies $P = [-b, b]$ for some $b \geq 0$ and $G = GL_1(\mathbb{Z}) = \{\pm 1\}$. One can identify the representation ring

$$\text{Rep}_{\mathbb{R}}(G) \cong \mathbb{Z}[\epsilon]/(\epsilon^2 - 1) = \text{span}_{\mathbb{Z}}\{1, \epsilon\},$$

where $1, \epsilon$ denote the isomorphism classes of the one-dimension trivial and nontrivial $\mathbb{R}[G]$ -modules, respectively. A calculation using $\mathbb{Z}^1 \cap mP = \{-mb, -mb + 1, \dots, mb - 1, mb\}$, similar to the one in the Introduction, shows that

$$\mathbf{I}(\mathbb{Z}^1 \cap mP) = \left(\prod_{i=-mb}^{+mb} (x - i) \right), \quad \text{gr } \mathbf{I}(\mathbb{Z}^1 \cap mP) = (x^{2mb+1}) \subset S = \mathbb{R}[x]$$

and consequently, one has

$$R(\mathbb{Z}^1 \cap mP) = S/(x^{2bm+1}) = \text{span}_{\mathbb{R}}\{1, \bar{x}, \bar{x}^2, \dots, \bar{x}^{2bm}\},$$

$$V_{\mathbb{Z}^1 \cap mP} = \ker \left(\left(\frac{\partial}{\partial y} \right)^{2bm+1} : \mathbb{D} \rightarrow \mathbb{D} \right) = \text{span}_{\mathbb{R}}\{1, y, y^2, \dots, y^{2bm}\} \subset \mathbb{D} = \mathbb{R}[y].$$

Since $G = \{\pm 1\}$ negates the variables x, y in S, \mathbb{D} , one concludes (with some algebra) that

$$(67) \quad E_P^G(t, q) = \sum_{m=0}^{\infty} (1 + q^2 + q^4 + \cdots + q^{2mb}) t^m + \sum_{m=0}^{\infty} (q + q^3 + q^5 + \cdots + q^{2mb-1}) \cdot \epsilon t^m$$

$$(68) \quad = \frac{1}{1 - q^2} \left[\frac{1}{1 - t} - \frac{q^2}{1 - tq^{2b}} + q\epsilon \left(\frac{1}{1 - t} - \frac{1}{1 - tq^{2b}} \right) \right] = \frac{1 + t(q^2[b-1]_{q^2} + q[b]_{q^2}\epsilon)}{(1 - t)(1 - tq^{2b})}$$

This expression (68) specializes to the non-equivariant series $E_P(t, q)$ by applying the ring homomorphism $\text{Rep}_{\mathbb{R}}(G) \rightarrow \mathbb{Z}$ that sets $\epsilon = 1$, giving an answer consistent with (57) at $v = \text{vol}_1(P) = 2b$:

$$E_P(t, q) = [E_P^G(t, q)]_{\epsilon=1} = \frac{1 + t(q^2[b-1]_{q^2} + q[b]_{q^2})}{(1 - t)(1 - tq^{2b})} = \frac{1 + tq[2b-1]_q}{(1 - t)(1 - tq^{2b})}.$$

On the other hand, setting $q = 1$ in (68) should also give Stapledon's equivariant Ehrhart series $E_P^G(t)$, which we calculate directly here. The G -orbit structure permuting

$$\mathbb{Z}^1 \cap mP = \{-mb, \dots, -1, 0, +1, \dots, mb\} = \{0\} \sqcup \bigsqcup_{i=1}^{mb} \{\pm i\}$$

has one fixed point orbit $\{0\}$ with class 1 in $\text{Rep}_{\mathbb{R}}[G]$ and mb free orbits $\{\pm i\}_{i=1,2,\dots,mb}$, each with class $1 + \epsilon$ in $\text{Rep}_{\mathbb{R}}[G]$. Hence one calculates that

$$E_P^G(t) := \sum_{m=0}^{\infty} [\mathbb{Z}^1 \cap mP] t^m = \sum_{m=0}^{\infty} (1 + (1 + \epsilon)mb) t^m = \frac{1}{1-t} + \frac{(1 + \epsilon)bt}{(1-t)^2} = \frac{1 + t((b-1) + b\epsilon)}{(1-t)^2},$$

which agrees with the expression for $E_P^G(t, q)$ in (68) at $q = 1$.

3.5. Examples: standard simplices. We compute the equivariant q -Ehrhart series for two families of simplices that carry an action of the symmetric group \mathfrak{S}_n on n letters, permuting the standard basis vectors $\mathbf{e}_1, \dots, \mathbf{e}_n$ in \mathbb{R}^n .

Definition 3.19. The *standard* $(n-1)$ -simplex in \mathbb{R}^n is

$$\Delta^{n-1} := \text{conv}\{\mathbf{e}_1, \dots, \mathbf{e}_n\},$$

lying inside the affine subspace of \mathbb{R}^n where $x_1 + \dots + x_n = 1$. There is also a full n -dimensional simplex in \mathbb{R}^n which is the *pyramid* $\text{Pyr}(\Delta^{n-1})$ having Δ^{n-1} as its base, and apex at the origin $\mathbf{0}$:

$$\text{Pyr}(\Delta^{n-1}) := \text{conv}\{\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_n\}.$$

We compute the equivariant q -Ehrhart series of $\Delta^{n-1}, \text{Pyr}(\Delta^{n-1})$. For this, it helps to have two facts. The first is an easy lemma on power series $F(q) = \sum_{i \geq 0} a_i q^i \in R[[q]]$ for commutative rings R , and their *truncations to $q^{\leq m}$* , defined as these polynomial partial sums in $R[q]$:

$$\text{trunc}_{\leq q^m} F(q) := \sum_{i=0}^m a_i q^i.$$

Lemma 3.20. *For any $F(q)$ in $R[[q]]$, one has $\sum_{m=0}^{\infty} t^m \cdot \text{trunc}_{\leq q^m} F(q) = \frac{F(tq)}{1-t}$ in $R[[t, q]]$.*

The second fact is a formula due to Lusztig and Stanley [42, Prop. 4.11] for the \mathfrak{S}_n -equivariant Hilbert series of the polynomial ring $S = \mathbb{R}[\mathbf{x}]$. Recall that the simple $\mathbb{R}[\mathfrak{S}_n]$ -modules $\{\mathbb{S}^\lambda\}$ are indexed by number partitions λ of n (written $\lambda \vdash n$), so that $\text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)$ has \mathbb{Z} -basis $\{\mathbb{S}^\lambda : \lambda \vdash n\}$.

Theorem 3.21. *(Lusztig, Stanley) The graded $\mathbb{R}[\mathfrak{S}_n]$ -module $S = \mathbb{R}[\mathbf{x}]$ has this class in $\text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)[[q]]$:*

$$[S]_q := \sum_{d=0}^{\infty} q^d [S_d] = \frac{1}{(1-q)(1-q^2)\cdots(1-q^n)} \sum_{\lambda \vdash n} f^\lambda(q) \cdot [\mathbb{S}^\lambda]$$

with $f^\lambda(q)$ the \mathfrak{S}_n fake-degree polynomial, having these sum, product expressions [46, Cor. 7.21.5]

$$(69) \quad f^\lambda(q) = \sum_T q^{\text{maj}(T)} = q^{b(\lambda)} \frac{[n]!_q}{\prod_{x \in \lambda} [h(x)]_q}$$

whose notations are explained below.

Here T in the summation in (69) runs over all *standard Young tableaux* of shape λ , and $\text{maj}(T)$ is the sum of all entries i in T for which $i + 1$ appears weakly southwest of i (using English notation for tableaux). The q -factorial is $[n]!_q := [n]_q [n-1]_q \cdots [2]_q [1]_q$. For a partition λ with ℓ parts, we have $b(\lambda) := \sum_{i=1}^{\ell} (i-1)\lambda_i$. For a box x in row (i, j) of the Ferrers diagram of λ (written $x \in \lambda$ in the product), the *hooklength* is $h(x) := \lambda_i + \lambda'_j - (i + j) + 1$, with λ' the *conjugate partition* to λ .

We first compute the equivariant q -Ehrhart series for the n -simplex $\text{Pyr}(\Delta^{n-1}) \subset \mathbb{R}^n$.

Proposition 3.22. *For the n -simplex $P := \text{Pyr}(\Delta^{n-1}) = \text{conv}\{\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_n\} \subset \mathbb{R}^n$, one has*

$$(70) \quad E_P^{\mathfrak{S}_n}(t, q) = \frac{1}{(1-t) \cdot \prod_{i=1}^n (1-t^i q^i)} \sum_{\lambda \vdash n} f^\lambda(tq) \cdot [\mathbb{S}^\lambda],$$

$$(71) \quad E_P(t, q) = \frac{1}{(1-t)(1-tq)^n}.$$

Proof. We start with (70). Note P and all of its dilates mP are antiblocking, with mP defined by the inequalities $\mathbf{x} \geq 0$ and $x_1 + \cdots + x_n \leq m$. Hence Lemma 3.8 shows that

$$R(\mathbb{Z}^n \cap mP) = S/S_{\geq m+1}.$$

Thus the graded $\mathbb{R}[\mathfrak{S}_n]$ -module $R(\mathbb{Z}^n \cap mP)$ is the same as the permutation module on the monomials of degree at most m in $S = \mathbb{R}[\mathbf{x}]$. Therefore in $\text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)[[q]]$ one has

$$[R(\mathbb{Z}^n \cap mP)]_q = \text{trunc}_{q \leq m} [S]_q.$$

Applying Lemma 3.20 then gives the equality marked (*) here

$$(72) \quad E_P^{\mathfrak{S}_n}(t, q) := \sum_{m=0}^{\infty} [R(\mathbb{Z}^n \cap mP)]_q \cdot t^m \\ \stackrel{(*)}{=} \frac{1}{1-t} \cdot [S]_{q \rightarrow tq} = \frac{1}{(1-t) \cdot \prod_{i=1}^n (1-t^i q^i)} \sum_{\lambda \vdash n} f^\lambda(tq) \cdot [\mathbb{S}^\lambda],$$

with the last equality substituting $q \mapsto tq$ in the formula from Theorem 3.21. This proves (70).

Although one could deduce (71) from (70), a perhaps simpler path repeats the same argument but replacing $[S]_q$ with the known non-equivariant Hilbert series $\text{Hilb}(S, q) = 1/(1-q)^n$. \square

Now it is easier to deal with the equivariant series for the $(n-1)$ -simplex $\Delta^{n-1} \subset \mathbb{R}^n$.

Proposition 3.23. *For $P := \Delta^{n-1} = \text{conv}\{\mathbf{e}_1, \dots, \mathbf{e}_n\} \subset \mathbb{R}^n$, one has*

$$(73) \quad E_P^{\mathfrak{S}_n}(t, q) = (1-tq) \cdot E_{\text{Pyr}(P)}^{\mathfrak{S}_n}(t, q) = \frac{1}{(1-t) \cdot \prod_{i=2}^n (1-t^i q^i)} \sum_{\lambda \vdash n} f^\lambda(tq) \cdot [\mathbb{S}^\lambda]$$

$$(74) \quad E_P(t, q) = (1-tq) \cdot E_{\text{Pyr}(P)}(t, q) = \frac{1}{(1-t)(1-tq)^{n-1}}.$$

Proof. We start by identifying $\text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP)$. Since $\mathbb{Z}^n \cap mP \subset \mathbb{Z}^n \cap m\text{Pyr}(P)$, one has

$$\mathbf{I}(\mathbb{Z}^n \cap mP) \supset \mathbf{I}(\mathbb{Z}^n \cap m\text{Pyr}(P)), \\ \text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP) \supset \text{gr } \mathbf{I}(\mathbb{Z}^n \cap m\text{Pyr}(P)) = S_{\geq m+1}.$$

On the other hand, since mP lies in the affine hyperplane $x_1 + \cdots + x_n = m$ within \mathbb{R}^n , one also has $x_1 + \cdots + x_n - m \in \mathbf{I}(\mathbb{Z}^n \cap mP)$ and $e_1 := x_1 + \cdots + x_n \in \text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP)$. Consequently,

$$\text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP) \supseteq (e_1) + S_{\geq m+1},$$

and we claim that this inclusion is actually an equality. To see this, note that the surjection

$$S/((e_1) + S_{\geq m+1}) \twoheadrightarrow S/\text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP) = R(\mathbb{Z}^n \cap mP)$$

is an isomorphism via dimension-counting: since mP is the dilation of a standard $(n-1)$ -simplex,

$$\dim_{\mathbb{R}} R(\mathbb{Z}^n \cap mP) = \#\mathbb{Z}^n \cap mP = \binom{m+n-1}{m},$$

and if one abbreviates the quotient ring $\overline{S} := S/(e_1) \cong \mathbb{R}[x_1, \dots, x_{n-1}]$, then

$$\dim_{\mathbb{R}} S/((e_1) + S_{\geq m+1}) = \dim_{\mathbb{R}} \overline{S}/\overline{S}_{\geq m+1} = \binom{m+n-1}{m}.$$

From this one concludes that in $\text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)[q]$ one has

$$[R(\mathbb{Z}^n \cap mP)]_q = \text{trunc}_{\leq q^m} [\overline{S}]_q$$

and hence, applying Lemma 3.20, one has

$$E_P^{\mathfrak{S}_n}(t, q) := \sum_{m=0}^{\infty} [R(\mathbb{Z}^n \cap mP)]_q t^m = \frac{1}{1-t} \cdot [\overline{S}]_{q \rightarrow tq}.$$

On the other hand, we claim that $[\overline{S}]_q = (1-q) \cdot [S]_q$. This holds because $e_1 = x_1 + \dots + x_n$ is an \mathfrak{S} -invariant nonzero divisor in S , so that multiplying by e_1 on S gives rise to an exact sequence of graded $\mathbb{R}[\mathfrak{S}_n]$ -modules $0 \rightarrow S(-1) \rightarrow S \rightarrow \overline{S} \rightarrow 0$. Comparing this with (72) then proves (73).

The proof of (74) is again a parallel, but easier computation. \square

Remark 3.24. We note here some notationally convenient reformulations of these results, for readers familiar with the *ring of symmetric functions* $\Lambda = \bigoplus_{n=0}^{\infty} \Lambda_n$ and the *Frobenius characteristic isomorphism* $\Lambda_n \cong \text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)$; see [34, 39, 46] for more background. One note of **caution**: multiplication in $\text{Rep}_{\mathbb{R}}(\mathfrak{S}_n)$ corresponds to the *internal* or *Kronecker product* on Λ_n , not the *external* or *induction product* corresponding to the multiplication $\Lambda_{n_1} \times \Lambda_{n_2} \rightarrow \Lambda_{n_1+n_2}$.

This Frobenius characteristic isomorphism maps the *Schur function* $s_{\lambda} \mapsto [\mathbb{S}^{\lambda}]$, and the *complete homogeneous symmetric function* $h_n \mapsto [1]$. Using the *plethystic notation*, explained in Bergeron [6, Ch. 3], Haglund [24], Loehr and Remmel [32], one can show (see [6, eqn. (7.7)]) that the class $[S]_q$ of the graded $\mathbb{R}[\mathfrak{S}_n]$ -module S corresponds to this element of $\Lambda_n[[q]]$:

$$[S]_q \mapsto h_n[X + qX + q^2X + \dots] = h_n \left[\frac{X}{1-q} \right].$$

The Lusztig-Stanley Theorem 3.21 expanding this plethysm in Schur functions is discussed in Haglund [24, (1.89)]. Consequently, one can recast Propositions 3.23 and 3.22 plethystically:

$$E_{\text{Pyr}^n}^{\mathfrak{S}_n}(t, q) = \frac{1}{1-t} \cdot h_n \left[\frac{X}{1-tq} \right] \quad \text{and} \quad E_{\Delta^{n-1}}^{\mathfrak{S}_n}(t, q) = \frac{1-tq}{1-t} \cdot h_n \left[\frac{X}{1-tq} \right].$$

Remark 3.25. The relation $E_P(t, q) = (1-tq) \cdot E_{\text{Pyr}(P)}(t, q)$ for $P = \Delta^{n-1}$ that appeared in Propositions 3.22, 3.23 anticipates the behavior of $E_P(t, q)$ under the pyramid operation $P \mapsto \text{Pyr}(P)$, and more generally under free joins $P * Q$ of lattice polytopes, as described in Theorem 1.2(iii). Pyramids are the special case $\text{Pyr}(P) = P * Q$ where Q is the one-point polytope of dimension zero.

3.6. Examples: cross-polytopes. The *cross-polytope* or *hyperoctahedron* $\diamond^n \subseteq \mathbb{R}^n$ is defined by

$$\diamond^n := \text{conv}\{\pm \mathbf{e}_1, \dots, \pm \mathbf{e}_n\}.$$

Its symmetries are the *hyperoctahedral group* B_n of all *signed permutation matrices*, acting by permuting and negating coordinates in \mathbb{R}^n . We first analyze the non-equivariant q -Ehrhart series for \diamond_n , and then incorporate the group action.

Proposition 3.26. *The q -Ehrhart series of the cross-polytope \diamond^n is given by*

$$(75) \quad E_{\diamond^n}(t, q) = \frac{1}{1-t} \left(\frac{1+qt}{1-q^2t} \right)^n.$$

Proof. Let $P := \diamond^n$. Then for each $m \geq 0$, the lattice points of the dilate mP are given by

$$(76) \quad \mathbb{Z}^n \cap mP = \{\mathbf{z} \in \mathbb{Z}^n : |z_1| + \dots + |z_n| \leq m\}.$$

We wish to identify the ideal $\text{gr } \mathbf{I}(\mathbb{Z}^n \cap m \cdot P) \subset S$. For this, it helps to note that every monomial in $S = \mathbb{R}[\mathbf{x}]$ can be expressed uniquely in the form that separates even and odd powers of variables

$$(77) \quad \mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B = x_1^{2a_1} \cdots x_n^{2a_n} \cdot \prod_{j \in B} x_j$$

for some pair (\mathbf{a}, B) with $\mathbf{a} \in \mathbb{N}^n$ and $B \subseteq [n]$. We then have the following claim.

Claim: $\text{gr } \mathbf{I}(\mathbb{Z}^n \cap m \cdot P)$ is the monomial ideal $I := \text{span}_{\mathbb{R}}\{\mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B : |\mathbf{a}| + \#B > m\}$.

To verify the Claim, we first show that for any such monomial $\mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B$ in I , so $|\mathbf{a}| + \#B > m$, if one defines the set $A := \text{supp}(\mathbf{a}) = \{j : a_j \geq 1\}$, then $\mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B$ is divisible by

$$\tau(f) = \prod_{j \in A \setminus B} x_j^{2a_j-1} \prod_{j \in B} x_j^{2a_j+1}$$

where f is the following polynomial in $\mathbf{I}(\mathbb{Z}^n \cap mP)$:

$$f(\mathbf{x}) := \prod_{j \in A \setminus B} \prod_{i_j = -(a_j-1)}^{+(a_j-1)} (x_j - i_j) \cdot \prod_{j \in B} \prod_{i_j = -a_j}^{+a_j} (x_j - i_j).$$

To see f has the appropriate vanishing to lie in $\mathbf{I}(\mathbb{Z}^n \cap mP)$, note that for any $\mathbf{z} \in \mathbb{Z}^n$ where $f(\mathbf{z}) \neq 0$, each coordinate z_j with $j \in A \setminus B$ must lie in $\mathbb{Z} \setminus [-(a_j-1), +(a_j-1)]$, and hence have $|z_j| \geq a_j$; similarly, each coordinate z_j with $j \in B$ must lie in $\mathbb{Z} \setminus [-a_j, +a_j]$, and hence have $|z_j| \geq a_j + 1$. But this shows that $\mathbf{z} \notin \mathbb{Z}^n \cap mP$, since

$$|\mathbf{z}| = \sum_{j=1}^n |z_j| \geq \sum_{j \in A \setminus B} a_j + \sum_{j \in B} (a_j + 1) = |\mathbf{a}| + \#B > m.$$

On the other hand, $\mathbb{Z}^n \cap mP$ has a bijection to the *complementary* set of monomials

$$\mathbb{Z}^n \cap mP = \{\mathbf{z} \in \mathbb{Z}^n : \sum_{i=1}^n |z_i| \leq m\} \longleftrightarrow \{\mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B : |\mathbf{a}| + \#B \leq m\},$$

sending $\mathbf{z} \mapsto \mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B$ with $B := \{j : z_j < 0\}$ and

$$a_j := \begin{cases} z_j & \text{if } z_j \geq 0, \\ |z_j| - 1 & \text{if } z_j < 0. \end{cases}$$

The inverse bijection sends $\mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B \mapsto \mathbf{z}$ defined by $z_j := a_j$ for $j \notin B$ and $z_j = -(a_j + 1)$ for $j \in B$. This shows the inclusion $\text{gr } \mathbf{I}(\mathbb{Z}^n \cap mP) \supseteq I$ is an equality, since the surjection

$$R(\mathbb{Z}^n \cap mP) = S/\text{gr } \mathbf{I}(\mathbb{Z}^n \cap m \cdot P) \twoheadrightarrow S/I$$

must be an isomorphism via dimension-counting.

To calculate $E_P(t, q)$, it helps to recast the unique expressions in (77) as a graded \mathbb{R} -vector space isomorphism. Define the polynomial subalgebra $\mathbb{R}[\mathbf{x}^2] = \mathbb{R}[x_1^2, \dots, x_n^2]$ within $S = \mathbb{R}[\mathbf{x}]$, and introduce the \mathbb{R} -linear subspace $U \subset S$ which is spanned by the squarefree monomials, that is, $U := \text{span}_{\mathbb{R}}\{\mathbf{x}_B : B \subseteq [n]\}$. One then has the following graded \mathbb{R} -vector space isomorphism:

$$(78) \quad \begin{aligned} \mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U &\longrightarrow S \\ \mathbf{x}^{2\mathbf{a}} \otimes \mathbf{x}_B &\longmapsto \mathbf{x}^{2\mathbf{a}} \cdot \mathbf{x}_B. \end{aligned}$$

Endow the tensor product on the left with a *bigrading* or \mathbb{N}^2 -grading in which

$$\text{deg}_{\mathbb{N}^2}(\mathbf{x}^{2\mathbf{a}} \otimes \mathbf{x}_B) = (|\mathbf{a}|, \#B).$$

Then the bigraded Hilbert series in variables (q_1, q_2) that tracks $\mathbf{x}^{2\mathbf{a}} \otimes \mathbf{x}_B$ via $q_1^{|\mathbf{a}|} q_2^{\#B}$ is

$$\text{Hilb}_{\mathbb{N}^2}(\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U, q_1, q_2) = \left(\frac{1 + q_2}{1 - q_1} \right)^n.$$

To recover the *usual* \mathbb{N} -grading, tracking $\mathbf{x}^{2\mathbf{a}} \otimes \mathbf{x}_B$ via $q^{2|\mathbf{a}| + \#B}$, one must set $q_1 = q^2$ and $q_2 = q$.

On the other hand, our analysis above shows that the map (78), followed by the quotient map $S \twoheadrightarrow S/I \cong R(\mathbb{Z}^n \cap mP)$, induces a graded \mathbb{R} -vector space isomorphism

$$(79) \quad (\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U)_{\substack{\leq q_1^a q_2^b \text{ with} \\ a+b \leq m}} \longrightarrow R(\mathbb{Z}^n \cap mP).$$

Consequently, using Lemma 3.20 at the equality labeled (*) below, one has

$$\begin{aligned} E_P(t, q) &:= \sum_{m=0}^{\infty} t^m \cdot \text{Hilb}(R(\mathbb{Z}^n \cap mP), q) = \left[\sum_{m=0}^{\infty} t^m \cdot \text{trunc}_{\substack{\leq q_1^a q_2^b \text{ with} \\ a+b \leq m}} \text{Hilb}(\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U, q_1, q_2) \right]_{\substack{q_1 \mapsto q^2 \\ q_2 \mapsto q}} \\ &\stackrel{(*)}{=} \left[\frac{1}{1-t} \left(\frac{1 + tq_2}{1 - tq_1} \right)^n \right]_{\substack{q_1 \mapsto q^2 \\ q_2 \mapsto q}} = \frac{1}{1-t} \left(\frac{1 + tq}{1 - tq^2} \right)^n. \quad \square \end{aligned}$$

To enhance Proposition 3.26 to a B_n -equivariant calculation, one must recall the parametrization of simple $\mathbb{R}[B_n]$ -modules, e.g., from Geissinger and Kinch [19], Macdonald [34, Ch. 1, App. B]. The simple $\mathbb{R}[B_n]$ -modules $\{\mathbb{S}^{(\lambda^+, \lambda^-)}\}_{(\lambda^+, \lambda^-)}$ are indexed by *ordered pairs of partitions*

$$(\lambda^+, \lambda^-) \text{ where } |\lambda^+| = n_+, |\lambda^-| = n_- \text{ with } n_+ + n_- = n.$$

One can construct $\mathbb{S}^{(\lambda^+, \lambda^-)}$ using the simple $\mathbb{R}[\mathfrak{S}_n]$ -modules $\{\mathbb{S}^\mu\}_{\mu \vdash n}$ as follows. Introduce

- the operation of *induction*

$$(U_+, U_-) \longmapsto (U_+ \otimes U_-) \uparrow_{B_{n_+} \times B_{n_-}}^{B_n},$$

for any pairs U_+, U_- of $\mathbb{R}[B_{n_+}], \mathbb{R}[B_{n_-}]$ -modules,

- the operation of *inflation* $V \longmapsto V \uparrow$ of $\mathbb{R}[\mathfrak{S}_n]$ -modules V to $\mathbb{R}[B_n]$ -modules, by precomposing with the group quotient map $\pi : B_n \rightarrow \mathfrak{S}_n$ that ignores the \pm signs, and

- the one-dimensional character $\chi_{\pm} : B_n \rightarrow \{\pm 1\}$ sending a signed permutation matrix w to the product of its ± 1 signs, that is, $\chi_{\pm}(w) := \det(w) / \det(\pi(w))$.

Then starting with the simple $\mathbb{R}[\mathfrak{S}_n]$ -modules $\{\mathbb{S}^{\mu}\}$, one builds the $\mathbb{R}[B_n]$ -module $\mathbb{S}^{(\lambda^+, \lambda^-)}$ as follows:

$$\mathbb{S}^{(\lambda^+, \lambda^-)} := \left(\mathbb{S}^{\lambda^+} \uparrow \quad \otimes \quad \chi_{\pm} \otimes (\mathbb{S}^{\lambda^-} \uparrow) \right) \uparrow_{B_{n_+} \times B_{n_-}}^{B_n}.$$

Proposition 3.27. *For the cross-polytope $P := \diamond^n \subset \mathbb{R}^n$, one has in $\text{Rep}_{\mathbb{R}}[B_n][q][[t]]$*

$$E_P^{B_n}(t, q) = \frac{1}{(1-t) \cdot \prod_{i=1}^n (1-t^i q^{2i})} \left(\sum_{\lambda \vdash n} f^{\lambda}(tq^2) \cdot [\mathbb{S}^{(\lambda, \emptyset)}] \right) \left(\sum_{i=0}^n (tq)^i \cdot [\mathbb{S}^{((n-i), (i))}] \right)$$

Proof. The group B_n acts on $S = \mathbb{R}[\mathbf{x}]$ by permuting and negating the variables x_1, \dots, x_n . As the isomorphism (79) is also an isomorphism of graded $\mathbb{R}[B_n]$ -modules, the key is to calculate the class

$$[\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U]_{q_1, q_2} = [\mathbb{R}[\mathbf{x}^2]]_{q_1} \cdot [U]_{q_2}$$

within the representation ring $\text{Rep}_{\mathbb{R}}[B_n][[q_1, q_2]]$ of \mathbb{N}^2 -graded $\mathbb{R}[B_n]$ -modules.

For the right tensor factor $U = \text{span}_{\mathbb{R}}\{\mathbf{x}_B : B \subseteq [n]\} \subset S$, we claim that its i^{th} -graded component U_i carries the simple $\mathbb{R}[B_n]$ -module $\mathbb{S}^{((n-i), (i))}$. This is because it is a direct sum of the $\binom{n}{i}$ lines which are the B_n -images of the line $L := \mathbb{R} \cdot x_1 x_2 \cdots x_i$. This line L is stabilized setwise by the subgroup $B_{n-i} \times B_i$, where the B_{n-i} factor acts trivially and the B_i factor acts via χ_{\pm} . Hence

$$[U]_{q_2} = \sum_{i=0}^n q_2^i \cdot [\mathbb{S}^{((n-i), (i))}].$$

To analyze $\mathbb{R}[\mathbf{x}^2]$ as a $\mathbb{R}\mathfrak{S}[B_n]$ -module, note that since all variables x_i appear squared as x_i^2 , the sign changes in B_n have no effect; B_n acts via inflation through the surjection $\pi : B_n \rightarrow \mathfrak{S}_n$. Consequently, one can obtain the class of the graded $\mathbb{R}[B_n]$ -module $\mathbb{R}[\mathbf{x}^2]$ from that of the graded $\mathbb{R}[\mathfrak{S}_n]$ -module $S = \mathbb{R}[\mathbf{x}]$ given in Theorem 3.21, simply by applying the inflation map

$$[\mathbb{S}^{\lambda}] \mapsto [\mathbb{S}^{\lambda} \uparrow] = [\mathbb{S}^{(\lambda, \emptyset)}].$$

The upshot is that

$$[\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U]_{q_1, q_2} = [\mathbb{R}[\mathbf{x}^2]]_{q_1} \cdot [U]_{q_2} = \left(\frac{\sum_{\lambda \vdash n} f^{\lambda}(q_1) \cdot [\mathbb{S}^{(\lambda, \emptyset)}]}{\prod_{i=1}^n (1-q_1^i)} \right) \cdot \left(\sum_{i=0}^n q_2^i \cdot [\mathbb{S}^{((n-i), (i))}] \right).$$

Finally, as in the proof of Proposition 3.26, $E_P^{B_n}(t, q)$ is obtained from $[\mathbb{R}[\mathbf{x}^2] \otimes_{\mathbb{R}} U]_{q_1, q_2}$ upon multiplying by $\frac{1}{1-t}$ and replacing $q_1 \mapsto tq^2$ and $q_2 \mapsto tq$. \square

4. MINKOWSKI CLOSURE AND THE PROOF OF THEOREM 1.3

Recall from the Introduction that we hope to approach Conjecture 1.1 through the new *harmonic algebra* \mathcal{H}_P of a lattice polytope defined in Section 5 below. The existence of this algebra structure relies on Theorem 1.3, asserting that for finite point loci $\mathcal{Z}, \mathcal{Z}' \subset \mathbf{k}^n$ (such as $\mathcal{Z} = \mathbb{Z}^n \cap mP$ and $\mathcal{Z}' = \mathbb{Z}^n \cap m'P \subset \mathbb{R}^n$), their harmonic spaces always satisfy

$$V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z} + \mathcal{Z}'}$$

The goal of this section is to prove this result. The first two subsections collect some preparatory material. This includes broadening the notion of the perp I^{\perp} or Macaulay inverse system to allow not only homogeneous ideals $I \subset S = \mathbf{k}[\mathbf{x}]$, but arbitrary ideals.

4.1. Vanishing ideals for finite point loci. For $\mathcal{Z} \subset \mathbf{k}^n$ with \mathbf{k} a field, the *vanishing ideal* is

$$I(\mathcal{Z}) := \{f(\mathbf{x}) \in \mathbf{k}[\mathbf{x}] : f(\mathbf{z}) = 0 \text{ for all } \mathbf{z} \in \mathcal{Z}\}.$$

It is convenient to have a concrete generating set for $I(\mathcal{Z})$ for \mathcal{Z} finite, even if it has redundant generators. The following proposition is well-known, but perhaps hard to find in the literature.

Lemma 4.1. *For any field \mathbf{k} and finite subset $\mathcal{Z} \subset \mathbf{k}^n$, its vanishing ideal has these descriptions:*

$$(80) \quad I(\mathcal{Z}) = \bigcap_{\mathbf{z} \in \mathcal{Z}} (x_1 - z_1, \dots, x_n - z_n)$$

$$(81) \quad = \prod_{\mathbf{z} \in \mathcal{Z}} (x_1 - z_1, \dots, x_n - z_n)$$

$$(82) \quad = \left(\prod_{\mathbf{z} \in \mathcal{Z}} (x_{p(\mathbf{z})} - z_{p(\mathbf{z})}) : \text{all functions } p : \mathcal{Z} \rightarrow [n] \right)$$

Proof. When $\#\mathcal{Z} = 1$ so that $\mathcal{Z} = \{\mathbf{z}\}$, the assertions all hold, and $I(\{\mathbf{z}\})$ is the maximal ideal $(x_1 - z_1, \dots, x_n - z_n)$. From this, (80) follows by definition.

Equality (81) then follows by induction on $\#\mathcal{Z}$, once we check that the two ideals $I = I(\{\mathbf{z}\})$ and $I' = I(\mathcal{Z} \setminus \{\mathbf{z}\})$ are coprime in the sense that $1 \in I + I'$: this implies the inclusion $I \cap I' \supseteq I \cdot I'$ becomes an equality. Coprimeness of I, I' follows from *multivariate Lagrange interpolation* (see e.g. [27, top of p. 13]) which provides the existence of a polynomial $f(\mathbf{x})$ that has $f(\mathbf{z}) = 1$ and $f(\mathbf{z}') = 0$ for all $\mathbf{z}' \in \mathcal{Z} \setminus \{\mathbf{z}\}$, so $f(\mathbf{x}) \in I'$: in the quotient ring $\mathbf{k}[\mathbf{x}]/I \cong \mathbf{k}$, one has $1 \equiv f(\mathbf{x}) \pmod{I}$, and hence $1 \in I + I'$.

The equality (82) also follows by induction on $\#\mathcal{Z}$, since the product $I \cdot I'$ of two ideals generated as $I = (f_p)_{p=1,2,\dots}, I' = (f'_q)_{q=1,2,\dots}$ can be generated as $I \cdot I' = (f_p f'_q)_{p,q=1,2,\dots}$. \square

4.2. Harmonic spaces for inhomogeneous ideals: completions and exponentials. It will be much easier to identify the harmonic spaces $\mathbf{I}(\mathcal{Z})^\perp$ of the *inhomogeneous* ideals $\mathbf{I}(\mathcal{Z}) \subset S = \mathbb{R}[\mathbf{x}]$, rather than their homogeneous deformations $\text{gr } \mathbf{I}(\mathcal{Z})$. However, the harmonic spaces I^\perp for inhomogeneous ideals $I \subset \mathbf{k}[\mathbf{x}]$ naturally live in the power series *completion* $\mathbb{R}[[\mathbf{y}]]$ of $\mathbb{R}[\mathbf{y}]$, or more generally, a completion $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ of the divided power algebra $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$. Within these completions, the harmonic spaces $\mathbf{I}(\mathcal{Z})^\perp$ will turn out to have a simple basis of “exponentials” indexed by $\mathbf{z} \in \mathcal{Z}$; see Lemma 4.4 below.

Recall the set-up from Sections 2.2, 2.3 for harmonic spaces of homogeneous ideals in $S = \mathbf{k}[\mathbf{x}]$. For \mathbf{k} a characteristic zero field, consider the *polynomial algebra*

$$\mathbb{D} := \mathbf{k}[\mathbf{y}] = \mathbf{k}[y_1, \dots, y_n] \text{ with } \mathbf{k}\text{-basis } \{\mathbf{y}^{\mathbf{a}} : \mathbf{a} \in \mathbb{N}^n\}.$$

More generally, over any field \mathbf{k} or any commutative ring with 1, consider the divided power algebra

$$\mathbb{D} := \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \text{ with } \mathbf{k}\text{-basis } \{\mathbf{y}^{(\mathbf{a})} = y_1^{(a_1)} \cdots y_n^{(a_n)} : \mathbf{a} \in \mathbb{N}^n\}, \text{ where } \mathbf{y}^{(\mathbf{a})} \mathbf{y}^{(\mathbf{b})} = \prod_{i=1}^n \binom{a_i + b_i}{a_i} \mathbf{y}^{(\mathbf{a} + \mathbf{b})}.$$

making the identification $y_1^{(a_1)} \cdots y_n^{(a_n)} := \frac{\mathbf{y}^{\mathbf{a}}}{a_1! \cdots a_n!}$ whenever \mathbf{k} has characteristic zero. One has an S -module structure on \mathbb{D} given by $\odot : S \times \mathbb{D} \rightarrow \mathbb{D}$ where x_i acts on \mathbb{D} as the derivation $\frac{\partial}{\partial y_i}$ (so that $x_i \odot y_j = \delta_{ij}$), with these formulas in the characteristic zero and arbitrary ring cases:

$$\mathbf{x}^{\mathbf{a}} \odot \mathbf{y}^{\mathbf{b}} = \begin{cases} \prod_{i=1}^n \frac{b_i!}{(b_i - a_i)!} \cdot \mathbf{y}^{\mathbf{b} - \mathbf{a}} & \text{if } a_i \leq b_i \text{ for } i = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

$$\mathbf{x}^{\mathbf{a}} \odot \mathbf{y}^{(\mathbf{b})} = \begin{cases} \mathbf{y}^{(\mathbf{b}-\mathbf{a})} & \text{if } a_i \leq b_i \text{ for } i = 1, \dots, n, \\ 0 & \text{otherwise.} \end{cases}$$

This leads to a \mathbf{k} -bilinear pairing $\langle -, - \rangle : S \times \mathbb{D} \rightarrow \mathbf{k}$ defined by

$$(83) \quad \langle f(\mathbf{x}), g(\mathbf{y}) \rangle := \text{the constant term of } f \odot g.$$

This $\langle -, - \rangle$ pairs the \mathbf{k} -bases $\{\mathbf{x}^{\mathbf{a}}\}_{\mathbf{a} \in \mathbb{N}^n}$ and $\{\mathbf{y}^{(\mathbf{a})}\}_{\mathbf{a} \in \mathbb{N}^n}$ as $\langle \mathbf{x}^{\mathbf{a}}, \mathbf{y}^{(\mathbf{b})} \rangle = \delta_{\mathbf{a}, \mathbf{b}}$, giving a *perfect pairing* in each degree, identifying $\mathbb{D}_d \cong S_d^*$ and $\mathbb{D}_{\leq d} \cong S_{\leq d}^*$, for all $d \geq 0$. In other words, S, \mathbb{D} are what are sometimes called *graded* (or *restricted*) \mathbf{k} -duals.

When working with *inhomogeneous* ideals $I \subseteq S = \mathbf{k}[\mathbf{x}]$, we will include \mathbb{D} in a larger ring $\hat{\mathbb{D}} = \varprojlim \mathbb{D}_{\leq m}$ which is the inverse limit of the projections $\mathbb{D}/\mathbb{D}_{\geq m} \rightarrow \mathbb{D}/\mathbb{D}_{\geq m-1}$. In other words, $\hat{\mathbb{D}} = \{ \sum_{\mathbf{a} \in \mathbb{N}^n} c_{\mathbf{a}} \mathbf{y}^{(\mathbf{a})} : c_{\mathbf{a}} \in \mathbf{k} \}$ with multiplication defined \mathbf{k} -linearly extending the rule in (35). We can therefore extend the \odot pairing and S -module structure to $\hat{\mathbb{D}}$ to $\odot : S \times \hat{\mathbb{D}} \rightarrow \hat{\mathbb{D}}$ via $f \odot g := \partial(f)(g)$, and extend the \mathbf{k} -bilinear pairing

$$(84) \quad \langle -, - \rangle : S \times \hat{\mathbb{D}} \rightarrow \mathbf{k}$$

via the same formula in which $\langle f, g \rangle$ is the constant term of $f \odot g$. This pairing $\langle -, - \rangle$ identifies $\hat{\mathbb{D}}$ isomorphically with the (unrestricted) \mathbf{k} -linear dual space:

$$\begin{aligned} S^* := \text{Hom}_{\mathbf{k}}(S, \mathbf{k}) &\longrightarrow \hat{\mathbb{D}} \\ \varphi &\longmapsto \sum_{\mathbf{a} \in \mathbb{N}^n} \varphi(\mathbf{x}^{\mathbf{a}}) \cdot \mathbf{y}^{(\mathbf{a})}. \end{aligned}$$

As a consequence, for any subspace $W \subset S$, its *annihilator or perp space*

$$W^{\perp} := \{g(\mathbf{y}) \in \hat{\mathbb{D}} : \langle f(\mathbf{x}), g(\mathbf{y}) \rangle = 0 \text{ for all } f(\mathbf{x}) \in W\}$$

has a \mathbf{k} -linear isomorphism

$$\begin{aligned} W^{\perp} &\longrightarrow (S/W)^* := \text{Hom}_{\mathbf{k}}(S/W, \mathbf{k}) \\ g(\mathbf{y}) &\longmapsto \langle -, g(\mathbf{y}) \rangle \end{aligned}$$

In particular, when \mathbf{k} is a field and S/W is finite-dimensional, then so is W^{\perp} , since $W^{\perp} \cong (S/W)^*$.

Definition 4.2. For any field \mathbf{k} , and for any (possibly inhomogeneous) ideal $I \subseteq S = \mathbf{k}[\mathbf{x}]$, define its *harmonic space* as $I^{\perp} \subseteq \hat{\mathbb{D}} = \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y}) (= \mathbf{k}[[\mathbf{y}]] \text{ if } \mathbf{k} \supseteq \mathbb{Q})$.

The preceding discussion shows that, for \mathbf{k} a field and whenever $\dim_{\mathbf{k}} S/I$ is finite, one has

$$(85) \quad \dim_{\mathbf{k}} I^{\perp} = \dim_{\mathbf{k}} (S/I)^* = \dim_{\mathbf{k}} S/I.$$

We can now identify a simple \mathbf{k} -basis for the harmonic space of the inhomogeneous ideals $I(\mathcal{Z})$.

Definition 4.3. For any field \mathbf{k} , given $\mathbf{z} \in \mathbf{k}^n$, define the *exponential* in the completion $\hat{\mathbb{D}}$

$$\exp(\mathbf{z} \cdot \mathbf{y}) := \sum_{d=0}^{\infty} (z_1 y_1 + \dots + z_n y_n)^{(d)} = \sum_{d=0}^{\infty} \sum_{\substack{\mathbf{d}=(d_1, \dots, d_n): \\ \sum_i d_i = d}} \mathbf{z}^{\mathbf{d}} \mathbf{y}^{(\mathbf{d})},$$

where the right equality above used (41). It can be rewritten more familiarly when $\mathbf{k} \supseteq \mathbb{Q}$ as

$$\exp(\mathbf{z} \cdot \mathbf{y}) := \sum_{d=0}^{\infty} \frac{(z_1 y_1 + \cdots + z_n y_n)^d}{d!} = \sum_{d=0}^{\infty} \sum_{\substack{\mathbf{d}=(d_1, \dots, d_n): \\ \sum_i d_i = d}} \mathbf{z}^{\mathbf{d}} \frac{\mathbf{y}^{\mathbf{d}}}{d_1! \cdots d_n!}.$$

It is not hard to check that one has

$$(86) \quad \exp(\mathbf{z} \cdot \mathbf{y}) = \exp(z \cdot y_1^{(1)}) \cdots \exp(z \cdot y_n^{(1)})$$

where for $z \in \mathbf{k}$ and $y \in \hat{\mathbb{D}}_1$, one has

$$\exp(z \cdot y^{(1)}) := \sum_{d=0}^{\infty} z^d y^{(d)} \left(= \exp(z \cdot y) = \sum_{d=0}^{\infty} z^d \frac{y^d}{d!} \text{ if } \mathbf{k} \supseteq \mathbb{Q}. \right).$$

Lemma 4.4. *For \mathbf{k} a field and $\mathcal{Z} \subset \mathbf{k}^n$ a finite subset, the harmonic space $\mathbf{I}(\mathcal{Z})^\perp \subseteq \hat{\mathbb{D}}$ has \mathbf{k} -basis*

$$\{\exp(\mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\}.$$

Proof. Note that (85) shows that the set in the lemma has the correct size

$$\#\mathcal{Z} = \dim_{\mathbf{k}} S/\mathbf{I}(\mathcal{Z}) = \dim_{\mathbf{k}} \mathbf{I}(\mathcal{Z})^\perp.$$

Hence it suffices to check that its elements lie in $\mathbf{I}(\mathcal{Z})^\perp$, and are \mathbf{k} -linearly independent.

To check that $\exp(\mathbf{z} \cdot \mathbf{y}) \in \mathbf{I}(\mathcal{Z})^\perp$ for each $\mathbf{z} = (z_1, \dots, z_n) \in \mathcal{Z}$, by Lemma 4.1 it suffices to check that for any $p : \mathcal{Z} \rightarrow [n]$ one has $f_p(\mathbf{x}) \odot \exp(\mathbf{z} \cdot \mathbf{y}) = 0$ where

$$f_p(\mathbf{x}) = \prod_{\mathbf{z}' \in \mathcal{Z}} (x_{p(\mathbf{z}')} - z'_{p(\mathbf{z}')}) = \left(\prod_{\mathbf{z}' \in \mathcal{Z} \setminus \{\mathbf{z}\}} (x_{p(\mathbf{z}')} - z'_{p(\mathbf{z}')}) \right) \cdot (x_{p(\mathbf{z})} - z_{p(\mathbf{z})}).$$

Thus it suffices to check $(x_{p(\mathbf{z})} - z_{p(\mathbf{z})}) \odot \exp(\mathbf{z} \cdot \mathbf{y}) = 0$. Letting $i := p(\mathbf{z})$, one has

$$\begin{aligned} (x_{p(\mathbf{z})} - z_{p(\mathbf{z})}) \odot \exp(\mathbf{z} \cdot \mathbf{y}) &= (x_i - z_i) \odot \left(\exp(z_1 y_1^{(1)}) \cdots \exp(z_n y_n^{(1)}) \right) \\ &= \left(\exp(z_1 y_1^{(1)}) \cdots \widehat{\exp(z_i y_i^{(1)})} \cdots \exp(z_n y_n^{(1)}) \right) \cdot (x_i - z_i) \odot \exp(z_i y_i^{(i)}). \end{aligned}$$

Thus in the end it suffices to check $(x_i - z) \odot \exp(z y_i^{(1)}) = 0$ for $z \in \mathbf{k}$, which is not hard:

$$x_i \odot \exp(z y_i^{(1)}) = \sum_{d=0}^{\infty} x_i \odot \left(z^d y_i^{(d)} \right) = \sum_{d=1}^{\infty} z^d y_i^{(d-1)} = z \sum_{d=1}^{\infty} z^{d-1} y_i^{(d-1)} = z \exp(z y_i^{(1)}).$$

We next check that $\{\exp(\mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\}$ are \mathbf{k} -linearly independent within $\hat{\mathbb{D}}$. We first deal with the $n = 1$ case. When $n = 1$, if we let $N := \#\mathcal{Z}$, so that $\mathcal{Z} = \{z_1, \dots, z_N\} \subset \mathbf{k}$, we note that inside $\hat{\mathbb{D}} = \hat{\mathbb{D}}_{\mathbf{k}}(y)$, one has

$$\begin{aligned} \exp(z_1 \cdot y) &= 1 + z_1 y^{(1)} + z_1^2 y^{(2)} + \cdots + z_1^{N-1} y^{(N-1)} + \cdots \\ \exp(z_2 \cdot y) &= 1 + z_2 y^{(1)} + z_2^2 y^{(2)} + \cdots + z_2^{N-1} y^{(N-1)} + \cdots \\ &\vdots \\ \exp(z_N \cdot y) &= 1 + z_N y^{(1)} + z_N^2 y^{(2)} + \cdots + z_N^{N-1} y^{(N-1)} + \cdots \end{aligned}$$

These exponentials are \mathbf{k} -linearly independent because $\{1, y^{(1)}, y^{(2)}, \dots, y^{(N-1)}\}$ are \mathbf{k} -linearly independent and the Vandermonde matrix $(z_i^{j-1})_{i,j=1,2,\dots,N}$ is invertible.

When $n \geq 2$, it is helpful to note that the factorization (86) lets one identify $\exp(\mathbf{z} \cdot \mathbf{x})$ with the element $\exp(zy_1^{(1)}) \otimes \dots \otimes \exp(zy_n^{(1)})$ lying within the proper³ subspace

$$\hat{\mathbb{D}}_{\mathbf{k}}(y_1) \otimes \dots \otimes \hat{\mathbb{D}}_{\mathbf{k}}(y_n) \subsetneq \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y}).$$

Letting $p_i : \mathbf{k}^n \rightarrow \mathbf{k}$ for $i = 1, 2, \dots, n$ denote the coordinate projections, the $n = 1$ case proven above shows each subset $\{\exp(zy_i^{(1)})\}_{z \in p_i(\mathcal{Z})} \subset \hat{\mathbb{D}}_{\mathbf{k}}(y_i)$ is \mathbf{k} -linearly independent. Therefore the set

$$(87) \quad \{\exp(z_1 y_1^{(1)}) \otimes \dots \otimes \exp(z_n y_n^{(1)}) : \mathbf{z} = (z_1, \dots, z_n) \in p_1(\mathcal{Z}) \times \dots \times p_n(\mathcal{Z})\}$$

is \mathbf{k} -linearly independent inside $\hat{\mathbb{D}}_{\mathbf{k}}(y_1) \otimes \dots \otimes \hat{\mathbb{D}}_{\mathbf{k}}(y_n)$. Since the set in (87) contains

$$\{\exp(z_1 y_1^{(1)}) \otimes \dots \otimes \exp(z_n y_n^{(1)}) : \mathbf{z} = (z_1, \dots, z_n) \in \mathcal{Z}\}$$

as a subset, the latter must therefore also be \mathbf{k} -linearly independent subset, as desired. \square

Recall the familiar formula

$$(88) \quad \exp(\mathbf{z} \cdot \mathbf{y}) \cdot \exp(\mathbf{z}' \cdot \mathbf{y}) = \exp((\mathbf{z} + \mathbf{z}') \cdot \mathbf{y})$$

which holds in the power series ring $\hat{\mathbb{D}} = \mathbf{k}[[\mathbf{y}]]$; it is an easy exercise using (40) to check that it remains valid in $\hat{\mathbb{D}} = \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})$ for any field \mathbf{k} . Combining (88) with Lemma 4.4 immediately gives the following easier and more precise *inhomogeneous ideal* version of Theorem 1.3.

Proposition 4.5. *For any field \mathbf{k} and finite subsets $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$, one has inside $\hat{\mathbb{D}}$ that*

$$(89) \quad \mathbf{I}(\mathcal{Z})^\perp \cdot \mathbf{I}(\mathcal{Z}')^\perp = \mathbf{I}(\mathcal{Z} + \mathcal{Z}')^\perp.$$

Our goal, Theorem 1.3 is a similar assertion replacing $\mathbf{I}(\mathcal{Z})^\perp$ with $\text{gr } \mathbf{I}(\mathcal{Z})^\perp$, that is,

$$\underbrace{\text{gr } \mathbf{I}(\mathcal{Z})^\perp}_{V_{\mathcal{Z}}} \cdot \underbrace{\text{gr } \mathbf{I}(\mathcal{Z}')^\perp}_{V_{\mathcal{Z}'}} \subseteq \underbrace{\text{gr } \mathbf{I}(\mathcal{Z} + \mathcal{Z}')^\perp}_{V_{\mathcal{Z} + \mathcal{Z}'}}.$$

The next subsection will employ a proof strategy that considers the *rescaled point loci* $\epsilon \mathcal{Z} \subset \mathbf{k}^n$ for choices of ϵ in the field \mathbf{k} , and shows that in a precise sense, one has $\lim_{\epsilon \rightarrow 0} \mathbf{I}(\epsilon \mathcal{Z}) = \text{gr } \mathbf{I}(\mathcal{Z})$. One then uses this to deduce Theorem 1.3 from Proposition 4.5 by taking perps and employing limits and continuity arguments. However, we must be careful to perform these perp operations and limits within *finite-dimensional* \mathbf{k} -vector spaces like $S_{\leq N}$ and $\mathbb{D}_{\leq N} := \mathbb{D}_{\mathbf{k}}(\mathbf{y})_{\leq N} = \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})_{\leq N}$ for some $N \geq 0$, which are \mathbf{k} -linear duals under the perfect pairing $\langle -, - \rangle : S_{\leq N} \times \mathbb{D}_{\leq N} \rightarrow \mathbf{k}$. For this reason, we introduce these \mathbf{k} -linear projection operators for each $N \geq 0$

$$\begin{aligned} \pi_N : S &\longrightarrow S_{\leq N} \\ \pi_N : \hat{\mathbb{D}} &\longrightarrow \hat{\mathbb{D}}_{\leq N} \end{aligned}$$

that send $\mathbf{x}^{\mathbf{a}}, \mathbf{y}^{(\mathbf{a})} \mapsto 0$ whenever $|\mathbf{a}| > N$. Note some simple properties:

$$(90) \quad \langle f, g \rangle = \langle f, \pi_N(g) \rangle \text{ whenever } f \in S_{\leq N},$$

$$(91) \quad \pi_N(g \cdot g') = \pi_N(\pi_N g \cdot \pi_N g') \text{ for any } g, g' \in \hat{\mathbb{D}}.$$

This leads to the next lemma, a finite-dimensional version of Lemma 4.4, Proposition 4.5.

Lemma 4.6. *Consider two finite subsets $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$.*

³Even in characteristic zero, one has $\mathbf{k}[[y_1]] \otimes \dots \otimes \mathbf{k}[[y_n]] \subsetneq \mathbf{k}[[\mathbf{y}]]$, without forming a *completed tensor product*.

(i) Whenever $N \geq \#\mathcal{Z} - 1$, the subspace $\pi_N(\mathbf{I}(\mathcal{Z})^\perp) \subseteq \mathbb{D}_{\leq N} = \mathbb{D}_{\mathbf{k}}(\mathbf{y})_{\leq N}$ has \mathbf{k} -basis

$$\{\pi_N \exp(\mathbf{z} \cdot \mathbf{y})\}_{\mathbf{z} \in \mathcal{Z}}.$$

(ii) Whenever $N \geq \#(\mathcal{Z} + \mathcal{Z}') - 1$,

$$\pi_N(\pi_N(\mathbf{I}(\mathcal{Z})^\perp) \cdot \pi_N(\mathbf{I}(\mathcal{Z}')^\perp)) = \pi_N(\mathbf{I}(\mathcal{Z} + \mathcal{Z}')^\perp).$$

Proof. Since $\mathbf{I}(\mathcal{Z})^\perp$ has \mathbf{k} -basis $\{\exp(\mathbf{z} \cdot \mathbf{y})\}_{\mathbf{z} \in \mathcal{Z}}$ by Proposition 4.5, it suffices to show that the set in assertion (i) is \mathbf{k} -linearly independent whenever $N \geq \#\mathcal{Z} - 1$. Note that, in proving this, we may assume that \mathbf{k} is infinite, by passing to an extension field of \mathbf{k} , such as its algebraic closure. After doing so, there exists a change of coordinates acting by an element of $GL_n(\mathbf{k})$ on \mathbf{k}^n so that the points $\mathbf{z} \in \mathcal{Z}$ all have *distinct* first coordinates z_1 . Letting $d := \#\mathcal{Z}$, so $N \geq d - 1$, consider the expansion coefficients of each $\pi_N \exp(\mathbf{z} \cdot \mathbf{y})$ with $\mathbf{z} = (z_1, \dots, z_n)$ only on the basis elements $1, y_1^{(1)}, y_1^{(2)}, \dots, y_1^{(d-1)}$ in $\mathbb{D}_{\leq N}$:

$$\pi_N \exp(\mathbf{z} \cdot \mathbf{y}) = \pi_N \exp(z_1 y_1^{(1)} + \dots + z_n y_n^{(1)}) = 1 + z_1 y_1^{(1)} + z_1^2 y_1^{(2)} + \dots + z_1^{d-1} y_1^{(d-1)} + \dots$$

As one varies over all $\mathbf{z} \in \mathcal{Z}$, the invertibility of the Vandermonde matrix $(z_1^j)_{\substack{\mathbf{z}=(z_1, \dots, z_n) \in \mathcal{Z} \\ j=1, 2, \dots, d}}$ again shows that these elements are \mathbf{k} -linearly independent. This proves assertion (i).

For assertion (ii), first note that when both $\mathcal{Z}, \mathcal{Z}' \neq \emptyset$, one always has $\#(\mathcal{Z} + \mathcal{Z}') \geq \#\mathcal{Z}, \#\mathcal{Z}'$. Thus the hypothesis $N \geq \#(\mathcal{Z} + \mathcal{Z}') - 1$ in (ii) allows one to apply (i) to both $\mathcal{Z}, \mathcal{Z}'$. Assertion (ii) then follows from this calculation using first (91) and then (88):

$$\pi_N(\pi_N \exp(\mathbf{z} \cdot \mathbf{y}) \cdot \pi_N \exp(\mathbf{z}' \cdot \mathbf{y})) = \pi_N(\exp(\mathbf{z} \cdot \mathbf{y}) \cdot \exp(\mathbf{z}' \cdot \mathbf{y})) = \pi_N(\exp((\mathbf{z} + \mathbf{z}') \cdot \mathbf{y})). \quad \square$$

Lemma 4.6 has another consequence that we will need.

Lemma 4.7. *For a finite subset $\mathcal{Z} \subset \mathbf{k}^n$ and for all $N \geq \#\mathcal{Z} - 1$, one has*

$$\pi_N(\mathbf{I}(\mathcal{Z})^\perp) = (\mathbf{I}(\mathcal{Z})_{\leq N})^\perp \cap \mathbb{D}_{\leq N}.$$

Proof. First let us check that for any $N \geq 0$ one has an inclusion from the left into the right side

$$(92) \quad \pi_N(\mathbf{I}(\mathcal{Z})^\perp) \subseteq (\mathbf{I}(\mathcal{Z})_{\leq N})^\perp \cap \mathbb{D}_{\leq N}.$$

Given any $g(\mathbf{y})$ in $\mathbf{I}(\mathcal{Z})^\perp$, its projection $\pi_N(g(\mathbf{y}))$ lies in $\mathbb{D}_{\leq N}$ by definition. But it also lies in $(\mathbf{I}(\mathcal{Z})_{\leq N})^\perp$, since for any $f(\mathbf{x}) \in \mathbf{I}(\mathcal{Z})_{\leq N}$, one has by (90) that $\langle f(\mathbf{x}), \pi_N(g(\mathbf{y})) \rangle = \langle f(\mathbf{x}), g(\mathbf{y}) \rangle = 0$.

When $N \geq \#\mathcal{Z} - 1$, we claim that the inclusion (92) must be an equality by dimension-counting. Since the \mathbf{k} -bilinear pairing $\langle -, - \rangle : S_{\leq N} \times \mathbb{D}_{\leq N} \rightarrow \mathbf{k}$ is perfect, one always has

$$\dim_{\mathbf{k}}(\mathbf{I}(\mathcal{Z})_{\leq N})^\perp \cap \mathbb{D}_{\leq N} = \dim_{\mathbf{k}} S_{\leq N} / \mathbf{I}(\mathcal{Z})_{\leq N} = \dim_{\mathbf{k}} S / \mathbf{I}(\mathcal{Z}) = \#\mathcal{Z},$$

where the second-to-last-inequality used Lemma 2.2. On the other hand, when $N \geq \#\mathcal{Z} - 1$, one also has from Lemma 4.6(i) that $\dim_{\mathbf{k}} \pi_N(\mathbf{I}(\mathcal{Z})^\perp) = \#\mathcal{Z}$. \square

4.3. Proof of Theorem 1.3. Recall the statement of the theorem.

Theorem 1.3 *For any pair of finite point loci $\mathcal{Z}, \mathcal{Z}'$ in \mathbf{k}^n over any field \mathbf{k} , one has*

$$V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z} + \mathcal{Z}'}$$

The proof ideas in the previous subsection suggest considering $\epsilon \mathcal{Z}$ for $\epsilon \in \mathbf{k}$ and continuity arguments with limits as $\epsilon \rightarrow 0$. To make sense of this, we will work in the following context.

Definition 4.8. A *metric topological field* is a field \mathbf{k} equipped with a metric $d(\cdot, \cdot) : \mathbf{k} \times \mathbf{k} \rightarrow \mathbb{R}_{\geq 0}$ whose field operations $+, \times : \mathbf{k} \times \mathbf{k} \rightarrow \mathbf{k}$ and $x \mapsto -x$ on \mathbf{k} , and $x \mapsto x^{-1}$ on \mathbf{k}^\times are continuous in the metric topology.

We will also need to take limits as ϵ approaches 0, so we will need the existence of nonzero sequences $\epsilon_1, \epsilon_2, \dots \in \mathbf{k}^\times$ that approach 0, meaning $\lim_{i \rightarrow \infty} d(\epsilon_i, 0) = 0$.

Example 4.9. The field $\mathbf{k} = \mathbb{R}$ with metric $d(z, z') = \|z - z'\|$ from its usual norm $\|\cdot\|$ is a metric topological field, with some nonzero sequences approaching zero, e.g., $\epsilon, \epsilon^2, \epsilon^3, \dots$ for $0 < \epsilon < 1$.

For the reader bothered by the metric field baggage, one can safely work throughout this proof with $\mathbf{k} = \mathbb{R}$, and remember that $\mathbb{D}_{\mathbb{R}}(\mathbf{y}) = \mathbb{R}[\mathbf{y}]$ via the identification of $y^{(d)} := \frac{y^d}{d!}$.

Example 4.10. For any field \mathbf{k} , the field of (*bounded below*) *Laurent series* in an indeterminate ϵ

$$\mathbf{k}((\epsilon)) := \left\{ f(\epsilon) = \sum_{m=-\infty}^{\infty} c_m \cdot \epsilon^m : c_m \in \mathbf{k}, c_m = 0 \text{ for } m \ll 0 \right\}$$

is a metric topological field. Its metric $d(z, z') = \|z - z'\|$ comes from an *ultrametric ϵ -adic norm* $\|\cdot\|$ defined as follows: for $f(\epsilon) = \sum_{m=-N}^{+\infty} c_m \epsilon^m$ with $c_{-N} \in \mathbf{k}^\times$, define $\|f(\epsilon)\| := 2^N$. Note that $\mathbf{k}((\epsilon))$ contains some nonzero sequences approaching zero, e.g., $\epsilon, \epsilon^2, \epsilon^3, \dots$

Example 4.10 shows that every field \mathbf{k} has an extension $K = \mathbf{k}((t))$ which is a metric topological field K having nonzero sequences that approach zero. This is helpful, due to the following fact.

Proposition 4.11. *Theorem 1.3 holds for a field \mathbf{k} if it holds for some field extension $K \supseteq \mathbf{k}$.*

Proof. For $\mathcal{Z} \subseteq \mathbf{k}^n \subseteq K^n$, use a field superscript to distinguish vanishing ideals, harmonic spaces

$$\begin{aligned} \mathbf{I}^{\mathbf{k}}(\mathcal{Z}) &\subseteq \mathbf{k}[\mathbf{x}] \\ \mathbf{I}^K(\mathcal{Z}) &\subseteq K[\mathbf{x}] \\ V_{\mathcal{Z}}^{\mathbf{k}} &= \text{gr } \mathbf{I}^{\mathbf{k}}(\mathcal{Z})^\perp \subseteq \mathbb{D}_{\mathbf{k}}(\mathbf{y}) \\ V_{\mathcal{Z}}^K &= \text{gr } \mathbf{I}^K(\mathcal{Z})^\perp \subseteq \mathbb{D}_K(\mathbf{y}). \end{aligned}$$

One can check that

$$V_{\mathcal{Z}}^{\mathbf{k}} = \text{gr } \mathbf{I}^{\mathbf{k}}(\mathcal{Z})^\perp = \text{gr } \mathbf{I}^K(\mathcal{Z})^\perp \cap \mathbb{D}_{\mathbf{k}}(\mathbf{y}) = V_{\mathcal{Z}}^K \cap \mathbb{D}_{\mathbf{k}}(\mathbf{y}).$$

Now assume Theorem 1.3 for K . For two finite loci $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n \subseteq K^n$, and $g(\mathbf{y}), g'(\mathbf{y}) \in V_{\mathcal{Z}}^{\mathbf{k}}, V_{\mathcal{Z}'}^{\mathbf{k}}$, respectively, regard them as lying in $V_{\mathcal{Z}}^K, V_{\mathcal{Z}'}^K$. Hence $g(\mathbf{y}) \cdot g'(\mathbf{y})$ lies in $V_{\mathcal{Z}+\mathcal{Z}'}^K$, by our assumption on K . However $g(\mathbf{y}) \cdot g'(\mathbf{y})$ also lies in $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$, and hence it lies in $V_{\mathcal{Z}+\mathcal{Z}'}^K \cap \mathbb{D}_{\mathbf{k}}(\mathbf{y}) = V_{\mathcal{Z}+\mathcal{Z}'}^{\mathbf{k}}$. \square

We next make precise how the inhomogeneous ideals $\mathbf{I}(\epsilon\mathcal{Z})$ will approach $\text{gr } \mathbf{I}(\mathcal{Z})$ as $\epsilon \rightarrow 0$.

Definition 4.12. Let \mathbf{k} be a metric topological field.

- Say $\{\mathbf{z}^\ell\}_{\ell=1,2,\dots}$ in \mathbf{k}^n *approach \mathbf{z}^∞ entrywise*, written $\{\mathbf{z}^\ell\} \xrightarrow{\text{ew}} \mathbf{z}^\infty$, if $\mathbf{z}_i^\ell \rightarrow \mathbf{z}_i^\infty$ for all i .
- For matrices $\{A^\ell\}_{\ell=1,2,\dots}$ in $\mathbf{k}^{m \times n}$, say $\{A^\ell\} \xrightarrow{\text{ew}} A^\infty$, if $A_{ij}^\ell \rightarrow A_{ij}^\infty$ for all i, j .
- Say \mathbf{k} -linear subspaces $\{U^\ell\}_{\ell=1,2,\dots}$ of \mathbf{k}^n *approach U^∞ strongly* if there exists a basis $\{\mathbf{u}_j^\infty\}_{j=1,\dots,m}$ for U^∞ and bases $\{\mathbf{u}_j^\ell\}_{j=1,\dots,m}$ for U^ℓ , having $\mathbf{u}_j^\ell \xrightarrow{\text{ew}} \mathbf{u}_j^\infty$ for each $j = 1, \dots, m$.

We point out one consequence of having subspaces $\{U^\ell\}_{\ell=1,2,\dots}$ of \mathbf{k}^n approach U^∞ strongly: for every vector \mathbf{u} in U^∞ , one can exhibit a sequence of vectors $\mathbf{u}^\ell \in U^\ell$ such that $\mathbf{u}^\ell \xrightarrow{\text{ew}} \mathbf{u}$: simply express $\mathbf{u} = \sum_{j=1}^m c_j \mathbf{u}_j^\infty$ uniquely in the given basis for U^∞ , and then let $\mathbf{u}^\ell := \sum_{j=1}^m c_j \mathbf{u}_j^\ell$.

We will also need some properties of *homogenization* of polynomials.

Definition 4.13. Recall the operation of *homogenization* with respect to a variable ϵ for an inhomogeneous polynomial $g(\mathbf{x}) = \sum_{m=0}^{\deg(g)} g_m(\mathbf{x})$ having homogeneous components $g_m(\mathbf{x})$:

$$g^\epsilon(\mathbf{x}) := \epsilon^{\deg(g)} g(\mathbf{x}/\epsilon) = \epsilon^{\deg(g)} g\left(\frac{x_1}{\epsilon}, \dots, \frac{x_n}{\epsilon}\right) = \tau(g(\mathbf{x})) + \sum_{m=0}^{\deg(g)-1} \epsilon^{\deg(g)-m} g_m(\mathbf{x}).$$

Note these two properties, when one specializes the homogenization variable ϵ to values in \mathbf{k}^\times :

$$(93) \quad \tau(g(\mathbf{x})) = \tau(g^\epsilon(\mathbf{x})) = \epsilon^{\deg(g)} \tau(g(\mathbf{x}/\epsilon)) \text{ for } \epsilon \in \mathbf{k}^\times,$$

$$(94) \quad \{g^{\epsilon_\ell}(\mathbf{x})\}_{\ell=1,2,\dots} \rightarrow \tau(g(\mathbf{x})) \text{ coefficientwise whenever } \epsilon_1, \epsilon_2, \dots \in \mathbf{k}^\times \text{ approach } 0.$$

Lemma 4.14. Fix a field \mathbf{k} and $I \subset S = \mathbf{k}[\mathbf{x}]$ a (possibly inhomogeneous) ideal.

(i) For ϵ in \mathbf{k}^\times , each of the rescaled ideals

$$I^\epsilon := (f(\mathbf{x}/\epsilon) : f(\mathbf{x}) \in I) = (f^\epsilon(\mathbf{x}) : f(\mathbf{x}) \in I)$$

inside $S = \mathbf{k}[\mathbf{x}]$ has the same associated homogeneous ideal $\text{gr } I^\epsilon = \text{gr } I$.

(ii) If $\epsilon_1, \epsilon_2, \dots$ in \mathbf{k}^\times approach 0 in a metric topological field \mathbf{k} , then for $N \geq 0$, one has \mathbf{k} -bases

$$\begin{aligned} &\{h_j^{\epsilon_\ell}(\mathbf{x})\}_{j=1,2,\dots,r} \text{ for the subspaces } I_{\leq N}^{\epsilon_\ell}, \\ &\{h_j(\mathbf{x})\}_{j=1,2,\dots,r} \text{ for the subspace } (\text{gr } I)_{\leq N}, \end{aligned}$$

with each sequence $\{h_j^{\epsilon_\ell}(\mathbf{x})\}$ coefficientwise approaching $h_j(\mathbf{x})$ for $j = 1, 2, \dots, r$.

In other words, inside $S_{\leq N}$, the subspaces $\{I_{\leq N}^{\epsilon_\ell}\}$ strongly approach the subspace $(\text{gr } I)_{\leq N}$.

Proof. Assertion (i) follows from (93). For assertion (ii), apply Proposition 2.7 to I , and produce a Gröbner basis G for I such that $\tau(G) := \{\tau(g(\mathbf{x})) : g(\mathbf{x}) \in G\}$ is a Gröbner basis for $\text{gr } I$. Since $(\text{gr } I)_{\leq N}$ is \mathbf{k} -spanned by the products $\{\mathbf{x}^{\mathbf{a}} \cdot \tau(g(\mathbf{x})) : |\mathbf{a}| + \deg(g) \leq N\}$, one can find a subset $\{h_j(\mathbf{x})\}_{j=1,2,\dots,r}$ giving a \mathbf{k} -basis for $(\text{gr } I)_{\leq N}$, where $r = \dim_{\mathbf{k}}(\text{gr } I)_{\leq N}$.

Now for each of these $h_j(\mathbf{x}) = \mathbf{x}^{\mathbf{a}} \cdot \tau(g(\mathbf{x}))$, if one defines $h_j^\epsilon(\mathbf{x}) := \mathbf{x}^{\mathbf{a}} \cdot g^\epsilon(\mathbf{x})$, one finds that $h_j^\epsilon(\mathbf{x})$ lies in $I_{\leq N}^\epsilon$ and $\tau(h_j^\epsilon(\mathbf{x})) = h_j(\mathbf{x})$. Picking $\epsilon_1, \epsilon_2, \dots$ approaching 0, the elements $\{h_j^{\epsilon_\ell}(\mathbf{x})\}$ coefficientwise approach $h_j(\mathbf{x})$ for $j = 1, 2, \dots, r$ by (94). This means that for ℓ sufficiently large, the sets $\{h_j^{\epsilon_\ell}(\mathbf{x})\}_{j=1,2,\dots,r}$ are \mathbf{k} -linearly independent, as they approach a \mathbf{k} -linearly independent set. But then they must give \mathbf{k} -bases for $I_{\leq N}^{\epsilon_\ell}$, since they each have the correct cardinality r :

$$\begin{aligned} r = \dim_{\mathbf{k}}(\text{gr } I)_{\leq N} &= \dim_{\mathbf{k}} S_{\leq N} - \dim_{\mathbf{k}} S_{\leq N}/(\text{gr } I)_{\leq N} \\ &= \dim_{\mathbf{k}} S_{\leq N} - \dim_{\mathbf{k}}(S/\text{gr } I)_{\leq N} \\ &= \dim_{\mathbf{k}} S_{\leq N} - \dim_{\mathbf{k}} S_{\leq N}/I_{\leq N}^{\epsilon_\ell} \quad \text{using (24)} \\ &= \dim_{\mathbf{k}} I_{\leq N}^{\epsilon_\ell}. \quad \square \end{aligned}$$

We will need that whenever a sequence of subspaces strongly approaches another subspace, the same holds for their perp spaces. For this, the following lemma is useful. One can regard it as saying the map $U \mapsto U^\perp$ between dual Grassmannians $\text{Gr}_r(\mathbf{k}^n) \rightarrow \text{Gr}_{n-r}((\mathbf{k}^n)^*)$ is continuous in the metric topology, whenever \mathbf{k} is a metric topological field; the idea is to use the duality between Plücker coordinates within an affine open chart. For this purpose, recall that an r -dimensional

subspace U of \mathbf{k}^n can be represented as the row space $\text{row}(A)$ for a matrix A in $\mathbf{k}^{r \times n}$ of full rank r . Also, identify $(\mathbf{k}^n)^* \cong \mathbf{k}^n$ via the standard \mathbf{k} -bilinear form $\langle \mathbf{z}, \mathbf{z}' \rangle := \sum_{i=1}^n z_i z'_i$.

Lemma 4.15. *Let \mathbf{k} be a metric topological field, and assume one has matrices A_1, A_2, \dots in $\mathbf{k}^{r \times n}$ approaching entrywise a matrix A_∞ of full rank r . Then there exists L sufficiently large and matrices B_L, B_{L+1}, \dots in $\mathbf{k}^{(n-r) \times n}$ of full rank $n-r$, entrywise approaching a matrix B_∞ , with*

$$\begin{aligned} \text{row}(B_\ell) &= \text{row}(A_\ell)^\perp \text{ for } \ell = L, L+1, \dots, \\ \text{row}(B_\infty) &= \text{row}(A_\infty)^\perp. \end{aligned}$$

Proof. By re-indexing the coordinates in \mathbf{k}^n , one may assume that the $r \times r$ submatrix P_∞ in the first r columns of A_∞ has full rank r . Then letting P_i denote the analogous submatrix of A_i , one must have P_L, P_{L+1}, \dots of full rank r for some sufficiently large L . Denoting the $r \times r$ identity matrix I_r , then for $\ell \geq L$ the matrices $P_\ell^{-1} A_\ell$ have the form $P_\ell^{-1} A_\ell = (I_r | C_\ell)$, and we claim that they entrywise approach the matrix $P_\infty^{-1} A_\infty = (I_r | C_\infty)$; the formula $P^{-1} = \frac{1}{\det P} \cdot \text{adj} P$ shows the entries of $P_\ell^{-1} A_\ell$ are rational functions of entries of A_ℓ , and ones whose denominator never vanishes.

Defining the matrices $B_\ell := (-C_\ell^t | I_{n-r})$, for $\ell = L, L+1, \dots$ and $B_\infty := (-C_\infty^t | I_{n-r})$, one can check that they have the desired relations $\text{row}(B_\ell) = \text{row}(A_\ell)^\perp$ for $\ell = L, L+1, \dots$, and $\text{row}(B_\infty) = \text{row}(A_\infty)^\perp$. One also has B_L, B_{L+1}, \dots entrywise approaching B_∞ , by construction. \square

Corollary 4.16. *Let \mathbf{k} be a metric topological field, and $\langle -, - \rangle : \mathbf{k}^n \times \mathbf{k}^n \rightarrow \mathbf{k}$ a perfect \mathbf{k} -bilinear pairing. Then for any \mathbf{k} -subspaces $\{U^\ell\}$ approaching U^∞ strongly within \mathbf{k}^n , one has the same for their perps: $\{(U^\ell)^\perp\}$ approach $(U^\infty)^\perp$ strongly within \mathbf{k}^n .*

Proof. Let $r = \dim_{\mathbf{k}} U^\infty = \dim_{\mathbf{k}} U^\ell$ for all ℓ . Apply Lemma 4.15 to a sequence of matrices A_1, A_2, \dots in $\mathbf{k}^{r \times n}$ whose rows are \mathbf{k} -bases for U^1, U^2, \dots that approach a \mathbf{k} -basis for U^∞ . \square

Lastly, we identify harmonic spaces $V_{\mathcal{Z}}$ of finite loci \mathcal{Z} as limits of better understood subspaces.

Lemma 4.17. *Let $\mathcal{Z} \subseteq \mathbf{k}^n$ be finite, and $N \geq \#\mathcal{Z} - 1$.*

- (i) *Then $V_{\mathcal{Z}} := \text{gr } \mathbf{I}(\mathcal{Z})^\perp$ within $\mathbb{D} := \mathbb{D}_{\mathbf{k}}(\mathbf{y})$ actually lies within $\mathbb{D}_{\leq N}$.*
- (ii) *Furthermore, if \mathbf{k} is a metric topological field with $\epsilon_1, \epsilon_2, \dots$ in \mathbf{k}^\times approaching 0, then the subspaces $\{\pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z})^\perp)\}$ strongly approach $V_{\mathcal{Z}}$ within $\mathbb{D}_{\leq N}$.*

Proof. For assertion (i), note Lemma 2.2 shows $(S/\text{gr } \mathbf{I}(\mathcal{Z}))_m = 0$ in degrees $m \geq \#\mathcal{Z}$. As (46) shows $V_{\mathcal{Z}} := \text{gr } \mathbf{I}(\mathcal{Z})^\perp$ and $S/\text{gr } \mathbf{I}(\mathcal{Z})$ are graded duals as \mathbf{k} -vector spaces, the same holds for $V_{\mathcal{Z}}$.

For assertion (ii), note that for any $\epsilon \in \mathbf{k}^\times$, one has that $\mathbf{I}(\epsilon \mathcal{Z})$ is the same as the rescaled ideal I^ϵ for $I := \mathbf{I}(\mathcal{Z})$. Therefore, Lemma 4.14 shows that inside $S_{\leq N}$, the subspaces $\{\mathbf{I}(\epsilon_\ell \mathcal{Z})_{\leq N}\}$ strongly approach the subspace $\text{gr } \mathbf{I}(\mathcal{Z})_{\leq N}$. Applying Corollary 4.16 one concludes that inside $\mathbb{D}_{\leq N}$,

$$\text{the subspaces } \{\mathbf{I}(\epsilon_\ell \mathcal{Z})_{\leq N}^\perp\} \text{ strongly approach the subspace } (\text{gr } \mathbf{I}(\mathcal{Z})_{\leq N})^\perp.$$

But assertion (i) here shows $(\text{gr } \mathbf{I}(\mathcal{Z})_{\leq N})^\perp = \text{gr } \mathbf{I}(\mathcal{Z})^\perp = V_{\mathcal{Z}}$. And Lemma 4.7 applied to each $\epsilon_\ell \mathcal{Z}$ shows that, for $N \geq \#\mathcal{Z} - 1 = \#(\epsilon_\ell \mathcal{Z}) - 1$, within $\mathbb{D}_{\leq N}$ one has $(\mathbf{I}(\epsilon_\ell \mathcal{Z})_{\leq N})^\perp = \pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z})^\perp)$. \square

We can now prove the theorem.

Theorem 1.3 *For any pair of finite point loci $\mathcal{Z}, \mathcal{Z}'$ in \mathbf{k}^n over any field \mathbf{k} , one has*

$$V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z} + \mathcal{Z}'}$$

Proof. Using Proposition 4.11 and Example 4.10, we may assume without loss of generality that \mathbf{k} is a metric topological field, containing sequences $\epsilon_1, \epsilon_2, \dots$ in \mathbf{k}^\times that approach 0. We may also assume without loss of generality that both $\mathcal{Z}, \mathcal{Z}'$ are nonempty, else there is nothing to prove.

We wish to choose N sufficiently large so that all of our various lemmas will apply to $\mathcal{Z}, \mathcal{Z}', \mathcal{Z} + \mathcal{Z}'$; it turns out that $N \geq \max\{\#(\mathcal{Z} + \mathcal{Z}') - 1, \#\mathcal{Z} + \#\mathcal{Z}' - 2\}$ will suffice.

Assume $g = g(\mathbf{y})$ in $V_{\mathcal{Z}}$ and $g' = g'(\mathbf{y})$ in $V_{\mathcal{Z}'}$. We wish to show that their product $g \cdot g'$ lies in $V_{\mathcal{Z} + \mathcal{Z}'}$. Without loss of generality, both $g, g' \neq 0$, and hence by Lemma 4.17, one has

$$(95) \quad \deg(g) \leq \#\mathcal{Z} - 1 \text{ and } \deg(g') \leq \#\mathcal{Z}' - 1.$$

Using Lemma 4.17, find sequences

$$\begin{aligned} g_1, g_2, \dots \text{ with } g_\ell \in \pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z})^\perp), \text{ and } \lim g_\ell = g \\ g'_1, g'_2, \dots \text{ with } g'_\ell \in \pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z}')^\perp) \text{ and } \lim g'_\ell = g'. \end{aligned}$$

Then one has the following chain of equalities:

$$(96) \quad g \cdot g' = \pi_N(g \cdot g') \quad (\text{ as (95) gives } \deg(g \cdot g') \leq (\#\mathcal{Z} - 1) + (\#\mathcal{Z}' - 1) \leq N)$$

$$(97) \quad = \pi_N(\lim g_\ell \cdot \lim g'_\ell) \quad (\text{ by our choice of } g_\ell, g'_\ell)$$

$$(98) \quad = \pi_N(\lim(g_\ell \cdot g'_\ell)) \quad (\text{ by continuity of multiplication } \mathbb{D}_{\leq N} \times \mathbb{D}_{\leq N} \rightarrow \mathbb{D}_{\leq 2N})$$

$$(99) \quad = \lim \pi_N(g_\ell \cdot g'_\ell) \quad (\text{ by continuity of } \pi_N : \mathbb{D}_{\leq 2N} \rightarrow \mathbb{D}_{\leq N}).$$

For each ℓ , one has

$$\pi_N(g_\ell \cdot g'_\ell) \in \pi_N(\pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z})^\perp) \cdot \pi_N(\mathbf{I}(\epsilon_\ell \mathcal{Z}')^\perp)) = \pi_N(\mathbf{I}(\epsilon_\ell(\mathcal{Z} + \mathcal{Z}'))^\perp)$$

where the last equality used Lemma 4.6 part (ii). Since $g \cdot g' = \lim \pi_N(g_\ell \cdot g'_\ell)$, applying Lemma 4.17 one more time shows that $g \cdot g'$ lies in $V_{\mathcal{Z} + \mathcal{Z}'}$, as desired. \square

We conclude this section with a few remarks. The first are some interesting examples of Theorem 1.3, including two small examples with point loci in \mathbb{R}^2 . Another example discusses the relation between Theorem 1.3 for point loci in \mathbf{k}^1 , and the *additive combinatorics of sumsets* and the *Cauchy-Davenport Theorem*. Following these examples, we remark on a shorter proof of Theorem 1.3 which avoids topology, and also on a similar-sounding result of F. Gundlach.

Example 4.18. One can easily have a *proper* inclusion $V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subsetneq V_{\mathcal{Z} + \mathcal{Z}'}$. E.g., inside \mathbb{R}^2 take

$$\begin{aligned} \mathcal{Z} &= \{(0, 0), (1, 0), (0, 1)\}, \\ \mathcal{Z}' &= \{(0, 0), (1, 0), (1, 1)\}, \\ \mathcal{Z} + \mathcal{Z}' &= \{(0, 0), (1, 0), (0, 1), (1, 1), (2, 0), (2, 1), (1, 2)\}. \end{aligned}$$

Although $\mathcal{Z} \neq \mathcal{Z}'$, the two loci $\mathcal{Z}, \mathcal{Z}'$ are equivalent under $\text{Aff}(\mathbb{Z}^2)$, with

$$\begin{aligned} \text{gr } \mathbf{I}(\mathcal{Z}) &= \text{gr } \mathbf{I}(\mathcal{Z}') = (x_1^2, x_1 x_2, x_2^2) \subseteq \mathbb{R}[x_1, x_2], \\ V_{\mathcal{Z}} &= V_{\mathcal{Z}'} = \text{span}_{\mathbb{R}}\{1, y_1, y_2\} \subset \mathbb{D}_{\mathbb{R}}(y_1, y_2) = \mathbb{R}[y_1, y_2], \\ V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} &= \text{span}_{\mathbb{R}}\{1, y_1, y_2, y_1^2, y_1 y_2, y_2^2\}. \end{aligned}$$

On the other hand, $V_{\mathcal{Z} + \mathcal{Z}'}$ has dimension 7 = $\#(\mathcal{Z} + \mathcal{Z}')$, and one can compute that

$$\begin{aligned} \text{gr } \mathbf{I}(\mathcal{Z} + \mathcal{Z}') &= (x_1^3, x_2^3, x_1 x_2^2), \\ V_{\mathcal{Z} + \mathcal{Z}'} &= \text{span}_{\mathbb{R}}\{1, y_1, y_2, y_1^2, y_1 y_2, y_2^2, y_1^2 y_2\} \end{aligned}$$

so that $V_{\mathcal{Z} + \mathcal{Z}'}$ properly contains the 6-dimensional space $V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'}$ in this case.


Example 4.19. A somewhat nontrivial instance of Theorem 1.3 occurs when

$$\mathcal{Z} = \mathcal{Z}' = \{(0, 0), (1, 1), (2, 1), (1, 2)\} = \mathbb{Z}^2 \cap P,$$

where P is the lattice triangle discussed in Remark 3.15. Here one can check that

$$\mathcal{Z} + \mathcal{Z}' = \{(0, 0), (1, 1), (2, 1), (1, 2), (2, 2), (3, 2), (2, 3), (4, 2), (3, 3), (2, 4)\} = \mathbb{Z}^2 \cap 2P,$$

as shown below:



Remark 3.15 already mentioned one can calculate by hand or Macaulay2 that

$$V_{\mathcal{Z}} (= V_{\mathcal{Z}'}) = \text{span}_{\mathbb{R}}\{1, \quad y_1, y_2, \quad y_1^2 + y_1y_2 + y_2^2\}.$$

with spacing to indicate segregation of \mathbb{R} -basis elements by degree. Similarly one can calculate that

$$V_{\mathcal{Z}+\mathcal{Z}'} = \text{span}_{\mathbb{R}}\{1, \quad y_1, y_2, \quad y_1^2, y_1y_2, y_2^2, \quad y_1^3, y_1^2y_2 + y_1y_2^2, y_2^3, \quad y_1^4 + 2y_1^3y_2 + 3y_1^2y_2^2 + 2y_1y_2^3 + y_2^4\}.$$

It can be checked that multiplying any two basis elements of $V_{\mathcal{Z}}$ and $V_{\mathcal{Z}'}$ results in an element of $V_{\mathcal{Z}+\mathcal{Z}'}$. In this example, one has equality $V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} = V_{\mathcal{Z}+\mathcal{Z}'}$.

Example 4.20. Let us see what Theorem 1.3 says in a 1-dimensional space \mathbf{k}^1 , about the cardinalities $r = \#\mathcal{Z}$, $r' = \#\mathcal{Z}'$ of two finite loci \mathcal{Z} , \mathcal{Z}' , versus the cardinality r'' of their *sumset* $\mathcal{Z} + \mathcal{Z}'$. As in the Example from the Introduction, one can easily check for finite $\mathcal{Z} \subset \mathbf{k}^1$ that one has

$$\begin{aligned} \mathbf{I}(\mathcal{Z}) &= \left(\prod_{z \in \mathcal{Z}} (x - z) \right) \subset S = \mathbf{k}[x], \quad \text{gr } \mathbf{I}(\mathcal{Z}) = (x^r) \subset S = \mathbf{k}[x], \\ V_{\mathcal{Z}} &= \text{span}_{\mathbf{k}}\{1, y^{(1)}, y^{(2)}, \dots, y^{(r-1)}\} \subset \mathbb{D} = \mathbb{D}_{\mathbf{k}}(y). \end{aligned}$$

From this one can calculate

$$\begin{aligned} V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} &= \text{span}_{\mathbf{k}}\{1, y^{(1)}, y^{(2)}, \dots, y^{(r-1)}\} \cdot \text{span}_{\mathbf{k}}\{1, y^{(1)}, y^{(2)}, \dots, y^{(r'-1)}\} \\ &= \text{span}_{\mathbf{k}}\left\{ y^{(k)} \cdot y^{(k')} = \binom{k+k'}{k, k'} y^{(k+k')} : 0 \leq k \leq r-1 \text{ and } 0 \leq k' \leq r'-1 \right\}. \end{aligned}$$

Consequently, the assertion $V_{\mathcal{Z}} \cdot V_{\mathcal{Z}'} \subseteq V_{\mathcal{Z}+\mathcal{Z}'}$ from Theorem 1.3 holds if and only if

$$r'' - 1 \geq \max \left\{ k + k' : \exists 0 \leq k \leq r-1 \text{ and } 0 \leq k' \leq r'-1 \text{ with } \binom{k+k'}{k, k'} \in \mathbf{k}^{\times} \right\}, \text{ or equivalently,}$$

$$r'' \geq \underbrace{\min \left\{ n : (u+v)^n = \sum_{k+k'=n} \binom{n}{k, k'} u^k v^{k'} \text{ lies in the ideal } (u^r, v^{r'}) \subset \mathbf{k}[u, v] \right\}}_{\text{call this function } \beta_{\mathbf{k}}(r, r')}.$$

In other words, Theorem 1.3 says that for $\mathcal{Z}, \mathcal{Z}' \subset \mathbf{k}^1$, one has

$$(100) \quad \#(\mathcal{Z} + \mathcal{Z}') \geq \beta_{\mathbf{k}}(\#\mathcal{Z}, \#\mathcal{Z}').$$

When \mathbf{k} has characteristic zero, one has $\beta_{\mathbf{k}}(r, r') = r + r' - 1$, and the sumset lower bound (100) is not hard to show directly; it is also easily seen to be sharp. When $\mathbf{k} = \mathbb{F}_{p^d}$ is a finite field, this lower bound is a result of Eliahou and Kervaire [14, Thm. 2.1], generalizing both the case $d = 1$ for

$\mathbf{k} = \mathbb{F}_p$ known as the *Cauchy-Davenport Theorem*, as well as the case $p = 2$ for $\mathbf{k} = \mathbb{F}_{2^a}$ due to work of Yuzvinsky [49] on quadratic forms. Eliahou and Kervaire show [15, Thm. 2.2] that this lower bound (100) is also sharp for $\mathbf{k} = \mathbb{F}_{p^a}$. Interestingly, their proof method for (100) uses associated graded ideals $\text{gr } I$ and is close in spirit to our results.

Remark 4.21. Some readers may be bothered by the above proof of Theorem 1.3, which has many lemmas, and relies on limits and continuity arguments within projections to finite-dimensional subspaces of $\mathbb{D} = \mathbb{D}_{\mathbf{k}}(\mathbf{y})$. There is a second proof that avoids these complications, at the expense of considering divided power algebras $\mathbb{D}_R(\mathbf{y})$ whose coefficients lie in any commutative ring with unit R , not just a field \mathbf{k} , such as the univariate polynomial ring $R = \mathbf{k}[\epsilon]$. In the second proof, the limits as $\epsilon \in \mathbf{k}^\times$ approaches 0 are replaced by the ring homomorphism $e : R = \mathbf{k}[\epsilon] \rightarrow \mathbf{k}$ sending $\epsilon \mapsto 0$, and the corresponding R -module map $e : \hat{\mathbb{D}}_R(\mathbf{y}) \rightarrow \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})$. We sketch this proof here, leaving the reader to consult Eisenbud [13, A2.4], Geramita [20, §9] for more on $\mathbb{D}_R(\mathbf{y})$ over a ring R .

As in Section 4.2, for R a commutative ring with 1, if we let $S := R[\mathbf{x}] = R[x_1, \dots, x_n]$, there is an S -module structure on the divided power algebra $\mathbb{D} := \mathbb{D}_R(\mathbf{y})$, extending to its completion $\hat{\mathbb{D}} := \hat{\mathbb{D}}_R(\mathbf{y})$ as $\odot : S \times \hat{\mathbb{D}} \rightarrow \hat{\mathbb{D}}$. This gives rise to an R -bilinear pairing $\langle -, - \rangle : S \times \hat{\mathbb{D}}_R(\mathbf{y}) \rightarrow R$ in which $\langle f(\mathbf{x}), g(\mathbf{y}) \rangle$ extracts the constant term of $f \odot g$. For $\mathcal{Z} \subseteq R^n$, we again define the *vanishing ideal* in $S = R[\mathbf{x}]$, and its *annihilator/perp* R -submodule of $\hat{\mathbb{D}}$:

$$\begin{aligned} \mathbf{I}_R(\mathcal{Z}) &:= \{f(\mathbf{x}) \in S = R[\mathbf{x}] : f(\mathbf{z}) = 0 \text{ for all } \mathbf{z} \in \mathcal{Z}\} \subseteq R[\mathbf{x}], \\ \mathbf{I}_R(\mathcal{Z})^\perp &:= \{g \in \hat{\mathbb{D}} = \hat{\mathbb{D}}_R(\mathbf{y}) : \langle f, g \rangle = 0 \text{ for all } f \in \mathbf{I}_R(\mathcal{Z})\} \subseteq \hat{\mathbb{D}}, \end{aligned}$$

Here we emphasize the dependence on the ring R via the subscript in the notation.

Now consider for \mathbf{k} a field and finite locus $\mathcal{Z} \subseteq \mathbf{k}^n$, the “rescaled locus” $\epsilon\mathcal{Z} \subseteq R^n$ for $R = \mathbf{k}[\epsilon]$, along with its vanishing ideal and annihilator/perp R -module

$$\epsilon\mathcal{Z} := \{\epsilon\mathbf{z} : \mathbf{z} \in \mathcal{Z}\} \subseteq R^n, \quad \mathbf{I}_R(\epsilon\mathcal{Z}) \subset R[\mathbf{x}], \quad \mathbf{I}_R(\epsilon\mathcal{Z})^\perp \subset \hat{\mathbb{D}}_R(\mathbf{y}).$$

Although the ideal $\mathbf{I}_R(\epsilon\mathcal{Z})$ annihilates $\exp(\epsilon\mathbf{z} \cdot \mathbf{y})$ under the \odot -action for $\mathbf{z} \in \mathcal{Z}$, the R -module $\mathbf{I}_R(\epsilon\mathcal{Z})^\perp$ contains elements which are not in the R -span of $\{\exp(\epsilon\mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\}$. In contrast with Lemma 4.4, to obtain the full R -module $\mathbf{I}_R(\epsilon\mathcal{Z})^\perp$ we must *saturate* with respect to ϵ as follows.

Lemma 4.22. *The above R -submodule $\mathbf{I}_R(\epsilon\mathcal{Z})^\perp \subseteq \hat{\mathbb{D}}_R(\mathbf{y})$ has this description:*

$$\mathbf{I}_R(\epsilon\mathcal{Z})^\perp = \{f \in \hat{\mathbb{D}}_R(\mathbf{y}) : \epsilon^M \cdot f \in \text{span}_R\{\exp(\epsilon\mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\} \text{ for some } M \geq 0\}.$$

Proof. (Sketch) Naming the right side in the lemma as $W_{\mathcal{Z}}$, the inclusion $\mathbf{I}_R(\epsilon\mathcal{Z})^\perp \supseteq W_{\mathcal{Z}}$ is not hard, and follows similarly to the proof of Lemma 4.4. To show $\mathbf{I}_R(\epsilon\mathcal{Z})^\perp \subseteq W_{\mathcal{Z}}$, one strengthens the saturation property of $W_{\mathcal{Z}}$, to show this

Claim: If $0 \neq h(\epsilon) \in R = \mathbf{k}[\epsilon]$ and $f \in \hat{\mathbb{D}}_R(\mathbf{y})$ have $h(\epsilon) \cdot f \in W_{\mathcal{Z}}$, then $f \in W_{\mathcal{Z}}$.

By replacing \mathbf{k} with its algebraic closure, it is enough to establish this claim when \mathbf{k} is algebraically closed. To do this, start with a relation of the form

$$(101) \quad \epsilon^M \cdot h(\epsilon) \cdot f = \sum_{\mathbf{z} \in \mathcal{Z}} c_{\mathbf{z}}(\epsilon) \cdot \exp(\epsilon\mathbf{z} \cdot \mathbf{y})$$

for some $c_{\mathbf{z}}(\epsilon) \in \mathbf{k}[\epsilon]$. If $\alpha \in \mathbf{k}^\times$ is a nonzero root of $h(\epsilon)$, substituting $\epsilon \rightarrow \alpha$ and using the \mathbf{k} -linear independence of $\{\exp(\alpha\mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\}$ we get $c_{\mathbf{z}}(\alpha) = 0$ for all $\mathbf{z} \in \mathcal{Z}$, so we may cancel a factor of $(\epsilon - \alpha)$ from both sides of (101). We reduce to the case where $h(\epsilon) = \epsilon^r$ for some r and the claim follows easily.

Given the claim, one extends scalars in $\hat{\mathbb{D}}_R(\mathbf{y})$ to $\hat{\mathbb{D}}_{\mathbf{k}(\epsilon)}(\mathbf{y})$ where $\mathbf{k}(\epsilon)$ is the fraction field of $R = \mathbf{k}[\epsilon]$. Lemma 4.4 applies to give the $\mathbf{k}(\epsilon)$ -basis $\{\exp(\epsilon \mathbf{z} \cdot \mathbf{y}) : \mathbf{z} \in \mathcal{Z}\}$ of $\mathbf{I}_{\mathbf{k}(\epsilon)}(\epsilon \mathcal{Z})^\perp$. The inclusion $\mathbf{I}_R(\epsilon \mathcal{Z})^\perp \subseteq W_{\mathcal{Z}}$ (and hence Lemma 4.22) is deduced from the containment $\mathbf{I}_R(\epsilon \mathcal{Z})^\perp \subseteq \mathbf{I}_{\mathbf{k}(\epsilon)}(\epsilon \mathcal{Z})^\perp$, clearing denominators, and applying the claim. \square

Note that in $\hat{\mathbb{D}}_R(\mathbf{y})$, one still has

$$\exp(\epsilon \mathbf{z} \cdot \mathbf{y}) \cdot \exp(\epsilon \mathbf{z}' \cdot \mathbf{y}) = \exp(\epsilon(\mathbf{z} + \mathbf{z}') \cdot \mathbf{y}),$$

and hence Lemma 4.22 immediately implies the following analogue of Proposition 4.5.

Corollary 4.23. *For any two finite loci $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$, as R -submodules of $\hat{\mathbb{D}}_R(\mathbf{y})$ one has*

$$(102) \quad \mathbf{I}_R(\epsilon \mathcal{Z})^\perp \cdot \mathbf{I}_R(\epsilon \mathcal{Z}')^\perp \subseteq \mathbf{I}_R(\epsilon(\mathcal{Z} + \mathcal{Z}'))^\perp.$$

We remark that one can produce small examples with $n = 1$ showing both that the ϵ -saturation in the statement of Lemma 4.22 is necessary, and in contrast to Proposition 4.5, that the inclusion in Corollary 4.23 can be proper.

The last technical lemma needed is the following analogue of Lemma 4.17(ii).

Lemma 4.24. *Let $e : \hat{\mathbb{D}}_R(\mathbf{y}) \rightarrow \hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})$ be the map which evaluates $\epsilon \rightarrow 0$. For any finite locus $\mathcal{Z} \subseteq \mathbf{k}^n$, the map e restricts to a surjection $e : \mathbf{I}_R(\epsilon \mathcal{Z})^\perp \rightarrow \text{gr } \mathbf{I}_{\mathbf{k}}(\mathcal{Z})^\perp$.*


Proof. (Sketch) One checks that e indeed maps $\hat{\mathbb{D}}_R(\mathbf{y})$ into $\hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})$. Row reduction over the field \mathbf{k} shows that the \mathbf{k} -span of $\{\exp(\epsilon \mathbf{z} \cdot \mathbf{y})\}_{\mathbf{z} \in \mathcal{Z}}$ contains $\#\mathcal{Z}$ elements $\{f_{\mathbf{z}}\}_{\mathbf{z} \in \mathcal{Z}}$ with the following property:

each $f_{\mathbf{z}} = \epsilon^{M_{\mathbf{z}}} g_{\mathbf{z}}$ for some nonnegative power $M_{\mathbf{z}}$ of ϵ times another element $g_{\mathbf{z}}$ (so $g_{\mathbf{z}}$ also lies in $\mathbf{I}_R(\epsilon \mathcal{Z})^\perp$ by Lemma 4.22) and the e -images $\{e(g_{\mathbf{z}})\}_{\mathbf{z} \in \mathcal{Z}}$ are \mathbf{k} -linearly independent in $\hat{\mathbb{D}}_{\mathbf{k}}(\mathbf{y})$.

Since the space $\text{gr } \mathbf{I}_{\mathbf{k}}(\mathcal{Z})^\perp$ has \mathbf{k} -dimension $\#\mathcal{Z}$, this shows that e surjects. \square

Second proof of Theorem 1.3. Given g, g' in $V_{\mathcal{Z}}, V'_{\mathcal{Z}'}$, one wishes to show that $g \cdot g'$ lies in $V_{\mathcal{Z} + \mathcal{Z}'}$. Lift them both using Lemma 4.24 to g_ϵ, g'_ϵ in $\mathbf{I}_R(\epsilon \mathcal{Z})^\perp, \mathbf{I}_R(\epsilon \mathcal{Z}')^\perp$ with $e(g_\epsilon) = g, e(g'_\epsilon) = g'$. Then Corollary 4.23 implies that $g_\epsilon \cdot g'_\epsilon$ lies in $\mathbf{I}_R(\epsilon(\mathcal{Z} + \mathcal{Z}'))^\perp$. Finally, $g \cdot g' = e(g_\epsilon) \cdot e(g'_\epsilon) = e(g_\epsilon \cdot g'_\epsilon)$ lies in $V_{\mathcal{Z} + \mathcal{Z}'}$, due to Lemma 4.24 once more. \square

There is a natural geometric interpretation for some of the operations appearing in this second proof. As mentioned in the Introduction, one can view $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow R(\mathcal{Z})$ as a flat deformation of the reduced subscheme $\mathcal{Z} \subseteq \mathbf{k}^n$ deforming linearly to a zero dimensional subscheme of degree $\#\mathcal{Z}$ supported at the origin.



The extension of scalars from \mathbf{k} to $R = \mathbf{k}[\epsilon]$ in the second proof corresponds to viewing $\epsilon \mathcal{Z} \subseteq \mathbb{A}_{\mathbf{k}}^n \times \mathbb{A}_{\mathbf{k}}^1$ as a flat family over $\mathbb{A}_{\mathbf{k}}^1 = \text{Spec}(\mathbf{k}[\epsilon])$ via $\epsilon \mathcal{Z} = \text{Spec}(\mathbf{k}[\epsilon][\mathbf{x}]/\mathbf{I}_{\mathbf{k}[\epsilon]}(\epsilon \mathcal{Z}))$, with generic fibers over $\epsilon = c \in \mathbf{k}^\times$ and special fiber over $\epsilon = 0$. The limit specialization $\epsilon \rightarrow 0$ corresponds to applying the map e or the functor $(-)\otimes_{\mathbf{k}[\epsilon]} \mathbf{k}[\epsilon]/(\epsilon)$, focusing on the special fiber. Passage to the fraction field

$\mathbf{k}(\epsilon)$, as in the proof of Lemma 4.22, corresponds to applying $(-)\otimes_{\mathbf{k}[\epsilon]} \mathbf{k}(\epsilon)$, which geometrically is localization over the general point $(0) \in \text{Spec}(\mathbf{k}[\epsilon])$.

Remark 4.25. Theorem 1.3 has a resemblance to another interesting observation of Gundlach [23]. Suppose \mathbf{k} has characteristic 0. For any finite point set $\mathcal{Z} \subseteq \mathbf{k}^n$, and any choice of monomial ordering \prec on $S = \mathbf{k}[\mathbf{x}]$, his result quoted as Proposition 2.9 earlier shows the set $\mathcal{S}_{\mathcal{Z}}^{\prec}$ of all \prec -standard monomials for $\mathbf{I}(\mathcal{Z})$ has the following alternate description: $\mathcal{S}_{\mathcal{Z}}^{\prec} = \{\text{sm}_{\prec}(f) : f \in \mathbf{k}^{\mathcal{Z}} \setminus \{0\}\}$. He then uses this to fairly quickly prove the following fact [23, eqn. (5)] : for any two finite point sets $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$, one has

$$(103) \quad \mathcal{S}_{\mathcal{Z}}^{\prec} \cdot \mathcal{S}_{\mathcal{Z}'}^{\prec} \subseteq \mathcal{S}_{\mathcal{Z}+\mathcal{Z}'}^{\prec}.$$

It is not clear how closely this is related to the similar-sounding Theorem 1.3.

5. HARMONIC ALGEBRAS

Our motivation for Theorem 1.3 was to define here the *harmonic algebra* \mathcal{H}_P as an approach toward our main Conjecture 1.1 generalizing the Classical Ehrhart Theorem. Before defining \mathcal{H}_P , we explain why a less algebraic approach to Conjecture 1.1 seems elusive.

5.1. The missing valuative property. Recall from the Introduction that the Classical Ehrhart Theorem asserts several properties for the *Ehrhart series* $E_P(t) := \sum_{m=0}^{\infty} i_P(m)t^m$ of a d -dimensional lattice polytope P in \mathbb{R}^n , where $i_P(m) := \#(mP \cap \mathbb{Z}^n)$. Specifically, one has its *rationality* $E_P(t) = \sum_{i=0}^d h_i^* t^i / (1-t)^{d+1}$ with a precise *denominator*, the *nonnegativity* of the numerator coefficients $\{h_i^*\}$, the combinatorial interpretation of the $\{h_i^*\}$ in terms of the semi-open parallelepiped (3) when P is a simplex, and the *reciprocity* relating $E_P(t^{-1})$ to $\overline{E}_P(t)$.

These properties were originally given elementary proofs that avoid any commutative algebra, via the following strategy: one first proves the result for lattice *simplices* (and semi-open simplices), and then generalizes to arbitrary lattice polytopes employing triangulations (and sometimes *shellings* of a triangulation) of P . The key tool in such proofs is the following *valuative property* that comes directly from the definitions of $i_P(m)$ and $E_P(t)$: when P, Q are lattice polytopes and their intersection $P \cap Q$ is a proper face of both P, Q , then

$$(104) \quad i_{P \cup Q}(m) = i_P(m) + i_Q(m) - i_{P \cap Q}(m)$$

$$(105) \quad E_{P \cup Q}(t) = E_P(t) + E_Q(t) - E_{P \cap Q}(t).$$

One might therefore hope to prove the analogous Conjecture 1.1 on

$$i_P(m; q) := \text{Hilb}(R(\mathbb{Z}^n \cap mP), q) = \text{Hilb}(V_{\mathbb{Z}^n \cap mP}, q), \text{ and}$$


$$E_P(t, q) := \sum_{m=0}^{\infty} i_P(m; q)t^m,$$

via similar valuative techniques. The following simple example illustrates some difficulty in identifying an analogous valuative property.

Example 5.1. Let $a, b \geq 1$ be coprime positive integers, and let $P, Q \subseteq \mathbb{R}^2$ be the lattice triangles

$$P := \text{conv}\{(0, 0), (0, b), (a, 0)\} \quad \text{and} \quad Q := \text{conv}\{(0, b), (a, 0), (a, b)\}.$$

Their union is the lattice rectangle $P \cup Q = [0, a] \times [0, b]$, and their intersection is the line segment $P \cap Q = \text{conv}\{(0, b), (a, 0)\}$. The figure below depicts the case $a = 5, b = 4$.



Since P, Q are equivalent under $\text{Aff}(\mathbb{Z}^2)$, and since P and $P \cup Q$ are both antiblocking polytopes, one can apply Corollary 3.12 to compute that

$$(106) \quad i_P(m; q) = i_Q(m; q) = \sum_{\substack{\mathbf{z}=(z_1, z_2) \\ \text{in } \mathbb{Z}^2 \cap mP}} q^{z_1+z_2} = \sum_{\substack{(z_1, z_2) \in \mathbb{Z}_{\geq 0}: \\ bz_1+az_2 \leq mab}} q^{z_1+z_2}$$

$$(107) \quad i_{P \cup Q}(m; q) = \sum_{\substack{\mathbf{z}=(z_1, z_2) \\ \text{in } \mathbb{Z}^2 \cap m(P \cup Q)}} q^{z_1+z_2} = [ma]_q \cdot [mb]_q$$

Since a, b are coprime, the set $P \cap Q \cap \mathbb{Z}^2 = \{(0, b), (a, 0)\}$ consists of only two lattice points, and its dilate $\mathbb{Z}^2 \cap m(P \cap Q)$ consists of $m+1$ collinear lattice points, so that

$$(108) \quad i_{P \cap Q}(m; q) = 1 + q + q^2 + \cdots + q^m = [m+1]_q.$$

It is not clear if there is a q -analogue of the valuative property (104) that applies here to relate (106), (107), (108). This subtlety is not surprising— for general finite loci $\mathcal{Z}, \mathcal{Z}' \subseteq \mathbf{k}^n$, it seems difficult to predict the structure of $\text{gr } \mathbf{I}(\mathcal{Z} \cup \mathcal{Z}')$ from that of $\text{gr } \mathbf{I}(\mathcal{Z})$, $\text{gr } \mathbf{I}(\mathcal{Z}')$ and $\text{gr } \mathbf{I}(\mathcal{Z} \cap \mathcal{Z}')$.

5.2. The harmonic algebra \mathcal{H}_P and a conjecture. As explained in the Introduction, work of Stanley (see, e.g., [45]) provided an alternative proof for the assertions in the Classical Ehrhart Theorem, avoiding valuative techniques, and substituting commutative algebra. For a d -dimensional lattice polytope $P \subset \mathbb{R}^n$, he considered the *affine semigroup ring* A_P inside the Laurent polynomial ring $\mathbf{k}[y_0, y_1^{\pm 1}, \dots, y_n^{\pm 1}]$ and its *interior ideal* \bar{A}_P , defined as follows:

$$\begin{aligned} A_P &:= \mathbf{k}[\mathbb{Z}^{n+1} \cap \text{cone}(P)] := \text{span}_{\mathbf{k}}\{y_0^m \mathbf{y}^{\mathbf{z}} : \mathbf{z} \in mP\}, \\ \bar{A}_P &:= \mathbf{k}[\mathbb{Z}^{n+1} \cap \text{int}(\text{cone}(P))] := \text{span}_{\mathbf{k}}\{y_0^m \mathbf{y}^{\mathbf{z}} : \mathbf{z} \in \text{int}(mP)\}, \end{aligned}$$

This ring A_P is the affine semigroup ring associated to the polyhedral $(d+1)$ -dimensional cone

$$\text{cone}(P) := \mathbb{R}_{\geq 0} \cdot (\{1\} \times P) \subset \mathbb{R}^{n+1},$$

whose lattice points in \mathbb{Z}^{n+1} form a semigroup under addition. The distinguished zeroth coordinate on $\mathbb{R}^{n+1} = \mathbb{R}^1 \times \mathbb{R}^n$ endows A_P with the structure of an \mathbb{N} -graded algebra of Krull dimension $d+1$, and one can re-interpret

$$(109) \quad E_P(t) = \text{Hilb}(A_P, t),$$

$$(110) \quad \bar{E}_P(t) = \text{Hilb}(\bar{A}_P, t).$$

As mentioned in the Introduction, Stanley explained the

- *rationality* of $E_P(t)$ via A_P being a finitely generated \mathbf{k} -algebra, shown by Gordan [21],
- *denominator* $(1-t)^{d+1}$ of $E_P(t)$ via Noether's Normalization Lemma [35],
- *nonnegativity* of the $\{h_i^*\}$ via Cohen-Macaulayness of A_P , a result of Hochster [26], and
- *reciprocity* via $\bar{A}_P \cong \Omega A_P$, the *canonical module* of A_P , work of Danilov [11], Stanley [44].

Motivated by this, we would like to eventually prove the q -analogous assertions of Conjecture 1.1, via the following commutative algebra.

Definition 5.2. Let P be a lattice polytope in \mathbb{R}^n , and introduce the polynomial ring

$$\mathbb{R}[y_0, \mathbf{y}] := \mathbb{R}[y_0, y_1, \dots, y_n] \quad (\cong \mathbb{R}[y_0] \otimes_{\mathbb{R}} \mathbb{D}_{\mathbb{R}}[\mathbf{y}]).$$

We will consider $\mathbb{R}[y_0, \mathbf{y}]$ as a *bigraded* or \mathbb{N}^2 -*graded* \mathbb{R} -algebra in which

$$\deg(y_0) = (1, 0), \quad \deg(y_1) = \dots = \deg(y_n) = (0, 1),$$

with Hilbert series in t, q tracking a monomial $y_0^m y_1^{z_1} \dots y_n^{z_n}$ by $t^m q^{z_1 + \dots + z_n}$. Occasionally we will specialize this to a single \mathbb{N} -grading by sending $q \mapsto 1$. Then define the *harmonic algebra* \mathcal{H}_P and its *interior ideal* $\overline{\mathcal{H}}_P$ as the following two \mathbb{N}^2 -homogeneous \mathbb{R} -linear subspaces of $\mathbb{R}[y_0, \mathbf{y}]$:

$$(111) \quad \mathcal{H}_P := \bigoplus_{m=0}^{\infty} \mathbb{R} \cdot y_0^m \otimes_{\mathbb{R}} V_{\mathbb{Z}^n \cap mP},$$

$$(112) \quad \overline{\mathcal{H}}_P := \bigoplus_{m=0}^{\infty} \mathbb{R} \cdot y_0^m \otimes_{\mathbb{R}} V_{\mathbb{Z}^n \cap \text{int}(mP)}.$$

We first justify the terms ‘‘algebra’’, ‘‘ideal’’ in the definition, starting with an easy observation.

Lemma 5.3. *For nested finite subsets $\mathcal{Z} \subseteq \mathcal{Z}' \subseteq \mathbf{k}^n$ over any field \mathbf{k} , one has $V_{\mathcal{Z}} \subseteq V_{\mathcal{Z}'}$.*

Proof. $\mathcal{Z} \subseteq \mathcal{Z}'$ implies the opposite inclusion $\mathbf{I}(\mathcal{Z}) \supseteq \mathbf{I}(\mathcal{Z}')$ within $S = \mathbb{R}[\mathbf{x}]$. This then implies $\text{gr } \mathbf{I}(\mathcal{Z}) \supseteq \text{gr } \mathbf{I}(\mathcal{Z}')$. Taking perps in $\mathbb{D}_{\mathbf{k}}(\mathbf{y})$ reverses inclusion: $V_{\mathcal{Z}} = \text{gr } \mathbf{I}(\mathcal{Z})^{\perp} \subseteq \text{gr } \mathbf{I}(\mathcal{Z}')^{\perp} = V_{\mathcal{Z}'}$. \square

Proposition 5.4. *Within the ring $\mathbb{R}[y_0, \mathbf{y}]$, the \mathbb{R} -linear subspace \mathcal{H}_P is an \mathbb{N}^2 -graded \mathbb{R} -subalgebra, and $\overline{\mathcal{H}}_P$ is an \mathbb{N}^2 -graded ideal of \mathcal{H}_P , with*

$$(113) \quad E_P(t, q) = \text{Hilb}(\mathcal{H}_P, t, q),$$

$$(114) \quad \overline{E}_P(t, q) = \text{Hilb}(\overline{\mathcal{H}}_P, t, q).$$

Furthermore, any finite subgroup G of $GL_n(\mathbb{Z})$ preserving P acts by \mathbb{N}^2 -graded automorphisms on \mathcal{H}_P . Thus in the representation ring $\text{Rep}(G)[[t, q]]$ of \mathbb{N}^2 -graded $\mathbb{R}G$ -modules, one has

$$E_P^G(t, q) = \sum_{m, j \geq 0} [(\mathcal{H}_P)_{m, j}] \cdot t^m q^j$$

Proof. For the first algebra assertion about \mathcal{H}_P , given $m, m' \geq 0$, one must check that

$$(\mathbb{R} \cdot y_0^m \otimes_{\mathbb{R}} V_{\mathbb{Z}^n \cap mP}) \cdot (\mathbb{R} \cdot y_0^{m'} \otimes_{\mathbb{R}} V_{\mathbb{Z}^n \cap m'P}) \subseteq \mathbb{R} \cdot y_0^{m+m'} \otimes_{\mathbb{R}} V_{\mathbb{Z}^n \cap (m+m')P}.$$

However, note that one can use Theorem 1.3 to deduce the first inclusion here

$$V_{\mathbb{Z}^n \cap mP} \cdot V_{\mathbb{Z}^n \cap m'P} \subseteq V_{(\mathbb{Z}^n \cap mP) + (\mathbb{Z}^n \cap m'P)} \subseteq V_{\mathbb{Z}^n \cap (m+m')P}.$$

The second inclusion follows from Proposition 5.3 together with this inclusion

$$(115) \quad (\mathbb{Z}^n \cap mP) + (\mathbb{Z}^n \cap m'P) \subseteq \mathbb{Z}^n \cap (m+m')P,$$

derived from $mP + m'P = (m+m')P$. The ideal assertion for $\overline{\mathcal{H}}_P$ is similar, replacing the last fact with $mP + \text{int}(m'P) = \text{int}((m+m')P)$. The remaining assertions follow from the definitions. \square

We conjecture the following properties for the harmonic algebra \mathcal{H}_P , analogous to those known for the affine semigroup ring A_P , which would explain most of Conjecture 1.1.

Conjecture 5.5. *Let P be a d -dimensional lattice polytope in \mathbb{R}^n , having V vertices.*

(i) *The harmonic algebra \mathcal{H}_P is a Noetherian (finitely generated) \mathbb{R} -subalgebra of $\mathbb{R}[y_0, \mathbf{y}]$.*

- (ii) There exist \mathbb{N}^2 -homogeneous elements $\{\theta_i\}_{1,2,\dots,\nu}$ in \mathcal{H}_P with $\dim(P) + 1 \leq \nu \leq V$, generating an \mathbb{R} -subalgebra $B \subseteq \mathcal{H}_P$ for which \mathcal{H}_P is a finite ring extension.
- (iii) The \mathbb{N}^2 -homogeneous generators for \mathcal{H}_P and the elements $\{\theta_i\}$ from (ii) can be chosen with with bidegrees (b_i, a_i) having $b_i \leq \#(P \cap \mathbb{Z}^n) - 1$.
- (iv) The algebra \mathcal{H}_P is Cohen-Macaulay.
- (v) Its canonical module $\Omega\mathcal{H}_P$ is isomorphic to the ideal $\overline{\mathcal{H}}_P$, up to a shift in \mathbb{N}^2 -grading.

5.3. Relation of Conjecture 5.5 to Conjecture 1.1. We explain here the implications of the algebraic Conjecture 5.5 on \mathcal{H}_P for the enumerative Conjecture 1.1 on its Hilbert series $E_P(t, q)$.

Note that when we specialize the \mathbb{N}^2 -grading of the harmonic algebra \mathcal{H}_P to the \mathbb{N} -grading, by setting $q = 1$, its Hilbert series $E_P(t, q)$ becomes the Ehrhart series $E_P(t)$, matching that of the affine semigroup ring A_P :

$$(116) \quad [\text{Hilb}(\mathcal{H}_P, t, q)]_{q=1} = [E_P(t, q)]_{q=1} = E_P(t) = \text{Hilb}(A_P, t),$$

where the first equality uses (113) and the last is (109). Consequently, if Conjecture 5.5(i) holds, so \mathcal{H}_P is a finitely generated algebra, we know \mathcal{H}_P has *Krull dimension* $d + 1$, the same that of A_P .

Also, one could then choose a finite set of \mathbb{N}^2 -homogeneous algebra generators, e.g., by taking the set of all \mathbb{N}^2 -homogeneous components from any finite list of algebra generators. Assume one has chosen such a list of \mathbb{N}^2 -homogeneous algebra generators for \mathcal{H}_P , or even just some $\{\theta_i\}$ satisfying the weaker condition that they generate a subalgebra $B \subseteq \mathcal{H}_P$ over which \mathcal{H}_P is a finite extension, as in Conjecture 5.5(ii). Name their \mathbb{N}^2 -degrees $\deg(\theta_i) = (b_i, a_i)$ for $i = 1, 2, \dots, \nu$, and create an \mathbb{N}^2 -graded polynomial ring $S := \mathbb{R}[X_1, \dots, X_\nu]$ with variables having $\deg(X_i) := (b_i, a_i)$. Then the map $S \rightarrow \mathcal{H}_P$ sending $X_i \mapsto \theta_i$ makes \mathcal{H}_P a finitely generated \mathbb{N}^2 -graded S -module. Hilbert's Syzygy Theorem therefore predicts the existence of a finite \mathbb{N}^2 -graded free S -resolution of \mathcal{H}_P

$$(117) \quad 0 \leftarrow \mathcal{H}_P \leftarrow F_0 \leftarrow F_1 \leftarrow F_2 \leftarrow \dots \leftarrow F_\nu \leftarrow 0.$$

Here each free S -module F_j in the resolution has the form

$$F_j = \bigoplus_{(b,a) \in \mathbb{N}^2} S(-(b, a))^{\beta_{j,(b,a)}},$$

for some \mathbb{N}^2 -graded Betti numbers $\beta_{j,(b,a)}$. That is, F_j has exactly $\beta_{j,(b,a)}$ of its S -basis elements of \mathbb{N}^2 -degree (b, a) . Taking the \mathbb{N}^2 -graded Euler characteristic of the resolution (117) implies

$$\begin{aligned} E_P(t, q) &= \text{Hilb}(\mathcal{H}_P, t, q) = \sum_{j=0}^{\nu} (-1)^j \text{Hilb}(S(-(b, a))^{\beta_{j,(b,a)}}, t, q) \\ &= \text{Hilb}(S, t, q) \cdot \sum_{j=0}^{\nu} (-1)^j \sum_{(b,a) \in \mathbb{N}^2} \beta_{j,(b,a)} q^b t^a \\ &= \frac{\sum_{j=0}^{\nu} (-1)^j \sum_{(b,a) \in \mathbb{N}^2} \beta_{j,(b,a)} q^b t^a}{\prod_{i=1}^{\nu} (1 - q^{a_i} t^{b_i})}. \end{aligned}$$

which shows the first rationality assertion in Conjecture 1.1 on $E_P(t, q)$, and also Conjecture 1.1(ii). This argument also works for the rationality assertion on $\overline{E}_P(t, q)$: the ideal $\overline{\mathcal{H}}_P$ in \mathcal{H}_P would be finitely generated, due to Noetherian-ness of \mathcal{H}_P , and hence also a finitely generated S -module. If one has finer information on the number ν of \mathbb{N}^2 -homogeneous generators of \mathcal{H}_P as in Conjecture 5.5(ii),(iii), then these would imply Conjecture 1.1(i),(iv).

If the Cohen-Macaulayness assertion Conjecture 5.5(iv) holds, this implies Conjecture 1.1(v) on lattice simplices by the following argument. Since lattice d -simplices have $V = d + 1$, which is the same as the Krull dimension of \mathcal{H}_P , Conjecture 5.5(ii) implies one can pick an \mathbb{N}^2 -homogeneous system of parameters $\{\theta_i\}_{i=1,2,\dots,d+1}$ for \mathcal{H}_P . Cohen-Macaulayness implies \mathcal{H}_P is a free S -module, so the resolution (117) stops at F_0 , and


$$E_P(t, q) = \text{Hilb}(F_0, t, q) = \frac{\sum_{(b,a) \in \mathbb{N}^2} \beta_{0,(b,a)} q^b t^a}{\prod_{i=1}^{\nu} (1 - q^{a_i} t^{b_i})}.$$

Lastly, Conjecture 5.5(v) could imply the q -reciprocity assertion Conjecture 1.1(vi), if the shift in \mathbb{N}^2 -grading for the isomorphism $\Omega\mathcal{H}_P \cong \overline{\mathcal{H}}_P$ works out correctly: one knows that the canonical module $\Omega\mathcal{H}_P$ for an \mathbb{N}^2 -graded Cohen-Macaulay ring \mathcal{H}_P of Krull dimension $d + 1$ has

$$\text{Hilb}(\Omega\mathcal{H}_P, t, q) = (-1)^{d+1} q^B t^A \cdot \text{Hilb}(\mathcal{H}_P, t^{-1}, q^{-1})$$

for some choice of \mathbb{N}^2 -degree (B, A) ; see Stanley [45, §I.12, p.49].

5.4. The singly-graded rings \mathcal{H}_P and A_P are generally not isomorphic. In light of the fact (116) that A_P and \mathcal{H}_P have the same singly-graded Hilbert series, one might wonder whether they are isomorphic as \mathbb{N} -graded algebras. This fails already for small lattice polygons, e.g., this triangle $P = \text{conv}\{(0, 0), (1, 2), (3, 1)\}$, for which a portion of $\mathbb{Z}^3 \cap \text{cone}(P)$ is shown below:



To see that $A_P \not\cong \mathcal{H}_P$ as \mathbb{N} -graded algebras, one can check that

$$(118) \quad (A_P)_1 \cdot (A_P)_1 = (A_P)_2, \text{ but}$$

$$(119) \quad (\mathcal{H}_P)_1 \cdot (\mathcal{H}_P)_1 \subsetneq (\mathcal{H}_P)_2.$$

Checking (118) means showing $(\mathbb{Z}^2 \cap P) + (\mathbb{Z}^2 \cap P) = (\mathbb{Z}^2 \cap 2P)$, which can be done directly. Meanwhile, the proper inclusion (119) follows from examining the first few terms of $E_P(t, q)$:

$$E_P(t, q) = 1 + (1 + 2q + 2q^2)t + (1 + 2q + 3q^2 + 4q^3 + 3q^4 + q^5)t^2 + o(t^2)$$

One sees $(\mathcal{H}_P)_1 \cdot (\mathcal{H}_P)_1$ has as its highest nonvanishing q -degree $q^2 \cdot q^2 = q^4$, but for $(\mathcal{H}_P)_2$ it is q^5 . This example seems related to the deformation $S/\mathbf{I}(\mathcal{Z}) \rightsquigarrow S/\text{gr } \mathbf{I}(\mathcal{Z})$ forgetting algebra structure.

Remark 5.6. In contrast to the above example, it may be interesting to note that there is a very similar-sounding algebra to \mathcal{H}_P which actually *is* isomorphic to A_P . Recall from Section 4.2 that for the inhomogeneous ideal $\mathbf{I}(\mathcal{Z})$ one can still define a perp space $\mathbf{I}(\mathcal{Z})^\perp$ inside the power series completion $\hat{\mathbb{D}} = \hat{\mathbb{D}}_{\mathbb{R}}[\mathbf{y}] \cong \mathbb{R}[[y_1, \dots, y_n]]$. The space $\mathbf{I}(\mathcal{Z})^\perp$ has \mathbb{R} -basis given in Lemma 4.4 by $\{\exp(\mathbf{z} \cdot \mathbf{y})\}_{\mathbf{z} \in \mathcal{Z}}$, and (88) asserts that these basis elements multiply via the usual rule

$$\exp(\mathbf{z} \cdot \mathbf{y}) \cdot \exp(\mathbf{z}' \cdot \mathbf{y}) = \exp((\mathbf{z} + \mathbf{z}') \cdot \mathbf{y}).$$

If one then considers the ring $\mathbb{R}[y_0][[y_1, \dots, y_n]]$ as \mathbb{N} -graded by $\deg(y_0^m) = 1$ and $\deg(y_i) = 0$ for $i = 1, 2, \dots, n$, one can compile the spaces $\mathbf{I}(\mathbb{Z}^n \cap mP)^\perp$ for $m \geq 0$ to define an \mathbb{N} -graded subalgebra

$$\hat{\mathcal{H}}_P := \bigoplus_{m=0}^{\infty} \mathbb{R} \cdot y_0^m \otimes \mathbf{I}(\mathbb{Z}^n \cap mP)^\perp \quad \text{inside } \mathbb{R}[y_0][[y_1, \dots, y_n]].$$

Since A_P has \mathbb{R} -basis $\{y_0^m \mathbf{y}^{\mathbf{z}} : \mathbf{z} \in \mathbb{Z}^n \cap mP\}$, with $y_0^m \mathbf{y}^{\mathbf{z}} \cdot y_0^{m'} \mathbf{y}^{\mathbf{z}'} = y_0^{m+m'} \mathbf{y}^{\mathbf{z}+\mathbf{z}'}$, the map

$$\begin{aligned} A_P &\longrightarrow \hat{\mathcal{H}}_P \\ y_0^m \mathbf{y}^{\mathbf{z}} &\longmapsto y_0^m \otimes \exp(\mathbf{z} \cdot \mathbf{y}) \end{aligned}$$

is an \mathbb{N} -graded \mathbb{R} -algebra isomorphism.

5.5. Example: antiblocking polytopes revisited. In spite of the example in Section 5.4, there is a subfamily of lattice polytopes P for which one has not only an \mathbb{N} -graded, but even an \mathbb{N}^2 -graded algebra isomorphism $\mathcal{H}_P \cong A_P$, and Conjecture 5.5 holds: the *antiblocking polytopes* of Section 3.3.

In order to discuss this, we first note that for a lattice polytope $P \subset \mathbb{R}^n$, the affine semigroup ring A_P has a finer $\mathbb{N} \times \mathbb{Z}^n$ -grading. This is because it is a homogeneous subalgebra of $\mathbb{R}[y_0, y_1^{\pm 1}, \dots, y_n^{\pm 1}]$ with respect to the $\mathbb{N} \times \mathbb{Z}^n$ -multigrading in which $\deg(y_i) = (0, \dots, 0, 1, 0, \dots, 0)$ is the i^{th} standard basis vector. One can then specialize this to a \mathbb{N}^2 -grading via $\deg(y_0) = (1, 0)$ and $\deg(y_i) = (0, 1)$ for $i = 1, 2, \dots, n$. In other words, this \mathbb{N}^2 -grading has $\deg(y_0^m y_1^{z_1} \dots y_n^{z_n}) = (m, z_1 + \dots + z_n)$, tracked by the Hilbert series monomial $t^m q^{z_1 + \dots + z_n}$.

Proposition 5.7. *For any antiblocking lattice polytope $P \subseteq \mathbb{R}^n$ one has an \mathbb{N}^2 -graded \mathbb{R} -algebra isomorphism $A_P \longrightarrow \mathcal{H}_P$ defined by the bijection on \mathbb{R} -bases $y_0^m \mathbf{y}^{\mathbf{z}} \longmapsto y_0^m \otimes \mathbf{y}^{\mathbf{z}}$ for $\mathbf{z} \in \mathbb{Z}^n \cap mP$. That is, identifying $\mathbb{R}[y_0] \otimes_{\mathbb{R}} \mathbb{D}_{\mathbb{R}}(\mathbf{y}) \cong \mathbb{R}[y_0, \mathbf{y}]$, one has equality $A_P = \mathcal{H}_P$ of the two subalgebras.*

Consequently, Conjecture 5.5 holds for antiblocking polytopes.


Proof. The isomorphism assertion follows from Lemma 3.8(iii).

To see Conjecture 5.5 holds, it was already mentioned that A_P is finitely-generated, Cohen-Macaulay, and with canonical module $\Omega_{A_P} \cong \bar{A}_P$ from results of Gordan, Hochster and of Danilov and Stanley, respectively. This proves Conjecture 5.5 parts (i),(iv),(v). For parts (ii),(iii), note that if the polytope P has vertices $\{\mathbf{z}^{(i)}\}_{i=1,2,\dots,V}$, then A_P contains \mathbb{N}^2 -homogeneous elements $\{\theta_i = y_0^1 \mathbf{y}^{\mathbf{z}^{(i)}}\}_{i=1,2,\dots,V}$ generating a subalgebra B for which $B \subseteq A_P$ is a finite extension. The proof of Gordan's result also shows that A_P is generated as a B -module by the elements

$$(120) \quad \left\{ y_0^m \mathbf{y}^{\mathbf{z}} : (m, \mathbf{z}) = \sum_{i=1}^V c_i \cdot (1, \mathbf{z}^{(i)}) \in \mathbb{Z}^{n+1} \text{ with } 0 \leq c_i < 1 \right\}.$$

These elements in (120) have \mathbb{N}^2 -degrees (b, a) with $0 \leq b \leq V - 1$, and together with $\{\theta_i\}_{i=1,2,\dots,V}$ generate A_P as an algebra. Since $V \leq \#\mathbb{Z}^n \cap P$, the degree bound in Conjecture 5.5(iii) holds. \square

5.6. Case study: an interesting triangle. For general lattice polytopes P , the structure of \mathcal{H}_P can be more complicated. We return to study the lattice triangle $P := \text{conv}\{\mathbf{0}, (1, 2), (2, 1)\} \subset \mathbb{R}^2$ discussed in Remark 3.15 and Example 4.19, with $P, 2P, 3P$ shown below:



This triangle P is not antiblocking, and is $\text{Aff}(\mathbb{Z}^2)$ -equivalent to the second lattice triangle of area 3 in Figure 1, with these Ehrhart and q -Ehrhart series

$$E_P(t) = \frac{1+t+t^2}{(1-t)^3} = 1 + 4t + 10t^2 + 19t^3 + \dots$$

$$E_P(t, q) = \frac{(1+qt)(1+qt+q^2t^2)}{(1-t)(1-q^2t)(1-q^3t^2)}$$

$$= 1 + (1+2q+q^2)t + (1+2q+3q^2+3q^3+q^4)t^2 + (1+2q+3q^2+4q^3+5^4+3q^5+q^6)t^3 + \dots$$

We explain below how one can use the harmonic algebra \mathcal{H}_P to prove the above calculation of $E_P(t, q)$ is correct. Before doing so, we note two interesting features of $E_P(t, q)$.

First, note $[E_P(t, q)]_{q=1} = E_P(t)$ requires canceling a numerator/denominator factor at $q = 1$.

Second, note that the bidegrees t, q^2t, q^3t^2 appearing in the denominator factors of $E_P(t, q)$ are *all different*, even though the triangle P has a $GL(\mathbb{Z}^2)$ -symmetry that swaps the two vertices $(2, 1), (1, 2)$. Thus one should not expect some simple and natural bijection between the vertices and the \mathbb{N}^2 -graded elements $\{\theta_1, \theta_2, \theta_3\}$ whose existence is predicted by Conjecture 5.5(ii).

Expecting that Conjecture 5.5 might hold, and examining $E_P(t, q)$, shown color-coded here,

$$(121) \quad E_P(t, q) = \frac{1 + 2qt + 2q^2t^2 + q^3t^3}{(1-t)(1-q^2t)(1-q^3t^2)}.$$

one might expect from the denominator that the harmonic algebra \mathcal{H}_P contains a homogeneous system of parameters $\theta_1, \theta_2, \theta_3$ of \mathbb{N}^2 -degrees $(0, 1), (1, 2), (2, 3)$. From the numerator one might expect to find six $\mathbb{R}[\theta_1, \theta_2, \theta_3]$ -basis elements whose \mathbb{N}^2 -degrees match $1 + 2qt + 2q^2t^2 + q^3t^3$. Using `Macaulay2`, one can compute \mathbb{N} -graded \mathbb{R} -bases for the harmonic spaces $V_{\mathbb{Z}^2 \cap mP}$ with $m = 0, 1, 2, 3$, and hence \mathbb{N}^2 -graded \mathbb{R} -bases for $(\mathcal{H}_P)_0, (\mathcal{H}_P)_1, (\mathcal{H}_P)_2, (\mathcal{H}_P)_3$. We then rewrote these bases to make the action of $G = \mathbb{Z}/2\mathbb{Z}$ apparent in each graded piece, and identified candidates for $\theta_1, \theta_2, \theta_3$ and the six $\mathbb{R}[\theta_1, \theta_2, \theta_3]$ -basis elements, color-coded in this table:

$(\mathcal{H}_P)_0$	1						
$(\mathcal{H}_P)_1$	$\theta_1 =$ y_0	y_0y_1 y_0y_2	$\theta_2 =$ $y_0(y_1^2 + y_1y_2 + y_2^2)$				
$(\mathcal{H}_P)_2$	y_0^2	$y_0^2y_1$ $y_0^2y_2$	$y_0^2y_1^2$ $y_0^2y_1y_2$ $y_0^2y_2^2$	$y_0^2y_1^3$ $\theta_3 = y_0^2(y_1^2y_2 + y_1y_2^2)$ $y_0^2y_2^3$	$y_0^2(y_1^4 + 2y_1^3y_2$ $+ 3y_1^2y_2^2 + 2y_1y_2^3$ $+ y_2^4)$		
$(\mathcal{H}_P)_3$	y_0^3	$y_0^3y_1$ $y_0^3y_2$	$y_0^3y_1^2$ $y_0^3y_1y_2$ $y_0^3y_2^2$	$y_0^3y_1^3$ $y_0^3(y_1^2y_2 + y_1y_2^2)$ $y_0^3(y_1^2y_2 - y_1y_2^2)$ $y_0^3y_2^3$	$y_0^3y_1^4$ $y_0^3y_1^3y_2$ $y_0^3y_1^2y_2^2$ $y_0^3y_1y_2^3$ $y_0^3y_2^4$	$y_0^3(y_1^5 + y_1^4y_2 + y_1^3y_2^2),$ $y_0^3(y_2^5 + y_1y_2^4 + y_1^2y_2^3)$ $y_0^3(y_1^4y_2 + 2y_1^3y_2^2$ $+ 2y_1^2y_2^3 + y_1y_2^4)$	$y_0^3(y_1^6 + 3y_1^5y_2$ $+ 6y_1^4y_2^2 + 7y_1^3y_2^3 + 6y_1^2y_2^4$ $+ 3y_1y_2^5 + y_2^6)$

With these candidates, one can use `Macaulay2` to verify (121) as follows. One can check that $\theta_1, \theta_2, \theta_3$, which have \mathbb{N} -degrees 1, 1, 2, do generate a subalgebra B of $\mathcal{H}_P \subset \mathbb{R}[y_0, y_1, y_2]$ having \mathbb{N} -graded Hilbert series $\frac{1}{(1-t)^2(1-t^2)}$. Thus $\theta_1, \theta_2, \theta_3$ are algebraically independent. And then one can check that together with the six elements colored brown in the table, they generate a subalgebra of \mathcal{H}_P having \mathbb{N} -graded Hilbert series $\frac{1+t+t^2}{(1-t)^3} = \frac{1+2t+2t^2+t^3}{(1-t)^2(1-t^2)}$. Since this matches $\text{Hilb}(\mathcal{H}_P, t) = E_P(t)$, these nine red and brown elements must generate *all* of \mathcal{H}_P . Furthermore, this shows that the six

brown elements must be free B -basis elements for \mathcal{H}_P , so it is Cohen-Macaulay, as predicted in Conjecture 5.5(iv), and \mathcal{H}_P is generated by elements of \mathbb{N}^2 -degrees (b, a) with $b \leq 3 = \#(P \cap \mathbb{Z}^2) - 1$, consistent with Conjecture 5.5(iii).

Remark 5.8. Note that this algorithm computed $E_P(t, q) = \text{Hilb}(\mathcal{H}_P, t, q)$ without finding generators for the ideals $\text{gr } \mathbf{I}(mP \cap \mathbb{Z}^n)$ for all $m \geq 0$. We only had to compute $V_{mP \cap \mathbb{Z}^n} = \text{gr } \mathbf{I}(mP \cap \mathbb{Z}^n)^\perp$ for $m \leq (\#\mathbb{Z}^n \cap P) - 1$. Whenever a lattice polytope P satisfies Conjecture 5.5(iii), this gives a finite algorithm to find a generating set for \mathcal{H}_P and compute $E_P(t, q)$.

Lastly, we use the above description of \mathcal{H}_P to compute the equivariant q -Ehrhart series $E_P^G(t, q)$ in $\text{Rep}_{\mathbb{R}}(G)[q][[t]]$ for the action of the group G of order two that swaps y_1, y_2 . As in Example 3.18, the representation ring $\text{Rep}_{\mathbb{R}}(G) \cong \mathbb{Z}[\epsilon]/(\epsilon^2 - 1) = \text{span}_{\mathbb{Z}}\{1, \epsilon\}$, where $1, \epsilon$ denote the isomorphism classes of the one-dimension trivial and nontrivial $\mathbb{R}[G]$ -modules, respectively. Examining the color-coded elements in the above table, one sees that they were chosen so that the red θ_i for $i = 1, 2, 3$ are all G -fixed, while the brown basis elements are either G -fixed (carrying the representation 1), or G -negated (carrying the representation ϵ), or swapped as a pair (carrying the regular representation $1 + \epsilon$). Consequently, one has

$$(122) \quad E_P^G(t, q) = \text{ch}_{t,q}(\mathcal{H}_P) = \frac{1 + (1 + \epsilon)qt + (1 + \epsilon)q^2t^2 + \epsilon q^3t^3}{(1 - t)(1 - q^2t)(1 - q^3t^2)}.$$

5.7. Some cautionary remarks. We give here a few further cautionary remarks regarding the definition of \mathcal{H}_P and Conjecture 5.5.

Remark 5.9. One might be tempted to define the harmonic algebra differently, in a more general context. Starting with any finite subset $\mathcal{Z} \subseteq \mathbf{k}^n$, over any field \mathbf{k} , one could introduce an \mathbb{N}^2 -graded \mathbf{k} -subalgebra of the ring $\mathbf{k}[y_0] \otimes_{\mathbf{k}} \mathbb{D}_{\mathbf{k}}(\mathbf{y})$ defined by

$$(123) \quad \mathcal{H}_{\mathcal{Z}} := \bigoplus_{m=0}^{\infty} \mathbf{k} \cdot y_0^m \otimes_{\mathbf{k}} \underbrace{V_{\mathcal{Z} + \dots + \mathcal{Z}}}_{m \text{ times}}$$


Here we adopt the convention that the 0-fold Minkowski sum of \mathcal{Z} with itself is $\{\mathbf{0}\}$, so that $1 = y_0^0 \otimes_{\mathbf{k}} \mathbf{y}^{\mathbf{0}} \in (\mathcal{H}_{\mathcal{Z}})_0$. Theorem 1.3 shows that this does defines a \mathbf{k} -subalgebra of $\mathbf{k}[y_0] \otimes_{\mathbf{k}} \mathbb{D}_{\mathbf{k}}(\mathbf{y})$. However, there are two issues with this definition, even when $\mathbf{k} = \mathbb{R}$.

The first issue is that when one takes $\mathcal{Z} = \mathbb{Z}^n \cap P$ for a lattice polytope $P \subset \mathbb{R}^n$, the above algebra $\mathcal{H}_{\mathcal{Z}}$ can be a *proper subalgebra* of the harmonic algebra \mathcal{H}_P . This is because the inclusion (115) can be strict, for lattice polytopes P failing to have the *integer decomposition property (IDP)*

$$\underbrace{(\mathbb{Z}^n \cap P) + \dots + (\mathbb{Z}^n \cap P)}_{m \text{ times}} = \mathbb{Z}^n \cap mP \quad \text{for all } m \geq 1.$$

See Braun [7] and Cox, Haase, Hibi and Higashitani [9] for more on the IDP.

Example 5.10. All lattice polytopes P with $\dim(P) \leq 2$ have the IDP. The smallest non-IDP lattice polytope is a *Reeve tetrahedron* $P = \text{conv}\{(0, 0, 0), (1, 0, 0), (0, 1, 0), (1, 1, 2)\}$:



It is not IDP because $(1, 1, 1) \in \mathbb{Z}^3 \cap 2P \setminus ((\mathbb{Z}^3 \cap 2P) + \mathbb{Z}^3 \cap 2P)$. One can compute for P that

$$E_P(t) = \frac{1+t^2}{(1-t)^4},$$


$$E_P(t, q) = \frac{(1+qt)(1+q^2t^2)(1+qt+q^2t^2)}{(1-t)(1-qt)(1-q^3t^2)(1-q^4t^3)}.$$

Although not every d -dimensional lattice polytope P has the IDP, a result of Cox, Haase, Hibi and Higashitani [9] shows that its dilation dP always has the IDP; see [9]. It follows that A_P is always generated by elements of \mathbb{N} -degree at most $d = \dim(P)$. However, the interesting lattice triangle P discussed in Section 5.6 required an algebra generator for \mathcal{H}_P of \mathbb{N} -degree $3 > 2 = d$, showing that sometimes we must go to higher degrees than d to generate \mathcal{H}_P .

A second issue with the above definition $\mathcal{H}_{\mathcal{Z}}$ is that it is not always finitely generated as an algebra, even when $\mathbf{k} = \mathbb{R}$ and $n = 1$. For example, let $\mathcal{Z} = \{0, 2, 3\} \subset \mathbb{R}^1$. It is not hard to check that the m -fold Minkowski sum $\mathcal{Z} + \dots + \mathcal{Z} = \{0, 2, 3, 4, \dots, 3m-1, 3m\}$ for all $m \geq 1$. This means that, after identifying $\mathbb{D}_{\mathbb{R}}(\mathbf{y})$ with $\mathbb{R}[y_1]$, one has $\underbrace{V_{\mathcal{Z} + \dots + \mathcal{Z}}}_{m \text{ times}} = \text{span}_{\mathbb{R}}\{1, y_1, y_1^2, \dots, y_1^{3m-1}\}$, and

$$(124) \quad \mathcal{H}_{\mathcal{Z}} = \text{span}_{\mathbb{R}}\{1\} \oplus \text{span}_{\mathbb{R}}\{y_0^m y_1^j : m \geq 0 \text{ and } 0 \leq j \leq 3m-1\}.$$

Therefore $\mathcal{H}_{\mathcal{Z}} \subset \mathbb{R}[y_0, y_1]$ is the semigroup ring for the additive subsemigroup of \mathbb{N}^2 that may be visualized as follows, with a dot at coordinates (m, j) representing $y_0^m y_1^j$ in $\mathcal{H}_{\mathcal{Z}}$:



This illustrates the lack of finite generation: minimal monomial \mathbb{R} -algebra generators for $\mathcal{H}_{\mathcal{Z}}$ are

$$\{y_0 y_1^2, y_0 y_1^3, y_0^2 y_1^5, y_0^3 y_1^8, y_0^4 y_1^{11}, \dots\} = \{y_0 y_1^2\} \cup \{y_0^m y_1^{3m-1}\}_{m=2,3,4,\dots}.$$

This does not violate Conjecture 5.5(i), since $\mathcal{H}_{\mathcal{Z}} \not\cong \mathcal{H}_P$ for any lattice polytope P . But proving Conjecture 5.5 must involve extra features of harmonic spaces of lattice points inside polytopes.

Remark 5.11. One can regard $P \rightsquigarrow \mathcal{H}_P$ as a functor $\mathcal{H} : \mathcal{C} \rightarrow \mathcal{D}$, where \mathcal{D} is the category of \mathbb{N}^2 -graded \mathbb{R} -algebras. Here \mathcal{C} is a category of lattice polytopes $P \subset \mathbb{R}^n$ whose morphisms $P \rightarrow P'$ are \mathbb{Z} -linear maps $\varphi : \mathbb{Z}^n \rightarrow \mathbb{Z}^{n'}$ with $f(P) \subseteq P'$. The fact that this induces an \mathbb{R} -algebra map $\mathcal{H}_P \rightarrow \mathcal{H}_{P'}$ stems from the fact that the point configurations $\mathcal{Z} := \mathbb{Z}^n \cap mP$ and $\mathcal{Z}' := \mathbb{Z}^{n'} \cap mP'$ have $f(\mathcal{Z}) \subseteq \mathcal{Z}'$. This implies that the map $\varphi^\sharp : \mathbb{R}[x_1, \dots, x_{n'}] \rightarrow \mathbb{R}[x_1, \dots, x_n]$ that precomposes $f \mapsto f \circ \varphi$ will have $\varphi^\sharp(\mathbf{I}(\mathcal{Z}')) \subseteq \mathbf{I}(\mathcal{Z})$, and hence $\varphi^\sharp(\text{gr } \mathbf{I}(\mathcal{Z}')) \subseteq \text{gr } \mathbf{I}(\mathcal{Z})$. Therefore the adjoint map $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^{n'}$, when extended to a ring map $\mathbb{R}[y_1, \dots, y_n] \rightarrow \mathbb{R}[y_1, \dots, y_{n'}]$ satisfies

$$\varphi(V_{\mathcal{Z}}) = \varphi(\text{gr } \mathbf{I}(\mathcal{Z})^\perp) \subseteq \text{gr } \mathbf{I}(\mathcal{Z}')^\perp = \varphi(V_{\mathcal{Z}'}).$$

If one instead considers $\mathcal{H} : P \rightsquigarrow \mathcal{H}_P$ as a functor into the category of \mathbb{N} -graded \mathbb{R} -algebras, one can compare it to the similar such functor $A : P \rightsquigarrow A_P$. These two functors $\mathcal{H}_{(-)}$ and $A_{(-)}$ are not isomorphic; see the example in Section 5.4. The authors do not know if they are equivalent.

Remark 5.12. One might hope to find some canonical \mathbb{R} -linear basis for \mathcal{H}_P . Although the authors are unaware of such a basis which is \mathbb{N}^2 -homogeneous, there is at least a canonical \mathbb{N} -homogeneous \mathbb{R} -basis, coming from such \mathbb{R} -bases for harmonic spaces $V_{\mathcal{Z}}$ of finite subsets $\mathcal{Z} \subset \mathbb{R}^n$.

To describe these, assume \mathbf{k} is a field of characteristic zero, and identify \mathbf{k}^n and its basis y_1, \dots, y_n with $(\mathbf{k}^n)^*$ and its basis x_1, \dots, x_n . Then mapping $y_n \mapsto x_n$ gives the middle isomorphism here

$$\mathbb{D} := \mathbb{D}_{\mathbf{k}}(\mathbf{y}) = \mathbf{k}[y_1, \dots, y_n] \cong \mathbf{k}[x_1, \dots, x_n] =: S.$$

This lets one view the subspace $V_{\mathcal{Z}} := \text{gr } \mathbf{I}(\mathcal{Z})^\perp \subset \mathbb{D}$ as a subspace of $S = \mathbf{k}[\mathbf{x}]$.

Proposition 5.13. *For \mathbf{k} a field of characteristic zero, and any finite point set $\mathcal{Z} \subset \mathbf{k}^n$, there is a unique basis $\{g_{\mathbf{z}}(\mathbf{x}) : \mathbf{z} \in \mathcal{Z}\}$ of the harmonic space $V_{\mathcal{Z}}$ such that $g_{\mathbf{z}}(\mathbf{z}') = \delta_{\mathbf{z}, \mathbf{z}'}$ for all $\mathbf{z}' \in \mathcal{Z}$.*

Proof. Since $V_{\mathcal{Z}} := \text{gr } \mathbf{I}(\mathcal{Z})^\perp$, one has a \mathbf{k} -vector space direct sum decomposition $S = V_{\mathcal{Z}} \oplus \text{gr } \mathbf{I}(\mathcal{Z})$. Consequently, the composite map $V_{\mathcal{Z}} \hookrightarrow S \twoheadrightarrow S/\text{gr } \mathbf{I}(\mathcal{Z})$ is a \mathbf{k} -vector space isomorphism. Thus any set of homogeneous polynomials that give a \mathbf{k} -basis for the (graded) subspace $V_{\mathcal{Z}} \subset S$ will descend to a \mathbf{k} -basis of $S/\text{gr } \mathbf{I}(\mathcal{Z})$. Proposition 2.1(ii) implies that the same set of polynomials will also descend to a basis of $S/\mathbf{I}(\mathcal{Z})$. Hence this composite is also a \mathbf{k} -vector space isomorphism:

$$(125) \quad V_{\mathcal{Z}} \hookrightarrow S \twoheadrightarrow S/\mathbf{I}(\mathcal{Z}).$$

Now multivariate Lagrange interpolation identifies

$$(126) \quad S/\mathbf{I}(\mathcal{Z}) \cong \{ \text{all functions } g : \mathcal{Z} \rightarrow \mathbf{k} \},$$

and the space on the right has a unique \mathbf{k} -basis of $\{g_{\mathbf{z}}\}_{\mathbf{z} \in \mathcal{Z}}$ defined by $g_{\mathbf{z}}(\mathbf{z}') = \delta_{\mathbf{z}, \mathbf{z}'}$. The assertion then follows from combining this with (125) and (126). \square

Note that Proposition 5.13 provides \mathbf{k} -bases $\{g_{\mathbf{z}}(\mathbf{x}) : \mathbf{z} \in \mathcal{Z}\}$ that are generally inhomogeneous. Nevertheless, given a lattice polytope P , and taking $\mathbf{k} = \mathbb{R}$, one can assemble all of these distinguished \mathbb{R} -bases for $V_{\mathbb{Z}^n \cap mP}$ into an \mathbb{N} -graded \mathbb{R} -basis $\{y_0^m g_{\mathbf{z}}(\mathbf{y}) : \mathbf{z} \in \mathbb{Z}^n \cap mP\}$ for \mathcal{H}_P . Note that, even though this \mathbb{R} -basis for \mathcal{H}_P is in bijection with an \mathbb{R} -basis for the semigroup ring A_P , the bijection does not respect multiplication. This is to be expected, since the example discussed in Section 5.4 shows that these two rings are not isomorphic as \mathbb{N} -graded \mathbb{R} -algebras.

6. DILATIONS, PRODUCTS, AND JOINS: PROOF OF THEOREM 1.2

Recall from the Introduction these three basic operations on polytopes:

- *dilation* by an positive integer factor d , sending P to dP ,
- *Cartesian product*, sending $P \subset \mathbb{R}^n$ and $Q \subset \mathbb{R}^m$ to $P \times Q \subset \mathbb{R}^{n+m}$,
- *free join*, sending P, Q to $P * Q \subseteq \mathbb{R}^{1+n+m}$ defined by

$$\begin{aligned} P * Q &:= \{(t, t\mathbf{p}, (1-t)\mathbf{q}) : 0 \leq t \leq 1, \mathbf{p} \in P, \mathbf{q} \in Q\} \\ &= \text{conv} (\quad \{1\} \times P \times \mathbf{0}_m \quad \sqcup \quad \{0\} \times \mathbf{0}_n \times Q \quad) \end{aligned}$$

We wish to understand how these operations interact with q -Ehrhart series $E_P(t, q)$. In the process we will see how they interact with harmonic algebras \mathcal{H}_P , by relating them to the commutative algebra constructions for \mathbb{N} -graded algebras of *Veronese subalgebras*, *Segre products* and *graded tensor products*; When applying these constructions to the \mathbb{N}^2 -graded algebra \mathcal{H}_P , we will always use the grading specialization $\deg(y_0^m \mathbf{y}^{\mathbf{z}}) = m$ to regard it as an \mathbb{N} -graded algebra. The analysis will also prove Theorem 1.2, whose statement we recall here.

Theorem 1.2. *Let P, Q be lattice polytopes.*

- (i) *For positive integers d , the dilation dP has $E_{dP}(t, q)$ given by*

$$E_{dP}(t, q) = \sum_{m=0}^{\infty} i_P(dm; q) t^m.$$

(ii) The Cartesian product $P \times Q$ has $E_{P \times Q}(t, q)$ given by the Hadamard product

$$E_{P \times Q}(t, q) = \sum_{m=0}^{\infty} i_P(m; q) \cdot i_Q(m; q) \cdot t^m.$$

(iii) The free join $P * Q$ has $E_{P * Q}(t, q)$ given by

$$E_{P * Q}(t, q) = \frac{1-t}{1-qt} \cdot E_P(t, q) \cdot E_Q(t, q).$$

We deal with each of the three parts of this theorem in the next three subsections.

6.1. Dilations and the Veronese construction. The first and simplest of these is given by dilation $P \mapsto dP$. If $A = \bigoplus_{m \geq 0} A_m$ is a graded algebra, the d^{th} Veronese subalgebra is

$$(127) \quad \text{Ver}_d(A) := \bigoplus_{m \geq 0} A_{dm}.$$

Then Theorem 1.2(i) along with the following result are immediate from the definitions.

Proposition 6.1. *For a lattice polytope P and any $d \geq 0$ one has $\mathcal{H}_{dP} \cong \text{Ver}_d(\mathcal{H}_P)$.*

6.2. Cartesian products and the Segre construction. The Cartesian product $P \times Q$ of lattice polytopes is again a lattice polytope. Since $m(P \times Q) = mP \times mQ$, it is easily seen that

$$i_{P \times Q}(m) = i_P(m) \cdot i_Q(m).$$

There is a similarly easy relation for their harmonic algebras, which comes from a general lemma on the behavior of orbit harmonic rings $R(\mathcal{Z}) = S/\text{gr } \mathbf{I}(\mathcal{Z})$ and harmonic spaces $V_{\mathcal{Z}}$ when one has a Cartesian product of point loci.

Lemma 6.2. *For a field \mathbf{k} and finite subsets $\mathcal{Z} \subseteq \mathbf{k}^n, \mathcal{Z}' \subseteq \mathbf{k}^{n'}$, one has graded isomorphisms*

$$\begin{aligned} R(\mathcal{Z} \times \mathcal{Z}') &\cong R(\mathcal{Z}) \otimes_{\mathbf{k}} R(\mathcal{Z}'), \\ V_{\mathcal{Z} \times \mathcal{Z}'} &\cong V_{\mathcal{Z}} \otimes_{\mathbf{k}} V_{\mathcal{Z}'}. \end{aligned}$$

Proof. Write $S_n, S_{n'}$ for the polynomial rings $\mathbf{k}[x_1, \dots, x_n], \mathbf{k}[x_1, \dots, x_{n'}]$ which contain $\mathbf{I}(\mathcal{Z}), \mathbf{I}(\mathcal{Z}')$, so that $\mathbf{I}(\mathcal{Z} \times \mathcal{Z}')$ will be an ideal in $S_n \otimes S_{n'}$. We claim one has the following equality of ideals

$$(128) \quad I := \text{gr } \mathbf{I}(\mathcal{Z} \times \mathcal{Z}') = \text{gr } \mathbf{I}(\mathcal{Z}) \otimes S_{n'} + S_n \otimes \text{gr } \mathbf{I}(\mathcal{Z}) =: J,$$

from which the remaining assertions will follow. To see this claim, first check the containment $I \supseteq J$: for $f \in \mathbf{I}(\mathcal{Z})$ and $f' \in \mathbf{I}(\mathcal{Z}')$, one has both $f \otimes 1$ and $1 \otimes f'$ lying in $\mathbf{I}(\mathcal{Z} \times \mathcal{Z}')$, and if $f, f' \neq 0$, their top degree components $\tau(f \otimes 1) = \tau(f) \otimes 1$ and $\tau(1 \otimes f') = 1 \otimes \tau(f')$ both lie in I . On the other hand, the inclusion $I \supseteq J$ must be an equality because the surjection

$$(S_n \otimes S_{n'}) / J = \twoheadrightarrow (S_n \otimes S_{n'}) / I$$

is an isomorphism via dimension-counting:

$$\dim_{\mathbf{k}}(S_n \otimes S_{n'}) / J \leq \dim_{\mathbf{k}} S_n / \text{gr } \mathbf{I}(\mathcal{Z}) \cdot \dim_{\mathbf{k}} S_{n'} / \text{gr } \mathbf{I}(\mathcal{Z}') = \#(\mathcal{Z} \times \mathcal{Z}') = \dim_{\mathbf{k}}(S_n \otimes S_{n'}) / I. \quad \square$$

For \mathbb{N} -graded \mathbf{k} -algebras A, B , their Segre product $\text{Segre}(A, B)$ is the graded algebra

$$(129) \quad \text{Segre}(A, B) = \bigoplus_{m=0}^{\infty} \underbrace{A_m \otimes_{\mathbf{k}} B_m}_{\text{Segre}(A, B)_m}$$

Proposition 6.3. *For any lattice polytopes P and Q , one has $\mathcal{H}_{P \times Q} \cong \text{Segre}(\mathcal{H}_P, \mathcal{H}_Q)$.*

Proof. This comes via Lemma 6.2 and the definitions, as $\mathbb{Z}^n \cap m(P \times Q) = (\mathbb{Z}^n \cap mP) \times (\mathbb{Z}^n \cap mQ)$ \square

6.3. Joins. Recall that for polytopes $P, Q \subseteq \mathbb{R}^n, \mathbb{R}^{n'}$, their (free) join $P * Q \subseteq \mathbb{R}^{1+n+n'}$ is

$$P * Q := \{(t, t\mathbf{p}, (1-t)\mathbf{q}) : 0 \leq t \leq 1, \mathbf{p} \in P, \mathbf{q} \in Q\} = \text{conv}(\{1\} \times P \times \mathbf{0}_{n'} \sqcup \{0\} \times \mathbf{0}_n \times Q)$$


If P and Q are lattice polytopes, the join $P * Q$ is again a lattice polytope. The effect of join on harmonic algebras is more complicated than dilation and Cartesian product. We will eventually see, in Theorem 6.6 below, that it is almost, but not quite, the \mathbb{N} -graded tensor product

$$A \otimes B = \bigoplus_{m=0}^{\infty} \underbrace{\bigoplus_{i+j=m} A_i \otimes B_j}_{(A \otimes B)_m}.$$

We first recall some facts on joins of lattice polytopes P, Q in $\mathbb{R}^n, \mathbb{R}^{n'}$. Let x_0 denote the extra first coordinate in the ambient space $\mathbb{R}^1 \times \mathbb{R}^n \times \mathbb{R}^{n'} = \mathbb{R}^{1+n+n'}$ for the join $P * Q$. Slicing by the hyperplanes $x_0 = r$ gives a disjoint decomposition of the dilates $m(P * Q)$ for $m \geq 0$:

$$(130) \quad \mathbb{Z}^{1+n+n'} \cap m(P * Q) = \bigsqcup_{\substack{(r,r'): \\ r+r'=m}} \{r\} \times (\mathbb{Z}^n \cap rP) \times (\mathbb{Z}^{n'} \cap r'Q).$$

We illustrate (130) for $m(P * Q)$ with $m = 1, 2, 3$, when $P = [0, 1] \subset \mathbb{R}^1$ and $Q = [0, 2] \subset \mathbb{R}^1$:



This decomposition (130) implies a well-known fact (see, e.g., Beck and Robins [4, Exer. 3.33]):

$$(131) \quad i_{P*Q}(m) = \sum_{\substack{(r,r'): \\ r+r'=m}} i_P(r) \cdot i_Q(r')$$

This fact (131) is easily seen to be equivalent to the following simple relation of Ehrhart series:

$$(132) \quad E_{P*Q}(t) = E_P(t) \cdot E_Q(t).$$

Recalling that $E_P(t, q) = \text{Hilb}(\mathcal{H}_P, t, q)$, the next result is the q -analogue of (132).

Theorem 6.4. *For lattice polytopes P, Q , one has the following relation among the \mathbb{N}^2 -graded Hilbert series for the harmonic algebras $\mathcal{H}_P, \mathcal{H}_Q, \mathcal{H}_{P*Q}$:*

$$E_{P*Q}(t, q) = \frac{1-t}{1-qt} \cdot E_P(t, q) \cdot E_Q(t, q).$$

As mentioned in Remark 3.25, if we take Q to be a single point, Theorem 6.4 says that the q -Ehrhart series of the pyramid $\text{Pyr}(P) := P * \{\text{pt}\}$ is $E_{\text{Pyr}(P)}(t, q) = (1-qt) \cdot E_P(t, q)$.

Proof. Letting $\mathcal{Z}_m := \mathbb{Z}^{1+n+n'} \cap m(P * Q) \subset \mathbb{R}^{1+n+n'}$, we start by describing the ideal

$$\mathrm{gr} \mathbf{I}(\mathcal{Z}_m) \subset \mathbb{R}[x_0] \otimes \mathbb{R}[x_1, \dots, x_n] \otimes \mathbb{R}[x_1, \dots, x_{n'}].$$

The product structure in each term of (130) leads to some natural elements of $\mathbf{I}(\mathcal{Z}_m)$ and $\mathrm{gr} \mathbf{I}(\mathcal{Z}_m)$. First translate P, Q , without loss of generality, so that $\mathbf{0}_n \in P$ and $\mathbf{0}_{n'} \in Q$, implying nestings

$$(133) \quad P \subseteq 2P \subseteq 3P \subseteq \dots \text{ and } Q \subseteq 2Q \subseteq 3Q \subseteq \dots$$

Suppose $\ell, \ell' \geq 0$ have $\ell + \ell' \leq m$, and one is given $f \in \mathbf{I}(\mathbb{Z}^n \cap \ell P)$ and $f' \in \mathbf{I}(\mathbb{Z}^{n'} \cap \ell' Q)$. Then we claim that the nestings (133) imply that the polynomial

$$(134) \quad \hat{f} := \prod_{r=\ell+1}^{m-(\ell'+1)} (x_0 - r) \otimes f \otimes f'$$

vanishes on \mathcal{Z}_ℓ . To see this, note that a typical point of $(r, \mathbf{z}, \mathbf{z}')$ of \mathcal{Z}_ℓ will either

- have $\ell + 1 \leq r \leq m - (\ell' + 1)$ so \hat{f} vanishes due to one of its factors $x_0 - r$,
- or else have $r \leq \ell$, and then \hat{f} vanishes due to the factor f , since $\mathbf{z} \in rP \subseteq \ell P$,
- or else have $m - r \leq \ell'$, and then \hat{f} vanishes due to the factor f' , since $\mathbf{z}' \in (m - r)P \subseteq \ell' P$

Thus \hat{f} lies in $\mathbf{I}(\mathcal{Z}_m)$, implying that $\mathrm{gr} \mathbf{I}(\mathcal{Z}_m)$ contains its top degree homogeneous component

$$(135) \quad \tau(\hat{f}) = x_0^{m-\ell-\ell'} \otimes \tau(f) \otimes \tau(f').$$

Note also that $x_0(x_0 - 1) \cdots (x_0 - m)$ vanishes on \mathcal{Z}_m . Hence $\mathrm{gr} \mathbf{I}(\mathcal{Z}_m)$ contains this ideal:

$$(136) \quad J_m := (x_0^{m+1} \otimes 1 \otimes 1) + \left((x_0^{m-\ell-\ell'}) \otimes \sum_{\ell+\ell' \leq m} \mathrm{gr} \mathbf{I}(\mathbb{Z}^n \cap \ell P) \otimes \mathrm{gr} \mathbf{I}(\mathbb{Z}^{n'} \cap \ell' Q) \right).$$

We wish to show that, in fact, this containment $J_m \subseteq \mathrm{gr} \mathbf{I}(\mathcal{Z}_m)$ is an equality. To this end, we calculate the perp space J_m^\perp within $\mathbb{R}[y_0] \otimes \mathbb{R}[y_1, \dots, y_n] \otimes \mathbb{R}[y_1, \dots, y_{n'}]$ for the ideal J_m . Abbreviate

$$(137) \quad V_\ell^P := \mathrm{gr} \mathbf{I}(\mathbb{Z}^n \cap \ell P)^\perp \quad \text{and} \quad V_{\ell'}^Q := \mathrm{gr} \mathbf{I}(\mathbb{Z}^{n'} \cap \ell' P)^\perp.$$

By Lemma 5.3 we have

$$(138) \quad \mathbb{R} \cdot 1 = V_0^P \subseteq V_1^P \subseteq \dots \subseteq V_\ell^P \quad \text{and} \quad \mathbb{R} \cdot 1 = V_0^Q \subseteq V_1^Q \subseteq \dots \subseteq V_{\ell'}^Q.$$

The following space V_m of polynomials is annihilated by J_m under the \odot -action, so lies in J_m^\perp :

$$(139) \quad V_m := \bigoplus_{s=0}^m \left(\sum_{\ell+\ell'=s} \mathbb{R} \cdot y_0^{m-s} \otimes V_\ell^P \otimes V_{\ell'}^Q \right).$$

By (144) in Lemma 6.5 below, V_m has the following dimension:

$$\begin{aligned} \dim_{\mathbb{R}} V_m &= \sum_{s=0}^m \dim \left(\sum_{\ell+\ell'=s} V_\ell^P \otimes V_{\ell'}^Q \right) = \sum_{r+r'=m} \dim(V_r^P) \cdot \dim(V_{r'}^Q) \\ &= \sum_{r+r'=m} i_P(r) \cdot i_Q(r') = i_{P*Q}(m) = \#\mathcal{Z}_m. \end{aligned}$$

Therefor these inequalities

$$\#\mathcal{Z}_m = \dim_{\mathbb{R}} V_m \leq \dim_{\mathbb{R}} J_m^\perp = \dim_{\mathbb{R}} S/J_m \leq \dim_{\mathbb{R}} S/\mathrm{gr} \mathbf{I}(\mathcal{Z}_m) = \#\mathcal{Z}_m$$

must all be equalities, implying

$$(140) \quad \text{gr } \mathbf{I}(\mathcal{Z}_m) = J_m,$$

$$(141) \quad V_{\mathcal{Z}_m} = \text{gr } \mathbf{I}(\mathcal{Z}_m)^\perp = J_m^\perp = V_m.$$

This now allows us to compute $\text{Hilb}(V_{\mathcal{Z}_m}, q)$, using (139), as

$$(142) \quad \text{Hilb}(V_{\mathcal{Z}_m}, q) = \text{Hilb}(V_m, q) = \sum_{s=0}^m q^{m-s} \text{Hilb} \left(\sum_{\ell+\ell'=s} V_\ell^P \otimes V_{\ell'}^Q, q \right).$$

On the other hand, direct calculation shows that the coefficient of t^m in $\frac{1-t}{1-qt} E_P(t, q) \cdot E_Q(t, q)$ is

$$\sum_{s=0}^m q^{m-s} \sum_{\ell+\ell'=s} \text{Hilb}(V_\ell^P \otimes V_{\ell'}^Q, q) - \sum_{s=0}^{m-1} q^{m-1-s} \sum_{\ell+\ell'=s} \text{Hilb}(V_\ell^P \otimes V_{\ell'}^Q, q).$$

This last expression coincides, by (143) in Lemma 6.5 below, with (142). In other words,

$$\frac{1-t}{1-qt} E_P(t, q) \cdot E_Q(t, q) = \sum_{m=0}^{\infty} t^m \cdot \text{Hilb}(V_{\mathcal{Z}_m}, q) = \text{Hilb}(\mathcal{H}_{P*Q}, t, q) = E_{P*Q}(t, q). \quad \square$$

The previous proof used the following technical lemma, on Hilbert series arising from graded tensor products $V \otimes V'$ of spaces V, V' equipped with homogeneous filtrations.

Lemma 6.5. *Let V, V' be graded vector spaces, each with nested sequences of homogeneous subspaces*

$$0 \subseteq V_1 \subseteq V_2 \subseteq \cdots \subseteq V \quad \text{and} \quad 0 \subseteq V'_1 \subseteq V'_2 \subseteq \cdots \subseteq V'$$

Then one has the following Hilbert series identity for each n :

$$(143) \quad \sum_{k=0}^n q^{n-k} \cdot \text{Hilb} \left(\sum_{a+b=k} V_a \otimes V'_b, q \right) = \sum_{k=0}^n q^{n-k} \sum_{a+b=k} \text{Hilb}(V_a \otimes V'_b, q) - \sum_{k=0}^{n-1} q^{n-k-1} \sum_{a+b=k} \text{Hilb}(V_a \otimes V'_b, q).$$

In particular, setting $q = 1$, one has

$$(144) \quad \sum_{k=0}^n \dim \left(\sum_{a+b=k} V_a \otimes V'_b \right) = \sum_{c+d=n} \dim(V_c) \cdot \dim(V'_d).$$

Proof. Since the spaces V_a and V'_b nest, the sum $\sum_{a+b=k} V_a \otimes V'_b$ of subspaces of $V \otimes V'$ is *not* a direct sum. However, after abbreviating

$$h_{ab} := \text{Hilb}(V_a/V_{a-1}, q) \cdot \text{Hilb}(V'_b/V'_{b-1}, q) = \text{Hilb}(V_a/V_{a-1} \otimes V'_b/V'_{b-1}, q),$$

the left side of the lemma can be rewritten

$$(145) \quad \sum_{k=0}^n q^{n-k} \cdot \sum_{a+b \leq k} h_{ab},$$

whose coefficient of h_{ab} is

$$\sum_{k=a+b}^n q^{n-k} = 1 + q + q^2 + \cdots + q^{n-(a+b)} = [\hat{n}]_q$$

abbreviating $\hat{n} := n - (a + b) + 1$. Meanwhile, the right side of the lemma can be rewritten

$$(146) \quad \sum_{k=0}^n q^{n-k} \sum_{a'+b'=k} \sum_{\substack{a:0 \leq a \leq a' \\ b:0 \leq b \leq b'}} h_{ab} - \sum_{k=0}^{n-1} q^{n-1-k} \sum_{a'+b'=k} \sum_{\substack{a:0 \leq a \leq a' \\ b:0 \leq b \leq b'}} h_{ab}.$$

Hence the coefficient of h_{ab} on the right side of the lemma is

$$\begin{aligned} & \sum_{\substack{(a',b'): \\ a' \geq a \\ b' \geq b \\ a'+b' \leq n}} q^{n-(a'+b')} - \sum_{\substack{(a',b'): \\ a' \geq a \\ b' \geq b \\ a'+b' \leq n-1}} q^{n-1-(a'+b')} \\ &= \hat{n}q^0 + (\hat{n}-1)q^1 + (\hat{n}-2)q^2 + \cdots + 2q^{\hat{n}-2} + q^{\hat{n}-1} \\ & \quad - ((\hat{n}-1)q^0 + (\hat{n}-2)q^1 + (\hat{n}-3)q^2 + \cdots + 2q^{\hat{n}-3} + q^{\hat{n}-2}) \\ &= q^0 + q^1 + \cdots + q^{\hat{n}-2} + q^{\hat{n}-1} = [\hat{n}]_q \quad \square \end{aligned}$$

We are ready to relate the harmonic algebras \mathcal{H}_{P^*Q} , \mathcal{H}_P , and \mathcal{H}_Q , starting with the following suggestive rewriting of Theorem 6.4:

$$(147) \quad (1 - qt) \cdot \text{Hilb}(\mathcal{H}_{P^*Q}, t, q) = (1 - t) \cdot \text{Hilb}(\mathcal{H}_P \otimes \mathcal{H}_Q, t, q).$$

To interpret this identity, it helps to name the ‘‘auxiliary’’ variables in \mathcal{H}_{P^*Q} , \mathcal{H}_P , \mathcal{H}_Q as y_{P^*Q} , y_P , y_Q , and to abbreviate the other variable sets $\mathbf{y}_n := (y_1, \dots, y_n)$, $\mathbf{y}_{n'} := (y_1, \dots, y_{n'})$. This means that the rings \mathcal{H}_{P^*Q} , $\mathcal{H}_P \otimes \mathcal{H}_Q$, are \mathbb{R} -subalgebras of the following ambient polynomial algebras

$$\begin{aligned} \mathcal{H}_{P^*Q} &\subset \mathbb{R}[y_{P^*Q}] \otimes \mathbb{R}[y_0] \otimes \mathbb{R}[\mathbf{y}_n] \otimes \mathbb{R}[\mathbf{y}_{n'}] \\ \mathcal{H}_P \otimes \mathcal{H}_Q &\subset \mathbb{R}[y_P, \mathbf{y}_n] \otimes \mathbb{R}[y_Q, \mathbf{y}_{n'}]. \end{aligned}$$

Their \mathbb{N}^2 -gradings have $\deg(y_P) = \deg(y_Q) = \deg(y_{P^*Q}) = (1, 0)$, and all other variables have $\deg(y_0) = \deg(y_i) = (0, 1)$. Since the ambient polynomial algebras are integral domains, so are the harmonic algebras inside them. Hence one concludes that

- $y_{P^*Q} \cdot y_0$ ($:= y_{P^*Q} \otimes y_0 \otimes 1 \otimes 1$) is a nonzero divisor inside \mathcal{H}_{P^*Q} , with \mathbb{N}^2 -degree $(1, 1)$, and
- $y_P \otimes 1 - 1 \otimes y_Q$ is a nonzero divisor inside $\mathcal{H}_P \otimes \mathcal{H}_Q$, with \mathbb{N}^2 -degree $(1, 0)$.

In a \mathbb{N}^2 -graded algebra A with a homogeneous nonzerodivisor θ of degree (a, b) , the exact sequence

$$0 \rightarrow A(-a, b) \xrightarrow{-\theta} A \rightarrow A/(\theta) \rightarrow 0$$

shows that one has the Hilbert series relation

$$\text{Hilb}(A/(\theta), t, q) = (1 - t^a q^b) \cdot \text{Hilb}(A, t, q).$$

Thus the left and right sides of (147) can be reinterpreted as Hilberts series for these quotient rings:

$$\begin{aligned} (1 - qt) \cdot \text{Hilb}(\mathcal{H}_{P^*Q}, t, q) &= \text{Hilb}(\mathcal{H}_{P^*Q}/(y_{P^*Q} \cdot y_0), t, q), \\ (1 - t) \cdot \text{Hilb}(\mathcal{H}_P \otimes \mathcal{H}_Q, t, q) &= \text{Hilb}((\mathcal{H}_P \otimes \mathcal{H}_Q)/(y_P \otimes 1 - 1 \otimes y_Q), t, q). \end{aligned}$$

In this way, the following result is an algebraic strengthening of Theorem 6.4.

Theorem 6.6. *For any lattice polytopes P and Q , one has an \mathbb{N}^2 -graded algebra isomorphism*

$$(\mathcal{H}_P \otimes \mathcal{H}_Q)/(y_P \otimes 1 - 1 \otimes y_Q) \cong \mathcal{H}_{P^*Q}/(y_{P^*Q} \cdot y_0)$$

Proof. In light of the preceding discussion on Hilbert series, it suffices by dimension-counting to exhibit an \mathbb{N}^2 -graded \mathbb{R} -algebra surjection from the left side to the right side in the theorem.

Consider the \mathbb{R} -linear map φ defined via

$$\begin{aligned} \mathbb{R}[y_P, \mathbf{y}_n] \otimes \mathbb{R}[y_Q, \mathbf{y}_{n'}] &\longrightarrow \mathbb{R}[y_{P*Q}] \otimes \mathbb{R}[y_0] \otimes \mathbb{R}[\mathbf{y}_n] \otimes \mathbb{R}[\mathbf{y}_{n'}] \\ y_P^\ell \cdot g(\mathbf{y}_n) \otimes y_Q^{\ell'} \cdot g'(\mathbf{y}_{n'}) &\longmapsto y_{P*Q}^{\ell+\ell'} \otimes 1 \otimes g(\mathbf{y}_n) \otimes g'(\mathbf{y}_{n'}). \end{aligned}$$

One can readily check that φ

- is actually a map of \mathbb{N}^2 -graded \mathbb{R} -algebras,
- that it restricts to a map $\mathcal{H}_P \otimes \mathcal{H}_Q \rightarrow \mathcal{H}_{P*Q}$, using (139), (141), and
- that $y_P \otimes 1 - 1 \otimes y_Q$ lies in its kernel.

Consequently, φ descends to a map on the quotient

$$(148) \quad \varphi : (\mathcal{H}_P \otimes \mathcal{H}_Q) / (y_P \otimes 1 - 1 \otimes y_Q) \longrightarrow \mathcal{H}_{P*Q}.$$

This map is not surjective. However, after post-composing it with the canonical projection map $\pi : \mathcal{H}_{P*Q} \twoheadrightarrow \mathcal{H}_{P*Q} / (y_{P*Q} y_0)$, then we claim that

$$(149) \quad \pi \circ \varphi : (\mathcal{H}_P \otimes \mathcal{H}_Q) / (y_P \otimes 1 - 1 \otimes y_Q) \longrightarrow \mathcal{H}_{P*Q} / (y_{P*Q} y_0)$$

actually *is* the surjection that we seek. To see this, note that the image of φ within \mathcal{H}_{P*Q} is

$$\bigoplus_{m=0}^{\infty} \sum_{\ell+\ell'=m} \mathbb{R} \cdot y_{P*Q}^m \cdot y_0^0 \otimes V_\ell^P \otimes V_{\ell'}^Q.$$

On the other hand, (139) shows the remaining part of \mathcal{H}_{P*Q} not lying in this image has the form

$$\bigoplus_{m=1}^{\infty} \bigoplus_{s=0}^{m-1} \left(\sum_{\ell+\ell'=s} \mathbb{R} \cdot y_{P*Q}^m \cdot y_0^{m-s} \otimes V_\ell^P \otimes V_{\ell'}^Q \right),$$

which is contained in the ideal $(y_{P*Q} y_0)$ and therefore vanishes in the quotient $\mathcal{H}_{P*Q} / (y_{P*Q} y_0)$. \square

REFERENCES

- [1] K. Akin, D. A. Buchsbaum, and J. Weyman. Schur functors and Schur complexes. *Adv. in Math.*, 44(3):207–278, 1982.
- [2] D. Armstrong, V. Reiner, and B. Rhoades. Parking spaces. *Advances in Mathematics*, 269:647–706, 2015.
- [3] G. Ballelli. Enumeration of lattice polytopes by their volume. *Discrete Comput. Geom.*, 65(4):1087–1122, 2021.
- [4] M. Beck and S. Robins. *Computing the continuous discretely. Integer-point enumeration in polyhedra. With illustrations by David Austin*. Undergraduate Texts Math. New York, NY: Springer, 2nd edition edition, 2015.
- [5] M. Beck and R. Sanyal. *Combinatorial reciprocity theorems*, volume 195 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2018. An invitation to enumerative geometric combinatorics.
- [6] F. Bergeron. *Algebraic combinatorics and coinvariant spaces*. CMS Treatises in Mathematics. Canadian Mathematical Society, Ottawa, ON; A K Peters, Ltd., Wellesley, MA, 2009.
- [7] B. Braun. Unimodality problems in Ehrhart theory. In *Recent trends in combinatorics*, volume 159 of *IMA Vol. Math. Appl.*, pages 687–711. Springer, [Cham], 2016.
- [8] F. Chapoton. q -analogues of Ehrhart polynomials. *Proc. Edinb. Math. Soc. (2)*, 59(2):339–358, 2016.
- [9] D. A. Cox, C. Haase, T. Hibi, and A. Higashitani. Integer decomposition property of dilated polytopes. *Electron. J. Combin.*, 21(4), 2012. #P4.28.
- [10] D. A. Cox, J. Little, and D. O’Shea. *Ideals, varieties, and algorithms. An introduction to computational algebraic geometry and commutative algebra*. Undergraduate Texts Math. Cham: Springer, 4th revised ed. edition, 2015.
- [11] V. I. Danilov. The geometry of toric varieties. *Uspekhi Mat. Nauk*, 33(2(200)):85–134, 247, 1978.
- [12] E. Ehrhart. Sur les polyèdres rationnels homothétiques à n dimensions. *CR Acad. Sci. Paris*, 254:616, 1962.

- [13] D. Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [14] S. Eliahou and M. Kervaire. Sumsets in vector spaces over finite fields. *Journal of Number Theory*, 71(1):12–39, 1998.
- [15] S. Eliahou and M. Kervaire. Sumsets in vector spaces over finite fields. *J. Number Theory*, 71(1):12–39, 1998.
- [16] D. R. Fulkerson. Blocking and anti-blocking pairs of polyhedra. *Math. Programming*, 1:168–194, 1971.
- [17] D. R. Fulkerson. Anti-blocking polyhedra. *J. Combinatorial Theory Ser. B*, 12:50–71, 1972.
- [18] A. M. Garsia and C. Procesi. On certain graded S_n -modules and the q -Kostka polynomials. *Advances in Mathematics*, 94(1):82–138, 1992.
- [19] L. Geissinger and D. Kinch. Representations of the hyperoctahedral groups. *J. Algebra*, 53(1):1–20, 1978.
- [20] A. V. Geramita. Inverse systems of fat points: Waring’s problem, secant varieties of Veronese varieties and parameter spaces for Gorenstein ideals. In *The Curves Seminar at Queen’s, Vol. X (Kingston, ON, 1995)*, volume 102 of *Queen’s Papers in Pure and Appl. Math.*, pages 2–114. Queen’s Univ., Kingston, ON, 1996.
- [21] P. Gordan. Ueber die Auflösung linearer Gleichungen mit reellen Coefficienten. *Math. Ann.*, 6(1):23–28, 1873.
- [22] S. Griffin. Ordered set partitions, Garsia-Procesi modules, and rank varieties. *Transactions of the American Mathematical Society*, 374(4):2609–2660, 2021.
- [23] F. Gundlach. Polynomials vanishing at lattice points in a convex set, 2021.
- [24] J. Haglund. *The q, t -Catalan numbers and the space of diagonal harmonics*, volume 41 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2008. With an appendix on the combinatorics of Macdonald polynomials.
- [25] J. Haglund, B. Rhoades, and M. Shimozono. Ordered set partitions, generalized coinvariant algebras, and the Delta Conjecture. *Advances in Mathematics*, 329:851–915, 2018.
- [26] M. Hochster. Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes. *Ann. of Math. (2)*, 96:318–337, 1972.
- [27] J. Huang and B. Rhoades. Ordered set partitions and the 0-Hecke algebra. *Algebraic Combinatorics*, 1(1):47–80, 2018.
- [28] J. Huang, B. Rhoades, and T. Scrimshaw. Hall-Littlewood polynomials and a Hecke action on ordered set partitions. *Proceedings of the American Mathematical Society*, 147(5):1839–1850, 2019.
- [29] B. Kostant. Lie group representations on polynomial rings. *Amer. J. Math.*, 85:327–404, 1963.
- [30] J. C. Lagarias and G. M. Ziegler. Bounds for lattice polytopes containing a fixed number of interior points in a sublattice. *Canad. J. Math.*, 43(5):1022–1035, 1991.
- [31] M. J. Liu. Viennot shadows and graded module structure in colored permutation groups. *arXiv preprint arXiv:2401.07850*, 2024.
- [32] N. A. Loehr and J. B. Remmel. A computational and combinatorial exposé of plethystic calculus. *J. Algebraic Combin.*, 33(2):163–198, 2011.
- [33] I. G. Macdonald. Polynomials Associated with Finite Cell-Complexes. *Journal of the London Mathematical Society*, 2(1):181–192, 1971.
- [34] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York, second edition, 2015. With contribution by A. V. Zelevinsky and a foreword by Richard Stanley, Reprint of the 2008 paperback edition [MR1354144].
- [35] E. Noether. Der endlichkeitsatz der invarianten endlicher linearer gruppen der charakteristik p . pages 28–35, 1926.
- [36] J. Oh and B. Rhoades. Cyclic sieving and orbit harmonics. *Mathematische Zeitschrift*, 300(1):639–660, 2022.
- [37] M. Reineke, B. Rhoades, and V. Tewari. Zonotopal algebras, orbit harmonics, and Donaldson–Thomas invariants of symmetric quivers. *International Mathematics Research Notices*, 2023(23):20169–20210, 2023.
- [38] B. Rhoades. Increasing subsequences, matrix loci, and viennot shadows. *arXiv preprint arXiv:2306.08718*, 2023.
- [39] B. E. Sagan. *The symmetric group*, volume 203 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001. Representations, combinatorial algorithms, and symmetric functions.
- [40] R. P. Stanley. Combinatorial reciprocity theorems. *Advances in Math.*, 14:194–253, 1974.
- [41] R. P. Stanley. Combinatorial reciprocity theorems. In *Combinatorics: Proceedings of the NATO Advanced Study Institute held at Nijenrode Castle, Breukelen, The Netherlands 8–20 July 1974*, pages 307–318. Springer, 1975.
- [42] R. P. Stanley. Invariants of finite groups and their applications to combinatorics. *Bull. Amer. Math. Soc. (N.S.)*, 1(3):475–511, 1979.
- [43] R. P. Stanley. Decompositions of rational convex polytopes. *Ann. Discrete Math*, 6(6):333–342, 1980.
- [44] R. P. Stanley. Linear Diophantine equations and local cohomology. *Invent. Math.*, 68(2):175–193, 1982.

- [45] R. P. Stanley. *Combinatorics and commutative algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, second edition, 1996.
- [46] R. P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
- [47] R. P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.
- [48] A. Stapledon. Equivariant Ehrhart theory. *Adv. Math.*, 226(4):3622–3654, 2011.
- [49] S. Yuzvinsky. Orthogonal pairings of Euclidean spaces. *Michigan Math. J.*, 28(2):131–145, 1981.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN, 55455
Email address: `reiner@umn.edu`

DEPARTMENT OF MATHEMATICS, UCSD, LA JOLLA, CA, 92093
Email address: `bprhoades@ucsd.edu`