

THE SHARP CONSTANTS IN THE REAL ANISOTROPIC LITTLEWOOD'S 4/3 INEQUALITY

NICOLÁS CARO-MONTOYA AND DIANA SERRANO-RODRÍGUEZ

Dedicated to the memory of Jhazaira Mantilla Pérez

ABSTRACT. The Real Anisotropic LITTLEWOOD'S 4/3 inequality is an extension of a famous result obtained in 1930 by J. E. LITTLEWOOD. It asserts that (see [17, Theorem 5.1]) for $a, b \in (0, \infty)$, the following assertions are equivalent:

- There is a constant $C_{a,b}^{\mathbb{R}} \geq 1$ such that

$$\left(\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_i, e_j)|^a \right)^{b \times \frac{1}{a}} \right)^{\frac{1}{b}} \leq C_{a,b}^{\mathbb{R}} \|A\|$$

for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{R}$.

- The exponents a, b satisfy $(a, b) \in [1, \infty) \times [1, \infty)$ with

$$\frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}.$$

Several authors have obtained optimal estimates of the best constant $C_{a,b}^{\mathbb{R}}$, for diverse pairs of values (a, b) .

In this paper we provide the optimal values of $C_{a,b}^{\mathbb{R}}$ for all admissible pair of values (a, b) . Furthermore, we provide new estimates for $C_{a,b}^{\mathbb{C}}$, which are optimal for several pairs of values (a, b) . As an application, we prove a variant (in the sense of [15]) of KHINCHIN'S inequality for STEINHAUS variables.

CONTENTS

1. Introduction	1
2. The proof of the main result	4
3. Remarks on the complex case	7
4. Variants of Khinchin's inequality for Steinhaus variables	8
References	13

1. INTRODUCTION

The origins of the theory of absolutely summing multilinear operators are linked to the so-called LITTLEWOOD'S 4/3 inequality [11]; for a detailed introduction to the theory of absolutely summing operators see [9]. On the other hand, the multilinear theory of absolutely summing operators has been recently explored in different contexts by several authors (see [3, 4, 5, 7, 13, 16, 17] and the references therein), with unexpected applications in other fields such as Quantum Information Theory (more precisely, quantum XOR games), Computational Learning Theory and results related to the study of DIRICHLET series (more precisely, questions concerning the n -dimensional BOHR radius; see [2, 4, 12, 20]). These applications were achieved, among other techniques, by the study

2020 *Mathematics Subject Classification.* 11Y60, 42B08, 46B09.

Key words and phrases. Bilinear forms; KHINCHIN'S inequality; LITTLEWOOD'S 4/3 inequality, Matrix norms.

of sharp constants for inequalities of the type LITTLEWOOD's 4/3 (and its multilinear version, namely BOHNENBLUST–HILLE's inequality). Driven by this, in the last decade several works have emerged that attempt to study generalizations of LITTLEWOOD's 4/3 inequality to the multilinear context and also to other sequence spaces (see [14, 18] and the references therein).

All along this paper \mathbb{K} stands for either \mathbb{R} or \mathbb{C} , and $c_0 = c_0(\mathbb{K})$, the Banach space of null sequences. By convention, when $a = \infty$, the sum $(\sum_j |x_j|^a)^{1/a}$ represents the supremum of the values $|x_j|$. In addition we define $f(\infty) := \lim_{s \rightarrow \infty} f(s)$ for any function f . We also define $1/0 = \infty$ and $1/\infty = 0$, and for $s \geq 1$ we denote by s^* the conjugate index of s , so that $\frac{1}{s} + \frac{1}{s^*} = 1$. Finally, we denote the canonical sequence of vectors of c_0 by $(e_k)_{k=1}^\infty$.

LITTLEWOOD's 4/3 inequality [11] asserts that there is a constant $L_{\mathbb{K}} \geq 1$ such that

$$\left(\sum_{i,j=1}^{\infty} |A(e_i, e_j)|^{\frac{4}{3}} \right)^{\frac{3}{4}} \leq L_{\mathbb{K}} \|A\|$$

for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{K}$, where

$$\|A\| := \sup \left\{ \left| \sum_{i,j=1}^{\infty} A(e_i, e_j) x_i y_j \right| : |x_i|, |y_j| \leq 1 \text{ for all } i, j \right\}$$

It is well known that the exponent 4/3 is optimal, and it is shown in [10] that the value $L_{\mathbb{R}} = \sqrt{2}$ is also optimal. For complex scalars we only know that $L_{\mathbb{C}} \leq 2/\sqrt{\pi}$.

However, the optimality feature of the exponent 4/3 is related to the specific configuration of this inequality. A key issue in the theory has been to investigate optimality ranges for summing inequalities involving anisotropic exponents. More specifically, the objective of study is to control

$$(1.1) \quad \left(\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_i, e_j)|^a \right)^{\frac{1}{a} \times b} \right)^{\frac{1}{b}}$$

for all norm-1 continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{K}$. This problem has been addressed through several approaches permeating the theory and the definitive answer was given over the past decade.

The following Anisotropic LITTLEWOOD's 4/3 inequality is proved in [1, Theorems 6.3]:

Theorem 1.1. *Let $a, b \in [1, 2]$. The following assertions are equivalent:*

- *There is a constant $C_{a,b}^{\mathbb{K}} \geq 1$ such that*

$$\left(\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_i, e_j)|^a \right)^{b \times \frac{1}{a}} \right)^{\frac{1}{b}} \leq C_{a,b}^{\mathbb{K}} \|A\|$$

for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{K}$.

- *The values a and b satisfy*

$$(1.2) \quad \frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}.$$

Observe that by taking $a = b = 4/3$ in Theorem 1.1 one recovers LITTLEWOOD's 4/3 inequality.

About the constants, for complex scalars, the determination of the exact values of the optimal constants involved is probably a difficult task. In this case, the only known

optimal estimates are (see [1, Remark 6.4] and [6, page 31]):

$$C_{1,2}^{\mathbb{C}} = C_{2,1}^{\mathbb{C}} = \frac{2}{\sqrt{\pi}};$$

$$C_{2,2}^{\mathbb{C}} = 1,$$

and, in general, if $1 \leq a, b \leq 2$ with $\frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}$, we have

$$1 \leq C_{a,b}^{\mathbb{C}} \leq \left(\frac{4}{\pi}\right)^{\frac{1}{a} + \frac{1}{b} - 1}.$$

For the case of real scalars we have:

Theorem 1.2 ([1, Theorem 6.3]). *Let $1 \leq a, b \leq 2$ with $\frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}$. Then for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{R}$ we have*

$$\left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a\right)^{b/a}\right)^{1/b} \leq 2^{\frac{1}{b} + \frac{1}{a} - 1} \|A\|.$$

Moreover, the constant $2^{\frac{1}{a} + \frac{1}{b} - 1}$ is optimal.

Condition “ $1 \leq a, b \leq 2$ ” in Theorem 1.1 can be dropped. In fact, [17, Theorem 5.1] provides the following definitive result:

Theorem 1.3. *Let $a, b \in (0, \infty)$. The following assertions are equivalent:*

- There is a constant $C_{a,b}^{\mathbb{K}} \geq 1$ such that

$$(1.3) \quad \left(\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_i, e_j)|^a\right)^{b \times \frac{1}{a}}\right)^{\frac{1}{b}} \leq C_{a,b}^{\mathbb{K}} \|A\|$$

for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{K}$.

- The exponents a, b satisfy $(a, b) \in [1, \infty) \times [1, \infty)$ with

$$\frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}.$$

Remark 1.4. We stress that Theorem 1.3 can be stated even in the case that a or b are equal to ∞ . This fact will be evident in the results we are going to demonstrate in the following section.

Remark 1.5. Observe that, by the inclusion of the ℓ_p spaces, since $C_{2,2}^{\mathbb{K}} = 1$, if $a \geq 2$ and $b \geq 2$, we have that the optimal constant $C_{a,b}^{\mathbb{K}}$ is 1 (see Figure 1).

These remarks, and the complete generality of Theorem 1.3, lead us to pose the following questions:

Problem 1 (see Figure 1). What are the optimal values of $C_{a,b}^{\mathbb{R}}$ in (1.3) in the cases $(a, b) \in [1, 2] \times [2, \infty]$ and $(a, b) \in [2, \infty] \times [1, 2]$?

Problem 2. What are the optimal values of $C_{a,b}^{\mathbb{C}}$ in (1.3), when the exponents a, b satisfy $(a, b) \in [1, \infty] \times [1, \infty]$ and $\frac{1}{a} + \frac{1}{b} \leq \frac{3}{2}$?

In this paper we solve Problem 1 (and partially Problem 2), thus providing a complete description of the constants in the Real Anisotropic LITTLEWOOD'S 4/3 inequality:




FIGURE 1. Illustration of Theorem 1.3: the curve passing through the points $(1, 2)$ and $(2, 1)$ is the hyperbola $b = \frac{2a}{3a-2}$. Region **0** corresponds to non-admissible exponents in the anisotropic LITTLEWOOD's 4/3 inequality. Region **I** correspond to Theorem 1.2, so that the value of the optimal constant is $2^{\frac{1}{a} + \frac{1}{b} - 1}$ in this case. Region **II** correspond to Remark 1.5; in this region the optimal constant has always value 1. Region **III** corresponds to points for which the optimal constant is unknown.

Theorem 1.6. *Let $a, b \in [1, \infty]$ with $\frac{1}{b} + \frac{1}{a} \leq \frac{3}{2}$. Then for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{R}$ we have*

$$\left(\sum_{k=1}^N \left(\sum_{j=1}^N |A(e_k, e_j)|^a \right)^{b/a} \right)^{1/b} \leq 2^{\max\{0, \frac{1}{b} + \frac{1}{a} - 1\}} \|A\|.$$

Moreover, the constant $2^{\max\{0, \frac{1}{b} + \frac{1}{a} - 1\}}$ is optimal.

In order to better understand this result, we divide the region of admissible exponents into the following 4 subregions (see Figure 2):

- (RI) Points (a, b) be such that $1 \leq b, a \leq 2$ and $\frac{1}{b} + \frac{1}{a} \leq \frac{3}{2}$.
- (RII) The infinite region of points $(a, b) \in [1, \infty] \times [a^*, \infty]$.
- (RIII) The infinite region of points $(a, b) \in [2, \infty] \times [1, a^*]$.
- (RIV) The infinite region of points $(a, b) \in [1, 2] \times [2, a^*]$.

This paper is organized as follows: in Section 2 we present the proof of our main result, Theorem 1.6, by using a variant of KHINCHIN's inequality [15] together with techniques of interpolation; this result gives a complete description of the constants in the Real Anisotropic LITTLEWOOD's 4/3 inequality, thus completely solving Problem 1. In Section 3 we extend the ideas of Section 2 to the setting of complex scalars, which allow us to partially solve Problem 2. Finally, as an application of the techniques from Section 3, we prove in Section 4 a variant (in the sense of [15]) of KHINCHIN's inequality for STEINHAUS variables.

2. THE PROOF OF THE MAIN RESULT

We need the following variant of the KHINCHIN inequality of [15]. The proof strongly uses the so-called Contraction Principle (see [9, Theorem 12.2]).




FIGURE 2. Illustration of Theorem 1.6: regions (RI), (RIII) and (RIV) correspond to the value $2^{\frac{1}{a} + \frac{1}{b} - 1}$ for the optimal constant, whereas region (RII) corresponds to points for which the optimal constant is 1.

Proposition 2.1 ([15, Lemma 1 and Proposition 1]). *Let N be a positive integer and $(a_i)_{i=1}^N \in \mathbb{K}^N$ be a sequence of scalars. If $r \in [2, \infty]$, then*

$$(2.1) \quad \left(\sum_{i=1}^N |a_i|^r \right)^{\frac{1}{r}} \leq 2^{\frac{1}{r}} \int_0^1 \left| \sum_{i=1}^N r_i(t) a_i \right| dt$$

and the estimate $2^{1/r}$ is optimal.

Above, as usual, $(r_i : [0, 1] \rightarrow \mathbb{R})_{i=1}^\infty$ is the sequence of independent and identically distributed random variables defined by

$$(2.2) \quad r_i(t) := \text{sign}(\sin(2^i \pi t)),$$

the so-called RADEMACHER functions.

In order to prove our main result, we start with the following extremal particular cases:

Lemma 2.2. *Let $a \in [2, \infty]$. Then for every continuous bilinear form $A : c_0 \times c_0 \rightarrow \mathbb{R}$ we have*

$$\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{1/a} \leq 2^{\frac{1}{a}} \|A\|.$$

Proof. Let N be a positive integer. Observe that, by Proposition 2.1, for all $a \geq 2$ we have

$$\sum_{k=1}^N \left(\sum_{j=1}^N |A(e_k, e_j)|^a \right)^{1/a} \leq 2^{\frac{1}{a}} \sum_{k=1}^N \int_0^1 \left| \sum_{j=1}^N r_j(t) A(e_k, e_j) \right| dt.$$

Moreover,

$$\begin{aligned}
(2.3) \quad 2^{\frac{1}{a}} \sum_{k=1}^N \int_0^1 \left| \sum_{j=1}^N r_j(t) A(e_k, e_j) \right| dt &\leq 2^{\frac{1}{a}} \int_0^1 \sum_{k=1}^N \left| \sum_{j=1}^N r_j(t) A(e_k, e_j) \right| dt \\
&= 2^{\frac{1}{a}} \int_0^1 \sum_{k=1}^N \left| A \left(e_k, \sum_{j=1}^N r_j(t) e_j \right) \right| dt \\
&\leq 2^{\frac{1}{a}} \int_0^1 \left\| A \left(\cdot, \sum_{j=1}^N r_j(t) e_j \right) \right\| dt \\
&\leq 2^{\frac{1}{a}} \|A\|. \quad \square
\end{aligned}$$

Lemma 2.3. *Let $a \in [2, \infty]$. Then for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{R}$ we have*

$$\left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{a^*/a} \right)^{1/a^*} \leq \|A\|.$$

Proof. Observe that the extreme case $a = \infty$ is given by Lemma 2.2. On the other hand, it is well known that for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{R}$

$$\left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^2 \right) \right)^{1/2} \leq \|A\|.$$

Let

$$\theta_0 := \frac{a-2}{a}.$$

Note that $\theta_0 \in (0, 1)$; since

$$\begin{aligned}
\frac{1}{a} &= \frac{\theta_0}{\infty} + \frac{1-\theta_0}{2}; \\
\frac{1}{a^*} &= \frac{\theta_0}{1} + \frac{1-\theta_0}{2},
\end{aligned}$$

a simple interpolation argument yields

$$\begin{aligned}
\left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{a^*/a} \right)^{1/a^*} &\leq \left(\sum_{k=1}^{\infty} \sup_j |A(e_k, e_j)| \right)^{\frac{a-2}{a}} \left(\left[\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^2 \right) \right]^{\frac{1}{2}} \right)^{\frac{2}{a}} \\
&\leq \|A\|. \quad \square
\end{aligned}$$

Now we are ready to prove the main result of this work.

Proof of Theorem 1.6. Let $b \in [1, \infty]$ and $a \in [1, \infty]$, with $\frac{1}{b} + \frac{1}{a} \leq \frac{3}{2}$. If $1 \leq b, a \leq 2$, the result follows from Theorem 1.2 (these points correspond to region (RI)).

Suppose at first that $1 \leq \frac{1}{b} + \frac{1}{a} \leq \frac{3}{2}$ (i.e., $\frac{1}{a^*} \leq \frac{1}{b} \leq \frac{3}{2} - \frac{1}{a}$). Thus, if $a \in [2, \infty]$, then we have $b \in [1, a^*]$ (these points correspond to region (RIII)). If

$$\theta_1 := 1 - \frac{a}{b^*},$$

then we have $\theta_1 \in (0, 1)$; moreover, since

$$\frac{1}{b} = \frac{\theta_1}{1} + \frac{1-\theta_1}{a^*},$$

an interpolation argument together with Lemmas 2.2 and 2.3 yields

$$(2.4) \quad \left\{ \begin{array}{l} \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{b/a} \right)^{1/b} \\ \leq \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{1/a} \right)^{1-\frac{a}{b^*}} \times \left(\left[\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{a^*/a} \right]^{1/a^*} \right)^{\frac{a}{b^*}} \\ \leq \left(2^{\frac{1}{a}} \|A\| \right)^{1-\frac{a}{b^*}} \cdot (\|A\|)^{\frac{a}{b^*}} \\ = \left(2^{\frac{1}{a}} \right)^{1-\frac{a}{b^*}} \|A\| \\ = 2^{\frac{1}{b} + \frac{1}{a} - 1} \|A\|. \end{array} \right.$$

Now, if $a \in [1, 2]$ (recall that $\frac{1}{a^*} \leq \frac{1}{b} \leq \frac{3}{2} - \frac{1}{a}$), then we have $b \in [2, a^*]$ (these points correspond to region (RIV)). Thus, if $b \neq \infty$, then from MINKOWSKI'S inequality and (2.4) we get

$$\begin{aligned} \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{b/a} \right)^{1/b} &\leq \left(\sum_{j=1}^{\infty} \left(\sum_{k=1}^{\infty} |A(e_k, e_j)|^b \right)^{a/b} \right)^{1/a} \\ &\leq 2^{\frac{1}{a} + \frac{1}{b} - 1} \|A\|. \end{aligned}$$

The extreme case $b = \infty$ and $a = 1$ follows trivially from

$$\sup_k \sum_{j=1}^{\infty} |A(e_k, e_j)| \leq \|A\|.$$

On the other hand, the bilinear form

$$A_0(x, y) = x_1 y_1 + x_1 y_2 + x_2 y_1 - x_2 y_2$$

satisfies $\|A_0\| = 2$ and

$$\left(\sum_{k=1}^2 \left(\sum_{j=1}^2 |A_0(e_k, e_j)|^a \right)^{b/a} \right)^{1/b} = 2^{\frac{1}{b} + \frac{1}{a} - 1} \|A_0\|$$

(see [10]), which shows that the value $2^{\frac{1}{b} + \frac{1}{a} - 1}$ cannot be improved.

Finally, for the case $b \in [1, \infty]$ and $a \in [1, \infty]$, with $\frac{1}{b} + \frac{1}{a} \leq 1$ (i.e., $\frac{1}{b} \leq \frac{1}{a^*}$), if we have $a \in [1, \infty]$ then $b \in [a^*, \infty]$ (these points correspond to region (RII)), and thus

$$\begin{aligned} \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{b/a} \right)^{1/b} &\leq \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{a^*/a} \right)^{1/a^*} \\ &\leq \|A\|. \end{aligned}$$

Thus, in this case the optimal constant is 1. \square

3. REMARKS ON THE COMPLEX CASE

In this section we partially replicate, for the case of complex scalars, the results obtained in the previous section. First, we have the following estimate:

Lemma 3.1. *Let $a \in [2, \infty]$. Then for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{C}$ we have*

$$\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{1/a} \leq \left(\frac{4}{\pi} \right)^{\frac{1}{a}} \|A\|.$$

Proof. Observe that by Proposition 2.1, for $a = \infty$:

$$\sum_{k=1}^{\infty} \sup_j |A(e_k, e_j)| \leq \|A\|.$$

Moreover, it is known that (see [6, page 31])

$$\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^2 \right)^{1/2} \leq \left(\frac{4}{\pi} \right)^{\frac{1}{2}} \|A\|.$$

Interpolating with $\theta = \frac{2}{a}$ we have

$$\begin{aligned} \sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{1/a} &\leq \left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^2 \right)^{1/2} \right)^{\theta} \left(\sum_{k=1}^{\infty} \sup_j |A(e_k, e_j)| \right)^{1-\theta} \\ &\leq \left(\left(\frac{4}{\pi} \right)^{\frac{1}{2}} \|A\| \right)^{\frac{2}{a}} \|A\|^{\frac{a-2}{a}} = \left(\frac{4}{\pi} \right)^{\frac{1}{a}} \|A\|. \quad \square \end{aligned}$$

By mimicking the proof of Lemma 2.3 and Theorem 1.6 we get, respectively:

Lemma 3.2. *Let $a \in [2, \infty]$. Then for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{C}$ we have*

$$\left(\sum_{k=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_k, e_j)|^a \right)^{a^*/a} \right)^{1/a^*} \leq \|A\|.$$

Theorem 3.3. *Let $a, b \in [1, \infty]$ with $\frac{1}{b} + \frac{1}{a} \leq \frac{3}{2}$. There is a constant $C_{a,b}^{\mathbb{C}} \geq 1$ such that*

$$\left(\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |A(e_i, e_j)|^a \right)^{b \times \frac{1}{a}} \right)^{\frac{1}{b}} \leq C_{a,b}^{\mathbb{C}} \|A\|$$

for every continuous bilinear form $A: c_0 \times c_0 \rightarrow \mathbb{C}$. Moreover, if $b \leq a$, then we have

$$1 \leq C_{b,a}^{\mathbb{C}} \leq C_{a,b}^{\mathbb{C}} \leq \left(\frac{4}{\pi} \right)^{\max\{0, \frac{1}{a} + \frac{1}{b} - 1\}}.$$

In particular, for $a, b \in [1, \infty]$ with $\frac{1}{b} + \frac{1}{a} \leq 1$ we have $C_{a,b}^{\mathbb{C}} = 1$.

4. VARIANTS OF KHINCHIN'S INEQUALITY FOR STEINHAUS VARIABLES

In this section, among other results, we obtain a version of Proposition 2.1 for STEINHAUS variables.

Observe that the optimal constant in Proposition 2.1 coincides with the optimal constants $C_{1,r}^{\mathbb{R}}$ and $C_{r,1}^{\mathbb{R}}$ for all $r \in [2, \infty]$.

KHINCHIN'S inequality for STEINHAUS variables plays a crucial role in the improvement of estimates for the constants in inequalities of the type LITTLEWOOD'S 4/3 (and their extended multilinear versions) for complex scalars (see [1, 8]). For a very recent approach to KHINCHIN'S inequality we refer to [19].




FIGURE 3. Illustration of Theorem 3.3: region II corresponds to points for which the optimal constant is 1. Region I corresponds to points for which the optimal constant (in the complex case) is unknown, except for $(a, b) = (1, 2)$ and $(2, 1)$, where we have $C_{a,b}^{\mathbb{C}} = \frac{2}{\sqrt{\pi}}$.

In the previous section we do not provide a version of Proposition 2.1 for STEINHAUS variables because its proof strongly uses the Contraction Principle. Such a principle is stated only for randomized sums (independent symmetric real-valued random variables), and we do not know if a similar principle holds for independent symmetric complex-valued random variables.

In contrast, as an application of the methods from the previous section, we obtain a version of Proposition 2.1 for STEINHAUS variables with optimal constant coinciding with the optimal constants $C_{1,r}^{\mathbb{C}}$ and $C_{r,1}^{\mathbb{C}}$, for all $r \in [2, \infty]$, as in the case of RADEMACHER functions. Such result is achieved by using ideas from [8], where a type of KHINCHINE'S inequality previously developed by R. BLEI is studied.

It is well known that, by considering the dyadic expansion of $2^N t$ for each $t \in [0, 1]$, one can prove the equality

$$\int_{[0,1]} \left| \sum_{i=1}^N r_i(t) a_i \right| dt = \frac{1}{2^N} \sum_{\eta \in \{1,-1\}^N} \left| \sum_{j=1}^N \eta_j a_j \right|,$$

where the functions r_i are given by (2.2). Thus, inequality (2.1) from Proposition 2.1 can be rewritten as

$$(4.1) \quad \left(\sum_{j=1}^N |a_j|^r \right)^{\frac{1}{r}} \leq 2^{\frac{1}{r}} \left(\frac{1}{2^N} \sum_{\eta \in \{1,-1\}^N} \left| \sum_{j=1}^N \eta_j a_j \right| \right)$$

for all $r \in [2, \infty]$, and for all sequences of scalars $(a_j)_{j=1}^N \in \mathbb{K}^N$ and all positive integers N .

The counterpart for the average

$$\frac{1}{2^N} \sum_{\eta \in \{1,-1\}^N} \left| \sum_{j=1}^N \eta_j a_j \right|$$

in the complex framework is

$$(4.2) \quad \left(\frac{1}{2\pi}\right)^N \int_0^{2\pi} \cdots \int_0^{2\pi} \left| \sum_{j=1}^N a_j e^{it_j} \right| dt_1 \cdots dt_N.$$

For the sake of simplicity we shall denote (4.2) by

$$(4.3) \quad \mathbb{E} \left| \sum_{j=1}^N a_j \varepsilon_j \right|,$$

where ε_j are STEINHAUS variables, that is, random variables uniformly distributed on the complex unit circle. Our goal is to prove the following version of Proposition 2.1 for STEINHAUS variables:

Theorem 4.1. *Let $r \in [2, \infty]$. There is a (optimal) constant $C_r \geq 1$ such that*

$$(4.4) \quad \left(\sum_{n=1}^N |a_n|^r \right)^{\frac{1}{r}} \leq C_r \left(\mathbb{E} \left| \sum_{n=1}^N a_n \varepsilon_n \right| \right)$$

for every positive integer N and for all sequences of scalars $(a_j)_{j=1}^N \in \mathbb{K}^N$, where $\varepsilon_1, \dots, \varepsilon_n$ are STEINHAUS variables. Moreover,

$$C_r = C_{1,r}^{\mathbb{C}} = C_{r,1}^{\mathbb{C}},$$

for all $r \in [2, \infty]$.

In what follows we state a KHINCHINE-type inequality that extends and unifies inequalities (1.1) and (4.4). First, we need to introduce some notation and results.

For each integer $M \geq 2$, we define

$$\begin{aligned} T_M &:= \left\{ \exp\left(\frac{2j\pi}{M}i\right) : j = 0, \dots, M-1 \right\}; \\ T_\infty &:= \left\{ \exp(ti) : t \in [0, 2\pi) \right\}, \end{aligned}$$

and

$$\begin{aligned} D_M &:= \text{conv}(T_M); \\ D_\infty &:= \text{conv}(T_\infty), \end{aligned}$$

where conv denotes the convex hull. Observe that D_∞ is the closed unit disk, and that $D_M \subseteq D_\infty$. Obviously, D_M is a convex and closed absorbing set in \mathbb{C} .

Let N and M be positive integers, with $M \geq 3$. For any bilinear form $A: \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{C}$ we define the norm

$$\|A\|_M := \sup\{|A(x, y)| : y \in T_M^N; |x_i| \leq 1 \text{ for } i = 1, \dots, N\}.$$

The following basic result provides estimates for the sup-norm $\|A\|$:

Theorem 4.2 ([8, Theorem 2.3]). *Let N, M be positive integers, with $M \geq 3$. Then*

$$\|A\|_M \leq \|A\| \leq R_M^{-1} \|A\|_M$$

for all bilinear forms $A: \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{C}$, where

$$R_M := \left[\frac{1}{2} + \frac{1}{2} \cos\left(\frac{2\pi}{M}\right) \right]^{\frac{1}{2}}.$$

For each $M \geq 2$, let

$$\Omega_M := \left\{ \frac{2j\pi}{M} : j = 0, \dots, M-1 \right\}.$$

Let $(a_n)_{n=1}^N \in \mathbb{K}^N$ be a sequence of scalars, and let $M \geq 2$. We define

$$E_M((a_n)_{n=1}^N) := \left(\frac{1}{M} \right)^N \sum_{\beta \in \Omega_M^N} \left| \sum_{n=1}^N a_n e^{i\beta_n} \right|.$$

Using the Dominated Convergence Theorem it is possible to prove that

$$\lim_{M \rightarrow \infty} E_M((a_n)_{n=1}^N) = \mathbb{E} \left| \sum_{n=1}^N a_n \varepsilon_n \right|.$$

We need the following auxiliary result:

Lemma 4.3 ([8, Lemma 2.4]). *Let $(a_n)_{n=1}^N \in \mathbb{K}^N$ be a sequence of scalars, and let $M \geq 2$. Then*

$$E_M((a_n)_{n=1}^N) = E_M((a_n e^{is_n})_{n=1}^N)$$

for all $s_1, \dots, s_N \in \Omega_M$.

Now we are ready to state and prove the announced extension of KHINCHINE's inequality, for $r \in [2, \infty]$, that extends and unifies inequalities (1.1) and (4.4). We emphasize that this result is already known for $r = 2$; see [6, Chapter II: Section 6].

Theorem 4.4 (BLEI–KHINCHINE inequality). *Let $r \in [2, \infty]$. For every $M \geq 2$, there is a (optimal) constant $\mathcal{B}_{M,r}$ such that*

$$(4.5) \quad \left(\sum_{n=1}^N |a_n|^r \right)^{1/r} \leq \mathcal{B}_{M,r} \cdot E_M((a_n)_{n=1}^N),$$

for every positive integer N and all sequences of scalars $(a_j)_{j=1}^N \in \mathbb{K}^N$. Moreover, for all $M \geq 3$ we have

$$(4.6) \quad \mathcal{B}_{M,r} \leq C_{1,r}^{\mathbb{C}} \cdot R_M^{-1},$$

where

$$R_M := \left[\frac{1}{2} + \frac{1}{2} \cos \left(\frac{2\pi}{M} \right) \right]^{\frac{1}{2}}.$$

Proof. The case $M = 2$ is precisely the inequality (4.1), so we only need to prove the case $M \geq 3$. Let $(a_n)_{n=1}^N \in \mathbb{K}^N$ be a sequence of scalars, such that $E_M((a_n)_{n=1}^N) = 1$. Then, by Lemma 4.3,

$$E_M((a_n e^{is_n})_{n=1}^N) = \left(\frac{1}{M} \right)^N \sum_{\beta \in \Omega_M^N} \left| \sum_{n=1}^N a_n e^{is_n} e^{i\beta_n} \right| = 1$$

for all $s_1, \dots, s_N \in \Omega_M$. Thus, for all $w_1, \dots, w_N \in T_M$ we have

$$\left(\frac{1}{M} \right)^N \sum_{t \in T_M^N} \left| \sum_{n=1}^N a_n w_n t_n \right| = 1.$$

Fix an enumeration of the set T_M^N , say $\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(M^N)}$, where $\boldsymbol{\tau}^{(i)} = (\tau_1^{(i)}, \dots, \tau_N^{(i)})$ for each i . In this way, the equality above can be rewritten as

$$(4.7) \quad \left(\frac{1}{M} \right)^N \sum_{i=1}^{M^N} \left| \sum_{n=1}^N a_n \tau_n^{(i)} w_n \right| = 1,$$

for all $w_1, \dots, w_N \in T_M$.

Consider the bilinear form $A: \mathbb{C}^{M^N} \times \mathbb{C}^N \rightarrow \mathbb{C}$ given by

$$A(e_i, e_n) = \frac{a_n \tau_n^{(i)}}{M^N} \quad (i \in \{1, \dots, M^N\}; n \in \{1, \dots, N\}).$$

We claim that $\|A\|_M \leq 1$. In fact, by using HÖLDER'S inequality together with equality (4.7) we get

$$\begin{aligned} \left| \sum_{i=1}^{M^N} \sum_{n=1}^N A(e_i, e_n) w_n z_i \right| &= \left| \sum_{i=1}^{M^N} \sum_{n=1}^N \frac{a_n \tau_n^{(i)}}{M^N} w_n z_i \right| \\ &\leq \left(\sum_{i=1}^{M^N} \left| \sum_{n=1}^N \frac{a_n \tau_n^{(i)}}{M^N} w_n \right| \right) \cdot \sup_{1 \leq i \leq M^N} |z_i| \\ &\leq \left(\frac{1}{M} \right)^N \sum_{i=1}^{M^N} \left| \sum_{n=1}^N a_n \tau_n^{(i)} w_n \right| \\ &= 1 \end{aligned}$$

for all $w_1, \dots, w_N \in T_M$ and all $z_1, \dots, z_{M^N} \in \mathbb{C}$ with $\sup_{1 \leq i \leq M^N} |z_i| \leq 1$. Therefore $\|A\|_M \leq 1$. Moreover, if $r \in [2, \infty]$, then by using Theorems 1.3 and 4.2 and the above norm estimate we obtain

$$\begin{aligned} \left(\sum_{n=1}^N |a_n|^r \right)^{1/r} &= \left[\sum_{n=1}^N \left(\sum_{i=1}^{M^N} \left| \left(\frac{1}{M} \right)^N a_n \tau_n^{(i)} \right|^r \right)^{\frac{1}{r}} \right] \\ &= \left[\sum_{n=1}^N \left(\sum_{i=1}^{M^N} |A(e_i, e_n)|^r \right)^{\frac{1}{r}} \right] \\ &\leq C_{1,r}^{\mathbb{C}} \|A\| \\ &\leq C_{1,r}^{\mathbb{C}} \cdot R_M^{-1} \|A\|_M \\ &\leq C_{1,r}^{\mathbb{C}} \cdot R_M^{-1}. \end{aligned}$$

Thus, the inequality follows and

$$\mathcal{B}_{M,r} \leq C_{1,r}^{\mathbb{C}} \cdot R_M^{-1},$$

for all $r \in [2, \infty]$, and $M \geq 3$, as claimed. \square

Now we can easily prove the main result of this section:

Proof of Theorem 4.1. Taking $M \rightarrow \infty$ in Theorem 4.4, and using that $R_M \rightarrow 1$, we obtain the inequality (4.4), with the estimate

$$C_r \leq C_{1,r}^{\mathbb{C}}$$

for all $r \in [2, \infty]$.

On the other hand, since $r > 1$, Theorem 3.3 implies

$$C_{1,r}^{\mathbb{C}} \leq C_{r,1}^{\mathbb{C}}$$

for all $r \in [2, \infty]$.

Finally, we can repeat the proof of Lemma 2.2 in the case of complex scalars, by just replacing (2.1) from Proposition 2.1 with (4.4) from Theorem 4.1. This yields

$$C_{r,1}^{\mathbb{C}} \leq C_r$$

for all $r \in [2, \infty]$. Putting together these inequalities we obtain the announced optimal estimates:

$$C_r = C_{1,r}^{\mathbb{C}} = C_{r,1}^{\mathbb{C}},$$

for all $r \in [2, \infty]$. □

REFERENCES

- [1] N. Albuquerque, F. Bayart, D. Pellegrino and J. Seoane-Sepúlveda, Sharp generalizations of the Bohnenblust–Hille inequality, *J. Funct. Analysis*, **266** (2014), 3726–3740. [2](#), [3](#), [8](#)
- [2] S. Arunachalam, A. Dutt, F. Escudero Gutiérrez and C. Palazuelos, Learning low-degree quantum objects. arXiv preprint arXiv:2405.10933, (2024). [1](#)
- [3] F. Bayart, Multiple summing maps: coordinatewise summability, inclusion theorems and p-Sidon sets. *J. Funct. Anal.* **274** (2018), no. 4, 1129–1154. [1](#)
- [4] F. Bayart, D. Pellegrino and J. B. Seoane-Sepúlveda, The Bohr radius of the n -dimensional polydisk is equivalent to $\sqrt{(\log n)/n}$, *Adv. Math.* **264** (2014), 726–746. [1](#)
- [5] A. Belacel, A. Bougoutaia, R. Macedo and P. Rueda, Intermediate classes of nuclear multilinear operators, *Bull. Braz. Math. Soc. (N.S.)* **54** (2023), <https://doi.org/10.1007/s00574-023-00365-5> [1](#)
- [6] R. Blei, *Analysis in Integer and Fractional Dimensions*. Cambridge Studies in Advanced Mathematics, 71. Cambridge University Press, Cambridge, 2001. xx+556 pp. [3](#), [8](#), [11](#)
- [7] G. Botelho, D. Freitas, Summing multilinear operators by blocks: The isotropic and anisotropic cases. *J. Math. Anal. Appl.* **490** (2020), no. 1, 124203, 21 pp. [1](#)
- [8] W.V. Cavalcante, D. Núñez-Alarcón, D.M. Pellegrino and P. Rueda, Optimal constants of the mixed Littlewood inequalities: the complex case. *Math. Inequal. Appl.* **25**, no. 1 (2022), 205–220. [8](#), [9](#), [10](#), [11](#)
- [9] J. Diestel, H. Jarchow, A. Tonge, *Absolutely summing operators*. Cambridge Studies in Advanced Mathematics, 43. Cambridge University Press, Cambridge, 1995. xvi+474 pp. [1](#), [4](#)
- [10] D. Diniz, G. A. Muñoz-Fernández, D. Pellegrino and J. B. Seoane-Sepúlveda, Lower Bounds for the constants in the Bohnenblust–Hille inequality: the case of real scalars, *Proc. Amer. Math. Soc.* **142** (2014), n. 2, 575–580. [2](#), [7](#)
- [11] J.E. Littlewood, On bounded bilinear forms in an infinite number of variables, *Q. J. Math. Oxford* **1** (1930) 164–174. [1](#), [2](#)
- [12] A. Montanaro, Some applications of hypercontractive inequalities in quantum information theory. *J. Math. Phys.* **53** (2012), no. **12**, 122206, 15 pp. [1](#)
- [13] D. Núñez-Alarcón, J. Santos, D. Serrano-Rodríguez, Unified Grothendieck’s and Kwapien’s Theorems for Multilinear Operators. *Bull Braz Math Soc, New Series* **55**, 3 (2024). [1](#)
- [14] D. Núñez-Alarcón, D. Serrano-Rodríguez and K. Teixeira, Sharp exponents for anisotropic Hardy–Littlewood type of inequalities, *Bull Braz Math Soc, To appear* (2024). [2](#)
- [15] D. Pellegrino, D. Santos and J. Santos, Optimal blow up rate for the constants of Khinchin type inequalities, *Quaest. Math.* **41**, (2018) doi:10.2989/16073606.2017.1380724 [1](#), [4](#), [5](#)
- [16] D. Pellegrino, J. Santos, J. Seoane-Sepúlveda, Some techniques on nonlinear analysis and applications. *Adv. Math.* **229** (2012), no. 2, 1235–1265. [1](#)
- [17] D. Pellegrino, J. Santos, D. Serrano and E. Teixeira, A regularity principle in sequence spaces and applications, *Bull. Sci. math.* **141**, (2017), doi:10.1016/j.bulsci.2017.09.001 [1](#), [3](#)
- [18] A. Raposo Jr, D. Serrano-Rodríguez, Coefficients of multilinear forms on sequence spaces, *Bull Braz Math Soc, New Series* **54**, 43 (2023). [2](#)
- [19] O. Ramaré, An Explicit Croot–Łaba–Sisask Lemma Free of Probabilistic Language, *Bull Braz Math Soc, New Series* **55**, 23 (2024). [8](#)
- [20] J. Slote, A. Volberg, and H. Zhang, Noncommutative Bohnenblust–Hille inequality in the Heisenberg–Weyl and Gell-Mann bases with applications to fast learning. arXiv preprint arXiv:2301.01438, (2023). [1](#)

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE PERNAMBUCO, AVENIDA JORNALISTA ANÍBAL FERNANDES, S/N - CIDADE UNIVERSITÁRIA, RECIFE/PE - BRASIL - 50740-560

Email address, N. Caro-Montoya: jorge.caro@ufpe.br

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, BOGOTÁ, COLOMBIA - 111321

Email address, D. Serrano-Rodríguez: diserranor@unal.edu.co