

REGULARIZATION BY NONLINEAR NOISE FOR PDES: WELL-POSEDNESS AND FINITE TIME EXTINCTION[†]

WEI HONG, SHIHU LI, WEI LIU¹

School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou, 221116, China

ABSTRACT. This work focuses on the regularization by nonlinear noise for a class of partial differential equations that may only have local solutions. In particular, we obtain the global existence, uniqueness and the Feller property for stochastic 3D Navier-Stokes equations, which provide positive answers to a longstanding open problem in this field.

Moreover, we discover a new phenomenon that for a potentially explosive deterministic system, an appropriate intervention of nonlinear noise can not only prevent blow-up but also lead to the finite time extinction of the associated stochastic system. Our main results have broad applications, including stochastic p -Laplace equations with heat sources, stochastic surface growth models and stochastic quasi-geostrophic equations.

Keywords: Stochastic PDEs; Regularization by noise; Well-posedness; 3D Navier-Stokes equations; Feller property; Finite time extinction.

Mathematics Subject Classification (2020): 60H15, 60H50, 35Q30, 35R60

CONTENTS

1. Introduction	2
1.1. Variational framework	3
1.2. Regularization by noise	5
1.3. Finite time extinction	8
1.4. Summary	8
2. Main results	9
2.1. Preliminaries	9
2.2. Well-posedness and Feller property	10
2.3. Finite time extinction	14
3. Examples/Applications	15
3.1. SDEs with superlinear drifts	16
3.2. Stochastic 3D Navier-Stokes equations	17
3.3. Stochastic singular p -Laplace equations with heat sources	20
3.4. Stochastic surface growth model	22
3.5. Stochastic quasi-geostrophic equation	23
3.6. Stochastic fast diffusion equations	27

[†] This work is supported by National Key R&D program of China (No. 2023YFA1010101). The research of W. Hong is also supported by NSFC (No. 12401177) and NSF of Jiangsu Province (No. BK20241048). The research of S. Li is also supported by NSFC (No. 12371147). The research of W. Liu is also supported by NSFC (No. 12171208, 12090011, 12090010) and the PAPD of Jiangsu Higher Education Institutions.

¹Corresponding author: weiliu@jsnu.edu.cn

4. Proof of main results	28
4.1. Energy estimates	29
4.2. Tightness of approximating solutions	32
4.3. Passage to the limit	37
4.4. Proof of existence and uniqueness	47
4.5. Proof of Feller property	48
4.6. Proof of finite time extinction	49
5. Appendix	53
References	53

1. INTRODUCTION

In the field of stochastic partial differential equations (SPDEs), there is great interest in investigating the impact and changes produced by random perturbations on deterministic systems, particularly, how the noise alters the well-posedness, stability and long-time asymptotic behavior of solutions. For example, it is interesting to observe that a deterministic system may lack uniqueness or global existence of solutions. However, when the system is perturbed by appropriate noise, it has a unique global solution. This phenomenon is commonly referred as *regularization by noise* in the literature. For a comprehensive discussion, we refer to e.g. the lecture notes by Flandoli [24].

The phenomenon of the regularization by noise for ordinary differential equation (ODE) has been studied very extensively. Zvonkin [74] first showed the existence of a unique strong solution to the following one-dimensional (1D) stochastic differential equations (SDEs) with only bounded measurable drift

$$(1.1) \quad dX_t = b(X_t)dt + dW_t, \quad X_0 = x,$$

where W_t is a 1D standard Wiener process. Then, this result was generalized by Veretenikov in [72] for the multi-dimensional case, and later it was further extended by Krylov and Röckner in [44] for the case of locally unbounded b under some integrability conditions. The key idea of the proof is to use a Zvonkin-type transformation [74] that allows either to eliminate the non-regular drift or to make it more regular.

On the other hand, noise not only promotes the uniqueness of solutions but also prevents blow-up. To illustrate this, we consider the following 1D nonlinear ODE

$$(1.2) \quad dX_t = X_t^2 dt, \quad X_0 = x.$$

If the initial data $x > 0$, the solution of Eq. (1.2) will blow up in finite time. However, by adding a suitable nonlinear stochastic perturbation, we observe that the following SDE is non-explosive

$$(1.3) \quad dX_t = X_t^2 dt + X_t^2 dW_t, \quad X_0 = x,$$

see Section 3.1 below for more details. This type of regularization by noise typically considers superlinear multiplicative noise, see e.g. [3].

In this work, we investigate the regularization by noise for a class of PDEs. More specifically, we aim to explore the effect of noise perturbation on possibly explosive PDEs.

We are particularly interested in what kind of noise perturbation can regularize the system, thereby preventing solutions from blowing up.

1.1. Variational framework. Let $(U, \langle \cdot, \cdot \rangle_U)$ and $(\mathbb{H}, \langle \cdot, \cdot \rangle_{\mathbb{H}})$ be separable Hilbert spaces, and \mathbb{H}^* be the dual space of \mathbb{H} . Let $(\mathbb{V}, \|\cdot\|_{\mathbb{V}})$ denote a reflexive Banach space such that the embedding

$$\mathbb{V} \subset \mathbb{H}$$

is continuous and dense. Identifying \mathbb{H} with its dual space in view of the Riesz isomorphism, we obtain a Gelfand triple

$$(1.4) \quad \mathbb{V} \subset \mathbb{H}(\simeq \mathbb{H}^*) \subset \mathbb{V}^*.$$

The dualization between \mathbb{V} and \mathbb{V}^* is denoted by ${}_{\mathbb{V}^*}\langle \cdot, \cdot \rangle_{\mathbb{V}}$. It is clear that

$${}_{\mathbb{V}^*}\langle \cdot, \cdot \rangle_{\mathbb{V}}|_{\mathbb{H} \times \mathbb{V}} = \langle \cdot, \cdot \rangle_{\mathbb{H}}.$$

Let $L_2(U, \mathbb{H})$ be the space of all Hilbert-Schmidt operators from U to \mathbb{H} . For the measurable maps

$$\mathcal{A} : [0, T] \times \mathbb{V} \rightarrow \mathbb{V}^*, \quad \mathcal{B} : [0, T] \times \mathbb{V} \rightarrow L_2(U, \mathbb{H}),$$

we consider the following SPDE

$$(1.5) \quad dX_t = \mathcal{A}(t, X_t)dt + \mathcal{B}(t, X_t)dW_t, \quad X_0 = x,$$

where $\{W_t\}_{t \in [0, T]}$ is an U -valued cylindrical Wiener process defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

In the deterministic case (i.e. $\mathcal{B} = 0$), the classical result states that Eq. (1.5) has a unique solution if the operator \mathcal{A} satisfies the monotonicity condition and the following coercivity condition (see e.g. [73, Theorem 30.A])

$$(1.6) \quad 2_{{}_{\mathbb{V}^*}\langle \mathcal{A}(t, u), u \rangle_{\mathbb{V}}} + \delta \|u\|_{\mathbb{V}}^\alpha \leq C \|u\|_{\mathbb{H}}^2 + C,$$

where $\delta, C > 0$. The theory of monotone operators originated from the work of Minty [54], and has been systematically studied by Browder [10, 11], Leray and Lions [46], and Hartman and Stampacchia [36].

As for SPDE, the variational approach was initially developed by Pardoux [57], Krylov and Rozovskii [43]. Later, Liu and Röckner [50] extended the classical variational framework to a class of locally monotone operators, which includes many examples that cannot be treated previously, such as stochastic Burgers equations and stochastic 2D Navier-Stokes equations. Very recently, by employing the technique of pseudo-monotone operators and the compactness argument, the variational framework was significantly generalized by Shang, Röckner and Zhang [63] to a more general fully local monotonicity condition. We refer interested readers to [39, 55] and references therein for more results on the variational framework.

It is important to note that the previous works, both in the deterministic and stochastic cases, require that the operator \mathcal{A} satisfies the standard coercivity condition (1.6), which ensures global-in-time bounds of solutions. The first purpose of this paper is to establish the well-posedness of SPDEs that satisfy the following generalized coercivity condition

$$(1.7) \quad 2_{{}_{\mathbb{V}^*}\langle \mathcal{A}(t, u), u \rangle_{\mathbb{V}}} + \delta \|u\|_{\mathbb{V}}^\alpha \leq g(\|u\|_{\mathbb{H}}^2) + C,$$

where $\delta, C > 0$, $g : [0, \infty) \rightarrow [0, \infty)$ is a non-decreasing continuous function. As stated before, the standard condition (1.6) can provide a priori bounds of solutions hence yield the global existence. However, under the generalized coercivity condition (1.7), the solution may blow up in finite time. In a related work [51], Liu and Röckner proved that Eq. (1.5) driven by additive noise (i.e. $\mathcal{B}(t, u) = \mathcal{B}(t)$) possesses a unique local solution under this generalized coercivity condition (1.7).

In this paper, we adopt the variational framework to investigate the regularization effect of noise and ensure the global well-posedness under the condition (1.7), which also differs from the approaches used in the aforementioned works. Specifically, we consider the regularization by nonlinear multiplicative noises that satisfy the following condition

$$(1.8) \quad g(\|u\|_{\mathbb{H}}^2) + \|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})}^2 \leq C(1 + \|u\|_{\mathbb{H}}^2) + \eta \frac{\|\mathcal{B}(t, u)^* u\|_U^2}{(1 + \|u\|_{\mathbb{H}}^2)},$$

where the constants $\eta \in (1, 2)$, $C > 0$ and the function g is the same as in (1.7). Intuitively, if $\mathcal{B} = 0$ (i.e. the deterministic case), the condition (1.8) implies that the function g is of linear growth. Therefore, the main idea of using this condition is to employ the fast enough growth of the nonlinear noise to prevent the potential singularity induced by other nonlinear terms (see Remark 2.3 for more details). Based on this condition, we can get the uniform energy estimates in probability of solutions by choosing an appropriate Lyapunov function.

The first two main results of this work, as presented in Theorems 2.1 and 2.2, establish the global existence, uniqueness and Feller property of solutions for various SPDEs. Notably, all the local or fully local monotone examples mentioned in [63], such as stochastic porous media equations, stochastic 2D Navier-Stokes equations, stochastic Cahn-Hilliard equations, stochastic liquid crystal models and stochastic Allen-Cahn-Navier-Stokes systems, can be encompassed within the more general framework presented in this paper. Additionally, regarding the aforementioned models, our results can be applied to deal with both linearly and superlinearly growing noises.

Moreover, our framework can also be applied to several important new models in geophysical fluid dynamics, such as stochastic p -Laplace equations with heat sources, stochastic 3D Navier-Stokes equations, stochastic quasi-geostrophic equations in both critical and subcritical cases, and stochastic surface growth models, which cannot be encompassed by existing works [1, 50, 55, 63]. It is worth noting that the global well-posedness of 3D Navier-Stokes equations and 1D surface growth models still remains to be open problems.

Now, we outline the main ideas presented in the proof. First, in order to construct a (probabilistically) weak solution of (1.5), we will combine the stochastic compactness argument with the technique from the theory of pseudo-monotone operators. Different from the proof in the existing works (cf. [50, 63]), under the generalized coercivity condition (1.7), we cannot guarantee that the Galerkin approximating solutions $\{X^{(n)}\}$ has the finite second moments. Instead, we establish the energy estimates of $\{X^{(n)}\}$ in probability by constructing a suitable Lyapunov function and utilizing the maximal supermartingale inequality. Then by employing the stopping time technique, we prove the tightness of sequence $\{X^{(n)}\}_{n \in \mathbb{N}}$ in space

$$\mathcal{Z}_T^1 := C([0, T]; \mathbb{V}^*) \cap L^\alpha([0, T]; \mathbb{H}) \cap L_w^\alpha([0, T]; \mathbb{V}),$$

where $L_w^\alpha([0, T]; \mathbb{V})$ denotes the space $L^\alpha([0, T]; \mathbb{V})$ endowed with the weak topology.

It should be pointed out that due to the lack of finite second moments for $\{X^{(n)}\}_{n \in \mathbb{N}}$, this will bring some nontrivial difficulties to the proof and we cannot follow the argument presented in [50, 63] to obtain the weak convergence of the sequence $\{\mathcal{A}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$ as well as the convergence of the sequence $\{\mathcal{B}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$. To this end, we first establish the tightness of $\{\mathcal{A}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$ in space

$$\mathcal{Z}_T^2 := L_w^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*).$$

Note that \mathcal{Z}_T^1 and \mathcal{Z}_T^2 are not Polish spaces. In this case, we apply the Jakubowski's beautiful generalization of the Skorokhod's representation theorem for nonmetric spaces, as presented by Brzeźniak and Ondreját [15], to show the almost sure convergence of both sequences $\{X^{(n)}\}_{n \in \mathbb{N}}$ and $\{\mathcal{A}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$ to certain elements \tilde{X} in \mathcal{Z}_T^1 and $\tilde{\mathcal{A}}(\cdot)$ in \mathcal{Z}_T^2 , respectively, on a new probability space. In this case, it is important to prove that \mathcal{Z}_T^1 and \mathcal{Z}_T^2 are considered as standard Borel spaces with respect to an appropriate topology (cf. Remark 4.1 and Lemma 4.8 for more details). Then by utilizing the truncation method, we can identify the strong convergence of $\mathcal{B}(\cdot, X^{(n)})$ to $\mathcal{B}(\cdot, \tilde{X})$. Additionally, by applying the pseudo-monotone property of $\mathcal{A}(t, \cdot)$, we establish the weak convergence of $\mathcal{A}(\cdot, X^{(n)})$ to $\mathcal{A}(\cdot, \tilde{X})$.

However, it is noted that we only have the almost sure estimates for \tilde{X} , rather than suitable moment estimates, thus the Itô's formula (cf. Theorem 4.2.5 in [52]) for $\|\tilde{X}_t\|_{\mathbb{H}}^2$, $t \in [0, T]$, cannot be directly applied. To solve this problem, we utilize the stopping time technique and the localization procedure. Then, based on the Itô's formula, we are able to show that $\tilde{X} \in C([0, T]; \mathbb{H})$. Hence, \tilde{X} is a probabilistically weak solution. The existence of probabilistically strong solutions follows from the pathwise uniqueness and the Yamada-Watanabe theorem.

1.2. Regularization by noise.

1.2.1. *Semi-linear case.* The phenomenon of regularization by noise has been extensively studied for SDEs, but it is also one of the most interesting research directions in the theory of SPDEs. Since the well-posedness of many deterministic systems are unknown in the literature, there is a considerable interest in exploring whether an appropriate stochastic perturbation can lead to well-posedness for the related stochastic equations. For example, due to the lack of global well-posedness for 3D Navier-Stokes equations, one want to know whether an appropriate random perturbation can ensure that the local strong solutions of 3D Navier-Stokes equations are non-explosive, or that the global weak solutions are unique (pathwise or in law). This topic has been a longstanding open problem in the field of SPDEs, as highlighted by Flandoli in [25, Page 242]:

- (i) *(Uniqueness by noise) Even if the deterministic Navier-Stokes equations do not have uniqueness of weak solutions (or the problem remains open), one would like to discover whether there exists a noise perturbation such that we have uniqueness (pathwise or in law) for the stochastic Navier-Stokes equations.*

- (ii) (*Noise prevents singularities*) *Maybe the deterministic Navier-Stokes equations develop singularities: from a regular initial condition $x \in H^1$, we have a local (unique) regular solution up to some time T^* , but then the regularity may deteriorate and we go back to weak solutions. Is it possible that a suitable random perturbation prevents the emergence of singularities.*

In recent years, the regularization effect of noise has been observed in various infinite-dimensional systems and in different settings. For example, Flandoli, Gubinelli and Priola [23] considered the linear transport equation with a globally Hölder continuous and bounded vector field. They demonstrated that a multiplicative stochastic perturbation of Brownian type is sufficient to render the equation well-posed, which is the first concrete example of a PDE related to fluid dynamics that may lack uniqueness, but is well-posed under a suitable noise perturbation. In [26], Flandoli and Luo showed that a multiplicative noise of transport type has regularization effect on 3D Navier-Stokes equations in vorticity form. They proved that, for a given large initial data, the equation admits a pathwise unique global solution with high probability. In [31], the authors obtained that adding a linear multiplicative noise to the 3D Euler equation leads to a regularization effect in the sense that the solution does not blow up with high probability if the initial data is sufficiently small or if the noise coefficient is sufficiently large.

In addition to linear noise, there have been some studies in literature focused on the regularization effect of nonlinear type noise. In a recent work by Tang and Wang [69], based on the Lyapunov function approach, they developed a general framework for solving singular SPDEs with applications to fluid models driven by pseudo-differential noises. Tang and Yang [70] investigated the stochastic Camassa-Holm equations and showed that the addition of “large noise” can prevent blow-up with probability one, which supports the notion that strong nonlinear noises have a regularization effect in preventing singularities. Bagnara et al. [4] considered a class of SPDEs with drift of superlinear growth, with applications to the stochastic Euler equation as a special case. They showed that a special superlinear noise can effectively prevent blow-up of solutions to these SPDEs. Crisan and Lang [19] recently proposed a framework for studying the regularization effect of superlinear noises. The advantage of [19] is that they can handle a family of compressible inviscid fluid dynamics models, including the inviscid 2D Burgers equation. We refer interested readers to [30, 60, 65] and references therein for the recent development on this topic.

In this work, we aim to establish a general result for the regularization effect using the variational framework, which is applicable to both semi-linear and quasi-linear SPDEs. As one specific application, we demonstrate that by adding a suitable nonlinear noise, the stochastic 3D Navier-Stokes equations have a unique global strong solution for any initial condition $x \in H^1$, which solves the aforementioned open problem (ii) (see Section 3.2 for the details).

It is worth noting that while the global existence and uniqueness of solutions for stochastic 3D Navier-Stokes equations can also be derived from the works of [19, 69], there are certain limitations in terms of the initial data and the regularity of the solutions. In [19], the initial data is required to belong to H^2 and the path of solutions is in the space $C([0, T], L^2)$. Similarly, in [69] the initial data is restricted to H^s for $s > \frac{9}{2}$. Therefore,

the results presented in this work for stochastic 3D Navier-Stokes equations are novel and cannot be encompassed by the aforementioned works. The main improvements lie in the following aspects. Firstly, we establish the existence of a unique global solution with H^1 -initial data, addressing the aforementioned problem (ii). Additionally, the path of solutions belongs to the space $C([0, T], H^1)$, which is consistent with the space where local solutions reside in the deterministic case. Moreover, in comparison to [19], the intensity of nonlinear noise is also different. In [19], the authors take $\mathcal{B}(t, u) = \theta \|u\|_{H^2}^m u$ for some $m > 0$ and $\theta > 0$, which depends on the structure of the operator \mathcal{A} . In our case, as a typical form of noise, we can take $\mathcal{B}(t, u) = \|u\|_{H^1}^m u$ (cf. Remark 2.7).

Furthermore, we establish the continuous dependence on the initial data (in space H^1 for stochastic 3D Navier-Stokes equations as a special case) in probability. In particular, this result implies that the corresponding Markov transition semigroup is Feller in $C_b(H^1)$. Notably, the Feller property of the stochastic 3D Navier-Stokes equations has not been obtained in any previous works. In the deterministic setting, even a selection of solutions depending continuously on the initial condition cannot be obtained. In the probabilistic counterpart, the Markov property of stochastic 3D Navier-Stokes equations is left open, and the existence of a Markov selection is of interest. The existence of Markov solutions for the stochastic 3D Navier-Stokes equations was established by Da Prato and Debussche [20] and Flandoli and Romito [27]. Due to the lack of continuity of solutions in the space of finite energy, the Markov property holds almost everywhere in time. These works also proved the Feller property of the selected Markov semigroup. Different from [20, 27], here we provide the first result concerning the Feller property of the corresponding Markov semigroup for the stochastic 3D Navier-Stokes equations, rather than regarding a Markov selection, which also solves an open problem in this field.

1.2.2. Quasi-linear case. As mentioned previously, one advantage of employing the variational framework is that it can handle a class of quasi-linear SPDEs, including stochastic porous media equations, stochastic fast diffusion equations and stochastic p -Laplace equations. To the best of our knowledge, there are no results available regarding the regularization effect of noise for the blow-up problem in quasi-linear systems. In particular, the existing frameworks such as [4, 19, 60, 69] for the regularization by noise are not applicable in this context.

For example, we consider the following 1D stochastic p -Laplace equation with nonlinear sources

$$(1.9) \quad dX_t = \operatorname{div}(|\nabla X_t|^{p-2} \nabla X_t) dt + \lambda X_t^2 dt + \mathcal{B}(t, X_t) dW_t,$$

where $\lambda = \pm 1$, Eq. (1.9) is degenerate if $p > 2$ or singular if $1 < p < 2$. Nonlinear parabolic equation like (1.9) appears in various applications. For instance, in combustion theory, the function X_t represents the temperature, the term $\operatorname{div}(|\nabla X_t|^{p-2} \nabla X_t)$ represents the thermal diffusion, and the nonlinear source X_t^2 is physically called the ‘‘hot source’’, while the source $-X_t^2$ is known as the ‘‘cool source’’. These different sources have completely different influences on the properties of solutions (cf. [21, 71]).

For the deterministic system (1.9) (i.e., $\mathcal{B}(t, u) = 0$), in the singular case where $1 < p < 2$ and $\lambda = 1$, the solution of the equation may blow up under certain initial condition, and therefore there is no global solution in general (see e.g. [47]). However, in this work, we

demonstrate that the addition of a suitable nonlinear noise can effectively prevent the blow-up of solutions. Consequently, the corresponding stochastic equation has a unique global solution for any initial condition $x \in L^2$. Remarkably, this seems to be the first concrete example of a quasi-linear PDE that becomes well-posed under the influence of a suitable nonlinear noise, see Section 3.3 for the details.

1.3. Finite time extinction. After we demonstrated the regularization effect of nonlinear noise on the global well-posedness of stochastic equations, one natural question is the influence of noise on the long-time behavior of stochastic systems, here we mainly focus on whether nonlinear noise can prevent or promote finite time extinction of solutions.

The problem of the finite time extinction is generally divided into the following three cases

- (i) $\mathbb{P}(X_t \text{ extinct in finite time}) > 0$ for small initial values;
- (ii) $\mathbb{P}(X_t \text{ extinct in finite time}) > 0$ for all initial values;
- (iii) $\mathbb{P}(X_t \text{ extinct in finite time}) = 1$ for all initial values.

There have been many studies in the literature on the finite time extinction of stochastic systems. For example, Barbu, Da Prato and Röckner [5] investigated the finite time extinction with positive probability for 1D self-organized criticality stochastic models driven by linear multiplicative noise. In a subsequent work [7], Barbu and Röckner established asymptotic extinction results with probability one for stochastic porous media equations in dimension $d \in \{1, 2, 3\}$. This result was further extended by Röckner and Wang [64] to more general cases. Specifically, they proved the finite time extinction with probability one for the Zhang model, and with positive probability for the Bak-Tang-Wiesenfeld (BTW) model. Notably, the finite-time extinction of the BTW model with probability one remained an open problem until it was resolved by Gess [29] for all dimensions $d \geq 1$. For more results of this topic, we refer interested readers to [6, 8, 37] and references therein.

It is important to note that all existing results on finite time extinction focused on systems perturbed by linear multiplicative noise. However, in this work, we make an advancement by proving that finite time extinction holds with probability one and with any initial value $x \in \mathbb{H}$ (i.e. Case (iii)) for a large class of quasi-linear SPDEs perturbed by nonlinear multiplicative noises, see Section 2.3 for more details.

As applications, our main result on finite time extinction is applicable to stochastic fast diffusion equations and stochastic singular p -Laplace equations (with nonlinear sources). More importantly, we discover a new phenomenon, i.e. for a deterministic model that may exhibit the blow-up of solutions, a suitable nonlinear noise intervention leads to the finite time extinction of the corresponding stochastic system. This result may shed some new insight on the impact of nonlinear noise on the long-time behavior of stochastic systems.

1.4. Summary. The main contributions of the present work can be summarized as follows.

- (i) The first main contribution lies in demonstrating that the addition of a suitable nonlinear noise ensures the global existence and uniqueness of solutions to the stochastic 3D Navier-Stokes equations for any initial data in H^1 , which provides a positive

answer to an open problem in the field of SPDEs. Furthermore, we establish the continuous dependence on the initial data in probability, as well as the Feller property of the corresponding Markov transition semigroup in $C_b(H^1)$ for the stochastic 3D Navier-Stokes equations. Note that the Feller property of the stochastic 3D Navier-Stokes equations is novel in the literature, which has not been obtained in any previous works.

- (ii) The second one is that we investigate the influence of nonlinear noise on the well-posedness and finite time extinction of quasi-linear systems that cannot be treated by the existing works [4, 19, 60, 69]. Our results also reveal an interesting phenomenon that for a deterministic model that may blow up, there exists an appropriate nonlinear noise intervention which can lead to the finite time extinction for the corresponding stochastic system.
- (iii) The third one is that we establish a general setting to study the global well-posedness of SPDEs under the conditions of fully local monotonicity and generalized coercivity. Furthermore, we prove the finite time extinction of solutions with probability one and any initial value $x \in \mathbb{H}$, instead of with positive probability and small initial value as demonstrated in [5, 6, 8].

Additionally, in comparison to [50, 63], our results can deal with superlinear type noise instead of merely linear growth. Furthermore, our framework can also be applied to some important new models, including but not limited to stochastic 3D Navier-Stokes equations, stochastic p -Laplace equations with heat sources, stochastic surface growth models and stochastic quasi-geostrophic equations.

The rest of paper is organized as follows. In Sect. 2, we introduce the main results about the existence, uniqueness, Feller property as well as finite time extinction in Theorems 2.1-2.3, respectively. Then in Sec. 3, we apply our general framework to concrete examples to illustrate the wide applicability of the main results. In Sect. 4, we give the proofs of Theorems 2.1-2.3. We also recall some useful lemmas in Appendix. Throughout this paper, C_p denotes some positive constant which may change from line to line, where the subscript p is used to emphasize that the constant depends on certain parameter p .

2. MAIN RESULTS

2.1. Preliminaries. We recall some definitions and necessary notations that are frequently used in the paper.

For any Banach space $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$, we denote by $\mathbb{C}_T(\mathbb{B}) := C([0, T]; \mathbb{B})$ the space of all continuous functions from $[0, T]$ to \mathbb{B} , which is a Banach space equipped with the uniform norm given by

$$\|u\|_{\mathbb{C}_T(\mathbb{B})} := \sup_{t \in [0, T]} \|u_t\|_{\mathbb{B}}, \quad u \in \mathbb{C}_T(\mathbb{B}).$$

Let $\mathcal{B}_b(\mathbb{B})$ (resp. $C_b(\mathbb{B})$) be the space of all bounded and Borel measurable (resp. continuous) functions on \mathbb{B} .

In this paper, we will employ the theory of pseudo-monotone operators. To this end, we first recall the definition of the pseudo-monotone operator. For abbreviation, we use the notation “ \rightharpoonup ” for weak convergence in a Banach space.

Definition 2.1. An operator \mathcal{A} from \mathbb{V} to \mathbb{V}^* is called to be a pseudo-monotone operator if $u_n \rightharpoonup u$ in \mathbb{V} and

$$\liminf_{n \rightarrow \infty} \mathbb{V}^* \langle \mathcal{A}(u_n), u_n - u \rangle_{\mathbb{V}} \geq 0,$$

then for any $v \in \mathbb{V}$,

$$\limsup_{n \rightarrow \infty} \mathbb{V}^* \langle \mathcal{A}(u_n), u_n - v \rangle_{\mathbb{V}} \leq \mathbb{V}^* \langle \mathcal{A}(u), u - v \rangle_{\mathbb{V}}.$$

Remark 2.1. Note that Browder [12] introduced a different definition of pseudo-monotone operator: An operator \mathcal{A} from \mathbb{V} to \mathbb{V}^* is called pseudo-monotone if $u_n \rightharpoonup u$ in \mathbb{V} and

$$\liminf_{n \rightarrow \infty} \mathbb{V}^* \langle \mathcal{A}(u_n), u_n - u \rangle_{\mathbb{V}} \geq 0$$

implies $\mathcal{A}(u_n) \rightharpoonup \mathcal{A}(u)$ in \mathbb{V}^* and

$$\lim_{n \rightarrow \infty} \mathbb{V}^* \langle \mathcal{A}(u_n), u_n \rangle_{\mathbb{V}} = \mathbb{V}^* \langle \mathcal{A}(u), u \rangle_{\mathbb{V}}.$$

This definition turns out to be equivalent to Definition 2.3 (cf. [52, Remark 5.2.12]).

2.2. Well-posedness and Feller property. In this part, we consider the well-posedness and Feller property of SPDE (1.5). To this end, we first recall the (probabilistically) weak and strong solutions to SPDE (1.5) as follows.

Definition 2.2. (Weak solution) A pair (X, W) is called a (probabilistically) weak solution to SPDE (1.5), if there exists a stochastic basis $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ such that X is an $\{\mathcal{F}_t\}$ -adapted process and W is an U -valued cylindrical Wiener process on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ and the following holds:

(i) $X \in \mathbb{C}_T(\mathbb{H})$, \mathbb{P} -a.s.;

(ii) $\int_0^T \|\mathcal{A}(s, X_s)\|_{\mathbb{V}^*} ds + \int_0^T \|\mathcal{B}(s, X_s)\|_{L_2(U; \mathbb{H})}^2 ds < \infty$, \mathbb{P} -a.s.;

(iii) $X_t = X_0 + \int_0^t \mathcal{A}(s, X_s) ds + \int_0^t \mathcal{B}(s, X_s) dW_s$, $t \in [0, T]$, \mathbb{P} -a.s. holds in \mathbb{V}^* .

Definition 2.3. (Strong solution) We say that there exists a (probabilistically) strong solution to (1.5) if for every probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ with an U -valued cylindrical Wiener process W , there exists an $\{\mathcal{F}_t\}$ -adapted process X such that properties (i)-(iii) in Definition 2.2 hold.

In this part, we suppose that there are some constants $\alpha > 1$, $\beta \geq 2$ and $C, \delta > 0$ such that the following conditions hold for a.e. $t \in [0, T]$.

(A₁) (Hemicontinuity) For any $u, v, w \in \mathbb{V}$, the maps

$$\mathbb{R}_+ \ni \lambda \mapsto \mathbb{V}^* \langle \mathcal{A}(t, u + \lambda v), w \rangle_{\mathbb{V}}$$

are continuous.

(A₂) (Local Monotonicity) For any $u, v \in \mathbb{V}$,

$$\begin{aligned} & 2_{\mathbb{V}^*} \langle \mathcal{A}(t, u) - \mathcal{A}(t, v), u - v \rangle_{\mathbb{V}} + \|\mathcal{B}(t, u) - \mathcal{B}(t, v)\|_{L_2(U; \mathbb{H})}^2 \\ & \leq (C + \rho(u) + \eta(v)) \|u - v\|_{\mathbb{H}}^2, \end{aligned}$$

where $\rho, \eta : \mathbb{V} \rightarrow [0, \infty)$ are measurable functions satisfying

$$(2.1) \quad \rho(u) + \eta(u) \leq C(1 + \|u\|_{\mathbb{V}}^\alpha)(1 + \|u\|_{\mathbb{H}}^\beta), \quad u \in \mathbb{V}.$$

(**A₃**) (Generalized Coercivity) For any $u \in \mathbb{V}$,

$$2_{\mathbb{V}^*} \langle \mathcal{A}(t, u), u \rangle_{\mathbb{V}} + \delta \|u\|_{\mathbb{V}}^{\alpha} \leq g(\|u\|_{\mathbb{H}}^2) + C,$$

where $g : [0, \infty) \rightarrow [0, \infty)$ is a non-decreasing continuous function.

(**A₄**) (Growth) For any $u \in \mathbb{V}$,

$$(2.2) \quad \|\mathcal{A}(t, u)\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} \leq C(1 + \|u\|_{\mathbb{V}}^{\alpha})(1 + \|u\|_{\mathbb{H}}^{\beta}).$$

(**A₅**) There exists a constant $\eta \in (1, 2)$ such that for any $u \in \mathbb{V}$,

$$(2.3) \quad g(\|u\|_{\mathbb{H}}^2) + \|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})}^2 \leq C(1 + \|u\|_{\mathbb{H}}^2) + \eta \frac{\|\mathcal{B}(t, u)^* u\|_U^2}{(1 + \|u\|_{\mathbb{H}}^2)},$$

where the function g is the same as in (**A₃**), and for any $u \in \mathbb{V}$,

$$(2.4) \quad \|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})}^2 \leq C(1 + \|u\|_{\mathbb{H}}^{\beta}).$$

Moreover, for any sequence $\{u_n\}_{n \in \mathbb{N}}$ and u in \mathbb{V} with $\|u_n - u\|_{\mathbb{H}} \rightarrow 0$,

$$(2.5) \quad \|\mathcal{B}(t, u_n) - \mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})} \rightarrow 0.$$

We state the main result concerning the global existence and uniqueness of (probabilistically) strong solutions and the Markov property to SPDEs (1.5).

Theorem 2.1. *Suppose that the embedding $\mathbb{V} \subset \mathbb{H}$ is compact and that (**A₁**)-(**A₅**) hold. For any initial data $x \in \mathbb{H}$, (1.5) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $p \geq 2$,*

$$(2.6) \quad \mathbb{P} \left(\|X\|_{\mathbb{C}_T(\mathbb{H})}^p + \int_0^T \|X_t\|_{\mathbb{V}}^{\alpha} dt < \infty \right) = 1.$$

Furthermore, if $\mathcal{A}(t, \cdot), \mathcal{B}(t, \cdot)$ are independent of $t \in [0, T]$, then the solution $(X_t)_{t \in [0, T]}$ of (1.5) is a time-homogenous Markov process.

Remark 2.2. (i) *There exist a large class of PDEs for which the global well-posedness is not available. Several examples of such equations arise from geophysical fluid dynamics, such as the 3D Navier-Stokes equations and surface growth models. Motivated by this observation, our current work focuses on a general coercivity condition (**A₃**), which was first introduced by the last named author and Röckner in [51, 52], where they proved that Eq. (1.5) has a unique local solution under this generalized coercivity condition in the case of the additive noise (i.e. $\mathcal{B}(t, u) = \mathcal{B}(t)$).*

This extension allows our main results to be applicable to various new stochastic models, which cannot be treated by previous works [1, 50, 55, 63] and may only have local solutions in their deterministic counterparts. Specifically, these models encompass stochastic 3D Navier-Stokes equations, stochastic 2D quasi-geostrophic equations in both critical and subcritical cases, as well as stochastic surface growth models. In particular, we can choose $g(x) = C_0 x^p$ with $C_0 > 0$ and $p > 1$ in these models (see Section 3 for details).

(ii) *Note that the main idea of the proof in [51, 52] based on a shift transformation to reduce SPDE (1.5) to a deterministic evolution equation with some random parameters.*

However, this strategy cannot handle the nonlinear multiplicative noise that plays an important role in this work. More importantly, comparing to [51, 52], we can establish the global existence and uniqueness of solutions, rather than merely local solutions.

Remark 2.3. The condition (2.3) in (\mathbf{A}_5) plays an essential role in the present work, which reflects the regularization effect of the nonlinear noise. The main idea is to utilize the fast enough growth of the nonlinear noise to prevent the potential singularity induced by other nonlinear terms, which enables to get the energy estimates in probability of solutions based on choosing a suitable Lyapunov function (see Lemma 4.1 below). More precisely, the condition (2.3) can be understood as follows:

- (i) if $g(x) = C_0x$, which corresponds to the classical coercivity condition (1.6), the diffusion coefficient $\mathcal{B}(t, \cdot)$ allows for both linear growth and superlinear growth.
- (ii) if $g(x) = C_0x^p$ with $p > 1$, the coefficient $\mathcal{B}(t, \cdot)$ needs to be of superlinear growth.

Remark 2.4. The classical variational framework (i.e. $g(x) = C_0x$ in (\mathbf{A}_3) , see e.g. [50, 63]) can be applied to PDEs/SPDEs that have been extensively studied in the literature, including (stochastic) porous medium equations, 2D Navier-Stokes equations, Cahn-Hilliard equations, and liquid crystal models. Our main results in this case also extend the very recent work [63], which investigated the well-posedness of SPDEs (1.5) with diffusion coefficients $\mathcal{B}(t, \cdot)$ satisfying the standard linear growth assumption. Instead of imposing the linear growth condition, our results can handle the case where the coefficient $\mathcal{B}(t, \cdot)$ exhibits superlinear growth, which is of independent interest.

Remark 2.5. As a consequence of the proof of Theorem 2.1, in fact we can obtain a more general result for the existence of weak solutions to (1.5) directly. More precisely, if the embedding $\mathbb{V} \subset \mathbb{H}$ is compact, (\mathbf{A}_3) - (\mathbf{A}_5) hold and $\mathcal{A}(t, \cdot)$ is pseudo-monotone from \mathbb{V} to \mathbb{V}^* for a.e. $t \in [0, T]$, then there exists a weak solution to (1.5) and the estimate (2.6) holds.

We remark that if the Gelfand triple (1.4) reduces to the case of finite dimensions, i.e. $\mathbb{V} = \mathbb{H} = \mathbb{R}^d$, the pseudo-monotonicity of $\mathcal{A}(t, \cdot)$ is equivalent to the continuity of $\mathcal{A}(t, \cdot)$. Thus, the above existence result of weak solutions is consistent with the classical theory in finite-dimensional SDEs (cf. [33, Theorem C.3]).

Based on the existence and uniqueness of solutions to (1.5), we intend to investigate the continuous dependence on the initial data in probability (in other word, the well-posedness of (1.5)) and the Feller property of the corresponding transition semigroup.

For any $\varphi \in \mathcal{B}_b(\mathbb{H})$, $t \geq 0$, we define a function $\mathcal{T}_t\varphi : \mathbb{H} \rightarrow \mathbb{R}$ by

$$\mathcal{T}_t\varphi(x) := \mathbb{E}\varphi(X_t(x)), \quad x \in \mathbb{H},$$

where $X_t(x)$ is the solution to (1.5) with the initial data x .

Remark 2.6. Based on Theorem 2.1, it is a direct consequence that $(\mathcal{T}_t)_{t \geq 0}$ is a stochastically continuous Markov semigroup on $\mathcal{B}_b(\mathbb{H})$.

Theorem 2.2. Suppose that the embedding $\mathbb{V} \subset \mathbb{H}$ is compact and that (\mathbf{A}_1) - (\mathbf{A}_5) hold. In addition, suppose that for any $u, v \in \mathbb{V}$, \mathcal{B} is locally Lipschitz in the sense that

$$(2.7) \quad \|\mathcal{B}(t, u) - \mathcal{B}(t, v)\|_{L_2(U, \mathbb{H})}^2 \leq (C + \rho(u) + \eta(v))\|u - v\|_{\mathbb{H}}^2,$$

where functions ρ, η are the same as in **(A₂)**. Let $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_{\mathbb{H}} \rightarrow 0$. Then

$$(2.8) \quad \|X(x_n) - X(x)\|_{\mathcal{C}_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

Furthermore, $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup, i.e., \mathcal{T}_t maps $C_b(\mathbb{H})$ into itself.

Remark 2.7. We would like to provide some examples for the diffusion coefficient \mathcal{B} that satisfies the aforementioned conditions **(A₅)** and (2.7). As we mentioned in Remark 2.2, we typically choose

$$g(x) = C_0 x^p \text{ with } p > 1, C_0 > 0$$

in our examples. Then the diffusion coefficient can be choose as follows.

(i) For $y \in U$, we take

$$(2.9) \quad \mathcal{B}(t, u)y := \sum_{k=1}^{\infty} b_k \|u\|_{\mathbb{H}}^m u \langle y, g_k \rangle_U,$$

where $\{g_1, g_2, \dots\}$ is an orthonormal basis of U , and the constants $b_k, k \geq 1$, satisfy

$$\gamma := \sum_{k=1}^{\infty} b_k^2 < \infty.$$

For $m = p - 1$, we assume that γ is a large constant depending on C_0 . For $m > p - 1$, we only need to assume that there is $k \geq 1$ such that $b_k \neq 0$.

For reader's convenience, we present the detailed proof that the example (2.9) satisfies the condition (2.3) as $m = p - 1$.

Proof. Note that

$$\begin{aligned} \|\mathcal{B}(t, u)^* u\|_U^2 &= \langle \mathcal{B}(t, u)^* u, \mathcal{B}(t, u)^* u \rangle_U \\ &= \langle u, \mathcal{B}(t, u) \mathcal{B}(t, u)^* u \rangle_{\mathbb{H}} \\ &= \sum_{k=1}^{\infty} b_k \|u\|_{\mathbb{H}}^{p-1} \langle u, u \rangle_{\mathbb{H}} \langle \mathcal{B}(t, u)^* u, g_k \rangle_U \\ &= \sum_{k=1}^{\infty} b_k \|u\|_{\mathbb{H}}^{p+1} \langle u, \sum_{j=1}^{\infty} b_j \|u\|_{\mathbb{H}}^{p-1} u \langle g_j, g_k \rangle_U \rangle_{\mathbb{H}} \\ &= \gamma \|u\|_{\mathbb{H}}^{2p+2}. \end{aligned}$$

For a large constant γ , we can obtain that there exists $\eta_0 \in (1, 2)$,

$$\begin{aligned} (g(\|u\|_{\mathbb{H}}^2) + \|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})}^2)(1 + \|u\|_{\mathbb{H}}^2) &\leq (C_0 \|u\|_{\mathbb{H}}^{2p} + \gamma \|u\|_{\mathbb{H}}^{2p})(1 + \|u\|_{\mathbb{H}}^2) \\ &\leq \eta_0 \gamma \|u\|_{\mathbb{H}}^{2p+2} + C, \end{aligned}$$

which implies that the condition (2.3) in **(A₅)** holds. \square

(ii) In particular, it is applicable to the 1D Wiener process \hat{W} . In fact, for $u \in \mathbb{H}$ and $y \in U$, by taking $b_k = c \delta_{1,k}$, where $c \neq 0$ is a constant, we have

$$(2.10) \quad \mathcal{B}(t, u)y := c \|u\|_{\mathbb{H}}^m u \langle y, g_1 \rangle_U.$$

For $m = p - 1$, we assume that c is a large constant depending on C_0 . For any $m > p - 1$, we can take $c = 1$ directly.

Remark 2.8. Since we have established the Feller property of the Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ associated with (1.5), a natural and important question arises investigating the existence of an invariant measure. One commonly employed technique for establishing the existence of an invariant measure is the Krylov-Bogolyubov procedure. According to the criterion presented by Maslowski and Seidler [53], we need to prove the following estimate: for any $\varepsilon > 0$ there exists $R > 0$ such that

$$(2.11) \quad \sup_{T \geq 1} \frac{1}{T} \int_0^T \mathbb{P}(\|X_t(x)\|_{\mathbb{H}} > R) dt < \varepsilon.$$

However, it is not sufficient to obtain (2.11) based on the estimate (4.2) below (see also (4.6)). This topic deserves further investigation in the future work.

2.3. Finite time extinction. The phenomenon of finite time extinction for the stochastic fast diffusion equations with linear multiplicative noises was rigorously derived by Barbu, Da Prato and Röckner [5, 6, 7, 8] and Gess [29], which displays a self-organized criticality (SOC) behavior. The purpose of this subsection is to provide a general setting for investigating finite time extinction of solutions with probability one for SPDEs (1.5) perturbed by nonlinear noises. Additionally, one may also anticipate that the regularization effect of a suitable nonlinear noise will eventually dominate the (additional) nonlinear source, which leads to finite time extinction.

Based on the existence and uniqueness results of solutions to (1.5) (i.e. Theorem 2.1), we shall suppose the following assumptions. Specifically, there are some constants $\alpha \in (1, 2)$ and $\delta > 0$ such that the following conditions hold for a.e. $t \in [0, T]$.

(**A₃**^{*}) (Enhanced Coercivity) For any $u \in \mathbb{V}$,

$$2_{\mathbb{V}^*} \langle \mathcal{A}(t, u), u \rangle_{\mathbb{V}} + \delta \|u\|_{\mathbb{V}}^{\alpha} \leq g(\|u\|_{\mathbb{H}}^2),$$

where the function g is the same as in (**A₃**) with $g(0) = 0$.

(**A₅**^{*}) For any $u \in \mathbb{V}$,

$$\|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})} = 0 \text{ if } \|u\|_{\mathbb{H}} = 0,$$

and

$$(2.12) \quad (g(\|u\|_{\mathbb{H}}^2) + \|\mathcal{B}(t, u)\|_{L_2(U, \mathbb{H})}^2) \|u\|_{\mathbb{H}}^2 \leq \alpha \|\mathcal{B}(t, u)^* u\|_U^2.$$

Remark 2.9. Note that assumption (2.12) is not comparable to assumption (2.3). However, if we choose $g(x) = C_0 x^p$, it is easy to select a nonlinear noise that satisfies assumptions (2.3) and (2.12) simultaneously, see Subsection 3.3 for details.

Let τ_e be the following extinction time

$$\tau_e := \inf \{t \geq 0 : \|X_t\|_{\mathbb{H}} = 0\},$$

where $(X_t)_{t \geq 0}$ is the solution to (1.5) given by Theorem 2.1 with initial value $x \in \mathbb{H}$.

Theorem 2.3. *Suppose that the embedding $\mathbb{V} \subset \mathbb{H}$ is compact and that (\mathbf{A}_1) , (\mathbf{A}_2) , (\mathbf{A}_3^*) , (\mathbf{A}_4) , (\mathbf{A}_5) with $\eta = \alpha$, and (\mathbf{A}_5^*) hold. Let the initial data $x \in \mathbb{H}$. Then for any $t \geq \tau_e$,*

$$(2.13) \quad \|X_t\|_{\mathbb{H}} = 0, \text{ } \mathbb{P}\text{-a.s..}$$

Moreover, we have

$$(2.14) \quad \mathbb{P}(\tau_e < \infty) = 1.$$

Remark 2.10. (i) *By applying our general setting, we are able to obtain the finite time extinction for a class of quasi-linear SPDEs driven by nonlinear multiplicative noises, including stochastic fast diffusion equations and stochastic singular p -Laplace equations (with hot sources), which seems new in the literature and demonstrates the regularization effect of nonlinear noise.*

(ii) *In the forthcoming work, we are interested in finite time extinction for the SOC models, e.g. the BTW model and the Zhang model, perturbed by nonlinear noises, which model dynamical systems that have a critical point as an attractor.*

3. EXAMPLES/APPLICATIONS

In this section, we will denote by $\Lambda \subseteq \mathbb{R}^d$ ($d \geq 1$) an open bounded domain with a smooth boundary. Let $C_0^\infty(\Lambda, \mathbb{R}^d)$ be the space of all infinitely differentiable functions from Λ to \mathbb{R}^d with compact support. For $p \geq 1$, let $L^p(\Lambda, \mathbb{R}^d)$ denote the vector valued L^p -space with the norm $\|\cdot\|_{L^p}$. For each integer $m \geq 0$, we use $W_0^{m,p}(\Lambda, \mathbb{R}^d)$ to denote the classical Sobolev space defined on Λ taking values in \mathbb{R}^d with the equivalent norm:

$$\|u\|_{W^{m,p}} := \left(\int_{\Lambda} |D^m u(x)|^p dx \right)^{\frac{1}{p}}.$$

In particular, we denote

$$\|u\|_m := \|u\|_{W^{m,2}}.$$

Below we recall the standard Gagliardo-Nirenberg interpolation inequality (cf. [56]) for the reader's convenience. If for any $1 \leq q, r < \infty$, and $0 \leq n < m$ satisfying

$$\frac{1}{p} = \frac{n}{d} + \theta \left(\frac{1}{r} - \frac{m}{d} \right) + (1 - \theta) \frac{1}{q}, \quad \frac{n}{m} \leq \theta < 1,$$

then there is a constant $C > 0$ such that

$$(3.1) \quad \|u\|_{W^{n,p}} \leq C \|u\|_{W^{m,r}}^\theta \|u\|_{L^q}^{1-\theta}, \quad u \in L^q(\Lambda, \mathbb{R}^d) \cap W^{m,2}(\Lambda, \mathbb{R}^d).$$

In the sequel, we first utilize a simple finite-dimensional example to illustrate the regularization effect by superlinear noises. Then, we present the main applications of our main results for infinite-dimensional systems.

3.1. SDEs with superlinear drifts. In this subsection, let $\mathbb{V} = \mathbb{H} = \mathbb{R}$. Consider the following nonlinear ODE on \mathbb{R}

$$(3.2) \quad \frac{dX_t}{dt} = X_t^2, \quad X_0 = x.$$

This classical equation has applications in many fields such as describing the variation of a population over time. Due to the superlinear coefficients of Eq. (3.2), if the initial data $x > 0$, the solution will blow up in finite time. In fact, it can be immediately seen that

$$X_t = \frac{1}{1/x - t}$$

is a solution of (3.2).

In this part, we are interested in ODE (3.2) with superlinear stochastic perturbation, i.e.,

$$(3.3) \quad dX_t = X_t^2 dt + c_0 X_t^m dW_t, \quad X_0 = x,$$

where $c_0 > 0, m \geq 1$, W_t is a standard 1D Wiener process defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

Applying our abstract result, we can see that there exists an appropriate nonlinear noise which effectively prevents the blow-up of solutions to Eq. (3.2).

Theorem 3.1. *Suppose either $m = \frac{3}{2}$ with $c_0 > \sqrt{2}$ or $m > \frac{3}{2}$. Then for any initial data $x \in \mathbb{R}$, (3.3) has a unique global strong solution.*

Proof. The conditions **(A₁)**, **(A₂)** and **(A₄)** can be easily proved with $\alpha = 2, \beta = 2m$ and

$$\rho(u) = \eta(u) := u + Cu^{m-1}.$$

Moreover, the condition **(A₃)** holds with

$$g(x) := c_1 x^{3/2} \text{ for } 2 < c_1 < c_0^2.$$


Concerning the condition **(A₅)**, if $m > \frac{3}{2}$, for any $\eta \in (1, 2)$ by Young's inequality we have

$$(3.4) \quad c_1 |u|^3 (1 + u^2) + c_0^2 u^{2m} (1 + u^2) \leq C + \eta c_0^2 u^{2m+2},$$


and if $m = \frac{3}{2}$, we can take $\eta \in (\frac{c_1}{c_0^2} + 1, 2)$ such that (3.4) holds. \square

In the following, we also present the numerical simulations to visually demonstrate the regularization effect of the nonlinear noise in Eq. (3.3).

Remark 3.1. *We first show the numerical simulations of solutions to ODE (3.2) and SDE (3.3) driven by linear multiplicative noise ($c_0 = 1, m = 1$) in the following figures*



Next, we present the numerical simulations of SDE (3.3) driven by superlinear multiplicative noise ($c_0 = 1, m = 2$), from which we can see the regularization effect of superlinear noise.



3.2. Stochastic 3D Navier-Stokes equations. 3D Navier-Stokes system is a classical model to describe the time evolution of an incompressible fluid, which is given as follows

$$(3.5) \quad \begin{cases} \partial_t u = \nu \Delta u - (u \cdot \nabla) u - \nabla p + f, \\ \operatorname{div}(u) = 0, u|_{\partial\Lambda} = 0, u(0) = u_0, \end{cases}$$

where u represents the velocity field of the fluid, $\nu > 0$ is the viscosity constant, p denotes the pressure and f is an external force field acting on the fluid.

Define

$$H^m := \left\{ v \in W_0^{m,2}(\Lambda, \mathbb{R}^3) : \operatorname{div}(v) = 0 \right\}.$$

Identifying H^1 with its dual space by the Riesz isomorphism, then we will use the following Gelfand triple

$$(3.6) \quad \mathbb{V} := H^2 \subseteq \mathbb{H} := H^1 \subseteq \mathbb{V}^*.$$

Let $\mathcal{P}_{\mathbb{H}}$ be the orthogonal projection operator on $L^2(\Lambda, \mathbb{R}^3)$ onto H^0 , which is called the Leray-Helmholtz projection. Then, the classical 3D Navier-Stokes system (3.5) can be reformulated in the following abstract form:

$$(3.7) \quad \partial_t u = Au + B(u) + F, \quad u(0) = u_0,$$

where operators

$$\begin{aligned} A : \mathbb{V} &\rightarrow \mathbb{V}^*, \quad Au := \nu \mathcal{P}_{\mathbb{H}} \Delta u, \\ B : \mathbb{V} \times \mathbb{V} &\rightarrow \mathbb{V}^*, \quad B(u, v) := \mathcal{P}_{\mathbb{H}}[(u \cdot \nabla)v], \quad B(u) := B(u, u) \\ F : [0, T] &\rightarrow H^0, \quad F_t = \mathcal{P}_{\mathbb{H}} f_t \end{aligned}$$

are well-defined.

In the past few decades, 3D Navier-Stokes equations have been intensively studied in the literature, but up to now, its global well-posedness is still a challenging open problem. Motivated by this, we are interested in the regularization by nonlinear noise for the following stochastic 3D Navier-Stokes equations

$$(3.8) \quad dX_t = [AX_t + B(X_t)]dt + \sum_{k=1}^{\infty} b_k \|X_t\|_1^m X_t d\beta_t^k,$$

where $m > 2$, $\sum_{k=1}^{\infty} b_k^2 \in (0, \infty)$, $\{\beta_t^k\}_{t \in [0, T]}$, $k \in \mathbb{N}$, are independent 1D Wiener processes defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

Theorem 3.2. *For any initial data $x \in \mathbb{H}$, (3.8) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $p \geq 2$, we have the following estimate*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|X_t\|_1^p + \int_0^T \|X_t\|_2^2 dt < \infty \right) = 1.$$

Furthermore, let $X_t(x)$ be the unique solution to (3.8) with the initial data x , $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_1 \rightarrow 0$, then

$$\|X(x_n) - X(x)\|_{\mathbb{C}_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In particular, the corresponding Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup in $C_b(\mathbb{H})$.

Proof. By Theorems 2.1 and 2.2, we only need to verify that the conditions (\mathbf{A}_1) - (\mathbf{A}_5) hold for equation (3.8).

Let

$$\mathcal{A}(t, u) := Au + B(u).$$

Then the conditions (\mathbf{A}_1) - (\mathbf{A}_4) hold for \mathcal{A} , whose proof is similar to Example 5.2.23 in [52], we include it here for completeness.

The hemicontinuity (\mathbf{A}_1) is obvious since B is a bilinear map.

By the Agmon inequality (see e.g. (A.29) in [28]),

$$(3.9) \quad \|u\|_{L^\infty} \leq C \|u\|_1 \|u\|_2, \quad u \in H^2,$$

we have

$$\begin{aligned} & \mathbb{V}^* \langle B(u) - B(v), u - v \rangle_{\mathbb{V}} \\ &= \langle B(u) - B(v), (-\Delta)(u - v) \rangle_{L^2} \\ &\leq \|u - v\|_2 \|(u \cdot \nabla)u - (v \cdot \nabla)v\|_{L^2} \\ &\leq \|u - v\|_2 (\|u\|_{L^\infty} \|\nabla u - \nabla v\|_{L^2} + \|u - v\|_{L^\infty} \|\nabla v\|_{L^2}) \\ &\leq \|u - v\|_2 \left(\|u\|_{L^\infty} \|u - v\|_1 + C \|u - v\|_2^{1/2} \|u - v\|_1^{1/2} \|v\|_1 \right) \\ &\leq \frac{\nu}{2} \|u - v\|_2^2 + C (\|u\|_2^2 \|u\|_1^2 + \|v\|_1^4) \|u - v\|_1^2, \quad u, v \in \mathbb{V}, \end{aligned}$$

then

$$\begin{aligned} & \mathbb{V}^* \langle Au + B(u) - Av - B(v), u - v \rangle_{\mathbb{V}} \\ &\leq -\frac{\nu}{2} \|u - v\|_2^2 + C (\|u\|_2^2 \|u\|_1^2 + \|v\|_1^4) \|u - v\|_1^2, \quad u, v \in \mathbb{V}, \end{aligned}$$

which implies the local monotonicity **(A₂)** holds with $\alpha = 2, \beta = m$ and

$$\rho(u) := \|u\|_2^2 \|u\|_1^2, \quad \eta(v) = \|v\|_1^4.$$

In particular, there exists a constant $C > 0$ such that

$$\mathbb{V}^* \langle Au + B(u), u \rangle_{\mathbb{V}} \leq -\frac{\nu}{2} \|u\|_2^2 + C(1 + \|u\|_1^6), \quad u \in \mathbb{V}.$$

Then, **(A₃)** holds with $g(x) = Cx^3$, i.e.,

$$\mathbb{V}^* \langle \mathcal{A}(t, u), v \rangle_{\mathbb{V}} \leq -\frac{\nu}{2} \|u\|_2^2 + C(1 + \|u\|_1^6), \quad u \in \mathbb{V}.$$

Concerning the growth condition, by (3.9) we have

$$\|B(u)\|_{\mathbb{V}^*}^2 \leq \|(u \cdot \nabla)u\|_{L^2}^2 \leq \|u\|_{L^\infty}^2 \|\nabla u\|_{L^2}^2 \leq C \|u\|_2 \|u\|_1^3 \leq C \|u\|_2^2 \|u\|_1^2, \quad u \in \mathbb{V}.$$

Hence, **(A₄)** holds.

Finally, as shown in Remark 2.7, we can see that condition **(A₅)** holds. The proof is completed. \square


Remark 3.2. (i) In the past several decades, the stochastic Navier-Stokes equations have attracted great interest in the literature (cf. [20, 24, 25, 26, 27, 32, 35, 45]). The existence of martingale solutions of stochastic 3D Navier-Stokes equations was first established by Flandoli and Gatarek in [22]. Da Prato and Debussche [20] proved the existence of Markov section for stochastic 3D Navier-Stokes equations, see also [27]. Recently, Hofmanová et al. [38] develop the convex integration method to prove the nonuniqueness in law for stochastic 3D Navier-Stokes equations. However, in this work, by applying our abstract result we shows that for any initial data in H^1 , there exists an appropriate noise perturbation such that the stochastic 3D Navier-Stokes equations admit a unique global strong solution. Here the “strong solution” is both in the analytical sense and in the probabilistic sense.

(ii) In [27], the authors also proved the Feller property of the selected Markov semigroup, but due to the lack of continuity of solutions in the space of finite energy, the Markov

property holds almost everywhere in time. In Theorem 3.2 we prove the Feller property of the corresponding Markov semigroup for the stochastic 3D Navier-Stokes equations, rather than regarding a Markov selection, which supports us to further investigate the long-time behaviour and the ergodicity property in the future work.

Remark 3.3. Throughout this section, for sake of simplicity, we only present an explicit form of noise term in the examples. However, we mention that our results can also be applied to more general type of noises, which satisfy the condition (\mathbf{A}_5) and (2.7).

3.3. Stochastic singular p -Laplace equations with heat sources. Before introducing the main results, we would like to present the numerical simulations for 1D ODE and SDE in the following figures to show the regularization effect of noise on the finite time extinction of solutions



Inspired by the above example, in this part, we are interested in the regularization by nonlinear noise on the finite time extinction for the singular p -Laplace equations with nonlinear sources. Consider

$$(3.10) \quad \partial_t u = \operatorname{div}(|\nabla u|^{p-2} \nabla u) + u^2, \quad u(0) = u_0,$$

where $1 < p < 2$. This model appears in the combustion theory, where u_t represents the temperature, the term $\operatorname{div}(|\nabla u|^{p-2} \nabla u)$ represents the thermal diffusion and the nonlinear source u^2 is physically called the “hot source” (cf. [21, 71]).

It is known that the solutions of Eq. (3.10) may blow up under certain initial condition, thus there is no global solution in general (see e.g. [47]). Taking account of random noises, we study the following 1D stochastic singular p -Laplace equations on the interval

$\Lambda := [0, L]$ as follows

$$(3.11) \quad \begin{cases} dX_t = \operatorname{div}(|\nabla X_t|^{p-2} \nabla X_t) dt + X_t^2 dt + \sum_{k=1}^{\infty} b_k \|X_t\|_{L^2}^{\frac{p}{3p-3}} X_t d\beta_t^k, \\ X_t|_{\partial\Lambda} = 0, \end{cases}$$

where $\gamma := \sum_{k=1}^{\infty} b_k^2$ is a large enough constant, $\{\beta_t^k\}_{t \in [0, T]}$, $k \in \mathbb{N}$, are independent 1D Wiener processes defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

Define the following Gelfand triple:

$$\mathbb{V} := W^{1,p}(\Lambda, \mathbb{R}) \subseteq \mathbb{H} := L^2(\Lambda, \mathbb{R}) \subseteq \mathbb{V}^*.$$

Theorem 3.3. *For any initial data $x \in \mathbb{H}$, (3.11) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $s \geq 2$,*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|X_t\|_{L^2}^s + \int_0^T \|X_t\|_{W^{1,p}}^p dt < \infty \right) = 1.$$

Furthermore, let $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_{L^2} \rightarrow 0$. Let $X_t(x)$ be the unique solution to (3.23) with the initial data x . Then

$$\|X(x_n) - X(x)\|_{\mathbb{C}_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In particular, the corresponding Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup in $C_b(\mathbb{H})$.

Proof. Let

$$\mathcal{A}(t, u) := \operatorname{div}(|\nabla u|^{p-2} \nabla u) + u^2.$$

For any $u, v \in \mathbb{V}$, by Young's inequality we have

$$\begin{aligned} \mathbb{V}^* \langle u^2 - v^2, u - v \rangle_{\mathbb{V}} &= \langle u^2 - v^2, u - v \rangle_{L^2} \\ &\leq C \int_{\Lambda} |u|(u-v)^2 dx + C \int_{\Lambda} |v|(u-v)^2 dx \\ &\leq C(\|u\|_{L^\infty} + \|v\|_{L^\infty}) \|u - v\|_{L^2}^2 \\ &\leq C(\|u\|_{W^{1,p}} + \|v\|_{W^{1,p}}) \|u - v\|_{L^2}^2, \end{aligned}$$

where we have used the Sobolev embedding $W_0^{1,p}(\Lambda) \subseteq L^\infty(\Lambda)$ for $p > d = 1$.

Then

$$\mathbb{V}^* \langle \mathcal{A}(t, u) - \mathcal{A}(t, v), u - v \rangle_{\mathbb{V}} \leq C(1 + \|u\|_{W^{1,p}}^p + \|v\|_{W^{1,p}}^p) \|u - v\|_{L^2}^2,$$

which implies the local monotonicity **(A₂)** holds with $\alpha = p$ and

$$\rho(u) = \eta(u) := \|u\|_{W^{1,p}}^p.$$

By interpolation inequality (3.1) and Young's inequality, for any $u \in \mathbb{V}$ there exists constants $C, C_0 > 0$ such that

$$(3.12) \quad \begin{aligned} \mathbb{V}^* \langle \mathcal{A}(u), u \rangle_{\mathbb{V}} &= - \|u\|_{W^{1,p}}^p + \|u\|_{L^3}^3 \\ &\leq - \|u\|_{W^{1,p}}^p + C \|u\|_{W^{1,p}}^{3\theta} \|u\|_{L^2}^{3(1-\theta)} \\ &\leq - \frac{1}{2} \|u\|_{W^{1,p}}^p + C_0 \|u\|_{L^2}^{\frac{8p-6}{3p-3}}, \end{aligned}$$

where $\theta = \frac{p}{9p-6} \in (0, 1)$. Thus, we can see that the condition **(A₃)** holds with $g(x) = C_0 x^{\frac{4p-3}{3p-3}}$.

For any $u, v \in \mathbb{V}$,

$$\begin{aligned} \mathbb{V}^* \langle \mathcal{A}(t, u), v \rangle_{\mathbb{V}} &= - \int_{\Lambda} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx + \int_{\Lambda} u^2 \cdot v dx \\ &\leq \left(\int_{\Lambda} |\nabla u|^p dx \right)^{\frac{p-1}{p}} \left(\int_{\Lambda} |\nabla v|^p dx \right)^{\frac{1}{p}} + C \|v\|_{L^\infty} \left(\int_{\Lambda} |u|^2 dx \right) \\ &\leq \|u\|_{W^{1,p}}^{p-1} \|v\|_{W^{1,p}} + C \|u\|_{L^2}^2 \|v\|_{W^{1,p}}. \end{aligned}$$

Therefore, the growth condition **(A₄)** holds, namely for any $u \in \mathbb{V}$,

$$\|\mathcal{A}(t, u)\|_{\mathbb{V}^*}^{\frac{p}{p-1}} \leq C (\|u\|_{W^{1,p}}^p + \|u\|_{L^2}^{\frac{2p}{p-1}}).$$

Finally, it is easy to show that **(A₁)** and **(A₅)** hold. We complete the proof by applying Theorems 2.1 and 2.2. \square

Let τ_e be the extinction time

$$(3.13) \quad \tau_e := \inf \{t \geq 0 : \|X_t\|_{L^2} = 0\},$$

where $(X_t)_{t \geq 0}$ is the solution to (3.11) with initial value $x \in \mathbb{H}$. Then, as a result of Theorems 2.3, we get the following finite time extinction result for Eq. (3.11).

Theorem 3.4. *Let the initial data $x \in \mathbb{H}$. Then for any $t \geq \tau_e$,*

$$\|X_t\|_{L^2} = 0, \quad \mathbb{P}\text{-a.s.}$$

Moreover, we have

$$\mathbb{P}(\tau_e < \infty) = 1.$$

Proof. By (3.12), we can see the condition **(A₃^{*})** holds. Moreover, by the proof in Remark 2.7, it is easy to see that condition **(A₅^{*})** holds. The proof is completed. \square

Remark 3.4. *Note that the deterministic singular p -Laplace equations (3.10) may blow up in finite time. We show that there exists an appropriate random intervention such that the solution of the corresponding stochastic equation is extinct in finite time. This phenomenon is new and interesting, indicating the regularization by nonlinear noise on the long-time behavior of quasi-linear equations.*

3.4. Stochastic surface growth model. In this part, we apply our main results to study the following surface growth model

$$(3.14) \quad \partial_t u = -\partial_x^4 u - \partial_x^2 u + \partial_x^2 (\partial_x u)^2, \quad u(0) = u_0,$$

where $\partial_x, \partial_x^2, \partial_x^4$ denote the first, second and fourth spatial derivatives, respectively. This model appears in the theory of growth of surfaces, which describes an amorphous material deposited on an initially flat surface in high vacuum (cf. [9, 59] and the references therein).

Due to the highly nonlinear characteristics in the model (3.14), the global well-posedness for the 1D surface growth model is still an open problem in both the deterministic and

stochastic case. In particular, it is known in the literature that this model has some similar features of difficulty as the 3D Navier-Stokes equation.

Taking account of random noises, the equation is formulated, on the interval $\Lambda := [0, L]$, as follows

$$(3.15) \quad \begin{cases} dX_t = [-\partial_x^4 X_t - \partial_x^2 X_t + \partial_x^2 (\partial_x X_t)^2] dt + \sum_{k=1}^{\infty} b_k \|X_t\|_2^m X_t d\beta_t^k, \\ X_t|_{\partial\Lambda} = 0, \end{cases}$$

where $m > 2$, $\sum_{k=1}^{\infty} b_k^2 \in (0, \infty)$, $\{\beta_t^k\}_{t \in [0, T]}$, $k \in \mathbb{N}$, are independent 1D Wiener processes defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

In this work, by applying our abstract result we demonstrate that there exists an appropriate noise perturbation such that the 1D surface growth model has a unique global strong solution. To this end, we take the following Gelfand triple

$$\mathbb{V} := W_0^{4,2}([0, L]) \subseteq \mathbb{H} := W^{2,2}([0, L]) \subseteq \mathbb{V}^*.$$

Theorem 3.5. *For any initial data $x \in \mathbb{H}$, (3.15) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $p \geq 2$,*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|X_t\|_2^p + \int_0^T \|X_t\|_4^2 dt < \infty \right) = 1.$$

Furthermore, let $X_t(x)$ be the unique solution to (3.15) with the initial data x , $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_2 \rightarrow 0$. Then

$$\|X(x_n) - X(x)\|_{\mathbb{C}_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In particular, the corresponding Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup in $C_b(\mathbb{H})$.

Proof. The conditions **(A₁)**-**(A₅)** hold for (3.15) with $\alpha = 2, \beta = m$ and $g(x) = Cx^3$, whose proof follows from Section 3.4 in [51], we omit the details. \square

Remark 3.5. *The solution obtained here for the stochastic surface growth model is a global strong solution both in the PDE and probability sense. We should remark that the local solution for the stochastic surface growth model has been established in [51, 65]. Blömker, Flandoli and Romito [9] proved the existence of weak martingale solutions and Markov selections for this model with space-time white noise. As far as we know, Theorem 3.5 is the first global well-posedness result of stochastic surface growth model.*

3.5. Stochastic quasi-geostrophic equation. Quasi-geostrophic equation is an important model in geophysical fluid dynamics, which is a special case of general quasi-geostrophic approximations for atmospheric and oceanic fluid flows with small Rossby and Ekman numbers. The following dissipative quasi-geostrophic equation in the 2D periodic domain \mathbb{T}^2 has attracted quite a lot of attention lately from various scholars

$$(3.16) \quad \begin{cases} \partial_t \theta = -(-\Delta)^\gamma \theta - (u \cdot \nabla) \theta, \\ u = (u_1, u_2) = (-R_2 \theta, R_1 \theta) = R^\perp \theta, \\ \theta(0) = \theta_0, \end{cases}$$

where $\gamma \in (0, 1)$, $\theta : \mathbb{T}^2 \rightarrow \mathbb{R}$ is a scalar function, R_1 and R_2 are the usual Riesz transforms in \mathbb{T}^2 .

When $\gamma > 1/2$ (the so-called subcritical case), the global well-posedness for Eq. (3.16) is well-known, see e.g. [61]. The critical case ($\gamma = 1/2$) exhibits similar features (singularities) as the 3D Navier-Stokes equations. The global well-posedness for the critical quasi-geostrophic equation is a challenging issue, which has made significant progress by Caffarelli and Vasseur [17]. They have provided the existence of a classical solution for $\gamma = 1/2$. Another highly significant result established by Kiselev et al. in [42], demonstrating that solutions for $\gamma = 1/2$ with periodic C^∞ initial data remain C^∞ for all times. In the remarkable work [18], Constantin and Nguyen proved existence of global weak L^2 solutions of the inviscid quasi-geostrophic equation in bounded domains. The blow-up or global regularity for $\gamma < 1/2$ (supercritical case) remains an open problem for the quasi-geostrophic equation.

Set $\Lambda := (-\Delta)^{1/2}$, define

$$\|f\|_s^2 := \|\Lambda^s f\|_{L^2}^2 = \sum_k |k|^{2s} \langle f, e_k \rangle^2, \quad s \geq 0,$$

where $\{e_k\}$ is the eigenbasis of $-\Delta$. We use H^s to denote the Sobolev space of all $f \in L^2(\mathbb{T}^2, \mathbb{R})$ with $\int_{\mathbb{T}^2} f dx = 0$ for which $\|f\|_s$ is finite.

Define the commutator

$$[\Lambda^s, f]g := \Lambda^s(fg) - f\Lambda^s g.$$

The following commutator estimate is very important for later use (see [41]).

Lemma 3.1. (*Commutator estimate*) *Suppose that $s > 0$, $p, p_2, p_3 \in (1, \infty)$ and $p_1, p_4 \in (1, \infty]$ satisfy*

$$(3.17) \quad \frac{1}{p} \geq \frac{1}{p_1} + \frac{1}{p_2}, \quad \frac{1}{p} \geq \frac{1}{p_3} + \frac{1}{p_4}.$$

Then we have

$$\|[\Lambda^s, f]g\|_{L^p} \leq C (\|\nabla f\|_{L^{p_1}} \|\Lambda^{s-1} g\|_{L^{p_2}} + \|\Lambda^s f\|_{L^{p_3}} \|g\|_{L^{p_4}}).$$

We recall the following product estimate (see e.g. [61, 66]).

Lemma 3.2. *Suppose that $s > 0$, $p, p_2, p_3 \in (1, \infty)$, $p_1, p_4 \in (1, \infty]$ satisfy (3.17). We have*

$$\|\Lambda^s(fg)\|_{L^p} \leq C (\|f\|_{L^{p_1}} \|\Lambda^s g\|_{L^{p_2}} + \|\Lambda^s f\|_{L^{p_3}} \|g\|_{L^{p_4}}).$$

Moreover, we also recall the following Sobolev embedding theorem (see e.g. [67]).

Lemma 3.3. *If $0 \leq s < \frac{d}{2}$ and $\frac{1}{p} + \frac{s}{d} = \frac{1}{q}$, then $H^s \subset L^p$. Moreover, there is a constant $C = C(s, p) > 0$ such that*

$$\|f\|_{L^p} \leq C \|\Lambda^s f\|_{L^q}.$$

For the quasi-geostrophic equation, we choose the following Gelfand triple

$$\mathbb{V} := H^2 \subseteq \mathbb{H} := H^{2-\gamma} \subseteq \mathbb{V}^*.$$

Now we consider the following stochastic quasi-geostrophic equation in \mathbb{T}^2

$$(3.18) \quad \begin{cases} dX_t = -(-\Delta)^\gamma X_t dt - (Y_t \cdot \nabla) X_t dt + \sum_{k=1}^{\infty} b_k \|X_t\|_{2-\gamma}^m X_t d\beta_t^k, \\ Y_t = (-R_2 X_t, R_1 X_t) = R^\perp X_t, \end{cases}$$

where $m > 1$, $\sum_{k=1}^{\infty} b_k^2 \in (0, \infty)$, $\{\beta_t^k\}_{t \in [0, T]}$, $k \in \mathbb{N}$, are independent 1D Wiener processes defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$.

Theorem 3.6. *Fix $\gamma \geq \frac{1}{2}$. For any initial data $x \in \mathbb{H}$, (3.18) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $p \geq 2$,*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|X_t\|_{2-\gamma}^p + \int_0^T \|X_t\|_2^2 dt < \infty \right) = 1.$$

Furthermore, let $X_t(x)$ be the unique solution to (3.18) with the initial data x , $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_{2-\gamma} \rightarrow 0$. Then

$$\|X(x_n) - X(x)\|_{C_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In particular, the corresponding Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup in $C_b(\mathbb{H})$.

Proof. We first check that the conditions (\mathbf{A}_1) - (\mathbf{A}_4) hold for

$$\mathcal{A}(t, u) := -(-\Delta)^\gamma u - (v \cdot \nabla)u,$$

where $v = (-R_2 u, R_1 u) = R^\perp u$.

The hemicontinuity (\mathbf{A}_1) is obvious since $(-\Delta)^\gamma u$ is linear and $(v \cdot \nabla)u$ is bilinear. For any $u_1, u_2 \in \mathbb{V}$,

$$(3.19) \quad \begin{aligned} \mathbb{V}^* \langle \mathcal{A}(t, u_1) - \mathcal{A}(t, u_2), u_1 - u_2 \rangle_{\mathbb{V}} &= -\|u_1 - u_2\|_2^2 + \mathbb{V}^* \langle v_1 \nabla u_1 - v_2 \nabla u_2, u_1 - u_2 \rangle_{\mathbb{V}} \\ &= -\|u_1 - u_2\|_2^2 + \mathbb{V}^* \langle (v_1 - v_2) \nabla u_2, u_1 - u_2 \rangle_{\mathbb{V}} \\ &\quad + \mathbb{V}^* \langle v_1 \nabla (u_1 - u_2), u_1 - u_2 \rangle_{\mathbb{V}}, \end{aligned}$$

where $v_i = (-R_2 u_i, R_1 u_i) = R^\perp u_i$, $i = 1, 2$.

By the singular integral theory of Calderón and Zygmund (cf. [67, Chapter 3]), for any $s \geq 0$, $p \in (1, \infty)$, there is a constant C_{sp} such that

$$(3.20) \quad \|\Lambda^s v\|_{L^p} \leq C_{sp} \|\Lambda^s u\|_{L^p},$$

where $v = (-R_2 u, R_1 u) = R^\perp u$.

Then by Lemmas 3.2-3.3, Young's inequality and (3.20), we have

$$\begin{aligned} &\mathbb{V}^* \langle (v_1 - v_2) \nabla u_2, u_1 - u_2 \rangle_{\mathbb{V}} \\ &= \langle \Lambda^{2-\gamma}((v_1 - v_2) \nabla u_2), \Lambda^{2-\gamma}(u_1 - u_2) \rangle_{L^2} \\ &\leq \|\Lambda^{2-2\gamma}((v_1 - v_2) \nabla u_2)\|_{L^2} \|\Lambda^2(u_1 - u_2)\|_{L^2} \\ &\leq C \left(\|\Lambda^{2-2\gamma}(v_1 - v_2)\|_{L^{p_1}} \|\nabla u_2\|_{L^{p_2}} \right. \\ &\quad \left. + \|v_1 - v_2\|_{L^\infty} \|\Lambda^{2-2\gamma}(\nabla u_2)\|_{L^2} \right) \|\Lambda^2(u_1 - u_2)\|_{L^2} \end{aligned}$$

$$\begin{aligned}
&\leq C \left(\|\Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \|\Lambda^{2-\gamma}u_2\|_{L^2} \right. \\
&\quad \left. + \|u_1 - u_2\|_{2-\gamma} \|\Lambda^2u_2\|_{L^2} \right) \|\Lambda^2(u_1 - u_2)\|_{L^2} \\
(3.21) \quad &\leq \frac{1}{4} \|u_1 - u_2\|_2^2 + C \|u_2\|_2^2 \|u_1 - u_2\|_{2-\gamma}^2,
\end{aligned}$$

where $\frac{1}{p_1} = \frac{1}{2} - \frac{\gamma}{2}$, $\frac{1}{p_2} = \frac{\gamma}{2}$. Also by Lemmas 3.1 and 3.3, Young's inequality and (3.20), we obtain that

$$\begin{aligned}
&\mathbb{V}^* \langle v_1 \nabla(u_1 - u_2), u_1 - u_2 \rangle_{\mathbb{V}} \\
&= \langle \Lambda^{2-\gamma}(v_1 \nabla(u_1 - u_2)) - v_1 \nabla \Lambda^{2-\gamma}(u_1 - u_2), \Lambda^{2-\gamma}(u_1 - u_2) \rangle_{L^2} \\
&\leq \|\Lambda^{2-\gamma}(v_1 \nabla(u_1 - u_2)) - v_1 \nabla \Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \|\Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \\
&\leq C \left(\|\nabla v_1\|_{L^{p_1}} \|\Lambda^{1-\gamma} \nabla(u_1 - u_2)\|_{L^{p_2}} \right. \\
&\quad \left. + \|\nabla(u_1 - u_2)\|_{L^{p_3}} \|\Lambda^{2-\gamma}(v_1)\|_{L^{p_4}} \right) \|\Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \\
&\leq C \left(\|\Lambda^{2-\gamma}(u_1)\|_{L^2} \|\Lambda^2(u_1 - u_2)\|_{L^2} \right. \\
&\quad \left. + \|\Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \|\Lambda^2u_1\|_{L^2} \right) \|\Lambda^{2-\gamma}(u_1 - u_2)\|_{L^2} \\
(3.22) \quad &\leq \frac{1}{4} \|u_1 - u_2\|_{\mathbb{V}}^2 + C \|u_1\|_2^2 \|u_1 - u_2\|_{2-\gamma}^2,
\end{aligned}$$

where $\frac{1}{p_1} = \frac{1}{p_3} = \frac{\gamma}{2}$, $\frac{1}{p_2} = \frac{1}{p_4} = \frac{1}{2} - \frac{\gamma}{2}$.

Collecting (3.19)-(3.22), we have

$$\begin{aligned}
&\mathbb{V}^* \langle \mathcal{A}(t, u_1) - \mathcal{A}(t, u_2), u_1 - u_2 \rangle_{\mathbb{V}} \\
&\leq -\frac{1}{2} \|u_1 - u_2\|_2^2 + C (\|u_1\|_2^2 + \|u_2\|_2^2) \|u_1 - u_2\|_{2-\gamma}^2, \quad u_1, u_2 \in \mathbb{V},
\end{aligned}$$

which implies that the local monotonicity **(A₂)** holds with $\alpha = 2$ and

$$\rho(u) = \eta(u) := \|u\|_2^2.$$

Similar to the proof of (3.22), we have that for any $u_1 \in \mathbb{V}$,

$$\begin{aligned}
\mathbb{V}^* \langle v_1 \nabla u_1, u_1 \rangle_{\mathbb{V}} &= \langle \Lambda^{2-\gamma}(v_1 \nabla u_1) - v_1 \nabla \Lambda^{2-\gamma}(u_1), \Lambda^{2-\gamma}(u_1) \rangle_{L^2} \\
&\leq C \|\Lambda^{2-\gamma}(u_1)\|_{L^2} \|\Lambda^2u_1\|_{L^2} \|\Lambda^{2-\gamma}(u_1)\|_{L^2} \\
&\leq \frac{1}{2} \|u_1\|_2^2 + C \|u_1\|_{2-\gamma}^4.
\end{aligned}$$

Then we can see that the generalized coercivity condition **(A₃)** holds with $g(x) = Cx^2$, i.e.,

$$\begin{aligned}
\mathbb{V}^* \langle \mathcal{A}(t, u), u \rangle_{\mathbb{V}} &= -\|u\|_2^2 + \mathbb{V}^* \langle v \nabla u, u \rangle_{\mathbb{V}} \\
&\leq -\frac{1}{2} \|u\|_2^2 + C \|u\|_{2-\gamma}^4, \quad u \in \mathbb{V}.
\end{aligned}$$

Finally, for any $u \in \mathbb{V}$, by Lemmas 3.2-3.3 we have

$$\|\mathcal{A}(t, u)\|_{\mathbb{V}^*} \leq \|\Lambda^{2\gamma}u\|_{2-2\gamma} + \|v \nabla u\|_{2-2\gamma}$$

$$\begin{aligned}
&\leq \|u\|_2 + C(\|\Lambda^{2-2\gamma}v\|_{L^{p_1}}\|\nabla u\|_{L^{p_2}} + \|\Lambda^{2-2\gamma}(\nabla u)\|_{L^2}\|v\|_{L^\infty}) \\
&\leq \|u\|_2 + C(\|\Lambda^{2-\gamma}u\|_{L^2}^2 + \|\Lambda^2u\|_{L^2}\|\Lambda^{2-\gamma}u\|_{L^2}),
\end{aligned}$$

where $\frac{1}{p_1} = \frac{1}{2} - \frac{\gamma}{2}$, $\frac{1}{p_2} = \frac{\gamma}{2}$. Hence, **(A₄)** holds, namely,

$$\|\mathcal{A}(t, u)\|_{\mathbb{V}^*}^2 \leq C\|u\|_2^2(1 + \|u\|_{2-\gamma}^2).$$

Finally, we can see from Remark 2.7 that the condition **(A₅)** holds. Therefore, the conclusion follows directly from Theorems 2.1 and 2.2. \square

Remark 3.6. (i) *The stochastic quasi-geostrophic equation, as an important fluid equation, has been extensively studied in recent years. We mention some known results concerning this model. Röckner, Zhu and Zhu [66] first proved the existence of weak martingale solutions and Markov selections for stochastic quasi-geostrophic equation in the 2D periodic domain \mathbb{T}^2 when $\gamma \in (0, 1)$. In the subcritical case $\gamma > \frac{1}{2}$, they established the existence and uniqueness of probabilistically strong solutions. Later, Brzeźniak and Motyl [14] proved the existence of martingale solutions in the whole space \mathbb{R}^d , and in the 2D subcritical case they also proves the pathwise uniqueness of solutions. A result on the regularization by noise with high probability for the stochastic quasi-geostrophic equation was proved in [65], see also Buckmaster et al. [16] for the regularization by noise with high probability for the inviscid quasi-geostrophic equation with sufficiently small smooth initial values.*

(ii) *Different from the aforementioned works, with the help of the superlinear noise, we can obtain the global well-posedness in the sense of probability one for the critical case, and we do not need to assume any small initial value condition.*

Instead of the aforementioned models, our results are applicable to all the examples mentioned in [52, 63] perturbed by both linear and superlinear noises. In the sequel, we only apply our general results to the stochastic fast diffusion equations to keep down the length of this paper.

3.6. Stochastic fast diffusion equations. Fast diffusion equation arises in the description of a large variety of physical phenomena and processes including fluid flows in porous media, diffusion processes in kinetic gas theory, heat transfer in plasmas and population dynamics (cf. [7, 29]).

In this part, we consider the stochastic fast diffusion equations, with either $d = 1, 2$, $r \in (0, 1)$ or $d \geq 3$, $r \in [\frac{d-2}{d+2}, 1)$,

$$(3.23) \quad \begin{cases} dX_t = \Delta(|X_t|^{r-1}X_t)dt + \sum_{k=1}^{\infty} b_k \|X_t\|_{-1}^m X_t d\beta_t^k, \\ X_t|_{\partial\Lambda} = 0, \end{cases}$$

where $m \geq 0$, $\sum_{k=1}^{\infty} b_k^2 \in (0, \infty)$, $\{\beta_t^k\}_{t \in [0, T]}$, $k \in \mathbb{N}$, are independent 1D Wiener processes defined on a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$, $\|\cdot\|_{-1}$ denotes the norm of dual space $(H_0^1(\Lambda, \mathbb{R}))^*$.

Define the following Gelfand triple

$$\mathbb{V} := L^{r+1}(\Lambda, \mathbb{R}) \subseteq \mathbb{H} := (H_0^1(\Lambda, \mathbb{R}))^* \subseteq \mathbb{V}^*.$$

We state the main result concerning the well-posedness of Eq. (3.23).

Theorem 3.7. *For any initial data $x \in \mathbb{H}$, (3.23) has a unique strong solution in the sense of Definition 2.3. Moreover, for any $p \geq 2$,*

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|X_t\|_{-1}^p + \int_0^T \|X_t\|_{L^{r+1}}^{r+1} dt < \infty \right) = 1.$$

Furthermore, let $\{x_n\}_{n \in \mathbb{N}}$ and x be a sequence with $\|x_n - x\|_{-1} \rightarrow 0$. Let $X_t(x)$ be the unique solution to (3.23) with the initial data x . Then

$$\|X(x_n) - X(x)\|_{\mathbb{C}_T(\mathbb{H})} \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In particular, the corresponding Markov semigroup $(\mathcal{T}_t)_{t \geq 0}$ is a Feller semigroup in $C_b(\mathbb{H})$.

Proof. It is known that the map $\mathcal{A}(t, u) := \Delta(|u|^{r-1}u)$ satisfies (\mathbf{A}_1) - (\mathbf{A}_4) with $g(x) = 0$, we refer to [52, Example 4.1.11] for some details. Moreover, it is easy to see that the conditions (\mathbf{A}_5) and (2.7) hold. By applying Theorems 2.1 and 2.2, the conclusions hold. \square

Next, we present the finite time extinction for stochastic fast diffusion equations. Let τ_e be the extinction time

$$\tau_e := \inf \{t \geq 0 : \|X_t\|_{-1} = 0\},$$

where $(X_t)_{t \geq 0}$ is the solution to (3.23) with initial value $x \in \mathbb{H}$. Then, by applying Theorems 2.3, we have the following result.

Theorem 3.8. *Let the initial data $x \in \mathbb{H}$. Then for any $t \geq \tau_e$,*

$$\|X_t\|_{-1} = 0, \text{ } \mathbb{P}\text{-a.s..}$$

Moreover, we have

$$\mathbb{P}(\tau_e < \infty) = 1.$$

Remark 3.7. *To the best of our knowledge, this is the first paper of investigating finite time extinction for stochastic fast diffusion equations driven by nonlinear noises. In addition, comparing to previous studies [5, 6, 7, 8], we can derive the extinction in finite time with probability one for all spatial dimensions (with the usual dimension-dependence restriction on r).*

4. PROOF OF MAIN RESULTS

In this part, we aim to prove Theorems 2.1-2.3. More precisely, in subsection 4.1, we consider the Galerkin approximation of SPDE (1.5), and derive a priori bounds in probability by choosing a suitable Lyapunov functions. In subsection 4.2, we present the tightness of the Galerkin approximating solutions by employing the stopping time technique. In subsection 4.3, we prove the existence of weak solutions to (1.5) combining the theory of pseudo-monotone operators, the stochastic compactness approach and the

Jakubowski's version of the Skorokhod theorem. In subsection 4.4, we show the pathwise uniqueness of solutions. Then, the existence and uniqueness of strong solutions to (1.5) follow from the infinite-dimensional version of Yamada-Watanabe theorem. In subsection 4.5, the continuous dependence on the initial data in probability and the Feller property of the Markov semigroup are derived. In subsection 4.6, we prove the finite time extinction of solutions to (1.5) by employing different Lyapunov functions.

4.1. Energy estimates. Let $\{e_1, e_2, \dots\} \subset \mathbb{V}$ be an orthonormal basis (ONB for short) on \mathbb{H} . Consider the maps $\mathcal{P}_n : \mathbb{V}^* \rightarrow \mathbb{H}_n$, $n \in \mathbb{N}$, given by

$$\mathcal{P}_n x := \sum_{i=1}^n \mathbb{V}^* \langle x, e_i \rangle_{\mathbb{V}} e_i, \quad x \in \mathbb{V}^*,$$

where $\mathbb{H}_n := \text{span}\{e_1, e_2, \dots, e_n\}$.

If we restrict \mathcal{P}_n to \mathbb{H} , denoted by $\mathcal{P}_n|_{\mathbb{H}}$, then it is an orthogonal projection onto \mathbb{H}_n on \mathbb{H} . Denote by $\{g_1, g_2, \dots\}$ the ONB of U . Let

$$W_t^{(n)} := \tilde{\mathcal{P}}_n W_t = \sum_{i=1}^n \langle W_t, g_i \rangle_U g_i, \quad n \in \mathbb{N},$$

where $\tilde{\mathcal{P}}_n$ is an orthonormal projection onto $U^n := \text{span}\{g_1, g_2, \dots, g_n\}$ on U .

For any $n \in \mathbb{N}$, we consider the following stochastic equation on \mathbb{H}_n ,

$$(4.1) \quad dX_t^{(n)} = \mathcal{P}_n \mathcal{A}(t, X_t^{(n)}) dt + \mathcal{P}_n \mathcal{B}(t, X_t^{(n)}) dW_t^{(n)},$$

with initial value $x^{(n)} := \mathcal{P}_n x$. Under **(A₁)**, **(A₃)**, **(A₄)** and the assumption (2.4), it is clear that there exists a weak solution to (4.1) in the sense of Definition 2.2 up to its life time. Furthermore, under the assumption (2.3), the solution is non-explosive (see Lemma 4.1), namely, there is a global weak solution to (4.1).

We have the following a priori estimates based on choosing a suitable Lyapunov function.

Lemma 4.1. *Suppose that the assumptions in Theorem 2.1 hold. For any $\varepsilon > 0$, there exists $\mathcal{K} > 0$ such that for any $p \geq 2$,*

$$(4.2) \quad \sup_{n \in \mathbb{N}} \mathbb{P} \left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}}^p + \int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq \mathcal{K} \right) \leq \varepsilon.$$

Proof. By Itô's formula for $\|\cdot\|_{\mathbb{H}}^2$, for any $t \in [0, T]$,

$$(4.3) \quad \begin{aligned} \|X_t^{(n)}\|_{\mathbb{H}}^2 &= \|x^{(n)}\|_{\mathbb{H}}^2 + \int_0^t \left(2_{\mathbb{V}^*} \langle \mathcal{P}_n \mathcal{A}(s, X_s^{(n)}), X_s^{(n)} \rangle_{\mathbb{V}} + \|\mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) \tilde{\mathcal{P}}_n\|_{L_2(U, \mathbb{H})}^2 \right) ds \\ &\quad + 2 \int_0^t \langle X_s^{(n)}, \mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) dW_s^{(n)} \rangle_{\mathbb{H}}. \end{aligned}$$

Then using Itô's formula for the Lyapunov function $V(r) := \log(1+r)$, by **(A₃)** we deduce that

$$\log(1 + \|X_t^{(n)}\|_{\mathbb{H}}^2)$$

$$\begin{aligned}
&\leq \log(1 + \|x^{(n)}\|_{\mathbb{H}}^2) - \delta \int_0^t \frac{\|X_s^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2} ds \\
&\quad + \int_0^t \frac{g(\|X_s^{(n)}\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s^{(n)})\|_{L_2(U, \mathbb{H})}^2 + C}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2} ds \\
&\quad - \int_0^t \frac{2(1 - \varepsilon_0)\|\mathcal{B}(s, X_s^{(n)}) * X_s^{(n)}\|_U^2}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} ds \\
&\quad - \int_0^t \frac{2\varepsilon_0\|\mathcal{B}(s, X_s^{(n)}) * X_s^{(n)}\|_U^2}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} ds + \mathcal{M}_t \\
&\leq C_T + \log(1 + \|x^{(n)}\|_{\mathbb{H}}^2) - \delta \int_0^t \frac{\|X_s^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2} ds \\
&\quad + \int_0^t \left\{ \frac{(g(\|X_s^{(n)}\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s^{(n)})\|_{L_2(U, \mathbb{H})}^2)(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} \right. \\
&\quad \left. - \frac{2(1 - \varepsilon_0)\|\mathcal{B}(s, X_s^{(n)}) * X_s^{(n)}\|_U^2}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} \right\} ds \\
&\quad + \left\{ \mathcal{M}_t - \int_0^t \frac{2\varepsilon_0\|\mathcal{B}(s, X_s^{(n)}) * X_s^{(n)}\|_U^2}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} ds \right\} \\
(4.4) \quad &=: C_T + \log(1 + \|x^{(n)}\|_{\mathbb{H}}^2) - \delta \int_0^t \frac{\|X_s^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2} ds + \text{(I)} + \text{(II)},
\end{aligned}$$

where $\varepsilon_0 \in (0, \frac{1}{2})$, and we denote

$$\mathcal{M}_t := 2 \int_0^t \frac{\langle X_s^{(n)}, \mathcal{B}(s, X_s^{(n)}) dW_s^{(n)} \rangle_{\mathbb{H}}}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2}.$$

In view of the assumption (2.3) in **(A₅)**, we can choose $\varepsilon_0 = 1 - \frac{\eta}{2}$ such that

$$\text{(I)} \leq C \int_0^t \frac{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2}{(1 + \|X_s^{(n)}\|_{\mathbb{H}}^2)^2} ds \leq C_T.$$

Thus, it follows from (4.4) that

$$(4.5) \quad \log(1 + \|X_t^{(n)}\|_{\mathbb{H}}^2) \leq C_T + \log(1 + \|x\|_{\mathbb{H}}^2) - \delta \int_0^t \frac{\|X_s^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_s^{(n)}\|_{\mathbb{H}}^2} ds + \text{(II)}.$$

On the one hand, by (4.5) we deduce that

$$\log(1 + \|X_t^{(n)}\|_{\mathbb{H}}^2) \leq C_T + \log(1 + \|x\|_{\mathbb{H}}^2) + \left(\mathcal{M}_t - \frac{\varepsilon_0}{2} \langle \mathcal{M} \rangle_t \right),$$

which implies that for any $R > 0$,

$$\begin{aligned}
& \mathbb{P}\left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}}^2 \geq R\right) \\
& \leq \mathbb{P}\left(\sup_{t \in [0, T]} \varepsilon_0 \log \|X_t^{(n)}\|_{\mathbb{H}}^2 \geq \varepsilon_0 \log R\right) \\
& \leq \mathbb{P}\left(\sup_{t \in [0, T]} \left(\varepsilon_0 \mathcal{M}_t - \frac{\varepsilon_0^2}{2} \langle \mathcal{M} \rangle_t\right) \geq \varepsilon_0 (\log R - C_T + \log(1 + \|x\|_{\mathbb{H}}^2))\right) \\
& \leq \mathbb{P}\left(\sup_{t \in [0, T]} \exp\left\{\varepsilon_0 \mathcal{M}_t - \frac{\varepsilon_0^2}{2} \langle \mathcal{M} \rangle_t\right\} \geq \exp\left\{\varepsilon_0 (\log R - C_T + \log(1 + \|x\|_{\mathbb{H}}^2))\right\}\right).
\end{aligned}$$

Therefore, using the maximal supermartingale inequality, we have

$$\begin{aligned}
(4.6) \quad & \mathbb{P}\left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}}^2 \geq R\right) \leq \exp\left\{-\varepsilon_0 (\log R - C_T + \log(1 + \|x\|_{\mathbb{H}}^2))\right\} \\
& \leq \frac{C_T}{R^{\varepsilon_0}},
\end{aligned}$$

which implies that for any $p \geq 2$,

$$(4.7) \quad \mathbb{P}\left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}}^p \geq R\right) \leq \frac{C_T}{R^{\frac{2\varepsilon_0}{p}}}.$$

Similarly, by (4.5) we also have

$$\begin{aligned}
& \mathbb{P}\left(\int_0^T \frac{\|X_t^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_t^{(n)}\|_{\mathbb{H}}^2} dt \geq R\right) \\
& \leq \mathbb{P}\left(\sup_{t \in [0, T]} \left(\varepsilon_0 \mathcal{M}_t - \frac{\varepsilon_0^2}{2} \langle \mathcal{M} \rangle_t\right) \geq \varepsilon_0 (\delta R - C_T + \log(1 + \|x\|_{\mathbb{H}}^2))\right) \\
& \leq \mathbb{P}\left(\sup_{t \in [0, T]} \exp\left\{\varepsilon_0 \mathcal{M}_t - \frac{\varepsilon_0^2}{2} \langle \mathcal{M} \rangle_t\right\} \geq \exp\left\{\varepsilon_0 (\delta R - C_T + \log(1 + \|x\|_{\mathbb{H}}^2))\right\}\right) \\
(4.8) \quad & \leq C_T \exp\left\{-\varepsilon_0 \delta R\right\}.
\end{aligned}$$

In order to get the estimate for $\int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt$, we set the following stopping time

$$\varrho_M^{(n)} := \inf\left\{t \in [0, T] : \|X_t^{(n)}\|_{\mathbb{H}} \geq M\right\} \wedge T, \quad M > 0,$$

with the convention $\inf \emptyset = \infty$. We deduce that

$$\begin{aligned}
& \mathbb{P}\left(\int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq R\right) \\
& \leq \mathbb{P}\left(\int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq R, \varrho_M^{(n)} \geq T\right) + \mathbb{P}(\varrho_M^{(n)} < T)
\end{aligned}$$

$$\begin{aligned}
&\leq \mathbb{P}\left(\int_0^{T \wedge \varrho_M^{(n)}} \frac{\|X_t^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_t^{(n)}\|_{\mathbb{H}}^2} \cdot (1 + \|X_t^{(n)}\|_{\mathbb{H}}^2) dt \geq R\right) + \mathbb{P}(\varrho_M^{(n)} < T) \\
(4.9) \quad &\leq \mathbb{P}\left(C_M \int_0^T \frac{\|X_t^{(n)}\|_{\mathbb{V}}^\alpha}{1 + \|X_t^{(n)}\|_{\mathbb{H}}^2} dt \geq R\right) + \mathbb{P}\left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}} \geq M\right).
\end{aligned}$$

In light of (4.7) and (4.8), we first take $R \rightarrow \infty$ and then $M \rightarrow \infty$ to obtain

$$(4.10) \quad \lim_{R \rightarrow \infty} \mathbb{P}\left(\int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq R\right) = 0.$$

Consequently, (4.2) follows from (4.7) and (4.10). We complete the proof. \square

Based on Lemma 4.1, we have the following bounds.

Lemma 4.2. *Suppose that the assumptions in Theorem 2.1 hold. For any $\varepsilon > 0$ there exists $\mathcal{K} > 0$ such that*

$$(4.11) \quad \sup_{n \in \mathbb{N}} \mathbb{P}\left(\int_0^T \|\mathcal{A}(t, X_t^{(n)})\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} dt \geq \mathcal{K}\right) \leq \varepsilon.$$

Proof. Set the following stopping time

$$(4.12) \quad \tau_M^{(n)} := \inf\left\{t \in [0, T] : \|X_t^{(n)}\|_{\mathbb{H}} + \int_0^t \|X_s^{(n)}\|_{\mathbb{V}}^\alpha ds \geq M\right\} \wedge T, \quad M > 0,$$

with the convention $\inf \emptyset = \infty$.

According to the assumption (2.2), we deduce that

$$\begin{aligned}
&\mathbb{P}\left(\int_0^T \|\mathcal{A}(t, X_t^{(n)})\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} dt \geq R\right) \\
&\leq \mathbb{P}\left(\int_0^T \|\mathcal{A}(t, X_t^{(n)})\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} dt \geq R, \tau_M^{(n)} \geq T\right) + \mathbb{P}(\tau_M^{(n)} < T) \\
&\leq \mathbb{P}\left(C \int_0^{T \wedge \tau_M^{(n)}} (1 + \|X_t^{(n)}\|_{\mathbb{V}}^\alpha)(1 + \|X_t^{(n)}\|_{\mathbb{H}}^\beta) dt \geq R\right) + \mathbb{P}(\tau_M^{(n)} < T) \\
&\leq \frac{C_M}{R} + \mathbb{P}\left(\sup_{t \in [0, T]} \|X_t^{(n)}\|_{\mathbb{H}} + \int_0^T \|X_s^{(n)}\|_{\mathbb{V}}^\alpha ds \geq M\right).
\end{aligned}$$

By Lemma 4.1, let $R \rightarrow \infty$ and then $M \rightarrow \infty$, we complete the assertion. \square

4.2. Tightness of approximating solutions. Set

$$\begin{aligned}
(4.13) \quad \mathcal{Z}_T^1 &:= \mathbb{C}_T(\mathbb{V}^*) \cap L^\alpha([0, T]; \mathbb{H}) \cap L_w^\alpha([0, T]; \mathbb{V}), \\
\mathcal{Z}_T^2 &:= L_w^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*),
\end{aligned}$$

where $L_w^\alpha([0, T]; \mathbb{V})$, $L_w^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)$ denote the spaces $L^\alpha([0, T]; \mathbb{V})$, $L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)$ endowed with the weak topology, respectively. In this subsection, we will show that $\{X^{(n)}\}_{n \in \mathbb{N}}$, $\{\mathcal{A}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$ are tight in \mathcal{Z}_T^1 , \mathcal{Z}_T^2 , respectively.

Remark 4.1. Here the intersection space \mathcal{Z}_T^1 takes the following intersection topology denoted by $\tau_{\mathcal{Z}_T^1}$: the class of open sets of \mathcal{Z}_T^1 are generated by the sets of the form $\mathcal{O}_1 \cap \mathcal{O}_2 \cap \mathcal{O}_3$, where \mathcal{O}_1 , \mathcal{O}_2 and \mathcal{O}_3 are the open sets in $\mathbb{C}_T(\mathbb{V}^*)$, $L^\alpha([0, T]; \mathbb{H})$ and $L_w^\alpha([0, T]; \mathbb{V})$, respectively. Let $\mathcal{B}(\tau_{\mathcal{Z}_T^1})$ be the corresponding Borel σ -algebra.

In the following, we formulate the compactness criterion for the space \mathcal{Z}_T^1 .

Lemma 4.3. Let \mathcal{K} be a subset of \mathcal{Z}_T^1 such that the following conditions hold

(i)

$$\sup_{X \in \mathcal{K}} \int_0^T \|X_t\|_{\mathbb{V}}^\alpha dt < \infty,$$

(ii)

$$\limsup_{\delta \rightarrow 0} \sup_{X \in \mathcal{K}} \sup_{s, t \in [0, T], |t-s| \leq \delta} \|X_t - X_s\|_{\mathbb{V}^*} = 0.$$

Then \mathcal{K} is relatively compact in \mathcal{Z}_T^1 .

Proof. The proof is inspired by Lemma 3.1 in [13]. Without loss of generality, we suppose that \mathcal{K} is $\tau_{\mathcal{Z}_T^1}$ -closed. It is known that the weak topology in $L_w^\alpha([0, T]; \mathbb{V})$ restricted in \mathcal{K} is metrizable. Therefore, the compactness of a subset of \mathcal{Z}_T^1 is equivalent to the sequential compactness.

Now, let $\{X^{(n)}\}_{n \in \mathbb{N}}$ denote a sequence in \mathcal{K} . It is sufficient to prove that there exists an element $X \in \mathcal{K}$ such that along a subsequence still denoted by $\{X^{(n)}\}_{n \in \mathbb{N}}$, we have

$$(4.14) \quad X^{(n)} \rightarrow X \text{ in } \mathcal{Z}_T^1 \text{ as } n \rightarrow \infty.$$

First, due to the Banach-Alaoglu theorem and the condition (i), it follows that

$$(4.15) \quad \mathcal{K} \text{ is compact in } L_w^\alpha([0, T]; \mathbb{V}).$$

On the other hand, in view of the condition (ii) we can deduce that there exists an dt -null set \mathcal{N} and a subsequence still denoted by $\{X^{(n)}\}_{n \in \mathbb{N}}$ such that for any $t \in [0, T] \setminus \mathcal{N}$,

$$\{\|X_t^{(n)}\|_{\mathbb{V}}\}_{n \in \mathbb{N}} \text{ is bounded.}$$

By the assumption of Theorem 2.1, the embedding $\mathbb{V} \subset \mathbb{H}$ is compact, we infer that $\mathbb{V} \subset \mathbb{V}^*$ is compact as well. Thus, the sequence $\{X_t^{(n)}\}_{n \in \mathbb{N}}$ contains a subsequence that is convergent in \mathbb{V}^* .

Denote $\{t_k\}_{k \in \mathbb{N}} \subset ([0, T] \setminus \mathcal{N}) \cap \mathbb{Q}$. Applying the diagonal method we can choose a subsequence still denoted by $\{X^{(n)}\}_{n \in \mathbb{N}}$ such that

$$(4.16) \quad \{X_{t_k}^{(n)}\}_{n \in \mathbb{N}} \text{ is convergent in } \mathbb{V}^* \text{ for all } k \in \mathbb{N}.$$

Next, we prove that the sequence $\{X^{(n)}\}_{n \in \mathbb{N}}$ is a Cauchy net in $\mathbb{C}_T(\mathbb{V}^*)$. For any $\varepsilon > 0$, by the condition (ii) there exists $\delta > 0$ such that

$$(4.17) \quad \sup_{X \in \mathcal{K}} \sup_{s, t \in [0, T], |t-s| \leq \delta} \|X_t - X_s\|_{\mathbb{V}^*} < \frac{\varepsilon}{3}.$$

Fix $t \in [0, T]$. We can find a $t_k \in ([0, T] \setminus \mathcal{N}) \cap \mathbb{Q}$ such that $|t_k - t| \leq \delta$. Combining (4.16)-(4.17), for large enough $n, m \in \mathbb{N}$ we deduce that

$$\|X_t^{(n)} - X_t^{(m)}\|_{\mathbb{V}^*} \leq \|X_t^{(n)} - X_{t_k}^{(n)}\|_{\mathbb{V}^*} + \|X_{t_k}^{(n)} - X_{t_k}^{(m)}\|_{\mathbb{V}^*} + \|X_{t_k}^{(m)} - X_t^{(m)}\|_{\mathbb{V}^*} \leq \varepsilon.$$

Since $t \in [0, T]$ is arbitrary, we can obtain that

$$\sup_{t \in [0, T]} \|X_t^{(n)} - X_t^{(m)}\|_{\mathbb{V}^*} \leq \varepsilon,$$

which yields that

$$(4.18) \quad \{X^{(n)}\}_{n \in \mathbb{N}} \text{ is a Cauchy net in } \mathbb{C}_T(\mathbb{V}^*).$$

Collecting (4.15) and (4.18), there exists a subsequence still denoted by $\{X^{(n)}\}_{n \in \mathbb{N}}$ and $X \in \mathbb{C}_T(\mathbb{V}^*) \cap L^\alpha([0, T]; \mathbb{V})$ such that

$$(4.19) \quad X^{(n)} \rightarrow X \text{ in } \mathbb{C}_T(\mathbb{V}^*) \cap L_w^\alpha([0, T]; \mathbb{V}) \text{ as } n \rightarrow \infty.$$

Hence, once we can prove

$$(4.20) \quad X^{(n)} \rightarrow X \text{ in } L^\alpha([0, T]; \mathbb{H}) \text{ as } n \rightarrow \infty,$$

then (4.14) follows. Since the embedding $\mathbb{V} \subset \mathbb{H}$ is compact, by the Lions lemma [49] for any $\varepsilon > 0$ there exists a constant $C_\varepsilon > 0$ such that for almost all $t \in [0, T]$,

$$\|X_t^{(n)} - X_t\|_{\mathbb{H}}^\alpha \leq \varepsilon \|X_t^{(n)} - X_t\|_{\mathbb{V}}^\alpha + C_\varepsilon \|X_t^{(n)} - X_t\|_{\mathbb{V}^*}^\alpha, \quad n \in \mathbb{N},$$

which also yields that

$$(4.21) \quad \|X^{(n)} - X\|_{L^\alpha([0, T]; \mathbb{H})}^\alpha \leq \varepsilon \|X^{(n)} - X\|_{L^\alpha([0, T]; \mathbb{V})}^\alpha + C_\varepsilon \|X^{(n)} - X\|_{L^\alpha([0, T]; \mathbb{V}^*)}^\alpha, \quad n \in \mathbb{N}.$$

Taking the upper limit as $n \rightarrow \infty$ in (4.21) and using the following bounds

$$\|X^{(n)} - X\|_{L^\alpha([0, T]; \mathbb{V})}^\alpha \leq C(\|X^{(n)}\|_{L^\alpha([0, T]; \mathbb{V})}^\alpha + \|X\|_{L^\alpha([0, T]; \mathbb{V})}^\alpha) \leq C,$$

we can conclude that

$$\lim_{n \rightarrow \infty} \|X^{(n)} - X\|_{L^\alpha([0, T]; \mathbb{H})}^\alpha = 0.$$

We complete the proof. \square

We recall the Aldous condition in the space \mathbb{V}^* .

Definition 4.1. *A sequence $\{X^{(n)}\}_{n \in \mathbb{N}}$ is said to satisfy the Aldous condition in \mathbb{V}^* iff for any $\varepsilon, \eta > 0$, there exists $\delta > 0$ such that for every stopping time sequence $(\zeta_n)_{n \in \mathbb{N}}$ with $\zeta_n \leq T$ one has*

$$\sup_{n \in \mathbb{N}} \sup_{0 \leq \Delta \leq \delta} \mathbb{P}(\|X_{\zeta_n + \Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*} \geq \eta) \leq \varepsilon.$$

The following lemma presents a tightness criterion for the laws of sequence $\{X^{(n)}\}_{n \in \mathbb{N}}$ on \mathcal{Z}_T^1 .

Lemma 4.4. *Let $\{X^{(n)}\}_{n \in \mathbb{N}}$ be a sequence of continuous $\{\mathcal{F}_t\}$ -adapted \mathbb{V}^* -valued processes such that*

(i)

$$\lim_{R \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{P} \left(\int_0^T \|X_t^{(n)}\|_{\mathbb{V}}^\alpha dt > R \right) = 0,$$

(ii) $\{X^{(n)}\}_{n \in \mathbb{N}}$ satisfies the Aldous condition in \mathbb{V}^* .

Let μ_n be the law of $X^{(n)}$ on the Borel σ -algebra $\mathcal{B}(\tau_{\mathcal{Z}_T^1})$. Then for every $\varepsilon > 0$ there exists a compact subset \mathcal{K}_ε of \mathcal{Z}_T^1 such that

$$\sup_{n \in \mathbb{N}} \mu_n(\mathcal{K}_\varepsilon) \geq 1 - \varepsilon.$$

Proof. In view of (i), for any $\varepsilon > 0$ there exists $R > 0$ such that

$$\sup_{n \in \mathbb{N}} \mathbb{P} \left(\int_0^T \|X_t^{(n)}\|_{\mathbb{V}} dt > R \right) \leq \frac{\varepsilon}{3}.$$

We denote

$$\mathcal{K} := \left\{ X^{(n)} \in \mathcal{Z}_T^1 : \int_0^T \|X_t^{(n)}\|_{\mathbb{V}} dt \leq R \right\}.$$

By Lemmas 3.6 and 3.8 in [13], in view of (ii) there exists a subset $\mathcal{A}_{\frac{\varepsilon}{3}} \subset \mathbb{C}_T(\mathbb{V}^*)$ such that

$$\mu_n(\mathcal{A}_{\frac{\varepsilon}{3}}^c) \leq \frac{\varepsilon}{3}$$

and

$$\lim_{\delta \rightarrow 0} \sup_{X \in \mathcal{A}_{\frac{\varepsilon}{3}}} \sup_{s, t \in [0, T], |t-s| \leq \delta} \|X_t - X_s\|_{\mathbb{V}^*} = 0.$$

Finally, we denote by \mathcal{K}_ε the closure of the set $\mathcal{K} \cap \mathcal{A}_{\frac{\varepsilon}{3}}$ in \mathcal{Z}_T^1 . Due to the compactness criterion presented in Lemma 4.3, we conclude that \mathcal{K}_ε is a compact set in \mathcal{Z}_T^1 . The proof is complete. \square

Based on Lemmas 4.1 and 4.4, in order to show the tightness of $\{X^{(n)}\}_{n \in \mathbb{N}}$ in \mathcal{Z}_T^1 it is sufficient to prove $\{X^{(n)}\}_{n \in \mathbb{N}}$ satisfies the Aldous condition in \mathbb{V}^* , which is given as follows.

Lemma 4.5. $\{X^{(n)}\}_{n \in \mathbb{N}}$ satisfies the Aldous condition in \mathbb{V}^* in the sense of Definition 4.1.

Proof. Recall Lemma 4.1, we know

$$(4.22) \quad \lim_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{P}(\tau_M^{(n)} < T) = 0,$$

where the stopping time $\tau_M^{(n)}$ is defined by (4.12). In addition, we have

$$\begin{aligned} & \mathbb{P} \left(\|X_{\zeta_n + \Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*} \geq \varepsilon \right) \\ & \leq \mathbb{P} \left(\|X_{\zeta_n + \Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*} \geq \varepsilon, \tau_M^{(n)} \geq T \right) + \mathbb{P}(\tau_M^{(n)} < T) \end{aligned}$$

$$(4.23) \quad \leq \frac{1}{\varepsilon^{\frac{\alpha}{\alpha-1}}} \mathbb{E} \|X_{\zeta_n+\Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} + \mathbb{P}(\tau_M^{(n)} < T).$$

Now we estimate the first term on the right hand side of (4.23). By (4.1) and applying B-D-G's inequality, it follows that

$$(4.24) \quad \begin{aligned} & \mathbb{E} \|X_{\zeta_n+\Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} \\ & \leq C \mathbb{E} \left\{ \int_{\zeta_n \wedge \tau_M^{(n)}}^{(\zeta_n+\Delta) \wedge \tau_M^{(n)}} \|\mathcal{P}_n \mathcal{A}(s, X_s^{(n)})\|_{\mathbb{V}^*} ds \right\}^{\frac{\alpha}{\alpha-1}} \\ & \quad + C \mathbb{E} \left\{ \int_{\zeta_n \wedge \tau_M^{(n)}}^{(\zeta_n+\Delta) \wedge \tau_M^{(n)}} \|\mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) \tilde{\mathcal{P}}_n\|_{L_2(U, \mathbb{H})}^2 ds \right\}^{\frac{\alpha}{2(\alpha-1)}} \\ & =: \text{(I)} + \text{(II)}. \end{aligned}$$

For (I), by **(A₄)** and Hölder's inequality, we have

$$(4.25) \quad \begin{aligned} \text{(I)} & \leq C |\Delta|^{\frac{1}{\alpha-1}} \cdot \mathbb{E} \left\{ \int_{\zeta_n \wedge \tau_M^{(n)}}^{(\zeta_n+\Delta) \wedge \tau_M^{(n)}} \|\mathcal{P}_n \mathcal{A}(s, X_s^{(n)})\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} ds \right\} \\ & \leq C |\Delta|^{\frac{1}{\alpha-1}} \cdot \mathbb{E} \left\{ \int_0^{T \wedge \tau_M^{(n)}} (1 + \|X_s^{(n)}\|_{\mathbb{V}}^\alpha) (1 + \|X_s^{(n)}\|_{\mathbb{H}}^\beta) ds \right\} \\ & \leq C_{M,T} |\Delta|^{\frac{1}{\alpha-1}}. \end{aligned}$$

For (II), we can get

$$(4.26) \quad \begin{aligned} \text{(II)} & \leq C \mathbb{E} \left\{ \int_{\zeta_n \wedge \tau_M^{(n)}}^{(\zeta_n+\Delta) \wedge \tau_M^{(n)}} (1 + \|X_s^{(n)}\|_{\mathbb{H}}^\beta) ds \right\}^{\frac{\alpha}{2(\alpha-1)}} \\ & \leq C_M |\Delta|^{\frac{\alpha}{2(\alpha-1)}}. \end{aligned}$$

Combining (4.24)-(4.26) gives

$$(4.27) \quad \limsup_{\Delta \rightarrow 0} \sup_{n \in \mathbb{N}} \mathbb{E} \|X_{\zeta_n+\Delta}^{(n)} - X_{\zeta_n}^{(n)}\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} = 0.$$

Finally, taking into account (4.22), (4.23) and (4.27) and letting $\Delta \rightarrow 0$ then $M \rightarrow \infty$ in (4.23), we conclude that the Aldous condition holds. The proof is completed. \square

We now give the tightness of $\{X^{(n)}\}_{n \in \mathbb{N}}$ in \mathcal{Z}_T^1 .

Lemma 4.6. $\{X^{(n)}\}_{n \in \mathbb{N}}$ is tight in \mathcal{Z}_T^1 .

Proof. Combining Lemmas 4.1, 4.4 and 4.5, the assertion follows. \square

The following lemma shows the tightness of $\{\mathcal{A}^{(n)}(\cdot) := \mathcal{A}(\cdot, X^{(n)})\}_{n \in \mathbb{N}}$ in \mathcal{Z}_T^2 .

Lemma 4.7. $\{\mathcal{A}^{(n)}(\cdot)\}_{n \in \mathbb{N}}$ is tight in \mathcal{Z}_T^2 .

Proof. The tightness of $\{\mathcal{A}^{(n)}(\cdot)\}_{n \in \mathbb{N}}$ in \mathcal{Z}_T^2 follows directly from the Banach-Alaoglu theorem and the estimate (4.11). \square

4.3. Passage to the limit. In this part, we aim to prove Theorems 2.1, where the Jakubowski's beautiful generalization of the Skorokhod's representation theorem, in the form presented by Brzeźniak and Ondreját, for nonmetric spaces (see Lemma 5.1) and the theory of pseudo-monotone operators play a crucial role.

In order to apply Jakubowski's version of the Skorokhod theorem, we first present the following lemma, whose proof is inspired by Theorem B.5 in [48].

Lemma 4.8. \mathcal{Z}_T^1 is a standard Borel space in the sense of Definition 5.2.

Proof. Let $\mathcal{Y} := L^2([0, T]; \mathbb{V}^*)$ with Borel σ -algebra $\mathcal{B}(\mathcal{Y})$. In order to show the assertion, according to Theorem 2.3 of Chapter V in [58] and the fact that \mathcal{Y} is a standard Borel space, it is sufficient to show the following three claims:

- (i) the embedding $\mathcal{Z}_T^1 \subset \mathcal{Y}$ is continuous;
- (ii) $\mathcal{Z}_T^1 \in \mathcal{B}(\mathcal{Y})$;
- (iii) $\mathcal{B}(\tau_{\mathcal{Z}_T^1}) = \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1$.

First, the claim (i) follows directly from $\mathbb{C}_T(\mathbb{V}^*) \subset \mathcal{Y}$ continuously. Next, we focus on proving claims (ii) and (iii).

Proof of (ii): By claim (i) we have

$$(4.28) \quad \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1 \subset \mathcal{B}(\tau_{\mathcal{Z}_T^1}).$$

Fix $N \in \mathbb{N}$. Let us denote

$$(L_N^\alpha([0, T]; \mathbb{V}))_w := \left\{ x \in L^\alpha([0, T]; \mathbb{V}) \mid \|x\|_{L^\alpha([0, T]; \mathbb{V})} \leq N \right\},$$

which is endowed with the weak topology on $L^\alpha([0, T]; \mathbb{V})$. Then $(L_N^\alpha([0, T]; \mathbb{V}))_w$ is a compact and metrizable, hence complete and separable, space with metric d_1 . Fix $N \in \mathbb{N}$. Set

$$\mathcal{Z}_T^{(N)} := (L_N^\alpha([0, T]; \mathbb{V}))_w \cap \mathbb{C}_T(\mathbb{V}^*) \cap L^\alpha([0, T]; \mathbb{H}),$$

which is a closed subset of \mathcal{Z}_T^1 . The metrics on $\mathbb{C}_T(\mathbb{V}^*)$ and $L^\alpha([0, T]; \mathbb{H})$ are denoted by d_2 and d_3 , respectively. Now, let $\mathcal{Z}_T^{(N)}$ be endowed with the metric $d := \max\{d_1, d_2, d_3\}$. Since the intersection of finite separable metric space (with the maximal metric) is a separable metric space, it follows that $\mathcal{Z}_T^{(N)}$ is a separable metric space. We intend to show that $\mathcal{Z}_T^{(N)}$ is complete. To this end, it is sufficient to show that for a sequence $\{x_k\}_{k \in \mathbb{N}}$ converging to $x^{(i)}$ in d_i , $i = 1, 2, 3$, we have

$$(4.29) \quad x^{(1)} = x^{(2)} = x^{(3)}.$$

Since $x_k \rightarrow x^{(1)}$ and $x_k \rightarrow x^{(3)}$ both in $L^\alpha([0, T]; \mathbb{H})$, it follows that

$$x^{(1)} = x^{(3)}.$$

In addition, $x_k \rightarrow x^{(2)}$ and $x_k \rightarrow x^{(3)}$ both in $L^\alpha([0, T]; \mathbb{V}^*)$, we can deduce that

$$x^{(2)} = x^{(3)}.$$

Hence, (4.29) follows, then $\mathcal{Z}_T^{(N)}$ is a complete separable metric space. Furthermore, the following embeddings are continuous

$$(4.30) \quad (\mathcal{Z}_T^{(N)}, d) \subset (\mathcal{Z}_T^1, \tau_{\mathcal{Z}_T^1}) \subset \mathcal{Y}.$$

Therefore, in view of Theorem 2.4 of Chapter V in [58] we can obtain

$$(4.31) \quad \mathcal{B}((\mathcal{Z}_T^{(N)}, d)) \subset \mathcal{B}(\mathcal{Y}).$$

Moreover, by (4.30) we have

$$(4.32) \quad \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^{(N)} \subset \mathcal{B}((\mathcal{Z}_T^{(N)}, d)).$$

Combining (4.31)-(4.32) we obtain

$$(4.33) \quad \mathcal{B}((\mathcal{Z}_T^{(N)}, d)) = \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^{(N)},$$

which yields that

$$\mathcal{Z}_T^1 = \bigcup_{N=1}^{\infty} \mathcal{Z}_T^{(N)} \in \mathcal{B}(\mathcal{Y}).$$

The proof of claim (ii) is completed.

Proof of (iii): Since $\mathcal{Z}_T^{(N)}$ is the closed subset of \mathcal{Z}_T^1 , it is clear that $\mathcal{Z}_T^{(N)} \in \mathcal{B}(\tau_{\mathcal{Z}_T^1})$. Then we can obtain

$$\mathcal{B}(\tau_{\mathcal{Z}_T^1}) \cap \mathcal{Z}_T^{(N)} = \{B \in \mathcal{B}(\tau_{\mathcal{Z}_T^1}) \mid B \subset \mathcal{Z}_T^{(N)}\}$$

and

$$\mathcal{B}(\tau_{\mathcal{Z}_T^1}) = \bigcup_{N=1}^{\infty} \{B \in \mathcal{B}(\tau_{\mathcal{Z}_T^1}) \mid B \subset \mathcal{Z}_T^{(N)}\}.$$

Moreover, the embedding $(\mathcal{Z}_T^{(N)}, d) \subset (\mathcal{Z}_T^{(N)}, \tau_{\mathcal{Z}_T^1} \cap \mathcal{Z}_T^{(N)})$ is continuous, which implies

$$(4.34) \quad \mathcal{B}(\tau_{\mathcal{Z}_T^1} \cap \mathcal{Z}_T^{(N)}) \subset \mathcal{B}((\mathcal{Z}_T^{(N)}, d)).$$

Suppose that A is an $\tau_{\mathcal{Z}_T^1}$ -closed subset of \mathcal{Z}_T^1 . Then $A \cap \mathcal{Z}_T^{(N)}$ is $\tau_{\mathcal{Z}_T^1}$ -closed. Then by (4.33) and (4.34) we have

$$(4.35) \quad \begin{aligned} A \cap \mathcal{Z}_T^{(N)} &\in \mathcal{B}(\tau_{\mathcal{Z}_T^1} \cap \mathcal{Z}_T^{(N)}) \\ &\subset \mathcal{B}((\mathcal{Z}_T^{(N)}, d)) \\ &= \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^{(N)}. \end{aligned}$$

Note that $\mathcal{Z}_T^{(N)} \in \mathcal{B}(\mathcal{Y})$, in view of claim (ii) we can get

$$\mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^{(N)} = \{B \in \mathcal{B}(\mathcal{Y}) \mid B \subset \mathcal{Z}_T^{(N)}\} \subset \{B \in \mathcal{B}(\mathcal{Y}) \mid B \subset \mathcal{Z}_T^1\} = \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1.$$

Then by (4.35) we have

$$A = \bigcup_{N=1}^{\infty} A \cap \mathcal{Z}_T^{(N)} \in \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1.$$

Thus, $\mathcal{B}(\tau_{\mathcal{Z}_T^1}) \subset \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1$, which together with (4.28) implies $\mathcal{B}(\tau_{\mathcal{Z}_T^1}) = \mathcal{B}(\mathcal{Y}) \cap \mathcal{Z}_T^1$. The proof is completed. \square

Now, we need to check that the spaces $\mathcal{Z}_T^1, \mathcal{Z}_T^2$ satisfy the conditions in Lemma 5.1. In fact, since $\mathbb{C}_T(\mathbb{V}^*)$ and $L^\alpha([0, T]; \mathbb{H})$ are separable Banach spaces, it is easy to see the conditions in Lemma 5.1 hold. Furthermore, for the space $L_w^\alpha([0, T]; \mathbb{V})$, it suffices to put

$$f_m(u) := \int_0^T \mathbb{V}^* \langle v_m(t), u(t) \rangle_{\mathbb{V}} dt \in \mathbb{R}, \quad u \in L_w^\alpha([0, T]; \mathbb{V}), \quad m \in \mathbb{N},$$

where $\{v_m\}_{m \geq 1}$ is a dense subset of $L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)$. Since $\{v_m\}_{m \geq 1}$ is dense in $L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)$, it separates points of $L^\alpha([0, T]; \mathbb{V})$. Moreover, since $(\mathcal{Z}_T^1, \mathcal{B}(\tau_{\mathcal{Z}_T^1}))$ is a standard Borel space (i.e. Lemma 4.8), the σ -algebra generated by the sequence of the above continuous functions, which separate the points in \mathcal{Z}_T^1 , is exactly $\mathcal{B}(\tau_{\mathcal{Z}_T^1})$ by Theorem 5.1. Thus all the conditions in Lemma 5.1 are satisfied for \mathcal{Z}_T^1 . From the same reason, we deduce that the conditions in Lemma 5.1 also hold for \mathcal{Z}_T^2 .

Set

$$\Xi_T := \mathcal{Z}_T^1 \times \mathcal{Z}_T^2 \times \mathbb{C}_T(U_1),$$

where U_1 is a Hilbert space such that the embedding $U \subset U_1$ is Hilbert-Schmidt. Since $\{X^{(n)}\}_{n \in \mathbb{N}}, \{\mathcal{A}^{(n)}(\cdot)\}_{n \in \mathbb{N}}$ are tight in $\mathcal{Z}_T^1, \mathcal{Z}_T^2$, respectively, it is clear that $\{(X^{(n)}, \mathcal{A}^{(n)}(\cdot), W)\}_{n \in \mathbb{N}}$ is also tight in Ξ_T . By the Jakubowski's version of the Skorokhod theorem, there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, and on this space, Ξ_T -valued random variables $(\tilde{X}^{(n)}, \tilde{\mathcal{A}}^{(n)}(\cdot), \tilde{W}^{(n)})$, $(\tilde{X}, \tilde{\mathcal{A}}(\cdot), \tilde{W})$ (here choosing a subsequence if necessary) such that

(i) the law of $(\tilde{X}^{(n)}, \tilde{\mathcal{A}}^{(n)}(\cdot), \tilde{W}^{(n)})$ under $\tilde{\mathbb{P}}$ is equivalent to the law of $(X^{(n)}, \mathcal{A}^{(n)}(\cdot), W)$ under \mathbb{P} ;

(ii) the following convergence hold

$$(4.36) \quad \tilde{X}^{(n)} \rightarrow \tilde{X} \text{ in } \mathcal{Z}_T^1, \tilde{\mathbb{P}}\text{-a.s., as } n \rightarrow \infty,$$

$$(4.37) \quad \tilde{\mathcal{A}}^{(n)}(\cdot) \rightharpoonup \tilde{\mathcal{A}}(\cdot) \text{ in } L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*), \tilde{\mathbb{P}}\text{-a.s., as } n \rightarrow \infty,$$

where “ \rightharpoonup ” stands for the weak convergence.

(iii) $\tilde{W}^{(n)} = \tilde{W}$ for any $n \in \mathbb{N}$, $\tilde{\mathbb{P}}$ -a.s..

Let $(\tilde{\mathcal{F}}_t)_{t \in [0, T]}$ be the filtration satisfying the usual conditions and generated by

$$\bigcup_{n \in \mathbb{N}} \left\{ \tilde{X}_s^{(n)}, \tilde{X}_s, \tilde{W}_s : s \in [0, t] \right\}.$$

Note that by the claims (i) and (iii) above, we know

$$\begin{aligned} & \mathbb{P}(W_t - W_s \in \cdot | \mathcal{F}_s) = \mathbb{P}(W_t - W_s \in \cdot) \\ \Rightarrow & \tilde{\mathbb{P}}(\tilde{W}_t^{(n)} - \tilde{W}_s^{(n)} \in \cdot | \tilde{\mathcal{F}}_s) = \tilde{\mathbb{P}}(\tilde{W}_t^{(n)} - \tilde{W}_s^{(n)} \in \cdot) \\ \Rightarrow & \tilde{\mathbb{P}}(\tilde{W}_t - \tilde{W}_s \in \cdot | \tilde{\mathcal{F}}_s) = \tilde{\mathbb{P}}(\tilde{W}_t - \tilde{W}_s \in \cdot). \end{aligned}$$

It follows that \tilde{W} is an $(\tilde{\mathcal{F}}_t)$ -cylindrical Wiener process on U . From (4.1) and the claims (i) and (iii), the following identity holds $\tilde{\mathbb{P}}$ -a.s.

$$(4.38) \quad \tilde{X}_t^{(n)} = x^{(n)} + \int_0^t \mathcal{P}_n \mathcal{A}(s, \tilde{X}_s^{(n)}) ds + \int_0^t \mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n d\tilde{W}_s, \quad t \in [0, T].$$

By the claim (i), the convergence (4.37) and the fact that the law of $\mathcal{A}(\cdot, \tilde{X}^{(n)})$ under $\tilde{\mathbb{P}}$ is equivalent to the law of $\mathcal{A}^{(n)}(\cdot)$ under \mathbb{P} , we can infer that (at least) along a subsequence still denoted by $\{n\}$ we have

$$(4.39) \quad \mathcal{A}(\cdot, \tilde{X}^{(n)}) \rightharpoonup \tilde{\mathcal{A}}(\cdot) \text{ in } L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*), \quad \tilde{\mathbb{P}}\text{-a.s.}, \text{ as } n \rightarrow \infty.$$

In addition, by Lemmas 4.1, we can obtain same bounds hold for $\tilde{X}^{(n)}$. More precisely, we have for any $\varepsilon > 0$ there exists $\mathcal{K} > 0$ such that for any $p \geq 2$,

$$(4.40) \quad \sup_{n \in \mathbb{N}} \tilde{\mathbb{P}} \left(\sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}}^p + \int_0^T \|\tilde{X}_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq \mathcal{K} \right) \leq \varepsilon.$$

Using the lower semicontinuity of norms $\|\cdot\|_{\mathbb{H}}$, $\|\cdot\|_{\mathbb{V}}$ in \mathbb{V}^* , by the convergence (4.36) and Fatou's lemma we can deduce for any $\varepsilon > 0$ we can choose \mathcal{K} such that

$$(4.41) \quad \begin{aligned} & \tilde{\mathbb{P}} \left(\sup_{t \in [0, T]} \|\tilde{X}_t\|_{\mathbb{H}}^p + \int_0^T \|\tilde{X}_t\|_{\mathbb{V}}^\alpha dt \geq \mathcal{K} \right) \\ & \leq \tilde{\mathbb{P}} \left(\sup_{t \in [0, T]} \liminf_{n \rightarrow \infty} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}}^p + \int_0^T \liminf_{n \rightarrow \infty} \|\tilde{X}_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq \mathcal{K} \right) \\ & \leq \tilde{\mathbb{P}} \left(\liminf_{n \rightarrow \infty} \left\{ \sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}}^p + \int_0^T \|\tilde{X}_t^{(n)}\|_{\mathbb{V}}^\alpha dt \right\} \geq \mathcal{K} \right) \\ & \leq \sup_{n \in \mathbb{N}} \tilde{\mathbb{P}} \left(\sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}}^p + \int_0^T \|\tilde{X}_t^{(n)}\|_{\mathbb{V}}^\alpha dt \geq \mathcal{K} \right) \\ & \leq \varepsilon. \end{aligned}$$

In the sequel, we investigate the convergence of (4.38), as $n \rightarrow \infty$, where we can choose a subsequence if necessary.

In the following lemma, we present the convergence of the stochastic integral in (4.38).

Lemma 4.9. $\int_0^\cdot \mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n d\tilde{W}_s \rightarrow \int_0^\cdot \mathcal{B}(s, \tilde{X}_s) d\tilde{W}_s$ in $L^\infty([0, T]; \mathbb{H})$ in probability as $n \rightarrow \infty$.

Proof. In light of Lemma 5.2 in [34] or Lemma 4.3 in [4] for the version of infinite dimensions, it suffices to show that for any $t \in [0, T]$,

$$(4.42) \quad \int_0^t \|\mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n - \mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 ds \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

Note that by (4.36) we know

$$(4.43) \quad \int_0^t \|\tilde{X}_t^{(n)} - \tilde{X}_t\|_{\mathbb{H}}^\kappa dt \rightarrow 0, \quad \tilde{\mathbb{P}}\text{-a.s.}, \quad \forall \kappa \in [1, \alpha).$$

Then, there exists a $\tilde{\mathbb{P}} \otimes dt$ -null set \mathcal{N} such that for any $(t, \omega) \in (\Omega \times [0, T]) \setminus \mathcal{N}$, along a subsequence still denoted by $\{n\}$ we have

$$(4.44) \quad \|\tilde{X}_t^{(n)}(\omega) - \tilde{X}_t(\omega)\|_{\mathbb{H}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Moreover, we know

$$\begin{aligned} & \|\mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n - \mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 \\ & \leq C \|\mathcal{P}_n(\mathcal{B}(s, \tilde{X}_s^{(n)}) - \mathcal{B}(s, \tilde{X}_s)) \tilde{\mathcal{P}}_n\|_{L_2(U, \mathbb{H})}^2 + C \|\mathcal{P}_n \mathcal{B}(s, \tilde{X}_s) \tilde{\mathcal{P}}_n - \mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 \\ & \leq C \|\mathcal{B}(s, \tilde{X}_s^{(n)}) - \mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 + C \|\mathcal{P}_n \mathcal{B}(s, \tilde{X}_s) \tilde{\mathcal{P}}_n - \mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 \\ & =: I_s^{(n)} + II_s^{(n)}. \end{aligned}$$

In what follows, we only focus on the convergence of $I^{(n)}$, since the convergence of $II^{(n)}$ follows directly from the property of orthonormal projections.

We intend to prove

$$\int_0^t I_s^{(n)} ds \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

In light of the continuity condition (4.42) and the convergence (4.44), it suffices to show

$$(4.45) \quad \int_0^t \|\mathcal{B}(s, \tilde{X}_s^{(n)})\|_{L_2(U, \mathbb{H})}^2 ds \rightarrow \int_0^t \|\mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 ds \text{ in probability as } n \rightarrow \infty.$$

Let $\chi_R \in C_c^\infty(\mathbb{R})$ be a cut-off function with

$$\chi_R(r) = \begin{cases} 1, & |r| \leq R \\ 0, & |r| > 2R. \end{cases}$$

Set

$$\begin{aligned} \Psi_R(t, w) &:= \int_0^t \|\mathcal{B}(s, w_s)\|_{L_2(U, \mathbb{H})}^2 \chi_R(\|w_s\|_{\mathbb{H}}) ds, \\ \Psi(t, w) &:= \int_0^t \|\mathcal{B}(s, w_s)\|_{L_2(U, \mathbb{H})}^2 ds. \end{aligned}$$

On the one hand, by (2.5), (4.44) and the continuity of χ_R we have as $n \rightarrow \infty$

$$\left| \|\mathcal{B}(s, \tilde{X}_s^{(n)})\|_{L_2(U, \mathbb{H})}^2 \chi_R(\|\tilde{X}_s^{(n)}\|_{\mathbb{H}}) - \|\mathcal{B}(s, \tilde{X}_s)\|_{L_2(U, \mathbb{H})}^2 \chi_R(\|\tilde{X}_s\|_{\mathbb{H}}) \right| \rightarrow 0, \quad \tilde{\mathbb{P}} \otimes dt\text{-a.e.},$$

which combining with the dominated convergence theorem yields

$$(4.46) \quad \Psi_R(t, \tilde{X}^{(n)}) \rightarrow \Psi_R(t, \tilde{X}), \quad \tilde{\mathbb{P}}\text{-a.s.}, \text{ as } n \rightarrow \infty.$$

On the other hand, by the definition of χ_R we obtain that for any $\varepsilon > 0$,

$$\begin{aligned} & \tilde{\mathbb{P}}\left(|\Psi(t, \tilde{X}^{(n)}) - \Psi_R(t, \tilde{X}^{(n)})| > \varepsilon\right) \\ &= \tilde{\mathbb{P}}\left(|\Psi(t, \tilde{X}^{(n)}) - \Psi_R(t, \tilde{X}^{(n)})| > \varepsilon, \sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}} \leq R\right) \\ & \quad + \tilde{\mathbb{P}}\left(|\Psi(t, \tilde{X}^{(n)}) - \Psi_R(t, \tilde{X}^{(n)})| > \varepsilon, \sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}} > R\right) \end{aligned}$$

$$\leq \sup_{n \in \mathbb{N}} \tilde{\mathbb{P}} \left(\sup_{t \in [0, T]} \|\tilde{X}_t^{(n)}\|_{\mathbb{H}} > R \right).$$

In view of (4.40), letting $n \rightarrow \infty$ and $R \rightarrow \infty$ we derive

$$(4.47) \quad |\Psi(t, \tilde{X}^{(n)}) - \Psi_R(t, \tilde{X}^{(n)})| \rightarrow 0 \text{ in probability.}$$

Applying similar argument, we also obtain

$$(4.48) \quad |\Psi(t, \tilde{X}) - \Psi_R(t, \tilde{X})| \rightarrow 0 \text{ in probability.}$$

Collecting (4.46)-(4.48), we conclude that (4.45) follows.

We complete the proof. \square

From Lemma 4.9, along a subsequence still denoted by $\{n\}$ we have that as $n \rightarrow \infty$,

$$(4.49) \quad \sup_{t \in [0, T]} \left\| \int_0^t \mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n d\tilde{W}_s - \int_0^t \mathcal{B}(s, \tilde{X}_s) d\tilde{W}_s \right\|_{\mathbb{H}} \rightarrow 0, \tilde{\mathbb{P}}\text{-a.s.}$$

Thus applying the convergence (4.36)-(4.39) and (4.49), for any $v \in \cup_{n \geq 1} \mathbb{H}_n(\subset \mathbb{V})$, $\varphi \in L^\infty([0, T] \times \Omega; \mathbb{R})$ we obtain

$$\begin{aligned} & \int_0^T \mathbb{V}^* \langle \tilde{X}_t, \varphi_t v \rangle_{\mathbb{V}} dt \\ &= \lim_{n \rightarrow \infty} \int_0^T \mathbb{V}^* \langle \tilde{X}_t^{(n)}, \varphi_t v \rangle_{\mathbb{V}} dt \\ &= \lim_{n \rightarrow \infty} \left(\mathbb{V}^* \langle x^{(n)}, v \rangle_{\mathbb{V}} \int_0^T \varphi_t dt + \int_0^T \int_0^t \mathbb{V}^* \langle \mathcal{P}_n \mathcal{A}(s, \tilde{X}_s^{(n)}), \varphi_t v \rangle_{\mathbb{V}} ds dt \right. \\ & \quad \left. + \int_0^T \langle \int_0^t \mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n d\tilde{W}_s, \varphi_t v \rangle_{\mathbb{H}} dt \right) \\ &= \lim_{n \rightarrow \infty} \left(\mathbb{V}^* \langle x^{(n)}, v \rangle_{\mathbb{V}} \int_0^T \varphi_t dt + \int_0^T \mathbb{V}^* \langle \mathcal{A}(s, \tilde{X}_s^{(n)}), \int_s^T \varphi_t dt \cdot v \rangle_{\mathbb{V}} ds \right. \\ & \quad \left. + \int_0^T \langle \int_0^t \mathcal{P}_n \mathcal{B}(s, \tilde{X}_s^{(n)}) \tilde{\mathcal{P}}_n d\tilde{W}_s, \varphi_t v \rangle_{\mathbb{H}} dt \right) \\ &= \int_0^T \mathbb{V}^* \langle x + \int_0^t \tilde{\mathcal{A}}(s) ds + \int_0^t \mathcal{B}(s, \tilde{X}_s) d\tilde{W}_s, \varphi_t v \rangle_{\mathbb{V}} dt, \tilde{\mathbb{P}}\text{-a.s.} \end{aligned}$$

Thus we define

$$(4.50) \quad \bar{X}_t := x + \int_0^t \tilde{\mathcal{A}}(s) ds + \int_0^t \mathcal{B}(s, \tilde{X}_s) d\tilde{W}_s, \quad t \in [0, T].$$

It is clear that

$$(4.51) \quad \tilde{X} = \bar{X}, \tilde{\mathbb{P}} \otimes dt\text{-a.e.}$$

In what follows, without loss of generality, we replace $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, $(\tilde{X}^{(n)}, \tilde{W}^{(n)})$, (\tilde{X}, \tilde{W}) , by $(\Omega, \mathcal{F}, \mathbb{P})$, $(X^{(n)}, W^{(n)})$, (X, W) , respectively, for sake of simplicity.

In order to characterize the limit of $\mathcal{A}^{(n)}(\cdot)$, we recall the following lemma concerning the pseudo-monotonicity (cf. [51]).

Lemma 4.10. *Assume that the embedding $\mathbb{V} \subset \mathbb{H}$ is compact, (\mathbf{A}_1) and (\mathbf{A}_2) hold. Then $\mathcal{A}(t, \cdot)$ is pseudo-monotone from \mathbb{V} to \mathbb{V}^* for a.e. $t \in [0, T]$.*

The next important step is to show $\tilde{\mathcal{A}}(\cdot) = \mathcal{A}(\cdot, X)$. To this end, we introduce the following lemma.

Lemma 4.11. *Assume that the embedding $\mathbb{V} \subset \mathbb{H}$ is compact, (\mathbf{A}_1) - (\mathbf{A}_4) hold. If*

- (i) $X^{(n)} \rightharpoonup X$ in $L^\alpha([0, T]; \mathbb{V})$, \mathbb{P} -a.s.,
- (ii) $\mathcal{A}(\cdot, X^{(n)}) \rightharpoonup \tilde{\mathcal{A}}$ in $L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)$, \mathbb{P} -a.s.,
- (iii) $\liminf_{n \rightarrow \infty} \int_0^T \mathbb{V}^* \langle \mathcal{A}(t, X_t^{(n)}), X_t^{(n)} \rangle_{\mathbb{V}} dt \geq \int_0^T \mathbb{V}^* \langle \tilde{\mathcal{A}}(t), X_t \rangle_{\mathbb{V}} dt$, \mathbb{P} -a.s.,

then $\tilde{\mathcal{A}}(\cdot) = \mathcal{A}(\cdot, X)$, $\mathbb{P} \otimes dt$ -a.e..

Proof. First, due to (\mathbf{A}_3) and (\mathbf{A}_4) , we can obtain that there exists $\delta_0 > 0$ such that

$$\begin{aligned}
& \mathbb{V}^* \langle \mathcal{A}(t, X_t^{(n)}), X_t^{(n)} - X_t \rangle_{\mathbb{V}} \\
& \leq -\delta \|X_t^{(n)}\|_{\mathbb{V}}^\alpha + C + g(\|X_t^{(n)}\|_{\mathbb{H}}^2) + \|\mathcal{A}(t, X_t^{(n)})\|_{\mathbb{V}^*} \|X_t\|_{\mathbb{V}} \\
& \leq -\delta \|X_t^{(n)}\|_{\mathbb{V}}^\alpha + C + g(\|X_t^{(n)}\|_{\mathbb{H}}^2) \\
& \quad + C(1 + \|X_t^{(n)}\|_{\mathbb{V}}^\alpha)^{\frac{\alpha-1}{\alpha}} (1 + \|X_t^{(n)}\|_{\mathbb{H}}^\beta)^{\frac{\alpha-1}{\alpha}} \|X_t\|_{\mathbb{V}} \\
(4.52) \quad & \leq -\delta_0 \|X_t^{(n)}\|_{\mathbb{V}}^\alpha + C + g(\|X_t^{(n)}\|_{\mathbb{H}}^2) + C(1 + \|X_t^{(n)}\|_{\mathbb{H}}^\beta)^{\alpha-1} \|X_t\|_{\mathbb{V}}^\alpha.
\end{aligned}$$

For convenience, we denote

$$\begin{aligned}
\phi^{(n)}(t, \omega) &:= \mathbb{V}^* \langle \mathcal{A}(t, X_t^{(n)}(\omega)), X_t^{(n)}(\omega) - X_t(\omega) \rangle_{\mathbb{V}}, \\
F^{(n)}(t, \omega) &:= C + g(\|X_t^{(n)}(\omega)\|_{\mathbb{H}}^2) + C(1 + \|X_t^{(n)}(\omega)\|_{\mathbb{H}}^\beta)^{\alpha-1} \|X_t(\omega)\|_{\mathbb{V}}^\alpha.
\end{aligned}$$

Then (4.52) reads as

$$(4.53) \quad \phi^{(n)}(t, \omega) \leq -\delta_0 \|X_t^{(n)}(\omega)\|_{\mathbb{V}}^\alpha + F^{(n)}(t, \omega).$$

The proof of this lemma is divided into the following four steps.

Step 1: In this step, we prove that for a.e. (t, ω) ,

$$(4.54) \quad \limsup_{n \rightarrow \infty} \phi^{(n)}(t, \omega) \leq 0.$$

According to (4.43), there exists a $\mathbb{P} \otimes dt$ -null set \mathcal{N} such that for any $(t, \omega) \in (\Omega \times [0, T]) \setminus \mathcal{N}$, along a subsequence still denoted by $\{n\}$ we have

$$(4.55) \quad \|X_t^{(n)}(\omega) - X_t(\omega)\|_{\mathbb{H}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

From now on, we fix $(t, \omega) \in (\Omega \times [0, T]) \setminus \mathcal{N}$ and suppose that

$$\limsup_{n \rightarrow \infty} \phi^{(n)}(t, \omega) > 0.$$

Thus we can take a subsequence $\{n_k\}_{k \in \mathbb{N}}$ such that

$$(4.56) \quad \lim_{k \rightarrow \infty} \phi^{(n_k)}(t, \omega) > 0.$$

It follows from (4.53) and (4.55) that

$$(4.57) \quad \left\{ \|X_t^{(n_k)}(\omega)\|_{\mathbb{V}}^\alpha \right\}_{k \in \mathbb{N}} \text{ is bounded.}$$

Therefore, there exists an element $z \in \mathbb{V}$ such that

$$X_t^{(n_k)}(\omega) \rightharpoonup z \text{ in } \mathbb{V} \text{ as } k \rightarrow \infty.$$

From the convergence (4.55), it is clear that $z = X_t(\omega)$ and

$$X_t^{(n_k)}(\omega) \rightarrow X_t(\omega) \text{ in } \mathbb{V} \text{ as } k \rightarrow \infty.$$

Using the fact that $\mathcal{A}(t, \cdot)$ is pseudo-monotone (cf. Lemma 4.10), we deduce that

$$(4.58) \quad \limsup_{k \rightarrow \infty} \phi^{(n_k)}(t, \omega) \leq 0,$$

which contradicts to (4.56). Hence, (4.54) holds.

Step 2: In this step, we prove that along a subsequence $\{n_k\}_{k \in \mathbb{N}}$ for a.e. (t, ω) ,

$$\lim_{k \rightarrow \infty} \phi^{(n_k)}(t, \omega) = 0.$$

First, by the conditions (ii)-(iii) in this lemma, the control (4.41), (4.55) and Fatou's lemma, we have for a.s. ω ,

$$\begin{aligned} 0 &\leq \liminf_{n \rightarrow \infty} \int_0^T \phi^{(n)}(t, \omega) dt \leq \limsup_{n \rightarrow \infty} \int_0^T \phi^{(n)}(t, \omega) dt \\ &\leq \int_0^T \limsup_{n \rightarrow \infty} \phi^{(n)}(t, \omega) dt \leq 0. \end{aligned}$$

Hence

$$(4.59) \quad \lim_{n \rightarrow \infty} \int_0^T \phi^{(n)}(t, \omega) dt = 0.$$

Then combining (4.54) and (4.59) and applying the dominated convergence theorem, it follows that

$$(4.60) \quad \lim_{n \rightarrow \infty} \int_0^T \phi_+^{(n)}(t, \omega) dt = 0,$$

where $\phi_+^{(n)}(t, \omega) := \max\{\phi^{(n)}(t, \omega), 0\}$.

By (4.59)-(4.60) and the fact that $|\phi^{(n)}| = 2\phi_+^{(n)} - \phi^{(n)}$, we have

$$\lim_{n \rightarrow \infty} \int_0^T |\phi^{(n)}(t, \omega)| dt = 0,$$

which implies that the claim follows.

Step 3: In this step, we prove that $\tilde{\mathcal{A}}(\cdot) = \mathcal{A}(\cdot, X)$, $\mathbb{P} \otimes dt$ -a.e..

Combining the claim of **Step 2**, (4.53) and (4.55), we can get (4.57) holds. Thus,

$$X_t^{(n_k)}(\omega) \rightharpoonup X_t(\omega) \text{ in } \mathbb{V} \text{ as } k \rightarrow \infty,$$

which combining with the pseudo-monotonicity of $\mathcal{A}(t, \cdot)$ implies that

$$\mathcal{A}(t, X_t^{(n_k)}(\omega)) \rightharpoonup \mathcal{A}(t, X_t(\omega)) \text{ in } \mathbb{V}^* \text{ as } k \rightarrow \infty,$$

Consequently, by the condition (ii) in the lemma and the uniqueness of the limit, it is clear that $\tilde{\mathcal{A}}(\cdot) = \mathcal{A}(\cdot, X)$, $\mathbb{P} \otimes dt$ -a.e.. \square

Now we have all the ingredients to prove the existence of weak solutions to (1.5).

Proof of existence of weak solutions: We aim to show that (X, W) obtained above is a weak solution to Eq. (1.5) in the sense of Definition 2.2. Combining (4.36), (4.39), (4.50), (4.51) and Lemma 4.11, we need to show that the condition (iii) in Lemma 4.11 holds and $X \in \mathbb{C}_T(\mathbb{H})$, \mathbb{P} -a.s.. The proof is separated into the following two steps.

Step 1: In this step, we prove that the condition (iii) in Lemma 4.11 holds. First, in view of (4.39) it implies that

$$\|\tilde{\mathcal{A}}(\cdot)\|_{L^{\frac{\alpha}{\alpha-1}}([0, T]; \mathbb{V}^*)} < \infty, \mathbb{P}\text{-a.s..}$$

Then we denote a stopping time

$$\tau_M^A := \inf \left\{ t \in [0, T] : \int_0^t \|\tilde{\mathcal{A}}(s)\|_{\mathbb{V}^*}^{\frac{\alpha}{\alpha-1}} ds \geq M \right\} \wedge T, \quad M > 0.$$

It is easy to see that

$$\lim_{M \rightarrow \infty} \tau_M^A = T, \mathbb{P}\text{-a.s..}$$

On the other hand, the estimate (4.41) yields

$$(4.61) \quad \sup_{t \in [0, T]} \|X_t\|_{\mathbb{H}} + \int_0^T \|X_t\|_{\mathbb{V}}^{\alpha} dt < \infty, \mathbb{P}\text{-a.s..}$$

By (4.36), we know that

$$(4.62) \quad X \in \mathbb{C}_T(\mathbb{V}^*), \mathbb{P}\text{-a.s..}$$

Since $\mathbb{H} \subset \mathbb{V}^*$ is dense, in view of (4.61) and (4.62) we deduce that X is weakly continuous in \mathbb{H} , so that $\|X\|_{\mathbb{H}}$ is lower semicontinuous. Therefore, we set

$$\tau_M^X := \inf \left\{ t \in [0, T] : \|X_t\|_{\mathbb{H}} + \int_0^t \|X_s\|_{\mathbb{V}}^{\alpha} ds \geq M \right\} \wedge T, \quad M > 0,$$

which is a stopping time and

$$\lim_{M \rightarrow \infty} \tau_M^X = T, \mathbb{P}\text{-a.s..}$$

Let us denote

$$\hat{\tau}_M := \tau_M^A \wedge \tau_M^X.$$

Recall Eq. (4.50), it is easy to see that

$$X_{t \wedge \hat{\tau}_M} = x + \int_0^t \mathbf{1}_{\{s \leq \hat{\tau}_M\}} \tilde{\mathcal{A}}(s) ds + \int_0^t \mathbf{1}_{\{s \leq \hat{\tau}_M\}} \mathcal{B}(s, X_s) dW_s, \quad t \in [0, T].$$

Let us denote

$$Y(t) := \mathbf{1}_{\{t \leq \hat{\tau}_M\}} \tilde{\mathcal{A}}(t), \quad Z(t) := \mathbf{1}_{\{t \leq \hat{\tau}_M\}} \mathcal{B}(t, X_t).$$

Then it is clear that

$$X_{\cdot \wedge \hat{\tau}_M} \in L^\alpha([0, T] \times \Omega; \mathbb{V}), \quad Y(\cdot) \in L^{\frac{\alpha}{\alpha-1}}([0, T] \times \Omega; \mathbb{V}^*), \quad Z(\cdot) \in L^2([0, T] \times \Omega; \mathbb{H})$$

and

$$\mathbb{E} \|X_{t \wedge \hat{\tau}_M}\|_{\mathbb{H}}^2 \leq C_M < \infty.$$

Therefore, according to Theorem 4.2.5 in [52], we can apply Itô's formula for $\|X_{t \wedge \hat{\tau}_M}\|_{\mathbb{H}}^2$ and deduce that

$$(4.63) \quad \begin{aligned} \|X_t\|_{\mathbb{H}}^2 &= \|x\|_{\mathbb{H}}^2 + \int_0^t \left(2_{\mathbb{V}^*} \langle \tilde{\mathcal{A}}(s), X_s \rangle_{\mathbb{V}} + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2 \right) ds \\ &\quad + 2 \int_0^t \langle \mathcal{B}(s, X_s) dW_s, X_s \rangle_{\mathbb{H}} \text{ on } \{t \leq \hat{\tau}_M\}. \end{aligned}$$

Note that $\lim_{M \rightarrow \infty} \hat{\tau}_M = T$, \mathbb{P} -a.s., which implies that the equality (4.63) holds for all $t \in [0, T]$.

Next, applying Itô's formula for $\|X_t^{(n)}\|_{\mathbb{H}}^2$, we have

$$(4.64) \quad \begin{aligned} \|X_t^{(n)}\|_{\mathbb{H}}^2 &= \|x^{(n)}\|_{\mathbb{H}}^2 + \int_0^t \left(2_{\mathbb{V}^*} \langle \mathcal{P}_n \mathcal{A}(s, X_s^{(n)}), X_s^{(n)} \rangle_{\mathbb{V}} + \|\mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) \tilde{\mathcal{P}}_n\|_{L_2(U, \mathbb{H})}^2 \right) ds \\ &\quad + 2 \int_0^t \langle \mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) dW_s^{(n)}, X_s^{(n)} \rangle_{\mathbb{H}}. \end{aligned}$$

The following lemma concerns the convergence of the martingale term in (4.64).

Lemma 4.12. *Along a subsequence still denoted by $\{n\}$, we have that as $n \rightarrow \infty$,*

$$(4.65) \quad \int_0^\cdot \langle \mathcal{P}_n \mathcal{B}(s, X_s^{(n)}) \tilde{\mathcal{P}}_n dW_s, X_s^{(n)} \rangle_{\mathbb{H}} \rightarrow \int_0^\cdot \langle \mathcal{B}(s, X_s) dW_s, X_s \rangle_{\mathbb{H}} \text{ in } L^\infty([0, T]; \mathbb{R}), \mathbb{P}\text{-a.s..}$$

Proof. The proof is exactly similar to that of Lemma 4.9, we omit the details. \square

Note that by the lower semicontinuity of norm $\|\cdot\|_{\mathbb{H}}$ in \mathbb{V}^* and the convergence (4.36), we have

$$(4.66) \quad \|X_t\|_{\mathbb{H}}^2 \leq \liminf_{n \rightarrow \infty} \|X_t^{(n)}\|_{\mathbb{H}}^2, \quad \mathbb{P}\text{-a.s..}$$

Finally, combining (4.36), (4.42), (4.63)-(4.66), it follows that

$$\liminf_{n \rightarrow \infty} \int_0^T \mathbb{V}^* \langle \mathcal{A}(t, X_t^{(n)}), X_t^{(n)} \rangle_{\mathbb{V}} dt \geq \int_0^T \mathbb{V}^* \langle \tilde{\mathcal{A}}(t), X_t \rangle_{\mathbb{V}} dt, \quad \mathbb{P}\text{-a.s..}$$

namely, the condition (iii) in Lemma 4.11 holds.

Step 2: In this step, we prove $X \in \mathbb{C}_T(\mathbb{H})$, \mathbb{P} -a.s.. In Step 1, we have show that X is weakly continuous in \mathbb{H} . Thus it suffices to prove $t \mapsto \|X_t\|_{\mathbb{H}}$ is continuous on $[0, T]$. Based on Step 1, we deduce that

$$(4.67) \quad \begin{aligned} \|X_t\|_{\mathbb{H}}^2 &= \|x\|_{\mathbb{H}}^2 + \int_0^t \left(2_{\mathbb{V}^*} \langle \mathcal{A}(s, X_s), X_s \rangle_{\mathbb{V}} + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2 \right) ds \\ &\quad + 2 \int_0^t \langle \mathcal{B}(s, X_s) dW_s, X_s \rangle_{\mathbb{H}}. \end{aligned}$$

Since the right-hand side of (4.67) is continuous on $[0, T]$, so must be its left-hand side.

We complete the proof of the existence of weak solutions. \square

4.4. Proof of existence and uniqueness. We first prove the pathwise uniqueness of solutions to (1.5).

Proof of pathwise uniqueness: Let X, Y be two solutions of (1.5) with same initial value $x \in \mathbb{H}$. Then, the difference process $Z := X - Y$ solves the following equation

$$Z_t = \int_0^t (\mathcal{A}(s, X_s) - \mathcal{A}(s, Y_s)) ds + \int_0^t (\mathcal{B}(s, X_s) - \mathcal{B}(s, Y_s)) dW_s, \quad t \in [0, T].$$

Set

$$\begin{aligned} \tau_M^X &:= \inf \left\{ t \in [0, T] : \|X_t\|_{\mathbb{H}} + \int_0^t \|X_s\|_{\mathbb{V}}^\alpha ds \geq M \right\} \wedge T, \quad M > 0, \\ \tau_M^Y &:= \inf \left\{ t \in [0, T] : \|Y_t\|_{\mathbb{H}} + \int_0^t \|Y_s\|_{\mathbb{V}}^\alpha ds \geq M \right\} \wedge T, \quad M > 0. \end{aligned}$$

It is clear that τ_M^X, τ_M^Y are stopping times and $\lim_{M \rightarrow \infty} \tau_M^X = T, \lim_{M \rightarrow \infty} \tau_M^Y = T$, \mathbb{P} -a.s..

Let $\tau_M := \tau_M^X \wedge \tau_M^Y$. Applying Itô's formula to $\|Z_t\|_{\mathbb{H}}^2$, which follows from the same argument as (4.63), and by **(A₂)** we derive

$$(4.68) \quad \begin{aligned} \|Z_t\|_{\mathbb{H}}^2 &= \int_0^t \left(2_{\mathbb{V}^*} \langle \mathcal{A}(s, X_s) - \mathcal{A}(s, Y_s), Z_s \rangle_{\mathbb{V}} + \|\mathcal{B}(s, X_s) - \mathcal{B}(s, Y_s)\|_{L_2(U, \mathbb{H})}^2 \right) ds \\ &\quad + 2 \int_0^t \langle (\mathcal{B}(s, X_s) - \mathcal{B}(s, Y_s)) dW_s, Z_s \rangle_{\mathbb{H}} \\ &\leq C \int_0^t (1 + \rho(X_s) + \eta(Y_s)) \|Z_s\|_{\mathbb{H}}^2 ds \\ &\quad + 2 \int_0^t \langle (\mathcal{B}(s, X_s) - \mathcal{B}(s, Y_s)) dW_s, Z_s \rangle_{\mathbb{H}}. \end{aligned}$$

Then, we have

$$\mathbb{E} \|Z_{t \wedge \tau_M}\|_{\mathbb{H}}^2 \leq C \mathbb{E} \int_0^{t \wedge \tau_M} (1 + \rho(X_s) + \eta(Y_s)) \|Z_s\|_{\mathbb{H}}^2 ds.$$

Applying stochastic Gronwall's lemma (cf. [32, Lemma 5.3]), it leads to

$$\mathbb{E} \|Z_{t \wedge \tau_M}\|_{\mathbb{H}}^2 \leq 0, \quad t \in [0, T],$$

which implies that

$$(4.69) \quad \mathbb{E}\|Z_t\|_{\mathbb{H}}^2 \leq \liminf_{M \rightarrow \infty} \mathbb{E}\|Z_{t \wedge \tau_M}\|_{\mathbb{H}}^2 \leq 0, \quad t \in [0, T].$$

Therefore, the pathwise uniqueness follows from (4.69) and the pathwise continuity in \mathbb{H} . \square

Now we have all the ingredients to verify Theorem 2.1.

Proof of Theorem 2.1: Combining Subsection 4.3 and the above pathwise uniqueness, the existence and uniqueness of strong solutions to (1.5) is a direct consequence of the infinite-dimensional version of Yamada-Watanabe theorem.

Finally, the Markov property follows from the standard arguments as in [52, Proposition 4.3.3] based on the above existence and uniqueness results. We complete the proof. \square

4.5. **Proof of Feller property.** Set the stopping time

$$\tau_M^n := \inf \left\{ t \in [0, T] : \|X_t(x_n)\|_{\mathbb{H}} + \|X_t(x)\|_{\mathbb{H}} + \int_0^t \|X_s(x_n)\|_{\mathbb{V}}^\alpha ds + \int_0^t \|X_s(x)\|_{\mathbb{V}}^\alpha ds \geq M \right\} \wedge T, \quad M > 0,$$

with the convention $\inf \emptyset = \infty$. According to the estimate (2.6) and the convergence of x_n , we can deduce that

$$(4.70) \quad \lim_{M \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{P}(\tau_M^n < T) = 0.$$

In the following, we first prove the continuous dependence on initial data in probability, and then we derive the Feller property of the transition semigroup.

Proof of Theorem 2.2: In view of the proof of (4.68), by B-D-G's inequality we have

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \in [0, T \wedge \tau_M^n]} \|X_t(x_n) - X_t(x)\|_{\mathbb{H}}^2 \right] \\ & \leq \|x_n - x\|_{\mathbb{H}}^2 + C \mathbb{E} \int_0^{T \wedge \tau_M^n} (1 + \rho(X_s(x_n)) + \eta(X_s(x))) \|X_s(x_n) - X_s(x)\|_{\mathbb{H}}^2 ds \\ & \quad + C \mathbb{E} \left(\int_0^{T \wedge \tau_M^n} \|X_s(x_n) - X_s(x)\|_{\mathbb{H}}^2 \|\mathcal{B}(s, X_s(x_n)) - \mathcal{B}(s, X_s(x))\|_{L_2(U, \mathbb{H})}^2 ds \right)^{\frac{1}{2}} \\ & \leq \|x_n - x\|_{\mathbb{H}}^2 + C \mathbb{E} \int_0^{T \wedge \tau_M^n} (1 + \rho(X_s(x_n)) + \eta(X_s(x))) \|X_s(x_n) - X_s(x)\|_{\mathbb{H}}^2 ds \\ & \quad + \frac{1}{2} \mathbb{E} \left[\sup_{t \in [0, T \wedge \tau_M^n]} \|X_t(x_n) - X_t(x)\|_{\mathbb{H}}^2 \right] \\ & \quad + C \mathbb{E} \int_0^{T \wedge \tau_M^n} \|\mathcal{B}(s, X_s(x_n)) - \mathcal{B}(s, X_s(x))\|_{L_2(U, \mathbb{H})}^2 ds \\ & \leq \|x_n - x\|_{\mathbb{H}}^2 + C \mathbb{E} \int_0^{T \wedge \tau_M^n} (1 + \rho(X_s(x_n)) + \eta(X_s(x))) \|X_s(x_n) - X_s(x)\|_{\mathbb{H}}^2 ds \end{aligned}$$

$$+\frac{1}{2}\mathbb{E}\left[\sup_{t\in[0,T\wedge\tau_M^n]}\|X_t(x_n)-X_t(x)\|_{\mathbb{H}}^2\right],$$

where we used the assumption (2.7) in the last step.

Applying stochastic Gronwall's lemma, we have

$$\mathbb{E}\left[\sup_{t\in[0,T\wedge\tau_M^n]}\|X_t(x_n)-X_t(x)\|_{\mathbb{H}}^2\right]\leq C_M\|x_n-x\|_{\mathbb{H}}^2.$$

Consequently, for any $\varepsilon > 0$,

$$\begin{aligned} & \mathbb{P}\left(\sup_{t\in[0,T]}\|X_t(x_n)-X_t(x)\|_{\mathbb{H}}>\varepsilon\right) \\ & \leq \mathbb{P}\left(\sup_{t\in[0,T]}\|X_t(x_n)-X_t(x)\|_{\mathbb{H}}>\varepsilon, T\leq\tau_M^n\right)+\mathbb{P}(\tau_M^n<T) \\ & \leq \frac{1}{\varepsilon^2}\mathbb{E}\left[\sup_{t\in[0,T\wedge\tau_M^n]}\|X_t(x_n)-X_t(x)\|_{\mathbb{H}}^2\right]+\mathbb{P}(\tau_M^n<T) \\ & \leq \frac{C_M}{\varepsilon^2}\|x_n-x\|_{\mathbb{H}}^2+\mathbb{P}(\tau_M^n<T). \end{aligned}$$

Taking (4.70) into account, we conclude that (2.8) holds.

Now we prove the Feller property of the transition semigroup. For any $t \geq 0$ and $\varphi \in C_b(\mathbb{H})$, we show $\mathcal{T}_t\varphi \in C_b(\mathbb{H})$, i.e.,

$$(4.71) \quad \mathbb{E}\varphi(X_t(x_n)) \rightarrow \mathbb{E}\varphi(X_t(x)), \text{ if } x_n \rightarrow x \text{ in } \mathbb{H}.$$

Without loss of generality, we assume $\varphi \in \text{Lip}_b(\mathbb{H})$ (the space of all bounded and Lipschitz continuous functions), since $\text{Lip}_b(\mathbb{H}) \subset C_b(\mathbb{H})$ densely. Note that by (2.8) it is easy to see that for any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(|\varphi(X_t(x_n)) - \varphi(X_t(x))| > \varepsilon\right) \leq \lim_{n \rightarrow \infty} \mathbb{P}\left(C_{\text{Lip}}\|X_t(x_n) - X_t(x)\|_{\mathbb{H}} > \varepsilon\right) = 0,$$

where C_{Lip} is the Lipschitz constant of φ . Thus, $\varphi(X_t(x_n)) \rightarrow \varphi(X_t(x))$ in probability, as $n \rightarrow \infty$. Since $\varphi(X_t(x_n)) \leq \|\varphi\|_{\infty}$ for all $n \in \mathbb{N}$, (4.71) follows from the Lebesgue dominated convergence theorem. We complete the proof of Theorem 2.2. \square

4.6. Proof of finite time extinction. In this part, we intend to prove Theorem 2.3. We first present the following lemma.

Lemma 4.13. *Suppose that the assumptions in Theorem 2.3 hold. Then there is a constant $C > 0$ such that for any $t \geq 0$,*

$$(4.72) \quad \mathbb{E}(1 + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} \leq e^{Ct}(1 + \|x\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}}.$$

Proof. Recall that for any $t \geq 0$,

$$\begin{aligned} \|X_t\|_{\mathbb{H}}^2 &= \|x\|_{\mathbb{H}}^2 + \int_0^t \left(2_{\mathbb{V}^*}\langle \mathcal{A}(s, X_s), X_s \rangle_{\mathbb{V}} + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2\right) ds \\ &\quad + 2 \int_0^t \langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}. \end{aligned}$$

Applying Itô's formula for the Lyapunov function $V(r) := (1+r)^{1-\frac{\alpha}{2}}$, by (\mathbf{A}_3^*) for any $t \geq 0$,

$$\begin{aligned}
& (1 + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} \\
& \leq (1 + \|x\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} - \delta(1 - \frac{\alpha}{2}) \int_0^t \frac{\|X_s\|_{\mathbb{V}}^\alpha}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds \\
& \quad + (1 - \frac{\alpha}{2}) \int_0^t \frac{g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds \\
& \quad - \frac{\alpha}{2}(1 - \frac{\alpha}{2}) \int_0^t \frac{2\|\mathcal{B}(s, X_s)^* X_s\|_U^2}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds + \mathcal{M}_t^0 \\
& = (1 + \|x\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} - \delta(1 - \frac{\alpha}{2}) \int_0^t \frac{\|X_s\|_{\mathbb{V}}^\alpha}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds + \mathcal{M}_t^0 \\
(4.73) \quad & + (1 - \frac{\alpha}{2}) \int_0^t \frac{(g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2)(1 + \|X_s\|_{\mathbb{H}}^2) - \alpha\|\mathcal{B}(s, X_s)^* X_s\|_U^2}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds,
\end{aligned}$$

where

$$\mathcal{M}_t^0 := 2(1 - \frac{\alpha}{2}) \int_0^t \frac{\langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}}{(1 + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}}.$$

Taking into account the assumption (2.3), by (4.73) we can get that

$$(1 + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} \leq (1 + \|x\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} + C \int_0^t (1 + \|X_s\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} ds + \mathcal{M}_t^0.$$

By the standard localization argument and the estimate (2.6), Gronwall' lemma implies

$$\mathbb{E}(1 + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} \leq e^{Ct} (1 + \|x\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}}.$$

We complete the proof. \square

The following lemma plays an important role for proving (2.13) and (2.14).

Lemma 4.14. *Suppose that the assumptions in Theorem 2.3 hold. Then there is $c^* > 0$ such that for any $0 \leq r < t$,*

$$\begin{aligned}
(4.74) \quad & \|X_t\|_{\mathbb{H}}^{2-\alpha} + \delta(c^*)^\alpha (1 - \frac{\alpha}{2}) \int_r^t \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \\
& \leq \|X_r\|_{\mathbb{H}}^{2-\alpha} + 2(1 - \frac{\alpha}{2}) \int_r^t \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} \frac{\langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}}{\|X_s\|_{\mathbb{H}}^\alpha}.
\end{aligned}$$

Proof. For any $\varepsilon > 0$, using Itô's formula for the Lyapunov function $V^\varepsilon(r) := (\varepsilon + r)^{1-\frac{\alpha}{2}}$, then we have by (\mathbf{A}_3^*) that for any $0 \leq r < t$,

$$\begin{aligned}
& (\varepsilon + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} \\
& \leq (\varepsilon + \|X_r\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} - \delta(1 - \frac{\alpha}{2}) \int_r^t \frac{\|X_s\|_{\mathbb{V}}^\alpha}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds
\end{aligned}$$

$$\begin{aligned}
& + (1 - \frac{\alpha}{2}) \int_r^t \frac{g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds \\
& - \frac{\alpha}{2} (1 - \frac{\alpha}{2}) \int_r^t \frac{2\|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2 \|X_s\|_{\mathbb{H}}^2}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds + \mathcal{M}_t \\
& = (\varepsilon + \|X_r\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} - \delta(1 - \frac{\alpha}{2}) \int_r^t \frac{\|X_s\|_{\mathbb{V}}^{\alpha}}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds \\
& + (1 - \frac{\alpha}{2}) \int_r^t \frac{(g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2)(\varepsilon + \|X_s\|_{\mathbb{H}}^2)}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds \\
(4.75) \quad & - (1 - \frac{\alpha}{2}) \int_r^t \frac{\alpha \|\mathcal{B}(s, X_s)^* X_s\|_{\mathbb{U}}^2}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds + \mathcal{M}_{r,t}^{\varepsilon},
\end{aligned}$$

where

$$\mathcal{M}_{r,t}^{\varepsilon} := 2(1 - \frac{\alpha}{2}) \int_r^t \frac{\langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}}.$$

Recall the fact that there is a constant $c^* > 0$ such that $\|u\|_{\mathbb{V}} \geq c^* \|u\|_{\mathbb{H}}$. Then, it follows that

$$\begin{aligned}
& (\varepsilon + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} + \delta(c^*)^{\alpha} (1 - \frac{\alpha}{2}) \int_r^t \frac{\|X_s\|_{\mathbb{H}}^{\alpha}}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} ds \\
& \leq (\varepsilon + \|X_r\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} + \mathcal{M}_{r,t}^{\varepsilon} \\
& + (1 - \frac{\alpha}{2}) \int_r^t \frac{(g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2)(\varepsilon + \|X_s\|_{\mathbb{H}}^2)}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds \\
& - (1 - \frac{\alpha}{2}) \int_r^t \frac{\alpha \|\mathcal{B}(s, X_s)^* X_s\|_{\mathbb{U}}^2}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} ds.
\end{aligned}$$

Then, due to (\mathbf{A}_5^*) we deduce that

$$\begin{aligned}
& (\varepsilon + \|X_t\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} + \delta(c^*)^{\alpha} (1 - \frac{\alpha}{2}) \int_r^t \frac{\|X_s\|_{\mathbb{H}}^{\alpha}}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \\
& \leq (\varepsilon + \|X_r\|_{\mathbb{H}}^2)^{1-\frac{\alpha}{2}} + 2(1 - \frac{\alpha}{2}) \int_r^t \frac{\langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}}} \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \\
& + (1 - \frac{\alpha}{2}) \int_r^t \frac{(g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2)(\varepsilon + \|X_s\|_{\mathbb{H}}^2)}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \\
(4.76) \quad & - (1 - \frac{\alpha}{2}) \int_r^t \frac{\alpha \|\mathcal{B}(s, X_s)^* X_s\|_{\mathbb{U}}^2}{(\varepsilon + \|X_s\|_{\mathbb{H}}^2)^{\frac{\alpha}{2}+1}} \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds.
\end{aligned}$$

Taking $\varepsilon \rightarrow 0$ on both sides of (4.76), we get

$$\begin{aligned}
& \|X_t\|_{\mathbb{H}}^{2-\alpha} + \delta(c^*)^{\alpha} (1 - \frac{\alpha}{2}) \int_r^t \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \\
& \leq \|X_r\|_{\mathbb{H}}^{2-\alpha} + \mathcal{M}_{r,t}
\end{aligned}$$

$$+(1 - \frac{\alpha}{2}) \int_r^t \frac{(g(\|X_s\|_{\mathbb{H}}^2) + \|\mathcal{B}(s, X_s)\|_{L_2(U, \mathbb{H})}^2) \|X_s\|_{\mathbb{H}}^2 - \alpha \|\mathcal{B}(s, X_s)^* X_s\|_U^2}{\|X_s\|_{\mathbb{H}}^{\alpha+2}} \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds.$$

where

$$\mathcal{M}_{r,t} := 2(1 - \frac{\alpha}{2}) \int_r^t \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} \frac{\langle X_s, \mathcal{B}(s, X_s) dW_s \rangle_{\mathbb{H}}}{\|X_s\|_{\mathbb{H}}^{\alpha}}.$$

According to the assumption (\mathbf{A}_5^*) , we have

$$\|X_t\|_{\mathbb{H}}^{2-\alpha} + \delta(c^*)^{\alpha} (1 - \frac{\alpha}{2}) \int_r^t \mathbf{1}_{\{\|X_s\|_{\mathbb{H}} > 0\}} ds \leq \|X_r\|_{\mathbb{H}}^{2-\alpha} + \mathcal{M}_{r,t}.$$

We complete the proof. \square

Proof of Theorem 2.3: We first prove (2.13). For any $0 \leq r < t$, by (4.72) and (4.74) we deduce

$$(4.77) \quad \mathbb{E}[\|X_t\|_{\mathbb{H}}^{2-\alpha} | \mathcal{F}_r] \leq \|X_r\|_{\mathbb{H}}^{2-\alpha} + \mathbb{E}[\mathcal{M}_{r,t} | \mathcal{F}_r] = \|X_r\|_{\mathbb{H}}^{2-\alpha},$$

which implies that the process

$$t \mapsto \|X_t\|_{\mathbb{H}}^{2-\alpha}$$

is an (\mathcal{F}_t) -nonnegative supermartingale. This combining with (4.72) yields that for every pair of stopping times $\tau^1 < \tau^2$,

$$\mathbb{E}\|X_{\tau^2}\|_{\mathbb{H}}^{2-\alpha} \leq \mathbb{E}\|X_{\tau^1}\|_{\mathbb{H}}^{2-\alpha}.$$

In particular, for any $t > \tau_e$, we have

$$\mathbb{E}\|X_t\|_{\mathbb{H}}^{2-\alpha} \leq \mathbb{E}\|X_{\tau_e}\|_{\mathbb{H}}^{2-\alpha} = 0.$$

Thus, it follows that for any $t \geq \tau_e$

$$\|X_t\|_{\mathbb{H}} = 0, \mathbb{P}\text{-a.s.}$$

As for (2.14), we set $r = 0$ and take expectation in (4.74), by a standard localization argument we obtain that for all $t \geq 0$,

$$\delta(c^*)^{\alpha} (1 - \frac{\alpha}{2}) \int_0^t \mathbb{P}(\tau_e > s) ds \leq \|x\|_{\mathbb{H}}^{2-\alpha}.$$

This implies

$$\mathbb{P}(\tau_e > t) \leq \|x\|_{\mathbb{H}}^{2-\alpha} / (\delta(c^*)^{\alpha} (1 - \frac{\alpha}{2})) t,$$

which yields

$$\mathbb{P}(\tau_e < \infty) = 1.$$

We complete the proof. \square

5. APPENDIX

The classical Skorokhod theorem can only be applied in metric space. In this work, we use the following Jakubowski's version of the Skorokhod theorem in the form presented by Brzeźniak and Ondreját [15].

Lemma 5.1. (*Skorokhod Theorem*) *Let \mathcal{Y} be a topological space such that there exists a sequence of continuous functions $f_m : \mathcal{Y} \rightarrow \mathbb{R}$ that separates points of \mathcal{Y} . Let us denote by \mathcal{S} the σ -algebra generated by the maps f_m . Then*

(i) *every compact subset of \mathcal{Y} is metrizable;*

(ii) *if (μ_m) is tight sequence of probability measures on $(\mathcal{Y}, \mathcal{S})$, then there exists a subsequence denoted also by (m) , a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with \mathcal{Y} -valued Borel measurable variables ξ_m, ξ such that μ_m is the law of ξ_m and ξ_m converges to ξ almost surely on Ω . Moreover, the law of ξ is a Random measure.*

We first recall the definitions of the countably generated Borel space and the standard Borel space in the sense of Parthasarathy (cf. [58, Chapter V, Definition 2.1 and 2.2]).

Definition 5.1. (*Countably generated Borel space*) *A Borel space $(\mathcal{X}, \mathcal{B})$ is said to be countably generated if there exists a denumerable class $\mathcal{D} \subset \mathcal{B}$ such that \mathcal{D} generates \mathcal{B} .*

Definition 5.2. (*Standard Borel space*) *A countably generated Borel space $(\mathcal{X}, \mathcal{B})$ is called standard if there exists a Polish space \mathcal{Y} such that the σ -algebras \mathcal{B} and \mathcal{Y} are σ -isomorphic.*

In order to apply Lemma 5.1, we recall the following result from [48].

Theorem 5.1. (*Theorem B.4 in [48]*) *Let $(\mathcal{X}, \mathcal{B})$ be any standard Borel space. Suppose that $\{f_m\}_{m \in \mathbb{N}}$ is an \mathcal{B} -measurable sequence from \mathcal{X} to \mathbb{R} , which separate the points of \mathcal{X} . Denote by $\sigma_0(\mathcal{X})$ the σ -algebra generated by $\{f_m\}_{m \in \mathbb{N}}$. Then $\sigma_0(\mathcal{X}) = \mathcal{B}$.*

Data availability Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Statements and Declarations On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [1] AGRESTI, A. and VERAAR, M. (2024). The critical variational setting for stochastic evolution equations. *Probab. Theory Relat. Fields* **188**, 957–1015.
- [2] ALDOUS, D. (1978). Stopping times and tightness, *Ann. Probab.* **6** (2), 335-340.
- [3] APPLEBY, J. A.D., MAO, X. and RODKINA, A (2008). Stabilization and destabilization of nonlinear differential equations by noise. *IEEE Trans. Automat. Control* **53**, 683-691.
- [4] BAGNARA, M., MAURELLI, M. and XU, F. (2023). No blow-up by nonlinear Itô noise for the Euler equations. arXiv:2305.09852.
- [5] BARBU, V. DA PRATO, G. and RÖCKNER, M. (2009). Stochastic porous media equations and self-organized criticality. *Comm. Math. Phys.* **285**, 901-923.
- [6] BARBU, V. DA PRATO, G. and RÖCKNER, M. (2009). Finite time extinction for solutions to fast diffusion stochastic porous media equations. *C. R. Math. Acad. Sci. Paris* **347**, 81-84.
- [7] BARBU, V. and RÖCKNER, M. (2012). Stochastic porous media equations and self-organized criticality: convergence to the critical state in all dimensions. *Comm. Math. Phys.* **311**, 539-555.

- [8] BARBU, V., RÖCKNER, M. and RUSSO, F. (2015). Stochastic porous media equations in \mathbb{R}^d . *J. Math. Pures Appl.* **103**, 1024-1052.
- [9] BLÖMKER, D., FLANDOLI F. and ROMITO, M. (2009). Markovianity and ergodicity for a surface growth PDE. *Ann. Probab.* **37**, 275-313.
- [10] BROWDER, F.E. (1963). Nonlinear elliptic boundary value problems. *Bull. Amer. Math. Soc.* **69**, 862-874.
- [11] BROWDER, F.E. (1964). Non-linear equations of evolution, *Ann. Math.* **80**, 485-523.
- [12] BROWDER, F. E. (1977). Pseudo-monotone operators and nonlinear elliptic boundary value problems on unbounded domains. *Proc. Nat. Acad. Sci.* **74**, 2659-2661.
- [13] BRZEŹNIAK, Z. and MOTYL, E. (2013). Existence of a martingale solution of the stochastic Navier-Stokes equations in unbounded 2D and 3D domains. *J. Differential Equations* **254**, 1627-1685.
- [14] BRZEŹNIAK, Z. and MOTYL, E. (2019). Fractionally dissipative stochastic quasi-geostrophic type equations on \mathbb{R}^d . *SIAM J. Math. Anal.* **51**, 2306-2358.
- [15] BRZEŹNIAK, Z. and ONDREJÁT, M. (2013). Stochastic geometric wave equations with values in compact Riemannian homogeneous spaces. *Ann. Probab.* **41**, 1938-1977.
- [16] BUCKMASTER, T., NAHMOD, A., STAFFILANI, G. and WIDMAYER, K. (2020). The surface quasi-geostrophic equation with random diffusion. *Int. Math. Res. Not. IMRN* **23**, 9370-9385.
- [17] CAFFARELLI, L. and VASSEUR, A. (2010). Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation. *Ann. of Math.* **171**, 1903-1930.
- [18] CONSTANTIN, P. and NGUYEN, H.Q. (2018). Global weak solutions for SQG in bounded domains. *Comm. Pure Appl. Math.* **71**, 2323-2333.
- [19] CRISAN, D. and LANG, O. (2024). Global solutions for stochastically controlled fluid dynamics models. *arXiv:2403.05923*.
- [20] DA PRATO, G. and DEBUSSCHE, A. (2003). Ergodicity for the 3D stochastic Navier-Stokes equations. *J. Math. Pure. Appl.* **82**, 877-947.
- [21] DIBENEDETTO, E. (1993). Degenerate Parabolic Equations, Springer, New York.
- [22] FLANDOLI, F. and GATAREK, D. (1995). Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Related Fields* **102**, 367-391.
- [23] FLANDOLI, F., GUBINELLI, M. and PRIOLA, E. (2010). Well-posedness of the transport equation by stochastic perturbation. *Invent. Math.* **180**, 1-53.
- [24] FLANDOLI, F. (2011). Random perturbation of PDEs and fluid dynamic models. *Lecture Notes in Mathematics*.
- [25] FLANDOLI, F. (2015). A stochastic view over the open problem of well-posedness for the 3D Navier-Stokes equations. Stochastic analysis: a series of lectures, *Progr. Probab.*, **68**, Birkhäuser/Springer, Basel, 221-246.
- [26] FLANDOLI, F. and LUO, D. (2021). High mode transport noise improves vorticity blow-up control in 3D Navier-Stokes equations. *Probab. Theory Related Fields* **180**, 309-363.
- [27] FLANDOLI, F. and ROMITO, M. (2008). Markov selections for the 3D stochastic Navier-Stokes equations. *Probab. Theory Related Fields* **140**, 407-458.
- [28] FOIAS, C., MANLEY, O., ROSA, R. and TEMAM, R. (2001). Navier-Stokes equations and turbulence (Vol. 83). Cambridge University Press.
- [29] GESS, B. (2015). Finite time extinction for stochastic sign fast diffusion and self-organized criticality. *Comm. Math. Phys.* **335**, 309-344.
- [30] GESS, B. (2018). Regularization and well-posedness by noise for ordinary and partial differential equations. Stochastic partial differential equations and related fields, 43-67, *Springer Proc. Math. Stat.* **229**, Springer, Cham.
- [31] GLATT-HOLTZ, N. and VICOL, V.C. (2014). Local and global existence of smooth solutions for the stochastic Euler equations with multiplicative noise. *Ann. Probab.* **42**, 80-145.
- [32] GLATT-HOLTZ, N. and ZIANE, M. (2009). Strong pathwise solutions of the stochastic Navier-Stokes system. *Adv. Differential Equations* **14**, 567-600.
- [33] GOLDYS, B., RÖCKNER, M. and ZHANG, X. (2009). Martingale solutions and Markov selections for stochastic partial differential equations. *Stochastic Process. Appl.* **119**, 1725-1764.

- [34] GYÖNGY, I. and MARTINEZ, T. (2001). On stochastic differential equations with locally unbounded drift. *Czechoslovak Math. J.* **51**, 763-783.
- [35] HAIRER, M. and MATTINGLY, J. C. (2006). Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing. *Ann. of Math. (2)* **164**, 993-1032.
- [36] HARTMAN, P. and STAMPACCHIA, G. (1966). On some nonlinear elliptic differential equations, *Acta Math.* **115**, 271-310.
- [37] HENSEL, S. (2021). Finite time extinction for the 1D stochastic porous medium equation with transport noise. *Stoch. Partial Differ. Equ. Anal. Comput.* **9**, 892-939.
- [38] HOFMANOVÁ, M., ZHU, R.-C. and ZHU, X.-C. (2024). Nonuniqueness in law of stochastic 3D Navier-Stokes equations. *J. Eur. Math. Soc. (JEMS)* **26**, 163-260.
- [39] HONG, W., HU, S. and LIU, W. (2024). McKean-Vlasov SDEs and SPDEs with locally monotone coefficients, *Ann. Appl. Probab.* **34**, 2136-2189.
- [40] JAKUBOWSKI A. (1986). On the Skorokhod topology. *Ann. Inst. H. Poincaré Probab. Statist.* **22**, 263-285.
- [41] KATO, T. and PONCE, G. (1988) Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.* **41**, 891-907.
- [42] KISELEV, A., NAZAROV, F. and VOLBERG, A. (2007). Global well-posedness for the critical 2D dissipative quasi-geostrophic equation. *Invent. Math.* **167**, 445-453.
- [43] KRYLOV, N.V. and ROZOVSKII, B.L. (1981). Stochastic evolution equations. *Translated from Itogi Nauki i Tekhniki, Seriya Sovremennye Problemy Matematiki. Plenum Publishing Corp.* **14**, 71-146.
- [44] KRYLOV, N.V. and RÖCKNER, M. (2005). Strong solutions of stochastic equations with singular time dependent drift. *Probab. Theory Related Fields* **131**, 154-196.
- [45] KUKSIN, S. NERSESYAN, V. and SHIRIKYAN, A. (2020). Exponential mixing for a class of dissipative PDEs with bounded degenerate noise. *Geom. Funct. Anal.* **30**, 126-187.
- [46] LERAY, J. and LIONS, J.L. (1965). Quelques résultats de Visik sur les problèmes elliptiques non linéaires par les méthodes de Minty-Browder, *Bull. Soc. Math. France* **93**, 97-107.
- [47] LI, Y. and XIE, C. (2003). Blow-up for p -Laplacian parabolic equations. *Electron. J. Differential Equations* **20**, 12 pp.
- [48] LIANG, S. (2021). Stochastic hypodissipative hydrodynamic equations: well-posedness, stationary solutions and ergodicity. PhD Thesis, Bielefeld University.
- [49] J.L. LIONS (1969). Quelques méthodes de résolution des problèmes aux limites non linéaires, Dunod, Paris.
- [50] LIU, W. and RÖCKNER, M. (2010). SPDE in Hilbert space with locally monotone coefficients. *J. Funct. Anal.* **259**, 2902-2922.
- [51] LIU, W. and RÖCKNER, M. (2013). Local and global well-posedness of SPDE with generalized coercivity conditions. *J. Differential Equations* **254**, 725-755.
- [52] LIU, W. and RÖCKNER, M. (2015). Stochastic Partial Differential Equations: An Introduction. Universitext, Springer.
- [53] MASLOWSKI, B. and SEIDLER, J. (1999). On sequentially weakly Feller solutions to SPDE's. *Rend. Lincei Mat. Appl.* **10**, 69-78.
- [54] MINTY, G. J. (1962). Monotone (non-linear) operators in Hilbert space. *Duke. Math. J.* **29**, 341-346.
- [55] NGUYEN, P., TAWRI, K. and TEMAM, R. (2021). Nonlinear stochastic parabolic partial differential equations with a monotone operator of the Ladyženskaya-Smagorinsky type, driven by a Lévy noise. *J. Funct. Anal.* **281**, Paper No. 109157, 74 pp.
- [56] NIRENBERG, L. (1959). On elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **13**, 115-162.
- [57] PARDOUX, E. (1975). Equations aux dérivées partielles stochastiques non linéaires monotones, *Ph.D. thesis*, Université Paris XI.
- [58] PARTHASARATHY, K.R. (1967). Probability Measures on Metric Spaces, Academic Press.
- [59] RAIBLE, M., LINZ, S. J. and HÄNGGI, P. (2000). Amorphous thin film growth: Minimal deposition equation. *Physical Review E* **62**, 1691-1705.

- [60] REN, P., TANG, H. and WANG, F.-Y. (2024). Distribution-path dependent nonlinear SPDEs with application to stochastic transport type equations. *Potential Anal.* **61**, 379-407.
- [61] RESNICK, S. (1995). Danamical Problems in Non-linear Advective Partial Differential Equations. Ph.D. Thesis, University of Chicago, Chicago.
- [62] RÖCKNER, M., SCHMULAND, B. and ZHANG, X. (2008). Yamada-Watanabe theorem for stochastic evolution equations in infinite dimensions. *Condens. Matter Phys.* **54**, 247-259.
- [63] RÖCKNER, M., SHANG, S. and ZHANG, T. (2024). Well-posedness of stochastic partial differential equations with fully local monotone coefficients. *Math. Ann.* **390**, 3419-3469.
- [64] RÖCKNER, M. and WANG, F.-Y. (2013). General extinction results for stochastic partial differential equations and applications. *J. Lond. Math. Soc.* **87**, 545-560
- [65] RÖCKNER, M., ZHU, R.-C. and ZHU, X.-C. (2014). Local existence and non-explosion of solutions for stochastic fractional partial differential equations driven by multiplicative noise. *Stochastic Process. Appl.* **124**, 1974-2002.
- [66] RÖCKNER, M., ZHU, R.-C. and ZHU, X.-C. (2015). Sub and supercritical stochastic quasi-geostrophic equation. *Ann. Probab.* **43**, 1202-1273.
- [67] STEIN, E. (1970). Singular Integrals and Differentiability Properties of Functions. *Princeton, NJ: Princeton University Press.*
- [68] SIMON J. (1987). Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl.* **164**, 65-96.
- [69] TANG, H. and WANG, F. Y. (2022). A general framework for solving singular SPDEs with applications to fluid models driven by pseudo-differential noise. *arXiv:2208.08312*.
- [70] TANG, H. and YANG, A. (2023). Noise effects in some stochastic evolution equations: global existence and dependence on initial data. *Ann. Inst. Henri Poincaré Probab. Stat.* **59**, 378-410.
- [71] TIAN, Y. and MU, C. (2008). Extinction and non-extinction for a p -Laplacian equation with nonlinear source. *Nonlinear Anal.* **69**, 2422-2431.
- [72] VERETENNIKOV, A. J. (1980). Strong solutions and explicit formulas for solutions of stochastic integral equations. *Mat. Sb. (N.S.)* **111**, 434-452, 480.
- [73] ZEIDLER, E. (1990). Nonlinear functional analysis and its applications, II/B, nonlinear monotone operators. *Springer-Verlag*, New York.
- [74] ZVONKIN, A. K. (1974). A transformation of the phase space of a diffusion process that will remove the drift. *Mat. Sb. (N.S.)* **93** (135), 129-149, 152.