

DECAY ESTIMATES FOR A CLASS OF DUNKL WAVE EQUATIONS

CHENG LUO, SHYAM SWARUP MONDAL, AND MANLI SONG

ABSTRACT. Let Δ_κ be the Dunkl Laplacian on \mathbb{R}^n and $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is a smooth function. The aim of this manuscript is twofold. First, we study the decay estimate for a class of dispersive semigroup of the form $e^{it\phi(\sqrt{-\Delta_\kappa})}$. We overcome the difficulty arising from the non-homogeneity of ϕ by frequency localization. As applications, in the next part of the paper, we establish Strichartz estimates for some concrete wave equations associated with the Dunkl Laplacian Δ_κ , which corresponds to $\phi(r) = r, r^2, r^2 + r^4, \sqrt{1+r^2}, \sqrt{1+r^4}$, and $r^\mu, 0 < \mu \leq 2, \mu \neq 1$. More precisely, we unify and simplify all the known dispersive estimates and extend to more general cases. Finally, using the decay estimates, we prove the global-in-time existence of small data Sobolev solutions for the nonlinear Klein-Gordon equation and beam equation with the power type nonlinearities.

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 2. Preliminaries | 5 |
| 2.1. Dunkl operators | 5 |
| 2.2. The Dunkl transform | 7 |
| 2.3. Dunkl convolution operator | 8 |
| 2.4. Dunkl Besov spaces | 8 |
| 2.5. Technical Lemmas | 9 |
| 3. Proof of main results | 11 |
| 4. Applications | 15 |
| 4.1. Wave equation | 16 |
| 4.2. Fractional Schrödinger equation | 18 |
| 4.3. Fourth-order Schrödinger equation | 20 |
| 4.4. Klein-Gordon equation | 21 |
| 4.5. Beam equation | 22 |
| 5. Well-posedness results for nonlinear Klein-Gordon and beam equations | 23 |
| Statements and Declarations | 28 |
| Acknowledgments | 28 |
| References | 28 |

1. INTRODUCTION

The main aim of this paper is to investigate decay estimates for a class of dispersive equations of the form

$$\begin{cases} i\partial_t u + \phi(\sqrt{-\Delta_\kappa})u = f, \\ u|_{t=0} = u_0, \end{cases} \quad (1.1)$$

Date: July 10, 2024.

2010 Mathematics Subject Classification. Primary 22E25, 33C45, 35H20, 35B40.

Key words and phrases. Dunkl Laplacian, Dunkl Bolev space, Wave equation, Decay estimate, Strichartz estimate.

where Δ_κ is the Dunkl Laplacian on \mathbb{R}^n , $f : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{C}$, and $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is a smooth function that satisfies some growth condition.

Dispersive inequalities for evolution equations, such as the Schrödinger and wave equations play a significant role in the study of semilinear and quasilinear problems, which naturally appear in several physical applications. Proving dispersion means estimating the decay in time on the evolution group associated with the free equation. In recent times, dispersive properties of evolution equations have become a crucial tool in the study of a variety of questions, including local and global existence for nonlinear equations, well-posedness of Cauchy problems for nonlinear equations in Sobolev spaces of low order, scattering theory, and many others. In many circumstances, the essential step in proving this time decay is based on the application of a stationary phase theorem on an (approximate) representation of the solution. Dispersion phenomena, when combined with an abstract functional analysis argument, famously known as the TT^* -argument (see [34]), offer a range of estimates involving space-time Lebesgue norms. These inequalities also known as Strichartz estimates, have evolved into a fundamental and amazing tool in the study of nonlinear partial differential equations over the last few decades, see [7] and references therein.

Strichartz estimates also be interpreted as Fourier restriction estimates, which have a vital role in classical harmonic analysis and are closely linked to arithmetic combinatorics. Considerable attention has been devoted by several researchers to deriving Strichartz estimates in different frameworks. For example, in the Euclidean setting, these estimates have been proved for several dispersive equations, such as the wave equation and Schrödinger equation, for instance, see the pioneering works [24, 34, 50]. The theory of Strichartz estimates has also been extended in the non-Euclidean frameworks. For example, see [4–6, 22, 23, 28, 35, 45–47] for Strichartz estimates on H-type groups and more generality of step 2 stratified Lie groups, [8, 10, 12, 16, 21, 36, 38, 39, 41, 42, 44] for metric measure spaces, [1–3] for hyperbolic spaces, [11, 13] for compact Riemannian manifolds, and [30] for bounded domains.

One of the essential ingredients in the derivation of Strichartz estimates is the dispersive estimates. Our aim of this paper is to establish a decay estimate for a class of dispersive equations as follows:

$$\begin{cases} i\partial_t u + \phi(\sqrt{-\Delta_\kappa})u = f, \\ u|_{t=0} = u_0, \end{cases} \quad (1.2)$$

where Δ_κ is the Dunkl Laplacian on \mathbb{R}^n , $f : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{C}$, and $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is a smooth function satisfying

(C1) There exists $m_1 > 0$ such that for any $\alpha \geq 2$ and $\alpha \in \mathbb{N}$,

$$|\phi'(r)| \sim r^{m_1-1}, \quad |\phi^{(\alpha)}(r)| \lesssim r^{m_1-\alpha}, \quad r \geq 1.$$

(C2) There exists $m_2 > 0$ such that for any $\alpha \geq 2$ and $\alpha \in \mathbb{N}$,

$$|\phi'(r)| \sim r^{m_2-1}, \quad |\phi^{(\alpha)}(r)| \lesssim r^{m_2-\alpha}, \quad 0 < r < 1.$$

(C3) There exists $\alpha_1 > 0$ such that

$$|\phi''(r)| \sim r^{\alpha_1-2}, \quad r \geq 1.$$

(C4) There exists $\alpha_2 > 0$ such that

$$|\phi''(r)| \sim r^{\alpha_2-2}, \quad 0 < r < 1.$$

Here we use a Fourier multiplier notation to define $\phi(\sqrt{-\Delta_\kappa})$ as in the following form

$$\phi(\sqrt{-\Delta_\kappa})f = \mathcal{F}_\kappa^{-1}\phi(|\xi|)\mathcal{F}_\kappa f,$$

where \mathcal{F}_κ denotes Dunkl transform, defined in (2.3), Section 2. Notice that the conditions (C1) and (C3) represent the homogeneous order of ϕ in high frequency, and the conditions (C2) and

(C4) reflect the homogeneous order of ϕ in low frequency. If ϕ satisfies conditions (C1) and (C3), then $\alpha_1 \leq m_1$. Similarly, if ϕ satisfies (C2) and (C4), then $\alpha_2 \geq m_2$.

Here, we want to emphasize that several dispersive wave equations associated with Dunkl Laplacian reduced to type (1.2). For instance, the Dunkl Schrödinger equation corresponds to $\phi(r) = r^2$, the Dunkl wave equation corresponds to $\phi(r) = r$, the fractional Dunkl Schrödinger equation corresponds to $\phi(r) = r^\mu$, $0 < \mu < 2, \mu \neq 1$, the fourth-order Dunkl Schrödinger equation corresponds to $\phi(r) = r^2 + r^4$, the beam equation $\phi(r) = \sqrt{1 + r^4}$, and the Klein-Gordon equation corresponds to $\phi(r) = \sqrt{1 + r^2}$, among others.

Dunkl operators are differential-difference operators that generalize the standard partial derivatives. These operators were introduced by Charles Dunkl [18] using a finite reflection group and a parameter function on its root system. Applications of Dunkl operators in mathematics and mathematical physics are numerous. For example, they play an important role in the integration of quantum many-body problems of the Calogero-Moser-Sutherland type and are employed in the study of probabilistic processes. Additionally, these operators have had a lasting impression on the study of special functions in one and multiple variables as well as orthogonal polynomials, see [20].

For $\kappa = 0$, as the the Dunkl Laplacian Δ_κ reduces to usual Laplacian Δ on \mathbb{R}^n , the system (1.2) reduces to

$$\begin{cases} i\partial_t u + \phi(\sqrt{-\Delta})u = f, \\ u|_{t=0} = u_0. \end{cases} \quad (1.3)$$

In 1977, Strichartz [50] derived the priori estimates for the solution to (1.3) in the space-time norm using the classical Fourier restriction theorem of Stein and Tomas [48, 53, 54]. However, later, it was improved by Ginibre-Velo [24] using a standard duality argument along with the dispersive estimate of the form

$$\left\| e^{it\phi(\sqrt{-\Delta})} u_0 \right\|_X \lesssim |t|^{-\theta} \|u_0\|_{X^*}, \quad (1.4)$$

where X^* is the dual space of X . In 2008, Guo-Peng-Wang [27] used a unified way to study the decay for a class of dispersive semigroup $e^{it\phi(\sqrt{-\Delta})}$ on \mathbb{R}^n for the system (1.3). They assumed that $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is smooth satisfying (C1)-(C4) and obtained several decay estimates in time for the dispersive semigroup $e^{it\phi(\sqrt{-\Delta})}$ by introducing Littlewood-Paley projector operator, see [27, Theorem 1].

Motivated by the recent works, in this paper, we aim to establish the decay estimate for a class of dispersive equations associated with Dunkl Laplacian (1.2). Note that, if ϕ is a homogeneous function of order m , namely, $\phi(\lambda r) = \lambda^m \phi(r)$ for all $\lambda > 0$, the dispersive estimate can be easily obtained by a theorem of Littman and dyadic decomposition. However, things become complicated when the function ϕ is not homogeneous because the scaling constants can not be effectively separated from the time. Since the phases we consider are not necessarily homogeneous or weakly dispersive, to overcome the difficulty, we use a Littlewood-Paley decomposition (see [27]) to more accurately capture the difference between the low and the high frequencies in different scales and catch the possible need for extra regularity.

Let S_0 and $\tilde{\Delta}_j$ be the Littlewood-Paley decomposition related to the Dunkl Laplacian Δ_κ and $N = 2\gamma + n$ (see Section 2 for detailed definition). Then the main results of this paper are as follows:

Theorem 1.1. *Assume $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}$ is smooth and $U_t = e^{it\phi(\sqrt{-\Delta_\kappa})}$. Then the subsequent outcomes are valid.*

(A) *For $j \geq 0$, assume that ϕ satisfies (C1), then*

$$\|U_t \tilde{\Delta}_j u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad 0 \leq \theta \leq \frac{N-1}{2}. \quad (1.5)$$

In addition, if ϕ satisfies (C3), then

$$\|U_t \tilde{\Delta}_j u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim |t|^{-\frac{N-1+\theta}{2}} 2^j \left(N - \frac{m_1(N-1+\theta)}{2} - \frac{\theta(\alpha_1 - m_1)}{2} \right) \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad 0 \leq \theta \leq 1. \quad (1.6)$$

(B) For $j < 0$, suppose ϕ satisfies (C2), then

$$\|U_t \tilde{\Delta}_j u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim |t|^{-\theta} 2^{j(N-m_2\theta)} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad 0 \leq \theta \leq \frac{N-1}{2}.$$

In addition, if ϕ satisfies (C4), then

$$\|U_t \tilde{\Delta}_j u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim |t|^{-\frac{N-1+\theta}{2}} 2^j \left(N - \frac{m_2(N-1+\theta)}{2} - \frac{\theta(\alpha_2 - m_2)}{2} \right) \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad 0 \leq \theta \leq 1.$$

(C) If ϕ satisfies (C2), then

$$\|U_t S_0 u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim (1 + |t|)^{-\theta} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad \theta = \min \left(\frac{N}{m_2}, \frac{N-1}{2} \right). \quad (1.7)$$

In addition, if ϕ satisfies (C4) and $\alpha_2 = m_2$, then

$$\|U_t S_0 u_0\|_{L^\infty(\mathbb{R}^n)} \lesssim (1 + |t|)^{-\theta} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \quad \theta = \min \left(\frac{N}{m_2}, \frac{N}{2} \right). \quad (1.8)$$

We prove the above result in Section 3. The main difficulty in the setup of Dunkl is that we don't have an explicit expression for the Dunkl transform compared to the classical Fourier transform. In this paper, we used Littlewood-Paley decomposition to better understand the regularity is needed at low and high frequencies in order to get a time decay. We also use the fact that Dunkl transform for radial functions can be expressed in terms of integrals of Bessel functions. In order to estimate these integrals, we rely on Van der Corput's Lemma and the decay properties of the Bessel functions.

Using the decay estimates in Theorem 1.1, we establish Strichartz estimates for the Dunkl Schrödinger equation (corresponds to $\phi(r) = r^2$), the Dunkl wave equation (corresponds to $\phi(r) = r$), the fractional Dunkl Schrödinger equation (corresponds to $\phi(r) = r^\mu$), $0 < \mu < 2, \mu \neq 1$, the fourth-order Dunkl Schrödinger equation (corresponds to $\phi(r) = r^2 + r^4$), the beam equation (corresponds to $\phi(r) = \sqrt{1 + r^4}$), and the Klein-Gordon equation (corresponds to $\phi(r) = \sqrt{1 + r^2}$). We also prove the global well-posedness for the nonlinear Klein-Gordon equation and beam equation for small initial data.

There are many situations where dispersion phenomena fail, for instance in compact manifolds, on some domains, on the Heisenberg group. However, in some cases, Strichartz estimates (in weak forms) or smoothing properties can be established using other approaches that are not based on dispersive inequality techniques. The Fourier restriction theorem and Kato smoothing effect techniques stand out among them. Recently, the authors in [32] proved Strichartz estimates for the Dunkl Schrödinger equation (which corresponds to $\phi(r) = r^2$) using the Fourier-Dunkl transform restriction theorem for paraboloid surfaces. See also [36, 38].

Remark 1.2. Since the Dunkl Laplacian Δ_κ reduces to the Euclidean Laplacian Δ for $\kappa = 0$, we recover the results proved in Guo-Peng-Wang [27].

Apart from the introduction, this paper is organized as follows: In Section 2, we recall harmonic analysis associated with the Dunkl operators, Dunkl transforms, and their certain properties. We also recall some useful technical lemmas for proving the decay estimates and Strichartz estimates. In Section 3, we prove our main result of this paper, i.e., Theorem 1.1. We provide Strichartz estimates for some concrete wave equations associated with the Dunkl Laplacian Δ_κ in Section 4. In Section 5, we prove global well-posedness for the nonlinear Klein-Gordon equation and beam equation for small initial data.

Notations:

- $A \lesssim B$ means that $A \leq CB$, and $A \sim B$ stands for $C_1B \leq A \leq C_2B$, where C, C_1, C_2 denote positive universal constants.
- Δ_κ denotes Dunkl-Laplacian on \mathbb{R}^n .
- S_0 and $\tilde{\Delta}_j$ be the Littlewood-Paley decomposition related to Δ_κ .
- J_ν denotes the Bessel function of order ν .
- $B_{p,q}^{s,\kappa}$ denotes the Dunkl Besov space.

2. PRELIMINARIES

In this section, we recall some basic definitions and important properties of the Dunkl operators and briefly overview the related harmonic analysis to make the paper self-contained. A complete account of harmonic analysis related to Dunkl operators can be found in [9, 10, 38, 40, 43, 52].

2.1. Dunkl operators. The basic ingredient in the theory of Dunkl operators are root systems and finite reflection groups. We start this section by the definition of root system.

Let $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean scalar product in \mathbb{R}^n . For $x \in \mathbb{R}^n$, we denote $|x|$ as $|x| = \langle x, x \rangle^{1/2}$. For $\alpha \in \mathbb{R}^n \setminus \{0\}$, we denote r_α as the reflection with respect to the hyperplane $\langle \alpha \rangle^\perp$, orthogonal to α and is defined by

$$r_\alpha(x) := x - 2 \frac{\langle \alpha, x \rangle}{|\alpha|^2} \alpha, \quad x \in \mathbb{R}^n.$$

A finite set \mathcal{R} in $\mathbb{R}^n \setminus \{0\}$ is said to be a root system if the following holds:

- (1) $r_\alpha(\mathcal{R}) = \mathcal{R}$ for all $\alpha \in \mathcal{R}$,
- (2) $\mathcal{R} \cap \mathbb{R}\alpha = \{\pm\alpha\}$ for all $\alpha \in \mathcal{R}$.

For a given root system \mathcal{R} , the subgroup $G \subset O(n, \mathbb{R})$ generated by the reflections $\{r_\alpha \mid \alpha \in \mathcal{R}\}$ is called the finite Coxeter group associated with \mathcal{R} . The dimension of $\text{span}\mathcal{R}$ is called the rank of \mathcal{R} . For a detailed study on the theory of finite reflection groups, we refer to [29]. Let $\mathcal{R}^+ := \{\alpha \in \mathcal{R} : \langle \alpha, \beta \rangle > 0\}$ for some $\beta \in \mathbb{R}^n \setminus \bigcup_{\alpha \in \mathcal{R}} \langle \alpha \rangle^\perp$, be a fix positive root system.

Some typical examples of such a system are the Weyl groups such as the symmetric group S_n for the type A_{n-1} root system and the hyperoctahedral group for the type B_n root system. In addition, H_3, H_4 (icosahedral groups) and, $I_2(n)$ (symmetry group of the regular n -gon) are also the Coxeter groups.

A multiplicity function for G is a function $\kappa : \mathcal{R} \rightarrow \mathbb{C}$ which is constant on G -orbits. Setting $\kappa_\alpha := \kappa(\alpha)$ for $\alpha \in \mathcal{R}$, from the definition of G -invariant, we have $\kappa_{g\alpha} = \kappa_\alpha$ for all $g \in G$. We say κ is non-negative if $\kappa_\alpha \geq 0$ for all $\alpha \in \mathcal{R}$. The \mathbb{C} -vector space of non-negative multiplicity functions on \mathcal{R} is denoted by \mathcal{K}^+ . Let us denote γ as $\gamma = \gamma(\kappa) := \sum_{\alpha \in \mathcal{R}^+} \kappa(\alpha)$. For convenience, we denote $N = 2\gamma + n$ throughout the paper.

For $\xi \in \mathbb{C}^n$ and $\kappa \in \mathcal{K}^+$, Dunkl in 1989 introduced a family of first-order differential-difference operators $T_\xi := T_\xi(\kappa)$ by

$$T_\xi(\kappa)f(x) := \partial_\xi f(x) + \sum_{\alpha \in \mathcal{R}^+} \kappa_\alpha \langle \alpha, \xi \rangle \frac{f(x) - f(r_\alpha x)}{\langle \alpha, x \rangle}, \quad f \in C^1(\mathbb{R}^n), \quad (2.1)$$

where ∂_ξ denotes the directional derivative corresponding to ξ . The operator T_ξ defined in (2.1), formally known as Dunkl operator and is one of the most important developments in the theory of special functions associated with root systems [17]. They commute pairwise and are skew-symmetric with respect to the G -invariant measure $h_\kappa(x)dx$, where the weight function

$$h_\kappa(x) := \prod_{\alpha \in \mathcal{R}^+} |\langle \alpha, x \rangle|^{2\kappa_\alpha}$$

is of homogeneous degree 2γ . Thanks to the G -invariance of the multiplicity function, the definition of the Dunkl operator is independent of the choice of the positive subsystem \mathcal{R}^+ . In [18], it is shown that for any $\kappa \in \mathcal{K}^+$, there is a unique linear isomorphism V_κ (Dunkl's intertwining operator) on the space $\mathcal{P}(\mathbb{R}^n)$ of polynomials on \mathbb{R}^n such that

- (1) $V_\kappa(\mathcal{P}_m(\mathbb{R}^n)) = \mathcal{P}_m(\mathbb{R}^n)$ for all $m \in \mathbb{N}$,
- (2) $V_\kappa|_{\mathcal{P}_0(\mathbb{R}^n)} = id$,
- (3) $T_\xi(\kappa)V_\kappa = V_\kappa\partial_\xi$,

where $\mathcal{P}_m(\mathbb{R}^n)$ denotes the space of homogeneous polynomials of degree m .

For any finite reflection group G and any $k \in \mathcal{K}^+$, Rösler in [43] proved that there exists a unique positive Radon probability measure ρ_x^κ on \mathbb{R}^n such that

$$V_\kappa f(x) = \int_{\mathbb{R}^n} f(\xi) d\rho_x^\kappa(\xi). \quad (2.2)$$

The measure ρ_x^κ depends on $x \in \mathbb{R}^n$ and its support is contained in the ball $B(\|x\|) := \{\xi \in \mathbb{R}^n : \|\xi\| \leq \|x\|\}$. In view of the Laplace type representation (2.2), Dunkl's intertwining operator V_κ can be extended to a larger class of function spaces.

Let $\{\xi_1, \xi_2, \dots, \xi_n\}$ be an orthonormal basis of $(\mathbb{R}^n, \langle \cdot, \cdot \rangle)$. Then the Dunkl Laplacian operator Δ_κ is defined as

$$\Delta_\kappa = \sum_{j=1}^n T_{\xi_j}^2(\kappa).$$

The definition of Δ_κ is independent of the choice of the orthonormal basis of \mathbb{R}^n . In fact, one can see that the operator Δ_κ also can be expressed as

$$\Delta_\kappa f(x) = \Delta f(x) + \sum_{\alpha \in \mathcal{R}^+} \kappa_\alpha \left\{ \frac{2\langle \nabla f(x), \alpha \rangle}{\langle \alpha, x \rangle} - |\alpha|^2 \frac{f(x) - f(r_\alpha x)}{\langle \alpha, x \rangle^2} \right\}, \quad f \in C^1(\mathbb{R}^n),$$

where ∇ and Δ are the usual gradient and Laplacian operator on \mathbb{R}^n , respectively. Observe that, for $\kappa \equiv 0$, the Dunkl Laplacian operator Δ_κ reduces to the classical Euclidean Laplacian Δ .

For every $y \in \mathbb{R}^n$, the system

$$\begin{cases} T_\xi u(x, y) = \langle y, \xi \rangle u(x, y), & \xi \in \mathbb{R}^n, \\ u(0, y) = 1, \end{cases}$$

admits a unique analytic solution on \mathbb{R}^n , which we denote as $K(x, y)$ and is known as the Dunkl kernel in general. The kernel has a unique holomorphic extension to $\mathbb{C}^n \times \mathbb{C}^n$ and is given by

$$K(x, y) = V_\kappa(e^{\langle \cdot, y \rangle})(x) = \int_{\mathbb{R}^n} e^{\langle \xi, y \rangle} d\rho_x^\kappa(\xi), \quad \forall x, y \in \mathbb{R}^n,$$

where the positive Radon probability measure ρ_x^κ depends on $x \in \mathbb{R}^n$ and its support is contained in the ball $B(\|x\|) := \{\xi \in \mathbb{R}^n : \|\xi\| \leq \|x\|\}$. When $\kappa \equiv 0$, the Dunkl kernel $K(x, y)$ reduces to the exponential $e^{\langle x, y \rangle}$. For $x, y \in \mathbb{R}^n$ and $z, w \in \mathbb{C}^n$, the Dunkl kernel satisfies the following properties:

- (i) $K(z, w) = K(w, z)$ and $K(\lambda z, w) = K(z, \lambda w)$, $\lambda \in \mathbb{C}$.
- (ii) For any $\nu \in \mathbb{N}^n$, we have

$$|D_z^\nu K(x, z)| \leq |x|^{|\nu|} e^{|\operatorname{Re}(z)|\|x\|},$$

where $D_z^\nu = \frac{\partial^{|\nu|}}{\partial z_1^{\nu_1} \cdots \partial z_n^{\nu_n}}$ and $|\nu| = \nu_1 + \cdots + \nu_n$. In particular, for any $x, y \in \mathbb{R}^n$, we have

$$|K(x, iy)| \leq 1.$$

- (iii) $K(ix, y) = \overline{K(-ix, y)}$ and $K(gx, gy) = K(x, y)$, $g \in G$.

2.2. The Dunkl transform. For $1 \leq p < \infty$, let $L_\kappa^p(\mathbb{R}^n)$ be the space of L^p -functions on \mathbb{R}^n with respect to the weight $h_\kappa(x)$ with the $L_\kappa^p(\mathbb{R}^n)$ -norm

$$\|f\|_{L_\kappa^p(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} |f(x)|^p h_\kappa(x) dx \right)^{\frac{1}{p}},$$

and $\|f\|_{L_\kappa^\infty(\mathbb{R}^n)} = \text{ess sup}_{x \in \mathbb{R}^n} |f(x)|$. The Dunkl transform of $f \in L_\kappa^1(\mathbb{R}^n)$ is defined by

$$\mathcal{F}_\kappa f(y) = \frac{1}{c_\kappa} \int_{\mathbb{R}^n} f(x) K(x, -iy) h_\kappa(x) dx, \quad (2.3)$$

where c_κ is the Mehta-type constant defined by

$$c_\kappa = \int_{\mathbb{R}^n} e^{-\frac{|x|^2}{2}} h_\kappa(x) dx.$$

When $\kappa \equiv 0$, the Dunkl transform coincides with the classical Fourier transform. The Dunkl transform plays the same role as the Fourier transform in classical Fourier analysis and enjoys properties similar to those of the classical Fourier transform. With $N = 2\gamma + n$, here we list some basic properties of the Dunkl transform (see [15, 19, 31]).

(i) For all $f \in L_\kappa^1(\mathbb{R}^n)$, we have

$$\|\mathcal{F}_\kappa f\|_{L_\kappa^\infty(\mathbb{R}^n)} \leq \frac{1}{c_\kappa} \|f\|_{L_\kappa^1(\mathbb{R}^n)}.$$

(ii) For any function f in the Schwartz space $\mathcal{S}(\mathbb{R}^n)$, we have

$$\mathcal{F}_\kappa(T_\xi f)(y) = i\langle \xi, y \rangle \mathcal{F}_\kappa f(y), \quad y, \xi \in \mathbb{R}^n.$$

In particular, it follows that

$$\mathcal{F}_\kappa(\Delta_\kappa f)(y) = -|y|^2 \mathcal{F}_\kappa f(y), \quad y \in \mathbb{R}^n.$$

(iii) For all $f \in L_\kappa^1(\mathbb{R}^n)$, we have

$$\mathcal{F}_\kappa(f(\cdot/\lambda))(y) = \lambda^N \mathcal{F}_\kappa f(\lambda y), \quad y \in \mathbb{R}^n,$$

for all $\lambda > 0$.

(iv) The Dunkl transform \mathcal{F}_κ is a homeomorphism of the Schwartz space $\mathcal{S}(\mathbb{R}^n)$ and its inverse is given by $\mathcal{F}_\kappa^{-1}g(x) = \mathcal{F}_\kappa g(-x)$, for all $g \in \mathcal{S}(\mathbb{R}^n)$. In addition, for all $f \in \mathcal{S}(\mathbb{R}^n)$, it satisfies

$$\int_{\mathbb{R}^n} |f(x)|^2 h_\kappa(x) dx = \int_{\mathbb{R}^n} |\mathcal{F}_\kappa f(y)|^2 h_\kappa(y) dy.$$

In particular, the Dunkl transform extends to an isometric isomorphism on $L_\kappa^2(\mathbb{R}^n)$. The Dunkl transform can be extended to the space of tempered distributions $\mathcal{S}'(\mathbb{R}^n)$ and is also a homeomorphism of $\mathcal{S}'(\mathbb{R}^n)$.

(v) For every $f \in L_\kappa^1(\mathbb{R}^n)$ such that $\mathcal{F}_\kappa f \in L_\kappa^1(\mathbb{R}^n)$, we have the inversion formula

$$f(x) = \frac{1}{c_\kappa} \int_{\mathbb{R}^n} \mathcal{F}_\kappa f(y) K(ix, y) h_\kappa(y) dy, \quad a.e. x \in \mathbb{R}^n.$$

(vi) If f is a radial function in $L_\kappa^1(\mathbb{R}^n)$ such that $f(x) = \tilde{f}(|x|)$, then

$$\mathcal{F}_\kappa f(y) = \frac{1}{\Gamma(N/2)} \int_0^\infty \tilde{f}(r) \frac{J_{\frac{N-2}{2}}(r|y|)}{(r|y|)^{\frac{N-2}{2}}} r^{N-1} dr, \quad (2.4)$$

where J_ν denotes the Bessel function of order $\nu > -\frac{1}{2}$.

2.3. Dunkl convolution operator. For given $x \in \mathbb{R}^n$, the Dunkl translation operator $f \mapsto \tau_x f$ is defined on $\mathcal{S}(\mathbb{R}^n)$ by

$$\mathcal{F}_\kappa(\tau_x f)(y) = K(x, iy)\mathcal{F}_\kappa f(y), \quad y \in \mathbb{R}^n.$$

At the moment, the explicit formula of the Dunkl translation operator is known only in two cases. One is when $f(x) = \tilde{f}(|x|)$ is a continuous radial function in $L_\kappa^2(\mathbb{R}^n)$, the Dunkl translation operator is represented by (see [14, 43])

$$\tau_x f(y) = V_\kappa[\tilde{f}(\sqrt{|y|^2 + |x|^2 - 2\langle y, \cdot \rangle})](x) = \int_{\mathbb{R}^n} \tilde{f}(\sqrt{|y|^2 + |x|^2 - 2\langle y, \xi \rangle}) d\rho_x^\kappa(\xi).$$

The Dunkl translation operator can also be extended to the space of tempered distributions $\mathcal{S}'(\mathbb{R}^n)$. It turns out to be rather difficult to extend important results in classical Fourier analysis to the setting of the Dunkl transform in general. One of the difficulties comes from the fact that the Dunkl translation operator is not positive in general. In fact, even $L_\kappa^p(\mathbb{R}^n)$ -boundedness of τ_x is not established in general. On the other hand, when f is a radial function in $L_\kappa^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, the following holds (see [25, 51])

$$\|\tau_x f\|_{L_\kappa^p(\mathbb{R}^n)} \leq \|f\|_{L_\kappa^p(\mathbb{R}^n)}.$$

Using the Dunkl translation operator, we define the Dunkl convolution of functions $f, g \in \mathcal{S}(\mathbb{R}^n)$ by

$$f *_\kappa g(x) = \int_{\mathbb{R}^n} \tau_x f(-y)g(y)h_\kappa(y)dy, \quad x \in \mathbb{R}^n.$$

The Dunkl convolution satisfies the following properties (see [51]).

- (1) $\mathcal{F}_\kappa(f *_\kappa g) = \mathcal{F}_\kappa(f)\mathcal{F}_\kappa(g)$ and $f *_\kappa g = g *_\kappa f$.
- (2) Young's inequality: Let $1 \leq p, q, r \leq \infty$ such that $1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}$. If $f \in L_\kappa^p(\mathbb{R}^n)$ and g is a radial function of $L_\kappa^q(\mathbb{R}^n)$, then $f *_\kappa g \in L_\kappa^r(\mathbb{R}^n)$ and we have

$$\|f *_\kappa g\|_{L_\kappa^r(\mathbb{R}^n)} \leq \|f\|_{L_\kappa^p(\mathbb{R}^n)} \|g\|_{L_\kappa^q(\mathbb{R}^n)}. \quad (2.5)$$

2.4. Dunkl Besov spaces. In this subsection, we recall the definition of the Dunkl Besov spaces using Littlewood-Paley decomposition related to the Dunkl Laplacian Δ_κ (see [33, 37]). Dunkl-Besov spaces will be a convenient tool to prove many of our results.

Fix a non-increasing smooth function R on $[0, +\infty)$ such that $R = 1$ in $[0, 1/2]$ and $R = 0$ in $[1, +\infty)$. Let $\psi(r) = R(r/2) - R(r)$ and $0 \leq \psi \leq 1$ such that $\text{supp } \psi \subseteq C_0 = [1/2, 2]$. Moreover, we observe that

$$\begin{aligned} R(r) + \sum_{j=0}^{\infty} \psi(2^{-j}r) &= 1, \quad r \geq 0, \\ \sum_{j \in \mathbb{Z}} \psi(2^{-j}r) &= 1, \quad r > 0. \end{aligned}$$

For $u \in \mathcal{S}'(\mathbb{R}^n)$, we define the Littlewood-Paley decomposition related to the Dunkl Laplacian by $S_0 u = R(\sqrt{-\Delta_\kappa})u = \mathcal{F}_\kappa^{-1}(R(|\cdot|)\mathcal{F}_\kappa u)$ and $\tilde{\Delta}_j u = \psi(2^{-j}\sqrt{-\Delta_\kappa})u = \mathcal{F}_\kappa^{-1}(\psi(2^{-j}|\cdot|)\mathcal{F}_\kappa u)$, for all $j \in \mathbb{Z}$. Moreover, we have

$$u = S_0 u + \sum_{j=0}^{\infty} \tilde{\Delta}_j u \in \mathcal{S}'(\mathbb{R}^n).$$

Moreover, we also have

$$S_0 u = S_0 \left(\sum_{j \in \mathbb{Z}} \tilde{\Delta}_j \right) u = \sum_{j \leq 0} S_0 \tilde{\Delta}_j u.$$

In addition, from Bernstein's inequalities [33], for any $\sigma \in \mathbb{R}$, $j \in \mathbb{Z}$, $1 \leq p, q \leq \infty$ and $u \in \mathcal{S}'(\mathbb{R}^n)$, we have

$$\left\| (-\Delta_\kappa)^{\frac{\sigma}{2}} \tilde{\Delta}_j u \right\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim 2^{j\sigma} \|\tilde{\Delta}_j u\|_{L_\kappa^p(\mathbb{R}^n)}, \quad (2.6)$$

$$\left\| \tilde{\Delta}_j u \right\|_{L_\kappa^q(\mathbb{R}^n)} \lesssim 2^{jN\left(\frac{1}{p}-\frac{1}{q}\right)} \|\tilde{\Delta}_j u\|_{L_\kappa^p(\mathbb{R}^n)}. \quad (2.7)$$

For $1 \leq p, q \leq \infty$ and $s \in \mathbb{R}$, the Dunkl Besov space $B_{p,q}^{s,\kappa}$ is defined as the set of tempered distributions $u \in \mathcal{S}'(\mathbb{R}^n)$ such that $\|u\|_{B_{p,q}^{s,\kappa}} < \infty$ with

$$\|u\|_{B_{p,q}^{s,\kappa}} = \|S_0 u\|_{L_\kappa^p(\mathbb{R}^n)} + \begin{cases} \left(\sum_{j=0}^{\infty} 2^{jsq} \|\tilde{\Delta}_j u\|_{L_\kappa^q(\mathbb{R}^n)}^q \right)^{\frac{1}{q}}, & 1 \leq q < \infty \\ \sup_{j \geq 0} 2^{js} \|\tilde{\Delta}_j u\|_{L_\kappa^p(\mathbb{R}^n)}, & q = \infty. \end{cases}$$

Also, for $1 \leq p \leq \infty$ and $s \in \mathbb{R}$, the Dunkl Sobolev space $H_p^{s,\kappa}(\mathbb{R}^n)$ is defined as the set of tempered distributions $u \in \mathcal{S}'(\mathbb{R}^n)$ such that $\mathcal{F}_\kappa^{-1} \left((1 + |\cdot|^2)^{\frac{s}{2}} \mathcal{F}_\kappa u \right) \in L_\kappa^p(\mathbb{R}^n)$ with the norm

$$\|u\|_{H_p^{s,\kappa}(\mathbb{R}^n)} = \|\mathcal{F}_\kappa^{-1} \left((1 + |\cdot|^2)^{\frac{s}{2}} \mathcal{F}_\kappa u \right)\|_{L_\kappa^p(\mathbb{R}^n)}.$$

Throughout the paper, we denote $H^{s,\kappa}(\mathbb{R}^n)$ for $H_2^{s,\kappa}(\mathbb{R}^n)$. Moreover, we have the equivalent property that

$$u \in H^{s,\kappa}(\mathbb{R}^n) \Leftrightarrow u \in B_{2,2}^{s,\kappa}(\mathbb{R}^n).$$

Now we list some important properties related to the Besov space $B_{p,q}^{s,\kappa}(\mathbb{R}^n)$ in the following proposition.

Proposition 2.1. [33] *Let $p, q \in [1, \infty]$ and $s \in \mathbb{R}$. Then*

- (1) *The space $B_{p,q}^{s,\kappa}(\mathbb{R}^n)$ is a Banach space with the norm $\|\cdot\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)}$.*
- (2) *The definition of $B_{p,q}^{s,\kappa}(\mathbb{R}^n)$ does not depend on the choice of the function R in the Littlewood-Paley decomposition.*
- (3) *The dual space of $B_{p,q}^{s,\kappa}(\mathbb{R}^n)$ is $B_{p',q'}^{-s,\kappa}(\mathbb{R}^n)$;*
- (4) *For any $\sigma > 0$, we have the continuous inclusion $B_{p,q}^{s+\sigma,\kappa}(\mathbb{R}^n) \subseteq B_{p,q}^{s,\kappa}(\mathbb{R}^n)$.*
- (5) *Let $u \in \mathcal{S}'(\mathbb{R}^n)$ and $\sigma \in \mathbb{R}$, then $u \in B_{p,q}^{s,\kappa}(\mathbb{R}^n)$ if and only if $(-\Delta_\kappa)^{\sigma/2} u \in B_{p,q}^{s-\sigma,\kappa}(\mathbb{R}^n)$.*
- (6) *For any $1 \leq p_1 \leq p_2 \leq \infty$, $1 \leq q_1 \leq q_2 \leq \infty$, and $s_1 \leq s_2 \in \mathbb{R}$ such that $s_1 - \frac{N}{p_1} = s_2 - \frac{N}{p_2}$, we also have the continuous inclusion $B_{p_1,q_1}^{s_1,\kappa}(\mathbb{R}^n) \subseteq B_{p_2,q_2}^{s_2,\kappa}(\mathbb{R}^n)$.*
- (7) *For all $p \in [2, \infty]$, we have the continuous inclusion $B_{p,2}^{0,\kappa}(\mathbb{R}^n) \subseteq L_\kappa^p(\mathbb{R}^n)$.*
- (8) $B_{2,2}^{0,\kappa}(\mathbb{R}^n) = L_\kappa^2(\mathbb{R}^n)$.

2.5. Technical Lemmas. In this subsection, we recall several important properties to prove the main result of this paper. From the Dunkl transform and Young's inequality, the $L_\kappa^1 - L_\kappa^\infty$ decay estimates are reduced to a list of oscillatory integrals related to Bessel functions. In order to estimate the oscillatory integrals, we recall the stationary phase lemmas and the properties of Bessel functions. We first recall the van der Corput lemma.

Lemma 2.2 (Van der Corput lemma [49]). *Let $g \in C^\infty([a, b])$ be real-valued such that $|g''(x)| \geq \delta$ for any $x \in [a, b]$ with $\delta > 0$. Then for any function $\psi \in C^\infty([a, b])$, there exists a constant C (does not depend on δ, a, b, g or ψ) such that*

$$\left| \int_a^b e^{ig(x)} \psi(x) dx \right| \leq C \delta^{-1/2} (\|\psi\|_\infty + \|\psi'\|_1).$$

Let J_ν be the Bessel function of order $\nu > -\frac{1}{2}$, defined as

$$J_\nu(r) = \frac{\left(\frac{r}{2}\right)^\nu}{\Gamma(\nu + \frac{1}{2})\pi^{\frac{1}{2}}} \int_{-1}^1 e^{ir\tau} (1 - \tau^2)^{\nu - \frac{1}{2}} d\tau.$$

We list some properties of J_ν in the following lemma.

Lemma 2.3 (see [26]). *For $r > 0$ and $\nu > -\frac{1}{2}$, we have*

$$(1) J_\nu(r) \leq C_\nu r^\nu, \quad (2.8)$$

$$(2) \frac{d}{dr} (r^{-\nu} J_\nu(r)) = -r^{-\nu} J_{\nu+1}(r), \quad (2.9)$$

$$(3) J_\nu(r) \leq C_\nu r^{-\frac{1}{2}}, \quad (2.10)$$

$$(4) J_\nu(r) = \frac{\left(\frac{r}{2}\right)^\nu}{\Gamma(\nu + \frac{1}{2})\Gamma(\frac{1}{2})} \left[ie^{-ir} \int_0^\infty e^{-rt} (t^2 + 2it)^{\nu-\frac{1}{2}} dt - ie^{ir} \int_0^\infty e^{-rt} (t^2 - 2it)^{\nu-\frac{1}{2}} dt \right]. \quad (2.11)$$

Remark 2.4. From (2.8) and (2.9) of Lemma 2.3, we can easily obtain that for any $0 \leq s \leq 2$ and $\beta \in \mathbb{N}_0$,

$$\left| \frac{d^\beta}{dr^\beta} \left(\psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} \right) \right| \leq C_\beta, \quad (2.12)$$

where ψ is the function in Section 2.4. Again from (2.11), we have the identity

$$\frac{J_{\frac{N-2}{2}}(r)}{r^{\frac{N-2}{2}}} = C \left(e^{ir} h(r) + e^{-ir} \overline{h(r)} \right), \quad (2.13)$$

where

$$h(r) = -i \int_0^\infty e^{-rt} (t^2 - 2it)^{\frac{N-3}{2}} dt.$$

Moreover, integrating by parts (if necessary) and using Van der Corput's Lemma, for any $\beta \in \mathbb{N}_0$, one can get

$$\left| \frac{d^\beta}{dr^\beta} h(r) \right| \leq C_\beta (1+r)^{-\frac{N-1}{2}-\beta}, \quad (2.14)$$

for all $\beta \in \mathbb{N}_0$. Thus, from (2.14), for any $s > 2$ and $\beta \in \mathbb{N}_0$, we have

$$\left| \frac{d^\beta}{dr^\beta} (\psi(r) h(rs) r^{N-1}) \right| \leq C_\beta s^{-\frac{N-1}{2}}. \quad (2.15)$$

We also exploit the following estimates, which can be easily proved by comparing the sums with the corresponding integrals.

Lemma 2.5. *Fix $\beta > 0$. There exists $C_\beta > 0$ such that for any $A > 0$, we have*

$$\begin{aligned} \sum_{j \in \mathbb{Z}, 2^j \leq A} 2^{j\beta} &\leq C_\beta A^\beta, \\ \sum_{j \in \mathbb{Z}, 2^j > A} 2^{-j\beta} &\leq C_\beta A^{-\beta}. \end{aligned}$$

Finally, we apply the following duality arguments (see [24]) to prove the Strichartz estimates for some concrete equations related to the Dunkl Laplacian.

Lemma 2.6. *Let H be a Hilbert space, X be a Banach space, X^* is the dual of X , and D be a vector space densely contained in X . Let $A \in \mathcal{L}_a(D, H)$ and $A^* \in \mathcal{L}_a(H, D_a^*)$ be its adjoint operator, defined by*

$$\langle A^*v, f \rangle_D = \langle v, Af \rangle_H, \quad \forall f \in D, \quad \forall v \in H,$$

where $\mathcal{L}_a(Y, Z)$ is the space of linear maps from a vector space Y to a vector space Z , D_a^* is the algebraic dual of D , $\langle \varphi, f \rangle_D$ is the pairing between D_a^* and D , and $\langle \cdot, \cdot \rangle_H$ is the scalar product in H . Then the following three conditions are equivalent:

(1) *There exists $0 \leq a \leq \infty$ such that for all $f \in D$,*

$$\|Af\| \leq a \|f\|_X.$$

(2) $\Re(A^*) \subset X^*$ and there exists $0 \leq a \leq \infty$, such that for all $v \in H$,

$$\|A^*v\|_{X^*} \leq a\|v\|.$$

(3) $\Re(A^*A) \subset X^*$, and there exists a , $0 \leq a \leq \infty$, such that for all $f \in D$,

$$\|A^*Af\|_{X^*} \leq a^2\|f\|_X,$$

where $\|\cdot\|$ denote the norm in H .

The constant a is the same in all three parts. If one of those conditions is satisfied, the operators A and A^*A extend by continuity to bounded operators from X to H and from X to X^* , respectively.

Lemma 2.7. Let H , D and two triplets (X_i, A_i, a_i) , $i = 1, 2$, satisfy any of the conditions of Lemma 2.6. Then for all choices of $i, j = 1, 2$, $\Re(A_i^*A_j) \subset X_i^*$, and for all $f \in D$, we have

$$\|A_i^*A_jf\|_{X_i^*} \leq a_i a_j \|f\|_{X_j}.$$

Lemma 2.8. Let H be a Hilbert space, I be an interval of \mathbb{R} , $X \subset \mathcal{S}'(I \times \mathbb{R}^n)$ be a Banach space which is stable under time restriction, and let X and A satisfy (any of) the conditions of Lemma 2.6. Then the operator A^*A is a bounded operator from $L_t^1(I, H)$ to X^* and from X to $L_t^\infty(I, H)$.

3. PROOF OF MAIN RESULTS

This section is devoted to proving Theorem 1.1, i.e., decay estimate for a class of dispersive semigroup $e^{it\phi(\sqrt{-\Delta_\kappa})}$ on \mathbb{R}^n .

Proof of Theorem 1.1. In order to prove part (A) first, we note that

$$U_t \tilde{\Delta}_j u_0 = \mathcal{F}_\kappa^{-1} \left(e^{it\phi(|\cdot|)} \psi(2^{-j}|\cdot|) \mathcal{F}_\kappa u_0 \right) = \mathcal{F}_\kappa^{-1} \left(e^{it\phi(|\cdot|)} \psi(2^{-j}|\cdot|) \right) *_\kappa u_0. \quad (3.1)$$

It follows from Young's inequality (2.5) that

$$\|U_t \tilde{\Delta}_j u_0\|_{L_\kappa^\infty(\mathbb{R}^n)} \leq \|\mathcal{F}_\kappa^{-1} \left(e^{it\phi(|\cdot|)} \psi(2^{-j}|\cdot|) \right)\|_{L_\kappa^\infty(\mathbb{R}^n)} \|u_0\|_{L_\kappa^1(\mathbb{R}^n)}.$$

Now our aim is to estimate $I_j(x) = \mathcal{F}_\kappa^{-1} \left(e^{it\phi(|\cdot|)} \psi(2^{-j}|\cdot|) \right) (x)$. From the Dunkl transform formula of a radial function (2.4), we have

$$\begin{aligned} I_j(x) &= \mathcal{F}_\kappa^{-1} \left(e^{it\phi(|\cdot|)} \psi(2^{-j}|\cdot|) \right) (x) \\ &= \frac{1}{\Gamma(N/2)} \int_0^\infty e^{it\phi(r)} \psi(2^{-j}r) \frac{J_{\frac{N-2}{2}}(r|x|)}{(r|x|)^{\frac{N-2}{2}}} r^{N-1} dr \\ &= \frac{2^{jN}}{\Gamma(N/2)} \int_0^\infty e^{it\phi(2^j r)} \psi(r) \frac{J_{\frac{N-2}{2}}(2^j r|x|)}{(2^j r|x|)^{\frac{N-2}{2}}} r^{N-1} dr \\ &= II_j(2^j s), \end{aligned} \quad (3.2)$$

where $s = |x|$ and

$$II_j(s) = \frac{2^{jN}}{\Gamma(N/2)} \int_0^\infty e^{it\phi(2^j r)} \psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} dr.$$

We will continue the discussion in two cases. First, we use the vanishing property at the origin (2.8) and the recurring property for the Bessel functions (2.9). On the other hand, we apply the decay property of the Bessel function (2.13).

Case 1. $s \leq 2$: By denoting $D_r = \frac{1}{i2^j t \phi'(2^j r)} \frac{d}{dr}$, we notice that

$$D_r \left(e^{it\phi(2^j r)} \right) = e^{it\phi(2^j r)}, \quad D_r^* f = \frac{i}{2^j t} \frac{d}{dr} \left(\frac{1}{\phi'(2^j r)} f \right).$$

For $\beta \in \mathbb{N}$ and $r \in [1/2, 2]$, it follows from the hypothesis (C1) that

$$\frac{d^\beta}{dr^\beta} \left(\frac{1}{\phi'(2^j r)} \right) \leq C_\beta 2^{-j(m_1-1)}. \quad (3.3)$$

Using integration by part, we have

$$\begin{aligned} II_j(s) &= \frac{2^{jN}}{\Gamma(N/2)} \int_0^\infty e^{it\phi(2^j r)} \psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} dr \\ &= \frac{2^{jN}}{\Gamma(N/2)} \int_0^\infty D_r \left(e^{it\phi(2^j r)} \right) \psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} dr \\ &= \frac{2^{jN}}{\Gamma(N/2)} \cdot \frac{i}{2jt} \int_0^\infty e^{it\phi(2^j r)} \frac{d}{dr} \left(\frac{1}{\phi'(2^j r)} \psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} \right) dr \\ &= \frac{2^{jN}}{\Gamma(N/2)} \cdot \left(\frac{i}{2jt} \right)^Q \sum_{\ell=0}^Q \sum_{(\beta_1, \beta_2, \dots, \beta_Q) \in \Lambda_Q^\ell} C_{Q, \ell} \int_0^\infty e^{it\phi(2^j r)} \\ &\quad \times \prod_{i=1}^Q \frac{d^{\beta_i}}{dr^{\beta_i}} \left(\frac{1}{\phi'(2^j r)} \right) \frac{d^{Q-\ell}}{dr^{Q-\ell}} \left(\psi(r) \frac{J_{\frac{N-2}{2}}(rs)}{(rs)^{\frac{N-2}{2}}} r^{N-1} \right) dr, \end{aligned}$$

for any $Q \in \mathbb{N}$, where $\Lambda_Q^\ell = \{(\beta_1, \beta_2, \dots, \beta_Q) \in \{0, 1, 2, \dots, \ell\}^Q : \sum_{i=1}^Q \beta_i = \ell\}$. Now using the recurring property for the Bessel functions (2.9), the vanishing property at the origin (2.8), and the estimate (3.3), we have

$$|II_j(s)| \leq C_{Q, N, k} |t|^{-Q} 2^{j(N-m_1Q)}, \quad (3.4)$$

for any $Q \in \mathbb{N}$. Using Riesz-Thorin interpolation between different $Q \in \mathbb{N}$ in (3.5), for any $\theta \geq 0$, we have

$$|II_j(s)| \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)}. \quad (3.5)$$

This completes the proof of (A) in this case.

Case 2. $s > 2$. In this case, we shall apply the decay property of Bessel functions. From the identity (2.13), we can write

$$\begin{aligned} II_j(s) &= C2^{jN} \int_0^\infty e^{it\phi(2^j r)} \psi(r) \left(e^{irs} h(rs) + e^{-irs} \overline{h(rs)} \right) r^{N-1} dr \\ &= C2^{jN} \int_0^\infty e^{i(t\phi(2^j r)+rs)} \psi(r) h(rs) r^{N-1} dr \\ &\quad + C2^{jN} \int_0^\infty e^{i(t\phi(2^j r)-rs)} \psi(r) \overline{h(rs)} r^{N-1} dr \\ &= II_j^+(s) + II_j^-(s), \end{aligned}$$

where

$$\begin{aligned} II_j^+(s) &= C2^{jN} \int_0^\infty e^{i(t\phi(2^j r)+rs)} \psi(r) h(rs) r^{N-1} dr = C2^{jN} \int_0^\infty e^{it\phi_+(2^j r)} \psi(r) h(rs) r^{N-1} dr, \\ II_j^-(s) &= C2^{jN} \int_0^\infty e^{i(t\phi(2^j r)-rs)} \psi(r) \overline{h(rs)} r^{N-1} dr = C2^{jN} \int_0^\infty e^{it\phi_-(2^j r)} \psi(r) \overline{h(rs)} r^{N-1} dr, \end{aligned}$$

with $\phi_\pm(r) = \phi(r) \pm \frac{rs}{2jt}$. Without loss of generality, we can assume that $t > 0$ and $\phi'(r) > 0$. Now we will estimate $II_j^+(s)$ and $II_j^-(s)$ separately.

For $II_j^+(s)$, note that $\phi'_+(r) \geq \phi'(r)$ and (3.3) also holds true if we replace ϕ by ϕ_+ . Then, proceeding similarly as **Case 1**, combined with (2.15), for any $\theta \geq 0$, we obtain

$$|II_j^+(s)| \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)}. \quad (3.6)$$

For $II_j^-(s)$, $\phi'_-(2^j r) = \phi'(2^j r) - \frac{s}{2^j t}$. Noticing that if $s = 2^j t \phi'(2^j r)$, then $\phi'_-(2^j r) = 0$. We continue our discussion by considering the following three sub-cases:

- When $s \geq 2 \sup_{r \in [1/2, 2]} 2^j t \phi'(2^j r)$:

In this case, we have $|\phi'_-(2^j r)| = \frac{1}{2^j t} (s - 2^j t \phi'(2^j r)) \geq \phi'(2^j r)$ and also the estimate (3.3) hold true if we replace ϕ by ϕ_- . Then, analogous to $II_j^+(s)$, for any $\theta \geq 0$, we obtain

$$|II_j^-(s)| \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)}. \quad (3.7)$$

- When $s \leq \frac{1}{2} \inf_{r \in [1/2, 2]} 2^j t \phi'(2^j r)$:

In this case, we have $\phi'_-(2^j r) = \frac{1}{2^j t} (2^j t \phi'(2^j r) - s) \geq \frac{1}{2} \phi'(2^j r)$ and (3.3) also hold true if we replace ϕ by ϕ_- . Analogous to $II_j^+(s)$, for any $\theta \geq 0$, we obtain

$$|II_j^-(s)| \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)}. \quad (3.8)$$

- When $\frac{1}{2} \inf_{r \in [1/2, 2]} 2^j t \phi'(2^j r) \leq s \leq 2 \sup_{r \in [1/2, 2]} 2^j t \phi'(2^j r)$:

In this case, we see that $s \sim t 2^{j m_1}$. Moreover, from (2.15), it follows that

$$\begin{aligned} |II_j^-(s)| &\lesssim 2^{jN} s^{-(N-1)/2} \\ &\lesssim |t|^{-(N-1)/2} 2^{j(N-m_1(N-1)/2)}. \end{aligned} \quad (3.9)$$

Now using the estimate (2.14) in $II_j^-(s)$, a simple calculation yields the following easy estimate

$$|II_j^-(s)| \lesssim 2^{jN}. \quad (3.10)$$

Applying interpolation between (3.9) and (3.10), we obtain

$$|II_j^-(s)| \lesssim |t|^{-\theta} 2^{j(N-m_1\theta)} \quad (3.11)$$

for all $0 \leq \theta \leq (N-1)/2$. Combining (3.11) and (3.5), we complete the proof of (1.5) in (A).

If (C3) holds in addition, we have $\left| \frac{d^2}{dr^2} (\phi_-(2^j r)) \right| = 2^{2j} |\phi''(2^j r)| \sim 2^{j\alpha_1}$ in the last sub-case discussed above. It follows from Lemma 2.2 and (2.15) that

$$|II_j^-(s)| \lesssim 2^{jN} |t 2^{j\alpha_1}|^{-1/2} s^{-(N-1)/2} \lesssim |t|^{-N/2} 2^{j(N-m_1 N/2 - (\alpha_1 - m_1)/2)}. \quad (3.12)$$

For any $0 \leq \theta \leq 1$, since $(N-1+\theta)/2 = (1-\theta)(N-1)/2 + \theta N/2$, by an interpolation between (3.11) (with $\theta = (N-1)/2$) and (3.12), we obtain

$$|II_j^-(s)| \lesssim |t|^{-(N-1+\theta)/2} 2^{j(N-m_1(N-1+\theta)/2 - \theta(\alpha_1 - m_1)/2)}. \quad (3.13)$$

Noting that $\alpha_1 \leq m_1$, from (3.5), (3.6), (3.7), (3.8) and (3.13), we have

$$|II_j(s)| \lesssim |t|^{-(N-1+\theta)/2} 2^{j(N-m_1(N-1+\theta)/2 - \theta(\alpha_1 - m_1)/2)}, \quad (3.14)$$

for all $0 \leq \theta \leq 1$. This completes the proof of (1.6) in part (A).

Now we will prove part (C), as the proof of part (B) is similar to part (A). To prove part (C), take any fixed θ satisfying $0 \leq \theta \leq \min\left(\frac{N}{m_2}, \frac{N-1}{2}\right)$. First, we consider the case when

$\theta = \frac{N-1}{2} < \frac{N}{m_2}$, i.e., $N - m_2\theta > 0$. Now applying part (B), we obtain

$$\begin{aligned} \|U_t S_0 u_0\|_{L^\infty_\kappa(\mathbb{R}^n)} &\lesssim \left\| \sum_{j \leq 0} U_t \tilde{\Delta}_j S_0 u_0 \right\|_{L^\infty_\kappa(\mathbb{R}^n)} \\ &\leq \sum_{j \leq 0} \left\| U_t \tilde{\Delta}_j S_0 u_0 \right\|_{L^\infty_\kappa(\mathbb{R}^n)} \\ &\lesssim \sum_{j \leq 0} |t|^{-\theta} 2^{j(N-m_2\theta)} \|S_0 u_0\|_{L^1_\kappa(\mathbb{R}^n)} \\ &\lesssim |t|^{-\theta} \|S_0 u_0\|_{L^1_\kappa(\mathbb{R}^n)} \lesssim |t|^{-\frac{(N-1)}{2}} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}. \end{aligned}$$

Therefore, it suffices to consider the case $\theta = \frac{N}{m_2} \leq \frac{N-1}{2}$. By (3.1) and (3.2), we have

$$U_t S_0 u_0 = \sum_{j \leq 0} U_t \tilde{\Delta}_j (S_0 u_0) = \left(\sum_{j \leq 0} II_j (2^j |\cdot|) \right) *_\kappa (S_0 u_0). \quad (3.15)$$

Argued similarly as (a) but for the proof of (b), we know that if $j_0 < 0$ and $s \sim |t| 2^{j_0 m_2} > 2$, then

$$\begin{aligned} |II_{j_0}(2^{j_0} s)| &\lesssim |t|^{-(N-1)/2} 2^{j_0(N-m_2(N-1)/2)} \\ &\lesssim |t|^{-(N-1)/2} \left| \frac{s}{t} \right|^{(N-m_2(N-1)/2) \frac{1}{m_2}} \\ &\lesssim |t|^{-\frac{N}{m_2}}. \end{aligned} \quad (3.16)$$

When $|j - j_0| > C \gg 1$, it holds true that

$$|II_j(2^j s)| \lesssim |t|^{-\alpha} 2^{j(N-m_2\alpha)}, \quad \forall \alpha \geq 0. \quad (3.17)$$

Hence, by (3.16) and (3.17), we get

$$\begin{aligned} &\sum_{j \leq 0} |II_j(2^j s)| \\ &\leq \left(\sum_{|j-j_0| \leq C} + \sum_{|j-j_0| > C, 2^j \leq |t|^{-\frac{1}{m_2}}} + \sum_{|j-j_0| > C, 2^j > |t|^{-\frac{1}{m_2}}} \right) |II_j(2^j s)| \\ &\lesssim |t|^{-\frac{N}{m_2}} + \sum_{2^j \leq |t|^{-\frac{1}{m_2}}} 2^{jN} + \sum_{2^j > |t|^{-\frac{1}{m_2}}} |t|^{-\alpha} 2^{j(N-m_2\alpha)} \\ &\lesssim |t|^{-\frac{N}{m_2}}, \end{aligned} \quad (3.18)$$

where the last inequality is obtained by applying Lemma 2.5 and choosing $\alpha > \frac{N}{m_2}$. By Young's inequality (2.5), combined with (3.15) and (3.18), we obtain

$$\begin{aligned} \|U_t S_0 u_0\|_{L^\infty_\kappa(\mathbb{R}^n)} &\leq \left\| \sum_{j \leq 0} II_j(2^j |\cdot|) \right\|_{L^\infty_\kappa(\mathbb{R}^n)} \|S_0 u_0\|_{L^1_\kappa(\mathbb{R}^n)} \\ &\lesssim |t|^{-\frac{N}{m_2}} \|S_0 u_0\|_{L^1_\kappa(\mathbb{R}^n)} \lesssim |t|^{-\frac{N}{m_2}} \|u_0\|_{L^1_\kappa(\mathbb{R}^n)}, \end{aligned}$$

which proves (1.7). In addition, if (C4) holds, we can argue similarly as previously. \square

4. APPLICATIONS

In this section, we provide several applications of Theorem 1.1 by considering some particular smooth function ϕ . In particular, we prove Strichartz estimates for some concrete wave equations related to the Dunkl Laplacian by using Theorem 1.1. Above all, we introduce the general Strichartz estimates for $U_t = e^{it\phi(\sqrt{-\Delta_k})}$. We start this section by recalling the following versions of the Hardy-Littlewood-Sobolev inequality:

Lemma 4.1 ([27]). *Assume $\theta_1, \theta_2 \in \mathbb{R}$. Let*

$$k(y) = \begin{cases} |y|^{-\theta_1}, & |y| \leq 1, \\ |y|^{-\theta_2}, & |y| > 1. \end{cases}$$

Assume that one of the following condition holds:

- (1) $0 < \theta_1 = \theta_2 < n$, $1 < p < q < \infty$ and $1 - \frac{1}{p} + \frac{1}{q} = \frac{\theta_1}{n}$,
- (2) $\theta_1 < \theta_2$, $0 < \theta_1 < n$, $1 < p < q < \infty$ and $1 - \frac{1}{p} + \frac{1}{q} = \frac{\theta_1}{n}$,
- (3) $\theta_1 < \theta_2$, $0 < \theta_2 < n$, $1 < p < q < \infty$ and $1 - \frac{1}{p} + \frac{1}{q} = \frac{\theta_2}{n}$,
- (4) $\theta_1 < \theta_2$, $1 \leq p \leq q \leq \infty$ and $\frac{\theta_1}{n} < 1 - \frac{1}{p} + \frac{1}{q} < \frac{\theta_2}{n}$.

We have

$$\left\| \int_{\mathbb{R}^n} f(\cdot - y)k(y)dy \right\|_{L^q(\mathbb{R}^n)} \lesssim \|f\|_{L^p(\mathbb{R}^n)}.$$

Definition 4.2. For $\theta_1, \theta_2 \in \mathbb{R}$ with $\theta_1 < \theta_2$, we say $r \in E(\theta_1, \theta_2)$ if one of the following conditions holds:

- (1) $0 < \theta_1 = \theta_2 < 1$ and $r = \frac{2}{\theta_1}$,
- (2) $\theta_1 < \theta_2$, $0 < \theta_1 < 1$ and $r = \frac{2}{\theta_1}$,
- (3) $\theta_1 < \theta_2$, $0 < \theta_2 < 1$ and $r = \frac{2}{\theta_2}$,
- (4) $\theta_1 < \theta_2$, $2 \leq r \leq \infty$ and $\theta_1 < \frac{2}{r} < \theta_2$.

For $2 \leq p \leq \infty$, $s = s(p) \in \mathbb{R}$ and $\theta_1 \leq \theta_2$, we assume that

$$\|U_t f\|_{B_{p,2}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t)\|f\|_{B_{p',2}^{-s,\kappa}(\mathbb{R}^n)}, \quad \text{where } k(t) = \begin{cases} |t|^{-\theta_1}, & |t| \leq 1, \\ |t|^{-\theta_2}, & |t| > 1. \end{cases} \quad (4.1)$$

Once we obtain this dispersive estimate with different decay rates between $|t| > 1$ and $|t| \leq 1$, we can prove the following proposition by using Lemma 4.1 and standard duality arguments in Lemma 2.6-2.8.

Proposition 4.3 (see [27, 34]). *Assume U_t satisfies (4.1). Then for $r \in E(\theta_1, \theta_2)$ and $T > 0$, we have*

$$\begin{aligned} & \|U_t f\|_{L^r(\mathbb{R}, B_{p,2}^{s,\kappa}(\mathbb{R}^n))} \lesssim \|f\|_{L_\kappa^2(\mathbb{R}^n)}, \\ & \left\| \int_0^t U_{t-\tau} g(\tau, \cdot) d\tau \right\|_{L^r([-T,T], B_{p,2}^{s,\kappa}(\mathbb{R}^n))} \lesssim \|g\|_{L^{r'}([-T,T], B_{p',2}^{-s,\kappa}(\mathbb{R}^n))}, \\ & \left\| \int_0^t U_{t-\tau} g(\tau, \cdot) d\tau \right\|_{L^\infty([-T,T], H^{s,\kappa}(\mathbb{R}^n))} \lesssim \|g\|_{L^{r'}([-T,T], B_{p',2}^{-s,\kappa}(\mathbb{R}^n))}, \\ & \left\| \int_0^t U_{t-\tau} g(\tau, \cdot) d\tau \right\|_{L^r([-T,T], B_{p,2}^{s,\kappa}(\mathbb{R}^n))} \lesssim \|g\|_{L^1([-T,T], L_\kappa^2(\mathbb{R}^n))}. \end{aligned}$$

Note that the endpoint case $\theta_i = 1$ also holds true and can be seen from Keel-Tao [34].

4.1. **Wave equation.** First, we consider the Dunkl-wave equation as

$$\begin{cases} \partial_t^2 u - \Delta_\kappa u = g, \\ u|_{t=0} = u_0, \\ \partial_t u|_{t=0} = u_1. \end{cases} \quad (4.2)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \frac{d\mathcal{W}_t}{dt} u_0 + \mathcal{W}_t u_1 - \int_0^t \mathcal{W}_{t-\tau} g(\tau, \cdot) d\tau,$$

where

$$\mathcal{W}_t = \frac{\sin(t\sqrt{-\Delta_\kappa})}{\sqrt{-\Delta_\kappa}}, \quad \frac{d\mathcal{W}_t}{dt} = \cos(t\sqrt{-\Delta_\kappa}).$$

This reduces to the semigroup $W_t = e^{it\sqrt{-\Delta_\kappa}}$, which corresponds to the case $\phi(r) = r$. Thus ϕ satisfies (C1) and (C2) with $m_1 = m_2 = 1$. Using Theorem 1.1, we obtain the following decay estimate for the wave operator.

Proposition 4.4. *Assume $2 \leq p \leq \infty$, $1 \leq q \leq \infty$, $\delta = \frac{1}{2} - \frac{1}{p}$ and $2s \leq -(N+1)\delta$, then we have*

$$\|W_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-2\max(s+N\delta,0)}, & |t| \leq 1, \\ |t|^{-(N-1)\delta}, & |t| \geq 1. \end{cases}$$

Proof. First, we prove the case $|t| \geq 1$. It follows from (1.7) of Theorem 1.1 by taking $\theta = \frac{N-1}{2}$ and Plancherel's identity that

$$\begin{aligned} \|W_t S_0 f\|_{L_\kappa^\infty(\mathbb{R}^n)} &\lesssim |t|^{-\frac{N-1}{2}} \|S_0 f\|_{L_\kappa^1(\mathbb{R}^n)}, \\ \|W_t S_0 f\|_{L_\kappa^2(\mathbb{R}^n)} &\leq \|S_0 f\|_{L_\kappa^2(\mathbb{R}^n)}. \end{aligned}$$

From (1.5) of Theorem 1.1 by taking $\theta = \frac{N-1}{2}$ and Plancherel's identity, for $j \geq 0$, we have

$$\begin{aligned} \|W_t \tilde{\Delta}_j f\|_{L_\kappa^\infty(\mathbb{R}^n)} &\lesssim |t|^{-\frac{N-1}{2}} 2^{j\frac{N+1}{2}} \|\tilde{\Delta}_j f\|_{L_\kappa^1(\mathbb{R}^n)}, \\ \|W_t \tilde{\Delta}_j f\|_{L_\kappa^2(\mathbb{R}^n)} &\leq \|\tilde{\Delta}_j f\|_{L_\kappa^2(\mathbb{R}^n)}. \end{aligned}$$

Applying the Riesz-Thorin interpolation theorem, for any $j \geq 0$, we have

$$\begin{aligned} \|W_t S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1)\delta} \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}, \\ \|W_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1)\delta} 2^{j(N+1)\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)} \leq |t|^{-(N-1)\delta} 2^{-2js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}, \end{aligned}$$

where in the last inequality we used the fact that $2s \leq -(N+1)\delta$ and $\delta = \frac{1}{2} - \frac{1}{p}$. Therefore

$$\|W_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-(N-1)\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad |t| \geq 1. \quad (4.3)$$

Next, we consider the case $|t| \leq 1$. Putting $\theta \geq 0$ in (1.7) of Theorem 1.1 and applying Plancherel's identity and Riesz-Thorin interpolation theorem, we get

$$\|W_t S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

Again for $j \geq 0$, from (1.5) of Theorem 1.1 with Plancherel's identity and Riesz-Thorin interpolation theorem, we obtain

$$\|W_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(N-\theta)\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)},$$

for $0 \leq \theta \leq \frac{N-1}{2}$. Therefore

$$2^{js} \|W_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(s+(N-\theta)\delta)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

If $-N\delta \leq s \leq -\frac{(N+1)\delta}{2}$, then we can choose $0 \leq \theta \leq \frac{N-1}{2}$ such that $s + N\delta = \theta\delta$. Furthermore, for $s < -N\delta$, we will choose $\theta = 0$. Combing all the scenarios, we have

$$2^{js} \|W_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\max(s+N\delta,0)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}$$

and hence

$$\|W_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-2\max(s+N\delta,0)} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad (4.4)$$

for $|t| \leq 1$. Thus from (4.3) and (4.4), we finally have

$$\|W_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-2\max(s+N\delta,0)}, & |t| \leq 1, \\ |t|^{-(N-1)\delta}, & |t| \geq 1. \end{cases}$$

□

Now using Proposition 4.3, we obtain the following estimates for the Dunkl-wave operator.

Theorem 4.5. *Let $N > 1$. For $i = 1, 2$, let $p_i, r_i \in [2, \infty]$ and $s_i \in \mathbb{R}$ such that*

$$(i) \quad \frac{2}{r_i} + \frac{N-1}{p_i} \leq \frac{N-1}{2},$$

$$(ii) \quad s_i \leq -\frac{N+1}{2} \left(\frac{1}{2} - \frac{1}{p_i} \right),$$

except for $(N, p_i, r_i) = (3, \infty, 2)$. Then for any $T > 0$, the Dunkl-wave operator W_t satisfies

$$\|W_t u_0\|_{L^{r_1}(\mathbb{R}, B_{p_1,2}^{s_1,\kappa}(\mathbb{R}^n))} \lesssim \|u_0\|_{L_\kappa^2(\mathbb{R}^n)},$$

$$\left\| \int_0^t W_{t-\tau} g(\tau, \cdot) d\tau \right\|_{L^{r_1}([-T,T], B_{p_1,2}^{s_1,\kappa}(\mathbb{R}^n))} \lesssim \|g\|_{L^{r'_2}([-T,T], B_{p_2,2}^{-s_2,\kappa}(\mathbb{R}^n))}.$$

It immediately reduces the Strichartz inequality for the solution of the Dunkl-wave equation (4.2).

Corollary 4.6. *Under the same hypotheses as in Theorem 4.5, the solution u of the Dunkl-wave equation (4.2) satisfies the following estimate*

$$\|u\|_{L^{r_1}([-T,T], B_{p_1,2}^{s_1+1,\kappa}(\mathbb{R}^n))} \lesssim \|u_0\|_{H^{1,\kappa}(\mathbb{R}^n)} + \|u_1\|_{L_\kappa^2(\mathbb{R}^n)} + \|g\|_{L^{r'_2}([-T,T], B_{p_2,2}^{-s_2,\kappa}(\mathbb{R}^n))}.$$

Finally, by the embedding relation between the Besov space and Lebesgue space in Proposition 2.1, we have the following Strichartz estimates on Lebesgue spaces.

Corollary 4.7. *Let $N > 1$. Suppose $r_i \geq 2$ and $2 \leq p_i < \infty$ such that*

$$\frac{1}{r_1} + \frac{N}{p_1} = \frac{N}{2} - 1, \quad \frac{2}{r_1} + \frac{N-1}{p_1} \leq \frac{N-1}{2} \quad \text{and} \quad \frac{1}{r_2} + \frac{N}{p_2} \leq \frac{N}{2},$$

except for $(N, p_1, r_1, p_2, r_2) = (3, \infty, 2, 3, 2)$. If u is the solution of the Dunkl-wave equation (4.2), for any $T > 0$, we have the following estimate

$$\|u\|_{L^{r_1}([-T,T], L_\kappa^{p_1}(\mathbb{R}^n))} \lesssim \|u_0\|_{H^{1,\kappa}(\mathbb{R}^n)} + \|u_1\|_{L_\kappa^2(\mathbb{R}^n)} + \|g\|_{L^{r'_2}([-T,T], L_\kappa^{p_2'}(\mathbb{R}^n))}.$$

Proof. From Corollary 4.6, it follows that

$$\|u\|_{L^{r_1}([-T,T], B_{\tilde{p}_1,2}^{s_1+1,\kappa}(\mathbb{R}^n))} \lesssim \|u_0\|_{H^{1,\kappa}(\mathbb{R}^n)} + \|u_1\|_{L_\kappa^2(\mathbb{R}^n)} + \|g\|_{L^{r'_2}([-T,T], B_{\tilde{p}_2,2}^{-s_2,\kappa}(\mathbb{R}^n))}$$

where $\tilde{p}_i, r_i \in [2, \infty]$ and

$$\frac{2}{r_1} + \frac{N-1}{\tilde{p}_1} = \frac{N-1}{2}, \quad \frac{2}{r_2} + \frac{N-1}{\tilde{p}_2} \leq \frac{N-1}{2}, \quad \text{and} \quad s_i = -\frac{N+1}{2} \left(\frac{1}{2} - \frac{1}{\tilde{p}_i} \right), \quad i = 1, 2,$$

except for $(N, \tilde{p}_i, r_i) = (3, \infty, 2)$. If $p_i, r_i \in [2, \infty], i = 1, 2$ with the sole exception of the pair $(N, p_1, r_1, p_2, r_2) = (3, \infty, 2, 3, 2)$ satisfies the condition

$$\frac{1}{r_1} + \frac{N}{p_1} = \frac{N}{2} - 1, \quad \frac{2}{r_1} + \frac{N-1}{p_1} \leq \frac{N-1}{2} \quad \text{and} \quad \frac{1}{r_2} + \frac{N}{p_2} \leq \frac{N}{2},$$

then we the following relations

$$s_1 + 1 \leq 0, \quad s_1 + 1 - \frac{N}{\tilde{p}_1} = 0 - \frac{N}{p_1},$$

and

$$-s_2 \geq 0, \quad -s_2 - \frac{N}{\tilde{p}_2} = 0 - \frac{N}{p_2}.$$

Therefore, using the embedding relation between the Besov space and Lebesgue space in Proposition 2.1, we obtain $B_{\tilde{p}_1, 2}^{s_1+1, \kappa}(\mathbb{R}^n) \subseteq L_{\kappa}^{p_1}(\mathbb{R}^n)$ and $L_{\kappa}^{p_2}(\mathbb{R}^n) \subseteq B_{\tilde{p}_2, 2}^{-s_2, \kappa}(\mathbb{R}^n)$ and this completes our desired result. \square

4.2. Fractional Schrödinger equation. Consider the fractional Dunkl-Schrödinger equation ($0 < \mu \leq 2$ and $\mu \neq 1$)

$$\begin{cases} i\partial_t u + (-\Delta_{\kappa})^{\frac{\mu}{2}} u = g, \\ u|_{t=0} = u_0. \end{cases} \quad (4.5)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \mathcal{S}_t^{\mu} u_0 - i \int_0^t \mathcal{S}_{t-\tau}^{\mu} g(\tau, \cdot) d\tau,$$

where $\mathcal{S}_t^{\mu} = e^{it(-\Delta_{\kappa})^{\frac{\mu}{2}}}$ and it corresponds to the case when $\phi(r) = r^{\mu}$. By a simple calculation, we see that

$$\phi'(r) = \mu r^{\mu-1}, \quad \phi''(r) = \mu(\mu-1)r^{\mu-2}.$$

This shows that ϕ satisfies (C1)-(C4) with $m_1 = \alpha_1 = m_2 = \alpha_2 = \mu$. Using Theorem 1.1, we obtain the following decay estimate for \mathcal{S}_t^{μ} .

Proposition 4.8. *Let $0 < \mu \leq 2$ and $\mu \neq 1$. Assume $2 \leq p \leq \infty, 1 \leq q \leq \infty, \delta = \frac{1}{2} - \frac{1}{p}$ and $s \leq -N(1 - \frac{\mu}{2})\delta$, then we have*

$$\|\mathcal{S}_t^{\mu} f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-\frac{2}{\mu} \max(s+N\delta, 0)}, & |t| \leq 1, \\ |t|^{-N\delta}, & |t| \geq 1. \end{cases}$$

Proof. First, we prove the case $|t| \geq 1$. From (1.8) of Theorem 1.1 with $\theta = \frac{N}{2}$, Plancherel's identity yields

$$\begin{aligned} \|\mathcal{S}_t^{\mu} S_0 f\|_{L_{\kappa}^{\infty}(\mathbb{R}^n)} &\lesssim |t|^{-\frac{N}{2}} \|S_0 f\|_{L_{\kappa}^1(\mathbb{R}^n)}, \\ \|\mathcal{S}_t^{\mu} S_0 f\|_{L_{\kappa}^2(\mathbb{R}^n)} &\leq \|S_0 f\|_{L_{\kappa}^2(\mathbb{R}^n)}. \end{aligned}$$

Again from (1.6) of Theorem 1.1 by taking $\theta = 1$, for $j \geq 0$, Plancherel's identity gives

$$\begin{aligned} \|\mathcal{S}_t^{\mu} \tilde{\Delta}_j f\|_{L_{\kappa}^{\infty}(\mathbb{R}^n)} &\lesssim |t|^{-\frac{N}{2}} 2^{jN(1-\frac{\mu}{2})} \|\tilde{\Delta}_j f\|_{L_{\kappa}^1(\mathbb{R}^n)}, \\ \|\mathcal{S}_t^{\mu} \tilde{\Delta}_j f\|_{L_{\kappa}^2(\mathbb{R}^n)} &\leq \|\tilde{\Delta}_j f\|_{L_{\kappa}^2(\mathbb{R}^n)}. \end{aligned}$$

Now applying the Riesz-Thorin interpolation theorem, for any $j \geq 0$, we obtain

$$\begin{aligned} \|\mathcal{S}_t^{\mu} S_0 f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-N\delta} \|S_0 f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}, \\ \|\mathcal{S}_t^{\mu} \tilde{\Delta}_j f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-N\delta} 2^{2jN(1-\frac{\mu}{2})\delta} \|\tilde{\Delta}_j f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)} \leq |t|^{-N\delta} 2^{-2js} \|\tilde{\Delta}_j f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}. \end{aligned}$$

This implies that

$$\|\mathcal{S}_t^\mu f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-N\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad |t| \geq 1.$$

Next, we discuss the case $|t| \leq 1$. For $\theta = 0$ in (1.8) of Theorem 1.1, applying $\theta = 0$, Plancherel's identity and Riesz-Thorin interpolation theorem, we get

$$\|\mathcal{S}_t^\mu S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

For $j \geq 0$, it follows from (1.5) and (1.6) of Theorem 1.1, Plancherel's identity and Riesz-Thorin interpolation theorem that for $0 \leq \theta \leq \frac{N}{2}$, we have

$$\|\mathcal{S}_t^\mu \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(N-\theta\mu)\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

This implies that

$$2^{js} \|\mathcal{S}_t^\mu \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(s+(N-\theta\mu)\delta)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

If $-N\delta \leq s \leq -N(1 - \frac{\mu}{2})\delta$, then we can choose $0 \leq \theta \leq \frac{N}{2}$ such that $s + N\delta = \mu\theta\delta$. If $s < -N\delta$, we take $\theta = 0$. Therefore

$$2^{js} \|\mathcal{S}_t^\mu \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-\frac{2}{\mu} \max(s+N\delta, 0)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}$$

and

$$\|\mathcal{S}_t^\mu f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-\frac{2}{\mu} \max(s+N\delta, 0)} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad |t| \leq 1,$$

which completes the proof of the proposition. \square

By Proposition 4.3, we obtain the following estimates for the fractional Schrödinger operator.

Theorem 4.9. *Let $0 < \mu \leq 2$ and $\mu \neq 1$. For $i = 1, 2$, let $p_i, r_i \in [2, \infty]$ and $s_i \in \mathbb{R}$ such that*

$$(i) \quad \frac{2}{r_i} + \frac{N}{p_i} \leq \frac{N}{2},$$

$$(ii) \quad s_i \leq -N \left(1 - \frac{\mu}{2}\right) \left(\frac{1}{2} - \frac{1}{p_i}\right),$$

except for $(N, p_i, r_i) = (2, \infty, 2)$. Then for any $T > 0$, the fractional Schrödinger operator \mathcal{S}_t^μ satisfies the estimates

$$\|\mathcal{S}_t^\mu u_0\|_{L^{r_1}(\mathbb{R}, B_{p_1,2}^{s_1,\kappa}(\mathbb{R}^n))} \lesssim \|u_0\|_{L_\kappa^2(\mathbb{R}^n)}, \quad (4.6)$$

$$\left\| \int_0^t \mathcal{S}_{t-\tau}^\mu g(\tau, \cdot) d\tau \right\|_{L^{r_1}([-T, T], B_{p_1,2}^{s_1,\kappa}(\mathbb{R}^n))} \lesssim \|g\|_{L^{r_2}'([-T, T], B_{p_2,2}^{-s_2,\kappa}(\mathbb{R}^n))}. \quad (4.7)$$

The above estimate immediately reduces the following Strichartz inequality for the solution of the fractional Schrödinger equation.

Corollary 4.10. *Under the same hypotheses as in Theorem 4.9, the solution u of the fractional Schrödinger equation (4.5) satisfies the following estimate*

$$\|u\|_{L^{r_1}([-T, T], B_{p_1,2}^{s_1,\kappa}(\mathbb{R}^n))} \lesssim \|u_0\|_{L_\kappa^2(\mathbb{R}^n)} + \|g\|_{L^{r_2}'([-T, T], B_{p_2,2}^{-s_2,\kappa}(\mathbb{R}^n))}.$$

Finally, by the embedding relation between the Besov space and Lebesgue space in Proposition 2.1, we have the Strichartz inequalities on Lebesgue spaces.

Corollary 4.11. *Let $0 < \mu < 2$ and $\mu \neq 1$. Suppose $p_i, r_i \geq 2$ such that*

$$\frac{1}{p_i} \geq \frac{1}{2} - \frac{2}{(2-\mu)N} \quad \text{and} \quad \frac{\mu}{r_i} + \frac{N}{p_i} \leq \frac{N}{2} - 1.$$

If u is the solution of the fractional Schrödinger equation (4.5), for any $T > 0$, we have the following estimate

$$\|u\|_{L^{r_1}([-T, T], L_{\kappa}^{p_1}(\mathbb{R}^n))} \lesssim \|u_0\|_{H^{1, \kappa}(\mathbb{R}^n)} + \|g\|_{L^{r'_2}([-T, T], L_{\kappa}^{p'_2}(\mathbb{R}^n))}.$$

In particular, when $\mu = 2$, fractional Schrödinger operator reduces to the Dunkl-Schrödinger equation

$$\begin{cases} i\partial_t u + \Delta_{\kappa} u = g, \\ u|_{t=0} = u_0. \end{cases} \quad (4.8)$$

By the embedding relation between the Besov space and Lebesgue space in Proposition 2.1 along with Theorem 4.9, we obtain the Strichartz inequalities on Lebesgue space for the Dunkl-Schrödinger equation considered in [32, 38, 39].

Corollary 4.12. *Suppose for $j = 1, 2$, $r_i, p_i \geq 2$, $(N, p_i, r_i) \neq (2, \infty, 2)$ and*

$$\frac{2}{r_i} + \frac{N}{p_i} \leq \frac{N}{2}.$$

If u is the solution of the Dunkl-Schrödinger equation (4.8), then for any $T > 0$, we have the following estimate

$$\|u\|_{L^{r_1}([-T, T], L_{\kappa}^{p_1}(\mathbb{R}^n))} \lesssim \|u_0\|_{L_{\kappa}^2(\mathbb{R}^n)} + \|g\|_{L^{r'_2}([-T, T], L_{\kappa}^{p'_2}(\mathbb{R}^n))}.$$

4.3. Fourth-order Schrödinger equation. Consider the fourth-order Schrödinger equation

$$\begin{cases} i\partial_t u + \Delta_{\kappa}^2 u - \Delta_{\kappa} u = g, \\ u|_{t=0} = u_0. \end{cases} \quad (4.9)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \mathcal{U}_t u_0 - i \int_0^t \mathcal{U}_{t-\tau} g(\tau, \cdot) d\tau,$$

where $\mathcal{U}_t = e^{it(\Delta_{\kappa}^2 - \Delta_{\kappa})}$ and it corresponds to the case when $\phi(r) = r^4 + r^2$. By a simple calculation, we see that

$$\phi'(r) = 4r^3 + 2r, \quad \phi''(r) = 12r^2 + 2.$$

Thus ϕ satisfies (C1)–(C4) with $m_1 = \alpha_1 = 4$, $m_2 = \alpha_2 = 2$. Using Theorem 1.1, we obtain the following decay estimate for \mathcal{U}_t .

Proposition 4.13. *Assume $2 \leq p \leq \infty$, $1 \leq q \leq \infty$, $\delta = \frac{1}{2} - \frac{1}{p}$ and $s \leq N\delta$.*

Then we have

$$\|\mathcal{U}_t f\|_{B_{p, q}^{s, \kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p', q}^{-s, \kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-\frac{1}{2} \max(s + N\delta, 0)}, & |t| \leq 1, \\ |t|^{-N\delta}, & |t| \geq 1. \end{cases}$$

Proof. First, we prove the case $|t| \geq 1$. For $\theta = \frac{N}{2}$ in (1.8) of Theorem 1.1, Plancherel's identity gives

$$\begin{aligned} \|\mathcal{U}_t S_0 f\|_{L_{\kappa}^{\infty}(\mathbb{R}^n)} &\lesssim |t|^{-\frac{N}{2}} \|S_0 f\|_{L_{\kappa}^1(\mathbb{R}^n)}, \\ \|\mathcal{U}_t S_0 f\|_{L_{\kappa}^2(\mathbb{R}^n)} &\leq \|S_0 f\|_{L_{\kappa}^2(\mathbb{R}^n)}. \end{aligned}$$

Applying Riesz-Thorin interpolation theorem, we have

$$\|\mathcal{U}_t S_0 f\|_{L_{\kappa}^p(\mathbb{R}^n)} \lesssim |t|^{-N\delta} \|S_0 f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}. \quad (4.10)$$

Again for $j \geq 0$, it follows from (1.5) and (1.6) of Theorem 1.1 that

$$\|\mathcal{U}_t \tilde{\Delta}_j f\|_{L_{\kappa}^{\infty}(\mathbb{R}^n)} \lesssim |t|^{-\theta} 2^{j(N-4\theta)} \|\tilde{\Delta}_j f\|_{L_{\kappa}^1(\mathbb{R}^n)}, \quad 0 \leq \theta \leq \frac{N}{2}. \quad (4.11)$$

Taking $\theta = \frac{N}{2}$ in (4.11), combined with Plancherel's identity and Riesz-Thorin interpolation theorem, it indicates

$$\|\mathcal{U}_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-N\delta} 2^{-2jN\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)} \leq |t|^{-N\delta} 2^{-2js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}}, \quad (4.12)$$

for any $j \geq 0$. From (4.10) and (4.12), we obtain our desired result of the proposition in this case.

Next, we discuss the case $|t| \leq 1$. For $\theta = 0$ in (1.7) of Theorem 1.1, Plancherel's identity and Riesz-Thorin interpolation theorem gives

$$\|\mathcal{U}_t S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

Again for $j \geq 0$, from (4.11), applying Plancherel's identity and Riesz-Thorin interpolation theorem, for $0 \leq \theta \leq \frac{N}{2}$, we obtain

$$\|\mathcal{U}_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(N-4\theta)\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)},$$

which reduces to

$$2^{js} \|\mathcal{U}_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(s+(N-4\theta)\delta)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

If $-N\delta \leq s \leq N\delta$, then we can choose $0 \leq \theta \leq \frac{N}{2}$ such that $s + N\delta = 4\theta\delta$. If $s < -N\delta$, we take $\theta = 0$. Therefore

$$2^{js} \|\mathcal{U}_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-\frac{1}{2} \max(s+N\delta, 0)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)},$$

and hence

$$\|\mathcal{U}_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-\frac{1}{2} \max(s+N\delta, 0)} \|f\|_{B_{p,q}^{-s,\kappa}(\mathbb{R}^n)}, \quad |t| \leq 1.$$

This completes the proof of the proposition. \square

4.4. Klein-Gordon equation. Moreover, we consider the Klein-Gordon equation

$$\begin{cases} \partial_t^2 u - \Delta_\kappa u + u = g, \\ u|_{t=0} = u_0, \\ \partial_t u|_{t=0} = u_1. \end{cases} \quad (4.13)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \frac{d\mathcal{K}_t}{dt} u_0 + \mathcal{K}_t u_1 - \int_0^t \mathcal{K}_{t-\tau} g(\tau, \cdot) d\tau,$$

where

$$\mathcal{K}_t = \frac{\sin(t\sqrt{I - \Delta_\kappa})}{\sqrt{I - \Delta_\kappa}}, \quad \frac{d\mathcal{K}_t}{dt} = \cos(t\sqrt{I - \Delta_\kappa}).$$

So we naturally introduce the operator $K_t = e^{it\sqrt{I - \Delta_\kappa}}$, which corresponds to the case when $\phi(r) = \sqrt{1 + r^2}$. By a simple calculation have

$$\phi'(r) = r(1 + r^2)^{-\frac{1}{2}}, \quad \phi''(r) = (1 + r^2)^{-\frac{3}{2}}.$$

This shows that ϕ satisfies (C1)–(C4) with $m_1 = 1$, $\alpha_1 = -1$, and $m_2 = \alpha_2 = 2$. Using Theorem 1.1, we obtain the following decay estimate for K_t .

Proposition 4.14. *Assume $2 \leq p \leq \infty$, $1 \leq q \leq \infty$, and $\delta = \frac{1}{2} - \frac{1}{p}$.*

(1) *Let $0 \leq \theta \leq 1$ and $2s = -(N + 1 + \theta)\delta$, then we have*

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-(N-1+\theta)\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}.$$

(2) *Let $0 \leq \theta \leq N - 1$ and $2s = -(N + 1 + \theta)\delta$, then we have*

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-(N-1-\theta)\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}.$$

(3) In particular, for $0 \leq \theta \leq 1$ and $2s \leq -(N+1+\theta)\delta$, we have

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-2\max(s+N\delta,0)}, & |t| \leq 1, \\ |t|^{-(N-1+\theta)\delta}, & |t| \geq 1. \end{cases}$$

Proof. First, we prove part (1). From (1.6) and (1.8) of Theorem 1.1 with Plancherel's identity and Riesz-Thorin interpolation theorem, for $0 \leq \theta \leq 1$, we obtain

$$\begin{aligned} \|K_t \tilde{\Delta}_j f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1+\theta)\delta} 2^{j(N+1+\theta)\delta} \|\tilde{\Delta}_j f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}, \quad j \geq 0, \\ \|K_t S_0 f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1+\theta)\delta} \|S_0 f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}. \end{aligned}$$

Hence, we complete the proof of part (1) from the fact that $2s = -(N+1+\theta)\delta$.

Next, we prove part (2). Again from (1.5) and (1.7) of Theorem 1.1 with the help of Plancherel's identity and Riesz-Thorin interpolation theorem, for $0 \leq \theta \leq N-1$, we get

$$\begin{aligned} \|K_t S_0 f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1-\theta)\delta} \|S_0 f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}, \\ \|K_t \tilde{\Delta}_j f\|_{L_{\kappa}^p(\mathbb{R}^n)} &\lesssim |t|^{-(N-1-\theta)\delta} 2^{j(N+1+\theta)\delta} \|\tilde{\Delta}_j f\|_{L_{\kappa}^{p'}(\mathbb{R}^n)}. \end{aligned}$$

From the fact that $2s = -(N+1+\theta)\delta$, we have

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-(N-1-\theta)\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}$$

and this completes the proof of part (2). To prove (3), we divide into the following three cases.

Case 1. $|t| \geq 1$. Following the proof of part (1), we immediately get the required estimates.

Case 2. $|t| \leq 1$ and $s+N\delta \geq 0$. In this case, our required estimate will follow immediately from part (2). Indeed, if $2s \leq -(N+1+\theta)\delta$ for some $0 \leq \theta \leq 1$, then $0 \leq 2(s+N\delta) \leq (N-1-\theta)\delta$, which also provides us $0 \leq \theta \leq N-1$. Therefore from part (2), we obtain

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim |t|^{-(N-1-\theta)\delta} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)} \leq |t|^{-2(s+N\delta)} \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}.$$

Case 3. $|t| \leq 1$ and $s+N\delta < 0$. In this case taking $\theta_1 = N-1$ in (2), for $s_1 = -N\delta > s$, we get

$$\|K_t f\|_{B_{p,q}^{s_1,\kappa}(\mathbb{R}^n)} \lesssim \|f\|_{B_{p',q}^{-s_1,\kappa}(\mathbb{R}^n)}.$$

Since $s_1 > s$, using the inclusion relation (4) between Besov spaces in Proposition 2.1, we immediately obtain

$$\|K_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \|K_t f\|_{B_{p,q}^{s_1,\kappa}(\mathbb{R}^n)} \lesssim \|f\|_{B_{p',q}^{-s_1,\kappa}(\mathbb{R}^n)} \leq \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}$$

and this completes the proof of part (3). \square

4.5. Beam equation. Finally, we consider the beam equation

$$\begin{cases} \partial_t^2 u + \Delta_{\kappa}^2 u + u = g, \\ u|_{t=0} = u_0, \\ \partial_t u|_{t=0} = u_1. \end{cases} \quad (4.14)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \frac{d\mathcal{B}_t}{dt} u_0 + \mathcal{B}_t u_1 - \int_0^t \mathcal{B}_{t-\tau} g(\tau, \cdot) d\tau,$$

where

$$\mathcal{B}_t = \frac{\sin(t\sqrt{I + \Delta_{\kappa}^2})}{\sqrt{I + \Delta_{\kappa}^2}}, \quad \frac{d\mathcal{B}_t}{dt} = \cos(t\sqrt{I + \Delta_{\kappa}^2}).$$

So we naturally introduce the operator $B_t = e^{it\sqrt{I+\Delta_k^2}}$, which corresponds to the case when $\phi(r) = \sqrt{1+r^4}$. By a simple calculation, we see that

$$\phi'(r) = 2r^3(1+r^4)^{-\frac{1}{2}}, \quad \phi''(r) = (6r^2 + 2r^6)(1+r^4)^{-\frac{3}{2}}.$$

This shows that ϕ satisfies (C1)–(C4) with $m_1 = \alpha_1 = 2, m_2 = \alpha_2 = 4$. Using Theorem 1.1, we obtain the following decay estimate for B_t .

Proposition 4.15. *Let $2 \leq p \leq \infty, 1 \leq q \leq \infty, \delta = \frac{1}{2} - \frac{1}{p}$, and $s \leq 0$. Then*

$$\|B_t f\|_{B_{p,q}^{s,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|f\|_{B_{p',q}^{-s,\kappa}(\mathbb{R}^n)}, \quad k(t) = \begin{cases} |t|^{-\max(s+N\delta,0)}, & |t| \leq 1, \\ |t|^{-\frac{N\delta}{2}}, & |t| \geq 1. \end{cases}$$

Proof. First, we prove the case $|t| \geq 1$. For $\theta = \frac{N}{4}$ in (1.8) of Theorem 1.1 with the help of Plancherel's identity and Riesz-Thorin interpolation theorem, we have

$$\|B_t S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-\frac{N\delta}{2}} \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}. \quad (4.15)$$

Again for $j \geq 0$, it follows from (1.5) and (1.6) of Theorem 1.1 that

$$\|B_t \tilde{\Delta}_j f\|_{L_\kappa^\infty(\mathbb{R}^n)} \lesssim |t|^{-\theta} 2^{j(N-2\theta)} \|\tilde{\Delta}_j f\|_{L_\kappa^1(\mathbb{R}^n)}, \quad 0 \leq \theta \leq \frac{N}{2}. \quad (4.16)$$

Taking $\theta = \frac{N}{2}$ in (4.16), combined with Plancherel's identity and Riesz-Thorin interpolation theorem, we obtain

$$\|B_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-N\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)} \leq |t|^{-\frac{N\delta}{2}} 2^{-2js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}, \quad (4.17)$$

for any $j \geq 0$. From (4.15) and (4.17), we get our desired result.

Next, we consider the case $|t| < 1$. By taking $\theta = 0$ in (1.8) of Theorem 1.1 along with Plancherel's identity and Riesz-Thorin interpolation theorem, we have

$$\|B_t S_0 f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim \|S_0 f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

For $j \geq 0$, it follows from (4.16), Plancherel's identity, and Riesz-Thorin interpolation theorem, for $0 \leq \theta \leq \frac{N}{2}$, we have

$$\|B_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(N-2\theta)\delta} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)},$$

and thus

$$2^{js} \|B_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-2\theta\delta} 2^{2j(s+(N-2\theta)\delta)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)}.$$

If $-N\delta \leq s \leq 0$, we can choose $0 \leq \theta \leq \frac{N}{2}$ such that $s + N\delta = 2\theta\delta$. If $s < -N\delta$, we can choose $\theta = 0$. Thus, we have

$$2^{js} \|B_t \tilde{\Delta}_j f\|_{L_\kappa^p(\mathbb{R}^n)} \lesssim |t|^{-\max(s+N\delta,0)} 2^{-js} \|\tilde{\Delta}_j f\|_{L_\kappa^{p'}(\mathbb{R}^n)},$$

which completes the proof of the proposition. \square

5. WELL-POSEDNESS RESULTS FOR NONLINEAR KLEIN-GORDON AND BEAM EQUATIONS

Using the decay estimates for the nonlinear Klein-Gordon and beam equation with the different decay rates between $|t| > 1$ and $|t| \leq 1$ in the previous section, we prove the global well-posedness results for nonlinear problems. First, we consider the Cauchy problem for the nonlinear Klein-Gordon equation

$$\begin{cases} \partial_t^2 u - \Delta_\kappa u + u = F(u), \\ u|_{t=0} = u_0, \\ \partial_t u|_{t=0} = u_1, \end{cases} \quad (5.1)$$

with a power-like nonlinearity $F(u)$ satisfies

$$F(0) = 0, \quad |F(u) - F(v)| \leq C(|u|^\alpha + |v|^\alpha)|u - v| \quad (5.2)$$

for some $\alpha > 0$, where C is a positive constant. By Duhamel's principle, the solution of (5.1) is formally given by

$$u = \frac{d\mathcal{K}_t}{dt}u_0 + \mathcal{K}_t u_1 - \int_0^t \mathcal{K}_{t-\tau} F(u) d\tau,$$

where $\mathcal{K}_t = \frac{\sin(t\sqrt{I-\Delta_\kappa})}{\sqrt{I-\Delta_\kappa}}$. Similarly, as in the Laplacian in Euclidean space, we find the critical index as

$$\alpha_K(N) = \frac{2 - N + \sqrt{N^2 + 12N + 4}}{2N}.$$

Using the classical fixed point argument, we obtain the global well-posedness theorem for the nonlinear Klein-Gordon equation.

Theorem 5.1. *Let $\alpha_K(N) < \alpha < \frac{4}{N}$, $\sigma(\alpha) = \frac{\alpha(N+2)}{2(\alpha+2)}$, and $(u_0, u_1) \in H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n) \times H^{\frac{\alpha+2}{\alpha+1} - 1, \kappa}(\mathbb{R}^n)$ with sufficient small norm. Then the equation (5.1) has a unique solution*

$$u \in C(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n)) \cap L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n)).$$

Proof. By Proposition 2.1 and 4.14, we have

$$\begin{aligned} \left\| \frac{d\mathcal{K}_t}{dt} u_0 \right\|_{L_k^{\alpha+2}(\mathbb{R}^n)} &\leq \left\| \frac{d\mathcal{K}_t}{dt} u_0 \right\|_{B_{\alpha+2,2}^{0,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_0\|_{B_{\frac{\alpha+2}{\alpha+1},2}^{\sigma(\alpha),\kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_0\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)}, \\ \|\mathcal{K}_t u_1\|_{L_k^{\alpha+2}(\mathbb{R}^n)} &\leq \|\mathcal{K}_t u_1\|_{B_{\alpha+2,2}^{0,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_1\|_{B_{\frac{\alpha+2}{\alpha+1},2}^{\sigma(\alpha)-1,\kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_1\|_{H^{\frac{\alpha+2}{\alpha+1} - 1, \kappa}(\mathbb{R}^n)}, \end{aligned} \quad (5.3)$$

where

$$k(t) = \begin{cases} |t|^{-\max(\frac{\alpha(N-2)}{2(\alpha+2)}, 0)}, & |t| \leq 1, \\ |t|^{-\frac{\alpha N}{2(\alpha+2)}}, & |t| \geq 1. \end{cases}$$

For $\alpha_K(N) < \alpha < \frac{4}{N}$, we notice that

$$(\alpha+1) \frac{\alpha(N-2)}{2(\alpha+2)} < 1, \quad (\alpha+1) \frac{\alpha N}{2(\alpha+2)} > 1, \quad \text{and } \sigma(\alpha) < 1.$$

This implies that $k \in L^{\alpha+1}(\mathbb{R})$ and

$$\begin{aligned} \left\| \frac{d\mathcal{K}_t}{dt} u_0 \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} &\leq C \|u_0\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)}, \\ \|\mathcal{K}_t u_1\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} &\leq C \|u_1\|_{H^{\frac{\alpha+2}{\alpha+1} - 1, \kappa}(\mathbb{R}^n)}. \end{aligned}$$

Therefore, from (5.3), Proposition 2.1, and Young's inequality, it follows that

$$\begin{aligned} \left\| \int_0^t \mathcal{K}_{t-\tau} F(u) d\tau \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} &\leq \left\| \int_0^t \|\mathcal{K}_{t-\tau} F(u)\|_{L_k^{\alpha+2}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})} \\ &\leq C \left\| \int_0^t k(t-\tau) \|F(u)\|_{H^{\frac{\alpha+2}{\alpha+1} - 1, \kappa}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})} \\ &\leq C \left\| \int_0^t k(t-\tau) \|F(u)\|_{L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})} \\ &\leq C \|k\|_{L^{\alpha+1}(\mathbb{R})} \|F(u)\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\ &\leq C \| |u|^{\alpha+1} \|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \end{aligned}$$

$$= C \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}^{\alpha+1}.$$

Taking $M = 2C(\|u_0\|_{H^{\frac{\sigma(\alpha), \kappa}{\alpha+2}}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\sigma(\alpha)-1, \kappa}{\alpha+2}}(\mathbb{R}^n)})$, we set the evolution space X as

$$X = \{u \in L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n)) : \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \leq M\}.$$

We say that a function u is a *mild solution* to (5.1) if u is a fixed point for the integral operator

$$\mathcal{T} : u \in X \mapsto \mathcal{T}u(t, x) := \frac{d\mathcal{K}_t}{dt}u_0 + A_t u_1 - \int_0^t \mathcal{K}_{t-\tau} F(u) d\tau.$$

We prove the global-in-time existence and uniqueness of small data Sobolev solutions of low regularity to the Cauchy problem (5.1) by finding a unique fixed point to the operator \mathcal{T} . It means that there exists a unique global solution u to the equation $\mathcal{T}u = u \in X$, which also gives the solution to (5.1). In order to prove that the operator \mathcal{T} has a uniquely determined fixed point, we use Banach's fixed point argument with respect to the norm on X as defined above.

Now for any $u, v \in X$, we have

$$\|\mathcal{T}u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \leq \frac{M}{2} + CM^{\alpha+1},$$

and

$$\begin{aligned} & \|\mathcal{T}u - \mathcal{T}v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\ &= \left\| \int_0^t \mathcal{K}_{t-\tau} (F(u) - F(v)) d\tau \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\ &\leq C \|F(u) - F(v)\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\ &\leq C \left\| (|u|^\alpha + |v|^\alpha) |u - v| \right\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\ &= C \left\| \left\| (|u|^\alpha + |v|^\alpha) |u - v| \right\|_{L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n)} \right\|_{L^1(\mathbb{R})} \\ &\leq C \left\| \left\| |u|^\alpha + |v|^\alpha \right\|_{L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n)} \left\| |u - v| \right\|_{L_k^{\alpha+2}(\mathbb{R}^n)} \right\|_{L^1(\mathbb{R})} \\ &\leq C \left\| |u|^\alpha + |v|^\alpha \right\|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} \left\| |u - v| \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\ &\leq C \left(\left\| |u|^\alpha \right\|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} + \left\| |v|^\alpha \right\|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} \right) \left\| |u - v| \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\ &= C \left(\left\| |u|^\alpha \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} + \left\| |v|^\alpha \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \right) \left\| |u - v| \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\ &\leq 2CM^\alpha \|u - v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}. \end{aligned}$$

Notice that, for $CM^\alpha < \frac{1}{2}$, $\mathcal{T} : X \rightarrow X$ is a contraction mapping. Applying Banach's fixed point theorem, there exists a global small data Sobolev solution u of the equation $u = \mathcal{T}u$ in X , which also gives the solution to the equation (5.1). In addition, by the embedding $H^{\frac{\sigma(\alpha), \kappa}{\alpha+2}}(\mathbb{R}^n) \subseteq H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n)$ and $H^{\frac{\sigma(\alpha)-1, \kappa}{\alpha+2}}(\mathbb{R}^n) \subseteq H^{-\frac{2}{\alpha+2}, \kappa}(\mathbb{R}^n)$, we have

$$\begin{aligned} \|u\|_{L^\infty(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n))} &\leq \left\| \frac{d\mathcal{K}_t}{dt}u_0 \right\|_{L^\infty(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n))} + \|\mathcal{K}_t u_1\|_{L^\infty(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n))} \\ &\quad + \left\| \int_0^t \mathcal{K}_{t-\tau} F(u) d\tau \right\|_{L^\infty(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n))} \\ &\leq \|u_0\|_{H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{-\frac{2}{\alpha+2}, \kappa}(\mathbb{R}^n)} + \|F(u)\|_{L^1(\mathbb{R}, H^{-\frac{2}{\alpha+2}, \kappa}(\mathbb{R}^n))} \end{aligned}$$

$$\begin{aligned}
&\lesssim \|u_0\|_{H^{\frac{\sigma(\alpha)}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\sigma(\alpha)-1}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|F(u)\|_{L^1(\mathbb{R}, H^{\frac{\sigma(\alpha)-1}{\alpha+1}, \kappa})} \\
&\lesssim \|u_0\|_{H^{\frac{\sigma(\alpha), \kappa}{\alpha+1}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\sigma(\alpha)-1, \kappa}{\alpha+1}(\mathbb{R}^n)} + \| |u|^{\alpha+1} \|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&= \|u_0\|_{H^{\frac{\sigma(\alpha), \kappa}{\alpha+1}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\sigma(\alpha)-1, \kappa}{\alpha+1}(\mathbb{R}^n)} + \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))},
\end{aligned}$$

which indicates that the unique solution also belongs to $L^\infty(\mathbb{R}, H^{\frac{\alpha}{\alpha+2}, \kappa}(\mathbb{R}^n))$. \square

Argued similarly as in the nonlinear Klein-Gordon equation (5.1), we also study the Cauchy problem for the nonlinear beam equation

$$\begin{cases} \partial_t^2 u + \Delta_\kappa^2 u + u = F(u), \\ u|_{t=0} = u_0, \\ \partial_t u|_{t=0} = u_1. \end{cases} \quad (5.4)$$

By Duhamel's principle, the solution is formally given by

$$u(t) = \frac{d\mathcal{B}_t}{dt} u_0 + \mathcal{B}_t u_1 - \int_0^t \mathcal{B}_{t-\tau} F(u) d\tau. \quad (5.5)$$

We also find the critical index

$$\alpha_B(N) = \frac{4 - N + \sqrt{N^2 + 24N + 16}}{2N}.$$

The following result is about the global well-posedness for the nonlinear beam equation (5.5).

Theorem 5.2. *Let $\alpha_B(N) < \alpha < \frac{8}{N}$, $\sigma(\alpha) = \frac{\alpha N}{\alpha+2} - \frac{2}{\alpha+1}$, $\sigma(\alpha) < s \leq 2$, $s_0 = s - \frac{\alpha N}{2(\alpha+2)}$ and $(u_0, u_1) \in H^{\frac{s, \kappa}{\alpha+1}}(\mathbb{R}^n) \times H^{\frac{s-2, \kappa}{\alpha+1}}(\mathbb{R}^n)$ with sufficient small norm. Then the equation (5.5) has a unique solution*

$$u \in C(\mathbb{R}, H^{s_0, \kappa}(\mathbb{R}^n)) \cap L^{\alpha+1}(\mathbb{R}, L_\kappa^{\alpha+2}(\mathbb{R}^n)).$$

Proof. From Proposition 2.1 and 4.15, we have

$$\begin{aligned}
\left\| \frac{d\mathcal{B}_t}{dt} u_0 \right\|_{L_k^{\alpha+2}(\mathbb{R}^n)} &\leq \left\| \frac{d\mathcal{B}_t}{dt} u_0 \right\|_{B_{\alpha+2, 2}^{0, \kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_0\|_{B_{\frac{\alpha+2}{\alpha+1}, 2}^{s, \kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_0\|_{H^{\frac{s, \kappa}{\alpha+1}}(\mathbb{R}^n)}, \\
\|\mathcal{B}_t u_1\|_{L_k^{\alpha+2}(\mathbb{R}^n)} &\leq \|\mathcal{B}_t u_1\|_{B_{\alpha+2, 2}^{0, \kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_1\|_{B_{\frac{\alpha+2}{\alpha+1}, 2}^{s-2, \kappa}(\mathbb{R}^n)} \lesssim k(t) \|u_1\|_{H^{\frac{s-2, \kappa}{\alpha+1}}(\mathbb{R}^n)},
\end{aligned} \quad (5.6)$$

where

$$k(t) = \begin{cases} |t|^{-\max(\frac{\alpha N}{2(\alpha+2)} - \frac{s}{2}, 0)}, & |t| \leq 1, \\ |t|^{-\frac{\alpha N}{4(\alpha+2)}}, & |t| \geq 1. \end{cases}$$

For $\alpha_B(N) < \alpha < \frac{8}{N}$ and $\sigma(\alpha) < s \leq 2$, we see that

$$(\alpha + 1) \left(\frac{\alpha N}{2(\alpha + 2)} - \frac{s}{2} \right) < 1 \quad \text{and} \quad (\alpha + 1) \frac{\alpha N}{4(\alpha + 2)} > 1,$$

which implies that $k \in L^{\alpha+1}(\mathbb{R})$ and

$$\begin{aligned}
\left\| \frac{d\mathcal{B}_t}{dt} u_0 \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} &\leq C \|u_0\|_{H^{\frac{s, \kappa}{\alpha+1}}(\mathbb{R}^n)}, \\
\|\mathcal{B}_t u_1\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} &\leq C \|u_1\|_{H^{\frac{s-2, \kappa}{\alpha+1}}(\mathbb{R}^n)}.
\end{aligned}$$

Therefore, from (5.6), Proposition 2.1, and Young's inequality, it follows that

$$\left\| \int_0^t \mathcal{B}_{t-\tau} F(u) d\tau \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \leq \left\| \int_0^t \|\mathcal{B}_{t-\tau} F(u)\|_{L_k^{\alpha+2}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})}$$

$$\begin{aligned}
&\leq C \left\| \int_0^t k(t-\tau) \|F(u)\|_{H^{\frac{s-2,\kappa}{\alpha+1}}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})} \\
&\leq C \left\| \int_0^t k(t-\tau) \|F(u)\|_{L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n)} d\tau \right\|_{L^{\alpha+1}(\mathbb{R})} \\
&\leq C \|k\|_{L^{\alpha+1}(\mathbb{R})} \|F(u)\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&\leq C \| |u|^{\alpha+1} \|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&= C \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}^{\alpha+1}.
\end{aligned}$$

Taking $M = 2C(\|u_0\|_{H^{\frac{s,\kappa}{\alpha+1}}(\mathbb{R}^n)} + \|u_1\|_{H_k^{\frac{s-2,\kappa}{\alpha+1}}(\mathbb{R}^n)})$, we set the evolution spaces

$$X = \{u \in L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n)) : \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \leq M\}.$$

Define the mapping

$$\mathcal{T} : u \mapsto \frac{d\mathcal{B}_t}{dt}u_0 + \mathcal{B}_t u_1 - \int_0^t \mathcal{B}_{t-\tau} F(u) d\tau, \quad \forall u \in X.$$

Then for any $u, v \in X$, we have

$$\|\mathcal{T}u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \leq \frac{M}{2} + CM^{\alpha+1}.$$

Thus

$$\begin{aligned}
&\|\mathcal{T}u - \mathcal{T}v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\
&= \left\| \int_0^t \mathcal{B}_{t-\tau} (F(u) - F(v)) d\tau \right\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\
&\leq C \|F(u) - F(v)\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&\leq C \|(|u|^\alpha + |v|^\alpha) |u - v|\|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&= C \left\| \|(|u|^\alpha + |v|^\alpha) |u - v|\|_{L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n)} \right\|_{L^1(\mathbb{R})} \\
&\leq C \left\| \|(|u|^\alpha + |v|^\alpha)\|_{L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n)} \|u - v\|_{L_k^{\alpha+2}(\mathbb{R}^n)} \right\|_{L^1(\mathbb{R})} \\
&\leq C \| |u|^\alpha + |v|^\alpha \|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} \|u - v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\
&\leq C \left(\| |u|^\alpha \|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} + \| |v|^\alpha \|_{L^{\frac{\alpha+1}{\alpha}}(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha}}(\mathbb{R}^n))} \right) \|u - v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\
&= C \left(\|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}^\alpha + \|v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}^\alpha \right) \|u - v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))} \\
&\leq 2CM^\alpha \|u - v\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}.
\end{aligned}$$

Notice that, for $CM^\alpha < \frac{1}{2}$, $\mathcal{T} : X \rightarrow X$ is a contraction mapping. Therefore, applying Banach's fixed point theorem, the equation (5.4) has a unique solution $u \in X$. In addition, by the embedding $H^{\frac{s,\kappa}{\alpha+1}}(\mathbb{R}^n) \subseteq H^{s_0,\kappa}(\mathbb{R}^n)$ and $H^{\frac{s-2,\kappa}{\alpha+1}}(\mathbb{R}^n) \subseteq H^{s_0-2,\kappa}(\mathbb{R}^n)$, we have

$$\begin{aligned}
&\|u\|_{L^\infty(\mathbb{R}, H^{s_0,\kappa}(\mathbb{R}^n))} \\
&\leq \left\| \frac{d\mathcal{B}_t}{dt}u_0 \right\|_{L^\infty(\mathbb{R}, H^{s_0,\kappa}(\mathbb{R}^n))} + \|\mathcal{B}_t u_1\|_{L^\infty(\mathbb{R}, H^{s_0,\kappa}(\mathbb{R}^n))} + \left\| \int_0^t \mathcal{B}_{t-\tau} F(u) d\tau \right\|_{L^\infty(\mathbb{R}, H^{s_0,\kappa}(\mathbb{R}^n))}
\end{aligned}$$

$$\begin{aligned}
&\leq \|u_0\|_{H^{s_0, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{s_0-2, \kappa}(\mathbb{R}^n)} + \|F(u)\|_{L^1(\mathbb{R}, H^{s_0-2, \kappa})} \\
&\lesssim \|u_0\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|F(u)\|_{L^1(\mathbb{R}, H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n))} \\
&\lesssim \|u_0\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \| |u|^{\alpha+1} \|_{L^1(\mathbb{R}, L_k^{\frac{\alpha+2}{\alpha+1}}(\mathbb{R}^n))} \\
&= \|u_0\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|u_1\|_{H^{\frac{\alpha+2}{\alpha+1}, \kappa}(\mathbb{R}^n)} + \|u\|_{L^{\alpha+1}(\mathbb{R}, L_k^{\alpha+2}(\mathbb{R}^n))}^\alpha,
\end{aligned}$$

which indicates that the unique solution also belongs to $L^\infty(\mathbb{R}, H^{s_0, \kappa}(\mathbb{R}^n))$. \square

STATEMENTS AND DECLARATIONS

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

Competing Interests: No potential competing of interest was reported by the author.

ACKNOWLEDGMENTS

C. Luo and M. Song are supported by the National Natural Science Foundation of China (Grant No. 11701452) and Guangdong Basic and Applied Basic Research Foundation (No. 2023A1515010656). S.S. Mondal is supported by the DST-INSPIRE Faculty Fellowship (Grant No. DST/INSPIRE/04/2023/002038).

REFERENCES

- [1] J. -P. Anker and V. Pierfelice, *Nonlinear Schrödinger equation on real hyperbolic spaces*, Ann. Inst. H. Poincaré (C) Non Linear Anal. **26**(5), 1853-1869 (2009). [2]
- [2] J. -P. Anker and V. Pierfelice, *Wave and Klein-Gordon equations on hyperbolic spaces*, Anal. PDE **7**, 953-995 (2014). [2]
- [3] J. -P. Anker, V. Pierfelice, and M. Vallarino, *The wave equation on hyperbolic spaces*, J. Differ. Equ. **252**(10), 5613-5661 (2012). [2]
- [4] H. Bahouri, D. Barilari, and I. Gallagher, *Strichartz estimates and Fourier restriction theorems on the Heisenberg group*, J. Fourier Anal. Appl. **27**(2) (2021). [2]
- [5] H. Bahouri, P. Gérard, and C. J. Xu, *Espaces de Besov et estimations de Strichartz généralisées sur le groupe de Heisenberg*, J. Anal. Math. **82**, 93-118 (2000). [2]
- [6] H. Bahouri, C. F. Kammerer, and I. Gallagher, *Dispersive estimates for the Schrödinger operator on step 2 stratified Lie groups*, Anal. PDE **9**, 545-574 (2016). [2]
- [7] H. Bahouri, J.-Y. Chemin, and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehrender Math. Wissenschaften 343, Springer, Heidelberg, 2011. [2]
- [8] S. Ben Saïd, *Strichartz estimates for Schrödinger-Laguerre operators*, Semigroup Forum **90**, 251-269 (2015). [2]
- [9] S. Ben Saïd, T. Kobayashi, and B. Orsted, *Laguerre semigroup and Dunkl operators*, Compos. Math. **148**, 1265-1336 (2012). [5]
- [10] S. Ben Saïd, A. K. Nandakumaran, and P. K. Ratnakumar, *Schrödinger propagator and the Dunkl Laplacian*, hal-00578446v1, (2011). [2, 5]
- [11] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear evolution equations I*, Geom. Funct. Anal. **3**, 107-156 (1993). [2]

- [12] T. A. Bui, P. D’Ancona, X. T. Duong, and D. Müller, *On the flows associated to self-adjoint operators on metric measure spaces*, Math. Ann. **375**, 1393-1426 (2019). [2]
- [13] N. Burq, P. Gérard, and N. Tzvetkov, *Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds*, Amer. J. Math. **126**, 569-605 (2004). [2]
- [14] F. Dai and H. Wang, *A transference theorem for the Dunkl transform and its applications*, J. Funct. Anal. **258** (Issue 12), 4052-4074 (2010). [8]
- [15] F. Dai and W. Ye, *Local restriction theorem and maximal Bochner-Riesz operators for the Dunkl transforms*, Trans. Amer. Math. Soc. **371**, 641–679 (2019). [7]
- [16] P. D’Ancona, V. Pierfelice, and F. Ricci, *On the wave equation associated to the Hermite and the twisted Laplacian*, J. Fourier Anal. Appl. **16**(2), 294-310 (2010). [2]
- [17] C. F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc. **311**, 167-183 (1989). [5]
- [18] C. F. Dunkl, *Integral kernels with reflection group invariance*, Canad. J. Math. **43**, 1213-1227 (1991). [3, 5]
- [19] C. F. Dunkl, *Hankel transforms associated to finite reflection groups*, Contemp. Math. **138**, 123-138 (1992). [7]
- [20] C. F. Dunkl and Y. Xu, *Orthogonal polynomials of several variables*(second edition), Cambridge University Press, 2014. [3]
- [21] G. Feng, M. Song, and H. Wu, *Decay estimates for a class of semigroups related to self-adjoint operators on metric measure spaces*, arXiv: 2308. 00388 (2023). [2]
- [22] G. Furioli, C. Melzi, and A. Veneruso, *Strichartz inequalities for the wave equation with the full Laplacian on the Heisenberg group*, Canad. J. Math. **59**(6), 1301-1322 (2007). [2]
- [23] G. Furioli and A. Veneruso, *Strichartz inequalities for the Schrödinger equation with the full Laplacian on the Heisenberg group*, Studia Math. **160**, 157-178 (2004). [2]
- [24] J. Ginibre and G. Velo, *Generalized Strichartz inequalities for the wave equation*, J. Funct. Anal. **133**, 50-68 (1995). [2, 3, 10]
- [25] D. V. Gorbachev, V. I. Ivanov, and S. Y. Tikhonov, *Positive L^p -bounded Dunkl-type generalized translation operator and its applications*, Constr. Approx. **49**, 555–605 (2019). [8]
- [26] L. Grafakos, *Classical Fourier Analysis*, 2nd ed., Grad. Texts in Math. **249**, Springer, New York, 2008. [10]
- [27] Z. Guo, L. Peng, and B. Wang, *Decay estimates for a class of wave equations*, J. Funct. Anal. **254**(6), 1642-1660 (2008). [3, 4, 15]
- [28] M. D. Hierro, *Dispersive and Strichartz estimates on H -type groups*, Studia Math. **169**, 1-20 (2005). [2]
- [29] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Stud. Adv. Math. 29, Cambridge University Press, Cambridge, 1990. [5]
- [30] O. Ivanovici, G. Lebeau, and F. Planchon, *Dispersion for the wave equation inside strictly convex domains I: the Friedlander model case*, Ann. Math. **180**, 323-380 (2014). [2]
- [31] M. F. E. de Jeu, *The Dunkl transform*, Invent. Math. **113**, 147–162 (1993). [7]

- [32] S. P. Jitendra Kumar, B. Pradeep, S. S. Mondal, and H. Mejjaoli, *Restriction theorem for Fourier-Dunkl transform II: Paraboloid, sphere, and hyperboloid surfaces*, J. Geom. Anal. **34**, Paper No. 74, 36pp (2024). [4, 20]
- [33] T. Kawazoe T and H. Mejjaoli, *Generalized Besov spaces and their applications*, Tokyo J Math. **35**(2), 297–320 (2012). [8, 9]
- [34] M. Keel and T. Tao, *Endpoints Strichartz estimates*, Amer. J. Math. **120**, 955-980 (1998). [2, 15]
- [35] H. Liu and M. Song, *Strichartz inequalities for the wave equation with the full Laplacian on H -type groups*, Abstr. Appl. Anal., Art. ID 219375, 10 pp (2014). [2]
- [36] H. Mejjaoli, *Strichartz estimates for the Dunkl wave equation and application*, J. Math. Anal. Appl. **346**(1), 41–54 (2008). [2, 4]
- [37] H. Mejjaoli, *Littlewood-Paley decomposition associated for the Dunkl operators and para-product operators*, J. Inequal. Pure Appl. Math. **9** (4), 1-25 (2008). [8]
- [38] H. Mejjaoli, *Dispersion phenomena in Dunkl-Schrödinger equation and applications*, Serdica Math. J. **35**(1), 25–60 (2009). [2, 4, 5, 20]
- [39] H. Majjaoli, *Dunkl-Schrödinger semigroups and applications*, Appl. Anal. **92**(8), 1597-1626 (2013). [2, 20]
- [40] S. S. Mondal and M. Song, *Orthonormal Strichartz inequalities for the (k, a) -generalized Laguerre operator and Dunkl operator*, Israel J. Math. (2023) (In press). [5]
- [41] A. K. Nandakumaran, and P. K. Ratnakumar, *Schrödinger equation and the oscillatory semigroup for the Hermite operator*, J. Funct. Anal. **224**(2), 371-385 (2005). [2]
- [42] P. K. Ratnakumar, *On Schrödinger Propagator for the Special Hermite Operator*, J. Fourier Anal. Appl. **14**(2), 286-300 (2008). [2]
- [43] R. Rösler, *Positivity of Dunkl's intertwining operator*, Duke Math. J. **98**, 445-463 (1999). [5, 6, 8]
- [44] V. K. Sohani, *Strichartz estimates for the Schrödinger propagator for the Laguerre operator*, Proc. Math. Sci. **123**, 525-537 (2013). [2]
- [45] M. Song, *Decay estimates for fractional wave equations on H -type groups*, J. Inequal. Appl. **2016**(246), 12pp (2016). [2]
- [46] M. Song and J. Yang, *Decay estimates for a class of wave equations on the Heisenberg group*, Ann. Mat. Pur. Appl. **202**, 2665–2685(2023). [2]
- [47] N. Song and J. Zhao, *Strichartz estimates on the quaternion Heisenberg group*, Bull. Sci. Math. **138**(2), 293-315 (2014). [2]
- [48] E. M. Stein, *Oscillatory integrals in Fourier analysis*. In: Beijing Lectures in Harmonic Analysis (Beijing, 1984), Annals of Mathematics Studies, vol. 112, Princeton University Press, Princeton, 1986. [3]
- [49] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality and oscillatory integrals*, Princeton Univ. Press, 1993. [9]
- [50] R. S. Strichartz, *Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations*, Duke Math. J. **44**, 705-714 (1977). [2, 3]

- [51] S. Thangavelu and Y. Xu, *Convolution operator and maximal function for Dunkl transform*, J. Anal. Math. **97**, 25-55 (2005). [8]
- [52] S. Thangavelu and Y. Xu, *Riesz transform and Riesz potentials for Dunkl transform*, J. Comput. Appl. Math. **199**, 181-195(2007). [5]
- [53] P. A. Tomas, *A restriction theorem for the Fourier transform*, Bull. Amer. Math. Soc. **81**, 477-478 (1975). [3]
- [54] P. A. Tomas, *Restriction theorems for the Fourier transform*, In: Proc. Symp. Pure Math., XXXV (1979). [3]

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHWESTERN POLYTECHNICAL UNIVERSITY, XI'AN, SHAANXI 710129, CHINA

Email address: caashmale@163.com

STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE KOLKATA, BT ROAD, BARANAGAR, KOLKATA 700108, INDIA

Email address: mondalshyam055@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHWESTERN POLYTECHNICAL UNIVERSITY, XI'AN, SHAANXI 710129, CHINA

Email address: mlsong@nwpu.edu.cn