

Superconducting states in metals with toroidal ordering

V.P.Mineev

Landau Institute for Theoretical Physics, 142432 Chernogolovka, Russia

(Dated: July 10, 2024)

The paper presents a theory of superconducting states in metals with toroidal symmetry.

I. INTRODUCTION

The possibility of a magnetoelectric effect in some antiferromagnetic crystals was first noted in the textbook by L.D. Landau and E.M. Lifshitz "Electrodynamics of Continuous Media" (Gostekhizdat, Moscow, 1958). This phenomenon is associated with a special symmetry of the material, in which the time reversal symmetry R and the inversion symmetry I are violated, although the product RI is still a symmetry operation. The thermodynamic potential of such a solid must contain terms proportional to the product of the first powers of the components of the electric and magnetic fields ($\Phi \sim EH$). I.E. Dzyaloshinskii^{1,2} gave a specific example of this type of antiferromagnet, namely, Cr_2O_3 , where, unlike ordinary two - or more sublattice antiferromagnets, the magnetic moments of Cr are in the same crystallographic unit cell. The point symmetry group of Cr_2O_3 $\mathbf{D}_{3d}(\mathbf{D}_3) = (E, C_3, C_3^2, 3u_2, 3\sigma_d R, 2S_6 R, IR)$ contains the product of time and space inversion, but does not include these operations separately. Soon after, D.N. Astrov³ discovered the magnetoelectric effect in this material.

Some of the magnetoelectric antiferromagnets exhibit optical nonreciprocity, which is the difference in reflectivity of left and right circularly polarized light (the Kerr effect) despite the absence of a net magnetic moment^{4,5}. In Cr_2O_3 the effect was observed experimentally by Krichevstov et al.^{6,7}. New interest in this type of phenomenon arose after spin-flip neutron scattering experiments on the underdoped high-temperature compound $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ ⁸, indicating a violation of time reversal that occurs without loss of translational symmetry. This was followed by the discovery of the Kerr effect on the same material by A. Kapitulnik's group⁹. The experiments were interpreted by K. Varma in terms of the "loop current model" (see recent review¹⁰).

At the same time, these experimental observations raised the problem of describing the superconducting state in materials with broken time reversal and inversion symmetry, but possessing symmetry with respect to the product of these operations, which are now usually called substances with toroidal order¹¹. Several issues related to this problem have been considered in¹²⁻¹⁴ in the framework of a single-band 2D model with d-wave superconducting ordering, and in the framework of a 1D model developed by S. Sumita and Y. Yanase¹⁵. Here we present a more general treatment of superconductivity in antiferromagnetic materials with a Cr_2O_3 -like structure, with magnetic moments located within a single unit cell, so that time and space inversion are broken, but the total magnetic moment is zero and symmetry with respect to the product of time inversion and time reversal operations holds. Several examples of structures of this type have been discussed by J. Orenstein¹⁶ in connection with the possible ordering of magnetic moments created by loop currents in YBCO.

Specific examples of superconductors of this type are currently unknown, although there are metals and semiconductors with toroidal magnetic ordering. See, for example, the paper by H.Amitsuka et al¹⁷ on UNi_4B and O.Fedchenko et al¹⁸ on Mn_2Au . With hope for future experimental discoveries, we develop here a general symmetry approach to the description of materials of this type. This will be done on a specific example of some orthorhombic antiferromagnet (see Fig. 1) with the point symmetry group $\mathbf{D}_{2h}(\mathbf{D}_2)$, which allows us to demonstrate the general structure of the




FIG. 1: The $y = b/2$ plane of elementary cell of an antiferromagnet with point symmetry group $\mathbf{D}_{2h}(\mathbf{D}_2)$.

theory of superconducting states in magnetoelectrics. Examples of other toroidal antiferromagnets with other point symmetry groups are listed in the Appendix.

The paper is organized as follows. The general symmetry properties are described in the next chapter. The third chapter is devoted to the development of the BCS-type theory in toroids. The results are collected in the Conclusion, where prospects for future research are also indicated. The Appendix demonstrates specific features of superconducting toroids with a point symmetry group different from that described in the main text.

II. STRUCTURE OF SUPERCONDUCTING STATES

We will consider 3D metal with orthorhombic crystal structure with point symmetry group $\mathbf{D}_{2h}(\mathbf{D}_2)$ including the following operations

$$E, C_{2x}, C_{2y}, C_{2z}, R\sigma_x, R\sigma_y, R\sigma_z, RI. \quad (1)$$

Each electron energy band consists of sum of even and odd functions of components of the wave vector \mathbf{k}

$$\varepsilon_{\mathbf{k}} = \varepsilon_{\mathbf{k}}^e + \varepsilon_{\mathbf{k}}^o, \quad \varepsilon_{\mathbf{k}}^e = f(k_x^2, k_y^2, k_z^2), \quad \varepsilon_{\mathbf{k}}^o = \gamma \sin k_x a \sin k_y b \sin k_z c \quad (2)$$

invariant in respect of all operations of the point symmetry group $\mathbf{D}_{2h}(\mathbf{D}_2)$. The Hamiltonian commutes with the product of time and space inversion operations RI . This means that to each energy $\varepsilon_{\mathbf{k}}$ corresponds two orthogonal each other spinor eigen-functions $\psi_{\mathbf{k}\alpha}(\mathbf{r})$ and $TI\hat{\psi}_{\mathbf{k}\alpha}(\mathbf{r})$.

The orthorhombic group $D_{2h}(D_2)$ has four one-dimensional representations. The absence symmetry in respect inversion means that all superconducting states consists of sum singlet and triplet states. The corresponding order parameters of superconducting states have the following form

$$\Delta_{\alpha\beta}(\mathbf{k}) = \Delta\Phi_{\alpha\beta}(\mathbf{k}) = \Delta \left[\phi_{\mathbf{k}}^s i\sigma_{\alpha\beta}^y + (\phi_{\mathbf{k}}^t \sigma_{\alpha\gamma}) i\sigma_{\gamma\beta}^y \right]. \quad (3)$$

Here, $\hat{\sigma} = (\hat{\sigma}^x, \hat{\sigma}^y, \hat{\sigma}^z)$ are the Pauli matrices. The functions $\phi_{\mathbf{k}}^s$ and $\phi_{\mathbf{k}}^t$ for all four representations $\Gamma=A, B_3, B_2, B_1$ are presented in the table

Γ	$\phi_{\mathbf{k}}^s$	$\phi_{\mathbf{k}}^t$
A	$a\hat{k}_x^2 + b\hat{k}_y^2 + c\hat{k}_z^2$	$i(\hat{k}_x\hat{x} + \hat{k}_y\hat{y} + \hat{k}_z\hat{z})$
B ₃	$i\hat{k}_y\hat{k}_z$	$\hat{k}_z\hat{y} + \hat{k}_y\hat{z}$
B ₂	$i\hat{k}_x\hat{k}_z$	$\hat{k}_z\hat{x} + \hat{k}_x\hat{z}$
B ₁	$i\hat{k}_x\hat{k}_y$	$\hat{k}_y\hat{x} + \hat{k}_x\hat{y}$

Here it is necessary to keep in mind that each term in each cell may include some amplitude (a real numerical factor or a real function of the squares of the wave vector components). For example, the vector function for the triplet state A is generally equal to $i(f\hat{k}_x\hat{x} + g\hat{k}_y\hat{y} + h\hat{k}_z\hat{z})$.

III. BCS THEORY FOR FERROELECTRIC METALS

The Hamiltonian including interaction of electrons with opposite momenta is

$$H = H_0 + H_{int} = \sum_{\mathbf{k}} (\xi_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^o) a_{\mathbf{k}\alpha}^+ a_{\mathbf{k}\alpha} + \frac{1}{2} \sum_{\mathbf{k}, \mathbf{k}'} V_{\alpha\beta, \lambda\mu}(\mathbf{k}, \mathbf{k}') a_{-\mathbf{k}\alpha}^+ a_{\mathbf{k}\alpha}^+ a_{\mathbf{k}'\lambda} a_{-\mathbf{k}'\mu}, \quad (4)$$

where

$$\xi_{\mathbf{k}} = \varepsilon_{\mathbf{k}}^e - \mu.$$

In the pairing interaction

$$V_{\alpha\beta, \lambda\mu}(\mathbf{k}, \mathbf{k}') = -V_{\Gamma} \Phi_{\alpha\beta}(\mathbf{k}) \Phi_{\lambda\mu}^{\dagger}(\mathbf{k}') \quad (5)$$

was left only term related to irreducible representation Γ corresponding to superconducting state with maximal critical temperature. The normal part of Hamiltonian can be rewritten as

$$H_0 = \frac{1}{2} \sum_{\mathbf{k}} (\xi_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^o) a_{\mathbf{k}\alpha}^+ a_{\mathbf{k}\alpha} - \frac{1}{2} \sum_{\mathbf{k}} (\xi_{-\mathbf{k}} + \varepsilon_{-\mathbf{k}}^o) a_{-\mathbf{k}\alpha} a_{-\mathbf{k}\alpha}^+ + \frac{1}{2} \sum_{\mathbf{k}\alpha} (\xi_{-\mathbf{k}} + \varepsilon_{-\mathbf{k}}^o) \quad (6)$$

After standard transformation in H_{int} according to mean field approach the Hamiltonian acquires the following form

$$H = \frac{1}{2} \sum_{\mathbf{k}} (\xi_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^o) a_{\mathbf{k}\alpha}^+ a_{\mathbf{k}\alpha} - \frac{1}{2} \sum_{\mathbf{k}} (\xi_{-\mathbf{k}} + \varepsilon_{-\mathbf{k}}^o) a_{-\mathbf{k}\alpha} a_{-\mathbf{k}\alpha}^+ + \frac{1}{2} \sum_{\mathbf{k}\alpha} (\xi_{-\mathbf{k}} + \varepsilon_{-\mathbf{k}}^o) + \frac{1}{2} \sum_{\mathbf{k}} \Delta_{\mathbf{k},\alpha\beta} a_{\mathbf{k}\alpha}^+ a_{-\mathbf{k}\beta}^+ + \frac{1}{2} \sum_{\mathbf{k}} \Delta_{\mathbf{k},\alpha\beta}^\dagger a_{-\mathbf{k}\alpha} a_{\mathbf{k}\beta} + \frac{1}{2} \sum_{\mathbf{k}} \Delta_{\mathbf{k},\alpha\beta} F_{\mathbf{k},\beta\alpha}^+ \quad (7)$$

where the matrix of the order parameter

$$\Delta_{\mathbf{k},\alpha\beta} = - \sum_{\mathbf{k}'} V_{\beta\alpha,\lambda\mu}(\mathbf{k}, \mathbf{k}') F_{\mathbf{k}',\lambda\mu} \quad (8)$$

is expressed through "anomalous average" $F_{\mathbf{k},\alpha\beta} = \langle a_{\mathbf{k}\alpha} a_{-\mathbf{k}\beta} \rangle$. Here, $\langle \dots \rangle$ means subsequent quantum mechanical and thermal averaging.

Introducing operators

$$A_{\mathbf{k},i}^+ = (a_{\mathbf{k}\alpha}^+, a_{-\mathbf{k}\alpha}), \quad A_{\mathbf{k},i} = \begin{pmatrix} a_{\mathbf{k}\alpha} \\ a_{-\mathbf{k}\alpha}^+ \end{pmatrix} \quad (9)$$

one can rewrite Eq.(7) in more compact form

$$H = \frac{1}{2} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k},ij} A_{\mathbf{k},i}^+ A_{\mathbf{k},j} + \frac{1}{2} \sum_{\mathbf{k}\alpha} (\xi_{\mathbf{k}} - \varepsilon_{\mathbf{k}}^o) + \frac{1}{2} \sum_{\mathbf{k}} \Delta_{\mathbf{k},\alpha\beta} F_{\mathbf{k},\beta\alpha}^+ \quad (10)$$

Here,

$$\varepsilon_{\mathbf{k},ij} = \begin{pmatrix} (\xi_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^o) \delta_{\alpha\beta} & \Delta_{\mathbf{k},\alpha\beta} \\ \Delta_{\mathbf{k},\alpha\beta}^\dagger & (-\xi_{\mathbf{k}} + \varepsilon_{\mathbf{k}}^o) \delta_{\alpha\beta} \end{pmatrix}. \quad (11)$$

Diagonalising Hamiltonian by means the Bogolubov transformation

$$A_{\mathbf{k},i} = U_{ij} B_{\mathbf{k},j}, \quad U_{ij} = \begin{pmatrix} u_{\mathbf{k},\alpha\beta} & v_{\mathbf{k},\alpha\beta} \\ v_{\mathbf{k},\alpha\beta}^\dagger & -u_{\mathbf{k},\alpha\beta} \end{pmatrix} \quad (12)$$

$$u_{\mathbf{k},\alpha\beta} = \frac{\xi_{\mathbf{k}} + E_{\mathbf{k}}^e}{\sqrt{(\xi_{\mathbf{k}} + E_{\mathbf{k}}^e)^2 + \Delta_{\mathbf{k}}^2}} \delta_{\alpha\beta}, \quad v_{\mathbf{k},\alpha\beta} = \frac{\Delta_{\alpha\beta}(\mathbf{k})}{\sqrt{(\xi_{\mathbf{k}} + E_{\mathbf{k}}^e)^2 + \Delta_{\mathbf{k}}^2}}, \quad (13)$$

$$E_{\mathbf{k}}^e = \sqrt{\xi_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2}, \quad \Delta_{\mathbf{k}}^2 = \frac{1}{2} \text{Tr} \hat{\Delta}^\dagger(\mathbf{k}) \hat{\Delta}(\mathbf{k}) \quad (14)$$

we obtain

$$\frac{1}{2} \sum_{\mathbf{k}} \varepsilon_{\mathbf{k},ij} A_{\mathbf{k},i}^+ A_{\mathbf{k},j} = \frac{1}{2} \sum_{\mathbf{k}} E_{\mathbf{k},ij} B_{\mathbf{k},i}^+ B_{\mathbf{k},j}, \quad (15)$$

where

$$E_{\mathbf{k},ij} = \begin{pmatrix} (\varepsilon_{\mathbf{k}}^o + E_{\mathbf{k}}^e) \delta_{\alpha\beta} & 0 \\ 0 & (\varepsilon_{\mathbf{k}}^o - E_{\mathbf{k}}^e) \delta_{\alpha\beta} \end{pmatrix}. \quad (16)$$

Thus, the energy of excitations is

$$E_{\mathbf{k}} = \varepsilon_{\mathbf{k}}^o + E_{\mathbf{k}}^e. \quad (17)$$

The corresponding density of states

$$N(E) = 2 \int \frac{d^3\mathbf{k}}{(2\pi)^3} \delta(E - E_{\mathbf{k}}) \quad (18)$$

at the Fermi surface is equal to zero $N(E = 0) = 0$ for the A-state only. Any other superconducting state is gapless $N(E = 0) \neq 0$. This property of superconductors with toroidal order in particular means nonzero specific heat ratio $(C(T)/T)_{T \rightarrow 0} \neq 0$ in completely pure metal without impurities and crystal imperfections.

The order parameter is determined by Eq.(8)

$$\Delta_{\mathbf{k},\alpha\beta} = - \sum_{\mathbf{k}'} V_{\beta\alpha,\lambda\mu}(\mathbf{k}, \mathbf{k}') \langle a_{\mathbf{k}\lambda} a_{-\mathbf{k}\mu} \rangle. \quad (19)$$

By application to this expression the Bogolubov transformation we obtain

$$\Delta_{\mathbf{k},\alpha\beta} = - \int \frac{d^3\mathbf{k}'}{(2\pi)^3} V_{\beta\alpha,\lambda\mu}(\mathbf{k}, \mathbf{k}') \frac{1 - f_{\mathbf{k}'} - f_{-\mathbf{k}'}}{2E_{\mathbf{k}}^e} \Delta_{\lambda\mu}(\mathbf{k}') = - \int \frac{d^3\mathbf{k}'}{(2\pi)^3} V_{\beta\alpha,\lambda\mu}(\mathbf{k}, \mathbf{k}') \frac{\tanh \frac{E_{\mathbf{k}'}}{2T} + \tanh \frac{E_{-\mathbf{k}'}}{2T}}{4E_{\mathbf{k}}^e} \Delta_{\lambda\mu}(\mathbf{k}'). \quad (20)$$

Here,

$$\langle b_{\mathbf{k}\alpha}^+ b_{\mathbf{k}\beta} \rangle = f_{\mathbf{k}} = f(E_{\mathbf{k}}) = \frac{1}{\exp((\varepsilon_{\mathbf{k}}^o + E_{\mathbf{k}}^e)/T) + 1} \quad (21)$$

is the Fermi distribution function.

At $T \rightarrow T_c$ one can neglect $\Delta_{\mathbf{k}}^2$ in $E_{\mathbf{k}}^e$ in Eq.(20). Estimating the integral with logarithmic accuracy we come to the expression for critical temperature similar to usual BCS formula

$$T_c \approx \varepsilon_0 \exp\left(-\frac{1}{\tilde{N}_0 V_{\Gamma}}\right), \quad (22)$$

where ε_0 is a cut-off for energy of pairing interaction and \tilde{N}_0 is the density of states averaged over the Fermi surface with a weight corresponding to the angular dependent functions of given irreducible representation.

IV. CONCLUSION

To summarise, we have developed a theory of superconductivity in metals with toroidal ordering. The treatment is general, valid for substances of any dimension, but it was limited to the case of single band metal with point symmetry groups with a one-dimensional representations. A generalisation for the multi-band metals and superconducting states corresponding to multi-dimensional representations can be developed without problems. Generalization to metals with more complex multipole order also does not present much of a problem.

Appendix A: Superconducting order parameters in metals with toroidal order

The structures with point symmetry groups: $\mathbf{D}_{2h}(\mathbf{C}_{2v})$, $\mathbf{C}_{2h}(\mathbf{C}_2)$, $\mathbf{C}_{2h}(\mathbf{C}_{\sigma_x})$ are sketched on Fig.2 a,b,c correspondingly. Below are written the functions of irreducible representations suitable for superconducting states in antiferromagnets with these symmetries.

1. Group $\mathbf{D}_{2h}(\mathbf{C}_{2v})$

This group consists of elements $E, C_{2y}, \sigma_x, \sigma_z, RC_{2x}, RC_{2z}, R\sigma_y, RI$. The odd part of electron band energy is

$$\varepsilon_{\mathbf{k}}^o = \gamma \sin k_y c.$$




FIG. 2: (a) The $y = b/2$ plane of elementary cell of an antiferromagnet with point symmetry group $\mathbf{D}_{2h}(\mathbf{D}_2)$; (b) The $y = b/2$ plane of elementary cell of an antiferromagnet with point symmetry group $\mathbf{C}_{2h}(\mathbf{C}_2)$; (c) The $y = b/2$ plane of elementary cell of an antiferromagnet with point symmetry group $\mathbf{C}_{2h}(\mathbf{C}_{\sigma_x})$. The moments are tipped out of the plane $y = b/2$ in the opposite directions.

The functions of irreducible representations for corresponding superconducting states are

Γ	$\phi_{\mathbf{k}}^s$	$\Phi_{\mathbf{k}}^t$
A	$a\hat{k}_x^2 + b\hat{k}_y^2 + c\hat{k}_z^2$	$i\hat{k}_x\hat{z}$
B ₃	$\hat{k}_y\hat{k}_z$	$i\hat{k}_x\hat{y}$
B ₂	$\hat{k}_x\hat{k}_z$	$i\hat{k}_x\hat{x}$
B ₁	$\hat{k}_x\hat{k}_y$	$i\hat{k}_y\hat{z}$

2. Group $\mathbf{C}_{2h}(\mathbf{C}_2)$

This group is subgroup of previous one. It consists of elements $E, C_{2y}, R\sigma_y, RI$. The odd part of electron band energy is

$$\varepsilon_{\mathbf{k}}^o = \gamma \sin k_y c.$$

The functions of irreducible representations for corresponding superconducting states are

Γ	$\phi_{\mathbf{k}}^s$	$\Phi_{\mathbf{k}}^t$
A	$a\hat{k}_x^2 + b\hat{k}_y^2 + c\hat{k}_z^2 + d\hat{k}_x\hat{k}_z$	$i\hat{k}_x\hat{z}$
B	$\hat{k}_y\hat{k}_z$	$i\hat{k}_y\hat{z}$

3. Group $\mathbf{C}_{2h}(\mathbf{C}_{\sigma_x})$

This group is another subgroup of the group $\mathbf{D}_{2h}(\mathbf{C}_{2v})$. It consists of elements E, σ_x, RC_{2x}, RI . The odd part of electron band energy is

$$\varepsilon_{\mathbf{k}}^o = \gamma \sin k_y c.$$

The functions of irreducible representations for corresponding superconducting states are

Γ	$\phi_{\mathbf{k}}^s$	$\Phi_{\mathbf{k}}^t$
A	$a\hat{k}_x^2 + b\hat{k}_y^2 + c\hat{k}_z^2 + d\hat{k}_y\hat{k}_z$	$i\hat{k}_y\hat{x}$
B	$\hat{k}_x\hat{k}_z$	$i\hat{k}_y\hat{z}$

¹ L.D.Landau and E.M.Lifshitz, Course of Theoretical Physics, Vol. 8: Electrodynamics of Continuous Media (Nauka, Moscow, 1982; Pergamon, Oxford, 1984).

- ² I.E. Dzyaloshinskii, J. Exptl. Theoret. Phys. **37**, 881 (1959) [Soviet Phys. JETP **10**, 628 (1960)].
- ³ D.N. Astrov, Zh. Eksp. Teor. Fiz. **40**, 1035 (1961) [Sov. Phys. JETP **13**, 729 (1961)].
- ⁴ W.F. Brown, Jr., S. Shtrikman, and D. Treves, J. Appl. Phys. **34**, 1233 (1963).
- ⁵ R. M. Hornreich and S. Shtrikman, Phys. Rev. **171**, 1065 (1968).
- ⁶ B.B. Krichevtsov, V.V. Pavlov, R.V. Pisarev, V.N. Gridnev, J. Phys. Cond. Matt. **5**, 8233 (1993).
- ⁷ B.B. Krichevtsov, V.V. Pavlov, R.V. Pisarev, V.N. Gridnev, Phys. Rev. Lett. **76**, 4628 (1996).
- ⁸ B. Fauque et al., Phys. Rev. Lett. **96**, 197001 (2006).
- ⁹ J. Xia et al., Phys. Rev. Lett. **100**, 127002 (2008).
- ¹⁰ C.M. Varma, Rev. Mod. Phys. **92**, 031001 (2020).
- ¹¹ S. Hayami, H. Kusunose, and Y. Motome, Phys. Rev. B **90**, 024432 (2014).
- ¹² E. Berg, C.-C. Chen, and S. A. Kivelson, Phys. Rev. Lett. **100**, 027003 (2008).
- ¹³ A. Allais and T. Senthil, Phys. Rev. B **86**, 045118 (2012).
- ¹⁴ Adil Amin, Hao Wu, T. Shishidou, and D.F. Agterberg, arXiv:2306.11218 [cond-mat.supr-con].
- ¹⁵ S. Sumita and Y. Yanase, Phys. Rev. B **93**, 224507 (2016).
- ¹⁶ J. Orenstein, Phys. Rev. Lett. **107**, 067002 (2011).
- ¹⁷ Hiraku Saito, Kenta Uenishi, Naoyuki Miura, Chihiro Tabata, Hiroyuki Hidaka, Tatsuya Yanagisawa, and Hiroshi Amitsuka, J. Phys. Soc. Jpn. **87**, 033702 (2018).
- ¹⁸ O. Fedchenko, L. Smejkal, M. Kallmayer, Y. Lytvynenko, K. Medjanik, S. Babenkov, D. Vasilyev, M. Klaeui, J. Demsar, G. Schönhense, et al., Journal of Physics: Condensed Matter **34**, 425501 (2022).