ON SOME CONJECTURAL DETERMINANTS OF SUN INVOLVING RESIDUES

RITUPARNA CHALIHA AND GAUTAM KALITA

ABSTRACT. For an odd prime p and integers d, k, m with $gcd(p, d) = 1$ and $2 \leq k \leq \frac{p-1}{2}$, we consider the determinant

$$
S_{m,k}(d,p) = |(\alpha_i - \alpha_j)^m|_{1 \le i,j \le \frac{p-1}{k}},
$$

where α_i are distinct k-th power residues modulo p. In this paper, we deduce some residue properties for the determinant $S_{m,k}(d, p)$ as a generalization of certain results of Sun. Using these, we further prove some conjectures of Sun related to

$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{2},2}(-1,p)}}{p}\right) \text{ and } \left(\frac{\sqrt{S_{3+\frac{p-1}{2},2}(-1,p)}}{p}\right).
$$

In addition, we investigate the number of primes p such that $p \mid S_{m+\frac{p-1}{k},k}(-1,p)$, and confirm another conjecture of Sun related to $S_{m+\frac{p-1}{2},2}(-1,p)$.

1. Introduction and statements of results

Let n be a positive integer and R a commutative ring with unity. For an $n \times n$ matrix $M = [a_{ij}]_{1 \le i,j \le n}$ with $a_{ij} \in R$, we denote the determinant by $|M|$ or $|[a_{ij}]_{1 \le i,j \le n}|$. Let p be an odd prime and χ_{ℓ} denotes a multiplicative character of order ℓ modulo p. For example, $\chi_2(\cdot) = \left(\frac{\cdot}{p}\right)$ is the usual Legendre symbol. In this paper, we study some conjectural determinants involving residues. Determinants with Legendre symbol entries were first considered by Lehmer [\[9\]](#page-17-0), where he used a general method to determine the characteristic roots of two classes of matrices to evaluate their determinants. Extending a result of Lehmer [\[9\]](#page-17-0), Carlitz [\[2\]](#page-16-0) obtained the characteristic polynomial of the matrix

$$
M = \left| \left(\frac{i - j}{p} \right) \right|_{1 \le i, j \le p - 1},
$$

and deduced that $|M| = p^{\frac{p-3}{2}}$. In [\[3\]](#page-16-1), Chapman studied some matrices that appeared in the investigation of lattices constructed from quadratic residue codes and their generalizations, and determined values of

$$
\left| \left(\frac{i+j-1}{p} \right) \right|_{1 \le i,j \le \frac{p-1}{2}} \text{ and } \left| \left(\frac{i+j-1}{p} \right) \right|_{1 \le i,j \le \frac{p+1}{2}}
$$

²⁰²⁰ Mathematics Subject Classification. 11A07, 11A15, 11C20, 15A15, 15B36, 11R11.

Key words and phrases. Residues, Legendre symbol, Determinants,

using quadratic Gauss sums. Following these, Vsemirnov [\[14,](#page-17-1) [15\]](#page-17-2) used a sophisticated matrix decomposition to confirm a challenging conjecture of Chapman [\[4\]](#page-16-2) on the determinant

$$
\left| \left(\frac{j-i}{p} \right) \right|_{1 \le i,j \le \frac{p+1}{2}}.
$$

In [\[11\]](#page-17-3), Sun concentrated on determinants of the form

$$
\left| \left(\frac{f(i,j)}{p} \right) \right|_{1 \le i,j \le \frac{p-1}{2}},
$$

where $f(x, y)$ is a quadratic form, and investigated their quadratic residue properties. In particular, for $p \nmid d$, Sun [\[11\]](#page-17-3) studied the determinant

$$
S(d, p) = \left| \left(\frac{i^2 + dj^2}{p} \right) \right|_{1 \le i, j \le \frac{p-1}{2}},
$$

and proved that

$$
\left(\frac{S(d,p)}{p}\right) = \begin{cases} \left(\frac{-1}{p}\right), & \text{if } \left(\frac{d}{p}\right) = 1; \\ 0, & \text{if } \left(\frac{d}{p}\right) = -1. \end{cases}
$$

In addition, Sun [\[11\]](#page-17-3) also posed a number of conjectures related to the determinant $S(d, p)$. In recent years, some of these conjectures and their generalizations have been proved by many mathematicians, such as Krachun et. al. [\[7\]](#page-17-4), Grinberg et. al. [\[5\]](#page-16-3), Wu [\[17\]](#page-17-5), and Wu-Wang [\[16\]](#page-17-6).

For $n, k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$, throughout the paper, let p be an odd prime such that $p \equiv 1 \pmod{k}$ and

$$
S_{n,k}(d,p) = |(\alpha_i + d\alpha_j)^n|_{1 \le i,j \le \frac{p-1}{k}},
$$

where α_i are distinct k-th residues modulo p. Note that $S_{\frac{p-1}{2},2}(d,p) \equiv S(d,p) \pmod{p}$. From [\[6,](#page-16-4) Theorem 1.8, it is known that for $p > 3$,

$$
\left(\frac{S_{\frac{p-3}{2},2}\left(d,p\right)}{p}\right)=\begin{cases} \left(\frac{d}{p}\right)^{\frac{p-1}{4}}, & \text{if } p\equiv 1\,(\text{mod}\,4); \\ \left(\frac{d}{p}\right)^{\frac{p-3}{4}}(-1)^{\left|\{0< k<\frac{p}{2}:\left(\frac{k}{p}\right)=-1\}\right|}, & \text{if } p\equiv 3\,(\text{mod}\,4). \end{cases}
$$

Moreover, for $n < \frac{p-3}{2}$, we have from [\[8,](#page-17-7) Lemma 9] that

$$
S_{n,2}(d,p) = 0.
$$

In [\[12\]](#page-17-8), Sun considered the determinant $S_{n,2}(d, p)$ for $n > \frac{p-1}{2}$, and proved a number of results related to them under the condition $\left(\frac{d}{p}\right) = -1$. Following these, Wu, She and Wang [\[18\]](#page-17-9) proved a conjecture of Sun [\[11,](#page-17-3) Conjecture 4.5] related the Legendre symbol $\left(\frac{S_{p+1,2}(d,p)}{p}\right)$ p . Recently, Ren and Sun [\[10\]](#page-17-10) studied the determinant $S_{n,2}(d, p)$ for $n > \frac{p-1}{2}$ under the condition $\left(\frac{d}{p}\right) = 1$. In the following theorems, we generalize some of the results of Ren and Sun [\[10\]](#page-17-10) to the determinant $S_{n,k}(d, p)$.

Theorem 1.1. Let p be an odd prime and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$ such that $p > 2k + 1$ and $p \equiv 1 \pmod{p}$ k). If $n \in \{\frac{p-1}{k}+1, \frac{p-1}{k}+2, \cdots, \frac{2(p-1)}{k}-1\}$ such that $n \equiv \frac{p-1}{k} \pmod{2}$ and $d \in \mathbb{Z}$ with $\chi_k(d) = -1$, then $S_{n,k}(d, p) \equiv 0 \pmod{p}.$

Remark 1.2. For $k = 2$, we obtain [\[10,](#page-17-10) Theorem 1.1] from Theorem [1.1.](#page-2-0)

Theorem 1.3. Let p be an odd prime and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$ such that $p > 2k + 1$ and $p \equiv 1 \pmod{p}$ 2k). Suppose $n \in \{\frac{p-1}{k} + 1, \frac{p-1}{k} + 2, \cdots, \frac{2(p-1)}{k} - 1\}$ is odd and $d \in \mathbb{Z}$ with $\chi_k(d) = 1$.

(a) If k is even, then
$$
\left(\frac{S_{n,k}(d,p)}{p}\right) \neq -1
$$
.
\n(b) If k is odd, then $\left(\frac{S_{n,k}(d,p)}{p}\right) \neq \begin{cases} -1, & \text{if } p \equiv 1 \pmod{4k}; \\ \left(\frac{d}{p}\right), & \text{if } p \equiv 2k+1 \pmod{4k}. \end{cases}$

Remark 1.4. For $k = 2$, Theorem [1.3](#page-2-1) readily provides [\[10,](#page-17-10) Theorem 1.2].

In [\[12\]](#page-17-8), Sun showed that

$$
S_{m+\frac{p-1}{2},2}(-1,p)\equiv 0\,(\mathrm{mod}\,p)
$$

when m is even and $p \equiv 3 \pmod{4}$, and $S_{m+\frac{p-1}{2},2}(-1,p)$ is an integer square modulo p when m is odd and $p \equiv 1 \pmod{4}$. In addition, Sun [\[12\]](#page-17-8) also posed a number of conjectures related to the determinant $S_{m+\frac{p-1}{2},2}(-1,p).$

Conjecture 1.5. [\[12,](#page-17-8) Conjecture 6.3] For any prime $p \equiv 1 \pmod{4}$, we have

$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{2},2}(-1,p)}}{p}\right) = (-1)^{|\{0 < k < \frac{p}{4}:\left(\frac{k}{p}\right) = -1\}|} \left(\frac{p}{3}\right).
$$

Conjecture 1.6. [\[12,](#page-17-8) Conjecture 6.4] For any prime $p \equiv 1 \pmod{4}$, we have

$$
\left(\frac{\sqrt{S_{3+\frac{p-1}{2},2}(-1,p)}}{p}\right) = (-1)^{|\{0 < k < \frac{p}{4}:\left(\frac{k}{p}\right) = -1\}|} \left(\frac{p}{4 + (-1)^{\frac{p-1}{4}}}\right).
$$

Conjecture 1.7. [\[12,](#page-17-8) Conjecture 6.5] For any positive odd integer m, the set

$$
E_2(m)=\{p:p\ is\ a\ prime\ with\ 4\ \left|\ p-1\ \ and\ p\ \right|\ S_{m+\frac{p-1}{2},2}(-1,p)\}
$$

is finite. In particular,

$$
E_2(5) = \{29\}, E_2(7) = \{13, 53\}, E_2(9) = \{13, 17, 29\},
$$

$$
E_2(11) = \{17, 29\} \text{ and } E_2(13) = \{17, 109, 401\}.
$$

Remark 1.8. Sun [\[12\]](#page-17-8) provided the list of $E_2(m)$ based on his calculations for primes < 1000 .

If m is odd and $p \equiv 1 \pmod{2k}$, then it is easy to see that $S_{m+\frac{p-1}{k},k}(-1,p)$ is a skew-symmetric matrix of even order, and hence $S_{m+\frac{p-1}{k},k}(-1,p)$ is an integral square. For certain values of m, we now investigate residue properties of $\sqrt{S_{m+\frac{p-1}{k},k}(-1,p)}$. For $a, e \in \mathbb{N}$, we define the e-factorial, denoted by $a!_{(e)}$, as

$$
a!_{(e)} = \begin{cases} a \cdot (a - e)!_{(e)}, & \text{if } a > e; \\ a, & \text{if } 1 \le a \le e. \end{cases}
$$

In particular, we denote $a!_{(2)}$ by $a!!$

Theorem 1.9. Let p be an odd prime and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$ such that $p \equiv 1 \pmod{2k}$. Suppose α_i are distinct k-th residues modulo p and $T(\frac{p-1}{k}) := \prod$ $1 \leq i < j \leq \frac{p-1}{k}$ $(\alpha_i - \alpha_j).$

(a) If
$$
p \equiv 1 \pmod{4k}
$$
, then
\n
$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{k},k}(-1,p)}}{p}\right) = \left(\frac{(k-1)(2k-1)}{p}\right)\left(\frac{\frac{p-1}{k}!!}{p}\right)\left(\frac{T(\frac{p-1}{k})}{p}\right).
$$

(b) If k is even and $p \equiv 2k + 1 \pmod{4k}$, then

$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{k},k}(-1,p)}}{p}\right) = \begin{cases} \left(\frac{T(\frac{p-1}{k})}{p}\right), & \text{if } p = 2k+1; \\ \left(\frac{k(2k-1)}{p}\right)\left(\frac{\frac{p-1}{k}-1)!!}{p}\right)\left(\frac{T(\frac{p-1}{k})}{p}\right), & \text{otherwise.} \end{cases}
$$

For $k = 2$, Theorem [1.9](#page-3-0) yields

$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{2},2}(-1,p)}}{p}\right) = \begin{cases}\n\left(\frac{3}{p}\right)\left(\frac{\frac{p-1}{2}!!}{p}\right)\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right), & \text{if } p \equiv 1 \text{ (mod 8)}; \\
\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right), & \text{if } p = 5; \\
\left(\frac{6}{p}\right)\left(\frac{\frac{p-3}{2}!!}{p}\right)\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right), & \text{if } p \equiv 5 \text{ (mod 8) and } p \neq 5.\n\end{cases}
$$
\n(1)

From Lemma [2.3,](#page-6-0) we note that

$$
\frac{2}{p} = \left(\frac{\frac{p-1}{2}!}{p}\right) = \left(\frac{\frac{p-1}{2}! \cdot \frac{p-3}{2}! \cdot \cdots}{p}\right),
$$
\n
$$
\left(\frac{\frac{p-1}{2}! \cdot \cdots}{p}\right) = \left(\frac{\frac{p-3}{2}! \cdot \cdots}{p}\right) \left(\frac{2}{p}\right).
$$
\n(2)

and hence

Moreover, Lemma [2.5](#page-6-1) and Lemma [2.3](#page-6-0) together provide

 $\sqrt{ }$

$$
\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right) = \left(\frac{2}{p}\right). \tag{3}
$$

Plugging [\(2\)](#page-3-1) and [\(3\)](#page-3-2) in [\(1\)](#page-3-3), and then using Lemma [2.4](#page-6-2) and Theorem [2.1,](#page-5-0) we immediately have the following corollary.

Corollary 1.10. Conjecture [1.5](#page-2-2) is true.

We also deduce another similar result in the following theorem.

Theorem 1.11. Let p be an odd primes and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$ such that $p \equiv 1 \pmod{2k}$. Suppose α_i are distinct k-th residues modulo p and $T\left(\frac{p-1}{k}\right) = \prod$ $1 \leq i < j \leq \frac{p-1}{k}$ $(\alpha_i - \alpha_j).$

(a) If $p \equiv 1 \pmod{4k}$, then

$$
\left(\frac{\sqrt{S_{3+}\frac{p-1}{k},k}(-1,p)}{p}\right) = \left(\frac{k(3k-1)(4k-1)(6k^3+(3k-1)(2k-1)(k-1))}{p}\right)
$$

$$
\left(\frac{\left(\frac{p-1}{k}-1\right)!!}{p}\right)\left(\frac{T\left(\frac{p-1}{k}\right)}{p}\right).
$$

(b) If k is even and $p \equiv 2k + 1 \pmod{4k}$, then

$$
\left(\frac{\sqrt{S_{3+\frac{p-1}{k},k}(-1,p)}}{p}\right) = \begin{cases} \left(\frac{T(\frac{p-1}{p})}{p}\right), & \text{if } p = 2k+1; \\ \left(\frac{(k-1)(2k-1)(4k-1)(6k^3+(3k-1)(2k-1)(k-1))}{p}\right) & \\ \left(\frac{\frac{p-1}{k}\right)!!}{p}\right), & \text{otherwise.} \end{cases}
$$

For $k = 2$, Theorem 1.[11](#page-4-0) implies

$$
\left(\frac{\sqrt{S_{3+\frac{p-1}{2},2(-1,p)}}}{p}\right) = \begin{cases} \left(\frac{10}{p}\right)\left(\frac{\frac{p-3}{2}!!}{p}\right)\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right), & \text{if } p \equiv 1 \text{ (mod 8)};\\ \left(\frac{T\left(\frac{p-1}{2}!!\right)}{p}\right), & \text{if } p = 5;\\ \left(\frac{3}{p}\right)\left(\frac{\frac{p-1}{2}!!}{p}\right)\left(\frac{T\left(\frac{p-1}{2}\right)}{p}\right), & \text{if } p \equiv 5 \text{ (mod 8) and } p \neq 5.\end{cases}
$$

Using now [\(2\)](#page-3-1), [\(3\)](#page-3-2), and Theorem [2.1,](#page-5-0) we have the following corollary.

Corollary 1.12. Conjecture [1.6](#page-2-3) is true.

From Theorem [1.9,](#page-3-0) it is easy to see that if $p \equiv 1 \pmod{2k}$ and $p \equiv 1 \pmod{4}$, then $p \nmid S_{1+\frac{p-1}{k},k}(-1,p)$ for any $2 \leq k \leq \frac{p-1}{2}$. In a similar way, Theorem [1.11](#page-4-0) implies that if $p \equiv 1 \pmod{2k}$ and $p \equiv 1 \pmod{4}$, then

$$
p \mid S_{3+\frac{p-1}{k},k}(-1,p)
$$

if and only if

$$
p | 6k3 + (3k - 1)(2k - 1)(k - 1).
$$

For a fixed k with $2 \le k \le \frac{p-1}{2}$, note that $6k^3 + (3k-1)((2k-1)(k-1))$ is a non-zero integer, and hence there are finite number of primes p such that $p \mid S_{3+\frac{p-1}{k},k}(-1,p)$. We now move our attention to the general determinant $S_{m+\frac{p-1}{k},k}(-1,p)$, and prove a similar result in the following theorem.

Theorem 1.13. Let p be an odd prime and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$. For any odd positive integer m, the set

 $E_k(m) = \{p : p \text{ is a prime with } 2k \mid (p-1) \text{ and } p \mid S_{m+\frac{p-1}{k},k}(-1,p)\}$

is finite.

Moreover, $E_k(1) = \phi$ and if $m \geq 3$ and $p > km + 1$, then $p \in E_k(m)$ if and only if $p \equiv 1 \pmod{2k}$ and

$$
p \mid \left((km)!_{(k)} + (km-1)!_{(k)} \right)
$$

or

$$
p \mid \left(\frac{(km-kl)!_{(k)}}{(kl)!(k)} + \frac{(km-kl-1)!(k)}{(kl-1)!(k)}\right)
$$

for $l = 1, \cdots, \frac{m-1}{2}$.

Putting $k = 2$ in Theorem [1.13,](#page-5-1) we obtain the following immediate corollary.

Corollary 1.14. Conjecture [1.7](#page-2-4) is true.

We now give an application of Theorem [1.13](#page-5-1) to obtain the set $E_2(13)$. Note that

$$
(2m)!! + (2m - 1)!! = 26!! + 25!! = 58917607974225 = 36 \cdot 52 \cdot 7 \cdot 11 \cdot 13 \cdot 109 \cdot 29629,
$$

which gives $109, 29629 \in E_2(13)$ due to Theorem [1.13.](#page-5-1) Moreover, we have

$$
\frac{(2m-2l)!!}{(2l)!!} + \frac{(2m-2l-1)!!}{(2l-1)!!} = \begin{cases} \frac{24!!}{2!!} + \frac{23!!}{1!!} = 3^6 \cdot 5^2 \cdot 7 \cdot 11 \cdot 924397, & \text{if } l = 1; \\ \frac{22!!}{4!!} + \frac{21!!}{3!!} = 3^5 \cdot 5^2 \cdot 7 \cdot 11 \cdot 31643, & \text{if } l = 2; \\ \frac{20!!}{6!!} + \frac{19!!}{5!!} = 3^4 \cdot 5 \cdot 7 \cdot 42703, & \text{if } l = 3; \\ \frac{18!!}{8!!} + \frac{17!!}{7!!} = 3^4 \cdot 5^2 \cdot 401, & \text{if } l = 4; \\ \frac{16!!}{10!!} + \frac{15!!}{9!!} = 3^3 \cdot 179, & \text{if } l = 5; \\ \frac{14!!}{12!!} + \frac{13!!}{11!!} = 3^3, & \text{if } l = 6. \end{cases}
$$

As a result, $401,924397 \in E_2(13)$. It is easy to calculate that $p = 17$ is the only prime less than 27 such that $p \mid S_{13+\frac{p-1}{2}}, 2(-1, p)$. Combining all these together, we obtain $E_2(13) = \{17, 109, 401, 29629, 924397\}.$

2. Preliminaries

In this section, we recall certain basic results that shall be used to prove our results. We begin with the Quadratic Reciprocity Law.

Theorem 2.1. [\[1,](#page-16-5) Theorem 9.9] If p and q are distinct odd primes, then

$$
\left(\frac{p}{q}\right)\left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{2}\cdot\frac{q-1}{2}}.
$$

The following result on determinant is known from [\[8\]](#page-17-7).

Lemma 2.2. [\[8,](#page-17-7) Lemma 10] Let R be a commutative ring with identity and let $P(x) =$ \sum^{n-1} $i=0$ $a_ix^i \in R[x]$. Then

$$
det[P(X_iY_j)]_{1\leq i,j\leq n} = a_0a_1\cdots a_{n-1} \prod_{1\leq i
$$

In [\[11\]](#page-17-3), Sun evaluated the following Legendre symbol values of certain factorials.

Lemma 2.3. [\[11,](#page-17-3) Lemma 2.3] Let $p \equiv 1 \pmod{4}$ be a prime, then

$$
\left(\frac{\frac{p-1}{2}!}{p}\right) = \left(\frac{2}{p}\right).
$$

Lemma 2.4. [\[11,](#page-17-3) Lemma 3.2] Let p be an odd prime. If $p \equiv 1 \pmod{4}$, then

$$
\left(\frac{\frac{p-3}{2}!!}{p}\right) = (-1)^{\left|\{0 < k < \frac{p}{4}:\left(\frac{k}{p}\right) = -1\right\}|}.
$$

We also restate a result of Sun from [\[13\]](#page-17-11).

Lemma 2.5. [\[13,](#page-17-11) (1.5)] For an odd prime p, we have

$$
\prod_{1 \le i < j \le \frac{p-1}{2}} (j^2 - i^2) \equiv \begin{cases} -\frac{p-1}{2}! \pmod{p}, & \text{if } p \equiv 1 \pmod{4} \\ 1 \pmod{p}, & \text{if } p \equiv 3 \pmod{4} \end{cases}
$$

Finally, we deduce a congruence relation for $S_{m,k}(d, p)$ modulo p using a method introduced by Ren and Sun [\[10\]](#page-17-10).

Lemma 2.6. Let p be an odd prime and $k \in \mathbb{N}$ with $2 \leq k \leq \frac{p-1}{2}$ such that $p \equiv 1 \pmod{k}$. Suppose $m, d \in \mathbb{Z}$ such that $p \nmid d$, $\chi_k(d) = \pm 1$, and $m \in \{\frac{p-1}{k} + 1, \frac{p-1}{k} + 2, \cdots, \frac{2(p-1)}{k} - 1\}$. If $p > 2k + 1$, then $S_{m,k}(d,p) \equiv a_{m,k}^2(d,p) \, b_{m,k}(d,p) \, (\text{mod } p),$

$$
where
$$

$$
a_{m,k}(d,p) = \prod_{l=0}^{\lfloor \frac{(k(m-1)-p+1)}{2k} \rfloor} \left[\binom{m}{l} + \chi_k(d) \binom{m}{m - \frac{p-1}{k} - l} \right] \prod_{0 \le l < \frac{p-1}{k} - 1 - \lfloor \frac{m}{2} \rfloor} \binom{m}{m - \frac{p-1}{k} + 1 + l}
$$

$$
\times \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)
$$

and

$$
b_{m,k}(d,p) = \begin{cases} \chi_{2k}(d)(-1)^{\frac{p-1}{2k}-1}(1+\chi_k(d))\left(\frac{m}{\frac{km-p+1}{2k}}\right)\left(\frac{m}{\frac{m}{2}}\right), & \text{if } 2 \mid m \text{ and } p \equiv 1 \pmod{2k}, \\ (\chi_k(d))^{\frac{m}{2}}(-1)^{\frac{p-k-1}{2k}}\left(\frac{m}{\frac{m}{2}}\right), & \text{if } 2 \mid m \text{ and } p \equiv k+1 \pmod{2k}, \\ (\chi_{2k}(d))^m(-1)^{\frac{p-1}{2k}}; & \text{if } 2 \nmid m \text{ and } p \equiv 1 \pmod{2k}, \\ (-1)^{\frac{p-k-1}{2k}}(1+\chi_k(d))\left(\frac{m}{\frac{km-p+1}{2k}}\right), & \text{if } 2 \nmid m \text{ and } p \equiv k+1 \pmod{2k}. \end{cases}
$$

Proof. Let g be a primitive root modulo p, then ${g^{ki}}_{i=1}^{p-1 \over k}$ is a permutation of all distinct k-th power residues modulo p. Therefore,

$$
\prod_{i=1}^{\frac{p-1}{k}} \alpha_i \equiv \prod_{i=1}^{\frac{p-1}{k}} g^{ki} = \{g^{\frac{p-1}{2}}\}^{\frac{p-1}{k}+1} \equiv (-1)^{\frac{p-1}{k}+1} \pmod{p}.
$$

Using this, we have

$$
S_{m,k}(d,p) = \prod_{j=1}^{\frac{p-1}{k}} \alpha_j^m \left| \left(\frac{\alpha_i}{\alpha_j} + d \right)^m \right|_{1 \le i,j \le \frac{p-1}{k}} \equiv (-1)^{m(\frac{p-1}{k}+1)} \left| \left(\frac{\alpha_i}{\alpha_j} + d \right)^m \right|_{1 \le i,j \le \frac{p-1}{k}} \pmod{p}.
$$
 (4)

Note that

$$
\left(\frac{\alpha_i}{\alpha_j} + d\right)^m = \sum_{l=0}^{m-\frac{p-1}{k}} \binom{m}{l} d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l + \sum_{m-\frac{p-1}{k}+1 \le l < \frac{p-1}{k}} \binom{m}{l} d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l
$$
\n
$$
+ \sum_{l=\frac{p-1}{k}}^{m} \binom{m}{l} d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l
$$
\n
$$
\equiv \sum_{l=0}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + \binom{m}{\frac{p-1}{k}+l} d^{-\frac{p-1}{k}} \right] d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l
$$
\n
$$
+ \sum_{m-\frac{p-1}{k}+1 \le l < \frac{p-1}{k}} \binom{m}{l} d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l \pmod{p}
$$
\n
$$
\equiv \sum_{l=0}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + \binom{m}{m-\frac{p-1}{k}-l} d^{-\frac{p-1}{k}} \right] d^{m-l} \left(\frac{\alpha_i}{\alpha_j}\right)^l
$$
\n
$$
+ \sum_{0 \le l < \frac{2(p-1)}{k}-m-1} \binom{m}{l+m-\frac{p-1}{k}+1} d^{\frac{p-1}{k}-1-l} \left(\frac{\alpha_i}{\alpha_j}\right)^{l+m-\frac{p-1}{k}+1} \pmod{p}.
$$

We now use Lemma [2.2](#page-6-3) to obtain

$$
\left| \left(\frac{\alpha_i}{\alpha_j} + d \right)^m \right|_{1 \le i,j \le \frac{p-1}{k}} = \prod_{l=0}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right] d^{m-l}
$$

$$
\prod_{0 \le l < \frac{2(p-1)}{k}-m-1} \binom{m}{l+m-\frac{p-1}{k}+1} d^{\frac{p-1}{k}-1-l} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j) \left(\frac{1}{\alpha_i} - \frac{1}{\alpha_j} \right) (\text{mod } p).
$$

Since

$$
\prod_{1 \leq i < j \leq \frac{p-1}{k}} \alpha_i \alpha_j = \prod_{1 \leq j \leq \frac{p-1}{k}} \alpha_j^{\frac{p-1}{k}-1} \equiv (-1)^{\frac{p-1}{k}+1} \, (\operatorname{mod} p),
$$

we have

$$
\prod_{1 \leq i < j \leq \frac{p-1}{k}} (\alpha_i - \alpha_j) \left(\frac{1}{\alpha_i} - \frac{1}{\alpha_j} \right) = (-1)^{\left(\frac{p-1}{2}\right)} \prod_{1 \leq i < j \leq \frac{p-1}{k}} \frac{(\alpha_i - \alpha_j)^2}{\alpha_i \alpha_j}
$$
\n
$$
\equiv (-1)^{\left\lfloor \frac{p-k-1}{2k} \right\rfloor} \prod_{1 \leq i < j \leq \frac{p-1}{k}} (\alpha_i - \alpha_j)^2 \, (\text{mod } p).
$$

As a result,

$$
\left| \left(\frac{\alpha_i}{\alpha_j} + d \right)^m \right|_{1 \le i,j \le \frac{p-1}{k}} \equiv (-1)^{\lfloor \frac{p-k-1}{2k} \rfloor} d^{\frac{\frac{p-1}{k}(2m - \frac{p-1}{k} + 1)}{2}} \prod_{l=0}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m - \frac{p-1}{k} - l} \right]
$$

$$
\prod_{0 \le l < \frac{2(p-1)}{k} - m - 1} \binom{m}{l + m - \frac{p-1}{k} + 1} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)^2 \, (\text{mod } p).
$$

Using this in [\(4\)](#page-7-0), we obtain

$$
S_{m,k}(d,p) \equiv (-1)^{m(\frac{p-1}{k}+1)+\lfloor \frac{p-k-1}{2k} \rfloor} d^{\frac{\frac{p-1}{k}(2m-\frac{p-1}{k}+1)}{2}} \prod_{l=0}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]
$$

$$
\prod_{0 \le l < \frac{2(p-1)}{k}-m-1} \binom{m}{l+m-\frac{p-1}{k}+1} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)^2 \, (\text{mod } p). \tag{5}
$$

It is easy to see that

$$
\prod_{0 \le l < \frac{2(p-1)}{k} - m - 1} \binom{m}{l + m - \frac{p-1}{k} + 1} = \begin{cases} \binom{m}{\frac{m}{2}} & \prod_{0 \le l < \frac{p-1}{k} - 1 - \frac{m}{2}} \binom{m}{l + m - \frac{p-1}{k} + 1}^2, & \text{if } 2 \mid m; \\ \prod_{0 \le l < \frac{p-1}{k} - 1 - \frac{m-1}{2}} \binom{m}{l + m - \frac{p-1}{k} + 1}^2, & \text{if } 2 \nmid m. \end{cases} \tag{6}
$$

Case I: Let $m \equiv \frac{p-1}{k} \pmod{2}$. Noting that $\chi_k(d) = d^{\frac{p-1}{k}} \equiv \pm 1 \pmod{p}$, we have

$$
\prod_{l=0}^{m-\frac{p-1}{k}} \left\{ \binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right\}
$$
\n
$$
= (1+d^{-\frac{p-1}{k}}) \left(\frac{m}{\frac{m-\frac{p-1}{k}}{2}} \right) \prod_{l=0}^{\frac{m-\frac{p-1}{k}}{2}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]
$$
\n
$$
\prod_{l=\frac{m-\frac{p-1}{k}}{2}+1}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]
$$
\n
$$
= (1+d^{-\frac{p-1}{k}}) (d^{-\frac{p-1}{k}})^{\frac{km-p+1}{2k}} \left(\frac{m}{\frac{m-\frac{p-1}{k}}{2}} \right)
$$
\n
$$
\prod_{l=0}^{\frac{m-\frac{p-1}{k}}{2}} -1 \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]^2 (\text{mod } p). \tag{7}
$$

Using (6) and (7) in (5) , and then using the facts that

$$
d^{\frac{p-1}{k}(2m-\frac{p-1}{k}+1)}(d^{-\frac{p-1}{k}})^{\frac{km-p+1}{2k}} \equiv d^{\frac{(m+1)(p-1)}{2k}}(\text{mod } p)
$$

$$
\equiv \begin{cases} (\chi_{2k}(d))^{m+1} \pmod{p}, & \text{if both } m, \frac{p-1}{k} \text{ are even;} \\ (\chi_k(d))^{\frac{m+1}{2}} \pmod{p}, & \text{if both } m, \frac{p-1}{k} \text{ are odd;} \end{cases}
$$

and

$$
(-1)^{m\left(\frac{p-1}{k}+1\right)+\left\lfloor\frac{p-k-1}{2k}\right\rfloor} = \begin{cases} (-1)^{\frac{p-1}{2k}-1}; & \text{if both } m, \frac{p-1}{k} \text{ are even}; \\ (-1)^{\frac{p-k-1}{2k}}; & \text{if both } m, \frac{p-1}{k} \text{ are odd}; \end{cases}
$$

we obtain the desired result.

Case II: Let $m \not\equiv \frac{p-1}{k} \pmod{2}$. In this case, we have

$$
\prod_{l=0}^{m-\frac{p-1}{k}} \left\{ \binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right\}
$$
\n
$$
= \prod_{l=0}^{\frac{m-\frac{p-1}{k}-1}{2}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]
$$
\n
$$
= \prod_{l=\frac{m-\frac{p-1}{k}+1}{2}}^{m-\frac{p-1}{k}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]
$$
\n
$$
= (d^{-\frac{p-1}{k}})^{\frac{k(m+1)-p+1}{2k}} \prod_{l=0}^{\frac{m-\frac{p-1}{k}-1}{2}} \left[\binom{m}{l} + d^{-\frac{p-1}{k}} \binom{m}{m-\frac{p-1}{k}-l} \right]^2 \pmod{p}.
$$
\n(8)

Putting (6) and (8) in (5) , we use the facts

$$
d^{\frac{\frac{p-1}{k}(2m-\frac{p-1}{k}+1)}{2}}(d^{-\frac{p-1}{k}})^{\frac{k(m+1)-p+1}{2k}} \equiv d^{\frac{m(p-1)}{2k}} \pmod{p}
$$

$$
\equiv \begin{cases} (\chi_k(d))^{\frac{m}{2}} \pmod{p}, & \text{if } m \text{ is even and } \frac{p-1}{k} \text{ is odd}; \\ (\chi_{2k}(d))^m \pmod{p}, & \text{if } m \text{ is odd and } \frac{p-1}{k} \text{ is even}; \end{cases}
$$

and

$$
(-1)^{m(\frac{p-1}{k}+1)+\lfloor \frac{p-k-1}{2k} \rfloor} = \begin{cases} (-1)^{\frac{p-k-1}{2k}}, & \text{if } m \text{ is even and } \frac{p-1}{k} \text{ is odd;} \\ (-1)^{\frac{p-1}{2k}}, & \text{if } m \text{ is odd and } \frac{p-1}{k} \text{ is even;} \end{cases}
$$

to complete the proof of the lemma.

3. Proof of the main results

In this section, we give proof of our main results.

Proof of Theorem [1.1.](#page-2-0) We have $n \equiv \frac{p-1}{k} \pmod{2}$. Therefore, both n and $\frac{p-1}{k}$ are even or odd. Since $\chi_k(d) = -1$, we must have from Lemma [2.6](#page-6-4) that $b_{n,k}(d, p) = 0$, and hence we complete the proof of the theorem. \Box

Proof of Theorem [1.3.](#page-2-1) From Lemma [2.6,](#page-6-4) we have

$$
\left(\frac{S_{n,k}(d,p)}{p}\right) = \left(\frac{a_{n,k}(d,p)}{p}\right)^2 \left(\frac{b_{n,k}(d,p)}{p}\right).
$$

Since *n* is odd and $p \equiv 1 \pmod{2k}$, Lemma [2.6](#page-6-4) provides

$$
\left(\frac{b_{n,k}(d,p)}{p}\right) = \left(\frac{d}{p}\right)^{\frac{p-1}{2k}} \left(\frac{\chi_k(d)}{p}\right)^{\frac{n-1}{2}} \left(\frac{-1}{p}\right)^{\frac{p-1}{2k}} = \left(\frac{-d}{p}\right)^{\frac{p-1}{2k}},
$$

and hence

$$
\left(\frac{S_{n,k}(d,p)}{p}\right) = \left(\frac{a_{n,k}(d,p)}{p}\right)^2 \left(\frac{-d}{p}\right)^{\frac{p-1}{2k}}.\tag{9}
$$

 (a) Let k be even. Since

$$
\chi_k(d) \equiv d^{\frac{p-1}{k}} \equiv 1 \, (\text{mod } p),
$$

we have

$$
d^{\frac{p-1}{2}} = (d^{\frac{p-1}{k}})^{\frac{k}{2}} \equiv 1 \, (\text{mod } p).
$$

As a result,

$$
\left(-\frac{d}{p}\right)^{\frac{p-1}{2k}}=1,
$$

and hence [\(9\)](#page-10-0) yields

$$
\left(\frac{S_{n,k}(d,p)}{p}\right) = \left(\frac{a_{n,k}(d,p)}{p}\right)^2 = \begin{cases} 1, & \text{if } a_{n,k}(d,p) \neq 0; \\ 0, & \text{if } a_{n,k}(d,p) = 0. \end{cases}
$$

Thus we obtain the desired result.

 (b) If k is odd, then it is easy to see that

$$
\left(-\frac{d}{p}\right)^{\frac{p-1}{2k}} = \begin{cases} 1, & \text{if } p \equiv 1 \, (\text{mod } 4k) \\ -\left(\frac{d}{p}\right), & \text{if } p \equiv 2k+1 \, (\text{mod } 4k). \end{cases}
$$

Using this in [\(9\)](#page-10-0) , and then noting that

$$
\left(\frac{S_{n,k}(d,p)}{p}\right) = 0
$$

if and only if

$$
\left(\frac{a_{n,k}(d,p)}{p}\right) = 0,
$$

we complete the proof of the theorem. $\hfill \square$

Proof of Theorem [1.9.](#page-3-0) We first note that $p \equiv 1 \pmod{4}$ in each case, and hence $\left(\frac{-1}{p}\right) = 1$. If $p=2k+1,$ then

$$
S_{1+\frac{p-1}{k},k}(-1,p) = S_{3,k}(-1,p) = \begin{vmatrix} 0 & (\alpha_1 - \alpha_2)^3 \\ (\alpha_2 - \alpha_1)^3 & 0 \end{vmatrix} = (\alpha_1 - \alpha_2)^6.
$$
 (10)

.

As a result,

$$
\left(\frac{\sqrt{S_{1+\frac{p-1}{k},k}(-1,p)}}{p}\right) = \left(\frac{\pm(\alpha_1-\alpha_2)^3}{p}\right) = \left(\frac{\alpha_1-\alpha_2}{p}\right) = \left(\frac{T(\frac{p-1}{k})}{p}\right)
$$

Suppose $p > 2k + 1$ such that $p \equiv 1 \pmod{2k}$. Note that $\frac{p-1}{k}$ is even and $\frac{p-1}{k} + 1$ is odd, and hence Lemma [2.6](#page-6-4) provides

$$
S_{\frac{p-1}{k}+1,k}(-1,p)\equiv a_{\frac{p-1}{k}+1,k}^2(-1,p)b_{\frac{p-1}{k}+1,k}(-1,p)\,(\operatorname{mod }p),
$$

where

$$
a_{\frac{p-1}{k}+1,k}(-1,p) = \left(\frac{p-1}{k}+2\right) \prod_{l=0}^{\frac{p-1}{2k}-2} {\binom{\frac{p-1}{k}+1}{2+l}} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)
$$

and

$$
b_{\frac{p-1}{k}+1,k}(-1,p) = 1.
$$

Therefore,

$$
\sqrt{S_{\frac{p-1}{k}+1,k}(-1,p)} \equiv a_{\frac{p-1}{k}+1,k}(-1,p) \pmod{p} \text{ or } -a_{\frac{p-1}{k}+1,k}(-1,p) \pmod{p}.
$$

Since $\left(\frac{-1}{p}\right) = 1$, we must have

$$
\left(\frac{\sqrt{S_{\frac{p-1}{k}+1,k}(-1,p)}}{p}\right) = \left(\frac{\left(\frac{p-1}{k}+2\right)\prod_{l=0}^{\frac{p-1}{2k}-2}\binom{\frac{p-1}{k}+1}{2+l}}{p}\right)\left(\frac{\prod_{1\leq i < j \leq \frac{p-1}{k}}(\alpha_i - \alpha_j)}{p}\right). \tag{11}
$$

Note that

$$
\prod_{l=0}^{\frac{p-1}{2k}-2} \binom{\frac{p-1}{k}+1}{2+l} = \left(\frac{p-1}{k}+1\right)^{\frac{p-1}{2k}-1} \prod_{r=1}^{\frac{p-1}{2k}-1} \frac{\left(\frac{p-1}{k}-r+1\right)^{\frac{p-1}{2k}-r}}{\left(\frac{p-1}{2k}-r+1\right)^r},\tag{12}
$$

Case I: Let $p \equiv 1 \pmod{4k}$, then [\(12\)](#page-11-0) yields

$$
\left(\frac{\prod_{l=0}^{\frac{p-1}{2k}-2}\binom{\frac{p-1}{k}+1}{2+l}}{p}\right)=\left(\frac{\left(\frac{p-1}{k}\right)!!}{p}\right)\left(\frac{\frac{p-1}{k}+1}{p}\right).
$$

Using this in [\(11\)](#page-11-1), we obtain

$$
\left(\frac{\sqrt{S_{\frac{p-1}{k}+1,k}(-1,p)}}{p}\right) = \left(\frac{(\frac{p-1}{k}+1)(\frac{p-1}{k}+2)}{p}\right)\left(\frac{\left(\frac{p-1}{k}\right)!!}{p}\right)\left(\frac{\prod\limits_{1 \leq i < j \leq \frac{p-1}{k}} (\alpha_i - \alpha_j)}{p}\right)
$$

Hence the result follows.

Case II: Let $p \equiv 2k + 1 \pmod{4k}$. In this case, we have from [\(12\)](#page-11-0) that

$$
\left(\frac{\prod_{l=0}^{\frac{p-1}{2k}-2}\binom{\frac{p-1}{k}+1}{2+l}}{p}\right) = \left(\frac{\left(\frac{p-1}{k}-1\right)!!}{p}\right).
$$

Using this in [\(11\)](#page-11-1), we obtain

$$
\left(\frac{\sqrt{S_{\frac{p-1}{k}+1,k}(-1,p)}}{p}\right) = \left(\frac{(\frac{p-1}{k}+2)}{p}\right)\left(\frac{(\frac{p-1}{k}-1)!!}{p}\right)\left(\frac{\prod\limits_{1 \leq i < j \leq \frac{p-1}{k}} (\alpha_i - \alpha_j)}{p}\right),
$$

completing proof of the theorem. $\hfill \square$

Proof of Theorem [1.11.](#page-4-0) It is easy to see that $p \equiv 1 \pmod{4}$ in each case, and hence $\left(\frac{-1}{p}\right) = 1$. If $p=2k+1,$ then

$$
S_{5,k}(-1,p) = \begin{vmatrix} 0 & (\alpha_1 - \alpha_2)^5 \\ (\alpha_2 - \alpha_1)^5 & 0 \end{vmatrix} = (\alpha_1 - \alpha_2)^{10},
$$

and hence

$$
\left(\frac{\sqrt{S_{5,k}(-1,p)}}{p}\right) = \left(\frac{\pm(\alpha_1-\alpha_2)^5}{p}\right) = \left(\frac{\alpha_1-\alpha_2}{p}\right).
$$

Suppose $p > 2k + 1$ such that $p \equiv 1 \pmod{2k}$. Noting that $\frac{p-1}{k}$ is even and $\frac{p-1}{k} + 3$ is odd, we have from Lemma [2.6](#page-6-4) that

$$
S_{\frac{p-1}{k}+3,k}(-1,p)\equiv a_{\frac{p-1}{k}+3,k}^2(-1,p)~b_{\frac{p-1}{k}+3,k}(-1,p)~(\text{mod}~p),
$$

where

$$
a_{\frac{p-1}{k}+3,k}(-1,p) = \prod_{l=0}^{1} \left[\binom{\frac{p-1}{k}+3}{l} + \binom{\frac{p-1}{k}+3}{3-l} \right] \prod_{0 \le l < \frac{p-1}{2k}-2} \binom{\frac{p-1}{k}+3}{4+l} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)
$$

and

$$
b_{\frac{p-1}{k}+3,k}(-1,p)=1.
$$

.

As a result, we have

$$
\sqrt{S_{\frac{p-1}{k}+3,k}(-1,p)}\equiv a_{\frac{p-1}{k}+3,k}(-1,p)\text{ or }-a_{\frac{p-1}{k}+3,k}(-1,p)\,(\mathrm{mod}\,p).
$$

Since $\left(\frac{-1}{p}\right) = 1$, we must have

$$
\left(\frac{\sqrt{S_{\frac{p-1}{k}+3,k}(-1,p)}}{p}\right) = \left(\frac{a_{\frac{p-1}{k}+3,k}(-1,p)}{p}\right).
$$

Noting that

$$
\prod_{l=0}^{1} \left[\binom{\frac{p-1}{k}+3}{l} + \binom{\frac{p-1}{k}+3}{3-l} \right] \equiv \left[\frac{6k^3 + (3k-1)(2k-1)(k-1)}{6k^3} \right] \left[\frac{(3k-1)(4k-1)}{2k^2} \right] \pmod{p},
$$

we have

$$
\left(\frac{\sqrt{S_{\frac{p-1}{k}+3,k}(-1,p)}}{p}\right) = \left(\frac{3k(3k-1)(4k-1)\{6k^3 + (3k-1)(2k-1)(k-1)\}}{p}\right)
$$
\n
$$
\left(\frac{\prod_{0 \le l < \frac{p-1}{2k}-2} {\binom{\frac{p-1}{k}+3}{4+l}}}{p}\right) \left(\frac{\prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)}{p}\right)
$$
\n(13)

Case I: Let $p \equiv 1 \pmod{4k}$. If $p = 4k + 1$, then

$$
\prod_{0 \le l < \frac{p-1}{2k} - 2} \binom{\frac{p-1}{k} + 3}{4 + l} = 1,
$$

and hence the result follows from [\(13\)](#page-13-0).

On the other hand, if $p > 4k + 1$, then

$$
\prod_{l=0}^{\frac{p-1}{2k}-3}\binom{\frac{p-1}{k}+3}{4+l}=\left(\frac{(\frac{p-1}{k}+3)(\frac{p-1}{k}+2)(\frac{p-1}{k}+1)\frac{p-1}{k}}{4\cdot 3\cdot 2\cdot 1}\right)^{\frac{p-1}{2k}-2\frac{p-1}{2k}-3}\prod_{r=1}^{\frac{p-1}{2k}-7}\frac{\left(\frac{p-1}{k}-r\right)^{\frac{p-1}{2k}-r-2}}{\left(\frac{p-1}{2k}-r+2\right)^r},
$$

and hence

$$
\left(\frac{\prod_{l=0}^{\frac{p-1}{2k}-3}\binom{\frac{p-1}{k}+3}{4+l}}{p}\right) = \left(\frac{\left(\frac{p-1}{k}-1\right)!!}{p}\right)\binom{3}{p}.
$$

Thus we obtain the desired result due to [\(13\)](#page-13-0).

Case II: Let $p \equiv 2k + 1 \pmod{4k}$. In this case, we have

$$
\prod_{l=0}^{\frac{p-1}{2k}-3}\binom{\frac{p-1}{k}+3}{4+l}=\left(\frac{(\frac{p-1}{k}+3)(\frac{p-1}{k}+2)(\frac{p-1}{k}+1)\frac{p-1}{k}}{4\cdot 3\cdot 2\cdot 1}\right)^{\frac{p-1}{2k}-2\frac{p-1}{2k}-3}\prod_{r=1}^{\frac{p-1}{2k}-7}\frac{\left(\frac{p-1}{k}-r\right)^{\frac{p-1}{2k}-r-2}}{\left(\frac{p-1}{2k}-r+2\right)^r},
$$

and hence

$$
\left(\frac{\prod_{l=0}^{\frac{p-1}{2k}-3}\binom{\frac{p-1}{k}+3}{4+l}}{p} = \left(\frac{\frac{p-1}{k}!!}{p}\right)\binom{3}{p}\left(\frac{\left(\frac{p-1}{k}+3\right)\left(\frac{p-1}{k}+2\right)\left(\frac{p-1}{k}+1\right)}{p}\right)
$$

$$
=\left(\frac{\frac{p-1}{k}!!}{p}\right)\left(\frac{3}{p}\right)\left(\frac{k(k-1)(2k-1)(3k-1)}{p}\right).
$$

Using this in [\(13\)](#page-13-0), we complete the proof of the theorem. \Box

Proof of Theorem [1.13.](#page-5-1) We first give a proof of the theorem for $m = 1$. If $p = 2k + 1$, then [\(10\)](#page-11-2) implies that

$$
p\nmid S_{1+\frac{p-1}{k},k}(-1,p),
$$

and hence $E_k(1) = \phi$.

On the other hand, if $p > 2k + 1$, then Lemma [2.6](#page-6-4) yields

$$
S_{1+\frac{p-1}{k},k}(-1,p)\equiv a_{1+\frac{p-1}{k},k}^2(-1,p)~b_{1+\frac{p-1}{k},k}(-1,p)~(\text{mod}~p),
$$

where

$$
a_{1+\frac{p-1}{k},k}(-1,p) \equiv \left(\frac{2k-1}{k}\right) \prod_{l=0}^{\frac{p-1}{2k}-2} {\binom{\frac{p-1}{k}+1}{2+l}} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j) \pmod{p}
$$

and

$$
b_{m+\frac{p-1}{k},k}(-1,p)=1.
$$

It is easy to see that $\frac{p-1}{k} < 1 + \frac{p-1}{k} < \frac{2(p-1)}{k}$ $\frac{b-1}{k}$. Therefore,

$$
p \nmid \prod_{l=0}^{\frac{p-1}{2k}-2} {\frac{p-1}{k}+1 \choose 2+l}
$$

.

As a result,

$$
p \nmid a_{m + \frac{p-1}{k},k}(-1,p),
$$

and hence

$$
p \nmid S_{1+\frac{p-1}{k},k}(-1,p),
$$

concluding that $E_k(1) = \phi$.

We now assume that $m \geq 3$ is odd and $p > km + 1$. Note that $\frac{p-1}{k} < m + \frac{p-1}{k} < \frac{2(p-1)}{k}$ $\frac{p-1}{k}$ and $\frac{p-1}{k}$ is even, and hence Lemma [2.6](#page-6-4) provides

$$
S_{m+\frac{p-1}{k},k}(-1,p)\equiv a_{m+\frac{p-1}{k},k}^2(-1,p)\,\,b_{m+\frac{p-1}{k},k}(-1,p)\,(\mathrm{mod}\,p),
$$

where

$$
a_{m+\frac{p-1}{k},k}(-1,p) = \prod_{l=0}^{\frac{m-1}{2}} \left[\binom{m+\frac{p-1}{k}}{l} + \binom{m+\frac{p-1}{k}}{m-l} \right] \prod_{0 \le l < \frac{p-1}{2k} - \frac{m+1}{2}} \binom{m+\frac{p-1}{k}}{m+1+l} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j)
$$

and

$$
b_{m+\frac{p-1}{k},k}(-1,p)=1.
$$

It is easy to see that

$$
p \nmid \prod_{0 \le l < \frac{p-1}{2k} - \frac{m+1}{2}} \binom{m + \frac{p-1}{k}}{m+1+l} \prod_{1 \le i < j \le \frac{p-1}{k}} (\alpha_i - \alpha_j).
$$

Therefore, we must have that

$$
p\mid S_{m+\frac{p-1}{k},k}(-1,p)
$$

if and only if

$$
p \mid \prod_{l=0}^{\frac{m-1}{2}} \left[\binom{m + \frac{p-1}{k}}{l} + \binom{m + \frac{p-1}{k}}{m-l} \right]. \tag{14}
$$

For $l = 0$, note that

$$
\binom{m+\frac{p-1}{k}}{l} + \binom{m+\frac{p-1}{k}}{m-l} = 1 + \binom{m+\frac{p-1}{k}}{m} \equiv \frac{(km)!(k) + (km-1)!(k)}{(km)!(k)} \pmod{p}.
$$
 (15)

Again, for $l = 1, 2, \ldots, \frac{m-1}{2}$, we have

$$
\binom{m+\frac{p-1}{k}}{l} + \binom{m+\frac{p-1}{k}}{m-l} = \frac{\left(m+\frac{p-1}{k}\right)\left(m+\frac{p-1}{k}-1\right)\cdots\left(m+\frac{p-1}{k}-(l-1)\right)}{l!} + \frac{\left(m+\frac{p-1}{k}\right)\left(m+\frac{p-1}{k}-1\right)\cdots\left(\frac{p-1}{k}+l+1\right)}{(m-l)!} + \frac{\left(km-1\right)\left(k(m-1)-1\right)\cdots\left(k(m-l+1)-1\right)}{k^l \cdot l!} + \frac{\left(km-1\right)\left(k(m-1)-1\right)\cdots\left(k(l+1)-1\right)}{k^{m-l} \cdot (m-l)!} \left(\text{mod } p\right)
$$

$$
\equiv \frac{\left(km-1\right)\cdots\left(k(m-l+1)-1\right)}{\left(km-kl\right)!\left(k\right)} + \frac{\left(km-kl-1\right)!\left(k\right)}{\left(kl\right)!\left(k\right)} + \frac{\left(km-kl-1\right)!\left(k\right)}{\left(kl-1\right)!\left(k\right)} \left(\text{mod } p\right). \tag{16}
$$

Noting that

$$
p \nmid (km-kl)!_{(k)}
$$

for $l = 0, 1, \dots, \frac{m-1}{2}$ and

$$
p \nmid (km-1)\cdots(k(m-l+1)-1)
$$

for $l = 1, 2, \dots, \frac{m-1}{2}$, we complete the proof the second part of the theorem because of [\(14\)](#page-15-0), [\(15\)](#page-15-1), and [\(16\)](#page-15-2).

For any fixed m and k, it is obvious that there are always finite number of primes $p \leq km + 1$ such that

$$
p \mid S_{m+\frac{p-1}{k},k}(-1,p).
$$

If $p > mk + 1$, then we have already deduced that

$$
p\mid S_{m+\frac{p-1}{k},k}(-1,p)
$$

if and only if

$$
p \mid \{(km)!(k) + (km-1)!(k)\}\prod_{l=1}^{\frac{m-1}{2}} \left[\frac{(km-kl)!_{(k)}}{(kl)!(k)} + \frac{(km-kl-1)!(k)}{(kl-1)!(k)}\right].
$$

Clearly,

$$
\{(km)!(k) + (km-1)!(k)\}\prod_{l=1}^{\frac{m-1}{2}} \left[\frac{(km-kl)!(k)}{(kl)!(k)} + \frac{(km-kl-1)!(k)}{(kl-1)!(k)}\right]
$$

is a non-zero integer that is independent of p , hence it must have finite number of prime divisors. Hence the result follows. \Box

4. Conflicts of interest/Competing interests

The authors do not have any conflict of interest.

5. Data availability

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

6. Acknowledgement:

The first author acknowledges the support received from Council of Scientific & Industrial Research (CSIR), India, through a Fellowship 09/1312(15921)/2022-EMR-I. The second author is supported by a project (CRG/2023/000482) of SERB, Department of Science and Technology, Goverment of India, under Core Research Grant.

REFERENCES

- [1] D. M. Burton, Elementary Number Theory, 7th Edition, Tata McGraw-Hill Publishing Company Limited, 2007.
- [2] L. Carlitz, Some cyclotomic matrices, Acta Arith. 5 (1959), 293–308.
- [3] R. Chapman, Determinants of Legendre symbol matrices, Acta Arith. 115 (2004), 231–244.
- [4] R. Chapman, My evil determinant problem, preprint, December 12, 2012, available from [https://empslocal.ex.ac.uk/people/staff/rjchapma/etc/evildet.pdf.](https://empslocal.ex.ac.uk/people/staff/rjchapma/etc/evildet.pdf)
- [5] D. Grinberg, Z.-W. Sun, and L. Zhao, Proof of three conjectures on determinants related to quadratic residues, Linear Multilinear Algebra 70 (2022), 3734–3746.
- [6] G. Kalita and R. Chaliha, On some determinants of generalized matrices involving Legendre symbol, under review.
- [7] D. Krachun, F. Petrov, Z.-W. Sun, and M. Vsemirnov, On some determinants involving Jacobi symbols, Finite Fields Appl. 64 (2020), Article 101672.
- [8] C. Krattenthaler, Advanced determinant calculus: a complement, Linear Algebra Appl. 411 (2005), 64–166.
- [9] D. H. Lehmer, On certain character matrices, Pacific J. Math. 6 (1956), 491–499.
- [10] C.-K. Ren and Z.-W. Sun, On some determinants arising from quadratic residues, arXiv:2404.11547 (2024).
- [11] Z.-W. Sun, On some determinants with Legendre symbol entries, Finite Fields Appl. 56 (2019), 285–307.
- [12] Z.-W. Sun, Some determinants involving quadratic residues modulo primes, arXiv:2401.14301 (2024).
- [13] Z.-W. Sun, Quadratic residues and related permutations and identities, Finite Fields Appl. 59 (2019), 246–283.
- [14] M. Vsemirnov, On the evaluation of R. Chapman's "evil determinant", Linear Algebra Appl. 436 (2012), 4101-4106.
- [15] M. Vsemirnov, On R. Chapman's "evil determinant": case $p \equiv 1 \pmod{4}$, Acta Arith. 159 (2013), 331–344.
- [16] H.-L. Wu and L.-Y. Wang, Application of Circulant matrices to determinants involving k-th power residues, Bull. Aust. Math. Soc. 106 (2022), 243-253.
- [17] H.-L.Wu, Determinants concerning Legendre symbols, C. R. Math .Acad. Sci. Paris 359 (2021), 651–655.
- [18] H.-L.Wu, Y.-F. She and L.-Y.Wang, Cyclotomic matrices and hypergeometric functions over finite fields, Finite Fields Appl. 82 (2022), Article 102054.

Department of Science and Mathematics, Indian Institute of Information Technology Guwahati, Kamrup-781015, Assam, INDIA

Email address: rituparnachaliha5@gmail.com

Department of Mathematics, Bhattadev University, Bajali, Pathsala, Assam-781325, INDIA Email address: haigautam@gmail.com