

# Long Orbit Empty Value (LOEV) principle in general\*

M. Ivanov<sup>†</sup>, D. Kamburova<sup>‡</sup>, N. Zlateva<sup>§</sup>

## Abstract

We explore LOEV method in spaces more general than metric.

## 1 Introduction

In the article [6] the LOEV method was introduced in complete metric spaces. It is an abstract Lemma which says that provided a multivalued map satisfies a property called (\*), certain conclusion can be made about the orbits of this map.

By specifying the multivalued map, many results can be derived in an unified fashion.

The purpose of this article is to explore generalizations of LOEV method in more general spaces, where a distance function with certain properties is available, but not a metric.

In Section 2 we lay the foundation in a general topological space. Although it is possible to work only with neighbourhoods, because the distance function is not mentioned in the claim and is, therefore, somewhat auxiliary, we prefer to still use a distance function for the sake of consistency. Note that unlike in [6] no completeness of any kind is assumed and, therefore, the main result is new even in metric space.

In Section 3 we move towards completeness. Since our spaces are uniform, completeness can be considered in the standard topological way, but then essentially nothing new, compared to Section 2, will be achieved. It is far more interesting, as spotted in [13], to work with sigma-semicompleteness, because we get new results, map the link to the correspondent Ekeland Variational Principle, and even characterize the space.

---

\*Research supported by European Union-NextGenerationEU, through the Bulgarian National Recovery and Resilience Plan, SUMMIT project BG-RRP-2.004-0008-C01.

<sup>†</sup>Radiant Life Technologies Ltd., Nicosia, Cyprus, e-mail:milen@radiant-life-technologies.com

<sup>‡</sup>Sofia University, Faculty of Mathematics and Informatics, 5 James Bourchier Blvd., 1164 Sofia, Bulgaria, e-mail:detelinak@fmi.uni-sofia.bg

<sup>§</sup>Sofia University, Faculty of Mathematics and Informatics, 5 James Bourchier Blvd., 1164 Sofia, Bulgaria, e-mail:zlateva@fmi.uni-sofia.bg

## 2 LOEV principle in a topological framework

First, we set the formal framework for the following study.

We work in a first countable topological space, usually denoted by  $(X, \tau)$ . We consider multivalued map  $S : X \rightrightarrows X$ . Its domain is:

$$\text{dom } S := \{x \in X : S(x) \neq \emptyset\}.$$

A (finite or infinite) succession of points in  $X$  satisfying  $x_{i+1} \in S(x_i)$  for  $i = 0, 1, 2, \dots$  is called an orbit of  $S$ , or  $S$ -orbit, starting at  $x_0$ . We say that a  $S$ -orbit ends at  $x \in X$  if the orbit is finite and  $x_n = x$ , or the orbit is infinite and the sequence  $\{x_i\}_{i=0}^{\infty}$  converges to  $x$ . If a  $S$ -orbit is infinite and the sequence  $\{x_i\}_{i=0}^{\infty}$  diverges, we say that it is a divergent  $S$ -orbit.

For convenience, we impose *non-stationarity* condition:

$$x \notin S(x), \quad \forall x \in X. \quad (1)$$

The meaning of this condition is that the stationary sequence  $x, x, x, \dots$  is never a  $S$ -orbit. Obviously, it can be easily ensured whenever necessary, by considering instead of  $S$  the map  $x \rightrightarrows S(x) \setminus \{x\}$ .

Recall, see [6], that  $S$  satisfies property  $(*)$  if  $S$  satisfies (1) and for each  $y \in S(x)$  and each sequence  $x_i \rightarrow x$ , there is a subsequence  $\{x_{i_k}\}_{k=1}^{\infty}$  such that  $y \in S(x_{i_k})$  for all  $k \in \mathbb{N}$ .

The property  $(*)$  is very practical, as can be seen from the examples in [6], but it can be easily relaxed in several directions, most obvious of which is to consider only sequences starting from a certain fixed point.

**Definition 2.1** *The multivalued map  $S : X \rightrightarrows X$ , where  $(X, \tau)$  is a first countable topological space, satisfies the property  $(*_1)$  for  $x_0 \in X$ , if  $S$  satisfies (1) and for each infinite  $S$ -orbit  $\{x_i\}_{i=0}^{\infty}$  starting at  $x_0$  and ending at  $x$  (that is,  $x_i \rightarrow x$ ) and each  $y \in S(x)$ , there is a subsequence  $\{x_{i_k}\}_{k=1}^{\infty}$  such that*

$$y \in S(x_{i_k}), \quad \forall k \in \mathbb{N}.$$

Next, we define what we require from the spaces we work on.

**Definition 2.2** *We will call the triple  $(X, \tau, h)$  a h-space, if  $(X, \tau)$  is first countable Hausdorff space and the distance function  $h : X \times X \rightarrow \mathbb{R}^+$  is such that*

- (i)  $h(x, y) = 0 \Leftrightarrow x = y$ ;
- (ii) if  $h(x, x_n) \rightarrow 0$ , then  $x_n \rightarrow x$ ;
- (iii) if  $x_n \rightarrow x$ , then  $h(x_{n+1}, x_n) \rightarrow 0$ .

What follows is our first main result, which is a far reaching generalization of LOEV method.

**Theorem 2.3** *Let  $(X, \tau, h)$  be a  $h$ -space. Let  $S : X \rightrightarrows X$  be a multi valued map satisfying  $(*_1)$  for  $x_0$ . Then at least one of (a) and (b) below is true:*

- (a) *There is a divergent  $S$ -orbit starting at  $x_0$ ;*
- (b) *There is a  $S$ -orbit starting at  $x_0$  and ending at  $x \in X$  such that  $S(x) = \emptyset$ .*

**Proof.** Assume that (a) is not true, that is each  $S$ -orbit starting at  $x_0$  is either finite or infinite and convergent. We can construct finite or infinite  $S$ -orbit by the following procedure: if  $x_0, x_1, \dots, x_i$  are already chosen, then

- if  $S(x_i) = \emptyset$  we are done,
- otherwise  $s_i := \min\{1, \sup_{y \in S(x_i)} h(y, x_i)\} > 0$  and we take  $x_{i+1} \in S(x_i)$  with

$$h(x_{i+1}, x_i) \geq \frac{s_i}{2}.$$

If we end with an infinite orbit, then since (a) is assumed false,  $\{x_i\}_{i \geq 0}$  is a convergent to some  $x \in X$  sequence. Observe that in this case by (iii)  $h(x_{i+1}, x_i) \rightarrow 0$ , and then  $s_i \rightarrow 0$ . We will show that  $S(x) = \emptyset$ . Assume the contrary, i.e., that there exists  $y \in S(x)$ ,  $y \neq x$ . By property  $(*_1)$  of  $S$  for  $x_0$ , there exists a subsequence  $\{x_{i_k}\}_{k \geq 1}$  of  $\{x_i\}_{i \geq 0}$  such that  $y \in S(x_{i_k})$  for all  $k \in \mathbb{N}$ .

Observe that for all  $k \in \mathbb{N}$ ,  $h(y, x_{i_k}) \leq s_{i_k}$  since  $y \in S(x_{i_k})$  and  $s_{i_k} \rightarrow 0$ . Then  $h(y, x_{i_k}) \rightarrow 0$  and by (ii)  $x_{i_k} \rightarrow y$ . But  $x_{i_k} \rightarrow x$  and  $x \neq y$ . Since  $X$  is a Hausdorff space this yields a contradiction. The proof is then completed.  $\square$

Often an intermediate property – between  $(*)$  and  $(*_1)$  – will be satisfied and it is worth considering.

**Definition 2.4** *The multivalued map  $S : X \rightrightarrows X$ , where  $(X, \tau)$  is a first countable topological space, satisfies the property  $(*_2)$  for  $x_0 \in X$ , if  $S$  satisfies (1) and for each infinite  $S$ -orbit  $\{x_i\}_{i=0}^\infty$  starting at  $x_0$  and each subsequence  $\{x_{i_k}\}_{k=1}^\infty$  converging to  $x$ , and each  $y \in S(x)$ , there is a subsequence  $\{x_{i_{k_j}}\}_{j=1}^\infty$  such that*

$$y \in S(x_{i_{k_j}}), \quad \forall j \in \mathbb{N}.$$

Obviously,  $(*) \Rightarrow (*_2) \Rightarrow (*_1)$ . More precisely,  $(*_2)$  for  $x_0$  implies  $(*_1)$  for  $x_0$ , while  $(*)$  implies  $(*_2)$  for all  $x$ .

If  $S$  is *monotone*, that is, if

$$S(S(x)) \subset S(x), \quad \forall x \in X, \tag{2}$$

then, of course, each subsequence of an infinite  $S$ -orbit is again an  $S$ -orbit, so  $(*_1) \iff (*_2)$  for a monotone  $S$ . Another way of defining the monotonicity of  $S$  is to say that

$$y \preceq x \iff y \in \{x\} \cup S(x)$$

defines a partial ordering on  $X$ . It will be noted later that for the most of applications found in the literature so far do rely on introducing some partial ordering.

We, on the other hand, need not impose monotonicity on  $S$ . Yet, this relation to ordering justifies the terms "monotone" that we choose in favor of the standard name for (2), that is *idempotent*.

Under  $(*_2)$  we can specify the nature of potential divergent orbit in Theorem 2.3.

**Proposition 2.5** *Let  $(X, h)$  be a first countable Hausdorff topological space. Let  $S : X \rightrightarrows X$  be a multi valued map satisfying  $(*_2)$  for  $x_0$ . Then at least one of the following is true.*

(a) *There is a  $S$ -orbit starting at  $x_0$ , say  $\{x_i\}_{i \geq 0}$ , such that*

$$\overline{\{x_i\}_{i \geq 0}} \not\subset \text{dom } S.$$

(b) *There is a divergent  $S$ -orbit starting at  $x_0$ , say  $\{x_i\}_{i=0}^\infty$ , such that*

$$\limsup_{i \rightarrow \infty} h(x_{i+1}, x_i) > 0.$$

(c) *There is a divergent  $S$ -orbit starting at  $x_0$  with no convergent subsequences.*

**Proof.** Let (a) be false, that is, there are no  $S$ -orbits starting at  $x_0$  and ending outside of the domain of  $S$  and, moreover, each subsequence of any  $S$ -orbit starting at  $x_0$  is either divergent or converges to a point in  $\text{dom } S$ .

Repeating the construction from the proof of Theorem 2.3 we get an infinite  $S$ -orbit  $\{x_i\}_{i=0}^\infty$  such that

$$h(x_{i+1}, x_i) \geq \min\{1, \sup_{y \in S(x_i)} h(y, x_i)\}/2.$$

If this orbit has no convergent subsequences, we are done, because (c) holds.

So, let  $x_{i_k} \rightarrow x$ , as  $k \rightarrow \infty$ . Since (a) is false,  $S(x) \neq \emptyset$ , so there is  $y \neq x$  such that  $y \in S(x)$ . From  $(*_2)$  there is further subsequence  $\{x_{i_{k_j}}\}_{j=1}^\infty$  such that  $y \in S(x_{i_{k_j}})$ . If (b) is not true, then

$$\min\{1, h(y, x_{i_{k_j}})\} \leq 2h(x_{i_{k_j}+1}, x_{i_{k_j}}) \rightarrow 0,$$

as  $j \rightarrow \infty$ . By (ii) this implies  $x_{i_{k_j}} \rightarrow y$ , as  $j \rightarrow \infty$ , and, therefore, because  $X$  is Hausdorff,  $x = y$ , contradiction.  $\square$

**Remark 2.6** *It is possible to characterize the topological spaces  $(X, \tau)$  on which there exists a distance function  $h$  turning  $(X, \tau, h)$  into a  $h$ -space.*

Indeed, consider a topological space  $(X, \tau)$  that has countable nested local base  $\{U_n(x)\}_{n \geq 1}$  at each point  $x \in X$  (that is,  $U_n(x) \in \tau$ ,  $U_{n+1}(x) \subset U_n(x)$  and  $\bigcap_n U_n(x) = \{x\}$ ) for which the following axioms hold:

(A1) for any  $\bar{x} \in X$ , and any  $n \in \mathbb{N}$  there exists  $m \in \mathbb{N}$  such that

$$U_m(x) \subset U_n(\bar{x}), \quad \forall x \in U_m(\bar{x});$$

(A2) if a sequence  $\{x_n\}_{n \geq 1}$  converges to  $x \in X$ , then  $\inf\{m \in \mathbb{N} : x_n \in U_m(x_{n+1})\}$  tends to infinity.

The function  $h : X \times X \rightarrow \mathbb{R}^+$  defined as

$$h(x, y) := \sup \left\{ \frac{1}{m} : y \in U_m(x) \right\}$$

satisfies (i)-(iii), so  $(X, \tau, h)$  is a  $h$ -space.

Let us observe that (A2) can be reformulated in terms of  $h$  as: if a sequence  $\{x_n\}_{n \geq 1}$  converges to  $x \in X$ , then  $h(x_{n+1}, x_n) \rightarrow 0$  which is (iii).

Obviously, (A1) yields (ii).

### 3 LOEV principle in $\Sigma_g$ semicomplete premetric space

Let  $X$  be a topological space. A distance function  $g : X \times X \rightarrow \mathbb{R}^+$  with the following properties:

(P1)  $g(x, x) = 0$  for any  $x \in X$  and  $g(x, y) > 0$  whenever  $x \neq y$ ;

(P2)  $g(x, \cdot)$  is continuous for every fixed  $x \in X$

is called a premetric function on  $X$ . If there exists a premetric function  $g$  on a topological space  $X$ , then necessarily  $X$  is a Hausdorff space. A topological space  $X$  with premetric function  $g$  on it will be called a premetric space and will be denoted by  $(X, g)$ .

**Example 3.1** *Let  $X$  be a completely regular space with countable local base at each point. Then  $X$  is a Hausdorff space. For every point  $x \in X$  and a closed set  $C \subset X$ ,  $x \notin C$ , there is a continuous function  $h : X \rightarrow [0, 1]$ , such that  $h(x) = 0$  and  $h|_C \equiv 1$ . Fix a point  $x \in X$  and let  $\{U_k\}_k$  be a countable (nested) family of open neighbourhoods of  $x$ . For any  $k \in \mathbb{N}$  let  $p_k : X \rightarrow [0, 1]$  be a continuous bounded function such that  $p_k(x) = 0$  and  $p_k|_{X \setminus U_k} \equiv 1$ . Set*

$$p_x(y) := \sum_{k=1}^{\infty} \frac{1}{2^k} p_k(y).$$

*The function  $g : X \times X \rightarrow [0, 1]$  defined as  $g(x, y) := p_x(y)$  is a premetric function. Moreover, if  $g(x, y_n) \rightarrow 0$ , then  $y_n \rightarrow x$ .*

$\Sigma$ -Cauchy sequences and  $\Sigma$  semicompleteness are considered in Suzuki [13]. We extend these notions in a premetric space  $(X, g)$  in the following way.

For a sequence  $\{x_n\}_{n \geq 0}$  in a premetric space  $(X, g)$ ,  $\sum_{i=0}^{\infty} g(x_{i+1}, x_i)$  can be considered as a  $g$ -length of the sequence. If a sequence  $\{x_n\}_{n \geq 0}$  in a premetric space  $(X, g)$  has a finite  $g$ -length, it will be called  $\Sigma_g$ -Cauchy sequence.

**Definition 3.2** *A first countable premetric space  $(X, g)$  is called  $\Sigma_g$  semicomplete if every  $\Sigma_g$ -Cauchy sequence in  $X$  has a subsequence convergent to an element of  $X$ .*

First we will prove in a  $\Sigma_g$  semicomplete space a variant of the Long Orbit Empty Value (LOEV) principle.

**Theorem 3.3** *Let  $(X, g)$  be  $\Sigma_g$  semicomplete space, let  $S : X \rightrightarrows X$  satisfy  $(*_2)$  for  $x_0 \in X$ . Then at least one of (a) and (b) below is true:*

- (a) *There is  $S$ -orbit starting at  $x_0$  with infinite  $g$ -length;*
- (b) *There is  $S$ -orbit starting at  $x_0$  and ending at  $x \notin \text{dom } S$ .*

**Proof.** Assume that (a) is not true, that is, each  $S$ -orbit, starting at  $x_0$  has finite  $g$ -length. We can construct finite or infinite  $S$ -orbit starting at the given  $x_0$  by the following procedure: if  $x_0, x_1, \dots, x_i$  are already chosen, then

- either  $S(x_i) = \emptyset$ , and we are done;
- or  $s_i := \min\{1, \sup g(y, x_i) : y \in S(x_i)\} > 0$ . Take  $x_{i+1} \in S(x_i)$  such that

$$g(x_{i+1}, x_i) \geq \frac{s_i}{2}. \quad (3)$$

If we end up with infinite  $S$ -orbit  $\{x_i\}_{i \geq 0}$  then, since (a) is assumed false,

$$\sum_{i=0}^{\infty} g(x_{i+1}, x_i) < \infty,$$

i.e.  $\{x_i\}_{i \geq 0}$  is a  $\Sigma_g$ -Cauchy sequence. From the  $\Sigma_g$  semicompleteness of the space  $(X, g)$ , it follows that there is a convergent subsequence  $\{x_{i_k}\}_k$ . Let  $x_{i_k} \rightarrow x \in X$ , as  $k \rightarrow \infty$ . Note that

$$\frac{1}{2} \sum_{i=0}^{\infty} s_i \leq \sum_{i=0}^{\infty} g(x_{i+1}, x_i) < \infty,$$

and since  $s_i > 0, \forall i \geq 0$ ,

$$\sum_{k=0}^{\infty} s_{i_k} \leq \sum_{i=0}^{\infty} s_i < \infty.$$

Therefore,

$$\{s_{i_k}\} \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (4)$$

From (6) and from the definition of  $s_{i_k}$  it follows that for sufficiently large  $k$ ,

$$1 > s_{i_k} = \sup\{g(y, x_{i_k}) : y \in S(x_{i_k})\} > 0. \quad (5)$$

Assume that  $S(x) \neq \emptyset$  and take  $y \in S(x)$ ,  $y \neq x$ . Since  $g$  has property (P1), we have that  $g(y, x) > 0$ . Moreover, by property  $(*_2)$  of  $S$  at  $x_0$ ,  $y \in S(x_{i_{k_j}})$  for  $j \in \mathbb{N}$  where  $x_{i_{k_j}}$  is a convergent to  $x$  subsequence. By (P2)  $g(y, \cdot)$  is continuous, so when  $j \rightarrow \infty$ ,  $g(y, x_{i_{k_j}}) \rightarrow g(y, x) = \delta > 0$ . Since

$$\sum_{k=0}^{\infty} s_{i_{k_j}} \leq \sum_{i=0}^{\infty} s_{i_k} < \infty,$$

we have that

$$s_{i_{k_j}} \rightarrow 0 \text{ as } j \rightarrow \infty. \quad (6)$$

On the other hand,  $s_{i_{k_j}} \geq g(y, x_{i_{k_j}}) \geq \delta/2$  for sufficiently large  $j$ . The latter means in particular that the sequence  $\{s_{i_{k_j}}\}_j$  does not converge to zero as  $j \rightarrow \infty$  which contradicts (6). The proof is then complete.  $\square$

**Remark 3.4** *If a sequence  $\{z_k\}_{k \geq 0}$  in a premetric space  $(X, g)$  has no convergent subsequences, then there exists its subsequence  $\{x_n\}_{n \geq 0}$ ,  $x_n = z_{k_n}$ , for all  $n \geq 0$  such that  $x_{n+1} \neq x_n$  for all  $n \geq 0$  with the same  $g$ -length. Indeed, we need only to omit the consecutive members of the sequence that repeat observing that they are necessarily finitely many and contribute with zero terms in the sum defining the length because of the property (P1) of  $g$ . Hence, if the former  $\{z_k\}_{k \geq 0}$  is  $\Sigma_g$ -Cauchy, the so constructed  $\{x_n\}_{n \geq 0}$  will be  $\Sigma_g$ -Cauchy too and with the same  $g$ -length. If we denote by  $M$  the set consisting of all points of  $\{x_n\}_{n \geq 0}$ , then  $M$  is a discrete set, i.e. any point in  $M$  is an isolated point.*

The next result shows that if LOEV principle holds in a premetric space  $(X, g)$ , then necessarily the space is  $\Sigma_g$  semicomplete.

**Theorem 3.5** *Let  $(X, g)$  be a first countable premetric space. If for any  $x_0 \in X$  and any  $S : X \rightrightarrows X$  that satisfies  $(*_2)$  for  $x_0$  at least one of (a) and (b) below is true:*

- (a) *There is  $S$ -orbit starting at  $x_0$  with infinite  $g$ -length;*
- (b) *There is  $x \in X$  such that  $S(x) = \emptyset$ ,*

*then the space  $(X, g)$  is  $\Sigma_g$  semicomplete.*

**Proof.** Suppose that  $(X, g)$  is not  $\Sigma_g$  semicomplete. Then there exists a  $\Sigma_g$ -Cauchy sequence  $\{x_n\}_{n \geq 0}$  that does not have any convergent subsequence,  $x_{n+1} \neq x_n$  for all  $n$  and the set  $M$  consisting of all points in the sequence is a discrete set, see Remark 3.4. Define the multi valued map  $S$  in the following way:

$$S(x) := \begin{cases} M, & \text{if } x \notin M; \\ x_{n+1}, & \text{if } x = x_n \text{ for some } x_n \in M. \end{cases}$$

Note that  $S(x) \neq \emptyset$  for all  $x \in X$ , therefore (b) does not hold. If  $x \notin M$ , since there is no cluster point of the sequence, then there is a neighbourhood  $N_x \ni x$  such that  $N_x \cap M = \emptyset$  and  $S(x') = S(x) = M$  for every  $x' \in N_x$ . If  $x = x_n \in M$ , again, there is a neighbourhood  $N_{x_n} \ni x_n$  such that  $N_{x_n} \cap M \setminus \{x_n\} = \emptyset$  and  $x_{n+1} = S(x_n) \subset S(x') = M$  for all  $x' \in N_{x_n} \setminus \{x_n\}$  because  $M$  is an discrete set. Therefore,  $S$  has property  $(*_2)$  for  $x_0$ . Since the unique  $S$ -orbit starting at  $x_0$  is the given sequence  $\{x_n\}_{n \geq 0}$  it is  $\Sigma_g$ -Cauchy. Hence, it is with finite  $g$ -length and (a) does not hold too. The contradiction completes the proof.  $\square$

The above result shows that the LOEV principle is, roughly speaking, equivalent to the  $\Sigma_g$  semicompleteness of the first countable premetric space.

Further we will prove variants of the Caristi theorem, Takahashi theorem and Ekeland variational principle in  $\Sigma_g$  semicomplete space and establish that they are all equivalent to the  $\Sigma_g$  semicompleteness.

We begin with a variant of the Caristi theorem, see [1].

**Theorem 3.6** *Let  $(X, g)$  be a  $\Sigma_g$  semicomplete space. Let the multi-valued map  $T : X \rightrightarrows X$  satisfy  $T(x) \neq \emptyset$  for all  $x \in X$ . Let the function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be proper, lower semicontinuous and bounded below. Assume that for any  $x \in X$  there exists  $y \in T(x)$  such that*

$$g(y, x) \leq f(x) - f(y). \quad (7)$$

*Then  $T$  has a fixed point, that is, there exists  $\bar{x} \in X$  such that  $\bar{x} \in T(\bar{x})$ .*

**Proof.** Assume that  $x \notin T(x)$  for all  $x \in X$ . Consider the map

$$S(x) := \{y \in X : 2^{-1}g(y, x) < f(x) - f(y)\}.$$

Since for each  $x \in X$  there is some  $y \in T(x)$ ,  $y \neq x$  satisfying (7) and by (P1),  $g(y, x) > 0$ ,  $S(x) \neq \emptyset$  for all  $x \in X$ . By property (P1) of  $g$ ,  $x \notin S(x)$  for all  $x \in X$ .  $S$  has the property  $(*_2)$  for any point  $x_0$  because of the lower semicontinuity of  $f$  and continuity of  $g(y, \cdot)$ , see (P2). We apply Theorem 3.3 to get a  $S$ -orbit  $\{x_i\}_{i \geq 0}$  with infinite  $g$ -length. From the the definition of  $S$  it follows that  $2^{-1}g(x_{i+1}, x_i) < f(x_i) - f(x_{i+1})$  for all  $i \geq 0$ . Summing the inequalities we obtain that

$$2^{-1} \sum_{i=0}^{n-1} g(x_{i+1}, x_i) < f(x_0) - f(x_n), \quad \forall n \geq 1.$$



Passing  $i$  to infinity, we get  $f(x_n) \rightarrow -\infty$  which contradicts boundedness below of  $f$ . The proof is then complete.  $\square$

Now we will prove that if the variant of Caristi theorem holds in a first countable premetric space  $(X, g)$ , then necessarily it is  $\Sigma_g$  semicomplete.

**Theorem 3.7** *Let  $(X, g)$  be a first countable premetric space. If every multi-valued map  $T : X \rightrightarrows X$  satisfying Caristi condition (7) with some proper lower semicontinuous and bounded below function  $f$  has a fixed point, then  $(X, g)$  is  $\Sigma_g$  semicomplete.*

**Proof.** Suppose that  $(X, g)$  is not  $\Sigma_g$  semicomplete. Then there exists a  $\Sigma_g$ -Cauchy sequence  $\{x_n\}_{n \geq 0}$  that does not have any convergent subsequence,  $x_{n+1} \neq x_n$  for all  $n$  and the set  $M$  consisting of all points of the sequence is a discrete set, see Remark 3.4. Define the function

$$f(x) := \begin{cases} \sum_{i=n}^{\infty} g(x_{i+1}, x_i), & \text{if } x = x_n \in M; \\ +\infty, & \text{otherwise} \end{cases} \quad (8)$$

It is obviously a bounded below by zero function such that  $\text{dom } f \equiv M$ , hence proper. Since  $M$  consists of isolated points,  $f$  is also lower semicontinuous.

Consider the map  $T(x) := \{y \in X : y \neq x, g(y, x) \leq f(x) - f(y)\}$ . If  $x \notin M$ , then  $M \in T(x)$ . If  $x = x_n$  for some  $x_n \in M$ , then  $x_{n+1} \in T(x_n)$ . Hence,  $T(x) \neq \emptyset$  for all  $x \in X$ . Since the map  $T$  satisfies the Caristi condition with the function  $f$ ,  $T$  has a fixed point by assumption, i.e., there is  $\bar{x} \in T(\bar{x})$ . This yields a contradiction, because  $x \notin T(x)$  for all  $x \in X$  according to the definition of  $T$ .  $\square$

In  $\Sigma_g$  semicomplete space we can prove also a variant of Takahashi theorem, see [14].

**Theorem 3.8** *Let  $(X, g)$  be a  $\Sigma_g$  semicomplete space. Let the function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be proper, lower semicontinuous and bounded below. Suppose that for each  $x \in X$  with  $f(x) > \inf_X f$  there exists  $y \in X$ ,  $y \neq x$ , such that*

$$g(y, x) \leq f(x) - f(y), \quad (9)$$

*Then, there exists  $v \in X$  such that  $f(v) = \inf_X f$ .*

**Proof.** Assume that the condition holds but  $f(x) > \inf_X f$  for all  $x \in X$ . This means that for any  $x \in X$  there exists  $y \in X$ ,  $y \neq x$ , such that

$$2^{-1}g(y, x) < g(y, x) \leq f(x) - f(y),$$

where the strict inequality follows from (P1). Then for all  $x \in X$ ,

$$S(x) := \{y \in X : y \neq x, 2^{-1}g(y, x) < f(x) - f(y)\} \neq \emptyset,$$

and  $x \notin S(x)$ . The lower semicontinuity of  $f$  and continuity of  $g$  of the second variable due to (P2) imply that  $S$  has the property  $(*_2)$  for any  $x_0 \in X$ . Hence, we can apply Theorem 3.3 to get a  $S$ -orbit  $\{x_n\}_{n \geq 0}$  with infinite  $g$ -length. From the definition of  $S$  it holds that  $2^{-1}g(x_{i+1}, x_i) < f(x_i) - f(x_{i+1})$  for all  $i \geq 0$  and summing the inequalities and passing to infinity, we obtain that  $f(x_n) \rightarrow -\infty$  which contradicts the boundedness below of  $f$ .  $\square$

As it is expected, if the variant of the Takahashi theorem holds in a first countable premetric space  $(X, g)$ , then necessarily the space is  $\Sigma_g$  semicomplete.

**Theorem 3.9** *Let  $(X, g)$  be a first countable premetric space. If for every proper, lower semicontinuous and bounded below function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  that satisfies (9) condition there exists  $v \in X$  such that  $f(v) = \inf_X f$ , then  $(X, g)$  is  $\Sigma_g$  semicomplete.*

**Proof.** Suppose that  $X$  is not  $\Sigma_g$  semicomplete. Then there exists a  $\Sigma_g$ -Cauchy sequence  $\{x_n\}_{n \geq 0}$  that does not have any convergent subsequence,  $x_{n+1} \neq x_n$  for all  $n$  and the set  $M$  consisting of all points in the sequence is a discrete set, see Remark 3.4. Consider the lower semicontinuous  $f : X \rightarrow \mathbb{R}^+$  with  $\text{dom } f = M$  and  $\inf_X f = 0$  defined by (8). Note that  $f(x_k) - f(x_m) = \sum_{i=k}^{m-1} g(x_{i+1}, x_i)$  whenever  $m > k \geq 0$ , and in particular,  $f(x_n) - f(x_{n+1}) = g(x_{n+1}, x_n)$  for all  $n \geq 0$ .

Takahashi condition holds for  $f$  because it is trivially fulfilled for points  $x \notin M$ , while for any point  $x \in M$  there is some  $x_n$  such that  $x = x_n$  and since  $x_{n+1} \neq x_n$ ,  $2^{-1}g(x_{n+1}, x_n) < g(x_{n+1}, x_n) = f(x_{n+1}) - f(x_n)$ . Hence, there exists  $v \in X$  such that  $0 = \inf_X f = f(v)$ . Note that necessarily  $v \in M$ . Therefore, there exists  $n \geq 0$  such that  $v = x_n$ . But having in mind that  $f(x_n) = 0$  and  $g(x_{n+1}, x_n) > 0$  (because of (P1) for  $g$ ) we get

$$0 \geq -f(x_{n+1}) = f(x_n) - f(x_{n+1}) = g(x_{n+1}, x_n) > 0,$$

contradiction.  $\square$

Now we will prove in a  $\Sigma_g$  semicomplete space a variant of Ekeland theorem [3].

**Theorem 3.10** *Let  $(X, g)$  be a  $\Sigma_g$  semicomplete space. Let the function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be proper, lower semicontinuous and bounded below. Then for every  $\varepsilon > 0$  there exists  $v \in X$ , such that*

$$f(v) \leq f(x) + \varepsilon g(x, v) \quad \text{for all } x \in X. \quad (10)$$

**Proof.** Assume that there is no point  $v \in X$  such that (10) holds. Then for every  $x \in X$  it should be some  $y \in X$  such that  $f(x) - \varepsilon g(y, x) > f(y)$ .

Consider the multi-valued map  $S$  defined as

$$S(x) := \{y \in X : f(y) < f(x) - \varepsilon g(y, x)\}.$$

By our assumption  $S(x) \neq \emptyset$  for all  $x \in X$ . By property (P1) of the function  $g$ ,  $x \notin S(x)$  for all  $x \in X$ . The lower semicontinuity of  $f$  and (P2) property of  $g$  imply that  $S$  satisfies  $(*_2)$  at any point  $x_0 \in X$ . So we can apply Theorem 3.3 to get a  $S$ -orbit  $\{x_n\}_{n \geq 0}$  with infinite  $g$ -length. But the definition of  $S$  implies that  $f(x_{i+1}) - f(x_i) < -\varepsilon g(x_{i+1}, x_i)$  for all  $i \geq 0$ . Summing the first  $n$  inequalities we get

$$f(x_n) - f(x_0) < -\varepsilon \sum_{i=0}^{n-1} g(x_{i+1}, x_i)$$

and passing  $n$  to infinity we get that  $f(x_n) \rightarrow -\infty$ . The latter contradicts the boundedness below of  $f$ .  $\square$

Conversely, if the variant of the Ekeland theorem holds in a first countable premetric space  $(X, g)$ , then necessarily the space is  $\Sigma_g$  semicomplete.

**Theorem 3.11** *Let  $(X, g)$  be a first countable premetric space. If for every  $\varepsilon > 0$  and every proper lower semicontinuous and bounded below function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  there exists  $v \in X$  such that (10) holds, then  $(X, g)$  is  $\Sigma_g$  semicomplete.*

**Proof.** Suppose that  $X$  is not  $\Sigma_g$  semicomplete. Then there exists a  $\Sigma_g$ -Cauchy sequence  $\{x_n\}_{n \geq 0}$  that does not have any convergent subsequence,  $x_{n+1} \neq x_n$  for all  $n$  and the set  $M$  consisting of all points in the sequence is a discrete set, see Remark 3.4. Consider the lower semicontinuous  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  with  $\text{dom } f \equiv M$  and  $\inf_X f = 0$  defined by (8) and recall that  $f(x_n) - f(x_{n+1}) = g(x_{n+1}, x_n)$  for all  $n \geq 0$ .

Then for every  $1 > \varepsilon > 0$  there exists  $v \in X$  such that for all  $x \in X$ ,

$$f(v) \leq f(x) + \varepsilon g(x, v).$$

Note that necessarily  $v \in M$ . Therefore, there exists  $n$  such that  $v = x_n$ . For  $x = x_{n+1}$ , we have that

$$f(x_n) \leq f(x_{n+1}) + \varepsilon g(x_{n+1}, x_n),$$

hence,

$$g(x_{n+1}, x_n) \leq \varepsilon g(x_{n+1}, x_n).$$

Since  $\varepsilon \in (0, 1)$ , this yields that  $g(x_{n+1}, x_n) = 0$ . But by (P1) of  $g$ , and  $x_{n+1} \neq x_n$ ,  $g(x_{n+1}, x_n) > 0$ , which yields a contradiction.  $\square$

## References

- [1] Caristi, J.: Fixed point theorems for mappings satisfying inwardness conditions. *Trans. Amer. Math. Soc.* 215, 241–251 (1976)
- [2] Cullen, H.F.: *Introduction to General Topology*. Heath, Boston (1967)
- [3] Ekeland, I.: On the variational principle. *J. Math. Anal. Appl.* 47(2), 324–353 (1974)
- [4] Gaumont, D., Kamburova, D., Revalski, J.P.: Perturbations of supinf problems with constraints. *Vietnam J. Math.* 47, 659–667 (2019)
- [5] Hamel, A. H.: Equivalent to Ekeland’s variational principle in uniform spaces. *Nonlinear Anal.* 62, 913–924 (2005)
- [6] Ivanov, M., Zlateva N.: Long orbit or empty value principle, fixed point and surjectivity theorems, *Compt. rend. Acad. bulg. Sci.* 69(5), 553–562 (2016).
- [7] Kenderov, P., Lucchetti, R.: Generic well-posedness of supinf problems. *Bull. Austr. Math. Soc.* 54, 5–25 (1996)
- [8] Kenderov, P., Revalski, J.P.: Dense existence of solutions of perturbed optimization problems and topological games. *C. R. Acad. Bulgare Sci.* 63(7), 937–942 (2010)
- [9] Kenderov, P., Revalski, J.P.: Variational principles for supinf problems. *C. R. Acad. Bulgare Sci.* 70(12), 1635–1642 (2017)
- [10] McLinden L.: An application of Ekeland’s theorem to minimax problems, *Nonlinear Anal.* 6(2), 189–196 (1982)
- [11] Oettli, W., Théra, M.: Equivalent to Ekeland’s principle. *Bull. Austr. Math. Soc.* 48(3), 385–392 (1993)
- [12] Sullivan F.: A characterization of complete metric spaces. *Proc. Amer. Math. Soc.* 83(2), 345–346 (1981)
- [13] Suzuki T.: Characterization of  $\Sigma$ -semicompleteness via Caristi’s fixed point theorem in semimetric spaces. *Journal of Function Spaces.* article ID 9435470, 1–7 (2018)
- [14] Takahashi, W.: Existence theorems generalizing fixed point theorems for multivalued mappings. In: Baillon, J.-B., Théra M. eds. *Fixed Point Theory and Applications*. Longman, 397–406 (1991)