

Trichotomy for the orbits of a hypercyclic operator on a Banach space

Jian Li

ABSTRACT. We obtain a trichotomy for the orbits of a hypercyclic operator T on a separable Banach space X : (1) every vector is mean asymptotic to zero; (2) generic vectors are absolutely mean irregular; (3) every hypercyclic vector is mean divergent to infinity. Examples of weighted backward shifts on ℓ^p show that all three cases can happen.

1. INTRODUCTION

Let $(X, \|\cdot\|)$ be a separable Banach space and $T: X \rightarrow X$ be a bounded linear operator. For a vector $x \in X$, the orbit of x is the set $\{T^n x: n \geq 0\}$, where T^n is the n th composition of T . The study of the behaviors of the orbits of an operator has attracted a lot of attention. Particularly, the Chapter III of the book [3] is entitled *The Orbit of a Linear Operator*. According to [3, Chapter III], by “regular” orbits, we mean for instance that the orbit may tend to infinity, or may tend to zero, or may stay inside a ball centered at 0, or may stay outside a ball centered at 0. Thanks to the Jordan decomposition theorem, every orbit for an operator on a finite-dimensional space is regular, see e.g. [8, Proposition 2.57]. The opposite case is the *irregular orbits*, that is which satisfy

$$\liminf_{n \rightarrow \infty} \|T^n x\| = 0 \text{ and } \limsup_{n \rightarrow \infty} \|T^n x\| = \infty.$$

A special case of irregular orbits are those of hypercyclic vectors. A vector $x \in X$ is called a *hypercyclic vector* if the orbit of x is dense in the whole space, and an operator T is called *hypercyclic* if there is some hypercyclic vector in X . In fact, a hypercyclic vector is called a transitive point in the context of topological dynamics. By the Birkhoff transitivity theorem (see e.g. [8, Theorem 1.16]), if T is hypercyclic then the collection of hypercyclic vectors is a dense G_δ subset of X .

In [4, Theorem 5], Bermúdez et al. characterized Li-Yorke chaos in terms of the existence of irregular vectors. Recall that an operator $T: X \rightarrow X$ is called *Li-Yorke chaotic* if there exists an uncountable subset S of X such that every pair $x, y \in S$ of distinct points we have

$$\liminf_{n \rightarrow \infty} \|T^n x - T^n y\| = 0 \text{ and } \limsup_{n \rightarrow \infty} \|T^n x - T^n y\| > 0.$$

Date: July 11, 2024.

2020 Mathematics Subject Classification. Primary: 47A16; Secondary: 37B05.

Key words and phrases. Hypercyclic operators, orbits, absolutely mean irregular vectors, mean Li-Yorke chaos.

In the study of distributional chaos of operators on Banach spaces [6], Bernardes et al. introduced the concept of absolutely mean irregular vectors, that is a vector $x \in X$ is an *absolutely mean irregular* if

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0 \text{ and } \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

Furthermore, in [5] Bernardes et al. characterized mean Li-Yorke chaos in terms of the existence of absolutely mean irregular vectors, and characterized dense mean Li-Yorke chaos in terms of the existence of a dense (or residual) set of absolutely mean irregular vectors. Recall that an operator $T: X \rightarrow X$ is called (*dense*) *mean Li-Yorke chaotic* if there exists a (dense) uncountable subset S of X such that every pair $x, y \in S$ of distinct points we have

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x - T^i y\| = 0 \text{ and } \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x - T^i y\| > 0.$$

In this paper, we study behaviors of orbits in the mean sense of a hypercyclic operator and obtain the following trichotomy.

Theorem 1.1. *Let T be a bounded linear operator on a separable Banach space X . If T is hypercyclic, then exactly one of three assertions holds:*

- (1) *every vector is mean asymptotic to zero, that is for every $x \in X$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0;$$

- (2) *generic vectors are absolutely mean irregular, that is there exists a residual subset X_0 of X such that for every $x \in X_0$,*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0$$

and

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty;$$

- (3) *every hypercyclic vector is mean divergent to infinity, that is for every hypercyclic vector $x \in X$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

Let $\mathcal{L}(X)$ be the collection of all bounded linear operators from X to itself. The *strong operator topology* on $\mathcal{L}(X)$ is defined as follows: every $T \in \mathcal{L}(X)$ has a neighborhood basis consisting of sets of the form

$$V_{x_1, \dots, x_n, \varepsilon, T} = \{S \in \mathcal{L}(X) : \|Sx_i - Tx_i\| < \varepsilon, i = 1, \dots, n\},$$

where $x_1, \dots, x_n \in X$ and $\varepsilon > 0$. It is not easy to see that, the Rolewicz's operator, $T: \ell^p(\mathbb{N}) \rightarrow \ell^p(\mathbb{N})$, $(x_1, x_2, \dots) \mapsto \lambda(x_2, x_3, \dots)$ with $|\lambda| > 1$, has a residual set of absolutely mean irregular vectors, see e.g. [5, Corollary 30] or Corollary 3.2

in Section 3. The following result reveals that there are plenty of hypercyclic operators with a residual set of absolutely mean irregular vectors.

Theorem 1.2. *If X is an infinite-dimensional separable Banach space, then the collection of hypercyclic operators with a residual set of absolutely mean irregular vectors is dense in $\mathcal{L}(X)$ with respect to the strong operator topology.*

It is shown in [5, Theorem 17] that if $T: X \rightarrow X$ is a bounded linear operator on a separable Banach space X then T is a densely mean Li-Yorke chaotic if and only if it has a residual set of absolutely mean irregular vectors. So we have the following consequence.

Corollary 1.3. *Every infinite-dimensional separable Banach space admits a dense mean Li-Yorke chaotic operator.*

Theorems 1.1 and 1.2 are proved in Sections 2 and 3 respectively. In Section 3 we also strengthen Theorem 1.2 a little bit for operators on a Hilbert space, see Theorem 3.5, and present some examples of weighted backward shifts on ℓ^p for the three cases in Theorem 1.1.

2. PROOF OF THEOREM 1.1

The aim of this section is to prove Theorem 1.1. To this end, we need some preparations.

For a (general) dynamical system, we mean a pair (X, T) , where (X, d) is a metric space and $T: X \rightarrow X$ is a continuous map. Following [10], we say that a dynamical system (X, T) is *mean equicontinuous* if for every $\varepsilon > 0$ there exists $\delta > 0$ such that for every $x, y \in X$ with $d(x, y) < \delta$,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n d(T^i x, T^i y) < \varepsilon,$$

and *mean sensitive* if there exists $\delta > 0$ such that for non-empty open subset U of X , there exist two points x and y in U satisfying

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n d(T^i x, T^i y) \geq \delta.$$

It is shown in [10, Corollary 5.5] that for a dynamical system on a compact metric space, if it is minimal then it is either mean equicontinuous or mean sensitive. The authors in [9] studied mean equicontinuity and mean sensitivity for endomorphisms of completely metrizable groups and obtained the following dichotomy.

Theorem 2.1 ([9, Theorem 4.9]). *Let T be a bounded linear operator on a Banach space X . Then either T is mean equicontinuous or mean sensitive.*

Combining [9, Lemma 4.5 and Proposition 4.33], we have the following result.

Proposition 2.2. *Let T be a bounded linear operator on a Banach space X . Then T is mean sensitive if and only if there exists a residual subset X_0 of X such that for every $x \in X_0$,*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

We have the following property about the orbits of mean equicontinuous operators.

Proposition 2.3. *Let T be a bounded linear operator on a Banach space X . Assume that T is mean equicontinuous.*

(1) *For a vector $x \in X$, if $\liminf_{n \rightarrow \infty} \|T^n x\| = 0$, then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

(2) *The collection of vectors x in X satisfying*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

is a closed subspace of X .

Proof. As T is mean equicontinuous, for every $\varepsilon > 0$ there exists $\delta > 0$ such that for any $u, v \in X$ with $\|u - v\| < \delta$,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i(u - v)\| < \varepsilon.$$

(1) Fix a vector $x \in X$ with $\liminf_{n \rightarrow \infty} \|T^n x\| = 0$. For the above δ , there exists $q \in \mathbb{N}$ such that $\|T^q x\| < \delta$. Then

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i(T^q x)\| < \varepsilon.$$

and

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| &= \limsup_{n \rightarrow \infty} \frac{1}{n+q} \sum_{i=1}^{n+q} \|T^i x\| \\ &= \limsup_{n \rightarrow \infty} \left(\frac{n}{n+q} \frac{1}{n} \sum_{i=1}^n \|T^i(T^q x)\| + \frac{1}{n+q} \sum_{i=1}^q \|T^i x\| \right) \\ &< \varepsilon. \end{aligned}$$

By the arbitrariness of ε , one has

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

(2) By the triangle inequality of norm, it is easy to see that

$$X_0 = \left\{ x \in X : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0 \right\}$$

is a subspace of X . Let $y \in X$ be an accumulation point of X_0 . For the above δ , there exists a vector $z \in X_0$ such that $\|z - y\| < \delta$. Then

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i y\| \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n (\|T^i(z - y)\| + \|T^i z\|) < \varepsilon.$$

By the arbitrariness of ε again, one has

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i y\| = 0.$$

This shows that X_0 is closed. □

We divide the proof of Theorem 1.1 into two parts. First by the dichotomy result of Theorem 2.1, we have the following dichotomy on the orbits, which implies the case (1) in Theorem 1.1 and its opposite side.

Proposition 2.4. *Let T be a bounded linear operator on a separable Banach space X . If T is hypercyclic, then either for every $x \in X$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0$$

or there exists a residual subset X_0 of X such that for every $x \in X_0$

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

Proof. By Theorem 2.1, T is either mean equicontinuous or mean sensitive. First assume that T is mean equicontinuous. For every hypercyclic vector $x \in X$, one has $\liminf_{n \rightarrow \infty} \|T^n x\| = 0$. Then by Proposition 2.3 (1),

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

As the collection of hypercyclic vectors is dense in X , by Proposition 2.3 (2) one has for every $y \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i y\| = 0.$$

Now assume that T is mean sensitive. Then by Proposition 2.2, there exists a residual subset X_0 of X such that for every $x \in X_0$

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

This ends the proof. □

By the proof of Proposition 2.4, we have the following consequence.

Corollary 2.5. *Let T be a bounded linear operator on a Banach space X . If T is hypercyclic then T is mean equicontinuous if and only if for every $x \in X$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

For a subset A of \mathbb{N} , the *upper density* of A is defined by

$$\overline{\text{dens}}(A) = \limsup_{n \rightarrow \infty} \frac{\#(A \cap [1, n])}{n},$$

and the *lower density* of A is

$$\underline{\text{dens}}(A) = \liminf_{n \rightarrow \infty} \frac{\#(A \cap [1, n])}{n},$$

where $\#(\cdot)$ denotes the number of elements of a finite set.

Let $T: X \rightarrow X$ be a bounded linear operator on a Banach space X . We say that T is *frequently hypercyclic* provided that there exists a vector $x \in X$, called a *frequently hypercyclic vector* for T , such that for any non-empty open subset U of X , $\{n \in \mathbb{N}: T^n x \in U\}$ has positive lower density, and *\mathcal{U} -frequently hypercyclic* provided that there exists a vector $x \in X$, called a *\mathcal{U} -frequently hypercyclic vector* for T , such that for any non-empty open subset U of X , $\{n \in \mathbb{N}: T^n x \in U\}$ has positive upper density.

Remark 2.6. Assume that $T: X \rightarrow X$ is \mathcal{U} -frequently hypercyclic. For any \mathcal{U} -frequently hypercyclic vector $x \in X$, one has

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| \geq \limsup_{n \rightarrow \infty} \frac{1}{n} \#(\{1 \leq i \leq n: T^i x \in X \setminus \{y \in X: \|y\| \leq 1\}\}) > 0.$$

By Proposition 2.4, there exists a residual subset X_0 of X such that for every $x \in X_0$,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

Now we consider the case (3) in Theorem 1.1 and its opposite side.

Proposition 2.7. *Let T be an operator on a separable Banach space X . If T is hypercyclic, then either there exists a residual subset X_0 of X such that for every $x \in X_0$*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0,$$

or for every hypercyclic vector $x \in X$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \infty.$$

Proof. Assume that there exists a hypercyclic vector $x \in X$ such that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| < \infty.$$

Let

$$M = \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\|.$$

Then for every $m \in \mathbb{N}$,

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i(T^m x)\| = M.$$

For every $k \in \mathbb{N}$, let

$$X_k = \left\{ y \in X : \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i y\| \leq \frac{M}{k} \right\}.$$

It is easy to see that

$$X_k = \bigcap_{N=1}^{\infty} \left\{ y \in X : \exists n \geq N \text{ s.t. } \frac{1}{n} \sum_{i=1}^n \|T^i y\| < \frac{M}{k} + \frac{1}{N} \right\}.$$

Then X_k is a G_δ subset of X . As x is hypercyclic, $\{T^m x : m \in \mathbb{N}\}$ is dense in X . Note that $\{\frac{1}{k} T^m x : m \in \mathbb{N}\} \subset X_k$. Therefore X_k is dense in X . As

$$X_0 := \left\{ y \in X : \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i y\| = 0 \right\} = \bigcap_{k=1}^{\infty} X_k.$$

we have that X_0 is residual in X . □

Now combining Propositions 2.4 and 2.7, we get Theorem 1.1.

3. PROOF OF THEOREM 1.2 AND SOME EXAMPLES

We say that a bounded linear operator T on a Banach space X satisfies the *Kitai criterion* if there exist two dense subsets E and F of X and a map $S : F \rightarrow F$ such that $TSy = y$, $T^k x \rightarrow 0$ and $S^k y \rightarrow 0$ as $k \rightarrow \infty$ for any $x \in E$ and $y \in F$, and a vector subset Y of X is an *absolutely mean irregular manifold* if every non-zero vector in Y is absolutely mean irregular for T .

Theorem 3.1 ([5, Theorem 29]). *Let T be a bounded linear operator on a separable Banach space X . If*

$$\left\{ x \in X : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0 \right\}$$

is dense in X , then either

- (1) *for every $x \in X$, $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0$, or*
- (2) *T admits a dense absolutely mean irregular manifold.*

Corollary 3.2. *Let T be a bounded linear operator on a separable Banach space X . If T satisfies the Kitai criterion and has a non-trivial periodic point then T admits a dense absolutely mean irregular manifold.*

Proof. Let E be the dense subset of X in the definition of the Kitai criterion. For every $x \in E$, $\lim_{n \rightarrow \infty} \|T^n x\| = 0$, then $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0$. As T has a non-trivial periodic point, the case (1) in Theorem 3.1 can not happen. Then T admits a dense absolutely mean irregular manifold. \square

By the proof of Theorem 1.1, it is easy to see that the collection of absolutely mean irregular vectors is G_δ . If an operator admits a dense absolutely mean irregular manifold, then the collection of absolutely mean irregular vectors is residual. Therefore, Theorem 1.2 is a directly consequence of the following result.

Theorem 3.3. *If X is an infinite-dimensional separable Banach space, then the collection of hypercyclic operators admitting a dense absolutely mean irregular manifold is dense in $\mathcal{L}(X)$ with respect to the strong operator topology.*

Proof. By the proof of Theorem 1.1 in [12], there exists a bounded linear operator $I+T$ on X satisfying the Kitai criterion. Then $I+T$ is mixing (see e.g. [8, Theorem 3.4]). In particular, $I+T$ is hypercyclic. Let x_0 be the point in the proof of Lemma 2.5 in [12]. Then $\|x_0\| = 1$ and $Tx_0 = 0$, which implies that x_0 is a non-trivial fixed point for $I+T$. By Corollary 3.2, $I+T$ admits a dense absolutely mean irregular manifold. By the proof of Theorem 8.18 in [8], the similarity orbit of $I+T$,

$$\mathcal{S}(I+T) = \{A^{-1}(I+T)A; A: X \rightarrow X \text{ invertible}\},$$

is dense in $\mathcal{L}(X)$ with respect to the strong operator topology. It is clear that the property of admitting a dense absolutely mean irregular manifold is invariant under similarity. So the collection of hypercyclic operators with a dense absolutely mean irregular manifold is dense in $\mathcal{L}(X)$ with respect to the strong operator topology. \square

We will need the following result about a sufficient condition for the case (2) in Theorem 1.1. We refer the reader to [1, Chapter 5] about ergodic measures of an operator.

Proposition 3.4. *Let T be a bounded linear operator on a separable Banach space X . If T admits an ergodic measure μ with full support such that $\int_X \|x\| d\mu(x) < \infty$, then there is a residual subset of absolutely mean irregular vectors.*

Proof. As the function $f: X \rightarrow \mathbb{R}$, $x \mapsto \|x\|$, is integrable, by the Birkhoff ergodic theorem (see e.g. [1, Theorem 5.3]) for μ -a.e. vector $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \int_X \|z\| d\mu(z) > 0.$$

As μ is ergodic with full support, the collection of hypercyclic vectors has full measure. In particular, there exists a hypercyclic vector $x \in X$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = \int_X \|z\| d\mu(z) > 0.$$

So cases (1) and (3) in Theorem 1.1 can not happen. This implies that there is a residual subset of absolutely mean irregular vectors. \square

If in addition X is a separable Hilbert space. The *strong* operator topology* on $\mathcal{L}(X)$ is defined as follows: any $T \in \mathcal{L}(X)$ has a neighborhood basis consisting of sets of the form

$$V_{x_1, \dots, x_n, \varepsilon, T}^* = \{S \in \mathcal{L}(X): \|Sx_i - Tx_i\| < \varepsilon, \|S^*x_i - T^*x_i\| < \varepsilon, i = 1, \dots, n\},$$

where $x_1, \dots, x_n \in X$, $\varepsilon > 0$ and T^* is the adjoint operator of T . It is clear that the strong operator topology is coarser than the strong* operator topology. For $M > 0$, the closed ball $\mathcal{L}_M(X) = \{T \in \mathcal{L}(X): \|T\| \leq M\}$ with respect to strong operator topology or strong* operator topology becomes a Polish (separable and completely metrizable) space, see e.g. [13, Section 4.6.2]. For the operators on a Hilbert space, we can strengthen Theorem 1.2 as follows.

Theorem 3.5. *Assume that X is an infinite-dimensional separable Hilbert space and $M > 1$. Then the intersection of the collection of hypercyclic operators with a residual subset of absolutely mean irregular vectors and $\mathcal{L}_M(X)$ is a dense G_δ subset of $\mathcal{L}_M(X)$ with respect to the strong* operator topology.*

Proof. Fix $M > 1$. We first show that the collection of hypercyclic operators with a residual subset of absolutely mean irregular vectors is a G_δ subset of $\mathcal{L}_M(X)$ with respect to the strong operator topology. Let \mathcal{U} be a countable topological basis of X consisting of non-empty open subsets. For every two sets U and V in \mathcal{U} , let

$$\mathcal{T}(U, V) = \{T \in \mathcal{L}(X): \exists n \in \mathbb{N} \text{ s.t. } T^n U \cap V \neq \emptyset\}$$

For every $n \in \mathbb{N}$, the map $\mathcal{L}_M(X) \rightarrow \mathcal{L}_M(X)$, $T \mapsto T^n$ is continuous with respect to the strong operator topology, see e.g. [7, Lemma 2.1]. Then $\mathcal{T}_M(U, V)$ is an open subset of $\mathcal{L}_M(X)$. By the Birkhoff transitivity theorem (see e.g. [8, Theorem 1.16]), the collection of hypercyclic operators is the set

$$\bigcap_{U, V \in \mathcal{U}} \mathcal{T}(U, V),$$

which is a G_δ subset of $\mathcal{L}_M(X)$.

For every set U in \mathcal{U} and $N \in \mathbb{N}$, let

$$\mathcal{M}(U, N) = \left\{ T \in \mathcal{L}_M(X): \exists n > N \text{ and } x \in U \text{ s.t. } \frac{1}{n} \sum_{i=1}^n \|T^i x\| < \frac{1}{N}, \right. \\ \left. \exists m > N \text{ and } y \in U \text{ s.t. } \frac{1}{m} \sum_{i=1}^m \|T^i y\| > N \right\}.$$

Then $\mathcal{M}(U, N)$ is an open subset of $\mathcal{L}_M(X)$. It is not hard to check that the collection of operators with a residual subset of absolutely mean irregular vectors is the set

$$\bigcap_{U \in \mathcal{U}, N \in \mathbb{N}} \mathcal{M}(U, N),$$

which is a G_δ subset of $\mathcal{L}_M(X)$.

Since the strong* operator topology is finer than the strong operator topology, this collection is also a G_δ subset of $\mathcal{L}_M(X)$ with respect to the strong* operator topology. By [7, Corollary 2.12] the collection $G\text{-MIX}(X)$ of operators in $\mathcal{L}_M(X)$ which admit a mixing invariant measure in the Gaussian sense with full support is dense in $\mathcal{L}_M(X)$ with respect to the strong* operator topology. By Fernique's integrability theorem (see e.g. [1, Exercise 5.5]), every Gaussian measure μ has finite moments of all orders, in particular, $\int_X \|x\| d\mu(x) < \infty$. According to Proposition 3.4, every operators in $G\text{-MIX}(X)$ admits a residual subset of absolutely mean irregular vectors. This end the proof. \square

To conclude this paper, we provide some examples for each case of Theorem 1.1. Following [11], we say that an operator $T: X \rightarrow X$ is *absolutely Cesàro bounded* if there exists a constant $C > 0$ such that

$$\sup_{n \in \mathbb{N}} \frac{1}{n} \sum_{i=1}^n \|T^i x\| \leq C \|x\|$$

for all $x \in X$. It is shown in [9, Theorem 4.32] that an operator is absolutely Cesàro bounded if and only if it mean equicontinuous.

Example 3.6. According to [11, Theorem 3.5 and Remark 3.6], for each $1 \leq p < \infty$ there exists a mixing unilateral weighted backward shift operator T on $\ell^p(\mathbb{N})$ which is absolutely Cesàro bounded. As it is also mean equicontinuous, by Corollary 2.5 for any $x \in \ell^p(\mathbb{N})$, one has

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| = 0.$$

This operator T satisfies the case (1) in the Theorem 1.1.

By Theorems 1.2 and 3.5, there are plenty of hypercyclic operators which satisfy the case (2) in the Theorem 1.1. The following example in [5] is more subtle.

Example 3.7. According to [5, Theorem 37], for each $1 \leq p < \infty$ there exists a bilateral shift T on $\ell^p(v, \mathbb{Z})$ such that T is hypercyclic and completely absolutely mean irregular, that is every non-zero vector is absolutely mean irregular.

Example 3.8. According to [2, Theorem 7], there exists a frequently hypercyclic weighted shift T on $c_0(\mathbb{Z})$ that is not distributionally chaotic. In fact, it is proved in subsection 6.3 of [2] that the orbit of every non-zero vector is not distributionally near 0, that is for every non-zero vector $x \in X$ there exists $\varepsilon = \varepsilon(x) > 0$ such that

$$\underline{\text{dens}}(\{n \in \mathbb{N} : \|T^n x\| \geq \varepsilon\}) > 0.$$

Then for every non-zero vector $x \in X$,

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|T^i x\| \geq \varepsilon \underline{\text{dens}}(\{n \in \mathbb{N} : \|T^n x\| \geq \varepsilon\}) > 0.$$

This shows that T satisfies the case (3) in the Theorem 1.1.

If one does not require the frequent hypercyclicity, there are relatively simple examples for the case (3) in the Theorem 1.1 as follows.

Proposition 3.9. *There exists a hypercyclic weighted backward shift B_w on $\ell^p(\mathbb{Z})$ with $1 \leq p < \infty$ such that for every non-zero vector $x \in \ell^p(\mathbb{Z})$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i x\| = \infty.$$

Proof. Let $a_i = 2^{2+i}$, $b_i = a_i + i - 1 + i$, and $c_i = b_i + 2i$ for $i \in \mathbb{N}$. Define a weight $w = (w_n)_{n \in \mathbb{Z}}$ as follows:

$$w_n = \begin{cases} 2, & n > 0; \\ \frac{1}{2^i}, & a_i + 1 \leq -n \leq b_i \text{ for some } i \in \mathbb{N}; \\ 2, & b_i + 1 \leq -n \leq c_i \text{ for some } i \in \mathbb{N}; \\ 1, & \text{otherwise,} \end{cases}$$

The product $\prod_{j=-n}^0 w_j$ is decreasing for $-n$ in $[a_i + 1, b_i]$ and increasing for $-n$ in $[b_i + 1, a_{i+1}]$. In particular, for every $i \in \mathbb{N}$,

$$\prod_{j=-b_i}^0 w_j = \frac{1}{2^i}, \text{ and } \prod_{j=-n}^0 w_j = 2^i \text{ for } c_i \leq -n \leq a_{i+1}.$$

Let B_w be the weighted backward shift on $\ell^p(\mathbb{Z})$ with the weight w . It is clear that

$$\lim_{i \rightarrow +\infty} \prod_{j=-b_i}^0 w_j = 0, \quad \lim_{n \rightarrow +\infty} \prod_{j=1}^n w_j = \infty.$$

As $\frac{1}{2} \leq |w_j| \leq 2$ for all $j \in \mathbb{Z}$, for any $k \in \mathbb{Z}$,

$$\lim_{i \rightarrow +\infty} \prod_{j=k-b_i}^k w_j = 0 \quad \lim_{i \rightarrow +\infty} \prod_{j=k+1}^{b_i+k} w_j = \infty.$$

By the hypercyclic characterization of weighted backward shifts (see e.g. [8, Example 4.5]), B_w is hypercyclic.

Note that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i e_0\| &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \prod_{j=-i+1}^0 w_j \\ &\geq \lim_{i \rightarrow \infty} \frac{1}{c_i} 2^{i-1} (a_i - c_{i-1}) = \infty. \end{aligned}$$

For all $j \in \mathbb{Z}$, as there exists a positive constant $\alpha = \alpha(j)$ such that $e_j = \alpha B_w^{-j} e_0$, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i e_j\| = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i (\alpha B_w^{-j} e_0)\| = \infty.$$

For every $x = (x_n)_{n \in \mathbb{Z}} \in \ell^p(\mathbb{Z})$, if $x \neq 0$, there exists $j \in \mathbb{Z}$ such that $x_j \neq 0$. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i x\| \geq \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \|B_w^i(x_j e_j)\| = \infty.$$

This ends the proof. \square

Acknowledgment. The author was supported in part by NSF of China (Grant nos. 12222110 and 12171298) and would like to thank Professors Quentin Menet and Xinseng Wang for their helpful comments. The author also expresses many thanks to the anonymous referee, whose suggestions have substantially improved this paper.

REFERENCES

- [1] Frédéric Bayart and Étienne Matheron, *Dynamics of linear operators*, Cambridge Tracts in Mathematics, vol. 179, Cambridge University Press, Cambridge, 2009. MR2533318 ↑8, 10
- [2] Frédéric Bayart and Imre Z. Ruzsa, *Difference sets and frequently hypercyclic weighted shifts*, Ergodic Theory Dynam. Systems **35** (2015), no. 3, 691–709, DOI 10.1017/etds.2013.77. MR3334899 ↑10
- [3] Bernard Beauzamy, *Introduction to operator theory and invariant subspaces*, North-Holland Mathematical Library, vol. 42, North-Holland Publishing Co., Amsterdam, 1988. MR0967989 ↑1
- [4] T. Bermúdez, A. Bonilla, F. Martínez-Giménez, and A. Peris, *Li-Yorke and distributionally chaotic operators*, J. Math. Anal. Appl. **373** (2011), no. 1, 83–93, DOI 10.1016/j.jmaa.2010.06.011. MR2684459 ↑1
- [5] N. C. Bernardes Jr., A. Bonilla, and A. Peris, *Mean Li-Yorke chaos in Banach spaces*, J. Funct. Anal. **278** (2020), no. 3, 108343, 31, DOI 10.1016/j.jfa.2019.108343. MR4030290 ↑2, 3, 7, 10
- [6] N. C. Bernardes Jr., A. Bonilla, A. Peris, and X. Wu, *Distributional chaos for operators on Banach spaces*, J. Math. Anal. Appl. **459** (2018), no. 2, 797–821, DOI 10.1016/j.jmaa.2017.11.005. MR3732556 ↑2
- [7] S. Grivaux, É. Matheron, and Q. Menet, *Linear dynamical systems on Hilbert spaces: typical properties and explicit examples*, Mem. Amer. Math. Soc. **269** (2021), no. 1315, v+147, DOI 10.1090/memo/1315. MR4238631 ↑9, 10
- [8] K.-G. Grosse-Erdmann and A. Peris Manguillot, *Linear chaos*, Universitext, Springer, London, 2011. MR2919812 ↑1, 8, 9, 11
- [9] Zhen Jiang and Jian Li, *Chaos for endomorphisms of completely metrizable groups and linear operators on Fréchet spaces* (2024). arXiv:2212.06304v2. ↑3, 10
- [10] Jian Li, Siming Tu, and Xiangdong Ye, *Mean equicontinuity and mean sensitivity*, Ergodic Theory Dynam. Systems **35** (2015), no. 8, 2587–2612, DOI 10.1017/etds.2014.41. MR3456608 ↑3
- [11] Lvlin Luo and Bingzhe Hou, *Some remarks on distributional chaos for bounded linear operators*, Turkish J. Math. **39** (2015), no. 2, 251–258, DOI 10.3906/mat-1403-41. MR3311688 ↑10
- [12] Stanislav Shkarin, *The Kitai criterion and backward shifts*, Proc. Amer. Math. Soc. **136** (2008), no. 5, 1659–1670, DOI 10.1090/S0002-9939-08-09179-X. MR2373595 ↑8
- [13] Gert K. Pedersen, *Analysis now*, Graduate Texts in Mathematics, vol. 118, Springer-Verlag, New York, 1989. MR0971256 ↑9

(J. Li) DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, 515821, GUANGDONG, CHINA
 Email address: lijian09@mail.ustc.edu.cn
 URL: <https://orcid.org/0000-0002-8724-3050>