## A Very Effective and Simple Diffusion Reconstruction for the Diluted Ising Model

Stefano Bae,<sup>1, \*</sup> Enzo Marinari,<sup>1, 2, †</sup> and Federico Ricci-Tersenghi<sup>1, 2, ‡</sup>

<sup>1</sup>Dipartimento di Fisica, Sapienza Università di Roma, P. A. Moro 5, 00185 Roma (Italy)

<sup>2</sup>Nanotec-CNR, Unità di Roma, and INFN-Sezione di Roma1, P. A. Moro 5, 00185 Roma (Italy)

(Dated: July 11, 2024)

Diffusion-based generative models are machine learning models that use diffusion processes to learn the probability distribution of high-dimensional data. In recent years, they have become extremely successful in generating multimedia content. However, it is still unknown if such models can be used to generate high-quality datasets of physical models. In this work, we use a Landau-Ginzburg-like diffusion model to infer the distribution of a 2D bond-diluted Ising model. Our approach is simple and effective, and we show that the generated samples reproduce correctly the statistical and critical properties of the physical model.

Introduction - Diffusion-based generative models (DM) are machine learning models designed to learn the probability distribution of high-dimensional data and to generate correctly new samples under such distribution [1, 2]. They are intensively applied in high-quality image and video generation [3, 4]. There have been several efforts to rephrase these models in the statistical mechanics framework and improve our theoretical understanding of its behavior. Relations with spontaneous symmetry breaking and phase transitions [5, 6] have been established, the curse of dimensionality, the characterization of time regimes [7, 8] and the connection with associative memory [9] have been analyzed. Also, the physical limits of the method and advantages and disadvantages with respect to standard Monte Carlo and Langevin sampling have been analyzed [10].

As the name suggests, DMs use a diffusion process to accomplish such tasks. The force field of the diffusion process that recursively transforms the initial data into a random noise (typically a Gaussian white noise) is a priori not known, and the relevant goal is to learn it. If one can do that, a time-reversed diffusion process can then be used to generate a new structured sample starting from what appears as pure noise. The recent work of [7, 8]defines a new approach of remarkable value. We propose here a new method that allows us to reconstruct the diluted Ising model (DIM). A model with quenched disorder has an intrinsic complexity, and it is a crucial test if a reconstruction method can analyze it. We will give this question a positive answer, and we will argue that an approach based on a simple structure can be very effective. We believe that these findings have a relevant impact on the more general question of the diffusion-based analysis.

We start from an initial training dataset  $\{\vec{a}^{\mu}\}_{\mu=1}^{P}$ . The *P* elements of the set are *N* dimensional vectors, and they are distributed according to an unknown initial distribution  $P_0(\vec{a})$ . These vectors evolve under an Ornstein-Uhlenbeck diffusion process, and the corresponding stochastic equation in the Ito formulation is:

$$\frac{dx_i(t)}{dt} = -x_i(t) + \eta_i(t) , \qquad (1)$$

where  $\eta_i(t)$  is a white Gaussian noise with correlation  $\langle \eta_i(t)\eta_j(t')\rangle = 2\delta_{ij}\delta(t-t')$ . The stationary distribution of this process is normal  $\mathcal{N}(\vec{x}; \vec{0}, \mathbb{I})$  centered at the origin and with an identity covariance matrix. The probability distribution of this forward process is

$$P\left(\vec{x},t\right) = \int P_0\left(\vec{a}\right) \,\mathcal{N}\left(\vec{x};\vec{a}e^{-t},\Delta\left(t\right)\mathbb{I}\right) d\vec{a} \,\,, \qquad (2)$$

where the mean of the normal distribution is  $\vec{a}e^{-t}$  and the variance is  $\Delta(t) = 1 - e^{-2t}$ . Ideally, given this  $P(\vec{x}, t)$ , one can derive the ideal force field, namely the score function

$$\mathcal{F}_i(\vec{x}, t) = \frac{\partial \log P(\vec{x}, t)}{\partial x_i} = -\frac{x_i - a_i e^{-t}}{\Delta(t)} , \qquad (3)$$

which can be used to define the backward process that will generate new samples [11]:

$$-\frac{dy_i}{dt} = y_i + 2T\mathcal{F}_i(\vec{y}, t) + \eta_i(t) .$$
(4)

However, the distribution  $P_0(\vec{a})$  of real-world data is not known, so in realistic applications, the score function must be learned from the data using a suitable approach (typically a deep neural network).

In our work, we choose to approximate the forward process distribution  $P(\vec{x},t) \approx Q(\vec{x},t;\theta)$  with a function to be discussed in detail below, where  $\theta$  are generic parameters. The approximated score is given by

$$S_i(\vec{x}, t; \theta) \equiv \frac{\partial \log Q(\vec{x}, t; \theta)}{\partial x_i} .$$
 (5)

Since we want to approximate the true score (3) with (5), we minimize the mean squared loss to find the optimal value of the parameters  $\theta$ :

$$\mathcal{L}(\theta) = \left\langle \sum_{k=1}^{N} \left( S_k(\vec{x}, t; \theta) - \mathcal{F}_k(\vec{x}, t) \right)^2 \right\rangle_{\vec{x}, \vec{a}}, \quad (6)$$

where  $\langle \cdot \rangle_{\vec{x},\vec{a}}$  is the average over data and forward trajectories.

Here we try, on the diluted Ising model, a very simple approach, that does not need any deep learning system, and we show that it works and that it is very effective. We define  $\log Q(\vec{x}, t)$  as a quadratic term plus a fourthorder term, and we obtain in this way a non-linear score function that is able to reproduce the statistical and critical properties of the diluted Ising model. This choice of the forward probability distribution, which is strongly reminiscent of a Landau-Ginzburg free energy, is able to generate the correct clustering of the data, reproducing the correct magnetization statistics. Analyzing the spatial correlation function and the magnetic susceptibility we find further and strong evidence that the generated samples show the same critical behavior as the original model. Finally, in light of [7], we discuss the generalization and the memorization regimes, and we characterize the speciation time, i.e. the time at which the trajectories start to fall in one of the free energy minima.

The diluted Ising model - We consider a link diluted Ising model on a 2D square lattice of linear size L and volume  $N = L^2$ . Given N Ising variables,  $a_i = \pm 1$ , the DIM Hamiltonian is defined as

$$\mathcal{H}(\vec{a}) = -\sum_{i,j} J_{ij} a_i a_j , \qquad (7)$$

where the couplings  $J_{ij}$  are null if  $|i - j| \neq 1$  and distributed according to  $p \,\delta(J_{ij} - 1) + (1 - p) \,\delta(J_{ij})$  if |i - j| = 1. The bond concentration p controls the probability of having a bond between nearest neighbors, and in the limit of  $p \rightarrow 1$  one recovers the usual 2D Ising Model. A set of i.i.d.  $\{J_{ij}\}$  extracted according to this probability distribution defines a quenched realization of the disorder, i.e. a disorder sample.

At equilibrium, the configurations of the variables  $\vec{a}$  are distributed according to the Boltzmann-Gibbs measure

$$P_{BG}(\vec{a}) = \frac{1}{\mathcal{Z}} e^{-\beta \mathcal{H}(\vec{a})} , \qquad (8)$$

where the partition function  $\mathcal{Z}$  normalizes  $P_{BG}(\vec{a})$ , and  $\beta = 1/T$  is the inverse temperature of the system. The order parameter that characterizes the different thermo-dynamical phases is the magnetization

$$m = \frac{1}{N} \sum_{i=1}^{N} a_i$$
 (9)

The value of the critical temperature  $T_c(p)$  that separates the two phases depends on the bond concentration.

Our goal is to approximate (8) and generate samples using a DM, to reproduce correctly the main properties of the DIM, such as the critical exponents and the distribution of the magnetization P(m). The DIM has been analyzed in detail, and there exists a large amount of results characterizing its critical and thermodynamical behavior [12–14]; this is very useful to fully characterize the consistency of our results.

*Our method* - The typical (and powerful) attitude adopted to reconstruct an effective score function that

will govern the DM is to use a strongly over-parametrized deep network, for example, U-Nets [4]. If on one side this can be a very useful approach, on the other side to fine-tuning and training such a complex network can be cumbersome, and it is natural to wonder if this is really needed. Even if reconstructing a bona fide statistical model with quenched disorder is not at all a trivial task, we focus on the possibility, that we judge valuable, to use a simple effective score function, and to succeed in the reconstruction in this way. We do indeed succeed, and in this way collect positive results, since we are able to backward reconstruct a non-trivial model with quenched disorder by using a very simple parametrization.

The simplest form one could select for implementing the forward process distribution would be Gaussian. This cannot lead to success: the score obtained in this way is linear, and once it is plugged in the backward Langevin process (4) it can only allow to obtain Gaussian distributed samples [8]. To try to overcome this problem, we use the simplest possible approach, and we add a fourthorder term to the Gaussian distribution, by setting

$$\log Q(\vec{x}, t) = -\frac{1}{2} \sum_{i,j}^{1,N} x_i A_{ij} x_j - \frac{g}{4N^2} \left( \sum_{i=1}^N x_i \right)^4 + \mathcal{C} ,$$
(10)

where A is a symmetric square matrix of dimension Nand g is the parameter that controls the effects of the fourth-order term. C is a normalization constant. The added term is the global magnetization, scaled by a factor  $\sqrt{N}$  and raised to the fourth power. The score corresponding to this forward distribution is:

$$\frac{\partial \log Q(\vec{x}, t)}{\partial x_k} = -\sum_{j=1}^N A_{kj} x_j - \frac{g}{N^2} \left(\sum_{i=1}^N x_i\right)^3 .$$
(11)

We can intuitively see that the term  $\left(\sum_{i=1}^{N} x_i\right)^3$  will help the variables to correlate in a coherent way. In fact, each sum is just the magnetization of the system and if we display it as  $\left(\sum_{i=1}^{N} x_i\right) \left(\sum_{i=1}^{N} x_i\right)^2$  we can see that the squared sum gives a measure of how data is correlated, while the first term also accounts for the direction of the  $x_i$ . In other words, if the variables exhibit correlations during the backward process, this term will amplify them by acting as an external field.

After the process of minimization of (6), which is described in detail in the Supplemental Information (SI), we obtain the following expressions for the matrix  $\boldsymbol{A}$  and the parameter g:

$$(\boldsymbol{A}(t))_{lm} = (\boldsymbol{C}(t)^{-1})_{lm} - \frac{g(t)}{N^2} \sum_{j=1}^{N} \omega_j^{(4)}(t) (\boldsymbol{C}(t)^{-1})_{jm} ,$$
(12)

$$g(t) = N \frac{\frac{\omega^{(4)}(t) - \tilde{\omega}^{(4)}(t)}{\Delta_t} - \gamma^{(0,4)}(t)}{\omega^{(6)}(t) - \gamma^{(4,4)}(t)} , \qquad (13)$$




FIG. 1. The smallest eigenvalue  $\lambda_{min}(t)$  of  $\mathbf{A}(t)$  for the DIM with L = 50 and p = 0.8. Each line is averaged over 20 samples. The different colors for different lines correspond to different temperatures linearly spaced in the range  $T \in [1.600, 2.075]$ . Darker lines are for lower values of T. For higher values of T,  $\lambda_{min}(t)$  never touches the 0 line, in agreement with the fact that for  $T > T_c$  there is only a paramagnetic minimum. When T decreases  $\lambda_{min}(t)$  crosses the x-axis, because of the presence of two minima. The two inset plots correspond to the distribution of the magnetization in time P(m, t) for  $t > t^*$  (red one) and  $t = t^*$  (blue one).

where the matrix C(t) is the correlation matrix of the forward process, which depends on the correlation of the training data  $C_0$ :

$$(\boldsymbol{C}(t))_{km} = \langle x_k x_m \rangle_{\vec{x},\vec{a}} = (\boldsymbol{C}_0)_{km} e^{-2t} + \Delta_t \delta_{km} \quad (14)$$

The other terms,  $\omega^{(4)}(t)$ ,  $\omega^{(6)}(t)$ ,  $\tilde{\omega}^{(4)}(t)$ ,  $\omega_j^{(4)}(t)$ ,  $\gamma^{(0,4)}(t)$ and  $\gamma^{(4,4)}(t)$ , are moments and combinations of moments of the forward process, whose details can also be found in the SI. They depend only on  $C_0$  and on the single sample initial magnetization  $m_0(\vec{a}^{\mu})$ .

Results - We have generated the training data for the DIM using the Wolff algorithm, an improved cluster Monte Carlo Markov Chain [15, 16]. We have used dilution p = 1.0 for studying the pure Ising model and p = 0.8 for analyzing a diluted model. In all cases, the DM was trained with  $P_{train} = 20000$  independent samples [17]. We eventually generated  $P_{qen} = 10000$  samples. Once the training samples have been generated, the parameters A(t) and g(t) can be estimated using (12) and (13). Then, using (4) new configurations of the system can be obtained. We stress that because of the nature of the diffusion process, the generated data are real instead of the original binary ones, and we worked starting from them. We eventually obtained binary spin values by applying a sign function (and a discussion of this procedure is given in the SI).




FIG. 2. The distribution of the magnetization P(m), with L = 100 and p = 0.8. The orange and blue lines correspond respectively to the original and the generated samples. From left to right we display the results for the low T broken phase (a), for the critical point (b), and for the paramagnetic phase (c). The samples generated by backward diffusion match the original distribution.

To develop a multimodal  $Q(\vec{x}, t)$  that correctly reproduces the true  $P(\vec{x},t)$ , we expect that the Hessian of the effective free energy in (10), A(t), should at some point develop a soft mode. Thus, we analyze the smallest eigenvalue of  $\mathbf{A}(t)$ ,  $\lambda_{min}(t)$ , for training sets at different temperatures. We show in Fig. 1 that there exists a time  $t^*$  at which the eigenvalue changes sign from positive to negative going backward in t. This means that the effective free energy is developing new minima that should correspond to the two minima of the DIM [6]. When the diffusion dynamics is at  $t > t^*$ , all the eigenvalues are positive, indicating that the free energy is convex and  $Q(\vec{x},t)$  is uni-modal. At  $t^* = 0$ , the free energy becomes flat along the direction corresponding to the eigenvector of  $\lambda_{min}(t^*)$ , and fluctuations along this direction are enhanced. For  $t < t^*$ , the uni-modal  $Q(\vec{x}, t)$  is no longer stable, and spontaneously evolves towards a  $Q(\vec{x}, t)$  with two peaks. Indeed, looking at the distribution of the magnetization P(m) shown in Fig. 2, we can see that in different thermodynamical phases — above, around, and below  $T_c(p)$  — the distribution of the magnetization of the generated samples is in agreement with that of the original data. Particularly remarkable is the observation in panel (a) that the peaks are reproduced correctly in the broken phase, notwithstanding the simplicity of our score function. It is also correct to notice that there is a slight abundance of  $m \approx 0$  configurations.

Our main goal is to show that our (simple) diffusion model is able to reconstruct in detail the critical properties of the DIM (and we have already shown in Fig. 2 that it reconstructs accurately the bimodal distribution of mbelow  $T_c$ ). We analyze next the x - x two points spatial correlation function C(r) and the fourth moment of the magnetization, namely the Binder parameter B(L, T).

The spatial correlation function C(r) is defined as

$$C(r) \equiv \frac{1}{S_r} \sum_{|i-j|=r} \left[ \langle x_i x_j \rangle_T - \langle x_i \rangle_T \langle x_j \rangle_T \right], \quad (15)$$

where the sum runs over all pairs of spins at distance r




FIG. 3. Panels (a) and (b): spatial correlation function C(r) at criticality for the original (orange above) and the reconstructed (blue below) data, with L = 100. We show the pure Ising model, p = 1.0, in (a), and the DIM with p = 0.8 in (b). C(r) has been averaged over 10 disorder samples. In both cases, the generated correlations show the same critical decay as the original one. Panels (c) and (d): DIM Binder parameter B(T, L) as a function of the temperature T, with  $L \in \{12, 25, 50, 100\}$  and p = 0.8, averaged over 10 disorder samples. The orange points and lines are for the original data, the blue ones are for the generated samples. The vertical dashed line is for the critical temperature  $T_c(p = 0.8) \simeq 1.650$  [18]. In panel (c) we show the intersection of the generated and original data at  $T_c(p = 0.8)$ . In panel (d) we show that the values of the Binder parameter rescaled using the critical exponents of the 2D Ising universality class collapse onto a single curve.

along the axis, being  $S_r$  their number.  $\langle \cdot \rangle_T$  is the average over the Gibbs measure (8) at temperature T, while  $[\cdot]$  is the average over the disorder (we have used few tens of samples). The Binder B(L,T) is defined as

$$B(L,T) \equiv 1 - \frac{\left[\langle m^4 \rangle_T\right]}{3\left[\langle m^2 \rangle_T\right]^2} .$$
 (16)

We show in Fig. 3 that the correlations that characterize the generated samples at the critical temperature decay as in the original, Boltzmann distributed samples: as one would expect the system is characterized by the same critical exponent  $\eta = \frac{1}{4}$  (but for size-dependent logarithmic corrections [14]) of the 2D pure Ising model universality class. By analyzing B(L,T) we also checked that the value  $\nu = 1$  is reproduced correctly by our backward reconstruction. We show in panels (c) and (d) of Fig. 3 the scaling behavior of B(L,T), derived by using the 2D Ising exponents (again ignoring logarithmic corrections), that confirms that the critical properties of the generated data reproduce with high accuracy the ones of the original data [14]. The intersection of the Binder parameter is used in literature to determine the critical temperature and the critical exponents of the model [19], and we can see that the analysis of the generated data leads to the same estimates as for the original data.

It is also important to understand if the DM is really generalizing from the data, thus producing new configurations, and it is not simply memorizing the dataset, and converging to one of the data points in the backward process. To answer in a quantitative way, we look at the distribution of the overlap between the generated data and the original data

$$q^{\mu} = \frac{1}{N} \sum_{i=1}^{N} a_i^0 \operatorname{sgn}[x_i^{\mu}(t=0)].$$
 (17)




FIG. 4. We show in panel (a) the distribution of the overlap between the generated and the original samples in the broken phase  $T = 1.6 < T_c$  with p = 0.8. The two symmetric peaks correspond to samples that have the same magnetization, while there is no peak at q = 1 suggesting that the model is typically not generating samples that are in the training set. In panel (b) we show the probabilities that two cloned trajectories at a time t fall in the same peak as a function of  $t/t^*$ . Different colors correspond to different system sizes.

As we have described before, because of the nature of the diffusion process, we generate real data and not binary data, and we eventually apply the sign function to make them binary again. In this way, if a generated configuration is present in the original training dataset, we will get a value of q close to one. In Fig. 4 we show the histogram of the overlap P(q). One can see that in the cold phase, it has two peaks that correspond to configurations that have aligned (positive peak) or opposite (negative peak) mean magnetization. There is no sizable peak at q = 1, which implies that our model is not memorizing the training data, but it is always running a bona fide generalization. This is because the only parameters we are learning from the data are the correlation  $C_0$  and the magnetization, so the model is able to reproduce its critical properties which depend on the

correlation of the data. As reported in [7], to reduce the collapse of the backward process of the training data, an exponential number of training samples in N is required. In modern applications, regularization can mitigate this problem [20], and in our case, we can think that the use of this particular form of the score function acts as a regularization.

Our last, important goal is to characterize the time at which the trajectories of the backward process cluster and form two separate peaks in the P(m). Following [7] we study the probability that two trajectories, cloned at time t, fall in the same cluster or peak at t = 0, denoted by  $\phi(t)$ . In [7] the speciation time  $t_s = \log(\Lambda)/2$ was introduced, where  $\Lambda$  is the largest eigenvalue of  $C_0$ . This proposal comes from the observation that close to  $t_s$  the fluctuations of the data become of the order of the Gaussian noise.

In our approach, the relevant fluctuations leading to a clustering of the trajectories are expected to arise close to the time  $t^*$  discussed above. We show in Fig. 4 that the  $\phi(t)$  curves do have a clean crossing when plotted as a function of  $t/t^*$  and the slope at the crossing point increases with the system size L, indicating that, for  $L \to \infty$ ,  $t^*$  plays the role of a threshold. In [7] the same crossing has been found plotting  $\phi(t)$  as a function of  $t/t_s$ , so both  $t^*$  and  $t_s$  belong to the critical region and define the same speciation time in the large L limit. Indeed we find that both  $t^*$  and  $t_s$  grow logarithmically with L, while their difference stays almost constant.

Conclusions - We have introduced a simple diffusion model that efficiently learns the high-dimensional probability distribution of configurations in a disordered model, namely the 2D diluted Ising model. Once trained, it can generate typical configurations above, below, and at the critical temperature, reconstructing accurately the magnetization and the critical exponents  $\eta$  and  $\nu$ . Using this simple model we have better characterized the speciation time  $t^*$  in terms of the effective Hessian  $\mathbf{A}(t)$ .

Our results clearly show that it is possible to use a diffusion model to reconstruct a statistical model with quenched disorder. This is potential very relevant consequences that will be explored in the future.

Acknowledgments - We warmly thank Giulio Biroli and Marc Mézard for the interesting conversations and the critical reading of the manuscript. EM acknowledges the hospitality of Mark Bowick at UCSB (CA, USA), where part of this work was done, and the support by grant NSF PHY-2309135 to the Kavli Institute for Theoretical Physics (KITP). This research has been supported by funding from the 2021 FIS (Fondo Italiano per la Scienza) funding scheme (FIS783 - SMaC - Statistical Mechanics and Complexity), from the 2022 PRIN funding scheme (2022LMHTET - Complexity, disorder and fluctuations: spin glass physics and beyond) from Italian MUR (Ministry of University and Research) and from ICSC - Italian Research Center on High-Performance Computing, Big Data, and Quantum Computing, funded by the European Union - NextGenerationEU.

- \* stefano.bae@uniroma1.it; First and corresponding author
- <sup>†</sup> enzo.marinari@uniroma1.it; The two last authors contributed equally.
- <sup>‡</sup> federico.ricci@uniroma1.it; The two last authors contributed equally.
- J. Sohl-Dickstein, E. Weiss, N. Maheswaranathan, and S. Ganguli, Deep unsupervised learning using nonequilibrium thermodynamics, in *Proceedings of the 32nd International Conference on Machine Learning*, Proceedings of Machine Learning Research, Vol. 37, edited by F. Bach and D. Blei (PMLR, Lille, France, 2015) pp. 2256–2265.
- [2] L. Yang, Z. Zhang, Y. Song, S. Hong, R. Xu, Y. Zhao, W. Zhang, B. Cui, and M.-H. Yang, Diffusion models: A comprehensive survey of methods and applications, ACM Computing Surveys 56, 1 (2023).
- [3] Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, and B. Poole, Score-based generative modeling through stochastic differential equations, arXiv preprint arXiv:2011.13456 (2020).
- [4] J. Ho, A. Jain, and P. Abbeel, Denoising diffusion probabilistic models, in Advances in Neural Information Processing Systems, Vol. 33, edited by H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin (Curran Associates, Inc., 2020) pp. 6840–6851.
- [5] L. Ambrogioni, The statistical thermodynamics of generative diffusion models, arXiv preprint arXiv:2310.17467 (2023).
- [6] G. Raya and L. Ambrogioni, Spontaneous symmetry breaking in generative diffusion models, Advances in Neural Information Processing Systems 36 (2024).
- [7] G. Biroli, T. Bonnaire, V. de Bortoli, and M. Mézard, (2024), arXiv:2402.18491.
- [8] G. Biroli and M. Mézard, J. Stat. Mech , 093402 (2023).
- [9] L. Ambrogioni, In search of dispersed memories: Generative diffusion models are associative memory networks, Entropy 26, 10.3390/e26050381 (2024).
- [10] D. Ghio, Y. Dandi, F. Krzakala, and L. Zdeborová, Sampling with flows, diffusion and autoregressive neural networks: A spin-glass perspective, arXiv preprint arXiv:2308.14085 (2023).
- [11] B. D. Anderson, Reverse-time diffusion equation models, Stochastic Processes and their Applications 12, 313 (1982).
- [12] R. B. Griffiths, Nonanalytic behavior above the critical point in a random ising ferromagnet, Physical Review Letters 23, 17 (1969).
- [13] A. B. Harris, Effect of random defects on the critical behaviour of ising models, Journal of Physics C: Solid State Physics 7, 1671 (1974).
- [14] H. G. Ballesteros, L. A. Fernández, V. Martín-Mayor, A. M. Sudupe, G. Parisi, and J. J. Ruiz-Lorenzo, Ising exponents in the two-dimensional site-diluted ising model, Journal of Physics A: Mathematical and General **30**, 8379 (1997).
- [15] U. Wolff, Collective monte carlo updating for spin systems, Physical Review Letters 62, 361 (1989).

- [16] M. E. J. Newman and G. T. Barkema, Monte Carlo methods in statistical physics (Clarendon Press, Oxford, 1999).
- [17] The Wolff algorithm has a small correlation time [16] and to generate independent samples we took measurements that were suitably spaced in time.
- [18] I. A. Hadjiagapiou, Monte carlo analysis of the critical properties of the two-dimensional randomly bond-diluted

ising model via wang–landau algorithm, Physica A: Statistical Mechanics and its Applications **390**, 1279 (2011).

- [19] K. Binder, Finite size scaling analysis of ising model block distribution functions, Zeitschrift für Physik B Condensed Matter 43, 119 (1981).
- [20] Z. Kadkhodaie, F. Guth, E. P. Simoncelli, and S. Mallat, Generalization in diffusion models arises from geometry-adaptive harmonic representation, arXiv preprint arXiv:2310.02557 (2023).

# **Supplementary Information**

A very effective and simple diffusion reconstruction for the diluted Ising model

## COMPUTATION OF THE APPROXIMATE SCORE FUNCTION

In this section we discuss a few details of our computations. The approximated distribution of the forward process and the corresponding score function are

$$\log Q(\vec{x},t) = -\frac{1}{2} \sum_{i,j}^{1,N} x_i A_{ij} x_j - \frac{g}{4N^2} \left( \sum_{i=1}^N x_i \right)^4 + \mathcal{C} , \qquad (18)$$

$$\frac{\partial \log Q(\vec{x}, t)}{\partial x_k} = -\sum_{j=1}^N A_{kj} x_j - \frac{g}{N^2} \left(\sum_{i=1}^N x_i\right)^3 \,. \tag{19}$$

To find the parameters A and g we minimize the mean squared loss:

$$\mathcal{L}(\boldsymbol{A},g) = \frac{1}{2} \left\langle \sum_{k=1}^{N} \left( \sum_{j=1}^{N} A_{kj} x_j + \frac{g}{N^2} \left( \sum_{i=1}^{N} x_i \right)^3 - \frac{x_k - a_k e^{-t}}{\Delta(t)} \right)^2 \right\rangle_{\vec{x},\vec{a}},\tag{20}$$

where the mean value  $\langle (\cdot) \rangle_{\vec{x},\vec{a}}$  is defined as:

$$\langle (\cdot) \rangle_{\vec{x},\vec{a}} = \sum_{\{\vec{a}\}} \int_{-\infty}^{\infty} d\vec{x} \, (\cdot) P(\vec{x},\vec{a}) = \sum_{\{\vec{a}\}} \int_{-\infty}^{\infty} d\vec{x} \, (\cdot) P_0(\vec{a}) \left(\frac{1}{2\pi\Delta(t)}\right)^{N/2} \exp\left(-\frac{(\vec{x} - \vec{a}e^{-t})^2}{2\Delta(t)}\right) \,. \tag{21}$$

We want to minimize (20) w.r.t. the matrix A and the scalar g. Taking the derivatives we obtain:

$$\frac{\partial \mathcal{L}(\boldsymbol{A},g)}{\partial A_{lm}} = \left\langle \sum_{k=1}^{N} A_{lk} x_k x_m + \frac{g}{N^2} \left( \sum_{i=1}^{N} x_i \right)^3 x_m - \frac{x_l x_m - a_l x_m e^{-t}}{\Delta(t)} \right\rangle_{\vec{x},\vec{a}} = 0 , \qquad (22)$$

$$\frac{\partial \mathcal{L}(\mathbf{A},g)}{\partial g} = \left\langle \frac{1}{N^2} \sum_{k,j}^{1,N} A_{kj} \left( \sum_{i=1}^N x_i \right)^3 x_j + \frac{g}{N^3} \left( \sum_{i=1}^N x_i \right)^6 - \sum_{k=1}^N \frac{\left( \sum_{i=1}^N x_i \right)^3 x_k - \left( \sum_{i=1}^N x_i \right)^3 a_k e^{-t}}{N^2 \Delta(t)} \right\rangle_{\vec{x},\vec{a}} = 0.$$
(23)

A, g and the mean values depend on time. However, to simplify the notation, we will not make the time dependence explicit.

#### Gaussian mean values

The next step is to get rid of the mean over  $\vec{x}$ . One can see from (21) that all the integrals in  $\vec{x}$  are about computing moments of a Gaussian. The simplest one is

$$\langle x_k x_m \rangle_{\vec{x},\vec{a}} = \langle a_k a_m \rangle_{\vec{a}} \ e^{-2t} + \Delta(t) \delta_{km} \ . \tag{24}$$

If  $C_0 = \langle a_k a_m \rangle_{\vec{a}}$  is the correlation of the training data we have that

$$(\boldsymbol{C}(t))_{km} = \langle x_k x_m \rangle_{\vec{x},\vec{a}} = (\boldsymbol{C}_0)_{km} e^{-2t} + \Delta(t) \delta_{km} .$$
<sup>(25)</sup>

The other mean values can be expressed as:

$$\omega^{(n)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^n \right\rangle_{\vec{x},\vec{a}},\tag{26}$$

8

$$\tilde{\omega}^{(n)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^{n-1} \sum_{j=1}^{N} a_i e^{-t} \right\rangle_{\vec{x},\vec{a}},\tag{27}$$

$$\omega_j^{(n)}(t) = \left\langle \left(\sum_{i=1}^N x_i\right)^{n-1} x_j \right\rangle_{\vec{x},\vec{a}}.$$
(28)

We have to consider the following terms:

$$\begin{split} &\omega^{(4)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^4 \right\rangle_{\vec{x},\vec{a}} = 3N^2 \Delta(t)^2 + 6N^3 \Delta(t) \langle m_0^2(a) \rangle_{\vec{a}} e^{-2t} + N^4 \langle m_0^4(a) \rangle_{\vec{a}} e^{-4t} , \\ &\omega_j^{(4)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^3 x_j \right\rangle_{\vec{x},\vec{a}} = 3N \Delta(t)^2 + 3N^2 \Delta(t) \langle (m_0^2(a) + m_0(a)a_j) \rangle_{\vec{a}} e^{-2t} + N^3 \langle m_0^3(a)a_j \rangle e^{-4t} , \\ &\tilde{\omega}^{(4)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^3 \sum_{j=1}^{N} a_j e^{-t} \right\rangle_{\vec{x},\vec{a}} = 3N^3 \Delta(t) \langle m_0^2(a) \rangle_{\vec{a}} e^{-2t} + N^4 \langle m_0^4(a) \rangle_{\vec{a}} e^{-4t} , \\ &\omega^{(6)}(t) = \left\langle \left(\sum_{i=1}^{N} x_i\right)^6 \right\rangle_{\vec{x},\vec{a}} = 15N^3 \Delta(t)^3 + 45N^4 \Delta(t)^2 \langle m_0^2(a) \rangle_{\vec{a}} e^{-2t} + 15N^5 \Delta(t) \langle m_0^4(a) \rangle_{\vec{a}} e^{-4t} + N^6 \langle m_0^6(a) \rangle_{\vec{a}} e^{-6t} . \end{split}$$

All these quantities depend on the correlation matrix  $(C_0)_{ij} = \langle a_i a_j \rangle_{\vec{a}}$  and the mean magnetization  $m_0(\vec{a}) = \frac{1}{N} \sum_{i=1}^{N} a_i$  of the training data.

## Expressions for $\boldsymbol{A}(t)$ and $\boldsymbol{g}(t)$

Using the relations that we have just found, equations (22) and (23) become:

...

$$\frac{\partial \mathcal{L}(\boldsymbol{A},g)}{\partial A_{lm}} = \sum_{k=1}^{N} A_{lk} C_{km} + \frac{g}{N^2} \omega_m^{(4)} - \delta_{lm} = 0 , \qquad (29)$$

$$\frac{\partial \mathcal{L}(\mathbf{A},g)}{\partial g} = \frac{1}{N^2} \sum_{k,j}^{1,N} A_{kj} \omega_j^{(4)} + \frac{g}{N^3} \omega^{(6)} - \frac{\omega^{(4)} - \tilde{\omega}^{(4)}}{N^2 \Delta(t)} = 0.$$
(30)

Now, inverting (29) we get

$$(\boldsymbol{A}(t))_{lm} = (\boldsymbol{C}(t)^{-1})_{lm} - \frac{g(t)}{N^2} \sum_{j=1}^{N} \omega_j^{(4)}(t) (\boldsymbol{C}(t)^{-1})_{jm} .$$
(31)

Putting this information in (30) we find

$$\frac{1}{N^2} \sum_{k,j}^{1,N} \left[ (\boldsymbol{C}(t)^{-1})_{kj} - \frac{g(t)}{N^2} \sum_{i=1}^{N} \omega_i^{(4)}(t) (\boldsymbol{C}(t)^{-1})_{ij} \right] \omega_j^{(4)}(t) + \frac{g(t)}{N^3} \omega^{(6)}(t) - \frac{\omega^{(4)}(t) - \tilde{\omega}^{(4)}(t)}{N^2 \Delta(t)} = 0 ,$$

from which we get

$$g(t) = N \frac{\frac{\omega^{(4)}(t) - \tilde{\omega}^{(4)}(t)}{\Delta(t)} - \gamma^{(0,4)}(t)}{\omega^{(6)}(t) - \gamma^{(4,4)}(t)} , \qquad (32)$$

where  $\gamma^{(0,4)}(t)$  and  $\gamma^{(4,4)}(t)$  are defined as

$$\gamma^{(0,4)}(t) = \sum_{k,j}^{1,N} (\mathbf{C}(t)^{-1})_{kj} \omega_j^{(4)}(t) ,$$
  
$$\gamma^{(4,4)}(t) = \sum_{i,j}^{1,N} \omega_i^{(4)}(t) (\mathbf{C}(t)^{-1})_{ij} \omega_j^{(4)}(t) .$$

#### THE CASE OF A LINEAR SCORE

In this section we will discuss what happens if one uses a linear score. A linear score function is obtained from a Gaussian distribution of the forward process

$$\log Q(\vec{x},t) = -\frac{1}{2} \sum_{i,j}^{1,N} x_i A_{ij} x_j , \qquad (33)$$

and

$$\frac{\partial \log Q(\vec{x}, t)}{\partial x_k} = -\sum_{j=1}^N A_{kj} x_j .$$
(34)

Using the same analysis of the previous section, the matrix A is defined by

$$\boldsymbol{A}(t) = \left(\boldsymbol{C}_0 e^{-2t} + \Delta(t) \mathbb{I}\right)^{-1} , \qquad (35)$$

and putting it in the backward equation we get

$$-\frac{dy_i}{dt} = y_i - 2T \sum_{j=1}^{N} (\mathbf{A}(t)_{ij}) y_j + \eta_i(t) .$$
(36)

Since the backward process is a Langevin equation with a linear force term, and the distribution at  $t \to \infty$  is a Gaussian, the only possible distribution at t = 0 will be a Normal distribution with covariance  $C_0$ . Therefore we expect that the generated data will have the same correlations as the original ones, but not the clustering of the data points around the states  $|\langle m \rangle| \neq 0$  in the cold phase.

Indeed, if we look at Fig. 5, where we show the smallest eigenvalue of  $\mathbf{A}(t)$  as a function of the time for the pure Ising model with system size L = 50, we can see that it never touches the x-axis, indicating that paramagnetic  $\langle m \rangle = 0$ state is always stable, and the distribution never develops new maxima. This is also clear from Fig. 6, where the distribution of the reconstructed magnetization is always Gaussian in all the thermodynamic phases.

However, the pure Gaussian approach can reconstruct the quantities that depend on the two-point correlation function C(r) in the high T phase. In Fig. 7 we show the correlation C(r) and the susceptibility  $\chi(T, L)$  for the pure Ising model. From panel (a) we can see that around  $T \approx T_c$  the correlation of the generated data decays as the original ones. In panel (c) we scale  $\chi(T, L)$  using Ising critical exponents, and we can see that for  $T > T_c$  both the original and generated samples fall into a single curve, while in the cold phase, they do not. This is due to the mean magnetization. In fact, where  $\langle m \rangle = 0$  for both the generated and original samples, the curves collapse  $(T > T_c)$ . As the temperature goes below  $T_c$  the original samples develop a non-zero magnetization while the generated ones maintain  $\langle m \rangle = 0$  giving no contribution to  $\chi(T, L)$ .

### ANALYSIS WITH BINARY VARIABLES

In this section, we discuss about the application of the sign function to the generated samples. Using the sign function at the end of the generative process at t = 0, it breaks randomly the correlation between the variables that are close to 0. This is due to the nature of the real variables, and variables that are slightly positive will become +1 regardless of the neighboring spins (the slightly negative ones become -1). Therefore the samples that are more sensitive to the sign operation will be the ones closest to the critical temperature.

First, we look at the magnetization distribution in Fig. 8. There are no qualitative differences with respect to the real variables, and we can observe that in the  $T \sim T_c$  and the  $T > T_c$  cases (central and right histograms green plots) the number of samples with  $m \approx 0$  increased because of the randomness added by the sign operation. Regarding the two-point correlation function C(r) there is no substantial difference. Indeed if we look the panels (a) and (b) of Fig. (9), we can observe that the decay rate is preserved even if the overall values are shifted downwards. The last, but not the least, analysis is the one of the behavior of the fourth-order cumulant, namely the Binder parameter [19]

$$B(T,L) = 1 - \frac{\langle m^4 \rangle_T}{3 \langle m^2 \rangle_T^2},$$
(37)




FIG. 5. Linear Gaussian reconstruction. The smallest eigenvalue  $\lambda_{min}(t)$  of  $\mathbf{A}(t)$  for the pure Ising model (L = 50 and p = 1). The different colors for different lines correspond to different temperatures linearly spaced in the range  $T \in [2.200, 2.485]$ . Darker lines are for lower values of T. The eigenvalue never reaches the value 0, indicating that the phase space remains convex: this approach cannot reconstruct the low T phase.




FIG. 6. Linear Gaussian reconstruction. The distribution of the magnetization P(m) for the pure Ising model, L = 50 and p = 1.0. The orange and blue lines correspond respectively to the original and to the generated samples. From left to right we display the results for the low T broken phase (a), for the critical point (b) and for the paramagnetic phase (c).

which is often used to characterize the critical temperature. In Figure 9 we show B(T, L) for a DIM with p = 0.8. In panel (c) the intersections of the different curves show that the critical temperature is around  $T_c(p = 0.8) \simeq 1.650$ . In panel (d) the rescaling with Ising critical exponents, makes all the data points collapse on a single curve. This shows that the sign function does not affect the critical properties of the generated samples. Also in terms of the magnetization, it does not change the overall behavior of the P(m) in the three thermodynamical phases.




FIG. 7. Linear Gaussian reconstruction. In Panel (a) we show the two-point correlation function for the pure Ising model, L = 50 and p = 1.0. The decay of the generated samples matches the original one. In Panel (b) we show  $\chi(T,L)$  for the pure Ising model as a function of the temperature for  $L = \{12, 25, 50\}$ . We use dots for the original data and the X's for the generated ones. The vertical dashed line is for the critical temperature  $T_c(p = 1.0) \simeq 2.27$ . In panel (c) we show the same plot scaled using the values  $\nu = 1$  and  $\gamma = \frac{7}{4}$  of the 2D Ising universality class. For  $T > T_c$  the generated and original data points collapse, while in the cold phase, they do not. This is due to the fact that the Gaussian reconstruction is not able to build correctly the mean magnetization, which stays equal to zero in the generated samples.




FIG. 8. Binary variables. The distribution of the magnetization P(m), L = 100 and p = 0.8. The orange and blue lines correspond respectively to the original and to the generated samples, while in green the line of the generated samples after applying the sign function. From left to right we display the results for the low T broken phase (a), for the critical point (b) and for the paramagnetic phase (c). The samples generated by backward diffusion match the original distribution. There is no substantial difference in the behavior of the distributions.




FIG. 9. Binary variables. Panels a) and b): DIM spatial correlation function C(r) for the original and the reconstructed data, with L = 100. (a) is for the pure Ising model p = 1.0 case. (b) is for p = 0.8, and here C(r) has been averaged over 10 disorder samples. In both cases the generated correlations show the same decay as the original one. Panels (c) and (d): DIM Binder parameter B(T, L) as a function of the temperature T, with  $L = \{12, 25, 50, 100\}$  and p = 0.8 averaged over 10 disorder samples. The vertical dashed line is for the critical temperature  $T_c(p = 0.8)$  [18]. In panel (d) we show that the values of the Binder parameter rescaled using the critical exponents of the 2D Ising universality class collapse onto a single curve.