

On the Berwald-Weyl Curvature

Zhongmin Shen* and Liling Sun†

Abstract

In this paper, we study the Berwald-Weyl curvature which is defined for a spray/Finsler metric with a volume form. We obtain some expressions for the Berwald-Weyl curvature. This quantity is a projective invariant with respect to a fixed volume form. We prove that for any spray of scalar curvature on a manifold of dimension $n \geq 3$, the Berwald-Weyl curvature vanishes with respect to any volume form. We also show that for any Finsler metric of constant Ricci curvature and constant S-curvature, the Berwald-Weyl curvature vanishes with respect to the Busemann-Hausdorff volume form. This study leads to a new notion of BWeyl-flat sprays/Finsler metrics.

Keywords: spray, Finsler metric, Berwald-Weyl curvature, BWeyl-flatness

Mathematics Subject Classification 2010: 53B40, 53C60

1 Introduction

In Finsler geometry, there are several well known projective invariants such as Douglas curvature and Weyl curvature. Finsler metrics with vanishing Weyl curvature are called Weyl metrics, that are exactly Finsler metrics of scalar flag curvature ([10]). In dimension $n \geq 3$, a spray/Finsler metric is locally projectively flat if and only if the Weyl curvature and the Douglas curvature vanish ([6]). In dimension two, the Weyl curvature always vanishes. There is another projective invariant, the Berwald-Weyl curvature. It is known that a two-dimensional spray/Finsler is locally projectively flat if and only if the Berwald-Weyl curvature and the Douglas curvature vanish ([1]). In this paper, we shall discuss the Berwald-Weyl curvature and its generalization to higher dimensions.

A spray on a manifold M is a vector field on the tangent bundle TM in the form $\mathbf{G} = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i}$. Fix a local coordinate system, one can construct a local projective spray $\tilde{\mathbf{G}}$ by setting $\tilde{G}^i = G^i - \frac{1}{n+1} \frac{\partial G^m}{\partial y^m} y^i$. Define a tensor \mathbf{W}^o

*supported in part by NSF China (NSFC No. 11671352)

†corresponding author, supported by NSF China (NSFC No. 12101259), Natural Science Foundation of Fujian Province(No. 2021J05173), Foundation of Fujian Educational Committee(No. JAT200274) and Research Foundation of Jimei University(No. ZQ2022032)

by

$$W_k^o := \bar{R}_{\parallel k} - \frac{1}{2} \bar{R}_{.k\parallel m} y^m,$$

where the covariant derivative “ \parallel ” is taken respect to $\bar{\mathbf{G}}$ ([7]) and $\bar{R} = \frac{1}{n-1} \mathbf{Ric}_{\bar{\mathbf{G}}}$ is the Ricci scalar of $\bar{\mathbf{G}}$. It is proved that in dimension two, \mathbf{W}^o is a globally defined quantity ([1] [7]). In higher dimensions, the above expression does not define a global quantity. To avoid this problem, one can use the globally defined associated projective spray $\hat{\mathbf{G}}$ using a volume form [7]. Given a volume form dV on M , we have the S-curvature $\mathbf{S} = \mathbf{S}_{(G,dV)}$ (2 .17). With the S-curvature, we can construct a new spray by

$$\hat{\mathbf{G}} := \mathbf{G} + \frac{2\mathbf{S}}{n+1} \mathbf{Y}, \quad (1 .1)$$

where $\mathbf{Y} = y^i \frac{\partial}{\partial y^i}$ denotes the canonical vertical field on TM . This spray is projectively invariant with respect to a fixed volume form. The spray $\hat{\mathbf{G}}$ is called the projective spray of (\mathbf{G}, dV) . Let $\hat{R} = \frac{1}{n-1} \mathbf{Ric}_{\hat{\mathbf{G}}}$ is the Ricci scalar of $\hat{\mathbf{G}}$. The Berwald-Weyl curvature $\mathbf{W}^o = W_k^o dx^k$ of (\mathbf{G}, dV) is defined by

$$W_k^o = \hat{R}_{\parallel k} - \frac{1}{2} \hat{R}_{.k\parallel m} y^m, \quad (1 .2)$$

where the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$ ([7]). The Berwald-Weyl curvature is projectively invariant with respect to a fixed volume form.

In general, the spray $\hat{\mathbf{G}}$ defined in (1 .1) depends on the volume form dV and so does the Berwald-Weyl curvature \mathbf{W}^o . In $\dim M \geq 3$, \mathbf{W}^o can be expressed in terms of the Weyl curvature.

Proposition 1.1 *Given a spray space (\mathbf{G}, dV) on the manifold M of dimension $n \geq 3$, the Berwald-Weyl curvature \mathbf{W}^o can be written as*

$$W_k^o = \frac{1}{n-2} W_{k\parallel m}^m, \quad (1 .3)$$

where the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$ and W_k^i is the Weyl curvature given in (2 .29).

It is well known that a spray is of scalar curvature if and only if its Weyl curvature \mathbf{W} vanishes, that is $W_k^i = 0$. By this fact and (1 .3), we conclude immediately the following

Theorem 1.2 *If \mathbf{G} is a spray of scalar curvature on a manifold M of $\dim M \geq 3$, then its Berwald-Weyl curvature $\mathbf{W}^o = 0$ for any given volume form dV .*

Since the Berwald -Weyl curvature is defined via the S-curvature, it is natural to express \mathbf{W}^o in terms of the geometric quantities of \mathbf{G} and the S-curvature with respect to the Berwald connection of \mathbf{G} .

Proposition 1.3 *Given a spray space (\mathbf{G}, dV) on the manifold M of dimension n , the Berwald-Weyl curvature \mathbf{W}^0 can be expressed as*

$$W_k^o = R_{|k} - \frac{1}{2}R_{.k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m - \frac{1}{n+1}W_k^m\mathbf{S}_{.m}, \quad (1.4)$$

where $R = \frac{1}{n-1}\mathbf{Ric}_G$ is the Ricci scalar of \mathbf{G} , χ_k is the non-Riemannian quantity given in (2.18), W_k^i is the Weyl curvature given in (2.29), $\mathbf{S} = \mathbf{S}_{(G,dV)}$ is the S-curvature of (\mathbf{G}, dV) given in (2.17) and the horizontal covariant derivatives “|” are taken with respect to \mathbf{G} .

We know that if a Finsler metric is of constant flag curvature k , then $R = kF^2$, $\chi_k = 0$ and $W_k^i = 0$. From (1.4), one can see that $W_k^o = 0$ for any volume form.

In dimension $n = 2$, since $\mathbf{W} = \mathbf{0}$. By this fact and (1.4), we see that \mathbf{W}^o is independent of the volume form. This fact is known [7].

Corollary 1.4 *For a two-dimensional spray (\mathbf{G}, dV) , the Berwald-Weyl curvature \mathbf{W}^o is independent of dV , given by*

$$W_k^o = R_{|k} - \frac{1}{2}R_{.k|m}y^m - \frac{1}{3}\chi_{k|m}y^m. \quad (1.5)$$

Hence it is a projective invariant of \mathbf{G} .

Every Finsler metric F induces a spray \mathbf{G}_F . Thus results on sprays are still true. However, for a Finsler metric F , we sometimes restrict the volume form dV to its Busemann-volume form dV_{BH} . In this case, the S-curvature of F is the S-curvature of (\mathbf{G}_F, dV_{BH}) . A Finsler metric F is of isotropic S-curvature if the S-curvature

$$\mathbf{S} = (n+1)cF,$$

where $c = c(x)$ is a scalar function on M . F is of constant S-curvature if the c is a constant. F is called an Einstein metric if

$$\mathbf{Ric} = (n-1)\sigma F^2,$$

where $\sigma = \sigma(x)$ is a scalar function on M . In particular, a Finsler metric F is of constant Ricci curvature if σ is a constant.

For a Finsler manifold (M, F) , if the S-curvature is constant with respect to the Busemann-Hausdorff volume form $dV = dV_{BH}$, then $\chi_k = 0$. Further, if the Ricci curvature is constant, then by (1.4), the Berwald-Weyl curvature vanishes

Theorem 1.5 *Let (M, F) be a Finsler manifold. If F is of constant Ricci curvature and constant S-curvature with respect to $dV = dV_{BH}$, then $\mathbf{W}^o = 0$.*

It is known that every Einstein Randers metric on a manifold M is of constant S-curvature[2]. Hence, we obtain the following

Corollary 1.6 For any Randers metric of constant Ricci curvature, $\mathbf{W}^o = 0$ for the $dV = dV_{BH}$.

We know that there exist Randers metrics of constant S-curvature and constant Ricci curvature which are not of scalar curvature [4]. Thus there exist Finsler metrics with vanishing Berwald-Weyl curvature which are not of scalar curvature. Below is another non-trivial example.

Example 1.7 Let $\alpha_1 = \sqrt{a_{ij}y^i y^j}$ and $\alpha_2 = \sqrt{\bar{a}_{ij}y^i y^j}$ be two Ricci-flat Riemannian metrics on the manifolds M_1 and M_2 , respectively. Consider the following 4-th root metric

$$F := \sqrt[4]{\alpha_1^4 + 2c\alpha_1^2\alpha_2^2 + \alpha_2^4},$$

where $0 < c \leq 1$ is a constant. It is a Riemannian metric when $c = 1$. This is a Ricci-flat Berwald metric on $M := M_1 \times M_2$. Note that the S-curvature $\mathbf{S}_{(F,dV)} = 0$ for the Busemann-Hausdorff volume form $dV = dV_{BH}$. Thus $\mathbf{W}^o = 0$.

2 Preliminaries

Let M be an n -dimensional manifold and TM its tangent bundle. We denote by $TM_0 = TM \setminus \{0\}$ the slit tangent bundle. Local coordinates on the base manifold M will be denoted by (x^i) , while the standard local coordinates on TM or TM_0 will be denoted by (x^i, y^i) .

A spray \mathbf{G} on M is a smooth vector field on TM_0 expressed in a standard local coordinate system (x^i, y^i) in TM as follows

$$\mathbf{G} = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i}, \quad (2.6)$$

where $G^i = G^i(x, y)$ are the local functions on TM satisfying $G^i(x, \lambda y) = \lambda^2 G^i(x, y)$ for $\forall \lambda > 0$. Every Finsler metric $F = F(x, y)$ on a manifold induces a spray with the *geodesic coefficients*

$$G^i = \frac{1}{4} g^{il} \{ [F^2]_{x^k y^l} y^k - [F^2]_{x^l} \}, \quad (2.7)$$

where $g^{ij} = (g_{ij})^{-1}$.

The vertical tangent bundle of TM_0 is expressed by

$$\mathcal{V}TM := \text{span} \left\{ \frac{\partial}{\partial y^1}, \dots, \frac{\partial}{\partial y^n} \right\}.$$

Put

$$\frac{\delta}{\delta x^k} := \frac{\partial}{\delta x^k} - N_k^l \frac{\partial}{\delta y^l}, \quad (2.8)$$

where $N_j^i = \frac{\partial G^i}{\partial y^j}$. Then the horizontal tangent bundle of TM_0 is expressed by

$$\mathcal{H}TM := \text{span} \left\{ \frac{\delta}{\delta x^1}, \dots, \frac{\delta}{\delta x^n} \right\}.$$

The tangent bundle $T(TM_0)$ of TM_0 is splitted into the direct sum $T(TM_0) = \mathcal{V}TM \oplus \mathcal{H}TM$.

Set $\omega^i = dx^i$ and $\omega^{n+i} := dy^i + N_j^i dx^j$. The local connection 1-forms of the Berwald connection are given by

$$\omega_j^i := \Gamma_{jk}^i dx^k,$$

where

$$\Gamma_{jk}^i = \frac{\partial N_j^i}{\partial y^k} = \frac{\partial^2 G^i}{\partial y^j \partial y^k}.$$

We have

$$d\omega^i = \omega^j \wedge \omega_j^i.$$

Then we obtain the curvature form:

$$\Omega_j^i = d\omega_j^i - \omega_j^l \wedge \omega_l^i. \quad (2.9)$$

Put

$$\Omega_j^i = \frac{1}{2} R_{j\ kl}^i \omega^k \wedge \omega^l - B_{j\ kl}^i \omega \wedge \omega^{n+l}. \quad (2.10)$$

The two curvature tensors $R_{j\ kl}^i$ and $B_{j\ kl}^i$ are called *Riemann curvature tensor* and *Berwald curvature tensor*, respectively.

The Riemann curvature $\mathbf{R}_y = R^i_k dx^k \otimes \frac{\partial}{\partial x^i}|_x : T_x M \rightarrow T_x M$ of a spray \mathbf{G} is a family of linear maps on tangent spaces which is defined by

$$R^i_k = y^j R_{j\ kl}^i y^l. \quad (2.11)$$

In a standard local coordinates, it is expressed by

$$R^i_k = 2 \frac{\partial G^i}{\partial x^k} - y^j \frac{\partial^2 G^i}{\partial x^j \partial y^k} + 2G^j \frac{\partial^2 G^i}{\partial y^j \partial y^k} - \frac{\partial G^i}{\partial y^s} \frac{\partial G^s}{\partial y^k}.$$

They are related by

$$R_{j\ kl}^i = \frac{1}{3} (R^i_{k\cdot l} - R^i_{l\cdot k})_{\cdot j}. \quad (2.12)$$

Here and hereafter, “ \cdot ” denotes the vertical derivatives with respect to y . For instance, $f_{\cdot k} = \frac{\partial f}{\partial y^k}$, $f_{\cdot k \cdot l} = \frac{\partial^2 f}{\partial y^k \partial y^l}$, etc. By (2.11) and (2.12), we see that the curvature tensors R^i_k and $R_{j\ kl}^i$ can represent each other. For this reason, they are all called *Riemann curvature tensor* if there is no confusion. Put

$$R^i_{kl} := y^j R_{j\ kl}^i. \quad (2.13)$$

By (2.12), one has

$$R^i_{kl} = \frac{1}{3} (R^i_{k\cdot l} - R^i_{l\cdot k}). \quad (2.14)$$

The Ricci curvature is the trace of \mathbf{R}_y defined by

$$\mathbf{Ric} := R^m_m = y^j R_{j\ ml}^m y^l \quad (2.15)$$

and $R := \frac{1}{n-1}\mathbf{Ric}$ is called the Ricci scalar of \mathbf{G} .

Define a linear map $\mathbf{T}_y : T_x M \rightarrow T_x M$ by

$$\mathbf{T}_y(u) = T^i_j(x, y)u^k \frac{\partial}{\partial x^i}, \quad u = u^k \frac{\partial}{\partial x^k},$$

where

$$T^i_j := R^i_j - (R\delta^i_j - \frac{1}{2}R_{.j}y^i). \quad (2.16)$$

A spray is said to be of *isotropic curvature* if $\mathbf{T} = 0$.

Given a volume form $dV = \sigma(x)dx^1 \cdots dx^n$ on a spray manifold (M, \mathbf{G}) . The S-curvature $\mathbf{S} = \mathbf{S}_{(G, dV)}$ of (\mathbf{G}, dV) is expressed by

$$\mathbf{S} := \frac{\partial G^m}{\partial y^m} - y^m \frac{\partial}{\partial x^m}(\ln \sigma). \quad (2.17)$$

There is a mysterious non-Riemannian quantities $\chi = \chi_i dx^i$. It is initially defined by the S-curvature \mathbf{S} in (2.17), but also given by the curvature T^i_j in (2.16)

$$\chi_i = -\frac{1}{3}T^m_{i.m} = \frac{1}{2}\{\mathbf{S}_{.i|m}y^m - \mathbf{S}_{|i}\}, \quad (2.18)$$

where the covariant derivative “|” is taken respect to \mathbf{G} .

Using $\mathbf{S} = \mathbf{S}_{(G, dV)}$, one can modify the spray \mathbf{G} to a new spray $\hat{\mathbf{G}}$ by

$$\hat{\mathbf{G}} := \mathbf{G} + \frac{2\mathbf{S}}{n+1}\mathbf{Y}, \quad (2.19)$$

where $\mathbf{Y} = y^i \frac{\partial}{\partial y^i}$ is the canonical vertical vector field. $\hat{\mathbf{G}}$ is a globally defined spray on M . We call $\hat{\mathbf{G}}$ the projective spray associated with \mathbf{G} on (M, dV) . In a standard local coordinates in TM , $\hat{\mathbf{G}} = y^i \frac{\partial}{\partial x^i} - 2\hat{G}^i \frac{\partial}{\partial y^i}$ is given by

$$\hat{G}^i = G^i - \frac{\mathbf{S}}{n+1}y^i.$$

By a direct computation, it is easy to see that the S-curvature $\hat{\mathbf{S}} = \mathbf{S}_{(\hat{\mathbf{G}}, dV)}$ always vanishes [7]. Thus the χ -curvature $\hat{\chi}$ of $\hat{\mathbf{G}}$ vanishes by (2.18). That is, for any (\mathbf{G}, dV) , the following holds

$$\hat{\mathbf{S}} = 0, \quad \hat{\chi} = 0. \quad (2.20)$$

Let N^i_k and \hat{N}^i_k be the non-linear connection coefficients of spray \mathbf{G} and $\hat{\mathbf{G}}$ and the Berwald connection coefficients are denoted by Γ^i_{jk} and $\hat{\Gamma}^i_{jk}$, respectively. It is easy to check that

$$\begin{aligned} \hat{N}^i_k &= N^i_k - \frac{\mathbf{S}}{n+1}\delta^i_k - \frac{\mathbf{S}_{.k}}{n+1}y^i \\ \hat{\Gamma}^i_{jk} &= \Gamma^i_{jk} - \frac{\mathbf{S}_{.k}}{n+1}\delta^i_j - \frac{\mathbf{S}_{.j}}{n+1}\delta^i_k - \frac{\mathbf{S}_{.k.j}}{n+1}y^i. \end{aligned} \quad (2.21)$$

Then by (2.8) and (2.21), we get

$$\frac{\hat{\delta}}{\delta x^k} = \frac{\delta}{\delta x^k} + \frac{\mathbf{S}_{\cdot k}}{n+1} \mathbf{Y} + \frac{\mathbf{S}}{n+1} \frac{\partial}{\partial y^k}, \quad (2.22)$$

where $\frac{\hat{\delta}}{\delta x^k}$ is defined in (2.8) by using $\hat{\mathbf{G}}$.

For the Riemann curvatures of \mathbf{G} and $\hat{\mathbf{G}}$, we have the following

Lemma 2.1 ([8]) *The Riemann curvatures and Ricci-scalar of \mathbf{G} and $\hat{\mathbf{G}}$ are related by*

$$\begin{aligned} \hat{R}^i_k &= R^i_k + \tau \delta_k^i - \frac{1}{2} \tau_{\cdot k} + \frac{3\chi_k}{n+1} y^i \\ \hat{R} &= R + \tau, \end{aligned} \quad (2.23)$$

where $\tau = (\frac{1}{n+1} \mathbf{S})^2 + \frac{1}{n+1} \mathbf{S}_{|m} y^m$ and $\chi_k = \frac{1}{2} (\mathbf{S}_{|m \cdot k} y^m - \mathbf{S}_{|k})$ and the covariant derivative “|” is taken respect to \mathbf{G} .

Let f be a function defined on TM . From (2.22), we have

$$f_{|k} = f_{|k} + \frac{\mathbf{Y}(f)}{n+1} \mathbf{S}_{\cdot k} + \frac{\mathbf{S}}{n+1} f_{\cdot k}, \quad (2.24)$$

where the covariant derivative “|” and “||” is taken respect to \mathbf{G} and $\hat{\mathbf{G}}$ respectively. Meanwhile, the Ricci identities are given by

$$f_{|m \cdot k} = f_{\cdot k |m}, \quad (2.25)$$

$$f_{|k |m} - f_{|m |k} = f_{\cdot l} R^l_{km}, \quad (2.26)$$

where R^l_{km} is defined in (2.14). Contracting (2.26) with y^m , we obtain

$$f_{|k |0} - f_{|0 |k} = f_{\cdot l} R^l_k. \quad (2.27)$$

The Weyl-type curvature of $\hat{\mathbf{G}}$ is given by

$$\hat{T}^i_k = \hat{R}^i_k - (\hat{R} \delta_k^i - \frac{1}{2} \hat{R}_{\cdot k} y^i).$$

It turns out that \hat{T}^i_k is independent of the volume form! Therefore it is a projective invariant. In fact, it is the well-known Weyl curvature

$$W^i_k = \hat{T}^i_k. \quad (2.28)$$

See Proposition 6.4 in [8]. Furthermore, one can rewrite (2.28) as

$$\begin{aligned} W^i_k &= T^i_k + \frac{3\chi_k}{n+1} y^i \\ &= R^i_k - (R \delta_k^i - \frac{1}{2} R_{\cdot k} y^i) + \frac{3\chi_k}{n+1} y^i. \end{aligned} \quad (2.29)$$

(2.29) is given in Lemma 3.3 in [8].

The Weyl-curvature \mathbf{W} can be used to characterize the sprays of scalar curvature.

Proposition 2.2 ([10][7]) *Let \mathbf{G} be a spray on a manifold. Then \mathbf{G} is of scalar curvature if and only if $\mathbf{W} = 0$.*

The following result proved by using the Bianchi identity will be used later.

Proposition 2.3 *For any spray \mathbf{G} on a n -dimensional manifold M , we have the following identity:*

$$R_{|p} - \frac{1}{2}R_{\cdot p|m}y^m - \frac{1}{n-1}(\chi_{p|m}y^m + R^m_{p|m}) = 0, \quad (2.30)$$

where the horizontal covariant derivative “|” is taken respect to \mathbf{G} .

Proof: The Bianchi identity says that

$$(R^i_{jkl|p} + B^i_{jpm}R^m_{kl}) + (R^i_{jlp|k} + B^i_{jlm}R^m_{pk}) + (R^i_{jpk|l} + B^i_{jkm}R^m_{lp}) = 0.$$

Contracting the above equation with y^j and using $y^j B^i_{jpm} = 0$, $y^j_{|p} = 0$ and $R^i_{kl} = y^j R^i_{jkl}$, it follows that

$$R^i_{kl|p} + R^i_{lp|k} + R^i_{pk|l} = 0.$$

Contracting it with y^l and using $R^i_k = R^i_{kl}y^l$ yields that

$$R^i_{k|p} - R^i_{p|k} + R^i_{pk|l}y^l = 0. \quad (2.31)$$

Then substituting (2.14) into (2.31), it follows that

$$R^i_{k|p} - R^i_{p|k} + \frac{1}{3}(R^i_{p\cdot k} - R^i_{k\cdot p})_{|l}y^l = 0. \quad (2.32)$$

Taking a trace of (2.32) over i and k , we get

$$R_{|p} - \frac{1}{3}R_{\cdot p|m}y^m = \frac{1}{n-1}(R^m_{p|m} - \frac{1}{3}R^m_{p\cdot m}). \quad (2.33)$$

Recall that the χ -curvature of \mathbf{G} is given by

$$\chi_i = -\frac{1}{6}[2R^m_{i\cdot m} + (n-1)R_{\cdot i}]$$

which means

$$R^m_{i\cdot m} = -3\chi_i - \frac{n-1}{2}R_{\cdot i}. \quad (2.34)$$

Substituting (2.34) into equality (2.33), we get the result. Q.E.D.

3 The Berwald-Weyl curvature \mathbf{W}^o

In this section, we shall discuss the Berwald-Weyl curvature. Let (\mathbf{G}, dV) be a spray on an n -manifold M . Let $\hat{\mathbf{G}}$ be the associated spray defined in (2.19). For a vector $y \in T_x M \setminus \{0\}$, the Berwald-Weyl curvature $\mathbf{W}_y^o : T_x M \rightarrow R$ is defined by

$$\mathbf{W}_y^o(u) := W_k^o(y)u^k, \quad u = u^k \frac{\partial}{\partial x^k},$$

where

$$W_k^o := \hat{R}_{\parallel k} - \frac{1}{2} \hat{R}_{\cdot k \parallel m} y^m. \quad (3.35)$$

Here $\hat{R} = \frac{1}{n-1} \hat{R}_m^m$ denotes the Ricci scalar of $\hat{\mathbf{G}}$ and the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$. By Ricci identity (2.25), we can rewrite W_k^o as

$$W_k^o = \frac{1}{2} \{3\hat{R}_{\parallel k} - (\hat{R}_{\parallel m} y^m)_{\cdot k}\}. \quad (3.36)$$

Since $\hat{\mathbf{G}}$ is a projective invariant with respect to a fixed volume form dV , \mathbf{W}^o is a projective invariant. By the homogeneity of \hat{R} and $y_{\parallel m}^k = 0$, it follows from (3.35) or (3.36) that

$$W_k^o y^k = 0.$$

Applying formula (2.30) to $\hat{\mathbf{G}}$ and using (2.20), we obtain

$$\hat{R}_{\parallel k} - \frac{1}{2} \hat{R}_{\cdot p \parallel m} y^m = \frac{1}{n-1} \hat{R}_{k \parallel m}^m, \quad (3.37)$$

where the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$ and $\hat{R}_{\cdot k}^m$ is the Riemann curvature of $\hat{\mathbf{G}}$. By (3.35) and (3.37), we immediately obtain another expression for W_k^o

$$W_k^o = \frac{1}{n-1} \hat{R}_{k \parallel m}^m.$$

Proof of Proposition 1.3: Firstly, according to (2.23) and (2.24), one gets

$$\hat{R}_{\parallel k} = (R + \tau)_{\parallel k} + \frac{2(R + \tau)}{n+1} \mathbf{S}_{\cdot k} + \frac{(R + \tau)_{\cdot k}}{n+1} \mathbf{S} \quad (3.38)$$

$$\hat{R}_{\parallel m} y^m = (R + \tau)_{\parallel m} y^m + \frac{4(R + \tau)}{n+1} \mathbf{S}. \quad (3.39)$$

Taking the derivative of (3.39) with respect to y^k yields that

$$(\hat{R}_{\parallel m} y^m)_{\cdot k} = (R + \tau)_{\parallel k} + (R + \tau)_{\parallel m \cdot k} y^m + \frac{4(R + \tau)}{n+1} \mathbf{S}_{\cdot k} + \frac{4(R + \tau)_{\cdot k}}{n+1} \mathbf{S} \quad (3.40)$$

Plugging (3.38) and (3.40) into (3.36), we obtain

$$\begin{aligned}
W_k^o &= R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{2(n+1)}[\mathbf{S}_{|l\cdot k|m}y^m y^l - \mathbf{S}_{|k|m}y^m] \\
&- \frac{3\mathbf{S}}{2(n+1)^2}(\mathbf{S}_{|m\cdot k}y^m - \mathbf{S}_{|k}) - \frac{1}{n+1}[\mathbf{S}_{|k|m}y^m - \mathbf{S}_{|m|k}y^m] \\
&+ \frac{R\mathbf{S}_{\cdot k}}{n+1} - \frac{\mathbf{S}R_{\cdot k}}{2(n+1)}. \tag{3.41}
\end{aligned}$$

Now, by using (2.18) and (2.26), W_k^o in (3.41) reduces to

$$\begin{aligned}
W_k^o &= R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m - \frac{3\mathbf{S}}{(n+1)^2}\chi_k \\
&- \frac{1}{n+1}R_k^l \mathbf{S}_{\cdot l} + \frac{R\mathbf{S}_{\cdot k}}{n+1} - \frac{\mathbf{S}R_{\cdot k}}{2(n+1)} \tag{3.42}
\end{aligned}$$

The last three terms in (3.42) are rewritten by

$$\begin{aligned}
&-\frac{1}{n+1}R_k^l \mathbf{S}_{\cdot l} + \frac{R\mathbf{S}_{\cdot k}}{n+1} - \frac{\mathbf{S}R_{\cdot k}}{2(n+1)} \\
&= -\frac{1}{n+1}[R_k^l - (R\delta_k^l - \frac{1}{2}R_{\cdot k}y^l)]\mathbf{S}_{\cdot l} \\
&= -\frac{1}{n+1}T_k^l \mathbf{S}_{\cdot l}, \tag{3.43}
\end{aligned}$$

where T_k^l is defined in (2.16). Taking (3.43) into (3.42), we obtain

$$W_k^o = R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{n+1}\chi_{k|j}y^j - \frac{3\mathbf{S}}{(n+1)^2}\chi_k - \frac{1}{n+1}T_k^l \mathbf{S}_{\cdot l}. \tag{3.44}$$

Finally, submitting (2.29) into (3.44), we obtain

$$W_k^o = R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m - \frac{1}{n+1}W_k^m \mathbf{S}_{\cdot m}, \tag{3.45}$$

This completes the proof.

Q.E.D.

Note that if a Finsler metric F has constant S-curvature, that is, the S-curvature of F with respect to $dV = dV_{BH}$ satisfies that $\mathbf{S} = (n+1)cF$, $c = \text{constant}$, then $\chi = 0$. By (3.45), we obtain Theorem 1.5. We can generalize Theorem 1.5 as follows.

Proposition 3.1 *Let F be a Finsler metric on an n -dimensional manifold M . If F is of constant S-curvature and is Einstein with $\mathbf{Ric} = (n-1)\sigma F^2$, then*

$$W_k^o = F^3 \left(\frac{\theta}{F} \right)_{\cdot k},$$

for the $dV = dV_{BH}$, where $\theta = \sigma_{x^m} y^m$.

Proof: By assumption, we have

$$\mathbf{S} = (n+1)cF, \quad R = \sigma F^2, \quad (3.46)$$

where c is a constant and $\sigma = \sigma(x)$ is a function on M . Then by (2.18), we have

$$\chi = 0, \quad \mathbf{S}_{.m} = (n+1)cF^{-1}y_m, \quad (3.47)$$

where $y_m := \frac{1}{2}(F^2)_{.m}$. By using (3.47) and (2.29), we get

$$W_k^m \mathbf{S}_{.m} = -(R\mathbf{S}_{.k} - \frac{1}{2}\mathbf{S}R_{.k}) = 0, \quad (3.48)$$

where we use $R_k^m y_m = 0$. On the other hand, by a direct calculation, we obtain

$$R_{|k} - \frac{1}{2}R_{.k|m}y^m = F^3\left(\frac{\theta}{F}\right)_{.k} \quad (3.49)$$

with $\theta = \sigma_{x^k}y^k$. Plugging (3.48) and (3.49) into (3.45), we complete the proof. Q.E.D.

It is known that if a Finsler metric is of constant flag curvature, then $\chi = 0$. By Corollary 1.4, we have

Corollary 3.2 *For any two-dimensional Finsler metric of constant flag curvature, $\mathbf{W}^o = 0$.*

Proof of Proposition 1.1: Taking the horizontal covariant derivatives “|” with respect to \mathbf{G} of (2.29), we have

$$W_{k|m}^m = R_{k|m}^m - (R_{|k} - \frac{1}{2}R_{.k|m}y^m) + \frac{3}{n+1}\chi_{k|m}y^m. \quad (3.50)$$

By (2.30), it follows that

$$R_{k|m}^m = (n-1)(R_{|k} - \frac{1}{2}R_{.k|m}y^m) - \chi_{k|m}y^m. \quad (3.51)$$

Substituting (3.51) into (3.50) yields that

$$W_{k|m}^m = (n-2)(R_{|k} - \frac{1}{2}R_{.k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m). \quad (3.52)$$

By (3.52) and the assumption $n \geq 3$, W_k^0 in (3.45) becomes

$$\begin{aligned} W_k^0 &= \frac{1}{n-2}W_{k|m}^m - \frac{1}{n+1}W_k^m \mathbf{S}_{.m} \\ &= \frac{1}{n-2}W_{k||m}^m, \end{aligned} \quad (3.53)$$

where the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$. Q.E.D.

We have another method to prove Proposition 1.1. Firstly, taking the horizontal covariant derivatives “ \parallel ” with respect to \mathbf{G} from (2 .16), we obtain

$$T_{k|m}^m = R_{k|m}^m - (R_{|k} - \frac{1}{2}R_{.k|m}y^m).$$

Plugging (3 .51) into preceding equation yields that

$$T_{k|m}^m = (n - 2)(R_{|k} - \frac{1}{2}R_{.k|m}y^m) - \chi_{k|m}y^m. \quad (3 .54)$$

Let \hat{T}_k^i be the Weyl-type curvature of projective spray $\hat{\mathbf{G}}$. Applying (3 .54) to $\hat{\mathbf{G}}$ and using (2 .20), we obtain

$$\hat{T}_{k|m}^m = (n - 2)(\hat{R}_{|k} - \frac{1}{2}\hat{R}_{.k|m}y^m), \quad (3 .55)$$

where the covariant derivative “ \parallel ” is taken respect to $\hat{\mathbf{G}}$. Therefore, according to (3 .35), (3 .55) and (2 .28), we obtain

$$W_k^o = \hat{R}_{|k} - \frac{1}{2}\hat{R}_{.k|m}y^m = \frac{1}{n-2}\hat{T}_{k|m}^m = \frac{1}{n-2}W_{k|m}^m.$$

4 BWeyl-flat

In this section, we shall define the notion of BWeyl-flat sprays and give three equivalent conditions for a spray to be BWeyl-flat when $\dim M \geq 3$.

Let \mathbf{G} be a spray on an n -dimensional manifold M of $n \geq 3$. By (3 .45), we know that the Berwald-Weyl curvature \mathbf{W}^o of \mathbf{G} depends on the volume form dV in general. We make the following

Definition 4.1 *A spray \mathbf{G} on an n -dimensional manifold M of $n \geq 3$ is said to be BWeyl-flat if there is a volume form dV on M such that Berwald-Weyl curvature \mathbf{W}^o vanishes, that is*

$$\mathbf{W}^o = \mathbf{0}.$$

A Finsler metric F on M is said to be BWeyl-flat if the induced spray $\mathbf{G} = \mathbf{G}_F$ is BWeyl-flat.

Remark 4.2 *We have obtained from Theorem1.2 that a spray of scalar curvature must have $\mathbf{W}^o = \mathbf{0}$ for any volume form. Therefore, a spray of scalar curvature must be BWeyl-flat in our sense.*

Let dV and $d\tilde{V}$ be volume forms on M with $dV = e^{(n+1)f}d\tilde{V}$, where $f = f(x)$ is a scalar function on M . Then the S-curvatures $\mathbf{S} = \mathbf{S}_{(G,dV)}$ and $\tilde{\mathbf{S}} = \mathbf{S}_{(G,d\tilde{V})}$ are related by

$$\mathbf{S} = \tilde{\mathbf{S}} - (n + 1)f_0, \quad (4 .56)$$

where $f_0 := f_{x^m}(x)y^m$. By (4.56), one sees that $\mathbf{S}_{(G,dV)} = (n+1)\phi_0$ if and only if $\mathbf{S}_{(G,d\tilde{V})} = (n+1)\tilde{\phi}_0$ with $\phi_0 = \tilde{\phi}_0 - f_0$.

Let W_k^o and \widetilde{W}_k^o be the Berwald-Weyl curvature of \mathbf{G} with respect to volume form dV and $d\tilde{V}$ respectively. It follows from (3.45) that

$$W_k^o = R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m - \frac{1}{n+1}W_k^m \mathbf{S}_{\cdot m}, \quad (4.57)$$

$$\widetilde{W}_k^o = R_{|k} - \frac{1}{2}R_{\cdot k|m}y^m - \frac{1}{n+1}\chi_{k|m}y^m - \frac{1}{n+1}W_k^m \tilde{\mathbf{S}}_{\cdot m}, \quad (4.58)$$

where “|” is the horizontal covariant derivative with respect to \mathbf{G} . It follows from (4.56), (4.57) and (4.58) that

$$\widetilde{W}_k^o = W_k^o - W_k^m f_m. \quad (4.59)$$

Now we give some equivalent conditions for a spray to be BWeyl-flat in $\dim M \geq 3$.

Theorem 4.3 *Let \mathbf{G} be a spray on an n -dimensional manifold M of $n \geq 3$. Then the following are equivalent:*

- (a) \mathbf{G} is of BWeyl-flat.
- (b) for any volume form dV on M there is a scalar function f on M such that

$$W_k^o = W_k^m f_m. \quad (4.60)$$

- (c) for any volume form dV on M there is a scalar function f on M such that

$$W_{k|m}^m = (n-2)W_k^m \Xi_{\cdot m}, \quad (4.61)$$

where “|” is the horizontal covariant derivative with respect to \mathbf{G} and $\Xi := \frac{\mathbf{S}}{n+1} + f_0$ with $f_0 = f_{x^m}y^m$ and $\mathbf{S} = \mathbf{S}_{(G,dV)}$.

Proof: (a) \Rightarrow (b). Assume that for some volume form $d\tilde{V}$,

$$\widetilde{\mathbf{W}}^o = 0.$$

Then by (4.59), for any volume form $dV = e^{(n+1)f}d\tilde{V}$,

$$W_k^o = W_k^m f_m. \quad (4.62)$$

(b) \Rightarrow (c). By assumption and (3.53), we have

$$\begin{aligned} W_{k|m}^m &= \frac{n-2}{n+1}W_k^m \mathbf{S}_{\cdot m} + (n-2)W_k^m f_m \\ &= (n-2)W_k^m \Xi_{\cdot m}, \end{aligned} \quad (4.63)$$

where $\Xi = \frac{\mathbf{S}}{n+1} + f_0$.

(c) \Rightarrow (a). Let dV be an arbitrary volume form on M . There is a scalar function f on M such that (4.61) holds. That is

$$W_{k|m}^m = (n-2)W_k^m \Xi_{.m}.$$

Taking $d\tilde{V} = e^{-(n+1)f}dV$, we have

$$\tilde{\mathbf{S}} = \mathbf{S} + (n+1)f_0 = (n+1)\Xi.$$

By (3.53), (4.59) and (4.63), we obtain

$$\widetilde{W}_k^o = W_k^o - W_k^m f_m = \frac{1}{n-2}W_{k|m}^m - \frac{1}{n+1}W_k^m \mathbf{S}_{.m} - W_k^m f_m = 0.$$

The condition on the S-curvature $\mathbf{S}_{(G,dV)} = (n+1)\phi_0$ being an exact 1-form, is actually a condition on the spray, independent of the choice of a particular volume form dV . If a spray \mathbf{G} satisfies that $\mathbf{S}_{(G,dV)} = (n+1)\phi_0$ for some scalar function ϕ on M , then $\Xi = 0$ for $f = -\phi$ holds, where $\Xi = \frac{\mathbf{S}}{n+1} + f_0$. Thus in view of (4.61), we obtain the following

Corollary 4.4 *Let \mathbf{G} be a spray on an n -dimensional manifold M of $n \geq 3$. Suppose that $W_{k|m}^m = 0$ and $\mathbf{S}_{(G,dV)} = (n+1)\phi_0$ for some scalar function ϕ , then \mathbf{G} is BWeyl-flat.*

A spray is said to be projectively Ricci-flat if there is a volume form dV on M such that $\hat{R} = \frac{1}{n-1}\mathbf{Ric}_{\hat{G}} = 0$. A Finsler metric F on M is said to be projectively Ricci-flat if the induced spray $\mathbf{G} = \mathbf{G}_F$ is projectively Ricci-flat[9]. Thus from (1.2), if a spray or a Finsler metric is projectively Ricci-flat, then it is BWeyl-flat.

Example 4.5 *Let $\alpha := (1+4|x|^2)\sqrt{\frac{(1+4|x|^2)|y|^2-4(x,y)}{1+4|x|^2}}$ and $\beta = 2\langle x, y \rangle$. It is shown that the square metric $F = \frac{(\alpha+\beta)^2}{\alpha}$ is projectively Ricci-flat[11]. Thus F is BWeyl-flat. On the other hand, by using Theorem 1.2 in [5], one can see that F is not of isotropic S-curvature by a direct computation.*

References

- [1] L. Berwald, Über systeme von gewöhnlichen differetialgleichungen zweiter ordnung, deren integralkurven mit dem system der geraden linien topologisch äquivalent sind, Ann. of Math. **48** (1947), 193-215.
- [2] D. Bao, C. Robles, On Ricci curvature and flag curvature in Finsler geometry, In: "A sampler of Finsler Geometry" MSRI series, Cambridge University Press, 2004.
- [3] S. S. Chern and Z. Shen, *Riemann Finsler Geometry, Nankai Tracts in Mathematics*, Vol. 6, World Scientific Co., Singapore, 2005.

- [4] X. Cheng and Z. Shen, *Randers metrics with special curvature properties*, Osaka J. Math. **40** (2003), 87-101.
- [5] X. Cheng and Z. Shen, *A class of Finsler metrics with isotropic S-curvature*, Israel J. Math. **169** (2009), 317-340.
- [6] J. Douglas, The general geometry of path, Ann. Math. **29** (1927-28), 143-168.
- [7] Z. Shen, *Differential Geometry of Spray and Finsler Spaces*, Kluwer Academic Publishers, 2001.
- [8] Z. Shen, *On sprays with vanishing χ -curvature*, International Journal of Mathematics, **32** (2021).
- [9] Z. Shen and L. Sun. On the projective Ricci curvature, Sci. China Math, **64** (2021), 1629-1636.
- [10] Z. Szabó, Ein Finslerscher Raum ist gerade dann von skalarer Krümmung, wenn seine Weyl sche ProjectivKrümmung verschwindet, Acta Sci. Math. **39** (1977), 163-168.
- [11] H. Zhu, On a class of projectively Ricci-flat Finsler metrics, J. Diff. Geom. **73** (2020)101680.

Zhongmin Shen
 Department of Mathematical Sciences,
 Indiana University-Purdue University Indianapolis, IN 46202-3216, USA.
 zshen@iupui.edu

Liling Sun
 School of Science,
 Jimei University, Xiamen 361021, P.R. China.
 sunliling@jmu.edu.cn