

ON THE STRUCTURE OF T - λ -SPHERICAL COMPLETIONS OF O-MINIMAL FIELDS.

PIETRO FRENI

ABSTRACT. Let T be the theory of an o-minimal field and T_0 a common reduct of T and T_{an} .

I adapt Mourgues' and Ressayre's constructions to deduce structure results for T_0 -reducts of T - λ -spherical completion of models of T_{convex} .

These in particular entail that whenever \mathbb{R}_L is a reduct of $\mathbb{R}_{an,exp}$ defining the exponential, every elementary extension of \mathbb{R}_L has an elementary truncation-closed embedding in \mathbf{No} . This partially answers a question in [3].

The main technical result is that certain expansion of Hahn fields by generalized power series interpreted as functions defined on the positive infinitesimal elements, have the property that truncation closed subsets generate truncation closed substructures. This leaves room for possible generalizations to the case in which T_0 is power bounded but not necessarily a reduct of T_{an} .

CONTENTS

1. Introduction	2
1.1. Motivation and background	2
1.2. Setting and Main Results	3
1.3. Structure of the paper	5
1.4. Acknowledgments	6
2. Preliminaries on T -convexity	6
2.1. T -convexity	6
2.2. Wim-constructible extensions and λ -spherical completions	7
3. Truncation closed expansions of Hahn fields	9
3.1. Well-ordered subsets of ordered Abelian groups	9
3.2. Rings of generalized series	11
3.3. Generalized power series	12
3.4. Infinitesimally convergent power series	16
4. Serial power-bounded structures	19
5. The Mourgues-Ressayre constructions revisited	21
References	26

2020 *Mathematics Subject Classification*. Primary 16W60, 03C64; Secondary 12J15, 06F20, 06F25.

1. INTRODUCTION

1.1. Motivation and background. Recall that given an (ordered) field \mathbb{K} and a linearly ordered group $(\mathfrak{M}, \cdot, <)$ the (ordered) *Hahn field* $\mathbb{K}((\mathfrak{M}))$ consists of the formal series $f := \sum_{\mathfrak{m}} k_{\mathfrak{m}} \mathfrak{m}$ whose support $\text{Supp}(f) := \{\mathfrak{m} \in \mathfrak{M} : k_{\mathfrak{m}} \neq 0\}$ is well ordered with respect to the opposite order on \mathfrak{M} . The field operations are given by the term-wise sum and Cauchy's product formula.

A distinctive feature of these objects is that they have an extra notion of *infinite sum* that is useful in applications, for example it allows for a natural interpretation of power series at infinitesimal elements (cf [12]).

Subfields of $\mathbb{K}((\mathfrak{M}))$ are usually referred to as *fields of generalized series* and can inherit the extra *summability structure* of the ambient $\mathbb{K}((\mathfrak{M}))$.

Fields of generalized series played an important role in the study of o-minimal fields (see for example [14], [21] or, more recently, [16]).

The study of their relation with o-minimal fields follows several line of investigation. The ones that will be addressed in this paper can be broadly synthesized in the following two loose questions:

- Q1 let \mathbb{K} be an o-minimal field, and $\mathbb{E} \subseteq \mathbb{K}((\mathfrak{M}))$ some field of generalized series containing \mathbb{K} , when does \mathbb{E} admit an expansion to an elementary extension of \mathbb{K} ?
- Q2 let \mathbb{K} be an o-minimal structure, and let $\mathbb{E} \subseteq \mathbb{K}((\mathfrak{M}))$ be given a structure of elementary extension of \mathbb{K} , what elementary extensions of \mathbb{K} admit *truncation closed* elementary embeddings into \mathbb{E} ?

By a truncation of $f \in \mathbb{K}((\mathfrak{M}))$ we mean an element of the form $f|_{\mathfrak{m}} := \sum_{\mathfrak{n} > \mathfrak{m}} k_{\mathfrak{n}} \mathfrak{n}$ and *truncation closed* means closed under taking truncations of the elements.

The fields of generalized series $\mathbb{E} \subseteq \mathbb{K}((\mathfrak{M}))$ for which the inherited summability structure is more relevant are obtained by restricting the family of allowed supports in the definition of Hahn field from all well ordered sets to some suitable ideal B of subsets of \mathfrak{M} (cf [1], [7]), such subfields will be denoted by $\mathbb{K}((\mathfrak{M}))_B$. These are the fields for which also the two questions above seem more relevant.

If B consists of the well ordered subsets with cardinality strictly less then some uncountable cardinal λ , $\mathbb{K}((\mathfrak{M}))_B$ is denoted by $\mathbb{K}((\mathfrak{M}))_{\lambda}$ and is called a λ -*bounded* Hahn field.

The first question is known to have positive answer in the case $\mathbb{E} = \mathbb{K}((\mathfrak{M}))$ and \mathbb{K} is power-bounded whenever \mathfrak{M} is a vector space over the exponents of \mathbb{K} (this is a consequence of the so called residue-valuation property, see [23, Sec.s 9 and 10], [18, Ch. 12 and 13] and [6], or [5, Sec. 4] for a self-contained treatment).

On the contrary, if \mathbb{K} is exponential, [8, Thm. 1], entails that no field of the form $\mathbb{K}((\mathfrak{M}))$ admits an expansion to an elementary extension of \mathbb{K} .

It is also known that for suitably constructed pairs (\mathfrak{M}, B) , $\mathbb{R}((\mathfrak{M}))_B$ can naturally be expanded to elementary extension of $\mathbb{R}_{an,exp}$: most notable

examples are the field of LE and EL transexponential series and the certain constructions of λ -bounded Hahn fields (cf [22], [9]).

Such naturally constructed extensions often enjoy the following two properties of compatibility of the exponential with the “serial” structure:

(D) $\log(\mathfrak{M}) \subseteq \mathbb{K}((\mathfrak{M}^{>1}))_B$ (cf [14])

(T4) for every sequence of monomials $(\mathfrak{m}_n)_{n \in \mathbb{N}}$ with $\mathfrak{m}_{n+1} \in \text{Supp}(\log \mathfrak{m}_n)$ for all $n \in \mathbb{N}$, there is $N \in \mathbb{N}$ such that for all $n \geq N$, $\log \mathfrak{m}_n = c_n \pm \mathfrak{m}_{n+1}$ for some c_n with $\text{Supp } c_n > \mathfrak{m}_{n+1}$ (cf [17]).

In that regard Q1 can be further specialized to the question of whether such structure can be chosen so that it satisfies this two extra conditions.

Answers to the second question have instead been given in [11] for real closed fields and in [4] and [3] respectively for $\mathbb{K} = \mathbb{R}_{an, \exp}$ and $\mathbb{K} = \mathbb{R}_{W, \exp}$ with W a Weierstrass system and $\mathbb{K}((\mathfrak{M}))_B = \mathbf{No}$ the (class-sized) field of surreal numbers with its natural (W, \exp) structure.

More recently, in [16], Rolin, Servi, and Speissegger, obtain related results for certain Generalized Quasianalytic Algebras as defined in [15]. More specifically, for $\mathcal{A} = an^*$ or \mathcal{A} a truncation closed and natural GQA containing the restricted exponential (see [16, Def. 3.4]), they explicitly construct a truncation-closed ordered differential field embedding of the Hardy field $\mathcal{H}(\mathbb{R}_{\mathcal{A}, \exp})$ of the o-minimal structure $\mathbb{R}_{\mathcal{A}, \exp}$, in the field \mathbb{T} of transseries. Such embedding is also an $L_{\mathcal{A}, \exp}$ -embedding if \mathbb{T} is given a suitable natural $L_{\mathcal{A}}$ -structure ([16, Sec. 4.2]).

In [5, Thm. B], the author showed that every T -convexly valued o-minimal field, admits for every cardinal λ a so-called T - λ -spherical completion, that is, a unique-up-to-non-unique-isomorphism elementary extension that is prime (i.e. weakly initial) among all the λ -spherically complete elementary extensions, and that such completion preserves the residue field.

This provides leverage toward further partial answers to Q1: it is not hard to see that given any T -convexly valued $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$, for λ large enough, the real closed reduct of a T - λ -spherical completion has the form $\mathbb{K}((\mathfrak{M}))_\lambda$ with \mathbb{K} the residue field of $(\mathbb{E}, \mathcal{O})$ and \mathfrak{M} some ordered group.

The present paper aims at providing further partial answers towards the two line of investigation Q1 and Q2. In that regard we will obtain, as corollaries of Theorem C, that in the case T defines an exponential:

C1 for λ large enough the reduct of the T - λ -spherical completion of $(\mathbb{E}, \mathcal{O})$ to the language of valued exponential field is isomorphic to a field of the form $\mathbb{K}((\mathfrak{M}))_\lambda$ endowed with an exponential satisfying D and T4;

C2 if T is a reduct of $T_{an, \exp}$ defining the exponential, then elementary extensions of the T -reduct \mathbb{R}_T of $\mathbb{R}_{an, \exp}$ admit truncation closed elementary embeddings into the field \mathbf{No} of surreal numbers.

1.2. Setting and Main Results. Let \mathbb{K} be an ordered field with powers from a field Λ and \mathfrak{M} be a multiplicatively written ordered Λ -vector space.

Recall that an n -varied non-singular generalized power series is a formal expression

$$f(\mathbf{x}) = \sum_{\gamma} c_{\gamma} \mathbf{x}^{\gamma}$$

with \mathbf{x} an n -tuple of variables, γ ranging over the n -tuples in Λ with non-negative entries and $c_{\gamma} \in \mathbb{K}$, such that $\{\gamma : c_{\gamma} \neq 0\}$ is a well-partial order in Λ^n (with the product order). Each such f can be interpreted as a function on the positive infinitesimals of the Hahn field $\mathbb{K}(\mathfrak{M})$.

The first main result of the paper is a purely formal fact about Hahn fields expanded with a set \mathcal{F} of such series. Namely:

Theorem A (3.20). *If the family \mathcal{F} is truncation-closed and closed under $f(x_0, \dots, x_{n-1}) \mapsto x_i \partial_{x_i} f(x_0, \dots, x_{n-1})$, and $X \subseteq \mathbb{K}(\mathfrak{M})$ is closed under truncations, then the smallest $(\mathcal{F}, +, \cdot)$ -structure generated by $X \cup \mathfrak{M} \cup \mathbb{K}$ is closed under truncations.*

The proof is by reducing this to a similar problem concerning classical power series with coefficients in $\mathbb{K}(\mathfrak{M})$ that are convergent on the set of infinitesimal elements (restricted power series). In fact we will deduce Theorem A from

Theorem B (3.30). *Let \mathcal{L} be a family of restricted power series with coefficients in $\mathbb{K}(\mathfrak{M})$, and suppose \mathcal{L} is closed under formal derivatives and coefficient-wise truncations, then the smallest ring containing $\mathbb{K} \cup \mathfrak{M}$ and closed under functions in \mathcal{L} is closed under truncations.*

We will then consider power-bounded o-minimal structures \mathbb{K} such that $\mathbb{K}(\mathfrak{M})$ expands naturally to an elementary extension $\mathbb{K}(\mathfrak{M})^{\mathcal{T}}$ of \mathbb{K} where the germs of enough functions definable in \mathbb{K} are interpreted on infinitesimals in $\mathbb{K}(\mathfrak{M})$ via generalized power series (*serial* structures, Definition 4.1).

The main example of serial structure is \mathbb{R}_{an} , but it is easy to show that all reducts of \mathbb{R}_{an} can be definitionally expanded to serial structures (Corollary 4.9). Although the definition of serial power bounded structure is modeled upon the definition of Generalized Quasianalytic Algebra (GQA) in [15], we leave open the question of whether all expansion of the reals by a GQA is interdefinable with a serial structure (4.11).

If the algebras of generalized power series used to interpret these germs are *closed under truncations*, Theorem A then ensures that each $\mathbb{K}(\mathfrak{M})^{\mathcal{T}}$, has the property that whenever $\mathbb{K} \preceq \mathbb{E} \preceq \mathbb{K}(\mathfrak{M})^{\mathcal{T}}$ is truncation closed and $x \in \mathbb{K}(\mathfrak{M})$ has all proper truncations in \mathbb{E} , the definable closure $\mathbb{E}\langle x \rangle$ of $\mathbb{E} \cup \{x\}$ is still truncation closed (Proposition 4.7).

This fact together with results from [5] allows to redo Mourgues' and Ressayre's constructions in [11] and [14] of truncation closed embeddings (satisfying (D) and (T4) in the case the models in question are expanded with a compatible exponential). We will need a minor modification of their construction to take into account the fact that we are aiming in general at embeddings into the λ -bounded versions of the Hahn fields.

The main results are best stated in terms of a structure theorem for the embedding of $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$ into its T - λ -spherical completion for suitably big λ .

Theorem C (5.11). *Let \mathbb{E} be a tame extension of \mathbb{K} , λ be a large enough cardinal and \mathbb{E}_λ the T - λ -spherical completion of $(\mathbb{E}, \text{CH}(\mathbb{K})) \models T_{\text{convex}}$ and let \mathfrak{M} be a section of the value group of \mathbb{E}_λ . Let \mathbb{K}_0 be a serial power-bounded reduct of \mathbb{K} , so that naturally $\mathbb{K}((\mathfrak{M})) \models \text{ED}(\mathbb{K}_0)$. Then there is a $\text{ED}(\mathbb{K}_0)$ -isomorphism $\eta : \mathbb{E}_\lambda \rightarrow \mathbb{K}((\mathfrak{M}))_\lambda$ such that $\text{exp}_* = \eta \circ \text{exp} \circ \eta^{-1}$ satisfies (D) and (T4) and $\eta(\mathbb{E})$ is truncation closed.*

As observed by Mantova, combined with [3, Thm 8.1], this implies C2 (Corollary 5.12).

Theorem C and the above mentioned [16, Main Theorem] are related, but differ in several regards. To explain the differences consider the specialization of Theorem C to the case in which T is the theory of the expansion $\mathbb{R}_{0,\text{exp}}$ by the unrestricted exponential, of a *serial* polynomially bounded structure \mathbb{R}_0 over the reals, already defining the restricted exponential. In that case, Theorem C says that every elementary extension $\mathbb{E} \succeq \mathbb{R}_{0,\text{exp}}$ has a truncation-closed *elementary* embedding (over $\mathbb{R}_{0,\text{exp}}$) in some elementary extension of the form $\mathbb{R}_{0,\text{exp}}((\mathfrak{M}))_\lambda$ where the functions definable in the structure \mathbb{R}_0 are interpreted in a natural way and the exponential is interpreted in such a way that (D) and (T4) are satisfied.

Notice in particular that Theorem C is conditional to the seriality hypothesis on \mathbb{R}_0 (which, for now, we only know to hold in the case \mathbb{R}_0 is a reduct of T_{an}), that there is no required compatibility with derivatives (as one could for example require if $(\mathbb{E}, \text{CH}(\mathbb{R}))$ is expanded to a T -convex T -differential field as defined in [6]), and that the group \mathfrak{M} arises from a rather abstract, and by no means explicit, completion machinery.

On the contrary, [16, Main Theorem], concerns the case $\mathbb{E} := \mathbb{R}_{\mathcal{A},\text{exp}}\langle t \rangle$ for some $t > \mathbb{R}$ and gives an *explicit* construction for a truncation-closed *differential* embedding with respect to the natural “derivation at t ”, into the (explicitly constructed) classical field of transseries \mathbb{T} . Moreover it does not require the seriality of $\mathbb{R}_{\mathcal{A}}$, and only requires an extra minor hypothesis (being an^* or being natural) on the GQA \mathcal{A} .

1.3. Structure of the paper. Section 2 reviews some preliminaries on T -convexity.

Section 3 is dedicated to the results on formal series: subsections 3.2 and 3.3 are dedicated to the main definitions and the statement of Theorem A; subsection 3.4 contains the reduction to the restricted power series together with the main proofs, both of Theorems A and B.

Section 4 defines the scope of applications of the results on formal series to o-minimal structure: it defines serial power-bounded structures and gives some examples and basic properties.

Finally Section 5 is dedicated the revisitation of Mourgues and Ressayre's constructions and to the proof of Theorem C.

1.4. Acknowledgments. The author is very grateful to his supervisors Vincenzo Mantova for his support, constant feedback and for pointing out Corollary 5.12, and Dugald Macpherson for his support and feedback.

This paper is part of a PhD project at the University of Leeds and is supported by a Scholarship of the School of Mathematics.

2. PRELIMINARIES ON T -CONVEXITY

In this section we briefly review T -convexity recalling the main results of [20], [19], and import some definitions and results from the preprint [5]. If T is the theory of an o-minimal field, $\mathbb{E} \prec \mathbb{E}_1 \models T$, and $S \subseteq \mathbb{E}_1$ we will denote by $\mathbb{E}\langle S \rangle_T := \text{dcl}_T(\mathbb{E}S)$ the T -definable closure of $\mathbb{E} \cup \{S\}$. We will omit the subscript T if it is clear from the context.

2.1. T -convexity. Let T be an o-minimal theory expanding RCF in a language L . If $\mathbb{E} \models T$, recall that a T -convex subring of \mathbb{E} is a convex subring of \mathbb{E} which is closed by continuous T -definable functions $f : \mathbb{E} \rightarrow \mathbb{E}$ (by T -definable we mean \emptyset -definable in T). It is said to be *non-trivial* if $\mathbb{E} \neq \mathcal{O}$. It is not hard to see that if $\mathbb{K} \preceq \mathbb{E}$, then the convex hull $\mathcal{O} := \text{CH}_{\mathbb{E}}(\mathbb{K})$ of \mathbb{K} in \mathbb{E} is a T -convex valuation subring.

Denote by L_{convex} the language obtained expanding L with a unary predicate \mathcal{O} and by T_{convex}^- the theory given by T together with an axiom scheme stating \mathcal{O} is a T -convex valuation ring. The theory T_{convex} is $T_{\text{convex}}^- \cup \{\exists x \notin \mathcal{O}\}$.

This notions were introduced by van den Dries and Lewenberg in [20] where they proved the following

Theorem 2.1 (van den Dries - Lewenberg, (3.10)-(3.15) in [20]). *The theory T_{convex} is complete and weakly o-minimal, moreover if T eliminates quantifiers and has a universal axiomatization (resp. is model-complete) in L , then T_{convex} eliminates quantifiers (resp. is model complete) in L_{convex} .*

Key-ingredient the fact that if $p(x)$ is a type over an o-minimal structure \mathbb{E} , it can extend in at most two ways to a type over the expanded structure $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}^-$.

Lemma 2.2 (van den Dries - Lewenberg, (3.6)-(3.7) in [20]). *If $p(x)$ is a unary type over $\mathbb{E} \models T$, $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}^-$, $x \models p$ and \mathcal{O}' is a T -convex valuation subring of $\mathbb{E}\langle x \rangle$ with $\mathcal{O}' \cap \mathbb{E} = \mathcal{O}$, then $\mathcal{O}' \in \{\mathcal{O}_x^-, \mathcal{O}_x\}$ where*

$$\mathcal{O}_x^- := \text{CH}_{\mathbb{E}\langle x \rangle}(\mathcal{O}), \quad \mathcal{O}_x := \{y \in \mathbb{E}\langle x \rangle : |y| < \mathbb{E}^{\succ \mathcal{O}}\}.$$

Remark 2.3. Of course if $\mathbb{E}\langle x \rangle$ does not contain any b with $\mathcal{O} < b < \mathbb{E}^{\succ \mathcal{O}}$, then $\mathcal{O}_x^- = \mathcal{O}_x$. If instead $\mathbb{E}\langle x \rangle$ contains such a b , by the exchange property, $\mathbb{E}\langle x \rangle = \mathbb{E}\langle b \rangle$ and $\mathcal{O}_x = \mathcal{O}_b$. For simplicity in the following we will denote $\mathcal{O}_x^- = \text{CH}_{\mathbb{E}\langle x \rangle}(\mathcal{O})$ by \mathcal{O} if there is no ambiguity.

Recall that an elementary extension $\mathbb{K} \preceq \mathbb{E}$ of models of T is said to be *tame* (notation $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$) if \mathbb{K} is definably Dedekind-complete in \mathbb{E} in the sense that for every \mathbb{E} -definable subset X of \mathbb{E} , if $X \cap \mathbb{K}$ is bounded, then it has a supremum in \mathbb{K} . It was proven by Marker and Steinhorn in [10] that $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$ if and only if for every tuple \bar{c} of elements in \mathbb{E} , $\text{tp}(\bar{c}/\mathbb{K})$ is a definable type (see also [13]).

Tame extensions of o-minimal structures are closely related to T -convex valuation rings in fact:

Fact 2.4 (van den Dries - Lewenberg, (2.12) in [20]). *If $\mathbb{E} \models T$, \mathcal{O} is a T -convex valuation ring then $\mathbb{K} \preceq \mathbb{E}$ is maximal among the elementary substructures of \mathbb{E} contained in \mathcal{O} if and only if $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$ and $\mathbb{K} + \mathfrak{o} = \mathcal{O}$.*

To every tame extension $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$, is associated a standard part map $\text{st}_{\mathbb{K}} : \text{CH}_{\mathbb{E}}(\mathbb{K}) \rightarrow \mathbb{K}$ uniquely defined by the property that for every $x \in \text{CH}_{\mathbb{E}}(\mathbb{K})$, $|\text{st}_{\mathbb{K}}(x) - x| < \mathbb{K}^{>0}$.

Theorem 2.5 (van den Dries, Sec. 1 in [19]). *If $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$, and $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$ with $\text{CH}(\mathbb{K}) = \mathcal{O}$, then $\text{st}_{\mathbb{K}} : \mathcal{O} \rightarrow \mathbb{K}$ induces an isomorphism between the induced structure on the imaginary sort \mathcal{O}/\mathfrak{o} and \mathbb{K} .*

In the following we will denote by $\mathbf{v}_{\mathcal{O}}$ the valuation associated with the valuation ring \mathcal{O} , and by $\mathbf{r}(\mathbb{E}, \mathcal{O})$ the residue field sort of a valued field $(\mathbb{E}, \mathcal{O})$.

Definition 2.6. Let T be the theory of an o-minimal field. For $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$, we will call

- *residue T -section* a $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$, such that $\mathcal{O} = \text{CH}(\mathbb{K})$;
- *\mathbb{K} -monomial group* a subgroup $\mathfrak{M} \subseteq (\mathbb{E}^{>0}, \cdot)$ stable under the action of $\text{Exponents}(\mathbb{E}) \cap \mathbb{K}$ and such that $\mathbf{v}_{\mathcal{O}}| : \mathfrak{M} \rightarrow \mathbf{v}_{\mathcal{O}}(\mathbb{E})$ is an isomorphism.

Remark 2.7. If T is power-bounded, then each exponent is \emptyset -definable, so $\text{Exponents}(\mathbb{E}) = \text{Exponents}(T)$ is the field of exponents of the theory and being a \mathbb{K} -monomial group does not depend on the T -residue section \mathbb{K} , so we will just say *monomial group*.

If T is exponential, then $\text{Exponents}(\mathbb{E}) = \mathbb{E}$ and a \mathbb{K} -monomial group is required to be closed under $\mathfrak{m} \mapsto \exp(k \log \mathfrak{m})$ for each $k \in \mathbb{K}$.

2.2. Wim-constructible extensions and λ -spherical completions. By an *embedding* of models of T_{convex}^- , $\iota : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}_1, \mathcal{O}_1)$ we will mean a T -elementary embedding $\iota : \mathbb{E} \rightarrow \mathbb{E}_1$ such that $\iota^{-1}(\mathcal{O}_1) = \mathcal{O}$.

Definition 2.8. Let $(\mathbb{U}, \mathcal{O}') \models T_{\text{convex}}^-$ and $\mathbb{E} \prec \mathbb{U}$, $\mathcal{O} = \mathbb{E} \cap \mathcal{O}'$ (so $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}^-$). We say that $x \in \mathbb{U} \setminus \mathbb{E}$ is *weakly immediate (wim)* (over \mathbb{E}) if its cut is an intersection of valuation balls and that it is *weakly immediately generated (wimg)* if $\mathbb{E}\langle x \rangle \setminus \mathbb{E}$ contains a weakly immediate element.

We say that x is *residual* if $(\mathbb{E}\langle x \rangle, \mathcal{O}' \cap \mathbb{E}\langle x \rangle)$ has a strictly larger residue field than $(\mathbb{E}, \mathcal{O})$.

We say that x is *purely valutional* if for every $y \in \mathbb{E}\langle x \rangle \setminus \mathbb{E}$ there is $c \in \mathbb{E}$ such that $\mathbf{v}_{\mathcal{O}'}(y - c) \notin \mathbf{v}_{\mathcal{O}'}(\mathbb{E})$.

A *principal* extension of models of T_{convex}^- is an embedding $\iota : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}_1, \mathcal{O}_1)$ of models of T -convex such that $\mathbb{E}_1 = (\iota\mathbb{E})\langle x \rangle_T$.

The principal extension ι will be said to be *wimg*, *residual* or *purely valutional* if x is respectively wimg, residual or purely valutional.

Remark 2.9. By [5, Thm. A] if x wimg it is not residual, hence every x is either wimg, or residual or purely valutional.

Remark 2.10. If $\mathbb{K} \models T$, then $(\mathbb{K}, \mathbb{K}) \models T_{\text{convex}}^-$ has no wimg principal extensions.

Remark 2.11. It follows from the residue-valuation property of power-bounded theories ([23, Sec.s 9 and 10], [18, Ch. 12 and 13], or see [5, Sec. 4]) that if T is power-bounded with field of exponents Λ , then

- (1) for every weakly immediate x , $(\mathbb{E}\langle x \rangle, \mathcal{O}' \cap \mathbb{E}\langle x \rangle)$ is an immediate extension of $(\mathbb{E}, \mathcal{O})$;
- (2) for every purely valutional x , there is $c \in \mathbb{E}$ such that $\mathbf{v}_{\mathcal{O}'}(\mathbb{E}\langle x \rangle) = \mathbf{v}_{\mathcal{O}'}(\mathbb{E}) + \Lambda \mathbf{v}_{\mathcal{O}'}(x - c)$.

In particular for T a power-bounded theory, principal extensions of models of T_{convex}^- are purely valutional if and only if they expand the value group (so in that context sometimes we will just say *valuational* instead of purely valutional).

Definition 2.12. Let $(\mathbb{E}, \mathcal{O}) \preceq (\mathbb{U}, \mathcal{O}') \models T_{\text{convex}}$ and let $x \in \mathbb{U}$ weakly immediate over $(\mathbb{E}, \mathcal{O})$. The cofinality of x is the cofinality of the set $\mathbb{E}^{<x}$ or equivalently of $-\mathbb{E}^{>x}$. We say that a weakly immediate x is λ -*bounded* if its cofinality is strictly smaller than λ .

Definition 2.13. Let $(\mathbb{E}, \mathcal{O}) \preceq (\mathbb{U}, \mathcal{O}') \models T_{\text{convex}}$. We call λ -*bounded wim-construction* a sequence $(x_i : i < \mu)$ of elements of some \mathbb{U} indexed by some ordinal μ , such that for all $j < \mu$, x_j is λ -bounded weakly immediate over $(\mathbb{E}_j := \mathbb{E}\langle x_i : i < j \rangle, \mathcal{O}' \cap \mathbb{E}_j)$. An extension $(\mathbb{E}^*, \mathcal{O}^*) \succeq (\mathbb{E}, \mathcal{O})$ is said to be λ -bounded wim-constructible if it generated by a λ -bounded wim-construction.

A *wim-construction* is a λ -bounded wim-construction for some λ . A *wim-constructible extension* is a λ -bounded wim-constructible extension for some λ .

Remark 2.14. If T is power-bounded, the wim-constructible extensions are precisely the immediate extensions.

The last section will heavily rely on the following

Theorem 2.15 (Thm. B in [5]). *Let $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$, and λ be an uncountable cardinal. There is a unique-up-to-non-unique-isomorphism λ -spherically complete λ -bounded wim-constructible extension $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ and it elementarily embeds in every λ -spherically complete extension of $(\mathbb{E}, \mathcal{O})$.*

Definition 2.16. We call such $(\mathbb{E}_\lambda, \mathcal{O}_\lambda) \succ (\mathbb{E}, \mathcal{O})$ the T - λ -spherical completion of $(\mathbb{E}, \mathcal{O})$.

Remark 2.17. Notice that a λ -bounded wim-constructible extension of $(\mathbb{E}, \mathcal{O})$ embeds in every λ -spherically complete extension of $(\mathbb{E}, \mathcal{O})$. In particular the T - λ -spherical completion is universal (i.e. weakly terminal) among the λ -bounded wim-constructible extensions.

3. TRUNCATION CLOSED EXPANSIONS OF HAHN FIELDS

3.1. Well-ordered subsets of ordered Abelian groups. We collect here some known results about well ordered subsets of ordered Abelian groups that are relevant for the rest of the paper;

Definition 3.1. Let $(\Gamma, <)$ be an ordered set. We call a partition \mathcal{P} of Γ a *segmentation* if every $P \in \mathcal{P}$ is order-convex. Given two partitions \mathcal{P} of Γ and \mathcal{Q} of Δ , we denote by $\mathcal{P} \otimes \mathcal{Q}$ the partition $\{P \times Q : P \in \mathcal{P}, Q \in \mathcal{Q}\}$ of $\Gamma \times \Delta$.

If $(\Gamma, <, +)$ is an ordered Abelian group, R is a binary relation on Γ , $X \subseteq \Gamma^n$ and $A \subseteq \Gamma$, we denote by X_{RA} the set

$$X_{RA} := \{x = (x_i)_{i < n} \in X : \forall a \in A, \Sigma(x)Ra\}, \quad \text{where } \Sigma(x) = \sum_{i < n} x_i.$$

If $a \in \Gamma$, we denote also by X_{Ra} the set $X_{R\{a\}}$.

The following Lemma is essentially a restatement of [11, Lem. 3.3].

Lemma 3.2 (Basic Segmentation Lemma). *Let $(\Gamma, +, <)$ be an ordered Abelian group and $S, T \subseteq \Gamma^{>0}$ be well ordered subsets and $U \subseteq \Gamma$ an upper subset of Γ , then there are finite segmentations \mathcal{S} of S and \mathcal{T} of T such that $\mathcal{S} \otimes \mathcal{T}$ refines the partition generated by $(S \times T)_{<U}$.*

Proof. If $\sigma(T \times S) \subseteq U$ the statement is trivial as $(S \times T)_{<U} = \emptyset$. If instead $\sigma(T \times S) \not\subseteq U$ then set $\delta_0 = \min\{t \in T : \exists s \in S, s + t < U\}$ and $\gamma_0 := \min\{s \in S : s + \delta_0 \in U\}$, so that $(S_{<\gamma_0} \times T)_{<U} = S_{<\gamma_0} \times T_{<\delta_0}$.

Now repeat the same argument for $S_0^+ := S_{\geq\gamma_0}$ and T and notice that either $\sigma(S_{\geq\gamma_0} \times T) \subseteq U$ or $\delta_1 := \min\{t \in T : \exists s \in S_0^+, s + t \in U\} < \delta_0$ by definition of γ_0 and S_0^+ .

The conclusion follows by induction and from the fact that T is well ordered. \square

Lemma 3.3 (Neumann's Lemma, from Thm. 1 and 2 in [12]). *Let $(\Gamma, +, <)$ be an ordered Abelian group and $S \subseteq \Gamma^{>0}$ be a well ordered subset, then $\Sigma : \bigcup_{k \in \mathbb{N}} S^k \rightarrow \Gamma$ has well ordered image and finite fibers.*

Proof. It suffices to show that there are no $k : \mathbb{N} \rightarrow \mathbb{N}$ and $(\bar{\gamma}(n) \in S^{k(n)} : n \in \mathbb{N})$ such that k is increasing and $(\Sigma \bar{\gamma}(n) : n \in \mathbb{N}) \in \Gamma^{\mathbb{N}}$ is weakly decreasing.

Suppose toward contradiction that $\bar{\gamma}$ is such a sequence and denote by $\mathbf{v} : \Gamma \rightarrow \mathbf{v}\Gamma$ the natural valuation on the ordered group Γ . Notice that

$\mathbf{v}\Sigma \bigcup_{k \in \mathbb{N}} S^k = \mathbf{v}S$, so in particular it is reverse-well ordered, therefore since $\mathbf{v}\Sigma \bar{\gamma}(n)$ is weakly increasing and $\mathbf{v}S$ is reverse well ordered it must eventually stabilize: so for some m , for all $n \geq m$, $\mathbf{v}\Sigma \bar{\gamma}(n) = \mathbf{v}\Sigma \bar{\gamma}(m) = v$. We can assume that the sequence $(\bar{\gamma}(n))_{n \in \mathbb{N}}$ was chosen so to get a maximal such v and so that each $\bar{\gamma}(n)$ is increasing qua finite sequence.

For each n , let $i(n) \leq k(n)$ be such that for each $j < i(n)$, $\mathbf{v}(\gamma_j(n)) > v$. Let $\mu = \min\{\gamma_j(n) : n \in \mathbb{N}, i(n) \leq j < k(n)\}$.

For some $N \in \mathbb{N}$, $N\mu > \Sigma(\bar{\gamma}(n))$ for all $n \geq m$, so for each $n \geq m$ there are at most N many js with $\mathbf{v}\gamma_j(n) = v$ so $i(n) \geq k(n) - N$. Up to possibly increasing m , we can also assume that $k(n) \geq N$ for all $n \geq m$.

The sequence $k'(n) := k(n + m)$ with $\gamma'(n) = \gamma(n + m)_{<k'(n)}$ contradicts the maximality of v . \square

The following Lemma is a curiosity and won't be used, so the reader may skip it. It somewhat represents a formal counterpart of Lemma 3.29.

Lemma 3.4 (Segmentation Lemma). *Let $(\Gamma, +, <)$ be an ordered Abelian group and $T \subseteq \Gamma^{\geq 0}$, $S \subseteq \Gamma^{> 0}$ well ordered subsets. Then for every $\gamma \in \Gamma$ and $n \in \mathbb{N}$, there are finite segmentations \mathcal{S} of S and \mathcal{T} of T such that, such that for all $m \in \mathbb{N}$, the partition $\mathcal{T} \otimes \mathcal{S}^{\otimes m}$ of $T \times (S^{\times m})$ refines the partition generated by $(T \times (S^{\times m}))_{<\gamma}$.*

Proof. Let $\mathbf{v} : \Gamma^{> 0} \rightarrow \mathbf{v}\Gamma$ be the natural valuation on Γ and $\prec, \preceq, \sim, \asymp$ the corresponding dominance, strict dominance and asymptotic equivalences. We prove the statement by induction on the pair of order types of S and T .

If the order type of S is 0, then $S = \emptyset$ and by Lemma 3.2, the statement holds for S , and for every T .

Fix $\gamma \in \Gamma$. Segment S as $S := S_{\succeq \gamma} \sqcup S_{\prec \gamma}$. Notice that there is $m_0 \in \mathbb{N}$ such that for all $m > m_0$, $((S_{\succeq \gamma})^{\times m})_{>[\gamma] \sim} = (S_{\succeq \gamma})^{\times m}$.

By Lemma 3.2, there is a finite segmentation \mathcal{S}_0 of $S_{\succeq \gamma}$ and a finite segmentation \mathcal{T}_0 of T such that for all $m \leq m_0$, $\mathcal{T}_0 \otimes \mathcal{S}_0^{\otimes m}$ refines the partition generated by $(T \times S_{\succeq \gamma}^{\times m})_{>[\gamma] \sim}$ and $(T \times S_{\succeq \gamma}^{\times m})_{<[\gamma] \sim}$.

This implies that $\mathcal{S}_1 := \mathcal{S}_0 \cup \{S_{\prec \gamma}\}$ is such that for all m , $\mathcal{T}_0 \otimes \mathcal{S}_1^{\otimes m}$ refines the partition generated by $(T \times S^{\times m})_{>[\gamma] \sim}$ and $(T \times S^{\times m})_{<[\gamma] \sim}$.

Now distinguish two cases

Case 1: $S_{\prec \gamma} \neq S$. Notice that for each $\bar{R} := (R_i)_{i < m} \in \mathcal{S}_0^{\times m}$ and $U \in \mathcal{T}_0$, $U \times \prod \bar{R} \subseteq (T \times S^{\times m})_{\sim \gamma}$ if and only if $\min(U) + \sum_{i < m} \min(R_i) \sim \gamma$.

Notice that the set $\mathcal{M} := \{(U, \bar{R}) : m \in \mathbb{N}, U \in \mathcal{T}_0, \bar{R} \in \mathcal{S}_0^m, U \times \prod \bar{R} \subseteq (T \times S^{\times m})_{\sim \gamma}\}$ is finite.

Given $(U, \bar{R}) \in \mathcal{M}$ we can apply the inductive hypothesis with $T' = \Sigma(U \times \prod \bar{R})$, $S' = S_{\prec \gamma}$, because the order type of S' is strictly smaller than the order type of S . This yields a finite segmentation \mathcal{T}' of T' and a finite segmentation \mathcal{S}' of S' such that for all $n \in \mathbb{N}$, $\mathcal{T}' \otimes \mathcal{S}'^{\otimes n}$ refines the partition generated by $(T' \times S')_{<\gamma}$. By Lemma 3.2 this implies there is a finite segmentation \mathcal{T}'' of $T'' = U \cup \bigcup \bar{R}$ such that $\{\Sigma(\bar{X}) : \bar{X} \in \mathcal{T}''^{\otimes m}\}$ refines \mathcal{T}' .

Thus refining \mathcal{T}_0 to \mathcal{T} so that $\mathcal{T}|_U$ refines $\mathcal{T}''|_U$ and \mathcal{S}_1 to \mathcal{S} so that $\mathcal{S}|_{R_i}$ refines $\mathcal{T}''|_{R_i}$ for all pairs (U, \bar{R}) , yields the required segmentations.

Case 2: $S_{\prec\gamma} = S$. For each $U \in \mathcal{T}_0$ with $U \subseteq [\gamma]_{\sim}$, either $\Sigma(U \times S^m) < \gamma$ for every n , or $T' = U - \min U$ and $S' = S$ are in the hypothesis of case 1 for $\gamma' = \gamma - \min U$. In any case by the inductive hypothesis we get a finite segmentation \mathcal{T}_U of U and \mathcal{S}_U of S such that for every $m \in \mathbb{N}$, $\mathcal{T}_U \otimes \mathcal{S}_U^{\otimes m}$ refines the partition generated by $(U \times S^{\times m})_{<\gamma}$. Refining \mathcal{T}_0 and \mathcal{S}_1 to \mathcal{T} and \mathcal{S} respectively so that for all $U \in \mathcal{T}_0$ with $U \subseteq [\gamma]_{\sim}$, \mathcal{S} refines \mathcal{S}_U and $\mathcal{T}|_U$ refines \mathcal{T}_U yields again the required finite segmentations. \square

3.2. Rings of generalized series. Let $(\mathfrak{M}, \cdot, 1, \leq)$ be a (possibly partially) ordered cancellative monoid. The ring of generalized series $\mathbb{K}((\mathfrak{M}))$ with coefficients from a ring \mathbb{K} and monomials from \mathfrak{M} is defined as the \mathbb{K} -module of \mathfrak{M} -tuples $(f_{\mathfrak{m}})_{\mathfrak{m} \in \mathfrak{M}} \in \mathbb{K}^{\mathfrak{M}}$ whose support $\text{Supp } f := \{\mathfrak{m} : f_{\mathfrak{m}} \neq 0\}$ is a well-partial order for the induced order by the reverse order \leq^{op} on \mathfrak{M} (that is, $\text{Supp } f$ does not contain infinite increasing sequences nor infinite antichains for $<$).

A family $(f_i)_{i \in I} \in \mathbb{K}((\mathfrak{M}))$ is said to be *summable* if for each $\mathfrak{m} \in \mathfrak{M}$, $\{i : (f_i)_{\mathfrak{m}} \neq 0\}$ is finite and $\bigcup_{i \in I} \text{Supp } f_i$ is still a well-partial order for \leq^{op} . If $(f_i)_{i \in I}$ is summable its sum is defined as the function

$$\left(\sum_{i \in I} f_i \right)_{\mathfrak{m}} := \sum_{i \in I} (f_i)_{\mathfrak{m}}.$$

Elements of \mathfrak{M} are regarded as elements of $\mathbb{K}((\mathfrak{M}))$ by identifying \mathfrak{m} with the function that is 1 at \mathfrak{m} and 0 at $\mathfrak{n} \neq \mathfrak{m}$. In that sense every element $f \in \mathbb{K}((\mathfrak{M}))$ can be regarded as the sum of the summable family $(f_{\mathfrak{m}} \mathfrak{m})_{\mathfrak{m} \in \mathfrak{M}}$, so

$$f = \sum_{\mathfrak{m} \in \mathfrak{M}} \mathfrak{m} f_{\mathfrak{m}}.$$

The product \cdot on $\mathbb{K}((\mathfrak{M}))$ is defined as the only \mathbb{K} -bilinear extension of the product on \mathfrak{M} which is strongly bilinear in the sense that for every pair $(g_i)_{i \in I}$ and $(f_j)_{j \in J}$ of summable families in $\mathbb{K}((\mathfrak{M}))$,

$$\left(\sum_{i \in I} f_i \right) \cdot \left(\sum_{j \in J} f_j \right) = \sum_{(i,j) \in I \times J} f_i \cdot f_j.$$

It is not hard to verify that defining the product as

$$\left(\left(\sum_{\mathfrak{m} \in \mathfrak{M}} a_{\mathfrak{m}} \mathfrak{m} \right) \cdot \left(\sum_{\mathfrak{n} \in \mathfrak{M}} b_{\mathfrak{n}} \mathfrak{n} \right) \right)_{\mathfrak{p}} := \sum_{\mathfrak{m}\mathfrak{n}=\mathfrak{p}} a_{\mathfrak{m}} b_{\mathfrak{n}}$$

yields the required property.

Definition 3.5 (Truncations and Segments). If S is a segment of \mathfrak{M} , and $f \in \mathbb{K}((\mathfrak{M}))$ we will call S -segment of f the series $f|_S = \sum_{\mathfrak{m} \in S} f_{\mathfrak{m}}$. If $\mathfrak{m} \in \mathfrak{M}$, the \mathfrak{m} -truncation of f , denoted by $f|\mathfrak{m}$ is the S -segment of f with

$S = \{\mathbf{n} : \mathbf{n} > \mathbf{m}\}$. A subset $X \subseteq \mathbb{K}(\mathfrak{M})$ will be said to be closed under truncations if for each $f \in X$ and each $\mathbf{m} \in \mathfrak{M}$, $f|\mathbf{m} \in X$, it will be said to be closed under segments if for every $f \in X$ and every segment S of \mathfrak{M} , $f|S \in X$.

Remark 3.6. If \mathfrak{M} is totally ordered and X is subgroup of $\mathbb{K}(\mathfrak{M})$, then X is closed under truncations if and only if it is closed under segments.

If $\mathfrak{N}_0, \mathfrak{N}_1, \dots, \mathfrak{N}_{n-1}$ are totally ordered Abelian groups and $\mathfrak{M} = \mathfrak{N}_0 \times \dots \times \mathfrak{N}_{n-1}$ is endowed with the partial order, every subgroup $X \subseteq \mathbb{K}(\mathfrak{M})$ is closed under segments if and only if it is closed under *partial truncations*, that is under taking segments $f|S$ for S of the form $S = \{(\mathbf{n}_0, \dots, \mathbf{n}_{n-1}) : \mathbf{n}_i > \mathbf{m}_i\}$.

3.3. Generalized power series. We fix for the rest of the paper an infinite set of formal variables Var . We will denote variables and sets of variables by x, y, z, \dots . If x is a set of variables, we will denote by S^x the set of assignments from S for x , i.e. the set of functions from x to S .

If x, y are disjoint sets of variables, I will denote by a slight abuse of notation (x, y) the set of variables $x \cup y$.

If x, y are disjoint sets of variables, and $a \in S^x, b \in S^y$ are assignments, I will denote by $(a, b) \in S^{(x, y)}$ the assignment which is a on x and b on y .

Definition 3.7. Let \mathbb{K} be a ring and $x = \{x_i : i < n\}$ a set of n distinct formal variables and Λ an ordered commutative ring. A *monomial* in x with exponents from Λ is an expression of the form x^γ for $\gamma \in \Lambda^x$. The set of monomials in x with exponents from Γ will be denoted by x^Λ and forms a partially ordered group $(x^\Lambda, \cdot, 1, \leq)$ with

$$x^\alpha x^\beta = x^{\alpha+\beta}, \quad x^\gamma \geq x^\beta \iff \gamma \leq \beta,$$

where Λ^x is given the product (partial) order. The intuition for reverting the order here is that positive infinitesimals will be assigned to the components of x .

Throughout the rest of the paper we will assume that \mathbb{K} is an ordered field with powers in Λ , that is that $(\mathbb{K}^{>0}, \cdot)$ has an ordered Λ -module structure written exponentially, that is $(\lambda, k) \mapsto k^\lambda$.

The ring $\mathbb{K}((x^\Lambda))$ is called *ring of generalized power series in x with coefficients from \mathbb{K} and exponents from Λ* . A *generalized power series* is an element of $\mathbb{K}((x^\Lambda))$, so it can be expressed as a formal sum

$$f = \sum_{\gamma \in \Lambda^n} c_\gamma x^\gamma$$

where $\{\gamma \in \Lambda^n : c_\gamma \neq 0\}$ is a *well-partial order* (that is, it has no infinite antichains and no infinite descending chains).

Such a series will be said to be

- *non-singular* if $c_\gamma = 0$ for all $\gamma \not\geq 0$ (i.e. if it lies in $\mathbb{K}(((x^\Lambda)^{\leq 1}))$)
- *normal* if $f = x^\alpha(k + h)$ with $k \in \mathbb{K}^{\neq 0}$ and $\text{Supp } h \subseteq \{x^\gamma : \gamma > 0\}$.
- *p-composable* if $f = x^\alpha(k + h)$ is normal with $\alpha > 0$ and $k > 0$.

Remark 3.8. If $x \subseteq y$, then every monomial x^γ is identified with the monomial in y given by y^β where $\beta \in \Lambda^y$ is given by $\beta(z) = \gamma(z)$ if for some $z \in y$ and $\beta(z) = 0$ otherwise. This induces natural inclusions

$$x^\Lambda \subseteq y^\Lambda, \quad \mathbb{K}((x^\Lambda)) \subseteq \mathbb{K}((y^\Lambda)).$$

Similarly if $\sigma : x \rightarrow y$ is an injection, it induces natural inclusions

$$\sigma : x^\Lambda \rightarrow y^\Lambda, \quad \sigma : \mathbb{K}((x^\Lambda)) \rightarrow \mathbb{K}((y^\Lambda)).$$

It is convenient to extend this reindexing to non-injective $\sigma : x \rightarrow y$, to take into account the operation of forming for example $f(x, x)$ out of $f(x, y)$. This is entirely possible because $f(x, x)$ will still be summable.

Definition 3.9. If $f \in \mathbb{K}((x, y)^\Lambda)$ is a generalized power series, the variables y are said to be *classical* in f if they appear only with integer exponents in f . If x is a single variable and y is a set of variables, and $f \in \mathbb{K}((x, y)^\Lambda)$, then $\partial_x f$ denotes the formal derivative of f in x : if $f = \sum_{\alpha, \beta} c_{\alpha, \beta} x^\alpha y^\beta$, then

$$\partial_x f := \sum_{\alpha, \beta} c_{\alpha, \beta} \alpha x^{\alpha-1} y^\beta.$$

A *family* of generalized power series \mathcal{F} is the datum for every $x \in \text{Var}_*$ of a subset $\mathcal{F}(x) \subseteq \mathbb{K}((x^\Lambda))$. A family \mathcal{F} is said to be *truncation-closed* if each $\mathcal{F}(x)$ is closed under *partial* truncations (see Remark 3.6).

We say that a family \mathcal{F} is a *language* of generalized power series if for any (non-necessarily injective) reindexing function $\sigma : x \rightarrow y$, $\sigma \mathcal{F}(x) \subseteq \mathcal{F}(y)$.

A language \mathcal{F} is said to be an *algebra* if $\mathcal{F}(x)$ is a \mathbb{K} -algebra for every x .

Remark 3.10. Notice that a family \mathcal{F} such that $\mathcal{F}(x)$ is a \mathbb{K} -algebra for every x , is an algebra for the above definition only if it also a language.

Lemma 3.11. *If a family \mathcal{F} is truncation-closed, then the algebra it generates is truncation-closed.*

Proof. Notice that by Lemma 3.2, the \mathbb{K} -algebra generated by each $\mathcal{F}(x)$ is closed under truncation provided that $\mathcal{F}(x)$ was. Thus it suffices to show that if \mathcal{F} is such that each $\mathcal{F}(x)$ is truncation-closed algebra then the language generated by \mathcal{F} is truncation-closed. To this end, it suffices to show that if $f \in \mathcal{F}(x, y, z)$, $x, y \in \text{Var}$, $z \in \text{Var}_N$ for some N , then for every $\alpha \in \Lambda$, there are $f_0, \dots, f_{n-1}, g_0, \dots, g_{n-1} \in \mathcal{F}(x, y, z)$, such that

$$f|\{x^\beta y^\gamma z^\delta : \gamma + \beta < \alpha\} = \sum_{i < n} f_i \cdot g_i$$

This again follows from the basic segmentation lemma (3.2). \square

Definition 3.12. For \mathfrak{M} a multiplicatively written totally ordered Λ -vector space, a generalized power series f in some set of variables x can be interpreted as a function $f : (\mathbb{K}((\mathfrak{M}^{<1}))^{>0})^x \rightarrow \mathbb{K}((\mathfrak{M}))$ by setting for $k \in (\mathbb{K}^{>0})^\Lambda$,

$\mathbf{m} \in (\mathfrak{M}^{<1})^x$ and $\varepsilon \in \mathbb{K}((\mathfrak{M}^{<1}))^x$

$$f(\mathbf{m}(k + \varepsilon)) := \sum_{m \in \mathbb{N}^n} \sum_{\gamma} \binom{\gamma}{m} c_{\gamma} k^{\gamma-m} \mathbf{m}^{\gamma} \varepsilon^m$$

where

$$\binom{\gamma}{m} := \prod_{i=1}^n \frac{1}{m_i!} \prod_{j=0}^{m_i-1} (\gamma_i - j) = \prod_{i=1}^n \binom{\gamma_i}{m_i}.$$

The family $(\mathbf{m}^{\gamma} \varepsilon^m)_{\gamma \in \text{Supp } f, m \in \mathbb{N}^n}$ is summable: in fact $(\varepsilon^m)_{m \in \mathbb{N}^n}$ is summable by Neumann's Lemma (3.3) and $(\mathbf{m}^{\gamma})_{\gamma \in \text{Supp } f}$ is summable by hypothesis.

Given $X \subseteq \mathbb{K}((\mathfrak{M}))$, we will denote by $\langle X \rangle_{\mathcal{F}}$, the $\mathcal{F} \cup \{+, \cdot\}$ -substructure of $\mathbb{K}((\mathfrak{M}))$ generated by X where the symbols in \mathcal{F} are interpreted as above.

If f is a series in (x, y) , $x \in \text{Var}$, $y \in \text{Var}_*$, $g(z) = z^{\gamma}(k + h(z))$ is a p -composable series in z , and $(xg(y), y) \in \mathbb{K}(((x, z, y)^{\Lambda}))^{(x, y)}$ is the assignment $x \mapsto xg(x, y)$, $y \mapsto y$ it is possible to define $f(xg(z), y)$ as

$$f(xg(z), y) = h := \sum_{\alpha, \beta, m} c_{\alpha, \beta} k^{\alpha-m} \binom{\alpha}{m} z^{\alpha\gamma} h(z)^m y^{\beta}.$$

Remark 3.13. The composition has the property that for each multiplicatively written totally ordered Λ -vector space \mathfrak{M} and each $x \in \mathbb{K}((\mathfrak{M}^{<1}))^{>0}$, $z \in (\mathbb{K}((\mathfrak{M}^{<1}))^{>0})^z$, $h(x, z, y) = f(xg(z), y)$.

Remark 3.14. If $f = \sum_{\alpha, \beta} c_{\alpha, \beta} x^{\alpha} y^{\beta} \in \mathbb{K}(((x, y)^{\Lambda}))$ and y are classical variables in f , then we can interpret them naively: in fact for $x = \mathbf{m}(k + \varepsilon) \in (\mathbb{K}((\mathfrak{M}^{<1}))^{>0})^x$ and every $y \in (\mathbb{K}((\mathfrak{M}^{<1}))^y)$

$$f(x, y) = \sum_{m, \beta} \sum_{\alpha} \binom{\alpha}{m} c_{\alpha, \beta} k^{\gamma-m} \mathbf{m}^{\gamma} \varepsilon^m y^{\beta}.$$

Definition 3.15. Let $\mathcal{F} := (\mathcal{F}(x))_{x \in \text{Var}_*}$ be an algebra of generalized power series with coefficients from \mathbb{K} and exponents from Λ . Consider the following closure properties:

- (1) \mathcal{F} is closed under partial truncations and contains the coordinate projections;
- (2) \mathcal{F} is closed under renormalized formal derivatives (i.e. for every $f \in \mathcal{F}(x, y)$, x is a single variable and y is a set of variables, the series $f \mapsto x \partial_x(f)$ is in $\mathcal{F}(x, y)$);
- (3) for $x \in \text{Var}$, $y \in \text{Var}^n$ and $k \in \mathbb{K}^{>0}$, if $f(x, y) = \sum_{\alpha, \beta} c_{\alpha, \beta} x^{\alpha} y^{\beta} \in \mathcal{F}(x, y)$

$$f(z_0(z_1 + k), y) := \sum_{\alpha, \beta, m} c_{\alpha, \beta} k^{\alpha-m} \binom{\alpha}{m} z_0^{\alpha} z_1^m y^{\beta} \in \mathcal{F}(z_0, z_1, y)$$

$$f(z_0 z_1, y) := \sum_{\alpha, \beta, m} c_{\alpha, \beta} z_0^{\alpha} z_1^{\alpha} y^{\beta} \in \mathcal{F}(z_0, z_1, y);$$

- (4) \mathcal{F} is closed under monomial division.

We will say that \mathcal{F} is *almost fine* if it satisfies (1) and (2). And we will say that it is *fine* if it satisfies (1), (2) and (3).

Remark 3.16. Points (1) and (3) together entail that $x\partial_x f \in \mathcal{F}(x, y)$. In particular an almost fine algebra is fine. If x is classical in f , points (3), (4) and (1) together entail that $\partial_x f \in \mathcal{F}(x, y)$.

Remark 3.17. If \mathcal{F} is closed under the operation of setting variables $= 0$ (so in particular if it is closed under partial truncations), point (3) can be restated by saying that \mathcal{F} is closed under right-composition with the set of p -composable polynomials $\mathbb{K}[x]_{\mathbf{p}}$ with positive coefficients. In particular, given any such algebra \mathcal{F} the smallest algebra containing \mathcal{F} and satisfying (3) is and is given by

$$\mathcal{F}_b(z) := \{f(p(z)) : p(z) \in \mathbb{K}[z]_{\mathbf{p}}^x, f(x) \in \mathcal{F}(x)\}.$$

Remark 3.18. If \mathcal{F} satisfies any (i) ($i < 5$), then the smallest family containing \mathcal{F} and closed under monomial division is still an algebra satisfying (i). This is clear for (1) and (4).

As for (2) observe that if x is a variable and $\partial_x f = x^\alpha g$, then $x\partial_x f = \alpha f + x^{\alpha+1}\partial_x g$ so $x^\alpha \partial_x g \in \mathcal{F}$.

As for (3) notice that $y^\beta | f(z_0(z_0 + k), y)$ if and only if $y^\beta | f(x, y)$ and that $z_0^\alpha | f(z_0(z_0 + k), y)$ if and only if $x^\alpha | f(x, y)$.

Lemma 3.19. *Suppose \mathcal{F} is an almost fine algebra. Then \mathcal{F}_b is a fine algebra.*

Proof. To show \mathcal{F}_b is fine, by Remark 3.16 we only need to show it is closed under truncations. For this in trun it suffices to show that for each $p(z) \in \mathbb{K}[z]_{\mathbf{p}}^x$, $\mathcal{F}(p(z)) = \{f(p(z)) : f \in \mathcal{F}(x)\}$ is a truncation-closed \mathbb{K} -algebra as then the algebra closure of the family $\bigcup_{z,p} \mathcal{F}(p(z))$ will be truncation closed by Lemma 3.11. In fact for the same reason it is enough to restrict to the tuples of polynomials that are compositions of polynomials of the form $z_0(k + z_1)$ with $k \in \mathbb{K}^{\geq 0}$. Thus the statement is reduced to proving that if $\mathcal{F}(x, y)$ is a subalgebra of $\mathbb{K}((x, y)^\Lambda)$ closed under truncations (with $x \in \text{Var}$ and $y \in \text{Var}_N$), then so is $\mathcal{F}(z_0(z_1 + k), y)$ for $k \geq 0$.

To this end notice that if $k > 0$, then

$$\begin{aligned} f(z_0(z_1 + k), y)|z_0^\alpha &= h(z_0(z_1 + k), y), & h(x, y) &= f(x, y)|x^\alpha \\ f(z_0(z_1 + k), y)|z_1^n &= \sum_{m < n} z_1^m k^{-m} h_m(z_0, y), & h_m(x, y) &= (x\partial_x)^m f(x, y) \end{aligned}$$

If instead $k = 0$, the statement is trivial. \square

In the next section we shall prove

Theorem 3.20. *Let \mathcal{F} be an almost fine algebra of generalized power series. Then for every $X \subseteq \mathbb{K}((\mathfrak{M}))$ closed under truncations, $\langle X \cup \mathfrak{M} \rangle_{\mathcal{F}}$ is closed under truncations.*

Remark 3.21. Notice that if \mathcal{F} contains the univariad series $\sum_{n \in \mathbb{N}} x^n$, then $\langle X \cup \mathfrak{M} \rangle_{\mathcal{F}}$ is a subfield.

3.4. Infinitesimally convergent power series. The main idea to prove Theorem 3.20 is that we can reduce to the study of *restricted* power series with coefficients in $\mathbb{K}(\mathfrak{M})$. The key step to accomplish this is Lemma 3.29.

Definition 3.22. Let $\mathbf{x} = \{x_i : i < n\}$ be a set of n distinct variables. An element $f \in \mathbb{K}(\mathfrak{M} \times \mathbf{x}^{\mathbb{N}})$ where $\mathfrak{M} \times \mathbf{x}^{\mathbb{N}}$ is given the product partial order, will be called an *infinitesimally convergent* or *restricted* power-series with coefficients in $\mathbb{K}(\mathfrak{M})$. It can be regarded as a power-series $f = \sum_{m \in \mathbb{N}^n} r_m \mathbf{x}^m \in (\mathbb{K}(\mathfrak{M}))[[\mathbf{x}]]$ with the property that

$$\text{Supp}_{\mathfrak{M}}(f) := \bigcup \{\text{Supp } r_m : m \in \mathbb{N}^n\}$$

is well ordered. We say that f is *composable* if $\text{Supp}_{\mathfrak{M}}(f) \leq 1$ and $r_0 \in \mathbb{K}(\mathfrak{M}^{<1})$. We define a coefficient-wise truncation $f \parallel \mathbf{m}$ for $f \in \mathbb{K}(\mathfrak{M} \times \mathbf{x}^{\mathbb{N}})$ and $\mathbf{m} \in \mathfrak{M}$ by

$$(f \parallel \mathbf{m})(\mathbf{x}) := \sum_m (r_m \parallel \mathbf{m}) \mathbf{x}^m.$$

Given an n -tuple of composable power series with generalized coefficients, $(g_i(z))_{i < n} \in \mathbb{K}(\mathfrak{M}^{\leq 1} \times z^{\mathbb{N}^l})$, and said $g \in \mathbb{K}(\mathfrak{M}^{\leq 1} \times z^{\mathbb{N}^l})^{\times}$ the assignment $x_i \mapsto g_i(z)$, we can define $f(g(z))$ as

$$f(g(z)) := \sum_{m \in \mathbb{N}^n} r_m g(z)^m = \sum_{m \in \mathbb{N}^n} r_m \prod_{i < n} g_i(z)^{m_i}.$$

In fact the family $(r_m g(z)^m)_{m \in \mathbb{N}^n}$ is summable in $\mathbb{K}(\mathfrak{M} \times z^{\mathbb{N}})$.

If \mathcal{L} is a set of infinitesimally convergent Laurent series with coefficients in $\mathbb{K}(\mathfrak{M})$, then we denote by $\mathcal{L}^\circ \subseteq \mathcal{L}$ the subset of composable series in \mathcal{L} . If $\mathcal{A}(x)$ is a subring of infinitesimally convergent power series in the variables x and $\mathcal{B}(y) = \mathcal{B}^\circ(y)$ is a set of composable series, then we denote

- by $\mathcal{A}(\mathcal{B}(y))$ the set of compositions of elements in \mathcal{A} with tuples in \mathcal{B} ;
- by $\mathcal{A}[\mathcal{B}(y)]$ the ring generated by the $\mathcal{A}(\mathcal{B}(y))$.

If \mathcal{L} is a set of infinitesimal convergent Laurent series with coefficients in $\mathbb{K}(\mathfrak{M})$ we denote by \mathcal{L}^* the smallest algebra containing \mathcal{L} that is closed under composition with composable series in \mathcal{L}^* .

Remark 3.23. A composable power series with generalized coefficients in an empty set of variables is just an element $a \in \mathbb{K}(\mathfrak{M}^{<1})$. So the evaluation of f at a tuple a from $\mathbb{K}(\mathfrak{M}^{<1})$ is a particular instance of composition.

Remark 3.24. If $f = \sum_{\gamma, l} k_{\gamma, l} x^\gamma y^l$ is a generalized power series all of whose classical variables are among the y variables and $x = \mathbf{m}(a + \varepsilon)$ with $a \in \mathbb{K}^{>0}$, then $f(x, y) = f_{am}(\varepsilon, y)$ where $f_{am}(z, y)$ is the infinitesimally convergent power given by

$$f_{am}(z, y) := \sum_{m, l} r_m z^m y^l, \quad \text{with} \quad r_m := \sum_{\gamma} \binom{\gamma}{m} k_{\gamma} a^{\gamma - m} \mathbf{m}^{\gamma}.$$

Definition 3.25. Given an almost fine \mathcal{F} and a multiplicatively written Λ -vector space \mathfrak{M} , we denote by $\mathcal{F}_{\mathfrak{M}}$ the set of power series with coefficients in $\mathbb{K}(\langle\mathfrak{M}\rangle)$ given by the series of the form $y^{-m}f(\mathbf{m}, y)$ where $f \in \mathcal{F}_b(\mathbf{x}, y)$ and $y^m|f(\mathbf{m}, y)$, \mathbf{x} is a tuple containing all non-classical variables of f and $\mathbf{m} \in (\mathfrak{M}^{<1})^{\times}$.

Remark 3.26. Notice that a subring of $\mathbb{K}(\langle\mathfrak{M}\rangle)$ containing \mathbb{K} and \mathfrak{M} is closed under \mathcal{F} if and only if it is closed under the family of infinitesimally convergent power series

$$\mathfrak{M}\mathcal{F}_{\mathfrak{M}} := \{\mathfrak{n}f(y) : \mathfrak{n} \in \mathfrak{M}, f \in \mathcal{F}_{\mathfrak{M}}\}.$$

Lemma 3.27. *Let \mathcal{F} be an almost fine set of composable generalized power series. Then $\mathfrak{M}\mathcal{F}_{\mathfrak{M}}$ is closed under $\|\cdot\|$ -truncations and formal derivatives.*

Proof. The fact that $\mathfrak{M}\mathcal{F}_{\mathfrak{M}}$ is closed under formal derivatives immediately descends from the fact that \mathcal{F}_b is closed under renormalized formal derivatives.

To see $\mathcal{F}_{\mathfrak{M}}$ is closed under $\|\cdot\|$ -truncations, it is enough to apply Lemma 3.2 and the hypothesis that \mathcal{F} is closed under partial truncations. \square

Lemma 3.28. *If $\mathcal{L}(\mathbf{x}) \subseteq \mathbb{K}(\langle\mathfrak{M} \times \mathbf{x}^{\mathbb{Z}}\rangle)$ is a subring (resp. subgroup, non-unital subring), then*

$$\{f \in \mathcal{L}(\mathbf{x}) : \forall \mathfrak{m} \in \mathfrak{M}, f|\mathfrak{m} \in \mathcal{L}(\mathbf{x})\}$$

is a subring (resp. subgroup, non-unital subring).

Proof. It suffices to show that if $f(\mathbf{x}) := \sum_m r_m \mathbf{x}^m$, $g(\mathbf{x}) := \sum_l s_l \mathbf{x}^l$, are elements of $\mathcal{L}(\mathbf{x})$ such that $f|\mathfrak{m}, g|\mathfrak{m} \in \mathcal{L}(\mathbf{x})$ for all $\mathfrak{m} \in \mathfrak{M}$, then $f(\mathbf{x}) \cdot g(\mathbf{x}) \in \mathcal{L}(\mathbf{x})$. Let $R = \text{Supp}_{\mathfrak{M}}(f)$ and $S = \text{Supp}_{\mathfrak{M}}(g)$. These are both well ordered so for fixed \mathfrak{m} , by Lemma 3.2 there are finite $\mathfrak{n}_0 < \dots < \mathfrak{n}_{a-1}$ in R and $\mathfrak{p}_0 > \dots > \mathfrak{p}_a$ in S such that for each $m \in R$, $l \in S$

$$(r_m \cdot s_l)|\mathfrak{m} = \sum_{j < a} (r_m|\mathfrak{n}_j) \cdot (s_l|\mathfrak{p}_{j+1} - s_l|\mathfrak{p}_j).$$

But then it suffices to observe that

$$\begin{aligned} (f \cdot g)|\mathfrak{m} &= \sum_p \left(\sum_{m+l=p} r_m s_l \right) \mathbf{x}^p = \\ &= \sum_{j < a} \sum_p \left(\sum_{m+l=p} (r_m|\mathfrak{n}_j) (s_l|\mathfrak{p}_{j+1} - s_l|\mathfrak{p}_j) \right) \mathbf{x}^p = \\ &= \sum_{j < a} (f|\mathfrak{n}_j)(\mathbf{x}) ((g|\mathfrak{p}_{j+1})(\mathbf{x}) - (g|\mathfrak{p}_j)(\mathbf{x})) \end{aligned}$$

which concludes the proof. \square

Lemma 3.29. *Assume $\mathcal{A}(x) \subseteq \mathbb{K}((\mathfrak{M} \times x^{\mathbb{Z}}))$ is a subring closed under $\|\cdot\|$ -truncations and formal derivatives and that $\mathcal{B}(y) \subseteq \mathbb{K}((\mathfrak{M} \times y^{\mathbb{Z}}))$ is closed under $\|\cdot\|$ -truncations. Then*

$$\mathcal{A}[\mathcal{B}^\circ(y)] := \mathbb{Z}[f(g(y)) : f(x) \in \mathcal{A}(x), g(y) \in \mathcal{B}^\circ(y)^x]$$

is closed under $\|\cdot\|$ -truncations.

Proof. Let $x = \{x_i : i < n\}$ have n distinct variables and notice that we can assume without loss of generality that $x \subseteq \mathcal{A}(x)$. Let $f(x) \in \mathcal{A}(x)$ and $g := (g_{x_i} := g_i)_{i < n} \in \mathcal{B}^\circ(y)^x$ be a x -tuple from $\mathcal{B}^\circ(y)$. We are going to prove by induction on the pair of order types respectively of $S := \bigcup_{i < n} \text{Supp}_{\mathfrak{M}}(g_i)$ and $R := \text{Supp}_{\mathfrak{M}}(f)$, that $f(g(y)) \in \mathcal{A}[\mathcal{B}^\circ(y)]$.

Fix \mathfrak{m} and let $v = \mathfrak{v}\mathfrak{m}$ be the natural valuation of \mathfrak{m} , and notice that

$$\tilde{f}(x, z) := f(x + z) = \sum_{h, l} \binom{h+l}{l} r_{h+lx^l} z^h = \sum_h \frac{1}{h!} (\partial_h f)(x) z^h.$$

We distinguish two cases.

Case 1. If $\mathfrak{v}S \not\geq \mathfrak{v}\mathfrak{m} = v$, then $\bigcup_{j < n} \text{Supp}_{\mathfrak{M}}(g_j \| v)$ has strictly smaller order type than S , so by the inductive hypothesis for all $\mathfrak{n} \in \mathfrak{M}$ and all multi-indexes h , $(\partial_h f)(g \| v) \| \mathfrak{n} \in \mathcal{L}^*$. We can rewrite $f(g(y))$ as

$$f(g) = \tilde{f}(g \| v, g - (g \| v)) = \sum_h (g - g \| v)^h \frac{1}{h!} (\partial_h f)(g \| v)$$

and observe that $(\partial_h f \| v)(g \| v) = ((\partial_h f)(g)) \| v$ and that for $|h|$ large enough, $((g - g \| v)^h) \| \mathfrak{m} = 0$, so for some large enough $M \in \mathbb{N}$,

$$f(g) \| \mathfrak{m} = \sum_{|h| < M} \left((g - g \| v)^h \frac{1}{h!} (\partial_h f)(g \| v) \right) \| \mathfrak{m}$$

and we can conclude.

Case 2. If $\mathfrak{v}S > \mathfrak{v}\mathfrak{m} = v$, then $g \| v = g$. Let for each $m \in \mathbb{N}^n$,

$$\mathfrak{m}_m := \begin{cases} \max\{\mathfrak{n} \in \text{Supp}(r_m) : \mathfrak{n} \sim \mathfrak{m}, g^m \| (\mathfrak{m}/\mathfrak{n}) \neq g^m\} & \text{if it exists} \\ \sup\{\mathfrak{n} \in \text{Supp}(r_m) : \mathfrak{n} < [\mathfrak{m}]_{\sim}\} & \text{otherwise.} \end{cases}$$

Notice that \mathfrak{m}_m is given by the first option whenever $\{\mathfrak{n} \in \text{Supp}(r_m) : \mathfrak{n} \sim \mathfrak{m}, g^m \| (\mathfrak{m}/\mathfrak{n}) \neq g^m\} \neq \emptyset$, and that in the second option the sup is in fact a maximum whenever $\{\mathfrak{n} \in \text{Supp}(r_m) : \mathfrak{n} < [\mathfrak{m}]_{\sim}\} \neq \emptyset$, and it is $-\infty$ otherwise. Set $\mathfrak{p} = \max_{m \in \mathbb{N}^n} \mathfrak{m}_m$ and notice that if $\mathfrak{p} \not\sim \mathfrak{m}$, then \mathfrak{m}_m is given by the second option for every m and $((r_m | \mathfrak{p}) \cdot g^m) \| \mathfrak{m} = (r_m | \mathfrak{p}) \cdot g^m$ so $f(g) \| \mathfrak{m} = (f | \mathfrak{p})(g)$. We may therefore assume that $R \ni \mathfrak{p} \sim \mathfrak{m}$ and set $f_1 := (f - f | \mathfrak{p}) / \mathfrak{p} \in \mathcal{A}/\mathfrak{p}$, so

$$(f - f | \mathfrak{p})(g) \| \mathfrak{m} = \mathfrak{p} \cdot (f_1(g) \| (\mathfrak{m}/\mathfrak{p}))$$

By construction g is now in the hypothesis of case (1) for $\mathfrak{m}/\mathfrak{p}$ and we can conclude that $(f - f | \mathfrak{p})(g) \| \mathfrak{m} \in \mathcal{A}[\mathcal{B}^\circ(y)]$.

On the other hand $(f||\mathfrak{p})(g) \in \mathcal{A}[\mathcal{B}^\circ(y)]$ by the inductive hypothesis because the order type of $\text{Supp}_{\mathfrak{M}}(f||\mathfrak{p})$ is strictly smaller than the order type of R . This concludes the proof. \square

Proposition 3.30. *If \mathcal{L} is an algebra of infinitesimally convergent power series with coefficients in $\mathbb{K}(\mathfrak{M})$ which is closed under truncations and formal derivatives then \mathcal{L}^* is*

- (1) closed under formal derivatives;
- (2) closed under $||$ -truncations, i.e. if $f(x) = \sum_m r_m x^m \in \mathcal{L}^*$ and $\mathfrak{m} \in \mathfrak{M}$, then $f||\mathfrak{m} := \sum_m (r_m|\mathfrak{m})x^m \in (\mathcal{L}_{\mathfrak{M}})^*$.

Proof. (1) is trivial by the chain rule.

(2) first observe that by Lemma 3.28, the set of $t \in \mathcal{L}^*$ such that $t||\mathfrak{M} \subseteq \mathcal{L}^*$ is a subring of \mathcal{L}^* , then apply Lemma 3.29 to conclude that since it contains \mathcal{L} and is closed under composition, it must be the whole \mathcal{L}^* . \square

Proof of Theorem 3.20. Suppose $X \subseteq \mathbb{K}(\mathfrak{M})$ is closed under truncations and contains \mathfrak{M} and \mathbb{K} . Then $\langle X \rangle_{\mathcal{F}}$ is the set of constants in $(\mathfrak{M}\mathcal{F}_{\mathfrak{M}} \cup X)^*$. By Lemma 3.27 the family $\mathfrak{M}\mathcal{F}_{\mathfrak{M}}$ is closed under truncations and formal derivatives and thus so is $X \cup \mathfrak{M}\mathcal{F}_{\mathfrak{M}}$. By Proposition 3.30 $(\mathfrak{M}\mathcal{F}_{\mathfrak{M}} \cup X)^*$ is closed under truncations, in particular its constants are. \square

4. SERIAL POWER-BOUNDED STRUCTURES

The following definition is partially modelled upon the definition of GQA in [15]. The idea is to require that the language L has both enough functions to represent all definable functions as terms and that the germs of the functions in L are somehow regular enough to be represented by generalized power series.

Definition 4.1. Let $\mathbb{K}_L = (\mathbb{K}, L)$ be a power-bounded o-minimal field in some functional language expansion L , consisting for each n and $r \in (\mathbb{K}^{>0})^n$ of a \mathbb{K} -algebra $L(r)$ of functions on $\prod_i(0, r_i)$ containing the coordinate projections. Suppose furthermore that f satisfies:

- (1) for every $f \in L(r)$, $x \in \prod_i(0, r_i)$ and $\sigma \in \{-1, 0, +1\}^n$, there is a $f_{x,\sigma}$ such that $f(x + \sigma z) = f_{x,\sigma}(z)$ for all small enough $z > 0$;
- (2) the \mathbb{K} algebra L_n of germs at the origin of functions in $L(r)$ for some $r \in (\mathbb{K}^{>0})^n$ is closed under renormalized partial derivatives, that is if $f \in L_n$, then $x_i \partial_i f$ where x_i is the germ of the projection on the i -th coordinate and $\partial_i f$ is the derivative in the i -th variable;
- (3) \mathbb{K}_L has a Skolem Theory.

Let Λ be the field of exponents of \mathbb{K}_L . Let for every n , \mathcal{T}_n be an injective algebra embedding $\mathcal{T}_n : L_n \rightarrow \mathbb{K}((x^\Lambda))$ satisfying $\mathcal{T}(x_i) = x_i$ and $\mathcal{T}_n(x_i \partial_i f) = x_i \partial_i \mathcal{T}(f)$ for any coordinate projection $x_i : \mathbb{K}^n \rightarrow \mathbb{K}$.

Given a multiplicatively written ordered Λ -vector space \mathfrak{M} , we will say that $(\mathbb{K}_L, \mathcal{T})$ is \mathfrak{M} -serial if \mathbb{K}_L satisfies (1), (2) and (3) and furthermore

- (4) interpreting $f \in L(r)$ by setting for every $x \in \prod_i(0, r_i)_{\mathbb{K}}$, and $\varepsilon \in \mathbb{K}((\mathfrak{M}^{<1}))$

$$f(x + \varepsilon) := \mathcal{T}_n(f_{x, \text{sgn}(\varepsilon)})(|\varepsilon|)$$

makes $\mathbb{K}((\mathfrak{M})) \supseteq \mathbb{K}$ an elementary extension of \mathbb{K} (we denote by $\mathbb{K}((\mathfrak{M}))^{\mathcal{T}}$, the field $\mathbb{K}((\mathfrak{M}))$ expanded with such interpretation of symbols in L).

We say that $(\mathbb{K}_L, \mathcal{T})$ is *serial* if it is \mathfrak{M} -serial for every multiplicatively written ordered Λ -vector space \mathfrak{M} . If the image of \mathcal{T}_n is truncation closed for every n , then we say that $(\mathbb{K}_L, \mathcal{T})$ is truncation closed. We denote by $\mathcal{T}(L)$ the (closure under reindexing of the variables of the) algebras series in the image of \mathcal{T} .

Remark 4.2. If each $L(r)$ contains (the restriction of) $1/x_i$ for each coordinate function x_i (as would usually be the case since \mathbb{K}_L is assumed to have a Skolem theory), then the condition on \mathcal{T}_n can be simplified to that of being a truncation closed differential algebra embedding.

Example 4.3. Every real closed field \mathbb{K} has an expansion by definable functions that is serial with a natural choice of \mathcal{T} . Consider for every $F(x, y) \in \mathbb{K}[x_0, \dots, x_{n-1}, y]$ such that $(\partial_y F)(0, 0) \neq 0$, the unique solution $f_F : \prod_{i < n}(0, r_i) \rightarrow \mathbb{K}$ of $F(x, f_F(x)) = 0$ defined for suitably small r_i s.

Then let for each $r \in (\mathbb{K}^{>0})^n$, $L(r)$ consist of closure under partial derivatives of the algebra of functions generated by the functions $f_F \circ g$ where f_F is as above for some F for which f_F is defined on $\prod_i(0, r_i)$ and $g = (g_0, \dots, g_{n-1})$ is a tuple of functions that are either constants in \mathbb{K} or of the form $x \mapsto x_i^q$ for some $q \in \mathbb{Q}$ and some $i < n$. Notice \mathcal{T}_n is defined naturally on the germs at the origin of the f_F s described above as they arise as implicit functions for polynomial equations. \mathcal{T}_n has then a natural extension to L_n .

Example 4.4. By [21, Thm. 2.14] the structure \mathbb{R}_{an} is interdefinable with a serial structure with the natural \mathcal{T} . In particular, by Corollary 4.9 so is every reduct of \mathbb{R}_{an} .

[23, Prop. 10.4], can be restated by saying that the structure $\mathbb{R}_{\mathcal{G}}$ is interdefinable with a \mathfrak{M} -serial structure for all \mathfrak{M} with finitely many Archimedean classes.

Remark 4.5. Notice that $\mathcal{T}(L)$ is an almost fine algebra of generalized series.

Lemma 4.6. *Suppose $(\mathbb{K}_L, \mathcal{T})$ is \mathfrak{M} -serial and truncation closed and $\mathbb{K} \cup \mathfrak{M} \subseteq \mathbb{E} \subseteq \mathbb{K}((\mathfrak{M}))^{\mathcal{T}}$ is truncation closed. Then \mathbb{E} is L -closed if and only if it is $\mathcal{T}(L)$ -closed.*

Proof. Suppose \mathbb{E} is $\mathcal{T}(L)$ -closed, since \mathbb{E} contains \mathbb{K} and \mathfrak{M} , for every $x \in \mathbb{E}$ and every $f \in L$, $f(x) \in \mathbb{E}$ by definition of $\mathbb{K}((\mathfrak{M}))^{\mathcal{T}}$.

Suppose \mathbb{E} is L closed, then it must clearly be $\mathcal{T}(L)$ -closed. \square

Proposition 4.7. *Suppose $(\mathbb{K}_L, \mathcal{T})$ is \mathfrak{M} -serial and truncation-closed and $\mathbb{K} \subseteq \mathbb{E} \subseteq \mathbb{K}((\mathfrak{M}))$ is a truncation closed elementary substructure. If $x \in$*

$\mathbb{K}(\langle \mathfrak{M} \rangle)$ is such that for all $\mathfrak{m} \in \mathfrak{M}$, $x|\mathfrak{m} \neq x \rightarrow x|\mathfrak{m} \in \mathbb{E}$, then $\mathbb{E}\langle x \rangle_L$ is truncation-closed.

Proof. Notice that $\mathfrak{N} := \mathfrak{M} \cap \mathbb{E}$ must be a subgroup of \mathfrak{M} and that $\mathbb{E} = \langle X \rangle_{\mathcal{T}(L)}$ for some truncation closed subset X (e.g. $X = \mathbb{E}$). We distinguish two cases. If $\text{Supp } x$ has a minimum \mathfrak{m} , then $\mathbb{E}\langle x \rangle_L = \mathbb{E}\langle \mathfrak{m} \rangle = \langle X \cup \mathfrak{N}' \rangle_{\mathcal{T}(L)}$ where $\mathfrak{N}' := \mathfrak{M}^\Delta$.

If instead $\text{Supp } x$ has no minimum, then $\mathbb{E}\langle x \rangle_L = \mathbb{E}\langle x \rangle_{\mathcal{T}(L)} = \langle X \cup \{x\} \rangle_{\mathcal{T}(L)}$ and we are done once again. \square

Lemma 4.8. *Suppose $(\mathbb{K}_L, \mathcal{T})$ is \mathfrak{M} -serial for every \mathfrak{M} in some class \mathcal{C} and $L_0 \subseteq L$. Then there is an expansion by definition $L_* \subseteq L$ such that $(\mathbb{K}_{L_*}, \mathcal{T}|_{L_*})$ is \mathfrak{M} -serial for every $\mathfrak{M} \in \mathcal{C}$.*

Proof. Notice \mathbb{K}_{L_0} can be Skolemized to \mathbb{K}_{L_1} using symbols in L , moreover, since Skolem functions are definable in o-minimal theories, L_1 can be built as an expansion by definitions. Similarly L_1 can be closed within L under renormalized derivatives of the germs to get some L_2 and in turn L_3 can be closed under condition (1) by still only choosing symbols from L to get some L_3 . Repeating this process ω many times and taking the union yields the required L_* . Such L_* still satisfies point (4) because elementary extensions are preserved under reducts. \square

Corollary 4.9. *If $(\mathbb{K}_L, \mathcal{T})$ is \mathfrak{M} -serial and truncation closed and the image of \mathcal{T}_n lies within $\bigcup_{n \in \mathbb{N}} \mathbb{K}(\langle z^{1/(n+1)\mathbb{Z}} \rangle)$, then every reduct of $(\mathbb{K}_L, \mathcal{T})$ has an expansion by definition that is serial and truncation closed.*

Remark 4.10. If $\mathcal{A} := \{\mathcal{A}_{m,n,r} : m, n \in \mathbb{N}, r \in (\mathbb{R}^{>0})^{m+n}\}$ is a generalized quasianalytic algebra as defined in [15, Sec. 2] and let Λ be the field of exponents of $\mathbb{R}_{\mathcal{A}}$. Defining L by setting for $r \in (\mathbb{R}^{>0})^n$ $A(r) := \mathcal{A}_{n,0,r} |_{\prod_{i < n} (0,r_i)}$ and $L(r) := \{f \circ x^\alpha : \alpha \in \Lambda^n, f \in A(r)\}$, should yield a power bounded structure \mathbb{R}_L satisfying (1), (2) and (3) of Definition 4.1 by [15, Cor. 5.3].

Question 4.11. *Let \mathcal{A} be a generalized quasianalytic algebra as defined in [15, Sec. 2]. Is $\mathbb{R}_{\mathcal{A}}$ interdefinable with a serial structure?*

Remark 4.12. Notice that if λ is greater than every ordinal embeddable in the field of exponents Λ then $\mathbb{K}(\langle \mathfrak{M} \rangle)_\lambda \subseteq \mathbb{K}(\langle \mathfrak{M} \rangle)^\mathcal{T}$ is an L -substructure. We denote the resulting expansion of $\mathbb{K}(\langle \mathfrak{M} \rangle)_\lambda$ to a L -structure by $\mathbb{K}(\langle \mathfrak{M} \rangle)_\lambda^\mathcal{T}$.

5. THE MORGUES-RESSAYRE CONSTRUCTIONS REVISITED

Definition 5.1. An *rv-sected model* of T_{convex}^- is a quadruple $\mathcal{E} := (\mathbb{E}, \mathcal{O}, \mathbb{K}, \mathfrak{M})$ where $(\mathbb{E}, \mathcal{O})$ is a model of T_{convex}^- , $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$ is a T -section for the residue field and \mathfrak{M} is a \mathbb{K} -monomial group. The rv-sected model of T_{convex}^- \mathcal{E} will be said to be *above* $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}^-$; $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}$ and \mathfrak{M} will be referred to respectively as the *residue section* and *monomial group* of \mathcal{E} .

An *embedding of rv-sected models* $\iota : \mathcal{E}_0 \rightarrow \mathcal{E}_1$ is an embedding $\iota : (\mathbb{E}_0, \mathcal{O}_0) \rightarrow (\mathbb{E}_1, \mathcal{O}_1)$ of the underlying models of T_{convex}^- such that $\iota(\mathbb{K}_0) \subseteq$

\mathbb{K}_1 and $\iota(\mathfrak{M}_0) \subseteq \mathfrak{M}_1$. In such a case we will also say that the rv-sections $\mathcal{E}_0, \mathcal{E}_1$ are *compatible* with $\iota : (\mathbb{E}_0, \mathcal{O}_0) \rightarrow (\mathbb{E}_1, \mathcal{O}_1)$.

For T a power-bounded theory, and embedding of models of T_{convex}^- , $\iota : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}_1, \mathcal{O}_1)$ will be said to be λ -*bounded* if it has the following property: for every rv-sected $\mathcal{E}, \mathcal{E}_1$ above $(\mathbb{E}, \mathcal{O})$ and $(\mathbb{E}_1, \mathcal{O}_1)$ respectively, such that $\iota : \mathcal{E} \rightarrow \mathcal{E}_1$ is an embedding of rv-sected models, $(\mathbb{E}_1, \mathcal{O}_1)$ is λ -bounded wim-constructible over $(\text{dcl}_T(\mathbb{E}\mathfrak{M}_1), \text{CH}(\mathbb{K}))$.

Remark 5.2. Notice that this does not create ambiguity with the previous definition of λ -bounded wim-constructible, as an extension of models of T_{convex}^- is λ -bounded wim-constructible if and only if it is both λ -bounded and wim-constructible as shown by the following

Lemma 5.3. *Let T be power-bounded with field of exponents Λ . The following are equivalent for an extension $\iota : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}', \mathcal{O}')$ of models of T_{convex}^- :*

- (1) *there is sequence $(\mathbb{E}_i, \mathcal{O}_i)$ such that $(\mathbb{E}_i, \mathcal{O}_i) = \bigcup_{j < i} (\mathbb{E}_{j+1}, \mathcal{O}_{j+1})$ and every $(\mathbb{E}_{i+1}, \mathcal{O}_{i+1})$ is a principal extension of $(\mathbb{E}_i, \mathcal{O}_i)$ which is either λ -bounded weakly immediate or valational;*
- (2) *ι is λ -bounded.*

Proof. (1) \Rightarrow (2), observe that a composition of λ -bounded extensions is λ -bounded, therefore it suffices to show that if $(\mathbb{E}, \mathcal{O}) \subseteq (\mathbb{E}\langle x \rangle, \mathcal{O}_1)$ is a principal extension which is either λ -bounded weakly immediate or valational, then the extension is λ -bounded. If it is λ -bounded weakly immediate there is nothing to prove, in the other case there is $c \in \mathbb{E}$ such that $\mathbf{v}_{\mathcal{O}_1}(x - c) \notin \mathbf{v}_{\mathcal{O}_1}\mathbb{E}$. Notice that if \mathfrak{M} is a monomial group for $(\mathbb{E}, \mathcal{O})$, and $\mathbf{v}_{\mathcal{O}}(y) \in \mathbf{v}_{\mathcal{O}_1}(\mathbb{E}_1) \setminus \mathbf{v}_{\mathcal{O}}(\mathbb{E})$, then by the rv-property $\mathfrak{M}_1 := \mathfrak{M}y^\Lambda$ is a monomial group for $(\mathbb{E}_1, \mathcal{O}_1)$ and one easily sees that $\mathbb{E}\langle x \rangle = \text{dcl}_T(\mathfrak{M}_1\mathbb{E})$.

(2) \Rightarrow (1) add one element at a time. \square

Lemma 5.4. *If T' is the theory of an on-minimal field, T is a power-bounded reduct and $(\mathbb{E}, \mathcal{O}) \prec (\mathbb{E}^*, \mathcal{O}^*) \models T'_{\text{convex}}$ is λ -wim constructible for some $\lambda \geq |T'|^+$, then the underlying extension of reducts to T_{convex} is λ -bounded.*

Proof. Since λ -bounded extensions of models of T are closed under composition, it suffices to show that every $(\mathbb{E}^*, \mathcal{O}^*) := (\mathbb{E}\langle x \rangle_{T'}, \mathcal{O}_x) \succ (\mathbb{E}, \mathcal{O}) \models T'_{\text{convex}}$ with x T'_{convex} -weakly immediate over $(\mathbb{E}, \mathcal{O})$, is λ -bounded qua extension of models of T .

By Lemma 5.3, it suffices to show that $(\mathbb{E}^*, \mathcal{O}^*) \succ (\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}$ is a composition of principal extensions that are either λ -bounded immediate or valational.

Since by Remark 2.9 $\mathbf{r}(\mathbb{E}^*, \mathcal{O}^*) = \mathbf{r}(\mathbb{E}, \mathcal{O})$, and T is power-bounded, given a dcl_T -basis $(x_i : i < \mu)$ of \mathbb{E}^* over \mathbb{E} , for every i , x_i is either valational or weakly immediate over $\mathbb{E}_i := \mathbb{E}\langle x_j : j < i \rangle$. So it suffices to show that whenever $\text{tp}(x_i/\mathbb{E}_i)$ is weakly immediate, it must have cofinality $< \lambda$.

Notice that by hypothesis $\mathbb{E}^{<x_i}$ has cofinality $< \lambda$ and that it must be $\mu \leq |T'|$, therefore $\mathbb{E}_i^{<x_i} \setminus \mathbb{E}^{<x_i}$ if non-empty has cofinality at most $|T| + |i| \leq |T'|$. It follows that $\mathbb{E}_i^{<x_i}$ has cofinality $< \lambda$. \square

Definition 5.5. Suppose $\mathbb{K}_L \models T$ is power-bounded with field of exponents Λ and that $(\mathbb{K}_L, \mathcal{T})$ is serial. For every multiplicatively written ordered Λ -vector space \mathfrak{M} , we will denote by $[\mathbb{K}((\mathfrak{M}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ the rv-sected model of T_{convex} $(\mathbb{K}((\mathfrak{M}))_\lambda^\mathcal{T}, \text{CH}(\mathbb{K}), \mathbb{K}, \mathfrak{M})$.

Given a rv-sected model of T_{convex} \mathcal{E} with residue section (T -isomorphic to) \mathbb{K} , a *t.c. embedding* is an embedding of rv-sected models of T_{convex}^- , $\iota : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ for some λ such that the image of ι is a truncation-closed subfield of $[\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$.

Let $(\mathbb{E}, \mathcal{O}) \preceq (\mathbb{E}_1, \mathcal{O}_1)$ and \mathcal{E} an rv-sected model above $(\mathbb{E}, \mathcal{O})$. A t.c. embedding $\iota : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ will be said to be *v-maximal within* $(\mathbb{E}_1, \mathcal{O}_1)$ if every proper extension of ι along some T_{convex} -extension $j : (\mathbb{E}, \mathcal{O}) \preceq (\mathbb{E}_2, \mathcal{O}_2)$ factoring the inclusion $(\mathbb{E}, \mathcal{O}) \preceq (\mathbb{E}_1, \mathcal{O}_1)$ is such that $\mathbf{v}_{\mathcal{O}_2}(\mathbb{E}) \neq \mathbf{v}_{\mathcal{O}_2}(\mathbb{E}_2)$.

Context: Througout the rest of the section $(\mathbb{K}_L, \mathcal{T})$ will be a serial power-bounded structure with field of exponents Λ , T will be the theory of $(\mathbb{K}_L, \mathcal{T})$ and T' will be the theory of an o-minimal expansion $\mathbb{K}_{L'}$ of \mathbb{K}_L .

Lemma 5.6. Assume $\eta : \mathcal{E}_0 \rightarrow \mathcal{E}_1$ is a λ -bounded extension of rv-sected models of T_{convex}^- with residue section \mathbb{K} ; \mathcal{E}_i be above \mathbb{E}_i and with monomial group \mathfrak{M}_i (for $i \in \{0, 1\}$). Assume furthermore $\eta|_{\mathbb{K}} = \text{id}$.

If $\iota : \mathcal{E}_0 \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ is a λ -bounded truncation closed embedding, and $j : \mathfrak{M}_1 \rightarrow \mathfrak{N}$ is an (ordered Λ -linear) extension of $\iota|_{\mathfrak{M}_0}$, then ι extends along η to a λ -bounded truncation closed embedding $\iota_1 : \mathcal{E}_1 \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ such that $\iota_1|_{\mathfrak{M}_1} = j$.

Proof. By Lemma 5.3, it suffices to prove the statement in the case two cases $\mathbb{E}_1 = \mathbb{E}_0\langle \mathfrak{m} \rangle$ with $\mathfrak{m} \in \mathfrak{M}_1 \setminus \eta\mathfrak{M}_0$ and $\mathbb{E}_1 = \mathbb{E}\langle x \rangle$ with x λ -bounded immediate.

In the first case, by Proposition 4.7, $(\iota\mathbb{E}_0)\langle j\mathfrak{m} \rangle_T$ is truncation closed and we are done.

If x is λ -bounded immediate and $(x_i)_{i < \mu} \in \mathbb{E}_0^\mu$ is a p.c.-sequence for x , then we can set $\iota(x)$ to be the only element of $(\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K}))$ which is a pseudolimit for $(\iota x_i)_{i < \mu}$ and is such that $\iota(x)|_{\mathfrak{m}} \neq \iota(x) \Rightarrow \iota(x)|_{\mathfrak{m}} \in \iota\mathbb{E}_0$. Again by Proposition 4.7 $(\iota\mathbb{E}_0)\langle \iota(x) \rangle_T$ is truncation closed and we are done. \square

Theorem 5.7. If $(\mathbb{E}, \mathcal{O}) \models T'_{\text{convex}}$, $\mathbb{K}_{L'} \preceq_{\text{tame}} \mathbb{E}$, $\mathcal{O} = \text{CH}(\mathbb{K})$, and λ is large enough, then there is an expansion $(\mathbb{M}, \mathcal{O}_1)$ of some $[\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ to a model of $(T')_{\text{convex}}^-$ and a λ -bounded wim-constructible T' -elementary embedding $\iota : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{M}, \mathcal{O}_1)$ with truncation-closed image.

Proof. Let $\varepsilon_\lambda : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ be an elementary embedding of $(\mathbb{E}, \mathcal{O})$ into its λ -bounded spherical completion. We can assume ε_λ is an inclusion and $\mathbb{K} \preceq_{\text{tame}} \mathbb{E}_\lambda$.

Let $\mathcal{E} := (\mathbb{E}, \mathcal{O}, \mathbb{K}, \mathfrak{M})$ and $\mathcal{E}_\lambda := (\mathbb{E}, \mathcal{O}, \mathbb{K}, \mathfrak{N})$ be rv-sections above the T_{convex}^- reducts of $(\mathbb{E}, \mathcal{O})$ and $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$, compatible with ε_λ .

By Lemma 5.6, if λ is large enough there is a λ -bounded truncation closed embedding $\iota' : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{M}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$.

Let ι'' be the composition of ι' with the natural inclusion

$$[\mathbb{K}((\mathfrak{M}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}} \hookrightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}.$$

Observe that ι'' also has a truncation closed image. Again by Lemma 5.6, ι'' can be extended along ε_λ to a λ -bounded truncation closed embedding $\eta : (\mathbb{E}_\lambda, \mathcal{O}_\lambda) \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$, now observe that since $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ is λ -spherically complete, η must be an isomorphism. The image of $\eta \circ \varepsilon_\lambda$ is the image of ι'' , so it is truncation closed. \square

Definition 5.8. Given a model of $(\mathbb{E}, \mathcal{O}) \models T_{\text{convex}}^-$. We say that an ordered exponential $\exp : \mathbb{E} \rightarrow \mathbb{E}^{>0}$ is compatible if $\exp(\mathcal{O}) \subseteq \mathcal{O}$.

In presence of a compatible exponential, we will say that a t.c. embedding $\iota : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda^T, \text{CH}(\mathbb{K})]_{\mathcal{T}}$, with $\mathcal{E} = (\mathbb{E}, \mathcal{O}, \mathbb{K}, \mathfrak{M})$ is

- *dyadic* (after Ressayre [14]) if $\iota : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ is such that $\iota \log \mathfrak{M} \subseteq \mathbb{K}((\mathfrak{N}^{>1}))_\lambda$;
- *T4* (after Schmeling [17]) if for every sequence of monomials $(\mathfrak{m}_n)_{n \in \mathbb{N}}$ with $\iota \mathfrak{m}_{n+1} \in \text{Supp}(\iota \log \mathfrak{m}_n)$ for all $n \in \mathbb{N}$, there is $N \in \mathbb{N}$ such that for all $n \geq N$, $\iota \log \mathfrak{m}_n = c_n \pm \iota \mathfrak{m}_{n+1}$ for some c_n with $\text{Supp} c_n > \iota \mathfrak{m}_{n+1}$.

We will call *R.S. embedding* (for Ressayre Schmeling) a t.c. embedding that is both dyadic and T4.

Lemma 5.9. *Let $j : (\mathbb{E}, \mathcal{O}) \preceq (\mathbb{E}_*, \mathcal{O}_*)$ be a λ -bounded extension of models of T_{convex}^- expanded with a compatible exponential. Let \mathfrak{N} be a multiplicative copy of the value group of $(\mathbb{E}_*, \mathcal{O}_*)$. Assume \mathcal{E} is an rv-section of $(\mathbb{E}, \mathcal{O})$ and $\iota : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$ is a R.S. embedding, maximal within $(\mathbb{E}_*, \mathcal{O}_*)$. Then there is an rv-section \mathcal{E}_* of $(\mathbb{E}_*, \mathcal{O}_*)$ extending \mathcal{E} , such that ι extends to a R.S. embedding $\iota_* : \mathcal{E}_* \rightarrow [\mathbb{K}((\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$.*

Proof. The same argument of Ressayre's [14] (see also [2]). Notice it suffices to show the statement with \mathfrak{N} a sufficiently saturated multiplicatively written ordered Λ -vector space.

Let \mathfrak{M} be the monomial group of the fixed rv-section \mathcal{E} on $(\mathbb{E}, \mathcal{O})$. By hypothesis $\iota \mathfrak{M} \subseteq \mathfrak{N}$ and $\log \iota \mathfrak{M} \subseteq \mathbb{K}((\mathfrak{N}^{>1}))$. Pick an $x \in \mathbb{E}_1 \setminus \mathbb{E}$, it suffices to show that we can extend ι to a R.S. embedding of some extension \mathcal{E}_ω of \mathcal{E} above some intermediate $(\mathbb{E}_\omega, \mathcal{O}_\omega)$, $(\mathbb{E}, \mathcal{O}) \preceq (\mathbb{E}_\omega, \mathcal{O}_\omega) \preceq (\mathbb{E}_*, \mathcal{O}_*)$ which is closed under exponentiation.

Consider $\mathbb{E}\langle x \rangle_T$. Since ι is \mathbf{v} -maximal within $(\mathbb{E}_*, \mathcal{O}_*)$, $\mathbf{wlog} \mathbf{v}(\mathbb{E}\langle x \rangle_T) \neq \mathbf{v}\mathbb{E}$, for otherwise by Lemma 5.6, this would contradict maximality. Thus

without loss of generality we can assume that $\mathbb{E}\langle x \rangle_T = \mathbb{E}\langle y \rangle_T$ where $\mathbf{v}(y) \notin \mathbf{v}(\mathbb{E})$ and $y \notin \mathcal{O}_*$.

We now inductively build a sequence $(y_i)_{i < \omega} \in \mathbb{E}_*$ such that $y_0 = y$ and for every i , $y_i \notin \mathcal{O}_*$, $\mathbf{v}_{\mathcal{O}_*}(y_i) \notin \mathbf{v}_{\mathcal{O}_*}\mathbb{E} + \sum_{j < i} \Lambda \mathbf{v}_{\mathcal{O}_*}(y_j)$, $\log |y_i| - y_{i+1} \in \mathbb{E}$ and $\iota(\log |y_i| - y_{i+1}) \in \mathbb{K}(\mathfrak{N}^{>1})$.

Given $(y_j)_{j < i+1}$, observe that $(\log |y_i| + \mathcal{O}_*) \cap \mathbb{E} = \emptyset$ because $y_i \notin \mathcal{O}_*$ and $\mathbf{v}_{\mathcal{O}_*}(y_i) \notin \mathbf{v}_{\mathcal{O}_*}(\mathbb{E})$. Notice that since $\log |y_i|$ is either weakly immediate λ -bounded or valuations over the extension $(\mathbb{E}, \text{CH}(\mathcal{O}))$, either $\mathbf{v}_{\mathcal{O}_*}(\log |y_i| - c_i) \notin \mathbf{v}_{\mathcal{O}_*}\mathbb{E}$ for some $c \in \mathbb{E}$ or $\log |y_i|$ is a pseudo-limit of a p.c. sequence, however since ι was \mathbf{v} -maximal within $(\mathbb{E}_*, \mathcal{O}_*)$ this second option would imply $\log |y_i| \in \mathbb{E}$, but this would contradict that \mathbb{E} is exp-closed. Thus there is $c_i \in \mathbb{E}$ such that $\mathbf{v}_{\mathcal{O}_*}(\log |y_i| - c_i) \notin \mathbf{v}_{\mathcal{O}_*}\mathbb{E}$.

Since $(\log |y_i| + \mathcal{O}_*) \cap \mathbb{E} = \emptyset$, it follows that $\log |y_i| - c_i \notin \mathcal{O}_*$, we can thus change the choice of c_i so that $c_i \in \mathbb{K}(\mathfrak{N}^{>1})$, maintaining the property $\mathbf{v}_{\mathcal{O}_*}(\log |y_i| - c_i) \notin \mathbf{v}_{\mathcal{O}_*}\mathbb{E}$.

Setting $y_{i+1} := \log |y_i| - c_i$ we have an extension of the sequence with the required properties: this is because by construction $1 \prec y_{i+1} \prec \log |y_i|$ so $1 \prec |y_{i+1}|^\Lambda \leq \exp(|y_{i+1}|) \prec y_i$, so in particular $\mathbf{v}_{\mathcal{O}_*} y_{i+1}$ is Λ -linearly independent over $\mathbf{v}_{\mathcal{O}_*}\mathbb{E}$ from $\{\mathbf{v}_{\mathcal{O}_*}(y_j) : j < i + 1\}$.

Let $\mathfrak{M}_1 := \mathfrak{M} \cdot \bigcup_{i < \omega} y_0^\Lambda \cdot \dots \cdot y_i^\Lambda$, $\mathbb{E}'_1 = \mathbb{E}(y_i : i < \omega)_T$ and $\mathcal{E}'_1 = (\mathbb{E}'_1, \mathcal{O}'_1, \mathbb{K}, \mathfrak{M}_1)$. Observe \mathcal{E}'_1 extends \mathcal{E} .

Let $h : \mathfrak{M}_1 \rightarrow \mathfrak{N}$ be an extension of $\iota|_{\mathfrak{M}}$ to \mathfrak{M}_1 . By Lemma 5.6 $\iota : \mathcal{E} \rightarrow (\mathbb{K}(\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K}))$ extends to a $\iota'_1 : \mathcal{E}'_1 \rightarrow (\mathbb{K}(\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K}))$ such that $\iota'_1|_{\mathfrak{M}_1} = h$.

Again by Lemma 5.6 ι'_1 extends to a $\iota_1 : \mathcal{E}_1 \rightarrow (\mathbb{K}(\mathfrak{N}))_\lambda, \text{CH}(\mathbb{K}))$ that is \mathbf{v} -maximal within $(\mathbb{E}_*, \mathcal{O}_*)$.

Notice $(\mathbb{E}_1, \mathcal{O}_1)$ is closed under logarithms and that:

- $\iota_1 \log \mathfrak{M}_1 \subseteq \mathbb{K}(\mathfrak{N}^{>1})$;
- $\text{Supp } \iota_1 \log |y_i| \subseteq \iota \mathfrak{M} \cup \{\iota_1 y_{i+1}\}$

Now inductively define a sequence $\iota_n : \mathcal{E}_n \rightarrow \mathbb{K}(\mathfrak{N})$ such that ι_n is maximal within $(\mathbb{E}_*, \mathcal{O}_*)$, $(\mathbb{E}_n, \mathcal{O}_n)$ is closed under logarithms, $\iota_n(\log \mathfrak{M}_n) \subseteq \mathbb{K}(\mathfrak{N}^{>1})$ and $\mathfrak{M}_{n+1} = \exp(\iota_n^{-1}(\mathbb{K}(\mathfrak{N}^{>1})))$. The base case is given by the $(\mathbb{E}_1, \mathcal{O}_1)$ constructed above and the inductive step is possible by Lemma 5.6 because $\mathfrak{M}_n \subseteq \mathfrak{M}_{n+1}$. Setting $\iota_\omega = \bigcup_n \iota_n$ we get the desired extension. \square

Corollary 5.10. *Suppose $(\mathbb{E}, \mathcal{O})$ is a λ -bounded extension of $(\mathbb{K}_{L'}, \mathbb{K}_{L'}) \models (T')_{\text{convex}}^-$ with T' exponential and \mathfrak{M} is a multiplicative copy of the value group of $(\mathbb{E}, \mathcal{O})$. Then there is some \mathcal{E} above $(\mathbb{E}, \mathcal{O})$ and a R.S. embedding $\mathcal{E} \rightarrow [\mathbb{K}(\mathfrak{N})]_\lambda, \text{CH}(\mathbb{K})_{\mathcal{T}}$.*

We have said enough to deduce Theorem C of the introduction.

Theorem 5.11. *Let $(\mathbb{E}, \mathcal{O}) \models (T')_{\text{convex}}$ with residue $\mathbb{K}_{L'}$, $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ its T' - λ -spherical completion and \mathfrak{N} a multiplicative copy of the value group of $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$. Suppose T' defines an exponential and λ is a large enough cardinal. Then there is a T_{convex} -isomorphism $\eta : (\mathbb{E}_\lambda, \mathcal{O}_\lambda) \rightarrow (\mathbb{K}(\mathfrak{N})_\lambda^T, \text{CH}(\mathbb{K}))$ such*

that $\eta(\mathbb{E})$ is truncation closed and the expansion of $(\mathbb{K}((\mathfrak{M}))_\lambda^T, \text{CH}(\mathbb{K}))$ by $\eta \circ \exp \circ \eta^{-1}$ satisfies (D) and (T4) in the introduction.

Proof. Let $\varepsilon_\lambda : (\mathbb{E}, \mathcal{O}) \rightarrow (\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ be an elementary embedding of $(\mathbb{E}, \mathcal{O})$ into its T - λ -spherical completion. Let \mathfrak{M} be a section of the value group of $(\mathbb{E}, \mathcal{O})$ and let $\mathcal{E} = (\mathbb{E}, \mathcal{O}, \mathbb{K}, \mathfrak{M})$. By Corollary 5.10, if λ is large enough there is a transserial T_{convex} -embedding $\iota' : \mathcal{E} \rightarrow [\mathbb{K}((\mathfrak{M}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$. By Lemma 5.9, ι' can be extended along ε_λ to a transserial embedding $\eta : \mathcal{E}_\lambda \rightarrow [\mathbb{K}((\mathfrak{M}))_\lambda, \text{CH}(\mathbb{K})]_{\mathcal{T}}$, for some \mathcal{E}_λ above $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ extending \mathcal{E} . Now observe that since $(\mathbb{E}_\lambda, \mathcal{O}_\lambda)$ is λ -spherically complete η must be an isomorphism. The map $\eta \circ \varepsilon_\lambda$ is then truncation-closed and $\eta \circ \exp \circ \eta^{-1}$ is an exponential on $(\mathbb{K}((\mathfrak{M}))_\lambda^T, \text{CH}(\mathbb{K}))$ satisfying (D) and (T4) in the introduction. \square

Corollary 5.12 (Mantova). *Let \mathbb{R}_L be a reduct of \mathbb{R}_{an} defining the restricted exponential, then every elementary extension of $\mathbb{R}_{L,\text{exp}}$ has an elementary truncation closed embedding in \mathbf{No} .*

Proof. Let $(\mathbb{R}_{An}, \mathcal{T})$ be the standard serial structure interdefinable with \mathbb{R}_{an} . By Example 4.4 and Lemma 4.8, we can assume without loss of generality that $(\mathbb{R}_L, \mathcal{T}|)$ is serial. Let $T = Th(\mathbb{R}_L)$. Notice that every elementary extension $\mathbb{E} \succeq \mathbb{R}_{L,\text{exp}}$ is tame, thus in particular the T_{convex}^- -reduct of $(\mathbb{E}, \text{CH}(\mathbb{R}))$ is a λ -bounded extension of $(\mathbb{R}_L, \mathbb{R}_L) \models T_{\text{convex}}^-$ for every large enough λ . Let $(\mathbb{E}_\lambda, \mathcal{O}_\lambda) \succeq (\mathbb{E}, \text{CH}(\mathbb{R}))$ and

$$\eta : (\mathbb{E}_\lambda, \mathcal{O}_\lambda) \rightarrow (\mathbb{R}((\mathfrak{M}))_\lambda^{T|}, \text{CH}(\mathbb{R}))$$

be as in Theorem 5.11.

Observe that since $(\mathbb{R}_L, \mathcal{T}|)$ is serial $(\mathbb{R}((\mathfrak{M}))_\lambda^{T|}, \text{CH}(\mathbb{R}))$ is an L -elementary substructure of $(\mathbb{R}((\mathfrak{M}))^{T|}, \text{CH}(\mathbb{R}))$. Now $(\mathbb{R}((\mathfrak{M}))^{T|}, \eta \circ \log \circ \eta^{-1})$ is a transserial Hahn field as defined in [3, Def. 6.2].

By [3, Thm. 8.1] there is a transserial initial embedding $\iota : \mathbb{R}((\mathfrak{M}))^{T|} \rightarrow \mathbf{No}$: that is, an embedding that is the identity on the reals, sends monomials to monomials and preserves logarithms and formal infinite sums. Since $(\mathbb{R}_L, \mathcal{T}|)$ is serial this entails that actually ι is L -elementary. On the other hand by [24, Thm. 3.2], $Th(\mathbb{R}_{L,\text{exp}})$ is still model complete, so

$$\iota| : (\mathbb{R}((\mathfrak{M}))_\lambda^T, \eta \circ \exp \circ \eta^{-1}) \rightarrow \mathbf{No}$$

is (L, exp) -elementary. Now $\eta(\mathbb{E})$ is truncation closed by Theorem 5.11, so $(\iota \circ \eta)(\mathbb{E})$ is truncation closed and elementary. \square

REFERENCES

- [1] A. Berarducci and P. Freni. On the value group of the transseries. *Pacific J. Math.*, 312(2):335–354, 2021.
- [2] P. D’Aquino, J. F. Knight, S. Kuhlmann, and K. Lange. Real closed exponential fields. *Fund. Math.*, 219(2):163–190, 2012.
- [3] P. Ehrlich and E. Kaplan. Surreal ordered exponential fields. *The Journal of Symbolic Logic*, 86(3):1066–1115, 2021.

- [4] A. Fornasiero. Initial embeddings in the surreal numbers of models of $t_{an}(exp)$.
- [5] P. Freni. T -convexity, weakly immediate types and t - λ -spherical completions of o-minimal structures. *arXiv preprint arXiv:2404.07646*, 2024.
- [6] E. Kaplan. T -convex T -differential fields and their immediate extensions. *Pacific J. Math.*, 320(2):261–298, 2022.
- [7] L. S. Krapp, S. Kuhlmann, and M. Serra. On Rayner structures. *Comm. Algebra*, 50(3):940–948, 2022.
- [8] F.-V. Kuhlmann, S. Kuhlmann, and S. Shelah. Exponentiation in power series fields. *Proceedings of the American Mathematical Society*, 125(11):3177–3183, 1997.
- [9] S. Kuhlmann and S. Shelah. κ -bounded exponential-logarithmic power series fields. *Annals of Pure and Applied Logic*, 136(3):284–296, 2005.
- [10] D. Marker and C. I. Steinhorn. Definable types in O-minimal theories. *J. Symbolic Logic*, 59(1):185–198, 1994.
- [11] M.-H. Mourgues and J. P. Ressayre. Every real closed field has an integer part. *J. Symbolic Logic*, 58(2):641–647, 1993.
- [12] B. H. Neumann. On ordered division rings. *Transactions of the American Mathematical Society*, 66(1):202–252, 1949.
- [13] A. Pillay. Definability of types, and pairs of O-minimal structures. *J. Symbolic Logic*, 59(4):1400–1409, 1994.
- [14] J.-P. Ressayre. Integer parts of real closed exponential fields (extended abstract). In *Arithmetic, proof theory, and computational complexity (Prague, 1991)*, volume 23 of *Oxford Logic Guides*, pages 278–288. Oxford Univ. Press, New York, 1993.
- [15] J.-P. Rolin and T. Servi. Quantifier elimination and rectilinearization theorem for generalized quasianalytic algebras. *Proc. Lond. Math. Soc. (3)*, 110(5):1207–1247, 2015.
- [16] J.-P. Rolin, T. Servi, and P. Speissegger. On transasymptotic expansions of o-minimal germs. *arXiv preprint arXiv:2402.12073*, 2024.
- [17] M. C. Schmeling. *Corps de transséries*. PhD thesis, Paris 7, 2001.
- [18] J. M. Tyne. *T-levels and T-convexity*. ProQuest LLC, Ann Arbor, MI, 2003. Thesis (Ph.D.)—University of Illinois at Urbana-Champaign.
- [19] L. van den Dries. T -convexity and tame extensions. II. *J. Symbolic Logic*, 62(1):14–34, 1997.
- [20] L. van den Dries and A. H. Lewenberg. T -convexity and tame extensions. *J. Symbolic Logic*, 60(1):74–102, 1995.
- [21] L. van den Dries, A. Macintyre, and D. Marker. The elementary theory of restricted analytic fields with exponentiation. *Ann. of Math. (2)*, 140(1):183–205, 1994.
- [22] L. van den Dries, A. Macintyre, and D. Marker. Logarithmic-exponential series. *Ann. Pure Appl. Logic*, 111(1-2):61–113, 2001.
- [23] L. van den Dries and P. Speissegger. The field of reals with multi-summable series and the exponential function. *Proc. London Math.*

Soc. (3), 81(3):513–565, 2000.

- [24] L. van den Dries and P. Speissegger. O-minimal preparation theorems. 11:87–116, 2002.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UNITED KINGDOM