Stochastic Gradient Descent for Two-layer Neural Networks[†]

Dinghao Cao¹, Zheng-Chu Guo¹ and Lei Shi²

School of Mathematical Sciences, Zhejiang University, Hangzhou 310058, P. R. China
 Shanghai Key Laboratory for Contemporary Applied Mathematics,
 School of Mathematical Sciences, Fudan University, Shanghai 200433, P. R. China

Abstract

This paper presents a comprehensive study on the convergence rates of the stochastic gradient descent (SGD) algorithm when applied to overparameterized two-layer neural networks. Our approach combines the Neural Tangent Kernel (NTK) approximation with convergence analysis in the Reproducing Kernel Hilbert Space (RKHS) generated by NTK, aiming to provide a deep understanding of the convergence behavior of SGD in overparameterized two-layer neural networks. Our research framework enables us to explore the intricate interplay between kernel methods and optimization processes, shedding light on the optimization dynamics and convergence properties of neural networks. In this study, we establish sharp convergence rates for the last iterate of the SGD algorithm in overparameterized two-layer neural networks. Additionally, we have made significant advancements in relaxing the constraints on the number of neurons, which have been reduced from exponential dependence to polynomial dependence on the sample size or number of iterations. This improvement allows for more flexibility in the design and scaling of neural networks, and will deepen our theoretical understanding of neural network models trained with SGD.

Keywords and phrases: Learning theory; Overparameterized two-layer neural network; Neural tangent kernel; Reproducing kernel Hilbert space; Stochastic gradient descent.

1 Introduction

Neural networks have emerged as powerful models capable of learning intricate patterns and making predictions based on input data. As parameterized models, obtaining optimal parameters for neural networks has become a highly relevant and extensively studied topic in both academic and industrial communities. Efficiently and accurately optimizing these parameters is crucial for the performance of neural networks. Among the various algorithms proposed for parameter optimization, stochastic gradient descent (SGD) stands out as one of the most

[†] The work by D. Cao and Z. C. Guo is partially supported by Zhejiang Provincial Natural Science Foundation of China [Project No. LR20A010001] and National Natural Science Foundation of China [Project No. U21A20426, No. 12271473]. The work by L. Shi is partially supported by the National Natural Science Foundation of China (Grants No. 12171039, No. 12061160462) and Shanghai Science and Technology Program [Project No. 21JC1400600]. All authors contributed equally to this work and are listed alphabetically. The corresponding author is Zheng-Chu Guo. Email addresses: caodinghao@zju.edu.cn (D. Cao), guozhengchu@zju.edu.cn (Z. C. Guo), leishi@fudan.edu.cn (L. Shi).

widely used and well-studied algorithms, which has been proven particularly effective in tackling large-scale data problems. However, despite its widespread adoption, theoretical aspects of the SGD algorithm in neural networks need to be further accomplished. It still needs to be thoroughly investigated under which conditions the algorithm converges, at what rate it converges, and how to adjust its parameters to achieve fast convergence. These questions have attracted significant attention and have been extensively explored recently. Studying the convergence behavior of the SGD algorithm has a profound impact on understanding the efficiency and effectiveness of neural network training.

Neural networks have become prevalent in various practical applications. Among the different types of neural networks, over-parameterized neural networks play a pivotal role in advancing deep learning. Over-parameterization refers to neural networks with more parameters than the amount of training data available. In fields like computer vision and natural language processing, over-parameterized neural networks such as ResNet and Transformer have already become the cutting-edge models. Understanding the convergence mechanism of the SGD algorithm applied to over-parameterized neural networks is crucial for effectively adjusting the parameters of neural networks.

In this paper, we focus on analyzing the convergence properties of over-parameterized two-layer neural networks trained with SGD, and studying its generalization performance under the framework of learning with Neural Tangent Kernel (NTK). The NTK has emerged as a powerful theoretical framework for comprehending the behavior of deep neural networks (DNNs). It was initially introduced by Jacot et al. in their influential work [18], where they established connections between neural networks with infinite-width and kernel methods. This groundbreaking research demonstrated that the forward pass of a randomly initialized neural network can be approximated by a kernel function. The work laid the foundation for subsequent studies on the NTK, driving further advancements in this field. By leveraging the NTK framework, researchers have gained valuable insights into the training dynamics and generalization properties of DNNs. The NTK serves as a bridge between neural network theory and practice, providing a deeper understanding of how neural networks learn and generalize. By investigating the properties of over-parameterized two-layer neural networks trained with SGD under the NTK regime, we aim to contribute to the understanding of their generalization capabilities. This analysis will shed light on the behavior of neural networks during training and their ability to generalize to unseen data. Ultimately, the insights gained from this paper can guide the parameter adjustment and application of neural networks, fostering advancements in deep learning.

The consistency property of the SGD algorithm in neural networks has been extensively studied. For instance, the work of [14] established the consistency of SGD to the global minimum when the square loss function is employed in a two-layer neural network with Gaussian inputs. Similarly, the consistency analysis for the hinge loss function was conducted in [6], while the cross-entropy loss function was studied in [21]. Moreover, specific network structures have also been considered. For instance, [22] demonstrated that applying the SGD algorithm to a two-layer ResNet with Gaussian inputs will converge to the global minimum. The generalization behavior of two-layer neural networks, incorporating the concept of algorithmic stability, was explored in [19]. The research investigated both gradient descent and stochastic gradient descent as training algorithms for the neural network and derived excess risk bounds that hold even under relaxed over-parameterization assumptions. In addition to studying the case of two-layer neural networks, extensive efforts have been made to understand the

consistency of the SGD algorithm in deep neural networks. The work of [1] established the convergence to the global minimum using the square loss function. Their research specifically addressed the consistency of SGD in deep convolutional neural networks and deep residual networks. For binary classification problems, global convergence of SGD was proven in [39]. Additionally, [2] demonstrated that when the square loss function is employed, SGD applied to a multi-layer recurrent neural network (RNN) converges to the global minimum at a linear convergence rate.

Compared with consistency, the convergence rate is an ingredient of more importance to characterize the dynamic behavior of SGD quantitively, which provides valuable insights into how fast the algorithm can converge to the optimal solution. In the work of Arora et al. [3], the convergence rate of the batch gradient descent algorithm was explored in over-parameterized two-layer neural networks with the square loss function and ReLU activation. Their research demonstrated a convergence rate of $O(T^{-\frac{1}{2}})$, where T represents the number of training samples. By utilizing the positive definiteness of the NTK, which implies the generated Gram matrix is strictly positive definite, they approximated the objective function of the neural network as a linear function within a small neighborhood. This approximation played a crucial role in analyzing the convergence rates of SGD in neural networks. However, subsequent studies have shown that as the number of training samples T increases, the eigenvalues of the NTK's Gram matrix tend to zero [32]. This observation challenges the positive definiteness assumption of the NTK utilized in previous works. Consequently, deriving convergence rates without relying on the positive definiteness assumption has become a challenging problem. Nitanda and Suzuki [26] proved that for overparameterized two-layer neural networks with smooth activation functions and the square loss function, the convergence rates of averaged SGD can reach $O\left(T^{-\frac{2r\beta}{2r\beta+1}}\right)$. Here, $\frac{1}{2} < r \le 1$ represents the regularity parameter of the target function (see Assumption 3), and $\beta > 1$ represents the capacity parameter of the hypothesis space (see Assumption 4). It is worth noting that achieving these fast convergence rates requires an exponential number of neurons in the network. While these results indicate the potential for fast convergences with SGD in certain cases, the practical limitation of exponentially many neurons suggests that achieving such fast rates may not be feasible or efficient in real-world scenarios with large-scale neural networks. Recent work by Nguyen and Mücke [25] addressed the regression problem by employing the batch gradient descent method to train the neural network. They demonstrated that the number of neurons required in the network can be reduced to a polynomial function of the number of iterations, providing a more practical approach to achieving convergence. In the context of classification problems, Cao and Gu [7] presented similar results regarding the convergence rate of SGD in overparameterized deep neural networks. They showed that the convergence rate of SGD can reach $O(T^{-\frac{1}{2}})$ (up to logarithmic terms).

Contributions. In this paper, we focus on analyzing the convergence rates of the last iterate of the SGD algorithm for an overparameterized two-layer neural network. We conduct a solid theoretical analysis under a framework that utilizes the NTK to approximate the objective function of the neural network within a small neighborhood. This approximation allows us to study the convergence properties of SGD and understand how the algorithm progresses towards the optimal solution. Furthermore, we establish a rigorous convergence analysis of SGD with NTK from a kernel-based learning perspective, successfully incorporating some a priori conditions from the convergence analysis of SGD in reproducing kernel Hilbert space (RKHS), such as characterizing the capacity of the space and the regularity

conditions on the smoothness of the objective function. The convergence analysis of SGD in RKHS has previously produced promising results [35, 13, 37, 17]. However, extending the existing kernel-based SGD framework in an RKHS to our current setting is not trivial. Firstly, the positive definite assumption of the NTK may not hold when the training data is sufficiently large, limiting the applicability of the method in [3] for large-scale datasets. Secondly, although the work of Nitanda and Suzuki [26] does not rely on the positive definiteness assumption of the NTK, they consider the Averaged SGD (ASGD) algorithm, which averages the outputs in each iteration and is different from our setting. Averaging helps to reduce variance and accelerate the convergence of SGD. However, their analysis requires that the width (i.e., the number of neurons) depends exponentially on the number of iterations or the sample size T, which differs significantly from practical applications, thus failing to demonstrate the algorithmic performance in real-world scenarios. Additionally, the standard last-step SGD algorithm, while widely used and effective in many practical applications, does not incorporate the variance reduction mechanism characteristic of the ASGD algorithm. Analyzing the convergence behavior of the non-averaged scheme in SGD is generally more challenging, as fluctuations in the gradients can significantly affect convergence properties. Consequently, understanding the convergence rates and behavior of the non-averaged SGD algorithm requires more intricate analysis techniques and estimations. To the best of our knowledge, our paper is the first to demonstrate the generalization performance of the last-step SGD applied to training neural networks. We develop an innovative and insightful analysis, achieving sharp convergence rates of $O(T^{-\frac{2r}{2r+1}})$ or $O(T^{-\frac{2r}{2r+1}+\epsilon})$ in two distinct scenarios, where $0 < \epsilon < \frac{2r}{2r+1}$ can be arbitrarily small. Moreover, we have made significant advancements in relaxing the constraints on the number of neurons or the width of the neural network. In previous works [26, 32], the number of neurons exhibited an exponential dependency. In our analysis, it now demonstrates a polynomial dependency on the sample size or the number of iterations.

The remainder of this paper is organized as follows. We introduce the two-layer neural network, SGD algorithm and NTK in Section 2. Main results and some related discussions are given in Section 3. The proofs of our main results are given in Section 4 and some useful lemmas and proofs are presented in Section 5.

2 Preliminary

In this paper, we focus on the regression problem, which involves learning a functional relationship from random samples in order to make predictions about future observations. Specifically, let $\mathcal{X} \subset \mathbb{R}^d$ be a compact metric space and $\mathcal{Y} \subseteq \mathbb{R}$, ρ be a Borel probability distribution on $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$. Given a pair $(x,y) \in \mathcal{Z}$, where x represents the input and y represents the corresponding output, we aim to learn a function $g: \mathcal{X} \to \mathcal{Y}$ that provides predictions for the output variable y given an input x. The prediction error incurred is characterized by the least-squares loss $(g(x) - y)^2$. The goal of the least square regression problem is to find an optimal prediction function g by minimizing the expected risk

$$\mathcal{E}(g) = \int_{\mathcal{X} \times \mathcal{Y}} (g(x) - y)^2 d\rho$$

over all measurable functions. This target function is known as the *regression function* and is defined as

$$g_{\rho}(x) = \int_{\mathcal{Y}} y d\rho(y|x), \quad x \in \mathcal{X},$$

where $\rho(\cdot|x)$ is the conditional distribution at x induced by the joint distribution ρ .

Within the framework of learning theory, the distribution ρ is often unknown, and the goal is to estimate g_{ρ} based on independently drawn samples from ρ . Regression has been extensively studied in the fields of machine learning and statistical inference, e.g., see [12, 31] and references therein. The performance of the estimator g of the regression function g_{ρ} is usually measured by the excess risk which is defined as

$$\mathcal{E}(g) - \mathcal{E}(g_{\rho}) = \|g - g_{\rho}\|_{\rho}^{2},$$

where $\|\cdot\|_{\rho}$ denotes the norm in the space $\mathcal{L}^2_{\rho_{\mathcal{X}}}$ induced by the inner product $\langle u, v \rangle_{\rho} = \int_{\mathcal{X}} u(x)v(x)d\rho_{\mathcal{X}}(x)$ for $u, v \in \mathcal{L}^2_{\rho_{\mathcal{X}}}$, here $\mathcal{L}^2_{\rho_{\mathcal{X}}}$ denotes the Hilbert space of functions from \mathcal{X} to \mathcal{Y} square-integrable with respect to the marginal distribution $\rho_{\mathcal{X}}$ of ρ . The excess risk quantifies the difference between the expected risk of the estimator g and the optimal expected risk achieved by the regression function g_{ρ} . In this paper, the estimator g of g_{ρ} is derived from a two-layer overparametrized neural network, which will be formally defined in the subsequent subsection.

2.1 Two-layer Neural Networks

Let $M \in \mathbb{N}_+$ be the network width (number of neurons). Let $a = (a_1, \dots, a_M)^\top \in \mathbb{R}^M$ be the parameters of the output layer, $B = (b_1^\top, \dots, b_M^\top)^\top \in \mathbb{R}^{dM}$ be the parameters of the input layer, and $c = (c_1, \dots, c_M)^\top \in \mathbb{R}^M$ be the bias parameters, for $r \in \mathbb{N}$ and $1 \le r \le M$, $a_r \in \mathbb{R}$, $b_r \in \mathbb{R}^d$ and $c_r \in \mathbb{R}$. We denote by $\Theta = (a, B, c) \in \mathbb{R}^{M(d+2)}$, and consider two-layer neural networks

$$g_{\Theta}(x) = \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r \sigma \left(b_r^{\top} x + \gamma c_r \right), \tag{1}$$

where $\sigma: \mathbb{R} \to \mathbb{R}$ is an activation function and $\gamma > 0$ is a scale of the bias terms.

Obviously, a neural network is a parametric model, and solving the model involves finding the optimal values for its parameters Θ . Iterative methods, such as the SGD algorithm that will be discussed in this paper, are commonly employed for this purpose. To use an iterative method, we start with an initial set of parameters and then update them iteratively according to a specific strategy until convergence is achieved. The initialization strategy employed in this paper is referred to as symmetric initialization [32]. The purpose of symmetric initialization is to ensure that the initial function $g_{\Theta^{(0)}}$ is set to zero. This choice is made for theoretical simplicity rather than any specific practical reason. Specifically, let $a^{(0)} = \left(a_1^{(0)}, \cdots, a_M^{(0)}\right)^{\top}$, $B^{(0)} = \left(b_1^{(0)^{\top}}, \cdots, b_M^{(0)^{\top}}\right)^{\top}$, $c^{(0)} = \left(c_1^{(0)}, \cdots, c_M^{(0)}\right)^{\top}$. Assume that the number of neurons M is even. The parameters for the output layer are initialized as

$$a_r^{(0)} = R, \quad r \in \left\{1, \dots, \frac{M}{2}\right\},$$

$$a_r^{(0)} = -R, \quad r \in \left\{\frac{M}{2} + 1, \dots, M\right\},$$

where R > 0. Let μ_0 be a uniform distribution on the sphere $\mathbb{S}^{d-1} = \{b \in \mathbb{R}^d : ||b||_2 = 1\} \subset \mathbb{R}^d$ used to initialize the parameters for the input layer. The parameters for the input layer are

initialized as

$$b_r^{(0)} = b_{r + \frac{M}{2}}^{(0)} \stackrel{i.i.d}{\sim} \mu_0, \quad r \in \left\{1, \dots, \frac{M}{2}\right\}.$$

The bias parameters are initialized as

$$c_r^{(0)} = 0, \quad r \in \{1, \cdots, M\}.$$

2.2 Stochastic Gradient Descent

We consider the following regularized expected risk minimization problem

$$\min_{\Theta} \left\{ \mathcal{E}\left(g_{\Theta}\right) + \lambda \left(\left\| a - a^{(0)} \right\|_{2}^{2} + \left\| B - B^{(0)} \right\|_{2}^{2} + \left\| c - c^{(0)} \right\|_{2}^{2} \right) \right\},\,$$

where regularization parameter $\lambda > 0$, $\|\cdot\|_2$ denotes the ℓ^2 norm for vectors in \mathbb{R}^d or \mathbb{R}^{dM} . We incorporate regularization into the expected risk because regularization helps to keep the iterations obtained by the optimization algorithm close to the initial value. This proximity to the initial value allows us to achieve better convergence properties during the optimization process [32].

In this paper, we optimize the parameters Θ using the SGD approach, which is outlined in Algorithm 1. Let $\Theta^{(t)} = (a^{(t)}, B^{(t)}, c^{(t)})$ denote the collection of t-th iterates of parameters. Let $\Theta^{(1)} = \Theta^{(0)}$, for $t \ge 1$ and $t \in \mathbb{N}$, assuming that we randomly sample data $(x_t, y_t) \sim \rho$ in the t-th iteration, the parameter update formula for the SGD algorithm, following the steps outlined in Algorithm 1, can be expressed as follows,

$$\begin{split} a_r^{(t+1)} - a_r^{(0)} &= (1 - \eta_t \lambda) \left(a_r^{(t)} - a_r^{(0)} \right) - \eta_t M^{-1/2} \left(g_{\Theta^{(t)}} \left(x_t \right) - y_t \right) \sigma \left(b_r^{(t) \top} x_t + \gamma c_r^{(t)} \right), \\ b_r^{(t+1)} - b_r^{(0)} &= (1 - \eta_t \lambda) \left(b_r^{(t)} - b_r^{(0)} \right) - \eta_t M^{-1/2} \left(g_{\Theta^{(t)}} \left(x_t \right) - y_t \right) a_r^{(t)} \sigma' \left(b_r^{(t) \top} x_t + \gamma c_r^{(t)} \right) x_t, \\ c_r^{(t+1)} - c_r^{(0)} &= (1 - \eta_t \lambda) \left(c_r^{(t)} - c_r^{(0)} \right) - \eta_t M^{-1/2} \left(g_{\Theta^{(t)}} \left(x_t \right) - y_t \right) a_r^{(t)} \gamma \sigma' \left(b_r^{(t) \top} x_t + \gamma c_r^{(t)} \right), \end{split}$$

where $\eta_t > 0$ is the step size (or learning rate). In practice, there are typically two common approaches for setting the step size: using a constant value or applying polynomial decay over time. In this paper, we set the step size to decrease over time according to the $\eta_t = \eta t^{-\theta}$, where η is a positive constant and $\frac{1}{2} < \theta < 1$.

Algorithm 1 Stochastic Gradient Descent

Input: Number of iterations T, width of network M, regularization parameter λ , step sizes $\{\eta_t\}_{t=1}^T$, initial parameters: $\Theta^{(1)} = \Theta^{(0)} = (a^{(0)}, B^{(0)}, c^{(0)})$ and bias parameter γ **Output:** Estimation for $\Theta^{(T+1)} = (a^{(T+1)}, B^{(T+1)}, c^{(T+1)})$

- 1: for t = 1 to T do
- 2: Randomly draw a sample $z_t = (x_t, y_t) \sim \rho$
- 3: $\Theta^{(t+1)} \leftarrow \Theta^{(t)} \eta_t \left(g_{\Theta^{(t)}}(x_t) y_t \right) \nabla_{\Theta} g_{\Theta}(x_t) \Big|_{\Theta = \Theta^{(t)}} \eta_t \lambda \left(\Theta^{(t)} \Theta^{(0)} \right)$
- 4: end for

2.3 Neural Tangent Kernel

The neural tangent kernel (NTK) is a kernel function that has gained significant attention in recent years due to its crucial role in the global convergence analysis of SGD training neural networks [18, 1, 3]. By capturing the relationship between the network's parameters and the loss landscape, the NTK provides insights into the dynamic behaviors of neural networks during training, which bridges the neural network models with kernel methods. Consequently, leveraging the power of RKHS and kernel methods can provide effective tools to aid in the analysis of various problems.

In our settings, NTK is defined as, $\forall x, x' \in \mathcal{X}$,

$$k_{\infty}\left(x,x'\right) = \mathbb{E}_{b^{(0)}}\left[\sigma\left(b^{(0)\top}x\right)\sigma\left(b^{(0)\top}x'\right)\right] + R^{2}\left(x^{\top}x' + \gamma^{2}\right)\mathbb{E}_{b^{(0)}}\left[\sigma'\left(b^{(0)\top}x\right)\sigma'\left(b^{(0)\top}x'\right)\right],\tag{2}$$

where $b^{(0)} \sim \mu_0$. In practical scenarios, the width of the neural network is finite. Therefore, it is common to approximate $k_{\infty}(x, x')$ using its empirical counterpart $k_M(x, x')$ defined as

$$k_{M}(x, x') = \frac{1}{M} \sum_{r=1}^{M} \sigma\left(b_{r}^{(0)\top}x\right) \sigma\left(b_{r}^{(0)\top}x'\right) + \frac{\left(x^{\top}x' + \gamma^{2}\right)}{M} \sum_{r=1}^{M} \sigma'\left(b_{r}^{(0)\top}x\right) \sigma'\left(b_{r}^{(0)\top}x'\right). \quad (3)$$

It can be easily proved that both k_{∞} and k_M are continuous, symmetric and positive definite kernel functions, thus they are Mercer kernel functions [11]. As a result, the RKHSs generated by these kernels can be defined as follows

$$\mathcal{H}_{\infty} = \overline{\operatorname{span}} \left\{ k_{\infty,x}(\cdot) = k_{\infty}(x,\cdot), \quad x \in \mathcal{X} \right\},$$

$$\mathcal{H}_{M} = \overline{\operatorname{span}} \left\{ k_{M,x}(\cdot) = k_{M}(x,\cdot), \quad x \in \mathcal{X} \right\}.$$

For \mathcal{H}_{∞} , the inner product between functions $k_{\infty,x}$ and $k_{\infty,x'}$ is denoted as $\langle k_{\infty,x}, k_{\infty,x'} \rangle_{\mathcal{H}_{\infty}}$ and is equal to $k_{\infty}(x,x')$. Similarly, the inner product between functions $k_{M,x}$ and $k_{M,x'}$ in \mathcal{H}_{M} is represented as $\langle k_{M,x}, k_{M,x'} \rangle_{\mathcal{H}_{M}} = k_{M}(x,x')$. Obviously, k_{∞} is a Mercer kernel (continuous and positive semi-definite), which uniquely induces an RKHS, and the reproducing property

$$f(x) = \langle f, k_{\infty, x} \rangle_{\mathcal{H}_{\infty}},\tag{4}$$

holds for any $x \in \mathcal{X}$ and $f \in \mathcal{H}_{\infty}$. Analogously, k_M is also a Mercer kernel of RKHS \mathcal{H}_M with $g(x) = \langle g, k_{M,x} \rangle_{\mathcal{H}_M}$ for any $x \in \mathcal{X}$ and $g \in \mathcal{H}_M$.

3 Main Results and Discussion

Without loss of generalization, we assume that there is no bias term c_r in this paper, since we can set $(x, \gamma) \to \bar{x}$, and $(b_r^\top, c_r) \to \bar{b}_r^\top$, then the neural network function (1) can be rewritten as

$$g_{\Theta}(x) = \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r \sigma \left(b_r^{\top} x + \gamma c_r \right) = \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r \sigma \left(\bar{b}_r^{\top} \bar{x} \right).$$

Therefore, we can assume there is no bias term c_r in the neural network function (1). And for $t \ge 1$, we define $\Theta^{(t)}$ as a vector

$$\Theta^{(t)} = (a_1^{(t)}, \cdots, a_M^{(t)}, b_1^{(t)^{\top}}, \cdots, b_M^{(t)^{\top}})^{\top} \in \mathbb{R}^{(d+1)M}.$$
 (5)

In this section, we will present the main theoretical results and outline the fundamental assumptions for our analysis. The activation function $\sigma(\cdot)$ is assumed to satisfy the following properties:

Assumption 1. There exists a constant $C_{\sigma} > 0$, such that $\|\sigma'\|_{\infty} \leq C_{\sigma}$, $\|\sigma''\|_{\infty} \leq C_{\sigma}$, and $|\sigma(u)| \leq 1 + |u|, \forall u \in \mathbb{R}$.

This assumption relies on the smoothness of the activation function used in the neural network. Commonly used activation functions such as Sigmoid, tanh, and the smooth approximation of ReLU all satisfy this requirement.

Assumption 2. Assume that $\rho_{\mathcal{X}}$ is non-degenerate and supported on $\mathcal{X} = \{x \in \mathbb{R}^d : ||x||_2 \leq 1\}$, $\mathcal{Y} \subset [-1,1], R = 1, \text{ and } \gamma \in [0,1].$

This assumption corresponds to the conventional boundedness assumption commonly used in convergence analysis. Without loss of generality, we assume all bounds to be 1, but these bounds can be relaxed to any bounded positive constants. Under Assumption 1 and 2, for any $x, x' \in \mathcal{X}$, we have

$$|k_M(x, x')| \le 4 + 2C_\sigma^2 =: \kappa^2.$$
 (6)

Let \mathcal{H}_{∞} be the RKHS generated by the NTK k_{∞} , L_{∞} be the integral operator in $\mathcal{L}^2_{\rho_{\mathcal{X}}}$ and $\{\mu_i\}_{i=1}^{\infty}$ be eigenvalues of L_{∞} sorted in non-ascending order. Where the integral operator $L_{\infty}: \mathcal{L}^2_{\rho_{\mathcal{X}}} \to \mathcal{L}^2_{\rho_{\mathcal{X}}}$ associated with NTK k_{∞} and $\rho_{\mathcal{X}}$ is defined as

$$L_{\infty}f(\cdot) = \int_{\mathcal{X}} f(x)k_{\infty}(x,\cdot) \, d\rho_{\mathcal{X}}(x), \, \forall f \in \mathcal{L}^{2}_{\rho_{\mathcal{X}}}.$$
 (7)

Given that k_{∞} is a Mercer kernel and \mathcal{X} is compact, the operator L_{∞} is self-adjoint, positive and trace class on $\mathcal{L}_{\rho_{\mathcal{X}}}^2$, as well as restricting on \mathcal{H}_{∞} . The compactness of L_{∞} implies the existence of an orthonormal eigensystem $\{\mu_k, \phi_k\}_{k \in \mathbb{N}}$ in $\mathcal{L}_{\rho_{\mathcal{X}}}^2$, where the eigenvalues $\{\mu_k\}_{k \in \mathbb{N}}$ (with geometric multiplicities) are non-negative and arranged in decreasing order. Then for any r > 0, we can define the r-th power of L_{∞} , denoted as L_{∞}^r , as follows $L_{\infty}^r \left(\sum_{k \geq 1} c_k \phi_k\right) = \sum_{k \geq 1} c_k \mu_k^r \phi_k$. It is worth noting that L_{∞}^r is a positive compact operator on $\mathcal{L}_{\rho_{\mathcal{X}}}^2$. Now let us introduce the notion of regularity, which is often interpreted as the smoothness or source condtion of the regression function g_{ρ} .

Assumption 3. There exists $r > \frac{1}{2}$, such that $g_{\rho} \in L_{\infty}^{r}\left(\mathcal{L}_{\rho_{\mathcal{X}}}^{2}\right)$, that is $\|L_{\infty}^{-r}g_{\rho}\|_{\mathcal{L}_{\rho_{\mathcal{X}}}^{2}} < \infty$.

This assumption is a measure of the smoothness of the regression function g_{ρ} , which is a standard requirement in learning theory. It implies that g_{ρ} belongs to the range space of L_{∞}^{r} expressed as

$$L_{\infty}^{r}\left(\mathcal{L}_{\rho_{\mathcal{X}}}^{2}\right) = \left\{ f \in \mathcal{L}_{\rho_{\mathcal{X}}}^{2} : \sum_{k \geq 1} \frac{\left\langle f, \phi_{k} \right\rangle_{\mathcal{L}_{\rho_{\mathcal{X}}}^{2}}^{2}}{\mu_{k}^{2r}} < \infty \right\},$$

Therefore, the larger r is, the faster the decay rate of the expansion coefficient of g_{ρ} under the eigenfunction basis of L_{∞} is, that is, the more smoothness that it may have. Besides, it is not difficult to see that $\forall r_1 \geq r_2, L_{\infty}^{r_1}(\mathcal{L}_{\rho_{\mathcal{X}}}^2) \subseteq L_{\infty}^{r_2}(\mathcal{L}_{\rho_{\mathcal{X}}}^2)$. Moreover, since $\rho_{\mathcal{X}}$ is non-degenerate,

from Theorem 4.12 in [12], $L_{\infty}^{1/2}$ is an isomorphism from $\overline{\mathcal{H}_{\infty}}$, the closure of \mathcal{H}_{∞} in $\mathcal{L}_{\rho_{\mathcal{X}}}^2$, to \mathcal{H}_{∞} , i.e., for each $f \in \overline{\mathcal{H}_{\infty}}$, $L_{\infty}^{1/2} f \in \mathcal{H}_{\infty}$ and

$$||f||_{\rho} = \left\| L_{\infty}^{1/2} f \right\|_{\mathcal{H}_{\infty}}.$$
 (8)

Therefore, $L_{\infty}^{1/2}(\mathcal{L}_{\rho_{\mathcal{X}}}^2) = \mathcal{H}_{\infty}$, and Assumption (3) implies $g_{\rho} \in \mathcal{H}_{\infty}$.

Next, we introduce a crucial assumption regarding the capacity of the RKHS, which is based on the integral operator L_{∞} .

Assumption 4. There exists $\beta > 1$, and positive constants b, c > 0, such that $\forall i \geq 1$, $bi^{-\beta} \leq \mu_i \leq ci^{-\beta}$.

It is worth noting that the upper bound part of this assumption always holds when $\beta = 1$. The capacity of the hypothesis space \mathcal{H}_{∞} is commonly measured using covering numbers or effective dimension $\mathcal{N}_{L_{\infty}}(\lambda) = \text{Tr}((L_{\infty} + \lambda I)^{-1}L_{\infty})$, where I denotes the identity operator. In [15], it has been proven that upper bound part of Assumption 4 is equivalent to the assumption of effective dimension $\mathcal{N}_{L_{\infty}}(\lambda) = O(\lambda^{-1/\beta})$. The decay condition of the eigenvalues often leads to sharp estimates.

Under the above assumptions, we prove the convergence rate of the SGD algorithm applied to an overparameterized two-layer neural network, which is the main content of the Theorem 1. The proof of the Theorem 1 will be given in Section 4. For $k \in \mathbb{N}_+$, let $\mathbb{E}_{z_1,\dots,z_k}$ denote taking expectation with respect to z_1,\dots,z_k , which is written as \mathbb{E}_{Z^k} for short.

Theorem 1. Suppose Assumptions 1, 2, 3 (with $\frac{1}{2} < r \le 1$) and 4 (with $\beta > 1$) hold. For any $\lambda > 0$, $T \in \mathbb{N}_+$. Run algorithm 1 with a polynomially decaying rate $\eta_t = \eta t^{-\theta}$ with $\theta \in (\frac{1}{2}, 1)$ and

$$0 < \eta < \left(\lambda + \frac{208(1+\kappa)^4}{(1-2^{\theta-1})(2\theta-1)} \left(\log \frac{8}{1-\theta} + \frac{1}{1-\theta}\right)\right)^{-1},$$

the smallest integer not less than $x \in \mathbb{R}$ then there exists $M_0(T, \lambda) = \left[\max\left\{T^{6-5\theta}, T^{2\theta}, \frac{T^{\theta}}{\lambda^2}\right\}\right] \in \mathbb{Z}_+$ such that for any $M \geq M_0(T, \lambda)$, the following inequalities holds with probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$

(a) if
$$\frac{1}{2} < \theta < \frac{2\beta - 1}{3\beta - 1}$$
,

$$\mathbb{E}_{Z^T} \left[\|g_{\Theta^{(T+1)}} - g_{\rho}\|_{\rho}^2 \right] \le \tilde{C} \left(T^{-2r(1-\theta)} + \exp\left\{ -H(\theta) T^{1-\frac{3\beta-1}{2\beta-1}\theta} \right\} + T^{-\theta} + \lambda^{2r} \right) \log \frac{2}{\delta},$$

(b) if
$$\frac{2\beta - 1}{3\beta - 1} \le \theta < 1$$
,

$$\mathbb{E}_{Z^T}\left[\|g_{\Theta^{(T+1)}} - g_\rho\|_\rho^2\right] \leq \tilde{C}\left(T^{-2r(1-\theta)} + \exp\left\{-H(\theta)\lambda T^{1-\theta}\right\} + T^{-\theta} + \lambda^{2r}\right)\log\frac{2}{\delta},$$

where $H(\theta) = \frac{2\eta}{1-\theta} 2^{1-\theta} \left(1 - \left(\frac{3}{4}\right)^{1-\theta}\right)$ and the constant \tilde{C} is independent of T and λ and will be given in the proof.

Remark. Theorem 1 establishes that, subject to suitable assumptions, the SGD algorithm applied to an overparameterized two-layer neural network can achieve global convergence towards the regression function. Furthermore, by appropriately selecting values for λ , θ , and M, we can attain the following precise and sharp convergence rate.

Corollary 1. Under the assumptions of Theorem 1, for any $M \ge M_0(T) = T^{1+\frac{5}{2r+1}}$ and $0 < \epsilon < \frac{2r}{2r+1}$, choose $\theta = \frac{2r}{2r+1}$ and

$$\lambda = \begin{cases} T^{-\frac{1}{2r+1}}, & \frac{1}{2} < r < 1 - \frac{1}{2\beta} \\ T^{-\frac{1}{2r+1} + \frac{\epsilon}{2r}}, & 1 - \frac{1}{2\beta} \le r \le 1 \end{cases},$$

the following inequality holds with probability at least $1-\delta$ over the random choice of $\Theta^{(0)}$

$$\mathbb{E}_{Z^T} \left[\|g_{\Theta^{(T+1)}} - g_{\rho}\|_{\rho}^2 \right] = \begin{cases} O\left(T^{-\frac{2r}{2r+1}}\right) \log \frac{2}{\delta}, & \frac{1}{2} < r < 1 - \frac{1}{2\beta} \\ O\left(T^{-\frac{2r}{2r+1} + \epsilon}\right) \log \frac{2}{\delta}, & 1 - \frac{1}{2\beta} \le r \le 1 \end{cases}.$$

Remark. The convergence rate of $O\left(T^{-\frac{2r\beta}{2r\beta+1}}\right)$ is commonly referred to as optimal in the minimax sense[8]. When Assumption 4 holds with $\beta=1$, we can observe from Corollary 1 that our convergence rate of $O\left(T^{-\frac{2r}{2r+1}+\epsilon}\right)$ with any $0<\epsilon<\frac{2r}{2r+1}$, can approach the capacity-independent optimal convergence rate of $O\left(T^{-\frac{2r}{2r+1}}\right)$ in the minimax sense. This means that with an arbitrarily small constant ϵ , we can achieve convergence rates that are arbitrarily close to the optimal capacity independent convergence rate.

Instead of considering the last iterate, Nitanda and Suzuki [26] investigated the averaging estimator of each iterate, i.e., $g_{\bar{\Theta}^{(T+1)}}$, with $\bar{\Theta}^{(T+1)} = \frac{1}{T+1} \sum_{t=1}^{T+1} \Theta^{(t)}$, where $\Theta^{(t)}$ is given by Algorithm 1. It shows that the minimax optimal rate can be derived under the same assumptions regarding the regularity of the regression function and the capacity of the RKHS, achieving this rate necessitates an exponential number of neurons. In other words, to obtain the convergence rate, a significantly larger number of neurons is required in [32]. We see that the convergence rates in Corollary 1 is slightly worse than that of [32], however, we have made significant progress in relaxing the constraints on the number of neurons, transitioning from exponential dependence $O(\exp(T))$ to polynomial dependence $O(T^{1+\frac{5}{2r+1}})$ on the sample size or number of iterations T. This advancement allows for more scalable and efficient learning algorithms in neural networks. The averaging scheme has the ability to reduce variance, which results in more robust estimators and improved convergence rates, such as [24, 28, 35]. However, experimental evidence suggests that a simpler approach, such as averaging only the last few iterates or even returning the last iterate alone, can yield excellent performance in practice [28, 29].

Many studies have been devoted to investigating the convergence rate of the SGD algorithm in the RKHS, while most of literature primarily focuses on the analysis involving a single RKHS [9, 15, 34, 10, 16, 20]. In contrast, our results consider the interplay between two kernels, namely k_M and k_∞ . For instance, Tarrès and Yao [33] investigated a regularized online learning algorithm in an RKHS \mathcal{H}_K associated with a Mercer kernel K. They considered the following iterative scheme, denoted as (9), which can be interpreted as the SGD method applied to solve the Tikhonov regularized problem in the RKHS \mathcal{H}_K . In this scheme,

the initial value is set as $f_0 = 0$, and at each iteration t, the update step is given by:

$$f_t = f_{t-1} - \eta_t [(f_{t-1}(x_t) - y_t) K_{x_t} + \lambda_t f_{t-1}], \quad \forall t \in \mathbb{N}_+,$$
(9)

where $z_t = (x_t, y_t)$ represents a sample, and the stochastic gradient is given by $2[(f_{t-1}(x_t)$ $y_t)K_{x_t} + \lambda_t f_{t-1}$. This stochastic gradient approximates the gradient of the objective function $\int (f(x)-y)^2 d\rho + \lambda ||f||_K^2$ evaluated at $f=f_{t-1}$ and $\lambda=\lambda_t$. Under the assumption that the regression function $f_{\rho} \in L_K^r\left(\mathcal{L}_{\rho_{\mathcal{X}}}^2\right)$ with $\frac{1}{2} < r \le 1$, Tarrès and Yao [33] showed that by setting $\eta_t = at^{-\frac{2r}{2r+1}}$ and $\lambda_t = \frac{1}{a}t^{-\frac{1}{2r+1}}$ for some constant a > 1, the algorithm achieves an error bound of $||f_T - f_\rho||_\rho^2 = O\left(T^{-\frac{2r}{2r+1}}(\log \delta/2)^4\right)$ holds with confidence at least $1 - \delta$. This convergence rate is capacity-independent optimal. It is worth noting that in our Corollary 1, the capacityindependent result with $\beta = 1$, denoted as $O\left(T^{-\frac{2r}{2r+1}+\epsilon}\right)$, can be arbitrarily close to the result obtained by Tarrès and Yao. Furthermore, assume that the regularization parameter λ remains constant throughout the iterations and does not vary with time t. Ying and Pontil [37] also studied the SGD algorithm (9) with $\lambda_t = \lambda$ under the same regularity condition on the regression function, i.e., $f_{\rho} \in L_K^r(\mathcal{L}_{\rho_{\mathcal{X}}}^2)$ with $\frac{1}{2} < r \le 1$. They used $\eta_t = t^{-\frac{2r}{2r+1}}$ and $\lambda = T^{-\frac{1}{2r+1} + \frac{\epsilon}{2r}}$ with any $0 < \epsilon < \frac{2r}{2r+1}$ and demonstrated convergence rates of $O\left(T^{-\frac{2r}{2r+1} + \epsilon}\right)$ in expectation. This result aligns with our findings when $\beta = 1$ in Corollary 1. Moreover, by employing Assumption 4 with $\beta > 1$, we can further enhance the result to achieve capacityindependent optimality.

Besides regularized online learning algorithm considered in the literature, the unregularized version (i.e., $\lambda=0$) has also received considerable attention in studies such as [37, 17, 13]. To gain a comprehensive understanding of the convergence rates of unregularized online learning algorithms in RKHS, we suggest referring to Section 3 of the paper [17]. The estimates and techniques developed in this paper, combined with the approximation and consistency results of deep neural networks [36, 38, 23], can be used to study the generalization capabilities of SGD algorithms in training deep network models. Additionally, we will consider extending the analysis in this paper to the framework of operator learning based on kernel methods [30] in our future work.

4 Proof of Main Results

In this section, we will prove our main results, i.e., Theorem 1 and Corollary 1.

4.1 Error Decomposition

Our goal is to estimate $\mathbb{E}_{Z^T}\left[\|g_{\Theta^{(T+1)}} - g_\rho\|_\rho^2\right]$. To simplify this estimation, we will decompose it into several individual quantities that are easier to estimate. In order to achieve this, we will introduce some essential tools and concepts related to convergence analysis in RKHS, which will be utilized repeatedly in subsequent proofs.

Let $M \in \mathbb{N}_+$, and $k_{M,x} = k_M(x,\cdot)$. Similar to (7), we define the integral operator L_M generated by the kernel function k_M as follows

$$L_M f(\cdot) = \int_{\mathcal{X}} f(x) k_M(x, \cdot) d\rho_{\mathcal{X}}(x), \quad \forall f \in \mathcal{L}^2_{\rho_{\mathcal{X}}}.$$

It is known that $L_M^{\frac{1}{2}}: \mathcal{L}_{\rho_{\mathcal{X}}}^2 \to \mathcal{H}_M$ is isometric isomorphism if $\operatorname{Ker} L_M = \{0\}$ [11], that is, for all $f, g \in \mathcal{L}_{\rho_{\mathcal{X}}}^2$, $\left\langle L_M^{1/2} f, L_M^{1/2} g \right\rangle_{\mathcal{H}_M} = \langle f, g \rangle_{\mathcal{L}_{\rho_{\mathcal{X}}}^2}$. Let $g_{M,\lambda}$ be the minimizer of the regularization risk in \mathcal{H}_M , that is

$$g_{M,\lambda} = \arg\min_{g \in \mathcal{H}_M} \left\{ \mathcal{E}(g) + \lambda \|g\|_{\mathcal{H}_M}^2 \right\}. \tag{10}$$

Since $\lambda > 0$, a solution to (10) exists and is unique, and given by [8]

$$g_{M,\lambda} = (L_M + \lambda I)^{-1} \mathbb{E}_{(X,Y)} [Y k_{M,X}] = (L_M + \lambda I)^{-1} L_M g_{\rho}.$$
 (11)

Similarly, we introduce the function $g_{\infty,\lambda}$ as

$$g_{\infty,\lambda} = \arg\min_{g \in \mathcal{H}_{\infty}} \left\{ \mathcal{E}(g) + \lambda \|g\|_{\mathcal{H}_{\infty}}^{2} \right\}, \tag{12}$$

and

$$g_{\infty,\lambda} = (L_{\infty} + \lambda I)^{-1} L_{\infty} g_{\rho}. \tag{13}$$

Next, let's consider the SGD algorithm in \mathcal{H}_M , which is commonly referred to as the *Reference SGD*. We will denote this algorithm as Algorithm 2, and its definition is as follows:

Algorithm 2 Reference SGD in \mathcal{H}_M

Input: Number of iterations: T, regularization parameter: λ , step sizes: $\{\eta_t\}_{t=1}^T$ **Output:** Estimator $g^{(T+1)}$

- 1: $q^{(1)} = 0$
- 2: for t = 1 to T do
- 3: Randomly draw a sample $z_t = (x_t, y_t) \sim \rho$
- 4: $q^{(t+1)} = (1 \eta_t \lambda) q^{(t)} \eta_t (q^{(t)}(x_t) y_t) k_{M,x_t}$
- 5: end for

Now, we can derive the following error decomposition

$$\|g_{\Theta^{(T+1)}} - g_{\rho}\|_{\rho}^{2} \le 4\left(\|g_{\Theta^{(T+1)}} - g^{(T+1)}\|_{\rho}^{2} + \|g^{(T+1)} - g_{M,\lambda}\|_{\rho}^{2} + \|g_{M,\lambda} - g_{\infty,\lambda}\|_{\rho}^{2} + \|g_{\infty,\lambda} - g_{\rho}\|_{\rho}^{2}\right). \tag{14}$$

The decomposition mentioned above has a highly intuitive interpretation.

- $\|g_{\Theta^{(T+1)}} g^{(T+1)}\|_{\rho}^2$: this term measures the similarity between the estimator obtained by the SGD algorithm in the neural network and that obtained by the SGD algorithm in \mathcal{H}_M , which is called *dynamics error*;
- $\|g^{(T+1)} g_{M,\lambda}\|_{\rho}^2$: this term is the convergence rate of the SGD algorithm in \mathcal{H}_M , which is called *convergence error*;
- $||g_{M,\lambda} g_{\infty,\lambda}||_{\rho}^2$: this term measures the similarity of the estimations of g_{ρ} in \mathcal{H}_M and \mathcal{H}_{∞} , which is called *random feature error*;
- $||g_{\infty,\lambda} g_{\rho}||_{\rho}^2$: this term measures the error between the estimation of g_{ρ} in \mathcal{H}_{∞} and the target function g_{ρ} , which is called *approximation error*.

Next, we will estimate these four errors separately and complete the proof of Theorem 1 and Corollary 1.

4.2 Estimation For Dynamics Error

The following proposition provides an estimation for the dynamics error, which states that for a two-layer neural network, as long as the network width is sufficiently large, the gap between the estimator acquired through the SGD algorithm in the neural network and that obtained by the SGD algorithm in \mathcal{H}_M can be sufficiently small.

Proposition 4.1. Suppose Assumptions 1 and 2 hold. For any $\lambda > 0$, let the step size be $\eta_t = \eta t^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and $0 < \eta < \frac{1-\theta}{5\kappa^2 + \lambda}$, when

$$M \ge M_1(T) = T^{6-5\theta}.$$

the following inequality holds

$$\left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\rho}^{2} \le \left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty}^{2} \le 25C_{\sigma}^{2}T^{-\theta}.$$

We will present the proof of Proposition 4.1 at the end of this subsection. In order to estimate the dynamics error $\|g_{\Theta^{(T+1)}} - g^{(T+1)}\|_{\rho}$, we introduce the following intermediate function $h_{\Theta}(x)$

$$h_{\Theta}(x) = \left\langle \nabla_{\Theta} g_{\Theta}(x) \Big|_{\Theta = \Theta^{(0)}}, \Theta - \Theta_{0} \right\rangle$$

$$= \frac{1}{\sqrt{M}} \sum_{r=1}^{M} \left(\left(a_{r} - a_{r}^{(0)} \right) \sigma \left(b_{r}^{(0) \top} x \right) + a_{r}^{(0)} \sigma' \left(b_{r}^{(0) \top} x \right) \left(b_{r} - b_{r}^{(0)} \right)^{\top} x \right), \tag{15}$$

which will be proved to be a good approximation of g_{Θ} , where $\Theta = (a_1, \dots, a_M, b_1^{\top}, \dots, b_M^{\top})^{\top} \in \mathbb{R}^{(d+1)M}$. Then we have the following error decomposition for the dynamic error

$$\left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\rho} \le \left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty} \le \left\| g_{\Theta^{(T+1)}} - h_{\Theta^{(T+1)}} \right\|_{\infty} + \left\| h_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty}, \tag{16}$$

here $||g||_{\infty} = \sup_{x \in \mathcal{X}} |g(x)|$. In the following, we will estimate the two terms on the right-hand side of (16). Before proceeding, let's establish the following bound for the network function g_{Θ} and its gradient.

Lemma 4.1. Suppose Assumptions 1 and 2 holds. For any Θ satisfying $\|\Theta - \Theta^{(0)}\|_2 \leq p$ for some p > 0, and $M \geq p^2$, there holds

$$\|g_{\Theta}\|_{\infty} \le 2(1+C_{\sigma})p, \quad \sup_{x \in \mathcal{X}} \left\|\nabla_{\Theta}g_{\Theta}(x) - \nabla_{\Theta}g_{\Theta}(x)\right\|_{\Theta=\Theta^{(0)}} \right\|_{2}^{2} \le \frac{3C_{\sigma}^{2}}{M}p^{2}.$$

Proof. By the definition of g_{Θ} , and recalling that $g_{\Theta^{(0)}}(x) = 0$, under Assumptions 1 and 2, we can derive

$$|g_{\Theta}(x)| = |g_{\Theta}(x) - g_{\Theta^{(0)}}(x)| = \left| \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r \sigma(b_r^{\top} x) - \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r^{(0)} \sigma(b_r^{(0)^{\top}} x) \right|$$
$$= \frac{1}{\sqrt{M}} \left| \sum_{r=1}^{M} \left[(a_r - a_r^{(0)}) \sigma(b_r^{(0)^{\top}} x) + (a_r - a_r^{(0)}) \left(\sigma(b_r^{\top} x) - \sigma(b_r^{(0)^{\top}} x) \right) \right|$$

$$+ a_{r}^{(0)} \left(\sigma(b_{r}^{\top}x) - \sigma(b_{r}^{(0)^{\top}}x) \right) \right]$$

$$\leq \frac{1}{\sqrt{M}} \sum_{r=1}^{M} \left[\left| a_{r} - a_{r}^{(0)} \right| \left(1 + \left\| b_{r}^{(0)} \right\|_{2} \|x\|_{2} \right) + C_{\sigma} |a_{r} - a_{r}^{(0)}| \left\| b_{r} - b_{r}^{(0)} \right\|_{2} \|x\|_{2} \right]$$

$$+ C_{\sigma} a_{r}^{(0)} \left\| b_{r} - b_{r}^{(0)} \right\|_{2} \|x\|_{2} \right]$$

$$\leq 2 \sqrt{\sum_{r=1}^{M} \left| a_{r} - a_{r}^{(0)} \right|^{2} + \frac{C_{\sigma}}{\sqrt{M}} \sqrt{\sum_{r=1}^{M} |a_{r} - a_{r}^{(0)}|^{2}} \sqrt{\sum_{r=1}^{M} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2}} + C_{\sigma} \sqrt{\sum_{r=1}^{M} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2}}$$

$$\leq 2 \left\| \Theta - \Theta^{(0)} \right\|_{2} + \frac{C_{\sigma}}{\sqrt{M}} \left\| \Theta - \Theta^{(0)} \right\|_{2}^{2} + C_{\sigma} \left\| \Theta - \Theta^{(0)} \right\|_{2}$$

$$\leq 2 (C_{\sigma} + 1)p,$$

where the first inequality holds due to the properties of the activation function $\sigma(\cdot)$ stated in Assumption 1, and the last inequality holds as a result of the conditions $M \geq p^2$ and $\|\Theta - \Theta^{(0)}\|_2 \leq p$.

And for the gradients, under Assumptions 1 and 2, we have

$$\begin{split} & \left\| \nabla_{\Theta} g_{\Theta}(x) - \nabla_{\Theta} g_{\Theta}(x) \right\|_{\Theta = \Theta^{(0)}}^{2} \right\|_{2}^{2} \\ &= \frac{1}{M} \sum_{r=1}^{M} \left[\left| \sigma(b_{r}^{\top} x) - \sigma(b_{r}^{(0)^{\top}} x) \right|^{2} + \left| a_{r} \sigma'(b_{r}^{\top} x) - a_{r}^{(0)} \sigma'(b_{r}^{(0)^{\top}} x) \right|^{2} \|x\|_{2}^{2} \right] \\ &= \frac{1}{M} \sum_{r=1}^{M} \left[\left| \sigma(b_{r}^{\top} x) - \sigma(b_{r}^{(0)^{\top}} x) \right|^{2} + \left| (a_{r} - a_{r}^{(0)}) \sigma'(b_{r}^{\top} x) + a_{r}^{(0)} \left(\sigma'(b_{r}^{\top} x) - \sigma'(b_{r}^{(0)^{\top}} x) \right) \right|^{2} \|x\|_{2}^{2} \right] \\ &\leq \frac{1}{M} \sum_{r=1}^{M} \left[C_{\sigma}^{2} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2} + 2C_{\sigma}^{2} \left| a_{r} - a_{r}^{(0)} \right|^{2} + 2C_{\sigma}^{2} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2} \right] \\ &\leq \frac{3C_{\sigma}^{2}}{M} \left\| \Theta - \Theta^{(0)} \right\|_{2}^{2} \\ &\leq \frac{3C_{\sigma}^{2}}{M} p^{2}, \end{split}$$

where the first inequality holds due to the properties of the activation function $\sigma(\cdot)$ stated in Assumption 1 and the fact that $||x||_2 \leq 1$. Then the proof is complete.

Next, we will prove that h_{Θ} is a good approximation of g_{Θ} .

Proposition 4.2. Suppose Assumptions 1 and 2 hold, for any Θ satisfying $\|\Theta - \Theta^{(0)}\|_2 \le p$ for some p > 0, we have

$$\|g_{\Theta} - h_{\Theta}\|_{\infty} \leq \frac{4C_{\sigma}}{\sqrt{M}}p^2.$$

Proof. Recall that $g_{\Theta^{(0)}}(x) = \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_r^{(0)} \sigma(b_r^{(0)^{\top}} x) = 0$ based on the definition of $g_{\Theta^{(0)}}$. Given the definitions of g_{Θ} and h_{Θ} , and assuming Assumption 1 holds for the activation function

 $\sigma(\cdot)$, for any $x \in \mathcal{X}$ and any Θ , we can deduce that

$$|g_{\Theta}(x) - h_{\Theta}(x)| = |g_{\Theta}(x) - g_{\Theta^{(0)}}(x) - h_{\Theta}(x)|$$

$$= \left| \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_{r} \sigma(b_{r}^{\top} x) - \frac{1}{\sqrt{M}} \sum_{r=1}^{M} a_{r}^{(0)} \sigma(b_{r}^{(0)^{\top}} x) - \frac{1}{\sqrt{M}} \sum_{r=1}^{M} \left(\left(a_{r} - a_{r}^{(0)} \right) \sigma\left(b_{r}^{(0)^{\top}} x \right) + a_{r}^{(0)} \sigma'\left(b_{r}^{(0)^{\top}} x \right) \left(b_{r} - b_{r}^{(0)} \right)^{\top} x \right) \right|$$

$$\leq \frac{1}{\sqrt{M}} \sum_{r=1}^{M} \left| \left(a_{r} - a_{r}^{(0)} \right) \left(\sigma\left(b_{r}^{\top} x \right) - \sigma(b_{r}^{(0)^{\top}} x) \right) + a_{r}^{(0)} \left(\sigma\left(b_{r}^{\top} x \right) - \sigma\left(b_{r}^{(0)^{\top}} x \right) - \sigma'\left(b_{r}^{(0)^{\top}} x \right) \left(b_{r} - b_{r}^{(0)} \right)^{\top} x \right) \right|$$

$$\leq \frac{1}{\sqrt{M}} \sum_{r=1}^{M} \left[2C_{\sigma} \left| a_{r} - a_{r}^{(0)} \right| \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2} + 2C_{\sigma} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2} \right]$$

$$\leq \frac{2C_{\sigma}}{\sqrt{M}} \sqrt{\sum_{r=1}^{M} \left| a_{r} - a_{r}^{(0)} \right|^{2}} \sqrt{\sum_{r=1}^{M} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2} + \frac{2C_{\sigma}}{\sqrt{M}} \sum_{r=1}^{M} \left\| b_{r} - b_{r}^{(0)} \right\|_{2}^{2}}$$

$$\leq \frac{4C_{\sigma}}{\sqrt{M}} \left\| \Theta - \Theta^{(0)} \right\|_{2}^{2}.$$

Then the desired result holds by $\|\Theta - \Theta^{(0)}\|_2 \le p$.

Now, we are prepared to establish a bound for $\|\Theta^{(t+1)} - \Theta^{(0)}\|_2$ for $t \in \mathbb{N}_T$, where $\mathbb{N}_T := \{1, \dots, T\}$.

Proposition 4.3. Let $\{\Theta^{(t)}\}_{t\geq 1}$ be defined by Algorithm 1. Assume that Assumptions 1 and 2 hold, set $\eta_i = \eta i^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda}$, and $M \geq T^{4(1-\theta)}$, then for $t \in \mathbb{N}_T$, we have

$$\left\|\Theta^{(t+1)} - \Theta^{(0)}\right\|_{2} \le t^{1-\theta}.$$
 (17)

Proof. We prove the lemma by induction. First, for t=1, since $\Theta^{(1)}=\Theta^{(0)}$ and $g_{\Theta^{(0)}}(x)=0$, we have

$$\begin{split} \left\| \Theta^{(2)} - \Theta^{(0)} \right\|_{2} &= \left\| (1 - \eta_{1} \lambda) (\Theta^{(1)} - \Theta^{(0)}) - \eta_{1} (g_{\Theta^{(1)}}(x_{1}) - y_{1}) \nabla_{\Theta} g_{\Theta}(x_{1}) \right|_{\Theta = \Theta^{(1)}} \right\|_{2} \\ &= \left\| 0 - \eta_{1} (0 - y_{1}) \nabla_{\Theta} g_{\Theta}(x_{1}) \right|_{\Theta = \Theta^{(1)}} \right\|_{2} \\ &= \eta_{1} |y_{1}| \left\| \nabla_{\Theta} g_{\Theta}(x_{1}) \right|_{\Theta = \Theta^{(1)}} \right\|_{2}. \end{split}$$

For $\|\nabla_{\Theta}g_{\Theta}(x_1)\|_{\Theta=\Theta^{(1)}}\|_2$, under Assumptions 1 and 2, we have

$$\begin{split} \left\| \nabla_{\Theta} g_{\Theta}(x_1) \right\|_{\Theta = \Theta^{(1)}} \right\|_2^2 &= \frac{1}{M} \sum_{r=1}^M \left[\left(\sigma(b_r^{(1)^{\top}} x_1) \right)^2 + \left(a_r^{(1)} \sigma' \left(b_r^{(1)^{\top}} x_1 \right) \right)^2 \|x_1\|_2^2 \right] \\ &\leq \frac{1}{M} \sum_{r=1}^M \left[\left(1 + |b_r^{(1)^{\top}} x_1| \right)^2 + C_{\sigma}^2 \right] \leq 4 + C_{\sigma}^2 \leq \kappa^2. \end{split}$$

This, combined with the conditions $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda} < \frac{1}{\kappa^2}$ and $|y_1| \le 1$, implies that

$$\left\| \Theta^{(2)} - \Theta^{(0)} \right\|_2 = \eta_1 |y_1| \left\| \nabla_{\Theta} g_{\Theta}(x_1) \right|_{\Theta = \Theta^{(1)}} \left\|_2 \le \eta \kappa < \kappa \frac{1 - \theta}{4\kappa^2 + \lambda} < \frac{1}{\kappa} \le 1,$$

which means that (17) holds for t = 1.

As the induction assumption, suppose (17) holds for all $t \le k \le T - 1$. In order to advance the induction, it is necessary to make an estimation for $\|\Theta^{(k+2)} - \Theta^{(0)}\|_2$, i.e., for t = k + 1. By the iteration equation of $\Theta^{(k+2)}$, we have

$$\begin{split} \Theta^{(k+2)} - \Theta^{(0)} &= (1 - \eta_{k+1} \lambda) (\Theta^{(k+1)} - \Theta^{(0)}) - \eta_{k+1} (g_{\Theta^{(k+1)}}(x_{k+1}) - y_{k+1}) \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(k+1)}} \\ &= (1 - \eta_{k+1} \lambda) (\Theta^{(k+1)} - \Theta^{(0)}) - \eta_{k+1} (g_{\Theta^{(k+1)}}(x_{k+1}) - y_{k+1}) \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \\ &- \eta_{k+1} (g_{\Theta^{(k+1)}}(x_{k+1}) - y_{k+1}) \left(\nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(k+1)}} - \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \right) \\ &= (1 - \eta_{k+1} \lambda) (\Theta^{(k+1)} - \Theta^{(0)}) - \eta_{k+1} h_{\Theta^{(k+1)}}(x_{k+1}) \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \\ &- \eta_{k+1} (g_{\Theta^{(k+1)}}(x_{k+1}) - h_{\Theta^{(k+1)}}(x_{k+1})) \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \\ &- \eta_{k+1} (g_{\Theta^{(k+1)}}(x_{k+1}) - y_{k+1}) \left(\nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(k+1)}} - \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \right) \\ &+ \eta_{k+1} y_{k+1} \nabla_{\Theta} g_{\Theta}(x_{k+1}) \Big|_{\Theta = \Theta^{(0)}} \\ &= (I - \eta_{k+1} (A_{k+1} + \lambda I)) (\Theta^{(k+1)} - \Theta^{(0)}) - \eta_{k+1} \alpha_{k+1}^{(1)} - \eta_{k+1} \alpha_{k+1}^{(2)} + \eta_{k+1} \alpha_{k+1}^{(3)} \\ &= - \sum_{i=1}^{k+1} \eta_i \prod_{j=i+1}^{k+1} (I - \eta_j (A_j + \lambda I)) (\alpha_i^{(1)} + \alpha_i^{(2)} - \alpha_i^{(3)}), \end{split}$$

where

$$A_{i} = \nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(0)}} \left(\nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(0)}} \right)^{T} \in \mathbb{R}^{[(d+1)M] \times [(d+1)M]},$$

$$\alpha_{i}^{(1)} = \left(g_{\Theta^{(i)}}(x_{i}) - h_{\Theta^{(i)}}(x_{i}) \right) \nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(0)}},$$

$$\alpha_{i}^{(2)} = \left(g_{\Theta^{(i)}}(x_{i}) - y_{i} \right) \left(\nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(i)}} - \nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(0)}} \right),$$

$$\alpha_{i}^{(3)} = y_{i} \nabla_{\Theta} g_{\Theta}(x_{i}) \Big|_{\Theta = \Theta^{(0)}}.$$

Since $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda} < \frac{1}{\kappa^2 + \lambda}$, we can establish the following decomposition

$$\left\| \Theta^{(k+2)} - \Theta^{(0)} \right\|_{2} = \left\| \sum_{i=1}^{k+1} \eta_{i} \prod_{j=i+1}^{k+1} \left(I - \eta_{j} \left(A_{j} + \lambda I \right) \right) \left(\alpha_{i}^{(1)} + \alpha_{i}^{(2)} - \alpha_{i}^{(3)} \right) \right\|_{2}$$

$$\leq \left\| \sum_{i=1}^{k+1} \eta_{i} \prod_{j=i+1}^{k+1} \left(I - \eta_{j} \left(A_{j} + \lambda I \right) \right) \alpha_{i}^{(1)} \right\|_{2} + \left\| \sum_{i=1}^{k+1} \eta_{i} \prod_{j=i+1}^{k+1} \left(I - \eta_{j} \left(A_{j} + \lambda I \right) \right) \alpha_{i}^{(2)} \right\|_{2}$$

$$+ \left\| \sum_{i=1}^{k+1} \eta_{i} \prod_{j=i+1}^{k+1} \left(I - \eta_{j} \left(A_{j} + \lambda I \right) \right) \alpha_{i}^{(3)} \right\|_{2}$$

$$\leq \sum_{i=1}^{k+1} \eta_{i} \left\| \alpha_{i}^{(1)} \right\|_{2} + \sum_{i=1}^{k+1} \eta_{i} \left\| \alpha_{i}^{(2)} \right\|_{2} + \sum_{i=1}^{k+1} \eta_{i} \left\| \alpha_{i}^{(3)} \right\|_{2}.$$

$$(18)$$

For the first term $\sum_{i=1}^{k+1} \eta_i \left\| \alpha_i^{(1)} \right\|_2$, by $\sup_{x \in \mathcal{X}} \left\| \nabla_{\Theta} g_{\Theta}(x) \right\|_{\Theta = \Theta^{(0)}} \right\|_2^2 = \sup_{x \in \mathcal{X}} k_M(x, x) \leq \kappa^2$, Proposition 4.2 and the induction assumption, we have

$$\begin{split} \sum_{i=1}^{k+1} \eta_i \left\| \alpha_i^{(1)} \right\|_2 &= \sum_{i=1}^{k+1} \eta_i \left\| (g_{\Theta^{(i)}}(x_i) - h_{\Theta^{(i)}}(x_i)) \nabla_{\Theta} g_{\Theta}(x_i) \right|_{\Theta = \Theta^{(0)}} \right\|_2 \\ &\leq \sum_{i=1}^{k+1} \eta_i \left\| g_{\Theta^{(i)}} - h_{\Theta^{(i)}} \right\|_{\infty} \left\| \nabla_{\Theta} g_{\Theta}(x_i) \right|_{\Theta = \Theta^{(0)}} \right\|_2 \\ &\leq \kappa \sum_{i=1}^{k+1} \eta_i \frac{4C_{\sigma}}{\sqrt{M}} (i-1)^{2(1-\theta)} \\ &\leq \frac{4C_{\sigma} \kappa \eta}{\sqrt{M}} \sum_{i=1}^{k+1} i^{2-3\theta} \\ &\leq \frac{4C_{\sigma} \kappa \eta}{3(1-\theta)} \frac{(k+1)^{3(1-\theta)}}{\sqrt{M}}. \end{split}$$

Employing $M \geq T^{4(1-\theta)}$, $\kappa^2 \geq 2C_\sigma^2$ and $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda} < \frac{1}{4C_\sigma\kappa}$, there holds

$$\sum_{i=1}^{k+1} \eta_i \left\| \alpha_i^{(1)} \right\|_2 \le \frac{1}{3} \frac{T^{2(1-\theta)}}{\sqrt{M}} (k+1)^{1-\theta} < \frac{1}{3} (k+1)^{1-\theta}.$$

For the second term $\sum_{i=1}^{k+1} \eta_i \|\alpha_i^{(2)}\|_2$, we can use the induction assumption along with Lemma 4.1 to derive

$$\begin{split} \sum_{i=1}^{k+1} \eta_{i} \left\| \alpha_{i}^{(2)} \right\|_{2} &= \sum_{i=1}^{k+1} \eta_{i} \left\| (g_{\Theta^{(i)}}(x_{i}) - y_{i}) \left(\nabla_{\Theta} g_{\Theta}(x_{i}) \right|_{\Theta = \Theta^{(i)}} - \nabla_{\Theta} g_{\Theta}(x_{i}) \right|_{\Theta = \Theta^{(0)}} \right) \right\|_{2} \\ &\leq \sum_{i=1}^{k+1} \eta_{i} (1 + \left\| g_{\Theta^{(i)}} \right\|_{\infty}) \sup_{x \in \mathcal{X}} \left\| \nabla_{\Theta} g_{\Theta}(x) \right|_{\Theta = \Theta^{(i)}} - \nabla_{\Theta} g_{\Theta}(x) \Big|_{\Theta = \Theta^{(0)}} \right\|_{2} \\ &\leq \sum_{i=1}^{k+1} \eta_{i} [1 + 2(1 + C_{\sigma})(i - 1)^{1 - \theta}] \cdot \frac{\sqrt{3}C_{\sigma}}{\sqrt{M}} (i - 1)^{1 - \theta} \\ &\leq \sum_{i=1}^{k+1} \eta_{i} [1 + 2(1 + C_{\sigma})] i^{1 - \theta} \cdot \frac{\sqrt{3}C_{\sigma}}{\sqrt{M}} i^{1 - \theta} \\ &\leq \frac{\eta \sqrt{3}C_{\sigma}(2C_{\sigma} + 3)}{3(1 - \theta)} \frac{(k + 1)^{3(1 - \theta)}}{\sqrt{M}} \\ &\leq \frac{1}{3} (k + 1)^{1 - \theta}, \end{split}$$

where the last inequality holds due to $M \ge T^{4(1-\theta)}$ and $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda} < \frac{1}{\sqrt{3}C_{\sigma}(2C_{\sigma}+3)}$. For the last term $\sum_{i=1}^{k+1} \eta_i \|\alpha_i^{(3)}\|_2$, we have

$$\sum_{i=1}^{k+1} \eta_i \left\| \alpha_i^{(3)} \right\|_2 = \sum_{i=1}^{k+1} \eta_i \left\| y_i \nabla_{\Theta} g_{\Theta}(x_i) \right|_{\Theta = \Theta^{(0)}} \left\|_2 \le \kappa \sum_{i=1}^{k+1} \eta_i \le \frac{\eta \kappa}{1 - \theta} (k+1)^{1-\theta} \le \frac{1}{3} (k+1)^{1-\theta},$$

where we use the condition that $0 < \eta < \frac{1-\theta}{4\kappa^2 + \lambda} < \frac{1}{3} \frac{1-\theta}{\kappa}$ for the last inequality.

Putting all these three estimates back into (18) yields that

$$\left\| \Theta^{(k+2)} - \Theta^{(0)} \right\|_2 \le (k+1)^{1-\theta}.$$

Hence, by mathematical induction, we have proven that (17) holds for all $t \in \mathbb{N}_T$. This completes the proof.

Now, let's estimate the second term $\|h_{\Theta^{(T+1)}} - g^{(T+1)}\|_{\infty}$ of the dynamic error in (16).

Proposition 4.4. Suppose Assumptions 1 and 2 hold. For any $\lambda > 0$ and $i \in \mathbb{N}_+$, let $\eta_i = \eta i^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and $0 < \eta < \frac{1}{\kappa^2 + \lambda}$, when

$$M > T^{6-5\theta}.$$

the following inequality holds

$$\left\| h_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty} \le \frac{5\kappa^2 \eta C_{\sigma}}{1 - \theta} T^{-\frac{\theta}{2}}.$$

Proof. By the definitions of $h_{\Theta^{(T+1)}},\,g^{(T+1)}$ and $\Theta^{(T+1)},$ we have

$$\begin{split} h_{\Theta^{(T+1)}} - g^{(T+1)} &= \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \Theta^{(T+1)} - \Theta^{(0)} \right\rangle \\ &- \left((1 - \eta_T \lambda) g^{(T)} - \eta_T \left(g^{(T)}(x_T) - y_T \right) k_{M,x_T} \right) \\ &= (1 - \eta_T \lambda) h_{\Theta^{(T)}} - \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(T)}} \right\rangle \\ &- \left((1 - \eta_T \lambda) g^{(T)} - \eta_T \left(g^{(T)}(x_T) - y_T \right) k_{M,x_T} \right) \\ &= (1 - \eta_T \lambda) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(T)}} \right\rangle \\ &+ \eta_T (g^{(T)}(x_T) - y_T) k_{M,x_T} \\ &= (1 - \eta_T \lambda) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(T)}} \right\rangle \\ &+ \eta_T (g^{(T)}(x_T) - h_{\Theta^{(T)}}(x_T)) k_{M,x_T} + \eta_T (h_{\Theta^{(T)}}(x_T) - y_T) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) \\ &- \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(0)}} \right\rangle + \eta_T (h_{\Theta^{(T)}}(x_T) - y_T) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) \\ &- \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(T)}} - \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(0)}} \right\rangle \\ &- \eta_T (g_{\Theta^{(T)}}(x_T) - y_T) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(T)}} - \nabla_{\Theta} g_{\Theta}(x_T) \Big|_{\Theta=\Theta^{(0)}} \right\rangle \\ &- \eta_T (g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x_T)) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T \left(g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x_T) \right) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T \left(g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x_T) \right) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T \left(g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x_T) \right) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T \left(g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x_T) \right) k_{M,x_T} \\ &= (I - \eta_T (k_{M,x_T} \otimes k_{M,x_T} + \lambda I)) \left(h_{\Theta^{(T)}} - g^{(T)} \right) - \eta_T \left(g_{\Theta^{(T)}}(x_T) - h_{\Theta^{(T)}}(x$$

where

$$\beta_t^{(1)} = \left(g_{\Theta^{(t)}}(x_t) - y_t\right) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta = \Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_t) \Big|_{\Theta = \Theta^{(t)}} - \nabla_{\Theta} g_{\Theta}(x_t) \Big|_{\Theta = \Theta^{(0)}} \right\rangle,$$

$$\beta_t^{(2)} = \left(g_{\Theta^{(t)}}(x_t) - h_{\Theta^{(t)}}(x_t)\right) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \Big|_{\Theta = \Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_t) \Big|_{\Theta = \Theta^{(0)}} \right\rangle.$$

Then we can proceed with the following error decomposition

$$\begin{split} & \left\| h_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty} = \left\| \sum_{t=1}^{T} \eta_{t} \prod_{i=t+1}^{T} \left(I - \eta_{i} (k_{M,x_{i}} \otimes k_{M,x_{i}} + \lambda I) \right) \left(\beta_{t}^{(1)} + \beta_{t}^{(2)} \right) \right\|_{\infty} \\ & \leq \left\| \sum_{t=1}^{T} \eta_{t} \prod_{i=t+1}^{T} \left(I - \eta_{i} (k_{M,x_{i}} \otimes k_{M,x_{i}} + \lambda I) \right) \beta_{t}^{(1)} \right\|_{\infty} + \left\| \sum_{t=1}^{T} \eta_{t} \prod_{i=t+1}^{T} \left(I - \eta_{i} (k_{M,x_{i}} \otimes k_{M,x_{i}} + \lambda I) \right) \beta_{t}^{(2)} \right\|_{\infty} \\ & \leq \sum_{t=1}^{T} \eta_{t} \left\| \beta_{t}^{(1)} \right\|_{\infty} + \sum_{t=1}^{T} \eta_{t} \left\| \beta_{t}^{(2)} \right\|_{\infty}. \end{split}$$

For the first term $\sum_{t=1}^{T} \eta_t \left\| \beta_t^{(1)} \right\|_{\infty}$, applying Lemma 4.1 to the bounds of $\sup_{x \in \mathcal{X}} \left\| \nabla_{\Theta} g_{\Theta}(x) \right\|_{\Theta = \Theta^{(t)}} - \nabla_{\Theta} g_{\Theta}(x) \Big|_{\Theta = \Theta^{(0)}} \|_{2}$ and $\left\| g_{\Theta^{(t)}} \right\|_{\infty}$, we can derive the following result

$$\begin{split} \sum_{t=1}^{T} \eta_{t} \left\| \beta_{t}^{(1)} \right\|_{\infty} &\leq \sum_{t=1}^{T} \eta_{t} \left\| \left(g_{\Theta^{(t)}}(x_{t}) - y_{t} \right) \left\langle \nabla_{\Theta} g_{\Theta}(\cdot) \right|_{\Theta = \Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_{t}) \Big|_{\Theta = \Theta^{(t)}} - \nabla_{\Theta} g_{\Theta}(x_{t}) \Big|_{\Theta = \Theta^{(0)}} \right\rangle \right\|_{\infty} \\ &\leq \kappa \sum_{t=1}^{T} \eta_{t} \left(1 + \left\| g_{\Theta^{(t)}} \right\|_{\infty} \right) \cdot \sup_{x \in \mathcal{X}} \left\| \nabla_{\Theta} g_{\Theta}(x) \right|_{\Theta = \Theta^{(t)}} - \nabla_{\Theta} g_{\Theta}(x) \Big|_{\Theta = \Theta^{(0)}} \right\|_{2} \\ &\leq \sqrt{3} \eta C_{\sigma} \kappa \left(1 + 2(1 + C_{\sigma}) \right) \frac{1}{\sqrt{M}} \sum_{t=1}^{T} t^{-\theta} t^{2(1-\theta)} \\ &\leq \frac{\sqrt{3} \eta C_{\sigma} \kappa \left(1 + 2(1 + C_{\sigma}) \right)}{3(1 - \theta)} \frac{T^{3(1-\theta)}}{\sqrt{M}} \\ &\leq \frac{5\sqrt{3} \kappa^{2} \eta C_{\sigma}}{3(1 - \theta)} T^{-\frac{\theta}{2}}, \end{split}$$

where the last inequality holds due to $M \ge T^{6-5\theta}$ and $\kappa^2 \ge C_\sigma^2$.

For the second term $\sum_{t=1}^{T} \eta_t \|\beta_t^{(2)}\|_{\infty}$, we can derive an upper bound for $\|g_{\Theta^{(t)}} - h_{\Theta^{(t)}}\|_{\infty}$ for $t \geq 1$ by utilizing Proposition 4.2. Consequently, we obtain the following bound

$$\sum_{t=1}^{T} \eta_{t} \|\beta_{t}^{(2)}\|_{\infty} \leq \sum_{t=1}^{T} \eta_{t} \|(g_{\Theta^{(t)}}(x_{t}) - h_{\Theta^{(t)}}(x_{t})) \langle \nabla_{\Theta} g_{\Theta}(\cdot) |_{\Theta=\Theta^{(0)}}, \nabla_{\Theta} g_{\Theta}(x_{t}) |_{\Theta=\Theta^{(0)}} \rangle \|_{\infty}$$

$$\leq \kappa^{2} \sum_{t=1}^{T} \eta_{t} \|g_{\Theta^{(t)}} - h_{\Theta^{(t)}}\|_{\infty}$$

$$\leq 4\kappa^{2} \eta C_{\sigma} \frac{1}{\sqrt{M}} \sum_{t=1}^{T} t^{2-3\theta}$$

$$\leq \frac{4\kappa^2 \eta C_{\sigma}}{3(1-\theta)} \frac{T^{3(1-\theta)}}{\sqrt{M}}$$
$$\leq \frac{4\kappa^2 \eta C_{\sigma}}{3(1-\theta)} T^{-\frac{\theta}{2}},$$

where the last inequality holds due to $M \geq T^{6-5\theta}$. Combining the previously established two bounds, we can conclude that

$$\left\| h_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\infty} \le \frac{(4 + 5\sqrt{3})\kappa^2 \eta C_{\sigma}}{3(1 - \theta)} T^{-\frac{\theta}{2}} \le \frac{5\kappa^2 \eta C_{\sigma}}{1 - \theta} T^{-\frac{\theta}{2}},$$

which finishes the proof.

Now we are ready to prove Proposition 4.1.

Proof of Proposition 4.1. Under Assumptions 1, 2, and $0 < \eta < \frac{1-\theta}{5\kappa^2 + \lambda} < \frac{1-\theta}{4\kappa^2 + \lambda}$, we can combine Proposition 4.2 and Proposition 4.3 to derive the following result

$$\|g_{\Theta^{(T+1)}} - h_{\Theta^{(T+1)}}\|_{\infty} \le \frac{4C_{\sigma}}{\sqrt{M}} T^{2(1-\theta)} \le 4C_{\sigma} T^{-\frac{\theta}{2}},$$
 (19)

where the last inequality holds when $M \geq T^{4-3\theta}$. Thus, when $M \geq \max\left\{T^{4(1-\theta)}, T^{4-3\theta}, T^{6-5\theta}\right\}$ = $T^{6-5\theta}$, we can substitute the bounds from (19) and Proposition 4.4 back into (16). By using the inequality $\frac{5\eta\kappa^2}{1-\theta} < \frac{5\kappa^2}{1-\theta} \cdot \frac{1-\theta}{5\kappa^2+\lambda} < 1$, we can derive the desired result.

4.3 Estimation For Convergence Error

In this part, we will analyze the convergence rate of Reference SGD, denoted as Algorithm 2, in the RKHS \mathcal{H}_M generated by the kernel k_M .

Proposition 4.5. Suppose Assumptions 1, 2 and 3 hold. For $T \in \mathbb{Z}^+$, $\delta \in (0,1)$, $\lambda > 0$, we consider a step size sequence defined as $\eta_i = \eta i^{-\theta}$ for $i \in \mathbb{N}_+$, where $\frac{1}{2} < \theta < 1$ and $0 < \eta < \left(\lambda + \frac{208(1+\kappa)^4}{(1-2^{\theta-1})(2\theta-1)} \left(\log \frac{8}{1-\theta} + \frac{1}{1-\theta}\right)\right)^{-1}$. When

$$M \ge M_2(T, \lambda) = \max \left\{ T^{2(1+r)(1-\theta)}, T^{2\theta}, \frac{1}{\lambda^2} \right\},$$

the following holds with high probability at least $1-\delta$ over the random choice of $\Theta^{(0)}$

(a) if
$$\frac{1}{2} < \theta < \frac{2\beta - 1}{3\beta - 1}$$
,

$$\mathbb{E}_{Z^T} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^2 \right] \le \left(V_1 T^{-2r(1-\theta)} + V_2 \left(\exp\left\{ -H(\theta) T^{1-\frac{3\beta-1}{2\beta-1}\theta} \right\} + T^{-\theta} \right) \right) \log \frac{2}{\delta},$$

(b) if
$$\frac{2\beta-1}{3\beta-1} \le \theta < 1$$
,

$$\mathbb{E}_{Z^T} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^2 \right] \le \left(V_1 T^{-2r(1-\theta)} + V_2 \left(\exp\left\{ -H(\theta)\lambda T^{1-\theta} \right\} + T^{-\theta} \right) \right) \log \frac{2}{\delta},$$

where

$$V_{1} = 2 \left[\frac{\eta \kappa^{2} \|g_{\rho}\|_{\rho}}{1 - \theta} + \left(\frac{r(1 - \theta)}{e\eta(1 - 2^{\theta - 1})} \right)^{r} \|L_{\infty}^{-r} g_{\rho}\|_{\rho} \right]^{2},$$

$$V_{2} = 2W_{2} \left(\frac{8\kappa^{4}\eta^{2}\theta}{2\theta - 1} + \frac{2\eta^{2}\theta}{2\theta - 1} \exp\left\{ 2H(\theta)\kappa^{2}\sqrt{2\log\frac{2}{\delta}} \right\} \frac{2\beta c^{2}}{2\beta - 1} + 6\kappa^{4}\eta \right),$$

$$W_{2} = 2 \left[1 + 12\kappa^{4r + 2} \|L_{\infty}^{-r} g_{\rho}\|_{\rho}^{2} + \frac{8\kappa^{4}\eta^{2}\theta}{2\theta - 1} \left(20\|g_{\rho}\|_{\rho}^{2} + 3\mathcal{E}\left(g_{\rho}\right) \right) \right],$$

$$H(\theta) = \frac{2\eta}{1 - \theta} 2^{1 - \theta} \left(1 - \left(\frac{3}{4} \right)^{1 - \theta} \right).$$

To prove Proposition 4.5, we begin by presenting the following error decomposition for the convergence error.

Lemma 4.2. Let $\{g^{(t)}\}$ and $g_{M,\lambda}$ be defined by Algorithm 2 and (10) respectively, then

$$\mathbb{E}_{Z^{T}} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^{2} \right] = \left\| \prod_{t=1}^{T} \left(I - \eta_{t} \left(L_{M} + \lambda I \right) \right) g_{M,\lambda} \right\|_{\rho}^{2} + \sum_{t=1}^{T} \eta_{t}^{2} \mathbb{E}_{Z^{t}} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \right],$$
(20)

where $\mathcal{B}_t = (y_t - g^{(t)}(x_t))k_{M,x_t} + L_M(g^{(t)} - g_\rho).$

Proof. Firstly, from Algorithm 2 and the expression (11) of $g_{M,\lambda}$, we get the following recursion

$$g^{(t+1)} - g_{M,\lambda} = (I - \eta_t (L_M + \lambda I))(g^{(t)} - g_{M,\lambda}) + \eta_t \mathcal{B}_t$$

$$= -\prod_{t=1}^T (I - \eta_t (L_M + \lambda I)) g_{M,\lambda} + \sum_{t=1}^T \eta_t \prod_{i=t+1}^T (I - \eta_i (L_M + \lambda I)) \mathcal{B}_t$$
(21)

where
$$\mathcal{B}_t = (y_t - g^{(t)}(x_t))k_{M,x_t} + L_M(g^{(t)} - g_\rho)$$
, and
$$\mathbb{E}_{z_t}[\mathcal{B}_t|z_1, \cdots, z_{t-1}, z_{t+1}, \cdots, z_T]$$

$$= \mathbb{E}_{z_t}\left[(y_t k_{M,x_t} - L_M g_\rho) + (L_M - k_{M,x_t} \otimes k_{M,x_t})g^{(t)} \middle| z_1, \cdots, z_{t-1}, z_{t+1}, \cdots, z_T \right]$$

$$= 0.$$

Then, it follows that

$$\begin{split} & \mathbb{E}_{Z^T} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^2 \right] \\ &= \mathbb{E}_{Z^T} \left[\left\| -\prod_{t=1}^T \left(I - \eta_t \left(L_M + \lambda I \right) \right) g_{M,\lambda} + \sum_{t=1}^T \eta_t \prod_{i=t+1}^T \left(I - \eta_i \left(L_M + \lambda I \right) \right) \mathcal{B}_t \right\|_{\rho}^2 \right] \\ &= \left\| \prod_{t=1}^T \left(I - \eta_t \left(L_M + \lambda I \right) \right) g_{M,\lambda} \right\|_{\rho}^2 + \mathbb{E}_{Z^T} \left[\left\| \sum_{t=1}^T \eta_t \prod_{i=t+1}^T \left(I - \eta_i \left(L_M + \lambda I \right) \right) \mathcal{B}_t \right\|_{\rho}^2 \right] \end{split}$$

$$-2\mathbb{E}_{Z^{T}}\left\langle \prod_{t=1}^{T}\left(I-\eta_{t}\left(L_{M}+\lambda I\right)\right)g_{M,\lambda},\sum_{t=1}^{T}\eta_{t}\prod_{i=t+1}^{T}\left(I-\eta_{i}\left(L_{M}+\lambda I\right)\right)\mathcal{B}_{t}\right\rangle _{\rho}$$

$$=\left\|\prod_{t=1}^{T}\left(I-\eta_{t}\left(L_{M}+\lambda I\right)\right)g_{M,\lambda}\right\|_{\rho}^{2}+\mathbb{E}_{Z^{T}}\left[\left\|\sum_{t=1}^{T}\eta_{t}\prod_{i=t+1}^{T}\left(I-\eta_{i}\left(L_{M}+\lambda I\right)\right)\mathcal{B}_{t}\right\|_{\rho}^{2}\right].$$

Furthermore, for the second term on the right-hand side of the above equation, we have

$$\mathbb{E}_{Z^{T}} \left[\left\| \sum_{t=1}^{T} \eta_{t} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \right] = \sum_{t=1}^{T} \eta_{t}^{2} \mathbb{E}_{Z^{t}} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \right]$$

$$+ \sum_{t=1}^{T} \sum_{s \neq t} \eta_{t} \eta_{s} \mathbb{E}_{Z^{T}} \left\langle \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t}, \prod_{i=s+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{s} \right\rangle_{\rho}$$

$$= \sum_{t=1}^{T} \eta_{t}^{2} \mathbb{E}_{Z^{t}} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \right].$$

This completes the proof.

Our next goal is to estimate the two terms in Lemma 4.2. To achieve this, we will utilize Proposition 4.6 to estimate the first term and Proposition 4.7 to estimate the second term.

Proposition 4.6. Under the assumptions of Proposition 4.5, the following inequality holds with probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$

$$\left\| \prod_{t=1}^{T} \left(I - \eta_t \left(L_M + \lambda I \right) \right) g_{M,\lambda} \right\|_{\rho}^{2} \le V_1 T^{-2r(1-\theta)} \log \frac{2}{\delta}, \tag{22}$$

where the constant V_1 is defined in Proposition 4.5.

Proof. From the expression (11) of $g_{M,\lambda}$, which is given by $g_{M,\lambda} = (L_M + \lambda I)^{-1} L_M g_{\rho}$, we can obtain

$$\left\| \prod_{t=1}^{T} (I - \eta_{t} (L_{M} + \lambda I)) g_{M,\lambda} \right\|_{\rho}^{2} = \left\| \prod_{t=1}^{T} (I - \eta_{t} (L_{M} + \lambda I)) (L_{M} + \lambda I)^{-1} L_{M} g_{\rho} \right\|_{\rho}^{2}$$

$$\leq \left\| \prod_{t=1}^{T} (I - \eta_{t} (L_{M} + \lambda I)) g_{\rho} \right\|_{\rho}^{2}.$$

If we define

$$h_M^{(t+1)} = (I - \eta_t(L_M + \lambda I))h_M^{(t)}, h_{\infty}^{(t+1)} = (I - \eta_t(L_{\infty} + \lambda I))h_{\infty}^{(t)},$$

with $h_M^{(0)} = h_\infty^{(0)} = g_\rho$, then we have $\left\|h_M^{(T+1)}\right\|_\rho^2 = \left\|\prod_{t=1}^T \left(I - \eta_t \left(L_M + \lambda I\right)\right) g_\rho\right\|_\rho^2$. Next, we need to bound $\left\|h_M^{(T+1)}\right\|_\rho$, to this end, we divide the term $\left\|h_M^{(T+1)}\right\|_\rho$ into the following two terms,

$$\left\| h_M^{(T+1)} \right\|_{\rho} \le \left\| h_M^{(T+1)} - h_{\infty}^{(T+1)} \right\|_{\rho} + \left\| h_{\infty}^{(T+1)} \right\|_{\rho}. \tag{23}$$

In the following, we will estimate the above two terms respectively.

(i) **Step 1** Bound $\|h_M^{(T+1)} - h_\infty^{(T+1)}\|_{\rho}$. By the definition of $h_M^{(T+1)}$ and $h_\infty^{(T+1)}$, we have $h_M^{(T+1)} - h_\infty^{(T+1)} = (I - \eta_T (L_M + \lambda I)) h_M^{(T)} - (I - \eta_T (L_\infty + \lambda I)) h_\infty^{(T)}$ $= (I - \eta_T (L_\infty + \lambda I)) (h_M^{(T)} - h_\infty^{(T)}) - \eta_T (L_M - L_\infty) h_M^{(T)}$ $= -\sum_{t=1}^T \eta_t \prod_{i=t+1}^T (I - \eta_i (L_\infty + \lambda I)) (L_M - L_\infty) h_M^{(t)}.$

Then by applying Lemma A.1 and Lemma A.2, we can assert that

$$\begin{split} \left\| h_{M}^{(T+1)} - h_{\infty}^{(T+1)} \right\|_{\rho} &= \left\| \sum_{t=1}^{T} \eta_{t} \prod_{i=t+1}^{T} (I - \eta_{i} (L_{\infty} + \lambda I)) (L_{M} - L_{\infty}) h_{M}^{(t)} \right\|_{\rho} \\ &\leq \sum_{t=1}^{T} \eta_{t} \left\| \prod_{i=t+1}^{T} (I - \eta_{i} (L_{\infty} + \lambda I)) \right\|_{op} \left\| L_{M} - L_{\infty} \right\|_{op} \left\| h_{M}^{(t)} \right\|_{\rho} \\ &\leq \sum_{t=1}^{T} \eta_{t} \left\| L_{M} - L_{\infty} \right\|_{op} \left\| \prod_{i=1}^{t-1} (I - \eta_{t} (L_{M} + \lambda I)) g_{\rho} \right\|_{\rho} \\ &\leq \left\| L_{M} - L_{\infty} \right\|_{op} \left\| g_{\rho} \right\|_{\rho} \sum_{t=1}^{T} \eta_{t} \\ &\leq \frac{\eta \kappa^{2} \|g_{\rho}\|_{\rho}}{1 - \theta} \frac{T^{1-\theta}}{\sqrt{M}} \sqrt{2 \log \frac{2}{\delta}}, \end{split}$$

holds with confidence at least $1 - \delta$. Moreover, when $M \ge T^{2(1+r)(1-\theta)}$

$$\left\| h_M^{(T+1)} - h_{\infty}^{(T+1)} \right\|_{\rho} \le \frac{\eta \kappa^2 \|g_{\rho}\|_{\rho}}{1 - \theta} T^{-r(1-\theta)} \sqrt{2 \log \frac{2}{\delta}}.$$

(ii) **Step 2** Bound $\|h_{\infty}^{(T+1)}\|_{\rho}$. By utilizing the definition of $h_{\infty}^{(T+1)}$ and the regularity condition of g_{ρ} (as stated in Assumption 3), we obtain

$$\left\| h_{\infty}^{(T+1)} \right\|_{\rho} = \left\| \prod_{t=1}^{T} (I - \eta_{t}(L_{\infty} + \lambda I)) g_{\rho} \right\|_{\rho} \leq \left\| \prod_{t=1}^{T} (I - \eta_{t}(L_{\infty} + \lambda I)) L_{\infty}^{r} \right\|_{\rho p} \|L_{\infty}^{-r} g_{\rho}\|_{\rho}.$$

Note that $\prod_{t=1}^{T} (I - \eta_t (L_{\infty} + \lambda I)) L_{\infty}^r$ is a compact self-adjoint operator, then

$$\left\| \prod_{t=1}^{T} \left(I - \eta_t \left(L_{\infty} + \lambda I \right) \right) L_{\infty}^r \right\|_{op} \leq \sup_{u \geq 0} \left[u^r \prod_{t=1}^{T} \left(1 - \eta_t (u + \lambda) \right) \right]$$

$$\leq \sup_{u \geq 0} \left[u^r \prod_{t=1}^{T} \left(1 - \eta_t u \right) \right] \leq \sup_{u \geq 0} \left[u^r \exp \left\{ -u \sum_{t=1}^{T} \eta_t \right\} \right]$$

$$= \left(\frac{r}{e} \right)^r \frac{1}{\left(\sum_{t=1}^{T} \eta_t \right)^r} \leq \left(\frac{r(1-\theta)}{e\eta(1-2^{\theta-1})} \right)^r T^{-r(1-\theta)},$$

where we use the fact that $h(u) = u^r \exp\left\{-u \sum_{t=1}^T \eta_t\right\}$ takes maxmium at $u = \frac{r}{\sum_{t=1}^T \eta_t}$, and the final inequality uses Lemma A.2.

Combining the above two estimations, we can derive that

$$\begin{split} \left\| h_{M}^{(T+1)} \right\|_{\rho} & \leq \left\| h_{M}^{(T+1)} - h_{\infty}^{(T+1)} \right\|_{\rho} + \left\| h_{\infty}^{(T+1)} \right\|_{\rho} \\ & \leq \left[\frac{\eta \kappa^{2} \|g_{\rho}\|_{\rho}}{1 - \theta} \sqrt{2 \log \frac{2}{\delta}} + \left(\frac{r(1 - \theta)}{e \eta (1 - 2^{\theta - 1})} \right)^{r} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho} \right] T^{-r(1 - \theta)} \\ & \leq \left[\frac{\eta \kappa^{2} \|g_{\rho}\|_{\rho}}{1 - \theta} + \left(\frac{r(1 - \theta)}{e \eta (1 - 2^{\theta - 1})} \right)^{r} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho} \right] T^{-r(1 - \theta)} \sqrt{2 \log \frac{2}{\delta}}, \end{split}$$

holds with probability at least $1 - \delta$ when $M \ge T^{2(1+r)(1-\theta)}$, the last inequality holds due to $2\log\frac{2}{\delta} = \log\frac{4}{\delta^2} > 1$ for $0 < \delta < 1$, which finishes the proof.

Before establishing a bound for the second term on the right hand side of (20), we introduce several lemmas that will be valuable in the proof. Lemma 4.3 and Lemma 4.5 will be proved in the appendix and Lemma 4.4 can be found in [4].

Lemma 4.3. For any $\lambda > 0$, let $\eta_i = \eta i^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and

$$0 < \eta < \left(\lambda + \frac{208(1+\kappa)^4}{(1-2^{\theta-1})(2\theta-1)} \left(\log \frac{8}{1-\theta} + \frac{1}{1-\theta}\right)\right)^{-1},\tag{24}$$

when $M \ge \frac{1}{\lambda^2}$ and $t \ge 2$, the following holds with high probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$

$$\mathbb{E}_{Z^{t-1}} \left[\left\| g^{(t)} \right\|_{\infty}^{2} \right] \leq 12\kappa^{4r+2} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho}^{2} \sqrt{2\log \frac{2}{\delta}} + \frac{8\kappa^{4}\eta^{2}\theta}{2\theta - 1} \left(20 \|g_{\rho}\|_{\rho}^{2} + 3\mathcal{E}\left(g_{\rho}\right) \right).$$

Lemma 4.4 ([4]). Let \mathcal{T} and \mathcal{S} be normal Hilbert-Schmidt operators and $\{t_i\}_{i=1}^{\infty}$ and $\{s_i\}_{i=1}^{\infty}$ be enumerations of their eigenvalues, then for any $\varepsilon > 0$, there exists a permutation π of \mathbb{N}_+ such that

$$\left[\sum_{i=1}^{\infty} \left| s_i - t_{\pi(i)} \right|^2 \right]^{\frac{1}{2}} \leq \|\mathcal{S} - \mathcal{T}\|_{HS} + \varepsilon.$$

Lemma 4.5. Assume that $\{\mu_j\}_{j=1}^{\infty}$ be the eigenvalues of L_{∞} sorted in non-ascending order, then

$$\sum_{j=1}^{\infty} \mu_j^2 \exp\left\{-H(\theta)(\lambda + \mu_j) T^{1-\theta}\right\} \leq \begin{cases} \frac{2\beta c^2}{2\beta - 1} \left(\exp\left\{-bH(\theta) T^{1 - \frac{3\beta - 1}{2\beta - 1}\theta}\right\} + T^{-\theta}\right), & \frac{1}{2} < \theta < \frac{2\beta - 1}{3\beta - 1} \\ \frac{2\beta c^2}{2\beta - 1} \exp\left\{-H(\theta) \lambda T^{1-\theta}\right\}, & \frac{2\beta - 1}{3\beta - 1} \leq \theta < 1 \end{cases}$$

Now we are ready to provide an estimation for the second term on the right hand side of (20).

Proposition 4.7. Under the assumptions of Proposition 4.5, the following inequalities holds with probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$

(a) if
$$\frac{1}{2} < \theta < \frac{2\beta - 1}{3\beta - 1}$$
,

$$\sum_{t=1}^{T} \eta_t^2 \mathbb{E}_{Z^t} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_i \left(L_M + \lambda I \right) \right) \mathcal{B}_t \right\|_{\rho}^2 \right] \le V_2 \left(\exp \left\{ -H(\theta) T^{1 - \frac{3\beta - 1}{2\beta - 1} \theta} \right\} + T^{-\theta} \right) \log \frac{2}{\delta}, \tag{25}$$

(b) if
$$\frac{2\beta - 1}{3\beta - 1} \le \theta < 1$$
,

$$\sum_{t=1}^{T} \eta_t^2 \mathbb{E}_{Z^t} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_i \left(L_M + \lambda I \right) \right) \mathcal{B}_t \right\|_{\rho}^2 \right] \le V_2 \left(\exp\left\{ -H(\theta) \lambda T^{1-\theta} \right\} + T^{-\theta} \right) \log \frac{2}{\delta}, \tag{26}$$

where constants V_2 and $H(\theta)$ are defined in Proposition 4.5.

Proof. Recall that $\mathcal{B}_t = (y_t - g^{(t)}(x_t))k_{M,x_t} + L_M(g^{(t)} - g_\rho)$, and $\mathbb{E}_{z_t}[\mathcal{B}_t] = 0$, we can derive

$$\mathbb{E}_{Z^{t}} \left[\left\| \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \right] = \mathbb{E}_{Z^{t}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\mathcal{H}_{M}}^{2} \right] \\
= \mathbb{E}_{Z^{t}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(\left(y_{t} - g^{(t)}(x_{t}) \right) k_{M, x_{t}} \right) \right\|_{\mathcal{H}_{M}}^{2} \right] \\
+ \mathbb{E}_{Z^{t-1}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(L_{M}(g^{(t)} - g_{\rho}) \right) \right\|_{\mathcal{H}_{M}}^{2} \right] \\
+ 2 \mathbb{E}_{Z^{t}} \left\langle L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(\left(y_{t} - g^{(t)}(x_{t}) \right) k_{M, x_{t}} \right), \right. \\
L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(L_{M}(g^{(t)} - g_{\rho}) \right) \right\rangle_{\mathcal{H}_{M}} \\
= \mathbb{E}_{Z^{t}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(L_{M}(g^{(t)} - g_{\rho}) \right) \right\|_{\mathcal{H}_{M}}^{2} \right] \\
- \mathbb{E}_{Z^{t-1}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(L_{M}(g^{(t)} - g_{\rho}) \right) \right\|_{\mathcal{H}_{M}}^{2} \right] \\
\leq \mathbb{E}_{Z^{t}} \left[\left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left(\left(y_{t} - g^{(t)}(x_{t}) \right) k_{M, x_{t}} \right) \right\|_{\mathcal{H}_{M}}^{2} \right].$$

Note that $||f||_{\mathcal{H}_M}^2 = \text{Tr}(f \otimes_{\mathcal{H}_M} f)$, we have

$$\mathbb{E}_{Z^{t}} \left\| L_{M}^{\frac{1}{2}} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \left((y_{t} - g^{(t)}(x_{t})) k_{M,x_{t}} \right) \right\|_{\mathcal{H}_{M}}^{2}$$

$$= \mathbb{E}_{Z^{t}} \operatorname{Tr} \left[L_{M} \prod_{i=t+1}^{T} (I - \eta_{i} (L_{M} + \lambda I))^{2} (y_{t} - g^{(t)}(x_{t}))^{2} (k_{M,x_{t}} \otimes_{\mathcal{H}_{M}} k_{M,x_{t}}) \right]$$

$$\leq 2 \left(1 + \mathbb{E}_{Z^{t-1}} \left[\left\| g^{(t)} \right\|_{\infty}^{2} \right] \right) \operatorname{Tr} \left[L_{M}^{2} \prod_{i=t+1}^{T} (I - \eta_{i} (L_{M} + \lambda I))^{2} \right].$$

Thus, according to Lemma 4.3, when $M \ge \frac{1}{\lambda^2}$, the following holds with confidence at least $1 - \delta$

$$\sum_{t=1}^{T} \eta_{t}^{2} \mathbb{E}_{Z^{t}} \left\| \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{t} \right\|_{\rho}^{2} \leq W_{2} \sqrt{2 \log \frac{2}{\delta}} \sum_{t=1}^{T} \eta_{t}^{2} \operatorname{Tr} \left[L_{M}^{2} \prod_{i=t+1}^{T} \left(I - \eta_{i} \left(L_{M} + \lambda I \right) \right)^{2} \right].$$

Our remaining task is to estimate the term $\sum_{t=1}^{T} \eta_t^2 \text{Tr} \left[L_M^2 \prod_{i=t+1}^T \left(I - \eta_i \left(L_M + \lambda I \right) \right)^2 \right]$. Let $\{\mu_{j,M}\}_{j=1}^{\infty}$ be the eigenvalues of L_M and $\{\mu_j\}_{j=1}^{\infty}$ be the eigenvalues of L_{∞} , both sorted in non-ascending order, then applying Lemma 4.4 to $\mathcal{T} = L_{\infty}$ and $\mathcal{S} = L_M$, and for $\varepsilon = \|L_M - L_{\infty}\|_{HS}$, there exists a permutation π of \mathbb{N}_+ such that

$$\sup_{j \ge 1} |\mu_j - \mu_{\pi(j),M}| \le 2 \|L_M - L_\infty\|_{HS}.$$

Furthermore, by Lemma A.1, the following holds with confidence at least $1 - \delta$,

$$\sup_{j>1} \left| \mu_j - \mu_{\pi(j),M} \right| \le 2 \|L_M - L_\infty\|_{HS} \le \frac{2\kappa^2}{\sqrt{M}} \sqrt{2\log\frac{2}{\delta}}.$$
 (27)

Using this result, we can now proceed with the error decomposition as follows

$$\sum_{t=1}^{T} \eta_t^2 \operatorname{Tr} \left[L_M^2 \prod_{i=t+1}^{T} \left(I - \eta_i \left(L_M + \lambda I \right) \right)^2 \right] = \sum_{j=1}^{\infty} \mu_{\pi(j),M}^2 \sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} \left(1 - \eta_i (\mu_{\pi(j),M} + \lambda) \right)^2 \\
= \sum_{j=1}^{\infty} (\mu_{\pi(j),M}^2 - \mu_j^2) \sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} \left(1 - \eta_i (\mu_{\pi(j),M} + \lambda) \right)^2 + \sum_{j=1}^{\infty} \mu_j^2 \sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} \left(1 - \eta_i (\mu_{\pi(j),M} + \lambda) \right)^2 \\
\leq \sum_{j=1}^{\infty} (\mu_{\pi(j),M}^2 - \mu_j^2) \sum_{t=1}^{T} \eta_t^2 + \sum_{j=1}^{\infty} \mu_j^2 \sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} \left(1 - \eta_i (\mu_{\pi(j),M} + \lambda) \right)^2. \tag{28}$$

For the first term on the right hand side of (28), we have

$$\sum_{j=1}^{\infty} (\mu_{\pi(j),M}^2 - \mu_j^2) \sum_{t=1}^{T} \eta_t^2 = \sum_{j=1}^{\infty} (\mu_{\pi(j),M} - \mu_j) (\mu_{\pi(j),M} + \mu_j) \sum_{t=1}^{T} \eta_t^2$$

$$\leq \frac{2\eta^2 \theta}{2\theta - 1} \sup_{j \geq 1} \left| \mu_j - \mu_{\pi(j),M} \right| \operatorname{Tr}(L_M + L_\infty)$$

$$\leq \frac{8\kappa^4 \eta^2 \theta}{2\theta - 1} \frac{1}{\sqrt{M}} \sqrt{2 \log \frac{2}{\delta}},$$

where the last inequality uses (27) and the equation $\text{Tr}(L_M) = \int_{\mathcal{X}} k_M(x,x) d\rho_{\mathcal{X}}(x) \leq \kappa^2$ and $\text{Tr}(L_\infty) = \int_{\mathcal{X}} k_\infty(x,x) d\rho_{\mathcal{X}}(x) \leq \kappa^2$. When $M \geq T^{2\theta}$, it can be further bounded by

$$\sum_{j=1}^{\infty} (\mu_{\pi(j),M}^2 - \mu_j^2) \sum_{t=1}^{T} \eta_t^2 \le \frac{8\kappa^4 \eta^2 \theta}{2\theta - 1} \sqrt{2\log \frac{2}{\delta}} T^{-\theta}.$$

For the second term on the right hand side of (28), using Lemma A.2 (c), we get

$$\sum_{j=1}^{\infty} \mu_{j}^{2} \sum_{t=1}^{T} \eta_{t}^{2} \prod_{i=t+1}^{T} (1 - \eta_{i}(\mu_{\pi(j),M} + \lambda))^{2}$$

$$\leq \sum_{j=1}^{\infty} \mu_{j}^{2} \left[\frac{2\eta^{2}\theta}{2\theta - 1} \exp\left\{ -H(\theta)(\lambda + \mu_{\pi(j),M})T^{1-\theta} \right\} + \frac{2^{\theta}\eta}{\lambda + \mu_{\pi(j),M}}T^{-\theta} \right]$$

$$\leq \frac{2\eta^{2}\theta}{2\theta - 1} \sum_{j=1}^{\infty} \mu_{j}^{2} \exp\left\{ -H(\theta)(\lambda + \mu_{j})T^{1-\theta} \right\} \cdot \exp\left\{ H(\theta) \sup_{j \geq 1} \left| \mu_{\pi(j),M} - \mu_{j} \right| T^{1-\theta} \right\}$$

$$+ 2^{\theta}\eta T^{-\theta} \sum_{j=1}^{\infty} \frac{\mu_{j}^{2}}{\lambda + \mu_{j}} \left(1 + \frac{\sup_{j \geq 1} \left| \mu_{\pi(j),M} - \mu_{j} \right|}{\lambda} \right)$$

$$\leq \frac{2\eta^{2}\theta}{2\theta - 1} \sum_{j=1}^{\infty} \mu_{j}^{2} \exp\left\{ -H(\theta)(\lambda + \mu_{j})T^{1-\theta} \right\} \cdot \exp\left\{ H(\theta) \frac{2\kappa^{2}}{\sqrt{M}} \sqrt{2\log\frac{2}{\delta}}T^{1-\theta} \right\}$$

$$+ 2^{\theta}\eta T^{-\theta} \sum_{j=1}^{\infty} \frac{\mu_{j}^{2}}{\lambda + \mu_{j}} \left(1 + \frac{2\kappa^{2}}{\sqrt{M}\lambda} \sqrt{2\log\frac{2}{\delta}} \right),$$

where the last inequality holds due to (27), moreover, when $M \ge \max \left\{ T^{2(1-\theta)}, \frac{1}{\lambda^2} \right\}$, and by $\sum_{i=1}^{\infty} \frac{\mu_j^2}{\lambda + \mu_j} \le \text{Tr}(L_{\infty}) \le \kappa^2$, the above inequality can be further bounded as follows

$$\sum_{j=1}^{\infty} \mu_j^2 \sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} (1 - \eta_i (\mu_{\pi(j),M} + \lambda))^2$$

$$\leq \frac{2\eta^2 \theta}{2\theta - 1} \exp\left\{ 2H(\theta) \kappa^2 \sqrt{2\log \frac{2}{\delta}} \right\} \sum_{j=1}^{\infty} \mu_j^2 \exp\left\{ -H(\theta) (\lambda + \mu_j) T^{1-\theta} \right\}$$

$$+ 6\eta \kappa^4 T^{-\theta} \sqrt{2\log \frac{2}{\delta}}.$$

Putting the above two bounds back into (28) yields

$$\sum_{t=1}^{T} \eta_t^2 \mathbb{E}_{Z^t} \left\| \prod_{i=t+1}^{T} \left(I - \eta_i \left(L_M + \lambda I \right) \right) \mathcal{B}_t \right\|_{\rho}^2$$

$$\leq W_2 \left(\frac{8\kappa^4 \eta^2 \theta}{2\theta - 1} T^{-\theta} + \frac{2\eta^2 \theta}{2\theta - 1} \exp\left\{ 2H(\theta)\kappa^2 \sqrt{2\log\frac{2}{\delta}} \right\} \sum_{j=1}^{\infty} \mu_j^2 \exp\left\{ -H(\theta)(\lambda + \mu_j) T^{1-\theta} \right\}$$

$$+ 6\kappa^4 \eta 12 T^{-\theta} \right) 2\log\frac{2}{\delta}.$$

Then the desired result can be obtained by applying Lemma 4.5 to the above inequality for the term $\sum_{j=1}^{\infty} \mu_j^2 \exp\left\{-H(\theta)(\lambda + \mu_j)T^{1-\theta}\right\}$.

At this stage, the proof of Proposition 4.5 becomes quite evident.

Proof of Proposition 4.5. By substituting the bounds (22), (25) and (26) back into the error decomposition of the convergence error (20), we obtain the desired result.

4.4 Estimation For Random Feature Error

In this subsection, our goal is to establish a bound for the random feature error $\|g_{M,\lambda} - g_{\infty,\lambda}\|_{\rho}^2$.

Proposition 4.8. Under Assumptions 1 and 2, for any $\lambda > 0$, let $\eta_i = i^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and $\eta > 0$, when

$$M \ge M_3(T,\lambda) := \frac{T^{\theta}}{\lambda^2},$$

the following inequality holds with confidence at least $1 - \delta$,

$$\|g_{M,\lambda} - g_{\infty,\lambda}\|_{\rho}^2 \le 2\kappa^4 \log \frac{2}{\delta} T^{-\theta}.$$

Proof. By the definition of $g_{M,\lambda}$ and $g_{\infty,\lambda}$, we have

$$g_{M,\lambda} - g_{\infty,\lambda} = ((L_M + \lambda I)^{-1} L_M - (L_\infty + \lambda I)^{-1} L_\infty) g_\rho$$

= $\lambda ((L_\infty + \lambda I)^{-1} - (L_M + \lambda I)^{-1}) g_\rho$
= $\lambda (L_\infty + \lambda I)^{-1} (L_M - L_\infty) (L_M + \lambda I)^{-1} g_\rho$.

Then it follows that

$$\begin{aligned} \|g_{M,\lambda} - g_{\infty,\lambda}\|_{\rho} &= \|\lambda (L_{\infty} + \lambda I)^{-1} (L_{M} - L_{\infty}) (L_{M} + \lambda I)^{-1} g_{\rho}\|_{\rho} \\ &\leq \|(L_{M} + \lambda I)^{-1}\|_{op} \|L_{M} - L_{\infty}\|_{op} \|g_{\rho}\|_{\rho} \\ &\leq \lambda^{-1} \|L_{M} - L_{\infty}\|_{op} \|g_{\rho}\|_{\rho}. \end{aligned}$$

Thus from Lemma A.1, we get when $M \ge \frac{T^{\theta}}{\lambda^2}$,

$$\|g_{M,\lambda} - g_{\infty,\lambda}\|_{\rho} \le \kappa^2 \sqrt{2\log\frac{2}{\delta}} T^{-\frac{\theta}{2}}$$

holds with confidence at least $1 - \delta$. This completes the proof.

4.5 Estimation For Approximation Error

The bound for the approximation error $||g_{\infty,\lambda} - g_{\rho}||_{\rho}^{2}$ is a well-known result, as shown in Proposition 3 of [8].

Proposition 4.9. Under Assumption 3 with $\frac{1}{2} < r \le 1$, then for any $\lambda > 0$,

$$\|g_{\infty,\lambda} - g_{\rho}\|_{\rho}^{2} \le \lambda^{2r} \|L_{\infty}^{-r} g_{\rho}\|_{\rho}^{2}.$$
 (29)

4.6 Proof of Main Results

Now we are in a position to prove our main results.

Proof of Theorem 1. We begin by comparing the assumption on the upper bound of η in Proposition 4.1 and Proposition 4.5. When $\frac{1}{2} < \theta < 1$, it can be observed that

$$\frac{1-\theta}{5\kappa^2 + \lambda} \le \left(\lambda + \frac{208(1+\kappa)^4}{(1-2^{\theta-1})(2\theta-1)} \left(\log \frac{8}{1-\theta} + \frac{1}{1-\theta}\right)\right)^{-1}.$$

Thus, for $i \in \mathbb{N}_+$ we select $\eta_i = \eta i^{-\theta}$ with $\frac{1}{2} < \theta < 1$ and $0 < \eta < \left(\lambda + \frac{208(1+\kappa)^4}{(1-2^{\theta-1})(2\theta-1)}\right) \left(\log \frac{8}{1-\theta} + \frac{1}{1-\theta}\right)^{-1}$. Let $M_1(T), M_2(T, \lambda), M_3(T, \lambda)$ be defined as given in Proposition 4.1, 4.5 and 4.8, respectively. For $\frac{1}{2} < \theta < 1$, we define

$$M_{0}(T,\lambda) = \max\{M_{1}(T), M_{2}(T,\lambda), M_{3}(T,\lambda)\} = \max\left\{T^{6-5\theta}, T^{2(1+r)(1-\theta)}, T^{2\theta}, \frac{1}{\lambda^{2}}, \frac{T^{\theta}}{\lambda^{2}}\right\}$$

$$= \max\left\{T^{6-5\theta}, T^{2\theta}, \frac{T^{\theta}}{\lambda^{2}}\right\}.$$
(30)

To establish the main result, we will analyze two cases based on the value of θ .

When $\frac{1}{2} < \theta < \frac{2\beta-1}{3\beta-1}$ and $M \ge M_0(T,\lambda)$. by substituting the bounds from Proposition 4.1, Proposition 4.5, Proposition 4.8 and Proposition 4.9 into the error decomposition (14), we conclude that

$$\begin{split} \mathbb{E}_{Z^T} \left[\left\| g_{\Theta^{(T+1)}} - g_{\rho} \right\|_{\rho}^2 \right] &\leq 4 \mathbb{E}_{Z^T} \left[\left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\rho}^2 \right] + 4 \mathbb{E}_{Z^T} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^2 \right] \\ &+ 4 \left\| g_{M,\lambda} - g_{\infty,\lambda} \right\|_{\rho}^2 + 4 \left\| g_{\infty,\lambda} - g_{\rho} \right\|_{\rho}^2 \\ &\leq 100 C_{\sigma}^2 T^{-\theta} + 4 \left(V_1 T^{-2r(1-\theta)} + V_2 \left(\exp\left\{ -H(\theta) T^{1-\frac{3\beta-1}{2\beta-1}\theta} \right\} + T^{-\theta} \right) \right) \log \frac{2}{\delta} + 16 \kappa^4 \log \frac{2}{\delta} \\ &+ 4 \lambda^{2r} \| L_{\infty}^{-r} g_{\rho} \|_{\rho}^2 \\ &\leq \tilde{C} \left(T^{-2r(1-\theta)} + \exp\left\{ -H(\theta) T^{1-\frac{3\beta-1}{2\beta-1}\theta} \right\} + T^{-\theta} + \lambda^{2r} \right) \log \frac{2}{\delta}. \end{split}$$

holds with confidence at least $1-\delta$, where constant $\tilde{C} = 200C_{\sigma}^2 + 4\left(V_1 + V_2\right) + 16\kappa^4 + 8\|L_{\infty}^{-r}g_{\rho}\|_{\rho}^2$.

Similarly, when $\frac{2\beta-1}{3\beta-1} \leq \theta < 1$ and $M \geq M_0(T,\lambda)$, with confidence at least $1-\delta$, there holds

$$\begin{split} \mathbb{E}_{Z^T} \left[\left\| g_{\Theta^{(T+1)}} - g_{\rho} \right\|_{\rho}^2 \right] &\leq 4 \mathbb{E}_{Z^T} \left[\left\| g_{\Theta^{(T+1)}} - g^{(T+1)} \right\|_{\rho}^2 \right] + 4 \mathbb{E}_{Z^T} \left[\left\| g^{(T+1)} - g_{M,\lambda} \right\|_{\rho}^2 \right] \\ &+ 4 \left\| g_{M,\lambda} - g_{\infty,\lambda} \right\|_{\rho}^2 + 4 \left\| g_{\infty,\lambda} - g_{\rho} \right\|_{\rho}^2 \\ &\leq 100 C_{\sigma}^2 T^{-\theta} + 4 \left(V_1 T^{-2r(1-\theta)} + V_2 \left(\exp\left\{ -H(\theta) \lambda T^{1-\theta} \right\} + T^{-\theta} \right) \right) \log \frac{2}{\delta} + 16 \kappa^4 \log \frac{2}{\delta} T^{-\theta} \\ &+ 4 \lambda^{2r} \| L_{\infty}^{-r} g_{\rho} \|_{\rho}^2 \\ &\leq \tilde{C} \left(T^{-2r(1-\theta)} + \exp\left\{ -H(\theta) \lambda T^{1-\theta} \right\} + T^{-\theta} + \lambda^{2r} \right) \log \frac{2}{\delta}. \end{split}$$

Then the proof is completed.

We will now establish the explicit convergence rates by considering the specific choice of $\theta = \frac{2r}{2r+1}$, where $\frac{1}{2} < r \le 1$.

Proof of Corollary 1. Since for all $\varepsilon > 0$, $\alpha > 0$ and $\tau > 0$, we have

$$\exp\{-\tau T^{\varepsilon}\} = O(T^{-\alpha}). \tag{31}$$

We obtain the convergence rate by applying Theorem 1 with $\theta = \frac{2r}{2r+1}$. We will consider two cases based on the value of θ , as outlined in Theorem 1.

Case 1: When $\frac{1}{2} < \theta = \frac{2r}{2r+1} < \frac{2\beta-1}{3\beta-1}$, which implies $1 - \frac{3\beta-1}{2\beta-1}\theta > 0$ and $\frac{1}{2} < r < 1 - \frac{1}{2\beta}$, using the property (31) with $\tau = H(\theta) = \frac{2\eta}{1-\theta}2^{1-\theta}\left(1-\left(\frac{3}{4}\right)^{1-\theta}\right) = 2\eta(2r+1)2^{\frac{1}{2r+1}}\left(1-\left(\frac{3}{4}\right)^{\frac{1}{2r+1}}\right) > 0$ and $\varepsilon = 1 - \frac{3\beta-1}{2\beta-1}\theta$, we can conclude that

$$\exp\left\{-H(\theta)T^{1-\frac{3\beta-1}{2\beta-1}\theta}\right\} = O(T^{-\frac{2r}{2r+1}}).$$

This together with the choice of $\lambda = T^{-\frac{1}{2r+1}}$, we have

$$M_0(T,\lambda) = \max\left\{T^{6-5\theta}, T^{2\theta}, \frac{T^{\theta}}{\lambda^2}\right\} = \max\left\{T^{\frac{2r+6}{2r+1}}, T^{\frac{4r}{2r+1}}, T^{\frac{2r+2}{2r+1}}\right\} = T^{\frac{2r+6}{2r+1}}.$$

Therefore, when $M \geq T^{\frac{2r+6}{2r+1}}$, we have

$$\mathbb{E}_{Z^T} \left[\|g_{\Theta^{(T+1)}} - g_{\rho}\|_{\rho}^2 \right] = O(T^{-\frac{2r}{2r+1}}) \log \frac{2}{\delta}, \tag{32}$$

holds with confidence at least $1 - \delta$.

Case 2: When $\frac{2\beta-1}{3\beta-1} \leq \theta < 1$, which implies $1-\frac{1}{2\beta} \leq r \leq 1$, we choose $\lambda = T^{-\frac{1}{2r+1}+\frac{\epsilon}{2r}}$ with $0 < \epsilon < \frac{2r}{2r+1}$. Under these conditions, we have $\lambda T^{1-\theta} = T^{-\frac{1}{2r+1}+\frac{\epsilon}{2r}} T^{1-\frac{2r}{2r+1}} = T^{\frac{\epsilon}{2r}}$, therefore, we conclude that

$$\exp\left\{-H(\theta)\lambda T^{1-\theta}\right\} = \exp\left\{-H(\theta)T^{\frac{\epsilon}{2r}}\right\} = O\left(T^{-\frac{2r}{2r+1}}\right). \tag{33}$$

Moreover, in this case,

$$M_0(T,\lambda) = \max\left\{T^{6-5\theta}, T^{2\theta}, \frac{T^{\theta}}{\lambda^2}\right\} = \max\left\{T^{\frac{2r+6}{2r+1}}, T^{\frac{4r}{2r+1}}, T^{\frac{2r+2}{2r+1} - \frac{\epsilon}{r}}\right\} = T^{\frac{2r+6}{2r+1}}.$$

Then we assert that when $M \geq T^{\frac{2r+6}{2r+1}}$, with confidence at least $1-\delta$, there holds

$$\mathbb{E}_{Z^T} \left[\| g_{\Theta^{(T+1)}} - g_{\rho} \|_{\rho}^2 \right] = O(T^{-\frac{2r}{2r+1} + \epsilon}) \log \frac{2}{\delta}.$$

This completes the proof of Corollary 1.

5 Appendix

In this section, we give some useful lemmas used in the proof of our main results.

A Useful Lemmas

Lemma A.1. For arbitrary $\delta \in (0,1)$ and $M \in \mathbb{Z}_+$, the following inequality holds with confidence at least $1-\delta$,

(a)
$$||L_M - L_\infty||_{op} \le ||k_M - k_\infty||_{L_\infty(\rho_X \times \rho_X)} \le \kappa^2 \sqrt{\frac{2\log\frac{2}{\delta}}{M}}$$
.

(b)
$$|Tr(L_M - L_\infty)| \le \kappa^2 \sqrt{\frac{2\log\frac{2}{\delta}}{M}}$$
.

$$(c) \|L_M - L_\infty\|_{HS} \le \kappa^2 \sqrt{\frac{2\log\frac{2}{\delta}}{M}}.$$

Proof. (a) Let's establish the proof for the first inequality. From the definition of operator norm, we get that

$$\begin{aligned} \|L_{M} - L_{\infty}\|_{op} &= \sup_{\|f\|_{\rho} = 1} \|L_{M}f - L_{\infty}f\|_{\rho} \\ &= \sup_{\|f\|_{\rho} = 1} \left\| \int_{\mathcal{X}} f(x)(k_{M,x} - k_{\infty,x})d\rho_{\mathcal{X}}(x) \right\|_{\rho} \\ &= \sup_{\|f\|_{\rho} = 1} \sqrt{\int_{\mathcal{X}} \left[\int_{\mathcal{X}} f(x)\left(k_{M}(x, x') - k_{\infty}(x, x')\right)d\rho_{\mathcal{X}}(x)\right]^{2} d\rho_{\mathcal{X}}(x')} \\ &\leq \sup_{\|f\|_{\rho} = 1} \sqrt{\int_{\mathcal{X}} \left[\left(\int_{\mathcal{X}} |f(x)|^{2} dx \right) \left(\int_{\mathcal{X}} |k_{M}(x, x') - k_{\infty}(x, x')|^{2} d\rho_{\mathcal{X}}(x) \right) \right] d\rho_{\mathcal{X}}(x')} \\ &\leq \|k_{M} - k_{\infty}\|_{L_{\infty}(\rho_{\mathcal{X}} \times \rho_{\mathcal{X}})}. \end{aligned}$$

Thus we get the first half of the inequality. For the second half, for any $x, x' \in \mathcal{X}$, define

$$\xi_r(x,x') = \sigma\left(b_r^{(0)\top}x\right)\sigma\left(b_r^{(0)\top}x'\right) + \left(x^\top x' + \gamma^2\right)\sigma'\left(b_r^{(0)\top}x\right)\sigma'\left(b_r^{(0)\top}x'\right),$$

then $|\xi_r(x,x')| \leq 4 + 2C_\sigma^2 = \kappa^2$ holds for any $r = 1, \dots, M$. Then, by applying the Hoeffding inequality (See e.g. Theorem 1 in [5]), we can assert that

$$|k_M(x, x') - k_{\infty}(x, x')| = \left| \frac{1}{M} \sum_{r=1}^{M} \left(\xi_r(x, x') - \mathbb{E}_{b^{(0)}} \xi(x, x') \right) \right| \le \kappa^2 \sqrt{\frac{2 \log \frac{2}{\delta}}{M}}.$$

holds with confidence at least $1 - \delta$, Thus $||k_M - k_\infty||_{L_\infty(\rho_X \times \rho_X)} \le \kappa^2 \sqrt{\frac{2\log\frac{2}{\delta}}{M}}$.

(b) This can be proved by utilizing the equation $Tr(L_{\infty}) = \int_{\mathcal{X}} k_{\infty}(x, x) d\rho_{\mathcal{X}}(x)$ and (a).

(c) According to the definition and properties of the norm of Hilbert-Schmidt operators, we can derive the following result

$$||L_M - L_\infty||_{HS}^2 = |\operatorname{Tr}[(L_M - L_\infty)^*(L_M - L_\infty)]| \le ||L_M - L_\infty||_{op} |\operatorname{Tr}(L_M - L_\infty)|.$$

Therefore, the desired result follows directly from (a) and (b).

Lemma A.2. Let $\eta_t = \eta t^{-\theta}$ with $\eta > 0$ and $\frac{1}{2} \le \theta < 1$, we have

$$(a) \frac{\eta(1-2^{\theta-1})}{1-\theta} T^{1-\theta} \le \sum_{t=1}^{T} \eta_t \le \frac{\eta}{1-\theta} T^{1-\theta}.$$

(b)
$$\sum_{t=1}^{T} \eta_t^2 \le \frac{2\eta^2 \theta}{2\theta - 1}$$
.

(c)
$$\forall \lambda > 0$$
, define $H(\theta) = \frac{2\eta}{1-\theta} 2^{1-\theta} \left(1 - \left(\frac{3}{4}\right)^{1-\theta}\right)$, then

$$\sum_{t=1}^{T} \eta_t^2 \prod_{i=t+1}^{T} (1 - \eta_i \lambda)^2 \le \frac{2\theta \eta^2}{2\theta - 1} \exp\left\{-H(\theta)\lambda T^{1-\theta}\right\} + \frac{\eta^2 \eta^2}{\lambda T^{\theta}}.$$

Proof. The proof of (a) and (b) is standard, see e.g. Lemma 4.3 in [17]. The proof of (c) can be found in Lemma 19 in [27].

B Proof of Lemmas in Section 4.3

We need the following lemma to establish the proof of Lemma 4.3.

Lemma B.1. For arbitrary $\lambda > 0$ and $0 < \delta < 1$, when $M \ge \frac{1}{\lambda^2}$, the following holds with probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$,

$$\|g_{M,\lambda}\|_{\infty} \leq \sqrt{2}\kappa^{2r+1} \|L_{\infty}^{-r}g_{\rho}\|_{\rho} \left(2\log\frac{2}{\delta}\right)^{\frac{1}{4}}.$$

Proof. By utilizing the definition of $g_{M,\lambda}(x)$ as $g_{M,\lambda} = L_M (L_M + \lambda I)^{-1} g_\rho$ and the reproducing property of k_M , for any $x \in \mathcal{X}$, we can conclude that

$$|g_{M,\lambda}(x)| = \left| \langle g_{M,\lambda}, k_{M,x} \rangle_{\mathcal{H}_M} \right| \le ||k_{M,x}||_{\mathcal{H}_M} ||g_{M,\lambda}||_{\mathcal{H}_M}$$

$$\le \kappa \left| \left| L_M \left(L_M + \lambda I \right)^{-1} g_\rho \right| \right|_{\mathcal{H}_M}$$

$$= \kappa \left| \left| L_M^{\frac{1}{2}} \left(L_M + \lambda I \right)^{-1} g_\rho \right| \right|_{0}.$$

Furthermore, under Assumption 3 with $\frac{1}{2} < r \le 1$, we have

$$\begin{split} & \left\| L_{M}^{\frac{1}{2}} \left(L_{M} + \lambda I \right)^{-1} g_{\rho} \right\|_{\rho} \leq \left\| L_{M}^{\frac{1}{2}} \left(L_{M} + \lambda I \right)^{-1} L_{\infty}^{r} \right\|_{op} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho} \\ & \leq \left\| L_{M}^{\frac{1}{2}} \left(L_{M} + \lambda I \right)^{-\frac{1}{2}} \right\|_{op} \left\| \left(L_{M} + \lambda I \right)^{-\frac{1}{2}} \left(L_{\infty} + \lambda I \right)^{\frac{1}{2}} \right\|_{op} \left\| \left(L_{\infty} + \lambda I \right)^{-\frac{1}{2}} L_{\infty}^{r} \right\|_{op} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho} \end{split}$$

$$\leq \kappa^{2r-1} \left\| (L_M + \lambda I)^{-\frac{1}{2}} (L_\infty + \lambda I)^{\frac{1}{2}} \right\|_{op} \left\| L_\infty^{-r} g_\rho \right\|_{\rho},$$

where the last inequality holds due to $\left\|L_M^{\frac{1}{2}}(L_M + \lambda I)^{-\frac{1}{2}}\right\|_{op} \le 1$ and $\left\|(L_\infty + \lambda I)^{-\frac{1}{2}}L_\infty^r\right\|_{op} \le \kappa^{2r-1}$. Moreover,

$$\begin{split} \left\| (L_M + \lambda I)^{-\frac{1}{2}} (L_\infty + \lambda I)^{\frac{1}{2}} \right\|_{op}^2 &= \left\| (L_M + \lambda I)^{-\frac{1}{2}} (L_\infty + \lambda I) (L_M + \lambda I)^{-\frac{1}{2}} \right\|_{op} \\ &= \left\| (L_M + \lambda I)^{-\frac{1}{2}} (L_\infty - L_M + L_M + \lambda I) (L_M + \lambda I)^{-\frac{1}{2}} \right\|_{op} \\ &\leq \frac{\| L_M - L_\infty \|_{op}}{\lambda} + 1, \end{split}$$

from Lemma A.1, when $M \geq \frac{1}{\lambda^2}$, we get

$$\left\| (L_M + \lambda I)^{-\frac{1}{2}} (L_\infty + \lambda I)^{\frac{1}{2}} \right\|_{op} \le \sqrt{\frac{\|L_M - L_\infty\|_{op}}{\lambda} + 1} \le \sqrt{\kappa^2 \sqrt{2\log\frac{2}{\delta}} + 1} \le \sqrt{\kappa^2 + 1} \left(2\log\frac{2}{\delta} \right)^{\frac{1}{4}},$$

holds with probability at least $1 - \delta$. In summary, we can establish that

$$\|g_{M,\lambda}\|_{\infty} \le \kappa^{2r} \|L_{\infty}^{-r} g_{\rho}\|_{\rho} \sqrt{\kappa^{2} + 1} \left(2\log\frac{2}{\delta}\right)^{\frac{1}{4}} \le \sqrt{2}\kappa^{2r+1} \|L_{\infty}^{-r} g_{\rho}\|_{\rho} \left(2\log\frac{2}{\delta}\right)^{\frac{1}{4}}.$$

holds with high probability at least $1 - \delta$ over the random choice of $\Theta^{(0)}$. This concludes the proof.

Proof of Lemma 4.3. First, we divide $\mathbb{E}_{Z^{t-1}} \|g^{(t)}\|_{\infty}^2$ into the following two part,

$$\mathbb{E}_{Z^{t-1}} \left\| g^{(t)} \right\|_{\infty}^{2} \leq 2 \mathbb{E}_{Z^{t-1}} \left\| g^{(t)} - g_{M,\lambda} \right\|_{\infty}^{2} + 2 \left\| g_{M,\lambda} \right\|_{\infty}^{2}.$$

Recall that $\mathcal{E}(f) = \mathbb{E}_{(x,y)\sim\rho}[f(x)-y]^2$, and $\mathbb{E}_{z_i}[\mathcal{B}_i] = 0$, which implies that

$$\mathbb{E}_{z_i|Z^{i-1}}\left[\left\|\mathcal{B}_i\right\|_{\mathcal{H}_M}^2\right] \leq \mathbb{E}_{z_i|Z^{i-1}}\left[\left\|\left(y_i - g^{(i)}\left(x_i\right)\right)k_{M,x_i}\right\|_{\mathcal{H}_M}^2\right] \leq \kappa^2 \mathcal{E}\left(g^{(i)}\right).$$

Then for the first term $\mathbb{E}_{Z^{t-1}} \|g^{(t)} - g_{M,\lambda}\|_{\infty}^2$, by (21) in Lemma 4.2, we have

$$\begin{split} \mathbb{E}_{Z^{t-1}} \left\| g^{(t)} - g_{M,\lambda} \right\|_{\infty}^{2} &\leq 2 \left\| \prod_{i=1}^{t-1} \left(I - \eta_{t} \left(L_{M} + \lambda I \right) \right) g_{M,\lambda} \right\|_{\infty}^{2} \\ &+ 2 \mathbb{E}_{Z^{t-1}} \left\| \sum_{i=1}^{t-1} \eta_{i} \prod_{j=i+1}^{t-1} \left(I - \eta_{j} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{i} \right\|_{\infty}^{2} \\ &\leq 2 \left\| g_{M,\lambda} \right\|_{\infty}^{2} + 2 \kappa^{2} \mathbb{E}_{Z^{t-1}} \left\| \sum_{i=1}^{t-1} \eta_{i} \prod_{j=i+1}^{t-1} \left(I - \eta_{j} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{i} \right\|_{\mathcal{H}_{M}}^{2} \\ &= 2 \left\| g_{M,\lambda} \right\|_{\infty}^{2} + 2 \kappa^{2} \sum_{i=1}^{t-1} \eta_{i}^{2} \mathbb{E}_{Z^{i}} \left\| \prod_{j=i+1}^{t-1} \left(I - \eta_{j} \left(L_{M} + \lambda I \right) \right) \mathcal{B}_{i} \right\|_{\mathcal{H}_{M}}^{2} \end{split}$$

$$\leq 2 \|g_{M,\lambda}\|_{\infty}^2 + 2\kappa^4 \sum_{i=1}^{t-1} \eta_i^2 \mathbb{E}_{Z^{i-1}} [\mathcal{E}(g^{(i)})].$$

By Proposition 2 in [37], when η satisfies the condition (24), we have

$$\mathbb{E}_{Z^{i-1}}[\mathcal{E}(g^{(i)})] \le 20 \|g_{\rho}\|_{\rho}^{2} + 3\mathcal{E}(g_{\rho}).$$

Combining Lemma A.2 (b), we can derive that

$$\mathbb{E}_{Z^{t-1}} \left\| g^{(t)} - g_{M,\lambda} \right\|_{\infty}^{2} \leq 2 \left\| g_{M,\lambda} \right\|_{\infty}^{2} + \frac{4\kappa^{4}\eta^{2}\theta}{2\theta - 1} \left(20 \|g_{\rho}\|_{\rho}^{2} + 3\mathcal{E}\left(g_{\rho}\right) \right).$$

Thus, from Lemma B.1, when $M \geq \frac{1}{\lambda^2}$, we have

$$\begin{split} \mathbb{E}_{Z^{t-1}} \left\| g^{(t)} \right\|_{\infty}^{2} &\leq 2 \left\| g_{M,\lambda} \right\|_{\infty}^{2} + 2 \mathbb{E}_{Z^{t-1}} \left\| g^{(t)} - g_{M,\lambda} \right\|_{\infty}^{2} \\ &\leq 6 \left\| g_{M,\lambda} \right\|_{\infty}^{2} + \frac{8 \kappa^{4} \eta^{2} \theta}{2\theta - 1} \left(20 \| g_{\rho} \|_{\rho}^{2} + 3 \mathcal{E} \left(g_{\rho} \right) \right) \\ &\leq 12 \kappa^{4r+2} \left\| L_{\infty}^{-r} g_{\rho} \right\|_{\rho}^{2} \sqrt{2 \log \frac{2}{\delta}} + \frac{8 \kappa^{4} \eta^{2} \theta}{2\theta - 1} \left(20 \| g_{\rho} \|_{\rho}^{2} + 3 \mathcal{E} \left(g_{\rho} \right) \right). \end{split}$$

This finishes the proof.

Proof of Lemma 4.5 To prove the result, we will consider two cases based on the value of θ .

(1) When $\frac{1}{2} < \theta < \frac{2\beta - 1}{3\beta - 1}$, from Assumption 4, $\forall j \ge 1$, we have $bj^{-\beta} \le \mu_j \le cj^{-\beta}$ with $\beta > 1$, then

$$\sum_{j=1}^{\infty} \mu_{j}^{2} \exp\left\{-H(\theta)(\lambda + \mu_{j})T^{1-\theta}\right\} \leq \sum_{j=1}^{\infty} \mu_{j}^{2} \exp\left\{-H(\theta)\mu_{j}T^{1-\theta}\right\}$$

$$\leq c^{2} \sum_{j=1}^{\infty} j^{-2\beta} \exp\left\{-bH(\theta)j^{-\beta}T^{1-\theta}\right\}$$

$$= c^{2} \sum_{j \leq \lfloor T^{\frac{\theta}{2\beta-1}} \rfloor} j^{-2\beta} \exp\left\{-bH(\theta)j^{-\beta}T^{1-\theta}\right\} + c^{2} \sum_{j \geq \lfloor T^{\frac{\theta}{2\beta-1}} \rfloor + 1} j^{-2\beta} \exp\left\{-bH(\theta)j^{-\beta}T^{1-\theta}\right\}$$

$$\leq c^{2} \sum_{j=1}^{\infty} j^{-2\beta} \exp\left\{-bH(\theta)T^{-\frac{\beta\theta}{2\beta-1}}T^{1-\theta}\right\} + \frac{c^{2}}{2\beta-1} \left(\lfloor T^{\frac{\theta}{2\beta-1}} \rfloor + 1\right)^{1-2\beta}$$

$$\leq c^{2} \exp\left\{-bH(\theta)T^{1-\frac{3\beta-1}{2\beta-1}\theta}\right\} \sum_{j=1}^{\infty} j^{-2\beta} + \frac{c^{2}}{2\beta-1}T^{-\theta}$$

$$\leq \frac{2\beta c^{2}}{2\beta-1} \left(\exp\left\{-bH(\theta)T^{1-\frac{3\beta-1}{2\beta-1}\theta}\right\} + T^{-\theta}\right).$$

(2) When $\frac{2\beta-1}{3\beta-1} \le \theta < 1$,

$$\sum_{j=1}^{\infty} \mu_j^2 \exp\left\{-H(\theta)(\lambda + \mu_j)T^{1-\theta}\right\} \le c^2 \sum_{j=1}^{\infty} j^{-2\beta} \exp\left\{-H(\theta)\lambda T^{1-\theta}\right\}$$
$$\le \frac{2\beta c^2}{2\beta - 1} \exp\left\{-H(\theta)\lambda T^{1-\theta}\right\}.$$

References

- [1] Z. Allen-Zhu, Y. Li, and Z. Song, A convergence theory for deep learning via over-parameterization, in International Conference on Machine Learning, PMLR, 2019, pp. 242–252.
- [2] —, On the convergence rate of training recurrent neural networks, Advances in Neural Information Processing Systems, 32 (2019).
- [3] S. Arora, S. Du, W. Hu, Z. Li, and R. Wang, Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks, in International Conference on Machine Learning, PMLR, 2019, pp. 322–332.
- [4] R. Bhatia and L. Elsner, The hoffman-wielandt inequality in infinite dimensions, in Proceedings of the Indian Academy of Sciences-Mathematical Sciences, vol. 104, Springer, 1994, pp. 483–494.
- [5] S. BOUCHERON, G. LUGOSI, AND O. BOUSQUET, *Concentration inequalities*, in Summer School on Machine Learning, Springer, 2003, pp. 208–240.
- [6] A. Brutzkus, A. Globerson, E. Malach, and S. Shalev-Shwartz, Sgd learns over-parameterized networks that provably generalize on linearly separable data, in International Conference on Learning Representations, 2018.
- [7] Y. CAO AND Q. Gu, Generalization bounds of stochastic gradient descent for wide and deep neural networks, Advances in Neural Information Processing Systems, 32 (2019).
- [8] A. CAPONNETTO AND E. DE VITO, Optimal rates for the regularized least-squares algorithm, Foundations of Computational Mathematics, 7 (2007), pp. 331–368.
- [9] L. Carratino, A. Rudi, and L. Rosasco, Learning with sgd and random features, Advances in Neural Information Processing Systems, 31 (2018).
- [10] X. Chen, B. Tang, J. Fan, and X. Guo, Online gradient descent algorithms for functional data learning, Journal of Complexity, 70 (2022), p. 101635.
- [11] F. CUCKER AND S. SMALE, On the mathematical foundations of learning, Bulletin of the American Mathematical Society, 39 (2002), pp. 1–49.
- [12] F. Cucker and D. X. Zhou, Learning theory: an approximation theory viewpoint, vol. 24, Cambridge University Press, 2007.
- [13] A. DIEULEVEUT AND F. BACH, Nonparametric stochastic approximation with large stepsizes, The Annals of Statistics, 44 (2016), pp. 1363–1399.
- [14] R. GE, J. D. LEE, AND T. MA, Learning one-hidden-layer neural networks with land-scape design, in International Conference on Learning Representations, 2018.
- [15] X. Guo, Z.-C. Guo, and L. Shi, Capacity dependent analysis for functional online learning algorithms, Applied and Computational Harmonic Analysis, 67 (2023), p. 101567.
- [16] Z.-C. Guo, A. Christmann, and L. Shi, *Optimality of robust online learning*, Foundations of Computational Mathematics, (2023), pp. 1–29.

- [17] Z.-C. Guo and L. Shi, Fast and strong convergence of online learning algorithms, Advances in Computational Mathematics, 45 (2019), pp. 2745–2770.
- [18] A. Jacot, F. Gabriel, and C. Hongler, Neural tangent kernel: Convergence and generalization in neural networks, Advances in Neural Information Processing Systems, 31 (2018).
- [19] Y. Lei, R. Jin, and Y. Ying, Stability and generalization analysis of gradient methods for shallow neural networks, Advances in Neural Information Processing Systems, 35 (2022).
- [20] Y. Lei, L. Shi, and Z.-C. Guo, Convergence of unregularized online learning algorithms, The Journal of Machine Learning Research, 18 (2017), pp. 6269–6301.
- [21] Y. Li and Y. Liang, Learning overparameterized neural networks via stochastic gradient descent on structured data, Advances in Neural Information Processing Systems, 31 (2018).
- [22] Y. LI AND Y. YUAN, Convergence analysis of two-layer neural networks with relu activation, Advances in Neural Information Processing Systems, 30 (2017).
- [23] S.-B. Lin, K. Wang, Y. Wang, and D.-X. Zhou, Universal consistency of deep convolutional neural networks, IEEE Transactions on Information Theory, 68 (2022), pp. 4610–4617.
- [24] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro, *Robust stochastic approximation approach to stochastic programming*, SIAM Journal on Optimization, 19 (2009), pp. 1574–1609.
- [25] M. NGUYEN AND N. MÜCKE, How many neurons do we need? a refined analysis for shallow networks trained with gradient descent, Journal of Statistical Planning and Inference, 233 (2024), p. 106169.
- [26] A. NITANDA AND T. SUZUKI, Optimal rates for averaged stochastic gradient descent under neural tangent kernel regime, in International Conference on Learning Representations, 2020.
- [27] L. PILLAUD-VIVIEN, A. RUDI, AND F. BACH, Exponential convergence of testing error for stochastic gradient methods, in Conference on Learning Theory, PMLR, 2018, pp. 250–296.
- [28] A. RAKHLIN, O. SHAMIR, AND K. SRIDHARAN, Making gradient descent optimal for strongly convex stochastic optimization, arXiv preprint arXiv:1109.5647, (2011).
- [29] S. SHALEV-SHWARTZ, Y. SINGER, AND N. SREBRO, Pegasos: Primal estimated subgradient solver for svm, in International Conference on Machine Learning, 2007, pp. 807– 814.
- [30] L. Shi and J.-Q. Yang, Learning operators with stochastic gradient descent in general hilbert spaces, arXiv preprint arXiv:2402.04691, (2024).
- [31] I. Steinwart and A. Christmann, Support vector machines, Springer Science & Business Media, 2008.

- [32] L. Su and P. Yang, On learning over-parameterized neural networks: A functional approximation perspective, Advances in Neural Information Processing Systems, 32 (2019).
- [33] P. Tarres and Y. Yao, Online learning as stochastic approximation of regularization paths: Optimality and almost-sure convergence, IEEE Transactions on Information Theory, 60 (2014), pp. 5716–5735.
- [34] Y.-G. WANG AND Z.-C. Guo, Differentially private sgd with random features, Applied Mathematics-A Journal of Chinese Universities, 39 (2024), pp. 1–23.
- [35] Y. YAO, On complexity issues of online learning algorithms, IEEE Transactions on Information Theory, 56 (2010), pp. 6470–6481.
- [36] D. Yarotsky, Error bounds for approximations with deep relu networks, Neural Networks, 94 (2017), pp. 103–114.
- [37] Y. Ying and M. Pontil, *Online gradient descent learning algorithms*, Foundations of Computational Mathematics, 8 (2008), pp. 561–596.
- [38] D.-X. Zhou, Universality of deep convolutional neural networks, Applied and Computational Harmonic Analysis, 48 (2020), pp. 787–794.
- [39] D. ZOU, Y. CAO, D. ZHOU, AND Q. GU, Gradient descent optimizes over-parameterized deep relu networks, Machine Learning, 109 (2020), pp. 467–492.