

OPTIMAL FUNCTION SPACES IN WEIGHTED SOBOLEV EMBEDDINGS WITH α -HOMOGENEOUS WEIGHTS

LADISLAV DRÁŽNÝ

ABSTRACT. We study weighted Sobolev inequalities on open convex cones endowed with α -homogeneous weights satisfying a certain concavity condition. We establish a so-called reduction principle for these inequalities and characterize optimal rearrangement-invariant function spaces for these weighted Sobolev inequalities. Both optimal target and optimal domain spaces are characterized. Abstract results are accompanied by general yet concrete examples of optimal function spaces. For these examples, the class of so-called Lorentz–Karamata spaces, which contains in particular Lebesgue spaces, Lorentz spaces, and some Orlicz spaces, is used.

1. INTRODUCTION

Various Sobolev spaces and inequalities have played important roles in mathematics for decades. Their applications include (but are not limited to) analysis of partial differential equations, calculus of variations, or harmonic analysis. In this paper, we study weighted Sobolev inequalities on open convex cones Σ in \mathbb{R}^n with α -homogeneous weights $w: \Sigma \rightarrow [0, \infty)$ and establish their optimal, in a sense, versions. Throughout the entire paper, we assume that $n \in \mathbb{N}$, the dimension of \mathbb{R}^n , is greater than 1. The weight w is a nonnegative continuous α -homogeneous function for some $\alpha > 0$ such that $w^{\frac{1}{\alpha}}$ is concave in Σ . We define the weighted measure μ on Σ as

$$\mu(E) = \int_E w(x) dx$$

for every Lebesgue measurable set $E \subseteq \Sigma$.

An important example of an admissible weight is the monomial weight defined as $w(x) = x_1^{A_1} \cdots x_k^{A_k}$ on the cone $\Sigma = \{x_i > 0, i = 1, \dots, k\}$ for some $1 \leq k \leq n$ and $A_i > 0$, which is $\alpha = A_1 + \cdots + A_k$ homogeneous. These monomial weights and corresponding weighted Sobolev inequalities

Date: July 11, 2024.

2020 Mathematics Subject Classification. 46E30, 46E35.

Key words and phrases. rearrangement-invariant function spaces, Sobolev embeddings, weighted Sobolev spaces, optimal function spaces, Lorentz–Karamata spaces, reduction principle.

This research was partly supported by grant No. 23-04720S of the Czech Science Foundation and by the Primus research programme PRIMUS/21/SCI/002 of Charles University.

have been quite fashionable recently (e.g., [8, 9, 21, 24, 49], to name a few). Their importance was observed in particular in [5], where they were used in connection with the regularity of stable solutions to certain planar reaction–diffusion problems, and they were studied in more detail in [6]. In particular, in the latter, they established a new isoperimetric inequality for the monomial weights. What is quite surprising and remarkable is that the isoperimetric quotient is minimized by intersections of balls with the cone Σ despite the fact that the monomial weights are not radially symmetric. Using their isoperimetric inequality, they also established the corresponding Sobolev inequality, namely for $p \in [1, D)$

$$\|u\|_{L^{p^*}(\Sigma, \mu)} \leq C_{D,p} \|\nabla u\|_{L^p(\Sigma, \mu)} \quad \text{for every } u \in C_c^1(\mathbb{R}^n),$$

where $D = n + k$ and $p^* = (Dp)/(n - p)$. Notice that the usual role of the dimension n is replaced by the sum of the dimension and the order of homogeneity of the weight. Furthermore, note that the presence of \mathbb{R}^n (instead of Σ) in $C_c^1(\mathbb{R}^n)$ is not a typo, because the functions are not required to vanish on the boundary of Σ .

Later, the isoperimetric inequality for the monomial weights was greatly generalized in [7] (see also [15]). In particular, their general isoperimetric inequality applies to the general setting of convex cones with α -homogeneous weights considered in this paper. Noteworthily, arguments based on symmetrization can often be successfully used even though the weights need not be radially symmetric. Besides the monomial weights, some examples of admissible weights and cones are the following. If $\Sigma = (0, \infty)^n$, then we can consider the weight $w(x) = (A_1 x_1^{\frac{1}{r}} + \dots + A_n x_n^{\frac{1}{r}})^{\alpha r}$ for $r \geq 1, A_i \geq 0$, and $\alpha > 0$. For $\Sigma \neq \mathbb{R}^n$, the weight $w(x) = \text{dist}(x, \partial\Sigma)^\alpha$, $\alpha > 0$ is admissible. More examples of admissible weights can be found in [7]. It is worth pointing out that it may happen that the weight w does not vanish on $\partial\Sigma$. Such weights can be obtained for instance by restricting an admissible weight w to a subcone of Σ .

In this paper, we study Sobolev inequalities of the form

$$\|u\|_{Y(\Sigma, \mu)} \leq C \|\nabla^m u\|_{X(\Sigma, \mu)} \quad \text{for every } u \in V_0^m X(\Sigma, \mu), \quad (1.1)$$

where (see Section 2 for precise definitions) $X(\Sigma, \mu)$ and $Y(\Sigma, \mu)$ are rearrangement-invariant function spaces on Σ endowed with the measure μ and $V_0^m X(\Sigma, \mu)$ is a suitable m th order ($1 \leq m < D$) Sobolev-type space built upon $X(\Sigma, \mu)$. We study the question of optimal function spaces in (1.1). On the one hand, for a given rearrangement-invariant function space $X(\Sigma, \mu)$, we will describe the optimal rearrangement-invariant function space $Y(\Sigma, \mu)$ with which the inequality (1.1) holds (see Theorem 3.8). The optimality is understood in the sense that $Y(\Sigma, \mu)$ cannot be replaced by a strictly smaller rearrangement-invariant function space. On the other hand, for a given $Y(\Sigma, \mu)$, we will describe the optimal rearrangement-invariant function space $X(\Sigma, \mu)$ with which the inequality (1.1) is valid (see Theorem 3.9).

This time, the optimality is understood in the sense that $X(\Sigma, \mu)$ cannot be replaced by a strictly larger rearrangement-invariant function space.

The point of departure for us is the suitable isoperimetric inequality obtained in [7], which we exploit to obtain a suitable weighted Pólya–Szegő inequality (see Proposition 3.1). By utilizing it and combining it (when $m > 1$) with the clever idea of iteration (see [13]), we prove a so-called reduction principle for the inequality (1.1) (see Theorem 3.2). To establish these results, we need to exploit a lot of different techniques developed and improved over time together with results from both classical and contemporary theory of (rearrangement-invariant) function spaces. The question of optimal rearrangement-invariant function spaces for a large number of various Sobolev inequalities has been intensively studied for more than two decades (e.g., [1, 11, 12, 13, 14, 17, 23, 28, 29]). Nevertheless, the setting considered in this paper appears not be covered, and so we aim to fill this gap in the actively developing setting of Sobolev inequalities on convex cones with α -homogeneous weights. Furthermore, abstract theorems are accompanied by several concrete (yet substantially general) examples of optimal function space in the inequality (1.1) (see Section 4).

The class of rearrangement-invariant function spaces is very general and contains a large number of classical function spaces. We will briefly informally introduce this class of function spaces here. Loosely speaking, rearrangement-invariant function spaces are often suitable for measuring integrability and, arguably, their most fundamental feature is that rearrangement-invariant function norms depend only on the measure of level sets. By that we mean that if u and v are two measurable functions such that the measures of the sets $\{x: |u(x)| > \lambda\}$ and $\{x: |v(x)| > \lambda\}$ are the same for every $\lambda > 0$, then their norms are equal. For example, the rearrangement invariance of the L^p norms follows from the well-known layer cake representation formula ([26, Theorem 1.13]). Apart from the Lebesgue spaces, other well-known examples of rearrangement-invariant function spaces are Lorentz spaces $L^{p,q}$ or Orlicz spaces L^A . Orlicz spaces naturally generalize Lebesgue spaces and their usefulness stems from their usage in analysis of nonlinear partial differential equations and variational problems whose nonlinearity is non-polynomial. Very loosely speaking, they measure integrability in a more fine-grained way by replacing power functions by general convex functions (i.e., $|f(x)|^p$ versus $A(|f(x)|)$), which allows for capturing non-polynomial growth. Lorentz spaces, which contain the famous weak Lebesgue spaces (corresponding to $q = \infty$), are not only intimately connected with the interpolation theory and harmonic analysis (e.g., [3, 20, 44]) but also with the theory of Sobolev spaces. For example, assuming $n > 1$, while a function whose weak gradient (locally) belongs to the Lebesgue space $L^n(\mathbb{R}^n)$ need not be (locally) bounded or be differentiable (in the classical sense) at any point, the situation changes dramatically when its weak gradient (locally) belongs to the Lorentz space $L^{n,1}(\mathbb{R}^n)$, which is a slightly (yet essentially) smaller function space than $L^n(\mathbb{R}^n)$. In the latter case, not only

is the function (locally) bounded, but it has a continuous representative that is differentiable almost everywhere (see [43]). Another well-known result is that the classical Sobolev embedding into the critical Lebesgue space L^{p^*} can be improved by replacing L^{p^*} with the Lorentz space $L^{p^*,p} \subsetneq L^{p^*}$ (e.g., [35, 37, 47]). This improvement can also be essential in analysis of partial differential equations and variational problems with critical growth (see [42]).

A considerably general (yet still reasonably concrete) subclass of rearrangement-invariant function spaces is constituted by so-called Lorentz–Karamata spaces (see [16, 17, 34, 39]). Among other more delicate function spaces, it contains not only Lebesgue spaces and Lorentz spaces but also a lot of Orlicz spaces (in particular, those of “exponential” and “logarithmic” type (see also [36])). We will consider this class of function spaces in Section 4, where we provide concrete examples of optimal function norms in (1.1). With this choice, not only can we describe the optimal function spaces explicitly, but it also enables us to capture delicate integrability properties in limiting situations (in particular, loosely speaking when X is “close to” $L^{\frac{D}{m}}$ (cf. [4, 22, 31, 48])).

2. PRELIMINARIES

In the whole paper we use the convention $\frac{1}{\infty} = 0$ and $0 \cdot \infty = 0$. When $E \subseteq (0, \infty)$ is Lebesgue measurable, we denote by $\lambda(E)$ its Lebesgue measure.

Let (R, μ) be a σ -finite nonatomic measure space. By $\mathcal{M}(R, \mu)$ we will denote the class of all μ -measurable functions on R whose values lie in $\mathbb{R} \cup \{-\infty, \infty\}$. We will denote the class of all μ -measurable functions on R whose values lie in $[0, \infty]$ by $\mathcal{M}^+(R, \mu)$. And the class of all functions in $\mathcal{M}(R, \mu)$ that are finite μ -almost everywhere in R will be denoted by $\mathcal{M}_0(R, \mu)$.

Now we introduce rearrangement-invariant Banach function spaces and their basic properties. The theory that is presented here except of the Sobolev spaces follows the first three chapters of [2].

Let $f \in \mathcal{M}(R, \mu)$. The *nonincreasing rearrangement* of the function f is the function $f_\mu^*: (0, \infty) \rightarrow [0, \infty]$ defined by

$$f_\mu^*(t) = \inf_{\lambda > 0} (\mu(\{x \in R; |f(x)| > \lambda\}) \leq t) \quad t \in (0, \infty).$$

The function f_μ^* is nonincreasing and right-continuous. If $f \in \mathcal{M}(R, \mu)$, $g \in \mathcal{M}(S, \nu)$ satisfy

$$f_\mu^*(t) = g_\nu^*(t), \quad t \in (0, \infty), \quad (2.1)$$

we say that f and g are *equimeasurable*. For instance, the functions f and f_μ^* are equimeasurable.

A mapping ρ on $\mathcal{M}^+(R, \mu)$ with values in $[0, \infty]$ we call a *rearrangement-invariant Banach function norm* if all the following properties are satisfied

for all $f, g \in \mathcal{M}^+(R, \mu)$, $\{f_k; k \in \mathbb{N}\} \subseteq \mathcal{M}^+(R, \mu)$, $c \in [0, \infty)$ and $A \subseteq R$ such that A is μ -measurable.

- (1) the *norm axiom*: $\rho(f) = 0$ if and only if $f = 0$ μ -almost everywhere in R , $\rho(cf) = c\rho(f)$, $\rho(f + g) \leq \rho(f) + \rho(g)$;
- (2) the *lattice axiom*: if $g \leq f$ μ -almost everywhere in R , then $\rho(g) \leq \rho(f)$;
- (3) the *Fatou axiom*: if $f_k \uparrow f$ μ -almost everywhere in R , then $\rho(f_k) \uparrow \rho(f)$;
- (4) the *nontriviality axiom*: if $\mu(A) < \infty$, then $\rho(\chi_A) < \infty$;
- (5) the *local embedding in L^1* : if $\mu(A) < \infty$, then

$$\int_A f d\mu \leq K_A \rho(f), \quad (2.2)$$

where $K_A \geq 0$ is a constant which may depend on A but which does not depend on f ;

- (6) the *rearrangement-invariance axiom*: if $f_\mu^* = g_\mu^*$, then $\rho(f) = \rho(g)$.

The collection of all functions $f \in \mathcal{M}(R, \mu)$ such that $\rho(|f|) < \infty$ is called a *rearrangement-invariant Banach function space*. We will denote it by $X(\rho)$, $X(R, \mu)$ or just by X .

As their name suggests, rearrangement-invariant Banach function spaces are Banach spaces. Textbook examples of rearrangement-invariant spaces are the Lebesgue spaces $L^p(R, \mu)$ for $p \in [1, \infty]$. Every rearrangement-invariant space contains simple functions (i.e., linear combinations of characteristic functions of μ -measurable sets of finite measure) and is contained in $\mathcal{M}_0(R, \mu)$.

Now we define suitable weighted Sobolev spaces built on rearrangement-invariant spaces. We start with some notation and conventions used in the rest of this paper. Throughout the rest of this paper, we assume that $n \in \mathbb{N}$, $n \geq 2$, is the dimension of \mathbb{R}^n . Furthermore, we assume that Σ is a nonempty open convex cone with vertex at the origin, i.e., $\Sigma \subseteq \mathbb{R}^n$ is a nonempty open convex set such that for every $x \in \Sigma$ and for every $r > 0$, we have $rx \in \Sigma$. We also assume that $w: \overline{\Sigma} \rightarrow [0, \infty)$ is a nonnegative continuous function that is not identically zero. Furthermore, we assume that w is α -homogeneous for some $\alpha > 0$, i.e., for every $x \in \overline{\Sigma}$ and for every $s > 0$, we have $w(sx) = s^\alpha w(x)$. Finally, we assume that the function $w^{\frac{1}{\alpha}}$ is concave in Σ . We set

$$D = n + \alpha. \quad (2.3)$$

We assume that $m \in \mathbb{N}$ is such that

$$1 \leq m < D. \quad (2.4)$$

We define the weighted measure μ on Σ as

$$\mu(E) = \int_E w(x) dx$$

for every Lebesgue measurable set $E \subseteq \Sigma$.

We now introduce the Sobolev spaces that we will work with. Let $k \in \mathbb{N}$ and let $u: \Sigma \rightarrow \mathbb{R}$ be k -times weakly differentiable function in Σ (i.e., it has all weak derivatives up to the k -th order). We denote by $\nabla^l u$, $l \in \{1, \dots, k\}$, the vector of all l -th order weak derivatives of u . We also set $\nabla^0 u = u$.

Let $X(\Sigma, \mu)$ be a rearrangement-invariant space. We say that u belongs to the space $V^k X(\Sigma, \mu)$ if

$$|\nabla^k u| \in X(\Sigma, \mu).$$

We say that u belongs to the space $V_0^k X(\Sigma, \mu)$ if $u \in V^k X(\Sigma, \mu)$ and for every $l \in \{0, 1, \dots, k-1\}$ and for every $\lambda > 0$ it holds that

$$\mu\left(\left\{x \in \Sigma; |\nabla^l u(x)| > \lambda\right\}\right) < \infty.$$

For short, we will write $\|\nabla^k u\|_{X(\Sigma, \mu)}$ instead of $\| |\nabla^k u| \|_{X(\Sigma, \mu)}$.

We will also encounter Sobolev space $W^{1,1}(\Sigma, \mu)$, which is a weighted counterpart of the classical Sobolev space $W^{1,1}(\Sigma)$. We say that a function u belongs to the space $W^{1,1}(\Sigma, \mu)$ if it is weakly differentiable in Σ , $u \in L^1(\Sigma, \mu)$ and $|\nabla u| \in L^1(\Sigma, \mu)$.

Now we turn back to the theory of rearrangement-invariant spaces. We present here their another important properties which we will need in what follows.

To every rearrangement-invariant space X , there is associated another rearrangement-invariant space, which is related to its continuous dual, but which is usually more useful in the theory of Banach function spaces. The mapping ρ' defined on $\mathcal{M}^+(R, \mu)$ by

$$\rho'(g) = \sup_{f \in \mathcal{M}^+(R, \mu), \rho(f) \leq 1} \int_R fg \, d\mu, \quad g \in \mathcal{M}^+(R, \mu), \quad (2.5)$$

is the *associate norm* of the function norm ρ . We say that the space $X(\rho')$ is the associate space to the space $X(\rho)$ and we denote this space by X' .

For example, when $X = L^p(R, \mu)$ for $p \in [1, \infty]$, then $X' = L^{p'}(R, \mu)$. Here $p' \in [1, \infty]$ is the dual index defined by $\frac{1}{p} + \frac{1}{p'} = 1$.

An important property of Banach function spaces is that, if X is a Banach function space, then $(X')' = X$.

For every Banach function norm ρ , the *Hölder inequality*

$$\int_R |fg| \, d\mu \leq \rho(f)\rho'(g)$$

holds for every $f, g \in \mathcal{M}(R, \mu)$.

For every $f \in \mathcal{M}(R, \mu)$ and $t \in (0, \infty)$, we have

$$\mu(\{x \in R; |f(x)| > f_\mu^*(t)\}) \leq t. \quad (2.6)$$

Furthermore, if $f_\mu^*(t) < \infty$ and $\mu(\{x \in R; |f(x)| > f_\mu^*(t) - \varepsilon\}) < \infty$ for some $\varepsilon > 0$, then

$$\mu(\{x \in R; |f(x)| \geq f_\mu^*(t)\}) \geq t. \quad (2.7)$$

The *fundamental function* $\varphi_X: [0, \mu(R)] \rightarrow [0, \infty]$ of the rearrangement-invariant space X is the mapping defined by

$$\varphi_X(t) = \|\chi_E\|_X, \quad t \in [0, \mu(R)],$$

where $E \subseteq R$ is an arbitrary set satisfying $\mu(E) = t$. The definition is correct since if $E, F \subseteq R$ are sets such that $\mu(E) = \mu(F)$, then their characteristic functions χ_E and χ_F are equimeasurable.

Furthermore, if $g \in \mathcal{M}(R, \mu)$ and if ρ is a rearrangement-invariant norm, we have

$$\rho'(g) = \sup_{f \in \mathcal{M}^+(R, \mu), \rho(f) \leq 1} \int_R f^* g^* d\mu. \quad (2.8)$$

The *Hardy–Littlewood inequality* is very important in the theory of rearrangement-invariant spaces. It states that

$$\int_R |fg| d\mu \leq \int_0^\infty f^*(t)g^*(t) dt$$

for every $f, g \in \mathcal{M}(R, \mu)$. In particular, by taking $g = \chi_E$, we have

$$\int_E |f| d\mu \leq \int_0^{\mu(E)} f^*(t) dt \quad (2.9)$$

for each μ -measurable set $E \subseteq R$.

Another important result in the theory of rearrangement-invariant spaces is the *Hardy–Littlewood–Pólya principle*. For every rearrangement-invariant norm ρ , if $f, g \in \mathcal{M}^+(R, \mu)$ are such that

$$\int_0^t f^*(\tau) d\tau \leq \int_0^t g^*(\tau) d\tau$$

for every $t \in (0, \infty)$, then

$$\rho(f) \leq \rho(g). \quad (2.10)$$

We will also need the following fact. For every $t \in (0, \mu(R))$ and for every $f \in \mathcal{M}(R, \mu)$, we have

$$\int_0^t f^*(\tau) d\tau = \sup \left(\left\{ \int_E |f| d\mu; E \subseteq R, E \text{ } \mu\text{-measurable, } \mu(E) = t \right\} \right). \quad (2.11)$$

Each rearrangement-invariant space on (R, μ) can be represented as a rearrangement-invariant space on $(0, \mu(R))$. More precisely, if $X(R, \mu)$ is a

rearrangement-invariant space, then there exists the unique rearrangement-invariant space $X(0, \mu(R))$ such that for every function $f \in X(R, \mu)$ it holds that

$$\|f\|_{X(R, \mu)} = \|f_\mu^*\|_{X(0, \mu(R))}.$$

The rearrangement-invariant space $X(0, \mu(R))$ is called *the representation space* of $X(R, \mu)$. For example, if $X(R, \mu) = L^p(R, \mu)$, then $X(0, \mu(R)) = L^p(0, \mu(R))$.

On the other hand, for every $f \in \mathcal{M}(0, \mu(R))$, there exists a function $u \in \mathcal{M}(R, \mu)$ such that $f_\lambda^*(t) = u_\mu^*(t)$ for every $t \in (0, \infty)$.

Closely related to the nonincreasing rearrangement is *the maximal non-increasing rearrangement*. The *maximal nonincreasing operator*

$$P_\mu: \mathcal{M}(R, \mu) \rightarrow \mathcal{M}^+(0, \infty)$$

is defined by

$$P_\mu(f)(t) = \frac{1}{t} \int_0^t f_\mu^*(\tau) d\tau, \quad f \in \mathcal{M}(R, \mu), t \in (0, \infty).$$

The image of a function $f \in \mathcal{M}(R, \mu)$ under the maximal nonincreasing operator P_μ is also commonly denoted by f_μ^{**} , and it is called *the maximal nonincreasing function*. The maximal nonincreasing function is nonincreasing and we have $f^* \leq f^{**}$.

If $X(0, \mu(R))$ is a rearrangement-invariant space and $h \in \mathcal{M}^+(0, \mu(R))$ is a nonincreasing function, we know thanks to (2.8) that

$$\|h\|_{X'(0, \mu(R))} = \sup_{g \in \mathcal{M}^+(0, \mu(R)), \|g\|_{X(0, \mu(R))} \leq 1} \int_0^{\mu(R)} h(t)g^*(t) dt.$$

In general, when $h \in \mathcal{M}^+(0, \mu(R))$ is not necessarily nonincreasing, we only have

$$\|h\|_{X'(0, \mu(R))} \geq \sup_{g \in \mathcal{M}^+(0, \mu(R)), \|g\|_{X(0, \mu(R))} \leq 1} \int_0^{\mu(R)} h(t)g^*(t) dt \quad (2.12)$$

owing to (2.5). However, it follows from [13, Theorem 9.5] and [38, Theorem 3.10] that

$$\|t^\alpha f^{**}(t)\|_{X'(0, \mu(R))} \leq 4 \sup_{g \in \mathcal{M}^+(0, \mu(R)), \|g\|_{X(0, \mu(R))} \leq 1} \int_0^{\mu(R)} t^\alpha f^{**}(t)g^*(t) dt \quad (2.13)$$

for every $\alpha \in [0, 1]$ and $f \in \mathcal{M}(R, \mu)$. Inequalities (2.12) and (2.13) mean that the norm of $t^\alpha f^{**}(t)$ can be approached, up to a multiplicative constant, by nonincreasing functions in this case even though the function $t^\alpha f^{**}(t)$ does not have to be nonincreasing.

We will continue by introducing the dilation operator. Let $\alpha \in (0, \infty)$. The *dilation operator* $D_\alpha: \mathcal{M}^+(0, \infty) \rightarrow \mathcal{M}^+(0, \infty)$ is defined by $(D_\alpha f)(t) = f(\alpha t)$ for each $f \in \mathcal{M}^+(0, \infty)$ and $t \in (0, \infty)$. This operator is bounded on

every rearrangement-invariant space over $(0, \infty)$. More precisely, there exists a constant $0 < C \leq \max\{1, \frac{1}{\alpha}\}$ such that

$$\|D_\alpha f\|_{X(0, \infty)} \leq C \|f\|_{X(0, \infty)} \quad (2.14)$$

for every $f \in \mathcal{M}^+(0, \infty)$, every $\alpha \in (0, \infty)$ and every rearrangement-invariant space $X(0, \infty)$.

We conclude this section by introduction the continuous embedding. We say that $X(R, \mu)$ is *continuously embedded* into $Y(R, \mu)$ if for every function $u \in X(R, \mu)$ it holds that $u \in Y(R, \mu)$ and that $\|u\|_{Y(R, \mu)} \leq C \|u\|_{X(R, \mu)}$, where C is a constant that does not depend on u . We denote the fact that $X(R, \mu)$ is continuously embedded into $Y(R, \mu)$ by $X(R, \mu) \hookrightarrow Y(R, \mu)$. In fact, inclusion between Banach function spaces is always continuous in the sense that $X(R, \mu) \hookrightarrow Y(R, \mu)$ if and only if $X(R, \mu) \subseteq Y(R, \mu)$. If $X(R, \mu), Y(R, \mu)$ are Banach function spaces, it holds that $X(R, \mu) \hookrightarrow Y(R, \mu)$ if and only if $Y'(R, \mu) \hookrightarrow X'(R, \mu)$.

In the rest of the paper, we will denote by C, K, C_i, K_i positive finite constants whose exact values are not important for our purposes.

3. REDUCTION PRINCIPLE AND OPTIMALITY

3.1. Reduction principle. The goal of this subsection is to prove a suitable reduction principle. To prove it we need to derive a variant of the Pólya–Szegő inequality. The prove of this theorem is at the end of this section.

Proposition 3.1 (Pólya–Szegő inequality). *Let X be a rearrangement-invariant space over (Σ, μ) and $u \in V_0^1 X(\Sigma, \mu)$. Then u_μ^* is a locally absolutely continuous function on the interval $(0, \infty)$, and it holds that*

$$\left\| t^{\frac{D-1}{D}} \frac{du_\mu^*}{dt}(t) \right\|_{X(0, \infty)} \leq C \|\nabla u\|_{X(\Sigma, \mu)}, \quad (3.1)$$

where C is a positive constant independent of u .

In the remaining part of this section we prove the reduction principle. Recall that the parameteres m and D were introduced in (2.3) and (2.4).

Theorem 3.2 (Reduction principle). *Let X and Y be rearrangement-invariant spaces over (Σ, μ) . Then the following two statements are equivalent.*

(1) *For all functions $v \in V_0^m X(\Sigma, \mu)$ it holds that*

$$\|v\|_{Y(\Sigma, \mu)} \leq C_1 \|\nabla^m v\|_{X(\Sigma, \mu)}. \quad (3.2)$$

(2) *For all functions $f \in \mathcal{M}^+(0, \infty)$ it holds that*

$$\left\| \int_t^\infty f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \leq C_2 \|f\|_{X(0, \infty)}. \quad (3.3)$$

Here, C_1 and C_2 are positive constants independent of v and of f respectively.

The proof of this theorem will be divided into two steps, Proposition 3.5 and Proposition 3.7.

Remark 3.3. As an easy consequence of the definition of the associate norm, we obtain the fact that (3.3) is equivalent to the following assertion:

(2*) For all functions $g \in \mathcal{M}^+(0, \infty)$ it holds that

$$\left\| t^{\frac{m}{D}} g^{**}(t) \right\|_{X'(0, \infty)} \leq C_2 \|g\|_{Y'(0, \infty)}. \quad (3.4)$$

Remark 3.4. We have

$$\begin{aligned} & \left\| \chi_{(0,1)}(t) \right\|_{Y(0, \infty)} \left\| t^{\frac{m}{D}-1} \chi_{(1, \infty)}(t) \right\|_{X'(0, \infty)} \\ &= \left\| \chi_{(0,1)}(t) \right\|_{Y(0, \infty)} \sup_{\|h\|_{X(0, \infty)}, h \geq 0} \int_1^\infty \tau^{\frac{m}{D}-1} h(\tau) d\tau \\ &\leq \sup_{\|h\|_{X(0, \infty)}, h \geq 0} \left\| \chi_{(0,1)}(t) \int_t^\infty \tau^{\frac{m}{D}-1} h(\tau) d\tau \right\|_{Y(0, \infty)}, \end{aligned}$$

so, (3.3) implies that

$$t^{\frac{m}{D}-1} \chi_{(1, \infty)}(t) \in X'(0, \infty).$$

Proposition 3.5. Let X, Y be rearrangement-invariant spaces over (Σ, μ) . Assume that there exists a positive constant C_1 such that for all functions $v \in V_0^m X(\Sigma, \mu)$ it holds that

$$\|v\|_{Y(\Sigma, \mu)} \leq C_1 \|\nabla^m v\|_{X(\Sigma, \mu)}.$$

Then there exists a positive constant C_2 such that for all functions $f \in \mathcal{M}^+(0, \infty)$ it holds that

$$\left\| \int_t^\infty f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \leq C_2 \|f\|_{X(0, \infty)}. \quad (3.5)$$

The constant C_2 depends only on the constant C_1 , on m and on D .

Owing to this proposition the n -dimensional part of the reduction principle ((3.2) in Theorem 3.2) implies the one-dimensional part ((3.3) in Theorem 3.2). In the proof of the proposition, we need to use the following technical lemma, whose proof is straightforward and omitted.

Lemma 3.6. Let $f \in \mathcal{M}^+(0, \infty) \cap L^\infty(0, \infty)$ be a function with a bounded support. Let $g: [0, \infty) \rightarrow [0, \infty)$ be the function defined by

$$g(t) = \int_t^\infty f(\tau) \tau^{\frac{m}{D}-m} (\tau-t)^{m-1} d\tau, \quad t \in [0, \infty). \quad (3.6)$$

Then $g \in \mathcal{C}^{m-1}(0, \infty)$ and

$$g^{(j)}(t) = (-1)^j \frac{(m-1)!}{(m-j-1)!} \int_t^\infty f(\tau) \tau^{\frac{m}{D}-m} (\tau-t)^{m-j-1} d\tau, \quad t \in (0, \infty)$$

for every $j \in \{1, \dots, m-1\}$. Moreover, $g^{(m-1)}$ is locally Lipschitz on $(0, \infty)$ and

$$g^{(m)}(t) = (-1)^m (m-1)! f(t) t^{\frac{m}{D}-m}$$

for almost every $t \in (0, \infty)$.

We denote by B_μ the μ -measure of the intersection of the unit ball in \mathbb{R}^n with Σ , i.e.,

$$B_\mu = \mu(\{x \in \Sigma; |x| \leq 1\}).$$

Now we prove Proposition 3.5.

Proof of Proposition 3.5. Choose an arbitrary function $f \in \mathcal{M}^+(0, \infty)$. Observe that it is enough to prove the theorem for $\|f\|_{X(0, \infty)} < \infty$. Firstly, assume that the function f belongs to $L^\infty(0, \infty)$ and that it has a bounded support. Define the function $v: \Sigma \rightarrow [0, \infty)$ by

$$v(x) = \int_{B_\mu |x|^D}^{\infty} f(\tau) \tau^{\frac{m}{D}-m} (\tau - B_\mu |x|^D)^{m-1} d\tau, \quad x \in \Sigma. \quad (3.7)$$

Now define the function $\sigma: \Sigma \rightarrow [0, \infty)$ by $\sigma(x) = B_\mu |x|^D$, $x \in \Sigma$. It holds that $\sigma \in \mathcal{C}^\infty(\Sigma)$. Clearly $v(x) = (g \circ \sigma)(x)$, $x \in \Sigma$, where g is the function from (3.6), so $v \in \mathcal{C}^{m-1}(\Sigma)$ owing to Lemma 3.6 and the m th order weak partial derivatives exist (see [25, Theorem 10.35]) and are linear combinations of the functins

$$x \mapsto \int_{B_\mu |x|^D}^{\infty} f(\tau) \tau^{\frac{m}{D}-m} (\tau - B_\mu |x|^D)^{m-l_1-1} d\tau |x|^{l_1(D-2)-2l_2} \prod_{j=1}^{2(l_1+l_2)-k} x_{i_j} \quad (3.8)$$

and

$$x \mapsto f(B_\mu |x|^D) |x|^{-m} \prod_{j=1}^m x_{i_j} \quad (3.9)$$

for the parametres l_1, l_2 satisfying $l_1 \in \mathbb{N}, l_1 \leq m-1, l_2 \in \{0, \dots, m\}, 2(l_1 + l_2) \geq m$. Since f has a bounded support, we also have

$$|\nabla^m v(x)| \leq K_1 \left(f(B_\mu |x|^D) + \sum_{l=1}^{m-1} \int_{B_\mu |x|^D}^{\infty} f(\tau) \tau^{\frac{m}{D}-l-1} d\tau |x|^{lD-m} \right) \quad (3.10)$$

for μ -almost every $x \in \Sigma$ thanks to (3.8) and (3.9).

For every $l \in \{1, \dots, m-1\}$, we now define the operator

$$F_l: (L^1 + L^\infty)(0, \infty) \rightarrow (L^1 + L^\infty)(0, \infty)$$

by

$$F_l(\varphi)(t) = t^{l-\frac{m}{D}} \int_t^{\infty} \varphi(\tau) \tau^{\frac{m}{D}-l-1} d\tau, \quad t \in (0, \infty), \quad \varphi \in (L^1 + L^\infty)(0, \infty).$$

For an arbitrary $l \in \{1, \dots, m-1\}$ we have $\|F_l\|_{L^\infty \rightarrow L^\infty} \leq \frac{D}{Dl-m}$ and $\|F_l\|_{L^1 \rightarrow L^1} \leq \frac{D}{Dl-m+D}$. So, we obtain the fact that the operator F_l is, owing to [2, Chapter 3, Theorem 2.2], bounded on the space $X(0, \infty)$.

Now define the functions $h: (0, \infty) \rightarrow [0, \infty)$ and $\omega: \Sigma \rightarrow [0, \infty)$. The function h is defined by

$$h(t) = f(t) + \sum_{l=1}^{m-1} F_l(f)(t), \quad t \in (0, \infty).$$

The function ω is defined by

$$\omega(x) = (h \circ \sigma)(x), \quad x \in \Sigma.$$

Owing to (3.10) we obtain

$$|\nabla^m v|_\mu^*(t) \leq K_2 \omega_\mu^*(t), \quad t \in (0, \infty).$$

The functions h and ω are equimeasurable. So, we have

$$|\nabla^m v|_\mu^*(t) \leq K_2 h^*(t), \quad t \in (0, \infty),$$

thanks to (2.1). Now we obtain

$$\begin{aligned} \|\nabla^m v\|_{X(\Sigma, \mu)} &\leq K_2 \|h\|_{X(0, \infty)} \\ &\leq K_2 \left(\|f\|_{X(0, \infty)} + \sum_{l=1}^{m-1} \|F_l(f)\|_{X(0, \infty)} \right) \leq K_3 \|f\|_{X(0, \infty)}. \end{aligned} \quad (3.11)$$

From (3.11) it follows that $v \in V^m X(\Sigma, \mu)$. Since the function f has a bounded support, we obtain the fact that $v \in V_0^m X(\Sigma, \mu)$ owing to (3.7).

Now, we know that the functions v and g are equimeasurable since the mapping σ is a measure-preserving transformation of the spaces (Σ, μ) and $([0, \infty), \lambda)$ (see [2, Section 2.7]). We obtain the fact that

$$\begin{aligned} \|v\|_{Y(\Sigma, \mu)} &= \|g\|_{Y(0, \infty)} \geq \left\| \int_{2t}^{\infty} f(\tau) \tau^{\frac{m}{D}-m} (\tau-t)^{m-1} d\tau \right\|_{Y(0, \infty)} \\ &= 2^{1-m} \left\| \int_{2t}^{\infty} f(\tau) \tau^{\frac{m}{D}-m} \tau^{m-1} d\tau \right\|_{Y(0, \infty)} = 2^{1-m} \left\| \int_{2t}^{\infty} f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)}. \end{aligned} \quad (3.12)$$

Finally, we have

$$\begin{aligned} \left\| \int_t^{\infty} f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} &\leq 2 \left\| \int_{2t}^{\infty} f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \leq 2^m \|v\|_{Y(\Sigma, \mu)} \\ &\leq 2^m C_1 \|\nabla^m v\|_{X(\Sigma, \mu)} \leq 2^m C_1 K_3 \|f\|_{X(0, \infty)}. \end{aligned}$$

The first inequality holds by virtue of (2.14). The second inequality holds thanks to (3.12). The last inequality holds owing to (3.11). So, we have proved the inequality (3.5) for bounded functions with bounded support. Now let f be general. Define a sequence $\{f_k\}_{k=1}^{\infty}$ of functions from $\mathcal{M}^+(0, \infty)$ by $f_k(t) = \min\{f(t), k\} \chi_{(0, k)}(t)$, $t \in (0, \infty)$, $k \in \mathbb{N}$. Since (3.5) holds for every f_k , $k \in \mathbb{N}$, we obtain the fact that (3.5) also holds for f thanks to the Fatou axiom of Banach function norms.

□

Now we prove the remaining implication in Theorem 3.2.

Proposition 3.7. *Let X, Y be rearrangement-invariant spaces over (Σ, μ) . Assume that there exists a positive constant C_2 such that for all functions $f \in \mathcal{M}^+(0, \infty)$ it holds that*

$$\left\| \int_t^\infty f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \leq C_2 \|f\|_{X(0, \infty)}. \quad (3.13)$$

Then there exists a positive constant C_1 such that for every $u \in V_0^m X(\Sigma, \mu)$ it holds that

$$\|u\|_{Y(\Sigma, \mu)} \leq C_1 \|\nabla^m u\|_{X(\Sigma, \mu)}.$$

The constant C_1 depends only on the constant C_2 , on m and on D .

Proof. The mapping $\sigma_{m, X}: \mathcal{M}^+(\Sigma, \mu) \rightarrow [0, \infty]$ defined by

$$\sigma_{m, X}(v) = \left\| t^{\frac{m}{D}} v_\mu^{**}(t) \right\|_{X'(0, \infty)}, \quad v \in \mathcal{M}^+(\Sigma, \mu),$$

is a rearrangement-invariant Banach function norm if and only if

$$t^{\frac{m}{D}-1} \chi_{(1, \infty)}(t) \in X'(0, \infty) \quad (3.14)$$

(see [13, Theorem 5.4] and [18, Theorem 4.4]). Since (3.13) holds we have the fact that (3.14) is true thanks to Remark 3.4. Hence $\sigma_{m, X}$ is a rearrangement-invariant Banach function norm. We denote the respective space by $Z_m X(\Sigma, \mu)$.

We want to prove that for every function $u \in V_0^m X(\Sigma, \mu)$ it holds that

$$\|u\|_{Y(\Sigma, \mu)} \leq C_2 \|u\|_{Z_m' X(\Sigma, \mu)} \leq C_2 K_m \|\nabla^m u\|_{X(\Sigma, \mu)}, \quad (3.15)$$

where K_m is a positive constant. The first inequality follows from (3.4). We prove the second inequality by induction on m . Firstly, we assume that $m = 1$. Choose an arbitrary function $u \in V_0^1 X(\Sigma, \mu)$. We have

$$\begin{aligned} \|u\|_{Z_1' X(\Sigma, \mu)} &= \|u_\mu^*\|_{Z_1' X(0, \infty)} = \left\| - \int_t^\infty \frac{du_\mu^*}{d\tau}(\tau) d\tau \right\|_{Z_1' X(0, \infty)} \\ &= \left\| \int_t^\infty \left(\tau^{\frac{D-1}{D}} \frac{du_\mu^*}{d\tau}(\tau) \right) \tau^{\frac{1-D}{D}} d\tau \right\|_{Z_1' X(0, \infty)} \leq \left\| t^{\frac{D-1}{D}} \frac{du_\mu^*}{dt}(t) \right\|_{X(0, \infty)} \\ &\leq C_3 \|\nabla u\|_{X(\Sigma, \mu)}. \end{aligned}$$

The second equality is true owing to the fact that u_μ is locally absolutely continuous on $(0, \infty)$ (see Proposition 3.1). The first inequality holds by virtue of (3.4) and the second one is true thanks to the Pólya–Szegő inequality (Proposition 3.1). So, we proved (3.15) for $m = 1$.

Now assume that $1 < m < D$ is arbitrary and that (3.15) holds for $m - 1$. Choose an arbitrary function $u \in V_0^m X(\Sigma, \mu)$ and $i \in \{1, \dots, n\}$. Then the

weak partial derivative $\frac{\partial u}{\partial x_i}$ belongs to $V_0^{m-1}X(\Sigma, \mu)$. So, we can use the induction hypothesis to obtain

$$\left\| \frac{\partial u}{\partial x_i} \right\|_{Z'_{m-1}X(\Sigma, \mu)} \leq K_{m-1} \left\| \nabla^{m-1} \frac{\partial u}{\partial x_i} \right\|_{X(\Sigma, \mu)} \leq K_{m-1} \|\nabla^m u\|_{X(\Sigma, \mu)}.$$

It means that

$$\|\nabla u\|_{Z'_{m-1}X(\Sigma, \mu)} \leq nK_{m-1} \|\nabla^m u\|_{X(\Sigma, \mu)} < \infty. \quad (3.16)$$

It follows that u belongs to $V_0^1 Z'_{m-1}X(\Sigma, \mu)$. Now it can be straightforwardly proved that

$$t^{\frac{1}{D}-1} \chi_{(1, \infty)}(t) \in Z_{m-1}X(0, \infty),$$

(for details see [28, Theorem 2.2]), which means that the mapping $u \mapsto \left\| t^{\frac{1}{D}} u_{\mu_D}^{**}(t) \right\|_{Z_{m-1}X(0, \infty)} = \|u\|_{Z_1(Z_{m-1}X)(0, \infty)}$ is a rearrangement-invariant Banach function norm (cf. (3.14)). From the case $m = 1$ it now follows that

$$\|u\|_{(Z_1(Z_{m-1}X))'(\Sigma, \mu)} \leq C_3 \|\nabla u\|_{Z'_{m-1}X(\Sigma, \mu)}. \quad (3.17)$$

Owing to [12, Theorem 3.4] and [30, Proposition 5.1] (cf. [13, Theorem 9.5]), we obtain

$$\begin{aligned} \|v\|_{Z_1(Z_{m-1}X)(\Sigma, \mu)} &= \left\| t^{\frac{m-1}{D}} \left(\tau^{\frac{1}{D}} v_{\mu}^{**}(\tau) \right)^{**}(t) \right\|_{X'(0, \infty)} \\ &\leq C_4 \left\| t^{\frac{m}{D}} v_{\mu}^{**}(t) \right\|_{X'(0, \infty)} = C_4 \|v\|_{Z_m X(\Sigma, \mu)} \end{aligned}$$

for every $v \in \mathcal{M}^+(\Sigma, \mu)$. From the previous inequality we obtain

$$\|v\|_{Z'_m X(\Sigma, \mu)} \leq C_5 \|v\|_{(Z_1(Z_{m-1}X))'(\Sigma, \mu)} \quad (3.18)$$

for every $v \in \mathcal{M}^+(\Sigma, \mu)$. By virtue of (3.16), (3.17) and (3.18), we obtain the fact that (3.15) holds for m . \square

3.2. Optimality. In this subsection we will describe the smallest target space in the inequality (3.2) among all spaces $Y(\Sigma, \mu)$ for a given space $X(\Sigma, \mu)$ and the largest domain space in the inequality (3.2) among all spaces $X(\Sigma, \mu)$ for a given space $Y(\Sigma, \mu)$.

Firstly we describe the optimal target space in the following theorem.

Theorem 3.8. *Let X be a rearrangement-invariant space over (Σ, μ) . Assume that*

$$t^{\frac{m}{D}-1} \chi_{(1, \infty)}(t) \in X'(0, \infty). \quad (3.19)$$

Then there exists a positive constant C , which depends only on m and on D , such that

$$\|u\|_{Z'_m X(\Sigma, \mu)} \leq C \|\nabla^m u\|_{X(\Sigma, \mu)}, \quad u \in V_0^m X(\Sigma, \mu). \quad (3.20)$$

Moreover, the space $Z'_m X(\Sigma, \mu)$ is the optimal space in the previous inequality among all rearrangement-invariant spaces in the following way. If $Y(\Sigma, \mu)$ is a rearrangement-invariant space satisfying

$$\|u\|_{Y(\Sigma, \mu)} \leq \tilde{C} \|\nabla^m u\|_{X(\Sigma, \mu)}, \quad u \in V_0^m X(\Sigma, \mu), \quad (3.21)$$

with a positive constant \tilde{C} that does not depend on u , then

$$Z'_m X(\Sigma, \mu) \hookrightarrow Y(\Sigma, \mu). \quad (3.22)$$

On the other hand, if (3.19) is not true, then the inequality (3.21) does not hold for any rearrangement-invariant space $Y(\Sigma, \mu)$.

Proof. The inequality (3.20) follows from the proof of Proposition 3.7. Assume that (3.19) holds and that $Y(\Sigma, \mu)$ is a rearrangement-invariant space satisfying (3.21). Owing to Proposition 3.5 we obtain the fact that $Y(\Sigma, \mu)$ satisfies the inequality (3.3). It means that we can use the first inequality in (3.15) to obtain the embedding (3.22).

Now, assume that (3.19) is not true. Then there does not exist any rearrangement-invariant space $Y(\Sigma, \mu)$ such that (3.3) holds thanks to Remark 3.4. Then by Proposition 3.5 there is not any rearrangement-invariant space $Y(\Sigma, \mu)$ such that (3.21) is true. \square

Let Y be a rearrangement-invariant space over (Σ, μ) . Assume that

$$\inf_{t \in [1, \infty)} \frac{t^{1-\frac{m}{D}}}{\varphi_Y(t)} > 0, \quad (3.23)$$

where φ_Y is the fundamental function of the space $Y(\Sigma, \mu)$. Then the mapping $\nu_{m,Y}: \mathcal{M}^+(\Sigma, \mu) \rightarrow [0, \infty]$ defined by

$$\nu_{m,Y}(u) = \sup_{v \sim u, v \geq 0} \left\| \int_t^\infty v(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)},$$

where the supremum is taken over all functions $v \in \mathcal{M}^+(0, \infty)$ equimeasurable with u , is a rearrangement-invariant Banach function norm (for the proof see [18, Theorem 4.1]). The respective space we will denote by $U_m Y(\Sigma, \mu)$.

Now we can finally describe the largest domain space.

Theorem 3.9. *Let Y be a rearrangement-invariant space over (Σ, μ) . Assume that the condition (3.23) is satisfied. Then there exists a positive constant C , which depends only on m and on D , such that*

$$\|u\|_{Y(\Sigma, \mu)} \leq C \|\nabla^m u\|_{U_m Y(\Sigma, \mu)}, \quad u \in V_0^m U_m Y(\Sigma, \mu). \quad (3.24)$$

Moreover, the space $U_m Y(\Sigma, \mu)$ is the optimal space in the previous inequality among all rearrangement-invariant spaces in the following way. If $X(\Sigma, \mu)$ is a rearrangement-invariant space satisfying

$$\|u\|_{Y(\Sigma, \mu)} \leq \tilde{C} \|\nabla^m u\|_{X(\Sigma, \mu)}, \quad u \in V_0^m X(\Sigma, \mu), \quad (3.25)$$

with a positive constant \tilde{C} that does not depend on u , then

$$X(\Sigma, \mu) \hookrightarrow U_m Y(\Sigma, \mu). \quad (3.26)$$

On the other hand, if (3.23) is not true, then the inequality (3.25) does not hold for any rearrangement-invariant space $X(\Sigma, \mu)$.

Proof. Let $f \in \mathcal{M}^+(0, \infty)$. Then

$$\begin{aligned} \left\| \int_t^\infty f(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} &\leq \sup_{g \sim f, g \geq 0} \left\| \int_t^\infty g(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \\ &= \|f\|_{U_m Y(0, \infty)}, \end{aligned}$$

so, the inequality (3.24) follows from Proposition 3.7.

Now, assume that $X(\Sigma, \mu)$ is a rearrangement-invariant space satisfying (3.25) and let $f, g \in \mathcal{M}^+(0, \infty)$ be equimeasurable. Then, by virtue of Proposition 3.5, we obtain

$$\begin{aligned} \|f\|_{U_m Y(0, \infty)} &= \sup_{g \sim f, g \geq 0} \left\| \int_t^\infty g(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} \leq K \sup_{g \sim f, g \geq 0} \|g\|_{X(0, \infty)} \\ &= K \|f\|_{X(0, \infty)}. \end{aligned}$$

It means that (3.26) is true.

On the other hand, if the condition (3.23) is not satisfied, then it can be proved analogously as in the proof of [18, Theorem 4.1] that for each $u \in \mathcal{M}^+(\Sigma, \mu)$ such that $u_\mu^* = \chi_{(0,1)}$ it holds that

$$\nu_{m, Y}(u) = \sup_{v \sim u, v \geq 0} \left\| \int_t^\infty v(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)} = \infty.$$

By the nontriviality axiom of the Banach function norm, we know that u belongs to an arbitrary rearrangement-invariant space $X(\Sigma, \mu)$. So, we obtain the fact that (3.3) does not hold for any rearrangement-invariant space $X(\Sigma, \mu)$. It means that the inequality (3.25) does not hold for any rearrangement-invariant space $X(\Sigma, \mu)$ owing to the reduction principle (Theorem 3.2). □

At the end of the section we prove Proposition 3.1. The proof is based on the proofs of theorems [10, Lemma 4.1], [11, Lemma 3.3] and [46, Lemma 1.E].

Proof of Proposition 3.1. Firstly, we prove the proposition for nonnegative u . We start with the proof of the local absolute continuity of the function u_μ^* . Let $\{(a_m, b_m)\}_{m \in M}$ be a countable system of pairwise disjoint nonempty bounded intervals. For each $m \in M$ define the function $f_m: \mathbb{R} \rightarrow \mathbb{R}$ in the

following way:

$$f_m(y) = \begin{cases} 0 & \text{if } y \leq u_\mu^*(b_m), \\ y - u_\mu^*(b_m) & \text{if } u_\mu^*(b_m) < y < u_\mu^*(a_m), \\ u_\mu^*(a_m) - u_\mu^*(b_m) & \text{if } u_\mu^*(a_m) \leq y. \end{cases}$$

For each $m \in M$ we now set $v_m = f_m \circ u$. It also holds that

$$\mu(\{x \in \Sigma; u(x) > u_\mu^*(b_m)\}) < \infty \quad (3.27)$$

since $u \in V_0^1 X(\Sigma, \mu)$. So, the function v_m is bounded and can be nonzero in a set of finite μ -measure only. We obtain the fact that $v_m \in L^1(\Sigma, \mu)$. Now we use the chain rule for Sobolev functions to obtain the fact that v_m is weakly differentiable in Σ and $\nabla v_m = \nabla u \chi_{\{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}}$ μ -almost everywhere in Σ . From this equality we get $|\nabla v_m| = 0$ μ -almost everywhere in the set $\{x \in \Sigma; u(x) \leq u_\mu^*(b_m)\}$. We know that the function $\nabla u \in X(\Sigma, \mu)$. So, by virtue of (2.2) it follows that $\nabla u \in L^1(E, \mu)$ for every μ -measurable set $E \subseteq \Sigma, \mu(E) < \infty$. We can now again use (3.27) to obtain $\nabla v_m \in L^1(\Sigma, \mu)$.

Now we can use the coarea formula (see [27], [45]) for the functions $v_m, m \in M$, and the isoperimetric inequality (see [7, Theorem 1.3]). We obtain

$$\begin{aligned} & \int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x) \\ & \geq C_{iso} \sum_{m \in M} \int_0^{u_\mu^*(a_m) - u_\mu^*(b_m)} \mu(\{x \in \Sigma; v_m(x) > t\})^{\frac{D-1}{D}} dt \\ & = C_{iso} \sum_{m \in M} \int_{u_\mu^*(b_m)}^{u_\mu^*(a_m)} \mu(\{x \in \Sigma; u(x) > t\})^{\frac{D-1}{D}} dt, \end{aligned} \quad (3.28)$$

where C_{iso} is the isoperimetric constant. Now we derive an upper estimate of

$$\int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x).$$

We obtain the fact that

$$\begin{aligned} & \int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x) \\ & \leq \int_0^{\sum_{m \in M} \mu(\{u_\mu^*(b_m) < u < u_\mu^*(a_m)\})} |\nabla u|_\mu^*(t) dt \leq \int_0^{\sum_{m \in M} (b_m - a_m)} |\nabla u|_\mu^*(t) dt. \end{aligned} \quad (3.29)$$

The first inequality holds by virtue of the Hardy—Littlewood inequality (2.9). We can verify the last inequality in the following way. We have

$$\begin{aligned} \mu(\{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}) & = \mu(\{u_\mu^*(b_m) < u\}) - \mu(\{u_\mu^*(a_m) \leq u\}) \\ & \leq b_m - a_m, \end{aligned}$$

where we used (2.6) and (2.7).

In the following part of the proof, we will assume that all the intervals $(a_m, b_m), m \in M$, are contained in an interval $[a, b] \subseteq (0, \infty)$. We prove that the function u_μ^* is absolutely continuous on the interval $[a, b]$. We can assume that $u_\mu^*(a) > 0$. We set

$$K = \mu(\{x \in \Sigma; u(x) \geq u_\mu^*(a)\}).$$

Then we have $K < \infty$ since $u \in V_0^1 X(\Sigma, \mu)$. Since $u_\mu^*(a) > 0$, we can use (2.7) to obtain $K > 0$. Owing to (3.28) we obtain the fact that

$$\begin{aligned} & \int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x) \\ & \geq C_{iso} \sum_{m \in M} \int_{u_\mu^*(b_m)}^{u_\mu^*(a_m)} \mu(\{x \in \Sigma; u(x) \geq u_\mu^*(a)\})^{\frac{D-1}{D}} dt \quad (3.30) \\ & = C_{iso} K^{\frac{D-1}{D}} \sum_{m \in M} (u_\mu^*(a_m) - u_\mu^*(b_m)). \end{aligned}$$

It follows that

$$\begin{aligned} & \sum_{m \in M} (u_\mu^*(a_m) - u_\mu^*(b_m)) \\ & \leq C_{iso}^{-1} K^{\frac{1-D}{D}} \int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x) \quad (3.31) \\ & \leq C_{iso}^{-1} K^{\frac{1-D}{D}} \int_0^{\sum_{m \in M} (b_m - a_m)} |\nabla u|_\mu^*(t) dt. \end{aligned}$$

The first inequality holds due to (3.30). The second inequality holds by virtue of (3.29).

Next we want to prove that

$$\int_0^t |\nabla u|_\mu^*(\tau) d\tau < \infty \quad (3.32)$$

for every $t \in (0, \infty)$. Since $|\nabla u| \in (L^1 + L^\infty)(\Sigma, \mu)$, it follows that $|\nabla u|_\mu^* \in L^1(0, t)$. So, (3.32) is true.

Since we know that $|\nabla u|_\mu^*$ is integrable over an arbitrary bounded interval $(0, t)$, we can use (3.31) to obtain the fact that the function u_μ^* is absolutely continuous on the interval $[a, b]$. It follows that it is locally absolutely continuous on the interval $(0, \infty)$, which is the desired result.

It remains to prove the inequality (3.1). From now we do not anymore assume that the intervals (a_m, b_m) are contained in $[a, b]$. Define the function $\phi: (0, \infty) \rightarrow [0, \infty)$ by $\phi(t) = -C_{iso} t^{\frac{D-1}{D}} \frac{du_\mu^*}{dt}(t)$, $t \in (0, \infty)$. We show that

$$\int_0^t \phi^*(\tau) d\tau \leq \int_0^t |\nabla u|_\mu^*(\tau) d\tau, \quad t \in (0, \infty). \quad (3.33)$$

Choose $t \in (0, \infty)$ arbitrarily. By virtue of (2.11), we know that it is enough to prove that for every measurable set $E \subseteq (0, \infty)$ such that $\lambda(E) = t$, it

holds that

$$\int_E \phi(\tau) d\tau \leq \int_0^t |\nabla u|_\mu^*(\tau) d\tau. \quad (3.34)$$

Now choose an arbitrary $m \in M$. We have

$$\int_{a_m}^{b_m} \phi(\tau) d\tau = - \int_{a_m}^{b_m} C_{iso} \tau^{\frac{D-1}{D}} \frac{du_\mu^*}{d\tau}(\tau) d\tau. \quad (3.35)$$

Since the function u_μ^* is absolutely continuous and nonincreasing on the interval $[a_m, b_m]$, and the function $\tau \mapsto \mu(\{x \in \Sigma; u(x) > \tau\})^{\frac{D-1}{D}}$, $\tau \in (0, \infty)$, is nonnegative on the interval (a_m, b_m) , we obtain the fact that (see [41, page 156])

$$\begin{aligned} & \int_{u_\mu^*(b_m)}^{u_\mu^*(a_m)} C_{iso} \mu(\{x \in \Sigma; u(x) > s\})^{\frac{D-1}{D}} ds \\ &= - \int_{a_m}^{b_m} C_{iso} \mu(\{x \in \Sigma; u(x) > u_\mu^*(\tau)\})^{\frac{D-1}{D}} \frac{du_\mu^*}{d\tau}(\tau) d\tau. \end{aligned} \quad (3.36)$$

Now we observe that

$$\begin{aligned} & \int_{a_m}^{b_m} C_{iso} \mu(\{x \in \Sigma; u(x) > u_\mu^*(\tau)\})^{\frac{D-1}{D}} \frac{du_\mu^*}{d\tau}(\tau) d\tau \\ &= \int_{a_m}^{b_m} C_{iso} \tau^{\frac{D-1}{D}} \frac{du_\mu^*}{d\tau}(\tau) d\tau. \end{aligned} \quad (3.37)$$

Owing to (2.6) we know that for every $\tau \in (a_m, b_m)$ it holds that

$$\mu(\{x \in \Sigma; u(x) > u_\mu^*(\tau)\}) \leq \tau.$$

Choose an arbitrary $\tau \in (a_m, b_m)$ such that the function u_μ^* is differentiable at τ . Assume that

$$\mu(\{x \in \Sigma; u(x) > u_\mu^*(\tau)\}) < \tau.$$

Then there exists $\delta > 0$, such that for every $s \in (\tau - \delta, \tau)$ it holds that

$$\mu(\{x \in \Sigma; u(x) > u_\mu^*(\tau)\}) < s.$$

From this inequality we obtain the fact that for each $s \in (\tau - \delta, \tau)$

$$\begin{aligned} u_\mu^*(s) &= \inf \{ \alpha > 0; \mu(\{x \in \Sigma; u(x) > \alpha\}) \leq s \} \\ &\leq u_\mu^*(\tau). \end{aligned}$$

Because the function u_μ^* is nonincreasing it holds that $u_\mu^*(s) = u_\mu^*(\tau)$ for all $s \in (\tau - \delta, \tau)$. Since the function u_μ^* is differentiable at τ , we have $\frac{du_\mu^*}{d\tau}(\tau) = 0$. So, we have just proved (3.37). By (3.35), (3.36) and (3.37) we have

$$\int_{a_m}^{b_m} \phi(\tau) d\tau = \int_{u_\mu^*(b_m)}^{u_\mu^*(a_m)} C_{iso} \mu(\{x \in \Sigma; u(x) > \tau\})^{\frac{D-1}{D}} d\tau. \quad (3.38)$$

Now we obtain

$$\begin{aligned}
\int_{\bigcup_{m \in M} (a_m, b_m)} \phi(\tau) d\tau &= \sum_{m \in M} \int_{u_\mu^*(b_m)}^{u_\mu^*(a_m)} C_{iso\mu}(\{x \in \Sigma; u(x) > \tau\})^{\frac{D-1}{D}} d\tau \\
&\leq \int_{\bigcup_{m \in M} \{u_\mu^*(b_m) < u < u_\mu^*(a_m)\}} |\nabla u(x)| d\mu(x) \\
&\leq \int_0^{\sum_{m \in M} (b_m - a_m)} |\nabla u|_\mu^*(\tau) d\tau.
\end{aligned} \tag{3.39}$$

The equality holds thanks to (3.38) and to the fact that all the intervals (a_m, b_m) , $m \in M$, are pairwise disjoint. The first inequality holds thanks to (3.28). The second inequality is true by virtue of (3.29). Now choose an arbitrary measurable set $E \subseteq (0, \infty)$, $\lambda(E) = t$. Then for every $\varepsilon > 0$, there exists a countable system $\{(a_m, b_m)\}_{m \in M}$ of pairwise disjoint nonempty bounded intervals such that

$$E \subseteq \bigcup_{m \in M} (a_m, b_m) \quad \text{and} \quad \lambda\left(\bigcup_{m \in M} (a_m, b_m) \setminus E\right) < \varepsilon.$$

So, choose an arbitrary $\varepsilon > 0$, and let $\{(a_m, b_m)\}_{m \in M}$ be such a system of intervals. From (3.39) we obtain

$$\int_E \phi(\tau) d\tau \leq \int_{\bigcup_{m \in M} (a_m, b_m)} \phi(\tau) d\tau \leq \int_0^{t+\varepsilon} |\nabla u|_\mu^*(\tau) d\tau.$$

We already know from (3.32) that for all $s \in (0, \infty)$ the function $|\nabla u|_\mu^*$ is integrable over $(0, s)$. So, we obtain

$$\int_E \phi(\tau) d\tau \leq \int_0^t |\nabla u|_\mu^*(\tau) d\tau.$$

This means that (3.34) is true. So, we have just proved (3.33). The inequality (3.1) now follows from the Hardy–Littlewood–Pólya principle (2.10).

We have proved the proposition for nonnegative u . Now let $u \in V_0^1 X(\Sigma, \mu)$ be general. It is true that $|u| \in V_0^1 X(\Sigma, \mu)$. So, the proposition holds for $|u|$. Since $u_\mu^* = |u|_\mu^*$, the proposition holds also for the function u . \square

4. EXAMPLES

In this section we will show some examples. We will describe the optimal space $Z'_m(\Sigma, \mu)$ when $X(\Sigma, \mu)$ is a Lorentz–Karamata space $L^{p,q,b}(\Sigma, \mu)$ with $p \in [1, \frac{D}{m}]$.

Now we introduce the theory of Lorentz–Karamata spaces. For proofs and more information see [16, 17, 34, 39]. The class of Lorentz–Karamata spaces contain a lot of classical function spaces, such as the Lebesgue spaces,

Lorentz spaces, Lorentz-Zygmund spaces or Orlicz spaces of the type L_{exp} or $L^p(\log L)^\alpha$.

We start with the concept of slowly varying functions. Let $f, g: (0, \infty) \rightarrow (0, \infty)$ be functions. We say that the function f is *equivalent* to the function g if there exist constants $C_1 > 0$ and $C_2 > 0$ such that $C_2 g(t) \leq f(t) \leq C_1 g(t)$ for every $t \in (0, \infty)$.

We say that a locally absolutely continuous function $b: (0, \infty) \rightarrow (0, \infty)$ is *slowly varying* if for every $\varepsilon > 0$ there exists a nondecreasing function φ_ε and a nonincreasing function $\varphi_{-\varepsilon}$ such that $t^\varepsilon b(t)$ is equivalent to $\varphi_\varepsilon(t)$ on $(0, \infty)$ and that $t^{-\varepsilon} b(t)$ is equivalent to $\varphi_{-\varepsilon}$ on $(0, \infty)$. A positive locally absolutely continuous function on $(0, \infty)$ that is equivalent to a positive constant function is a trivial example of a slowly varying function. Functions of logarithmic type constitute less trivial and very important examples of slowly varying functions. For $k \in \mathbb{N}$, the function $\ell_k: (0, \infty) \rightarrow (0, \infty)$ defined as

$$\ell_k(t) = \begin{cases} 1 + |\log t| & \text{if } k = 1, \\ 1 + \log \ell_{k-1}(t) & \text{if } k > 1, \end{cases}$$

$t \in (0, \infty)$, is slowly varying. More generally, the function $\ell_k^\mathbb{A}: (0, \infty) \rightarrow (0, \infty)$ defined as

$$\ell_k^\mathbb{A}(t) = \begin{cases} \ell_k^{\alpha_0}(t) & \text{if } t \in (0, 1), \\ \ell_k^{\alpha_\infty}(t) & \text{if } t \in [1, \infty), \end{cases}$$

where $\mathbb{A} = (\alpha_0, \alpha_\infty) \in \mathbb{R}^2$, is slowly varying.

Note that by b^{-1} we will denote the function $\frac{1}{b}$.

Now we finally introduce Lorentz–Karamata spaces. Let b be a slowly varying function. Let $p, q \in [1, \infty]$. We define the Lorentz–Karamata functionals by

$$\|f\|_{p,q,b} = \left\| t^{\frac{1}{p} - \frac{1}{q}} b(t) f_\mu^*(t) \right\|_{L^q(0,\infty)}$$

and by

$$\|f\|_{(p,q,b)} = \left\| t^{\frac{1}{p} - \frac{1}{q}} b(t) f_\mu^{**}(t) \right\|_{L^q(0,\infty)}$$

for every $f \in \mathcal{M}(\Sigma, \mu)$. The *Lorentz–Karamata spaces* are defined as

$$L^{p,q,b}(\Sigma, \mu) = \left\{ f \in \mathcal{M}(\Sigma, \mu); \|f\|_{p,q,b} < \infty \right\}$$

and as

$$L^{(p,q,b)}(\Sigma, \mu) = \left\{ f \in \mathcal{M}(\Sigma, \mu); \|f\|_{(p,q,b)} < \infty \right\}.$$

If we take $p = q$ and $b \equiv 1$, we obtain the Lebesgue spaces. More generally, we obtain the Lorentz spaces $L^{p,q}(\Sigma, \mu)$ and $L^{(p,q)}(\Sigma, \mu)$ by taking $b \equiv 1$. Furthermore, Lorentz–Karamata spaces also include the Lorentz–Zygmund spaces, which were thoroughly studied in [36]. We obtain them by taking slowly varying functions of logarithmic type as above.

Even though we refer to Lorentz–Karamata spaces as spaces, they are not always rearrangement-invariant spaces.

The space $L^{(p,q,b)}(R, \mu)$ is a rearrangement-invariant Banach function space if and only if $q \in [1, \infty]$ and one of the following conditions holds:

- (1) $p \in (1, \infty)$,
- (2) $p = 1$ and $\left\| t^{-\frac{1}{q}} b(t) \chi_{(1, \infty)}(t) \right\|_{L^q(0, \infty)} < \infty$,
- (3) $p = \infty$ and $\left\| t^{-\frac{1}{q}} b(t) \chi_{(0, 1)}(t) \right\|_{L^q(0, \infty)} < \infty$.

The Lorentz–Karamata functional $\|\cdot\|_{p,q,b}$ is equivalent to a rearrangement-invariant Banach function norm if and only if $q \in [1, \infty]$ and one of the following conditions is satisfied:

- (1) $p \in (1, \infty)$,
- (2) $p = q = 1$ and b is equivalent to a nonincreasing function on $(0, \infty)$,
- (3) $p = \infty$ and $\left\| t^{-\frac{1}{q}} b(t) \chi_{(0, 1)}(t) \right\|_{L^q(0, \infty)} < \infty$.

In what follows we will also use the fact that if $p > 1$, then the spaces $L^{p,q,b}(\Sigma, \mu)$ and $L^{(p,q,b)}(\Sigma, \mu)$ are equivalent.

Now we describe the optimal target space for the Lorentz–Karamata space $L^{p,q,b}(\Sigma, \mu)$.

Theorem 4.1. *Let $p \in [1, \frac{D}{m}]$, let $q \in [1, \infty]$ and let b be a slowly varying function. Assume that $X(\Sigma, \mu) = L^{p,q,b}(\Sigma, \mu)$. Then we have the following description of the optimal target space $Z'_m X(\Sigma, \mu)$.*

- (1) $Z'_m X(\Sigma, \mu) = L^{\frac{Dp}{D-mp}, q, b}(\Sigma, \mu)$ if $p \in (1, \frac{D}{m})$ or $p = q = 1$ and b is equivalent to a nonincreasing function on $(0, \infty)$.
- (2) $Z'_m X(\Sigma, \mu) = L^{\infty, q, a}(\Sigma, \mu)$ if $p = \frac{D}{m}$, $q \in (1, \infty]$ and it holds that

$$\left\| t^{-\frac{1}{q'}} b^{-1}(t) \right\|_{L^{q'}(1, \infty)} < \infty. \quad (4.1)$$

Here, the function a is defined by

$$a(t) = \frac{b^{1-q'}(t)}{\int_t^\infty \tau^{-1} b^{-q'}(\tau) d\tau}, \quad t \in (0, \infty).$$

- (3) $Z'_m X(\Sigma, \mu) = \Lambda^1(d')(\Sigma, \mu)$ if $p = \frac{D}{m}$, $q = 1$, (4.1) is satisfied and $\lim_{t \rightarrow 0_+} d(t) = 0$. Here, the function d is defined by

$$d(t) = \inf_{\tau \in [t, \infty)} b(\tau), \quad t \in (0, \infty). \quad (4.2)$$

- (4) $Z'_m X(\Sigma, \mu) = \Lambda^1(d')(\Sigma, \mu) \cap L^\infty(\Sigma, \mu)$ if $p = \frac{D}{m}$, $q = 1$, (4.1) is satisfied and $\lim_{t \rightarrow 0_+} d(t) > 0$. The function d is again defined by (4.2).

The function space $\Lambda^1(d')(\Sigma, \mu)$ is an example of a classical Lorentz space (e.g., see [40, Chapter 10]). It is defined as the collection of all functions $f \in \mathcal{M}(\Sigma, \mu)$ such that $\|d'(t) f_\mu^*(t)\|_{L^1(0, \infty)} < \infty$.

Proof. First of all we prove that the space $Z'_m X(\Sigma, \mu)$ is the optimal space. Owing to [39, Section 3.7] we obtain the fact that the space $(L^{p,q,b}(\Sigma, \mu))'$ is equivalent to the space $L^{p',q',b^{-1}}(\Sigma, \mu)$. So, in the view of Theorem 3.8, it is enough to prove that

$$\left\| t^{\frac{m}{D}-1} \chi_{(1,\infty)}(t) \right\|_{L^{p',q',b^{-1}}(0,\infty)} = \left\| t^{\frac{1}{p'}-\frac{1}{q'}} (t+1)^{\frac{m}{D}-1} b^{-1}(t) \right\|_{L^{q'}(0,\infty)} < \infty. \quad (4.3)$$

If $p \in [1, \frac{D}{m})$, then (4.3) follows from the basic properties of slowly varying functions (see [39, Lemma 2.16]). If $p = \frac{D}{m}$ then (4.3) follows from (4.1) and from Theorem 3.8.

It remains to describe the space $Z'_m X(\Sigma, \mu)$. Firstly, assume that $p \in (1, \frac{D}{m})$. Owing to [39, Section 3.5] the maximal nonincreasing operator P_μ is bounded on the space $L^{p,q,b}(0, \infty)$. Now we show that the norm of the space $Z_m X(\Sigma, \mu)$ is equivalent to the functional

$$u \mapsto \left\| t^{\frac{1}{q}-\frac{1}{p}} b^{-1}(t) \int_t^\infty u_\mu^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{q'}(0,\infty)}, \quad u \in \mathcal{M}^+(\Sigma, \mu). \quad (4.4)$$

By virtue of [39, Section 3.7] we can rewrite the functional in the form

$$\begin{aligned} & \left\| \int_t^\infty u_\mu^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{(L^{p,q,b})'(0,\infty)} \\ &= \sup_{\|g\|_{L^{p,q,b}(0,\infty)} \leq 1} \int_0^\infty g^*(t) \int_t^\infty u_\mu^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau dt \\ &= \sup_{\|g\|_{L^{p,q,b}(0,\infty)} \leq 1} \int_0^\infty u_\mu^{**}(\tau) \tau^{\frac{m}{D}} g^{**}(\tau) d\tau. \end{aligned}$$

By virtue of (2.13) we obtain the first inequality

$$\begin{aligned} \|u\|_{Z_m X(\Sigma, \mu)} &= \left\| t^{\frac{m}{D}} u_\mu^{**}(t) \right\|_{(L^{p,q,b})'(0,\infty)} \\ &\leq 4 \sup_{\|g\|_{L^{p,q,b}(0,\infty)} \leq 1} \int_0^\infty g^{**}(\tau) \tau^{\frac{m}{D}} u_\mu^{**}(\tau) d\tau, \end{aligned}$$

while the second inequality follows directly from the boundedness of the operator P_μ . The norm of the associate space $(L^{\frac{Dp}{D-mp},q,b}(\Sigma, \mu))'$ is equivalent to the functional

$$u \mapsto \left\| t^{\frac{1}{q}-\frac{1}{p}+\frac{m}{D}} b^{-1}(t) u_\mu^{**}(t) \right\|_{L^{q'}(0,\infty)}, \quad u \in \mathcal{M}^+(\Sigma, \mu), \quad (4.5)$$

owing to [39, Section 3.7]. It remains to prove that the functionals (4.4) and (4.5) are equivalent. Since

$$t^{\frac{m}{D}} = \frac{m}{D \left(1 - 2^{-\frac{m}{D}}\right)} \int_{\frac{t}{2}}^t \tau^{\frac{m}{D}-1} d\tau,$$

we obtain

$$\begin{aligned} t^{\frac{m}{D}} u_{\mu}^{**}(t) &\leq \frac{m}{D \left(1 - 2^{-\frac{m}{D}}\right)} \int_{\frac{t}{2}}^t u_{\mu}^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \\ &\leq \frac{m}{D \left(1 - 2^{-\frac{m}{D}}\right)} \int_{\frac{t}{2}}^{\infty} u_{\mu}^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \end{aligned} \quad (4.6)$$

for every $u \in \mathcal{M}^+(\Sigma, \mu)$ and for every $t \in (0, \infty)$. By virtue of (4.6) and [39, Lemma 2.16], we obtain existence of a positive constant C_1 such that

$$t^{\frac{1}{q}-\frac{1}{p}+\frac{m}{D}} b^{-1}(t) u_{\mu}^{**}(t) \leq \frac{2^{\frac{1}{q}-\frac{1}{p}} m C_1}{D \left(1 - 2^{-\frac{m}{D}}\right)} \left(\frac{t}{2}\right)^{\frac{1}{q}-\frac{1}{p}} b^{-1}\left(\frac{t}{2}\right) \int_{\frac{t}{2}}^{\infty} u_{\mu}^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \quad (4.7)$$

for every $u \in \mathcal{M}^+(\Sigma, \mu)$ and for every $t \in (0, \infty)$. Owing to (2.14) and (4.7) we obtain existence of a positive constant C_2 such that

$$\left\| t^{\frac{1}{q}-\frac{1}{p}+\frac{m}{D}} b^{-1}(t) u_{\mu}^{**} \right\|_{L^{q'}(0, \infty)} \leq C_2 \left\| t^{\frac{1}{q}-\frac{1}{p}} b^{-1}(t) \int_t^{\infty} u_{\mu}^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{q'}(0, \infty)}$$

for every $u \in \mathcal{M}^+(\Sigma, \mu)$. Now we prove the opposite inequality, i.e., we prove that there exists a constant $K_1 > 0$ such that

$$\begin{aligned} &\left\| t^{\frac{1}{q}-\frac{1}{p}} b^{-1}(t) \int_t^{\infty} u_{\mu}^{**}(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{q'}(0, \infty)} \\ &\leq K_1 \left\| t^{\frac{1}{q}-\frac{1}{p}+\frac{m}{D}} b^{-1}(t) u_{\mu}^{**}(t) \right\|_{L^{q'}(0, \infty)}, \end{aligned} \quad (4.8)$$

for every $u \in \mathcal{M}^+(\Sigma, \mu)$. By virtue of the weighted Hardy inequality ([32, Theorem 2]), the inequality (4.8) is valid if and only if

$$\sup_{t \in (0, \infty)} \left\| \tau^{\frac{1}{q}-\frac{1}{p}} b^{-1}(\tau) \right\|_{L^{q'}(0, t)} \left\| \tau^{\frac{1}{p}-\frac{1}{q}-1} b(\tau) \right\|_{L^q(t, \infty)} < \infty,$$

which is not hard to verify (see [39, Lemma 2.16]).

It remains to derive the description of the space $Z'_m X(\Sigma, \mu)$ in case of $p = q = 1$ and b is equivalent to a nonincreasing function on $(0, \infty)$ or $p = \frac{D}{m}$ and $q \in [1, \infty]$. Thanks to [39, Section 3.5 and Section 3.7] it is sufficient to prove that $Z_m X(\Sigma, \mu)$ with the norm satisfying

$$\|u\|_{Z_m X(\Sigma, \mu)} = \left\| t^{\frac{m}{D}} u_{\mu}^{**}(t) \right\|_{L^{p', q', b^{-1}}(0, \infty)}$$

for each $u \in \mathcal{M}^+(\Sigma, \mu)$ is equivalent to $L^{(r, q', b^{-1})}(\Sigma, \mu)$, where $r = 1$ if $p = \frac{D}{m}$ and $r = \frac{D}{m}$ if $p = 1$. Note that in case of $p = \frac{D}{m}, q = 1$, we also exploit the fact that d^{-1} is locally absolutely continuous on $(0, \infty)$. Now, for every

$u \in \mathcal{M}^+(\Sigma, \mu)$ we have

$$\begin{aligned} \|u\|_{Z_m X(\Sigma, \mu)} &= \left\| t^{\frac{m}{D}} u_\mu^{**}(t) \right\|_{L^{p', q', b^{-1}}(0, \infty)} \leq \left\| \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} u_\mu^{**}(\tau) \right\|_{L^{p', q', b^{-1}}(0, \infty)} \\ &= \left\| t^{\frac{1}{p'} - \frac{1}{q'}} b^{-1}(t) \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} u_\mu^{**}(\tau) \right\|_{L^{q'}(0, \infty)} \\ &\leq K_2 \left\| t^{\frac{1}{r} - \frac{1}{q'}} b^{-1}(t) u_\mu^{**}(t) \right\|_{L^{q'}(0, \infty)} = \|u\|_{L^{(r, q', b^{-1})}(\Sigma, \mu)}, \end{aligned}$$

where K_2 is a positive constant that does not depend on u . If $p = q = 1$, then the last inequality follows from the fact that b^{-1} is equivalent to a nonincreasing function on $(0, \infty)$. If $p = \frac{D}{m}, q = 1$, then the last inequality follows from the fact that $t^{1 - \frac{m}{D}} b^{-1}(t)$ is equivalent to a nonincreasing function on $(0, \infty)$. If $q > 1$, then we use [19, Theorem 3.2] to prove the last inequality. Owing to this theorem the last inequality is valid if and only if for every $t \in (0, \infty)$ it is true that

$$t^{\frac{m}{D}} \left\| \tau^{\frac{D-m}{D} - \frac{1}{q'}} b^{-1}(\tau) \right\|_{L^{q'}(0, t)} \leq K \left\| \tau^{1 - \frac{1}{q'}} b^{-1}(\tau) \right\|_{L^{q'}(0, t)},$$

where $K > 0$ is a constant that does not depend on t , which is satisfied thanks to [39, Lemma 2.16]. So, we have proved the first inequality. For every $u \in \mathcal{M}^+(\Sigma, \mu)$ we have now

$$\begin{aligned} \|u\|_{L^{(r, q', b^{-1})}(\Sigma, \mu)} &= \left\| t^{\frac{1}{r} - \frac{1}{q'}} b^{-1}(t) u_\mu^{**}(t) \right\|_{L^{q'}(0, \infty)} \\ &\leq \left\| t^{\frac{1}{r} - \frac{m}{D} - \frac{1}{q'}} b^{-1}(t) \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} u_\mu^{**}(\tau) \right\|_{L^{q'}(0, \infty)} \\ &= \left\| \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} u_\mu^{**}(\tau) \right\|_{L^{p', q', b^{-1}}(0, \infty)} \leq C \left\| t^{\frac{m}{D}} u_\mu^{**}(t) \right\|_{L^{p', q', b^{-1}}(0, \infty)} \\ &= C \|u\|_{Z_m X(\Sigma, \mu)}, \end{aligned}$$

where C is a positive constant that does not depend on u . To prove the second inequality, we exploit [18, Lemma 4.10] since the function $t \mapsto t^{\frac{m}{D}}, t \in [0, \infty)$, is quasiconcave. \square

Remark 4.2. If the function b is in addition nonincreasing on $(0, 1]$ and constant on $[1, \infty)$, then we obtain the fact that d is constant on $(0, \infty)$, so $d' \equiv 0$ on $(0, \infty)$. It means that the optimal space $Z'_m X(\Sigma, \mu)$ is equivalent to $L^\infty(\Sigma, \mu)$. For instance, if $d \equiv 1$ on $(0, \infty)$, then we obtain the Lorentz space $L^{\frac{D}{m}, 1}(\Sigma, \mu)$ (cf. [43]).

Finally, we describe the optimal domain space in the view of Theorem 3.9 for a given Lorentz–Karamata space.

Theorem 4.3. *Let $p \in \left[\frac{D}{D-m}, \infty\right]$, let $q \in [1, \infty]$ and let b be a slowly varying function. Assume that $Y(\Sigma, \mu) = L^{p,q,b}(\Sigma, \mu)$. Then we have the following description of the optimal domain space $U_m Y(\Sigma, \mu)$.*

- (1) $U_m Y(\Sigma, \mu) = L^{\frac{Dp}{D+mp}, q, b}(\Sigma, \mu)$ if $p \in \left(\frac{D}{D-m}, \infty\right)$.
- (2) $U_m Y(\Sigma, \mu) = L^{1,1,b}(\Sigma, \mu)$ if $p = \frac{D}{D-m}$, $q = 1$ and b is equivalent to a nonincreasing function on $(0, \infty)$.
- (3) $U_m Y(\Sigma, \mu)$ is given by

$$\|f\|_{U_m Y(\Sigma, \mu)} = \left\| t^{-\frac{1}{q}} b(t) \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^q(0, \infty)}$$

if $p = \infty$ and

$$\left\| t^{-\frac{1}{q}} b(t) \right\|_{L^q(0,1)} < \infty.$$

Proof. For every $f \in \mathcal{M}(0, \infty)$, we define the operator $T_{\frac{m}{D}}(f)$ by

$$T_{\frac{m}{D}}(f)(t) = t^{-\frac{m}{D}} \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} f_\mu^*(\tau), \quad t \in (0, \infty).$$

Owing to [18, Theorem 4.7 and Remark 4.8], we obtain the fact that the operator $T_{\frac{m}{D}}: Y'(0, \infty) \rightarrow Y'(0, \infty)$ is bounded if and only if the norm of the space $W_m Y(\Sigma, \mu)$ is equivalent to

$$\left\| \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{Y(0, \infty)}. \quad (4.9)$$

So, we prove the fact that the operator $T_{\frac{m}{D}}$ is bounded on the space $\left(L^{p,q,b}(0, \infty)\right)'$. Firstly, assume that $p < \infty$ or $q < \infty$. We use [39, Section 3.7] to describe the space $\left(L^{p,q,b}(0, \infty)\right)'$. If $p \in \left[\frac{D}{D-m}, \infty\right)$, then we obtain the fact that $\left(L^{p,q,b}(\Sigma, \mu)\right)'$ is equivalent to $L^{(p', q', b^{-1})}(\Sigma, \mu)$. If $p = \infty$ and $q \in [1, \infty)$, then $\left(L^{\infty, q, b}(\Sigma, \mu)\right)'$ is equivalent to $L^{(1, q', a)}(\Sigma, \mu)$, where a is the slowly varying function given by

$$a(t) = \left(\int_0^t \tau^{-1} b^q(\tau) d\tau \right)^{-1} b^{q-1}(t), \quad t \in (0, \infty).$$

In what follows we will write c instead of b^{-1} and a since the estimates are the same in both cases. Then, for every $f \in \mathcal{M}(0, \infty)$, we obtain

$$\begin{aligned}
& \left\| T_{\frac{m}{D}}(f) \right\|_{L^{(p', q', c)}(0, \infty)} = \left\| t^{\frac{1}{p'} - \frac{1}{q'}} c(t) \left(T_{\frac{m}{D}}(f) \right)_{\mu}^{**}(t) \right\|_{L^{q'}(0, \infty)} \\
& \leq C_2 \left\| t^{\frac{1}{p'} - \frac{1}{q'}} c(t) T_{\frac{m}{D}}(f_{\mu}^{**})(t) \right\|_{L^{q'}(0, \infty)} \\
& = C_2 \left\| t^{\frac{1}{p'} - \frac{1}{q'}} c(t) t^{-\frac{m}{D}} \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} f_{\mu}^{**}(\tau) \right\|_{L^{q'}(0, \infty)} \\
& \leq C_3 \left\| t^{\frac{1}{p'} - \frac{1}{q'}} c(t) f_{\mu}^{**}(t) \right\|_{L^{q'}(0, \infty)} = C_3 \|f\|_{L^{(p', q', c)}(0, \infty)}.
\end{aligned} \tag{4.10}$$

The first inequality is true thanks to [33, Lemma 4.1]. Now we prove the second inequality. If $q > 1$, then we use [19, Theorem 3.2] to prove it. Owing to this theorem the second inequality is valid if and only if for every $t \in (0, \infty)$ it is true that

$$t^{\frac{m}{D}} \left\| \tau^{\frac{1}{p'} - \frac{1}{q'} - \frac{m}{D}} c(\tau) \right\|_{L^{q'}(0, t)} \leq K \left\| \tau^{\frac{1}{p'} - \frac{1}{q'}} c(\tau) \right\|_{L^{q'}(0, t)}, \tag{4.11}$$

where $K > 0$ is a constant that does not depend on t , which is not hard to verify (see [39, Lemma 2.16]). If $q = 1$, then $q' = \infty$, so the second inequality in (4.10) is of the form

$$\begin{aligned}
& C_2 \sup_{t \in (0, \infty)} t^{\frac{1}{p'} - \frac{m}{D}} c(t) \sup_{\tau \in [t, \infty)} \tau^{\frac{m}{D}} f_{\mu}^{**}(\tau) = C_2 \sup_{\tau \in (0, \infty)} \tau^{\frac{m}{D}} f_{\mu}^{**}(\tau) \sup_{t \in (0, \tau)} t^{\frac{1}{p'} - \frac{m}{D}} c(t) \\
& \leq C_3 \sup_{t \in (0, \infty)} t^{\frac{1}{p'}} c(t) f_{\mu}^{**}(t) = C_3 \|f\|_{L^{(p', \infty, c)}(0, \infty)}.
\end{aligned}$$

Note that we used the fact that b^{-1} is equivalent to a nondecreasing function on $(0, \infty)$ if $p = \frac{D}{D-m}$. It means that (4.10) is true. So, the operator $T_{\frac{m}{D}}$ is bounded on the space $(L^{p, q, b}(0, \infty))'$. If $p = q = \infty$, then we use [39, Proposition 3.3 and Theorem 3.31] to obtain the fact that $(L^{\infty, \infty, b}(\Sigma, \mu))'$ is equivalent to $L^{1, 1, h^{-1}}(\Sigma, \mu)$, where h is the slowly varying function given by $h(t) = \sup_{\tau \in (0, t)} b(\tau)$, $t \in (0, \infty)$. The boundedness of the operator $T_{\frac{m}{D}}$ in this case can be proved similarly as in the preceding cases hence we omit the proof. Now it follows from the proof of [18, Theorem 4.1] that the condition (3.23) is satisfied. It means that the norm of the optimal space $U_m Y(\Sigma, \mu)$ given by Theorem 3.9 is equivalent to (4.9). So, we have proved the assertion (3) of the theorem.

It remains to prove that (4.9) is equivalent to the norm of the space $U_m Y(\Sigma, \mu)$ in the first two cases. For each $f \in \mathcal{M}(0, \infty)$ we obtain

$$\begin{aligned} & \left\| \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{p,q,b}(0,\infty)} = \left\| t^{\frac{1}{p}-\frac{1}{q}} b(t) \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{p,q,b}(0,\infty)} \\ & \geq \left\| t^{\frac{1}{p}-\frac{1}{q}} b(t) \int_t^{2t} f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{p,q,b}(0,\infty)} \geq C_4 \left\| t^{\frac{1}{p}-\frac{1}{q}+\frac{m}{D}} b(t) f_\mu^*(2t) \right\|_{L^q(0,\infty)} \\ & \geq C_5 \left\| t^{\frac{1}{p}-\frac{1}{q}+\frac{m}{D}} b(t) f_\mu^*(t) \right\|_{L^q(0,\infty)} = C_5 \|f\|_{L^{\frac{Dp}{D+mp},q,b}(0,\infty)}. \end{aligned}$$

In the third inequality, we used the boundedness of the dilation operator (see (2.14)). On the other hand, for every $f \in \mathcal{M}(0, \infty)$ we have

$$\begin{aligned} & \left\| \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^{p,q,b}(0,\infty)} = \left\| t^{\frac{1}{p}-\frac{1}{q}} b(t) \int_t^\infty f_\mu^*(\tau) \tau^{\frac{m}{D}-1} d\tau \right\|_{L^q(0,\infty)} \\ & \leq C_6 \left\| t^{\frac{1}{p}-\frac{1}{q}+\frac{m}{D}} b(t) f_\mu^*(t) \right\|_{L^q(0,\infty)} = C_6 \|f\|_{L^{\frac{Dp}{D+mp},q,b}(0,\infty)}. \end{aligned}$$

By virtue of the weighted Hardy inequality ([32, Theorem 2]), the preceding inequality is valid if and only if

$$\sup_{t \in (0,\infty)} \left\| \tau^{\frac{1}{p}-\frac{1}{q}} b(\tau) \right\|_{L^q(0,t)} \left\| \tau^{\frac{1}{q}-\frac{1}{p}-1} b^{-1}(\tau) \right\|_{L^{q'}(t,\infty)} < \infty,$$

which is not hard to verify (see [39, Lemma 2.16]). \square

Remark 4.4. It is a natural question what we can say about the optimal domain space for the Lorentz–Karamata space $Y(\Sigma, \mu) = L^{p,q,b}(\Sigma, \mu)$ in the case $p = \frac{D}{D-m}, q \in (1, \infty]$. In this case we have

$$\frac{t^{1-\frac{m}{D}}}{\varphi_Y(t)} = \frac{t^{1-\frac{m}{D}}}{\left\| \chi_{(0,t)} \right\|_{L^{\frac{D}{D-m},q,b}(0,\infty)}} = \frac{t^{1-\frac{m}{D}}}{\left\| \tau^{1-\frac{m}{D}-\frac{1}{q}} b(\tau) \right\|_{L^q(0,t)}}. \quad (4.12)$$

It can be proved by [39, Lemma 2.16] and elementary computations that the right-hand side of (4.12) is equivalent to b^{-1} on $(0, \infty)$. It means that the condition (3.23) is satisfied if and only if b is equivalent to a nonincreasing function on $(1, \infty)$. By Theorem 3.9 we conclude that the optimal domain space for $Y(\Sigma, \mu) = L^{\frac{D}{D-m},q,b}(\Sigma, \mu)$ exists if and only if b is equivalent to a nonincreasing function on $(1, \infty)$. Furthermore, if this is not the case, then there is no domain space X with which (3.25) is valid.

On the other hand, it can be proved by [19, Theorem 3.2] that the operator $T_{\frac{m}{D}}$ is not bounded on the space $\left(L^{\frac{D}{D-m},q,b} \right)'(0, \infty)$, which is equivalent to $L^{\frac{D}{m},q',b^{-1}}(0, \infty)$ owing to [39, Section 3.7]. In particular, notice that the condition (4.11) is not satisfied, which is not hard to verify thanks to [39, Lemma 2.16]). The unboundedness of $T_{\frac{m}{D}}$ on $\left(L^{\frac{D}{D-m},q,b} \right)'(0, \infty)$ means that

we cannot simplify the norm of the optimal space $U_m Y(\Sigma, \mu)$ ([18, Theorem 4.7 and Remark 4.8]).

REFERENCES

- [1] A. Alberico, A. Cianchi, L. Pick, and L. Slavíková. Sharp Sobolev type embeddings on the entire Euclidean space. *Commun. Pure Appl. Anal.*, 17(5):2011–2037, 2018. ISSN 1534-0392,1553-5258. doi: 10.3934/cpaa.2018096. URL <https://doi.org/10.3934/cpaa.2018096>.
- [2] C. Bennett and R. Sharpley. *Interpolation of operators*, volume 129 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1988. ISBN 0-12-088730-4.
- [3] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*, volume No. 223 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin-New York, 1976.
- [4] H. Brézis and S. Wainger. A note on limiting cases of Sobolev embeddings and convolution inequalities. *Comm. Partial Differential Equations*, 5(7):773–789, 1980. ISSN 0360-5302,1532-4133. doi: 10.1080/03605308008820154. URL <https://doi.org/10.1080/03605308008820154>.
- [5] X. Cabré and X. Ros-Oton. Regularity of stable solutions up to dimension 7 in domains of double revolution. *Comm. Partial Differential Equations*, 38(1):135–154, 2013. ISSN 0360-5302,1532-4133. doi: 10.1080/03605302.2012.697505. URL <https://doi.org/10.1080/03605302.2012.697505>.
- [6] X. Cabré and X. Ros-Oton. Sobolev and isoperimetric inequalities with monomial weights. *J. Differential Equations*, 255(11):4312–4336, 2013. ISSN 0022-0396,1090-2732. doi: 10.1016/j.jde.2013.08.010. URL <https://doi.org/10.1016/j.jde.2013.08.010>.
- [7] X. Cabré, X. Ros-Oton, and J. Serra. Sharp isoperimetric inequalities via the ABP method. *J. Eur. Math. Soc. (JEMS)*, 18(12):2971–2998, 2016. ISSN 1435-9855,1435-9863. doi: 10.4171/JEMS/659. URL <https://doi.org/10.4171/JEMS/659>.
- [8] H. Castro. Hardy-Sobolev-type inequalities with monomial weights. *Ann. Mat. Pura Appl. (4)*, 196(2):579–598, 2017. ISSN 0373-3114,1618-1891. doi: 10.1007/s10231-016-0587-2. URL <https://doi.org/10.1007/s10231-016-0587-2>.
- [9] H. Castro. Extremals for Hardy-Sobolev type inequalities with monomial weights. *J. Math. Anal. Appl.*, 494(2):Paper No. 124645, 31, 2021. ISSN 0022-247X,1096-0813. doi: 10.1016/j.jmaa.2020.124645. URL <https://doi.org/10.1016/j.jmaa.2020.124645>.
- [10] A. Cianchi and L. Pick. Sobolev embeddings into BMO, VMO, and L_∞ . *Ark. Mat.*, 36(2):317–340, 1998. ISSN 0004-2080,1871-2487. doi: 10.1007/BF02384772. URL <https://doi.org/10.1007/BF02384772>.

- [11] A. Cianchi and L. Pick. Optimal Gaussian Sobolev embeddings. *J. Funct. Anal.*, 256(11):3588–3642, 2009. ISSN 0022-1236,1096-0783. doi: 10.1016/j.jfa.2009.03.001. URL <https://doi.org/10.1016/j.jfa.2009.03.001>.
- [12] A. Cianchi and L. Pick. Optimal Sobolev trace embeddings. *Trans. Amer. Math. Soc.*, 368(12):8349–8382, 2016. ISSN 0002-9947,1088-6850. doi: 10.1090/tran/6606. URL <https://doi.org/10.1090/tran/6606>.
- [13] A. Cianchi, L. Pick, and L. Slavíková. Higher-order Sobolev embeddings and isoperimetric inequalities. *Adv. Math.*, 273:568–650, 2015. ISSN 0001-8708,1090-2082. doi: 10.1016/j.aim.2014.12.027. URL <https://doi.org/10.1016/j.aim.2014.12.027>.
- [14] A. Cianchi, L. Pick, and L. Slavíková. Sobolev embeddings, rearrangement-invariant spaces and Frostman measures. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 37(1):105–144, 2020. ISSN 0294-1449,1873-1430. doi: 10.1016/j.anihpc.2019.06.004. URL <https://doi.org/10.1016/j.anihpc.2019.06.004>.
- [15] E. Cinti, F. Glaudo, A. Pratelli, X. Ros-Oton, and J. Serra. Sharp quantitative stability for isoperimetric inequalities with homogeneous weights. *Trans. Amer. Math. Soc.*, 375(3):1509–1550, 2022. ISSN 0002-9947,1088-6850. doi: 10.1090/tran/8525. URL <https://doi.org/10.1090/tran/8525>.
- [16] D. E. Edmunds and W. D. Evans. *Hardy operators, function spaces and embeddings*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004. ISBN 3-540-21972-2. doi: 10.1007/978-3-662-07731-3. URL <https://doi.org/10.1007/978-3-662-07731-3>.
- [17] D. E. Edmunds, R. Kerman, and L. Pick. Optimal Sobolev imbeddings involving rearrangement-invariant quasinorms. *J. Funct. Anal.*, 170(2):307–355, 2000. ISSN 0022-1236,1096-0783. doi: 10.1006/jfan.1999.3508. URL <https://doi.org/10.1006/jfan.1999.3508>.
- [18] D. E. Edmunds, Z. Mihula, V. Musil, and L. Pick. Boundedness of classical operators on rearrangement-invariant spaces. *J. Funct. Anal.*, 278(4):108341, 56, 2020. ISSN 0022-1236,1096-0783. doi: 10.1016/j.jfa.2019.108341. URL <https://doi.org/10.1016/j.jfa.2019.108341>.
- [19] A. Gogatishvili, B. Opic, and L. Pick. Weighted inequalities for Hardy-type operators involving suprema. *Collect. Math.*, 57(3):227–255, 2006. ISSN 0010-0757,2038-4815.
- [20] L. Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014. ISBN 978-1-4939-1193-6; 978-1-4939-1194-3. doi: 10.1007/978-1-4939-1194-3. URL <https://doi.org/10.1007/978-1-4939-1194-3>.
- [21] P. Gurka and D. Hauer. More insights into the Trudinger-Moser inequality with monomial weight. *Calc. Var. Partial Differential Equations*, 60(1):Paper No. 16, 27, 2021. ISSN 0944-2669,1432-0835. doi: 10.1007/s00526-020-01890-7. URL <https://doi.org/10.1007/s00526-020-01890-7>.

- [22] K. Hansson. Imbedding theorems of Sobolev type in potential theory. *Math. Scand.*, 45(1):77–102, 1979. ISSN 0025-5521,1903-1807. doi: 10.7146/math.scand.a-11827. URL <https://doi.org/10.7146/math.scand.a-11827>.
- [23] R. Kerman and L. Pick. Optimal Sobolev imbeddings. *Forum Math.*, 18(4):535–570, 2006. ISSN 0933-7741,1435-5337. doi: 10.1515/FORUM.2006.028. URL <https://doi.org/10.1515/FORUM.2006.028>.
- [24] N. Lam. Sharp Trudinger-Moser inequalities with monomial weights. *NoDEA Nonlinear Differential Equations Appl.*, 24(4):Paper No. 39, 21, 2017. ISSN 1021-9722,1420-9004. doi: 10.1007/s00030-017-0456-8. URL <https://doi.org/10.1007/s00030-017-0456-8>.
- [25] G. Leoni. *A first course in Sobolev spaces*, volume 181 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2017. ISBN 978-1-4704-2921-8. doi: 10.1090/gsm/181. URL <https://doi.org/10.1090/gsm/181>.
- [26] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001. ISBN 0-8218-2783-9. doi: 10.1090/gsm/014. URL <https://doi.org/10.1090/gsm/014>.
- [27] J. Malý, D. Swanson, and W. P. Ziemer. The co-area formula for Sobolev mappings. *Trans. Amer. Math. Soc.*, 355(2):477–492, 2003. ISSN 0002-9947,1088-6850. doi: 10.1090/S0002-9947-02-03091-X. URL <https://doi.org/10.1090/S0002-9947-02-03091-X>.
- [28] Z. Mihula. Embeddings of homogeneous Sobolev spaces on the entire space. *Proc. Roy. Soc. Edinburgh Sect. A*, 151(1):296–328, 2021. ISSN 0308-2105,1473-7124. doi: 10.1017/prm.2020.14. URL <https://doi.org/10.1017/prm.2020.14>.
- [29] Z. Mihula. Poincaré-Sobolev inequalities with rearrangement-invariant norms on the entire space. *Math. Z.*, 298(3-4):1623–1640, 2021. ISSN 0025-5874,1432-1823. doi: 10.1007/s00209-020-02652-z. URL <https://doi.org/10.1007/s00209-020-02652-z>.
- [30] Z. Mihula. Optimal behavior of weighted Hardy operators on rearrangement-invariant spaces. *Math. Nachr.*, 296(8):3492–3538, 2023. ISSN 0025-584X,1522-2616.
- [31] J. Moser. A sharp form of an inequality by N. Trudinger. *Indiana Univ. Math. J.*, 20:1077–1092, 1971. ISSN 0022-2518,1943-5258. doi: 10.1512/iumj.1971.20.20101. URL <https://doi.org/10.1512/iumj.1971.20.20101>.
- [32] B. Muckenhoupt. Hardy’s inequality with weights. *Studia Math.*, 44:31–38, 1972. ISSN 0039-3223,1730-6337. doi: 10.4064/sm-44-1-31-38. URL <https://doi.org/10.4064/sm-44-1-31-38>.
- [33] V. Musil and R. Olhava. Interpolation theorem for Marcinkiewicz spaces with applications to Lorentz gamma spaces. *Math. Nachr.*, 292(5):1106–1121, 2019. ISSN 0025-584X,1522-2616. doi: 10.1002/mana.201700452. URL <https://doi.org/10.1002/mana.201700452>.

- [34] J. S. Neves. Lorentz-Karamata spaces, Bessel and Riesz potentials and embeddings. *Dissertationes Math. (Rozprawy Mat.)*, 405:46, 2002. ISSN 0012-3862,1730-6310. doi: 10.4064/dm405-0-1. URL <https://doi.org/10.4064/dm405-0-1>.
- [35] R. O’Neil. Convolution operators and $L(p, q)$ spaces. *Duke Math. J.*, 30:129–142, 1963. ISSN 0012-7094,1547-7398. URL <http://projecteuclid.org/euclid.dmj/1077374532>.
- [36] B. Opic and L. Pick. On generalized Lorentz-Zygmund spaces. *Math. Inequal. Appl.*, 2(3):391–467, 1999. ISSN 1331-4343,1848-9966. doi: 10.7153/mia-02-35. URL <https://doi.org/10.7153/mia-02-35>.
- [37] J. Peetre. Espaces d’interpolation et théorème de Soboleff. *Ann. Inst. Fourier (Grenoble)*, 16:279–317, 1966. ISSN 0373-0956,1777-5310. URL http://www.numdam.org/item?id=AIF_1966__16_1_279_0.
- [38] D. Peša. Reduction principle for a certain class of kernel-type operators. *Math. Nachr.*, 293(4):761–773, 2020. ISSN 0025-584X,1522-2616. doi: 10.1002/mana.201800510. URL <https://doi.org/10.1002/mana.201800510>.
- [39] D. Peša. Lorentz–Karamata spaces, preprint April 2023. arXiv:2006.14455v4.
- [40] L. Pick, A. Kufner, O. John, and S. Fučík. *Function spaces. Vol. 1*, volume 14 of *De Gruyter Series in Nonlinear Analysis and Applications*. Walter de Gruyter & Co., Berlin, extended edition, 2013. ISBN 978-3-11-025041-1; 978-3-11-025042-8.
- [41] W. Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987. ISBN 0-07-054234-1.
- [42] B. Ruf. Lorentz spaces and nonlinear elliptic systems. In *Contributions to nonlinear analysis*, volume 66 of *Progr. Nonlinear Differential Equations Appl.*, pages 471–489. Birkhäuser, Basel, 2006. ISBN 978-3-7643-7149-4; 3-7643-7149-8. doi: 10.1007/3-7643-7401-2_32. URL https://doi.org/10.1007/3-7643-7401-2_32.
- [43] E. M. Stein. Editor’s note: the differentiability of functions in \mathbf{R}^n . *Ann. of Math. (2)*, 113(2):383–385, 1981. ISSN 0003-486X.
- [44] E. M. Stein and G. Weiss. *Introduction to Fourier analysis on Euclidean spaces*, volume No. 32 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1971.
- [45] D. Swanson. Area, coarea, and approximation in $W^{1,1}$. *Ark. Mat.*, 45(2):381–399, 2007. ISSN 0004-2080,1871-2487. doi: 10.1007/s11512-007-0051-z. URL <https://doi.org/10.1007/s11512-007-0051-z>.
- [46] G. Talenti. Inequalities in rearrangement invariant function spaces. In *Nonlinear analysis, function spaces and applications, Vol. 5 (Prague, 1994)*, pages 177–230. Prometheus, Prague, 1994. ISBN 80-85849-69-0.
- [47] L. Tartar. Imbedding theorems of Sobolev spaces into Lorentz spaces. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8)*, 1(3):479–500, 1998. ISSN 0392-4041.

- [48] N. S. Trudinger. On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.*, 17:473–483, 1967. doi: 10.1512/iumj.1968.17.17028. URL <https://doi.org/10.1512/iumj.1968.17.17028>.
- [49] J. Wang. Weighted Hardy-Sobolev, log-Sobolev and Moser-Onofri-Beckner inequalities with monomial weights. *Potential Anal.*, 58(2):225–240, 2023. ISSN 0926-2601,1572-929X. doi: 10.1007/s11118-021-09938-9. URL <https://doi.org/10.1007/s11118-021-09938-9>.

LADISLAV DRÁŽNÝ, CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF MATHEMATICAL ANALYSIS, SOKOLOVSKÁ 49/83, 186 75 PRAHA 8, CZECH REPUBLIC; CZECH TECHNICAL UNIVERSITY IN PRAGUE, FACULTY OF ELECTRICAL ENGINEERING, DEPARTMENT OF MATHEMATICS, TECHNICKÁ 2, 166 27 PRAHA 6, CZECH REPUBLIC

Email address: drazny@karlin.mff.cuni.cz

ORCID: 0009-0003-1807-6003