## arXiv:2407.07826v1 [cond-mat.soft] 10 Jul 2024

## Using Activity to Compartmentalize Binary Mixtures

Nicholas J Lauersdorf, Ehssan Nazockdast,<sup>\*</sup> and Daphne Klotsa<sup>†</sup>

Department of Applied Physical Sciences, University of North Carolina, Chapel Hill, North Carolina 27515, USA.

We computationally study suspensions of slow and fast active Brownian particles that have undergone motility induced phase separation and are at steady state. Such mixtures, of varying non-zero activity, remain largely unexplored even though they are relevant for a plethora of systems and applications ranging from cellular biophysics to drone swarms. Our mixtures are modulated by their activity ratios ( $Pe^R$ ), which we find to encode information by giving rise to three regimes, each of which display their unique emergent behaviors. Specifically, we found non-monotonic behavior of macroscopic properties, *e.g.* density and pressure, as a function of activity ratio, microphase separation of fast and slow particle domains, increased fluctuations on the interface and severe avalanche events compared to monodisperse active systems. Our approach of simultaneously varying the two activities of the particle species allowed us to discover these behaviors and explain the microscopic physical mechanisms that drive them.

DOI: Secondary publications and information retrieval purposes.

Active matter describes nonequilibrium Introduction. systems comprised of components that locally consume energy to move, such as swarms of bacteria and clusters of self-propelled colloids. Past studies on monodisperse systems and minimal models have led to the discovery of a range of remarkable emergent phenomena including motility-induced phase separation (MIPS) [1–10], swarming and flocking [11–19], and pattern formation [20– 34]. Adding complexity, studies of polydisperse mixtures have included components of different sizes [35, 36], shapes [37, 38], interactions [15, 28, 39-45], and propulsion mechanisms [15, 16, 37, 46-51]. The relative activities of the particles have mostly been varied in the limiting case of active/passive mixtures [52–62], passive particles in active baths [32, 63–77], and active particles with static obstacles [40, 78–91]. However, mixtures of particles with distinct nonzero active driving forces remain largely unexplored, even though they are relevant in a variety of systems ranging from fundamental physics to applications, as we explain next. Recent studies are an exception and show there is growing interest [92–97].

Studying active/active mixtures is interesting from a fundamental physics standpoint, since it is a simple nonequilibrium model system that challenges our understanding of what collective behavior to expect, and what quantities equilibrate; still it is sufficiently simple to allow theoretical progress from first principles. We often think of activity as a temperature analogue. But while fast particles can effectively exchange activity with slower ones when in contact, as has been observed in active/passive [69–71, 82, 98–109] and slow/fast [96] systems, they do not reach a uniform activity and so the answer is unclear. In biology there are many active matter systems of importance that are relevant to biophysics and medicine, and they are typically polydisperse including in their motility speeds [110–118], such as bacteria and sperm cells. Furthermore in synthetic systems, limitations in fabrication techniques give rise to a distribution of swim speeds in self-propelled colloids [119–122], but there is limited understanding of how that affects the emergent behavior. Finally from a soft-matter and materials physics perspective, tuning the relative activities of particles opens an enormous parameter space with potentially a wealth of physics providing opportunities for more control in designing dynamic complex assemblies. Active matter mixtures exhibit behaviors akin to those found in living systems *e.g.* nonequilibrium transitions, microphase separation, and bistable states, that have not been observed before.

In this paper, we performed simulations of binary active mixtures of fast and slow Active Brownian Particles (ABPs), that have undergone motility induced phase separation (MIPS) and are in steady state. We studied the properties of the phase separated system as a function of the ratio of the slow to fast particle activities,  $0 \leq \mathrm{Pe}^{\mathrm{R}} \leq 1$ , and discovered three regimes corresponding to small, intermediate and large activity ratios. At large activity ratios the fast and slow particles are uniformly mixed and the behavior is analogous to monodisperse ABP suspensions, even when the slow activity is nearly a third of the fast. At intermediate and small activity ratios, when the particles are increasingly heterogeneous in their activities, is where we saw the most interesting behavior: *Microscopically*, the system exhibits microphase separation, increased avalanche events and fluctuations, and active herding (slow particles pushed by the fast ones). We also found nonmonotonic behavior in macroscopic quantities, including cluster pressure, density and compressibility. To obtain a deeper understanding, we developed a coarse-grained continuum model, which provided further insight and whose predictions were in good agreement with the simulation results. We thus propose a physical mechanism that links microscopic and macroscopic behavior and explains the observed emergent phenomena.

Model & Methods: Using HOOMD-Blue [123–125], we

<sup>\*</sup> ehssan@email.unc.edu

 $<sup>^\</sup>dagger$ dklotsa@unc.edu



FIG. 1. a) A simulation snapshot of a monodisperse ABP system at steady state after MIPS with Pe = 500. Phases are labeled bulk (green), interface (orange), and gas (purple) and average particle orientation is shown as arrows. b) Bulk (cluster) (i) interparticle pressure, (ii) number density, and (iii) compressibility normalized by their respective values in monodisperse (same activity,  $Pe^{R} = 1$ ) case, as a function of activity ratio (Pe<sup>R</sup>). c) The steady-state fast particle fraction in the bulk phase ( $\chi_{\rm B}^{\rm F}$ , green circle) and the interface ( $\chi_{\rm I}^{\rm F}$ , orange cross) as a function of activity ratio (Pe<sup>R</sup>). The solid black line represents the 1D-continuum model predictions. In (b) and (c), three regimes, corresponding to three range of activity ratios, are visualized by background color and separated by dashed black lines. d) Number density of slow (blue) and fast (red) particles for our ABP simulation (dots) and 1D continuum model (solid line) as a function of distance from the cluster's center of mass (r) normalized by the cluster radius  $(r_c)$  for  $Pe^R = 0.1, 0.3$ , and 1.0.

simulated ABPs of diameter  $\sigma$  that translate and rotate over time (measured in units of the Brownian diffusion time,  $\tau_{\rm B}$ ) in accordance with overdamped Langevin dynamics and interact via the Weeks-Chandler-Anderson (WCA) potential with a soft repulsive well depth ( $\epsilon = 1$ ). Each particle is subject to an active force,  $\mathbf{F}_{\rm a} = \xi v_{\rm p} \hat{\mathbf{p}}$ , where  $v_{\rm p}$  is the swim speed in the absence of collisions and it modulates the activity, quantified through the Peclet number (Pe =  $3v_{\rm p}\tau_{\rm r}/\sigma$  where  $\tau_{\rm r}$  is the characteristic reorientation time). We simulate mixtures of fast (Pe<sup>F</sup>) and slow (Pe<sup>S</sup>) ABPs whose activities are varied independently between 0-500 with particle fractions  $\chi^{\rm F}$  and  $\chi^{\rm S}$  respectively, where  $\chi^{\rm F} = 1.0 - \chi^{\rm S}$ , varied between 0.2-0.8 at  $\phi = 0.6$ . We ran a range of relative fractions (see SI §1); in the main paper for simplicity we show results for  $\chi^{\rm F} = \chi^{\rm S} = 0.5$ , but the findings remain the same. We define the activity ratio  $Pe^{R} = \frac{Pe^{S}}{Pe^{F}}$ , ranging from a monodisperse active system ( $Pe^{R} = 1$ ) to the most heterogeneous ( $Pe^{R} = 0$ ) which is a mixture of passive and active particles (Fig. S1). It is useful to also define the net activity,  $Pe^{Net} = \chi^{S}Pe^{S} + \chi^{F}Pe^{F}$  [92], varied between 0-500. More details about the model and the simulation parameters can be found in the SI §1.

We studied systems that have undergone MIPS and have reached steady state. We ran simulations using a variety of initial conditions, e.g. random gas and differently-instantiated clusters, and for very long times  $(600\tau_B)$ ; unlike previous works [52] we found that all initializations eventually led to the same steady-states for  $Pe^{R} \geq 0.2$ . For  $Pe^{R} < 0.2$ , the large fluctuations in the cluster size prolong MIPS and make it much less likely; hence, for low activity ratios we instantiated the clusters (and then observed them melt or persist over long times). We provide a detailed discussion in SI §2. We distinguish three phases shown in Fig. 1(a): 1) a dilute *qas* phase with random alignment of active forces; 2) a bulkwith uniform high density and also random alignment of active forces; and 3) the *interface*, that sits between the bulk and the gas, and is characterized by a transition from high to low density and a net alignment of particles' active forces and a net compressive force towards the center of the cluster [126] (SI §3).

Results: We begin by studying the macroscopic mechanical properties of the bulk phase, specifically the average pressure as a function of the activity ratio, Pe<sup>R</sup>. In monodisperse systems, pressure has a nearly linear dependence with activity [126, 127]. To remove this dependence and focus on the activity ratio, for each system we normalize the pressure by the computed pressure at activity ratio 1, denoted  $\Pi_{\rm B}^{0}$ . This non-dimensionalization collapses all the data from different net activities into a single curve (Figs. 1(b)(i), S2). As shown, the dimensionless pressure displays nonmonotonic variations with activity ratio with a minimum around  $Pe^{R} \approx 0.35$ , followed by an increase and then a plateau at small activity ratios,  $\text{Pe}^{\text{R}} < 0.175$ . The number density of particles in the bulk phase  $(n_{\text{B}})$  and compressibility  $(\beta_{\text{B}} = \frac{1}{n_{\text{B}}} \frac{\partial n_{\text{B}}}{\partial \Pi_{\text{B}}})$  also show non-monotonic behavior with Pe<sup>R</sup> similar to the pressure, (Fig.1(b)(ii) and (iii), and S2). From these observations, we can define three regimes in terms of activity ratio: large  $0.35 < \text{Pe}^{\text{R}} \le 1$ , intermediate  $0.175 < \text{Pe}^{\text{R}} < 0.35$ and small  $Pe^{R} < 0.175$ . These regimes emerge naturally and consistently from our results obtained from independently measured quantities besides pressure, density and compressibility, as will be shown in this paper.

To explore the microscopic basis of this nonmonotonic behavior, we computed the relative composition of fast and slow particles in the cluster (Figs. 1(c) and S3). Note that similar to the dimensionless pressure, the fraction of fast (and slow) particles is only a function of activity ratio and independent of the net activity. We observe that as  $Pe^{R}$  decreases the fast particle fraction increases in the bulk and the interface, particularly for intermediate



FIG. 2. a) Simulation snapshots of an active/passive mixture zoomed-in on the interface showing the formation of a segregated domain of passive particles from the interface (top row) and its incorporation into the bulk (two bottom rows). b) The degree of segregation for slow-slow, slow-fast, and fast-fast neighbor pairs within the bulk as a function of activity ratio (Pe<sup>R</sup>). The inset shows segregation at the interface; see also SI section §5. c) Avalanche parameter ( $\kappa$ , green circle) and compression force fluctuations ( $H(F_{\rm I})$  (defined in SI §6), orange cross) normalized by their respective values in monodisperse (same activity, Pe<sup>R</sup> = 1) case, vs ratio (Pe<sup>R</sup>).

and low  $\text{Pe}^{\text{R}}$  (Figs. 1(c), S3). At the interface, because the active forces are aligned, an increase of the number of fast particles leads to a larger compression force pressing towards the bulk. Hence, the otherwise counterintuitive uptick in the bulk pressure at intermediate  $\text{Pe}^{\text{R}}$ (Fig. 1(b)(i)) may result from this increase in compression force. However, it remains unclear why there are more fast particles in the cluster in the intermediate and low  $\text{Pe}^{\text{R}}$  regimes. To search for answers, next we considered the spatial distribution of fast and slow particles.

As shown in Fig. 1(d), for large activity ratios, including the monodisperse case  $Pe^{R} = 1$ , the number density of fast and slow particles is nearly equal and increases across the interface from a low uniform value in the gas to a large uniform value in the bulk. For intermediate and low activity ratios, however, we see a split in how fast and slow particles are distributed. At the interface, the particle fraction radially transitions from majority slow particles nearest to the gas to majority fast nearest to the bulk, similar to active/passive MIPS [57] and slow and fast ABPs assembling on a rigid wall [97]. This observation can be understood by thinking of the dense cluster as a rigid boundary, on which fast and slow particles assemble. In the simplest case of a dilute suspension of fast and slow point particles, the density variations with distance from the rigid boundary (bulk surface), x, simplify to:  $n^{\text{S,F}} \approx n_0^{\text{S,F}} + E \exp(-2x \text{Pe}^{(\text{S,F})}/\sigma)$ , where  $n_0$  is the net (average) density of slow or fast particles in the gas phase, and E is an integration constant determined by boundary conditions; see SI §4 for details. As a result, we expect the fast particles to concentrate on the interior of the cluster and form a thinner boundary layer, compared to slow particles, which is in agreement with simulation results in Fig 1(d).

A close inspection of simulation movies revealed that instead of two distinct radial layers, we observe microphase separated domains of fast and slow particles at the interface, which get integrated into the bulk as well (Fig 2(a)). To quantify microphase separation as a function of activity ratio, we computed the pair probability density of observing slow-slow, slow-fast and fastfast pairs as nearest neighbors in both the bulk and interface normalized by the average slow or fast particle fraction of the respective phase (Fig 2(b)). At large activity ratios, the particles are spatially uniform and bulk or interface segregation goes to 1. At intermediate and low activity ratios, we observe a significant increase in the probability of observing slow-slow and to a lesser degree fast-fast pairs and, therefore, lower probability of observing slow-fast pairs. We are thus showing that microphase separation, reported previously mainly for active/passive [35, 36, 52, 55, 57, 58, 92], is a function of activity ratio and begins much earlier in the intermediate  $Pe^{R}$  regime.

Simulation results indicate that microphase separated domains start at the interface and move into the bulk (Fig. 2(a)). To test this, we developed a 1D coarsegrained model that solves for the time-dependent variations in the number density and average alignment of two different active species in ABP suspensions. The only ingredients of the model are thermal diffusion, activity, and interparticle forces; details are provided in the SI §4. Interestingly, the model predicted that during the early stages of MIPS, at intermediate activity ratios, the interface and the bulk phase are primarily composed of fast particles. With time, the density of fast particles is reduced, while the density of slow particles is increased. This supports that microphase separation starts at the interface and gets incorporated into the bulk, see also (SI §4). Furthermore, the results of the model are in agreement with the computed splitting of slow and fast particles' number density at the interface for low and in-



FIG. 3. Schematic of the proposed mechanism showing the process of active herding, microphase separation, and enhanced avalanche events.

termediate activity ratios shown in Fig 1(a)(i) and (ii). The model also correctly predicts the enrichment of fast particles in the dense phase at low and intermediate activity ratios.

In monodisperse systems the active particles leave the interface when they orient away from the cluster. The desorption timescale is, thus, determined by the rotational diffusion timescale of the particles. However, in simulations it has been shown that the desorption of a particle typically coincides with large groups of particles collectively escaping from the cluster via *avalanche-like events* [2, 97, 126, 128–131]. We expect that in our active mixtures the presence of microphase-separated domains especially at the interface would lead to stronger fluctuations in the compression force (that holds the cluster together) and thus avalanche events.

To test this hypothesis, we turned to our ABP simulations and computed the desorption rate of particles from the cluster at different activity ratios. The avalanche parameter is defined as the ratio of the desorption rate from simulations to that predicted from rotational diffusion time, i.e. larger values denote stronger avalanche events. We also computed the fluctuations around the average compression force at the interface, integrated over the thickness of the interface and over time, at different activity ratios (Fig 2(c)). Both quantities are normalized by their corresponding values in the monodisperse system to study the effect of activity ratio. We find that both the avalanche parameter and the compression force fluctuations trace each other over the entire range of activity ratios. More importantly, both parameters closely follow the trend observed in the bulk (and interface) segregation parameter: they remain nearly unchanged at larger activity ratios and show a sharp increase at intermediate and low activity ratios. These observations support the following mechanism: microphase separation at the interface leads to increases in the fluctuations of compression and much stronger avalanche events, where up to half of the particles in the cluster escape (Fig. S4). In contrast, at high activity ratios fewer than 2% of the cluster particles escape during avalanche events (Fig. S4).

The steady state microstructure is determined by the balance of adsorption and desorption of each particle species. Thus, we expect the strong avalanche events at intermediate and low  $Pe^{R}$  (i.e. high desorption), to be balanced by enhanced adsorption. Indeed, simulations show that fast particles push and deposit slow particles from the gas onto the interface, which we call *active herd-ing* (SI §7). As they do so, the fast particles push through the slow layer at the interface and assemble closer to the bulk boundary or move into the bulk phase.

Based on these observations, we propose the following mechanism for the steady state of binary active mixtures at intermediate and low Pe<sup>R</sup>: MIPS begins with the nucleation of small clusters of fast particles. These clusters act as rigid boundaries on which slow and fast particles adsorb. Fast particles herd slow particles leading to slow enhanced adsorption. The cluster overall has more fast particles than slow, while the net density remains roughly unchanged with activity ratio. This leads to the nonmonotonic behavior in the macroscopic quantities (pressure etc). The difference in activities leads to microphase separation of slow and fast particles at the interface, with fast particles assembling at the inner layer closer to the bulk and the slow particles assembling at the outer layer of the interface closer to the gas phase. As more particles assemble around the existing interface, the cluster grows in size and the phase-separated slow and fast particles that were making up the interface at earlier times get integrated to the cluster. The microphaseseparated domains at the interface lead to larger fluctuations in the compression force which in turn leads to enhanced avalanche events. The process repeats with the re-adsorption of fast and slow particles from the gas onto the interface through active herding.

We have shown that our approach of continuously varying the activity of particle species revealed the existence of three regimes of Péclet ratio, previously unknown. The regimes that emerge are robust and demonstrate several distinct structural and dynamical features in micro- and macro-scale. Thus the behavior of mixtures is not a simple interpolation between the active monodisperse case and active/passive; rather there is a richness in between the extremes. Based on our results, we propose a mechanism that describes the dynamic steadystate of the system at low and intermediate Pe<sup>R</sup> through a series of steps that include interesting nonequilibrium phenomena including active herding, microphase separation, active force fluctuations at the interface and strong avalanche events. We emphasize the role of the interface, which compartmentalizes the bulk, acting like a semipermeable membrane that selectively filters out slow particles and allows fast particles in. In other words, the interior structure of the cluster is modulated by the structure and mechanics of the interface.

Acknowledgments. This work was generously made possible by the Department of Defense (DoD) National Defense Science and Engineering Graduate (NDSEG) Research Fellowship.

- I. Theurkauff, C. Cottin-Bizonne, J. Palacci, C. Ybert, and L. Bocquet, Dynamic clustering in active colloidal suspensions with chemical signaling, Physical Review Letters 108, 1 (2012), arXiv:1202.6264.
- [2] G. S. Redner, M. F. Hagan, and A. Baskaran, Structure and Dynamics of a Phase-Separating Active Colloidal Fluid, Physical Review Letters 110, 1 (2013).
- [3] M. E. Cates and J. Tailleur, Motility-Induced Phase Separation, Annual Review of Condensed Matter Physics 6, 219 (2015), arXiv:1406.3533.
- [4] W. J. Ridgway, M. P. Dalwadi, P. Pearce, and S. J. Chapman, Motility-Induced Phase Separation Mediated by Bacterial Quorum Sensing, Physical review letters 131, 228302 (2023).
- [5] H. Zhao, A. Košmrlj, and S. S. Datta, Chemotactic Motility-Induced Phase Separation, Physical Review Letters 131, 1 (2023), arXiv:2301.12345.
- [6] L. Caprini, U. Marini Bettolo Marconi, and A. Puglisi, Spontaneous Velocity Alignment in Motility-Induced Phase Separation, Physical Review Letters **124**, 78001 (2020), arXiv:1912.03168.
- [7] M. N. Van Der Linden, L. C. Alexander, D. G. Aarts, and O. Dauchot, Interrupted Motility Induced Phase Separation in Aligning Active Colloids, Physical Review Letters 123, 98001 (2019), arXiv:1902.08094.
- [8] P. Digregorio, D. Levis, A. Suma, L. F. Cugliandolo, G. Gonnella, and I. Pagonabarraga, Full Phase Diagram of Active Brownian Disks: From Melting to Motility-Induced Phase Separation, Physical Review Letters 121, 98003 (2018), arXiv:1805.12484.
- [9] S. Ro, B. Guo, A. Shih, T. V. Phan, R. H. Austin, D. Levine, P. M. Chaikin, and S. Martiniani, Model-Free Measurement of Local Entropy Production and Extractable Work in Active Matter, Physical Review Letters 129, 220601 (2022).
- [10] C. B. Caporusso, P. Digregorio, D. Levis, L. F. Cugliandolo, and G. Gonnella, Motility-Induced Microphase and Macrophase Separation in a Two-Dimensional Active Brownian Particle System, Physical Review Letters 125, 178004 (2020).
- [11] T. Vicsek, A. Czirok, E. Ben-Jacob, I. Cohen, and O. Shochet, Novel Type of Phase Transition in a System of Self-Driven Particles, Physical Review Letters 75, 1226 (1995).
- [12] P. Romanczuk, U. Erdmann, H. Engel, and L. Schimansky-Geier, Beyond the Keller-Segel model, European Physical Journal: Special Topics 157, 61 (2008).
- [13] N. Sarkar, A. Basu, and J. Toner, Swarming Bottom Feeders: Flocking at Solid-Liquid Interfaces, Physical Review Letters 127, 268004 (2021), arXiv:2101.10195.
- [14] B. Zhang and A. Snezhko, Hyperuniform Active Chiral Fluids with Tunable Internal Structure, Physical Review Letters 128, 218002 (2022), arXiv:2205.12384.
- [15] J. Chen, X. Lei, Y. Xiang, M. Duan, X. Peng, and H. P. Zhang, Emergent Chirality and Hyperuniformity in an Active Mixture with Nonreciprocal Interactions, Physical Review Letters 132, 118301 (2024).
- [16] L. Caprini and H. Löwen, Flocking without Alignment Interactions in Attractive Active Brownian Particles, Physical Review Letters 130, 148202 (2023),

arXiv:2303.07746.

- [17] Q. Yu and Y. Tu, Energy Cost for Flocking of Active Spins: The Cusped Dissipation Maximum at the Flocking Transition, Physical Review Letters **129**, 278001 (2022).
- [18] S. Maity and A. Morin, Spontaneous Demixing of Binary Colloidal Flocks, Physical Review Letters 131, 178304 (2023), arXiv:2306.15614.
- [19] B. Benvegnen, O. Granek, S. Ro, R. Yaacoby, H. Chaté, Y. Kafri, D. Mukamel, A. Solon, and J. Tailleur, Metastability of Discrete-Symmetry Flocks, Physical Review Letters 131, 1 (2023), arXiv:2306.01156.
- [20] B. Liebchen and D. Levis, Collective Behavior of Chiral Active Matter: Pattern Formation and Enhanced Flocking, Physical Review Letters 119, 1 (2017).
- [21] A. Tiribocchi, R. Wittkowski, D. Marenduzzo, and M. E. Cates, Active Model H: Scalar Active Matter in a Momentum-Conserving Fluid, Physical Review Letters 115, 1 (2015), arXiv:1504.07447.
- [22] S. Dey, D. Das, and R. Rajesh, Spatial structures and giant number fluctuations in models of active matter, Physical Review Letters 108, 1 (2012).
- [23] R. Kürsten and T. Ihle, Dry Active Matter Exhibits a Self-Organized Cross Sea Phase, Physical Review Letters 125, 188003 (2020), arXiv:2002.03198.
- [24] M. Linkmann, G. Boffetta, M. C. Marchetti, and B. Eckhardt, Phase Transition to Large Scale Coherent Structures in Two-Dimensional Active Matter Turbulence, Physical Review Letters **122**, 214503 (2019).
- [25] C. Dombrowski, L. Cisneros, S. Chatkaew, R. E. Goldstein, and J. O. Kessler, Self-concentration and largescale coherence in bacterial dynamics, Physical Review Letters 93, 2 (2004).
- [26] J. Dunkel, S. Heidenreich, K. Drescher, H. H. Wensink, M. Bär, and R. E. Goldstein, Fluid dynamics of bacterial turbulence, Physical Review Letters **110**, 1 (2013), arXiv:1302.5277.
- [27] J. Gachelin, A. Rousselet, A. Lindner, and E. Clement, Collective motion in an active suspension of Escherichia coli bacteria, New Journal of Physics 16, 10.1088/1367-2630/16/2/025003 (2014).
- [28] Y. Duan, J. Agudo-Canalejo, R. Golestanian, and B. Mahault, Dynamical Pattern Formation without Self-Attraction in Quorum-Sensing Active Matter: The Interplay between Nonreciprocity and Motility, Physical Review Letters 131, 148301 (2023), arXiv:2306.07904.
- [29] B. Chakrabarti, M. J. Shelley, and S. Fürthauer, Collective Motion and Pattern Formation in Phase-Synchronizing Active Fluids, Physical Review Letters 130, 128202 (2023).
- [30] E. H. Colombo, C. López, and E. Hernández-García, Pulsed Interaction Signals as a Route to Biological Pattern Formation, Physical Review Letters 130, 58401 (2023).
- [31] H. Reinken, S. Heidenreich, M. Bär, and S. H. Klapp, Ising-like Critical Behavior of Vortex Lattices in an Active Fluid, Physical Review Letters **128**, 48004 (2022), arXiv:2107.11130.
- [32] Y. Peng, L. Lai, Y. S. Tai, K. Zhang, X. Xu, and X. Cheng, Diffusion of ellipsoids in bacterial suspensions, Physical Review Letters **116**, 1 (2016),

arXiv:1509.05893.

- [33] T. Albers, S. Delnoij, N. Schramma, and M. Jalaal, Billiards with Spatial Memory, Physical Review Letters 132, 157101 (2023), arXiv:2307.01734.
- [34] M. K. Faluweki, J. Cammann, M. G. Mazza, and L. Goehring, Active Spaghetti: Collective Organization in Cyanobacteria, Physical review letters 131, 158303 (2023), arXiv:2301.11667.
- [35] X. Yang, M. L. Manning, and M. C. Marchetti, Aggregation and segregation of confined active particles, Soft Matter 10, 6477 (2014), arXiv:1403.0697.
- [36] S. R. McCandlish, A. Baskaran, and M. F. Hagan, Spontaneous segregation of self-propelled particles with different motilities, Soft Matter 8, 2527 (2012), arXiv:1110.2479.
- [37] N. H. P. Nguyen, D. Klotsa, M. Engel, and S. C. Glotzer, Emergent collective phenomena in a mixture of hard shapes through active rotation, Physical Review Letters 112, 1 (2014).
- [38] S. E. Moran, P. W. Schönhöfer, and S. C. Glotzer, Shape-driven, emergent behavior in active particle mixtures, New Journal of Physics 24, 10.1088/1367-2630/ac7161 (2022).
- [39] J. Bartnick, M. Heinen, A. V. Ivlev, and H. Löwen, Structural correlations in diffusiophoretic colloidal mixtures with nonreciprocal interactions, Journal of Physics Condensed Matter 28, 10.1088/0953-8984/28/2/025102 (2016), arXiv:1510.01870.
- [40] O. Chepizhko and F. Peruani, Diffusion, subdiffusion, and trapping of active particles in heterogeneous media, Physical Review Letters 111, 1 (2013), arXiv:1310.0830.
- [41] J. Agudo-Canalejo and R. Golestanian, Active Phase Separation in Mixtures of Chemically Interacting Particles, Physical Review Letters 123, 18101 (2019), arXiv:1901.09022.
- [42] A. Dinelli, J. O'Byrne, A. Curatolo, Y. Zhao, P. Sollich, and J. Tailleur, Non-reciprocity across scales in active mixtures, Nature Communications 14, 10.1038/s41467-023-42713-5 (2023), arXiv:2203.07757.
- [43] H. Alston, L. Cocconi, and T. Bertrand, Irreversibility across a Nonreciprocal P T -Symmetry-Breaking Phase Transition, Physical Review Letters 131, 258301 (2023), arXiv:2304.08661.
- [44] A. V. Ivlev, J. Bartnick, M. Heinen, C. R. Du, V. Nosenko, and H. Löwen, Statistical mechanics where newton's third law is broken, Physical Review X 5, 1 (2015).
- [45] S. Saha, J. Agudo-Canalejo, and R. Golestanian, Scalar Active Mixtures: The Nonreciprocal Cahn-Hilliard Model, Physical Review X 10, 41009 (2020), arXiv:2005.07101.
- [46] D. Levis and B. Liebchen, Simultaneous phase separation and pattern formation in chiral active mixtures, Physical Review E 100, 1 (2019), arXiv:1901.08496.
- [47] K. Yeo, E. Lushi, and P. M. Vlahovska, Collective dynamics in a binary mixture of hydrodynamically coupled microrotors, Physical Review Letters 114, 1 (2015), arXiv:1410.2878.
- [48] D. Bárdfalvy, S. Anjum, C. Nardini, A. Morozov, and J. Stenhammar, Symmetric Mixtures of Pusher and Puller Microswimmers Behave as Noninteracting Suspensions, Physical Review Letters 125, 18003 (2020), arXiv:2004.06938.
- [49] B. Q. Ai, Z. G. Shao, and W. R. Zhong, Mixing and

demixing of binary mixtures of polar chiral active particles, Soft Matter 14, 4388 (2018), arXiv:1805.12389.

- [50] S. J. Kole, G. P. Alexander, S. Ramaswamy, and A. Maitra, Layered Chiral Active Matter: Beyond Odd Elasticity, Physical Review Letters 126, 248001 (2021).
- [51] P. Arora, A. K. Sood, and R. Ganapathy, Emergent stereoselective interactions and self-recognition in polar chiral active ellipsoids, Science Advances 7, 1 (2021), arXiv:2009.00963.
- [52] J. Stenhammar, R. Wittkowski, D. Marenduzzo, and M. E. Cates, Activity-induced phase separation and selfassembly in mixtures of active and passive particles, Physical Review Letters **114**, 1 (2015), arXiv:1408.5175.
- [53] S. C. Takatori and J. F. Brady, A theory for the phase behavior of mixtures of active particles, Soft Matter 11, 7920 (2015).
- [54] R. Wittkowski, J. Stenhammar, and M. E. Cates, Nonequilibrium dynamics of mixtures of active and passive colloidal particles, New Journal of Physics 19, 10.1088/1367-2630/aa8195 (2017), arXiv:1705.07479.
- [55] P. Dolai, A. Simha, and S. Mishra, Phase separation in binary mixtures of active and passive particles, Soft Matter 14, 6137 (2018), arXiv:1706.02968.
- [56] B. Van Der Meer, V. Prymidis, M. Dijkstra, and L. Filion, Predicting the phase behavior of mixtures of active spherical particles, Journal of Chemical Physics 152, 10.1063/5.0002279 (2020), arXiv:1609.03867.
- [57] A. Wysocki, R. G. Winkler, and G. Gompper, Propagating interfaces in mixtures of active and passive Brownian particles, New Journal of Physics 18, 10.1088/1367-2630/aa529d (2016).
- [58] N. K. Agrawal and P. S. Mahapatra, Alignmentmediated segregation in an active-passive mixture, Physical Review E 104, 1 (2021).
- [59] D. Rogel Rodriguez, F. Alarcon, R. Martinez, J. Ramírez, and C. Valeriani, Phase behaviour and dynamical features of a two-dimensional binary mixture of active/passive spherical particles, Soft Matter 16, 1162 (2020).
- [60] S. Bhattacharyya and J. M. Yeomans, Phase Separation Driven by Active Flows, Physical Review Letters 130, 238201 (2023), arXiv:2302.00153.
- [61] L. Hecht, S. Mandal, H. Löwen, and B. Liebchen, Active Refrigerators Powered by Inertia, Physical Review Letters 129, 178001 (2022), arXiv:2209.07227.
- [62] F. Caballero and M. C. Marchetti, Activity-Suppressed Phase Separation, Physical Review Letters **129**, 268002 (2022), arXiv:2206.12574.
- [63] A. Morozov and D. Marenduzzo, Enhanced diffusion of tracer particles in dilute bacterial suspensions, Soft Matter 10, 2748 (2014).
- [64] D. O. Pushkin and J. M. Yeomans, Fluid mixing by curved trajectories of microswimmers, Physical Review Letters 111, 1 (2013), arXiv:1307.6025.
- [65] Z. Lin, J. L. Thiffeault, and S. Childress, Stirring by squirmers, Journal of Fluid Mechanics 669, 167 (2011), arXiv:1007.1740.
- [66] T. Ishikawa, J. T. Locsei, and T. J. Pedley, Fluid particle diffusion in a semidilute suspension of model microorganisms, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 82, 1 (2010).
- [67] P. T. Underhill, J. P. Hernandez-Ortiz, and M. D. Graham, Diffusion and spatial correlations in suspensions of swimming particles, Physical Review Letters 100, 1

(2008), arXiv:0805.3784.

- [68] Z. Preisler and M. Dijkstra, Configurational entropy and effective temperature in systems of active Brownian particles, Soft Matter 12, 6043 (2016).
- [69] O. Granek, Y. Kafri, and J. Tailleur, Anomalous Transport of Tracers in Active Baths, Physical Review Letters 129, 38001 (2022), arXiv:2108.11970.
- [70] Y. Wang, Z. Shen, Y. Xia, G. Feng, and W. Tian, Phase separation and super diffusion of binary mixtures of active and passive particles, Chinese Physics B 29, 10.1088/1674-1056/ab81f4 (2020).
- [71] L. Angelani, C. Maggi, M. L. Bernardini, A. Rizzo, and R. Di Leonardo, Effective interactions between colloidal particles suspended in a bath of swimming cells, Physical Review Letters 107, 1 (2011).
- [72] A. Maitra and R. Voituriez, Enhanced Orientational Ordering Induced by an Active yet Isotropic Bath, Physical Review Letters 124, 48003 (2020), arXiv:1909.12014.
- [73] L. Angelani, R. Di Leonardo, and G. Ruocco, Selfstarting micromotors in a bacterial bath, Physical Review Letters 102, 1 (2009), arXiv:0812.2375.
- [74] P. Liu, S. Ye, F. Ye, K. Chen, and M. Yang, Constraint Dependence of Active Depletion Forces on Passive Particles, Physical Review Letters 124, 158001 (2020).
- [75] P. Malgaretti and H. Stark, Szilard Engines and Information-Based Work Extraction for Active Systems, Physical Review Letters **129**, 228005 (2022), arXiv:2203.13075.
- [76] T. K. Saha, J. Ehrich, M. Gavrilov, S. Still, D. A. Sivak, and J. Bechhoefer, Information Engine in a Nonequilibrium Bath, Physical Review Letters 131, 1 (2023), arXiv:2208.00288.
- [77] C. Maes, Fluctuating Motion in an Active Environment, Physical Review Letters 125, 208001 (2020), arXiv:2005.13462.
- M. J. Huang, J. Schofield, and R. Kapral, Transport in active systems crowded by obstacles, Journal of Physics A: Mathematical and Theoretical 50, 10.1088/1751-8121/50/7/074001 (2017).
- [79] S. Nayak, S. Das, P. Bag, T. Debnath, and P. K. Ghosh, Driven transport of active particles through arrays of symmetric obstacles, Journal of Chemical Physics 159, 10.1063/5.0176523 (2023), arXiv:2310.09272.
- [80] A. Duzgun and J. V. Selinger, Active Brownian particles near straight or curved walls: Pressure and boundary layers, Physical Review E 97, 32606 (2018), arXiv:1712.09627.
- [81] S. Das and R. Chelakkot, Morphological transitions of active Brownian particle aggregates on porous walls, Soft Matter 16, 7250 (2020), arXiv:2004.11436.
- [82] P. Forgács, A. Libál, C. Reichhardt, and C. J. Reichhardt, Active matter shepherding and clustering in inhomogeneous environments, Physical Review E 104, 1 (2021), arXiv:2107.12508.
- [83] N. Nikola, A. P. Solon, Y. Kafri, M. Kardar, J. Tailleur, and R. Voituriez, Active Particles with Soft and Curved Walls: Equation of State, Ratchets, and Instabilities, Physical Review Letters 10.1103/Phys-RevLett.117.098001 (2016), arXiv:1512.05697.
- [84] J. Codina, B. Mahault, H. Chaté, J. Dobnikar, I. Pagonabarraga, and X. Q. Shi, Small Obstacle in a Large Polar Flock, Physical Review Letters **128**, 1 (2022), arXiv:2112.08410.
- [85] C. D. Schimming, C. J. Reichhardt, and C. Reichhardt,

Vortex Lattices in Active Nematics with Periodic Obstacle Arrays, Physical Review Letters **132**, 18301 (2024), arXiv:2309.07886.

- [86] L. R. Leite, D. Lucena, F. Q. Potiguar, and W. P. Ferreira, Depletion forces on circular and elliptical obstacles induced by active matter, Physical Review E 94, 1 (2016), arXiv:1605.05873.
- [87] A. D. Borba, J. L. Domingos, E. C. Moraes, F. Q. Potiguar, and W. P. Ferreira, Controlling the transport of active matter in disordered lattices of asymmetrical obstacles, Physical Review E 101, 1 (2020), arXiv:1908.05942.
- [88] D. M. van Roon, G. Volpe, M. M. Telo da Gama, and N. A. Araújo, The role of disorder in the motion of chiral active particles in the presence of obstacles, Soft Matter 18, 6899 (2022).
- [89] R. Martinez, F. Alarcon, J. L. Aragones, and C. Valeriani, Trapping flocking particles with asymmetric obstacles, Soft Matter 16, 4739 (2020), arXiv:1911.13064.
- [90] S. E. Spagnolie, G. R. Moreno-Flores, D. Bartolo, and E. Lauga, Geometric capture and escape of a microswimmer colliding with an obstacle, Soft Matter 11, 3396 (2015), arXiv:1412.4598.
- [91] R. Martinez, F. Alarcon, D. R. Rodriguez, J. L. Aragones, and C. Valeriani, Collective behavior of Vicsek particles without and with obstacles, European Physical Journal E 41, 10.1140/epje/i2018-11706-8 (2018), arXiv:1911.10374.
- [92] T. Kolb, T. Kolb, and D. Klotsa, Active binary mixtures of fast and slow hard spheres, Soft Matter 16, 1967 (2020), arXiv:1909.02919.
- [93] P. De Castro, F. M. Rocha, S. Diles, R. Soto, and P. Sollich, Diversity of self-propulsion speeds reduces motilityinduced clustering in confined active matter, Soft Matter 17, 9926 (2021).
- [94] P. de Castro, S. Diles, R. Soto, and P. Sollich, Active mixtures in a narrow channel: motility diversity changes cluster sizes, Soft Matter 17, 2050 (2021).
- [95] M. Rojas-Vega, P. de Castro, and R. Soto, Mixtures of self-propelled particles interacting with asymmetric obstacles, European Physical Journal E 46, 1 (2023).
- [96] D. Debnath, P. K. Ghosh, V. R. Misko, Y. Li, F. Marchesoni, and F. Nori, Enhanced motility in a binary mixture of active nano/microswimmers, Nanoscale 12, 9717 (2020), arXiv:2004.11584.
- [97] M. Rojas-Vega, P. De Castro, and R. Soto, Wetting dynamics by mixtures of fast and slow selfpropelled particles, Physical Review E 107, 14608 (2023), arXiv:2301.01856.
- [98] L. G. Wilson, V. A. Martinez, J. Schwarz-Linek, J. Tailleur, G. Bryant, P. N. Pusey, and W. C. Poon, Differential dynamic microscopy of bacterial motility, Physical Review Letters **106**, 7 (2011), arXiv:1004.4764.
- [99] D. T. Chen, A. W. Lau, L. A. Hough, M. F. Islam, M. Goulian, T. C. Lubensky, and A. G. Yodh, Fluctuations and rheology in active bacterial suspensions, Physical Review Letters **99**, 1 (2007), arXiv:0709.1465.
- [100] X. L. Wu and A. Libchaber, Particle diffusion in a quasitwo-dimensional bacterial bath, Physical Review Letters 84, 3017 (2000).
- [101] G. L. Miño, J. Dunstan, A. Rousselet, E. Clément, and R. Soto, Induced diffusion of tracers in a bacterial suspension: Theory and experiments, Journal of Fluid Mechanics **729**, 423 (2013), arXiv:1210.7704.

- [102] A. Jepson, V. A. Martinez, J. Schwarz-Linek, A. Morozov, and W. C. Poon, Enhanced diffusion of nonswimmers in a three-dimensional bath of motile bacteria, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 88, 3 (2013), arXiv:1307.1274.
- [103] C. Valeriani, M. Li, J. Novosel, J. Arlt, and D. Marenduzzo, Colloids in a bacterial bath: Simulations and experiments, Soft Matter 7, 5228 (2011).
- [104] K. C. Leptos, J. S. Guasto, J. P. Gollub, A. I. Pesci, and R. E. Goldstein, Dynamics of Enhanced Tracer Diffusion in Suspensions of Swimming Eukaryotic Microorganisms, Physical Review Letters 103, 1 (2009).
- [105] H. Kurtuldu, J. S. Guasto, K. A. Johnson, and J. P. Gollub, Enhancement of biomixing by swimming algal cells in two-dimensional films, Proceedings of the National Academy of Sciences of the United States of America 108, 10391 (2011).
- [106] T. V. Kasyap, D. L. Koch, and M. Wu, Hydrodynamic tracer diffusion in suspensions of swimming bacteria, Physics of Fluids 26, 10.1063/1.4891570 (2014).
- [107] L. Restrepo-Pérez, L. Soler, C. S. Martínez-Cisneros, S. Sánchez, and O. G. Schmidt, Trapping self-propelled micromotors with microfabricated chevron and heartshaped chips, Lab on a Chip 14, 1515 (2014).
- [108] G. Miño, T. E. Mallouk, T. Darnige, M. Hoyos, J. Dauchet, J. Dunstan, R. Soto, Y. Wang, A. Rousselet, and E. Clement, Enhanced diffusion due to active swimmers at a solid surface, Physical Review Letters 106, 1 (2011), arXiv:1012.4624.
- [109] Y. Baek, A. P. Solon, X. Xu, N. Nikola, and Y. Kafri, Generic Long-Range Interactions between Passive Bodies in an Active Fluid, Physical Review Letters 120, 58002 (2018), arXiv:1709.02281.
- [110] A. Gough, A. Stern, J. Maier, T. Lezon, T.-Y. Shun, C. Chennubhotla, M. Haney, S. Schurdak, and D. Taylor, Biologically Relevant Heterogeneity: Metrics and Practical Insights, 2015 IEEE Summer Topicals Meeting Series, SUM 2015 10, 1 (2018), arXiv:15334406.
- [111] D. J. Knight and K. J. Girling, Gut flora in health and disease [9], Lancet 361, 1831 (2003).
- [112] K. L. Heck and T. A. Thoman, Experiments on predator-prey interactions in vegetated aquatic habitats, Journal of Experimental Marine Biology and Ecology 53, 125 (1981).
- [113] J. W. Jolles, N. J. Boogert, V. H. Sridhar, I. D. Couzin, and A. Manica, Consistent Individual Differences Drive Collective Behavior and Group Functioning of Schooling Fish, Current Biology 27, 2862 (2017).
- [114] J. W. Jolles, N. Weimar, T. Landgraf, P. Romanczuk, J. Krause, and D. Bierbach, Group-level patterns emerge from individual speed as revealed by an extremely social robotic fish, Biology Letters 16, 10.1098/rsbl.2020.0436 (2020).
- [115] P. P. Klamser, L. Gómez-Nava, T. Landgraf, J. W. Jolles, D. Bierbach, and P. Romanczuk, Impact of Variable Speed on Collective Movement of Animal Groups, Frontiers in Physics 9, 1 (2021), arXiv:2106.00959.
- [116] B. Christensen, Predator Foraging Capabilities and Prey Antipredator Behaviours : Pre- versus Postcapture Constraints on Size-Dependent Predator-Prey Interactions Author (s): Bent Christensen Published by : Wiley on behalf of Nordic Society Oikos Stable URL

: https://w, Oikos 76, 368 (1996).

- [117] M. Forget, S. Adiba, L. G. Brunnet, and S. De Monte, Heterogeneous individual motility biases group composition in a model of aggregating cells, Frontiers in Ecology and Evolution 10, 10.3389/fevo.2022.1052309 (2022).
- [118] E. R. Jerison and S. R. Quake, Heterogeneous T cell motility behaviors emerge from a coupling between speed and turning in vivo, eLife 9, 1 (2020).
- [119] J. Palacci, C. Cottin-Bizonne, C. Ybert, and L. Bocquet, Sedimentation and effective temperature of active colloidal suspensions, Physical Review Letters 105, 1 (2010).
- [120] J. Palacci, S. Sacanna, A. P. Steinberg, D. J. Pine, and P. M. Chaikin, Living Crystals of Light-Activated Colloidal Surfers, Science **339**, 936 (2013).
- [121] S. Ebbens, R. A. Jones, A. J. Ryan, R. Golestanian, and J. R. Howse, Self-assembled autonomous runners and tumblers, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 82, 6 (2010).
- [122] T. C. Lee, M. Alarcón-Correa, C. Miksch, K. Hahn, J. G. Gibbs, and P. Fischer, Self-propelling nanomotors in the presence of strong Brownian forces, Nano Letters 14, 2407 (2014).
- [123] J. A. Anderson, J. Glaser, and S. C. Glotzer, HOOMD-blue: A Python package for highperformance molecular dynamics and hard particle Monte Carlo simulations, Computational Materials Science 10.1016/j.commatsci.2019.109363 (2020), arXiv:1308.5587.
- [124] J. Glaser, T. D. Nguyen, J. A. Anderson, P. Lui, F. Spiga, J. A. Millan, D. C. Morse, and S. C. Glotzer, Strong scaling of general-purpose molecular dynamics simulations on GPUs, Computer Physics Communications **192**, 97 (2015), arXiv:1412.3387.
- [125] J. A. Anderson, C. D. Lorenz, and A. Travesset, General purpose molecular dynamics simulations fully implemented on graphics processing units, Journal of Computational Physics 227, 5342 (2008).
- [126] N. Lauersdorf, T. Kolb, M. Moradi, E. Nazockdast, and D. Klotsa, Phase behavior and surface tension of soft active Brownian particles, Soft Matter 17, 6337 (2021).
- [127] A. K. Omar, Y. Wu, Z. G. Wang, and J. F. Brady, Swimming to Stability: Structural and Dynamical Control via Active Doping, ACS Nano 13, 560 (2019), arXiv:1902.02423.
- [128] G. S. Redner, A. Baskaran, and M. F. Hagan, Reentrant phase behavior in active colloids with attraction, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 88, 1 (2013).
- [129] R. Soto and R. Golestanian, Run-and-tumble dynamics in a crowded environment: Persistent exclusion process for swimmers, Physical Review E - Statistical, Nonlinear, and Soft Matter Physics 89, 1 (2014), arXiv:1306.0481.
- [130] C. J. Reichhardt and C. Reichhardt, Avalanche dynamics for active matter in heterogeneous media, New Journal of Physics 20, 10.1088/1367-2630/aaa392 (2018), arXiv:1709.06121.
- [131] C. Reichhardt and C. J. Reichhardt, Clogging and depinning of ballistic active matter systems in disordered media, Physical Review E 97, 1 (2018), arXiv:1803.08992.