APPROXIMATE UNITARY k-DESIGNS FROM SHALLOW, LOW-COMMUNICATION CIRCUITS

NICHOLAS LARACUENTE AND FELIX LEDITZKY

ABSTRACT. Random unitaries are useful in quantum information and related fields but hard to generate with limited resources. An approximate unitary k-design is an ensemble of unitaries and measure over which the average is close to a Haar (uniformly) random ensemble up to the first k moments. A particularly strong notion of approximation bounds the distance from Haar randomness in relative error: the approximate design can be written as a convex combination involving an exact design and vice versa. We construct multiplicative-error approximate unitary k-design ensembles for which communication between subsystems is O(1) in the system size. These constructions use the alternating projection method to analyze overlapping Haar twirls, giving a bound on the convergence speed to the full twirl with respect to the 2-norm. Using the von Neumann subalgebra indices to replace system dimension, the 2-norm distance converts to relative error without introducing any additional dependence on system size. Via recursion on these constructions, we construct a scheme yielding relative error designs in $O((k \log k +$ $\log m + \log(1/\epsilon) k$ polylog(k)) depth, where m is the dimension of the complete system and ϵ the approximation error. This sublinear depth construction answers one variant of [Harrow and Mehraban 2023, Open Problem 1]. Moreover, entanglement generated by the sublinear depth scheme follows area laws on spatial lattices up to corrections logarithmic in the full system size.

1. Introduction

Uniformly random unitary and state ensembles are useful in quantum information [12, 23, 29] and fundamental physics [1, 37]. Such unitaries are however difficult to construct via quantum circuits [35]. More accessible constructions approximate the properties of random unitaries [5, 10, 33]. The unitary k-design is one such notion: a measure over the unitary group that is indistinguishable from Haar random in the first k statistical moments. Equivalently, any protocol that samples a unitary from a k-design k or fewer times would be unable to distinguish it from the Haar ensemble by any test using k or fewer copies with probability better than randomly guessing.

As exact designs are still challenging to construct, approximate designs often take their place. There are several notions of approximate designs. A particularly strong notion considered in [3] and [7] posits that an ensemble is an approximate design in multiplicative or relative error if the average over that ensemble can be written using a convex combination including the Haar average, and the Haar average in turn can be written using a convex combination including the considered ensemble. Multiplicative error designs are secure against queries by an adaptive adversary trying to distinguish it from Haar ensembles [6]. This relative notion implies a weaker, additive notion requiring that ensemble averages over k-fold applications of unitaries from the

design and Haar ensemble are close in diamond norm distance. Designs in diamond norm distance are secure against non-adaptive adversaries, and the parameter k is a lower-bound on the circuit complexity required for typical unitaries in such a design [4]. Moreover, a weaker but related notion replaces the diamond norm distance by the two-norm distance in what is called a tensor product expander (TPE). These notions are briefly reviewed in Section 2.1.

Many previous works have studied how random circuits converge to k-designs in bounded depth [21, 3, 26, 22, 19]. Recent results have yielded efficient constructions in required circuit depth on 1-dimensional connectivity [18, 30, 6]. Even more recently, it was shown that random 2-local circuits on one-dimensional connectivity yield ϵ -approximate relative error k-designs on m-qudit systems in depth $O((mk + \log(1/\epsilon))\operatorname{polylog}(k))$, and random local circuits on all-to-all connectivity do so with gate count $O(m(mk + \log(1/\epsilon))\operatorname{polylog}(k))$ [7].

Nevertheless, many questions have remained open (see [22, Section 1.5] and [7, Section 1.4]). In particular, we highlight [22, Section 1.5, Open Question 1], which asks whether one can construct a multiplicative error k-design on a system of m qubits using random circuits with depth sublinear in m. Our results imply that this is possible, even in one-dimensional spatial lattice connectivity, if the circuits are allowed to follow a layer-dependent architecture as detailed in Section 5.2.

We initially approach the problem by asking a different question: how much quantum communication do two parties need to jointly construct an approximate unitary k-design ensemble? Some inspiration comes from pseudoentangled state ensembles [2]. Based on widely believed complexity-theoretic assumptions, these ensembles are hard to distinguish from uniformly random state ensembles for any polynomial-time quantum algorithm. Nonetheless, the entanglement across any cut of a pseudoentangled state ensemble need only scale slightly faster than logarithmic in the subsystem size. As noted in [2, p.1], earlier results rule out an analog of pseudoentanglement for exact or highly precise unitary k-design ensembles, as statistically approximating the Haar measure to exponential precision requires near-maximal entanglement [28, 9]. Nonetheless, these barrier results left open whether smaller amounts of entanglement could suffice for an ϵ -approximate k-design with constant ϵ in the system size.

Summary of Main Results. We show that for k not too large, it is indeed possible to construct unitary k-designs with relatively little quantum communication between subsystems, hence generating little entanglement. Specifically:

- (1) **Twirl-Swap-Twirl:** Let A and B be systems of the same size. Consider a protocol that locally applies k-designs to $A^{\otimes k}$ and $B^{\otimes k}$ respectively, then exchanges the same ℓ qudits between each copy of A and B respectively, then again applies local k-designs. Such a protocol achieves an ϵ -approximate relative k-design when $\ell = O(k \log k + \log(1/\epsilon))$. In particular, this bound is *independent* of the sizes of A and B as long as each contains at least $\omega(k \log k + \log(1/\epsilon))$ qudits. The full technical version of this result with explicit constants is stated in Theorem 4.3.
- (2) **Twirl-Crosstwirl:** Let A_1, \ldots, A_P be subsystems of a multipartite system A. Consider the following protocol: (1) locally apply a k-design to each A_p for $p = 1, \ldots, P$; (2) apply

a "crosstwirl" design across a joint system combining ℓ qudits from each A_p . Assume each A_p is sufficiently large. Then this protocol achieves an ϵ -approximate relative k-design again when

$$\ell \ge 2Pk\log_a k + 2\log_a k + \log_a P + \log_a(1/\epsilon) + \log_a 5. \tag{1.1}$$

This result is stated in Theorem 4.8. Shown as Corollary 4.7, the protocol yields an ϵ -approximate k-TPE when

$$\ell \ge 2\log_a k + \log_a P + \log_a(1/\epsilon) + \log_a(5/2)$$
 (1.2)

Compared with the $k \log k$ dependence for designs, the k-TPE is achieved with logarithmic k dependence.

(3) **Recursive Crosstwirl:** Consider an m-qudit system with connectivity given by a lattice in spatial dimension D. We give a construction of an ϵ -approximate relative k-design using unitaries of circuit depth

$$O(k(k \log k + \log m + \log(1/\epsilon))\operatorname{polylog}(k))$$
. (1.3)

Moreover, for any spatially contiguous region S, the design requires only

$$O(\#(\partial S) \times (k \log k + \log m + \log(1/\epsilon))^{1/D} + k \log k + \log m + \log(1/\epsilon))$$
(1.4)

qudits of quantum communication and generates only that much entanglement on any product state input, where $\#(\partial S)$ is the number of qudits on the boundary of S. The technical version is stated as Theorem 5.8, and in Section 5.2 the explicit architecture is constructed. This result addresses [22, Section 1.5, Open Question 1], showing that with a specific, layered architecture, random circuits produce relative error k-designs in sublinear depth.

Discussion. The Twirl-Swap-Twirl and Twirl-Crosstwirl results are primarily about the entanglement or quantum communication needed to produce a design. One may replace the local k-designs in Twirl-Swap-Twirl by δ -approximate relative k-designs to achieve error $(1+\epsilon)(1+\delta)^4-1$. Similarly, one may replace the local and crossing k-designs in Twirl-Crosstwirl by δ -approximate relative k-designs, achieving error $(1+\epsilon)(1+\delta)^{P+1}-1$. Via the results of [7], if A and B are each m-qudit systems, this means these two schemes are approximately implementable in local circuit depth $O((mk + \log(1/\epsilon))\operatorname{polylog}(k))$ on a one-dimensional line of qudits. The main technical methods in both these results are 1) a new subspace angle approach using Schur-Weyl duality to show efficient TPE bounds, and 2) a more efficient conversion from TPE to relative error that replaces system dimension by von Neumann subalgebra indices.

Recursive Crosstwirl is more directly analogous to the "across any cut" notion of pseudoentanglement as in [2]. In this case, up to a logarithmic dependence in the total system size m, the entanglement scales like an area law. Area-law versus volume-law entanglement is an important concept in condensed matter and high-energy physics [13]. Moreover, Recursive Crosstwirl achieves a new record for scaling of circuit depth in the system size. Though Recursive Crosstwirl is not implemented in lattice brickwork, it can be implemented using 2-qudit Haar random

unitaries. In Remark 5.9, we discuss how Recursive Crosstwirl can be implemented on an architecture that closely resembles brickwork in one dimension.

In general, these quantum communication and entanglement bounds are primarily intended for the regime in which the dimension of each system is larger than k!. A single iteration only works if subsystems support ℓ -qudit exchanges, placing a lower bound on dimension. If k! is too large for the subsystems, then it is still possible to obtain designs by repeating our constructions. For large k, however, the quantum communication required is no longer smaller than that of general random circuits, and the entanglement entropy resembles a volume law. Nonetheless, that Recursive Crosstwirl achieves sublinear depth in m may suggest further investigation of potential efficiencies of recursive constructions.

Open questions. The convergence in the construction of both the Twirl-Swap-Twirl and the Twirl-Crosstwirl protocol is determined by the operator norm of certain matrices $\mathbf{X} = a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}$ and $\mathbf{Y} = cg\mathbb{1}_{k!^2} \odot \mathbf{N}_\ell^T\mathbf{N}_\ell - b\mathbf{N}^T\mathbf{N}$, respectively. Here, a,b,c,g are constants defined in (4.58), and the matrices $\mathbf{M}, \mathbf{N}, \mathbf{N}_\ell$ are defined in (4.23), (4.24) and (4.62), respectively. The elements of these matrices are functions of permutation operators. While the operator norm of \mathbf{Y} can be controlled quite transparently using the generalized Perron-Frobenius Theorem, the different structure of the matrix \mathbf{X} defies this approach for Twirl-Swap-Twirl, requiring a coarser analysis that leads to a weaker bound on the convergence of the design construction in this case. A more refined analysis of $\|\mathbf{X}\|$ may lead to an improved bound on this convergence mirroring that for the Twirl-Crosstwirl protocol. Another remaining question is whether standard brickwork random circuits in one dimension or any dimension produce relative error designs in sublinear depth.

Structure of this paper. The remainder of this paper is structured as follows. In Section 2 we give some background on approximate k-designs, von Neumann algebras and index theory, and Schur-Weyl duality. In Section 3 we discuss how tensor product expander (TPE) bounds in the $(2 \to 2)$ -norm can be converted into efficient relative error bounds using the algebraic structure of von Neumann subalgebras. In Section 4 we prove TPE bounds for the Twirl-Swap-Twirl and Twirl-Crosstwirl protocols mentioned above using the alternating projection method. In Section 5 we discuss recursive design constructions, in particular focusing on spatial lattice geometries to construct our the Recursive Crosstwirl protocol. Finally, Appendix A contains the proof of the multipartite Twirl-Crosstwirl protocol.

Acknowledgments. We acknowledge helpful email exchanges with Aram Harrow during the development of these results. FL acknowledges support by National Science Foundation grant no. 2137953. Part of this work was completed during the workshop "Beyond i.i.d. in Information Theory 11" hosted by the University of Tübingen from July 31 to August 4, 2023.

Note added: We thank Thomas Schuster, Jonas Haferkamp, and Hsin-Yuan Huang for sharing results from their concurrent and independent work on low-depth designs and pseudorandom unitaries [38]. Their work also obtains approximate design ensembles via $O(\log m)$ circuit depth and improves the communication scaling in k to $O(\log k)$ via a different technical analysis. Our

work uses a different extension to higher-dimensional lattices and tree-like geometries for a closer approximation to area law entanglement (see Remark 5.13).

2. Background

A (finite-dimensional) quantum channel is a completely positive map on a space of matrices. Following the physics convention, we apply quantum channels in the Schrödinger picture, such that a channel Φ is considered to act on states, and its adjoint on observables. We call a quantum channel trace-symmetric if it is its own adjoint under the trace, $\operatorname{tr}(\Phi(x)y) = \operatorname{tr}(x\Phi(y))$.

The Schatten p-norm of a matrix ρ is defined as $\|\rho\|_p = \operatorname{tr}(|\rho|^p)^{1/p}$ with $|\rho| := \sqrt{\rho^{\dagger}\rho}$. We recall the $p \to q$ distance

$$\|\Phi - \Psi\|_{p \to q} = \sup_{\rho \neq 0} \frac{\|(\Phi - \Psi)(\rho)\|_q}{\|\rho\|_p}$$
 (2.1)

and its completely bounded strengthening,

$$\|\Phi - \Psi\|_{p \to p, \text{cb}} = \sup_{d \in \mathbb{N}} \|(\Phi - \Psi) \otimes \operatorname{Id}_{(d)}\|_{p \to p}, \qquad (2.2)$$

where $\mathrm{Id}_{(d)}$ is the identity channel in dimension d. The diamond norm is defined as $\|\cdot\|_{\Diamond} := \|\cdot\|_{1\to 1,\mathrm{cb}}$. Furthermore, we recall the complete semidefinite ordering on a pair of channels Φ, Ψ , in which $\Phi \prec \Psi$ iff $\Psi \otimes \mathrm{Id}_{(d)} - \Phi \otimes \mathrm{Id}_{(d)}$ is a completely positive map (quantum channel). The notation \succ is defined analogously.

2.1. **Designs.** We briefly recall notions of approximate k-designs and related concepts. For a unitary measure μ on U(d), let

$$\Phi_{\mu,k}(\rho) := \int U^{\otimes k} \rho U^{\otimes k\dagger} d\mu \tag{2.3}$$

for every input state ρ on a system of dimension d. Let $\Phi_{\text{Haar},k}$ denote such a construction with respect to the Haar measure on U(d). A measure μ on U(d) is an ϵ -approximate...

• ...tensor product expander (TPE) if

$$\|\Phi_{\mu,k} - \Phi_{\text{Haar},k}\|_{2\to 2} < \epsilon . \tag{2.4}$$

 \bullet ...additive k-design if

$$\|\Phi_{\mu,k} - \Phi_{\text{Haar},k}\|_{\Diamond} \le \epsilon . \tag{2.5}$$

 \bullet ...multiplicative or relative error k-design if

$$(1 - \epsilon)\Phi_{\text{Haar},k} \prec \Phi_{\mu,k} \prec (1 + \epsilon)\Phi_{\text{Haar},k} . \tag{2.6}$$

Multiplicative error is a stronger criterion than and implies additive error, which in turn applies the TPE condition. As a more general criterion that we may use in intermediate results, we say that a measure is a design with multiplicative error (ϵ, δ) if

$$(1 - \epsilon)\Phi_{\text{Haar}\,k} \prec \Phi_{\mu\,k} \prec (1 + \delta)\Phi_{\text{Haar}\,k} \,. \tag{2.7}$$

For a k-design μ , we refer to $\Phi_{\mu,k}$ as the corresponding k-design channel. For a pair of measures μ and ν , we denote by $\mu * \nu$ the convolution, defined as the measure for which $\Phi_{\mu*\nu} = \Phi_{\mu,k} \circ \Phi_{\nu,k}$ for all $k \in \mathbb{N}$.

2.2. Von Neumann Algebras and Entropy. Particularly in Section 3, some proofs involve finite-dimensional von Neumann algebra index theory. By $\mathbb{B}(\mathcal{H})$ we denote the bounded operators on Hilbert space \mathcal{H} . Within finite dimension, this is equivalent to the space \mathbb{M}_d of matrices acting on d-dimensional Hilbert space. A von Neumann subalgebra $\mathcal{N} \subseteq \mathbb{M}_d$ of the finite-dimensional matrix algebra \mathbb{M}_d consists of all matrices with the block diagonal form

$$\mathcal{N} = \bigoplus_{\lambda} \mathcal{A}_{\lambda} \otimes \mathbb{1}_{B_{\lambda}} , \qquad (2.8)$$

where each \mathcal{A}_{λ} is itself a matrix subalgebra on subsystem A_{λ} , and each and B_{λ} corresponds to subsystem on which an effective Haar unitary average has been performed [34, 16]. The notation " $\mathbb{1}_{B_{\lambda}}$ " distinguishes that this is the algebra of multiples of the identity matrix, also sometimes denoted \mathbb{C} , as it is equivalent to the algebra of complex scalars. The dimension of such a mixed system is often referred to as a multiplicity, since the block A_{λ} is repeated a number of times equal to the dimension of $\mathbb{1}_{B_{\lambda}}$, and then B_{λ} is called a multiplicity space. By \mathcal{M}_{*} we denote the predual of the von Neumann algebra \mathcal{M} . Although within finite dimension $\mathcal{M}_{*} \cong \mathcal{M}$, one conventionally thinks of states as being elements of the predual and observables being elements of the algebra. By $|\mathcal{M}|$ we denote the of \mathcal{M} the vector space on which $|\mathcal{M}|$ acts. Note that the usable, effective dimension of \mathcal{M} is generally smaller, as multiplicities contribute to the space's dimension without contributing to the amount of information that could potentially be stored in a system represented by \mathcal{M} .

With each von Neumann algebra \mathcal{N} comes a unique, unital, trace-preserving conditional expectation $\mathcal{E}_{\mathcal{N}}$. The action of $\mathcal{E}_{\mathcal{N}}$ on a matrix can be broken into two steps. First, remove off-diagonal blocks, taking $\mathbb{M}_d \to \bigoplus_{\lambda} \mathcal{A}_{\lambda} \otimes B_{\lambda}$. Second, perform a complete mixing of each B_{λ} system. A subalgebra $\mathcal{N} \subseteq \mathcal{M}$ is a subset of \mathcal{N} that is also a von Neumann algebra. When $\mathcal{N} \subseteq \mathcal{M}$, it follow that $\mathcal{E}_{\mathcal{M}} \mathcal{E}_{\mathcal{N}} = \mathcal{E}_{\mathcal{N}} \mathcal{E}_{\mathcal{M}} = \mathcal{E}_{\mathcal{N}}$.

As simple examples, the completely depolarizing channel is a conditional expectation to the subalgebra denoted $\mathbb{1}$ or \mathbb{C} . We may for instance take a bipartite system $A \otimes B$, then write $\mathcal{E}_{A \otimes \mathbb{C}}$ as the conditional expectation that completely depolarizes the B system. Another somewhat complementary example is the pinching to a basis: this removes all of the off-diagonal terms (with respect to that basis) of its input matrix. A more sophisticated example appears in the following Section 2.3.

The complete Pimsner-Popa index [36, 16] of a von Neumann algebra inclusion $\mathcal{N} \subseteq \mathcal{M}$ is defined as

$$C(\mathcal{M}: \mathcal{N}) := \inf\{c \in \mathbb{R} \mid c\mathcal{E}_{\mathcal{N}}(\rho) \le \rho \,\forall \rho \in \mathcal{M}_*\}$$
 (2.9)

$$C_{\rm cb}(\mathcal{M}:\mathcal{N}) := \inf\{c \in \mathbb{R} \mid c\mathcal{E}_{\mathcal{N}} \prec \mathcal{E}_{\mathcal{M}}\}$$
(2.10)

$$= \sup_{d \in \mathbb{N}} C(\mathcal{M} \otimes \mathbb{M}_d : \mathcal{N} \otimes \mathbb{M}_d) . \tag{2.11}$$

The quantum relative entropy [39] is defined by

$$D(\rho \| \sigma) := \operatorname{tr}(\rho(\log \rho - \log \sigma)) \tag{2.12}$$

and is taken to be infinite if the support of ρ is not contained in that of σ . Conventions vary as to what base the logarithm is taken with, but for inequalities that only compare entropies to other entropies, the base does not matter. In the rest of this section, we will assume that the base of any logarithms for which is is unspecified is equal to that of the relative entropies in those equations. The more familiar von Neumann entropy is defined as $H(\rho) = -\operatorname{tr}(\rho \log \rho) = -D(\rho || 1)$. An identity known as the chain rule of relative entropy (see, e.g., [27]) states that for von Neumann algebras $\mathcal{N} \subseteq \mathcal{M} \subseteq \mathcal{L}$,

$$D(\mathcal{E}_{\mathcal{M}}(\rho)||\mathcal{E}_{\mathcal{L}}(\rho)) = D(\mathcal{E}_{\mathcal{M}}(\rho)||\mathcal{E}_{\mathcal{N}}(\rho)) + D(\mathcal{E}_{\mathcal{N}}(\rho)||\mathcal{E}_{\mathcal{L}}(\rho)). \tag{2.13}$$

Furthermore, for any von Neumann algebras $\mathcal{M} \subseteq \mathcal{N}$,

$$D(\mathcal{E}_{\mathcal{M}}(\rho)||\mathcal{E}_{\mathcal{N}}(\rho)) = H(\mathcal{E}_{\mathcal{N}}(\rho)) - H(\mathcal{E}_{\mathcal{M}}(\rho)). \tag{2.14}$$

A more recent result [16, Theorem 3.1] relating the index to relative entropy states that for every $\alpha \in (1/2, \infty)$,

$$\log C(\mathcal{M}: \mathcal{N}) = \max_{\rho \in \mathcal{M}_*} D_{\alpha}(\rho \| \mathcal{E}_{\mathcal{N}}(\rho)) = \max_{\rho \in \mathcal{M}_*} \min_{\sigma \in \mathcal{N}_*} D(\rho \| \sigma)$$
(2.15)

for any $\alpha \in [1/2, \infty]$, where D_{α} denotes the sandwiched Rényi divergence with $\lim_{\alpha \to 1} D_{\alpha}(\cdot \| \cdot) = D(\cdot \| \cdot)$ [40, 32]. We will use a special case of this result. Note that taking the supremum over states in \mathcal{M}_{*} is equivalent to taking the supremum over states in \mathbb{M}_{d} , then applying $\mathcal{E}_{\mathcal{M}}$ to the first input. Therefore,

$$\log C_{\rm cb}(\mathcal{M}:\mathcal{N}) = \sup_{d \in \mathbb{N}} \max_{\rho \in \mathcal{M}_* \otimes \mathbb{M}_d} D((\mathcal{E}_{\mathcal{M}} \otimes \operatorname{Id}_{(d)})(\rho) \| (\mathcal{E}_{\mathcal{N}} \otimes \operatorname{Id}_{(d)})(\rho)) . \tag{2.16}$$

2.3. Schur-Weyl Duality. For a (finite-dimensional) Hilbert space \mathcal{H} we denote by $\mathcal{L}(\mathcal{H})$ or $\operatorname{End}(\mathcal{H})$ the space of linear operators acting on \mathcal{H} . We recall Schur-Weyl duality in the general setting. We fix a dimension $d \geq 2$ and a positive integer $k \in \mathbb{N}$ and consider the following unitary representations of the symmetric group \mathfrak{S}_k and the unitary group \mathcal{U}_d on the state space $\mathcal{H} = (\mathbb{C}^d)^{\otimes k}$:

$$\varphi \colon \mathfrak{S}_k \longrightarrow \mathcal{U}(\mathcal{H}), \quad \pi \longmapsto (|\phi_1\rangle \otimes \cdots \otimes |\phi_k\rangle \mapsto |\phi_{\pi^{-1}(1)}\rangle \otimes \cdots \otimes |\phi_{\pi^{-1}(k)}\rangle)$$
 (2.17)

$$\psi \colon \mathcal{U}_d \longrightarrow \mathcal{U}(\mathcal{H}), \quad U \longmapsto U^{\otimes k}$$
 (2.18)

It is straightforward to check that the two actions on \mathcal{H} commute, $[\psi(U), \varphi(\pi)] = 0$ for all $U \in \mathcal{U}_d$ and $\pi \in \mathfrak{S}_k$. This implies that the algebra $\langle \varphi(\pi) : \pi \in \mathfrak{S}_k \rangle \subseteq \operatorname{End}(\mathcal{H})$ spanned by the representation (2.17) is contained in the commutant of the algebra $\langle Q(U) : U \in \mathcal{U}_d \rangle \subseteq \operatorname{End}(\mathcal{H})$ spanned by the representation (2.18), and vice versa. Schur-Weyl duality (see, e.g., [14]) states that, in fact, these two algebras are equal to each other's commutant. It follows that, if an operator $X \in \mathcal{L}(\mathcal{H})$ commutes with all $\psi(U)$ for $U \in \mathcal{U}_d$, it can be written as

$$X = \sum_{\pi \in \mathfrak{S}_k} x_{\pi} \varphi(\pi) \tag{2.19}$$

for some coefficients $x_{\pi} \in \mathbb{C}$. Together with the complete reducibility of the representations (2.17) and (2.18), Schur-Weyl duality furthermore yields the following useful decompositions of state space and representations:

$$\mathcal{H} \cong \bigoplus_{\lambda \vdash_d k} \mathcal{V}_{\lambda}^d \otimes \mathcal{W}_{\lambda} \tag{2.20}$$

$$\varphi(\pi) \cong \bigoplus_{\lambda \vdash_d k} \mathbb{1}_{\mathcal{V}_{\lambda}^d} \otimes \varphi_{\lambda}(\pi)$$
 (2.21)

$$\psi(U) \cong \bigoplus_{\lambda \vdash_d k} \psi_{\lambda}^d(U) \otimes \mathbb{1}_{\mathcal{W}_{\lambda}}. \tag{2.22}$$

In all three equations the direct sums run over partitions $\lambda = (\lambda_1, \dots, \lambda_d) \vdash_d k$ of k into d parts, i.e., $\lambda_1 \geq \dots \geq \lambda_d \geq 0$ and $\sum_{i=1}^d \lambda_i = k$. These partitions label the irreducible representations (irreps) $(\mathcal{V}_{\lambda}^d, \psi_{\lambda})$ of \mathcal{U}_d and $(\mathcal{W}_{\lambda}, \varphi_{\lambda})$ of \mathfrak{S}_k . A consequence of Schur-Weyl duality is that the \mathfrak{S}_k -irrep \mathcal{W}_{λ} is the multiplicity space of the \mathcal{U}_d -irrep $(\mathcal{V}_{\lambda}^d, \psi_{\lambda}^d)$, and vice versa. By Schur's Lemma, any operator $X \in \mathcal{L}(\mathcal{H})$ commuting with all $\psi(U)$ is of the form

$$X = \bigoplus_{\lambda \vdash_d k} \mathbb{1}_{\mathcal{V}_{\lambda}^d} \otimes X_{\lambda} \tag{2.23}$$

for some $X_{\lambda} \in \mathcal{L}(W_{\lambda})$, and the analogous statement holds for operators commuting with all $\varphi(\pi)$. The dimensions of the irreps appearing in (2.20) are given by the following formulae:

$$d_{\lambda} := \dim \mathcal{W}_{\lambda} = \frac{k!}{\prod_{(i,j)\in\lambda} h(i,j)}$$
 (2.24)

$$m_{\lambda}^{d} := \dim \mathcal{V}_{\lambda}^{d} = \prod_{1 \le i < j \le d} \frac{\lambda_{i} - \lambda_{j} + j - i}{j - i}.$$
 (2.25)

In Eq. (2.24) the hook length h(i,j) of a box $(i,j) \in \lambda$ is defined as the number of boxes to the right and below of (i,j) plus the box itself.

3. TPE TO RELATIVE ERROR DESIGNS

A common strategy for proving approximate design bounds is to first prove a tensor product expander (TPE) bound, that is, a $(2 \to 2)$ -norm bound on the distance from the approximate to an exact design, which can then be converted to diamond norm, semidefinite order inequality, or other stronger criteria. Conventionally, this conversion involves a factor of the dimension of the space, usually q^{mk} up to some constants in the exponential. It is however known that, when restricting states and channels to von Neumann subalgebras of the full Hilbert space, one may often replace the space's dimension by a lower effective dimension that excludes subspace multiplicity.

¹Partitions are often graphically represented as Young diagrams, a left-aligned arrangement of k boxes with λ_i boxes in the i-th row. For example, the Young diagram corresponds to the partition $\lambda = (3,1)$.

Lemma 3.1. Let $f(\cdot)$ be any convex function on quantum states on a given finite-dimensional space. Then for any von Neumann algebra $\mathcal{M} = \bigoplus_{\lambda} (\mathcal{A}_{\lambda} \otimes \mathbb{1}_{B_{\lambda}})$ and $d \in \mathbb{N}$, the supremum $\sup_{\rho} f(\mathcal{E}_{\mathcal{M} \otimes \mathbb{M}_d}(\rho))$ is achieved by a state of the form $\rho^{A_{\lambda_0} \otimes \mathbb{M}_d} \otimes \hat{1}/|B_{\lambda_0}|$ for some block label λ_0 , where

$$|\rho\rangle^{\tilde{A}_{\lambda_0}\otimes\mathbb{M}_d} \cong \sum_i \alpha_i |i\rangle^{\tilde{A}_{\lambda_0}} \otimes |i\rangle^{\mathbb{M}_d} .$$
 (3.1)

for an appropriate choice of basis and up to a unitary rotation on the auxiliary system.

Proof. Because f is convex, $\sup_{\rho} f(\rho)$ is attained within finite dimension by a pure input state ρ . Because the argument to f is always an output of $\mathcal{E}_{\mathcal{M}\otimes\mathbb{M}_d}$, we may assume that the input state ρ is already in \mathcal{M}_* . Therefore, a maximum-attaining ρ is supported on a single λ , which we call λ_0 . Furthermore, any input ρ that is correlated between $A_{\lambda_0}\otimes\mathbb{M}_d$ and B_{λ_0} will be replaced by $\operatorname{tr}_{B_{\lambda_0}}(\rho)\otimes \hat{1}/|B_{\lambda_0}|$, where $\operatorname{tr}_{B_{\lambda_0}}(\rho)$ is mixed. Again by convexity, a supremum-attaining input will have the form $\rho^{A_{\lambda_0}\otimes\mathbb{M}_d}\otimes\hat{1}/|B_{\lambda_0}|$, where $\rho^{A_{\lambda_0}\otimes\mathbb{M}_d}$ is pure. Schmidt decomposition completes the Lemma.

Lemma 3.2. Let $\mathcal{L} \subseteq \mathcal{N} \subseteq \mathcal{M}$ as von Neumann algebra inclusions. Then

$$\sup_{\rho} D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{N}}(\rho)) \le \sup_{\rho} D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{L}}(\rho)) , \qquad (3.2)$$

where both suprema are taken over the set of density matrices.

Proof. Via the chain rule of relative entropy in (2.13),

$$\sup_{\rho} D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{L}}(\rho)) = \sup_{\rho} \left(D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{N}}(\rho)) + D(\mathcal{E}_{\mathcal{N}}(\rho) \| \mathcal{E}_{\mathcal{L}}(\rho)) \right)$$
(3.3)

$$\geq D(\mathcal{E}_{\mathcal{M}}(\omega) \| \mathcal{E}_{\mathcal{N}}(\omega)) + D(\mathcal{E}_{\mathcal{N}}(\omega) \| \mathcal{E}_{\mathcal{L}}(\omega))$$
(3.4)

$$= \sup_{\rho} D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{N}}(\rho)) + D(\mathcal{E}_{\mathcal{N}}(\omega) \| \mathcal{E}_{\mathcal{L}}(\omega))$$
(3.5)

$$\geq \sup_{\rho} D(\mathcal{E}_{\mathcal{M}}(\rho) \| \mathcal{E}_{\mathcal{N}}(\rho)) , \qquad (3.6)$$

where ω in the first inequality is chosen to maximize $D(\mathcal{E}_{\mathcal{M}}(\omega)||\mathcal{E}_{\mathcal{N}}(\omega))$, and the second inequality follows from positivity of the relative entropy on states.

Lemma 3.3. Let \mathcal{M} be a von Neumann algebra in finite dimension with block decomposition $\mathcal{M} = \bigoplus_{\lambda} \mathcal{A}_{\lambda} \otimes \mathbb{C}^{|B_{\lambda}|}$. Then for any auxiliary dimension $d \geq 0$,

$$\sup_{\rho} D(\mathcal{E}_{\mathcal{M} \otimes \mathbb{M}_d}(\rho) \| \mathcal{E}_{\mathbb{C} \otimes \mathbb{M}_d}(\rho)) \le \max_{\lambda} \log \left(\frac{|A_{\lambda}| |\mathcal{M}|}{|B_{\lambda}|} \right). \tag{3.7}$$

Proof. Since relative entropy is jointly convex, we may assume by Lemma 3.1 that the supremum on the left-hand side of (3.7) is achieved on a state of the form $\rho^{A_{\lambda_0}\otimes\mathbb{M}_d}\otimes \hat{1}/|B_{\lambda_0}|$, where $\rho^{A_{\lambda_0}\otimes\mathbb{M}_d}\equiv |\rho\rangle\langle\rho|^{A_{\lambda_0}\otimes\mathbb{M}_d}$ is a pure state with

$$|\rho\rangle^{\tilde{A}_{\lambda_0}\otimes\mathbb{M}_d} \cong \sum_i \alpha_i |i\rangle^{\tilde{A}_{\lambda_0}} \otimes |i\rangle^{\mathbb{M}_d}$$
 (3.8)

for some block label λ_0 and Schmidt coefficients α_i . Since $\mathcal{E}_{\mathbb{C}}$ does not directly act on the auxiliary system, we may project \mathbb{M}_d to a subsystem of dimension $|A_{\lambda_0}|$ in both arguments without decreasing the relative entropy. Using the chain rule of quantum relative entropy with respect to a conditional expectation,

$$D\left(\mathcal{E}_{\mathcal{M}\otimes\mathbb{M}_{|A_{\lambda_0}|}}(\rho)\Big\|\mathcal{E}_{\mathbb{C}\otimes\mathbb{M}_{|A_{\lambda_0}|}}(\rho)\right) = H\left(\mathcal{E}_{\mathbb{C}\otimes\mathbb{M}_{|A_{\lambda_0}|}}(\rho)\right) - H\left(\mathcal{E}_{\mathcal{M}\otimes\mathbb{M}_{|A_{\lambda_0}|}}(\rho)\right) . \tag{3.9}$$

For any choice of λ_0 , the maximum value of the first term is $\log(|A_{\lambda_0}||\mathcal{M}|)$, and the minimum value of the second term is $\log |B_{\lambda_0}|$.

Lemma 3.4. For any finite-dimensional von Neumann algebra $\mathcal{M} = \bigoplus_{\lambda} \mathcal{A}_{\lambda} \otimes \mathbb{1}_{B_{\lambda}}$ and subalgebra $\mathcal{N} \subseteq \mathcal{M}$,

$$C_{\rm cb}(\mathcal{M}:\mathcal{N}) \le |\mathcal{M}| \max_{\lambda} \frac{|A_{\lambda}|}{|B_{\lambda}|}$$
 (3.10)

Proof. First recall Equation (2.16), stating that

$$\log C_{\mathrm{cb}}(\mathcal{M}:\mathcal{N}) = \sup_{d \in \mathbb{N}} \max_{\rho \in \mathcal{M}_* \otimes \mathbb{M}_d} D((\mathcal{E}_{\mathcal{M}} \otimes \mathrm{Id}_{(d)})(\rho) \| (\mathcal{E}_{\mathcal{N}} \otimes \mathrm{Id}_{(d)})(\rho)) . \tag{3.11}$$

Via Lemma 3.2.

$$D((\mathcal{E}_{\mathcal{M}} \otimes \operatorname{Id}_{(d)})(\rho) \| (\mathcal{E}_{\mathcal{N}} \otimes \operatorname{Id}_{(d)})(\rho)) \le D((\mathcal{E}_{\mathcal{M}} \otimes \operatorname{Id}_{(d)})(\rho) \| (\mathcal{E}_{\mathbb{C}} \otimes \operatorname{Id}_{(d)})(\rho))$$
(3.12)

for each value of $d \in \mathbb{N}$, so it must hold for the supremum over $d \in \mathbb{N}$. Via 3.3, noting that \mathbb{C} is a subalgebra of every finite-dimensional von Neumann algebra,

$$\sup_{\rho} D(\mathcal{E}_{\mathcal{M} \otimes \mathbb{M}_d}(\rho) \| \mathcal{E}_{\mathbb{C} \otimes \mathbb{M}_d}(\rho)) \le \max_{\lambda} \log \left(\frac{|A_{\lambda}| |\mathcal{M}|}{|B_{\lambda}|} \right)$$
(3.13)

for every $d \in \mathbb{N}$, so again it must hold for the supremum over $d \in \mathbb{N}$.

Lemma 3.5. In the Schur-Weyl decomposition $\mathbb{C}_d^{\otimes k} \cong \bigoplus_{\lambda \vdash_d k} \mathcal{V}_\lambda^d \otimes \mathcal{W}_\lambda$ of k copies of a complex Hilbert space of dimension d, we have the following bound for each partition $\lambda \vdash_d k$:

$$|\mathcal{V}_{\lambda}^{d}| \ge d^{k} \frac{|\mathcal{W}_{\lambda}|}{k!} \left(1 - \frac{k^{2}}{d}\right). \tag{3.14}$$

Proof. By [8, Theorem 1.16] or its later restatement as [30, Lemma 2.11] we have

$$|\mathcal{V}_{\lambda}^{d}| = \frac{|\mathcal{W}_{\lambda}|}{k!} \prod_{(i,j)\in\lambda} (d+j-i) , \qquad (3.15)$$

where $(i, j) \in \lambda$ means that i and j respectively denote the row index and column index of a cell in the Young tableau. We have $i, j \leq k$ since a Young tableau has k cells in total, and hence

$$\prod_{(i,j)\in\lambda} (d+j-i) \ge (d-k)^k = d^k (1-k/d)^k \ge d^k (1-k^2/d) , \qquad (3.16)$$

where the final inequality is Bernoulli's inequality.

Lemma 3.6. Let A be a quantum system with subsystem decomposition $A = A_1 \otimes \cdots \otimes A_r$. Denote by \mathcal{T}_C the k-fold twirl, averaging over the unitary group on $C^{\otimes k}$ for a (sub)system C, and let $\mathcal{M}_{(n)}$ be the von Neumann subalgebra onto which \mathcal{T}_{A_n} projects. For any subalgebra $\mathcal{L} \subseteq \bigotimes_{n=1}^r \mathcal{M}_{(n)}$,

$$C_{\text{cb}}\left(\bigotimes_{n=1}^{r} \mathcal{M}_{(n)} : \mathcal{L}\right) \le k!^{r} \prod_{n=1}^{r} \min\left\{\left(1 - \frac{k^{2}}{|A_{n}|}\right)^{-1}, \frac{|A_{n}|^{k}}{k!}\right\}.$$
 (3.17)

Proof. Let $\mathcal{M}_{(n),k}$ denote the von Neumann subalgebra corresponding to the invariant subspace of the k-fold Haar unitary twirl \mathcal{T}_{A_n} on a system $A_n^{\otimes k}$. Recalling Schur-Weyl duality and the block structure (2.23) of operators commuting with any unitary (which a Haar unitary twirl enforces), the invariant subalgebra of $\bigotimes_{n=1}^r \mathcal{T}_{A_n}$ has the form

$$\bigotimes_{n=1}^{r} \mathcal{M}_{(n),k} := \bigotimes_{n=1}^{r} \left(\bigoplus_{\lambda} \mathbb{B} \left(\mathbb{C}^{|\mathcal{V}_{\lambda}^{|A_{n}|}|} \right) \otimes \mathbb{B}(\mathcal{W}_{\lambda}) \right) . \tag{3.18}$$

By the block decomposition product, each block of $\bigotimes_{n=1}^r \mathcal{M}_{(n),k}$ has the product of multiplicities of its constituents. Let $\vec{\lambda} = (\lambda_1, \dots, \lambda_r)$ index each block of $\bigotimes_{n=1}^r \mathcal{M}_{(n),k}$. By Lemma 3.5, each multiplicity $b_{\vec{\lambda}}$ in $\bigotimes_{n=1}^r \mathcal{M}_{(n),k}$ obeys

$$b_{\vec{\lambda}} \ge \min\left\{\frac{1}{k!^r} \prod_{n=1}^r \left(|A_n|^k |\mathcal{W}_{\lambda_n}^{|A_n|}| \left(1 - \frac{k^2}{|A_n|}\right) \right), 1 \right\}. \tag{3.19}$$

Each untraced block with index $\vec{\lambda}$ has size $a_{\vec{\lambda}} := |\mathcal{W}_{\lambda_1}^{|A_1|}| \dots |\mathcal{W}_{\lambda_n}^{|A_n|}|$. Recall also that $|A|^k = |A_1|^k \dots |A_r|^k$. Lemma 3.4 together with the fact that a multiplicity is always at least 1 (because there are no systems of dimension less than 1) then yields that

$$C_{\text{cb}}\left(\bigotimes_{n=1}^{r} \mathcal{M}_{(n)} : \mathcal{L}\right) \le |A|^k \max_{\vec{\lambda}} \frac{a_{\vec{\lambda}}}{b_{\vec{\lambda}}} \le \prod_{n=1}^{r} \min\left\{k! \left(1 - \frac{k^2}{|A_n|}\right)^{-1}, |A_n|^{2k}\right\},$$
 (3.20)

which concludes the proof.

Lemma 3.7. Let A be a quantum system with subsystem decomposition $A = A_1 \otimes \cdots \otimes A_r$. Denote by \mathcal{T}_C the k-fold twirl, averaging over the unitary group on $C^{\otimes k}$ for a (sub)system C. Let the von Neumann algebra \mathcal{N} be a subalgebra of $\bigotimes_{n=1}^r \mathcal{T}_{A_n}$, and let Φ be any trace-symmetric quantum channel on $A^{\otimes k}$ for which \mathcal{N} is a fixed-point subalgebra. If

$$\epsilon := \left\| \left(\bigotimes_{n=1}^{r} \mathcal{T}_{A_{n}} \right) \left(\Phi - \mathcal{E}_{\mathcal{N}} \right) \left(\bigotimes_{n=1}^{r} \mathcal{T}_{A_{n}} \right) \right\|_{2 \to 2} \times k!^{r} \prod_{n=1}^{r} \min \left\{ \left(1 - \frac{k^{2}}{|A_{n}|} \right)^{-1}, \frac{|A_{n}|^{2k}}{k!} \right\} < 1,$$
(3.21)

then
$$(1 - \epsilon)\mathcal{E}_{\mathcal{N}} \prec \Phi \prec (1 + \epsilon)\mathcal{E}_{\mathcal{N}}$$
.

Proof. We recall a standard result that relates a system's completely bounded return time (the time it takes for a particular cp-order inequality to hold) to its spectral gap. This result is shown explicitly as [17, Lemma A.1] based on similar results and methods from [15], [16], and [27]. For

any trace-symmetric quantum channel Φ that is an \mathcal{N} -bimodule map, that is an \mathcal{N} -bimodule map (implied by the channel being absorbed by $\mathcal{E}_{\mathcal{N}}$), if

$$\gamma := \|\mathcal{E}_{\mathcal{M}} \Phi \mathcal{E}_{\mathcal{M}} - \mathcal{E}_{\mathcal{N}}\|_{2 \to 2} < 1, \qquad (3.22)$$

then for every $n \in \mathbb{N}$ such that $n \geq \ln C_{\text{cb}}(\mathcal{M} : \mathcal{N}) / \ln(1/\gamma)$, the desired cb-order inequality holds with $\epsilon = \gamma^n C_{\text{cb}}(\mathcal{M} : \mathcal{N})$. Combining this fact with Lemma 3.6 yields the desired inequality, in this case setting n = 1.

4. Analyzing approximate designs using the alternating projection method

We will use the following notation in this section. We consider k copies of systems A and B each consisting of m qudits of local dimension q. The full system is thus A^kB^k , and we sometimes write this as $A_1 \ldots A_kB_1 \ldots B_k$ or $A_1B_1 \ldots A_kB_k$ depending on the locality cut. In each case, $A_i \cong A$ and $B_i \cong B$ for all i. We denote by \mathfrak{S}_k the symmetric group of degree k, and we consider its action (2.17) on a tensor space $X_1 \ldots X_k$ by permuting tensor factors. The corresponding permutation operator for $\pi \in \mathfrak{S}_k$ will be denoted by $\pi_X \in \mathcal{L}(X_1 \ldots X_k)$.

For $X \in \{A, B, AB\}$ we consider twirling with respect to the representation (2.18) of the unitary group $\mathcal{U}(X)$ acting on X,

$$\mathcal{T}_X(\cdot) = \int_{\mathcal{U}(X)} dU_X \, U_X^{\otimes k} \cdot (U_X^{\dagger})^{\otimes k},\tag{4.1}$$

with dU_X denoting the Haar measure on $\mathcal{U}(X)$. Such twirls give rise to orthogonal projections on the Hilbert space $\mathcal{L}(X)$ of operators acting on X, equipped with the Hilbert-Schmidt (or Frobenius) inner product $\langle X,Y\rangle := \operatorname{tr}(X^{\dagger}Y)$. In the proofs below we will make repeated use of the following fact:

Lemma 4.1. Let X be a Hilbert space and denote by $\mathcal{L}(X)$ its Hilbert space of operators equipped with the Hilbert-Schmidt inner product. Let $\mathcal{P}, \mathcal{Q}, \mathcal{R}$ be orthogonal projections acting on $\mathcal{L}(X)$, and assume that

$$PR = RP = R = RQ = QR. \tag{4.2}$$

Then for any $X \in \mathcal{L}(X)$ we have

$$\|\mathcal{QP}(X) - \mathcal{R}(X)\|_2^2 = \langle X, \mathcal{PQP}(X) - \mathcal{R}(X) \rangle. \tag{4.3}$$

Proof. Using the fact that $||X||_2^2 = \langle X, X \rangle$, we can expand the 2-norm as follows:

$$\|\mathcal{QP}(X) - \mathcal{R}(X)\|_2^2 = \langle \mathcal{QP}(X) - \mathcal{R}(X), \mathcal{QP}(X) - \mathcal{R}(X) \rangle \tag{4.4}$$

$$=\langle \mathcal{QP}(X), \mathcal{QP}(X)\rangle - \langle \mathcal{QP}(X), \mathcal{R}(X)\rangle$$

$$-\langle \mathcal{R}(X), \mathcal{QP}(X)\rangle + \langle \mathcal{R}(X), \mathcal{R}(X)\rangle. \tag{4.5}$$

Because of the self-adjointness of \mathcal{P} , \mathcal{Q} and \mathcal{R} with respect to $\langle \cdot, \cdot \rangle$, we have

$$\langle \mathcal{QP}(X), \mathcal{QP}(X) \rangle = \langle X, \mathcal{PQ}^2 \mathcal{P}(X) \rangle = \langle X, \mathcal{PQP}(X) \rangle$$
 (4.6)

$$\langle \mathcal{R}(X), \mathcal{R}(X) \rangle = \langle X, \mathcal{R}^2(X) \rangle = \langle X, \mathcal{R}(X) \rangle.$$
 (4.7)

Moreover, because of the assumptions in (4.2),

$$\langle \mathcal{QP}(X), \mathcal{R}(X) \rangle = \langle X, \mathcal{PQR}(X) \rangle = \langle X, \mathcal{R}(X) \rangle,$$
 (4.8)

and similarly $\langle \mathcal{R}(X), \mathcal{QP}(X) \rangle = \langle X, \mathcal{R}(X) \rangle$. Using these identities in (4.5) finishes the proof. \square

In the following, we analyze two different protocols. The first one alternates between individual twirls on A^k and B^k and swapping the first ℓ qudits between each of the A_i and B_i blocks. The second protocol alternates between individual twirls and a twirl on $A_i^{\ell}B_i^{\ell}$, where A_i^{ℓ} consists of the first ℓ qudits in system A_i , and similarly for B_i^{ℓ} . For both protocols we show convergence to the full twirl \mathcal{T}_{AB} and derive bounds on the corresponding convergence rates.

4.1. **Twirl-Swap-Twirl.** Here we show that a protocol alternating between a) individual twirls on A^k and B^k and b) swapping the first ℓ qudits between each of the A_i and B_i blocks converges to the full twirl on \mathcal{T}_{AB} , thus implementing an approximate k-design on A^kB^k that converges to an exact design in the limit. We will give a bound on the convergence rate of this protocol using the alternate projection method [34].

To this end, we define the following operation that swaps the first ℓ qudits between A and B in each of the k blocks:

$$SWAP_{\ell} := \bigotimes_{i=1}^{k} \mathbb{F}_{A_i^{\ell} B_i^{\ell}} \cdot \mathbb{F}_{A_i^{\ell} B_i^{\ell}}.$$
(4.9)

Note that $SWAP_{\ell}^2 = id$.

Consider now the following maps defined in terms of SWAP_{ℓ} and the twirls $\mathcal{T}_X(\cdot)$ for $X \in \{A, B, AB\}$ from (4.1):

$$\mathcal{P} = \mathcal{T}_A \otimes \mathcal{T}_B \tag{4.10}$$

$$Q_{sw} = SWAP_{\ell} \circ \mathcal{P} \circ SWAP_{\ell} \tag{4.11}$$

$$\mathcal{R} = \mathcal{T}_{AB}.\tag{4.12}$$

Our protocol consists of a repeated application of \mathcal{P} followed by \mathcal{Q}_{sw} .

The maps $\mathcal{P}, \mathcal{Q}_{sw}, \mathcal{R}$ defined in (4.10)–(4.12) are orthogonal projections in the space of operators acting on $A^k B^k$. For example, \mathcal{P} satisfies $\mathcal{P}^{\dagger} = \mathcal{P}$ (with respect to the Frobenius inner product $\langle X, Y \rangle = \operatorname{tr}(X^{\dagger}Y)$ and $\mathcal{P}^2 = \mathcal{P}$, and similarly for \mathcal{Q}_{sw} and \mathcal{R} . By left- and right-invariance of the Haar measure, we also have the 'dominance relations'

$$\mathcal{PR} = \mathcal{RP} = \mathcal{R} = \mathcal{RQ}_{sw} = \mathcal{Q}_{sw} \mathcal{R}. \tag{4.13}$$

We claim that for $\ell < m$ images of the projections in (4.10)–(4.12) satisfy

$$\operatorname{im} \mathcal{P} \cap \operatorname{im} \mathcal{Q}_{sw} = \operatorname{im} \mathcal{R},$$
 (4.14)

which implies via von Neumann's alternating projection theorem [34] that

$$\lim_{n \to \infty} (\mathcal{Q}_{sw} \mathcal{P})^n = \mathcal{R}. \tag{4.15}$$

To see that (4.14) is true, note that any operator $\mathcal{P}(X)$ for $X \in \mathcal{L}(A^k B^k)$ is invariant under $U_A^{\otimes k} \otimes V_B^{\otimes k}$ for any $U_A \in \mathcal{U}(A)$ and $V_B \in \mathcal{U}(B)$. Schur-Weyl duality in the form (2.19) then says

that we have $\mathcal{P}(X) = \sum_{\sigma,\tau \in \mathfrak{S}_k} x_{\sigma,\tau} \sigma_A \otimes \tau_B$ for some coefficients $x_{\sigma,\tau} \in \mathbb{C}$, where $\sigma_A = \varphi(\sigma)_A \in \mathcal{L}(A^k)$ denotes the permutation operator corresponding to the permutation $\sigma \in \mathfrak{S}_k$ acting on the k copies of the A-system, and similarly for τ_B . On the other hand, by (4.119) in Sec. 4.3 we have SWAP $_{\ell}(\sigma_A \otimes \tau_B) = \tau_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \otimes \sigma_{B^{\ell}} \otimes \tau_{B^{m-\ell}}$. The intersection of im \mathcal{P} and im \mathcal{Q} thus consists of operators that can be written as a linear combination of permutation operators $\pi_{AB} = \pi_A \otimes \pi_B$ for $\pi \in \mathfrak{S}_k$, and this space is equal to im \mathcal{R} by another application of (2.19).

Our goal is to obtain a bound on the convergence rate in (4.15), for which we follow the exposition in [11]. The *subspace angle* $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{sw})$ can be expressed as [11, Lemma 9.5]

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{sw}}) := \|\mathcal{Q}_{\operatorname{sw}} \mathcal{P} - \mathcal{R}\|_{2 \to 2} = \sup_{X \neq 0} \frac{\|\mathcal{Q}_{\operatorname{sw}} \mathcal{P}(X) - \mathcal{R}(X)\|_{2}}{\|X\|_{2}}, \tag{4.16}$$

The following bound in terms of the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{sw}})$ then holds for all $X \in \mathcal{L}(A^k B^k)$ and $n \in \mathbb{N}$ [11, Theorem 9.8]:

$$\|(\mathcal{Q}_{sw}\mathcal{P})^n(X) - \mathcal{R}(X)\|_2 \le c(\operatorname{im}\mathcal{P}, \operatorname{im}\mathcal{Q}_{sw})^{2n-1}\|X\|_2.$$
 (4.17)

Our primary technical result in this section is a bound on the subspace angle of \mathcal{P} and \mathcal{Q} :

Proposition 4.2. Let A and B each consist of k blocks of m qudits of local dimension q, and let ℓ be the number of qudits swapped between A and B as defined in (4.9), where $\ell \leq m/2$ without loss of generality.² We assume that $k \leq q^m$ and set $a = (1 - \varepsilon)^{-2}$ with $\varepsilon = \frac{k^2}{2q^m}$. Then the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{sw})$ corresponding to the projections \mathcal{P} and \mathcal{Q}_{sw} defined in (4.10) and (4.11), respectively, satisfies

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{sw}) \le \sqrt{9ak^2q^{-m} + 4a^2k!^3q^{-2\ell} + 2a^2k!^5q^{-4\ell}}$$
 (4.18)

From Proposition 4.2, we obtain the following main Theorem on designs:

Theorem 4.3. Let A and B each consist of k blocks of m qudits of local dimension q. Let \mathcal{T}_A and \mathcal{T}_B be exact k-design channels respectively on $A^{\otimes k}$ and $B^{\otimes k}$. Let SWAP_{ℓ} denote a unitary that exchanges an arbitrary ℓ -qubit subsystem of A with one of B in each of the k blocks for $\ell \leq m/2$. Furthermore, assume that $k^2 < 2q^m$. Then $(\mathcal{T}_A \otimes \mathcal{T}_B)\mathrm{SWAP}_{\ell}(\mathcal{T}_A \otimes \mathcal{T}_B)$ is an ϵ -approximate multiplicative error k-design channel when

$$\ell \geq \frac{7}{2} \log_q k! + \log_q \left(\frac{1}{\epsilon}\right) - 4 \log_q \left(1 - \frac{k^2}{q^m}\right) + \log_q 4.$$

Proof. By Proposition 4.2,

$$\|\mathcal{P}\mathcal{Q}_{sw} - \mathcal{T}_{AB}\|_{2\to 2} \le \sqrt{9ak^2q^{-m} + 4a^2k!^3q^{-2\ell} + 2a^2k!^5q^{-4\ell}}$$

To simplify, assume that $q^m \ge 2q^\ell \ge 2k!$ and that $k \ge 2$ and $a^2k!^3q^{-2\ell} \ge a^2k!^5q^{-4\ell} \ge 8k^2q^{-m}$. If k=1, then the design is exact, as local mixings yield complete, global mixture, so we ignore that case. Therefore,

$$\|\mathcal{P}\mathcal{Q}_{sw} - \mathcal{T}_{AB}\|_{2\to 2} \le 3ak!^{3/2}q^{-\ell}$$
.

²The case $m/2 < \ell < m$ follows from this by a simple relabeling of A and B.

In Lemma 3.7, we therefore have

$$\epsilon = 3q^{-\ell}ak!^{7/2} \Big(1 - \frac{k^2}{q^m}\Big)^{-2} \leq 3q^{-\ell}k!^{7/2} \Big(1 - \frac{k^2}{q^m}\Big)^{-4} \; .$$

We solve this equation for ℓ to obtain the bound stated in the theorem. Note that another multiplicative contribution comes from pre- and post-compositions of $\mathcal{T}_A \otimes \mathcal{T}_B$. Finally, note that while Proposition 4.2 explicitly defines the swaps as on the first qudits by index, the definition of the first qudits is arbitrary.

We now give the proof of the technical result in Proposition 4.2.

Proof of 4.2. We first rewrite the squared numerator on the right-hand side of (4.16) by expanding the 2-norm, using Lemma 4.1 and the relations (4.13):

$$\sup_{X \neq 0} \frac{\|\mathcal{Q}_{sw}\mathcal{P}(X) - \mathcal{R}(X)\|_2^2}{\|X\|_2^2} = \sup_{X \neq 0} \frac{\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X) - \mathcal{R}(X) \rangle}{\langle X, X \rangle}.$$
 (4.19)

The relations (4.13) furthermore show that the supremum in (4.19) is achieved on some $X = \mathcal{P}(X)$, which according to (2.19) can be written as

$$X = \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, \sigma_A \otimes \tau_B \tag{4.20}$$

for some coefficients $x_{\sigma,\tau} \in \mathbb{C}$. Note that $\sigma_A \otimes \tau_B$ is invariant under the projections $\mathcal{P}, \mathcal{Q}_{sw}, \mathcal{R}$ whenever $\sigma = \tau$. Following [20], we denote by \mathbf{G}_S the $(k! \times k!)$ -Gram matrix of permutation operators acting on k copies of a system S with elements $(\mathbf{G}_S)_{\sigma,\tau} = \langle \sigma_S, \tau_S \rangle_{\mathbf{n}}$ for $\sigma, \tau \in \mathfrak{S}_k$, where

$$\langle X, Y \rangle_{\rm n} = |S|^{-k} \operatorname{tr}(X^{\dagger}Y)$$
 (4.21)

denotes the normalized Frobenius product and $|S| = \dim \mathcal{H}_S$.

Because of the quotient of inner products in (4.19), we can replace $\langle \cdot, \cdot \rangle$ by the normalized inner product $\langle \cdot, \cdot \rangle_n$. Using the assumption $X = \mathcal{P}(X)$ together with Lemma 4.5(i), (ii) and (iv) below, the expression (4.19) is then equal to

$$\frac{\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X) - \mathcal{R}(X) \rangle}{\langle X, X \rangle} = \frac{\langle \mathbf{x}, \left[\mathbf{M} \left(\mathbf{G}_{A}^{-1} \otimes \mathbf{G}_{B}^{-1} \right) \mathbf{M} - \mathbf{N}^{T} \mathbf{G}_{AB}^{-1} \mathbf{N} \right] \mathbf{x} \rangle}{\langle \mathbf{x}, \left(\mathbf{G}_{A} \otimes \mathbf{G}_{B} \right) \mathbf{x} \rangle}, \tag{4.22}$$

where the inner product $\langle \cdot, \cdot \rangle$ on the right-hand side (and in the following) is now the usual standard inner product on $\mathbb{C}^{k!^2}$. The matrices \mathbf{M} of size $(k!^2 \times k!^2)$ and \mathbf{N} of size $(k! \times k!^2)$ have the following coefficients for $\pi, \rho, \omega, \chi \in \mathfrak{S}_k$ (cf. Lemma 4.5):

$$(\mathbf{M})_{(\pi,\rho),(\omega,\chi)} = \langle \pi_A, \chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \rho_B, \omega_{B^{\ell}} \otimes \chi_{B^{m-\ell}} \rangle_{\mathbf{n}}$$

$$(4.23)$$

$$(\mathbf{N})_{\pi,(\omega,\chi)} = \langle \pi_{AB}, \omega_A \otimes \chi_B \rangle_{\mathbf{n}}. \tag{4.24}$$

Here, for $\chi, \omega \in \mathfrak{S}_k$ we write χ_{A^ℓ} for the permutation operator acting on the first $\ell \leq m$ qudits within each of the k blocks (of size m each) in A, and $\omega_{A^{m-\ell}}$ for the permutation operator acting on the remaining $m-\ell$ qudits within each block. The same conventions hold for permutations acting on the blocks in B.

The rest of the proof is concerned with bounding the expression (4.22) for all $\mathcal{L}(A^k B^k) \ni X \neq 0$ (or equivalently $\mathbb{C}^{k!^2} \ni \mathbf{x} \neq 0$), in turn giving a bound on the supremum over $\mathbf{x} \neq 0$ and thus the 2-norm in (4.16) that we seek to estimate. To this end, we employ the following inequalities proved in [20, Lemma 1], with λ_{\min} and λ_{\max} denoting the smallest and largest eigenvalue of a Hermitian matrix, respectively:

$$\lambda_{\min}(\mathbf{G}_A) \ge 1 - \frac{k^2}{2q^m} \equiv 1 - \varepsilon$$
 (4.25)

$$\lambda_{\max}(\mathbf{G}_{AB}) \le \exp\left(\frac{k^2}{2q^{2m}}\right) \equiv \exp(\varepsilon'),$$
 (4.26)

where we set $\varepsilon := \frac{k^2}{2q^m}$ and $\varepsilon' := \frac{k^2}{2q^{2m}}$. From (4.25) we get $\lambda_{\max}\left(\mathbf{G}_A^{-1}\right) \leq (1-\varepsilon)^{-1}$ and hence

$$\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1} \le (1 - \varepsilon)^{-2} \, \mathbb{1}_{k!^2}. \tag{4.27}$$

We also have $\lambda_{\max}\left(-\mathbf{G}_{AB}^{-1}\right) \leq -\exp(-\varepsilon')$ from (4.26), which leads to

$$-\mathbf{G}_{AB}^{-1} \le -\exp(-\varepsilon')\mathbb{1}_{k!}.\tag{4.28}$$

With $a := (1 - \varepsilon)^{-2}$ and $b := \exp(-\varepsilon')$, these operator inequalities yield

$$\langle \mathbf{x}, \mathbf{M} \left(\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1} \right) \mathbf{M} - \mathbf{N}^T \mathbf{G}_{AB}^{-1} \mathbf{N} \mathbf{x} \rangle \le \langle \mathbf{x}, \left(a \mathbf{M}^2 - b \mathbf{N}^T \mathbf{N} \right) \mathbf{x} \rangle.$$
 (4.29)

The bound (4.25) further gives $\mathbf{G}_A \otimes \mathbf{G}_B \geq a^{-1} \mathbb{1}_{k!^2} = (1 - \varepsilon)^2 \mathbb{1}_{k!^2}$ and thus

$$\langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B) \mathbf{x} \rangle^{-1} \le a \langle \mathbf{x}, \mathbf{x} \rangle^{-1}.$$
 (4.30)

Using the inequalities (4.29) and (4.30) in (4.22) gives

$$\sup_{X \neq 0} \frac{\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X) - \mathcal{R}(X) \rangle}{\langle X, X \rangle} \le a \sup_{\mathbf{x} \neq 0} \frac{\langle \mathbf{x}, (a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}) \mathbf{x} \rangle}{\langle \mathbf{x}, \mathbf{x} \rangle} = a \|a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}\|, \quad (4.31)$$

where the equality follows from the fact that \mathbf{M} , and thus also $a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}$, is Hermitian by Lemma 4.5.

We now focus on bounding the operator norm of $a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}$. The main idea is that both \mathbf{M}^2 and $\mathbf{N}^T\mathbf{N}$ can be written as a sum of two matrices: a diagonal matrix \mathbf{M}_0 with operator norm 1 that is identical for both \mathbf{M}^2 and $\mathbf{N}^T\mathbf{N}$, and a matrix with small coefficients provided q^ℓ is much larger than certain powers of k!. Using simple norm estimates, $a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}$ can then be written as a sum of matrices with small operator norm.

Inspecting the definition of \mathbf{M} in (4.23), we see that $(\mathbf{M})_{(\pi,\pi),(\pi,\pi)}$ is equal to 1 for any $\pi \in \mathfrak{S}_k$. On the other hand, for a quantum system S with dimension |S| we have $\langle \pi_S, \rho_S \rangle_{\mathbf{n}} = |S|^{-|\pi^{-1}\rho|}$, where $|\sigma|$ denotes the minimum number of terms needed to write $\sigma \in \mathfrak{S}_k$ as a product of transpositions [20]. Thus, $\langle \pi_S, \rho_S \rangle_{\mathbf{n}} \leq |S|^{-1}$ if $\pi \neq \rho$, which implies that $(\mathbf{M})_{(\pi,\rho),(\omega,\chi)} \leq q^{-2\ell}$ if

at least one of π, ρ, ω, χ is distinct from the rest. For example, for $\chi \neq \pi$ we have

$$(\mathbf{M})_{(\pi,\pi),(\pi,\chi)} = \langle \pi_A, \chi_{A^{\ell}} \otimes \pi_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \pi_B, \pi_{B^{\ell}} \otimes \chi_{B^{m-\ell}} \rangle_{\mathbf{n}}$$
(4.32)

$$= \langle \pi_{A^{\ell}}, \chi_{A^{\ell}} \rangle_{n} \langle \pi_{B^{m-\ell}}, \chi_{B^{m-\ell}} \rangle_{n} \tag{4.33}$$

$$\leq q^{-\ell}q^{-(m-\ell)} \tag{4.34}$$

$$\leq q^{-2\ell}.
\tag{4.35}$$

We can thus write

$$\mathbf{M} = \mathbf{M}_0 + q^{-2\ell} \mathbf{M}_1, \tag{4.36}$$

where \mathbf{M}_1 is a $(k!^2 \times k!^2)$ -matrix with each coefficient bounded from above by 1, and \mathbf{M}_0 is a $(k!^2 \times k!^2)$ -matrix with coefficients $(\mathbf{M}_0)_{(\pi,\rho),(\omega,\chi)} = \delta_{\pi,\rho}\delta_{\pi,\omega}\delta_{\pi,\chi}$. In other words, \mathbf{M}_0 is a diagonal matrix with 1's in positions $((\pi,\pi),(\pi,\pi))$ for $\pi \in \mathfrak{S}_k$, and 0's elsewhere. Squaring \mathbf{M} and using (4.36), we then have

$$\mathbf{M}^2 = \left(\mathbf{M}_0 + q^{-2\ell}\mathbf{M}_1\right)^2 \tag{4.37}$$

$$= \mathbf{M}_0^2 + q^{-2\ell} (\mathbf{M}_0 \mathbf{M}_1 + \mathbf{M}_1 \mathbf{M}_0) + q^{-4\ell} \mathbf{M}_1^2$$
(4.38)

$$= \mathbf{M}_0 + 2q^{-2\ell}\mathbf{M}_2 + k!^2q^{-4\ell}\mathbf{M}_3 \tag{4.39}$$

where \mathbf{M}_2 , \mathbf{M}_3 are again matrices with coefficients bounded by 1, and we used that $\mathbf{M}_0^2 = \mathbf{M}_0$. We now apply the same argument to $\mathbf{N}^T \mathbf{N}$: Inspecting the definition of \mathbf{N} in (4.112) reveals that $(\mathbf{N})_{\pi,(\sigma,\tau)}$ is 1 if $(\sigma,\tau) = (\pi,\pi)$, and bounded from above by q^{-m} otherwise. Hence, we write

$$\mathbf{N} = \mathbf{N}_0 + q^{-m} \mathbf{N}_1, \tag{4.40}$$

where \mathbf{N}_0 is a $(k! \times k!^2)$ -matrix with coefficients $(\mathbf{N}_0)_{\pi,(\sigma,\tau)} = \delta_{\pi,\sigma}\delta_{\pi,\tau}$, and \mathbf{N}_1 is a matrix whose coefficients are bounded from above by 1. Note that we have $\mathbf{N}_0^T\mathbf{N}_0 = \mathbf{M}_0$, and hence

$$\mathbf{N}^T \mathbf{N} = (\mathbf{N}_0 + q^{-m} \mathbf{N}_1)^T (\mathbf{N}_0 + q^{-m} \mathbf{N}_1)$$

$$\tag{4.41}$$

$$= \mathbf{M}_0 + q^{-m} \left(\mathbf{N}_1^T \mathbf{N}_0 + \mathbf{N}_0^T \mathbf{N}_1 \right) + q^{-2m} \mathbf{N}_1^T \mathbf{N}_1$$
 (4.42)

$$= \mathbf{M}_0 + 2q^{-m}\mathbf{M}_4 + k!q^{-2m}\mathbf{M}_5, \tag{4.43}$$

with matrices M_4, M_5 whose coefficients are bounded by 1 as before.

Using (4.39) and (4.43), we get the following bound on the operator norm of $a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}$:

$$||a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}|| \tag{4.44}$$

$$= \left\| a\mathbf{M}_0^2 + 2aq^{-2\ell}\mathbf{M}_2 + ak!^2q^{-4\ell}\mathbf{M}_3 - b\mathbf{M}_0 - 2bq^{-m}\mathbf{M}_4 - bk!q^{-2m}\mathbf{M}_5 \right\|$$
(4.45)

$$\leq |a - b| \|\mathbf{M}_0\| + 2aq^{-2\ell} \|\mathbf{M}_2\| + ak!^2 q^{-4\ell} \|\mathbf{M}_3\| + 2bq^{-m} \|\mathbf{M}_4\| + bk! q^{-2m} \|\mathbf{M}_5\|, \tag{4.46}$$

where we used the triangle inequality in the last inequality. We have $\|\mathbf{M}_0\| = 1$ by definition, and \mathbf{M}_i for i = 2, ..., 5 are $(k!^2 \times k!^2)$ -matrices with coefficients bounded above by 1. Using the

norm inequality $||X|| \leq \sqrt{d} \max_{1 \leq i \leq d} \sum_{j=1}^{d} |x_{ij}|$ valid for any $(d \times d)$ -matrix X, we can bound the operator norms of \mathbf{M}_i for $i = 2, \ldots, 5$ by $k! \cdot k!^2 = k!^3$, giving

$$\|\mathbf{L}\| \le a - b + 4ak!^3 q^{-2\ell} + 2ak!^5 q^{-4\ell},\tag{4.47}$$

where we also used $a = (1 - \varepsilon)^{-2} \ge \exp(-\varepsilon') = b$ and the assumption $2\ell \le m$. Note that both $a, b \to 1$ for $q^{2m} \ll k^2$, and hence also $a - b \to 0$ in this limit.

Using the bound on $||a\mathbf{M}^2 - b\mathbf{N}^T\mathbf{N}||$ from (4.47) in (4.29), and substituting the result in (4.22), we arrive at

$$\frac{\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X) - \mathcal{R}(X) \rangle}{\langle X, X \rangle} \le \left(a - b + 4ak!^3 q^{-2\ell} + 2ak!^5 q^{-4\ell} \right) \frac{\langle \mathbf{x}, \mathbf{x} \rangle}{\langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B) \mathbf{x} \rangle}$$
(4.48)

$$\leq a(a-b) + 4a^2k!^3q^{-2\ell} + 2a^2k!^5q^{-4\ell}, \tag{4.49}$$

where we used (4.30) in the second inequality. Since the left-hand side of (4.49) bounds the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{sw})^2$ from above via (4.16) and (4.19),

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{sw})^2 \le a(a-b) + 4a^2k!^3q^{-2\ell} + 2a^2k!^5q^{-4\ell}.$$
 (4.50)

Note that $a=1/(1-\varepsilon)^2 \le 1/(1-2\varepsilon)$. Using the geometric series expansion for a and that $\exp(\varepsilon') \ge 1-\varepsilon'$,

$$a-b \le 1 + \varepsilon(2-\varepsilon) + \varepsilon \frac{(2-\varepsilon)}{(1-\varepsilon)^2} - 1 + \varepsilon'$$
.

Assuming $\varepsilon \leq 1/2$ and $\varepsilon' \leq \varepsilon$ then gives $a - b \leq 9k^2/q^m$, which concludes the proof.

4.2. **Twirl-Crosstwirl.** In this section we consider a different iterative protocol to implement a unitary design on A^kB^k . The protocol alternates between two maps: individual twirls on A and B, and a twirl on the first ℓ qudits within each block A and B (as opposed to a simple swapping of those qudits as in Section 4.1). Note that this protocol is a simplified bipartite version of a much more general multipartite protocol that we discuss later in Section 4.4. In addition to considering P parties, the multipartite version furthermore allows for different numbers m_p of total qudits in each block, and different numbers ℓ_p of qudits participating in the "crosstwirl". However, for the sake of exposing the main proof ideas, we restrict the discussion in this section to a simple bipartite setting where the blocks in A and B consist of the same number m of qudits, and the crosstwirl acts on the same number ℓ of qudits.

We consider the following projections:

$$\mathcal{P} = \mathcal{T}_A \otimes \mathcal{T}_B \tag{4.51}$$

$$Q_{\rm ct} = \mathcal{T}_{A^{\ell}B^{\ell}},\tag{4.52}$$

where A^{ℓ} denotes the first ℓ qudits in a block A_i , and similarly for B^{ℓ} . We also denote by $A^{m-\ell}$ and $B^{m-\ell}$ the remaining $m-\ell$ qudits within A_i and B_i , respectively. We will once again show that iteratively applying \mathcal{P} and \mathcal{Q}_{ct} converges to the full twirl

$$\mathcal{R} = \mathcal{T}_{AB}.\tag{4.53}$$

As before, we have the dominance relations

$$PR = RP = R = RQ_{ct} = Q_{ct}R$$
(4.54)

because of left- and right-invariance of the Haar measure, as well as the relation

$$\operatorname{im} \mathcal{P} \cap \operatorname{im} \mathcal{Q}_{\operatorname{ct}} = \operatorname{im} \mathcal{R} \tag{4.55}$$

which follows from (2.19) and similar arguments as those used to prove (4.55). The alternating projection method hence guarantees that $\lim_{n\to\infty} (\mathcal{Q}_{\operatorname{ct}}\mathcal{P})^n = \mathcal{R}$ as in (4.17), with the convergence speed controlled by the subspace angle

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{ct}}) := \|\mathcal{Q}_{\operatorname{ct}} \mathcal{P} - \mathcal{R}\|_{2 \to 2} = \sup_{X \neq 0} \frac{\|\mathcal{Q}_{\operatorname{ct}} \mathcal{P}(X) - \mathcal{R}(X)\|_{2}}{\|X\|_{2}}.$$
 (4.56)

We derive the following bound on $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{ct}})$.

Proposition 4.4. Let A and B each consist of k blocks of m qudits of local dimension q, and let $\ell < m$ be the number of qudits within each of A and B on which the crosstwirl \mathcal{Q}_{ct} defined in (4.52) acts. We assume that $k \leq \min\{q^{2\ell}, q^m\}$ and $k^2 < 2q^{2\ell}$. Then the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{ct})$ corresponding to the projections \mathcal{P} and \mathcal{Q}_{sw} defined in (4.51) and (4.52), respectively, satisfies

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{ct}})^2$$

$$\leq acgq^{-4\ell k}k!^2\binom{q^{2\ell}+k-1}{k}^2-2abq^{-4mk}k!^2\binom{q^{2m}+k-1}{k}^2+abq^{-4mk}k!^4\binom{q^m+k-1}{k}^4, \tag{4.57}$$

with constants

$$a = \left(1 - \frac{k^2}{2q^m}\right)^{-2} \quad b = \exp\left(-\frac{k^2}{2q^{2m}}\right) \quad c = \left(1 - \frac{k^2}{2q^{2\ell}}\right)^{-1} \quad g = \exp\left(\frac{k^2}{q^{m-\ell}}\right). \tag{4.58}$$

Proposition 4.4 could be converted to a relative error bound using results of Section 3. We forgo that discussion, however, as the bound would be entirely subsumed by the more general multipartite protocol in Proposition 4.8. Instead, we present a proof of Proposition 4.4 here, as it more simply illustrates the subpace angle technique than does the corresponding component of that of Theorem 4.8.

Proof of Proposition 4.4. Using the relations (4.54) together with Lemma 4.1, we can once again rewrite the square of the argument on the right-hand side of (4.56) in terms of the normalized Frobenius inner product $\langle \cdot, \cdot \rangle_n$ defined in (4.21):

$$\sup_{X \neq 0} \frac{\|\mathcal{Q}_{ct}\mathcal{P}(X) - \mathcal{R}(X)\|_{2}^{2}}{\|X\|_{2}^{2}} = \sup_{X \neq 0} \frac{\langle X, \mathcal{P}\mathcal{Q}_{ct}\mathcal{P}(X) - \mathcal{R}(X)\rangle_{n}}{\langle X, X\rangle_{n}},$$
(4.59)

and due to (4.54) we may assume that the supremum in (4.59) is achieved on an operator

$$X = \mathcal{P}(X) = \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, \sigma_A \otimes \tau_B \tag{4.60}$$

for some coefficients $x_{\sigma,\tau} \in \mathbb{C}$, which again follows from Schur-Weyl duality and (2.19). Using Lemma 4.5(i), (iii) and (iv), we can rewrite the right-hand side of (4.59) as

$$\frac{\langle X, \mathcal{PQ}_{ct}\mathcal{P}(X) - \mathcal{R}(X) \rangle_{n}}{\langle X, X \rangle_{n}} = \frac{\langle \mathbf{x}, \left[(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot \left(\mathbf{N}_{\ell}^{T} \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell} \right) - \mathbf{N}^{T} \mathbf{G}_{AB}^{-1} \mathbf{N} \right] \mathbf{x} \rangle}{\langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B}) \mathbf{x} \rangle}, \quad (4.61)$$

where the matrices on the right-hand side have coefficients (cf. Lemma 4.5)

$$(\mathbf{N}_{\ell})_{\pi,(\sigma,\tau)} = \langle \pi_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}} \tag{4.62}$$

$$(\mathbf{N})_{\pi,(\sigma,\tau)} = \langle \pi_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}}. \tag{4.63}$$

The rest of the proof is dedicated to bounding the supremum over $\mathbf{x} \neq 0$ of the expression in (4.61). In a first step, we eliminate the Gram matrices using bounds on their eigenvalues from [20]. We have $-\mathbf{G}_{AB}^{-1} \leq -b\mathbb{1}_{k!}$ with $b = \exp\left(-\frac{k^2}{2q^{2m}}\right)$ by (4.28), and thus

$$-\langle \mathbf{x}, \mathbf{N}^T \mathbf{G}_{AB}^{-1} \mathbf{N} \mathbf{x} \rangle \le -b\langle \mathbf{x}, \mathbf{N}^T \mathbf{N} \mathbf{x} \rangle. \tag{4.64}$$

Second, $\langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B) \mathbf{x} \rangle^{-1} \leq a \langle \mathbf{x}, \mathbf{x} \rangle^{-1}$ with $a = \left(1 - \frac{k^2}{2q^m}\right)^{-2}$ by (4.30). For the matrix $(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell})$, we recall the following properties of the Hadamard product: $X \odot Y \geq 0$ for $X, Y \geq 0$ [25, Thm. 5.2.1] and $(Z - X) \odot Y = Z \odot Y - X \odot Y$. We use these relations together with the following operator inequalities following from [20, Lemma 1]:

$$\mathbf{G}_{\ell}^{-1} \le c \mathbb{1}_{k!} \quad \text{with } c = \left(1 - \frac{k^2}{2q^{2\ell}}\right)^{-1}$$
 (4.65)

$$\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}} \le g \mathbb{1}_{k!^2} \quad \text{with } g = \exp\left(\frac{k^2}{q^{m-\ell}}\right),$$
 (4.66)

and hence

$$(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^{T} \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell}) \le c(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^{T} \mathbf{N}_{\ell})$$
(4.67)

$$\leq cg \mathbb{1}_{k!^2} \odot \left(\mathbf{N}_{\ell}^T \mathbf{N}_{\ell} \right); \tag{4.68}$$

so that

$$\langle \mathbf{x}, (\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell}) \mathbf{x} \rangle \le cg \langle \mathbf{x}, \mathbb{1}_{k!^2} \odot (\mathbf{N}_{\ell}^T \mathbf{N}_{\ell}) \mathbf{x} \rangle. \tag{4.69}$$

Collecting all of these bounds, we obtain

$$\sup_{\mathbf{x}\neq 0} \frac{\left\langle \mathbf{x}, \left[(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot \left(\mathbf{N}_{\ell}^{T} \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell} \right) - \mathbf{N}^{T} \mathbf{G}_{AB}^{-1} \mathbf{N} \right] \mathbf{x} \right\rangle}{\left\langle \mathbf{x}, \left(\mathbf{G}_{A} \otimes \mathbf{G}_{B} \right) \mathbf{x} \right\rangle}$$

$$\leq a \sup_{\mathbf{x}\neq 0} \frac{\left\langle \mathbf{x}, \left(cg \mathbb{1}_{k!^{2}} \odot \mathbf{N}_{\ell}^{T} \mathbf{N}_{\ell} - b \mathbf{N}^{T} \mathbf{N} \right) \mathbf{x} \right\rangle}{\left\langle \mathbf{x}, \mathbf{x} \right\rangle}$$

$$= a \left\| cg \mathbb{1}_{k!^{2}} \odot \mathbf{N}_{\ell}^{T} \mathbf{N}_{\ell} - b \mathbf{N}^{T} \mathbf{N} \right\|.$$

$$(4.70)$$

Note that $\mathbb{1}_{k!^2} \odot \mathbf{N}_{\ell}^T \mathbf{N}_{\ell}$ is a diagonal matrix with the corresponding entries from $\mathbf{N}_{\ell}^T \mathbf{N}_{\ell}$ on its diagonal. Let us thus write $cg\mathbb{1}_{k!^2} \odot \mathbf{N}_{\ell}^T \mathbf{N}_{\ell} - b\mathbf{N}^T \mathbf{N} \equiv \mathbf{D} - \mathbf{K}$ with matrices

$$(\mathbf{D})_{(\pi,\rho),(\sigma,\tau)} = \begin{cases} (cg\mathbf{N}_{\ell}^T\mathbf{N}_{\ell} - b\mathbf{N}^T\mathbf{N})_{(\pi,\rho),(\pi,\rho)} & \text{if } \pi = \sigma \text{ and } \rho = \tau \\ 0 & \text{otherwise;} \end{cases}$$
(4.72)

$$(\mathbf{K})_{(\pi,\rho),(\sigma,\tau)} = \begin{cases} 0 & \text{if } \pi = \sigma \text{ and } \rho = \tau \\ b(\mathbf{N}^T \mathbf{N})_{(\pi,\rho),(\sigma,\tau)} & \text{otherwise.} \end{cases}$$
(4.73)

That is, **D** is a diagonal matrix combining the diagonals of $\mathbf{N}_{\ell}^T \mathbf{N}_{\ell}$ and $\mathbf{N}^T \mathbf{N}$, while **K** is the off-diagonal part of $\mathbf{N}^T \mathbf{N}$; this is a non-negative irreducible³ matrix with zeroes on the diagonal. In terms of these new matrices,

$$\sup_{\mathbf{x}\neq 0} \frac{\left\langle \mathbf{x}, \left[(\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot \left(\mathbf{N}_{\ell}^{T} \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell} \right) - \mathbf{N}^{T} \mathbf{G}_{AB}^{-1} \mathbf{N} \right] \mathbf{x} \right\rangle}{\left\langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B}) \mathbf{x} \right\rangle} \leq a \|\mathbf{D} - \mathbf{K}\|$$
(4.74)

$$\leq a\|\mathbf{D}\| + a\|\mathbf{K}\|. \tag{4.75}$$

We first bound the operator norm of $\|\mathbf{K}\|$. To this end, we recall from above that \mathbf{K} is a non-negative irreducible matrix, and thus the generalized Perron-Frobenius Theorem [24, Ch. 1] implies that $\|\mathbf{K}\| \leq \max_{(\pi,\rho)} \sum_{\sigma,\tau} (\mathbf{K})_{(\pi,\rho),(\sigma,\tau)}$. As before, let (π^*, ρ^*) label the row achieving this maximum. Then we have the following bound:

$$\|\mathbf{K}\| \le \sum_{\sigma,\tau} (\mathbf{K})_{(\pi^*,\rho^*),(\sigma,\tau)} \tag{4.76}$$

$$\leq \sum_{\rho} \sum_{\sigma,\tau} (\mathbf{K})_{(\pi^*,\rho),(\sigma,\tau)} \tag{4.77}$$

$$= b \sum_{\rho} \left(\sum_{\sigma,\tau} (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\sigma,\tau)} - (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\pi^*,\rho)} \right)$$
(4.78)

$$= b \sum_{\rho,\sigma,\tau} (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\sigma,\tau)} - b \sum_{\rho} (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\pi^*,\rho)}$$

$$(4.79)$$

Writing out the first sum gives

$$\sum_{\rho,\sigma,\tau} (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\sigma,\tau)} = \sum_{\rho,\sigma,\tau,\omega} \langle \pi_A^* \otimes \rho_B, \omega_{AB} \rangle_{\mathbf{n}} \langle \omega_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}}$$
(4.80)

$$= \sum_{\omega} \langle \pi_A^*, \omega_A \rangle_{\mathbf{n}} \sum_{\rho} \langle \rho_B, \omega_B \rangle_{\mathbf{n}} \sum_{\sigma} \langle \omega_A, \sigma_A \rangle_{\mathbf{n}} \sum_{\tau} \langle \omega_B, \tau_B \rangle_{\mathbf{n}}$$
(4.81)

³A non-negative $(n \times n)$ -matrix M is called irreducible if the directed graph $G_M = (V, E)$ with vertices $V = \{1, \ldots, n\}$ and edges $E = \{i \to j : (M)_{ij} \neq 0\}$ is strongly connected, i.e., every vertex in G_M can be reached from every other vertex. Since every off-diagonal element of \mathbf{K} is strictly positive, the resulting directed graph $G_{\mathbf{K}}$ is strongly connected, and hence \mathbf{K} is an irreducible matrix. Most statements from the Perron-Frobenius Theorem still apply to a non-negative irreducible matrix, in particular the inequality $\lambda_{\max}(M) \leq \max_i \sum_j (M)_{ij}$ on its largest eigenvalue [24, Ch. 1].

To evaluate the sums in (4.81), we recall from [20] that for any $\omega \in \mathfrak{S}_k$ and $d \geq 1$ we have

$$\sum_{\omega} d^{c(\omega)} = k! \binom{d+k-1}{k}. \tag{4.82}$$

With $d := q^m$, we can use this identity to evaluate the right-most sum in (4.81) as follows:

$$\sum_{\tau} \langle \omega_B, \tau_B \rangle_{\mathbf{n}} = d^{-k} \sum_{\tau} d^{c(\omega^{-1}\tau)} = d^{-k} \sum_{\tau} d^{c(\tau)} = d^{-k} k! \binom{d+k-1}{k}, \tag{4.83}$$

where the second equality holds since $\tau \mapsto \omega^{-1}\tau$ for fixed ω is a bijection on \mathfrak{S}_k . This argument allows us to compute the four sums in (4.81) one by one, starting from the right-most one, giving

$$\sum_{\rho,\sigma,\tau} (\mathbf{N}^T \mathbf{N})_{(\pi^*,\rho),(\sigma,\tau)} = q^{-4mk} k!^4 \binom{q^m + k - 1}{k}^4.$$
 (4.84)

Similarly, for the second sum in (4.79), we get

$$\sum_{\rho} (\mathbf{N}^T \mathbf{N})_{(\pi^*, \rho), (\pi^*, \rho)} = \sum_{\rho, \omega} \langle \pi_A^* \otimes \rho_B, \omega_{AB} \rangle_{\mathbf{n}} \langle \omega_{AB}, \pi_A^* \otimes \rho_B \rangle_{\mathbf{n}}$$
(4.85)

$$= \sum_{\rho,\omega} \langle \pi_A^*, \omega_A \rangle_{\mathbf{n}}^2 \langle \rho_B, \omega_B \rangle_{\mathbf{n}}^2$$
 (4.86)

$$= q^{-4mk} \sum_{\omega} (q^{2m})^{c((\pi^*)^{-1}\omega)} \sum_{\rho} (q^{2m})^{c(\rho^{-1}\omega)}$$
 (4.87)

$$=q^{-4mk}k!^2\binom{q^{2m}+k-1}{k}^2, (4.88)$$

where the second sum in (4.87) can be evaluated by noting that $\rho \mapsto \rho^{-1}\omega$ is a bijection on \mathfrak{S}_k . Using the identities (4.84) and (4.88) in (4.79) yields

$$\|\mathbf{K}\| \le bq^{-4mk}k!^4 \binom{q^m + k - 1}{k}^4 - bq^{-4mk}k!^2 \binom{q^{2m} + k - 1}{k}^2. \tag{4.89}$$

We now bound the operator norm of the diagonal matrix **D**. The entries of **D** are positive: First note that, using $\exp(x) \ge 1 + x$ and $(1 - x)^{-1} \ge 1 + x$ for $x \in [0, 1)$,

$$\frac{cg}{b} = \frac{\exp\left(\frac{k^2}{q^{m-\ell}}\right) \exp\left(\frac{k^2}{2q^{2m}}\right)}{1 - \frac{k^2}{2q^{2\ell}}} \ge \left(1 + \frac{k^2}{q^{m-\ell}}\right) \left(1 + \frac{k^2}{2q^{2m}}\right) \left(1 + \frac{k^2}{2q^{2\ell}}\right) \ge 1,\tag{4.90}$$

and hence $cg \geq b$. Moreover, recall from [20] that for $\sigma, \tau \in \mathfrak{S}_k$ the normalized Frobenius inner product of two permutation operators σ_X, τ_X on a space X is equal to

$$\langle \sigma_X, \tau_X \rangle_{\mathbf{n}} = d^{c(\sigma^{-1}\tau)-k},$$
 (4.91)

where d = |X| is the dimension of X, and $c(\pi)$ denotes the number of cycles in $\pi \in \mathfrak{S}_k$. Since $c(\pi) \leq k$ for any $\pi \in \mathfrak{S}_k$, it then follows that, for two spaces X, Y with $|X| \leq |Y|$,

$$\langle \sigma_X, \tau_X \rangle_{\mathbf{n}} = |X|^{c(\sigma^{-1}\tau)-k} \ge |Y|^{c(\sigma^{-1}\tau)-k} = \langle \sigma_Y, \tau_Y \rangle_{\mathbf{n}}.$$
 (4.92)

For $\pi, \rho \in \mathfrak{S}_k$, we then have

$$cg(\mathbf{N}_{\ell}^T \mathbf{N}_{\ell})_{(\pi,\rho),(\pi,\rho)} \ge b(\mathbf{N}_{\ell}^T \mathbf{N}_{\ell})_{(\pi,\rho),(\pi,\rho)} \tag{4.93}$$

$$= b \sum_{\omega} \langle \pi_{A^{\ell}} \otimes \rho_{B^{\ell}}, \omega_{A^{\ell}B^{\ell}} \rangle_{n} \langle \omega_{A^{\ell}B^{\ell}}, \pi_{A^{\ell}} \otimes \rho_{B^{\ell}} \rangle_{n}$$

$$(4.94)$$

$$= b \sum_{\alpha} \langle \pi_{A^{\ell}}, \omega_{A^{\ell}} \rangle_{\mathbf{n}} \langle \rho_{B^{\ell}}, \omega_{B^{\ell}} \rangle_{\mathbf{n}} \langle \omega_{A^{\ell}}, \pi_{A^{\ell}} \rangle_{\mathbf{n}} \langle \omega_{B^{\ell}}, \rho_{B^{\ell}} \rangle_{\mathbf{n}}$$
(4.95)

$$\geq b \sum_{\omega} \langle \pi_A, \omega_A \rangle_{\mathbf{n}} \langle \rho_B, \omega_B \rangle_{\mathbf{n}} \langle \omega_A, \pi_A \rangle_{\mathbf{n}} \langle \omega_B, \rho_B \rangle_{\mathbf{n}}$$
(4.96)

$$= b \sum_{\omega} \langle \pi_A \otimes \rho_B, \omega_{AB} \rangle_{\mathbf{n}} \langle \omega_{AB}, \pi_A \otimes \rho_B \rangle_{\mathbf{n}}$$

$$(4.97)$$

$$= b(\mathbf{N}^T \mathbf{N})_{(\pi,\rho),(\pi,\rho)},\tag{4.98}$$

where we used (4.92) with $|A^{\ell}| = |B^{\ell}| = q^{\ell} \le q^m = |A| = |B|$ in the inequality.

Thus, $(\mathbf{D})_{(\pi,\rho),(\pi,\rho)} \geq 0$ for all $\pi, \rho \in \mathfrak{S}_k$, and the operator norm $\|\mathbf{D}\|$ is equal to the largest diagonal element. Let this element be $(\mathbf{D})_{(\pi^*,\rho^*),(\pi^*,\rho^*)}$. Then we can bound the operator norm as follows:

$$\|\mathbf{D}\| = (\mathbf{D})_{(\pi^*, \rho^*), (\pi^*, \rho^*)} \tag{4.99}$$

$$\leq \sum_{\rho} (\mathbf{D})_{(\pi^*,\rho),(\pi^*,\rho)} \tag{4.100}$$

$$= \sum_{\rho} (cg \mathbf{N}_{\ell}^T \mathbf{N}_{\ell} - b \mathbf{N}^T \mathbf{N})_{(\pi^*, \rho), (\pi^*, \rho)}$$

$$\tag{4.101}$$

$$= cgq^{-4\ell k}k!^2 {q^{2\ell} + k - 1 \choose k}^2 - bq^{-4mk}k!^2 {q^{2m} + k - 1 \choose k}^2,$$
(4.102)

where we used (4.88) twice with local dimensions q^{ℓ} for \mathbf{N}_{ℓ} and q^{m} for \mathbf{N} .

Together, the bounds (4.102) on $\|\mathbf{D}\|$ and (4.89) on $\|\mathbf{K}\|$ yield

$$a\|\mathbf{D} - \mathbf{K}\|$$

$$\leq a\|\mathbf{D}\| + a\|\mathbf{K}\|$$

$$\leq acgq^{-4\ell k}k!^{2} {q^{2\ell} + k - 1 \choose k}^{2} - 2abq^{-4mk}k!^{2} {q^{2m} + k - 1 \choose k}^{2} + abq^{-4mk}k!^{4} {q^{m} + k - 1 \choose k}^{4}$$

$$(4.103)$$

Substituting this bound subsequently in (4.75), (4.61) and (4.59) finally proves the claim of the proposition.

4.3. Auxiliary lemma for rewriting Frobenius inner products. In this section we prove a lemma that is used repeatedly in the proofs of Propositions 4.2 and 4.4. It expresses the inner products of operators on A^kB^k appearing in the formulas (4.19) and (4.59) for the subspace angle in terms of certain matrices acting on vectors in $\mathbb{C}^{k!^2}$. Part (i) of the lemma generalizes

[20, Eq. (18)]. In the statement and proof of the lemma, we denote by $\langle X, Y \rangle_{\mathbf{n}} = (q^{2m})^{-k} \operatorname{tr}(X^{\dagger}Y)$ the normalized Frobenius product on $\mathcal{L}(A^k B^k)$, and by $\langle \mathbf{x}, \mathbf{y} \rangle$ the standard inner product on $\mathbb{C}^{k!^2}$.

Lemma 4.5. Let k, q, m, ℓ be as in Propositions 4.2 and 4.4, and consider operators $X, Z \in \mathcal{L}(A^k B^k)$ of the form

$$X = \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, \sigma_A \otimes \tau_B \quad \text{for some } x_{\sigma, \tau} \in \mathbb{C}, \tag{4.105}$$

$$Z = \sum_{\sigma, \tau \in \mathfrak{S}_k} z_{\sigma, \tau} \, \sigma_A \otimes \tau_B \quad \text{for some } z_{\sigma, \tau} \in \mathbb{C}. \tag{4.106}$$

Denote by $\mathbf{x}, \mathbf{z} \in \mathbb{C}^{k!^2}$ the vectors with coefficients $(\mathbf{x})_{\sigma,\tau} = x_{\sigma,\tau}$ and $(\mathbf{z})_{\sigma,\tau} = z_{\sigma,\tau}$ for $\sigma, \tau \in \mathfrak{S}_k$, respectively.

- (i) $\langle X, Z \rangle_{\rm n} = \langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B) \mathbf{z} \rangle$.
- (ii) For $k \leq q^m$ we have

$$\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X)\rangle_{n} = \langle \mathbf{x}, \mathbf{M} \left(\mathbf{G}_{A}^{-1} \otimes \mathbf{G}_{B}^{-1} \right) \mathbf{M} \mathbf{x} \rangle,$$
 (4.107)

where **M** is a Hermitian $(k!^2 \times k!^2)$ -matrix with coefficients

$$(\mathbf{M})_{(\pi,\rho),(\omega,\chi)} = \langle \pi_A, \chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \rho_B, \omega_{B^{\ell}} \otimes \chi_{B^{m-\ell}} \rangle_{\mathbf{n}}. \tag{4.108}$$

(iii) For $k \leq \min\{q^{2\ell}, q^m\}$ we have

$$\langle X, \mathcal{P}Q_{\mathrm{ct}}\mathcal{P}(X)\rangle_{\mathrm{n}} = \langle \mathbf{x}, (\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell}) \mathbf{x}\rangle,$$
 (4.109)

where $\mathbf{G}_{\ell} \equiv \mathbf{G}_{A^{\ell}B^{\ell}}$ and \mathbf{N}_{ℓ} is a $(k! \times k!^2)$ -matrix with coefficients

$$(\mathbf{N}_{\ell})_{\pi,(\sigma,\tau)} = \langle \omega_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}}. \tag{4.110}$$

(iv) For $k \leq q^{2m}$ we have

$$\langle X, \mathcal{R}(X) \rangle_{n} = \langle \mathbf{x}, \mathbf{N}^{T} \mathbf{G}_{AB}^{-1} \mathbf{N} \mathbf{x} \rangle,$$
 (4.111)

where **N** is a $(k! \times k!^2)$ -matrix with coefficients

$$(\mathbf{N})_{\pi,(\sigma,\tau)} = \langle \pi_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}}. \tag{4.112}$$

Proof of Lemma 4.5(i). Using the assumptions on the operators X and Z, we calculate:

$$\langle X, Z \rangle_{\rm n} = \frac{1}{q^{2mk}} \operatorname{tr}(X^{\dagger} Z)$$
 (4.113)

$$= \frac{1}{q^{2mk}} \sum_{\sigma,\tau,\pi,\rho \in \mathfrak{S}_k} x_{\sigma,\tau}^* z_{\pi,\rho} \operatorname{tr}\left((\sigma_A \otimes \tau_B)^{\dagger} (\pi_A \otimes \rho_B) \right)$$
(4.114)

$$= \frac{1}{q^{2mk}} \sum_{\sigma,\tau,\pi,\rho \in \mathfrak{S}_k} x_{\sigma,\tau}^* z_{\pi,\rho} \operatorname{tr}\left(\sigma_A^{\dagger} \pi_A\right) \operatorname{tr}\left(\tau_B^{\dagger} \rho_B\right)$$
(4.115)

$$= \sum_{\sigma,\tau \in \mathfrak{S}_k} x_{\sigma,\tau}^* \sum_{\pi,\rho \in \mathfrak{S}_k} (\mathbf{G}_A \otimes \mathbf{G}_B)_{(\sigma,\tau),(\pi,\rho)} z_{\pi,\rho}$$

$$(4.116)$$

$$= \langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B) \mathbf{z} \rangle, \tag{4.117}$$

where we used the definition $(\mathbf{G}_A)_{\sigma,\pi} = \frac{1}{q^{mk}} \operatorname{tr}(\sigma_A^{\dagger} \pi_A)$ for the elements of the Gram matrix \mathbf{G}_A following [20], and similarly for \mathbf{G}_B .

Proof of Lemma 4.5(ii). To compute the operator $\mathcal{PQ}_{sw}\mathcal{P}(X)$, recall that we have $X = \mathcal{P}(X)$ by assumption, and $\mathcal{Q}_{sw} = SWAP_{\ell} \circ \mathcal{P} \circ SWAP_{\ell}$ with $SWAP_{\ell}$ defined in (4.9). The action of $SWAP_{\ell}$ on X is equal to

$$SWAP_{\ell}(X) = \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} SWAP_{\ell} (\sigma_A \otimes \tau_B)$$
(4.118)

$$= \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, \tau_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \otimes \sigma_{B^{\ell}} \otimes \tau_{B^{m-\ell}}, \tag{4.119}$$

where we write $\tau_{A^{\ell}}$ for the permutation operator corresponding to $\tau \in \mathfrak{S}_k$ acting on the first $\ell \leq m$ qudits within each of the k blocks (of size m each) in A. Similarly, $\sigma_{A^{m-\ell}}$ acts on the remaining $m-\ell$ qudits within each block. The same conventions hold for the blocks in B.

Another application of the projection $\mathcal{P} = \mathcal{T}_A \otimes \mathcal{T}_B$ to the operator in (4.119) gives

$$\mathcal{P} \circ \text{SWAP}_{\ell} \circ \mathcal{P}(X) = \sum_{\sigma, \tau \in \mathfrak{S}_{k}} x_{\sigma, \tau} \, \mathcal{T}_{A} \left(\tau_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \right) \otimes \mathcal{T}_{B} \left(\sigma_{B^{\ell}} \otimes \tau_{B^{m-\ell}} \right) \tag{4.120}$$

$$= \sum_{\sigma,\tau \in \mathfrak{S}_k} x_{\sigma,\tau} \sum_{\omega \in \mathfrak{S}_k} a_{\omega}^{\sigma,\tau} \,\omega_A \otimes \sum_{\chi \in \mathfrak{S}_k} a_{\chi}^{\tau,\sigma} \,\chi_B, \tag{4.121}$$

$$= \sum_{\omega, \chi \in \mathfrak{S}_k} \left(\sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, a_{\omega}^{\sigma, \tau} \, a_{\chi}^{\tau, \sigma} \right) \omega_A \otimes \chi_B \tag{4.122}$$

$$\equiv \sum_{\omega,\chi \in \mathfrak{S}_k} y_{\omega,\chi} \,\omega_A \otimes \chi_B,\tag{4.123}$$

where the second line uses the fact that the twirl \mathcal{T}_A projects an operator onto the span of permutation operators $\{\omega_A \colon \omega \in \mathfrak{S}_k\}$, and hence $\mathcal{T}_A (\tau_{A^\ell} \otimes \sigma_{A^{m-\ell}})$ can be expanded as a linear combination of these operators with coefficients $a_\omega^{\sigma,\tau}$. The same argument applies to $\mathcal{T}_B (\sigma_{B^\ell} \otimes \tau_{B^{m-\ell}})$ which defines the coefficients $a_\chi^{\tau,\sigma}$. In the last line we defined $y_{\omega,\chi} \coloneqq \sum_{\sigma,\tau \in \mathfrak{S}_k} x_{\sigma,\tau} a_\omega^{\sigma,\tau} a_\chi^{\tau,\sigma}$.

Since $\mathcal{PQ}_{sw}\mathcal{P} = \mathcal{P} \circ SWAP_{\ell} \circ \mathcal{P} \circ SWAP_{\ell} \circ \mathcal{P}$, we repeat the above argument by applying $\mathcal{P} \circ SWAP_{\ell}$ once again to the operator in (4.123), giving

$$\mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X) = \mathcal{P} \circ SWAP_{\ell} \left(\mathcal{P} \circ SWAP_{\ell} \circ \mathcal{P}(X) \right) \tag{4.124}$$

$$= \sum_{\omega, \chi \in \mathfrak{S}_{L}} y_{\omega, \chi} \mathcal{P} \circ \text{SWAP}_{\ell} (\omega_{A} \otimes \chi_{B})$$

$$(4.125)$$

$$= \sum_{\pi,\rho\in\mathfrak{S}_k} \sum_{\omega,\chi\in\mathfrak{S}_k} y_{\omega,\chi} \, a_{\pi}^{\omega,\chi} \, a_{\rho}^{\chi,\omega} \, \pi_A \otimes \rho_B \tag{4.126}$$

$$= \sum_{\pi,\rho\in\mathfrak{S}_k} z_{\pi,\rho} \,\pi_A \otimes \rho_B,\tag{4.127}$$

where we defined $z_{\pi,\rho} = \sum_{\omega,\chi \in \mathfrak{S}_k} y_{\omega,\chi} a_{\pi}^{\omega,\chi} a_{\rho}^{\chi,\omega}$. Defining a vector **z** with components $(\mathbf{z})_{\pi,\rho} = z_{\pi,\rho}$ for $\pi, \rho \in \mathfrak{S}_k$, part (i) of the lemma then gives

$$\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X)\rangle_{n} = \langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B}) \mathbf{z} \rangle.$$
 (4.128)

We now express the vector \mathbf{z} with coefficients $(\mathbf{z})_{\pi,\rho} = z_{\pi,\rho} = \sum_{\omega,\chi\in\mathfrak{S}_k} y_{\omega,\chi} a_{\pi}^{\omega,\chi} a_{\rho}^{\chi,\omega}$ in terms of \mathbf{x} . Using $y_{\omega,\chi} \coloneqq \sum_{\sigma,\tau\in\mathfrak{S}_k} x_{\sigma,\tau} a_{\omega}^{\sigma,\tau} a_{\chi}^{\tau,\sigma}$ from (4.123), the coefficients $z_{\pi,\rho}$ are given by

$$z_{\pi,\rho} = \sum_{\omega,\chi \in \mathfrak{S}_k} y_{\omega,\chi} \, a_{\pi}^{\omega,\chi} \, a_{\rho}^{\chi,\omega} \tag{4.129}$$

$$= \sum_{\omega, \chi, \sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, a_{\omega}^{\sigma, \tau} \, a_{\chi}^{\tau, \sigma} \, a_{\pi}^{\omega, \chi} \, a_{\rho}^{\chi, \omega} \tag{4.130}$$

$$= \sum_{\sigma, \tau \in \mathfrak{S}_{k}} \left(\sum_{\omega, \chi \in \mathfrak{S}_{k}} a_{\omega}^{\sigma, \tau} a_{\chi}^{\tau, \sigma} a_{\pi}^{\omega, \chi} a_{\rho}^{\chi, \omega} \right) x_{\sigma, \tau} \tag{4.131}$$

Our goal is to interpret the sum in parentheses as a $(k!^2 \times k!^2)$ -matrix with coefficients indexed by pairs of permutations $((\pi, \rho), (\sigma, \tau))$. Those coefficients can in turn be expressed in terms of the matrix **M** in the statement of the lemma and the Gram matrices **G**.

To this end, recall from the calculation leading to (4.123) that the $a_{\pi}^{\omega,\chi}$ are defined via

$$\mathcal{T}_A(\chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}}) = \sum_{\pi \in \mathfrak{S}_k} a_{\pi}^{\omega, \chi} \pi_A. \tag{4.132}$$

We take the inner product with π'_A on both sides. For the left-hand side, we have

$$\left\langle \pi_A', \mathcal{T}_A(\chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}}) \right\rangle_{\mathbf{n}} = \left\langle \mathcal{T}_A(\pi_A'), \chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}} \right\rangle_{\mathbf{n}} = \left\langle \pi_A', \chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}} \right\rangle_{\mathbf{n}} =: v_{\pi'}^{\omega, \chi}, \quad (4.133)$$

since π'_A is invariant under \mathcal{T}_A . Hence,

$$v_{\pi'}^{\omega,\chi} = \left\langle \pi'_A, \mathcal{T}_A(\chi_{A^{\ell}} \otimes \omega_{A^{m-\ell}}) \right\rangle_{\mathbf{n}} = \sum_{\pi \in \mathfrak{S}_k} a_{\pi}^{\omega,\chi} \left\langle \pi'_A, \pi_A \right\rangle_{\mathbf{n}} = \sum_{\pi \in \mathfrak{S}_k} (\mathbf{G}_A)_{\pi',\pi} a_{\pi}^{\omega,\chi}, \tag{4.134}$$

or equivalently $\mathbf{v}^{\omega,\chi} = \mathbf{G}_A \mathbf{a}^{\omega,\chi}$ where $(\mathbf{v}^{\omega,\chi})_{\pi} = v_{\pi}^{\omega,\chi}$ and $(\mathbf{a}^{\omega,\chi})_{\pi} = a_{\pi}^{\omega,\chi}$, and the same relation holds with \mathbf{G}_B .

The Gram matrices \mathbf{G}_A and \mathbf{G}_B are invertible if and only if $k \leq q^m$ [20, Lemma 1], in which case $\mathbf{a}^{\omega,\chi} = \mathbf{G}_A^{-1} \mathbf{v}^{\omega,\chi}$. Recall that we try to interpret the sum in parentheses in (4.131) as a

⁴Since |A| = |B| it is not strictly necessary to distinguish between \mathbf{G}_A and \mathbf{G}_B , but it does help in parsing the somewhat cumbersome expressions appearing in the sums.

matrix with coefficients indexed by $((\pi, \rho), (\sigma, \tau))$. Using the formula for $\mathbf{a}^{\omega, \chi}$, we compute:

$$\sum_{\omega,\chi\in\mathfrak{S}_{k}} a_{\omega}^{\sigma,\tau} a_{\chi}^{\tau,\sigma} a_{\pi}^{\omega,\chi} a_{\rho}^{\chi,\omega} \tag{4.135}$$

$$= \sum_{\omega,\chi} (\mathbf{G}_A^{-1} \mathbf{v}^{\sigma,\tau})_{\omega} (\mathbf{G}_B^{-1} \mathbf{v}^{\tau,\sigma})_{\chi} (\mathbf{G}_A^{-1} \mathbf{v}^{\omega,\chi})_{\pi} (\mathbf{G}_B^{-1} \mathbf{v}^{\chi,\omega})_{\rho}$$

$$(4.136)$$

$$= \sum_{\omega,\chi} \sum_{\omega',\chi'\pi',\rho'} (\mathbf{G}_{A}^{-1})_{\omega,\omega'} v_{\omega'}^{\sigma,\tau} (\mathbf{G}_{B}^{-1})_{\chi,\chi'} v_{\chi'}^{\tau,\sigma} (\mathbf{G}_{A}^{-1})_{\pi,\pi'} v_{\pi'}^{\omega,\chi} (\mathbf{G}_{B}^{-1})_{\rho,\rho'} v_{\rho'}^{\chi,\omega}$$
(4.137)

$$= \sum_{\pi',\rho'} (\mathbf{G}_A^{-1})_{\pi,\pi'} (\mathbf{G}_B^{-1})_{\rho,\rho'} \sum_{\omega,\chi} v_{\pi'}^{\omega,\chi} v_{\rho'}^{\chi,\omega} \sum_{\omega',\chi'} (\mathbf{G}_A^{-1})_{\omega,\omega'} (\mathbf{G}_B^{-1})_{\chi,\chi'} v_{\omega'}^{\sigma,\tau} v_{\chi'}^{\tau,\sigma}$$

$$(4.138)$$

$$= \sum_{\pi',\rho'} (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1})_{(\pi,\rho),(\pi',\rho')} \sum_{\omega,\chi} v_{\pi'}^{\omega,\chi} v_{\rho'}^{\chi,\omega} \sum_{\omega',\chi'} (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1})_{(\omega,\chi),(\omega',\chi')} v_{\omega'}^{\sigma,\tau} v_{\chi'}^{\tau,\sigma}. \tag{4.139}$$

Defining a matrix **M** with coefficients $(\mathbf{M})_{(\pi,\rho),(\omega,\chi)} = v_{\pi}^{\omega,\chi} v_{\rho}^{\chi,\omega}$, we thus see that

$$\sum_{\omega,\chi\in\mathfrak{S}_k} a_{\omega}^{\sigma,\tau} a_{\chi}^{\tau,\sigma} a_{\pi}^{\omega,\chi} a_{\rho}^{\chi,\omega} = \left((\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1}) \mathbf{M} (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1}) \mathbf{M} \right)_{(\pi,\rho),(\sigma,\tau)}, \tag{4.140}$$

and together with (4.131) we obtain the vector equation

$$\mathbf{z} = (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1}) \mathbf{M} (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1}) \mathbf{M} \mathbf{x}. \tag{4.141}$$

Using this in (4.128) finally gives

$$\langle X, \mathcal{P}\mathcal{Q}_{sw}\mathcal{P}(X)\rangle_{n} = \langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B}) \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{M}(\mathbf{G}_{A}^{-1} \otimes \mathbf{G}_{B}^{-1}) \mathbf{M} \mathbf{x} \rangle,$$
 (4.142)

which concludes the proof of Lemma 4.5(ii).

Proof of Lemma 4.5(iii). Applying Q_{ct} to X in (4.60) gives an operator

$$Q_{\rm ct}(X) = Q_{\rm ct}(\mathcal{P}(X)) = \sum_{\sigma, \tau \in \mathfrak{S}_k} x_{\sigma, \tau} \, Q_{\rm ct}(\sigma_A \otimes \tau_B) \tag{4.143}$$

$$= \sum_{\sigma,\tau,\omega\in\mathfrak{S}_k} x_{\sigma,\tau} y_{\omega}^{\sigma,\tau} \,\omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \otimes \omega_{B^{\ell}} \otimes \tau_{B^{m-\ell}}, \tag{4.144}$$

where we define the $y_{\omega}^{\sigma,\tau}$ through

$$Q_{\rm ct}(\sigma_{A^{\ell}} \otimes \tau_{B^{\ell}}) = \sum_{\omega} y_{\omega}^{\sigma,\tau} \omega_{A^{\ell}B^{\ell}}. \tag{4.145}$$

Applying \mathcal{P} once more to this operator, we then obtain

$$\mathcal{PQ}_{\mathrm{ct}}\mathcal{P}(X) = \sum_{\sigma,\tau,\omega\in\mathfrak{S}_k} x_{\sigma,\tau} \, y_{\omega}^{\sigma,\tau} \, \mathcal{P}(\omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \otimes \omega_{B^{\ell}} \otimes \tau_{B^{m-\ell}}) \tag{4.146}$$

$$= \sum_{\sigma, \tau, \omega \in \mathfrak{S}_{k}} x_{\sigma, \tau} y_{\omega}^{\sigma, \tau} \mathcal{T}_{A}(\omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}}) \otimes \mathcal{T}_{B}(\omega_{B^{\ell}} \otimes \tau_{B^{m-\ell}})$$
(4.147)

$$= \sum_{\sigma,\tau,\omega,\pi,\rho\in\mathfrak{S}_k} x_{\sigma,\tau} y_{\omega}^{\sigma,\tau} z_{\pi}^{\omega,\sigma} z_{\rho}^{\omega,\tau} \pi_A \otimes \rho_B$$
 (4.148)

$$=\sum_{\pi,\rho}\bar{x}_{\pi,\rho}\pi_A\otimes\rho_B,\tag{4.149}$$

where the $z_{\pi}^{\omega,\sigma}$ in the third line are defined through

$$\mathcal{T}_X(\omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}}) = \sum_{\pi} z_{\pi}^{\omega, \sigma} \pi_X \quad \text{for } X \in \{A, B\},$$
 (4.150)

and in the fourth line we defined coefficients $\bar{x}_{\pi,\rho} = \sum_{\sigma,\tau,\omega} x_{\sigma,\tau} y_{\omega}^{\sigma,\tau} z_{\pi}^{\omega,\sigma} z_{\rho}^{\omega,\tau}$.

For the vectors $\mathbf{x}, \bar{\mathbf{x}} \in \mathbb{C}^{k!^2}$ with coefficients $(\mathbf{x})_{\pi,\rho} = x_{\pi,\rho}$ and $(\bar{\mathbf{x}})_{(\pi,\rho)} = \bar{x}_{\pi,\rho}$, part (i) of the lemma then yields

$$\langle X, \mathcal{P}Q_{\mathrm{ct}}\mathcal{P}(X)\rangle_{\mathrm{n}} = \langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B)\bar{\mathbf{x}}\rangle,$$
 (4.151)

and introducing the $(k!^2 \times k!^2)$ -matrix **L** with coefficients $(\mathbf{L})_{(\pi,\rho),(\sigma,\tau)} = \sum_{\omega} y_{\omega}^{\sigma,\tau} z_{\mu}^{\omega,\sigma} z_{\rho}^{\omega,\tau}$, we see that $\bar{\mathbf{x}} = \mathbf{L} \cdot \mathbf{x}$. Hence, we can rewrite the right-hand side of (4.151) as

$$\langle X, \mathcal{PQ}_{ct}\mathcal{P}(X)\rangle_{n} = \langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B})\bar{\mathbf{x}}\rangle = \langle \mathbf{x}, (\mathbf{G}_{A} \otimes \mathbf{G}_{B})\mathbf{L}\mathbf{x}\rangle.$$
 (4.152)

We continue manipulating this expression by rewriting the coefficients $z_{\pi}^{\omega,\sigma}$ and $y_{\omega}^{\sigma,\tau}$ appearing in **L**. For the $z_{\pi}^{\omega,\sigma}$, we take the inner product with π'_A on both sides of (4.150) and use the fact that π'_A is invariant under the twirl \mathcal{T}_A :

$$v_{\pi'}^{\omega,\sigma} := \langle \pi'_A, \omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \rangle_{p} = \langle \pi'_A, \mathcal{T}_A(\omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}}) \rangle_{p} \tag{4.153}$$

$$= \sum_{\pi} z_{\pi}^{\omega,\sigma} \left\langle \pi_A', \pi_A \right\rangle_{\mathbf{n}} \tag{4.154}$$

$$= \sum_{\pi} z_{\pi}^{\omega,\sigma}(\mathbf{G}_A)_{\pi',\pi}. \tag{4.155}$$

Defining vectors $\mathbf{v}^{\omega,\sigma}, \mathbf{z}^{\omega,\sigma} \in \mathbb{C}^{k!}$ with coefficients $(\mathbf{v}^{\omega,\sigma})_{\pi} = v_{\pi}^{\omega,\sigma}$ and $(\mathbf{z}^{\omega,\sigma})_{\pi} = z_{\pi}^{\omega,\sigma}$, we can rewrite the above equation as $\mathbf{v}^{\omega,\sigma} = \mathbf{G}_A \mathbf{z}^{\omega,\sigma}$, or $\mathbf{z}^{\omega,\sigma} = \mathbf{G}_A^{-1} \mathbf{v}^{\omega,\sigma}$ if $k \leq q^m$ such that \mathbf{G}_A is invertible [20, Lemma 1]. The same assertion also holds for \mathbf{G} . Since also $k \leq q^{2\ell}$ by assumption, an analogous argument applied to the coefficients $y_{\omega}^{\sigma,\tau}$ defined in (4.145) shows that $\mathbf{y}^{\sigma,\tau} = \mathbf{G}_{\ell}^{-1} \mathbf{w}^{\sigma,\tau}$. Here we defined the Gram matrix $\mathbf{G}_{\ell} \coloneqq \mathbf{G}_{A^{\ell}B^{\ell}}$ and vectors $\mathbf{y}^{\sigma,\tau}, \mathbf{w}^{\sigma,\tau} \in \mathbb{C}^{k!}$ with coefficients $(\mathbf{y}^{\sigma,\tau})_{\pi} = y_{\pi}^{\sigma,\tau}$ and

$$(\mathbf{w}^{\sigma,\tau})_{\pi} = \langle \pi_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}} =: w_{\pi}^{\sigma,\tau}. \tag{4.156}$$

Substituting these expressions in the definition of (L) above, we then obtain:

$$(\mathbf{L})_{(\pi,\rho),(\sigma,\tau)} = \sum_{\omega} y_{\omega}^{\sigma,\tau} z_{\pi}^{\omega,\sigma} z_{\rho}^{\omega,\tau}$$

$$(4.157)$$

$$= \sum_{\omega} (\mathbf{G}_{\ell}^{-1} \mathbf{w}^{\sigma,\tau})_{\omega} (\mathbf{G}_{A}^{-1} \mathbf{v}^{\omega,\sigma})_{\pi} (\mathbf{G}_{B}^{-1} \mathbf{v}^{\omega,\tau})_{\rho}$$

$$(4.158)$$

$$= \sum_{\omega,\omega',\pi',\rho'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} w_{\omega'}^{\sigma,\tau} (\mathbf{G}_{A}^{-1})_{\pi,\pi'} v_{\pi'}^{\omega,\sigma} (\mathbf{G}_{B}^{-1})_{\rho,\rho'} v_{\rho'}^{\omega,\tau}$$

$$(4.159)$$

$$= \sum_{\omega,\omega',\pi',\rho'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} w_{\omega'}^{\sigma,\tau} v_{\pi'}^{\omega,\sigma} v_{\rho'}^{\omega,\tau} (\mathbf{G}_{A}^{-1} \otimes \mathbf{G}_{B}^{-1})_{(\pi,\rho),(\pi',\rho')}$$
(4.160)

$$= \sum_{\pi',\rho'} (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1})_{(\pi,\rho),(\pi',\rho')} \sum_{\omega,\omega'} (\mathbf{G}_\ell^{-1})_{\omega,\omega'} v_{\pi'}^{\omega,\sigma} v_{\rho'}^{\omega,\tau} w_{\omega'}^{\sigma,\tau}, \tag{4.161}$$

which we can rewrite as a matrix product

$$\mathbf{L} = (\mathbf{G}_A^{-1} \otimes \mathbf{G}_B^{-1})\overline{\mathbf{L}} \tag{4.162}$$

where $\overline{\mathbf{L}}$ is a $(k!^2 \times k!^2)$ -matrix with coefficients

$$(\overline{\mathbf{L}})_{(\pi,\rho),(\sigma,\tau)} = \sum_{\omega,\omega'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} v_{\pi}^{\omega,\sigma} v_{\rho}^{\omega,\tau} w_{\omega'}^{\sigma,\tau}. \tag{4.163}$$

Plugging in the definitions $v_{\pi}^{\omega,\sigma} = \langle \pi_A, \omega_{A^{\ell}} \otimes \sigma_{A^{m-\ell}} \rangle_{\mathbf{n}}$ and $w_{\omega'}^{\sigma,\tau} = \langle \omega'_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}}$, we obtain

$$(\overline{\mathbf{L}})_{(\pi,\rho),(\sigma,\tau)} \tag{4.164}$$

$$= \sum_{\omega,\omega'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} v_{\pi}^{\omega,\sigma} v_{\rho}^{\omega,\tau} w_{\omega'}^{\sigma,\tau}$$

$$(4.165)$$

$$= \sum_{\omega,\omega'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} \langle \pi_A, \omega_{A\ell} \otimes \sigma_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \rho_B, \omega_{B\ell} \otimes \tau_{B^{m-\ell}} \rangle_{\mathbf{n}} \langle \omega'_{A\ell B\ell}, \sigma_{A\ell} \otimes \tau_{B\ell} \rangle_{\mathbf{n}}$$
(4.166)

$$= \sum_{\omega,\omega'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} \langle \pi_{A^{\ell}}, \omega_{A^{\ell}} \rangle_{\mathbf{n}} \langle \pi_{A^{m-\ell}}, \sigma_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \rho_{B^{\ell}}, \omega_{B^{\ell}} \rangle_{\mathbf{n}} \langle \rho_{B^{m-\ell}}, \tau_{B^{m-\ell}} \rangle_{\mathbf{n}} \langle \omega'_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}}$$

$$(4.167)$$

$$= \langle \pi_{A^{m-\ell}}, \sigma_{A^{m-\ell}} \rangle_{\mathbf{n}} \langle \rho_{B^{m-\ell}}, \tau_{B^{m-\ell}} \rangle_{\mathbf{n}} \sum_{\omega} \langle \pi_{A^{\ell}} \otimes \rho_{B^{\ell}}, \omega_{A^{\ell}B^{\ell}} \rangle_{\mathbf{n}} \sum_{\omega'} (\mathbf{G}_{\ell}^{-1})_{\omega,\omega'} \langle \omega'_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}}$$

$$(4.168)$$

$$= (\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}})_{(\pi,\rho),(\sigma,\tau)} \left(\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell} \right)_{(\pi,\rho),(\sigma,\tau)}, \tag{4.169}$$

where in the last step we defined the $(k! \times k!^2)$ -matrix \mathbf{N}_{ℓ} with coefficients

$$(\mathbf{N}_{\ell})_{\omega,(\sigma,\tau)} = \langle \omega_{A^{\ell}B^{\ell}}, \sigma_{A^{\ell}} \otimes \tau_{B^{\ell}} \rangle_{\mathbf{n}}. \tag{4.170}$$

We see that (4.169) describes the coefficients of a Hadamard product, and thus

$$\overline{\mathbf{L}} = (\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell}). \tag{4.171}$$

Substituting this expression in (4.162) and then (4.152), we finally get

$$\langle X, \mathcal{P}Q_{\mathrm{ct}}\mathcal{P}(X)\rangle_{\mathrm{n}} = \langle \mathbf{x}, (\mathbf{G}_A \otimes \mathbf{G}_B)\mathbf{L}\mathbf{x}\rangle$$
 (4.172)

$$= \langle \mathbf{x}, \overline{\mathbf{L}} \mathbf{x} \rangle \tag{4.173}$$

$$= \langle \mathbf{x}, (\mathbf{G}_{A^{m-\ell}} \otimes \mathbf{G}_{B^{m-\ell}}) \odot (\mathbf{N}_{\ell}^T \mathbf{G}_{\ell}^{-1} \mathbf{N}_{\ell}) \mathbf{x} \rangle, \tag{4.174}$$

which concludes the proof of Lemma 4.5(iii).

Proof of Lemma 4.5(iv). Recall once more that we may assume $X = \mathcal{P}(X) = \sum_{\sigma,\tau \in \mathfrak{S}_k} x_{\sigma,\tau} \sigma_A \otimes \tau_B$ for some $x_{\sigma,\tau} \in \mathbb{C}$. For the full twirl $\mathcal{R} = \mathcal{T}_{AB}$, we then have

$$\mathcal{R}(X) = \sum_{\sigma,\tau} x_{\sigma,\tau} \mathcal{R}(\sigma_A \otimes \tau_B) = \sum_{\sigma,\tau} x_{\sigma,\tau} \sum_{\pi} r_{\pi}^{\sigma,\tau} \pi_{AB}$$
 (4.175)

for some coefficients $r_{\pi}^{\sigma,\tau} \in \mathbb{C}$, and

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \sum_{\sigma, \tau, \sigma', \tau', \pi} x_{\sigma', \tau'}^* x_{\sigma, \tau} r_{\pi}^{\sigma, \tau} \left\langle \sigma_A' \otimes \tau_B', \pi_{AB} \right\rangle_{\mathbf{n}}$$
(4.176)

$$= \sum_{\sigma,\tau,\sigma',\tau',\pi} x_{\sigma',\tau'}^* x_{\sigma,\tau} r_{\pi}^{\sigma,\tau} t_{\pi}^{\sigma',\tau'}, \tag{4.177}$$

where we defined $t_{\pi}^{\sigma,\tau} := \langle \pi_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}}$.

Note that $\mathcal{T}_{AB}(\pi_{AB}) = \pi_{AB}$ for any $\pi \in \mathfrak{S}_k$, and hence

$$t_{\pi}^{\sigma,\tau} = \langle \pi_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}} = \langle \mathcal{T}(\pi_{AB}), \sigma_A \otimes \tau_B \rangle_{\mathbf{n}} = \langle \pi_{AB}, \mathcal{T}_{AB}(\sigma_A \otimes \tau_B) \rangle_{\mathbf{n}}. \tag{4.178}$$

Thus, we may express $r_{\pi}^{\sigma,\tau}$ in terms of $t_{\pi}^{\sigma,\tau}$ as

$$t_{\pi}^{\sigma,\tau} = \langle \pi_{AB}, \mathcal{T}_{AB}(\sigma_A \otimes \tau_B) \rangle \tag{4.179}$$

$$= \sum_{\pi'} r_{\pi'}^{\sigma,\tau} \langle \pi_{AB}, \pi'_{AB} \rangle \tag{4.180}$$

$$= \sum_{\pi'} (\mathbf{G}_{AB})_{\pi,\pi'} r_{\pi'}^{\sigma,\tau}. \tag{4.181}$$

Defining vectors $\mathbf{t}^{\sigma,\tau}, \mathbf{r}^{\sigma,\tau} \in \mathbb{C}^{k!}$ with $(\mathbf{t}^{\sigma,\tau})_{\pi} = t_{\pi}^{\sigma,\tau}$ and $(\mathbf{r}^{\sigma,\tau})_{\pi} = r_{\pi'}^{\sigma,\tau}$, the identity (4.181) can be rewritten as the vector equation $\mathbf{t}^{\sigma,\tau} = \mathbf{G}_{AB}\mathbf{r}^{\sigma,\tau}$, which is equivalent to $\mathbf{r}^{\sigma,\tau} = \mathbf{G}_{AB}^{-1}\mathbf{t}^{\sigma,\tau}$ in the regime $k \leq q^{2m}$ such that \mathbf{G}_{AB} is invertible [20, Lemma 1].

Using $\mathbf{r}^{\sigma,\tau} = \mathbf{G}_{AB}^{-1} \mathbf{t}^{\sigma,\tau}$ in (4.177), we obtain

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \sum_{\sigma, \tau, \sigma', \tau', \pi} x_{\sigma', \tau'}^* x_{\sigma, \tau} r_{\pi}^{\sigma, \tau} t_{\pi}^{\sigma', \tau'}$$

$$\tag{4.182}$$

$$= \sum_{\sigma,\tau,\sigma',\tau',\pi} x_{\sigma',\tau'}^* x_{\sigma,\tau} (\mathbf{G}_{AB}^{-1} \mathbf{t}^{\sigma,\tau})_{\pi} t_{\pi}^{\sigma',\tau'}$$

$$\tag{4.183}$$

$$= \sum_{\sigma,\tau,\sigma',\tau',\pi,\pi'} x_{\sigma',\tau'}^* x_{\sigma,\tau} (\mathbf{G}_{AB}^{-1})_{\pi,\pi'} t_{\pi'}^{\sigma,\tau} t_{\pi'}^{\sigma',\tau'}$$
(4.184)

$$= \sum_{\sigma',\tau'} x_{\sigma',\tau'}^* \sum_{\pi} t_{\pi}^{\sigma',\tau'} \sum_{\pi'} (\mathbf{G}_{AB}^{-1})_{\pi,\pi'} \sum_{\sigma,\tau} t_{\pi'}^{\sigma,\tau} x_{\sigma,\tau}. \tag{4.185}$$

Defining the $(k! \times k!^2)$ -matrix **N** with coefficients $(\mathbf{N})_{\pi,(\sigma,\tau)} = t_{\pi}^{\sigma,\tau} = \langle \pi_{AB}, \sigma_A \otimes \tau_B \rangle_{\mathbf{n}}$, the last equation can be written as

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \langle \mathbf{x}, \mathbf{N}^T \mathbf{G}_{AB}^{-1} \mathbf{N} \mathbf{x} \rangle,$$
 (4.186)

which concludes the proof of Lemma 4.5(iv).

4.4. **Multipartite setting.** In this section we analyze a multipartite version of the Twirl-Crosstwirl protocol from Section 4.2, which is depicted in Figure 1. We consider P parties A_1, \ldots, A_P that are indexed by a lower-case $p = 1, \ldots, P$. As before, each party A_p consists of K subsystems $A_{p,1}, \ldots, A_{p,K}$, which we index by a lower-case $k = 1, \ldots, K$. To be as general as possible we allow for differing dimensions among the P parties. That is, each $A_{p,k}$ consists of m_p qudits of local dimension q so that $|A_{p,k}| = q^{m_p}$.

We now consider a protocol that alternates between independent twirls across the 'rows' and twirls of the first $\ell_p \leq m_p$ qudits within each block $A_{p,k}$ across the 'columns' (see Fig. 1 which illustrates the row-column terminology). To be more precise, we define the following orthogonal projections:

$$\mathcal{P} = \bigotimes_{p=1}^{P} \mathcal{T}_{A_p} \tag{4.187}$$

$$Q_{\text{mct}} = \mathcal{T}_{A_{1}^{\ell_{1}} A_{2}^{\ell_{2}} \dots A_{P}^{\ell_{P}}}, \tag{4.188}$$

$$\mathcal{R} = \mathcal{T}_{A_1 \dots A_P}.\tag{4.189}$$


Here, $\mathcal{T}_{A_p} = \int dU \, U_{A_{p,*}}^{\otimes K}(\cdot) (U_{A_{p,*}}^{\dagger})^{\otimes K}$ with $U_{A_{p,*}}^{\otimes K} \equiv \bigotimes_{k=1}^K U_{A_{p,k}}$. The systems $A_{p,k}^{\ell_p}$ consist of the first ℓ_p qudits within each block $A_{p,k}$, and the twirl $\mathcal{Q}_{\mathrm{mct}}$ defined in (4.188) applies a random unitary $U_{A_1^{\ell_1}...A_p^{\ell_p}}^{\otimes K}$ across the K columns.

As in the previous sections, we have the relations

$$\mathcal{PR} = \mathcal{RP} = \mathcal{R} = \mathcal{RQ}_{\text{mct}} = \mathcal{Q}_{\text{mct}} \mathcal{R} \qquad \text{im } \mathcal{P} \cap \text{im } \mathcal{Q}_{\text{mct}} = \text{im } \mathcal{R}, \qquad (4.190)$$

and thus a repeated alternating application of \mathcal{P} and \mathcal{Q}_{mct} converges to the full twirl \mathcal{R} :

$$\lim_{n \to \infty} (\mathcal{Q}_{\text{mct}} \mathcal{P})^n = \mathcal{R}. \tag{4.191}$$



(A) Twirl \mathcal{P} consists of individual twirls \mathcal{T}_{A_p} over the P collections of systems $A_{p,1}, \ldots, A_{p,K}$ for $p = 1, \ldots, P$.

(B) Twirl $\mathcal{Q}_{\mathrm{mct}}$ is a 'crosstwirl' $\mathcal{T}_{A_1^{\ell_1}...A_P^{\ell_P}}$ over the first ℓ_p systems in each of the systems $A_{1,k},\ldots,A_{P,k}$ for $k=1,\ldots,K$.

FIGURE 1. Multipartite Twirl-Crosstwirl protocol described in Section 4.4, alternating between (A) the twirl \mathcal{P} defined in (4.187) and (B) the twirl \mathcal{Q}_{mct} defined in (4.188). Note that the different blocks $A_{p,k}$ for $p = 1, \ldots, P$ and $k = 1, \ldots, K$ can have different sizes m_p and a different number ℓ_p of qudits participating in the crosstwirl \mathcal{Q}_{mct} in (B). In this example, $(\ell_1, m_1) = (3, 9)$, $(\ell_2, m_2) = (2, 6)$, and $(\ell_P, m_P) = (1, 7)$.

We have the following bound on the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})$ controlling the convergence speed in (4.191):

Proposition 4.6. Let $K, q, P, \ell_1, \ldots, \ell_P, m_1, \ldots, m_P$ be as above, and set $L = \ell_1 + \cdots + \ell_P$ and $M = m_1 + \cdots + m_P$. We assume that $K \leq q^{m_p}$ for all $p = 1, \ldots, P$ as well as $K \leq q^L$ and $K^2 < 2q^L$. Then the subspace angle $c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})$ of the two projections \mathcal{P} and $\mathcal{Q}_{\operatorname{mct}}$ defined in (4.187) and (4.188), respectively, satisfies

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})^{2} \leq acgq^{-2LK}K!^{P} \prod_{p=1}^{P} {q^{2\ell_{p}} + K - 1 \choose K} - 2abq^{-2MK}K!^{P} \prod_{p=1}^{P} {q^{2m_{p}} + K - 1 \choose K} + abq^{-2MK}K!^{2P} \prod_{p=1}^{P} {q^{m_{p}} + K - 1 \choose K}^{2},$$

$$(4.192)$$

with constants

$$a = \prod_{p=1}^{P} \left(1 - \frac{K^2}{2q^{m_p}} \right)^{-1} \qquad b = \exp\left(-\frac{K^2}{2q^M} \right)$$

$$c = \left(1 - \frac{K^2}{2q^L} \right)^{-1} \qquad g = \prod_{p=1}^{P} \exp\left(\frac{K^2}{2q^{m_p - \ell_p}} \right).$$
(4.193)

The proof of Proposition 4.6 is a rather straightforward generalization to the multipartite setting of the proof of Proposition 4.4 and given in Appendix A.

For later use, we rephrase Proposition 4.6 as a simplified bound in terms of TPEs:

Corollary 4.7. Let $\mathcal{P}, \mathcal{Q}_{\text{mct}}, K, q, P, \ell_1, \dots, \ell_P, m_1, \dots, m_P$ be as in Proposition 4.6, and furthermore assume that $K^2 \sum_p q^{\ell_p} \leq 1$ and that $m_p \geq 3\ell_p$ for each p. Then

$$\|\mathcal{P}Q_{ct} - \mathcal{R}\|_{2\to 2} \le 5K\sqrt{\sum_{p} \frac{1}{q^{2\ell_p}}}$$
 (4.194)

As noted in the introduction, a direct implication of Corollary 4.7 is that, if $\ell_p = \ell$ for each p, then $\ell = \log_q K + \log_q(5) + (1/2) \log_q(P) + \log_q(1/\epsilon)$ suffices to obtain TPE error ϵ . We leave the proof of this Corollary to the end of the section, as it primarily involves algebraic manipulation, and instead illustrate its main consequences.

Theorem 4.8. Let $A = \bigotimes_{p=1}^{P} A_p$. Let ν be an exact K-design on a system defined as the tensor product over $p = 1, \ldots, P$ of any ℓ_p qudits of local dimension q from each A_p . Assume that $4K^2 \times \sum_p q^{-\ell_p} \leq 1$ and that $m_p \geq 3\ell_p$ for each p. Let each μ_p be an exact K-design on the system A_p . Then $\nu * \bigotimes_p \mu_p$ is an ϵ -approximate relative K-design when ϵ given by

$$\epsilon = 25K!^{2P}K\sqrt{\sum_{p} \frac{1}{q^{2\ell_p}}}$$
 (4.195)

satisfies $\epsilon < 1$. Moreover, each unitary in $\nu * \bigotimes_p \mu_p$ requires at most $2\ell_p$ qudits of quantum communication between A_p and the rest of A to implement, generating at most $2\ell_p \log_2 q$ ebits of entanglement entropy from a product state input.

Proof. For each p = 1, ..., P, let \tilde{A}_p denote the subsystem of A_p consisting of the ℓ_p qudits affected by \mathcal{Q}_{mct} , and let \tilde{A}'_p be its complement within A_p . The fixed point subalgebra of $\bigotimes_{n=1}^P \mathcal{T}_{A_n}$ is a subalgebra of that of $(\bigotimes_{n=1}^P \mathcal{T}_{\tilde{A}_p} \otimes \mathcal{T}_{\tilde{A}'_p})$. Observe that

$$Q_{\text{mct}} \circ \bigotimes_{p=1}^{P} \mathcal{T}_{A_{p}} = \left(\bigotimes_{p=1}^{P} \mathcal{T}_{\tilde{A}_{p}} \otimes \mathcal{T}_{\tilde{A}'_{p}}\right) \circ Q_{\text{mct}} \circ \bigotimes_{p=1}^{P} \mathcal{T}_{A_{p}} \circ \left(\bigotimes_{p=1}^{P} \mathcal{T}_{\tilde{A}_{p}} \otimes \mathcal{T}_{\tilde{A}'_{p}}\right) , \quad (4.196)$$

since each $\mathcal{T}_{\tilde{A}'_p}$ commutes with \mathcal{Q}_{mct} and is absorbed by \mathcal{Q}_{mct} , and since each $\mathcal{T}_{\tilde{A}_p}$ commutes with \mathcal{T}_{A_p} and is also absorbed by \mathcal{Q}_{mct} . Lemma 3.7 then implies we should set the norm conversion

factor to

$$K!^{2P} \left(\prod_{p=1}^{P} \min \left\{ \left(1 - \frac{K^2 q^{\ell_p}}{|A_p|} \right)^{-1}, \frac{|A_p|^K}{q^{\ell_p K} K!} \right\} \min \left\{ \left(1 - \frac{K^2}{q^{\ell_p}} \right)^{-1}, \frac{q^{\ell_p K}}{K!} \right\} \right). \tag{4.197}$$

Assuming that $m_p \geq \ell_p$ simplifies the above, yielding the desired results with

$$\epsilon = 5K!^{2P}K\sqrt{\sum_{p} \frac{1}{q^{2\ell_{p}}}} \prod_{p=1}^{P} \left(1 - \frac{K^{2}q^{\ell_{p}}}{|A_{p}|}\right)^{-1} \left(1 - \frac{K^{2}}{q^{\ell_{p}}}\right)^{-1}.$$
 (4.198)

By the assumptions of the Theorem and Remark 4.9,

$$\prod_{p=1}^{P} \left(1 - \frac{K^2 q^{\ell_p}}{|A_p|} \right)^{-1} \left(1 - \frac{K^2}{q^{\ell_p}} \right)^{-1} \le \prod_{p=1}^{P} \left(1 - \frac{K^2}{q^{\ell_p}} \right)^{-2} \le \left(1 + 2 \sum_{p} K^2 / q^{\ell_p} \right)^2 \tag{4.199}$$

$$\leq 1 + 8K^2 \sum_{p} \frac{1}{q^{\ell_p}} \leq 5.$$
(4.200)

We now proceed to prove Corollary 4.7, for which we need the following remark and lemma.

Remark 4.9. For any $n \in \mathbb{N}$ and $\epsilon_1, \ldots, \epsilon_n \in \mathbb{R}$ such that each $\epsilon_i > -1$, and all ϵ_i having the same sign, the generalized Bernoulli's inequality [31, eq. (7.1)] states that

$$\prod_{j=1}^{n} (1 + \epsilon_j) \ge 1 + \sum_{j=1}^{n} \epsilon_j$$

Moreoever,

$$\prod_{j=1}^{n} (1 + \epsilon_j) \le \prod_{j=1}^{n} e^{\epsilon_j} \le \exp\left(\sum_{j} \epsilon_j\right).$$

Furthermore, $e^x \le 1 + x + x^2$ for every $0 \le 1 \le x$, and $e^{-x} \le 1 - x + x^2$ for every $-1 \le x \le 1$. Therefore,

$$\prod_{j=1}^{n} (1 + \epsilon_j) \le 1 + \sum_{j=1}^{n} \epsilon_j + \left(\sum_{j=1}^{n} \epsilon_j\right)^2 \le 1 + 2\sum_{j=1}^{n} \epsilon_j$$

if $|\sum_{j=1}^{n} \epsilon_j| \le 1$. Also note that for any $\delta \in [0, 1/2]$,

$$\frac{1}{1-\delta} = 1 + \delta + \frac{\delta^2}{1-\delta} \le 1 + 2\delta . \tag{4.201}$$

Lemma 4.10. Let $q, m, k \in \mathbb{N}$. Then

$$1 + \frac{k(k-1)}{2q^m} \le q^{-mk}k! \binom{q^m + k - 1}{k} \le \exp\left(\frac{k(k-1)}{2q^m}\right), \tag{4.202}$$

and if $k(k-1)/2q^m \le 1.7$, then

$$\dots \le 1 + \frac{k(k-1)}{2a^m} + \frac{k^2(k-1)^2}{4a^{2m}}$$
.

Proof. Expanding the binomial,

$$q^{-mk}k!\binom{q^m+k-1}{k} = q^{-mk}\frac{(q^m+k-1)!}{(q^m-1)!},$$
(4.203)

as the k! factors immediately cancel. Expanding factorials,

$$q^{-mk}\frac{(q^m+k-1)!}{(q^m-1)!} = q^{-mk}\prod_{j=1}^k (q^m+j-1) = \prod_{j=1}^k \left(1 + \frac{j-1}{q^m}\right). \tag{4.204}$$

Remark 4.9 completes this Lemma along with the fact that $(1+x)^r \leq e^{rx}$ for $x \in \mathbb{R}, r \geq 0$.

Proof of Corollary 4.7. Most of the proof follows from successively applying Remark 4.9. Applying Lemma 4.10 to Proposition 4.6,

$$\|\mathcal{PQ}_{ct} - \mathcal{R}\|_{2\to 2}^2 \le acg \exp\left(\sum_{p} \frac{K^2}{2q^{\ell_p}}\right) - 2ab\left(1 + \sum_{p} \frac{K^2}{q^{2m_p}}\right) + ab \exp\left(1 + \sum_{p} \frac{K^2}{q^{m_p}}\right)$$
(4.205)

$$\leq acg\left(1 + \sum_{p} \frac{K^2}{q^{\ell_p}}\right) + ab\left(1 + \sum_{p} \frac{2K^2}{q^{m_p}}\right) - 2ab\left(1 + \sum_{p} \frac{K^2}{q^{2m_p}}\right). \tag{4.206}$$

Using that $1 \ge b \ge 1 - K^2/2q^M$,

$$\|\mathcal{PQ}_{ct} - \mathcal{R}\|_{2\to 2}^2 \le a \left(cg \left(1 + \sum_{p} \frac{K^2}{q^{\ell_p}} \right) + \sum_{p} \frac{2K^2}{q^{m_p}} - 1 + \frac{K^2}{2q^M} \right). \tag{4.207}$$

Expanding the first term,

$$cg\left(1 + \sum_{p} \frac{K^2}{q^{\ell_p}}\right) \le \left(1 + \frac{K^2}{q^L}\right)\left(1 + \sum_{p} \frac{K^2}{q^{m_p - \ell_p}}\right)\left(1 + \sum_{p} \frac{K^2}{q^{2\ell_p}}\right)$$
 (4.208)

$$\leq 1 + 2K^2 \left(\sum_{p} \frac{1}{q^{m_p - \ell_p}} + \sum_{p} \frac{1}{q^{2\ell_p}} + \frac{1}{q^L} \right).$$
 (4.209)

Therefore,

$$\|\mathcal{P}\mathcal{Q}_{ct} - \mathcal{R}\|_{2\to 2}^2 \le aK^2 \left(2\sum_p \left(\frac{1}{q^{m_p}} + \frac{1}{q^{2\ell_p}} + \frac{1}{q^{m_p - \ell_p}}\right) + \frac{2}{q^L} + \frac{1}{2q^M}\right). \tag{4.210}$$

With the assumption that $m_p \geq 3\ell_p$ for each p,

$$\|\mathcal{PQ}_{ct} - \mathcal{R}\|_{2\to 2}^2 \le 9aK^2 \sum_{p} \frac{1}{q^{2\ell_p}}.$$
 (4.211)

Noting that $a \leq 2$ by the Lemma's assumptions and taking the square root,

$$\|\mathcal{PQ}_{ct} - \mathcal{R}\|_{2\to 2} \le 5K\sqrt{\sum_{p} \frac{1}{q^{2\ell_p}}}$$
 (4.212)

5. Recursive Crosstwirls

As shown by Proposition 4.6, one may use the crosstwirl as a superchannel that takes a few approximate k-design channels in tensor product and returns an approximate k-design channel on the full system. In this section, we study schemes that build up large k-designs from small k-designs by iterating the crosstwirl.

The following Lemma shows that when iterating approximate designs in relative error, the combined relative error is to first order approximately additive in its constituents. This Lemma will be valuable in subsequent sections.

Lemma 5.1. Consider two families of channels $(\Phi_n)_{n=1}^N$ and $(\Psi_n)_{n=1}^N$ such that

$$(1 - \epsilon_n)\Psi_n \prec \Phi_n \prec (1 + \epsilon_n)\Psi_n \tag{5.1}$$

for each n = 1 ... N. With $\epsilon = \prod_n (1 + \epsilon_n) - 1$,

$$(1 - \epsilon) \prod_{n} \Psi_n \prec \prod_{n} \Phi_n \prec (1 + \epsilon) \prod_{n} \Psi_n$$
 (5.2)

Furthermore, assume that $(1 - \delta)\Gamma \prec \prod_n \Psi_n \prec (1 + \delta)\Gamma$ for some channel Γ . Then $(1 + \delta)(1 + \epsilon)$ upper bounds the total relative error of Φ with respect to Γ .

Proof. Recall that for each n, $\Phi_n = (1 - \epsilon)\Psi_n + \epsilon\Theta_n$ for some channel Θ_n , and $(1 + \epsilon)^{-1}\Psi_n = (1 - (1 + \epsilon)^{-1})\Phi_n + \epsilon\Theta'_n$ for some channel Θ'_n . Therefore,

$$\prod_{n} \Phi_{n} = \prod_{n} (1 - \epsilon_{n}) \Psi_{n} + \left(1 - \prod_{n} (1 - \epsilon_{n})\right) \Theta$$
(5.3)

for some channel Θ that is a convex combination of Θ_n for different n. Similarly,

$$\prod_{n} \Psi_{n} = \prod_{n} (1 + \epsilon_{n})^{-1} \Phi_{n} + \left(1 - \prod_{n} (1 + \epsilon_{n})^{-1}\right) \Theta'$$
(5.4)

for some channel Θ' . Therefore,

$$\prod_{n} (1 - \epsilon_n) \Psi_n \prec \prod_{n} \Phi_n \prec \prod_{n} (1 + \epsilon_n) \Psi_n . \tag{5.5}$$

Using Remark 4.9,

$$\prod_{n} (1 - \epsilon_n) \ge 1 - \sum_{n} \epsilon_n \ge 1 - \left(1 - \prod_{n} (1 + \epsilon_n)\right). \tag{5.6}$$

Therefore, one obtains that with $\epsilon' = 1 - \prod_n (1 + \epsilon_n) \le 1 + \sum_n \epsilon_n \le \prod_n (1 + \epsilon_n)$,

$$(1 - \epsilon') \prod_{n} \Psi_n \prec \prod_{n} \Phi_n \prec (1 + \epsilon) \prod_{n} \Psi_n \tag{5.7}$$

The final part of the Lemma follows from expanding $\prod_n \Phi_n$ as its convex combination involving $\prod_n \Psi_n$.

5.1. **General recursive schemes.** Consider a system A divided into P_1 subsystems $A = A_1 \otimes \cdots \otimes A_{P_1}$, where each subsystem is further divided into P_2 subsystems, $A_i = A_{i,1} \otimes \cdots \otimes A_{i,P_{2,i}}$, and so on until some depth $r \in \mathbb{N}$. We may denote such a system using a tree data structure.

Definition 5.2 (Recursive Crosstwirl & Tree). Let A be a system of m qudits. A **recursive** crosstwirl tree is defined inductively as

$$R_{\vec{n}} = (A_{(\vec{n})}, ((S_{\vec{n},1}, R_{\vec{n} \oplus (1)}), \dots, (S_{\vec{n},t_{\vec{n}}}, R_{\vec{n} \oplus (t_{\vec{n}})}))), \qquad (5.8)$$

where \vec{n} is a vector index indicating the full path from the root to that node. Each $A_{(\vec{n})}$ is a subsystem of A composed of some qudits. $R_{\vec{n}\oplus(1)},\ldots,R_{\vec{n}\oplus(t_{\vec{n}})}$ are the node's children, where $t_{\vec{n}}$ can be any positive integer. For each $R_{\vec{n}\oplus(i)}$, we require that $A_{\vec{n}\oplus(i)}\subseteq A_{(\vec{n})}$. Moreover, $A_{(\vec{n})}=\bigotimes_i A_{(\vec{n})\oplus(i)}$. For the root node \vec{n} is the unique zero-dimensional vector with corresponding subsystem $A_{()}=A$. Each $S_{\vec{n},i}$ for $i=1,\ldots,t_{\vec{n}}$ is a set of qudit indices $1,\ldots,m$ within the subsystem corresponding to $R_{t_{\vec{n}}\oplus(i)}$ that will be involved in a crosstwirl on all of the node's children. By $S_{\vec{n}}$ we denote the vector $(S_1,\ldots,S_{\vec{n},t_n})$.

For each j = 0, ..., depth(R) - 1, we denote the jth layer of a crosstwirl tree R by the subset of vector indices $\text{La}_j[R]$, where each $\vec{n} \in \text{La}_j[R]$ is a vector of length j specifying a node in the tree. For each j = 0, ..., depth(R) - 1, let

$$\Psi_{j}[R] := \bigotimes_{\vec{n} \in \text{La}_{j}[R]} \begin{cases} \mathcal{T}[\vec{S}_{\vec{n}}] & \text{node } \vec{n} \text{ is not a leaf} \\ \mathcal{T}_{A_{(\vec{n})}} & \text{node } \vec{n} \text{ is a leaf}, \end{cases}$$
(5.9)

where $\mathcal{T}[\vec{S}_{\vec{n}}]$ is a crosstwirl applying a Haar-random unitary average to the tensor product system of all qudits selected by $\mathcal{T}[\vec{S}_{\vec{n}}]$. A **recursive crosstwirl** is defined as the channel $\Psi[R] = \Psi_0[R] \circ \cdots \circ \Psi_{\text{depth}(R)}[R]$.

Lemma 5.3 (Entanglement Union Bound). Consider a quantum system A initially in tensor product with an environment E. If unitaries U_1, \ldots, U_n are applied to $A \otimes E$ in sequence, each j-th differing from the identity only on l_j qudits within A of local dimension q, then the number of qudits of quantum communication needed to implement the sequence is at most $2\sum_{j=1}^{n} l_j \log q$, which also upper-bounds the final entanglement entropy of A with E.

Proof. For each $j \in 1 ... n$, the two parties may implement U_j by transmitting the l_j qudits from A to the environment, performing the full unitary there, and sending them back to A. This operation requires at most $2l_j$ qudits of quantum communication.

Remark 5.4. Let R be a recursive crosstwirl tree, and for each node index \vec{n} let $A_{(\vec{n})}$ denote the system corresponding to that node. Let S be a given set of node indices for which $C \subseteq \bigotimes_{\vec{n} \in S} A_{(\vec{n})}$, and each $R_{\vec{n}}$ for $\vec{n} \in S$ is a leaf. We denote by ∂S the boundary of S, that is, the set of node indices \vec{n} for which $R_{\vec{n}}$ is a leaf, $C \cap A_{(\vec{n})} \neq \emptyset$, and $A_{(\vec{n})} \not\subseteq C$. Let \tilde{S} be a set of pairs (\vec{r}, \vec{w}) , containing a pair for each crosstwirl defined by $\vec{l}_{\vec{r}}$ for which $\vec{l}_{\vec{r},\vec{w}}$ qudits of C are involved. If $C \otimes B$ for an auxiliary system B is initially unentangled with C', the complement of C in A_1 for

another auxiliary system B'

$$\sum_{\vec{n} \in \partial S} \log_2 |A_{(\vec{n})}| + 2 \sum_{\vec{r}, \vec{w} \in \tilde{S}} \vec{l}_{\vec{r}, \vec{n}} \log_2 q$$
 (5.10)

qubits of quantum communication between $C \otimes B$ and $C' \otimes B'$ are needed to implement the recursive crosstwirl Ψ .

Definition 5.5. We say that a recursive crosstwirl (tree) parallelizes or is a parallel recursive crosstwirl (tree) if each crosstwirl involved acts on a distinct set of qudits.

Remark 5.6. Let R be a recursive crosstwirl tree. Then the recursive crosstwirl $\Psi[R]$ is implemented via unitaries of depth at most

$$\sum_{i=1}^{\operatorname{depth}(R)} \max_{n \in \operatorname{La}_{j}[R]} \operatorname{depth}(\Phi_{n})$$
(5.11)

where depth($\Phi_{\vec{n}}$) denotes the maximum depth needed to implement a unitary in the ensemble for $\Phi_{\vec{n}}$. If R parallelizes, then the required depth is at most

$$\max_{\vec{n} \in \text{Leaves}(R)} \operatorname{depth}(\Phi_{\vec{n}}) + \max_{\vec{n} \in \text{Non-leaves}(R)} \operatorname{depth}(\Phi_{\vec{n}}) . \tag{5.12}$$

To summarize the results of this section so far, recursive crosstwirls can form relative error k-designs on m qudits in depth not much larger than that needed for constituent designs on leaf subsystems or crosstwirls, also requiring a similar amount of quantum communication. However, as these results are quite general, they leave out the specifics of these bounds. The subsequent Subsection 5.2 fills that gap.

5.2. **Spatial Lattices.** A common paradigm in approximate k-design ensembles considers qudits interacting on a spatial lattice, such as in [22]. This scenario is also common in applications of entanglement entropy, with particular emphasis on entanglement entropy between subregions in these lattices [13]. Here we consider recursive crosstwirls on these geometries; low-dimensional examples are depicted in Figure 2 (for D = 1) and Figure 3 (for D = 2).

Definition 5.7. Let $A = \bigotimes_{p=1}^P = \mathbb{M}_q^{\otimes m}$ be such that P is a natural power of 2^D , and each $|A_p| \geq q^\ell$ for some $\ell \in \mathbb{N}$. Let R be a complete 2^D -ary recursive crosstwirl tree of depth $\frac{1}{D} \log_2 P$. For each non-leaf node index \vec{n} , we interpret the child nodes $R_{\vec{n}\oplus 1}\dots R_{\vec{n}\oplus 2^D}$ to be arranged in a D-dimensional hypercube. Each $S_{\vec{n},i}$ for $i=1,\dots,2^D$ is then chosen such that the qudits indexed by $S_{\vec{n},1} \cup \dots \cup S_{\vec{n},2^D}$ form a contiguous system of $2^D \ell$ qudits on a D-dimensional lattice. We call R a **D-ary lattice crosstwirl tree** (D-LCT) and the corresponding recursive crosstwirl Ψ a **D-ary lattice crosstwirl** (D-LC).

Our main theorem in this section is the following result:

Theorem 5.8. Let A be a system of M connected qubits on a hypercube lattice such that $M = 2^x$ and is sufficiently large, with $x \in \mathbb{N}$ being the side length. For any $\epsilon > 0$ and $K \in \mathbb{N}$, we may




FIGURE 2. (A) A 1-LCT in which the vertical direction represents layers. At the bottom, each of 8 leaf subsystems is in a K-design. By the top, the whole system is in an approximate design. (B) First, a twirl is applied to each of the 8 leaf subsystems. Then blue, then red, then green crosstwirls are applied in successive layers, although they may parallelize. Each colored rectangle represents the portion of the system invoved in a crosstwirl at that layer.




FIGURE 3. (A) A 2-LCT in which the horizontal direction represents layers. At the left, each of 16 leaf subsystems is in a K-design. By the right, the whole system is an approximate design. (B) First, Haar twirls are applied to each of the 16 leaf subsystems. Then blue, then red crosstwirls are applied in successive layers, although they may parallelize. Each colored square represents the portion of the system invovled in a crosstwirl at that layer.

construct an approximate, parallelizable lattice crosstwirl yielding an ϵ -approximate relative error K-design for which

$$\ell \le 150M \times K!^{2^D} K \times 2^{D/2} \times \frac{1}{\epsilon} \,, \tag{5.13}$$

and

• The maximum circuit depth to implement any unitary is $O(\ell K \operatorname{polylog}(K))$, and the implementation can be constructed using Haar random 2-local unitaries.

• Quantum communication between any contiguous subregion S and its complement in A is at most $\#(\partial S) \times (\sqrt[D]{2^{D+1}\ell} + 1/(2^{D+1}\ell)^{(D-1)/D} \sqrt[D]{\ell}) + (2^{D+1}+1)\ell$ quaits.

Proof. The Theorem follows from Propositions 5.11 and 5.12 proved below. In particular, Proposition 5.11 takes a design scheme with a depth function $d(r, q, K, \eta)$, where r is the qudit number, q the local dimension, K as in this Theorem, and η the error, yielding a total depth of $2d(2^{D+3}\ell, q, K, \frac{\epsilon}{2M})$. To complete the construction, we invoke the scheme from [7] yielding depth $(rK + \log(1/\epsilon))$ polylog(K), which translates to this Theorem's depth after substituting ℓ for the parameter r and $\epsilon/2M$ for the parameter η . Moreover, that scheme uses one-dimensional brickwork, an ensemble constructed from 2-local random unitaries. Proposition 5.12 determines the required amount of quantum communication.

Note that Theorem 5.8 does not rely qualitatively on [7] and could be proven using the older results of [3] as the underlying design construction. Such would yield a polynomially larger dependence on K but maintain the same logarithmic dependencies on M and ϵ .

Remark 5.9 (Almost Brickwork). A recent result [7] showed that brickwork random circuits on a one-dimensional lattice are k-designs with depth $O(mk \operatorname{polylog}(K))$. On the 1-hypercube, the steps used in Theorem 5.8 can be implemented this way, yielding the Theorem statement with $g(k) = O(\operatorname{polylog}(K))$. In this case, layers of the construction consist of patches of brickwork random circuits on different subregions. Moreover, recall that composing random unitaries to a subsystem that would already be output as a k-design state does not remove the K-design property. Therefore, one may apply 2-local random unitaries to most of the system at each step. To satisfy the criteria of Theorem 5.8, the only deviation from brickwork is that random unitaries not cross boundaries separating different subsystems indexed by the 1-LCT.

The rest of this Section is devoted to proving intermediate results needed for Theorem 5.8.

Lemma 5.10. A D-LCT is parallelizable if $|A_p| \ge 2\ell$ for every leaf index p = 1, ..., P.

Proof. The proof is inductive, the base case being that any leaf node has at least 2ℓ qudits. For any $j=2,\ldots,\frac{1}{D}\log P$, assume for each $n\in\mathrm{La}_j[R]$ that $A_{(n)}$ has at least 2ℓ qudits not involved in any prior crosstwirl. When each $A_{(n)}$ is crosstwirled with its 2^D-1 siblings, ℓ qudits of $A_{(n)}$ and $2^D\ell$ qudits in total will be involved in the crosstwirl without overlapping any prior crosstwirl. Let $A_{(i)}$ be the subsystem corresponding to the parent of the n-th node. In a D-dimensional hypercube of 2^D subcubes, each cube corresponds to one exposed corner. Hence there are 2^D subsequent crosstwirls possibly involving $A_{(i)}$, and $2^D\ell$ qudits remaining.

Proposition 5.11. Let A be a system of M connected qudits on a hypercube lattice such that $M = 2^{Dx}$, with $x \in \mathbb{N}$ being the side length of the hypercube in units of qudits.

Let $\Phi[r,q,k,\delta]$ be a K-design channel scheme on this connectivity with circuit depth $d(r,q,k,\delta)$, where r is the number of qudits acted on in the given connectivity, q is the local qudit dimension, K the number of copies in the design, and δ the relative error tolerance.

For any $\epsilon > 0$ and $K \in \mathbb{N}$, we may construct an approximate, parallelizable D-LCT R on $A^{\otimes K} = \bigotimes_{p=1}^{P} A_p^{\otimes K}$ yielding an ϵ -approximate K-design in relative error with depth at most

 $2d(2^{D+3}\ell,q,K,\frac{\epsilon}{5M})$ such that

$$\ell = \log_q \left(150M \times K!^{2^D} K \times 2^{D/2} \times \frac{1}{\epsilon} \right). \tag{5.14}$$

Proof. Assume that we will construct a D-LC with P leaves. For some ℓ to be determined as the number of qudits involved in each crosstwirl, define a logarithmic leaf side length as $s := \lceil \frac{1}{D} \log_2(8\ell) \rceil$. Then the number of leaves P is at most $2^{D(x-s)}$. The number of qudits in each leaf system is then between 2ℓ and $2^{D+3}\ell$ if $M \ge 2^{D+3}\ell$. If m is not this large, then we simply perform a full, approximate design twirl on M and are finished.

The leaf-level approximate designs contribute a multiplicative total error of $(1+\delta)^P$ in depth at most $d(2^{D+3}\ell, q, k, \delta)$. The non-leaf design approximations used to perform the crosstwirls then contribute a multiplied total error of $(1+\delta)^{P-1}$ in depth at most $d(2^{D+1}\ell, q, k, \varepsilon)$. Then combining error contributions from Theorem 4.8 using Lemma 5.1,

$$(1 + \epsilon_{\text{tot}}) \le (1 + \delta)^{2P - 1} \left(1 + 25K!^{2^D} K \times 2^{D/2} \times \frac{1}{q^{\ell}} \right)^{P - 1}. \tag{5.15}$$

Now set

$$\ell = \log_q \left(25 \times K!^{2D} K \times 2^{D/2} \times \frac{6P}{\epsilon} \right) \tag{5.16}$$

for some $c \in \mathbb{R}^+$. Then using Remark 4.9 with the Proposition's assumptions,

$$(1 + \epsilon_{\text{tot}}) \le (1 + \delta)^{2P - 1} \left(1 + \frac{\epsilon}{6P} \right)^{P - 1} \le \left(1 + \frac{2P - 1}{5P} \epsilon \right) \left(1 + \frac{P - 1}{3P} \right) \le \epsilon ,$$
 (5.17)

setting $\delta = \epsilon/10P$. We replace the P in ℓ by M recalling the assumption that $P \leq 2M$, so that the final result is stated only in terms of constants given. Since the D-LC is parallelizable by Lemma 5.10, Remark 5.6 yields the depth bound.

Proposition 5.12. Let A be a D-hypercubical system of |A| qudits, each with local dimension q, where $M = 2^{Dx}$ for side length x. Let $S \subseteq A$ be a subsystem corresponding to a geometrically contiguous subset of those qudits. Let ∂S denote the boundary of S, the set of qudits that are part of S with neighbors not in S. Let $\#(\partial S)$ be the number of qudits on that boundary. The D-LC constructed in Proposition 5.11 requires at most

$$\left[\frac{\#(\partial S)}{(2^{D+1}\ell)^{(D-1)/D}}\right](2^{D+1}+1)\ell\log_2 q \tag{5.18}$$

ebits of quantum communication between S and its complement in A.

Proof. For the amount of entanglement across subregions, recall Remark 5.4. The first term therein is the contribution from the boundary potentially cutting through leaf systems. The entanglement between parts of those leaf systems would contribute to the entanglement across the boundary. While entanglement between parts of a leaf system is upper bounded by the smaller of the volumes on either side of the cut, the portion of the boundary occupied by a single leaf system is proportional to that system's area. Though leaf subsystem sizes vary beween

 2ℓ and $2^{D+1}\ell$, larger leaf systems contribute proportionally more entanglement. Therefore, the worst case leaf subsystem contribution to entanglement entropy is upper bounded by

$$\left[\frac{\#(\partial S)}{(2^{D+1}\ell)^{(D-1)/D}}\right] 2^{D+1}\ell \ . \tag{5.19}$$

The second, added contribution to total entanglement entropy as in Lemma 5.3 comes from the crosstwirls. To determine this contribution, we make the following geometric observations:

- Each crosstwirl is geometrically contiguous by construction.
- Therefore, each crosstwirl occurs at a corner of each involved subsystem. Once a corner has been in a crosstwirl, it becomes the center of the higher-level subsystem.
- Hence only one corner from that original subsystem remains exposed for a potential, later crosstwirl. Once both corners have been crosstwirled, that subsystem has no more exposed corners or boundaries. Hence each leaf subsystem is involved in at most 2 crosstwirls.

To get from one crosstwirl location to another, a cut through the full system must traverse the rest of a full leaf system. Therefore, as in the entanglement contribution from leaf systems, the boundary length is divided by the leaf subsystem size. Hence the total contribution from crosstwirls is upper bounded by

$$\left[\frac{\#(\partial \mathcal{S})}{(2^{D+1}\ell)^{(D-1)/D}}\right]\ell, \qquad (5.20)$$

which concludes the proof.

Remark 5.13. A concurrent work [38] also shows that designs can be constructed on any geometry in logarithmic depth. That result uses a Hamiltonian path construction, in which any connected graph can efficiently approximate a one-dimensional layout. Via a detailed technical analysis, the other work also obtains an improved dependence on k. Straightforwardly counting the points at which a Hamiltonian path may cross a boundary suggests that subregion entanglement in such a construction scales as $\#(\partial S) \times O(\log K + \log m + \log \epsilon)$. Our recursive construction as in Theorem 5.8 exploits partial cancellation between the small block length and boundary length to show that on lattices, entanglement may scale as $\#(\partial S) \times O((K \log K + \log m + \log \epsilon)^{1/D}) + O(K \log K + \log m + \log \epsilon)$. A simple combination of schemes would be expected to achieve scaling of $\#(\partial S) \times O((\log K + \log m + \log \epsilon)^{1/D}) + O(\log K + \log m + \log \epsilon)$.

APPENDIX A. PROOF OF THE SUBSPACE ANGLE BOUND FOR MULTIPARTITE CROSSTWIRL PROTOCOL

In this appendix we give a full proof of Proposition 4.6 giving a bound on the subspace angle for the multipartite crosstwirl protocol described in Section 4.4.

Proof of Proposition 4.6. We first rewrite the subspace angle in terms of the normalized Frobenius inner product $\langle \cdot, \cdot \rangle_n$ using Lemma 4.1 and (4.190):

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})^{2} = \sup_{X \neq 0} \frac{\|\mathcal{Q}_{\operatorname{mct}} \mathcal{P}(X) - \mathcal{R}(X)\|^{2}}{\|X\|^{2}} = \sup_{X \neq 0} \frac{\langle X, \mathcal{P} \mathcal{Q}_{\operatorname{mct}} \mathcal{P}(X) - \mathcal{R}(X) \rangle_{n}}{\langle X, X \rangle_{n}}, \quad (A.1)$$

and the supremum is achieved on an element $X = \mathcal{P}(X)$, which by (2.19) can be written as

$$X = \mathcal{P}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \, \sigma_{A_1}^1 \otimes \dots \otimes \sigma_{A_P}^P \tag{A.2}$$

for some coefficients $x_{\vec{\sigma}} \in \mathbb{C}$, where we write $\vec{\sigma} = (\sigma^1, \dots, \sigma^P)$ with $\sigma^p \in \mathfrak{S}_k$ for $p = 1, \dots, P$. Using Lemma A.1 proved below, we express the right-hand side of (A.1) as

$$\sup_{X \neq 0} \frac{\langle X, \mathcal{P}Q_{\text{mct}}\mathcal{P}(X) - \mathcal{R}(X) \rangle_{n}}{\langle X, X \rangle_{n}}$$

$$= \sup_{\mathbf{x} \neq 0} \frac{\left\langle \mathbf{x}, \left[\left(\bigotimes_{p=1}^{P} \mathbf{G}_{A_{p}^{m_{p}-\ell_{p}}} \right) \odot \left(\mathbf{N}_{\ell_{1}, \dots, \ell_{P}}^{T} \mathbf{G}_{\ell_{1}, \dots, \ell_{P}}^{-1} \mathbf{N}_{\ell_{1}, \dots, \ell_{P}} \right) - \mathbf{N}_{A_{1} \dots A_{P}}^{T} \mathbf{G}_{A_{1} \dots A_{P}}^{-1} \mathbf{N}_{A_{1} \dots A_{P}} \right] \mathbf{x} \right\rangle}{\left\langle \mathbf{x}, \left(\bigotimes_{p=1}^{P} \mathbf{G}_{A_{p}} \right) \mathbf{x} \right\rangle}, \tag{A.3}$$

where $\mathbf{G}_{\ell_1,\dots,\ell_P} \equiv \mathbf{G}_{A_1^{\ell_1}\dots A_P^{\ell_P}}$ is the Gram matrix of permutation operators acting on $A_1^{\ell_1}\dots A_P^{\ell_P}$, and the $(K!\times K!^P)$ -matrices $\mathbf{N}_{\ell_1,\dots,\ell_P}$ and $\mathbf{N}_{A_1\dots A_P}$ are defined as

$$(\mathbf{N}_{\ell_1,\dots,\ell_P})_{\pi,\vec{\sigma}} = \left\langle \pi_{A_1^{\ell_1}\dots A_P^{\ell_P}}, \sigma_{A_1^{\ell_1}}^1 \otimes \dots \otimes \sigma_{A_P^{\ell_P}}^P \right\rangle_{\mathbf{n}}$$
(A.4)

$$(\mathbf{N}_{A_1...A_P})_{\pi,\vec{\sigma}} = \langle \pi_{A_1...A_P}, \sigma_{A_1}^1 \otimes \cdots \otimes \sigma_{A_P}^P \rangle_{\mathbf{n}}. \tag{A.5}$$

Just as in the proofs of Theorems 4.2 and 4.4, we first eliminate the various Gram matrices in this expression. To this end, we set $M=m_1+\cdots+m_P$ and $L=\ell_1+\cdots+\ell_P$, and define constants

$$a = \prod_{p=1}^{P} \left(1 - \frac{K^2}{2q^{m_p}} \right)^{-1} \qquad b = \exp\left(-\frac{K^2}{2q^M} \right)$$
 (A.6)

$$c = \left(1 - \frac{K^2}{2q^L}\right)^{-1} \qquad g = \prod_{p=1}^{P} \exp\left(\frac{K^2}{2q^{m_p - \ell_p}}\right). \tag{A.7}$$

We then use various operator inequalities that follow from bounds on the largest and smallest eigenvalues of Gram matrices in [20]. First, $\mathbf{G}_{A_p} \geq \left(1 - \frac{K^2}{2q^{m_p}}\right) \mathbb{1}_{K!}$ and hence

$$\left\langle \mathbf{x}, \left(\bigotimes_{p=1}^{P} \mathbf{G}_{A_p} \right) \mathbf{x} \right\rangle^{-1} \le a \langle \mathbf{x}, \mathbf{x} \rangle^{-1}.$$
 (A.8)

⁵Note that the superscript in σ^p is an index and not a power.

Second, $\mathbf{G}_{A_1...A_P} \leq \exp\left(\frac{K^2}{2q^M}\right) \mathbb{1}_{K!}$ and hence

$$-\left\langle \mathbf{x}, \mathbf{N}_{A_{1}...A_{P}}^{T} \mathbf{G}_{A_{1}...A_{P}}^{-1} \mathbf{N}_{A_{1}...A_{P}} \mathbf{x} \right\rangle \leq -b \left\langle \mathbf{x}, \mathbf{N}_{A_{1}...A_{P}}^{T} \mathbf{N}_{A_{1}...A_{P}} \mathbf{x} \right\rangle. \tag{A.9}$$

To deal with the Hadamard product term, we use the same argument as in the proof of Proposition 4.4: From $\mathbf{G}_{\ell_1,\dots,\ell_P} \geq \left(1 - \frac{K^2}{2q^L}\right) \mathbb{1}_{K!}$ we get

$$\mathbf{N}_{\ell_{1},...,\ell_{P}}^{T}\mathbf{G}_{\ell_{1},...,\ell_{P}}^{-1}\mathbf{N}_{\ell_{1},...,\ell_{P}} \leq c\mathbf{N}_{\ell_{1},...,\ell_{P}}^{T}\mathbf{N}_{\ell_{1},...,\ell_{P}},$$
(A.10)

and $\mathbf{G}_{A_p^{m_p-\ell_p}} \leq \exp\left(\frac{K^2}{2q^{m_p-\ell_p}}\right)\mathbbm{1}_{K!}$ gives

$$\bigotimes_{p=1}^{P} \mathbf{G}_{A_p^{m_p-\ell_p}} \le g \mathbb{1}_{K!P}. \tag{A.11}$$

Now $X \odot Y \ge 0$ if $X,Y \ge 0$ [25, Ch. 5] and $(X+Z) \odot Y = X \odot Z + X \odot Y$, which we can use together with (A.10) and (A.11) to obtain

$$\left\langle \mathbf{x}, \left(\bigotimes_{p=1}^{P} \mathbf{G}_{A_{p}^{m_{p}-\ell_{p}}} \right) \odot \left(\mathbf{N}_{\ell_{1},\dots,\ell_{P}}^{T} \mathbf{G}_{\ell_{1},\dots,\ell_{P}}^{-1} \mathbf{N}_{\ell_{1},\dots,\ell_{P}} \right) \mathbf{x} \right\rangle$$

$$\leq cg \left\langle \mathbf{x}, \left(\mathbb{1}_{K!^{P}} \odot \mathbf{N}_{\ell_{1},\dots,\ell_{P}}^{T} \mathbf{N}_{\ell_{1},\dots,\ell_{P}} \right) \mathbf{x} \right\rangle. \quad (A.12)$$

Note that $\mathbb{1}_{K!^P} \odot \mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P}$ is a diagonal matrix.

We use the above bounds in (A.3) to get

$$\sup_{X \neq 0} \frac{\langle X, \mathcal{P}Q_{\text{mct}}\mathcal{P}(X) - \mathcal{R}(X) \rangle_{n}}{\langle X, X \rangle_{n}} \\
\leq a \sup_{\mathbf{x} \neq 0} \frac{\langle \mathbf{x}, \left(cg \mathbb{1}_{K!^{P}} \odot \mathbf{N}_{\ell_{1}, \dots, \ell_{P}}^{T} \mathbf{N}_{\ell_{1}, \dots, \ell_{P}} - b \mathbf{N}_{A_{1} \dots A_{P}}^{T} \mathbf{N}_{A_{1} \dots A_{P}} \right) \mathbf{x} \rangle}{\langle \mathbf{x}, \mathbf{x} \rangle} \\
= a \left\| cg \mathbb{1}_{K!^{P}} \odot \mathbf{N}_{\ell_{1}, \dots, \ell_{P}}^{T} \mathbf{N}_{\ell_{1}, \dots, \ell_{P}} - b \mathbf{N}_{A_{1} \dots A_{P}}^{T} \right\|. \tag{A.14}$$

Our strategy to bound this operator norm is the same as in the proof of Proposition 4.4. Let us define two $(K!^P \times K!^P)$ -matrices **D** and **K** via

$$(\mathbf{D})_{\vec{\sigma},\vec{\omega}} = \begin{cases} \left(cg \mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} - b \mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P} \right)_{\vec{\sigma},\vec{\sigma}} & \text{if } \vec{\sigma} = \vec{\omega}, \\ 0 & \text{otherwise;} \end{cases}$$
(A.15)

$$(\mathbf{K})_{\vec{\sigma},\vec{\omega}} = \begin{cases} 0 & \text{if } \vec{\sigma} = \vec{\omega}, \\ \left(b\mathbf{N}_{A_1...A_P}^T \mathbf{N}_{A_1...A_P}\right)_{\vec{\sigma},\vec{\omega}} & \text{otherwise.} \end{cases}$$
(A.16)

These matrices satisfy

$$\mathbf{D} - \mathbf{K} = cg \mathbb{1}_{K!^P} \odot \mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} - b \mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P}, \tag{A.17}$$

and we have

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})^{2} = \sup_{X \neq 0} \frac{\|\mathcal{Q}_{\operatorname{mct}} \mathcal{P}(X) - \mathcal{R}(X)\|^{2}}{\|X\|^{2}}$$
(A.18)

$$= \sup_{X \neq 0} \frac{\langle X, \mathcal{PQ}_{\text{mct}} \mathcal{P}(X) - \mathcal{R}(X) \rangle_{\text{n}}}{\langle X, X \rangle_{\text{n}}}$$
(A.19)

$$\leq a\|\mathbf{D} - \mathbf{K}\| \tag{A.20}$$

$$\leq a\|\mathbf{D}\| + a\|\mathbf{K}\|. \tag{A.21}$$

We first bound the operator norm of **K**. This is a non-negative irreducible matrix, for which the generalized Perron-Frobenius Theorem [24, Ch. 1] states that $\|\mathbf{K}\| \leq \max_{\vec{\sigma}} \sum_{\vec{\omega}} (\mathbf{K})_{\vec{\sigma},\vec{\omega}}$. Let $\vec{\sigma} = (\sigma^1, \dots, \sigma^P)$ denote the vector of permutations achieving this maximum. Then we upperbound this expression by summing over all $\sigma^2, \dots, \sigma^P \in kS_k$. The reason to do this is that we can calculate the latter sum explicitly as follows:

$$\|\mathbf{K}\| \le \sum_{\vec{\omega}} (\mathbf{K})_{\vec{\sigma}, \vec{\omega}} \tag{A.22}$$

$$\leq \sum_{\sigma^2, \dots, \sigma^P} \sum_{\vec{\omega}} (\mathbf{K})_{\vec{\sigma}, \vec{\omega}} \tag{A.23}$$

$$= \sum_{\sigma^2,\dots,\sigma^P} \left(\sum_{\vec{\omega}} (b\mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P})_{\vec{\sigma},\omega} - (b\mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P})_{\vec{\sigma},\vec{\sigma}} \right)$$
(A.24)

$$= b \sum_{\sigma^2, \dots, \sigma^P} \sum_{\vec{\omega}} (\mathbf{N}_{A_1 \dots A_P}^T \mathbf{N}_{A_1 \dots A_P})_{\vec{\sigma}, \omega} - b \sum_{\sigma^2, \dots, \sigma^P} (\mathbf{N}_{A_1 \dots A_P}^T \mathbf{N}_{A_1 \dots A_P})_{\vec{\sigma}, \vec{\sigma}}$$
(A.25)

For the first sum,

$$\sum_{\sigma^{2},\dots,\sigma^{P}} \sum_{\vec{\omega}} \left(\mathbf{N}_{A_{1}\dots A_{P}}^{T} \mathbf{N}_{A_{1}\dots A_{P}} \right)_{\vec{\sigma},\vec{\omega}}
= \sum_{\sigma^{2},\dots,\sigma^{P}} \sum_{\vec{\omega}} \sum_{\pi} \prod_{p=1}^{P} \left\langle \sigma_{A_{p}}^{p}, \pi_{A_{p}} \right\rangle_{\mathbf{n}} \prod_{p=1}^{P} \left\langle \pi_{A_{p}}, \omega_{A_{p}}^{p} \right\rangle_{\mathbf{n}}
= \sum_{\pi} \left\langle \sigma_{A_{1}}^{1}, \pi_{A_{1}} \right\rangle_{\mathbf{n}} \sum_{\sigma^{2}} \left\langle \sigma_{A_{2}}^{2}, \pi_{A_{2}} \right\rangle_{\mathbf{n}} \cdots \sum_{\sigma^{P}} \left\langle \sigma_{A_{P}}^{P}, \pi_{A_{P}} \right\rangle_{\mathbf{n}} \sum_{\omega^{1}} \left\langle \pi_{A_{1}}, \omega_{A_{1}}^{1} \right\rangle_{\mathbf{n}} \cdots \sum_{\omega^{P}} \left\langle \pi_{A_{P}}, \omega_{A_{P}}^{P} \right\rangle_{\mathbf{n}}
(A.26)$$

$$=q^{-2MK}K!^{2P}\prod_{p=1}^{P}\binom{q^{m_p}+K-1}{K}^2,$$
(A.28)

where we used the identity (4.83) 2P times.

For the second sum in (A.25),

$$\sum_{\sigma^2,\dots,\sigma^P} (\mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P})_{\vec{\sigma},\vec{\sigma}} = \sum_{\sigma^2,\dots,\sigma^P} \sum_{\pi} \prod_{p=1}^P \left\langle \pi_{A_p}, \sigma_{A_p}^p \right\rangle_{\mathbf{n}}^2$$
(A.29)

$$= \sum_{\pi} \left\langle \pi_{A_1}, \sigma_{A_1}^1 \right\rangle_n^2 \sum_{\sigma^2} \left\langle \pi_{A_2}, \sigma_{A_2}^2 \right\rangle_n^2 \cdots \sum_{\sigma^P} \left\langle \pi_{A_P}, \sigma_{A_P}^P \right\rangle_n^2 \qquad (A.30)$$

$$= q^{-2MK} K!^P \prod_{p=1}^P \binom{q^{2m_p} + K - 1}{K}, \tag{A.31}$$

where we used the generalization of the identity (4.88) to P parties. Substituting (A.28) and (A.31) in (A.25) gives the bound

$$\|\mathbf{K}\| \le bq^{-2MK}K!^{2P} \prod_{p=1}^{P} {q^{m_p} + K - 1 \choose K}^2 - bq^{-2MK}K!^P \prod_{p=1}^{P} {q^{2m_p} + K - 1 \choose K}.$$
 (A.32)

To bound the operator norm of the diagonal matrix \mathbf{D} , we first note that all diagonal entries are positive: We have $gc \geq b$ (just as in (4.90)) using the assumption $K^2 < 2q^L$, and hence

$$(\mathbf{D})_{\vec{\sigma},\vec{\sigma}} = \left(cg\mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} - b\mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P}\right)_{\vec{\sigma},\vec{\sigma}}$$
(A.33)

$$\geq b \left(\mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} - \mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P} \right)_{\vec{\sigma},\vec{\sigma}} \tag{A.34}$$

$$>0, (A.35)$$

where the last strict inequality follows from the calculation

$$\left(\mathbf{N}_{\ell_{1},\ldots,\ell_{P}}^{T}\mathbf{N}_{\ell_{1},\ldots,\ell_{P}}\right)_{\vec{\sigma},\vec{\sigma}} = \sum_{\pi} \left\langle \bigotimes_{p=1}^{P} \sigma_{A_{p}^{\ell_{p}}}^{p}, \pi_{A_{1}^{\ell_{1}}\ldots A_{P}^{\ell_{P}}} \right\rangle_{\mathbf{n}} \left\langle \pi_{A_{1}^{\ell_{1}}\ldots A_{P}^{\ell_{P}}}, \bigotimes_{p=1}^{P} \sigma_{A_{p}^{\ell_{p}}}^{p} \right\rangle_{\mathbf{n}}$$
(A.36)

$$= \sum_{\pi} \prod_{p=1}^{P} \left\langle \sigma_{A_p^{\ell_p}}^p, \pi_{A_p^{\ell_p}} \right\rangle_{\mathbf{n}} \prod_{p=1}^{P} \left\langle \pi_{A_p^{\ell_p}}, \sigma_{A_p^{\ell_p}}^p \right\rangle_{\mathbf{n}} \tag{A.37}$$

$$= \sum_{p=1} \left\langle \bigotimes_{p=1}^{P} \sigma_{A_p}^p, \pi_{A_1 \dots A_P} \right\rangle_{\mathbf{n}} \left\langle \pi_{A_1 \dots A_P}, \bigotimes_{p=1}^{P} \sigma_{A_p}^p \right\rangle_{\mathbf{n}} \tag{A.39}$$

$$= \left(\mathbf{N}_{A_1 \dots A_P}^T \mathbf{N}_{A_1 \dots A_P} \right)_{\vec{\sigma} \ \vec{\sigma}}, \tag{A.40}$$

where we used (4.92) for the inequality, which is strict if at least one $\ell_p < m_p$.

The operator norm of the diagonal matrix **D** is thus equal to the largest diagonal entry, say, in row $\vec{\sigma}$. Similar to above, we then sum over $\sigma^2, \ldots, \sigma^P$ to get an explicit upper bound on $\|\mathbf{D}\|$:

$$\|\mathbf{D}\| = (\mathbf{D})_{\vec{\sigma}, \vec{\sigma}} \tag{A.41}$$

$$\leq \sum_{\sigma^2, \dots, \sigma^P} (\mathbf{D})_{\vec{\sigma}, \vec{\sigma}} \tag{A.42}$$

$$= \sum_{\sigma^2,\dots,\sigma^P} \left(cg \mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} - b \mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P} \right)_{\vec{\sigma},\vec{\sigma}}$$
(A.43)

$$= cg \sum_{\sigma^2,\dots,\sigma^P} \left(\mathbf{N}_{\ell_1,\dots,\ell_P}^T \mathbf{N}_{\ell_1,\dots,\ell_P} \right)_{\vec{\sigma},\vec{\sigma}} - b \sum_{\sigma^2,\dots,\sigma^P} \left(\mathbf{N}_{A_1\dots A_P}^T \mathbf{N}_{A_1\dots A_P} \right)_{\vec{\sigma},\vec{\sigma}}$$
(A.44)

$$= cgq^{-2LK}K!^{P} \prod_{p=1}^{P} {q^{2\ell_{p}} + K - 1 \choose K} - bq^{-2MK}K!^{P} \prod_{p=1}^{P} {q^{2m_{p}} + K - 1 \choose K}, \quad (A.45)$$

where we used the identity (A.31) for the second sum in (A.44), and with the replacement $m_p \to \ell_p$ for the first sum therein.

Substituting (A.32) and (A.45) in the bound (A.21) on the subspace angle finally gives

$$c(\operatorname{im} \mathcal{P}, \operatorname{im} \mathcal{Q}_{\operatorname{mct}})^{2} \leq acgq^{-2LK}K!^{P} \prod_{p=1}^{P} \binom{q^{2\ell_{p}} + K - 1}{K} - 2abq^{-2MK}K!^{P} \prod_{p=1}^{P} \binom{q^{2m_{p}} + K - 1}{K} + abq^{-2MK}K!^{2P} \prod_{p=1}^{P} \binom{q^{m_{p}} + K - 1}{K}^{2},$$

$$(A.46)$$

which concludes the proof.

The following auxiliary lemma is used in the proof of Theorem 4.6. It expresses the Frobenius inner products of operators appearing therein in terms of inner products of vectors in $\mathbb{C}^{K!^P}$ acted upon by certain (Gram) matrices whose elements are inner products of permutation operators.

Lemma A.1. Let $K, q, P, \ell_1, \ldots, \ell_P, m_1, \ldots, m_P$ be as in Theorem 4.6, and consider operators $X, Z \in \mathcal{L}(A_1^K \ldots A_P^K)$ of the form

$$X = \sum_{\sigma^1, \dots, \sigma^P \in \mathfrak{S}_k} x_{\sigma^1, \dots, \sigma^P} \, \sigma^1_{A_1} \otimes \dots \otimes \sigma^P_{A_P} \quad \text{for some } x_{\sigma^1, \dots, \sigma^P} \in \mathbb{C}, \tag{A.47}$$

$$Z = \sum_{\sigma^1, \dots, \sigma^P \in \mathfrak{S}_k} z_{\sigma^1, \dots, \sigma^P} \, \sigma^1_{A_1} \otimes \dots \otimes \sigma^P_{A_P} \quad \text{for some } z_{\sigma^1, \dots, \sigma^P} \in \mathbb{C}. \tag{A.48}$$

Denote by $\mathbf{x}, \mathbf{z} \in \mathbb{C}^{K!^P}$ the vectors with coefficients $(\mathbf{x})_{\sigma^1,\dots,\sigma^P} = x_{\sigma^1,\dots,\sigma^P}$ and $(\mathbf{z})_{\sigma^1,\dots,\sigma^P} = z_{\sigma^1,\dots,\sigma^P}$, respectively.

(i)
$$\langle X, Z \rangle_{n} = \langle \mathbf{x}, (\mathbf{G}_{A_{1}} \otimes \cdots \otimes \mathbf{G}_{A_{P}}) \mathbf{z} \rangle$$

(ii) If
$$K \leq |A_{p,k}| = q^{m_p}$$
 for all $p = 1, ..., P$ and $K \leq |A_1^{\ell_1} ... A_p^{\ell_p}| = q^{\ell_1 + ... + \ell_P}$,
$$\langle X, \mathcal{P} \mathcal{Q}_{\text{mct}} \mathcal{P}(X) \rangle_{\text{n}} = \left\langle \mathbf{x}, \left(\mathbf{G}_{A_1^{m_1 - \ell_1}} \otimes \cdots \otimes \mathbf{G}_{A_p^{m_P - \ell_P}} \right) \odot \left(\mathbf{N}_{\ell_1, ..., \ell_P}^T \mathbf{G}_{\ell_1, ..., \ell_P}^{-1} \mathbf{N}_{\ell_1, ..., \ell_P} \right) \mathbf{x} \right\rangle, \tag{A.49}$$

where $\mathbf{G}_{\ell_1,\dots,\ell_P} \equiv \mathbf{G}_{A_1^{\ell_1}\dots A_P^{\ell_P}}$ is the Gram matrix of permutation operators acting on the systems $A_1^{\ell_1}\dots A_P^{\ell_P}$, and $\mathbf{N}_{\ell_1,\dots,\ell_P}$ is a $(K!\times K!^P)$ -matrix with coefficients

$$(\mathbf{N}_{\ell_1,\dots,\ell_P})_{\tau,(\sigma^1,\dots,\sigma^P)} = \left\langle \tau_{A_1^{\ell_1}\dots A_P^{\ell_P}}, \sigma_{A_1^{\ell_1}}^1 \otimes \dots \otimes \sigma_{A_P^{\ell_P}}^P \right\rangle_{\mathbf{n}}.$$
 (A.50)

(iii) If $K \leq |A_1 \dots A_P| = q^{m_1 + \dots + m_P}$, then

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \left\langle \mathbf{x}, \mathbf{N}_{A_1 \dots A_P}^T \mathbf{G}_{A_1 \dots A_P}^{-1} \mathbf{N}_{A_1 \dots A_P} \mathbf{x} \right\rangle,$$
 (A.51)

where $\mathbf{N}_{A_1...A_P}$ is a $(K! \times K!^P)$ -matrix with coefficients

$$(\mathbf{N}_{A_1...A_P})_{\tau,(\sigma^1,...,\sigma^P)} = \langle \tau_{A_1...A_P}, \sigma^1_{A_1} \otimes \cdots \otimes \sigma^P_{A_P} \rangle_{\mathbf{n}}. \tag{A.52}$$

For the proof of Lemma A.1 we introduce the following shorthand notation to streamline the presentation. For $\sigma \in \mathfrak{S}_k$ and $p = 1, \ldots, P$, we write

$$\sigma_p \equiv \sigma_{A_p}$$
 $\sigma_{\ell_p} \equiv \sigma_{A_p^{\ell_p}}$ $\sigma_{m_p - \ell_p} \equiv \sigma_{A_n^{m_p - \ell_p}}.$ (A.53)

In the proofs below we will index different permutations like σ^p for $p=1,\ldots,P$. For the corresponding permutation operators $\sigma^p_{A_p}$ or $\sigma^p_{A_p^{\ell_p}}$ we then write σ^p_p and $\sigma^p_{\ell_p}$, respectively. Note that in σ^p_p the superscript p labels the permutation, while the subscript p indicates the p-th collection of A-systems (or 'column' in the pictorial language of Fig. 1) on which σ^p_p acts. For multipartite permutation operators we often skip system labels too, e.g.,

$$\sigma_{A_1...A_P} \equiv \sigma_{1...P} \qquad \sigma_{A_1^{\ell_1}...A_P^{\ell_P}} \equiv \sigma_{\ell_1,...,\ell_P} \qquad \sigma_{A_1^{m_1-\ell_1}...A_P^{m_P-\ell_P}} \equiv \sigma_{m_1-\ell_1,...,m_P-\ell_P}$$
(A.54)

We also write $\vec{\sigma} = (\sigma^1, \dots, \sigma^P)$ for a tuple of permutations $\sigma^p \in \mathfrak{S}_k$ and $p = 1, \dots, P$.

Proof of Lemma A.1(i). This is a straightforward generalization of the proof of Lem. 4.5(i). With the assumptions on X and Z, we calculate:

$$\langle X, Z \rangle_{\mathbf{n}} = \sum_{\vec{\omega}, \vec{\sigma}} x_{\vec{\omega}}^* z_{\vec{\sigma}} \langle \omega_1^1 \otimes \cdots \otimes \omega_P^P, \sigma_1^1 \otimes \cdots \otimes \sigma_P^P \rangle_{\mathbf{n}}$$
(A.55)

$$= \sum_{\vec{\omega}, \vec{\sigma}} x_{\vec{\omega}}^* z_{\vec{\sigma}} \left\langle \omega_1^1, \sigma_1^1 \right\rangle_{\mathbf{n}} \dots \left\langle \omega_P^P, \sigma_P^P \right\rangle_{\mathbf{n}}$$
(A.56)

$$= \sum_{\vec{\omega},\vec{\sigma}} x_{\vec{\omega}}^* z_{\vec{\sigma}} (\mathbf{G}_{A_1})_{\omega^1,\sigma^1} \dots (\mathbf{G}_{A_P})_{\omega^P,\sigma^P}$$
(A.57)

$$= \sum_{\vec{\sigma}} x_{\vec{\omega}}^* \sum_{\vec{\sigma}} (\mathbf{G}_{A_1} \otimes \cdots \otimes \mathbf{G}_{A_P})_{\vec{\omega}, \vec{\sigma}} z_{\vec{\sigma}}$$
(A.58)

$$= \langle \mathbf{x}, (\mathbf{G}_{A_1} \otimes \cdots \otimes \mathbf{G}_{A_P}) \mathbf{z} \rangle, \tag{A.59}$$

which concludes the proof.

Proof of Lemma A.1(ii). Recall that we have

$$X = \mathcal{P}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \, \sigma_1^1 \otimes \dots \otimes \sigma_P^P \tag{A.60}$$

for some coefficients $x_{\vec{\sigma}} \in \mathbb{C}$. Applying the crosstwirl \mathcal{Q}_{mct} to an operator $\sigma^1_{\ell_1} \otimes \cdots \otimes \sigma^P_{\ell_P}$ gives

$$Q_{\text{mct}}\left(\sigma_{\ell_1}^1 \otimes \cdots \otimes \sigma_{\ell_P}^P\right) = \sum_{\tau} y_{\tau}^{\vec{\sigma}} \, \tau_{\ell_1,\dots,\ell_P} = \sum_{\tau} y_{\tau}^{\vec{\sigma}} \, \tau_{\ell_1} \otimes \cdots \otimes \tau_{\ell_P} \tag{A.61}$$

for some coefficients $y_{\tau}^{\vec{\sigma}}$. Applying the relation (A.61) in (A.2), we obtain

$$Q_{\text{mct}}\mathcal{P}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} Q_{\text{mct}} \left(\sigma_{\ell_1}^1 \otimes \cdots \otimes \sigma_{\ell_P}^P \right) \otimes \sigma_{m_1 - \ell_1}^1 \otimes \dots \sigma_{m_P - \ell_P}^P$$
(A.62)

$$= \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} \tau_{\ell_1} \otimes \sigma_{m_1 - \ell_1}^1 \otimes \cdots \otimes \tau_{\ell_P} \otimes \sigma_{m_P - \ell_P}^P. \tag{A.63}$$

Another application of \mathcal{P} to this operator yields

$$\mathcal{PQ}_{\mathrm{mct}}\mathcal{P}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} \mathcal{P}\left(\tau_{\ell_{1}} \otimes \sigma_{m_{1}-\ell_{1}}^{1} \otimes \cdots \otimes \tau_{\ell_{P}} \otimes \sigma_{m_{P}-\ell_{P}}^{P}\right)$$
(A.64)

$$= \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} \mathcal{T}_{A_1} \left(\tau_{\ell_1} \otimes \sigma_{m_1 - \ell_1}^1 \right) \otimes \cdots \otimes \mathcal{T}_{A_P} \left(\tau_{\ell_P} \otimes \sigma_{m_P - \ell_P}^P \right)$$
(A.65)

$$= \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} \sum_{\vec{\omega}} z_{\omega^{1}}^{\tau,\sigma^{1}} \dots z_{\omega^{P}}^{\tau,\sigma^{P}} \omega_{1}^{1} \otimes \dots \otimes \omega_{P}^{P}$$
(A.66)

$$= \sum_{\vec{\omega}} \left(\sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} z_{\omega^{1}}^{\tau,\sigma^{1}} \dots z_{\omega^{P}}^{\tau,\sigma^{P}} \right) \omega_{1}^{1} \otimes \dots \otimes \omega_{P}^{P}, \tag{A.67}$$

with coefficients $z_*^{*,*}$ defined just as in the proof of Lemma 4.5(iii) via

$$\mathcal{T}_{A_p}\left(\tau_{\ell_p}\otimes\sigma_{m_p-\ell_p}^p\right) = \sum_{\omega^p} z_{\omega^p}^{\tau,\sigma^p} \omega_p^p. \tag{A.68}$$

Defining vectors $\mathbf{s},\mathbf{x}\in\mathbb{C}^{K!^p}$ via

$$(\mathbf{x})_{(\vec{\sigma})} = x_{\vec{\sigma}} \qquad (\mathbf{s})_{(\vec{\omega})} = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\tau} y_{\tau}^{\vec{\sigma}} z_{\omega^{1}}^{\tau, \sigma^{1}} \dots z_{\omega^{P}}^{\tau, \sigma^{P}}, \qquad (A.69)$$

part (i) of the lemma gives

$$\langle X, \mathcal{PQ}_{\text{mct}}\mathcal{P}(X)\rangle = \langle \mathbf{x}, (\mathbf{G}_{A_1} \otimes \cdots \otimes \mathbf{G}_{A_P})\mathbf{s}\rangle = \langle \mathbf{x}, (\mathbf{G}_{A_1} \otimes \cdots \otimes \mathbf{G}_{A_P})\mathbf{L}\mathbf{x}\rangle, \tag{A.70}$$

where the inner product on the right is the standard one on $\mathbb{C}^{K!^p}$, and we defined the $(K!^p \times K!^p)$ matrix **L** with coefficients

$$(\mathbf{L})_{(\vec{\omega}),(\vec{\sigma})} = \sum_{\tau} y_{\tau}^{\vec{\sigma}} z_{\omega^1}^{\tau,\sigma^1} \dots z_{\omega^P}^{\tau,\sigma^P}. \tag{A.71}$$

The Gram matrices \mathbf{G}_{A_p} are invertible since $K \leq |A_p| = q^{m_p}$ for all p by assumption [20, Lem. 1]. Defining vectors $\mathbf{z}^{\tau,\sigma^p}, \mathbf{v}^{\tau,\sigma^p} \in \mathbb{C}^{K!}$ via $(\mathbf{z}^{\tau,\sigma^p})_{\omega^p} = z_{\omega^p}^{\tau,\sigma^p}$ and

$$(\mathbf{v}^{\tau,\sigma^p})_{\omega^p} = \left\langle \omega_p^p, \tau_{\ell_p} \otimes \sigma_{m_p - \ell_p}^p \right\rangle_{\mathbf{n}}, \tag{A.72}$$

we have the following vector equations for all p = 1, ..., P, which follow just as in the proof of Lemma 4.5(iii):

$$\mathbf{z}^{\tau,\sigma^p} = \mathbf{G}_{A_p}^{-1} \mathbf{v}^{\tau,\sigma^p} \tag{A.73}$$

We also rewrite the coefficients $y_{\tau}^{\vec{\sigma}}$ defined via (A.61):

$$w_{\chi}^{\vec{\sigma}} := \left\langle \chi_{\ell_1 \dots \ell_P}, \sigma_{\ell_1}^1 \otimes \dots \otimes \sigma_{\ell_P}^P \right\rangle_{\mathbf{n}} \tag{A.74}$$

$$= \left\langle \chi_{\ell_1 \dots \ell_P}, \mathcal{Q}_{\text{mct}} \left(\sigma_{\ell_1}^1 \otimes \dots \otimes \sigma_{\ell_P}^P \right) \right\rangle_{\mathbf{n}} \tag{A.75}$$

$$= \sum_{\tau} y_{\tau}^{\vec{\sigma}} \langle \chi_{\ell_1} \otimes \cdots \otimes \chi_{\ell_P}, \tau_{\ell_1} \otimes \cdots \otimes \tau_{\ell_P} \rangle_{\mathbf{n}}$$
(A.76)

$$= \sum_{\tau} (\mathbf{G}_{\ell_1,\dots,\ell_P})_{\chi,\tau} y_{\tau}^{\vec{\sigma}}, \tag{A.77}$$

where the second equality follows since $\chi_{\ell_1...\ell_P}$ is invariant under $\mathcal{Q}_{\mathrm{mct}}$, and $\mathbf{G}_{\ell_1,...,\ell_P} \equiv \mathbf{G}_{A_1^{\ell_1}...A_P^{\ell_P}}$ is the Gram matrix of permutation operators acting on $A_1^{\ell_1}...A_P^{\ell_P}$. If $K \leq |A_1^{\ell_1}...A_P^{\ell_P}| = q^{\ell_1+\cdots+\ell_P}$, then this matrix is also invertible, and we have

$$\mathbf{y}^{\vec{\sigma}} = \mathbf{G}_{\ell_1,\dots,\ell_P}^{-1} \mathbf{w}^{\vec{\sigma}} \tag{A.78}$$

for the vectors $(\mathbf{y}^{\vec{\sigma}})_{\tau} = y_{\tau}^{\vec{\sigma}}$ and $(\mathbf{w}^{\vec{\sigma}})_{\tau} = w_{\tau}^{\vec{\sigma}}$.

Using the expressions (A.73) and (A.78) in the definition (A.71) of the matrix \mathbf{L} and employing the same arguments as in the proof of Lemma 4.5(iii), we get

$$\mathbf{L} = \left(\mathbf{G}_{A_1}^{-1} \otimes \cdots \otimes \mathbf{G}_{A_P}^{-1}\right) \overline{\mathbf{L}},\tag{A.79}$$

where $\overline{\mathbf{L}}$ is a $(K!^P \times K!^P)$ -matrix with coefficients

 $(\overline{\mathbf{L}})_{(\vec{\omega}),(\vec{\sigma})}$

$$= \sum_{\tau,\chi} (\mathbf{G}_{\ell_1,\dots,\ell_P}^{-1})_{\tau,\chi} v_{\omega^1}^{\tau,\sigma^1} \dots v_{\omega^P}^{\tau,\sigma^P} w_{\chi}^{\vec{\sigma}}$$
(A.80)

$$= \sum_{\tau,\chi} (\mathbf{G}_{\ell_1,\dots,\ell_P}^{-1})_{\tau,\chi} \left(\prod_{p=1}^P \left\langle \omega_p^p, \tau_{\ell_p} \otimes \sigma_{m_p-\ell_p}^p \right\rangle_{\mathbf{n}} \right) \left\langle \chi_{\ell_1\dots\ell_P}, \sigma_{\ell_1}^1 \otimes \dots \otimes \sigma_{\ell_P}^P \right\rangle_{\mathbf{n}}$$
(A.81)

$$= \sum_{\tau,\chi} (\mathbf{G}_{\ell_1,\dots,\ell_P}^{-1})_{\tau,\chi} \left(\prod_{p=1}^P \left\langle \omega_{\ell_p}^p, \tau_{\ell_p} \right\rangle_{\mathbf{n}} \left\langle \omega_{m_p-\ell_p}^p, \sigma_{m_p-\ell_p}^p \right\rangle_{\mathbf{n}} \right) \left\langle \chi_{\ell_1\dots\ell_P}, \sigma_{\ell_1}^1 \otimes \dots \otimes \sigma_{\ell_P}^P \right\rangle_{\mathbf{n}} \quad (A.82)$$

$$= \prod_{p=1}^{P} \left\langle \omega_{m_{p}-\ell_{p}}^{p}, \sigma_{m_{p}-\ell_{p}}^{p} \right\rangle_{\mathbf{n}} \sum_{\tau} \left(\prod_{p=1}^{P} \left\langle \omega_{\ell_{p}}^{p}, \tau_{\ell_{p}} \right\rangle_{\mathbf{n}} \right) \sum_{\chi} (\mathbf{G}_{\ell_{1}, \dots, \ell_{P}}^{-1})_{\tau, \chi} \left(\prod_{p=1}^{P} \left\langle \chi_{\ell_{p}}, \sigma_{\ell_{p}}^{p} \right\rangle_{\mathbf{n}} \right). \tag{A.83}$$

This shows that $\overline{\mathbf{L}}$ is equal to the following Hadamard product:

$$\overline{\mathbf{L}} = \left(\mathbf{G}_{A_1^{m_1 - \ell_1}} \otimes \cdots \otimes \mathbf{G}_{A_p^{m_P - \ell_P}} \right) \odot \left(\mathbf{N}_{\ell_1, \dots, \ell_P}^T \mathbf{G}_{\ell_1, \dots, \ell_P}^{-1} \mathbf{N}_{\ell_1, \dots, \ell_P} \right), \tag{A.84}$$

where we defined the $(K! \times K!^P)$ -matrix $\mathbf{N}_{\ell_1,\dots,\ell_P}$ with coefficients

$$(\mathbf{N}_{\ell_1,\dots,\ell_P})_{\tau,\vec{\sigma}} = \left\langle \tau_{\ell_1,\dots,\ell_P}, \sigma^1_{\ell_1} \otimes \dots \otimes \sigma^P_{\ell_P} \right\rangle_{\mathbf{n}}. \tag{A.85}$$

Using (A.84) and (A.79) in (A.70), we obtain

$$\langle X, \mathcal{PQ}_{\text{mct}}\mathcal{P}(X) \rangle = \left\langle \mathbf{x}, \left(\mathbf{G}_{A_1^{m_1 - \ell_1}} \otimes \cdots \otimes \mathbf{G}_{A_P^{m_P - \ell_P}} \right) \odot \left(\mathbf{N}_{\ell_1, \dots, \ell_P}^T \mathbf{G}_{\ell_1, \dots, \ell_P}^{-1} \mathbf{N}_{\ell_1, \dots, \ell_P} \right) \mathbf{x} \right\rangle, \tag{A.86}$$

which proves the claim.

Proof of Lemma A.1(iii). Once again, let $X = \mathcal{P}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sigma_1^1 \otimes \cdots \otimes \sigma_P^P$ for some coefficients $x_{\vec{\sigma}} \in \mathbb{C}$ by assumption of the lemma. We define coefficients $r_{\vec{\sigma}}^{\vec{\sigma}}$ via

$$\mathcal{R}\left(\sigma_1^1 \otimes \dots \otimes \sigma_P^P\right) = \sum_{\pi} r_{\pi}^{\vec{\sigma}} \, \pi_{1\dots P},\tag{A.87}$$

so that we can write

$$\mathcal{R}(X) = \sum_{\vec{\sigma}} x_{\vec{\sigma}} \sum_{\pi} r_{\pi}^{\vec{\sigma}} \pi_{1...P}.$$
 (A.88)

Taking the normalized inner product with X gives

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \sum_{\vec{\sigma}, \vec{\sigma}, \pi} x_{\vec{\sigma}}^* x_{\vec{\sigma}} r_{\pi}^{\vec{\sigma}} \langle \omega_1^1 \otimes \cdots \otimes \omega_P^P, \pi_{1...P} \rangle_{\mathbf{n}}$$
(A.89)

$$= \sum_{\vec{\omega}, \vec{\sigma}, \pi} x_{\vec{\omega}}^* x_{\vec{\sigma}} r_{\pi}^{\vec{\sigma}} t_{\pi}^{\vec{\omega}}, \tag{A.90}$$

where we defined the coefficients

$$t_{\pi}^{\vec{\omega}} = \left\langle \pi_{1\dots P}, \omega_1^1 \otimes \dots \otimes \omega_P^P \right\rangle_{\mathbf{n}} \tag{A.91}$$

$$= \left\langle \mathcal{R}(\pi_{1\dots P}), \omega_1^1 \otimes \dots \otimes \omega_P^P \right\rangle_{\mathbf{n}} \tag{A.92}$$

$$= \left\langle \pi_{1\dots P}, \mathcal{R} \left(\omega_1^1 \otimes \dots \otimes \omega_P^P \right) \right\rangle_{\mathbf{n}} \tag{A.93}$$

$$= \sum_{\rho} r_{\rho}^{\vec{\omega}} \langle \pi_{1\dots P}, \rho_{1\dots P} \rangle_{\mathbf{n}} \tag{A.94}$$

$$= \sum_{\rho} (\mathbf{G}_{A_1...A_P})_{\pi,\rho} r_{\rho}^{\vec{\omega}}. \tag{A.95}$$

Here, we used the invariance of $\pi_{1...P}$ under \mathcal{R} in the second equality, and the definition (A.87) of the r_*^* coefficients in fourth equality.

Defining vectors $\mathbf{r}^{\vec{\sigma}}$ with $(\mathbf{r}^{\vec{\sigma}})_{\pi} = r_{\pi}^{\vec{\sigma}}$ and $\mathbf{t}^{\vec{\sigma}}$ with $(\mathbf{t}^{\vec{\sigma}})_{\pi} = t_{\pi}^{\vec{\sigma}}$, we then have $\mathbf{t}^{\vec{\sigma}} = \mathbf{G}_{A_1...A_P} \mathbf{r}^{\vec{\sigma}}$. If $K \leq |A_1...A_P| = q^{m_1+...m_P}$, then $\mathbf{G}_{A_1...A_P}$ is invertible [20], and hence $\mathbf{r}^{\vec{\sigma}} = \mathbf{G}_{A_1...A_P}^{-1} \mathbf{t}^{\vec{\sigma}}$.

Using this in (A.90), we have

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \sum_{\vec{\omega}, \vec{\sigma}, \pi} x_{\vec{\omega}}^* x_{\vec{\sigma}} r_{\pi}^{\vec{\sigma}} t_{\pi}^{\vec{\omega}}$$
(A.96)

$$= \sum_{\vec{\omega}, \vec{\sigma}, \pi, \rho} x_{\vec{\omega}}^* x_{\vec{\sigma}} (\mathbf{G}_{A_1 \dots A_P}^{-1})_{\pi, \rho} t_{\rho}^{\vec{\sigma}} t_{\pi}^{\vec{\omega}}$$
(A.97)

$$= \sum_{\vec{\omega}} x_{\vec{\omega}}^* \sum_{\pi} t_{\pi}^{\vec{\omega}} \sum_{\rho} (\mathbf{G}_{A_1 \dots A_P}^{-1})_{\pi, \rho} \sum_{\vec{\sigma}} t_{\rho}^{\vec{\sigma}} x_{\vec{\sigma}}, \tag{A.98}$$

which can be rewritten as

$$\langle X, \mathcal{R}(X) \rangle_{\mathbf{n}} = \left\langle \mathbf{x}, \mathbf{N}_{A_1 \dots A_P}^T \mathbf{G}_{A_1 \dots A_P}^{-1} \mathbf{N}_{A_1 \dots A_P} \mathbf{x} \right\rangle$$
 (A.99)

in terms of the $(K! \times K!^P)$ -matrix $\mathbf{N}_{A_1...A_P}$ with coefficients

$$(\mathbf{N}_{A_1...A_P})_{\pi\vec{\sigma}} = t_{\pi}^{\vec{\sigma}} = \langle \pi_{1...P}, \sigma_1^1 \otimes \cdots \otimes \sigma_P^P \rangle_{\mathbf{n}}. \tag{A.100}$$

This concludes the proof.

References

- [1] S. Aaronson. "The Complexity of Quantum States and Transformations: From Quantum Money to Black Holes". arXiv preprint (July 2016). arXiv: 1607.05256 [quant-ph].
- [2] S. Aaronson, A. Bouland, B. Fefferman, S. Ghosh, U. Vazirani, C. Zhang, and Z. Zhou. "Quantum Pseudoentanglement". arXiv preprint (Apr. 2023). arXiv: arXiv: 2211.00747 [quant-ph].
- [3] F. G. S. L. Brandão, A. W. Harrow, and M. Horodecki. "Local Random Quantum Circuits are Approximate Polynomial-Designs". *Communications in Mathematical Physics* 346.2 (Sept. 2016), pp. 397–434. arXiv: 1208.0692 [quant-ph].
- [4] F. G. Brandão, W. Chemissany, N. Hunter-Jones, R. Kueng, and J. Preskill. "Models of Quantum Complexity Growth". PRX Quantum 2.3 (July 29, 2021), p. 030316. arXiv: 1912.04297 [quant-ph].
- [5] W. Brown and O. Fawzi. "Decoupling with Random Quantum Circuits". Communications in Mathematical Physics 340.3 (Dec. 2015), pp. 867–900. arXiv: 1307.0632 [quant-ph].
- [6] C.-F. Chen, A. Bouland, F. G. S. L. Brandão, J. Docter, P. Hayden, and M. Xu. "Efficient unitary designs and pseudorandom unitaries from permutations". arXiv preprint (Apr. 2024). arXiv: 2404.16751 [quant-ph].
- [7] C.-F. Chen, J. Haah, J. Haferkamp, Y. Liu, T. Metger, and X. Tan. "Incompressibility and spectral gaps of random circuits". arXiv preprint (2024). arXiv: 2406.07478 [quant-ph].
- [8] M. Christandl. "The structure of bipartite quantum states-insights from group theory and cryptography". PhD thesis. University of Cambridge, 2006. arXiv: quant-ph/0604183.
- [9] J. Cotler, N. Hunter-Jones, and D. Ranard. "Fluctuations of subsystem entropies at late times". *Physical Review A* 105.2 (Feb. 2022), p. 022416. arXiv: 2010.11922 [quant-ph].

REFERENCES 53

- [10] A. Deshpande, P. Niroula, O. Shtanko, A. V. Gorshkov, B. Fefferman, and M. J. Gullans. "Tight Bounds on the Convergence of Noisy Random Circuits to the Uniform Distribution". PRX Quantum 3.4 (Dec. 2022), p. 040329. arXiv: 2112.00716 [quant-ph].
- [11] F. Deutsch. *Best Approximation in Inner Product Spaces*. New York, NY: Springer New York, 2001.
- [12] F. Dupuis. "The decoupling approach to quantum information theory". PhD thesis. Université de Montréal, 2010. arXiv: 1004.1641 [quant-ph].
- [13] J. Eisert, M. Cramer, and M. B. Plenio. "Colloquium: Area laws for the entanglement entropy". Reviews of Modern Physics 82 (1 Feb. 2010), pp. 277–306. arXiv: 0808.3773 [quant-ph].
- [14] W. Fulton and J. Harris. Representation theory: a first course. Vol. 129. Springer Science & Business Media, 2013.
- [15] L. Gao, M. Junge, and N. LaRacuente. "Fisher Information and Logarithmic Sobolev Inequality for Matrix-Valued Functions". *Annales Henri Poincaré* 21.11 (Nov. 2020), pp. 3409–3478. arXiv: 1807.08838 [quant-ph].
- [16] L. Gao, M. Junge, and N. LaRacuente. "Relative entropy for von Neumann subalgebras". International Journal of Mathematics 31.06 (June 2020), p. 2050046. arXiv: 1909.01906 [quant-ph].
- [17] L. Gao and C. Rouzé. "Complete Entropic Inequalities for Quantum Markov Chains". Archive for Rational Mechanics and Analysis 245.1 (July 1, 2022), pp. 183–238. arXiv: 2102.04146 [quant-ph].
- [18] J. Haah, Y. Liu, and X. Tan. "Efficient approximate unitary designs from random Pauli rotations". arXiv preprint (2024). arXiv: 2402.05239 [quant-ph].
- [19] J. Haferkamp and N. Hunter-Jones. "Improved spectral gaps for random quantum circuits: Large local dimensions and all-to-all interactions". *Physical Review A* 104.2 (Aug. 2021), p. 022417. arXiv: 2012.05259 [quant-ph].
- [20] A. W. Harrow. "Approximate orthogonality of permutation operators, with application to quantum information". Letters in Mathematical Physics 114.1 (Dec. 2023), p. 1. arXiv: 2309.00715 [quant-ph].
- [21] A. W. Harrow and R. A. Low. "Random Quantum Circuits are Approximate 2-designs". Communications in Mathematical Physics 291.1 (Oct. 1, 2009), pp. 257–302. arXiv: 0802. 1919 [quant-ph].
- [22] A. W. Harrow and S. Mehraban. "Approximate Unitary t-Designs by Short Random Quantum Circuits Using Nearest-Neighbor and Long-Range Gates". Communications in Mathematical Physics 401.2 (July 2023), pp. 1531–1626. arXiv: 1809.06957 [quant-ph].
- [23] P. Hayden, M. Horodecki, A. Winter, and J. Yard. "A Decoupling Approach to the Quantum Capacity". *Open Systems & Information Dynamics* 15.01 (Mar. 2008), pp. 7–19. arXiv: quant-ph/0702005.
- [24] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, 2012.
- [25] R. A. Horn and C. R. Johnson. *Topics in Matrix Analysis*. 1st ed. Cambridge University Press, Apr. 1991.

54 REFERENCES

- [26] N. Hunter-Jones. "Unitary designs from statistical mechanics in random quantum circuits". arXiv preprint (May 2019). arXiv: 1905.12053 [quant-ph].
- [27] N. LaRacuente. "Quasi-factorization and multiplicative comparison of subalgebra-relative entropy". *Journal of Mathematical Physics* 63.12 (Dec. 2022), p. 122203. arXiv: 1912.00983 [quant-ph].
- [28] R. A. Low. "Large deviation bounds for k -designs". Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences 465.2111 (Nov. 2009), pp. 3289–3308.
- [29] A. A. Mele. "Introduction to Haar Measure Tools in Quantum Information: A Beginner's Tutorial". Quantum 8 (May 8, 2024), p. 1340. arXiv: 2307.08956 [quant-ph].
- [30] T. Metger, A. Poremba, M. Sinha, and H. Yuen. "Simple constructions of linear-depth t-designs and pseudorandom unitaries". arXiv preprint (Apr. 2024). arXiv: 2404.12647 [quant-ph].
- [31] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink. "Bernoulli's Inequality". *Classical and New Inequalities in Analysis*. Dordrecht: Springer Netherlands, 1993, pp. 65–81.
- [32] M. Müller-Lennert, F. Dupuis, O. Szehr, S. Fehr, and M. Tomamichel. "On quantum Rényi entropies: A new generalization and some properties". *Journal of Mathematical Physics* 54.12, 122203 (2013). arXiv: 1306.3142 [quant-ph].
- [33] Y. Nakata, C. Hirche, M. Koashi, and A. Winter. "Efficient Quantum Pseudorandomness with Nearly Time-Independent Hamiltonian Dynamics". *Physical Review X* 7.2 (Apr. 2017), p. 021006. arXiv: 1609.07021 [quant-ph].
- [34] J. V. Neumann. "On Rings of Operators. Reduction Theory". The Annals of Mathematics 50.2 (Apr. 1949), p. 401.
- [35] M. A. Nielsen and I. L. Chuang. *Quantum computation and quantum information*. eng. 10th anniversary edition. Cambridge: Cambridge university press, 2010.
- [36] M. Pimsner and S. Popa. "Entropy and index for subfactors". Annales scientifiques de l'École Normale Supérieure 19.1 (1986), pp. 57–106.
- [37] L. Piroli, C. Sünderhauf, and X.-L. Qi. "A Random Unitary Circuit Model for Black Hole Evaporation". *Journal of High Energy Physics* 2020.4 (Apr. 2020), p. 63. arXiv: 2002.09236 [hep-th].
- [38] T. Schuster, J. Haferkamp, and H.-Y. Huang. "Random unitaries in extremely low depth". arXiv preprint (2024).
- [39] H. Umegaki. "Conditional expectation in an operator algebra, IV (entropy and information)". Kodai Mathematical Seminar Reports. Vol. 14. 2. Department of Mathematics, Tokyo Institute of Technology. 1962, pp. 59–85.
- [40] M. M. Wilde, A. Winter, and D. Yang. "Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy". *Communications in Mathematical Physics* 331.2 (2014), pp. 593–622. arXiv: 1306.1586 [quant-ph].

REFERENCES 55

Nicholas La
Racuente, Department of Computer Science, University of Indiana Bloomington,
700 N Woodlawn Ave, Bloomington, IN, 47408, USA

Email address: nlaracu@iu.edu

Felix Leditzky, Department of Mathematics and IQUIST, University of Illinois Urbana-Champaign, 1409 W Green St, Urbana, IL 61801, USA

 $Email\ address {\tt : leditzky@illinois.edu}$