

Uncovering Emergent Spacetime Supersymmetry with Rydberg Atom Arrays

Chengshu Li,^{1,*} Shang Liu,^{2,*} Hanteng Wang,¹ Wenjun Zhang,³ Zi-Xiang Li,^{4,5} Hui Zhai,^{1,6} and Yingfei Gu^{1,†}

¹*Institute for Advanced Study, Tsinghua University, Beijing, 100084, China*

²*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, California 93106, USA*

³*Department of Physics and State Key Laboratory of Low Dimensional Quantum Physics, Tsinghua University, Beijing, 100084, China*

⁴*Beijing National Laboratory for Condensed Matter Physics and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*

⁵*University of Chinese Academy of Sciences, Beijing 100049, China*

⁶*Hefei National Laboratory, Hefei 230088, China*

(Dated: July 11, 2024)

In the zoo of emergent symmetries in quantum many-body physics, the previously unrealized emergent spacetime supersymmetry (SUSY) is particularly intriguing. Although it was known that spacetime SUSY could emerge at the (1+1)d tricritical Ising transition, an experimental realization is still absent. In this letter, we propose to realize the tricritical Ising transition with Rydberg atom arrays, taking advantage of the reconfigurability of these systems. In such systems, the spacetime SUSY manifests itself in the respective correlation functions of a bosonic mode and its fermionic partner. However, the correlation function of the fermionic mode inevitably involves a string operator, making direct measurement challenging in the conventional setting. Here, we utilize the analog-digital hybrid nature of the Rydberg atom arrays, which can simulate a physical Hamiltonian and perform a digital quantum circuit on the same platform, to measure the correlation function of the fermionic mode. This hybridized protocol provides an experimentally feasible way to reveal the hidden structure of the spacetime SUSY that emerges at the tricritical Ising transition.

Introduction. Spacetime supersymmetry (SUSY) was proposed in the '70s as a new type of symmetry of spacetime and fundamental particles [1, 2]. It exchanges bosons and fermions and provides appealing solutions to long-standing problems in particle physics, such as the hierarchy problem [3]. However, the experimental evidence of SUSY has not been observed up to the highest energy we can reach in particle accelerators so far. An opposite strategy is to realize SUSY as an emergent symmetry at low energy in quantum many-body systems – a fascinating phenomenon where the low-energy states possess larger symmetries than the physical Hamiltonian. This phenomenon often occurs when the system is fine-tuned to a quantum critical point [4].

Early in the '80s, it was known that spacetime SUSY could emerge at (1+1) dimensional tricritical Ising transition [5, 6]. From the Landau–Ginzburg perspective, the conventional second-order Ising transition is described by the sign change of the quadratic term in a ϕ^4 theory, where ϕ is the order parameter. For a more general free energy functional $\alpha\phi^2 + \beta\phi^4 + \phi^6$, a first-order transition could occur at a critical line with $\alpha > 0$ and $\beta < 0$. The aforementioned tricritical point refers to the meeting point of the above two transitions, i.e. $\alpha = \beta = 0$.

Continuous efforts have been made to realize the tricritical Ising transition [7–14] (or other form of SUSY theories [15–19]). However, an experimental identification of spacetime SUSY is still missing. In addition to the challenges in fine-tuning the model to the tricritical point where the SUSY could emerge, there is another fundamental difficulty in the characterization: the fermionic modes that emerge from the underlying bosonic physi-

cal degrees of freedom inevitably involve non-local string operators, whose detection often requires unconventional measurement schemes [20–23].

In the present letter, we propose to overcome these challenges with the Rydberg atom array that has shown to be powerful in simulating quantum phases of matter [24–30] and quantum information processing [31–37]. We demonstrate that the geometric reconfigurability of the Rydberg atom array facilitates the realization of a tricritical Ising point. Furthermore, we point out that the analog–digital hybrid nature [33] of the platform is ideal for revealing the hidden structure of the emergent spacetime SUSY. To be clearer, the hybrid nature here refers to the ability to run analog quantum simulation and digital quantum circuits on the same platform. More explicitly, the analog mode is based on the pseudo-spin doublet consisting of one of the hyperfine levels in the electronic ground state manifold and one Rydberg state. The tricritical Ising transition will be realized in this mode. While in the digital mode, the data qubit is encoded into two hyperfine levels, with the Rydberg state only used to mediate the entangling gates [31]. This hybrid mode allows us to efficiently generate the desired quantum many-body wave function via analog mode and coherently extract the quantum correlation in the complex wave function with a digital quantum circuit. In particular, we show that this hybrid protocol can be used to measure non-local fermionic correlations to verify emergent SUSY.

Model. We consider a two-leg ladder geometry of atom arrays as shown in Fig. 1(a) where the horizontal and vertical separations between atoms are denoted by r_1 and

r_2 respectively. The two colors denote the two sets (A and B) of atoms subjected to different Rydberg excitation lasers. This scheme is readily realized in dual-species atom arrays [38–41], where the resonance frequencies of different Rydberg states are largely separated so that crosstalk between different excitations is negligible. For simplicity, although not necessary, we assume that two Rydberg excitations share equal coupling strength Ω and equal detuning Δ . In the following, we set the Rabi frequency $\Omega = 1$ as the energy unit. The interaction between two Rydberg atoms is the van der Waals interaction given by C_{XY}/r^6 ($X, Y = A$ or B). Here, we set $C_{AA} > 0$, and we define a length unit r_0 by $C_{AA}/r_0^6 = \Omega$.

The Hamiltonian of this two-leg ladder two-sets Rydberg array is given by

$$\hat{H} = \sum_i \left(\frac{\Omega}{2} \hat{\sigma}_i^x - \Delta \hat{n}_i \right) + \sum_{ij} \frac{C_{XY}}{r_{ij}^6} \hat{n}_i \hat{n}_j. \quad (1)$$

It acts on the pseudo-spin states $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ corresponding to the ground state $|g\rangle$ and the Rydberg state $|r\rangle$ of atom i . $\hat{n}_i = (1 + \hat{\sigma}_i^z)/2$ is the number of Rydberg excitations on site i . The index $i = (i_x, i_y)$ labels the position of each atom with $i_y = 1$ or 2 denoting the upper or lower leg, respectively. For i_x being odd or even, the atoms belong to A or B subsets. Depending on different quantum numbers of Rydberg states, the van der Waals interaction between them can be either repulsive or attractive.

Phase Diagram. This two-leg ladder system has a $\mathbb{Z}_2$ symmetry as the exchanging symmetry between the upper and the lower legs. Correspondingly, the ground state of this system could exhibit two phases distinguished by whether the $\mathbb{Z}_2$ symmetry is broken, with a first and/or second-order transition between them. If both transitions are present, the intersection of the two transition lines will be the desired tricritical Ising point we aim for.

To show that the above conception can be realized in our model with realistic experimental parameters, we set $C_{BB} = 1.6C_{AA}$ and $C_{AB} = C_{BA} = -0.97C_{AA}$ as an illustrating example [42] and obtain the phase diagram shown in Fig. 1(b). In the computation, we have also fixed the spacing between the upper and lower legs to be $r_2 = 0.5r_0$ so that the repulsive interaction between two Rydberg states with the same i_x is much larger than Ω and therefore blockades the double occupancy of Rydberg states in the same column. The left-over tuning parameters are detuning Δ and horizontal spacing r_1 , both are accurately accessible in the reconfigurable atom array platform.

In the phase diagram, the $\mathbb{Z}_2$ symmetry-breaking phase (dubbed “ferromagnetic”) occurs in the large Δ and small r_1 regime: the larger Δ induces more Rydberg excitation, while the smaller r_1 leads to a stronger attractive interaction between horizontal neighbors (note that $C_{AB} < 0$), which favors the configuration that all Rydberg excitations stay in the same row, resulting in the




FIG. 1. (a) Layout of our model. Two colors denote two sets of atoms subjected to different Rydberg excitation lasers. r_1 and r_2 denote the nearest separation of atoms along the horizontal and vertical directions. (b) The phase diagram computed for two-leg ladder geometry with 2×40 sites. The solid line denotes a second-order transition, and the dashed line denotes a first-order transition, with “TCI” denoting the tricritical point. “Ferro” and “Disordered” denote the symmetry-breaking phase and the symmetric phase. Here, the $\mathbb{Z}_2$ symmetry is the exchanging symmetry between the upper and the lower legs. Δ is in unit of Ω and r_1 is in unit of r_0 defined in the main text. Dots with a circle denote atoms in the Rydberg state. (c) The central charge c obtained from the entanglement entropy at different given values of r_1 . $c = 7/10$ is the central charge for the tricritical Ising point. The inset shows the fitting of the subsystem entanglement entropy $S(j)$ as a function of the subsystem size j using the Calabrese–Cardy formula. The phase diagram is obtained by the DMRG calculation with bond dimension 200.

$\mathbb{Z}_2$ symmetry breaking phase. Now, starting from the symmetry breaking phase, there are two ways to restore symmetry: (1) shifting toward a more negative value for detuning Δ , which suppresses the Rydberg excitations and ultimately arrives at a symmetric state without Rydberg excitation; (2) shifting towards larger horizontal separation r_1 , which reduces the attractive interaction and therefore unlocks the correlating effects between different columns. Along this direction, the ground state

approximates the product of a symmetric superposition of one Rydberg excitation on the upper row and one on the lower row. It turns out that the transition along path 1 tends to be first order while path 2 tends to be second order. Therefore, the turning point of these two phase boundaries realizes a tricritical Ising transition point on the (r_1, Δ) parameter space.

Regarding the methodology, the phase diagram Fig. 1(b) is determined by a combination of the order parameter and the subsystem entanglement entropy, both extracted from the ground state wavefunction numerically calculated by the density-matrix renormalization group (DMRG) method [43]. Here the order parameter we choose is the “magnetization” $n^- = n_1 - n_2$, where $n_{i_y} = L^{-1} \sum_{i_x=1}^L n_i$ with $i = (i_x, i_y)$ and $i_y = 1$ or 2. $n_1 - n_2 \neq 0$ indicates the $\mathbb{Z}_2$ symmetry breaking. The first-order transition is signaled by an abrupt drop in the order parameter. Therefore its location can be precisely determined. In contrast, across the second-order transition, the order parameter vanishes continuously and therefore does not trigger a sharp signal in numerics. To precisely determine the location of the second-order phase transition, we compute the central charge by fitting the entanglement entropy of the subsystem $S(j)$ to the Calabrese–Cardy formula [44] $S(j, L) = \frac{c}{3} \ln \left[\frac{L}{\pi} \sin \left(\frac{\pi j}{L} \right) \right] + S_0$ with j denoting the subsystem size and L the total size. A typical example is shown in the inset of Fig. 1(c). The central charge c obtained from the fitting exhibits a maximum at the second-order phase boundary. Fig. 1(c) shows the central charge c along the transition line. One can see that c approaches $1/2$ at the Ising transition line, consistent with the prediction of Ising conformal field theory (CFT), and c drops toward zero at the first-order transition. In between, c peaked at the tricritical point with the value 0.7, consistent with the prediction of supersymmetric CFT.

SUSY Manifestation. A key physical consequence of the emergent SUSY at the critical point is that the scaling dimensions of a low-lying bosonic mode and its fermionic superpartner differ by $1/2$. For a motivating and simplest example, let us consider a D dimensional (space-time dimension) supersymmetric theory of a free massless scalar paired with a massless free fermion, with Lagrangian $\mathcal{L}_{\text{SUSY}} = \mathcal{L}_B[\phi] + \mathcal{L}_F[\psi]$. From the bosonic part $\mathcal{L}_B = \int d^D \mathbf{r} (\partial \phi)^2$, we get the dimension of $[\phi]$ from equation $2[\phi] = D - 2$. Therefore, by dimensional analysis $\langle \phi(\mathbf{r}_i) \phi(\mathbf{r}_j) \rangle \sim 1/|\mathbf{r}_i - \mathbf{r}_j|^{D-2}$. In parallel, the fermionic partner ψ with $\mathcal{L}_F = \int d^D \mathbf{r} \bar{\psi} i \not{\partial} \psi$ satisfies $2[\psi] = D - 1$ and, consequently, $\langle \psi(\mathbf{r}_i) \psi(\mathbf{r}_j) \rangle \sim 1/|\mathbf{r}_i - \mathbf{r}_j|^{D-1}$. The point is that the exponents of these two decayed spatial correlation functions of power law differ by 1, which is a measurable consequence of the SUSY we will investigate in the following.

In contrast to the aforementioned free theory example, where SUSY is inherent, our microscopic model (1) is constructed based on pseudo-spin degrees of freedom,




FIG. 2. η_ε and η_ψ (see text for definition) along the second-order transition line is plotted for different r_1 . The inset plots the spatial correlation functions $\langle \varepsilon_i \varepsilon_j \rangle$ and $-(\langle \gamma'_i \gamma_j \rangle + \langle \gamma_i \gamma'_j \rangle)$ as a function of $|r_i - r_j|$, which is a linear function under the log-log plot. The slope of these two linear functions defines η_ε and η_ψ . The dashed line indicates that these two numbers differ by unit at the ending point of the second-order transition at $r_1 \approx 0.884$.

with SUSY emerging only at the tricritical point. In such a scenario, it is generally challenging to identify the SUSY pair of fields that result from a non-trivial renormalization group flow [45]. In this letter, we exploit the aforementioned $\mathbb{Z}_2$ symmetry to identify the SUSY pair in our model. In the field-theoretic description of the tricritical Ising transition, there is a $\mathbb{Z}_2$ -even real bosonic field ε representing the “energy”, and a natural candidate in our microscopic model is the sum of Rydberg excitation numbers on each rung, by an abuse of notation let us denote $\varepsilon_i = \hat{n}_i^+ = \hat{n}_{(i,1)} + \hat{n}_{(i,2)}$. Under SUSY, this $\mathbb{Z}_2$ -even operator should be related to a pair of $\mathbb{Z}_2$ -odd real fermion operators. A simple $\mathbb{Z}_2$ -odd lattice operator is $\hat{n}_i^- = \hat{n}_{(i,1)} - \hat{n}_{(i,2)}$. However, $\hat{n}_i^-$ itself is not yet a fermionic operator. To finish the construction, we need to attach a Jordan–Wigner string so that distant fermion operators anticommute. This gives two natural candidates:

$$\hat{\gamma}_i = \left(\prod_{j=1}^{i-1} \text{SWAP}_j \right) \hat{n}_i^-, \quad \hat{\gamma}'_i = \left(\prod_{j=1}^i \text{SWAP}_j \right) \hat{n}_i^-, \quad (2)$$

where the swap operator SWAP_j applies an exchange between the upper and the lower legs at site- j onto the quantum state. Their linear combination corresponds to the time reversal pair of fermion fields in the continuum field theory, i.e. $\psi = \hat{\gamma} - i\hat{\gamma}'$ and $\bar{\psi} = \hat{\gamma} + i\hat{\gamma}'$ with abuse of notation.

Now, to establish the SUSY relation, we need to check the exponents of the decayed spatial correlation function of the bosonic ε and fermionic ψ (or $\bar{\psi}$, as they have the same scaling dimension). For the former, we need to measure the correlation function of the local operator

$\varepsilon_i = \hat{n}_{(i,1)} + \hat{n}_{(i,2)}$, that is,

$$\langle \varepsilon_i \varepsilon_j \rangle \sim 1/|\mathbf{r}_i - \mathbf{r}_j|^{\eta_\varepsilon}, \quad (3)$$

and extract the exponent η_ε . For the fermionic operator ψ , we need to extract the exponent η_ψ from the following correlation function

$$\langle \psi_i \psi_j \rangle = -i (\langle \gamma'_i \gamma_j \rangle + \langle \gamma_i \gamma'_j \rangle) \sim 1/|\mathbf{r}_i - \mathbf{r}_j|^{\eta_\psi}, \quad (4)$$

where in the first step we have used the property that $\langle \gamma_i \gamma_j \rangle = \langle \gamma'_i \gamma'_j \rangle = 0$ enforced by the time-reversal symmetry. Note the correlation function $\langle \gamma'_i \gamma_j \rangle$ or $\langle \gamma_i \gamma'_j \rangle$ involves a non-local string operator, for instance

$$-\langle \gamma'_i \gamma_j \rangle = \langle \hat{n}_i^- \left(\prod_{k=i+1}^{j-1} \text{SWAP}_k \right) \hat{n}_j^- \rangle \quad (5)$$

The SUSY in our model manifests itself as $\eta_\psi - \eta_\varepsilon = 1$.

We show in Fig. 2 the numerical results for the equal-time correlation functions $\langle \varepsilon_i \varepsilon_j \rangle$ and $-(\langle \gamma'_i \gamma_j \rangle + \langle \gamma_i \gamma'_j \rangle)$. The inset confirms the power-law behavior of the two correlation functions at the tricritical point. From the slope of the log-log plot, we can extract the exponent as shown in Fig. 2. It verifies clearly that at the end point of the second-order transition line, these two powers η_ε and η_ψ differ by 1, which is direct evidence of the emergent SUSY. Away from the tricritical point, the relation fails.

Measuring fermionic correlation on Rydberg atom array. Measuring the non-local string operator in (5) with a set of SWAP is generally difficult. The situation is similar to the measurement of the second Renyi entropy where a many-body interference between two identical copies of quantum states is performed [46, 47]. Here, we need to interfere with two subsets of our single quantum state. To accomplish the measurement, we can utilize the hybrid nature of the Rydberg atom array platform to perform a quantum version of the “analog-to-digital converter” and then process the quantum data via a measurement circuit consisting of entangling gates between the upper and lower chains.

A quantum Analog-to-Digital Converter (qADC) and its partner quantum Digital-to-Analog Converter (qDAC) map between the “analog basis” $\{|g\rangle, |r\rangle\}$ (“ground state” and “Rydberg state”) and the “digital basis” $\{|0\rangle, |1\rangle\}$. In the Rydberg atom, these two Hilbert spaces are embedded in the rich atomic levels. For instance, in the ^{87}Rb atom, the digital basis can be chosen as the two hyperfine spin levels of the ground state, and one of them, say $|0\rangle$ can be used for the ground state $|g\rangle$ in the analog basis as well. For this three-level system, the qADC can be realized by stimulated Raman adiabatic passage (STIRAP) [48] where an additional laser is applied to coherently transfer the Rydberg state $|r\rangle$ to the hyperfine $|1\rangle$ state, while keeping the $|g\rangle = |0\rangle$ state invariant. In this protocol, an optimization between the qADC pulse and the Rydberg laser in the analog mode is




FIG. 3. (a) $|0\rangle$ and $|1\rangle$ are two hyperfine levels of the electronic ground state, and $|r\rangle$ is a Rydberg state. $|0\rangle = |g\rangle$ and $|r\rangle$ are used for quantum simulation and $|0\rangle$ and $|1\rangle$ form a basis for digital quantum computation. The blue line denotes the Rydberg coupling Ω in the many-body Hamiltonian (1), and the orange line represents the qADC coupling that transfers a many-body wave function into the quantum computation basis. (b) Schematic of the time sequence of turning off the Rydberg coupling and the AtoD pulse. (c) After the qADC, we use the quantum circuit shown in the box to process the quantum many-body wave function to measure the fermionic correlation function (4). The tables in the last row summarize the correspondence between the measurement outcome and the eigenvalue of SWAP and n_- .

expected to mitigate the dephasing error caused by the turning off of the Rydberg laser that couples the analog basis. Alternatively, one can also realize the qADC via a two-step process, named the coherent mapping protocol in Ref. [33], where the ground state $|g\rangle$ (identified as $|1\rangle$ in this protocol) is transferred to digital $|0\rangle$ and then $|r\rangle$ is transferred to $|1\rangle$.

Following the qADC, we can utilize the quantum computation architecture to process the many-body wave function in the digital basis where $|0\rangle$ represents the empty states and $|1\rangle$ represents the states with Rydberg excitation. For each rung in the ladder, we have three allowed states $|00\rangle$, $\frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$, $\frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$ with

eigenvalue 1, 1, -1 under SWAP respectively. To single out the last state, we run a measurement circuit shown as Fig. 3 (c) and collect the instance with measurement outcome $|11\rangle$ in the computation basis. Then the fermionic correlator (4) is obtained as

$$\langle (-1)^{\#|11\rangle \text{ pairs between } (i,j)} \hat{n}_i^- \hat{n}_j^- \rangle. \quad (6)$$

Namely, the measurement of $\langle \psi_i \psi_j \rangle$ amounts to averaging the measurements of two local operators $\hat{n}_i^- \hat{n}_j^-$, with an additional phase -1 assigned to all the instances with odd number of $|11\rangle$ between site i and j .

Summary and Outlook. In this work, we show that a two-leg ladder Rydberg array structure with two sets of Rydberg excitations is sufficient to realize the tricritical Ising transition where the spacetime SUSY emerges. This proposed setup is especially suitable for the dual-species Rydberg atom platforms where the two Rydberg excitations can be implemented without cross-talk. We would like to remark that unlike the single chain model discussed in literature [24, 49, 50] where the Ising symmetry is realized as a translation by one lattice spacing, here the Ising $\mathbb{Z}_2$ symmetry is realized as the exchanging between two rails of the ladder. Combining this geometric realization of the internal symmetry with the reconfigurability of the atom array, one can further study how various boundary conditions can be realized geometrically [51].

To manifest the emergent SUSY encoded in the many-body wavefunction, we leverage the analog-digital hybrid nature of the Rydberg atom platform. This involves directly loading the “experimental data” prepared by the analog mode to the “quantum computer” in the digital mode via a quantum Analog-to-Digital converter (qADC). By processing the data through quantum circuits, we can access the information that is difficult to retrieve in conventional experiments where the data are processed classically [52–54].

Additionally, we envision that a quantum Digital-to-Analog Converter (qDAC) holds potential in the quantum simulation of many-body physics. Using the quantum circuit module, more interesting initial states can be prepared and sent to the quantum simulator, opening new possibilities for exploring many-body quantum dynamics.

Acknowledgments. We thank Wenlan Chen, Xun Gao, Jiazhong Hu, Xiaodong Liu, Xiao-Liang Qi, Kunpeng Wang, Peng Xu and Tao Zhang for discussions. We acknowledge financial support from the Innovation Program for Quantum Science and Technology 2021ZD0302005, the NSFC Grant No. 12042505 and U23A6004, the National Key R&D Program of China 2023YFA1406702, and the Tsinghua University Dushi Program. S.L. is supported by the Gordon and Betty Moore Foundation under Grant No. GBMF8690 and the National Science Foundation under Grant No. NSF PHY-1748958. Z.X.L. is supported by the NSFC under Grant

No. 12347107. H.Z. acknowledges support from the XPLOER Prize. Y.G. acknowledges support from the DAMO Academy Young Fellow program. The DMRG calculations are performed using the ITensor library (v0.3) [55].

* These two authors contributed equally.

† guyingfei@tsinghua.edu.cn

- [1] J.-L. Gervais and B. Sakita, Field Theory Interpretation of Supergauges in Dual Models, *Nucl. Phys. B* **34**, 632 (1971).
- [2] J. Wess and B. Zumino, Supergauge Transformations in Four-Dimensions, *Nucl. Phys. B* **70**, 39 (1974).
- [3] S. Dimopoulos and H. Georgi, Softly Broken Supersymmetry and SU(5), *Nucl. Phys. B* **193**, 150 (1981).
- [4] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, 2011).
- [5] D. Friedan, Z. Qiu, and S. Shenker, Superconformal invariance in two dimensions and the tricritical Ising model, *Phys. Lett. B* **151**, 37 (1985).
- [6] A. B. Zamolodchikov, Conformal symmetry and multi-critical points in two-dimensional quantum field theory, *Sov. J. Nucl. Phys.* **44**, 529 (1986).
- [7] M. Blume, Theory of the First-Order Magnetic Phase Change in UO₂, *Phys. Rev.* **141**, 517 (1966).
- [8] H. Capel, On the possibility of first-order phase transitions in Ising systems of triplet ions with zero-field splitting, *Physica* **32**, 966 (1966).
- [9] F. C. Alcaraz, J. R. Drugowich de Felício, R. Köberle, and J. F. Stilck, Hamiltonian studies of the Blume-Emery-Griffiths model, *Phys. Rev. B* **32**, 7469 (1985).
- [10] T. Grover, D. N. Sheng, and A. Vishwanath, Emergent Space-Time Supersymmetry at the Boundary of a Topological Phase, *Science* **344**, 280 (2014).
- [11] A. Rahmani, X. Zhu, M. Franz, and I. Affleck, Emergent Supersymmetry from Strongly Interacting Majorana Zero Modes, *Phys. Rev. Lett.* **115**, 166401 (2015).
- [12] Z.-X. Li, Y.-F. Jiang, and H. Yao, Edge Quantum Criticality and Emergent Supersymmetry in Topological Phases, *Phys. Rev. Lett.* **119**, 107202 (2017).
- [13] E. O’Brien and P. Fendley, Lattice Supersymmetry and Order-Disorder Coexistence in the Tricritical Ising Model, *Phys. Rev. Lett.* **120**, 206403 (2018).
- [14] C. Li, H. Ebisu, S. Sahoo, Y. Oreg, and M. Franz, Coupled wire construction of a topological phase with chiral tricritical Ising edge modes, *Phys. Rev. B* **102**, 165123 (2020).
- [15] S.-S. Lee, Emergence of supersymmetry at a critical point of a lattice model, *Phys. Rev. B* **76**, 075103 (2007).
- [16] Y. Yu and K. Yang, Simulating the Wess-Zumino Supersymmetry Model in Optical Lattices, *Phys. Rev. Lett.* **105**, 150605 (2010).
- [17] S.-K. Jian, Y.-F. Jiang, and H. Yao, Emergent Spacetime Supersymmetry in 3D Weyl Semimetals and 2D Dirac Semimetals, *Phys. Rev. Lett.* **114**, 237001 (2015).
- [18] S.-K. Jian, C.-H. Lin, J. Maciejko, and H. Yao, Emergence of Supersymmetric Quantum Electrodynamics, *Phys. Rev. Lett.* **118**, 166802 (2017).
- [19] Z.-X. Li, A. Vaezi, C. B. Mendl, and H. Yao, Numerical observation of emergent spacetime supersymmetry at

- quantum criticality, *Sci. Adv.* **4**, eaau1463 (2018).
- [20] M. Endres, M. Cheneau, T. Fukuhara, C. Weitenberg, P. Schauß, C. Gross, L. Mazza, M. C. Bañuls, L. Pollet, I. Bloch, and S. Kuhr, Observation of Correlated Particle-Hole Pairs and String Order in Low-Dimensional Mott Insulators, *Science* **334**, 200 (2011).
- [21] T. A. Hilker, G. Salomon, F. Grusdt, A. Omran, M. Boll, E. Demler, I. Bloch, and C. Gross, Revealing hidden antiferromagnetic correlations in doped Hubbard chains via string correlators, *Science* **357**, 484 (2017).
- [22] C. S. Chiu, G. Ji, A. Bohrdt, M. Xu, M. Knap, E. Demler, F. Grusdt, M. Greiner, and D. Greif, String patterns in the doped Hubbard model, *Science* **365**, 251 (2019).
- [23] S. de Léséleuc, V. Lienhard, P. Scholl, D. Barredo, S. Weber, N. Lang, H. P. Büchler, T. Lahaye, and A. Browaeys, Observation of a symmetry-protected topological phase of interacting bosons with Rydberg atoms, *Science* **365**, 775 (2019).
- [24] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing many-body dynamics on a 51-atom quantum simulator, *Nature* **551**, 579 (2017).
- [25] A. Browaeys and T. Lahaye, Many-body physics with individually controlled Rydberg atoms, *Nat. Phys.* **16**, 132 (2020).
- [26] S. Ebadi, T. T. Wang, H. Levine, A. Keesling, G. Semeghini, A. Omran, D. Bluvstein, R. Samajdar, H. Pichler, W. W. Ho, S. Choi, S. Sachdev, M. Greiner, V. Vuletić, and M. D. Lukin, Quantum phases of matter on a 256-atom programmable quantum simulator, *Nature* **595**, 227 (2021).
- [27] P. Scholl, M. Schuler, H. J. Williams, A. A. Eberharter, D. Barredo, K.-N. Schymik, V. Lienhard, L.-P. Henry, T. C. Lang, T. Lahaye, A. M. Läuchli, and A. Browaeys, Quantum simulation of 2D antiferromagnets with hundreds of Rydberg atoms, *Nature* **595**, 233 (2021).
- [28] G. Semeghini, H. Levine, A. Keesling, S. Ebadi, T. T. Wang, D. Bluvstein, R. Verresen, H. Pichler, M. Kalinowski, R. Samajdar, A. Omran, S. Sachdev, A. Vishwanath, M. Greiner, V. Vuletić, and M. D. Lukin, Probing topological spin liquids on a programmable quantum simulator, *Science* **374**, 1242 (2021).
- [29] F. Fang, K. Wang, V. S. Liu, Y. Wang, R. Cimmino, J. Wei, M. Bintz, A. Parr, J. Kemp, K.-K. Ni, and N. Y. Yao, Probing critical phenomena in open quantum systems using atom arrays (2024), [arXiv:2402.15376 \[quant-ph\]](https://arxiv.org/abs/2402.15376).
- [30] Y. Cheng and H. Zhai, Emergent Gauge Theory in Rydberg Atom Arrays (2024), [arXiv:2401.07708 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2401.07708).
- [31] M. Saffman, T. G. Walker, and K. Mølmer, Quantum information with Rydberg atoms, *Rev. Mod. Phys.* **82**, 2313 (2010).
- [32] M. Morgado and S. Whitlock, Quantum simulation and computing with Rydberg-interacting qubits, *AVS Quantum Sci.* **3**, 023501 (2021).
- [33] D. Bluvstein, H. Levine, G. Semeghini, T. T. Wang, S. Ebadi, M. Kalinowski, A. Keesling, N. Maskara, H. Pichler, M. Greiner, V. Vuletić, and M. D. Lukin, A quantum processor based on coherent transport of entangled atom arrays, *Nature* **604**, 451 (2022).
- [34] S. Ma, G. Liu, P. Peng, B. Zhang, S. Jandura, J. Claes, A. P. Burgers, G. Pupillo, S. Puri, and J. D. Thompson, High-fidelity gates and mid-circuit erasure conversion in an atomic qubit, *Nature* **622**, 279 (2023).
- [35] P. Scholl, A. L. Shaw, R. B.-S. Tsai, R. Finkelstein, J. Choi, and M. Endres, Erasure conversion in a high-fidelity Rydberg quantum simulator, *Nature* **622**, 273 (2023).
- [36] K. Singh, C. E. Bradley, S. Anand, V. Ramesh, R. White, and H. Bernien, Mid-circuit correction of correlated phase errors using an array of spectator qubits, *Science* **380**, 1265 (2023).
- [37] D. Bluvstein, S. J. Evered, A. A. Geim, S. H. Li, H. Zhou, T. Manovitz, S. Ebadi, M. Cain, M. Kalinowski, D. Hangleiter, J. P. Bonilla Ataides, N. Maskara, I. Cong, X. Gao, P. Sales Rodriguez, T. Karolyshyn, G. Semeghini, M. J. Gullans, M. Greiner, V. Vuletić, and M. D. Lukin, Logical quantum processor based on reconfigurable atom arrays, *Nature* **626**, 58 (2024).
- [38] Y. Zeng, P. Xu, X. He, Y. Liu, M. Liu, J. Wang, D. J. Papoula, G. V. Shlyapnikov, and M. Zhan, Entangling Two Individual Atoms of Different Isotopes via Rydberg Blockade, *Phys. Rev. Lett.* **119**, 160502 (2017).
- [39] C. Sheng, J. Hou, X. He, K. Wang, R. Guo, J. Zhuang, B. Mamat, P. Xu, M. Liu, J. Wang, and M. Zhan, Defect-Free Arbitrary-Geometry Assembly of Mixed-Species Atom Arrays, *Phys. Rev. Lett.* **128**, 083202 (2022).
- [40] K. Singh, S. Anand, A. Pocklington, J. T. Kemp, and H. Bernien, Dual-Element, Two-Dimensional Atom Array with Continuous-Mode Operation, *Phys. Rev. X* **12**, 011040 (2022).
- [41] S. Anand, C. E. Bradley, R. White, V. Ramesh, K. Singh, and H. Bernien, A dual-species Rydberg array (2024), [arXiv:2401.10325 \[quant-ph\]](https://arxiv.org/abs/2401.10325).
- [42] These coefficients are based on the existing dual-species platform, e.g. the Rydberg states of A and B can be the $^{70}S_{1/2}$ and $^{73}S_{1/2}$ states of ^{85}Rb and ^{87}Rb respectively [56]. Similar phase diagrams can be obtained for other choices of atoms, for instance, the Rb-Cs mixture.
- [43] The codes for our numerical calculations are available at <https://github.com/chengshul/RydbergSUSY>.
- [44] P. Calabrese and J. Cardy, Entanglement entropy and quantum field theory, *J. Stat. Mech.: Theory Exp.* **2004** (06), P06002.
- [45] Y. Zou and G. Vidal, Emergence of conformal symmetry in quantum spin chains: Antiperiodic boundary conditions and supersymmetry, *Phys. Rev. B* **101**, 045132 (2020).
- [46] A. J. Daley, H. Pichler, J. Schachenmayer, and P. Zoller, Measuring entanglement growth in quench dynamics of bosons in an optical lattice, *Phys. Rev. Lett.* **109**, 020505 (2012).
- [47] R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, Measuring entanglement entropy in a quantum many-body system, *Nature* **528**, 77 (2015).
- [48] N. V. Vitanov, A. A. Rangelov, B. W. Shore, and K. Bergmann, Stimulated Raman adiabatic passage in physics, chemistry, and beyond, *Rev. Mod. Phys.* **89**, 015006 (2017).
- [49] P. Fendley, K. Sengupta, and S. Sachdev, Competing density-wave orders in a one-dimensional hard-boson model, *Phys. Rev. B* **69**, 075106 (2004).
- [50] K. Slagle, D. Aasen, H. Pichler, R. S. K. Mong, P. Fendley, X. Chen, M. Endres, and J. Alicea, Microscopic characterization of Ising conformal field theory in Rydberg

- chains, *Phys. Rev. B* **104**, 235109 (2021).
- [51] C. Li *et al.*, Geometric Realization of Boundary Conditions in Reconfigurable Rydberg Arrays, to appear.
 - [52] D. Aharonov, J. Cotler, and X.-L. Qi, Quantum algorithmic measurement, *Nat. Commun.* **13**, 887 (2022).
 - [53] H.-Y. Huang, M. Broughton, J. Cotler, S. Chen, J. Li, M. Mohseni, H. Neven, R. Babbush, R. Kueng, J. Preskill, and J. R. McClean, Quantum advantage in learning from experiments, *Science* **376**, 1182 (2022).
 - [54] J. Z. Lu, L. Jiao, K. Wolinski, M. Kornjača, H.-Y. Hu, S. Cantu, F. Liu, S. F. Yelin, and S.-T. Wang, Digital-analog quantum learning on Rydberg atom arrays (2024), [arXiv:2401.02940](https://arxiv.org/abs/2401.02940) [quant-ph].
 - [55] M. Fishman, S. R. White, and E. M. Stoudenmire, The ITensor Software Library for Tensor Network Calculations, *SciPost Phys. Codebases*, 4 (2022).
 - [56] N. Šibalić, J. Pritchard, C. Adams, and K. Weatherill, ARC: An open-source library for calculating properties of alkali Rydberg atoms, *Comput. Phys. Commun.* **220**, 319 (2017).