# Wasserstein k-Centres Clustering for Distributional Data

Ryo Okano<sup>1</sup> and Masaaki Imaizumi<sup>1,2</sup>

<sup>1</sup>The University of Tokyo <sup>2</sup>RIKEN Center for Advanced Intelligence Project

July 17, 2024

#### Abstract

We develop a novel clustering method for distributional data, where each data point is regarded as a probability distribution on the real line. For distributional data, it has been challenging to develop a clustering method that utilizes the mode of variation of data because the space of probability distributions lacks a vector space structure, preventing the application of existing methods for functional data. In this study, we propose a novel clustering method for distributional data on the real line, which takes account of difference in both the mean and mode of variation structures of clusters, in the spirit of the k-centres clustering approach proposed for functional data. Specifically, we consider the space of distributions equipped with the Wasserstein metric and define a geodesic mode of variation of distributional data using geodesic principal component analysis. Then, we utilize the geodesic mode of each cluster to predict the cluster membership of each distribution. We theoretically show the validity of the proposed clustering criterion by studying the probability of correct membership. Through a simulation study and real data application, we demonstrate that the proposed distributional clustering method can improve cluster quality compared to conventional clustering algorithms. *Keywords:* Clustering: distributional data; geodesic and convex principal component analysis; Wasserstein distance.

## 1 Introduction

Cluster analysis is one of the fundamental tools in statistics used to search for homogeneous subgroups of individuals within a data set. Conventional methods, such as hierarchical clustering (Ward Jr, 1963), the k-means clustering algorithm (MacQueen, 1967), and model-based clustering (Banfield and Raftery, 1993), have been widely used for vector-valued multivariate data. These

1

<sup>&</sup>lt;sup>1</sup>R.Okano was supported by JSPS Grant-in-Aid for JSPS Fellows (22KJ1067). M.Imaizumi was supported by JSPS KAKENHI (21K11780), JST CREST (JPMJCR21D2), and JST FOREST (JPMJFR216I).

methods have also been extended to the clustering of more complex data, such as functional data (Abraham et al., 2003; James and Sugar, 2003; Serban and Wasserman, 2005; Chiou and Li, 2007; Jacques and Preda, 2013), and finite-dimensional manifold-valued data (Dhillon and Modha, 2001; Banerjee et al., 2005; Mardia et al., 2022).

This study focuses on distributional data on the real line, which is a type of complex data. Distributional data arises when each data point can be regarded as a probability distribution, and its analysis is gaining increasing attention in statistics (Petersen et al., 2022). Examples of distributional data sets include age distributions of countries and house price distributions of cities. Because the space of probability distributions does not have a vector space structure, distributional data cannot be analyzed with existing methods devised for multivariate or functional data. Moreover, distributional data cannot be analyzed with methods for finite-dimensional manifold-valued data because the space of probability distributions usually does not have a finite-dimensional tangent space. A common approach for analyzing distributional data is using the geometry provided by the Wasserstein metric in optimal transport (Villani, 2008). This approach treats each distributional data point as a point in the Wasserstein space (Panaretos and Zemel, 2020), a nonlinear metric space of probability distributions equipped with the Wasserstein metric. With this approach, statistical methodologies for analyzing distributional data have recently been developed, particularly on the real line, such as principal component analysis (Bigot et al., 2017; Cazelles et al., 2018; Campbell and Wong, 2022), regression models (Chen et al., 2023; Ghodrati and Panaretos, 2022; Okano and Imaizumi, 2024), and autoregressive models for distributional time series (Zhang et al., 2022; Zhu and Müller, 2023).

Several clustering approaches have been developed to handle distributional data. Using the Wasserstein metric, Verdinelli and Wasserman (2019) proposed a fast k-means type clustering method for distributional data based on a modified Wasserstein distance. Del Barrio et al. (2019) proposed a robust clustering method for distributional data based on trimmed k-barycenters in the Wasserstein space. Zhuang et al. (2022) defined two versions of the k-means clustering formulations in the Wasserstein space, namely centroid-based and distance-based formulations, and provided evidence for pitfalls (irregularity and non-robustness) of the centroid-based formulation. The authors also considered a semidefinite program relaxation of the distance-based formulation and showed its exact recovery property for clustering Gaussian distributions. Without using the Wasserstein metric, Calò et al. (2014) proposed a method for clustering probability density functions based on a hierarchical mixture modeling of data. Chazal et al. (2021) proposed a method for clustering of distributional data on Euclidean space with general dimensions.

One challenge in working with distributional data is clustering in a way that incorporates the differences in the mode of variation structures of clusters. While clusters of distributional data usually have different means, they also have different modes or patterns of variation. For example,

consider two clusters of distributional data: one cluster consists of population age distributions of districts in a country for men, and the other cluster consists of those for women. These clusters typically not only have different means, but also different modes of variation. However, this difference in the modes of variation may not be captured by k-means clustering, which previous studies have shown focuses only on the differences in the mean structures of clusters. While Chiou and Li (2007) developed k-centres clustering for functional data to utilize the mode of variation of clusters, it cannot handle distributional data due to the lack of a formal definition of the mode of variation in this setup. Specifically, since distributional data do not have a linear form of basis functions, it is not possible to describe their mode of variation by the ordinary principal component analysis required for k-centers clustering.

In this paper, we propose k-centres distributional clustering, which takes account of difference in the mode of variation structures of clusters of distributions on the real line. To this aim, we employ the following steps. First, to define the modes of variations of clusters of distributional data, we use geodesic principal component analysis for distributional data proposed by Bigot et al. (2017). Specifically, we consider the Wasserstein space of distributions and extract the mode of variation of distributional data using a geodesic in this space, which we call a geodesic mode of variation of distributional data. Second, we measure the discrepancy between a distribution as a data point and the geodesic mode of each cluster, and then determine the cluster membership of each distribution by minimizing the discrepancy.

Our method has several merits. First, the k-centres clustering approach can improve cluster quality by considering the difference in the mode of variation structures of clusters, in contrast to the k-means approach, which only takes account of the difference in the mean structures. This point is demonstrated in our experiment as well as in the work of Chiou and Li (2007) in the context of functional data. Second, our approach does not rely on any distributional assumptions on data, unlike several model-based clustering approaches that assume a Gaussian model, as seen in Banfield and Raftery (1993) for multivariate data clustering, and James and Sugar (2003); Jacques and Preda (2013) for functional data clustering. Third, it provides visual insights into clusters by exploring the mean and mode of variation structures of each cluster. Furthermore, our theory validates that the proposed method appropriately controls the probability of correct assignment of data to clusters. The experiments confirm these merits.

The remainder of this paper is organized as follows. In Section 2, we provide background on the Wasserstein space and geodesic principal component analysis. In Section 3, we introduce the fundamental principles of the proposed clustering method and present its specific algorithm. In Section 4, we theoretically demonstrate the validity of our clustering method. The practical performance of the proposed method is illustrated through a simulation study in Section 5 and a real data application in Section 6. We conclude with a brief discussion in Section 7. The codes implementing the proposed method and data set used in Section 6 are available at https: //github.com/RyoOkano21/kCentresDIstributionalClustering.

# 2 Background

#### 2.1 The Wasserstein Space

We provide some minimal background on the Wasserstein space needed to present our clustering method. For more details, see, for example, Ambrosio et al. (2008); Villani (2008); Bigot et al. (2017); Panaretos and Zemel (2020).

Let  $\Omega = [a, b]$  be a compact interval in  $\mathbb{R}$ , and  $\mathcal{P}(\Omega)$  be the set of Borel probability measures on  $\Omega$ . The 2-Wasserstein distance between  $\mu_1, \mu_2 \in \mathcal{P}(\Omega)$  is defined by

$$d_W(\mu_1,\mu_2) = \left\{ \int_0^1 [F_1^{-1}(u) - F_2^{-1}(u)]^2 du \right\}^{1/2},$$

where  $F_1^{-1}$  and  $F_2^{-1}$  are the quantile functions of  $\mu_1$  and  $\mu_2$ , respectively. It can be shown that the 2-Wasserstein distance  $d_W$  is a metric on  $\mathcal{P}(\Omega)$ , and the resulting metric space  $(\mathcal{P}(\Omega), d_W)$  is called as the Wasserstein space of probability distributions.

We provide several concepts of the Wasserstein space  $(\mathcal{P}(\Omega), d_W)$ , which is a generalization of those of Riemannian manifolds. In the following, we fix a reference measure  $\mu_* \in \mathcal{P}(\Omega)$ , which is assumed to be absolutely continuous with respect to the Lebesgue measure on  $\Omega$ . We define the tangent space of  $\mathcal{P}(\Omega)$  at  $\mu_*$  as the Hilbert space  $\mathcal{L}^2_{\mu_*}(\Omega)$  of real-valued,  $\mu_*$ -square-integrable functions on  $\Omega$ . This space is equipped with an inner product  $\langle \cdot, \cdot \rangle_{\mu_*}$  defined by  $\langle g_1, g_2 \rangle_{\mu_*} = \int_{\Omega} g_1 g_2 d\mu_*$  and a norm  $\|\cdot\|_{\mu_*}$  defined by  $\|g\|_{\mu_*} = \langle g, g \rangle^{1/2}_{\mu_*}$ . The exponential map  $\operatorname{Exp}_{\mu_*} : \mathcal{L}_{\mu_*}(\Omega) \to \mathcal{P}(\Omega)$  is then defined by

$$\operatorname{Exp}_{\mu_*}g = (g + \operatorname{id}) \# \mu_*,$$

where id denotes the identity function on  $\Omega$ , and for a measurable function  $h: \Omega \to \mathbb{R}$ ,  $h \# \mu_*$  is the push-forward measure such that  $h \# \mu_*(A) = \mu_*(h^{-1}(A))$  for any Borel set  $A \subset \mathbb{R}$ . Moreover, the logarithmic map  $\operatorname{Log}_{\mu_*}: \mathcal{P}(\Omega) \to \mathcal{L}^2_{\mu_*}(\Omega)$  is defined by

$$\operatorname{Log}_{\mu_*}\mu = F^{-1} \circ F_* - \operatorname{id},\tag{1}$$

where  $F_*$  and  $F^{-1}$  are the distribution function of  $\mu_*$  and quantile function of  $\mu$ , respectively. We denote the range of the logarithmic map as  $V_{\mu_*}(\Omega)$ . The following proposition summarizes the properties of the Wasserstein space.

**Proposition 1** (Bigot et al. (2017), Theorem 2.2, Remark 2.2 and Proposition 2.1). *The following statements hold:* 

(i) The restriction of the exponential map  $\operatorname{Exp}_{\mu_*}$  to  $V_{\mu_*}(\Omega)$  is an isometric homeomorphism, with inverse  $\operatorname{Log}_{\mu_*}$ . Hence, we have  $d_W(\mu_1, \mu_2) = \|\operatorname{Log}_{\mu_*}\mu_1 - \operatorname{Log}_{\mu_*}\mu_2\|_{\mu_*}$  for all  $\mu_1, \mu_2 \in \mathcal{P}(\Omega)$ .

- (ii)  $V_{\mu_*}(\Omega)$  is the set of functions  $g \in \mathcal{L}^2_{\mu_*}(\Omega)$  such that g+id is  $\mu_*$ -a.e. increasing and (g+id) $(x) \in \Omega$  for all  $x \in \Omega$ .
- (iii) The set  $V_{\mu_*}(\Omega)$  is closed and convex in  $\mathcal{L}^2_{\mu_*}(\Omega)$ .

**Remark 1** (Connection to the optimal transport map). From a viewpoint of optimal transport theory, transforming with the logarithmic map is interpreted as the transformation of a distribution to the corresponding optimal transport map. Specifically, for the probability measures  $\mu_*$  and  $\mu$ , any map  $T: \Omega \to \Omega$  that minimizes Monge's problem  $\inf_{T\#\mu_*=\mu} \int_{\Omega} [T(x) - x]^2 d\mu_*(x)$  is called as an optimal transport map from  $\mu_*$  to  $\mu$ . In our setting, such optimal transport map uniquely exists and can be expressed as  $T = F^{-1} \circ F_*$ , where  $F_*$  and  $F^{-1}$  are the distribution function of  $\mu_*$  and quantile function of  $\mu$ , respectively. Therefore, the function obtained by the logarithmic map (1) is identical to the optimal transport map up to subtraction of the identity function.

We further define the notion of a geodesic in the Wasserstein space. In general, a metric space (S, d) is said to be geodesic if for every  $x, y \in S$ , there exists a shortest path between x and y. Given a geodesic metric space (S, d), a set  $T \subset S$  is called geodesic if for every  $x, y \in T$ , the shortest path between them is totally contained in T. The following proposition summarizes the properties of geodesics in the Wasserstein space.

Proposition 2 (Bigot et al. (2017), Remark 2.3 and Corollary 2.1). The following statements hold:

- (i) The Wasserstein space  $(\mathcal{P}(\Omega), d_W)$  is a geodesic metric space.
- (ii) A set  $G \subset \mathcal{P}(\Omega)$  is geodesic if and only if  $\operatorname{Log}_{\mu_*}(G)$  is convex in  $\mathcal{L}^2_{\mu_*}(\Omega)$ .

For a geodesic set  $G \subset \mathcal{P}(\Omega)$ , its dimension  $\dim(G)$  is defined as the dimension of the convex set  $\operatorname{Log}_{\mu_*}(G)$  in  $\mathcal{L}^2_{\mu_*}(\Omega)$ , that is, the dimension of the smallest affine subspace of  $\mathcal{L}^2_{\mu_*}(\Omega)$  containing  $\operatorname{Log}_{\mu_*}(G)$ . Note that  $\dim(G)$  does not depend on the choice of reference measure  $\mu_*$  (Remark 2.4, Bigot et al., 2017).

#### 2.2 Geodesic Principal Component Analysis in the Wasserstein Space

We review a formulation of geodesic principal component analysis (PCA) in the Wasserstein space, which was originally introduced by Bigot et al. (2017). Let  $\boldsymbol{\mu}$  be a random element taking its values in the Wasserstein space ( $\mathcal{P}(\Omega), d_W$ ), which is assumed to be square-integrable in the sense that  $\mathbb{E}[d_W^2(\boldsymbol{\mu}, \boldsymbol{\mu})] < \infty$ , for some (thus for all)  $\boldsymbol{\mu} \in \mathcal{P}(\Omega)$ . In applications,  $\boldsymbol{\mu}$  usually follows an empirical measure of observed samples  $\mu_1, ..., \mu_n \in \mathcal{P}(\Omega)$  such that  $\mathbb{P}(\boldsymbol{\mu} \in A) = n^{-1} \sum_{i=1}^n I_A(\mu_i)$  for any Borel set  $A \subset \mathcal{P}(\Omega)$ . Here,  $I_A$  is the indicator function of a set A. The Fréchet mean or barycentre of  $\boldsymbol{\mu}$ in the Wasserstein space is defined by

$$\mu_{\oplus} \in \underset{\mu \in \mathcal{P}(\Omega)}{\operatorname{arg\,min}} \mathbb{E}[d_W^2(\boldsymbol{\mu}, \mu)], \tag{2}$$

which uniquely exists (Proposition 4.1, Bigot et al., 2017).

To formulate geodesic PCA in the Wasserstien space, we introduce the following. For a point  $\mu \in \mathcal{P}(\Omega)$  and nonempty set  $G \subset \mathcal{P}(\Omega)$ , we define the distance between them as  $d_W(\mu, G) = \inf_{\lambda \in G} d_W(\mu, \lambda)$ . Note that if G is a closed geodesic subset, there exists a unique point  $\lambda \in G$  such that  $d_W(\nu, \lambda) = d_W(\nu, G)$  by the combination of Proposition 1 and the Hilbert projection theorem. We define  $K_W(G) = \mathbb{E}[d_W^2(\mu, G)]$  for any nonempty set  $G \subset \mathcal{P}(\Omega)$ . For any integer  $j \geq 1$ , we denote by  $\mathrm{CG}_{\mu \oplus, j}$  the family of nonempty, closed and geodesic subsets  $G \subset \mathcal{P}(\Omega)$ , such that  $\dim(G) \leq j$  and  $\mu_{\oplus} \in G$ .

Now, we state a notion of geodesic PCA in the Wasserstein space. Precisely, we focus on the nested version of geodesic PCA by Bigot et al. (2017), which has advantages in computation and interpretation.

**Definition 1** (Nested geodesic PCA in the Wassersetin space). Let  $G_0 = {\mu_{\oplus}}$ . For a given integer  $M \ge 1$ , we say that a set  $G_M \subset \mathcal{P}(\Omega)$  is an  $(M, \mu_{\oplus})$ -nested principal geodesic of  $\mu$  if there exist sets  $G_0, ..., G_{M-1}$  such that  $G_0 \subset G_1 \subset \cdots \subset G_{M-1} \subset G_M$ , and for each j = 1, ..., M,

$$G_j \in \underset{G \in CG_{\mu_{\oplus},j}, G \supset G_{j-1}}{\operatorname{arg\,min}} K_W(G).$$
(3)

In our setting, the nested geodesic PCA problem has nonempty solutions for any  $M \ge 1$ (Theorem 4.1, Bigot et al., 2017). Furthermore, a solution of nested geodesic PCA problem in the Wasserstein space is obtained by solving a convex PCA problem in the tangent space. We illustrate this relationship in Figure 1. We give a review of a formulation of convex PCA problem and its link to geodesic PCA problem in Section S.1 in the Supplementary Material.

Based on the result of geodesic PCA, as with the ordinal PCA, we can define a notion of lowdimensional representation of  $\mu$ . Given an  $(M, \mu_{\oplus})$ -nested principal geodesic  $G_M \subset \mathcal{P}(\Omega)$  of  $\mu$ , we define an *M*-dimensional representation of  $\mu$  as

$$\Pi_{G_M} \boldsymbol{\mu} \in \operatorname*{arg\,min}_{\boldsymbol{\mu} \in G_M} d_W^2(\boldsymbol{\mu}, \boldsymbol{\mu}), \tag{4}$$

which uniquely exists.

## 3 Method

#### **3.1** Basic Principle

We explain the basic principle of the k-centres distributional clustering, which is our proposal. We consider the Wasserstein space  $(\mathcal{P}(\Omega), d_W)$ , where  $\Omega = [a, b]$  is a compact interval in  $\mathbb{R}$ . Let  $\boldsymbol{\nu}^{(c)}$  be a  $\mathcal{P}(\Omega)$ -valued random element for each cluster c = 1, ..., K, and  $\boldsymbol{\nu}$  be a  $\mathcal{P}(\Omega)$ -valued random element defined as a mixture of random variables  $\boldsymbol{\nu}^{(c)}$  associated with a random cluster variable C on  $\{1, ..., K\}$ . Rigorously, we have  $\mathbb{P}(\boldsymbol{\nu} \in A | C = c) = \mathbb{P}(\boldsymbol{\nu}^{(c)} \in A)$  for any Borel set  $A \subset \mathcal{P}(\Omega)$ . We will



Figure 1: Illustration of the relationship between geodesic PCA in the Wasserstein space and convex PCA in the tangent space (see Section S.1 for details). The star points in the Wasserstein space  $\mathcal{P}(\Omega)$  are realizations of  $\boldsymbol{\mu}$ , while the the black points in the tangent space  $\mathcal{L}^2_{\mu_*}(\Omega)$  are those of  $\log_{\mu_*}\boldsymbol{\mu}$ . The reference measure  $\mu_*$  is chosen as the Fréchet mean  $\mu_{\oplus}$  of  $\boldsymbol{\mu}$ . The black segment in  $\mathcal{L}^2_{\mu_*}(\Omega)$  is an  $(M, \overline{g})$ -nested principal convex component  $C_M$  of  $\log_{\mu_*}\boldsymbol{\mu}$ , and the curve line  $G_M$  in  $\mathcal{P}(\Omega)$  is defined as  $G_M = \operatorname{Exp}_{\mu_*}(C_M)$ . Then  $G_M$  is an  $(M, \nu_{\oplus})$ -nested principal geodesic of  $\boldsymbol{\mu}$ .

observe independent realizations of  $\nu$  in which each individual cluster membership is determined by the random variable C, and consider their clustering.

We describe the basic principle of our clustering method. Let  $\nu$  be an observed realization of the random element  $\nu$ . For c = 1, ..., K, we denote with  $\nu_{\oplus}^{(c)}$  the Fréchet mean of the random element  $\nu^{(c)}$  defined as in (2). Additionally, we denote with  $G_M^{(c)}$  an  $(M, \nu_{\oplus}^{(c)})$ -nested principle geodesic of  $\nu^{(c)}$  defined as in (3), and we call this principal geodesic  $G_M^{(c)}$  as the geodesic mode of variation of cluster c. Then, for each c = 1, ..., K, we consider a model  $\tilde{\nu}^{(c)}$  of  $\nu$  such that

$$\tilde{\nu}^{(c)} = \prod_{G^{(c)}} \nu, \tag{5}$$

where  $\Pi_{G_M^{(c)}}\nu$  is the *M*-dimensional representation of  $\nu$  defined as in (4). If the cluster membership of  $\nu$  actually belongs to cluster *c*, that is, on an event that the random variable *C* takes the value *c*, then the model  $\tilde{\nu}^{(c)}$  is a good approxiation of  $\nu$  in the sense of the Wasserstein distance. Otherwise, discrepancies exist between  $\tilde{\nu}^{(c)}$  and  $\nu$ . With this principle, we predict cluster membership of  $\nu$  based on the Wasserstein distance between  $\nu$  and the model  $\tilde{\nu}^{(c)}$ . Specifically, we propose to determine the cluster membership of  $\nu$  by the criterion

$$c^{*}(\nu) = \underset{c \in \{1, ..., K\}}{\arg\min} d_{W}(\nu, \tilde{\nu}^{(c)}).$$
(6)

We illustrate this basic principle of our clustering method in Figure 2.



Figure 2: Illustration of the basic principle of the proposed clustering method. For simplicity, we consider the case of two clusters (K = 2). The two curves are nested principal geodesics  $G_M^{(1)}, G_M^{(2)}$  as the geodesic mode of the variations, and the star and triangular points around them are typical realizations of  $\boldsymbol{\nu}^{(1)}$  and  $\boldsymbol{\nu}^{(2)}$ , respectively. The black point is an observed data  $\nu$ , and the cross mark on each curve indicates its model  $\tilde{\nu}^{(c)}$ . In this case, we have  $d_W(\nu, \tilde{\nu}^{(1)}) < d_W(\nu, \tilde{\nu}^{(2)})$ , and thus  $\nu$  is classified into the cluster 1.

**Remark 2** (The dimensions of the geodesic modes). In the proposed method, the same value M is used for the dimensions of the geodesic modes of all clusters. This is in construct to the k-centres functional clustering (Section 2.2.2, Chiou and Li, 2007), where the number of principal components is allowed to vary from cluster to cluster. We mention that it would be desirable to use the same values of dimension for all clusters for a fair comparison of cluster structures.

**Remark 3** (Clustering on the tangent space). The proposed clustering for distributional data is equivalent to the k-centres clustering for elements in the tangent space with convex PCA. As stated in Remark 1, the transformation with the logarithmic map is essentially identical to the transformation to the corresponding optimal transport map. Hence, the proposed clustering for distributional data is almost equivalent to the k-centres functional clustering for corresponding optimal transport maps.

## 3.2 Clustering Procedure

Setting Suppose  $\nu_1, ..., \nu_n \in \mathcal{P}(\Omega)$  are *n* independent realizations of the mixture random element  $\boldsymbol{\nu}$ , and we aim to classify these *n* distributions into *K* groups. Throughout this paper, we assume the number of clusters *K* is known or predetermined. If *K* is not predetermined, we determine it by performing clustering for different numbers of clusters, and evaluating their performance by some criterion. A representative option is using the silhouette method (Rousseeuw, 1987), which

can be applied to our method since it can be calculated using the Wasserstein distance between distributions.

In preparation, we define the transformations of the distributional data. Let  $\hat{\mu}_* \in \mathcal{P}(\Omega)$  be an absolutely continuous reference measure and transform the distributional data as  $g_i = \text{Log}_{\hat{\mu}_*} \nu_i, i = 1, ..., n$ . The reference measure  $\hat{\mu}_*$  is typically chosen as the empirical Fréchet mean of  $\nu_1, ..., \nu_n$  in the Wasserstein space,

$$\hat{\nu}_{\oplus} \in \operatorname*{arg\,min}_{\mu \in \mathcal{P}(\Omega)} \frac{1}{n} \sum_{i=1}^{n} d_W^2(\nu_i, \mu),\tag{7}$$

which uniquely exists (Proposition 4.1, Bigot et al., 2017), or the uniform distribution on  $\Omega$ . We denote by  $\overline{g} = n^{-1} \sum_{i=1}^{n} g_i$  the empirical mean of the transformed data.

**Remark 4** (Observation of distributional data). In practical analysis, the distributions  $\nu_i$  are seldom directly observed; instead, we observe independent samples from them. In this case, we need to estimate the distributions from the samples before implementing the clustering method. See Section S.3 in the Supplementary Material for a description of the estimation procedure.

We describe the procedure of our clustering method. It consists of three steps: the selection of the dimension M of geodesic modes, initial clustering and reclassification.

Selection of the Dimension of Geodesic Modes. First of all, we select the dimension M of the geodesic modes for our method. To this end, we define a notion of explained variation for geodesic PCA in the Wasserstein space. Specifically, let  $\hat{\nu}_{\oplus}$  be the empirical Fréchet mean of the n distributions  $\nu_1, ..., \nu_n$  defined by (7), and for any  $M' \in \mathbb{N}$ , let  $\hat{G}_{M'} \subset \mathcal{P}(\Omega)$  be an  $(M', \hat{\nu}_{\oplus})$ -nested principal geodesic of the n distributions. The cumulative proportion of variation explained by  $\hat{G}_{M'}$  is defined as

$$EV(\hat{G}_{M'}) = \frac{n^{-1} \sum_{i=1}^{n} d_W^2(\Pi_{\hat{G}_{M'}}\nu_i, \hat{\nu}_{\oplus})}{TV}.$$

where the total variation is defined as  $TV = n^{-1} \sum_{i=1}^{n} d_W^2(\nu_i, \hat{\nu}_{\oplus})$ . Then, we propose to select M as

$$M = \min\{M' \in \mathbb{N} : EV(\hat{G}_{M'}) \ge \tau\}.$$
(8)

Here,  $\tau \in (0, 1)$  is a pre-selected threshold value, and setting  $\tau = 0.9$  or  $\tau = 0.8$  works reasonably well in our numerical experience.

Though we use geodesic PCA in the Wasserstein space for selecting the dimension M of geodesic modes, we can select the same M by using convex PCA in the tangent space (see Section S.4 for details). This is a reasonable alternative in practice, because convex PCA is easier to implement than geodesic PCA.

**Initial Clustering.** In this step, we classify the *n* distributions  $\nu_1, ..., \nu_n$  into *K* groups as initialization. Specifically, we apply the *k*-means clustering for *M*-dimensional vectors obtained by applying convex PCA to the transformed data. This approach is analogous to the initial clustering step of the *k*-centres functional clustering method (Section 2.2.1, Chiou and Li, 2007).

We describe its details. Using the selected M in the previous step, we compute an  $(M, \overline{g})$ nested principal convex component of  $g_1, ..., g_n$ . Based on it, for each i = 1, ..., n we obtain the
convex principal component scores of  $g_i$ ,  $\hat{\xi}_i = (\hat{\xi}_{i1}, ..., \hat{\xi}_{iM}) \in \mathbb{R}^M$  (see Section S.1.2 for details).
The initial cluster membership is determined by the k-means method to the M-dimensional vectors  $\hat{\xi}_i, i = 1, ..., n$ . Let  $h_i^{(0)} \in \{1, ..., K\}$  be the label of the cluster membership for the *i*-th distribution
at this initialization step.

**Reclassification.** With the initial clustering results, we reclassify each distribution into the best predicted cluster with the principle described in Section 3.1. Let  $h_i^{(t)} \in \{1, ..., K\}$  be the label of cluster membership for the *i*-th distribution at the *t*-th iteration. Given the set of clustering results  $\mathcal{H}^{(t)} = \{h_i^{(t)} : i = 1, ..., n\}$ , we obtain for each individual *i* and cluster *c* the estimates of the Fréchet mean  $\hat{\nu}_{\oplus}^{(c)}$  and an  $(M, \hat{\nu}_{\oplus}^{(c)})$ -principal geodesic  $\hat{G}_M^{(c)}$  based on the observed distributions  $\nu_k$  with  $h_k^{(t)} = c$  for all  $k \neq i$ , leaving out the *i*-th observed distribution. Here, *M* is the value of dimension that was selected according to the criterion (8) in the first step. Given these estimates, we obtain the *i*-th predicted model for each cluster *c*,

$$\tilde{\nu}_{(i)}^{(c)} = \prod_{\hat{G}_M^{(c)}} \nu_i$$

as in (5). The *i*-th distribution  $\nu_i$  is then classified into cluster  $h_i^{(t+1)}$  such that

$$h_i^{(t+1)} = \underset{c \in \{1, \dots, K\}}{\arg\min} d_W(\nu_i, \tilde{\nu}_{(i)}^{(c)})$$

as in criterion (6). This step is performed for all i, which leads to the updated set of results  $\mathcal{H}^{(t+1)} = \{h_i^{(t+1)} : i = 1, ..., n\}$ . This updating procedure is iteratively implemented until no more data can be reclassified.

## 4 Theory

In this section, we investigate the theoretical properties of the proposed distributional clustering method. Specifically, we demonstrate that the membership criterion (6) gives a correct clustering for distributional data. For full proofs, see Section S.2 in the Supplementary Material.

#### 4.1 Setup and Assumption

Let assume that there are two clusters labeled c and d in the Wasserstein space  $(\mathcal{P}(\Omega), d_W)$ , and  $\mathcal{P}(\Omega)$ -valued random elements  $\boldsymbol{\nu}^{(c)}$  and  $\boldsymbol{\nu}^{(d)}$  are defined according to the clusters. We denote the

the Fréchet means of  $\boldsymbol{\nu}^{(c)}$  and  $\boldsymbol{\nu}^{(d)}$  by  $\nu_{\oplus}^{(c)}$  and  $\nu_{\oplus}^{(d)}$ , respectively. Also, we denote  $(1, \nu_{\oplus}^{(c)})$  and  $(1, \nu_{\oplus}^{(d)})$ -nested principal geodesics of  $\boldsymbol{\nu}^{(c)}$  and  $\boldsymbol{\nu}^{(d)}$  by  $G^{(c)}$  and  $G^{(d)}$ , respectively. Then for the random element  $\boldsymbol{\nu}^{(c)}$  of cluster c, stochastic models  $\tilde{\boldsymbol{\nu}}_{c}^{(c)}$  and  $\tilde{\boldsymbol{\nu}}_{c}^{(d)}$  are defined as

$$\tilde{\boldsymbol{\nu}}_c^{(c)} = \Pi_{G^{(c)}} \boldsymbol{\nu}^{(c)} \quad \text{and} \quad \tilde{\boldsymbol{\nu}}_c^{(d)} = \Pi_{G^{(d)}} \boldsymbol{\nu}^{(c)},$$

respectively.

Our theoretical interest is whether the proposed criterion (6) correctly determines the cluster membership of  $\nu^{(c)}$ . Rigorously, if the condition

$$d_W(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_c^{(c)}) < d_W(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_c^{(d)})$$
(9)

holds, then the cluster membership of  $\boldsymbol{\nu}^{(c)}$  is correctly determined by the proposed criterion (6). Otherwise, the cluster membership of  $\boldsymbol{\nu}^{(c)}$  is wrongly determined by the proposed criterion. A goal of our theory is to show that the condition (9) holds with a high probability in several situations.

We state the assumptions. Let  $\mu_* \in \mathcal{P}(\Omega)$  be an absolutely continuous reference measure, and define  $\mathcal{L}^2_{\mu_*}(\Omega)$ -valued random elements  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$  as  $\mathbf{g}^{(c)} = \mathrm{Log}_{\mu_*} \boldsymbol{\nu}^{(c)}$  and  $\mathbf{g}^{(d)} = \mathrm{Log}_{\mu_*} \boldsymbol{\nu}^{(d)}$ , respectively. Then we make the following assumption.

Assumption 1. Assume that the  $\mathcal{L}^2_{\mu_*}(\Omega)$ -valued random elements  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$  have expansions

$$\mathbf{g}^{(c)} = m^{(c)} + \sum_{j=1}^{J} \xi_j^{(c)} \rho_j^{(c)} \quad and \quad \mathbf{g}^{(d)} = m^{(d)} + \sum_{j=1}^{J} \xi_j^{(d)} \rho_j^{(d)},$$

respectively. Here, J is a positive integer,  $m^{(c)}$  and  $m^{(d)}$  are some vectors in the range of the logarithmic map  $V_{\mu_*}(\Omega)$ , and  $\{\rho_j^{(c)}\}_{j=1}^J$  and  $\{\rho_j^{(d)}\}_{j=1}^J$  are some orthonormal vectors in  $\mathcal{L}^2_{\mu_*}(\Omega)$ .  $\{\xi_j^{(c)}\}_{j=1}^J$  and  $\{\xi_j^{(d)}\}_{j=1}^J$  are respectively uncorrelated random variables with zero means such that  $\operatorname{Var}(\xi_1^{(c)}) \geq \operatorname{Var}(\xi_2^{(c)}) \geq \cdots \geq \operatorname{Var}(\xi_J^{(c)}) \geq 0$  and  $\operatorname{Var}(\xi_1^{(d)}) \geq \operatorname{Var}(\xi_2^{(d)}) \geq \cdots \geq \operatorname{Var}(\xi_J^{(d)}) \geq 0$ . It is also assumed that for all  $\ell = 1, ..., J$ , the truncated expansions  $m^{(c)} + \sum_{j=1}^{\ell} \xi_j^{(c)} \rho_j^{(c)}$  and  $m^{(d)} + \sum_{j=1}^{\ell} \xi_j^{(d)} \rho_j^{(d)}$  are in  $V_{\mu_*}(\Omega)$  with probability equal to 1.

Under Assumption 1,  $m^{(c)}$  and  $m^{(d)}$  are the mean functions of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$ , and  $\rho_j^{(c)}$  and  $\rho_j^{(d)}$ are the *j*-th convex principal directions of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$ , respectively. We also assume that the orthogonal projection of  $\mathbf{g}^{(c)}$  onto the space  $m^{(d)} + \text{span}\{\rho_1^{(d)}\}$  is in  $V_{\mu_*}(\Omega)$ , which is formally stated as follows.

Assumption 2. Assume that the condition

$$m^{(d)} + \langle \mathbf{g}^{(c)} - m^{(d)}, \rho_1^{(d)} \rangle_{\mu_*} \rho_1^{(d)} \in V_{\mu_*}(\Omega)$$

holds with probability equal to 1.

#### 4.2 Correct Membership Probability

In this section, we consider two cases that the two clusters have either the same mean or same covariance, then we study the probability that the condition (9) holds. Specifically, we express the probabilities by using the random element  $\mathbf{g}^{(c)}$  takes values. Furthermore, in the subsequent remarks, we evaluate those probabilities more explicitly by imposing a Gaussian distributional assumption on the coefficients of  $\mathbf{g}^{(c)}$ .

**Common Mean Case.** We consider that the random elements  $\boldsymbol{\nu}^{(c)}$  and  $\boldsymbol{\nu}^{(d)}$  from the clusters have the same mean structures, i.e.,  $\nu_{\oplus}^{(c)} = \nu_{\oplus}^{(d)}$  holds. Then, we obtain the following result: **Proposition 3.** Assume  $\nu_{\oplus}^{(c)} = \nu_{\oplus}^{(d)}$  and define a vector  $m \in \mathcal{L}^2_{\mu_*}(\Omega)$  by  $m = \text{Log}_{\mu_*}\nu_{\oplus}^{(c)} = \text{Log}_{\mu_*}\nu_{\oplus}^{(d)}$ . Then under Assumptions 1 and 2, it holds that

$$\mathbb{P}(d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(c)}) < d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(d)})) 
= \mathbb{P}(\langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} > 0, \langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} > 0) 
+ \mathbb{P}(\langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} < 0, \langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} < 0).$$
(10)

Proposition 3 implies that when the mean structures of the two clusters are same, the probability that the condition (9) holds is equal to the probability that the zero-mean random variable  $\mathbf{g}^{(c)} - m$  takes values in regions  $S_1$  or  $S_2$ , where

$$S_1 = \{ g \in \mathcal{L}^2_{\mu_*}(\Omega) : \langle g, \rho_1^{(c)} + \rho_1^{(d)} \rangle_{\mu_*} > 0, \langle g, \rho_1^{(c)} - \rho_1^{(d)} \rangle_{\mu_*} > 0 \}$$

and

$$S_2 = \{ g \in \mathcal{L}^2_{\mu_*}(\Omega) : \langle g, \rho_1^{(c)} + \rho_1^{(d)} \rangle_{\mu_*} < 0, \langle g, \rho_1^{(c)} - \rho_1^{(d)} \rangle_{\mu_*} < 0 \},$$

respectively. As illustrated in Figure 3, the vectors  $\rho_1^{(c)} + \rho_1^{(d)}$  and  $\rho_1^{(c)} - \rho_1^{(d)}$  are orthogonal, and if  $\rho_1^{(c)} \neq \rho_1^{(d)}$ , the first convex principal direction  $\rho_1^{(c)}$  of  $\mathbf{g}^{(c)}$  satisfies  $\rho_1^{(c)} \in S_1$  and  $-\rho_1^{(c)} \in S_2$ . From these observations, we can expect that the condition (9) holds with a high probability if  $\rho_1^{(c)} \neq \rho_1^{(d)}$ .

**Remark 5.** If we impose a Gaussian distributional assumption on the random coefficients  $\xi_1^{(c)}, ..., \xi_J^{(c)}$  of  $\mathbf{g}^{(c)}$ , we can express the probability (10) more explicitly. Note that since Gaussian distributions have unbounded supports, imposing Gaussianity on the coefficients of  $\mathbf{g}^{(c)}$  may violate the assumption that  $m^{(c)} + \sum_{j=1}^{\ell} \xi_j^{(c)} \rho_j^{(c)} \in V_{\mu_*}(\Omega)$  holds with probability 1 for all  $\ell = 1, ..., J$ . In this remark, we impose a Gaussian assumption instead of the above assumption for the purpose of understanding the probability (10). Let assume that the J-dimensional random vector  $(\xi_1^{(c)}, ..., \xi_J^{(c)})$  follows the zero-mean Gaussian distribution with covariance matrix

$$\Sigma = \begin{pmatrix} \operatorname{Var}(\xi_1^{(c)}) & 0 \\ & \ddots & \\ 0 & \operatorname{Var}(\xi_J^{(c)}) \end{pmatrix}.$$
(11)



Figure 3: Illustration of the probabilities in Proposition 3. The contour lines indicate the distribution of the random variable  $\mathbf{g}^{(c)}$ , and the solid arrows indicate the first convex principal directions  $\rho_1^{(c)}$  and  $\rho_1^{(d)}$  of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$ . The dashed arrows indicate the vectors  $\rho_1^{(c)} + \rho_1^{(d)}$  and  $\rho_1^{(c)} - \rho_1^{(d)}$ , and the filled squares indicate the regions  $S_1 = \{g \in \mathcal{L}^2_{\mu_*}(\Omega) : \langle g, \rho_1^{(c)} + \rho_1^{(d)} \rangle_{\mu_*} > 0, \langle g, \rho_1^{(c)} - \rho_1^{(d)} \rangle_{\mu_*} > 0\}$  and  $S_2 = \{g \in \mathcal{L}^2_{\mu_*}(\Omega) : \langle g, \rho_1^{(c)} + \rho_1^{(d)} \rangle_{\mu_*} < 0, \langle g, \rho_1^{(c)} - \rho_1^{(d)} \rangle_{\mu_*} < 0\}$ . As we have  $\rho_1^{(c)} \in S_1$  and  $-\rho_1^{(c)} \in S_2$ , it is expected that  $\mathbf{g}^{(c)} - m$  takes values in the regions  $S_1$  or  $S_2$  with a high probability.

Then with a two-dimensional zero-mean Gaussian random vector  $(W_1, W_2)$  with covariance matrix

$$\begin{pmatrix} \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}}^{2} & \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} \\ \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} & \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} \end{pmatrix}$$

the probability (10) is expressed as  $\mathbb{P}(W_1 > 0, W_2 > 0) + \mathbb{P}(W_1 < 0, W_2 < 0)$ , which is equal to  $0.5 + \pi^{-1} \operatorname{arcsin} \operatorname{Corr}(W_1, W_2)$  (see Section 15.10 in Stuart and Ord (2010) for this equality). Furthermore, assuming  $\operatorname{span}\{\rho_1^{(c)}, \dots, \rho_J^{(c)}\} = \operatorname{span}\{\rho_1^{(d)}, \dots, \rho_J^{(d)}\}$ , there exists an index  $\ell \in \{1, \dots, J\}$  such that  $\rho_\ell^{(c)} = \rho_1^{(d)}$  and  $\rho_j^{(c)} \neq \rho_1^{(d)}$  for  $j \neq \ell$ . If  $\ell \geq 2$  we have

$$\begin{split} \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}}^{2} &= \operatorname{Var}(\xi_{1}^{(c)}) + \operatorname{Var}(\xi_{\ell}^{(c)}), \\ \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} &= \operatorname{Var}(\xi_{1}^{(c)}) - \operatorname{Var}(\xi_{\ell}^{(c)}), \\ \sum_{j=1}^{J} \operatorname{Var}(\xi_{j}^{(c)}) \langle \rho_{j}^{(c)}, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}}^{2} &= \operatorname{Var}(\xi_{1}^{(c)}) + \operatorname{Var}(\xi_{\ell}^{(c)}). \end{split}$$

This implies the correlation between  $W_1$  and  $W_2$  is given by  $\{\operatorname{Var}(\xi_1^{(c)}) - \operatorname{Var}(\xi_\ell^{(c)})\}/\{\operatorname{Var}(\xi_1^{(c)}) + \operatorname{Var}(\xi_\ell^{(c)})\}$ , and the probability (10) is eventually expressed as

$$0.5 + \frac{1}{\pi} \arcsin\left(\frac{\operatorname{Var}(\xi_1^{(c)}) - \operatorname{Var}(\xi_\ell^{(c)})}{\operatorname{Var}(\xi_1^{(c)}) + \operatorname{Var}(\xi_\ell^{(c)})}\right).$$
(12)

Since we assumed  $\operatorname{Var}(\xi_1^{(c)}) \geq \operatorname{Var}(\xi_\ell^{(c)})$ , the value (12) is guaranteed to be 0.5 or higher. Also, as illustrated in Figure 4, as  $\operatorname{Var}(\xi_1^{(c)})$  gets larger, the value (12) approaches to 1.

**Common Covariance Case.** We consider that the random elements  $\boldsymbol{\nu}^{(c)}$  and  $\boldsymbol{\nu}^{(d)}$  from the clusters have the same covariance structures, i.e., the *j*-th convex principal directions  $\rho_j^{(c)}$  and  $\rho_j^{(d)}$  are identical for j = 1, ..., J. Then, we obtain the following result:

**Proposition 4.** Assume  $\rho_j^{(c)} = \rho_j^{(d)}$  holds for j = 1, ..., J and denote the common vector by  $\rho_j$ . Then under Assumptions 1 and 2, we have

$$\mathbb{P}(d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(c)}) < d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(d)})) = \mathbb{P}(\langle \mathbf{g}^{(c)} - m^{(c)}, \psi \rangle_{\mu_{*}} > \langle m^{(c)} - m^{(d)}, \rho_{1} \rangle_{\mu_{*}}^{2} - \|m^{(c)} - m^{(d)}\|_{\mu_{*}}^{2}),$$
(13)

where  $\psi = 2m^{(c)} - 2m^{(d)} - 2\langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*} \rho_1$ .

Proposition 4 implies that when the covariance structures of the two clusters are same, the probability that the condition (9) holds is equal to the probability that the zero-mean random



Figure 4: Probability (12) as a function of  $\operatorname{Var}(\xi_1^{(c)})$ .  $\operatorname{Var}(\xi_\ell^{(c)})$  is set as 1. We see that as  $\operatorname{Var}(\xi_1^{(c)})$  gets larger, the probability approaches to 1.

variable  $\langle \mathbf{g}^{(c)} - m^{(c)}, \psi \rangle_{\mu_*}$  is more than  $\langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*}^2 - \|m^{(c)} - m^{(d)}\|_{\mu_*}^2$ . Since the value  $\langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*}^2 - \|m^{(c)} - m^{(d)}\|_{\mu_*}^2$  is non-positive by the Cauchy–Schwarz inequality, we can expect that the condition (9) holds with a high probability.

**Remark 6.** As in Remark 5, we can evaluate the probability (13) more explicitly by imposing the Gaussian distributional assumption on the coefficients of  $\mathbf{g}^{(c)}$ . Let assume the random vector  $(\xi_1^{(c)},...,\xi_J^{(c)})$  follows the zero-mean Gaussian distribution with covariance matrix (11). Then with a zero-mean Gaussian random variable V with variance  $\sum_{j=1}^{J} \operatorname{Var}(\xi_j^{(c)}) \langle \rho_j, \psi \rangle_{\mu_*}^2$ , the probability (13) is expressed as  $\mathbb{P}(V > \langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*}^2 - \|m^{(c)} - m^{(d)}\|_{\mu_*}^2)$ . In the following, we assume  $J \ge 2$ . If  $m^{(c)} - m^{(d)} \in \operatorname{span}\{\rho_2,...,\rho_J\}$ , then the variance of V is bounded above by  $4\operatorname{Var}(\xi_1^{(c)})\|m^{(c)} - m^{(d)}\|_{\mu_*}^2$ , and  $\langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*}^2 - \|m^{(c)} - m^{(d)}\|_{\mu_*}^2$  reduces to  $-\|m^{(c)} - m^{(d)}\|_{\mu_*}^2$ , which implies the probability (13) is bounded below by

$$\Phi\left(\frac{\|m^{(c)} - m^{(d)}\|_{\mu_*}}{2\sqrt{\operatorname{Var}(\xi_1^{(c)})}}\right).$$
(14)

Here,  $\Phi$  is the distribution function of the standard normal distribution, and  $\operatorname{Var}(\xi_1^{(c)})$  is assumed to be positive. Since  $\|m^{(c)} - m^{(d)}\|_{\mu_*} \geq 0$ , the value (14) is guaranteed to be 0.5 or higher. Also, as illustrated in Figure 5, as  $\|m^{(c)} - m^{(d)}\|_{\mu_*}$  gets larger, the value (14) approaches to 1. On the other hand, if  $m^{(c)} - m^{(d)} \in \operatorname{span}\{\rho_1\}$ , then we have  $\tilde{\boldsymbol{\nu}}_c^{(c)} = \tilde{\boldsymbol{\nu}}_c^{(d)}$  with probability 1. Under this situation, the proposed clustering criterion does not correctly determine the cluster membership of  $\boldsymbol{\nu}^{(c)}$ . This situation is analogous to a situation called as the non-identifiable situation in Chiou and Li (2007), where two models via subspace projection are not distinguished well (Theorem 1, Chiou and Li, 2007).



Figure 5: Probability (14) as a function of  $||m^{(c)} - m^{(d)}||_{\mu_*}$ .  $\operatorname{Var}(\xi_{\ell}^{(c)})$  is set as 1. We see that as  $||m^{(c)} - m^{(d)}||_{\mu_*}$  gets larger, the probability approaches to 1.

# 5 Simulation Study

In this section, we investigate the practical performance of the proposed method through a simulation study.

## 5.1 Simulation Setup

We define the following model  $\nu_i^{(c)} \in \mathcal{P}(\Omega)$  for each individual  $i \in \{1, ..., n\}$  and cluster  $c \in \{1, 2\}$ :

$$\nu_i^{(c)} = \text{Exp}_{\mu_*} g_i^{(c)}, \tag{15}$$

where  $g_i^{(c)}$  is a function defined as

$$g_i^{(c)}(x) = m^{(c)}(x) + \xi_{i1}^{(c)}\phi_1^{(c)}(x) + \xi_{i2}^{(c)}\phi_2^{(c)}(x), \quad x \in \Omega.$$

Here, the reference measure  $\mu_*$  is set to be the uniform measure on  $\Omega$ , and the mean function  $m^{(c)} \in V_{\mu_*}(\Omega)$  and set of principal directions  $\mathcal{S}^{(c)} = \{\phi_1^{(c)}, \phi_2^{(c)}\} \subset \mathcal{L}^2_{\mu_*}(\Omega)$  are given by the simulation designs. The random coefficients  $\xi_{i1}^{(c)}$  and  $\xi_{i2}^{(c)}$  are independently generated from the uniform distributions  $U[-\lambda_1^{(c)}, \lambda_1^{(c)}]$  and  $U[-\lambda_2^{(c)}, \lambda_2^{(c)}]$ , respectively. The values  $\lambda_1^{(c)}$  and  $\lambda_2^{(c)}$  are set such that  $\lambda_1^{(c)} \geq \lambda_2^{(c)} > 0$  and  $m^{(c)} + \xi_{i1}^{(c)} \phi_1^{(c)} + \xi_{i2}^{(c)} \phi_2^{(c)} \in V_{\mu_*}(\Omega)$  with probability equal to 1, and we denote  $\lambda^{(c)} = (\lambda_1^{(c)}, \lambda_2^{(c)})$ .

We consider various simulation designs with the combinations of  $m^{(c)}, S^{(c)}$  and  $\lambda^{(c)}$  between clusters that are analogous to those in Chiou and Li (2007). Specifically, they are summarized in Table 1 with the following notation:  $f_1(x) = \Phi_{[0,1]}^{-1}(x; 0.75, 0.3) - x, f_2 = \Phi_{[0,1]}^{-1}(x; 0.75, 0.25) -$ 

Moon functions	Principa	al directions
Mean functions	$\mathcal{S}^{(1)} = \mathcal{S}^{(2)} = E_1$	$\mathcal{S}^{(1)} = E_1, \mathcal{S}^{(2)} = E_2$
(1)(2) $t$		(I): $\lambda^{(1)} = \theta_1/2, \lambda^{(2)} = \theta_2$
$m^{(j)} \equiv m^{(j)} \equiv f_1$		(II): $\lambda^{(1)} = \theta_1, \lambda^{(2)} = \theta_2$
$m^{(1)} = f_1, m^{(2)} = f_2$	(III): $\lambda^{(1)} = \lambda^{(2)} = \theta_1$	(IV): $\lambda^{(1)} = \theta_1, \lambda^{(2)} = \theta_2$
$m^{(1)} = f_1, m^{(2)} = f_3$	(V): $\lambda^{(1)} = \lambda^{(2)} = \theta_1$	(VI): $\lambda^{(1)} = \theta_1, \lambda^{(2)} = \theta_2$
$m^{(1)} = \phi_{11}/10, m^{(2)} = \phi_{11}/15$	(VII): $\lambda^{(1)} = \lambda^{(2)} = \theta_1$	(VIII): $\lambda^{(1)} = \theta_1, \lambda^{(2)} = \theta_2$

Table 1: Simulation designs

 $x, f_3 = \Phi_{[0,1]}^{-1}(x; 0.65, 0.25) - x$ , where  $\Phi_{[0,1]}^{-1}(\cdot; \mu, \sigma)$  denotes the quantile function of the normal distribution with mean  $\mu$  and variance  $\sigma^2$  truncated on [0,1];  $E_1 = \{\phi_{11}, \phi_{12}\}$ , where  $\phi_{11}(x) = \sqrt{2}\sin(2\pi x), \phi_{12}(x) = \sqrt{2}\sin(8\pi x); E_2 = \{\phi_{21}, \phi_{22}\}$ , where  $\phi_{21}(x) = \sqrt{2}\sin(4\pi x), \phi_{22}(x) = \sqrt{2}\sin(6\pi x); \theta_1 = (0.04, 0.001), \theta_2 = (0.02, 0.0133)$ . In Figures 6 and 7, we illustrate 50 realizations of  $\nu_i^{(c)}$  for each cluster under each simulation design. We note that designs (I) and (II) have the same means but different principal directions, designs (III) and (V) have the same principal directions but different means, and designs (IV), (VI) and (VIII) have different means and principal directions. Under design (VII), it holds that span $\mathcal{S}^{(1)} \subset \operatorname{span}\mathcal{S}^{(2)}$  and  $m^{(1)}, m^{(2)} \in \operatorname{span}\mathcal{S}^{(2)}$ . This situation is called the non-identifiable situation by Chiou and Li (2007), where it was reported that k-centres functional clustering method was outperformed by other methods.

Under each simulation design, we generate a synthetic data set as follows. First, for each individual i = 1, ..., n, we independently generate cluster membership variable  $c_i$  with equal probabilities between  $\{1, 2\}$ . Then, we generate distributional data  $\nu_i$  from the model (15) for cluster  $c_i$ . In other words, if  $c_i = 1$ , we generate  $\nu_i$  from the model  $\nu_i^{(1)}$  in (15), and if  $c_i = 2$ , we generate  $\nu_i$  from the model  $\nu_i^{(2)}$  in (15). Furthermore, we generate a set of independent N samples  $Y_i = \{Y_{il}\}_{l=1}^N$ from the distribution  $\nu_i$ . Our aim is to recover the true cluster memberships  $\{c_i, i = 1, ..., n\}$  based on the synthetic data set  $\{Y_i, i = 1, ..., n\}$ .

# 5.2 Other Methods for Comparison, Implementation and Measures of Cluster Quality

The following methods are applied to the synthetic data set for simulation comparisons: the k-means clustering of convex principal component scores (CPCA) that is described in the initial clustering in Section 3.2; the k-centres distributional clustering method (kCDC) which is our proposal; the Wasserstein k-means method (WkM) studied by Papayiannis et al. (2021) and Zhuang et al. (2022), for example; the k-means method with the trimmed Wasserstein distance (WkM<sub> $\delta$ </sub>) by Verdinelli and Wasserman (2019) with a trimming constant  $\delta \in \{0.01, 0.05, 0.1\}$ .

For selecting the dimension M of geodesic modes, we set the threshold value  $\tau = 0.9$ . In the initial clustering and reclassification steps, we use a computational algorithm developed by



Figure 6: Illustration of 50 realizations of distributions  $\nu_i^{(c)}$  for each cluster under simulation designs (I), (II), (III) and (IV). The solid lines are the densities of realizations, and the dotted lines are the densities of the population Fréchet means of clusters.



Figure 7: Illustration of 50 realizations of distributions  $\nu_i^{(c)}$  for each cluster under simulation designs (V), (VI), (VII) and (VIII). The solid lines are the densities of realizations, and the dotted lines are the densities of the population Fréchet means of clusters.

Campbell and Wong (2022) for implementations of convex and geodesic PCA.

To compare performance of the clustering methods, we use two measures of cluster quality. The first is the correct classification rate (cRate), which is defined as the maximal possible ratio of correctly classified objects to the total number of objects to be clustered. By definition, cRate takes a value between 0 and 1, and a larger cRate indicates a better clustering quality. The second measure of clustering quality is the adjusted Rand index (aRand) (Hubert and Arabie, 1985), which is a corrected form of the Rand index (Rand, 1971). The Rand index measures the agreement between two partitions by counting the number of paired objects that are either in the same group or in different groups in both partitions. If the two partitions are an external criterion and a clustering result, then the Rand index can be viewed as the quality of the clustering. aRand is a form of the Rand index that has an expected value 0 and is bounded above by 1. As with cRate, a larger aRand indicates a better clustering quality.

#### 5.3 Result

Table 2 summarizes simulation results with n = 100 and N = 2000. The reported cluster quality outcome is the average of 100 replications for each simulation design. When the two clusters have different principal directions in designs (II), (IV) and (VIII), the performance of the proposed method is better than the other methods. On the other hands, when the two clusters have the same principal directions in designs (III) and (V), or they are in the non-identifiable situation in design (VII), the proposed method is outperformed by the other methods. We note that the proposed method still shows a comparative performance with the other methods in design (V).

# 6 Real Data Analysis

## 6.1 Data

We use a real dataset consisting of population age distributions of districts in Austria for comparing results of several clustering methods. The raw data are available from STATcube – Statistical Database of Statistics Austria (https://www.statistik.at/en/databases/statcubestatistical-database). For a given year, district and sex, this database provides a cross-sectional table in which the number of people at each age is recorded. From this table, one can compute a histgram representing the relative frequency by age, and we call this histgram as a population age distribution. This kind of population age distribution data is often used in the literature of distributional data analysis (Delicado, 2011; Hron et al., 2016; Bigot et al., 2017; Del Barrio et al., 2019). In this study, we use population age distributions for men and women in the 42 districts of Upper and Lower Austria for the year 2020. We remove the data for people that are more than 100 years old, which implies each distribution is supported on the interval  $\Omega = [0, 100]$ . Figure 8 plots their densities obtained by smoothing the histgrams. Our goal is to classify these  $84(= 42 \times 2)$ 

Design						
	CPCA	kCDC	WkM	$\mathrm{WkM}_{0.01}$	$\mathrm{WkM}_{0.05}$	$\mathrm{WkM}_{0.1}$
(I)						
cRate	0.549	0.529	0.547	0.548	0.548	0.538
aRand	0.003	-0.005	0.002	0.002	0.002	-0.003
(II)						
cRate	0.694	0.876	0.694	0.694	0.694	0.694
aRand	0.147	0.575	0.147	0.147	0.147	0.147
(III)						
cRate	0.684	0.552	0.684	0.684	0.684	0.657
aRand	0.133	0.009	0.133	0.133	0.133	0.098
(IV)						
cRate	0.807	0.894	0.807	0.807	0.808	0.779
aRand	0.378	0.636	0.378	0.378	0.381	0.316
(V)						
cRate	0.781	0.713	0.781	0.781	0.781	0.781
aRand	0.447	0.340	0.447	0.447	0.447	0.447
(VI)						
cRate	1.000	0.988	1.000	1.000	1.000	1.000
aRand	1.000	0.954	1.000	1.000	1.000	1.000
(VII)						
cRate	0.699	0.547	0.698	0.698	0.698	0.698
aRand	0.156	0.022	0.154	0.154	0.154	0.154
(VIII)						
cRate	0.799	0.835	0.800	0.799	0.800	0.799
aRand	0.356	0.465	0.358	0.356	0.358	0.356

Table 2: Simulation results. The greater the index cRate and cRand, the higher the quality of the clustering. The proposed method is k-centres distributional clustering (kCDC).



Figure 8: Densities of population age distributions for men and women in the 42 districts in Upper and Lower Austria for the year 2020. The black solid lines are for men, and the red dashed lines are for women.

distributions into two groups with several clustering methods and determine whether the clustering results reflect gender differences.

#### 6.2 Setup and Result

We compare the results of the six clustering methods used in the simulation study. For selecting the dimension M of geodesic modes of proposed method, we set the threshold value  $\tau = 0.8$ . By this criterion, M = 1 is selected. We use a computational algorithm developed by Campbell and Wong (2022) for implementations of convex and geodesic PCA. The correct classification rate (cRate) and adjusted Rand index (aRand) are used as measures of cluster quality.

Table 3 summarizes the results. We see that the proposed method outperforms the other methods in distinguishing the gender groups. We further investigate the structure of each cluster by visualizing the estimated Fréchet mean and mode of variation. Figure 9 shows the estimated Fréchet means of the two clusters. We see that the Fréchet mean of cluster 2 (the females group) has a heavier tail on the age range (60, 100) than that of cluster 1 (the males group). This result reasonably reflects the fact that female persons generally live longer than male persons. Figure 10

	CPCA	kCDC	WkM	$\mathrm{WkM}_{0.01}$	$\mathrm{WkM}_{0.05}$	$\mathrm{WkM}_{0.1}$
cRate	0.797	0.928	0.797	0.809	0.797	0.797
aRand	0.346	0.731	0.346	0.375	0.346	0.346

Table 3: Cluster qualities for the population age distribution data



Figure 9: Densities of the estimated Fréhcet means of the two clusters. The black solid line is for cluster 1 (the males group), and the red dashed line is for cluster 2 (the females group).

illustrates the modes of variations of the two clusters based on the estimated principal geodesics. For cluster 1 (the males group), we see that the largest variability occurs in the range (20, 40) and a large variability occurs in the range (70, 80). For cluster 2 (the female groups), we see that two large variabilities occur in the ranges (10, 40) and (70, 80). In summary, the proposed method identifies the gender groups with the highest cluster quality, and it also provides a visual insight into the clusters.

## 7 Discussion

In this paper, we propose a novel clustering method for distributional data on the real line. This method performs clustering for distributional data in the spirit of the k-centres clustering approach of Chiou and Li (2007) for functional data. We use geodesic PCA in the Wasserstein space to define the geodesic mode of the variation of clusters, which is used in the reclassification step



Figure 10: Illustrations of the modes of variations of the two clusters based on the estimated principal geodesics. For each cluster  $c \in \{1, 2\}$  and value  $\alpha \in \mathbb{R}$ , we define a distribution  $\hat{\nu}^{(c)}(\alpha) \in \mathcal{P}(\Omega)$  by  $\tilde{\nu}^{(c)}(\alpha) = \operatorname{Exp}_{\mu_*}(\hat{g}^{(c)} + \alpha \sqrt{\hat{\lambda}_1^{(c)}} \hat{\phi}_1^{(c)})$ . Here,  $\mu_* \in \mathcal{P}(\Omega)$  is some reference measure,  $\hat{g}^{(c)} \in \mathcal{L}^2_{\mu_*}(\Omega)$  is the estimated mean in the tangent space,  $\hat{\phi}_1^{(c)} \in \mathcal{L}^2_{\mu_*}(\Omega)$  is the estimated first convex principal direction and  $\hat{\lambda}_1^{(c)}$  is the variance of the first convex principal component scores. For each cluster, the black solid line is the density corresponding to  $\alpha = 0$ , while the boundaries of the shaded region are densities corresponding to  $\alpha = \pm 1$ .

for determining the cluster membership of each distribution. In contrast to conventional clustering methods like the k-means type clustering methods, the proposed method takes account of difference in both the mean and mode of variation structues of clusters, potentially improving cluster quality. Our theory demonstrates the validity of our clustering method, which we illustrate through a simulation study and real data analysis.

We note that the proposed clustering method lacks a theoretical support of consistency. For classical k-means clustering, where the cluster centre is only the cluster mean, Pollard (1981) established the almost sure convergence of the set of cluster centres. However, for k-centres type clustering, where the cluster centres include both the cluster mean and principal component directions, such consistency results do not seem to have been established yet, and further technical development is needed.

There are several directions for future extensions. First, one can consider a clustering method for distributional data in the spirit of the correlation-based functional clustering of (Chiou and Li, 2008), which groups curves with similar shapes. Second, we can consider applying clustering methods to multivariate distributional data. The Wasserstein space of probability distributions on a multi-dimensional Euclidean space can be endowed with the basic concepts of Riemannian manifolds, such as the tangent space, the exponential map, and the logarithmic map (see Section S.5.1 in the Supplementary Material for details). However, in the multivariate setting, the logarithmic map generally does not possess the isometric property. This implies that a clear connection between geodesic PCA in the Wasserstein space and convex PCA in the tangent space is not available. Additionally, the Wasserstein distance and logarithmic map generally do not have closed-form expressions in the multivariate setting, which brings about difficulties in numerical computation. These factors need to be considered when extending the proposed clustering method to the multivariate setting. As a special case, for multivariate Gaussian distributions, the Wasserstein distance and the logarithmic map have closed-form expressions. Based on this fact, we present a k-centres clustering method for multivariate Gaussian distributions in Section S.5 of the Supplementary Material.

# Supplementary Material

# S.1 Convex Principal Component Analysis

## S.1.1 Formulation

Convex PCA was originally introduced by Bigot et al. (2017) to analyze geodesic PCA in the Wasserstein space. In this section, we review a general formulation of convex PCA and its link to geodesic PCA in the Wasserstein space. Let H be a separable Hilbert space over  $\mathbb{R}$ , with an inner product  $\langle \cdot, \cdot \rangle_H$  and norm  $\|\cdot\|_H$ , and X be a nonempty closed convex subset of H. Let  $\mathbf{x}$  be an X-valued random variable, assumed to be square-integrable in the sense that  $\mathbb{E}[\|\mathbf{x}\|_H^2] < \infty$ . The mean of  $\mathbf{x}$  is defined as the unique element in  $\arg \min_{y \in X} \mathbb{E}[\|\mathbf{x} - y\|_H^2]$ , and we denote it as  $\overline{x}$ .

To formulate convex PCA, we introduce the following. For  $x, y \in H$  and nonempty  $E \subset H$ , we define  $d(x, y) = ||x - y||_H$  and  $d(x, E) = \inf_{z \in E} d(x, z)$ . Note that if E is closed and convex, there exists an unique point  $z^* \in E$  such that  $d(x, z^*) = d(x, E)$  by the Hilbert projection theorem. We define  $K(C) = \mathbb{E}[d^2(\mathbf{x}, C)]$  for any nonempty set  $C \subset X$ . For  $A \subset H$ , its dimension, dim(A), is defined as the dimension of the smallest affine subspace of H containing A. For any integer  $j \geq 1$ , we denote by  $CC_{\overline{x},j}$  the family of nonempty, closed and convex subsets  $C \subset X$ , such that  $\dim(C) \leq j$  and  $\overline{x} \in C$ .

Now we state a notion of convex PCA, nested convex PCA.

**Definition 2** (Nested convex PCA). Let  $C_0 = \{\overline{x}\}$ . For a given integer  $M \ge 1$ , we say that a set  $C_M \subset X$  is an  $(M, \overline{x})$ -nested principal convex component of  $\mathbf{x}$  if there exist sets  $C_0, ..., C_{M-1}$  such that  $C_0 \subset C_1 \subset \cdots \subset C_{M-1} \subset C_M$ , and for each j = 1, ..., M,

$$C_j \in \operatorname*{arg\,min}_{C \in CC_{\overline{x},j}, C \supset C_{j-1}} K(C).$$

When X is compact, the nested convex PCA problem has nonempty solutions for any  $M \ge 1$ (Theorem 3.1, Bigot et al., 2017). Furthermore, a solution of the nested convex PCA problem can be obtained by constructing a sequence of orthogonal vectors in H as follows. For  $\phi \in H$ , we let

$$V(\phi) = \mathbb{E}\left[\min_{z \in (\overline{x} + \operatorname{span}\{\phi\}) \cap X} \|\mathbf{x} - z\|_{H}^{2}\right].$$

Define

$$\phi_1^* \in \operatorname*{arg\,min}_{\phi \in H: \|\phi\|_H = 1} V(\phi),$$

and for  $j \geq 2$ ,

$$\phi_j^* \in \operatorname*{arg\,min}_{\phi \in H: \phi \in P_{j-1}^\top, \|\phi\|_H = 1} V(\phi),$$

where  $P_{j-1} = \operatorname{span}\{\phi_1, ..., \phi_{j-1}\}$ . Then if we set

$$C_j = (\overline{x} + \operatorname{span}\{\phi_1^*, \dots, \phi_j^*\}) \cap X$$
(16)

for j = 1, ..., M, the sets  $\{C_j\}_{j=1}^M$  form a sequence of nested principal convex components of **x** (Proposition 3.4, Bigot et al., 2017). We call the above vector  $\phi_j^*$  as the *j*-th convex principal direction of **x**.

## S.1.2 Low-Dimensional Representation and Explained Variation

Based on the result of the nested convex PCA problem, a notion of low-dimensional representation or dimension reduction of the random variable  $\mathbf{x}$  can be defined. Given an  $(M, \overline{x})$ -nested principal convex component  $C_M \subset X$  of  $\mathbf{x}$ , we define the *M*-dimensional representation of  $\mathbf{x}$  as

$$\Pi_{C_M} \mathbf{x} \in \operatorname*{arg\,min}_{z \in C_M} \|\mathbf{x} - z\|_H^2,\tag{17}$$

which uniquely exists. Especially, if  $C_M$  is of the form (16), there exist coefficients  $\xi_1, ..., \xi_M \in \mathbb{R}$  such that

$$\Pi_{C_M} \mathbf{x} = \overline{x} + \sum_{j=1}^M \xi_j \phi_j^*.$$
(18)

We call the scalar  $\xi_j$  as the *j*-th convex principal component score of  $\mathbf{x}$ . Note that in contrast to the ordinal PCA, the *M*-dimensional representation  $\Pi_{C_M} \mathbf{x}$  does not always obtained by orthogonally projecting  $\mathbf{x}$  onto the space  $\overline{x} + \text{span}\{\phi_1^*, ..., \phi_M^*\}$ , and thus the convex principal component score  $\xi_j$  in (18) is not necessarily equal to the inner product  $\langle \mathbf{x} - \overline{x}, \phi_j^* \rangle_H$ .

As with the ordinal PCA, a notion of explained variation can be defined for convex PCA (Campbell and Wong, 2022). The cumulative proportion of variation explained by the principal convex component  $C_M$  is defined as

$$EV_c(C_M) = \frac{\mathbb{E}[\|\Pi_{C_M} \mathbf{x} - \overline{x}\|_H^2]}{TV_c},$$

where the total variation  $TV_c$  is defined as  $\mathbb{E}[\|\mathbf{x} - \overline{x}\|_H^2]$ . Obviously,  $0 \leq EV_c(C_M) \leq 1$  holds for any  $M \geq 1$ , and  $EV_c(C_M)$  is monotonic in M:  $EV_c(C_M) \leq EV_c(C_{M+1})$  for any  $M \geq 1$ . For further detail of this notion of variation explained, see Section 2 of Campbell and Wong (2022).

## S.1.3 Link to Geodesic PCA in the Wasserstein Space

The notion of convex PCA is strongly linked to geodesic PCA in the Wasserstein space. Let  $\boldsymbol{\mu}$  be a square-integrable  $\mathcal{P}(\Omega)$ -valued random variable. With an absolutely continuous reference measure  $\mu_*$ , define a  $V_{\mu_*}(\Omega)$ -valued random variable  $\mathbf{g} = \text{Log}_{\mu_*}\boldsymbol{\mu}$ , and denote its mean as  $\overline{g}$ . Let  $C_M \subset V_{\mu_*}(\Omega)$  be an  $(M, \overline{g})$ -nested principal convex component of  $\mathbf{g}$  obtained by applying the nested

convex PCA to  $H = \mathcal{L}^2_{\mu_*}(\Omega), X = V_{\mu_*}(\Omega)$  and  $\mathbf{x} = \mathbf{g}$ . Then if we set  $G_M = \operatorname{Exp}_{\mu_*}(C_M)$ , the set  $G_M$  is an  $(M, \nu_{\oplus})$ -nested principal geodesic of  $\boldsymbol{\mu}$  (Proposition 4.4, Bigot et al., 2017). In Figure 1, swe illustrated this relationship between geodesic PCA in the Wasserstein space and convex PCA in the tangent space.

# S.2 Proofs

For the proofs, we define stochastic models of the random element  $\mathbf{g}^{(c)}$ . Under Assumption 1,  $m^{(c)}$  and  $m^{(d)}$  are the mean functions of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$ , and  $\rho_j^{(c)}$  and  $\rho_j^{(d)}$  are the *j*-th convex principal directions of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$ , respectively. Hence,  $(1, m^{(c)})$  and  $(1, m^{(d)})$ -nested principal convex components of  $\mathbf{g}^{(c)}$  and  $\mathbf{g}^{(d)}$  are defined as  $C^{(c)} = (m^{(c)} + \operatorname{span}\{\rho_1^{(c)}\}) \cap V_{\mu_*}(\Omega)$  and  $C^{(d)} = (m^{(d)} + \operatorname{span}\{\rho_1^{(d)})\} \cap V_{\mu_*}(\Omega)$ , respectively. Then for the random variable  $\mathbf{g}^{(c)}$ , we define stochastic models  $\tilde{\mathbf{g}}_c^{(c)}$  and  $\tilde{\mathbf{g}}_c^{(d)}$  as

$$\tilde{\mathbf{g}}_c^{(c)} = \Pi_{C^{(c)}} \mathbf{g}^{(c)} \quad \text{and} \quad \tilde{\mathbf{g}}_c^{(d)} = \Pi_{C^{(d)}} \mathbf{g}^{(c)},$$

respectively. Under Assumption 1, we have  $\tilde{\mathbf{g}}_{c}^{(c)} = m^{(c)} + \xi_{1}^{(c)} \rho_{1}^{(c)}$ . Additionally, under Assumptions 1 and 2, we have  $\tilde{\mathbf{g}}_{c}^{(d)} = m^{(d)} + \langle \mathbf{g}^{(c)} - m^{(d)}, \rho_{1}^{(d)} \rangle_{\mu_{*}} \rho_{1}^{(d)}$ .

Proof of Proposition 3. By the isometric property of the logarithmic map, we have

$$d_W(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_c^{(c)}) = \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_c^{(c)}\|_{\mu_*} \quad \text{and} \quad d_W(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_c^{(d)}) = \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_c^{(d)}\|_{\mu_*}.$$
 (19)

Note that under Assumption 1, it holds that  $\xi_1^{(c)} = \langle \mathbf{g}^{(c)} - m^{(c)}, \rho_1^{(c)} \rangle_{\mu_*}$ . Thus, we have

$$\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}}^{2} = \|\mathbf{g}^{(c)} - m^{(c)} - \langle \mathbf{g}^{(c)} - m^{(c)}, \rho_{1}^{(c)} \rangle_{\mu_{*}} \rho_{1}^{(c)}\|_{\mu_{*}}^{2}$$
$$= \|\mathbf{g}^{(c)} - m^{(c)}\|_{\mu_{*}}^{2} - \langle \mathbf{g}^{(c)} - m^{(c)}, \rho_{1}^{(c)} \rangle_{\mu_{*}}^{2}.$$
(20)

Additionally, under Assumptions 1 and 2, we have

$$\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}}^{2} = \|\mathbf{g}^{(c)} - m^{(d)} - \langle \mathbf{g}^{(c)} - m^{(d)}, \rho_{1}^{(d)} \rangle_{\mu_{*}} \rho_{1}^{(d)}\|_{\mu_{*}}^{2}$$
$$= \|\mathbf{g}^{(c)} - m^{(d)}\|_{\mu_{*}}^{2} - \langle \mathbf{g}^{(c)} - m^{(d)}, \rho_{1}^{(d)} \rangle_{\mu_{*}}^{2}.$$
(21)

Combing (19), (20), (21), under the assumption  $m^{(c)} = m^{(d)} = m$ , we obtain

$$\begin{split} P(d_{W}(\boldsymbol{\nu}^{(c)},\tilde{\boldsymbol{\nu}}_{c}^{(c)}) &< d_{W}(\boldsymbol{\nu}^{(c)},\tilde{\boldsymbol{\nu}}_{c}^{(d)})) \\ &= P(\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}}^{2} < \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}}^{2}) \\ &= P(\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}}^{2} < \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}}^{2}) \\ &= P(\langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} \rangle_{\mu_{*}}^{2} - \langle \mathbf{g}^{(c)} - m, \rho_{1}^{(d)} \rangle_{\mu_{*}}^{2} > 0) \\ &= P(\langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} > 0, \langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} > 0) \\ &+ P(\langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} + \rho_{1}^{(d)} \rangle_{\mu_{*}} < 0, \langle \mathbf{g}^{(c)} - m, \rho_{1}^{(c)} - \rho_{1}^{(d)} \rangle_{\mu_{*}} < 0). \end{split}$$

Proof of Proposition 4. As in the proof of Proposition 3, we have the equations (19) and  $\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}}^{2}$  and  $\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}}^{2}$  are expressed as (20) and (21), respectively. For the first and second terms in (21), we have

$$\|\mathbf{g}^{(c)} - m^{(d)}\|_{\mu_*}^2 = \|\mathbf{g}^{(c)} - m^{(c)}\|_{\mu_*}^2 + \|m^{(c)} - m^{(d)}\|_{\mu_*}^2 + 2\langle \mathbf{g}^{(c)} - m^{(c)}, m^{(c)} - m^{(d)} \rangle_{\mu_*}$$

and

$$\langle \mathbf{g}^{(c)} - m^{(d)}, \rho_1^{(d)} \rangle_{\mu_*}^2 = \langle \mathbf{g}^{(c)} - m^{(c)}, \rho_1^{(d)} \rangle_{\mu_*}^2 + \langle m^{(c)} - m^{(d)}, \rho_1^{(d)} \rangle_{\mu_*}^2 + 2 \langle \mathbf{g}^{(c)} - m^{(c)}, \rho_1^{(d)} \rangle_{\mu_*} \langle m^{(c)} - m^{(d)}, \rho_1^{(d)} \rangle_{\mu_*}$$

respectively. Thus, with  $\psi = 2m^{(c)} - 2m^{(d)} - 2\langle m^{(c)} - m^{(d)}, \rho_1 \rangle_{\mu_*} \rho_1$ , we have

$$\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}^{(d)}_{c}\|^{2}_{\mu_{*}} = \|\mathbf{g}^{(c)} - m^{(c)}\|^{2}_{\mu_{*}} - \langle \mathbf{g}^{(c)} - m^{(d)}, \rho_{1}^{(d)} \rangle^{2}_{\mu_{*}} + \langle \mathbf{g}^{(c)} - m^{(c)}, \psi \rangle_{\mu_{*}} + \|m^{(c)} - m^{(d)}\|^{2}_{\mu_{*}} - \langle m^{(c)} - m^{(d)}, \rho_{1}^{(d)} \rangle^{2}_{\mu_{*}}.$$

$$(22)$$

Combining (19), (20) and (22), under the assumption  $\rho_j^{(c)} = \rho_j^{(d)} = \rho_j$  for j = 1, ..., J, we obtain

$$P(d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(c)}) < d_{W}(\boldsymbol{\nu}^{(c)}, \tilde{\boldsymbol{\nu}}_{c}^{(d)}))$$

$$= P(\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}} < \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}})$$

$$= P(\|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(c)}\|_{\mu_{*}}^{2} < \|\mathbf{g}^{(c)} - \tilde{\mathbf{g}}_{c}^{(d)}\|_{\mu_{*}}^{2})$$

$$= P(\langle \mathbf{g}^{(c)} - m^{(c)}, \psi \rangle_{\mu_{*}} > \langle m^{(c)} - m^{(d)}, \rho_{1} \rangle_{\mu_{*}}^{2} - \|m^{(c)} - m^{(d)}\|_{\mu_{*}}^{2}).$$

# S.3 Estimation of Distributional Data

Let assume that the *n* distributional data  $\nu_1, ..., \nu_n \in \mathcal{P}(\Omega)$  are not directly observed, and instead we observe for each i = 1, ..., n, a collection of independent measurements  $\{Y_{il}\}_{l=1}^{N_i}$  sampled from  $\nu_i$ . Here,  $N_i$  is the sample size which may vary across distributions. In this case, we estimate the distribution  $\nu_i$  from  $\{Y_{il}\}_{l=1}^{N_i}$  before implementing the clustering method. One option is estimating the distribution function  $F_i$  of  $\nu_i$  with the empirical distribution function  $\hat{F}_i$  of  $\{Y_{il}\}_{l=1}^{N_i}$ ,

$$\hat{F}_i(x) = \frac{1}{N_i} \sum_{l=1}^{N_i} I_{(-\infty,x]}(Y_{il})$$

The quantile function  $F_i^{-1}$  of  $\nu_i$  is then estimated by converting  $\hat{F}_i$  to a quantile function  $\hat{F}_i^{-1}$  by left continuous inversion,

$$\hat{F}_i^{-1}(u) = \inf\{x \in \mathbb{R} : \hat{F}_i(x) \ge u\}.$$

Another option is estimating the density function of  $\nu_i$  from  $\{Y_{il}\}_{l=1}^{N_i}$  and then computing the distribution and quantile functions by integration and inversion.

# S.4 Selection of the Dimension of Geodesic Modes by Convex PCA

As described in Section 3.2, we can select the dimension M of the geodesic modes by using convex PCA, as an alternative to geodesic PCA. Specifically, for any  $M' \in \mathbb{N}$ , let  $\hat{C}_{M'} \subset \mathcal{L}^2_{\hat{\mu}_*}(\Omega)$  be an  $(M', \overline{g})$ -nested principal convex component of the *n* transformed data  $g_1, \ldots, g_n$ . The cumulative proportion of variation explained by  $\hat{C}_{M'}$  is defined as

$$EV_c(\hat{C}_{M'}) = \frac{n^{-1} \sum_{i=1}^n \|\Pi_{\hat{C}_{M'}} g_i - \overline{g}\|_{\mu_*}^2}{TV_c},$$

where the total variation is written as  $TV_c = n^{-1} \sum_{i=1}^n ||g_i - \overline{g}||^2_{\mu_*}$ , and  $\Pi_{\hat{C}_{M'}}g_i$  is the projection of  $g_i$  onto  $\hat{C}_{M'}$ . By the isometric property of the logarithmic map, it can be shown that  $EV_c(\hat{C}_{M'}) = EV(\hat{G}_{M'})$  for any  $M' \in \mathbb{N}$ . Hence, value of M selected as (8) is equal to the value selected as

$$M = \min\{M' \in \mathbb{N} : EV_c(\hat{C}_{M'}) \ge \tau\}$$

## S.5 k-Centres Clustering for Multivariate Gaussian Distributions

## S.5.1 Optimal Transport of Gaussian Distributions on $\mathbb{R}^d$

In this subsection, we provide some background on optimal transport of Gaussian distributions on  $\mathbb{R}^d$ , for  $d \geq 1$ . Firstly, we explain optimal transport of general probability distributions on  $\mathbb{R}^d$  (Ambrosio et al., 2008; Villani, 2008; Panaretos and Zemel, 2020). Let  $\mathcal{P}_2(\mathbb{R}^d)$  be the set of Borel probability measures on  $\mathbb{R}^d$  with finite second moments. The 2-Wasserstein distance between  $\mu_1, \mu_2 \in \mathcal{P}_2(\mathbb{R}^d)$  is defined by

$$d_W(\mu_1, \mu_2) = \left(\inf_{\pi \in \Gamma(\mu_1, \mu_2)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|^2 d\pi(x, y)\right)^{1/2},$$

where  $\|\cdot\|$  denotes the Euclidean norm on  $\mathbb{R}^d$ , and the infimum is taken over the set  $\Gamma(\mu_1, \mu_2)$ of all couplings of  $\mu_1$  and  $\mu_2$ . The 2-Wasserstein distance  $d_W$  is a metric on  $\mathcal{P}_2(\mathbb{R}^d)$ , and the metric space  $(\mathcal{P}_2(\mathbb{R}^d), d_W)$  is called the Wasserstein space. For two given measures  $\mu_*, \mu \in \mathcal{P}_2(\mathbb{R}^d)$ , any map  $\mathbf{t} : \mathbb{R}^d \to \mathbb{R}^d$  that minimizes Monge's problem  $\min_{\mathbf{t} \neq \mu_* = \mu} \int_{\mathbb{R}^d} ||\mathbf{t}(x) - x||^2 d\mu_*(x)$  is called an optimal transport map from  $\mu_*$  to  $\mu$ . If  $\mu_*$  is absolutely continuous with respect to Lebesgue measure on  $\mathbb{R}^d$ , such optimal transport map uniquely exists, and we denote it as  $\mathbf{t}_{\mu_*}^{\mu}$ . We note that when d = 1, the optimal transport map is given by  $\mathbf{t}_{\mu_*}^{\mu} = F^{-1} \circ F_*$ , where  $F_*$  and  $F^{-1}$  are the distribution function of  $\mu_*$  and the quantile function of  $\mu$ , respectively. Analogously to the case of d = 1, the notions of tangent space, exponential map and logarithmic map at  $\mu_*$  can be defined. Specifically, the tangent space is the Hilbert space  $\mathcal{L}^2_{\mu_*}(\mathbb{R}^d)$  of  $\mathbb{R}^d$ -valued functions g on  $\mathbb{R}^d$  that are  $\mu_*$ -square-integrable in the sense  $\int_{\mathbb{R}^d} ||g||^2 d\mu_* < \infty$ . This space  $\mathcal{L}^2_{\mu_*}(\mathbb{R}^d)$  is equipped with an inner product  $\langle \cdot, \cdot \rangle_{\mu_*}$  defined by  $\langle g_1, g_2 \rangle_{\mu_*} = \int_{\mathbb{R}^d} g_1^\top g_2 d\mu_*$  and norm  $\|\cdot\|_{\mu_*}$  defined by  $\|g\|_{\mu_*} = \langle g, g \rangle_{\mu_*}^{1/2}$ . The exponential map  $\operatorname{Exp}_{\mu_*} : \mathcal{L}^2_{\mu_*}(\mathbb{R}^d) \to \mathcal{P}_2(\mathbb{R}^d)$  is defined by  $\operatorname{Exp}_{\mu_*}g = (g + \mathrm{id})\#\mu_*$ , and the logarithmic map  $\operatorname{Log}_{\mu_*} : \mathcal{P}_2(\mathbb{R}^d) \to \mathcal{L}^2_{\mu_*}(\mathbb{R}^d)$  is defined by  $\operatorname{Log}_{\mu_*}\mu = \mathbf{t}^{\mu}_{\mu_*} - \mathrm{id}$ .

Next, we explain optimal transport of multivariate Gaussian measures (see Section 1.6.3 of Panaretos and Zemel (2020) for further details). We simplify the setting by focusing on centred Gaussian measures. Let  $\mathcal{G}_0(d)$  be the set of Gaussian measures on  $\mathbb{R}^d$  with zero means. For Gaussian measures  $\mu_1 = N(0, \Sigma_1)$  and  $\mu_2 = N(0, \Sigma_2)$  in  $\mathcal{G}_0(d)$ , the Wasserstein distance between them is explicitly expressed as

$$d_W(\mu_1, \mu_2) = \sqrt{\operatorname{tr}[\Sigma_1 + \Sigma_2 - 2(\Sigma_1^{1/2} \Sigma_2 \Sigma_1^{1/2})^{1/2}]}.$$

Here, for any positive semidefinite and symmetric matrix  $\Sigma$ , we denote its principal square root as  $\Sigma^{1/2}$ . Also, for two given Gaussian measures  $\mu_* = N(0, \Sigma_*)$  and  $\mu = N(0, \Sigma)$  in  $\mathcal{G}_0(d)$ , where  $\mu_*$  is assumed to be non-singular, the optimal transport map from  $\mu_*$  to  $\mu$  is expressed as  $\mathbf{t}^{\mu}_{\mu_*}(x) = \Sigma_*^{-1/2} [\Sigma_*^{1/2} \Sigma \Sigma_*^{1/2}]^{1/2} \Sigma_*^{-1/2} x$ .

Now, we consider the notions of tangent space, exponential and logarithmic maps of the space of Gaussian distributions  $(\mathcal{G}_0(d), d_W)$  at the non-singular reference measure  $\mu_* = N(0, \Sigma_*) \in \mathcal{G}_0(d)$ . Let Sym(d) denote the set of  $d \times d$  symmetric matrices, and denote with  $g_V$  the linear transformation on  $\mathbb{R}^d$  represented by a matrix  $V \in \text{Sym}(d)$ . Then, for any  $\mu = N(0, \Sigma)$  in  $\mathcal{G}_0(d)$ , we have  $\text{Log}_{\mu*}\mu = \mathbf{t}_{\mu*}^{\mu} - \text{id} = g_{\Sigma_*^{-1/2}[\Sigma_*^{1/2}\Sigma\Sigma_*^{1/2}]^{1/2}\Sigma_*^{-1/2}} - g_{I_d} = g_{\Sigma_*^{-1/2}[\Sigma_*^{1/2}\Sigma\Sigma_*^{-1/2}-I_d]}$ , where  $I_d$  denotes the identity matrix of size d. Moreover, we have  $\langle g_{V_1}, g_{V_2} \rangle_{\mu*} = \int_{\mathbb{R}^d} (V_1 x)^\top (V_2 x) d\mu_*(x) = \text{tr}[V_1 \Sigma_* V_2]$ for any  $V_1, V_2 \in \text{Sym}(d)$ , and  $\text{Exp}_{\mu*}g_V = (g_V + \text{id})\#\mu_* = N(0, (V + I_d)\Sigma_*(V + I_d))$  for any  $V \in \text{Sym}(d)$ . Based on these observations and the identification of the linear transformation  $g_V$ with the matrix V, we again define the tangent space, exponential and logarithmic maps of the space  $(\mathcal{G}_0, d_W)$  as follows. The tangent space at the reference measure  $\mu_* = N(0, \Sigma_*)$  is defined as a finite-dimensional inner product space  $(\text{Sym}(d), \langle \cdot, \cdot \rangle_{\Sigma_*})$ , where  $\langle \cdot, \cdot \rangle_{\Sigma_*}$  is an inner product defined by  $\langle V_1, V_2 \rangle_{\Sigma_*} = \text{tr}(V_1 \Sigma_* V_2)$ . We denote the norm induced by this inner product as  $\| \cdot \|_{\Sigma_*}$ . The exponential map  $\text{Exp}_{\Sigma_*}: \text{Sym}(d) \to \mathcal{G}_0(d)$  is then defined by

$$\operatorname{Exp}_{\Sigma_*} V = N(0, (V + I_d)\Sigma_*(V + I_d)),$$

and the logarithmic map  $\operatorname{Log}_{\Sigma_*} : \mathcal{G}_0(d) \to \operatorname{Sym}(d)$  is defined by

$$\mathrm{Log}_{\Sigma_*}\mu = \Sigma_*^{-1/2} [\Sigma_*^{1/2} \Sigma \Sigma_*^{1/2}]^{1/2} \Sigma_*^{-1/2} - I_d,$$

for  $\mu = N(0, \Sigma) \in \mathcal{G}_0(d)$ . We denote the range of the logarithmic map  $\operatorname{Log}_{\Sigma_*}$  as  $V_{\Sigma_*}(d) \subset \operatorname{Sym}(d)$ . It can be easily checked that  $V_{\Sigma_*}(d)$  is expressed as

 $V_{\Sigma_*}(d) = \{ V \in \operatorname{Sym}(d) : V + I_d \text{ is positive semidefinite} \},\$ 

and hence  $V_{\Sigma_*}(d)$  is closed and convex in Sym(d).

In contrast to the one-dimensional case, the space  $(\mathcal{G}_0(d), d_W)$  is not isometric to the set  $V_{\Sigma_*}(d)$  in Sym(d). The following proposition shows that the logarithmic map  $\mathrm{Log}_{\Sigma_*}$  has an isometric property on a specific class of Gaussian distributions.

**Proposition S1** (cf. Panaretos and Zemel (2020), Section 2.3.2). Let  $\mathscr{C} \subset \mathscr{G}_0(d)$  be a class of Gaussian distributions such that  $\mu_* \in \mathscr{C}$  and  $\Sigma_1 \Sigma_2 = \Sigma_2 \Sigma_1$  for any  $\mu_1 = N(0, \Sigma_1), \mu_2 = N(0, \Sigma_2)$  in  $\mathscr{C}$ . Then we have

$$d_W(\mu_1, \mu_2) = \| \operatorname{Log}_{\Sigma_*} \mu_1 - \operatorname{Log}_{\Sigma_*} \mu_2 \|_{\Sigma_*}$$

for any  $\mu_1, \mu_2 \in \mathscr{C}$ .

#### S.5.2 Clustering Procedure

Setting We propose a clustering method for multivariate Gaussian distributions based on the k-centres clustering approach. Suppose there are n Gaussian distributions  $\nu_1 = N(0, \Sigma_1), ..., \nu_n = N(0, \Sigma_n)$  in  $\mathcal{G}_0(d)$ , and we aim to classify them into K groups. As with the case of univariate distributions, the distributions  $\nu_i$  may be not directly observed, and instead we observe a collection of independent measurements  $\{Y_{il}\}_{l=1}^{N_i}$  sampled from  $\nu_i$ . In this case, we need to estimate the Gaussian distribution  $\nu_i$ , especially its covariance matrix  $\Sigma_i$ , from the measurements. A natural estimator is the empirical covariance matrix

$$\hat{\Sigma}_i = \frac{1}{N_i} \sum_{l=1}^{N_i} (Y_{il} - \overline{Y}_i) (Y_{il} - \overline{Y}_i)^\top,$$

where  $\overline{Y}_i = N_i^{-1} \sum_{l=1}^{N_i} Y_{il}$  is the empirical mean. To keep notations simple, we will use the same notation  $\nu_i$  for the estimated distribution. The procedure of the proposed clustering method is as follows.

**Initial Clustering** As with the case of univariate distributions, we initially classify the *n* distributions into *K* groups by performing dimension reduction based on the stochastic structure of overall data and then applying a conventional clustering algorithm on the resulting low-dimensional vectors. Let  $\hat{\mu}_* \in \mathcal{G}_0(d)$  be a reference Gaussian distribution with a covariance matrix  $\hat{\Sigma}_*$  and transform the Gaussian distributions  $\nu_i$  as  $V_i = \text{Log}_{\hat{\Sigma}_*}\nu_i, i = 1, ..., n$ . A typical choice of  $\hat{\mu}_*$  is the empirical Fréchet mean of  $\nu_1, ..., \nu_n$  in the space  $(\mathcal{G}_0(d), d_W)$ ,

$$\hat{\nu}_{\oplus} = \operatorname*{arg\,min}_{\mu \in \mathcal{G}_0(d)} \frac{1}{n} \sum_{i=1}^n d_W^2(\nu_i, \mu)$$

which can be computed efficiently by using the steepest descent algorithm (Zemel and Panaretos, 2019). Let M be an integer such that  $1 \leq M \leq d(d+1)/2$ . We have the M convex principal directions  $\{\hat{\Phi}_1^*, ..., \hat{\Phi}_M^*\} \subset \text{Sym}(d)$  by applying convex PCA to the data  $V_1, ..., V_n$  with the Hilbert

space  $(\text{Sym}(d), \langle \cdot, \cdot \rangle_{\Sigma_*})$  and convex set  $V_{\hat{\Sigma}_*}(d)$ . Then the set  $\hat{C}_M = (\overline{V} + \text{span}\{\hat{\Phi}_1, ..., \hat{\Phi}_M\}) \cap V_{\hat{\Sigma}_*}(d)$ is an  $(M, \overline{V})$ -principal convex component of  $V_1, ..., V_n$ . Based on this result, for each i = 1, ..., n, we obtain the convex principal component scores of  $V_i, \hat{\xi}_i = (\hat{\xi}_{i1}, ..., \hat{\xi}_{iM}) \in \mathbb{R}^M$ . The initial clustering membership is determined by applying a conventional algorithm such as the *k*-means method to the *M*-dimensional vectors  $\hat{\xi}_i, i = 1, ..., n$ . As with the case of univariate distributions, we use the notion of explained variation for convex PCA to choose the dimension *M* of the principal convex component.

**Reclassification** With the initial clustering results, we use the mean and covariance structures of the clusters to reclassify each data into a best predicted cluster. Specifically, let  $h_i^{(t)} \in \{1, ..., K\}$ be the label of cluster membership for the *i*-th distribution at the *t*-th iteration. Given the set of clustering results  $\mathcal{H}^{(t)} = \{h_i^{(t)} : i = 1, ..., n\}$ , we obtain for each individual *i* and cluster *c* the estimates of mean  $\overline{V}^{(c)}$  and an  $(M, \overline{V}^{(c)})$ -nested principal convex component  $\hat{C}_M^{(c)}$  based on the matrices  $V_k$  with  $h_k^{(t)} = c$  for all  $k \neq i$ , leaving out the *i*-th matrix. Here, *M* is the value of dimension that was chosen in the initial clustering step. Given these estimates, we obtain the *i*-th predicted model for each cluster *c*,

$$\tilde{V}_{(i)}^{(c)} = \prod_{\hat{C}_M^{(c)}} V_i,$$

where  $\Pi_{\hat{C}_{M}^{(c)}}V_{i}$  is the *M*-dimensional representation of  $V_{i}$  defined as in (17). The *i*-th individual is classified into cluster  $h_{i}^{(t+1)}$  such that

$$h_i^{(t+1)} = \underset{c \in \{1, \dots, K\}}{\operatorname{arg\,min}} \| V_i - \tilde{V}_{(i)}^{(c)} \|_{\hat{\Sigma}_*}.$$

This step is performed for all *i*, which leads to an updated set of results  $\mathcal{H}^{(t+1)} = \{h_i^{(t+1)} : i = 1, ..., n\}$ . The updating procedure is iteratively implemented until no more data can be reclassified.

## References

- Abraham, C., P.-A. Cornillon, E. Matzner-Løber, and N. Molinari (2003). Unsupervised curve clustering using B-splines. Scandinavian journal of statistics 30(3), 581–595.
- Ambrosio, L., N. Gigli, and G. Savaré (2008). Gradient Flows: in Metric Spaces and in the Space of Probability Measures. Springer.
- Banerjee, A., I. S. Dhillon, J. Ghosh, S. Sra, and G. Ridgeway (2005). Clustering on the Unit Hypersphere using von Mises-Fisher Distributions. *Journal of Machine Learning Research* 6(9), 1345–1382.
- Banfield, J. D. and A. E. Raftery (1993). Model-based Gaussian and non-Gaussian clustering. Biometrics 49(3), 803–821.

- Bigot, J., R. Gouet, T. Klein, and A. López (2017). Geodesic PCA in the Wasserstein space by convex PCA. Annales de l'Institut Henri Poincaré, Probabilités et Statistiques 53(1), 1–26.
- Calò, D. G., A. Montanari, and C. Viroli (2014). A hierarchical modeling approach for clustering probability density functions. *Computational statistics & data analysis 71*, 79–91.
- Campbell, S. and T.-K. L. Wong (2022). Efficient Convex PCA with applications to Wasserstein geodesic PCA and ranked data. arXiv preprint arXiv:2211.02990.
- Cazelles, E., V. Seguy, J. Bigot, M. Cuturi, and N. Papadakis (2018). Geodesic PCA versus log-PCA of histograms in the Wasserstein space. SIAM Journal on Scientific Computing 40(2), B429–B456.
- Chazal, F., C. Levrard, and M. Royer (2021). Clustering of measures via mean measure quantization. *Electronic Journal of Statistics* 15, 2060–2104.
- Chen, Y., Z. Lin, and H.-G. Müller (2023). Wasserstein regression. *Journal of the American Statistical Association* 118(542), 869–882.
- Chiou, J.-M. and P.-L. Li (2007). Functional clustering and identifying substructures of longitudinal data. Journal of the Royal Statistical Society Series B: Statistical Methodology 69(4), 679–699.
- Chiou, J.-M. and P.-L. Li (2008). Correlation-based functional clustering via subspace projection. Journal of the American Statistical Association 103(484), 1684–1692.
- Del Barrio, E., J. A. Cuesta-Albertos, C. Matrán, and A. Mayo-Iscar (2019). Robust clustering tools based on optimal transportation. *Statistics and Computing* 29, 139–160.
- Delicado, P. (2011). Dimensionality reduction when data are density functions. *Computational Statistics & Data Analysis* 55(1), 401–420.
- Dhillon, I. S. and D. S. Modha (2001). Concept decompositions for large sparse text data using clustering. *Machine learning* 42, 143–175.
- Ghodrati, L. and V. M. Panaretos (2022). Distribution-on-distribution regression via optimal transport maps. *Biometrika* 109(4), 957–974.
- Hron, K., A. Menafoglio, M. Templ, K. Hruzová, and P. Filzmoser (2016). Simplicial principal component analysis for density functions in Bayes spaces. *Computational Statistics & Data Analysis 94*, 330–350.
- Hubert, L. and P. Arabie (1985). Comparing partitions. Journal of classification 2, 193–218.
- Jacques, J. and C. Preda (2013). Funclust: A curves clustering method using functional random variables density approximation. *Neurocomputing* 112, 164–171.

- James, G. M. and C. A. Sugar (2003). Clustering for sparsely sampled functional data. *Journal of the American Statistical Association* 98(462), 397–408.
- MacQueen, J. (1967). Some methods for classification and analysis of multivariate observations. In Proceedings of the fifth Berkeley symposium on mathematical statistics and probability, Volume 1, pp. 281–297. Oakland, CA, USA.
- Mardia, K. V., H. Wiechers, B. Eltzner, and S. F. Huckemann (2022). Principal component analysis and clustering on manifolds. *Journal of Multivariate Analysis 188*, 104862.
- Okano, R. and M. Imaizumi (2024). Distribution-on-distribution regression with Wasserstein metric: Multivariate Gaussian case. *Journal of Multivariate Analysis*, 105334.
- Panaretos, V. M. and Y. Zemel (2020). An Invitation to Statistics in Wasserstein Space. Springer Nature.
- Papayiannis, G., G. Domazakis, D. Drivaliaris, S. Koukoulas, A. Tsekrekos, and A. Yannacopoulos (2021). On clustering uncertain and structured data with Wasserstein barycenters and a geodesic criterion for the number of clusters. *Journal of Statistical Computation and Simulation 91*(13), 2569–2594.
- Petersen, A., C. Zhang, and P. Kokoszka (2022). Modeling probability density functions as data objects. *Econometrics and Statistics* 21, 159–178.
- Pollard, D. (1981). Strong consistency of k-means clustering. The annals of statistics, 135–140.
- Rand, W. M. (1971). Objective criteria for the evaluation of clustering methods. Journal of the American Statistical association 66 (336), 846–850.
- Rousseeuw, P. J. (1987). Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. *Journal of computational and applied mathematics* 20, 53–65.
- Serban, N. and L. Wasserman (2005). CATS: clustering after transformation and smoothing. Journal of the American Statistical Association 100(471), 990–999.
- Stuart, A. and K. Ord (2010). Kendall's advanced theory of statistics, distribution theory, Volume 1. John Wiley & Sons.
- Verdinelli, I. and L. Wasserman (2019). Hybrid Wasserstein distance and fast distribution clustering. *Electronic Journal of Statistics* 13(2), 5088–5119.
- Villani, C. (2008). Optimal Transport: Old and New, Volume 338. Springer Science & Business Media.

- Ward Jr, J. H. (1963). Hierarchical grouping to optimize an objective function. Journal of the American statistical association 58(301), 236–244.
- Zemel, Y. and V. M. Panaretos (2019). Fréchet means and Procrustes analysis in Wasserstein space. *Bernoulli* 25(2), 932–976.
- Zhang, C., P. Kokoszka, and A. Petersen (2022). Wasserstein autoregressive models for density time series. Journal of Time Series Analysis 43(1), 30–52.
- Zhu, C. and H.-G. Müller (2023). Autoregressive optimal transport models. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 85(3), 1012–1033.
- Zhuang, Y., X. Chen, and Y. Yang (2022). Wasserstein K-means for clustering probability distributions. Advances in Neural Information Processing Systems 35, 11382–11395.