

BLOCH FUNCTIONS WITH WILD BOUNDARY BEHAVIOUR IN \mathbb{C}^N

STÉPHANE CHARPENTIER, NICOLAS ESPOULLIER, RACHID ZAROUF

ABSTRACT. We prove the existence of functions f in the Bloch space of the unit ball \mathbb{B}_N of \mathbb{C}^N with the property that, given any measurable function φ on the unit sphere \mathbb{S}_N , there exists a sequence $(r_n)_n$, $r_n \in (0, 1)$, converging to 1, such that for every $w \in \mathbb{B}_N$,

$$f(r_n(\zeta - w) + w) \rightarrow \varphi(\zeta) \text{ as } n \rightarrow \infty, \text{ for almost every } \zeta \in \mathbb{S}_N.$$

The set of such functions is residual in the little Bloch space. A similar result is obtained for the Bloch space of the polydisc.

1. INTRODUCTION AND STATEMENTS OF THE RESULTS

Our purpose is to show the existence of functions in the Bloch space of the unit ball and of the polydisc of \mathbb{C}^N that carry a quite erratic non-tangential boundary behaviour.

Throughout the paper, we will denote by $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$,

$$\mathbb{B}_N := \{z = (z_1, \dots, z_N) \in \mathbb{C}^N : |z|^2 := |z_1|^2 + \dots + |z_N|^2 < 1\}$$

and $\mathbb{D}^N := \mathbb{D} \times \dots \times \mathbb{D}$ (with $N \geq 1$) the open unit disc of the complex plane \mathbb{C} , the open unit ball and the polydisc of \mathbb{C}^N , respectively. The notations \mathbb{T} , \mathbb{S}_N and \mathbb{T}^N will stand for the unit circle, the unit sphere and the distinguished boundary of \mathbb{D}^N . Without possible confusions, we shall indifferently denote by $|\cdot|$ the modulus of a complex number and the euclidean norm in \mathbb{C}^N .

Given a domain $D \subset \mathbb{C}^N$, we denote by $H(D)$ the Fréchet space of all functions holomorphic in \mathbb{D} , endowed with the topology of locally uniform convergence. For $f \in H(\mathbb{B}_N)$, the radial derivative of f is defined by

$$\mathcal{R}(f) = \sum_{k=1}^N z_k \frac{\partial f}{\partial z_k}.$$

The Bloch space $\mathcal{B}(\mathbb{B}_N)$ of \mathbb{B}_N consists in all functions f holomorphic on \mathbb{B}_N such that

$$\sup_{z \in \mathbb{B}_N} (1 - |z|^2) |\mathcal{R}(f)(z)| < \infty.$$

Endowed with the norm $\|f\|_{\mathcal{B}(\mathbb{B}_N)} := |f(0)| + \sup_{z \in \mathbb{B}_N} (1 - |z|^2) |\mathcal{R}(f)(z)|$, it is a Banach space. In particular, a function $f \in H(\mathbb{D})$ belongs to the Bloch space $\mathcal{B}(\mathbb{D})$ of the unit disc provided that $\sup_{|z| < 1} (1 - |z|^2) |f'(z)| < \infty$. We then define the Bloch space $\mathcal{B}(\mathbb{D}^N)$ of the polydisc as the subspace of $H(\mathbb{D}^N)$ consisting of those f for which

$$\sup_{z \in \mathbb{D}^N} \sum_{k=1}^N (1 - |z_k|^2) \left| \frac{\partial f}{\partial z_k}(z) \right| < \infty.$$

The quantity $\|f\|_{\mathcal{B}(\mathbb{D}^N)} := |f(0)| + \sup_{z \in \mathbb{D}^N} \sum_{k=1}^N (1 - |z_k|^2) \left| \frac{\partial f}{\partial z_k}(z) \right|$ is a norm that turns $\mathcal{B}(\mathbb{D}^N)$ into a Banach space.

2020 *Mathematics Subject Classification.* 32A40, 32A18, 32A10.

Key words and phrases. Bloch function, radial boundary behaviour, universal function.

The little Bloch spaces $\mathcal{B}_0(\mathbb{D})$, $\mathcal{B}_0(\mathbb{B}_N)$ and $\mathcal{B}_0(\mathbb{D}^N)$ of the unit disc, the unit ball and the polydisc, respectively, are defined as the closure of all polynomials in the corresponding Bloch space. For any $N \geq 1$, $\mathcal{B}_0(\mathbb{B}_N)$ coincides with the space of all functions f in $H(\mathbb{B}_N)$ such that

$$\lim_{|z| \rightarrow 1^-} (1 - |z|^2) |\mathcal{R}(f)(z)| = 0.$$

It is known that a similar assertion fails for $\mathcal{B}_0(\mathbb{D}^N)$, $N \geq 2$ [27]. When the context will be clear, we will refer to a function in one of the Bloch spaces introduced above as a *Bloch function*. For equivalent definitions of these spaces and their classical properties, we refer the reader to [29, Chapter 3] and [26, 27].

The search for Bloch functions with *irregular* radial boundary behaviour is a rather classical topic. In the unit disc, it is rather easy to build a function in $\mathcal{B}(\mathbb{D})$ with finite radial limit at no point of \mathbb{T} . Indeed, the lacunary series $\sum_k z^{2^k}$ is a Bloch function that has finite radial limits at no points of \mathbb{T} , by a tauberian theorem due to Hardy and Littlewood [16]. In passing, by Plessner's theorem, this implies that the image by such functions of almost all radii in \mathbb{D} is dense in \mathbb{C} . In the unit ball of \mathbb{C}^N , proving the existence of Bloch functions with radial limits at no point of \mathbb{S}_N is far more involved. Ryll and Wojtaszczyk introduced the so-called Ryll-Wojtaszczyk homogeneous polynomials in order to build a function in $\mathcal{B}(\mathbb{B}_N)$ with infinite radial limits at *almost every* point of \mathbb{S}_N [24]. Later, using modifications of those polynomials by Aleksandrov [1], Ullrich built a function $f \in \mathcal{B}(\mathbb{B}_N)$ such that for *every* $\zeta \in \mathbb{S}_N$, $|f(r\zeta)| \rightarrow \infty$, as $r \rightarrow 1$ [28]. Note that, by [4, Theorem 3], for any function f in $\mathcal{B}(\mathbb{B}_N)$ with finite radial limit at no point of \mathbb{S}_N , there exists a dense set of points ζ in \mathbb{S}_N such that $|f(r\zeta)| \rightarrow \infty$ as $r \rightarrow 1$.

The main contribution of this note is to prove that *quasi-all* functions in $\mathcal{B}_0(\mathbb{B}_N)$ or $\mathcal{B}_0(\mathbb{D}^N)$ (where quasi-all should be understood in the sense of Baire category theorem) carry an even more erratic non-tangential boundary behaviour, near *almost every* point of \mathbb{S}_N or \mathbb{T}^N . We recall that a subset of a separable Fréchet space is said residual if it contains a dense countable intersection of open sets. Let us denote by m_N the normalised Lebesgue measure on \mathbb{S}_N and by $\sigma_N := m_1 \times \dots \times m_1$ the normalised Lebesgue measure on \mathbb{T}^N .

Main theorem. *Let $(r_n)_n$ be a sequence of real numbers in $(0, 1)$, converging to 1 as $n \rightarrow \infty$.*

- (1) *There exists a residual subset of $\mathcal{B}_0(\mathbb{B}_N)$ consisting of functions f that satisfy the following property:*

Given any m_N -measurable function φ on \mathbb{S}_N , there exists a sequence $(n_k)_k$, $n_k \in \mathbb{N}$, $(\mathcal{Q}_{\mathbb{B}_N})$: such that for any $w \in \mathbb{B}_N$ and m_N -a.e. $\zeta \in \mathbb{S}_N$,

$$f(r_{n_k}(\zeta - w) + w) \rightarrow \varphi(\zeta) \text{ as } k \rightarrow \infty.$$

- (2) *There exists a residual subset of $\mathcal{B}_0(\mathbb{D}_N)$ consisting of functions f that satisfy the following property:*

Given any σ_N -measurable function φ on \mathbb{T}_N , there exists a sequence $(n_k)_k$, $n_k \in \mathbb{N}$, $(\mathcal{Q}_{\mathbb{D}_N})$: such that for any $w \in \mathbb{D}_N$ and σ_N -a.e. $\zeta \in \mathbb{T}_N$,

$$f(r_{n_k}(\zeta - w) + w) \rightarrow \varphi(\zeta) \text{ as } k \rightarrow \infty.$$

Similar results were already stated in other spaces of holomorphic functions. For example, Bayart proved (1) where the Bloch space $\mathcal{B}_0(\mathbb{B}_N)$ is replaced with the Fréchet space $H(\mathbb{B}_N)$, or by any *little growth space* $H_{w,0}^\infty$ of the ball [5]. Given a (continuous) weight function $w(t) : [0, 1) \rightarrow (0, +\infty)$ with $w(0^+) = 0$, let us recall that the growth space H_w^∞ consists of all functions f holomorphic in \mathbb{B}_N such that

$$\sup_{z \in \mathbb{B}_N} w(1 - |z|) |f(z)| < \infty.$$

The space $H_{w,0}^\infty$ can be defined as the closure of set of polynomials in H_w^∞ , and it can be checked that it coincides with the set of those f for which $w(1 - |z|)|f(z)| \rightarrow 0$ as $|z| \rightarrow 1$. Since there exist weights w for which H_w^∞ is contained in all Bergman spaces A^p , $p \geq 1$, the space $\bigcap_{p \geq 1} A^p$ contains a (lot of) function(s) satisfying the property ($\mathcal{Q}_{\mathbb{B}_N}$). In [10], an extension of this result to $H(D)$, where D is a pseudoconvex domain of \mathbb{C}^N , is obtained. We observe that in the space $H(\mathbb{D})$, it is known [9] that quasi-all functions satisfy a much stronger property. Finally, we mention that our theorem answers Question 5.7 in [11].

Let us now comment on the proof. The above statement typically falls within the theory of universality [6, 14], and it is now well-understood that most of results of this kind follow from a Baire category argument and eventually reduce to a suitable *simultaneous approximation*. The general idea is that, if we are given a (reasonable) separable Fréchet space X continuously embedded in $H(\mathbb{B}_N)$ (the strategy is the same for the polydisc), in order to build a function f in X with some prescribed boundary behaviour m_N -a.e. on \mathbb{S}_N , it is enough, roughly speaking, to find a function f in X that simultaneously approximates 0 in X and any given continuous function on some *large* subset of \mathbb{S}_N . We point out that simultaneous polynomial approximation in spaces of holomorphic functions is an independent active topic of research. We refer to the seminal thesis work of Khrushchev [18] and to the recent paper [22] (see also the references therein).

Following this strategy, in order to prove the assertion (1) in our main theorem, we will prove a slightly more precise version of the following simultaneous approximation lemma (see Lemma 2.1).

Lemma (Simultaneous approximation in the Bloch space of the ball). *For any $\varepsilon > 0$, any $g \in \mathcal{B}_0(\mathbb{B}_N)$, and any function φ continuous on \mathbb{S}_N , there exist a measurable set $E \subset \mathbb{S}_N$, with $m_N(E) \geq 1 - \varepsilon$, and a polynomial f such that*

$$\|f - g\|_{\mathcal{B}(\mathbb{B}_N)} \leq \varepsilon \quad \text{and} \quad \sup_{z \in E} |f(z) - \varphi(z)| \leq \varepsilon.$$

A similar statement will be proved for \mathbb{D}^N (see Lemma 2.5). For any holomorphic function space X on \mathbb{B}_N , that contains polynomials, if the set of all polynomials is dense in X and contained in the pointwise multipliers of X , then proving the above lemma reduces to the case where $g = 0$ and $\varphi = 1$. Now, a general approach to simultaneously approximate 0 in X and 1 on a *large* subset of \mathbb{S}_N consists in selecting the approximant f as equal to $P \circ I$, where P is a polynomial (usually given by Oka-Weyl theorem - or Runge's theorem in \mathbb{C}) that simultaneously approximate uniformly 1 on arbitrarily *large* proper compact subset of \mathbb{S}_N and 0 in a compact subset of \mathbb{B}_N , and where I is an inner mapping chosen so that $P \circ I$ is small in X norm, and so that it *transports* the approximation property of P on (another) *large* subset of \mathbb{S}_N . For example, in [5], the author made use of a similar simultaneous approximation result for $H_{w,0}^\infty$, that was due to Iordan [17]. Therein, the polynomial P is chosen using a Mergelyan's type result for the ball [15], and the inner mapping is $z\Phi^k(z)$, $z \in \mathbb{B}_N$, where ϕ is any inner function vanishing at 0, and the power k large enough (depending on the weight w).

This approach is followed by Limani [22, Section 4.2], who basically proves the above lemma for $N = 1$. The main difficulty consists in showing the existence of an inner function I such that $\|P \circ I\|_{\mathcal{B}}$ is arbitrarily small, which essentially reduces to the existence of an inner function I with arbitrarily small Bloch norm. It turns out that this can be achieved using a deep result of Aleksandrov, Anderson and Nicolau [2]. We mention that this allows Limani to prove the existence of so-called Menshov universal functions in the Bloch space of \mathbb{D} , answering a question posed in [7]. We also refer to [19, 20] for results of a similar nature in other Banach spaces of holomorphic functions on \mathbb{D} .

In order to prove our simultaneous approximation lemma for the Bloch space of the ball, a first difficulty is that one does not have a proper Mergelyan's theorem, and the second one is that inner functions are usually much more difficult to construct. However, a combination of simple geometric ideas, that served already in [10], and a partial generalisation of Aleksandrov-Anderson-Nicolau's result, due to Doubtsov [12], will allow us to obtain the main theorem.

Let us say a few words on the case of the polydisc, which slightly differs from the case of the ball ($N \geq 2$). Indeed, the only multipliers of $\mathcal{B}_0(\mathbb{D}^N)$ are the constant functions [3, Corollary 3.6] and thus we cannot directly follow the strategy described for \mathbb{B}_N . However, this can be overcome working in the unit disc and building our approximating polynomials as a product of polynomials of one variable.

The paper is organised as follows: the second section is dedicated to the results of simultaneous approximation in the Bloch space of the ball and the polydisc. The proof of our main theorem is presented in Section 3. A final section is devoted to commenting briefly on possible extensions and variations.

Notations. The notation $A \lesssim_P B$ shall be used when there exists a positive constant C , depending on the parameter P , such that $A \leq CB$. If the constant C does not depend on any parameter relevant in the context, we shall simply use the symbol \lesssim .

2. SIMULTANEOUS APPROXIMATION IN THE BLOCH SPACE

For any compact set E in \mathbb{C}^N , we will denote by $C(E)$ the Banach space of all continuous functions on E , endowed with the supremum norm, denoted by $\|\cdot\|_{\infty, E}$.

2.1. Simultaneous approximation in the Bloch space of the ball. In this part, our aim is to prove the following approximation lemma.

Lemma 2.1. *Let $\varepsilon > 0$, let L be a compact subset of \mathbb{B}_N , let $g \in \mathcal{B}_0(\mathbb{B}_N)$, and let $\varphi \in C(\mathbb{S}_N)$. There exist a set $E \subset \mathbb{S}_N$ with $m_N(E) \geq 1 - \varepsilon$ and a polynomial $f \in \mathcal{B}_0(\mathbb{B}_N)$, such that*

- (i) $\|f - g\|_{\mathcal{B}} < \varepsilon$;
- (ii) $\|f - \varphi\|_{\infty, E} < \varepsilon$.

Since the polynomials are dense in $\mathcal{B}_0(\mathbb{B}_N)$, it is enough to prove the lemma for $g = 0$, upon replacing φ by $\varphi - Q$, with Q close to g in $\mathcal{B}_0(\mathbb{B}_N)$. The proof is based on two results. The first one is geometric and follows immediately from [10, Lemma 3.6], for e.g.

Lemma 2.2. *Let $\varepsilon > 0$ and let U be a domain in \mathbb{C}^N such that $U \cap \mathbb{S}_N \neq \emptyset$. There exists a measurable subset E of $U \cap \mathbb{S}_N$ such that:*

- (i) $m_N(U \cap \mathbb{S}_N \setminus E) < \varepsilon$;
- (ii) for any compact set $L \subset \mathbb{B}_N$, the set $E \cup L$ is polynomially convex.

The second one deals with the existence of inner functions in the ball with a specific behaviour at the boundary. It is a combination of two results by Aleksandrov-Anderson-Nicolau [2] and Doubtsov [12].

Theorem 2.3. *Let $\eta > 0$ be fixed. There exists a non-constant inner function $I : \mathbb{B}_N \rightarrow \mathbb{C}$ such that*

$$\frac{(1 - |z|^2)|\mathcal{R}I(z)|}{1 - |I(z)|^2} < \eta, \quad z \in \mathbb{B}_N.$$

In particular, $\|I\|_{\mathcal{B}} < \eta$.

Proof. Let us fix $\eta > 0$. Let μ be a pluriharmonic probability measure such that the slices μ_ζ are uniformly symmetric and singular for all $\zeta \in \mathbb{S}_N$. Such measures exist, by Corollary 2.3 in [12]. Then define the non-constant singular inner function

$$\Theta(z) = \exp \left(- \int_{\mathbb{S}_N} \left(\frac{2}{(1 - \langle z, \zeta \rangle)^n} - 1 \right) d\mu(\zeta) \right).$$

Then, again by [12], one has

$$\lim_{|z| \rightarrow 1} \frac{(1 - |z|^2) |\mathcal{R}\Theta(z)|}{1 - |\Theta(z)|^2} = 0.$$

In particular, there exists a positive constant $M < \infty$ such that

$$\frac{(1 - |z|^2) |\mathcal{R}\Theta(z)|}{1 - |\Theta(z)|^2} \leq M, \quad z \in \mathbb{B}_N.$$

By Theorem 1 in [2] applied with $\phi(t) = \eta t / (2M)$, there exists an inner function $g : \mathbb{D} \rightarrow \mathbb{D}$ such that

$$\frac{(1 - |w|^2) |g'(w)|}{1 - |g(w)|^2} \leq \frac{\eta}{2M}, \quad w \in \mathbb{D}.$$

Let us set $I = g \circ \Theta$. Then I is a non-constant inner function on \mathbb{B}_N such that, for any $z \in \mathbb{B}_N$,

$$\frac{(1 - |z|^2) |\mathcal{R}I(z)|}{1 - |I(z)|^2} = \frac{(1 - |z|^2) |g'(\Theta(z))| |\mathcal{R}\Theta(z)|}{1 - |g(\Theta(z))|^2} \leq \frac{\eta}{2M} \frac{(1 - |z|^2) |\mathcal{R}\Theta(z)|}{1 - |\Theta(z)|^2} < \eta.$$

□

We are now ready to prove Lemma 2.1.

Proof of Lemma 2.1 with $g = 0$. Let $\varepsilon > 0$, let L be a compact subset of \mathbb{B}_N , and let $\varphi \in C(\mathbb{S}_N)$.

First, by uniform continuity of φ , there exist open domains U_1, \dots, U_l in \mathbb{C}^N , pairwise disjoint, with $U_j \cap \mathbb{S}_N \neq \emptyset$ for any j , and complex numbers c_1, \dots, c_l , such that

- (a) $m_N(\mathbb{S}_N \setminus (\overline{U_1} \cup \dots \cup \overline{U_l})) \leq \varepsilon/5$;
- (b) $|\varphi(z) - c_k| \leq \varepsilon/2$ for any $z \in \overline{U_k} \cap \mathbb{S}_N$ and any $k \in \{1, \dots, l\}$.

Now, applying Lemma 2.2 to each U_k , we choose a compact set $F \subset \mathbb{S}_N \cap (U_1 \cup \dots \cup U_l)$ with $\sigma(F) \geq 1 - \varepsilon/4$, such that $F \cup \{0\}$ is polynomially convex. Since the continuous function equal to 0 at 0 and to c_k on U_k , $k \in \{1, \dots, l\}$, extends holomorphically to an open neighborhood of $\{0\} \cup \overline{U_1} \cup \dots \cup \overline{U_l}$, the Oka-Weyl theorem (see [25], e.g.) ensures the existence of a polynomial Q such that

$$|Q(z) - \varphi(z)| < \varepsilon/2, \quad z \in F.$$

Let also P be a polynomial such that

$$(1) \quad |P(0)| < \frac{\varepsilon}{2\|Q\|_{\infty, \mathbb{B}_N}} \quad \text{and} \quad |P(z) - 1| < \frac{\varepsilon}{2\|Q\|_{\infty, \mathbb{B}_N}}, \quad z \in F.$$

Let now I be given by Theorem 2.3 for some $\eta > 0$ whose value will be fixed later. We define the mapping $J : z = (z_1, \dots, z_n) \mapsto I(z)z = (z_1 I(z), \dots, z_n I(z))$, $z \in \mathbb{B}_N$. It is clear that J is an inner mapping from \mathbb{B}_N to \mathbb{B}_N that vanishes at 0. Now, a simple calculation shows that

$$\mathcal{R}(P \circ J)(z) = (\mathcal{R}(I)(z) + I(z)) \sum_{i=1}^n z_i \frac{\partial P}{\partial z_i}(J(z)),$$

Hence, since P is a polynomial, we get for any $z \in \mathbb{B}_N$,

$$\begin{aligned} |\mathcal{R}(P \circ J)(z)| &\leq \sum_{i=1}^n \left| \frac{\partial P}{\partial z_i}(J(z)) \right| (|I(z)| + |\mathcal{R}(I)(z)|) \\ &\lesssim_P |I(z)| + |\mathcal{R}(I)(z)|. \end{aligned}$$

Moreover, as it is well-known,

$$|I(z)| \lesssim \|I\|_{\mathcal{B}(\mathbb{B}_N)} \log\left(\frac{1}{1-|z|^2}\right), \quad z \in \mathbb{B}_N.$$

Then for any $z \in \mathbb{B}_N$,

$$(1-|z|^2)|\mathcal{R}(P \circ J)(z)| \lesssim_P (1-|z|^2)|\mathcal{R}(I)(z)| + (1-|z|^2) \log\left(\frac{1}{1-|z|^2}\right) \|I\|_{\mathcal{B}(\mathbb{B}_N)}.$$

By Theorem 2.3, the function $P \circ J$ belongs to $\mathcal{B}_0(\mathbb{B}_N)$, and one has $\|I\|_{\mathcal{B}(\mathbb{B}_N)} < \eta$. Therefore

$$(1-|z|^2)|\mathcal{R}(P \circ J)(z)| \lesssim_P \eta \left(1 + (1-|z|^2) \log\left(\frac{1}{1-|z|^2}\right) \right), \quad z \in \mathbb{B}_N,$$

hence $(1-|z|^2)|\mathcal{R}(P \circ J)(z)| \lesssim_P \eta$, $z \in \mathbb{B}_N$.

We set $f = Q.(P \circ J)$. Using the fact that the multiplication by Q is a bounded linear operator on $\mathcal{B}_0(\mathbb{B}_N)$ [29, Theorem 3.21], and upon choosing η small enough, we deduce by the previous that

$$(1-|z|^2)|\mathcal{R}(f)(z)| < \varepsilon/2, \quad z \in \mathbb{B}_N.$$

Moreover, since J vanishes at 0, by (1), we have $|f(0)| < \varepsilon/2$. All in all we have proven the inequality $\|f\|_{\mathcal{B}(\mathbb{B}_N)} < \varepsilon$.

To finish the proof, observe that, by (1) and the definition of Q , for any $z \in J^{-1}(F) \cap F$, we have

$$|f(z) - \varphi(z)| \leq |Q(z)| |(P(J(z)) - 1)| + |Q(z) - \varphi(z)| < \varepsilon.$$

Since $J = Iz$, with I inner, Loewner's lemma (see [23, Example 14.2]) implies

$$m_N(J^{-1}(F)) = \sigma(F) \geq 1 - \varepsilon/4,$$

hence $m_N(J^{-1}(F) \cap F) \geq 1 - \varepsilon/2$. We set $E = J^{-1}(F) \cap F$.

To finish, it remains to explain why f can be chosen as a polynomial. To do so, let us recall that H^∞ is continuously embedded into \mathcal{B}_0 and that $\|f_r - f\|_{\mathcal{B}} \rightarrow 0$ as $r \rightarrow 1$ for any $f \in \mathcal{B}_0$, where f_r denote the dilate of f defined by $f_r(z) = f(rz)$, $z \in \mathbb{B}_N$ [29]. Therefore, we may choose $r \in (0, 1)$ close enough to 1 and approximate f_r by a polynomial uniformly on $\overline{\mathbb{B}_N}$, to get our conclusion. \square

For $\eta > 0$, we will denote by $I_\eta : \mathbb{B}_N \rightarrow \mathbb{B}_N$ any inner mapping given by Theorem 2.3, and let $J_\eta(z) = I(z)z$, $z \in \mathbb{B}_N$. The following corollary is a straightforward consequence of the proof of Lemma 2.1.

Corollary 2.4. *Let $d \geq 1$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$ and let $\varphi \in C(\mathbb{S}_n)$. There exist two polynomials P and Q such that, for any $\eta > 0$, there exists a compact set $E \subset \mathbb{S}_N$, with $m_N(E) \geq 1 - \tilde{\varepsilon}_2$, such that, if we set $f := Q(P \circ J_\eta)$, then the following assertions hold:*

- (1) $|f(0)| < \varepsilon_1$;
- (2) $\|f\|_{\mathcal{B}(\mathbb{B}_N)} < \eta + |f(0)|$;
- (3) $\|f - \varphi\|_{E, \infty} < \varepsilon_1$.

In the next paragraph, we shall use this corollary to obtain a simultaneous approximation lemma for the polydisc.

2.2. Simultaneous approximation in the Bloch space of the polydisc. We shall prove the following Bloch simultaneous approximation lemma for the polydisc.

Lemma 2.5. *Let $\varepsilon > 0$, let $g \in \mathcal{B}_0(\mathbb{D}^N)$ and let $\varphi \in C(\mathbb{T}^N)$. There exist a set $E \subset \mathbb{T}^N$ with $\sigma_N(E) \geq 1 - \varepsilon$ and a polynomial f , such that*

- (1) $\|f - g\|_{\mathcal{B}(\mathbb{D}^N)} \leq \varepsilon$;
- (2) $\|f - \varphi\|_{E, \infty} \leq \varepsilon$.

Proof. Since the polynomials are dense in $\mathcal{B}_0(\mathbb{D}^N)$, we may assume that g is identically equal to 0. Let ε , L , and φ be fixed as in the statement. By the Stone-Weierstrass Theorem, there exist $M > 0$ and functions $\varphi_{j,l} \in C(\mathbb{T})$, $j = 1, \dots, N$ and $1 \leq l \leq M$, such that

$$\|\varphi - (\varphi_{1,1} \dots \varphi_{N,1} + \dots + \varphi_{1,M} \dots \varphi_{N,M})\|_{\infty, \mathbb{T}^N} < \varepsilon.$$

We shall use Corollary 2.4 to approximate each product $\varphi_{1,l} \dots \varphi_{N,l}$, $1 \leq l \leq M$, by some suitable polynomial. Let us fix $l = 1$ and set

$$C := \prod_{j=1}^N (\|\varphi_{j,1}\|_{\infty, \mathbb{T}} + 1),$$

Let also $\tilde{\varepsilon} > 0$ whose value will be fixed later. We apply Corollary 2.4 to each $\varphi_{j,1}$, $1 \leq j \leq N$, with $d = 1$, $\varepsilon_1 = \min(1, \frac{\varepsilon}{2M}, \frac{\varepsilon}{CNM})$ and $\varepsilon_2 = \tilde{\varepsilon}$. This gives us N pairs of polynomials $(P_{j,1}, Q_{j,1})$, $1 \leq j \leq N$, such that if we fix an $\eta > 0$ satisfying

$$(2) \quad \eta < \frac{\varepsilon}{2NM \max_{1 \leq k \leq N} \sup_{(z_1, \dots, z_N) \in \mathbb{D}^N} \prod_{j \neq k} |f_{j,1}(z_j)|},$$

there exist compact sets $E_{j,1}$, $1 \leq j \leq N$, with $\sigma_1(E_{j,1}) \geq 1 - \tilde{\varepsilon}$, such that setting $f_{j,1} := Q_{j,1}(P_{j,1} \circ J_\eta)$, we have:

- (i) $|f_{j,1}(0)| < \frac{\varepsilon}{2M}$;
- (ii) $\|f_{j,1}\|_{\mathcal{B}(\mathbb{D}^N)} < \eta + |f_{j,1}(0)|$;
- (iii) $\|f_{j,1} - \varphi_{j,1}\|_{\infty, E_{j,1}} < \frac{\varepsilon}{CNM}$.

Furthermore, as explained at the very end of the proof of Lemma 2.1, we may and shall assume that each function $f_{j,1}$, $1 \leq j \leq N$, is a polynomial.

Now, we define the polynomial f_1 of N complex variables by

$$f_1(z_1, \dots, z_N) := f_{1,1}(z_1) \dots f_{N,1}(z_N),$$

and the compact subset $E_1 := E_{1,1} \times \dots \times E_{N,1}$ of \mathbb{T}^N . We shall check the following two assertions:

- (a) $\|f_1\|_{\mathcal{B}(\mathbb{D}^N)} < \varepsilon/M$;
- (b) $\|f_1 - \varphi_{1,1} \dots \varphi_{N,1}\|_{\infty, E(1)} < \varepsilon/M$.

First, notice that $|f_1(0)| < \varepsilon/(2M)$, by (i). Moreover, by (ii) and the choice of η , for any $(z_1, \dots, z_N) \in \mathbb{D}^N$,

$$\begin{aligned} \sup_{z \in \mathbb{D}^N} \sum_{k=1}^N (1 - |z_k|^2) \left| \frac{\partial f_1}{\partial z_k}(z) \right| &= \sup_{z \in \mathbb{D}^N} \sum_{k=1}^N (1 - |z_k|^2) |f'_{k,1}(z_k)| \prod_{j \neq k} |f_{j,1}(z_j)| \\ &\leq \eta N \max_{1 \leq k \leq N} \sup_{(z_1, \dots, z_N) \in \mathbb{D}^N} \prod_{j \neq k} |f_{j,1}(z_j)| < \frac{\varepsilon}{2M} \end{aligned}$$

which gives (a).

Next, by (iii), for any $z = (z_1, \dots, z_N) \in E_1$, we have

$$|\varphi_{1,1} \dots \varphi_{N,1}(z) - f_1(z)|$$

$$\begin{aligned}
&\leq \sum_{j=1}^N |f_{1,1}(z_1) \cdots f_{j-1,1}(z_{j-1})(f_{j,1}(z_j) - \varphi_{j,1}(z_j))\varphi_{j+1,1}(z_{j+1}) \cdots \varphi_{N,1}(z_N)| \\
&\leq \frac{\varepsilon}{CNM} \sum_{j=1}^N (\|\varphi_{1,1}\|_{\infty, \mathbb{T}} + \varepsilon) \cdots (\|\varphi_{j-1,1}\|_{\infty, \mathbb{T}} + \varepsilon) \|\varphi_{1,1}\|_{\infty, \mathbb{T}} \cdots \|\varphi_{j-1,1}\|_{\infty, \mathbb{T}} \\
&\leq \frac{\varepsilon}{CM} \prod_{j=1}^N (\|\varphi_{j,1}\|_{\infty, \mathbb{T}} + 1) = \frac{\varepsilon}{M},
\end{aligned}$$

whence assertion (b).

Further, we repeat the previous argument for the other products $\varphi_{1,l} \cdots \varphi_{N,l}$, $2 \leq l \leq M$. At the end, this provides us with M polynomials f_1, \dots, f_M and M subsets E_1, \dots, E_M such that, if we set $E := E_1 \cap \dots \cap E_M$ and $f := f_1 + \dots + f_M$, then

- (1) $\|f\|_{\mathcal{B}(\mathbb{D}^N)} < \varepsilon$;
- (2) $\|f - (\varphi_{1,1} \cdots \varphi_{N,1} + \dots + \varphi_{1,M} \cdots \varphi_{N,M})\|_{\infty, E} < \varepsilon$.

Moreover, upon choosing $\tilde{\varepsilon}$ small enough, we may assume $\sigma_N(E) \geq 1 - \varepsilon$. The lemma is proven, since $\|\varphi - (\varphi_{1,1} \cdots \varphi_{N,1} + \dots + \varphi_{1,M} \cdots \varphi_{N,M})\|_{\infty, \mathbb{T}^N} < \varepsilon$. \square

We note that Lemma 2.1 and Lemma 2.5 coincide when $N = 1$.

3. WILD BLOCH FUNCTIONS IN THE BALL AND THE POLYDISC

As it has been said in the introduction, showing the existence of functions with universal boundary behaviour reduces to proving simultaneous approximation. It turns out that this fact is a quite general one in the theory of universality [6]. For the seek of brevity, we shall present an abstract result that, combined with Lemma 2.1, will immediately imply our main results.

Let (X, d_X) be a Fréchet space and let Y be a separable topological vector space, metrizable with a translation-invariant metric d_Y . Let also $T_n : X \rightarrow Y$, $n \in \mathbb{N}$, be continuous linear maps. We say that the sequence $\mathcal{T} := (T_n)_n$ is universal if there exists $x \in X$ provided the set $\{T_n(x) : n \in \mathbb{N}\}$ is dense in Y . More generally, we will say that a family $\mathcal{T}_{i \in I}$ of sequences $(T_n^i)_n$, $i \in I$, of continuous linear maps from X to Y , is universal *uniformly with respect to* $i \in I$, if there exists $x \in X$ such that, for any $\varepsilon > 0$ and any $y \in Y$, there exists $n \in \mathbb{N}$ such that

$$\sup_{i \in I} d(T_n^i(x), y) < \varepsilon.$$

We denote by $\mathcal{U}(\mathcal{T})$ the set of all universal vectors for \mathcal{T} and by $\mathcal{U}(\mathcal{T}_{i \in I})$ the set of all vectors that are universal for $\mathcal{T}_{i \in I}$, uniformly with respect to $i \in I$. The following theorem is a very small improvement of a particular case of [6, Theorem 27],

Theorem 3.1. *Let $\mathcal{T}_{i \in I}$, be a family of sequences $(T_n^i)_n$, $i \in I$, of continuous linear maps from X to Y . Assume that there exists a dense subset G of X such that, for any $x \in G$, there exists $y \in Y$ such that $(T_n^i(x))_n$ converges to y , uniformly with respect to $i \in I$. If the following two conditions hold:*

- (1) *For every $\varepsilon > 0$ and every $y \in Y$, the set $\{x \in X : \sup_{i \in I} d_Y(T_n^i(x), y) < \varepsilon\}$ is open in X ;*
- (2) *For every $\varepsilon > 0$ and every $y \in Y$, there exists $n \in \mathbb{N}$ and $x \in X$, such that*

$$d_X(x, 0) < \varepsilon \quad \text{and} \quad \sup_{i \in I} d_Y(T_n^i(x), y) < \varepsilon,$$

then $\mathcal{U}(\mathcal{T}_{i \in I})$ is residual in X .

Proof. Even if it is very similar to some part of that of [6, Theorem 27], we include it here. We first prove that (2) implies the following apparently stronger property:

$$(2)' : \quad \text{For every } \varepsilon > 0, y \in Y, w \in X, \text{ there exist } n \in \mathbb{N} \text{ and } x \in X \text{ such that} \\ d_X(x, w) < \varepsilon \quad \text{and} \quad \sup_{i \in I} d_Y(T_n^i(x), y) < \varepsilon.$$

Let $\varepsilon > 0, y \in Y, w \in X$ be fixed. By assumption, there exists $w_0 \in G, v \in Y$ and $N \in \mathbb{N}$, such that $d_X(w, w_0) < \varepsilon/2$ and $\sup_{i \in I} d_Y(T_n^i(w_0), v) < \varepsilon/2$, for any $n \geq N$. By (2), there exists $n \in \mathbb{N}$ and $x_0 \in X$ such that $d_X(x_0, 0) < \varepsilon/2$ and $\sup_{i \in I} d_Y(T_n^i(x_0), y - v) < \varepsilon/2$, where n can be chosen with $n \geq N$. Then, setting $x = x_0 + w_0 \in X$, since the metrics are invariant by translation, we get $d_X(x, w) < \varepsilon$ and $\sup_{i \in I} d_Y(T_n^i(x), w) < \varepsilon$.

To conclude, let us fix a dense sequence $(y_l)_l$ in Y . We observe that

$$\mathcal{U}(\mathcal{T}_{i \in I}) \supset \bigcap_{k, l \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} \left\{ x \in X : \sup_{i \in I} d_Y(T_n^i(x), y_l) < \frac{1}{k} \right\}.$$

Now the proof follows by the Baire Category theorem. Indeed, every set $\bigcup_{n \in \mathbb{N}} \left\{ x \in X : \sup_{i \in I} d_Y(T_n^i(x), y_l) < \frac{1}{k} \right\}$ is open and dense in X , by (1) and (2)', respectively. \square

3.1. Wild Bloch functions in the ball. We shall see that Lemma 2.1 implies that (2) in Theorem 3.1 holds, for suitable choices of X, Y , and \mathcal{T} . In the whole subsection, let $X = \mathcal{B}_0(\mathbb{B}_N)$ be the little Bloch space of \mathbb{B}_N , and let Y be the space $L^0(\mathbb{S}_N)$ of all (equivalent classes of) functions m_N -measurable on \mathbb{S}_N (with identification of two functions that coincide off a set of m -measure 0), endowed with the translation-invariant metric

$$d(g, h) = \int_{\mathbb{S}_N} \min(1, |g - h|) dm_N.$$

We recall that a sequence $(g_n)_n$ in $L^0(\mathbb{S}_N)$ converges in measure to $g \in L^0(\mathbb{S}_N)$ if and only if every subsequence of $(g_n)_n$ has a subsequence that converges m_N -a.e. to g on \mathbb{S}_N . In particular, this implies that the set \mathcal{P} of all polynomials is dense in $L^0(\mathbb{S}_N)$.

Let us fix a sequence $(r_n)_n, r_n \in (0, 1)$, converging to 1. For $w \in \mathbb{B}_N$, let us define \mathcal{T}_w as the sequence $(T_n^w)_n$ where

$$T_n^w := \begin{cases} \mathcal{B}_0(\mathbb{B}_N) & \rightarrow L^0(\mathbb{S}_N) \\ f & \mapsto f(r_n(\cdot - w) + w). \end{cases}$$

Note that if $L \subset \mathbb{B}_N$ is compact then, by uniform continuity, the set

$$\{f \in \mathcal{B}_0(\mathbb{B}_N) : \sup_{w \in L} d(T_n^w(f)(z), f_0) < \varepsilon\}$$

is open, for every $f_0 \in \mathcal{B}_0(\mathbb{B}_N)$, every $\varepsilon > 0$ and every $n \in \mathbb{N}$. In particular, condition (1) of Theorem 3.1 is always satisfied in this setting. The same argument shows that for every $P \in \mathcal{P}$, the sequence $(T_n^w(P))_n$ converges to $P \in L^0(\mathbb{S}_N)$, uniformly with respect to $w \in L$.

In order to apply Theorem 3.1, it thus remains to check condition (2). The latter is a consequence of Lemma 2.1. Indeed, fix $\varepsilon > 0$ and $\varphi \in L^0(\mathbb{S}_N)$. By Lusin's theorem, every measurable function on \mathbb{S}_N can be approximated m_N -a.e. by continuous functions, so we can assume that $\varphi \in C(\mathbb{S}_N)$. We can now apply Lemma 2.1, with $g = 0$, to get a compact set $E \subset \mathbb{S}_N$, with $m_N(E) \geq 1 - \varepsilon/2$, and a polynomial $f \in \mathcal{B}_0(\mathbb{B}_N)$ such that

$$\|f\|_{\mathcal{B}_0(\mathbb{B}_N)} < \varepsilon \quad \text{and} \quad \|f - \varphi\|_{\infty, E} < \varepsilon/2.$$

Since f is uniformly continuous on $\overline{\mathbb{B}_N}$, the second inequality implies the existence of some $n \in \mathbb{N}$ such that

$$\sup_{w \in L} \|T_n^w(f) - \varphi\|_{\infty, E} < \varepsilon/2$$

To see that condition (2) of Theorem 3.1 holds, it suffices to observe that

$$\begin{aligned} \sup_{w \in L} d(T_n^w(f), \varphi) &= \sup_{w \in L} \int_{\mathbb{S}_N} \min(1, |T_n^i(f) - \varphi|) dm_N \\ &\leq \sup_{w \in L} \int_E \min(1, |T_n^w(f) - \varphi|) dm_N + m_N(\mathbb{S}_N \setminus E) < \varepsilon. \end{aligned}$$

By Theorem 3.1, we deduce that the set $\mathcal{U}(\mathcal{T}_{w \in L})$ is residual in $\mathcal{B}_0(\mathbb{B}_N)$. Now, considering an exhaustion $(L_n)_n$ of \mathbb{B}_N by compact sets, we obtain that the set

$$\bigcap_{n \in \mathbb{N}} \mathcal{U}(\mathcal{T}_{w \in L_n})$$

is residual in $\mathcal{B}_0(\mathbb{B}_N)$. Using again that a sequence $(g_n)_n$ in $L^0(\mathbb{S}_N)$ converges in measure to $g \in L^0(\mathbb{S}_N)$ if and only if every subsequence of $(g_n)_n$ has a subsequence that converges m -a.e. to g on \mathbb{S}_N , we deduce that the previous implies the assertion (1) of the main theorem, announced in the introduction. More precisely,

Theorem 3.2. *Let $(r_n)_n$ be a sequence of real numbers in $(0, 1)$, converging to 1. There exists a function f in $\mathcal{B}_0(\mathbb{B}_N)$ with the following property ($\mathcal{Q}_{\mathbb{B}_N}$): given any m -measurable function φ on \mathbb{S}_N , there exists a sequence $(n_k)_k$ of integers such that for any $w \in \mathbb{B}_N$ and m_N -a.e. $\zeta \in \mathbb{S}_N$,*

$$f(r_{n_k}(\zeta - w) + w) \rightarrow \varphi(\zeta) \text{ as } k \rightarrow \infty.$$

The set of such functions is residual in $\mathcal{B}_0(\mathbb{B}_N)$.

3.2. Wild Bloch functions in the polydisc. Similarly to the previous subsection, we will now derive from Lemma 2.5 and Theorem 3.1 a way to exhibit a large set of function in $\mathcal{B}_0(\mathbb{D}^N)$ with universal boundary behaviour. We set $X = \mathcal{B}_0(\mathbb{D}^N)$, $Y = L^0(\mathbb{T}^N)$, that is the space of all (equivalent classes of) functions σ_N -measurable on \mathbb{T}^N (with equivalence relation identifying two functions that coincide off a set of σ_N -measure 0), endowed with the translation-invariant metric

$$d(g, h) = \int_{\mathbb{T}^N} \min(1, |g - h|) d\sigma_N.$$

As recalled in Subsection 3.1, a sequence $(g_n)_n$ in $L^0(\mathbb{T}^N)$ converges in measure to $g \in L^0(\mathbb{T}^N)$ if and only if every subsequence of $(g_n)_n$ has a subsequence that converges σ_N -a.e. to g on \mathbb{T}^N . Thus the set \mathcal{P} of all polynomials is dense in $L^0(\mathbb{T}^N)$.

For a sequence $(r_n)_n$, $r_n \in (0, 1)$, converging to 1 and $w \in \mathbb{B}_N$, let \mathcal{T}_w be the sequence $(T_n^w)_n$ where

$$T_n^w := \begin{cases} \mathcal{B}_0(\mathbb{D}^N) & \rightarrow L^0(\mathbb{T}^N) \\ f & \mapsto f(r_n(\cdot - w) + w). \end{cases}$$

For $L \subset \mathbb{D}$ compact, let us write $L^N = L \times \dots \times L$ (N times). By uniform continuity, the set

$$\{f \in \mathcal{B}_0(\mathbb{D}^N) : \sup_{w \in L^N} d(T_n^w(f)(z), f_0) < \varepsilon\}$$

is open, for every $f_0 \in \mathcal{B}_0(\mathbb{D}^N)$, every $\varepsilon > 0$ and every $n \in \mathbb{N}$. Thus condition (1) of Theorem 3.1 is valid in this setting. For the same reason, for every $P \in \mathcal{P}$, the sequence $(T_n^w(P))_n$ converges to $P \in L^0(\mathbb{T}^N)$, uniformly with respect to $w \in L^N$.

Therefore we may apply Theorem 3.1, whenever we have proven that condition (2) holds true as well. This is where Lemma 2.5 comes into play. First, Lusin's theorem ensures that we may assume that $\varphi \in C(\mathbb{T}^N)$. Then, if $\varepsilon > 0$ and $\varphi \in C(\mathbb{T}^N)$ are given, Lemma 2.5

applied with $g = 0$, provides us with a compact set $E \subset \mathbb{T}^N$ and a polynomial $f \in \mathcal{B}_0(\mathbb{D}^N)$ such that $\sigma_N(E) \geq 1 - \varepsilon/2$ and

$$\|f\|_{\mathcal{B}_0(\mathbb{D}^N)} < \varepsilon \quad \text{and} \quad \|f - \varphi\|_{\infty, E} < \varepsilon/2.$$

By uniform continuity of f on $\overline{\mathbb{D}^N}$, we get from the second inequality the existence of some $n \in \mathbb{N}$ such that

$$\sup_{w \in L^N} \|T_n^w(f) - \varphi\|_{E, \infty} < \varepsilon/2.$$

Next, observe that

$$\begin{aligned} \sup_{w \in L^N} d(T_n^w(f), \varphi) &= \sup_{w \in L^N} \int_{\mathbb{T}^N} \min(1, |T_n^w(f) - \varphi|) d\sigma_N \\ &\leq \sup_{w \in L^N} \int_E \min(1, |T_n^w(f) - \varphi|) d\sigma_N + \sigma_N(\mathbb{T}^N \setminus E) < \varepsilon, \end{aligned}$$

which is condition (2) of Theorem 3.1.

Finally, Theorem 3.1 tells us that the set $\mathcal{U}(\mathcal{T}_{w \in L^N})$ is residual in $\mathcal{B}_0(\mathbb{D}^N)$. Let $(L_n)_n$ be an exhaustion of \mathbb{D} by compact sets. Then $(L_n^N)_n$ is an exhaustion by \mathbb{D}^N of compact sets and, by the Baire theorem, we conclude that

$$\bigcap_{n \in \mathbb{N}} \mathcal{U}(\mathcal{T}_{w \in L_n^N})$$

is residual in $\mathcal{B}_0(\mathbb{D}^N)$. We derive the following theorem.

Theorem 3.3. *Let $(r_n)_n$ be a sequence of real numbers in $(0, 1)$, converging to 1. There exists a function f in $\mathcal{B}_0(\mathbb{D}^N)$ satisfying the following property ($\mathcal{Q}_{\mathbb{D}^N}$): given any σ_N -measurable function φ on \mathbb{T}^N , there exists a sequence $(n_k)_k$ of integers such that for any $w \in \mathbb{D}^N$ and σ_N -a.e. $\zeta \in \mathbb{T}^N$,*

$$f(r_{n_k}(\zeta - w) + w) \rightarrow \varphi(\zeta) \quad \text{as } k \rightarrow \infty.$$

Moreover, the set of such functions is residual in $\mathcal{B}_0(\mathbb{D}^N)$.

Note that Theorem 3.2 and Theorem 3.3 gives the same statement when $N = 1$.

4. FURTHER EXTENSIONS AND VARIATIONS

4.1. Weighted Bloch spaces. Let $\omega : (0, 1) \rightarrow (0, +\infty)$ be a non-decreasing weight (that is $\omega(t) \rightarrow 0$ as $t \rightarrow 0^+$). The weighted Bloch space $\mathcal{B}_\omega(\mathbb{D})$ associated with ω is defined as the space of all functions f holomorphic in \mathbb{D} , such that

$$\|f\|_{\mathcal{B}_\omega(\mathbb{D})} := |f(0)| + \sup_{|z| < 1} \frac{1 - |z|^2}{\omega(1 - |z|)} |f'(z)| < \infty,$$

and the little weighted Bloch space $\mathcal{B}_{0,\omega}(\mathbb{D})$ as the closure of all polynomials in $\mathcal{B}_\omega(\mathbb{D})$ or, equivalently, as the closed subspace consisting of all f such that

$$\frac{1 - |z|}{\omega(1 - |z|)} |f'(z)| \rightarrow 0 \quad \text{as } |z| \rightarrow 1^-.$$

Further, we can also define the weighted Bloch space of \mathbb{D}^N as the Banach space of all functions $f \in H(\mathbb{D}^N)$ that satisfy

$$\|f\|_{\mathcal{B}_\omega(\mathbb{D}^N)} := |f(0)| + \sup_{|z| < 1} \sum_{k=1}^N \frac{1 - |z_k|^2}{\omega(1 - |z_k|^2)} \left| \frac{\partial f}{\partial z_k}(z) \right| < \infty,$$

and the little Bloch space $\mathcal{B}_{0,\omega}(\mathbb{D}^N)$ as the closure of the set of polynomials in $\mathcal{B}_\omega(\mathbb{D}^N)$.

Thus, replacing Theorem 2.3 by [2, Theorem 5.2], we may obtain a version of Corollary 2.4 for $N = 1$, with $\mathcal{B}_\omega(\mathbb{D})$ instead of $\mathcal{B}(\mathbb{D})$, and then obtain the same statement as Lemma 2.5 for $\mathcal{B}_{0,\omega}(\mathbb{D}^N)$. The same lines as those leading to Theorem 3.3 in Subsection 3.2 may finally lead to the following improvement.

Theorem 4.1. *Let ω be a non-decreasing weight such that, for some $\varepsilon > 0$, the function $\omega(t)/t^{1-\varepsilon}$ is decreasing. If*

$$\int_x^1 \frac{\omega(t)^2}{t} dt = +\infty,$$

then there exists a residual set of functions in $\mathcal{B}_{0,\omega}(\mathbb{D}^N)$ satisfying the property $(\mathcal{Q}_{\mathbb{D}^N})$ (see Theorem 3.3).

Interestingly, the integral condition in the previous result is sharp. Indeed, Proposition 1.1 in [13] shows that if $\int_x^1 \frac{\omega(t)^2}{t} dt < +\infty$, then every function in $\mathcal{B}_\omega(\mathbb{D})$ has radial limits almost everywhere. Since the latter result holds true for the unit ball \mathbb{B}_N as well, it is natural to wonder whether Theorem 4.1 also holds for $\mathcal{B}_0(\mathbb{B}_N)$, $N \geq 2$ (for some weight function ω defined on \mathbb{B}_N).

4.2. Cluster sets of Bloch functions along curves. For $\zeta \in \mathbb{S}_N$, let us say that a (continuous) path $\gamma : [0, 1) \rightarrow \mathbb{B}_N$ *terminates non-tangentially* at ζ if $\gamma(t) \rightarrow \zeta$ as $t \rightarrow 1$, and there exists a compact set $L \subset \mathbb{B}_N$ such that for any $t \in [0, 1)$, $\gamma(t)$ belongs to the set

$$\{r(\zeta - w) + w : w \in L, 0 \leq r \leq 1\}.$$

In Subsection 3.1, it is shown the existence of a residual set of functions f with the property that, given any $\varepsilon > 0$, any $\varphi \in C(\mathbb{S}_N)$ and any compact set $L \subset \mathbb{B}_N$, there exists $n \in \mathbb{N}$ and $1 - \varepsilon < r < 1$, such that for m_N -a.e. $\zeta \in \mathbb{S}_N$,

$$\sup_{w \in L} |f(r(z - w) + w) - \varphi(z)| < \varepsilon.$$

Since for any path γ terminating non-tangentially at some $\zeta \in \mathbb{S}_N$, one has $\gamma([0, 1)) \cap \{r(\zeta - w) + w : w \in L\} \neq \emptyset$, for any $r \in [0, 1)$ large enough, we immediately deduce the following corollary.

Corollary 4.2. *There exists a residual set in $\mathcal{B}_0(\mathbb{B}_N)$ consisting of functions f that satisfy the following property: there exists a set $E \subset \mathbb{S}_N$, with $m_N(E) = 1$, such that for any path γ terminating non-tangentially at a point of E , the cluster set of f along γ is maximal - that is the set $f(\gamma([0, 1)))$ is dense in \mathbb{C} .*

This result complements the fact that, for every function f in $\mathcal{B}(\mathbb{B}_N)$, there exists a dense set of points in \mathbb{S}_N at which f admits (finite or infinite) radial limit [4]. For $N = 1$, since every Bloch function is normal, it turns out that every radial limit of a non-constant function in $\mathcal{B}(\mathbb{D})$ is also a non-tangential limit [21]. In contrast, we recall that there exists a residual set in $H(\mathbb{B}_N)$ of functions having a maximal cluster set along any path terminating at a point of \mathbb{S}_N , with finite length [10]. For the case of the disc, the assumption *with finite length* can be dropped (see [8] or [9]).

A formulation of Corollary 4.2 for the polydisc could be naturally formulated. For the sake of brevity, we do not do it here and leave it to the interested reader.

REFERENCES

1. A. B. Aleksandrov, *Proper holomorphic mappings from the ball to the polydisk*, Dokl. Akad. Nauk SSSR **286** (1986), no. 1, 11–15.
2. A. B. Aleksandrov, J. M. Anderson, and A. Nicolau, *Inner functions, Bloch spaces and symmetric measures*, Proc. London Math. Soc. (3) **79** (1999), no. 2, 318–352.

3. R. F. Allen and F. Colonna, *Multiplication operators on the Bloch space of bounded homogeneous domains*, *Comput. Methods Funct. Theory* **9** (2009), no. 2, 679–693.
4. F. Bagemihl and W. Seidel, *Koebe arcs and Fatou points of normal functions*, *Comment. Math. Helv.* **36** (1961), 9–18.
5. F. Bayart, *Universal radial limits of holomorphic functions*, *Glas. Math. J.* **47** (2005), no. 2, 261–267.
6. F. Bayart, K.-G. Grosse-Erdmann, V. Nestoridis, and C. Papadimitropoulos, *Abstract theory of universal series and applications*, *Proc. London Math. Soc.* (3) **96** (2008), 417–463.
7. H. P. Beise and J. Müller, *Generic boundary behaviour of Taylor series in Hardy and Bergman spaces*, *Math. Z.* **284** (2016), 1185–1197.
8. A. Boivin, P. M. Gauthier, and P. V. Paramonov, *Approximation on closed sets by analytic or meromorphic solutions of elliptic equations and applications*, *Canad. J. Math* **54** (2002), no. 5, 945–969.
9. S. Charpentier, *Holomorphic functions with universal boundary behaviour*, *J. Approx. Theory* **254** (2020), 105391.
10. S. Charpentier and L. Kosiński, *Wild boundary behaviour of holomorphic functions in domains of \mathbb{C}^N* , *Indiana Univ. Math. J.* **70** (2021), no. 6, 2351–2367.
11. S. Charpentier, M. Manolaki, and K. Maronikolakis, *Abel universal functions: boundary behaviour and Taylor polynomials*, arXiv:2310.05611, 2023.
12. E. Doubtsov, *Little Bloch functions, symmetric pluriharmonic measures, and Zygmund’s dichotomy*, *J. Funct. Anal.* **170** (2000), no. 2, 286–306.
13. ———, *Weighted Bloch spaces and quadratic integrals*, *J. Math. Anal. Appl.* **412** (2014), no. 1, 269–276.
14. K.-G. Grosse-Erdmann, *Universal families and hypercyclic operators*, *Bull. Amer. Math. Soc.* **36** (1999), no. 3, 345–381.
15. M. Hakim and N. Sibony, *Boundary properties of holomorphic functions in the ball of \mathbb{C}^n* , *Math. Ann.* **276** (1987), no. 4, 549–555.
16. G. H. Hardy and J. E. Littlewood, *A Further Note on the Converse of Abel’s Theorem*, *Proc. London Math. Soc.* (2) **25** (1926), 219–236.
17. A. Iordan, *On the radial cluster sets for holomorphic functions in the unit ball of \mathbb{C}^n* , *Math. Z.* **202** (1989), no. 2, 199–206.
18. S.V. Khrushchev, *The problem of simultaneous approximation and of removal of the singularities of Cauchy type integrals*, *Trudy Mat. Inst. Steklov.* **130** (1978), 124–195, 223, *Spectral theory of functions and operators*.
19. ———, *A continuous function with universal Fourier series on a given closed set of Lebesgue measure zero*, *J. Approx. Theory* **252** (2020), 105361, 6.
20. ———, *Uniformly convergent fourier series with universal power parts on closed subsets of measure zero*, *J. Approx. Theory* **287** (2023), 105864.
21. O. Lehto and K. I. Virtanen, *Boundary behaviour and normal meromorphic functions*, *Acta Math.* **97** (1957), 47–65.
22. A. Limani, *Asymptotic polynomial approximation in the Bloch space*, arXiv:2403.08723, 2024.
23. W. Rudin, *New constructions of functions holomorphic in the unit ball of \mathbb{C}^n* , *CBMS Regional Conference Series in Mathematics*, vol. 63, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1986.
24. J. Ryll and P. Wojtaszczyk, *On homogeneous polynomials on a complex ball*, *Trans. Amer. Math. Soc.* **276** (1983), no. 1, 107–116.
25. E.L. Stout, *Polynomial convexity*, *Progress in Mathematics*, vol. 261, Birkhäuser Boston, Inc., Boston, MA, 2007.
26. R. M. Timoney, *Bloch functions in several complex variables. I*, *Bull. London Math. Soc.* **12** (1980), no. 4, 241–267.
27. ———, *Bloch functions in several complex variables. II*, *J. Reine Angew. Math.* **319** (1980), 1–22.
28. D. C. Ullrich, *A Bloch function in the ball with no radial limits*, *Bull. London Math. Soc.* **20** (1988), no. 4, 337–341.
29. K. Zhu, *Spaces of holomorphic functions in the unit ball*, *Graduate Texts in Mathematics*, vol. 226, Springer-Verlag, New York, 2005.

NICOLAS ESPOULLIER, AIX-MARSEILLE UNIV, CNRS, I2M, MARSEILLE, FRANCE

Email address: nicolas.espoullier@etu.univ-amu.fr

RACHID ZAROUF, AIX MARSEILLE UNIV, LAB ADEF, CAMPUS UNIV ST JEROME, 52 AVE ESCADRILLE
NORMANDIE, F-13013 MARSEILLE, FRANCE

Email address: rachid.zarouf@univ-amu.fr