

Multivariate root-n-consistent smoothing parameter free matching estimators and estimators of inverse density weighted expectations

Hajo Holzmann*

Fachbereich Mathematik und Informatik
 Philipps-Universität Marburg
 holzmann@mathematik.uni-marburg.de

Alexander Meister

Institut für Mathematik
 Universität Rostock
 alexander.meister@uni-rostock.de

July 12, 2024

Abstract

Expected values weighted by the inverse of a multivariate density or, equivalently, Lebesgue integrals of regression functions with multivariate regressors occur in various areas of applications, including estimating average treatment effects, nonparametric estimators in random coefficient regression models or deconvolution estimators in Berkson errors-in-variables models. The frequently used nearest-neighbor and matching estimators suffer from bias problems in multiple dimensions. By using polynomial least squares fits on each cell of the K^{th} -order Voronoi tessellation for sufficiently large K , we develop novel modifications of nearest-neighbor and matching estimators which again converge at the parametric \sqrt{n} -rate under mild smoothness assumptions on the unknown regression function and without any smoothness conditions on the unknown density of the covariates. We stress that in contrast to competing methods for correcting for the bias of matching estimators, our estimators do not involve nonparametric function estimators and in particular do not rely on sample-size dependent smoothing parameters. We complement the upper bounds with appropriate lower bounds derived from information-theoretic arguments, which show that some smoothness of the regression function is indeed required to achieve the parametric rate. Simulations illustrate the practical feasibility of the proposed methods.

Keywords: Average treatment effects; Berkson errors in variables models; bias correction; inverse density weighted expectations; matching estimators; random coefficients; Voronoi tessellation

1 Introduction

We consider estimation of inverse density weighted expectations, or, equivalently, of Lebesgue integrals of regression functions, with multivariate regressors. As we shall discuss in detail, such functionals occur in various areas of applications, including estimating average treatment effects (Abadie and Imbens, 2006, 2011), random coefficient regression models (Hoderlein et al., 2017), (Gaillac and Gautier, 2022), Berkson errors in variables models (Delaigle et al., 2006) or transfer learning under covariate shift (Kouw and Loog, 2019; Portier et al., 2023). Frequently used nearest-neighbor as well as matching estimators involve a bias of order $n^{-1/d}$ in d -dimensions (Abadie and Imbens,

*Corresponding author. Prof. Dr. Hajo Holzmann, Fachbereich Mathematik und Informatik, Philipps-Universität Marburg, Hans-Meerweinstr. 6, 35043 Marburg, Germany

2006). Therefore, methods for bias correction have been proposed in the treatment effect literature (Abadie and Imbens, 2011; Lin et al., 2023). These however involve nonparametric estimates and rates of convergence for regression functions and their derivatives, potentially under the strong smoothness assumptions.

In this paper we introduce simple modifications of nearest-neighbor and matching estimators for such functionals which again converge at the parametric \sqrt{n} -rate under mild smoothness assumptions on the unknown regression function. These novel estimators do not involve nonparametric function estimators and in particular do not rely on sample-size dependent smoothing parameters. Our upper bounds are complemented by a discussion of lower bounds which show that some smoothness is required for \sqrt{n} -consistent estimation to be possible.

More precisely, consider a d -vector of covariates Z supported on some compact and convex set $S \subseteq \mathbb{R}^d$ and having a strictly positive density f_Z on S . Further let $S^* \subseteq S$ be some given non-empty Borel set. We consider the following two closely related estimation problems, in which we focus on the multivariate setting $d \geq 2$.

First we intend to estimate from i.i.d. data (U_j, Z_j) , $j = 1, \dots, n$, distributed as (U, Z) , for some given real-valued measurable function $g(u, z)$ the functional

$$\Psi := \mathbb{E} \left[g(U, Z) \frac{1_{S^*}(Z)}{f_Z(Z)} \right] = \int_{S^*} G(z) dz, \quad (1.1)$$

where

$$G(z) := \mathbb{E} [g(U, Z) \mid Z = z] \quad (1.2)$$

is the regression function of Z on $g(U, Z)$. The issue in estimating Ψ in (1.1) in the representation as expected value is the unknown density f_Z which occurs in the denominator.

Second, from independent and identically distributed observations (Y_i, Z_i) , $i = 1, \dots, n$, where Y_i is real-valued, we consider estimation of the functional

$$\Phi = \mathbb{E} \left[Y 1_{S^*}(Z) \frac{f(Z)}{f_Z(Z)} \right]. \quad (1.3)$$

Here f is an additional Lebesgue density in \mathbb{R}^d . If f is known, by putting $g(y, z) = y f(z)$, this is a special case of the functional Ψ with Y taking the role of U . We shall devise an estimator in case of unknown f , from which an additional sample X_1, \dots, X_m , independent of the (Y_i, Z_i) , is supposed to be available.

In dimension $d = 1$, estimating Ψ in (1.1) has been studied in Lewbel and Schennach (2007). Based on the theory of order statistics and spacings they construct estimators which do not require nonparametric tuning parameters and converge at the parametric \sqrt{n} -rate. Further, they even show how to make their estimator asymptotically efficient by letting the order of the spacings diverge. However, as noted in Lewbel and Schennach (2007) their method does not easily extend to multiple dimensions. Estimating functionals of the form Ψ occurs frequently in applications which require $d \geq 2$. Examples include random coefficient regression models (Hoderlein et al., 2017; Gaillac and Gautier, 2022; Gautier and Kitamura, 2013) or series estimators which involve inner products of the regression function and basis functions with respect to the Lebesgue measure, such as nonparametric deconvolution estimators in the Berkson errors-in-variables models (Delaigle et al., 2006). We construct an estimator of Ψ in (1.1) which achieves the parametric \sqrt{n} -rate with neither dividing by a nonparametric

density estimate nor estimating the regression function G nonparametrically. Our methods involve least squares polynomial fits on cells of the K^{th} -order Voronoi tessellation, that is on the K -nearest neighbor partition induced by the sample Z_1, \dots, Z_n . We stress that our method does not have tuning parameters which must be chosen depending on the sample size and on the unknown smoothness of G , which would be typical of nonparametric methods. Compared to the analysis of the one-dimensional case $d = 1$ in Lewbel and Schennach (2007), higher-order smoothness assumptions on G are required to achieve the parametric rate, however, no smoothness conditions on the density f_Z of the covariates are needed in our analysis. Moreover, using information-theoretic arguments we show that some smoothness of the regression function G is indeed required in order to achieve the parametric rate. Furthermore we extend the methodology to construct a \sqrt{n} -consistent estimator for the functional Φ in (1.3) in case of unknown density f . We discuss how this estimator can be applied for average treatment effects, and show that for the special case of constant approximations it coincides with the classical matching estimator from the treatment effect literature. Using polynomial least squares fits of appropriate order depending on the dimension, we achieve a bias correction resulting in \sqrt{n} -consistent estimators under much milder smoothness assumptions than in Abadie and Imbens (2011) and without invoking nonparametric regression estimates. The functional Φ also occurs in transfer learning under covariate shift, when we have at our disposal labeled data from a source distribution but only unlabeled data from the target distribution. In such settings our method allows for a novel approach to estimating importance-weighted averages.

In our proofs we obtain results on the moments of the volume of higher-order Voronoi cells, as well as on properties of design matrices for multivariate polynomial regression under random design which might be of some independent interest. Let us briefly discuss some further methodologically related work. Devroye et al. (2017) study the asymptotic distribution for the volume of a first-order Voronoi cells, and Sharpnack (2023) analyzes matching estimators by relying on properties of first-order Voronoi cells. Kallus (2020) investigate weighted matching estimators with emphasis on weights from reproducing kernels, and Samworth (2012) provides comprehensive analysis of weighted nearest neighbor estimators. Finally, Hu et al. (2022) study nonparametric estimates of a function of bounded total variation by functions which are constant on elements of the first-order Voronoi tessellation.

The structure of the paper is as follows. In Section 2.1 we first motivate our method and the need for higher-order bias correction by considering and briefly analyzing a simple special case. Then in Section 2.2 we introduce our general method and state our main theoretical result for estimating Ψ , that is, the parametric rate of convergence under smoothness assumptions on the regression function G . This is extended in Section 2.3 to a \sqrt{n} -consistent estimator of Φ . Section 2.4 complements these upper bounds by providing lower bounds for estimating Ψ , which show that some smoothness of G (or f_Z) is generally required to estimate at the parametric rate. In Section 3 we discuss in detail how to apply the methodology to series estimators and in particular the Berkson errors-in-variables problem, to random coefficient regression models, to the estimation of average treatment effects and to estimating importance-weighted averages in transfer learning under covariate shift. Section 4 contains a simulation study, while Section 5 concludes. The main proofs of the upper bounds are given in Section 6, while additional technical arguments as well as further simulations results are deferred to the supplementary appendix.

We provide a brief overview of some important notation: the Euclidean norm is denoted by $\|\cdot\|$, the dimension by d ; and we write λ for the d -dimensional Lebesgue measure; for $\kappa = (\kappa_1, \dots, \kappa_d)^\top \in \mathbb{N}_0^d$ we denote $|\kappa| = \kappa_1 + \dots + \kappa_d$; and, for a function $f(z)$ in $z = (z_1, \dots, z_d)$, we denote its partial

derivatives by

$$\partial_\kappa f(z) = \frac{\partial^{|\kappa|}}{\partial z_1^{\kappa_1} \dots \partial z_d^{\kappa_d}} f(z).$$

2 Estimation method and rate of convergence

In this section we propose estimators for the functionals Ψ in (1.1) and Φ in (1.3). We start in Section 2.1 with a special case of our method which illustrates the need for higher-order bias correction in multiple dimensions. Smoothing-parameter free, \sqrt{n} -consistent estimators of the functionals Ψ and Φ are introduced in Sections 2.2 and 2.3, respectively. These are complemented in Section 2.4 with a discussion of lower bounds.

2.1 Motivation

To stress the need for a higher-order bias correction, let us start by considering and briefly analyzing a simple special case for the estimator of the functional Ψ in (1.1). Here we approximate the integral in (1.1) by a Riemann sum based on the first-order Voronoi tessellation of S^* . Consider the first order Voronoi cells

$$C_j := \{z \in S^* \mid \|z - Z_j\| < \|z - Z_k\|, \forall k \neq j\}.$$

An estimator of Ψ is then given by

$$\hat{\Psi} := \sum_{j=1}^n g(U_j, Z_j) \cdot \lambda(C_j), \quad (2.1)$$

where λ denotes the d -dimensional Lebesgue measure. Denoting by σ_Z the σ -field generated by Z_1, \dots, Z_n we obtain that

$$\mathbb{E}[\hat{\Psi} \mid \sigma_Z] = \sum_{j=1}^n G(Z_j) \cdot \lambda(C_j),$$

a discretized version of (1.1). Here G is the regression function defined in (1.2).

To show why this simple estimator does not achieve the parametric rate in dimensions $d > 3$ and even has issues for $d = 2$, consider the expected conditional variance and expected squared conditional bias decomposition of the mean squared error

$$\mathbb{E}[|\hat{\Psi} - \Psi|^2] = \mathbb{E}[\text{var}(\hat{\Psi} \mid \sigma_Z)] + \mathbb{E}[|\mathbb{E}[\hat{\Psi} \mid \sigma_Z] - \Psi|^2]. \quad (2.2)$$

Under the Assumptions 1 and 2 below, if the conditional variance $\text{var}(g(U_1, Z_1) \mid Z_1)$ is uniformly bounded, say by the constant $C_V > 0$, we have that

$$\mathbb{E}[\text{var}(\hat{\Psi} \mid \sigma_Z)] = \mathbb{E}\left[\sum_{j=1}^n \lambda^2(C_j) \cdot \text{var}(g(U_j, Z_j) \mid Z_j)\right] \leq n C_V \mathbb{E}[\lambda^2(C_1)] \lesssim n^{-1},$$

since the $\lambda^2(C_j)$ are identically distributed and Lemma 6.2 in Section 6.1.1 implies that $\mathbb{E}[\lambda^2(C_1)] \lesssim n^{-2}$. However, for the squared conditional bias, if G is Lipschitz continuous with constant $C_H > 0$ we

have that

$$\begin{aligned} \mathbb{E} \left[\left| \mathbb{E}[\hat{\Psi} | \sigma_Z] - \Psi \right|^2 \right] &= \mathbb{E} \left[\left| \sum_{j=1}^n \int_{C_j} (G(Z_j) - G(z)) \, dz \right|^2 \right] \\ &\leq C_H^2 \sum_{j,k=1}^n \iint_{(S^*)^2} \mathbb{E} \left[1_{C_j}(z) 1_{C_k}(z') \|z - Z_{(1)}(z)\| \|z' - Z_{(1)}(z')\| \right] \, dz \, dz' \lesssim n^{-2/d}, \end{aligned}$$

where $Z_{(1)}(z)$ denotes the first-nearest neighbor of z in Z_1, \dots, Z_n , and the final bound follows from Lemma 6.4 in the Section 6.1.2. Thus in dimensions $d \geq 3$ the order of the bias is slower than the parametric rate, and even for $d = 2$ it is not negligible compared to the variance.

To improve the rate of the bias, instead of the constant approximation to G on each Voronoi cell, we use polynomial approximation. In order to make this well-defined and computable we use higher-order Voronoi cells.

2.2 Bias corrected, \sqrt{n} -consistent estimation of Ψ

Our general method for estimating Ψ in (1.1) uses polynomial approximation of degree L and the K -th order Voronoi tessellation of S^* for sufficiently large integers $K, L \in \mathbb{N}$ depending on the dimension d . For each $J \subseteq \{1, \dots, n\}$ with $\#J = K$ we set

$$C(J) := \{z \in S^* \mid \|z - Z_j\| < \|z - Z_k\| \quad \forall j \in J, k \in \{1, \dots, n\} \setminus J\}. \quad (2.3)$$

Then for $z \in C(J)$ we let $\hat{G}(z)$ be the least squares polynomial of degree L in d variables when regressing Z_j on $g(U_j, Z_j)$ for indices $j \in J$. Let \mathcal{J}_K be the collection of all subsets of $\{1, \dots, n\}$ with exactly K elements. We set

$$\hat{\Psi} = \hat{\Psi}(L, K) := \sum_J \int_{C(J)} \hat{G}(z) \, dz. \quad (2.4)$$

The parameters K and L need not be chosen depending on the sample size n and hence do not take the role of smoothing parameters. They simply have to be sufficiently large depending on the dimension d , see Remark 1. Also note that $\hat{G}(z)$ is a weighted average of $g(U_j, Z_j)$ for those j corresponding to the K -nearest neighbors of z in Z_1, \dots, Z_n . Specifically, for $L = 0$ we obtain

$$\hat{G}(z) = \frac{1}{K} \sum_{j \in J} g(U_j, Z_j) \quad \text{if } z \in C(J), \quad \text{and} \quad \hat{\Psi} = \sum_{J \in \mathcal{J}_K} \left(\frac{1}{K} \sum_{j \in J} g(U_j, Z_j) \right) \lambda(C(J)). \quad (2.5)$$

For general L , more formally we introduce the notation

$$\begin{aligned} \mathcal{K} &:= \mathcal{K}(d, L) := \{\kappa \in \mathbb{N}_0^d \mid |\kappa| \leq L\}, \\ \xi_\kappa(z, w) &:= \prod_{k=1}^d (w_k - z_k)^{\kappa_k}, \quad \kappa \in \mathcal{K}, \quad w \in \mathbb{R}^d, \quad \xi(z, w) = (\xi_\kappa(z, w))_{\kappa \in \mathcal{K}}. \end{aligned}$$

Monomials of degree $\leq L$ are indexed by elements in \mathcal{K} , so that K^* is the number of free parameters of a polynomial in d variables of total degree $\leq L$. We consider $\xi_\kappa(z, w)$ as a monomial function in w , centered at z . Then $\xi(z, w)$ is a vector of functions in w of dimension K^* .

For $z \in S^*$ let $J(z)$ denote that $J \in \mathcal{J}_K$ which satisfies $z \in C(J)$, and note that Lebesgue-almost all $z \in S^*$ belong to some $C(J)$, since the boundaries of the $C(J)$ are sets of measure 0.

The least squares polynomial $\hat{G}(z)$ of degree L in d variables at z when regressing Z_j on $g(U_j, Z_j)$ for indices $j \in J$ is obtained as the first entry $\gamma_{(0, \dots, 0)}$ of the vector

$$\operatorname{argmin}_{\gamma \in \mathbb{R}^{\mathcal{K}}} \sum_{j \in J} (g(U_j, Z_j) - \gamma^\top \cdot \xi(z, Z_j))^2 = \mathcal{M}(z)^{-1} \hat{\mathcal{G}}(z),$$

where

$$\begin{aligned} \mathcal{M}(z) &:= \sum_{j \in J(z)} \xi(z, Z_j) \xi(z, Z_j)^\top = \left(\sum_{j \in J(z)} \xi_{\kappa + \kappa'}(z, Z_j) \right)_{\kappa, \kappa' \in \mathcal{K}}, \\ \hat{\mathcal{G}}(z) &:= \sum_{j \in J(z)} g(U_j, Z_j) \cdot \xi(z, Z_j), \end{aligned}$$

and the matrix $\mathcal{M}(z)$ is invertible and, hence, strictly positive definite almost surely for sufficiently large K , making the estimator $\hat{G}(z)$ uniquely defined, see Section 6.1.1. Thus, we can write

$$\hat{G}(z) := \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \hat{\mathcal{G}}(z) = \sum_{j \in J(z)} g(U_j, Z_j) \cdot \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) \quad (2.6)$$

with the unit row vector \mathbf{e}_0 whose component for $\kappa = \mathbf{0}$ is 1.

Our main result requires the following two assumptions.

Assumption 1 (Support and design density). The support S of Z is a compact and convex subset of $S \subseteq \mathbb{R}^d$. The diameter of S is called ρ'' . The distribution of Z has a Lebesgue-density f_Z that is bounded away from zero by some $\rho > 0$ on S and bounded from above by $\bar{\rho}$.

Assumption 2 (Exclusion of boundary effects). We assume that the Euclidean balls $B_{\rho'}(z)$ with the center z , for all $z \in S^*$, and the radius $\rho' > 0$ are included in S as a subset.

Let $B_{\rho'}(S^*) = \bigcup_{z \in S^*} B_{\rho'}(z)$ denote the open ρ' -neighborhood of S^* . We assume that the restriction of the regression function G to $B_{\rho'}(S^*)$ belongs to the Hölder class $\mathcal{G}(l, \beta, C_H)$ with parameters $l \in \mathbb{N}_0$, $0 < \beta \leq 1$, $C_H > 0$ of functions on $B_{\rho'}(S^*)$ defined as follows:

$$\begin{aligned} \mathcal{G}(l, \beta, C_H) &= \left\{ h : B_{\rho'}(S^*) \rightarrow \mathbb{R} \mid h \text{ has partial derivatives of order } \leq l, \right. \\ &\quad \text{for each } \kappa \in \mathbb{N}_0^d, \text{ with } |\kappa| = l \text{ we have that} \\ &\quad \left. \left| \partial_\kappa h(z) - \partial_\kappa h(z') \right| \leq C_H \|z - z'\|^\beta, \quad z, z' \in B_{\rho'}(S^*) \right\}. \end{aligned} \quad (2.7)$$

Furthermore we set

$$K^* = K^*(d, L) = \#\mathcal{K} = \sum_{l=0}^L \binom{d+l-1}{l}, \quad D = D(d, L) = \sum_{l=1}^L l \binom{d+l-1}{l}. \quad (2.8)$$

Theorem 1. Consider estimation of the functional Ψ in (1.1) in dimensions $d \geq 2$ under the Assumptions 1 and 2. Suppose that the regression function G in (1.2) belongs to the Hölder class $\mathcal{G}(l, \beta, C_H)$ in (2.7) with parameters $l \in \mathbb{N}_0$, $0 < \beta \leq 1$, $C_H > 0$. Moreover assume that the conditional variance $\operatorname{var}(g(U_1, Z_1) | Z_1)$ is uniformly bounded by a constant $C_V > 0$. In the estimator $\hat{\Psi}(L, K)$ in (2.4),

choose $L = l$. If $L = l = 0$ take any fixed any $K \geq 1$, while for $L \geq 1$ choose some $K \geq 2 + (2D+1)K^*$, with K^* and D defined in (2.8). Then for the mean conditional variance and the mean squared conditional bias given the σ -field σ_Z generated by Z_1, \dots, Z_n we have that

$$\sup_{G \in \mathcal{G}(l, \beta, C_H)} \mathbb{E} [\text{var}(\hat{\Psi}|\sigma_Z)] \leq C_V \cdot C \cdot n^{-1}, \quad \sup_{G \in \mathcal{G}(l, \beta, C_H)} \mathbb{E} [|\mathbb{E}[\hat{\Psi}|\sigma_Z] - \Psi|^2] \leq C_H^2 \cdot C \cdot n^{-\frac{2(l+\beta)}{d}}, \quad (2.9)$$

where the constant $C > 0$ only depends on $L, K, \rho, \rho', \rho'', \bar{\rho}, d$ and $\lambda(S^*)$. Thus, if $l + \beta \geq d/2$,

$$\sup_{G \in \mathcal{G}(l, \beta, C_H)} \mathbb{E} [|\hat{\Psi} - \Psi|^2] \lesssim n^{-1},$$

and the contribution of the squared bias is negligible if $l + \beta > d/2$.

The main steps of the proof are provided in Section 6.1, with some technical details being deferred to Section B in the supplementary appendix.

Remark 1 (Choice of L and K). For given dimension d and order of the polynomial approximation L , the theorem specifies a minimal choice $K \geq 2 + (2D + 1)K^*$, K^* in (2.8), under which the upper risk bounds are theoretically guaranteed. Specifically, the minimal value of K for $L = 1$ is $K = 2 + (2d + 1)(1 + d)$. For $d = 2$ the bias is negligible for $L = l = 1$ if $\beta > 0$, and are result requires $K = 17$. Similarly, for $d = 3$ the bias is negligible for $L = l = 1$ if $\beta > 1/2$, and we need $K = 30$ in our theoretical result. In the simulations, we illustrate that smaller values of K often seem to suffice.

Remark 2 (Higher moments of the volume of $C(J)$). For the derivation of upper bounds as in Theorem 1, the asymptotic properties of the d -dimensional Lebesgue measure $\lambda(C(J))$ of the Voronoi cells as $n \rightarrow \infty$ are essential. Since

$$\lambda(S^*) = \sum_{J \in \mathcal{J}_K} \lambda(C(J))$$

and the $C(J)$ are identically distributed we have that $\mathbb{E}[\lambda(C(J))] = \mathcal{O}(n^{-K})$. Hence $\lambda(C(J)) = \mathcal{O}_{\mathbb{P}}(n^{-K})$ which implies $\lambda(C(J))^\ell = \mathcal{O}_{\mathbb{P}}(n^{-\ell K})$. As a surprising phenomenon the higher moments of this term show an unusual behaviour since $\mathbb{E}[\lambda(C(J))^\ell] \simeq n^{-\ell - K + 1}$. The upper bound is provided in Lemma 6.2 for $\ell = 2$, and for the lower bound see Section B in the supplementary appendix.

2.3 Bias corrected matching estimators for Φ

Now let us turn to estimating the functional Φ in (1.3) in the setting where f is unknown. Suppose that we have an additional independent sample X_1, \dots, X_m , each X_j with density f , independent of $(Y_1, Z_1), \dots, (Y_n, Z_n)$. One possibility to estimate (1.3) is to use a nonparametric estimate \hat{f} of f from X_1, \dots, X_m , and then to employ (2.1) with the estimated function $\hat{g}(y, z) = y \hat{f}(z)$. Alternatively, we can avoid a nonparametric estimate of f and proceed as follows.

Consider the Voronoi cells as in (2.3), and for $J \subseteq \{1, \dots, n\}$ with $\#J = K$, that is $J \in \mathcal{J}_K$, and $z \in C(J)$ we let $\hat{G}(z)$ be the least squares polynomial fit of order L as in (2.6), with the observed Y_j replacing $g(U_j, Z_j)$. Then we set

$$\hat{\Phi} = \hat{\Phi}(L, K) = \frac{1}{m} \sum_{k=1}^m \sum_{J \in \mathcal{J}_K} 1(X_k \in C(J)) \hat{G}(X_k). \quad (2.10)$$

Consider the class of functions

$$\mathcal{G}(l, \beta, C_H, C_G) = \left\{ h : B_{\rho'}(S^*) \rightarrow \mathbb{R} \mid h \in \mathcal{G}(l, \beta, C_H), |h(z)| \leq C_G, \forall z \in B_{\rho'}(S^*) \right\}.$$

Theorem 2. Consider estimation of the functional Φ in (1.3) in dimensions $d \geq 2$ under the Assumptions 1 and 2. Moreover assume that the conditional variance $\text{var}(Y_1|Z_1)$ is uniformly bounded by a constant $C_V > 0$, and that the density f is bounded from above by C_f . In the estimator $\hat{\Phi}(L, K)$ in (2.10), choose $L = l$. If $L = l = 0$ take any fixed $K \geq 1$, while for $L \geq 1$ choose some $K \geq 2 + (2D + 1)K^*$, with K^* and D defined in (2.8). Then

$$\sup_{G \in \mathcal{G}(l, \beta, C_H, C_G)} \mathbb{E} [|\hat{\Phi} - \Phi|^2] \leq C \left((C_V + C_G^2) m^{-1} + C_V n^{-1} + C_H^2 n^{-\frac{2(l+\beta)}{d}} \right), \quad (2.11)$$

where the constant $C > 0$ only depends on $L, K, \rho, \rho', \rho'', \bar{\rho}, d, \lambda(S^*)$ and C_f . Thus, if $l + \beta \geq d/2$,

$$\sup_{G \in \mathcal{G}(l, \beta, C_H, C_G)} \mathbb{E} [|\hat{\Phi} - \Phi|^2] \lesssim n^{-1} + m^{-1},$$

and the contribution of the squared bias is negligible if $l + \beta > d/2$.

The proof is provided in Section 6.2.

Remark 3 (Matching estimators). For $L = 0$ the estimator in (2.10) corresponds to the matching estimator from the treatment effect literature (Abadie and Imbens, 2006; Imbens, 2004): For $k = 1, \dots, m$ let $\{Z_j \mid j \in J_k\}$, $J_k \subseteq \{1, \dots, n\}$ with $\#J_k = K$ be the set of K -nearest neighbors of X_k in Z_1, \dots, Z_n , and set $\hat{Y}_k = K^{-1} \sum_{j \in J_k} Y_j$. Then (2.10) can be written as

$$\hat{\Phi} = \frac{1}{m} \sum_{k=1}^m \sum_{J \in \mathcal{J}_K} 1(X_k \in C(J)) \frac{1}{K} \sum_{j \in J} Y_j = \hat{\Phi} = \frac{1}{m} \sum_{k=1}^m \hat{Y}_k.$$

See Section 3.3 for further discussion of estimating average treatment effects.

2.4 Discussion of lower bounds

In the oracle setting when f_Z is perfectly known the usual parametric rate is always attained by the estimator

$$\hat{\Psi}_{\text{oracle}} := \frac{1}{n} \sum_{j=1}^n g(U_j, Z_j) / f_Z(Z_j),$$

under the Assumptions 1 and 2 and the condition that $\text{var}(g(U_1, Z_1)|Z_1)$ is bounded, where no smoothness constraints are required.

On the other hand we can show, inspired by techniques from quadratic functional estimation (e.g. Tsybakov (2009)), that, under unknown f_Z , the parametric rate cannot be kept when no smoothness conditions are imposed. This occurs even in the univariate setting, as demonstrated in the following theorem.

Theorem 3. Consider the classical univariate nonparametric regression model $U_j = h(Z_j) + \varepsilon_j$, $j = 1, \dots, n$. Impose that both the design density f_Z is contained in the class $\mathcal{F} = \mathcal{F}(\rho, \bar{\rho})$ of all measurable functions on $[0, 1]$ which are bounded from above by $\bar{\rho} > 1$ and from below by $\rho > 0$; the regression function h lies in the class $\mathcal{H} = \mathcal{H}(\tilde{\rho})$ of all measurable functions bounded by $\tilde{\rho} > 0$; and the regression errors ε_j are standard Gaussian and independent of Z_j . Then,

$$\liminf_{n \rightarrow \infty} n^{1/2} \cdot \sup_{f_Z \in \mathcal{F}, h \in \mathcal{H}} \mathbb{E}_{f_Z, h} \left| \hat{H}_n - \int_{1/4}^{3/4} h(x) dx \right|^2 > 0,$$

for any sequence of estimators \hat{H}_n based on the data (U_j, Z_j) , $j = 1, \dots, n$.

The proof is given in Section A. Note that the regression model in Theorem 3 is included in the general framework by putting $g(u, z) = u$ and the conditional density $f_{U|Z}(u|z) = f_\varepsilon(u - h(z))$ when h denotes the regression function and f_ε stands for the standard normal density.

Remark 4 (Continuous design density and regression functions). We mention that the claim of Theorem 3 still holds when the classes \mathcal{F} and \mathcal{H} are changed to the corresponding sub-classes \mathcal{F}^0 and \mathcal{H}^0 which contain only the continuous functions in \mathcal{F} and \mathcal{H} , respectively, as \mathcal{F}^0 and \mathcal{H}^0 are dense subsets with respect to the L_1 -metric.

Remark 5 (Fixed design). Linear functional estimation in multivariate regression with fixed design is not included in the general framework of this work. In particular, Assumption 1 is violated. Still we mention that, in this setting, no estimator can attain faster convergence rates than $n^{-2(l+\beta)/d}$ as they occur as an upper bound in Theorem 1. This can be seen as follows. Assume that the design points Z_1, \dots, Z_n form an equidistant grid of some d -dimensional cube S so that $\|Z_j - Z_k\| \geq c \cdot n^{-1/d}$ for all $j \neq k$ and some fixed constant $c > 0$. Consider the competing regression functions $h_0 \equiv 0$ and $h_n(z) := \sum_{j=1}^n b^{l+\beta} K(\|z - Z_j\|/b)$ for some non-negative kernel function K which is infinitely often differentiable on \mathbb{R} , supported on $[0, 1]$ and satisfies $K(0) = 0$ as well as $K(1/2) > 0$, while $b = (c/2) \cdot n^{-1/d}$. Note that h_0 and h_n lead to the same distribution of the data, whereas the (non-squared) distance between $\int_{S^*} h_0$ and $\int_{S^*} h_n$ is bounded from below by the rate $n \cdot b^{l+\beta} \cdot b^d \asymp n^{-(l+\beta)/d}$ for any open non-void subset S^* of S . Note that these arguments cannot be applied to the random design case as, then, the competing regression functions must not depend on the design variables Z_1, \dots, Z_n . The rate $n^{-(l+\beta)/d}$ also occurs in Kohler (2014) for the L_1 -risk in noiseless regression.

3 Applications

In this section we present various applications of our methodology. In Section 3.1 we discuss series estimators in a given orthogonal system with respect to the Lebesgue measure, where for random design the Fourier coefficients are of the form (1.1). We give an explicit analysis in the nonparametric Berkson errors in variables problem (Delaigle et al., 2006), where our estimate is an ingredient of a Fourier series deconvolution estimator. In Section 3.2 we indicate how estimating functionals as in (1.1) arise in non- and semiparametric estimators in random coefficients regression models.

Sections 3.3 on estimating average treatment effects and 3.4 on transfer learning involve the functional (1.3) with unknown density f_0 , for which as in Section 2.3 an additional sample is available. For the average treatment effect on the treated and for the average treatment effect over a subset of the covariate space we achieve a bias correction resulting in \sqrt{n} -consistent estimators without involving nonparametric estimates of the regression functions and under much milder smoothness assumptions than in Abadie and Imbens (2011).

3.1 Series estimators and Berkson errors in variables models

Series estimators of a regression function $h(x) = \mathbb{E}[Y|X = x]$ in a given orthogonal basis (ϕ_n) or more generally a frame in $L_2(\mu)$ for some measure μ require estimation of

$$\langle h, \phi_n \rangle_\mu = \int h \phi_n \, d\mu.$$

Often in the mathematical analysis it is simply assumed that μ is the distribution of the covariates X , so that $\langle m, \phi_n \rangle = \mathbb{E}[Y \phi_n(X)]$ can directly be estimated by an average. But if μ is a given measure such as the Lebesgue measure λ on some compact domain determined by an a-priori choice of the (ϕ_n) as e.g. the Fourier basis, wavelets (Kerkycharian and Picard, 2004) or Gabor frames from time-frequency analysis (Dahlke et al., 2022), this may not be a realistic assumption for random X . Then, assuming X has Lebesgue density f_X , we rather have

$$\langle h, \phi_n \rangle_\lambda = \mathbb{E} \left[\frac{Y \phi_n(X)}{f_X(X)} \right], \quad (3.1)$$

leading to the estimation problem considered in (1.1) when putting $g(u, z) = u \cdot \phi_n(z)$.

Let us illustrate this further in the nonparametric Berkson errors-in-variables model, in which one observes (U, Z) according to

$$U = h(Z + \delta) + \epsilon, \quad (3.2)$$

where ϵ, δ and Z are independent and ϵ is centered and square integrable. When estimating h by deconvolution methods, Delaigle et al. (2006) use a local linear estimator of the calibrated regression function $g(z) = \mathbb{E}[U|Z = z]$ and numerically compute its Fourier coefficients. Meister (2009, Section 3.4.2) shows how to avoid nonparametric estimation of g in one dimension $d = 1$ by using spacings. Here we extend these results to higher dimensions $d \geq 2$ using our novel approach.

We choose the Lebesgue measure $\mu = \lambda$ and the Fourier basis

$$\phi_{\mathbf{j}}(\mathbf{x}) = \exp \left(i \sum_{k=1}^d j_k x_k \right), \quad \mathbf{x} = (x_1, \dots, x_d)$$

and the multi-index $\mathbf{j} = (j_1, \dots, j_d)$ where all components j_k , $k = 1, \dots, d$, are integers satisfying $|j_k| \leq J_n$ for some smoothing parameter J_n . Assume that δ has a known d -dimensional Lebesgue density and let \tilde{f}_δ denote the density of $-\delta$. We derive our result under the following set of assumptions.

Assumption 3. Impose that h and f_δ are supported on some fix compact subset of $(-\pi, \pi)^d$; that the support of the convolution $h * \tilde{f}_\delta$ is included in some fixed compact set S^* , which is contained in the interior of the compact and convex support S of f_Z as a subset; that f_Z is bounded away from zero on S (such that the Assumptions 1 and 2 are satisfied); that the known error density f_δ is ordinary smooth, i.e. its Fourier coefficients $f_\delta^{\text{ft}}(\mathbf{j}) := \langle f_\delta, \phi_{\mathbf{j}} \rangle_\lambda$ satisfy

$$c_\delta \cdot (1 + \|\mathbf{j}\|)^{-\gamma} \leq |f_\delta^{\text{ft}}(\mathbf{j})| \leq C_\delta \cdot (1 + \|\mathbf{j}\|)^{-\gamma},$$

for all \mathbf{j} and some constants $0 < c_\delta < C_\delta$ and $\gamma > d/2$, which guarantees that f_δ is squared integrable on the whole of \mathbb{R}^d .

Now apply our estimator $\hat{\Psi} = \hat{\Psi}_{\mathbf{j}}$ in (2.4) for

$$g(u, z) = g_{\mathbf{j}}(u, z) = u \cdot \phi_{\mathbf{j}}(z),$$

so that $G(z) = G_{\mathbf{j}}(z) = \phi_{\mathbf{j}}(z) \cdot [h * \tilde{f}_\delta](z)$, and S^* as in Assumption 3. Finally we consider the estimator

$$\hat{h} = (2\pi)^{-d} \sum_{\mathbf{j} \in \mathbf{J}^{(n)}} \hat{\Psi}_{\mathbf{j}} \cdot \phi_{-\mathbf{j}} / f_\delta^{\text{ft}}(-\mathbf{j})$$

of the regression function h , where $\mathbf{J}^{(n)} := \{-J_n, \dots, J_n\}^d$.

Theorem 4. Consider the Berkson errors in variables model (3.2) under Assumption 3 for $d \geq 2$. Furthermore suppose that the regression function h fulfills the Sobolev condition

$$\sum_{\mathbf{j}} |h^{\text{ft}}(\mathbf{j})|^2 (1 + \|\mathbf{j}\|^{2\alpha}) \leq C_\alpha, \quad (3.3)$$

for constants $\alpha, C_\alpha > 0$, where α is assumed to be sufficiently large such that

$$4(\alpha - 1)(\alpha + \gamma) \geq d(2\alpha + 2\gamma + d).$$

Then under the smoothing regime $J_n \asymp n^{1/(2\alpha+2\gamma+d)}$ we obtain uniformly over the class (3.3)

$$\mathbb{E} \left[\int_{[-\pi, \pi]^d} |\hat{h}(x) - h(x)|^2 dx \right] \lesssim n^{-2\alpha/(2\alpha+2\gamma+d)}. \quad (3.4)$$

The proof is provided in Section C.

3.2 Random coefficient regression

Random coefficient regression models form another field of application of our methods. We consider the linear random coefficient regression model

$$Y = \alpha + \beta^\top X, \quad (3.5)$$

where Y is the observed scalar response, X is a d -dimensional covariate vector, and $\beta \in \mathbb{R}^d$ and $\alpha \in \mathbb{R}$ are random regression coefficients, with (α, β) being independent of X . Note that α contains the deterministic intercept β_0 and the centered random errors ϵ , that is $\alpha = \beta_0 + \epsilon$. The joint density $f_{\alpha, \beta}$ is of interest and to be estimated. The model (3.5) contains heterogeneity in a linear equation, as is often required in econometric applications.

Gaillac and Gautier (2022) propose an estimator which allows for compactly supported covariates X . They use the identified relation

$$\mathbb{E}[\exp(itY)|X = x] = f_{\alpha, \beta}^{\text{ft}}(t, tx),$$

where $f_{\alpha, \beta}^{\text{ft}}$ is the $(d + 1)$ - dimensional Fourier transform of $f_{\alpha, \beta}$. Their estimator involves a series estimator of a partial Fourier transform of $f_{\alpha, \beta}$ in the first coordinate using the prolate spheroidal wave functions as basis. Thus, estimates of a form related to (3.1) from the previous section are required, to which our method can be applied.

Hoderlein et al. (2017) consider a triangular model with random coefficients. In case of additional exogenous covariates their semiparametric estimation method uses the simple estimator of the form (2.1) without higher-order bias correction to estimate the contrast function.

In their analysis of model (3.5), Hoderlein et al. (2010) normalize the covariates to the sphere (upper hemisphere) by setting $S = (1, X)/\|(1, X)\|$ and $U = Y/\|(1, X)\|$. Setting $(\alpha, \beta^\top) =: \bar{\beta}^\top$ this leads to the equation

$$U = S^\top \bar{\beta},$$

in which S and $\bar{\beta}^\top$ remain independent. A regularized approximation to $f_{\bar{\beta}}$ is then given by

$$\bar{f}_{\bar{\beta}}(b; h) = \int_{S^d} \mathbb{E} [K_{h,p}(S^\top b - U)|S = s] d\mu(s) \quad (3.6)$$

where μ is the Lebesgue measure on the d -dimensional sphere \mathcal{S}^d , $h > 0$ is a bandwidth parameter, and $K_{h,d}$ is a kernel tailored to the estimation problem. See Hoderlein et al. (2010) for the formula. (3.6) is of similar form to (1.1), but replacing the integral over Euclidean space with an integral over the sphere. Similar expressions involving integrals of regression function over the sphere occur in Dunker et al. (2019) or in the analysis of the binary choice model with random coefficients in Gautier and Kitamura (2013). An extension of our estimation approach to such situations would therefore be of some interest.

3.3 Average treatment effects

Let us discuss how the bias-corrected matching estimator in Section 2.3 can be applied to estimate average treatment effects in the potential outcome framework. Our methods complement the seminal work of Abadie and Imbens (2006, 2011); Imbens (2004) in several aspects. First we show that bias correction can be achieved without involving nonparametric estimators. Further we quantify a precise finite amount of smoothness of the regression functions to achieve \sqrt{n} -consistent estimation, without assuming infinite smoothness as in Abadie and Imbens (2011). Moreover, our results on lower bounds show that some amount of smoothness of design densities or regression functions is required for \sqrt{n} -consistent estimation to be possible. Recently, Lin et al. (2023) further analyzed the bias-corrected estimator of Abadie and Imbens (2011) in case of diverging K and under milder smoothness assumptions (Assumptions 4.4 and 4.5 in Lin et al. (2023)) which seem to be closer to but still stronger than those imposed in Theorem 2. Further, Lin et al. (2023) related to the recent literature on debiased/ doubly robust machine learning methods for treatment effects (Wang and Shah, 2024; Chernozhukov et al., 2018, 2024).

Consider the potential outcome framework: We have i.i.d. observations (Y_i, Z_i, D_i) , $i = 1, \dots, n$, distributed as the generic (Y, Z, D) . Here Y is the real-valued target quantity, Z a covariate vector, and D the binary treatment indicator. Suppose that the random variables $Y(0)$ and $Y(1)$ denote the potential outcomes, and that the observed outcome is generated as $Y = Y(D)$.

First consider estimation of the average treatment effect on the treated (ATET)

$$\tau^t = \mathbb{E} [Y(1) - Y(0) | D = 1].$$

Assuming that $Y(0)$ and D are independent given Z (conditional ignorability), and that $\mathbb{P}(p(Z) < 1) = 1$, where $p(Z) = \mathbb{P}(D = 1|Z)$ is the propensity score, the ATET is identified as (Chernozhukov et al., 2024, Section 5.C)

$$\tau^t = \mathbb{E} [Y|D = 1] - \frac{1}{\pi} \mathbb{E} \left[\mathbf{1}(D = 0) Y \frac{p(Z)}{1 - p(Z)} \right] := \tau_1^t - \tau_0^t,$$

where $\pi = \mathbb{P}(D = 1)$. Let $n_1 = \#\{i | D_i = 1\}$, $n_0 = n - n_1$. We estimate τ_1^t by the simple average $\hat{\tau}_1^t = n_1^{-1} \sum_{i=1}^n Y_i \mathbf{1}(D_i = 1)$. Next assume that the conditional distribution of Z given $D = i$ has Lebesgue density f_i , $i = 0, 1$. Then we can write

$$\tau_0^t = \mathbb{E} \left[Y \frac{f_1(Z)}{f_0(Z)} | D = 0 \right].$$

We estimate this parameter by $\hat{\tau}_0^t$ with the estimator (2.10), using those n_0 -observations (Y_i, Z_i) for which $D_i = 0$, and as sample from f_1 (which has the role of f in (1.3)) we take those Z_i for which $D_i = 1$. Finally set

$$\hat{\tau}^t = \hat{\tau}_1^t - \hat{\tau}_0^t.$$

Assumption 4. The density f_0 satisfies Assumption 1 for f_Z . Further, the support \mathcal{S}^* of f_1 is contained in the support \mathcal{S} as in Assumption 2. Moreover, f_1 bounded, the conditional variance $\text{var}(Y|Z = z, D = 0)$ is uniformly bounded.

Under Assumption 4, if the regression function $G_0(z) = \mathbb{E}[Y|Z = z, D = 0]$ is contained in $\mathcal{G}(l, \beta, C_H, C_G)$, when choosing K and L as in Theorem 2 in $\hat{\tau}_0^t$, we obtain, conditionally on the treatment variables, that

$$\mathbb{E} [(\hat{\tau}^t - \tau^t)^2 | \sigma_D] \lesssim n_1^{-1} + n_0^{-1} + n_0^{-\frac{2(l+\beta)}{d}},$$

where $\sigma_D = \sigma\{D_1, \dots, D_n\}$. If $l + \beta \geq d/2$ we obtain a bound of order $n_1^{-1} + n_0^{-1}$. If n_0 is large compared to n_1 , $n_0^{-2(l+\beta)/d}$ might be small compared to n_1 even for lower smoothness of G_0 . In an asymptotic analysis, this would require the distribution of D to depend on n with $\mathbb{P}(D = 1)$ tending to zero, see Assumption 3' in Abadie and Imbens (2006).

Now let us turn briefly to the average treatment effect (ATE). Here we require conditional ignorability for both $Y(1)$ and $Y(0)$, that is $Y(d)$ and D are independent given Z , $d = 0, 1$, and that $\mathbb{P}(0 < p(Z) < 1) = 1$. Let \mathcal{S}^* be a subset of the support of the covariate Z . Then the average treatment effect over \mathcal{S}^* is

$$\tau_{\mathcal{S}^*} = \mathbb{E} [Y(1) \mathbf{1}(Z \in \mathcal{S}^*)] - \mathbb{E} [Y(0) \mathbf{1}(Z \in \mathcal{S}^*)]$$

The average potential outcome of $Y(1)$ over \mathcal{S}^* is identified as

$$\begin{aligned} \mathbb{E} [Y(1) \mathbf{1}(Z \in \mathcal{S}^*)] &= \mathbb{E} \left[Y \mathbf{1}(D = 1) \frac{\mathbf{1}(Z \in \mathcal{S}^*)}{p(Z)} \right] \\ &= \mathbb{E} [Y \mathbf{1}(D = 1) \mathbf{1}(Z \in \mathcal{S}^*)] + (1 - \pi) \mathbb{E} \left[Y \frac{f_0(X)}{f_1(X)} \mathbf{1}(Z \in \mathcal{S}^*) | D = 1 \right], \end{aligned} \quad (3.7)$$

with a similar expression for $\mathbb{E} [Y(0) \mathbf{1}(Z \in \mathcal{S}^*)]$. As for ATET we can apply the estimator (2.10) to the second expression and under Assumption 4 for both f_0 and f_1 obtain the \sqrt{n} - rate of convergence.

3.4 Reweighting in transfer learning under covariate shift

In the transfer learning problem for classification, we have the target problem from a distribution (X, Y) as well as the source problem (X^S, Y^S) . Here Y and Y^S are binary variables, which are to predicted from the covariate Z respectively Z^S . Interest focuses on the classification problem for (X, Y) , but labeled data are mainly or only available from the source problem (Z_i^S, Y_i^S) , $i = 1, \dots, n_S$, together with unlabeled data Z_i , $i = 1, \dots, n$ from the target sample, that is with distribution P_Z . Under covariate shift it is assumed that the conditional distributions $P_{Y|X}$ and $P_{Y^S|X^S}$ are equal, and only the marginal distributions of covariates, P_X and P_{X^S} differ. See Kouw and Loog (2019); Portier et al. (2023); Sugiyama et al. (2007)

If $\ell(\hat{y}, y)$ is a loss function, the goal would be to train a classifier h which minimizes the expected loss under the target distribution $\mathbb{E}[\ell(h(X), Y)]$, which is however not directly accessible due to insufficient trained data from the target classification problem. Suppose that the distributions P_X and P_{X^S} have Lebesgue densities f_X and f_{X^S} , respectively. Then using the equality of the conditional distributions $P_{Y|X}$ and $P_{Y^S|X^S}$ we have that the average loss for the target problem can be computed as a importance-weighted average of the source problem,

$$\mathbb{E}[\ell(h(X), Y)] = \mathbb{E} \left[\frac{f_X(X^S)}{f_{X^S}(X^S)} \ell(h(X^S), Y^S) \right], \quad (3.8)$$

where the second expected value is over the source distribution, for which a labeled sample is available. (3.8) is of the form (1.3), and we can apply the matching estimator (2.10).

4 Simulations

In a brief simulation study we illustrate the practical feasibility of our methods. To numerically compute the estimator $\hat{\Psi}$ in (2.4), we simulate x_1^*, \dots, x_m^* from the uniform distribution on S^* for large m , and then effectively compute the estimator $\hat{\Phi}$ in (2.10) with $f = 1$ and the x_j^* taking the role of observed X_j 's, and normalize by $\lambda(S^*)$. Fast implementations to compute the K -nearest neighbors in Z_1, \dots, Z_n for a given set x_1^*, \dots, x_m^* of m points are available, also for moderately high values of K . We use the function `kNN` from the R-library `dbscan`, consider regressors in $d = 2$ and $d = 3$, and focus on the orders $L = 0$ and $L = 1$ for polynomial approximation. Computing the estimator once for $n = 1000$ and $m = 10000$ takes about 1 second on a computer with i7-10700 CPU, 2.90 GHz and 32GB RAM.

In a simulation in $d = 2$, we choose the coordinates $Z_{i,1}$ and $Z_{i,2}$ of the regressors as independent and Beta-distributed with parameters $\alpha = \beta = 3$ and $S^* = [0.2, 0.8]^2$, and as regression function we take

$$f_1(z_1, z_2) = 5 z_1 (z_2^2 + 1) + 10 \cos(z_1/z_2).$$

The DGP then is $Y_i = f_1(Z_{i,1}, Z_{i,2}) + 0.1 \varepsilon_i$, with ε_i independent standard normal. We use sample sizes $n \in \{100, 1000\}$, $m = 10000$ for the generated sample x_1^*, \dots, x_m^* and $N = 1000$ repetitions in each scenario. We compute the true value of the functional by numerical integration and obtain $\Psi_1 = 2.4981$. In Table 1 we report the values of the empirical bias, standard deviation and root mean squared error over the $N = 1000$ repetitions, all scaled by \sqrt{n} . Overall, while the reduction in MSE from $L = 0$ to $L = 1$ is not that large in this setting, there seems to be some reduction in asymptotic bias when using the linear approximation over Voronoi cells.

The method does not seem to depend sensitively on the choice of K for $L = 1$ for $n = 1000$, while for $n = 100$ a small choice of $K = 5$ performs best in this simulation setting. In the supplementary appendix, Section D we present another setting in $d = 2$ where higher values of K are required for optimal performance.

For a simulation with $d = 3$ we again choose coordinates $Z_{i,1}, Z_{i,2}$ and $Z_{i,3}$ of regressors as independent and Beta-distributed with parameters $\alpha = \beta = 3$, and let $S^* = [0.2, 0.8]^3$. The regression function is chosen as

$$f_2(z_1, z_2, z_3) = \exp(2 \cos(7 z_1) \sin(7 z_2)) \cdot (4 - 8(z_3 - 0.5)^2),$$

and as DGP we take $Y_i = f_2(Z_{i,1}, Z_{i,2}, Z_{i,3}) + 0.4 \varepsilon_i$, with ε_i independent standard normally distributed. Again we use $N = 1000$ repetitions in each scenario, and sample sizes $n \in \{100, 1000\}$ and $m = 10000$ for the generated sample x_1^*, \dots, x_m^* . The results are displayed in Table 2. The true value in this setting is $\Psi_3 = 1.4176$. While, for the smaller sample size $n = 100$, using $L = 0$ is preferable, for $n = 1000$ using $L = 1$ leads to a substantial reduction in the bias.

5 Concluding remarks

In contrast to other proposals in literature our estimators for expected values weighted by the inverse of a multivariate density do not rely on nonparametric function estimators and therefore do not require

$n = 100$	$L = 0$	K	1	3	4	6	8	10
		\sqrt{n} ·BIAS	0.1744	0.3074	0.3703	0.4858	0.5875	0.6814
		\sqrt{n} ·STDV	0.3498	0.4536	0.4976	0.5740	0.6388	0.6934
		\sqrt{n} ·RMSE	0.3908	0.5479	0.6202	0.7520	0.8679	0.9722
	$L = 1$	K	4	5	6	8	10	12
		\sqrt{n} ·BIAS	0.0188	0.0136	0.0029	-0.0150	-0.0345	-0.0606
		\sqrt{n} ·STDV	0.4072	0.3862	0.4075	0.4534	0.4992	0.5420
		\sqrt{n} ·RMSE	0.4076	0.3865	0.4075	0.4536	0.5004	0.5454
$n = 1000$	$L = 0$	K	1	3	4	6	8	10
		\sqrt{n} ·BIAS	0.0541	0.1064	0.1336	0.1872	0.2397	0.2933
		\sqrt{n} ·STDV	0.5869	0.5843	0.5841	0.5876	0.5920	0.5960
		\sqrt{n} ·RMSE	0.5894	0.5939	0.5992	0.6167	0.6387	0.6643
	$L = 1$	K	4	5	6	8	10	12
		\sqrt{n} ·BIAS	0.0203	0.0250	0.0314	0.0451	0.0593	0.0727
		\sqrt{n} ·STDV	0.5856	0.5797	0.5792	0.5785	0.5783	0.5785
		\sqrt{n} ·RMSE	0.5859	0.5802	0.5800	0.5802	0.5813	0.5831

Table 1: Simulation scenario with f_1 in $d = 2$

$n = 100$	$L = 0$	K	1	3	6	8	10	12
		\sqrt{n} ·BIAS	0.3265	0.5211	0.7081	0.7912	0.8535	0.9004
		\sqrt{n} ·STDV	0.7086	0.8674	1.0261	1.1015	1.1625	1.2141
		\sqrt{n} ·RMSE	0.7802	1.0119	1.2467	1.3562	1.4421	1.5116
	$L = 1$	K	5	6	8	10	12	14
		\sqrt{n} ·BIAS	-0.4611	-0.5416	-0.6972	-0.8296	-0.9426	-1.0446
		\sqrt{n} ·STDV	0.7106	0.6545	0.6654	0.7009	0.7417	0.7827
		\sqrt{n} ·RMSE	0.8471	0.8495	0.9637	1.0861	1.1995	1.3053
$n = 1000$	$L = 0$	K	1	3	6	8	10	12
		\sqrt{n} ·BIAS	0.2910	0.4584	0.6606	0.7731	0.8766	0.9726
		\sqrt{n} ·STDV	0.5137	0.5482	0.6058	0.6386	0.6715	0.7009
		\sqrt{n} ·RMSE	0.5904	0.7146	0.8963	1.0028	1.1043	1.1988
	$L = 1$	K	5	6	8	10	12	14
		\sqrt{n} ·BIAS	-0.1133	-0.1336	-0.1844	-0.2320	-0.2798	-0.3284
		\sqrt{n} ·STDV	0.5013	0.4336	0.4330	0.4338	0.4360	0.4382
		\sqrt{n} ·RMSE	0.5139	0.4537	0.4707	0.4919	0.5180	0.5476

Table 2: Simulation scenario with f_2 in $d = 3$

the choice of data-dependent smoothing parameters. In our analysis we focus on risk bounds and the amount of smoothness of the regression functions required for the parametric rate. A further novel contribution, previously not available in the literature, are the lower bounds which we provide and which show that, for unknown design density, some smoothness is required for the parametric rate to be attainable. Our analysis is thus complementary to much of the literature which focuses on asymptotic normality, testing and efficiency issues.

Assumption 1 is sometimes called the strong density assumption in the statistics literature (Audibert and Tsybakov, 2007). For nonparametric density estimation in random coefficient models in $d = 1$, Holzmann and Meister (2020) consider the effect on the minimax rates of a design density which tends to zero on the boundary of the support. Their method could be adapted to study estimation of the functionals Ψ and Φ in $d = 1$ under such weaker design assumptions. Multivariate extensions of such an analysis would be of some interest. A point of departure could be the case $d = 2$, in which removal of the bias with polynomial approximation is not required at least to obtain the parametric rates. A further extension of some interest would be to estimate functionals Ψ in which the integral is over more abstract compact manifolds, such as the unit sphere equipped with spherical Lebesgue measure, with the polynomial approximation being taken locally in the tangent space.

6 Proofs

6.1 Proof of Theorem 1

We use the notation introduced at the beginning of Section 2.2. Recall the estimator (2.4),

$$\hat{\Psi} = \sum_{J \in \mathcal{J}_K} \int_{C(J)} \hat{G}(z) dz,$$

where the Voronoi cells $C(J)$ are given in (2.3), and $\hat{G}(z)$ in (2.6). Again consider the expected conditional variance and expected squared conditional bias decomposition of the mean squared error (2.2),

$$\mathbb{E} [|\hat{\Psi} - \Psi|^2] = \mathbb{E} [\text{var}(\hat{\Psi}|\sigma_Z)] + \mathbb{E} [|\mathbb{E}(\hat{\Psi}|\sigma_Z) - \Psi|^2],$$

where σ_Z is the σ -field generated by Z_1, \dots, Z_n

We provide the main steps in the proof of the bound (2.9) in Theorem 1 for the expected conditional variance term in Section 6.1.1, and for the expected squared conditional bias in Section 6.1.2. Section B contains further technical details.

6.1.1 Expected conditional variance in (2.9)

We show the first inequality in (2.9), and proceed in the three steps.

In Step 1 we make use of conditional independence of the Y_1, \dots, Y_n given σ_Z , of the independence of two random Voronoi cells $C(J)$ and $C(J')$ for $J \cap J' = \emptyset$, as well as of the assumption of a bounded conditional variance $\text{var}(g(U, Z)|Z)$.

In Step 2 we derive a tight bound on $\mathbb{E} [\lambda(C(J)) \lambda(C(J'))]$ for $J \cap J' \neq \emptyset$, and conclude in the case $L = 0$.

Finally, in Step 3 we use a bound on the conditional moments of $(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)$ from Lemma 6.3, proven in Section B, to extend the bound to $L \geq 1$. Here, the invertability of $\mathcal{M}(z)$ almost surely for sufficiently large K follows from Lemma B.3 in Section B in the supplementary appendix.

Step 1: Using conditional independence

Note that if $J(z)$ and $J(z')$ are disjoint, the estimates $\widehat{G}(z)$ and $\widehat{G}(z')$ are independent conditionally on σ_Z . Therefore

$$\begin{aligned} & \mathbb{E} [\text{var}(\widehat{\Psi}|\sigma_Z)] \\ &= \mathbb{E} \left[\sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \int_{C(J)} \int_{C(J')} \text{cov}(\widehat{G}(z), \widehat{G}(z') | \sigma_Z) dz dz' \right] \\ &\leq \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \mathbb{E} \left[\int_{C(J)} \{\text{var}(\widehat{G}(z)|\sigma_Z)\}^{1/2} dz \cdot \int_{C(J')} \{\text{var}(\widehat{G}(z')|\sigma_Z)\}^{1/2} dz' \right] \end{aligned} \quad (6.1)$$

by applying the Cauchy-Schwarz inequality conditionally on σ_Z in the second step. By conditional independence of the $g(U_j, Z_j)$, $j = 1, \dots, n$ given σ_Z , observing (2.6) for all $z \in S^*$ we obtain

$$\begin{aligned} \text{var}(\widehat{G}(z)|\sigma_Z) &= \sum_{j \in J(z)} \text{var}(g(U_j, Z_j)|\sigma_Z) \cdot \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) \xi(z, Z_j)^\top \mathcal{M}(z)^{-1} \mathbf{e}_0 \\ &\leq C_V \cdot \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0. \end{aligned} \quad (6.2)$$

Therefore, (6.1) can be upper bounded by

$$\begin{aligned} & \mathbb{E} [\text{var}(\widehat{\Psi}|\sigma_Z)] \\ &\leq C_V \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \iint_{(S^*)^2} \mathbb{E} \left[\mathbf{1}_{C(J)}(z) \mathbf{1}_{C(J')}(z') (\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}(z')^{-1} \mathbf{e}_0)^{1/2} \right] dz dz'. \end{aligned} \quad (6.3)$$

To deal with the inner expected value, for $J \in \mathcal{J}_K$ and $z \in S^*$ let us introduce

$$\mathcal{M}_J(z) := \sum_{j \in J} \xi(z, Z_j) \xi(z, Z_j)^\top = \left(\sum_{j \in J} \xi_{\kappa+\kappa'}(z, Z_j) \right)_{\kappa, \kappa' \in \mathcal{K}}, \quad (6.4)$$

so that $\mathcal{M}_{J(z)}(z) = \mathcal{M}(z)$. Then for all $z, z' \in S^*$ and $J, J' \in \mathcal{J}_K$, since $\mathbf{1}_{C(J)}(z) = 1$ if and only if $J = J(z)$ we have that

$$\begin{aligned} & \mathbb{E} [\mathbf{1}_{C(J)}(z) \mathbf{1}_{C(J')}(z') (\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}(z')^{-1} \mathbf{e}_0)^{1/2}] \\ &= \mathbb{E} [\mathbf{1}_{C(J)}(z) \mathbf{1}_{C(J')}(z') (\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2}] \\ &= \mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2} \mathbb{P}(z \in C(J), z' \in C(J') | Z_k, k \in J \cup J') \right]. \end{aligned} \quad (6.5)$$

Further,

$$\begin{aligned} & \mathbb{P}(z \in C(J), z' \in C(J') | Z_k, k \in J \cup J') \\ &= \mathbb{P}(\|z - Z_j\| < \|z - Z_k\|, \|z' - Z_{j'}\| < \|z' - Z_{k'}\|, \forall j \in J, j' \in J', k \notin J, k' \notin J' | Z_k, k \in J \cup J') \\ &\leq \mathbb{P}(\|z - Z_j\| < \|z - Z_k\|, \|z' - Z_{j'}\| < \|z' - Z_{k'}\|, \forall j \in J, j' \in J', k, k' \notin J \cup J' | Z_k, k \in J \cup J') \\ &= \left(1 - \mathbb{P}_Z(B_d(z, \max_{j \in J} \|z - Z_j\|) \cup B_d(z', \max_{j' \in J'} \|z' - Z_{j'}\|)) \right)^{n - \#J \cup J'}, \end{aligned} \quad (6.6)$$

when we write \mathbb{P}_Z for the image measure of Z_1 , and $B_d(x, r)$ denotes the d -dimensional Euclidean ball around x with the radius r . For fixed $z, z' \in S^*$ and $J, J' \in \mathcal{J}_K$, put

$$\alpha := \alpha(z, z', J, J') := \max \left\{ \max_{j \in J} \|z - Z_j\|, \max_{j \in J'} \|z' - Z_j\| \right\}. \quad (6.7)$$

Then using Assumptions 1 and 2, (6.6) is bounded from above by

$$\exp \left(- (n - \#J \cup J') \cdot \rho \cdot \pi^{d/2} \cdot \min\{\rho', \alpha\}^d / \Gamma(d/2 + 1) \right),$$

and using the Cauchy-Schwarz inequality (6.5) is smaller or equal to

$$\begin{aligned} & \left(\mathbb{E} \left[\left(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0 \right) \exp \left(- (n - \#J \cup J') \cdot \rho \cdot \pi^{d/2} \cdot \min\{\rho', \alpha\}^d / \Gamma(d/2 + 1) \right) \right] \right)^{1/2} \\ & \cdot \left(\mathbb{E} \left[\left(\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0 \right) \exp \left(- (n - \#J \cup J') \cdot \rho \cdot \pi^{d/2} \cdot \min\{\rho', \alpha\}^d / \Gamma(d/2 + 1) \right) \right] \right)^{1/2}. \end{aligned} \quad (6.8)$$

To further bound (6.8) we make use of the fact that the non-zero terms in (6.3) have $J \cap J' \neq \emptyset$. If J and J' are not disjoint there exists some $j \in J \cap J'$ so that

$$\alpha \geq (\|z - Z_j\| + \|z' - Z_j\|) / 2 \geq \|z - z'\| / 2.$$

On the other hand,

$$\alpha \geq \max \left\{ \max_{j \in J} \|z - Z_j\|, \max_{j \in J'} \|z - Z_j\| - \|z - z'\| \right\} \geq \max_{j \in J \cup J'} \|z - Z_j\| - \|z - z'\|,$$

so that

$$\begin{aligned} \alpha & \geq \max \left\{ \|z - z'\| / 2, \max_{j \in J \cup J'} \|z - Z_j\| - \|z - z'\| \right\} \\ & \geq \max \left\{ \|z - z'\| / 2, \max_{j \in J \cup J'} \|z - Z_j\| / 3 \right\} \\ & =: \alpha^* = \alpha^*(z, z', J, J'). \end{aligned} \quad (6.9)$$

Let us summarize the bounds obtained so far in the following lemma.

Lemma 6.1. *Under the assumptions of Theorem 1, setting $\mathcal{M}_J(z)$ as in (6.4), $\alpha^* = \alpha^*(z, z', J, J')$ as in (6.9) and $C_\alpha = \rho \cdot \pi^{d/2} / \Gamma(d/2 + 1)$ we have the bound*

$$\begin{aligned} & \mathbb{E} \left[\text{var}(\hat{\Psi} | \sigma_Z) \right] \\ & \leq C_V \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \iint_{(S^*)^2} \left(\mathbb{E} \left[\left(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0 \right) \exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] \right)^{1/2} \\ & \quad \left(\mathbb{E} \left[\left(\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0 \right) \exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] \right)^{1/2} dz dz'. \end{aligned} \quad (6.10)$$

In particular, for $J, J' \in \mathcal{J}_K$ with $J \cap J' \neq \emptyset$,

$$\mathbb{E} \left[\lambda(C(J)) \lambda(C(J')) \right] \leq \iint_{(S^*)^2} \mathbb{E} \left[\exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] dz dz'. \quad (6.11)$$

Step 2: Concluding for $L = 0$.

For $L = 0$ we can drop $\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0 = 1/K$ from the expected values, so that we can insert (6.11) into (6.10). In the third step we show that for $L \geq 1$,

$$\begin{aligned} & \mathbb{E} \left[\left(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0 \right) \exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] \\ & \leq C_3 \mathbb{E} \left[\exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right], \end{aligned} \quad (6.12)$$

for some constant $C_3 > 0$. Then, by symmetry of (6.10) in z , J and z' and J' we obtain that the argument that we give here also applies to $L \geq 1$.

Writing

$$n^* := (n - 2K) C_\alpha \asymp n$$

and using the inequality $\#J \cup J' \leq 2K$, we have that

$$\mathbb{E} \left[\exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] \leq \exp \left(- n^* \cdot (\rho')^d \right) + \mathbb{E} \left[\exp \left(- n^* \cdot (\alpha^*)^d \right) \right]. \quad (6.13)$$

Since the first term decreases exponentially in n , we focus on the second. It holds that

$$\begin{aligned} \mathbb{E} \left[\exp \left(- n^* \cdot (\alpha^*)^d \right) \right] &= \int_0^1 \mathbb{P} \left[\alpha^* < \left(- (\log t) / n^* \right)^{1/d} \right] dt = \int_0^\infty \mathbb{P} \left[\alpha^* < (s/n^*)^{1/d} \right] \exp(-s) ds \\ &= \int_{s > n^* \|z - z'\|^d / 2^d} \mathbb{P} \left[\|z - Z_j\| < 3(s/n^*)^{1/d}, \forall j \in J \cup J' \right] \exp(-s) ds \\ &= \int_{s > n^* \|z - z'\|^d / 2^d} \mathbb{P}_Z \left(B_d(z, 3(s/n^*)^{1/d}) \right)^{\#J \cup J'} \exp(-s) ds \\ &\leq \left\{ \bar{\rho} \pi^{d/2} 3^d / (n^* \Gamma(d/2 + 1)) \right\}^{\#J \cup J'} \int_{s > n^* \|z - z'\|^d / 2^d} s^{\#J \cup J'} \exp(-s) ds. \end{aligned}$$

As $\max\{1, s^{2K}\} \exp(-s) \leq (4K)^{2K} \exp(-s/2)$ for all $s > 0$, it follows that

$$\begin{aligned} \int_{s > n^* \|z - z'\|^d / 2^d} s^{\#J \cup J'} \exp(-s) ds &\leq (4K)^{2K} \int_{s > n^* \|z - z'\|^d / 2^d} \exp(-s/2) ds \\ &= 2(4K)^{2K} \cdot \exp \left(- n^* \|z - z'\|^d / 2^{d+1} \right). \end{aligned}$$

Moreover,

$$\begin{aligned} \iint_{(S^*)^2} \exp \left(- n^* \|z - z'\|^d / 2^{d+1} \right) dz dz' &\leq \iint_{S^* \times \mathbb{R}^d} \exp \left(- n^* \|w\|^d / 2^{d+1} \right) dw dz' \\ &= \lambda_d(S^*) \cdot \int \exp \left(- \|w\|^d / 2^{d+1} \right) dw / n^*. \end{aligned}$$

Observing (6.11) in Lemma 6.1, we obtain the first part of the following lemma.

Lemma 6.2. *Under the assumptions of Theorem 1, for $J, J' \in \mathcal{J}_K$ with $J \cap J' \neq \emptyset$,*

$$\begin{aligned} \mathbb{E} \left[\lambda(C(J)) \lambda(C(J')) \right] &\leq \iint_{(S^*)^2} \mathbb{E} \left[\exp \left(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d \right) \right] dz dz' \\ &\leq \text{const.} \cdot n^{-\#J \cup J' - 1}, \end{aligned} \quad (6.14)$$

where the constant only depends on K , d , ρ , $\bar{\rho}$, ρ' and ρ'' .

In particular, for $L = 0$ we obtain

$$\mathbb{E} \left[\text{var}(\hat{\Psi} | \sigma_Z) \right] \lesssim n^{-1}.$$

For the second part, using Lemma 6.1 and the previous bound (6.14) we obtain

$$\mathbb{E} [\text{var}(\hat{\Psi}|\sigma_Z)] \leq \text{const. } C_V n^{-1} \cdot \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} n^{-\#J \cup J'}$$

Results from combinatorics provide that \mathcal{J}_K contains exactly $\binom{n}{K}$ elements; and that, for each $J \in \mathcal{J}_K$, there exist exactly $\binom{K}{\ell} \binom{n-K}{K-\ell}$ sets $J' \in \mathcal{J}_K$ with $\#(J \cap J') = \ell$. Therefore,

$$\begin{aligned} \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} n^{-\#J \cup J'} &= \sum_{\ell=1}^K \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{\#J \cap J' = \ell\} n^{\ell-2K} \\ &\leq \frac{n^K}{K!} \sum_{\ell=1}^K \binom{K}{\ell} \frac{1}{(K-\ell)!} \cdot n^{K-\ell+\ell-2K} \leq (2^K - 1)/K!, \end{aligned} \quad (6.15)$$

which proves the second part of the lemma.

Step 3: Reducing the case $L \geq 1$ to $L = 0$

We need to prove (6.12), then the calculation from the second step for $L = 0$ applies to $L \geq 1$ as well. This requires a quite sophisticated conditioning argument.

By \hat{j} we denote the smallest $j \in J$ such that $\|z - Z_j\| \geq \|z - Z_k\|$ for all $k \in J$; moreover we define

$$\alpha' := \max_{j \in J \setminus \hat{j}} \|z - Z_j\|.$$

The σ -fields generated by \hat{j} , $Z_{\hat{j}}$, α' , on the one hand, and by \hat{j} , $Z_{\hat{j}}$, on the other hand, are called \mathfrak{A}' and \mathfrak{A} , respectively. Note that with this notation we can write α^* in (6.9) as

$$\alpha^* = \max \{ \max\{\|z - Z_{\hat{j}}\|, \alpha'\}/3, \|z - z'\|/2 \},$$

and that the random variable α^* is measurable with respect to \mathfrak{A}' . Then the first factor in (6.8) has the upper bound

$$\begin{aligned} &\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \exp(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \mid \mathfrak{A}' \right] \exp(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \mid \mathfrak{A} \right] \exp(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha^*\}^d) \right], \end{aligned} \quad (6.16)$$

where in the last step we used that the random vector $(\mathbf{e}_0^\top \mathcal{M}_J(z) \mathbf{e}_0, \hat{j}, Z_{\hat{j}})$ and α' are independent.

Lemma 6.3. *Under the assumptions of Theorem 1, given $\eta \geq 1$ for $K \geq 1 + (\lfloor 2\eta D \rfloor + 1)K^*$ we have that*

$$\mathbb{E} [(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^\eta \mid \mathfrak{A}] \leq C_3,$$

almost surely, where the deterministic constant $C_3 \in (0, \infty)$ only depends on η , d , L , ρ , $\bar{\rho}$, ρ' , ρ'' .

The proof of the Lemma is deferred to Section B. This concludes the proof of the first inequality in (2.9). \square

6.1.2 Expected squared conditional bias in (2.9)

To prove the second inequality in (2.9) we proceed in two steps. First, for the case $L = 0$ we use a bound on expected integrated distances of points in S^* to their K -nearest neighbors provided in Lemma 6.4 below. Second, we use a Taylor approximation and again the bound for the conditional moments of $(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)$ from Lemma 6.3 to cover the case $L \geq 1$.

Step 1: The case $L = 0$.

Suppose that $L = 0$ and that $G \in \mathcal{G}(0, \beta, C_H)$. Then

$$\begin{aligned} \mathbb{E} [|\mathbb{E}[\hat{\Psi}|\sigma_Z] - \Psi|^2] &= \mathbb{E} \left[\left| \sum_{J \in \mathcal{J}_K} \int_{C(J)} \frac{1}{K} \sum_{j \in J} (G(Z_j) - G(z)) dz \right|^2 \right] \\ &\leq C_H^2 \sum_{J, J' \in \mathcal{J}_K} \int_{S^*} \int_{S^*} \mathbb{E} [1_{C(J)}(z) 1_{C(J')}(z') \|z - Z_{(K)}(z)\|^\beta \|z' - Z_{(K)}(z')\|^\beta] dz dz', \end{aligned}$$

where $Z_{(K)}(z)$ is the K -nearest neighbor of z in Z_1, \dots, Z_n .

Lemma 6.4. *Under the Assumptions 1 and 2 we have for $\beta > 0$ (not necessarily ≤ 1) and $K \in \mathbb{N}$ that*

$$\sum_{J, J' \in \mathcal{J}_K} \int_{S^*} \int_{S^*} \mathbb{E} [1_{C(J)}(z) 1_{C(J')}(z') \|z - Z_{(K)}(z)\|^\beta \|z' - Z_{(K)}(z')\|^\beta] dz dz' \lesssim n^{-\frac{2\beta}{d}}.$$

Proof of Lemma 6.4. For $\alpha = \alpha(z, z', J, J')$ as in (6.7), with the argument leading to (6.8) we have that

$$\begin{aligned} &\mathbb{E} [1_{C(J)}(z) 1_{C(J')}(z') \|z - Z_{(K)}(z)\|^\beta \|z' - Z_{(K)}(z')\|^\beta] \\ &\leq \mathbb{E} [\alpha^{2\beta} \exp(- (n - \#J \cup J') \cdot C_\alpha \cdot \min\{\rho', \alpha\}^d)] \\ &\leq (\rho'')^{2\beta} \exp(-n^* (\rho')^d) + \mathbb{E} [\alpha^{2\beta} \exp(-n^* \cdot \alpha^d)], \end{aligned}$$

where as before $C_\alpha = \rho \cdot \pi^{d/2} / \Gamma(d/2 + 1)$ and $n^* = (n - 2K) C_\alpha$. Now

$$\begin{aligned} &\mathbb{E} [\alpha^{2\beta} \exp(-n^* \cdot \alpha^d)] \\ &= \int_{S^{J \cup J'}} \max_{j \in J \cup J'} \|z - z_j\|^{2\beta} \cdot \exp(-n^* \max_{j \in J \cup J'} \|z - z_j\|^d) \prod_{j \in J \cup J'} f_Z(z_j) dz_j \\ &\leq (\bar{\rho})^{2K} \int_{(\mathbb{R}^d)^{J \cup J'}} \max_{j \in J \cup J'} \|u_j\|^{2\beta} \cdot \exp(-n^* \max_{j \in J \cup J'} \|u_j\|^d) \prod_{j \in J \cup J'} du_j \\ &\leq (\bar{\rho} d C_d)^{2K} \int_{(0, \infty)^{J \cup J'}} \max_{j \in J \cup J'} r_j^{2\beta} \cdot \exp(-n^* \max_{j \in J \cup J'} r_j^d) \prod_{j \in J \cup J'} r_j^{d-1} dr_j \\ &= (\bar{\rho} d C_d)^{2K} (n^*)^{-2\beta/d} (n^*)^{-\#(J \cup J')(d-1)/d} (n^*)^{-\#(J \cup J')/d} \\ &\quad \cdot \int_{(0, \infty)^{J \cup J'}} \max_{j \in J \cup J'} r_j^{2\beta} \cdot \exp(-\max_{j \in J \cup J'} r_j^d) \prod_{j \in J \cup J'} r_j^{d-1} dr_j \\ &\leq \text{const.} \cdot n^{-2\beta/d} n^{-\#(J \cup J')}, \end{aligned}$$

and observing as in (6.15),

$$\sum_{J, J' \in \mathcal{J}_K} n^{-\#J \cup J'} \leq 2^K / K!$$

concludes the proof of the lemma. \square

Step 2: Taylor expansion for $L = l \geq 1$

Now suppose that $G \in \mathcal{G}(l, \beta, C_H)$, $l \geq 1$ and that $L = l$.

The Taylor expansion of G in z , evaluated at Z_j up to order L can be written as

$$G(Z_j) = \xi(z, Z_j)^\top \mathcal{G}(z) + \mathcal{R}(z, Z_j),$$

where

$$\begin{aligned} \mathcal{G}(z) &:= (\partial_\kappa G(z) / \kappa!)_{\kappa \in \mathcal{K}}, \\ \mathcal{R}(z, Z_j) &:= L \sum_{|\kappa'|=L} \frac{\xi_{\kappa'}(z, Z_j)}{\kappa'!} \cdot \int_0^1 (1-s)^{L-1} (\partial_{\kappa'} G(z + s(Z_j - z)) - \partial_{\kappa'} G(z)) ds. \end{aligned}$$

Therefore, observing $\mathbf{e}_0^\top \mathcal{G}(z) = G(z)$ and (2.6) we obtain that

$$\mathbb{E}[\hat{G}(z) | \sigma_Z] = \sum_{j \in J(z)} \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) G(Z_j) = G(z) + \sum_{j \in J(z)} \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) \mathcal{R}(z, Z_j)$$

Therefore, it holds that

$$\mathbb{E}[\hat{\Psi} | \sigma_Z] - \Psi = \sum_{J \in \mathcal{J}_K} \int_{C(J)} \sum_{j \in J} \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) \mathcal{R}(z, Z_j) dz,$$

almost surely, where we have used that the $C(J)$, $J \in \mathcal{J}_K$, are pairwise disjoint and form an almost-partition of S^* ; concretely, $S^* \setminus \bigcup_{J \in \mathcal{J}_K} C(J)$ is included in

$$\bigcup_{j, k=1}^n \{z \in S^* \mid \|z - Z_k\| = \|z - Z_j\|\},$$

as a subset, which is a d -dimensional Borel set with the Lebesgue measure zero as Z_1 has a Lebesgue density f_Z . Using the Cauchy-Schwarz inequality and the multinomial theorem twice we deduce that

$$\begin{aligned} & \left| \sum_{j \in J(z)} \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) \mathcal{R}(z, Z_j) \right| \\ & \leq C_H \left\{ \sum_{j \in J(z)} \|z - Z_j\|^{2\beta} \left(\sum_{|\kappa'|=L+1} \frac{1}{\kappa'!} \xi_{\kappa'}(z, Z_j)^2 \right) \sum_{|\kappa'|=L+1} \frac{1}{\kappa'!} \right\}^{1/2} \\ & \quad \cdot \left(\sum_{j \in J(z)} |\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j)|^2 \right)^{1/2} \\ & \leq C_H \frac{K^{L/2+1/2}}{(L+1)!} \left(\sum_{j \in J(z)} \|Z_j - z\|^{2(L+\beta)} \right)^{1/2} (\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0)^{1/2} \\ & \leq C_H \frac{K^{L/2+1}}{(L+1)!} \|Z_{(K)}(z) - z\|^{L+\beta} (\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0)^{1/2}. \end{aligned}$$

Thus the expected squared conditional bias is bounded from above as follows.

$$\begin{aligned}
& \mathbb{E} \left[\left| \mathbb{E}(\hat{\Psi} | \sigma_Z) - \Psi \right|^2 \right] \\
& \leq C_H^2 \frac{K^{L+2}}{(L+1)!^2} \sum_{J, J' \in \mathcal{J}_K} \iint_{(S^*)^2} \mathbb{E} \left[\mathbf{1}_{C(J)}(z) \mathbf{1}_{C(J')}(z') \cdot (\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} \right. \\
& \quad \left. (\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2} \|Z_{(K)}(z) - z\|^{L+\beta} \|Z_{(K)}(z') - z'\|^{L+\beta} \right] dz dz' \\
& \leq C_H^2 \frac{K^{L+2}}{(L+1)!^2} \sum_{J, J' \in \mathcal{J}_K} \iint_{(S^*)^2} \mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2} \right. \\
& \quad \left. \alpha^{2\beta} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] dz dz' \tag{6.17}
\end{aligned}$$

with $\alpha = \alpha(z, z', J, J')$ as in (6.7), and again $n^* = (n - 2K)C_\alpha$ and using the argument leading to (6.8). Using the Cauchy Schwarz inequality,

$$\begin{aligned}
& \mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} (\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2} \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \\
& \leq \left(\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2} \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \right)^{1/2} \\
& \quad \cdot \left(\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_{J'}(z')^{-1} \mathbf{e}_0)^{1/2} \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \right)^{1/2}
\end{aligned}$$

Now the conditioning argument from step 3 in Section 6.1.1 can again be applied to eliminate the terms $(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^{1/2}$ and reduce to the setting with $L = 0$:

Let \hat{j} denote the smallest $j \in J$ such that $\|z - Z_j\| \geq \|z - Z_k\|$ for all $k \in J$ and define

$$\alpha' := \max_{j \in J' \setminus J} \|z - Z_j\|,$$

where now possibly $J \cap J' = \emptyset$. The σ -fields generated by $\hat{j}, Z_{\hat{j}}, \alpha'$, on the one hand, and by $\hat{j}, Z_{\hat{j}}$, on the other hand, are again called \mathfrak{A}' and \mathfrak{A} , respectively. Note that with this notation we can write α as

$$\alpha = \max\{\|z - Z_{\hat{j}}\|, \alpha'\}.$$

and that the random variable α is measurable with respect to \mathfrak{A}' . Then as in (6.16),

$$\begin{aligned}
& \mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \\
& = \mathbb{E} \left[\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \mid \mathfrak{A}' \right] \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \\
& = \mathbb{E} \left[\mathbb{E} \left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0) \mid \mathfrak{A} \right] \alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right] \\
& \leq C_3 \mathbb{E} \left[\alpha^{2(L+\beta)} \exp(-n^* \cdot \min\{\rho', \alpha\}^d) \right]
\end{aligned}$$

where in the second step we used that the random vector $(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0, \hat{j}, Z_{\hat{j}})$ and α' are independent, and in the last step we applied Lemma 6.3 with $\eta = 1$. The argument is completed as in step 1. \square

6.2 Proof of Theorem 2

Let $\sigma_{X,Z}$ denote the σ -field generated by $X_1, \dots, X_m, Z_1, \dots, Z_n$, and let σ_Z denote the σ -field generated by Z_1, \dots, Z_n only. Then we have the decomposition

$$\mathbb{E} \left[\left| \hat{\Phi} - \Phi \right|^2 \right] = \mathbb{E} \left[\text{var}(\hat{\Phi} | \sigma_{X,Z}) \right] + \mathbb{E} \left[\text{var}(\mathbb{E}[\hat{\Phi} | \sigma_{X,Z}] | \sigma_Z) \right] + \mathbb{E} \left[\left| \mathbb{E}[\hat{\Phi} | \sigma_Z] - \Phi \right|^2 \right]. \tag{6.18}$$

We shall show the following bounds:

$$\sup_{G \in \mathcal{G}(l, \beta, C_H, C_G)} \mathbb{E} [\text{var}(\widehat{\Phi} | \sigma_{X,Z})] \leq C_V \cdot C \cdot (n^{-1} + m^{-1}), \quad (6.19)$$

$$\sup_{G \in \mathcal{G}(l, \beta, C_H, C_G)} \mathbb{E} [\text{var}(\mathbb{E}[\widehat{\Phi} | \sigma_{X,Z}] | \sigma_Z)] \leq C_G^2 \cdot C \cdot m^{-1}, \quad (6.20)$$

$$\sup_{G \in \mathcal{G}(l, \beta, C_H, C_G)} \mathbb{E} [|\mathbb{E}[\widehat{\Phi} | \sigma_Z] - \Phi|^2] \leq C_H^2 \cdot C \cdot n^{-\frac{2(l+\beta)}{d}}, \quad (6.21)$$

where the constant $C > 0$ only depends on $L, K, \rho, \rho', \rho'', \bar{\rho}, d, \lambda(S^*)$ and C_f . These bounds imply (2.11), and hence the statement of the theorem.

Proof of (6.19): By conditional independence if $J \cap J' = \emptyset$,

$$\text{var}(\widehat{\Phi} | \sigma_{X,Z}) = \frac{1}{m^2} \sum_{k, k'=1}^m \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}(J \cap J' \neq \emptyset) \mathbf{1}(X_k \in C(J), X_{k'} \in C(J')) \text{cov}(\widehat{G}(X_k), \widehat{G}(X_{k'}) | \sigma_{X,Z}).$$

Then

$$\begin{aligned} \mathbb{E} [\text{var}(\widehat{\Phi} | \sigma_{X,Z}) | \sigma_Z] &= \frac{1}{m} \int_{S^*} \text{var}(\widehat{G}(z) | \sigma_Z) f(z) dz \\ &+ \frac{m(m-1)}{m^2} \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}(J \cap J' \neq \emptyset) \int_{C(J)} \int_{C(J')} \text{cov}(\widehat{G}(z), \widehat{G}(z') | \sigma_Z) f(z) f(z') dz dz'. \end{aligned} \quad (6.22)$$

Since

$$\begin{aligned} &\sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \int_{C(J)} \int_{C(J')} \text{cov}(\widehat{G}(z), \widehat{G}(z') | \sigma_Z) f(z) f(z') dz dz' \\ &\leq C_f^2 \sum_{J, J' \in \mathcal{J}_K} \mathbf{1}\{J \cap J' \neq \emptyset\} \int_{C(J)} \{\text{var}(\widehat{G}(z) | \sigma_Z)\}^{1/2} dz \cdot \int_{C(J')} \{\text{var}(\widehat{G}(z') | \sigma_Z)\}^{1/2} dz', \end{aligned} \quad (6.23)$$

the expected value of the second term in (6.22) can be upper bounded as the conditional variance term in (2.9), see (6.1). For the first term in (6.22), we use the bound (6.2) on $\text{var}(\widehat{G}(z) | \sigma_Z)$ as well as Lemma 6.3 to obtain that $\mathbb{E}[\text{var}(\widehat{G}(z) | \sigma_Z)]$ is bounded from above uniformly in z . This proves (6.19).

Proof of (6.20) and (6.21):

Write

$$\bar{G}(z; Z) = \mathbb{E} [\widehat{G}(z) | \sigma_Z] = \sum_{j \in J(z)} \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j) G(Z_j). \quad (6.24)$$

Then

$$\mathbb{E} [\widehat{G}(X_k) | \sigma_{X,Z}] = \bar{G}(X_k; Z),$$

and therefore

$$\mathbb{E}[\widehat{\Phi} | \sigma_{X,Z}] = \frac{1}{m} \sum_{k=1}^m \sum_{J \in \mathcal{J}_K} \mathbf{1}(X_k \in C(J)) \bar{G}(X_k; Z).$$

Further,

$$|\mathbb{E}[\widehat{\Phi} | \sigma_Z] - \Phi| = \left| \int_{S^*} (\mathbb{E}[\widehat{G}(z) | \sigma_Z] - G(z)) f(z) dz \right| \leq C_f \int_{S^*} |\mathbb{E}[\widehat{G}(z) | \sigma_Z] - G(z)| dz,$$

which can be bounded as the conditional bias term in (2.9) so that (6.21) follows. Moreover,

$$\begin{aligned} \text{var}(\mathbb{E}[\widehat{\Phi}|\sigma_{X,Z}]|\sigma_Z) &= \frac{1}{m} \text{var}\left(\sum_{J \in \mathcal{J}_K} 1(X_1 \in C(J)) \bar{G}(X_1; Z)|\sigma_Z\right) \\ &\leq \frac{1}{m} \int_{S^*} (\mathbb{E}[\hat{G}(z)|\sigma_Z])^2 f(z) \, dz. \end{aligned}$$

Using the Cauchy-Schwarz inequality in (6.24) we obtain

$$(\mathbb{E}[\hat{G}(z)|\sigma_Z])^2 \leq K C_G^2 \cdot \sum_{j \in J(z)} |\mathbf{e}_0^\top \mathcal{M}(z)^{-1} \xi(z, Z_j)|^2 = K C_G^2 \cdot \mathbf{e}_0^\top \mathcal{M}(z)^{-1} \mathbf{e}_0.$$

Using Lemma 6.3 proves (6.20) and hence concludes the proof of the Theorem. \square

References

- Abadie, A. and Imbens, G. W. (2006). Large sample properties of matching estimators for average treatment effects. *Econometrica*, 74(1):235–267.
- Abadie, A. and Imbens, G. W. (2011). Bias-corrected matching estimators for average treatment effects. *Journal of Business & Economic Statistics*, 29(1):1–11.
- Audibert, J.-Y. and Tsybakov, A. B. (2007). Fast learning rates for plug-in classifiers. *Ann. Stat.*, 35(2):608–633.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *Econom. J.*, 21(1):c1–c68.
- Chernozhukov, V., Hansen, C., Kallus, N., Spindler, M., and Syrgkanis, V. (2024). Applied causal inference powered by ml and ai. *arXiv preprint arXiv:2403.02467*.
- Dahlke, S., Heuer, S., Holzmann, H., and Tafo, P. (2022). Statistically optimal estimation of signals in modulation spaces using gabor frames. *IEEE Transactions on Information Theory*, 68(6):4182–4200.
- Delaigle, A., Hall, P., and Qiu, P. (2006). Nonparametric methods for solving the Berkson errors-in-variables problem. *J. R. Stat. Soc., Ser. B, Stat. Methodol.*, 68(2):201–220.
- Devroye, L., Györfi, L., Lugosi, G., and Walk, H. (2017). On the measure of voronoi cells. *Journal of Applied Probability*, 54(2):394–408.
- Dunker, F., Eckle, K., Proksch, K., and Schmidt-Hieber, J. (2019). Tests for qualitative features in the random coefficients model. *Electron. J. Stat.*, 13(2):2257–2306.
- Gaillac, C. and Gautier, E. (2022). Adaptive estimation in the linear random coefficients model when regressors have limited variation. *Bernoulli*, 28(1):504–524.
- Gautier, E. and Kitamura, Y. (2013). Nonparametric estimation in random coefficients binary choice models. *Econometrica*, 81(2):581–607.

- Hoderlein, S., Holzmann, H., and Meister, A. (2017). The triangular model with random coefficients. *Journal of Econometrics*, 201(1):144–169.
- Hoderlein, S., Klemelä, J., and Mammen, E. (2010). Analyzing the random coefficient model non-parametrically. *Econom. Theory*, 26(3):804–837.
- Holzmann, H. and Meister, A. (2020). Rate-optimal nonparametric estimation for random coefficient regression models. *Bernoulli*, 26(4):2790–2814.
- Hu, A. J., Green, A., and Tibshirani, R. J. (2022). The voronoigram: minimax estimation of bounded variation functions from scattered data. *arXiv preprint arXiv: 2212.14514*.
- Imbens, G. W. (2004). Nonparametric estimation of average treatment effects under exogeneity: A review. *Review of Economics and Statistics*, 86(1):4–29.
- Kallus, N. (2020). Generalized optimal matching methods for causal inference. *Journal of Machine Learning Research*, 21(62):1–54.
- Kerkycharian, G. and Picard, D. (2004). Regression in random design and warped wavelets. *Bernoulli*, 10(6):1053–1105.
- Kohler, M. (2014). Optimal global rates of convergence for noiseless regression estimation problems with adaptively chosen design. *Journal of Multivariate Analysis*, 132:197–208.
- Kouw, W. M. and Loog, M. (2019). An introduction to domain adaptation and transfer learning. *arXiv preprint arXiv:1812.11806*.
- Lewbel, A. and Schennach, S. M. (2007). A simple ordered data estimator for inverse density weighted expectations. *Journal of Econometrics*, 136(1):189–211.
- Lin, Z., Ding, P., and Han, F. (2023). Estimation based on nearest neighbor matching: from density ratio to average treatment effect. *Econometrica*, 91(6):2187–2217.
- Meister, A. (2009). *Deconvolution problems in nonparametric statistics*. Lecture Notes in Statistics, Springer.
- Portier, F., Truquet, L., and Yamane, I. (2023). Scalable and hyper-parameter-free non-parametric covariate shift adaptation with conditional sampling. *arXiv preprint arXiv:2312.09969*.
- Samworth, R. J. (2012). Optimal weighted nearest neighbour classifiers. *Ann. Statist.*, 40(5):2733–2763.
- Sharpnack, J. (2023). On L^2 -consistency of nearest neighbor matching. *IEEE Trans. Inf. Theory*, 69(6):3978–3988.
- Sugiyama, M., Nakajima, S., Kashima, H., Buenau, P., and Kawanabe, M. (2007). Direct importance estimation with model selection and its application to covariate shift adaptation. *Advances in neural information processing systems*, 20.
- Tsybakov, A. (2009). *Introduction to nonparametric estimation*. Springer.

Wang, Y. and Shah, R. D. (2024). Debiased inverse propensity score weighting for estimation of average treatment effects with high-dimensional confounders. *Ann. Statist.*, to appear; *arXiv preprint arXiv:2011.08661*.

A Proof of Theorem 3

Proof of Theorem 3: Set $I'_k := [k/M, (k+1/2)/M]$, $I''_k := ((k+1/2)/M, (k+1)/M)$, $k = 0, \dots, M-1$ for some integer $M > 1$. Moreover, let

$$\varphi_k := (1 + \theta_k \alpha) 1_{I'_k} + (1 - \theta_k \alpha) 1_{I''_k},$$

for some $\alpha > 0$ and $\theta = (\theta_0, \dots, \theta_{M-1}) \in \{0\} \cup \{-1, 1\}^M$; and

$$f_\theta := \sum_{k=0}^{M-1} \varphi_k.$$

Now write $f^{(n)}$ for the n -fold product density of a density f ; and let $\hat{\theta} = (\hat{\theta}_0, \dots, \hat{\theta}_{M-1})$ be a random vector with i.i.d. components which satisfy $P[\hat{\theta}_k = 1] = P[\hat{\theta}_k = -1] = 1/2$ for all $k = 0, \dots, M-1$. The densities $f_{\hat{\theta}}$ and f_0 are used as candidates for the design density f_Z where α is chosen sufficiently small such that $f_{\hat{\theta}}, f_0 \in \mathcal{F}$ are guaranteed. Put $x = (x_1, \dots, x_n)$ and

$$\begin{aligned} N_k^+(x) &:= \sum_{j=1}^n 1_{\{1\}}(\hat{\theta}_k) \cdot 1_{I'_k}(x_j) + 1_{\{-1\}}(\hat{\theta}_k) \cdot 1_{I''_k}(x_j), \\ N_k^-(x) &:= \sum_{j=1}^n 1_{\{1\}}(\hat{\theta}_k) \cdot 1_{I''_k}(x_j) + 1_{\{-1\}}(\hat{\theta}_k) \cdot 1_{I'_k}(x_j). \end{aligned}$$

Consider that

$$f_{\hat{\theta}}^{(n)}(x) = \prod_{k=0}^{M-1} (1 + \alpha)^{N_k^+(x)} \cdot (1 - \alpha)^{N_k^-(x)}.$$

The likelihood function $\eta_{\hat{\theta}}^{(n)}$ equals

$$\eta_{\hat{\theta}}^{(n)}(\mathbf{u}, \mathbf{z}) = \prod_{j=1}^n f_\theta(z_j) f_\varepsilon(u_j - h_\theta(z_j)),$$

with $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{z} = (z_1, \dots, z_n)$, where we choose the regression function $h_\theta = \beta \cdot F(f_\theta)$ for the regression function where the function F satisfies $F(1) = 1$ but, apart from that, remains to be selected; and f_ε denotes the standard normal density. The parameter $\beta > 0$ is chosen sufficiently small such that $h_{\hat{\theta}}, h_0 \in \mathcal{H}$. Writing

$$I_k^+ := \begin{cases} I'_k, & \text{if } \hat{\theta}_k = 1, \\ I''_k, & \text{if } \hat{\theta}_k = -1, \end{cases}$$

and

$$I_k^- := \begin{cases} I'_k, & \text{if } \hat{\theta}_k = -1, \\ I''_k, & \text{if } \hat{\theta}_k = 1, \end{cases}$$

we deduce that

$$|\mathbb{E} \eta_{\hat{\theta}}^{(n)}(\mathbf{u}, \mathbf{z})|^2$$

$$\begin{aligned}
&= [f_\varepsilon^{(n)}(\mathbf{u})]^2 \cdot \left| \prod_{k=0}^{M-1} \mathbb{E}(1+\alpha)^{N_k^+(\mathbf{z})} (1-\alpha)^{N_k^-(\mathbf{z})} \exp \left\{ -\frac{1}{2} \beta^2 [F^2(1+\alpha)N_k^+(\mathbf{z}) + F^2(1-\alpha)N_k^-(\mathbf{z})] \right\} \right. \\
&\quad \left. \cdot \exp \left\{ \beta F(1+\alpha) \sum_{j=1}^n 1_{I_k^+}(z_j) u_j + \beta F(1-\alpha) \sum_{j=1}^n 1_{I_k^-}(z_j) u_j \right\} \right|^2 \\
&= [f_\varepsilon^{(n)}(\mathbf{u})]^2 \cdot \prod_{k=0}^{M-1} \left\{ \frac{1}{4} (1+\alpha)^{2N_k'(\mathbf{z})} (1-\alpha)^{2N_k''(\mathbf{z})} \exp \left[-\beta^2 [F^2(1+\alpha)N_k'(\mathbf{z}) + F^2(1-\alpha)N_k''(\mathbf{z})] \right] \right. \\
&\quad \cdot \exp \left[2\beta F(1+\alpha) \sum_{j=1}^n 1_{I_k'}(z_j) u_j + 2\beta F(1-\alpha) \sum_{j=1}^n 1_{I_k''}(z_j) u_j \right] \\
&\quad + \frac{1}{4} (1+\alpha)^{2N_k''(\mathbf{z})} (1-\alpha)^{2N_k'(\mathbf{z})} \exp \left[-\beta^2 [F^2(1+\alpha)N_k''(\mathbf{z}) + F^2(1-\alpha)N_k'(\mathbf{z})] \right] \\
&\quad \cdot \exp \left[2\beta F(1+\alpha) \sum_{j=1}^n 1_{I_k''}(z_j) u_j + 2\beta F(1-\alpha) \sum_{j=1}^n 1_{I_k'}(z_j) u_j \right] \\
&\quad + \frac{1}{2} (1-\alpha^2)^{N_k(\mathbf{z})} \exp \left[-\frac{\beta^2}{2} (F^2(1+\alpha) + F^2(1-\alpha)) \cdot N_k(\mathbf{z}) \right] \\
&\quad \left. \cdot \exp \left[\beta [F(1+\alpha) + F(1-\alpha)] \sum_{j=1}^n 1_{I_k}(z_j) u_j \right] \right\},
\end{aligned}$$

where $N_k'(x) := \sum_{j=1}^n 1_{I_k'}(x_j)$, $N_k''(x) := \sum_{j=1}^n 1_{I_k''}(x_j)$ and $N_k(x) := N_k'(x) + N_k''(x)$. On the other hand,

$$\eta_0^{(n)}(\mathbf{u}, \mathbf{z}) = f_\varepsilon^{(n)}(\mathbf{u}) \cdot \exp(-\beta^2 n/2) \cdot \exp\left(\beta \sum_{j=1}^n u_j\right),$$

on $\mathbf{z} \in [0, 1]^n$ so that

$$\begin{aligned}
&\int |\mathbb{E} \eta_\theta^{(n)}(\mathbf{u}, \mathbf{z})|^2 / \eta_0^{(n)}(\mathbf{u}, \mathbf{z}) d\mathbf{u} \\
&= \exp(\beta^2 n/2) \cdot \prod_{k=0}^{M-1} \left\{ \frac{1}{4} (1+\alpha)^{2N_k'(\mathbf{z})} (1-\alpha)^{2N_k''(\mathbf{z})} \exp \left[-\beta^2 [F^2(1+\alpha)N_k'(\mathbf{z}) + F^2(1-\alpha)N_k''(\mathbf{z})] \right] \right. \\
&\quad \cdot \exp \left[\frac{\beta^2}{2} [2F(1+\alpha) - 1]^2 N_k'(\mathbf{z}) + \frac{\beta^2}{2} [2F(1-\alpha) - 1]^2 N_k''(\mathbf{z}) \right] \\
&\quad + \frac{1}{4} (1+\alpha)^{2N_k''(\mathbf{z})} (1-\alpha)^{2N_k'(\mathbf{z})} \exp \left[-\beta^2 [F^2(1+\alpha)N_k''(\mathbf{z}) + F^2(1-\alpha)N_k'(\mathbf{z})] \right] \\
&\quad \cdot \exp \left[\frac{\beta^2}{2} [2F(1+\alpha) - 1]^2 N_k''(\mathbf{z}) + \frac{\beta^2}{2} [2F(1-\alpha) - 1]^2 N_k'(\mathbf{z}) \right] \\
&\quad + \frac{1}{2} (1-\alpha^2)^{N_k(\mathbf{z})} \exp \left[-\frac{\beta^2}{2} (F^2(1+\alpha) + F^2(1-\alpha)) \cdot N_k(\mathbf{z}) \right] \\
&\quad \left. \cdot \exp \left[\frac{\beta^2}{2} (F(1+\alpha) + F(1-\alpha) - 1)^2 \cdot N_k(\mathbf{z}) \right] \right\}.
\end{aligned}$$

Now we integrate this term with respect to \mathbf{z} . Thus consider \mathbf{Z} as an n -dimensional random vector which consists of i.i.d. components that are uniformly distributed on $[0, 1]$. Given $(N_0(\mathbf{Z}), \dots, N_{M-1}(\mathbf{Z}))$, the random variable $N_k'(\mathbf{Z})$ are conditionally independent and conditionally binomially distributed

with the parameters $N_k(\mathbf{Z})$ and $1/2$. Therefore,

$$\begin{aligned}
& \iint |\mathbb{E}\eta_{\hat{\theta}}^{(n)}(\mathbf{u}, \mathbf{z})|^2 / \eta_0^{(n)}(\mathbf{u}, \mathbf{z}) d\mathbf{u} d\mathbf{z} \\
&= \mathbb{E} \prod_{k=0}^{M-1} (\gamma_0^{N_k(\mathbf{Z})} / 2 + \gamma_1^{N_k(\mathbf{Z})} / 2) \\
&= 2^{-M} \sum_{\mathcal{N} \subseteq \{0, \dots, M-1\}} \mathbb{E} \gamma_0^{\sum_{k \in \mathcal{N}} N_k(\mathbf{Z})} \gamma_1^{n - \sum_{k \in \mathcal{N}} N_k(\mathbf{Z})} \\
&= \mathbb{E} \left\{ \gamma_0 \frac{B}{M} + \gamma_1 \left(1 - \frac{B}{M}\right) \right\}^n, \tag{A.1}
\end{aligned}$$

where $B \sim \mathbf{B}(M, 1/2)$ and

$$\begin{aligned}
\gamma_0 &:= \frac{1}{2}(1 + \alpha)^2 \exp \{ \beta^2 [1 - F(1 + \alpha)]^2 \} + \frac{1}{2}(1 - \alpha)^2 \exp \{ \beta^2 [1 - F(1 - \alpha)]^2 \}, \\
\gamma_1 &:= (1 - \alpha^2) \exp \{ \beta^2 [1 - F(1 + \alpha)] \cdot [1 - F(1 - \alpha)] \}.
\end{aligned}$$

Then, (A.1) equals

$$\begin{aligned}
& \left(\frac{1}{2} \gamma_0 + \frac{1}{2} \gamma_1 \right)^n \cdot \mathbb{E} \left\{ 1 + \frac{2\gamma_0 - 2\gamma_1}{\gamma_0 + \gamma_1} \cdot \left(\frac{B}{M} - \frac{1}{2} \right) \right\}^n \\
& \leq \left(\frac{1}{2} \gamma_0 + \frac{1}{2} \gamma_1 \right)^n \cdot \exp \left\{ \frac{(\gamma_0 - \gamma_1)^2}{2(\gamma_0 + \gamma_1)^2} \cdot \frac{n^2}{M} \right\}.
\end{aligned}$$

By Taylor expansion we obtain that

$$\begin{aligned}
\gamma_0/2 + \gamma_1/2 &= \frac{1}{4} \sum_{k=0}^{\infty} \frac{1}{k!} \beta^{2k} \left((1 + \alpha)[1 - F(1 + \alpha)]^k + (1 - \alpha)[1 - F(1 - \alpha)]^k \right)^2, \\
\gamma_0 - \gamma_1 &= \frac{1}{2} \sum_{k=0}^{\infty} \frac{1}{k!} \beta^{2k} \left((1 + \alpha)[1 - F(1 + \alpha)]^k - (1 - \alpha)[1 - F(1 - \alpha)]^k \right)^2.
\end{aligned}$$

Thus the χ^2 -distance between $\mathbb{E}\eta_{\hat{\theta}}^{(n)}$ and $\eta_0^{(n)}$ admits the following upper bound

$$\chi^2(\mathbb{E}\eta_{\hat{\theta}}^{(n)}, \eta_0^{(n)}) \leq \exp(c + c^2/16) - 1,$$

for all n, M whenever

$$\sum_{k=1}^{\infty} \frac{1}{k!} \beta^{2k} \left((1 + \alpha)[1 - F(1 + \alpha)]^k + (1 - \alpha)[1 - F(1 - \alpha)]^k \right)^2 \leq c/n, \tag{A.2}$$

$$\sum_{k=0}^{\infty} \frac{1}{k!} \beta^{2k} \left((1 + \alpha)[1 - F(1 + \alpha)]^k - (1 - \alpha)[1 - F(1 - \alpha)]^k \right)^2 \leq c \cdot n^{-1} M^{1/2}, \tag{A.3}$$

for all n, M , where the constant is $c > 0$ sufficiently small.

Impose that M is an integer multiple of 4. Then,

$$\left| \int_{1/4}^{3/4} h_{\hat{\theta}} - \int_{1/4}^{3/4} h_0 \right|^2 = \frac{1}{16} \beta^2 (F(1 + \alpha) + F(1 - \alpha) - 2)^2, \tag{A.4}$$

holds true almost surely. Now we specify that $F(x) = 1/x$, which provides the lower bound $\beta^2\alpha^4$ on (A.4) (up to a constant factor). Furthermore the left side of the inequality (A.2) is bounded from above by $\alpha^4\beta^4$ (again up to some constant factor), while (A.3) can always be satisfied by selecting M sufficiently large. Note that there are no smoothness restrictions. Choose α as a positive constant sufficiently small; and $\beta = \beta_n \asymp n^{-1/4}$ with a sufficiently small constant factor.

Using Le Cam's inequality, the consideration

$$\begin{aligned}
& \sup_{f_Z \in \mathcal{F}, h \in \mathcal{H}} \mathbb{E}_{f_Z, h} \left| \hat{H}_n - \int_{1/4}^{3/4} h(x) dx \right|^2 \\
& \geq \frac{1}{2} \iint \mathbb{E} \left| \hat{H}_n(\mathbf{u}, \mathbf{z}) - \int_{1/4}^{3/4} h_{\hat{\theta}}(x) dx \right|^2 \eta_{\hat{\theta}}^{(n)}(\mathbf{u}, \mathbf{z}) d\mathbf{u} d\mathbf{z} \\
& \quad + \frac{1}{2} \iint \left| \hat{H}_n(\mathbf{u}, \mathbf{z}) - \int_{1/4}^{3/4} h_0(x) dx \right|^2 \eta_0^{(n)}(\mathbf{u}, \mathbf{z}) d\mathbf{u} d\mathbf{z} \\
& \geq \frac{1}{64} \beta^2 (F(1+\alpha) + F(1-\alpha) - 2)^2 \cdot \iint \min \{ \mathbb{E} \eta_{\hat{\theta}}^{(n)}(\mathbf{u}, \mathbf{z}), \eta_0^{(n)}(\mathbf{u}, \mathbf{z}) \} d\mathbf{u} d\mathbf{z} \\
& \geq \frac{1}{64} \beta^2 (F(1+\alpha) + F(1-\alpha) - 2)^2 \cdot \left\{ 1 - \left[1 - \left(1 - \frac{1}{2} \chi^2(\mathbb{E} \eta_{\hat{\theta}}^{(n)}, \eta_0^{(n)}) \right)^2 \right]^{1/2} \right\},
\end{aligned}$$

concludes the proof of the theorem. \square

B Proof of Lemma 6.3, and further technical lemmas

Recall that for $J \in \mathcal{J}_K$ we take \hat{j} as the smallest $j \in J$ for which $\|z - Z_j\| \geq \|z - Z_k\|$ for all $k \in J$, and denote by \mathfrak{A} the σ -field generated by $(\hat{j}, Z_{\hat{j}})$. Further we set $\hat{J} := J \setminus \{\hat{j}\}$ and in the following denote

$$u_0 = \mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0$$

We need to show that for $\eta \geq 1$ and $K \geq 1 + (\lfloor 2\eta D \rfloor + 1)K^*$ we have that

$$\mathbb{E}[u_0^\eta | \mathfrak{A}] \leq C_3. \quad (\text{B.1})$$

Note that u_0 is the first component of the column vector \mathbf{u} which satisfies $\mathcal{M}_J(z)\mathbf{u} = \mathbf{e}_0$. Multiplying this system of linear equation by \mathbf{u}^\top from the left side we arrive at

$$u_0 = \sum_{j \in J} \left(\sum_{\kappa \in \mathcal{K}} u_\kappa \xi_\kappa(z, Z_j) \right)^2.$$

As we are seeking for a positive upper bound on u_0 we may assume that $u_0 > 0$. Dividing by u_0^2 yields that

$$\begin{aligned}
u_0 &= 1 / \left\{ \sum_{j \in J} \left(\sum_{\kappa \in \mathcal{K}} (u_\kappa / u_0) \xi_\kappa(z, Z_j) \right)^2 \right\} \\
&\leq 1 / \left\{ \inf_{P \in \mathcal{P}_1} \sum_{j \in J} P^2(z - Z_j) \right\} \\
&\leq 1 / \left\{ \inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}} P^2(z - Z_j) \right\}, \quad (\text{B.2})
\end{aligned}$$

where \mathcal{P}_1 denotes the set of all d -variate polynomials with the degree $\leq L$ which take on the value 1 at 0, and we used the notation $\hat{J} := J \setminus \{\hat{j}\}$ introduced above.

Lemma B.1. *Given \mathfrak{A} , the random variables Z_j , $j \in \widehat{\mathcal{J}}$, are conditionally independent and each Z_j has the conditional Lebesgue density*

$$f_{Z|\mathfrak{A}}(u) = f_Z(u) \cdot \mathbf{1}_{[0, \|z - Z_{\widehat{\mathcal{J}}}\|]}(\|z - u\|) / \int f_Z(u') \cdot \mathbf{1}_{[0, \|z - Z_{\widehat{\mathcal{J}}}\|]}(\|z - u'\|) du', \quad u \in \mathbb{R}^d.$$

Proof of Lemma B.1. Let Φ and Ω denote two probe functions and consider that

$$\begin{aligned} \mathbb{E}[\Phi(Z_j, j \in \widehat{\mathcal{J}}) \cdot \Omega(\widehat{j}, Z_{\widehat{\mathcal{J}}})] &= \sum_{j' \in \widehat{\mathcal{J}}} \mathbb{E} \mathbf{1}_{\{\widehat{j} = j'\}} \cdot \Phi(Z_k, k \in \widehat{\mathcal{J}} \setminus \{j'\}) \cdot \Omega(j', Z_{j'}) \\ &= \sum_{j' \in \widehat{\mathcal{J}}} \mathbb{E} \int \Omega(j', u) f_Z(u) \int \cdots \int \Phi(z_1, \dots, z_{K-1}) \prod_{k \in \widehat{\mathcal{J}} \setminus \{j'\}} f_Z(z_k) \mathbf{1}_{[0, \|u - z\|]}(\|z_k - z\|) dz_1 \cdots dz_{K-1} du. \end{aligned}$$

Now, putting $\Phi \equiv 1$ and changing Ω to $\Omega \cdot \tilde{\Phi}$, we deduce that

$$\mathbb{E}[\tilde{\Phi}(\widehat{j}, Z_{\widehat{\mathcal{J}}}) \cdot \Omega(\widehat{j}, Z_{\widehat{\mathcal{J}}})] = \sum_{j' \in \widehat{\mathcal{J}}} \mathbb{E} \int \Omega(j', u) \tilde{\Phi}(j', Z_{j'}) f_Z(u) \left(\int f_Z(\zeta) \mathbf{1}_{[0, \|u - z\|]}(\|\zeta - z\|) d\zeta \right)^{K-1} du,$$

so that

$$\begin{aligned} \mathbb{E}\{\Phi(Z_j, j \in \widehat{\mathcal{J}}) \mid \mathfrak{A}\} &= \int \cdots \int \Phi(z_1, \dots, z_{K-1}) \\ &\quad \cdot \prod_{k \in \widehat{\mathcal{J}} \setminus \{j'\}} \left(f_Z(z_k) \mathbf{1}_{[0, \|z - Z_{\widehat{\mathcal{J}}}\|]}(\|z_k - z\|) / \int f_Z(\zeta) \mathbf{1}_{[0, \|z - Z_{\widehat{\mathcal{J}}}\|]}(\|\zeta - z\|) d\zeta \right) dz_1 \cdots dz_{K-1}, \end{aligned}$$

holds almost surely, from what follows the claim of the lemma. \square

As an immediate consequence of Lemma B.1, the random variables $V_j := (z - Z_j) / \|z - Z_{\widehat{\mathcal{J}}}\|$, $j \in \widehat{\mathcal{J}}$, are conditionally i.i.d. and have the conditional Lebesgue density

$$f_{V|\mathfrak{A}}(v) = \mathbf{1}_{B_d(0,1)}(v) \cdot f_Z(z - v \|z - Z_{\widehat{\mathcal{J}}}\|) / \int_{B_d(0,1)} f_Z(z - v' \|z - Z_{\widehat{\mathcal{J}}}\|) dv', \quad v \in \mathbb{R}^d, \quad (\text{B.3})$$

given \mathfrak{A} . Furthermore, continuing in (B.2) we have that

$$\inf_{P \in \mathcal{P}_1} \sum_{j \in \widehat{\mathcal{J}}} P^2(z - Z_j) = \inf_{P \in \mathcal{P}_1} \sum_{j \in \widehat{\mathcal{J}}} P^2(V_j), \quad (\text{B.4})$$

since $P \in \mathcal{P}_1$ implies that $x \mapsto P(x / \|z - Z_{\widehat{\mathcal{J}}}\|) \in \mathcal{P}_1$ and vice versa. Now fix any $J^* \subseteq \widehat{\mathcal{J}}$ whose cardinality is denoted by $K^* < K$. The joint conditional density $f_{V|J^*|\mathfrak{A}}^*$ of $(V_j, j \in J^*)$ given \mathfrak{A} equals

$$f_{V|J^*|\mathfrak{A}}^*(v) = \prod_{j \in J^*} f_{V|\mathfrak{A}}(v_j), \quad v = (v_j, j \in J^*) \in (\mathbb{R}^d)^{K^*}.$$

Preparatory to further consideration we are interested in the conditional distribution of $\|V - w\|$ given \mathfrak{A} and $W := (V - w) / \|V - w\|$ when V has the conditional density $f_{V|J^*|\mathfrak{A}}^*$ given \mathfrak{A} ; and $w = (w_j, j \in J^*) \in (\mathbb{R}^d)^{K^*}$ is deterministic.

Lemma B.2. (a) If V has the conditional density $f_{V|\mathfrak{A}}^*$ given \mathfrak{A} , then the conditional Lebesgue density of $\|V - w\|$ given \mathfrak{A} and W equals

$$f_{\|V-w\|\mathfrak{A},W}^*(r) = \mathbf{1}_{(0,\infty)}(r) \cdot f_{V|\mathfrak{A}}^*(w + rW) r^{dK^*-1} / \int_0^\infty f_{V|\mathfrak{A}}^*(w + sW) s^{dK^*-1} ds, \quad r \in \mathbb{R}.$$

(b) Writing

$$\begin{aligned} \beta_1 &:= \min\{1/2, \rho'/(2\rho'')\}, \\ \beta_2 &:= \sqrt{2K^* + 2\beta_1^2}, \\ \beta_3 &:= \beta_2^{dK^*-1} \cdot \bar{\rho}^{K^*} \cdot d \cdot K^* / \{\rho^{K^*} \cdot \beta_1^{dK^*}\}, \end{aligned}$$

we assume that $\|w\| \leq \beta_1$. Then, the support of $f_{V,w|\mathfrak{A},W}^*$ is included in the interval $[0, \beta_2]$ and $f_{V,w|\mathfrak{A},W}^*$ is bounded from above by β_3 .

Proof of Lemma B.2. (a) For two probe functions Φ and Ω , we consider that

$$\begin{aligned} \mathbb{E}[\Phi(\|V - w\|)\Omega(W) | \mathfrak{A}] &= \int \Phi(\|v - w\|)\Omega((v - w)/\|v - w\|) f_{V|\mathfrak{A}}^*(v) dv \\ &= \mathbb{E}[\Phi(\|U - w\|)\Omega(\tilde{U}) f_{V|\mathfrak{A}}^*(U)] \cdot \pi^{dK^*/2} R^{dK^*} / \Gamma(dK^*/2 + 1) \\ &= \mathbb{E}\left[\Omega(\tilde{U}) \int_0^R \Phi(r) f_{V|\mathfrak{A}}^*(w + r\tilde{U}) r^{dK^*-1} dr\right] \cdot d \cdot K^* \cdot \pi^{dK^*/2} / \Gamma(dK^*/2 + 1), \end{aligned} \quad (\text{B.5})$$

where $\tilde{U} := (U - w)/\|U - w\|$; and $R > 0$ is sufficiently large such that the dK^* -dimensional ball around w with the radius R contains the closure of $B_d(0, 1)^{(K^*)}$ – and, hence, the support of $f_{V|\mathfrak{A}}^*$ – as a subset; and the random vector U is uniformly distributed on this ball. Therein we have used that $\|U - w\|$ and \tilde{U} are independent. Now evaluating (B.5) for $\Phi \equiv 1$ and Ω being replaced by $\tilde{\Phi} \cdot \Omega$, we deduce that

$$\tilde{\Phi}(\tilde{U}) = \int_0^R \Phi(r) f_{V|\mathfrak{A}}^*(w + r\tilde{U}) r^{dK^*-1} dr / \int_0^R f_{V|\mathfrak{A}}^*(w + s\tilde{U}) s^{dK^*-1} ds, \quad \text{a.s.},$$

so that

$$\mathbb{E}(\Phi(\|V - w\|) | W, \mathfrak{A}) = \tilde{\Phi}(W) = \int_0^R \Phi(r) f_{V|\mathfrak{A}}^*(w + rW) r^{dK^*-1} dr / \int_0^R f_{V|\mathfrak{A}}^*(w + sW) s^{dK^*-1} ds,$$

holds true almost surely. As the left side does not depend on R the equality remains valid for $R \rightarrow \infty$. That completes the proof of part (a).

(b) Part (a) yields that $\|w_j + rW_j\| \leq 1$ for all $j \in J^*$ with $W = (W_j, j \in J^*)$ whenever $f_{V,w|\mathfrak{A},W}^*(r) > 0$ since the support of $f_{V|\mathfrak{A}}^*$ is included in $B_d(0, 1)^{(K^*)}$. Thus, $r\|W_j\| \leq 1 + \|w_j\|$ holds for all $j \in J^*$ so that

$$r^2 = \sum_{j \in J^*} r^2 \|W_j\|^2 \leq \sum_{j \in J^*} (2 + 2\|w_j\|^2) = 2K^* + 2\beta_1^2.$$

Therefore the support of $f_{V,w|\mathfrak{A},W}^*$ is included in the interval $[0, \beta_2]$. Moreover, we have

$$\begin{aligned}
f_{V,w|\mathfrak{A},W}^*(r) &= \mathbf{1}_{[0,\beta_2]}(r) \left\{ \prod_{j \in J^*} \mathbf{1}_{B_d(0,1)}(w_j + rW_j) f_Z(z - (w_j + rW_j) \|z - Z_{\hat{J}}\|) \right\} r^{dK^*-1} \\
&/ \int_0^\infty \left\{ \prod_{j \in J^*} \mathbf{1}_{B_d(0,1)}(w_j + sW_j) f_Z(z - (w_j + sW_j) \|z - Z_{\hat{J}}\|) \right\} s^{dK^*-1} ds \\
&\leq \beta_2^{dK^*-1} \cdot \bar{\rho}^{K^*} / \left(\rho^{K^*} \int_0^{\beta_1} s^{dK^*-1} ds \right) \\
&= \beta_3,
\end{aligned}$$

where we have used that, for $s \in [0, \beta_1)$, it holds that $\|w_j + sW_j\| < 1$; that $\|w_j + sW_j\| \cdot \|z - Z_{\hat{J}}\| < \rho'$; and, hence, $z - (w_j + sW_j) \|z - Z_{\hat{J}}\| \in S$ as $z \in S^*$ and $Z_{\hat{J}} \in S$, for all $j \in J^*$. \square

These properties of $f_{V,w|\mathfrak{A},W}^*$ are essential to establish the following result.

Lemma B.3. *As above consider $J^* \subseteq \hat{J}$. Assume that $d \geq 2$; that $L \geq 1$; and that $K^* = \#\mathcal{K}(d, L) < K$. Define D as in (2.8), that is*

$$D := \sum_{\ell=1}^L \ell \cdot \#\{\kappa \in \mathbb{N}_0^d \mid \kappa_1 + \dots + \kappa_d = \ell\}.$$

By $\Xi_{d,L}(\mathbf{x})$ we denote the square matrix which consists of the row vectors $(\xi_\kappa(0, \mathbf{x}_j))_{\kappa \in \mathcal{K}(d,L)}$ for $j = 1, \dots, K^*$ where $\mathbf{x} := (\mathbf{x}_1, \dots, \mathbf{x}_{K^*}) \in (\mathbb{R}^d)^{K^*}$. Let us write $\vartheta_{d,L}(\mathbf{x})$ for the determinant of $\Xi_{d,L}(\mathbf{x})$. Let V_j , $j \in J^*$ be a conditionally i.i.d. random sample, drawn from $f_{V|\mathfrak{A}}^*$, given \mathfrak{A} . Write V for the dK^* -dimensional row vector $V := (V_j, j \in J^*)$. Then, the random matrix $\Xi_{d,L}(V)$ is invertible almost surely, and

$$\mathbb{P}[|\vartheta_{d,L}(V)| \leq \varepsilon \mid \mathfrak{A}] \leq C_1 \cdot \varepsilon^{1/D}, \quad \forall \varepsilon \in (0, \varepsilon_1),$$

almost surely for deterministic positive constants C_1 and ε_1 which only depend on $d, L, \rho, \bar{\rho}, \rho', \rho''$.

Proof of Lemma B.3. Clearly the function $\vartheta_{d,L}$ forms a polynomial in dK^* variables (namely the components of \mathbf{x}) with the degree of at most D . Note that $\vartheta_{d,L}$ is non-constant since it contains exactly $K^*! \geq 2$ distinct monomials where each of them has the coefficient of either 1 or -1 . Hence the zero set of $\vartheta_{d,L}$ has the dK^* -dimensional Lebesgue measure zero. It follows from there that $\vartheta_{d,L}(V)$ does not vanish and, thus, $\Xi_{d,L}(V)$ is invertible almost surely as the random vector V has the dK^* -dimensional Lebesgue density $f_{V|\mathfrak{A}}^*$.

We quantify the small ball probabilities of $|\vartheta_{d,L}(V)|$ around zero in terms of upper bounds. By $M > 0$ we denote the maximum of $|\vartheta_{d,L}|$ on the closure of the dK^* -dimensional ball with the center 0 and the radius $\beta_1/2$ where β_1, β_2 and β_3 are as in Lemma B.2. We specify w from Lemma B.2 as some element of that closed ball in which this maximum is taken on. Note that $\|w\| \leq \beta_1$; and that any non-negative, non-constant and continuous function takes on at least one maximum on any compact domain. Consider that

$$\mathbb{P}[|\vartheta_{d,L}(V)| \leq \varepsilon \mid \mathfrak{A}] = \mathbb{E}\{\mathbb{P}[|\vartheta_{d,L}(w + \|V - w\|W)| \leq \varepsilon \mid W, \mathfrak{A}] \mid \mathfrak{A}\}, \quad (\text{B.6})$$

where $W = (V - w)/\|V - w\|$. We apply the fundamental theorem of algebra to the univariate polynomial $r \mapsto \vartheta_{d,L}(w + rW)$ with the degree $D^* \leq D$; and we exploit that the absolute value of this polynomial takes on the value M at 0 so that

$$|\vartheta_{d,L}(w + rW)| \geq M \cdot \prod_{k=1}^{D^*} |1 - r/|\zeta_j||,$$

for all $r \geq 0$ where $\zeta_j, j = 1, \dots, D^*$, are the complex roots of the polynomial. Hence,

$$\begin{aligned} \mathbb{P}[|\vartheta_{d,L}(w + \|V - w\|W)| \leq \varepsilon \mid W, \mathfrak{A}] &\leq \mathbb{P}\left[\prod_{k=1}^{D^*} |1 - \|V - w\|/|\zeta_j|| \leq \varepsilon/M \mid W, \mathfrak{A}\right] \\ &\leq \sum_{k=1}^{D^*} \mathbb{P}\left[|\zeta_j| \cdot (1 + (\varepsilon/M)^{1/D^*}) \geq \|V - w\| \geq |\zeta_j| \cdot (1 - (\varepsilon/M)^{1/D^*}) \mid W, \mathfrak{A}\right], \end{aligned} \quad (\text{B.7})$$

for all $\varepsilon \in (0, M/2)$. If $|\zeta_j| > \beta_2/(1 - (1/2)^{1/D^*})$ then

$$\mathbb{P}\left[|\zeta_j| \cdot (1 + (\varepsilon/M)^{1/D^*}) \geq \|V - w\| \geq |\zeta_j| \cdot (1 - (\varepsilon/M)^{1/D^*}) \mid W, \mathfrak{A}\right] = 0.$$

Otherwise,

$$\mathbb{P}\left[|\zeta_j| \cdot (1 + (\varepsilon/M)^{1/D^*}) \geq \|V - w\| \geq |\zeta_j| \cdot (1 - (\varepsilon/M)^{1/D^*}) \mid W, \mathfrak{A}\right] \leq 2\beta_2 \beta_3 (\varepsilon/M)^{1/D} / (1 - (1/2)^{1/D}).$$

Therefore, the term (B.7) is bounded from above by

$$2D \beta_2 \beta_3 (\varepsilon/M)^{1/D} / (1 - (1/2)^{1/D}),$$

for all $\varepsilon \in (0, M/2)$. Inserting this upper bound into (B.6) finally completes the proof. \square

The inequality from Lemma B.3 can be extended to the term u_0 .

Lemma B.4. *Grant the assumptions and the notation from Lemma B.3. Then,*

$$\mathbb{P}\left[\inf_{P \in \mathcal{P}_1} \sum_{j \in J^*} P^2(V_j) \leq \varepsilon \mid \mathfrak{A}\right] \leq C_2 \cdot \varepsilon^{1/(2D)}, \quad \forall \varepsilon \in (0, \varepsilon_2),$$

for deterministic positive constants C_2 and ε_2 which only depend on $d, L, \rho, \bar{\rho}, \rho', \rho''$.

Proof of Lemma B.4. Consider that

$$\inf_{P \in \mathcal{P}_1} \sum_{j \in J^*} P^2(V_j) = \inf_{\alpha \in \mathbb{R}^{\mathcal{K}}, \alpha_0=1} \|\Xi_{d,L}(V)\alpha\|^2 = \inf_{\alpha \in \mathbb{R}^{\mathcal{K}}, \alpha_0=1} \alpha^\top \Xi_{d,L}(V)^\top \Xi_{d,L}(V)\alpha,$$

so that the above term is larger or equal to the smallest eigenvalue of the symmetric and positive definite matrix $\Xi_{d,L}(V)^\top \Xi_{d,L}(V)$. On the other hand, the largest eigenvalue of $\Xi_{d,L}(V)^\top \Xi_{d,L}(V)$ is bounded from above by the product of the spectral norms of $\Xi_{d,L}(V)^\top$ and $\Xi_{d,L}(V)$ and, thus, by $\|\Xi_{d,L}(V)\|_F^2$ where $\|\cdot\|_F$ denotes the Frobenius norm. We have

$$\begin{aligned} \|\Xi_{d,L}(V)\|_F^2 &= \sum_{j \in J^*} \sum_{\kappa \in \mathcal{K}} \xi_\kappa^2(0, V_j) \leq \sum_{j \in J^*} \sum_{\ell=0}^L \|V_j\|^{2\ell} \sum_{|\kappa|=\ell} \binom{\ell}{\kappa_1, \dots, \kappa_d} \prod_{k=1}^d (V_{j,k}^2 / \|V_j\|^2)^{\kappa_k} \\ &= \sum_{j \in J^*} \sum_{\ell=0}^L \|V_j\|^{2\ell} \leq (L+1) \cdot K^*, \end{aligned}$$

as $\|V_j\| \leq 1$ a.s., see (B.3). As the determinant of a symmetric matrix equals the product of all of its eigenvalues it follows that

$$\vartheta_{d,L}^2(V) = \det(\Xi_{d,L}(V)^\top \Xi_{d,L}(V)) \leq ((L+1)K^*)^{K^*-1} \cdot \inf_{P \in \mathcal{P}_1} \sum_{j \in J^*} P^2(V_j).$$

Then, Lemma B.3 yields that

$$\begin{aligned} \mathbb{P}\left[\inf_{P \in \mathcal{P}_1} \sum_{j \in J^*} P^2(V_j) \leq \varepsilon \mid \mathfrak{A}\right] &\leq \mathbb{P}\left[|\vartheta_{d,L}(V)| \leq \varepsilon^{1/2} ((L+1)K^*)^{1/2-K^*/2} \mid \mathfrak{A}\right] \\ &\leq C_1 ((L+1)K^*)^{1/(2D)-K^*/(2D)} \cdot \varepsilon^{1/(2D)}, \end{aligned}$$

for all $\varepsilon \in (0, \varepsilon_2)$ where the constant $\varepsilon_2 > 0$ only depends on $d, L, \rho, \bar{\rho}, \rho'$ and ρ'' . \square

Now we ready to bound the conditional moments of u_0 as stated in Lemma 6.3.

Proof of Lemma 6.3. Combining (B.2) and (B.4), we deduce that

$$\mathbb{E}\left[(\mathbf{e}_0^\top \mathcal{M}_J(z)^{-1} \mathbf{e}_0)^\eta \mid \mathfrak{A}\right] \leq \int_0^\infty \mathbb{P}\left[\inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}} P^2(V_j) < t^{-1/\eta} \mid \mathfrak{A}\right] dt. \quad (\text{B.8})$$

There exist pairwise disjoint subsets $\hat{J}_k, k = 1, \dots, m$, of \hat{J} with $\#\hat{J}_k = K^*$ for all $k = 1, \dots, m$, where $m > 2\eta D$. Clearly,

$$\inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}} P^2(V_j) \geq \sum_{k=1}^m \inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}_k} P^2(V_j) \geq \max_{k=1, \dots, m} \inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}_k} P^2(V_j),$$

where the random variables $\inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}_k} P^2(V_j), k = 1, \dots, m$, are i.i.d. (conditionally on \mathfrak{A}). By Lemma B.4, the right side of (B.8) is smaller or equal to

$$\begin{aligned} &\int_0^\infty \mathbb{P}\left[\max_{k=1, \dots, m} \inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}_k} P^2(V_j) < t^{-1/\eta} \mid \mathfrak{A}\right] dt \\ &\leq \varepsilon_2^{-\eta} + \int_{t > \varepsilon_2^{-\eta}} \left\{ \mathbb{P}\left[\inf_{P \in \mathcal{P}_1} \sum_{j \in \hat{J}_1} P^2(V_j) < t^{-1/\eta} \mid \mathfrak{A}\right] \right\}^m dt \\ &\leq \varepsilon_2^{-\eta} + C_2^m \int_{t > \varepsilon_2^{-\eta}} t^{-m/(2\eta D)} dt \\ &\leq \varepsilon_2^{-\eta} + C_2^m \frac{2\eta D}{m - 2\eta D} \varepsilon_2^{-\eta + m/(2D)}, \end{aligned}$$

so that the lemma has been shown. \square

Proof of Remark 2: For any $J \in \mathcal{J}_K$ and integer $\ell \geq 1$, we deduce by Fubini's theorem that

$$\begin{aligned} \mathbb{E}\lambda^\ell(C(J)) &= \mathbb{E}\left(\int_{S^*} \mathbf{1}_{C(J)}(x) dx\right)^\ell \\ &= \int_{S^*} \cdots \int_{S^*} \mathbb{P}[x_1, \dots, x_\ell \in C(J)] dx_1 \cdots dx_\ell \\ &= \int_{S^*} \cdots \int_{S^*} \mathbb{P}[\|x_m - Z_j\| < \|x_m - Z_k\|, \forall m = 1, \dots, \ell, j \in J, k \notin J] dx_1 \cdots dx_\ell \\ &= \int_{S^*} \cdots \int_{S^*} \mathbb{E} \mathbb{P}_Z^K\left(\bigcap_{m=1}^{\ell} B_d(x_m, \min_{k \notin J} \|x_m - Z_k\|)\right) dx_1 \cdots dx_\ell, \end{aligned}$$

where $B(c, r)$ denotes the Euclidean ball around the center c with the radius r ; and \mathbb{P}_Z stands for the image measure of Z_1 . For some deterministic sequence $(\alpha_n)_n \downarrow 0$ we consider that

$$\begin{aligned} \mathbb{P}\left[\min_{m=1,\dots,\ell} \min_{k \notin J} \|x_m - Z_k\| > \alpha_n\right] &= \mathbb{P}[\|x_m - Z_k\| > \alpha_n, \forall m = 1, \dots, \ell, k \notin J] \\ &= \mathbb{P}_Z^{n-K} \left(\bigcap_{m=1}^{\ell} \mathbb{R}^d \setminus B(x_m, \alpha_n) \right) = \left\{ 1 - \mathbb{P}_Z \left(\bigcup_{m=1}^{\ell} B(x_m, \alpha_n) \right) \right\}^{n-K} \\ &\geq \left\{ 1 - \sum_{m=1}^{\ell} \mathbb{P}_Z(B(x_m, \alpha_n)) \right\}^{n-K} \geq \left\{ 1 - \ell \cdot \bar{\rho} \cdot \frac{\pi^{d/2}}{\Gamma(d/2 + 1)} \cdot \alpha_n^d \right\}^{n-K}, \end{aligned}$$

for n sufficiently large. We write $\mathcal{E}_n(x)$, $x = (x_1, \dots, x_\ell)$, for the event that $\min_{m=1,\dots,\ell} \min_{k \notin J} \|x_m - Z_k\| > \alpha_n$. We fix that $\alpha_n \asymp n^{-1/d}$ so that

$$\liminf_{n \rightarrow \infty} \inf_{x \in (S^*)^\ell} P(\mathcal{E}_n(x)) > 0.$$

Furthermore we introduce the set

$$\mathcal{S}_n := \{x = (x_1, \dots, x_\ell) \in (S^*)^\ell : \|x_1 - x_m\| < \alpha_n/2, \forall m = 1, \dots, \ell\}.$$

On the event $\mathcal{E}_n(x)$ the ball $B(x_1, \alpha_n/2)$ is included in $\bigcap_{m=1}^{\ell} B_d(x_m, \min_{k \notin J} \|x_m - Z_k\|)$ and in S (for n sufficiently large) as a subset for any $x \in \mathcal{S}_n$. Therefore,

$$\begin{aligned} \mathbb{E} \lambda^\ell(C(J)) &\geq \left(\frac{\pi^{d/2}}{2^d \Gamma(d/2 + 1)} \right)^K \cdot \rho^K \cdot \alpha_n^{Kd} \cdot \int \cdots \int_{\mathcal{S}_n} \mathbb{P}(\mathcal{E}_n(x)) dx_1 \cdots dx_\ell \\ &\geq \left(\frac{\pi^{d/2}}{2^d \Gamma(d/2 + 1)} \right)^K \cdot \rho^K \cdot \alpha_n^{Kd} \cdot \inf_{x \in (S^*)^\ell} \mathbb{P}(\mathcal{E}_n(x)) \cdot \int \cdots \int_{\mathcal{S}_n} dx_1 \cdots dx_\ell. \end{aligned}$$

The ℓd -dimensional Lebesgue measure of \mathcal{S}_n shall be bounded from below by $\text{const.} \cdot \alpha_n^{d(\ell-1)}$ as n tends to ∞ . Then, unfortunately, the ℓ th moment of $\lambda(C(J))$ cannot attain the desired rate $n^{-\ell K}$ but only $n^{-\ell-K+1}$. \square

C Proofs for Section 3

Proof of Theorem 4. We show below that the estimators $\hat{\Psi}_j$ of the Fourier coefficients $\langle h * \tilde{f}_\delta, \phi_j \rangle$ resulting from (2.4) satisfy for each integer $l \in \mathbb{N}$ that

$$\sup_{j \in \mathbf{J}^{(n)}} \mathbb{E}[|\hat{\Psi}_j - \Psi_j|^2] \lesssim n^{-1} + (J_n \cdot n^{-1/d})^{2l}, \quad (\text{C.1})$$

If we take l large enough, for our choice of J_n the first term dominates as $J_n \lesssim n^{1/d-1/(2l)}$ and $l > d/2$. By Parseval's identity, standard risk bounds for series estimators under the Sobolve condition (3.3) lead to

$$\mathbb{E} \left[\int_{[-\pi, \pi]^d} |\hat{h}(x) - h(x)|^2 dx \right] \lesssim J_n^{-2\alpha} + J_n^d \cdot J_n^{2\gamma} \cdot n^{-1},$$

from which the rate (3.4) follows upon inserting our choice of J_n .

Concerning (C.1): Since $\|\phi_{\mathbf{j}}\|_{\infty} \leq 1$ the constant C_V in Theorem 1 is set equal to the variance $\text{var } \epsilon$; whereas

$$\|\partial_{i_1} \cdots \partial_{i_l} G_{\mathbf{j}}\|_{\infty} \leq \sum_{M \subseteq \{1, \dots, l\}} \|\partial_{i_M} (h * \tilde{f}_{\delta})\|_{\infty} \cdot J_n^{l - \#M},$$

for all $\mathbf{j} \in \mathbf{J}^{(n)} := \{-J_n, \dots, J_n\}^d$ where $\partial_{i_1}, \dots, \partial_{i_l}$, $i_1, \dots, i_l \in \{1, \dots, d\}$, denote the partial differential operators with respect to the components z_{i_1}, \dots, z_{i_l} where we allow for coincidence of the i_1, \dots, i_l , and ∂_{i_M} stands for the chaining of all ∂_{i_k} , $k \in M$. Therein note the symmetry of the partial derivatives. By the Cauchy-Schwarz inequality and the Fourier representation of the Sobolev norm, we consider that

$$\|\partial_{i_M} (h * \tilde{f}_{\delta})\|_{\infty} = \|(\partial_{i_M} h) * \tilde{f}_{\delta}\|_{\infty} \leq \left(\int |\partial_{i_M} h|^2 \right)^{1/2} \cdot \left(\int |f_{\delta}|^2 \right)^{1/2} \leq (2\pi)^{-d/2} \cdot C_{\alpha}^{1/2} \cdot \left(\int |f_{\delta}|^2 \right)^{1/2},$$

when $l \leq \alpha$, so that

$$\|\partial_{i_1} \cdots \partial_{i_l} G_{\mathbf{j}}\|_{\infty} \leq C_H = C_H(\mathbf{j}, l) := (2\pi)^{-d/2} \cdot C_{\alpha}^{1/2} \cdot \left(\int |f_{\delta}|^2 \right)^{1/2} \cdot (1 + J_n)^l,$$

for $\mathbf{j} \in \mathbf{J}^{(n)}$ when $l \leq \lceil \alpha \rceil - 1$. Then, $G_{\mathbf{j}} \in \mathcal{G}(l - 1, 1, C_H(\mathbf{j}, l))$ and Theorem 1 yields (C.1). \square

D Additional simulations

In $d = 2$ we consider a further simulation setting with regressions function

$$f_3(z_1, z_2) = 5 z_1 (z_2^2 + 1) + 10 \cos(z_1/z_2)$$

and $Y_i = f_j(Z_{i,1}, Z_{i,2}) + 0.2 \varepsilon_i$, otherwise the parameters are the same as before. Results are displayed in Table 3.

Higher values of K perform better in this setting, but the method does not seem to depend sensitively on the choice of K for $L = 1$ as long as K is not chosen too small.

$n = 100$	$L = 0$	K	1	3	4	6	8	10
		\sqrt{n} · BIAS	0.0140	0.0280	0.0343	0.0457	0.0557	0.0650
		\sqrt{n} · STDV	0.0988	0.0949	0.0961	0.0995	0.1035	0.1073
		\sqrt{n} · RMSE	0.0998	0.0990	0.1021	0.1095	0.1175	0.1254
	$L = 1$	K	4	8	10	12	14	16
		\sqrt{n} · BIAS	-0.0002	-0.0011	-0.0018	-0.0035	-0.0060	-0.0091
		\sqrt{n} · STDV	0.1539	0.0950	0.0942	0.0949	0.0962	0.0978
		\sqrt{n} · RMSE	0.1539	0.0950	0.0942	0.0949	0.0964	0.0982
$n = 1000$	$L = 0$	K	1	3	4	6	8	10
		\sqrt{n} · BIAS	0.0121	0.0145	0.0170	0.0221	0.0267	0.0320
		\sqrt{n} · STDV	0.1242	0.1160	0.1145	0.1126	0.1122	0.1116
		\sqrt{n} · RMSE	0.1248	0.1169	0.1157	0.1148	0.1153	0.1161
	$L = 1$	K	4	8	10	12	14	16
		\sqrt{n} · BIAS	0.0157	0.0098	0.0111	0.0124	0.0138	0.0153
		\sqrt{n} · STDV	0.1758	0.1165	0.1139	0.1123	0.1111	0.1103
		\sqrt{n} · RMSE	0.1765	0.1169	0.1145	0.1130	0.1120	0.1114

Table 3: Simulation scenario with f_3 in $d = 2$