

CONVERGENCE RATE OF THE Q -CURVATURE FLOW

PAK TUNG HO AND SANGHOON LEE

ABSTRACT. Carlotto, Chodosh and Rubinstein have studied the convergence rate of the Yamabe flow. Inspired by their result, we study the convergence rate of the Q -curvature flow in this paper. In particular, we provide an example of a slowly converging Q_6 -curvature flow in dimension 6, in contrast to the dimension 2 case, where the Q -curvature flow always converges exponentially.

1. INTRODUCTION

Let (M, g_0) be a closed (i.e. compact without boundary) Riemannian manifold of dimension $n \geq 3$. As a generalization of the Uniformization Theorem, the *Yamabe problem* is to find a metric conformal to g_0 such that its scalar curvature R_g is constant. This was solved by Aubin [2], Trudinger [45] and Schoen [40].

The *Yamabe flow* is a geometric flow introduced to study the Yamabe problem. It is defined as

$$(1.1) \quad \frac{\partial}{\partial t} g(t) = -(R_{g(t)} - \bar{R}_{g(t)})g(t),$$

where $\bar{R}_{g(t)}$ is the average of the scalar curvature of $g(t)$:

$$\bar{R}_{g(t)} = \frac{\int_M R_{g(t)} dV_{g(t)}}{\int_M dV_{g(t)}}.$$

The existence and convergence of the Yamabe flow has been studied in [6, 7, 15, 42, 48]. See also [3, 13, 16, 21, 22, 27, 37, 41] and references therein for results related to the Yamabe flow.

In [9], Carlotto, Chodosh and Rubinstein studied the rate of convergence of the Yamabe flow (1.1). They proved the following: (see Theorem 1 in [9])

Theorem 1.1. *Assume $g(t)$ is a solution of the Yamabe flow (1.1) that converges in $C^{2,\alpha}(M, g_\infty)$ to g_∞ as $t \rightarrow \infty$ for some $\alpha \in (0, 1)$. Then there is a $\delta > 0$ depending only on g_∞ such that:*

(i) *If g_∞ is an integrable critical point, then the convergence occurs at an exponential rate, that is*

$$\|g(t) - g_\infty\|_{C^{2,\alpha}(M, g_\infty)} \leq Ce^{-\delta t}$$

for some constant $C > 0$ depending on $g(0)$.

(ii) *In general, the rate of convergence cannot be worse than polynomial, that is*

$$\|g(t) - g_\infty\|_{C^{2,\alpha}(M, g_\infty)} \leq C(1+t)^{-\delta}$$

for some constant $C > 0$ depending on $g(0)$.

Date: 5th April, 2024.

2000 Mathematics Subject Classification. Primary 53E99; Secondary 53C18, 35R01.

Theorem 1.2. *Assume that g_∞ is a nonintegrable critical point of the Yamabe energy with order of integrability $p \geq 3$. If g_∞ satisfies the Adams-Simon positive condition AS_p , then there exists metric $g(0)$ conformal to g_∞ such that the solution $g(t)$ of the Yamabe flow (1.1) starting from $g(0)$ exists for all time and converges in $C^\infty(M, g_\infty)$ to g_∞ as $t \rightarrow \infty$. The convergence occurs “slowly” in the sense that*

$$C^{-1}(1+t)^{-\frac{1}{p-1}} \leq \|g(t) - g_\infty\|_{C^{2,\alpha}(M, g_\infty)} \leq C(1+t)^{-\frac{1}{p-2}}$$

for some constant $C > 0$.

We refer the readers to [9, Definition 8] and [9, Definition 10] respectively for the precise definitions of integrable critical point and Adams-Simon positive condition AS_p . In [9], examples of g_∞ satisfying the AS_p , for some $p \geq 3$, have been found. Hence, there exist examples of the Yamabe flow (1.1) which does not converge exponentially. To the best of the authors’ knowledge, this is the first geometric flow found not to have exponential convergence. Inspired by [9], the convergence rate of other geometric flows have been studied. In [28], the convergence rates of the Yamabe flow with boundary and the conformal mean curvature flow on manifolds with boundary have been studied. In [29], the convergence rate of the weighted Yamabe flow on the smooth metric measure spaces has been studied.

Suppose that (M, g) is a closed 4-dimensional Riemannian manifold. The Q -curvature is defined by

$$(1.2) \quad Q_g = -\frac{1}{6}(\Delta_g R_g - R_g^2 + 3|Ric_g|^2),$$

where R_g is the scalar curvature and Ric_g is the Ricci curvature tensor. The Paneitz operator is defined by

$$(1.3) \quad P_g f = \Delta_g^2 f + \delta \left[\left(\frac{2}{3} R_g g - 2 Ric_g \right) df \right]$$

for $f \in C^\infty(M)$. Here δ denotes the divergence and d the differential. Under a conformal change of the metric $\tilde{g} = e^{2w}g$, the Q -curvature transforms according to

$$(1.4) \quad P_g w + Q_g = Q_{\tilde{g}} e^{4w}.$$

The Chern-Gauss-Bonnet theorem asserts that

$$\int_M Q_g dV_g + \int_M \frac{1}{4} |W_g|^2 dV_g = 8\pi^2 \chi(M)$$

where $\chi(M)$ is the Euler-characteristics of M , and W_g is the Weyl tensor. Note that W_g is pointwise conformally invariant in the sense that $|W_{\tilde{g}}|^2 dV_{\tilde{g}} = |W_g|^2 dV_g$ whenever $\tilde{g} = e^{2w}g$. Hence, the total Q -curvature

$$\int_M Q_g dV_g$$

is conformally invariant.

Now suppose that (M, g) is a closed Riemannian manifold of dimension n , where n is even. In [19, 20], Fefferman and Graham constructed a conformally invariant formally invariant self-adjoint operator P_g with the leading term $(-\Delta_g)^{\frac{n}{2}}$. Moreover, there is a curvature quantity Q_g which transforms according to

$$(1.5) \quad P_g w + Q_g = Q_{\tilde{g}} e^{nw}$$

whenever $\tilde{g} = e^{2w}g$. This implies that the total Q -curvature

$$\int_M Q_g dV_g$$

is conformally invariant. Indeed, we have

$$P_g = -\Delta_g \quad \text{and} \quad Q_g = K_g$$

when $n = 2$, where K_g is the Gauss curvature. In particular, it follows from the Gauss-Bonnet theorem that

$$\int_M K_g dV_g = 2\pi\chi(M)$$

is a conformally invariant. Moreover, when $n = 4$, the operator P_g and the curvature quantity Q_g reduce to the Paneitz operator and the Q -curvature defined in (1.3) and (1.2) respectively.

Uniformization theorem for Q -curvature and the problem of prescribing Q -curvature have been studied. See [8, 10, 12, 17, 23, 24, 26, 38, 39, 47] and the references therein for the related results. In particular, the Q -curvature flow has been defined to study the Uniformization theorem for Q -curvature:

$$(1.6) \quad \frac{\partial}{\partial t}g(t) = -(Q_{g(t)} - \overline{Q}_{g(t)})g(t),$$

where $Q_{g(t)}$ is the Q -curvature of $g(t)$, and $\overline{Q}_{g(t)}$ is its average:

$$(1.7) \quad \overline{Q}_{g(t)} = \frac{\int_M Q_{g(t)} dV_{g(t)}}{\int_M dV_{g(t)}}.$$

The Q -curvature (1.6) on the sphere S^n has been studied in [5] when $n = 4$ and in [25] for general n . In particular, Brendle proved in [8] the following:

Theorem 1.3. *Assume that the Paneitz operator of (M, g) is weakly positive with kernel consisting of the constant functions. Moreover, assume that*

$$\int_M Q_g dV_g < (n-1)\omega_n$$

where ω_n is the volume of standard sphere S^n . Then the Q -curvature flow (1.6) exists for all times and converges to a metric of constant Q -curvature.

We remark that [8, Theorem 1.1] is more general than Theorem 1.3. Indeed, Brendle in [8] proved the existence and convergence of the prescribed Q -curvature flow under the same assumptions in Theorem 1.3.

Inspired by the results of Carlotto, Chodosh and Rubinstein about the convergence rate of Yamabe flow, i.e. Theorems 1.1 and 1.2 mentioned above, we study in this paper the convergence rate of the Q -curvature flow (1.6).

The following theorems are the main results in this paper:

Theorem 1.4. *Assume that $g(t)$ is a solution to the Q -curvature flow that is converging in $C^{n,\alpha}(M, g_\infty)$ to g_∞ as $t \rightarrow \infty$ for some $\alpha \in (0, 1)$. Then, there is $\delta > 0$ depending only on f and g_∞ such that*

(i) *If g_∞ is an integrable critical point, then the convergence occurs at an exponential rate*

$$\|g(t) - g_\infty\|_{C^{n,\alpha}(M, g_\infty)} \leq Ce^{-\delta t},$$

for some constant $C > 0$ depending on $g(0)$.

(ii) In general, the convergence cannot be worse than a polynomial rate

$$\|g(t) - g_\infty\|_{C^{n,\alpha}(M,g_\infty)} \leq C(1+t)^{-\delta},$$

for some constant $C > 0$ depending on $g(0)$.

Theorem 1.5. *Assume that g_∞ is a non-integrable critical point of the functional E with order of integrability $p \geq 3$. If g_∞ satisfies the Adam-Simon positivity condition AS_p , then there exists a metric $g(0)$ conformal to g_∞ such that the Q -curvature flow $g(t)$ starting from $g(0)$ exists for all time and converges in $C^\infty(M, g_\infty)$ to g_∞ as $t \rightarrow \infty$. The convergence occurs “slowly” in the sense that*

$$C(1+t)^{-\frac{1}{p-2}} \leq \|g(t) - g_\infty\|_{C^n(M,g_\infty)} \leq C(1+t)^{-\frac{1}{p-2}}$$

for some constant $C > 0$.

The precise definitions of integrable critical point and the Adam-Simon positivity condition AS_p can be found in Section 2. In Section 3, we prove Theorem 1.4, In Section 4, we prove Theorem 1.5. In Section 5, we construct an example of Riemannian manifold that satisfies the condition AS_3 . This allows us to conclude that there exists a Q -curvature flow converges exactly at a polynomial rate described in Theorem 1.5. In Section 6, we compare low dimensional cases when $n = 2, 4, 6$.

2. THE Q -CURVATURE FLOW

Suppose that n is even. Let M be a compact n -dimensional manifold equipped with the Riemannian metric g_∞ such that its volume satisfies $\int_M dV_{g_\infty} = 1$. Along the Q -curvature flow (1.6), the volume $\int_M dV_{g(t)}$ is preserved. Indeed, it follows from (1.6) and (1.7) that

$$(2.1) \quad \frac{d}{dt} \left(\int_M dV_{g(t)} \right) = \int_M \frac{\partial}{\partial t} (dV_{g(t)}) = -\frac{n}{2} \int_M (Q_{g(t)} - \bar{Q}_{g(t)}) dV_{g(t)} = 0.$$

Hence, if we assume $\int_M dV_{g(0)} = 1$, then

$$\int_M dV_{g(t)} = 1 \quad \text{for all } t \geq 0.$$

Since the flow (1.6) preserves the conformal structure, we can write $g(t) = e^{2u(t)}g_\infty$. Therefore, (1.6) can be written as

$$(2.2) \quad \frac{\partial}{\partial t} u(t) = -\frac{1}{2}(Q_{g(t)} - \bar{Q}_{g(t)}).$$

Since $g(t) = e^{2u(t)}g_\infty$, it follows from (1.5) that

$$(2.3) \quad P_{g_\infty} u(t) + Q_{g_\infty} = Q_{g(t)} e^{nu(t)}.$$

We define the functional

$$E(w) = \int_M \left(\frac{n}{2} w P_{g_\infty} w + n Q_{g_\infty} w \right) dV_{g_\infty} - \left(\int_M Q_{g_\infty} dV_{g_\infty} \right) \log \left(\int_M e^{nw} dV_{g_\infty} \right).$$

Using these, we find

$$(2.4) \quad \frac{d}{dt} E(u(t)) = -\frac{n}{2} \int_M (Q_{g(t)} - \bar{Q}_{g(t)})^2 dV_{g(t)}$$

along the Q -curvature flow.

Consider the *unit volume conformal class* associated to g_∞ :

$$[g_\infty]_1 = \left\{ e^{2w} g_\infty : w \in C^{n,\alpha}(M), \int_M e^{nw} dV_{g_\infty} = 1 \right\}.$$

For $k \in \mathbb{N}$, we denote the k -th differential of the functional E on $[g_\infty]_1$ at the point w in the directions v_1, \dots, v_k by

$$D^k E(w)[v_1, \dots, v_k].$$

As we will see below, the functional $v \mapsto D^k E(w)[v_1, \dots, v_{k-1}, v]$ is in the image of $L^2(M, g_\infty)$ under the natural embedding onto $C^{n,\alpha}(M, g_\infty)'$. Therefore, we will also write

$$D^k E(w)[v_1, \dots, v_{k-1}]$$

for this element of $L^2(M, g_\infty)$. When $k = 1$, we will drop the (second) brackets, and thus consider $Df(w) \in L^2(M, g_\infty)$.

We may write the differential of E restricted to $[g_\infty]_1$ as

$$\begin{aligned} \frac{1}{2} DE(w)[v] &= n \int_M (P_{g_\infty} w + Q_{g_\infty} - \overline{Q}_{g_\infty} e^{nw}) v dV_{g_\infty} \\ &= n \int_M (Q_{g_w} - \overline{Q}_{g_w}) e^{nw} v dV_{g_\infty} \end{aligned}$$

where $g_w = e^{2w} g_\infty$, where we have used the facts that $\int_M dV_{g_\infty} = \int_M dV_{g_w} = 1$ and the total Q -curvature is conformally invariant:

$$\int_M Q_{g_\infty} dV_{g_\infty} = \int_M Q_{g_w} dV_{g_w}.$$

Thus, a metric g_∞ is a critical point for the energy E restricted to $[g_\infty]_1$ exactly when g_∞ has constant Q -curvature. Regarded as an element of $L^2(M, g_\infty)$, we have that

$$(2.5) \quad \frac{1}{2} DE(w) = P_{g_\infty} w + Q_{g_\infty} - \overline{Q}_{g_\infty} e^{nw}.$$

From now on, we assume that g_∞ has constant Q -curvature. We denote by \mathcal{CQC}_1 the set of unit volume, constant Q -curvature metrics in $[g_\infty]_1$. We find

$$(2.6) \quad \frac{1}{2} D^2 E(g_\infty)[v, w] = \frac{1}{2} \frac{d}{dt} (DE(tv)[w]) \Big|_{t=0} = n \int_M (P_{g_\infty} v - nQ_{g_\infty} v) w dV_{g_\infty}.$$

Therefore, if we define \mathcal{L}_∞ by means of the formula

$$\frac{1}{n} \int_M w \mathcal{L}_\infty v dV_{g_\infty} := \frac{1}{2} D^2 E(g_\infty)[v, w],$$

then, by (2.6), we can obtain

$$\mathcal{L}_\infty v = P_{g_\infty} v - nQ_{g_\infty} v.$$

We define $\Lambda_0 := \ker \mathcal{L}_\infty \subset L^2(M, g_\infty)$.

Note that $P_{g_\infty} - nQ_{g_\infty}$ is a self-adjoint operator with leading term $(-\Delta_{g_\infty})^{\frac{n}{2}}$. It follows from a classical theorem of spectral theory (cf. [33]) that Λ_0 is finite dimensional, since it is the eigenspace of the operator $P_{g_\infty} - nQ_{g_\infty}$ for the zero eigenvalue. We will write Λ_0^\perp for the $L^2(M, g_\infty)$ -orthogonal complement.

It is crucial throughout this work that the functional E is an analytic map in the sense of [49, Definition 8.8]. More precisely, fix a metric g_∞ then the functional E is an analytic functional on $C^{n,\alpha}(M, g_\infty)$ in the sense that for each $w_0 \in C^{n,\alpha}(M, g_\infty)$, there is an $\epsilon > 0$ and bounded multilinear operators

$$E^{(k)} : C^{n,\alpha}(M, g_\infty)^{\times k} \rightarrow \mathbb{R} \text{ for each } k \geq 0$$

such that if $\|w - w_0\|_{C^{n,\alpha}} < \epsilon$, then $\sum_{k=0}^{\infty} \|E^{(k)}\| \cdot \|w - w_0\|_{C^{2n,\alpha}}^k < \infty$ and

$$E(w) = \sum_{k=0}^{\infty} E^{(k)}(\underbrace{w - w_0, \dots, w - w_0}_{k\text{-times}}) \text{ in } C^{n,\alpha}(M, g_\infty).$$

It is not hard to verify this, by simply expanding the denominator of E in a power series around $\log\left(\int_M e^{nw_0} dV_{g_\infty}\right)$ and noting that the other terms in E are already linear or bilinear functions in w .

We need the following Proposition from [44, Section 3]; which can be established with the help of the implicit function theorem:

Proposition 2.1. *There is $\epsilon > 0$ and an analytic map $\Phi : \Lambda_0 \cap \{v : \|v\|_{L^2} < \epsilon\} \rightarrow C^{n,\alpha}(M, g_\infty) \cap \Lambda_0^\perp$ such that $\Phi(0) = 0$, $D\Phi(0) = 0$,*

$$(2.7) \quad \sup_{\substack{\|v\|_{L^2} < \epsilon, \\ \|w\|_{L^2} \leq 1}} \|D\Phi(v)[w]\|_{L^2} < 1,$$

and so that defining $\Psi(v) = v + \Phi(v)$, we have that $\int_M e^{2\Psi(v)} dV_{g_\infty} = 1$ and

$$\text{proj}_{\Lambda_0^\perp}[DE(\Psi(v))] = \text{proj}_{\Lambda_0^\perp}\left[\left(Q_{e^{2\Psi(v)}g_\infty} - \overline{Q}_{e^{2\Psi(v)}g_\infty}\right)e^{n\Psi(v)}\right] = 0.$$

Furthermore

$$\text{proj}_{\Lambda_0}[DE(\Psi(v))] = \text{proj}_{\Lambda_0}\left[\left(Q_{e^{2\Psi(v)}g_\infty} - \overline{Q}_{e^{2\Psi(v)}g_\infty}\right)e^{n\Psi(v)}\right] = DF,$$

where $F : \Lambda_0 \cap \{v : \|v\|_{L^2} \leq \epsilon\} \rightarrow \mathbb{R}$ is defined by $F(v) = E(\Psi(v))$. Finally, the intersection of \mathcal{CQC}_1 with a small $C^{n,\alpha}(M, g_\infty)$ -neighborhood of 0 coincides with

$$\mathcal{S}_0 := \{\Psi(v) : v \in \Lambda_0, \|v\|_{L^2} < \epsilon, DF(v) = 0\},$$

which is a real analytic subvariety (possibly singular) of the following $(\dim \Lambda_0)$ -dimensional real analytic submanifold of $C^{n,\alpha}(M, g_\infty)$:

$$\mathcal{S} : \{\Psi(v) : v \in \Lambda_0, \|v\|_{L^2} < \epsilon\}.$$

We will refer to \mathcal{S} as the *natural constraint* for the problem.

Definition 2.2. For $g_\infty \in \mathcal{CQC}_1$, we say that g_∞ is *integrable* if for all $v \in \Lambda_0$, there is a path $w(t) \in C^n((-\epsilon, \epsilon) \times M, g_\infty)$ such that $e^{2w(t)}g_\infty \in \mathcal{CQC}_1$ and $w(0) = 0$, $w'(0) = v$. Equivalently, g_∞ is integrable if and only if \mathcal{CQC}_1 agrees with \mathcal{S} in a small neighborhood of 0 in $C^{n,\alpha}(M, g_\infty)$.

We remark that the integrability defined in Definition 2.2 is equivalent to the functional F (as defined in Proposition 2.1) being constant in a neighborhood of 0 inside Λ_0 [1, Lemma 1].

Definition 2.3. If $\Lambda_0 = 0$, i.e. if \mathcal{L}_∞ is injective, then we call g_∞ a *nondegenerate critical point*. On the other hand, if Λ_0 is nonempty, we call g_∞ *degenerate*.

Note that if g_∞ is a nondegenerate critical point, then g_∞ is automatically integrable in the above sense.

Now suppose that g_∞ is a nonintegrable critical point. Because $F(v) = E(\Psi(v))$, defined in Proposition 2.1, is analytic, we may expand it in a power series

$$F(v) = F(0) + \sum_{j \geq p} F_j(v)$$

where F_j is a degree- j homogeneous polynomial on Λ_0 and p is chosen so that F_p is nonzero. We will call p the *order of integrability* of g_∞ . We will also need a further hypothesis for nonintegrable critical points introduced in [1].

Definition 2.4. We say that g_∞ satisfies the *Adams-Simon positivity condition*, AS_p for short (here p is the order of integrability of g_∞), if it is nonintegrable and $F_p|_{\mathbb{S}^k}$ attains a positive maximum for some $\hat{v} \in \mathbb{S}^k \subset \Lambda_0$. Recall that F_p is the lowest-degree nonconstant term in the power series expansion of $F(v)$ around 0 and \mathbb{S}^k is the unit sphere in Λ_0 .

An important observation is that when the order of integrability p is odd, the Adams-Simon positivity condition is always satisfied. Moreover the order of integrability (at a critical point of E) always satisfies $p \geq 3$. In the next Proposition, we compute F_3 term.

Proposition 2.5. *We have the following identity:*

$$(2.8) \quad F_3(v) = -2n^2 \int_M Q_{g_\infty} v^3 dV_{g_\infty}.$$

Proof. We follow the computation in [9, Appendix A]. It is easy to see that $F_1(v) = F_2(v) = 0$. To compute $D^3F(0)$, we may in fact compute $D^3\tilde{F}(0)$, where $\tilde{F} : \Lambda_0 \rightarrow \mathbb{R}$ is defined by $\tilde{F}(v) = E(v)$. We first compute D^3F :

$$(2.9) \quad \begin{aligned} D^3F(w)[v, u, z] &= D^3E(\Psi(w))[D\Psi(w)[v], D\Psi(w)[u], D\Psi(w)[z]] \\ &\quad + D^2E(\Psi(w))[D^2\Psi(w)[u, z], D\Psi(w)[v]] \\ &\quad + D^2E(\Psi(w))[D\Psi(w)[u], D^2\Psi(w)[v, z]] \\ &\quad + D^2E(\Psi(w))[D\Psi(w)[z], D^2\Psi(w)[v, u]] \\ &\quad + \langle DE(\Psi(w)), D^3\Psi(w)[v, u, z] \rangle. \end{aligned}$$

Setting $w = 0$, and using similar considerations as before (in particular noting that $D^2E(0)[\cdot]$ is self-adjoint), we obtain $D^3F(0)[v, u, z] = D^3E(0)[v, u, z]$. Performing the same computation for $D^3\tilde{F}(0)$ yields the same result. Next, we compute $D^3\tilde{F}(0)$. Recall from (2.5) that

$$\frac{1}{2}DE(w)[v] = \int_M (P_{g_\infty} w + Q_{g_\infty} - \overline{Q}_{g_\infty} e^{nw}) v dV_{g_\infty}.$$

When computing the third derivative of E at 0, the first two terms will vanish. Hence, we have

$$(2.10) \quad D^3E(0)[v, u, z] = -2n^2 \int_M Q_{g_\infty} vuz dV_{g_\infty}.$$

This together with (2.9) implies that

$$D^3F(0)[v, u, z] = -2n^2 \int_M Q_{g_\infty} vuz dV_{g_\infty},$$

which proves (2.8). \square

3. PROOF OF THEOREM 1.4

One of the tools for controlling the rate of convergence of the Q -curvature flow will be the Łojasiewicz-Simon inequality.

Proposition 3.1. *Suppose that $g_\infty \in CQC_1$. There is $\theta \in (0, \frac{1}{2}]$, $\epsilon > 0$ and $C > 0$ (both depending only on n and g_∞) such that for $u \in C^{n,\alpha}(M, g_\infty)$ with $\|u\|_{C^{n,\alpha}(M, g_\infty)} < \epsilon$, then*

$$|E(u) - E(0)|^{1-\theta} \leq C \|DE(u)\|_{L^2(M, g_\infty)}.$$

If g_∞ is an integrable critical point, then $\theta = \frac{1}{2}$. If g_∞ is non-integrable, then this holds for some $\theta \in (0, \frac{1}{p}]$, where p is the order of integrability of g_∞ .

Proof. One may follow the argument in [9]. \square

Proof of Theorem 1.4. We consider the Q -curvature flow $g(t) = e^{2u(t)}g_\infty$ which converges to g_∞ in $C^n(M, g_\infty)$ as $t \rightarrow \infty$. In Proposition 3.1, we have shown that there is a Łojasiewicz-Simon inequality near g_∞ for some $\theta \in (0, \frac{1}{2}]$. We emphasize that if we regard $DE(g(t))$ as an element of $L^2(M, g_\infty)$, then

$$(3.1) \quad DE(g(t)) = 2(Q_{g(t)} - \bar{Q}_{g(t)})e^{nu(t)}.$$

Since $g(t) \rightarrow g_\infty$ as $t \rightarrow \infty$, we can choose t_0 so that for $t \geq t_0$, $\|u(t)\|_{C^0(M, g_\infty)} \leq \frac{1}{2}$. This together with (2.4) and Proposition 3.1 implies that

$$(3.2) \quad \begin{aligned} \frac{d}{dt}(E(u(t)) - E(0)) &= -\frac{n}{2} \int_M (Q_{g(t)} - \bar{Q}_{g(t)})^2 e^{nu(t)} dV_{g_\infty} \\ &\leq -c \int_M (Q_{g(t)} - \bar{Q}_{g(t)})^2 e^{2nu(t)} dV_{g_\infty} \\ &= -c \|DE(u(t))\|_{L^2(M, g_\infty)}^2 \\ &\leq -c |E(u(t)) - E(0)|^{2-2\theta}, \end{aligned}$$

where $c > 0$ is a constant depending only on n and g_∞ (that we let change from line to line). Let us first assume that the Łojasiewicz-Simon inequality is satisfied with $\theta = \frac{1}{2}$, i.e. we are in the integrable case. Then (3.2) yields $E(u(t)) - E(0) \leq Ce^{-2\delta t}$, for some $\delta > 0$ depending only on n and g_∞ , and $C > 0$ depending on $g(0)$ (chosen so that this actually holds for all $t \geq 0$). On the other hand, if Łojasiewicz-Simon inequality holds with $\theta \in (0, \frac{1}{2})$, then the same argument shows that $E(u(t)) - E(0) \leq C(1+t)^{\frac{1}{2\theta-1}}$.

Exploiting the fact that the flow converges in C^n , we may use the Łojasiewicz-Simon inequality to compute

$$\begin{aligned} \frac{d}{dt}(E(u(t)) - E(0))^\theta &= \theta(E(u(t)) - E(0))^{\theta-1} \frac{d}{dt}(E(u(t)) - E(0)) \\ &\leq -c\theta(E(u(t)) - E(0))^{\theta-1} \|DE(u(t))\|_{L^2(M, g_\infty)}^2 \\ &\leq -c\theta \|DE_f(u(t))\|_{L^2(M, g_\infty)} \\ &\leq -c\theta \left\| \frac{\partial u(t)}{\partial t} \right\|_{L^2(M, g_\infty)} \end{aligned}$$

where we have used (2.2) and (3.1) in the last equality. Thus, if $\theta = \frac{1}{2}$ (recall $\lim_{t \rightarrow \infty} u(t) = 0$), then

$$\begin{aligned} \|u(t)\|_{L^2(M, g_\infty)} &\leq \int_t^\infty \left\| \frac{\partial u(s)}{\partial s} \right\|_{L^2(M, g_\infty)} ds \\ &\leq -c \int_t^\infty \frac{d}{ds} \left[(E(u(s)) - E(1))^{\frac{1}{2}} \right] ds \\ &= c(E(u(t)) - E(1))^{\frac{1}{2}} \leq C e^{-\delta t}. \end{aligned}$$

A similar computation if $\theta \in (0, \frac{1}{2})$ yields $\|u(t)\|_{L^2(M, g_\infty)} \leq C(1+t)^{-\frac{\theta}{1-2\theta}}$.

To obtain C^n estimates, we may interpolate between $L^2(M, g)$ and $W^{k,2}(M, g)$ for k large enough: interpolation [4, Theorem 6.4.5] and Sobolev embedding yields some constant $\eta \in (0, 1)$ so that

$$\|u(t)\|_{C^{n,\alpha}(M, g_\infty)} \leq \|u(t)\|_{L^2(M, g_\infty)}^\eta \|u(t)\|_{W^{k,2}(M, g_\infty)}^{1-\eta}.$$

Because $u(t)$ converges to 0 in $C^{n,\alpha}$ (and thus in C^∞ by parabolic Schauder estimates and bootstrapping), the second term is uniformly bounded. Thus, exponential (polynomial) decay of the L^2 norm gives exponential (polynomial) decay of the $C^{n,\alpha}$ norm as well. \square

4. SLOWLY CONVERGING Q -CURVATURE FLOW

In this section, we show that, given a nonintegrable critical point g_∞ satisfying a particular hypothesis, there exists a Q -curvature flow $g(t)$ such that $g(t)$ converges to g_∞ exactly at a polynomial rate.

This section is organized as follows: In section 4.1, we show that the Q -curvature flow can be represented by two different flows. To be more specific, we will project the flow equation to the kernel Λ_0 of \mathcal{L}_∞ and its orthogonal complement Λ_0^\perp , respectively. In section 4.2, we solve the kernel-projected flow. In section 4.3, we solve the kernel-orthogonal projected flow. In section 4.4, we combine all the previous results to prove Theorem 1.5.

4.1. Projecting the Q -curvature flow with estimates. Here and in the sequel we will always use $f'(t)$ to denote the time derivative of a function $f(t)$. We will skip the proof of the following lemma, for its proof is the same as that of [9, Lemma 15].

Lemma 4.1. *Assume that g_∞ satisfies AS_p as defined in Definition 2.4, i.e. $F_p|_{\mathbb{S}^k}$ achieves a positive maximum for some point \hat{v} in the unit sphere $\mathbb{S}^k \subset \Lambda_0$. Then, for any fixed $T \geq 0$, the function*

$$(4.1) \quad \varphi(t) := \varphi(t, T) = (T+t)^{-\frac{1}{p-2}} \left(\frac{n}{p(p-2)F_p(\hat{v})} \right)^{\frac{1}{p-2}} \hat{v}$$

solves $n\varphi' + DF_p(\varphi) = 0$.

In the next result and subsequently in this section, we will always denote by $\|f(t)\|_{C^{k,\alpha}}$ the parabolic $C^{k,\alpha}$ norm on $(t, t+1) \times M$. More precisely, for $\alpha \in (0, 1)$, we define the seminorm

$$|f(t)|_{C^{0,\alpha}} = \sup_{\substack{(s_i, x_i) \in (t, t+1) \times M \\ (s_1, x_1) \neq (s_2, x_2)}} \frac{|f(s_1, x_1) - f(s_2, x_2)|}{(d_{g_\infty}(x_1, x_2)^n + |t_1 - t_2|)^{\frac{\alpha}{n}}}$$

and for $k \geq 0$ and $\alpha \in (0, 1)$, we define the norm

$$(4.2) \quad \|f(t)\|_{C^{k,\alpha}} = \sum_{|\beta|+nj \leq k} \sup_{(t,t+1) \times M} |D_x^\beta D_t^j f| + \sum_{|\beta|+nj=k} |D_x^\beta D_t^j f|_{C^{0,\alpha}}$$

where the norm and derivatives in the sum are taken with respect to g_∞ . When we mean an alternative norm, we will always indicate the domain.

Lemma 4.2. *For the functional E , there holds*

$$(4.3) \quad \|D^3 E(w)[u, v]\|_{C^{0,\alpha}} \leq C \|u\|_{C^{n,\alpha}} \|v\|_{C^{n,\alpha}}$$

for some uniform constant $C > 0$. Furthermore, for w_1, w_2 such that $\|w_i\|_{C^{n,\alpha}} < 1$, we have

$$\begin{aligned} \|D^3 E(w_1)[v, v] - D^3 E(w_2)[u, u]\|_{C^{0,\alpha}} &\leq C (\|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}}) \\ &\quad \times (\|u\|_{C^{n,\alpha}} + \|v\|_{C^{n,\alpha}}) \|u - v\|_{C^{n,\alpha}} \end{aligned}$$

for some uniform constant $C > 0$.

Proof. This follows from (2.10). \square

Lemma 4.3. *There exists $T_0 > 0$, $\epsilon_0 > 0$ and $c > 0$, all depending on g_∞ and \hat{v} , such that the following holds: Fix $T > T_0$. Then, for $\varphi(t)$ as in Lemma 4.1 and $w \in C^{n,\alpha}(M \times [0, \infty))$, and $u := \varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp$, the function*

$$E_0^\top(w) := \text{proj}_{\Lambda_0} [DE(u)e^{-nu} - DE(u)]$$

satisfies

$$\begin{aligned} \|E_0^\top(w)\|_{C^{0,\alpha}} &\leq c \left\{ (T+t)^{-\frac{p-1}{p-2}} + \|w^\top\|_{C^{0,\alpha}}^{p-1} + \|w^\perp\|_{C^{n,\alpha}} \right\} \left\{ (T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right\}, \\ \|E_0^\top(w_1) - E_0^\top(w_2)\|_{C^{0,\alpha}} &\leq c \left\{ (T+t)^{-\frac{p-1}{p-2}} + \|w_1^\top\|_{C^{0,\alpha}}^{p-1} + \|w_2^\top\|_{C^{0,\alpha}}^{p-1} + \|w_1^\perp\|_{C^{n,\alpha}} + \|w_2^\perp\|_{C^{n,\alpha}} \right\} \\ &\quad \times \|w_1 - w_2\|_{C^{n,\alpha}} \\ &+ c \left\{ (T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right\} \left(\|w_1^\top\|_{C^{0,\alpha}}^{p-2} + \|w_2^\top\|_{C^{0,\alpha}}^{p-2} \right) \\ &\quad \times \|w_1^\top - w_2^\top\|_{C^{0,\alpha}} \\ &+ c \left\{ (T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right\} \|w_1^\perp - w_2^\perp\|_{C^{n,\alpha}}. \end{aligned}$$

Identical estimates hold for $E_0^\perp(w) := \text{proj}_{\Lambda_0^\perp} [DE(u)e^{-nu} - DE(u)]$. Here, we are using the parabolic Hölder norms on $(t, t+1) \times M$ as defined above; the bounds hold for each fixed $t \geq 0$, with the constants independent of T and t .

Proof. First, we have

$$e^{-nu} = 1 - n \int_0^1 e^{-snu} u ds.$$

where $u = \varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp$. So, letting $E_0(w) := DE(u)e^{-nu} - DE(u)$, we have

$$(4.4) \quad \|E_0(w)\|_{C^{0,\alpha}} \leq c \|DE(u)\|_{C^{0,\alpha}} \left((T+t)^{-\frac{1}{p-2}} + \|w^\top\|_{C^{0,\alpha}} + \|w^\perp\|_{C^{0,\alpha}} \right),$$

where we used the fact that $\Phi(0) = 0$ and Φ is an analytic map. The rest of the proof is similar to that [9, Lemma 16] using Taylor's theorem, Proposition 2.1, and Lemma 4.2. \square

The next result reduces the Q -curvature flow to two flows, one on Λ_0 and the other on Λ_0^\perp .

Proposition 4.4. *There exists $T_0 > 0$, $\epsilon_0 > 0$ and $c > 0$, all depending on g_∞ and \hat{v} , such that the following holds: Fix $T > T_0$. Then, for $\varphi(t)$ as in Lemma 4.1 and $w \in C^{n,\alpha}(M \times [0, \infty))$, there are functions $E^\top(w)$ and $E^\perp(w)$ such that $u := \varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp$ is a solution to the Q -curvature flow if and only if*

$$(4.5) \quad n(w^\top)' + D^2 F_p(\varphi)w^\top = E^\top(w),$$

$$(4.6) \quad (w^\perp)' + \frac{1}{2}\mathcal{L}_\infty w^\perp = E^\perp(w).$$

Here, as long as $\|w\|_{C^{n,\alpha}} \leq \epsilon_0$, the error terms E^\top and E^\perp satisfy

$$\begin{aligned} \|E^\top(w)\|_{C^{0,\alpha}} &\leq c \left((T+t)^{-\frac{p-1}{p-2}} + \|w^\top\|_{C^{0,\alpha}}^{p-1} + \|w^\perp\|_{C^{n,\alpha}} \right) \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \\ &\quad + c(T+t)^{-\frac{p}{p-2}} + c(T+t)^{-\frac{p-1}{p-2}} \|w^\top\|_{C^{0,\alpha}} + c(T+t)^{-\frac{p-3}{p-2}} \|w^\top\|_{C^{0,\alpha}}^2 \\ &\quad + c\|w^\top\|_{C^{0,\alpha}}^{p-1} + c \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \|w^\perp\|_{C^{n,\alpha}}, \end{aligned}$$

$$\begin{aligned} &\|E^\perp(w_1) - E^\perp(w_2)\|_{C^{0,\alpha}} \\ &\leq c \left((T+t)^{-\frac{p-1}{p-2}} + \|w_1^\top\|_{C^{0,\alpha}}^{p-1} + \|w_2^\top\|_{C^{0,\alpha}}^{p-1} + \|w_1^\perp\|_{C^{n,\alpha}} + \|w_2^\perp\|_{C^{n,\alpha}} \right) \\ &\quad \times \|w_1 - w_2\|_{C^{n,\alpha}} \\ &\quad + c \left((T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right) (\|w_1^\top\|_{C^{0,\alpha}}^{p-2} + \|w_2^\top\|_{C^{0,\alpha}}^{p-2}) \\ &\quad \times \|w_1^\top - w_2^\top\|_{C^{0,\alpha}} \\ &\quad + c \left((T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right) \|w_1^\perp - w_2^\perp\|_{C^{n,\alpha}} \\ &\quad + c \left((T+t)^{-\frac{p-3}{p-2}} (\|w_1^\top\|_{C^{0,\alpha}} + \|w_2^\top\|_{C^{0,\alpha}}) + \|w_1^\top\|_{C^{0,\alpha}}^{p-2} + \|w_2^\top\|_{C^{0,\alpha}}^{p-2} \right) \\ &\quad \times \|w_1^\top - w_2^\top\|_{C^{0,\alpha}} \\ &\quad + c(T+t)^{-\frac{p-1}{p-2}} \|w_1^\top - w_2^\top\|_{C^{0,\alpha}}, \end{aligned}$$

$$\begin{aligned} &\|E^\perp(w)\|_{C^{0,\alpha}} \\ &\leq c \left((T+t)^{-\frac{p-1}{p-2}} + \|w^\top\|_{C^{0,\alpha}}^{p-1} + \|w^\perp\|_{C^{n,\alpha}} \right) \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \\ &\quad + c \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \|w^\perp\|_{C^{n,\alpha}} \\ &\quad + c \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \left((T+t)^{-\frac{p-1}{p-2}} + \|w'\|_{C^{0,\alpha}} \right) \end{aligned}$$

$$\begin{aligned}
& \|E^\perp(w_1) - E^\perp(w_2)\|_{C^{0,\alpha}} \\
& \leq c \left((T+t)^{-\frac{p-1}{p-2}} + \|w_1^\top\|_{C^{0,\alpha}}^{p-1} + \|w_2^\top\|_{C^{0,\alpha}}^{p-1} + \|w_1^\perp\|_{C^{n,\alpha}} + \|w_2^\perp\|_{C^{n,\alpha}} \right) \\
& \quad \times \|w_1 - w_2\|_{C^{n,\alpha}} \\
& + c \left((T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right) (\|w_1^\top\|_{C^{0,\alpha}}^{p-2} + \|w_2^\top\|_{C^{0,\alpha}}^{p-2}) \\
& \quad \times \|w_1^\top - w_2^\top\|_{C^{0,\alpha}} \\
& + c \left((T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right) \|w_1^\perp - w_2^\perp\|_{C^{n,\alpha}} \\
& + c \left((T+t)^{-\frac{1}{p-2}} + \|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}} \right) \|w_1' - w_2'\|_{C^{0,\alpha}} \\
& + c \left((T+t)^{-\frac{p-1}{p-2}} + \|w_1'\|_{C^{0,\alpha}} + \|w_2'\|_{C^{0,\alpha}} \right) \|w_1 - w_2\|_{C^{n,\alpha}}.
\end{aligned}$$

Here we are using the parabolic Hölder norms on $(t, t+1) \times M$ as defined above; the bounds hold for each fixed $t \geq 0$, with the constants independent of T and t .

Proof. We follow the proof of [9, Proposition 17]. It follows from (2.2) and (2.5) that u is a solution to the Q -curvature flow if and only if

$$(4.7) \quad \frac{\partial u}{\partial t} = -\frac{1}{n} DE(u)e^{-nu}.$$

We now project the Q -curvature flow equation onto Λ_0 and Λ_0^\perp , so u solves the Q -curvature flow if and only if the following two equations are satisfied:

$$\begin{aligned}
(4.8) \quad & n(\varphi + w^\top)' = -\text{proj}_{\Lambda_0} [DE(\varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp)] - E_0^\top(w), \\
& n(\Phi(\varphi + w^\top) + w^\perp)' = -\text{proj}_{\Lambda_0^\perp} [DE(\varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp)] - E_0^\perp(w),
\end{aligned}$$

where $E_0(w)$ is defined as in Lemma 4.3. The rest of the proof is similar except for that the Taylor's theorem gives us

$$\begin{aligned}
(4.9) \quad & \text{proj}_{\Lambda_0} DE(\varphi + w^\top + \Phi(\varphi + w^\top) + w^\perp) \\
& = \text{proj}_{\Lambda_0} DE(\varphi + w^\top + \Phi(\varphi + w^\top)) + E_1^\top(w),
\end{aligned}$$

with slightly different bounds

$$\begin{aligned}
(4.10) \quad & \|E_1^\top(w)\|_{C^{0,\alpha}} \leq c \left((T+t)^{-\frac{1}{p-2}} + \|w\|_{C^{n,\alpha}} \right) \|w^\perp\|_{C^{n,\alpha}}, \\
& \|E_1^\top(w_1) - E_1^\top(w_2)\|_{C^{0,\alpha}} \leq c (\|w_1\|_{C^{n,\alpha}} + \|w_2\|_{C^{n,\alpha}}) \|w_1^\perp - w_2^\perp\|_{C^{n,\alpha}}.
\end{aligned}$$

□

4.2. Solving the kernel-projected flow with polynomial decay estimates.

In this subsection we solve the kernel-projected flow (4.5). First, from the definition of φ in (4.1) and the fact that D^2F_p is homogeneous of degree $p-2$,

$$D^2F_p(\varphi) = (T+t)^{-1} \underbrace{\left(\frac{n}{p(p-2)F_p(\hat{v})} \right)}_{:=\mathcal{D}} D^2F_p(\hat{v}).$$

Let μ_1, \dots, μ_k be the eigenvalues of \mathcal{D} and e_i the corresponding orthonormal basis in which \mathcal{D} is diagonalized. Then the kernel-projected flow is equivalent to the following system of ODEs for $v_i := w^\top \cdot e_i$,

$$(4.11) \quad n(w^\top)' + \frac{\mu_i}{T+t} v_i = E_i^\top := E^\top \cdot e_i, \quad i = 1, \dots, k.$$

Fix for the rest of this subsection a number γ with $\gamma \notin \{\frac{1}{n}\mu_1, \dots, \frac{1}{n}\mu_k\}$. Define the following weighted norms:

$$\|u\|_{C_\gamma^{0,\alpha}} := \sum_{t>0} [(T+t)^\gamma \|u(t)\|_{C^{0,\alpha}}] \quad \text{and} \quad \|u\|_{C_{1,\gamma}^{0,\alpha}} := \|u\|_{C_\gamma^{0,\alpha}} + \|u'\|_{C_{1+\gamma}^{0,\alpha}}.$$

We recall that these Hölder norms are space-time norms on the interval $(t, t+1) \times M$, as defined in (4.2).

Given γ as above, we define $\Pi_0 = \Pi_0(\gamma)$ by

$$(4.12) \quad \Pi_0 := \text{span}\{v \in \Lambda_0 : \mathcal{D}v = \mu v, \text{ and } \mu > n\}.$$

Moreover, let $\text{proj}_{\Pi_0} : \Lambda \rightarrow \Pi_0$ be the corresponding linear projector.

Lemma 4.5. *For any $T > 0$ such that $\|E^\top\|_{C_{1+\gamma}^{0,\alpha}} < \infty$, there is a unique u with $u(t) \in \Lambda_0$, $t \in [0, \infty)$, satisfying $\|u\|_{C_\gamma^0} < \infty$, $\text{proj}_{\Pi_0}(u(0)) = 0$, and such that $v_i := u \cdot e_i$ solves the system (4.11). Furthermore, we have the bound*

$$\|u\|_{C_{1,\gamma}^{0,\alpha}} \leq C \|E^\top\|_{C_{1+\gamma}^{0,\alpha}}.$$

Proof. See [9, Lemma 18]. □

4.3. Solving the kernel-orthogonal projected flow. Define the weighted norms

$$\|u\|_{L_q^2} = \sup_{t \in [0, \infty)} [(T+t)^q \|u(t)\|_{L^2(M)}],$$

where the L^2 norm is the spatial norm of $u(t)$ on M , taken with respect to g_∞ , and

$$\|u\|_{C_q^{n,\alpha}} = \sup_{t \geq 0} [(T+t)^q \|u(t)\|_{C^{n,\alpha}}],$$

where, as usual, the Hölder norms are the space-time norms defined in (4.2). Also, let

$$\begin{aligned} \Lambda_\downarrow &:= \overline{\text{span}\{\varphi \in C^\infty(M) : \mathcal{L}_\infty \varphi + \delta \varphi = 0, \delta > 0\}}^{L^2}, \\ \Lambda_\uparrow &:= \text{span}\{\varphi \in C^\infty(M) : \mathcal{L}_\infty \varphi + \delta \varphi = 0, \delta < 0\}. \end{aligned}$$

From the spectral theory, $L^2(M, g_\infty) = \Lambda_\uparrow \oplus \Lambda_0 \oplus \Lambda_\downarrow$ and Λ_\uparrow and Λ_0 are finite dimensional. Write the nonnegative integers as an ordered union $\mathbb{N} = K_\uparrow \cup K_0 \cup K_\downarrow$, where the ordering of the indices from an ordering of the eigenfunctions of the \mathcal{L}_∞ and the partitioning of \mathbb{N} corresponds to which of Λ_\downarrow , Λ_0 , or Λ_\uparrow the k -th eigenfunction of \mathcal{L}_∞ lies in.

Lemma 4.6. *For any $T > 0$ and $q < \infty$ such that $\|E^\perp\|_{L_q^2} < \infty$, there is a unique $u(t)$ with $u(t) \in \Lambda_0^\perp$, $t \in [0, \infty)$, satisfying $\|u\|_{L_q^2} < \infty$, $\text{proj}_{\Lambda_\downarrow}(u(0)) = 0$ and*

$$(4.13) \quad u' = \mathcal{L}_\infty u + E^\perp.$$

Furthermore, $\|u\|_{L_q^2} \leq C \|E^\perp\|_{L_q^2}$ and $\|u\|_{C_q^{n,\alpha}} \leq C \|E^\perp\|_{C_q^{0,\alpha}}$.

Proof. We now consider the $C_q^{n,\alpha}$ bounds for u . By interior parabolic Schauder estimates (see [18] for example) and Arzelà-Ascoli theorem, we have that for $t \geq 1$,

$$\|u(t)\|_{C^{n,\alpha}} \leq C \left(\sup_{s \in (t-1, t+1)} \|u(s, x)\|_{L^2(M)} + \|E^\perp\|_{C^{0,\alpha}((t-1, t+1) \times M)} \right) + C\epsilon \|u(t)\|_{C^{0,\alpha}((t-1, t+1) \times M)}.$$

Multiplying it by $(T+t)^q$ and taking the supremum over $t \geq 1$ yields

$$(4.14) \quad \sup_{t \geq 1} [(T+t)^q \|u(t)\|_{C^{n,\alpha}}] \leq C \|E^\perp\|_{C_q^{0,\alpha}} + C\epsilon \|u\|_{C_q^{0,\alpha}}$$

using $\|u\|_{L_q^2} \leq C \|E^\perp\|_{L_q^2}$. To finish the proof, it remains to extend the supremum up to $t = 0$. The global Schauder estimates (c.f. [18]) shows that

$$(4.15) \quad \|u(t)\|_{C^{n,\alpha}((0,1) \times M)} \leq C \left(\sup_{s \in (0,1)} \|u(s, x)\|_{L^2(M)} + \epsilon \|u\|_{C^{0,\alpha}((0,1) \times M)} + \|E^\perp\|_{C^{0,\alpha}((0,1) \times M)} + \|u(0)\|_{C^{n,\alpha}(M)} \right).$$

The remaining parts of proof is analogous to that of [9, Lemma 19]. \square

4.4. Construction of a slowly converging flow. To proceed further, we define the norm

$$\|f\|_\gamma^* := \|\text{proj}_{\Lambda_0} f\|_{C_{1,\gamma}^{0,\alpha}} + \|\text{proj}_{\Lambda_0^\perp} f\|_{C_{1+\gamma}^{n,\alpha}}.$$

For γ to be specified below, the Banach space X is defined as

$$(4.16) \quad X := \{w : \|w\|_\gamma^* < \infty\}.$$

Proposition 4.7. *Assume that g_∞ satisfies AS_p . We may thus fix a point where $F_p|_{\mathbb{S}^{k-1}}$ achieves a positive maximum and denote it by \hat{v} . Define*

$$\varphi(t) = (T+t)^{-\frac{1}{p-2}} \left(\frac{n}{p(p-2)F_p(\hat{v})} \right)^{\frac{1}{p-2}} \hat{v}$$

as in Lemma 4.1. Then, there exists $C > 0$, $T > 0$, $\frac{1}{p-2} < \gamma < \frac{2}{p-2}$ and $u(t) \in C^\infty(M \times (0, \infty))$ such that $u(t) > 0$ for all $t > 0$, $g(t) := e^{2u(t)}g_\infty$ is a solution to the Q -curvature flow and

$$\|w^\top(t) + \Phi(\varphi(t) + w^\top(t)) + w^\perp(t)\|_\gamma^* = \|u(t) - \varphi(t)\|_\gamma^* \leq C.$$

Proof. The proposition follows from the contraction mapping principle, as applied [9, Proposition 20]. \square

Now we are ready to prove Theorem 1.5.

Proof of Theorem 1.5. From Proposition 4.7, we have constructed $\varphi(t)$ and $u(t)$ so that

$$\varphi(t) = (T+t)^{-\frac{1}{p-2}} \left(\frac{n}{p(p-2)F_p(\hat{v})} \right)^{\frac{1}{p-2}} \hat{v},$$

$e^{2u(t)}g_\infty$ is a solution to the Q -curvature flow, and

$$u(t) = \varphi(t) + \tilde{w}(t) := \varphi(t) + w^\top(t) + \Phi(\varphi(t) + w^\top(t)) + w^\perp(t),$$

where $\tilde{w}(t)$ satisfies $\|\tilde{w}\|_{C^0} \leq C(1+t)^{-\gamma}$ for some $C > 0$ and all $t \geq 0$. We have arranged that $\gamma > 1/(p-2)$, which implies that $\varphi(t)$ is decaying slower than $\tilde{w}(t)$. Thus

$$\|u(t)\|_{C^0} \geq C(1+t)^{-\frac{1}{p-2}}$$

as $t \rightarrow \infty$. From this, the assertion follows \square

5. AN EXAMPLE SATISFYING AS_3

In this section, we provide an example of Riemannian metric satisfying AS_3 for the 6th order Paneitz operator P_6 and Q_6 -curvature in dimension $n = 6$. Moreover, we will see that this Riemannian metric satisfies the assumptions in Theorem 1.3. This allows us to conclude the existence of slowly converging Q -curvature flows.

We will omit the subscript g in curvature tensors and differential operators in this section, as it will not cause any ambiguity. We introduce some notations: $J = \frac{R}{10}$, δ denotes the adjoint operator of exterior derivative d , and A denotes the Schouten tensor $A = \frac{1}{4}(Ric - Jg)$. The Bach tensor \mathcal{B} is defined as

$$(5.1) \quad \mathcal{B}_{ij} = \Delta A_{ij} - \nabla^k \nabla_j A_{ik} + A^{kl} W_{ikjl}$$

We recall following formulas for the P_6 and Q_6 in dimension 6 from [30, Theorem 6.10.3] and [31, Theorem 10.1]. Note that the sign of P_6 differs from [31] because we want P_6 to have $(-\Delta)^3$ as the leading order term. Additionally, the sign of the Riemann curvature tensor is different from [30].

$$P_6 = -\Delta^3 - 4\Delta\delta(J - 2A)d - 4\delta(J - 2A)d\Delta - 2\Delta(J\Delta) \\ - 8\delta(-J^2 + |A|^2 + 4JA - 6A^2 - \mathcal{B})d$$

$$Q_6 = \{16tr(A^3) - 24J|A|^2 + 8J^3 + 8\langle \mathcal{B}, A \rangle\} \\ + (\Delta^2 J - 8\delta(AdJ)) + 4\Delta(|A|^2 - J^2)$$

Our goal is to find a λ such that $(S^2(\lambda) \times \mathbb{C}P^2, g_{prod})$ with the product metric g_{prod} satisfies the AS_3 condition, where $S^2(\lambda)$ is the standard sphere with metric g_λ having the sectional curvature λ and $\mathbb{C}P^2$ is equipped with the Fubini-Study metric g_{FS} .

There is a canonical projection from $S^2(\lambda) \times \mathbb{C}P^2$ to each of $S^2(\lambda)$ and $\mathbb{C}P^2$ denoted by π_1 and π_2 , respectively. We denote the pullback of the metric and the curvature quantities by using the superscript (1) and (2). For example, $g^{(1)} = \pi_1^* g_\lambda$, $g^{(2)} = \pi_2^* g_{FS}$, $Rm^{(1)} = \pi_1^* Rm_{g_\lambda}$ and $Rm^{(2)} = \pi_2^* Rm_{g_{FS}}$. We also denote $\Delta^{(1)}$, $\Delta^{(2)}$ Laplacians acting on the first component and the second component of the product structure, respectively.

It is clear that

$$g_{prod} = \pi_1^* g_\lambda \oplus \pi_2^* g_{FS} = g^{(1)} \oplus g^{(2)} \\ Rm = \pi_1^* Rm_{g_\lambda} \oplus \pi_2^* Rm_{g_{FS}} = Rm^{(1)} \oplus Rm^{(2)} \\ \Delta = \Delta^{(1)} + \Delta^{(2)}.$$

Since $(\mathbb{C}P^2, g_{FS})$ is an Einstein manifold with Einstein constant 6, we have

$$Ric = \lambda g^{(1)} \oplus 6g^{(2)}, R = 24 + 2\lambda \\ J = \frac{\lambda + 12}{5}, A = \left(\frac{\lambda - 3}{5}\right)g^{(1)} \oplus \left(\frac{18 - \lambda}{20}\right)g^{(2)}.$$

We recall the following decomposition of the Riemann curvature tensor:

$$Rm = W + A \otimes g$$

where \otimes is the Kulkarni-Nomizu product defined as $(S \otimes T)_{ijkl} = S_{ik}T_{jl} + S_{jl}T_{ik} - S_{il}T_{jk} - S_{jk}T_{il}$. We observe the following basic formulas for the product metric.

Lemma 5.1. *The following identities hold for components $g^{(1)}$ and $g^{(2)}$ of g_{prod} .*

- (1) $(g^{(1)})^{kl}(g^{(1)} \otimes g^{(1)})_{ikjl} = 2g_{ij}^{(1)}$.
- (2) $(g^{(2)})^{kl}(g^{(2)} \otimes g^{(2)})_{ikjl} = 6g_{ij}^{(2)}$.
- (3) $(g^{(1)})^{kl}(g^{(1)} \otimes g^{(2)})_{ikjl} = 2g_{ij}^{(2)}$.
- (4) $(g^{(2)})^{kl}(g^{(1)} \otimes g^{(2)})_{ikjl} = 4g_{ij}^{(1)}$.

Proof. We prove the first and third identity by observing $(g^{(1)})^{kl}g_{ik}^{(2)} = 0$. For the first identity:

$$(g^{(1)})^{kl}(g^{(1)} \otimes g^{(1)})_{ikjl} = 2(g^{(1)})^{kl}(g_{ij}^{(1)}g_{kl}^{(1)} - g_{il}^{(1)}g_{jk}^{(1)}) = (4 - 2)g_{ij}^{(1)} = 2g_{ij}^{(1)}.$$

For the third identity:

$$(g^{(1)})^{kl}(g^{(1)} \otimes g^{(2)})_{ikjl} = (g^{(1)})^{kl}(g_{ij}^{(1)}g_{kl}^{(2)} + g_{kl}^{(1)}g_{ij}^{(2)} - g_{il}^{(1)}g_{jk}^{(2)} - g_{jk}^{(1)}g_{il}^{(2)}) = 2g_{ij}^{(2)}.$$

□

Now we are ready to compute Q_6 -curvature and P_6 operator for the product metric g .

Proposition 5.2. *For $(S^2(\lambda) \times \mathbb{C}P^2, g_{prod})$, we have*

$$\begin{aligned} Q_6 &= 0.48\lambda^3 - 6.72\lambda^2 + 63.36\lambda - 34.56 \\ P_6 &= -\Delta^3 - 2(\lambda - 12)\Delta^{(1)}\Delta + 2\lambda\Delta^{(2)}\Delta \\ &\quad + 8\left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25}\right)\Delta^{(1)} + 8\left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25}\right)\Delta^{(2)}. \end{aligned}$$

Proof. Due to the product structure of the metric g_{prod} , it is easy to see that the formula for Q_6 -curvature is simplified as

$$(5.2) \quad Q_6 = 6tr(A^3) - 24J|A|^2 + 8J^3 + 8\langle \mathcal{B}, A \rangle.$$

It remains for us to compute the Bach tensor \mathcal{B} . Using the formula (5.1) and noting that both $(S^2(\lambda), g_\lambda)$ and $(\mathbb{C}P^2, g_{FS})$ are Einstein, it is easy to see that $\mathcal{B}_{ij} = A^{kl}W_{ikjl}$.

To compute $A^{kl}W_{kijl}$, we use the identity

$$\begin{aligned} W + A \otimes g &= Rm = Rm^{(1)} \oplus Rm^{(2)} \\ &= \left(\frac{\lambda}{2}g^{(1)} \otimes g^{(1)}\right) \oplus (W^{(2)} + A^{(2)} \otimes g^{(2)}) \end{aligned}$$

to see that

$$\begin{aligned}
\mathcal{B}_{ij} &= A^{kl}W_{ikjl} \\
&= A^{kl} \left[- (A \otimes g) \oplus \left(\frac{\lambda}{2}g^{(1)} \otimes g^{(1)} \right) \oplus (W^{(2)} + A^{(2)} \otimes g^{(2)}) \right]_{ikjl} \\
&= A^{kl} \left[- (A \otimes g) \oplus \left(\frac{\lambda}{2}g^{(1)} \otimes g^{(1)} \right) \oplus (g^{(2)} \otimes g^{(2)}) \right]_{ikjl} \\
&= \left(\frac{\lambda+2}{20} \right) \left[\left(\frac{\lambda-3}{5} \right)g^{(1)} \oplus \left(\frac{18-\lambda}{20} \right)g^{(2)} \right]^{kl} (6g^{(1)} \otimes g^{(1)} \oplus g^{(2)} \otimes g^{(2)} - 3g^{(1)} \otimes g^{(2)})_{ikjl} \\
&= \frac{3(\lambda+2)(\lambda-6)}{40} (2g^{(1)} - g^{(2)})
\end{aligned}$$

where we used identities from Lemma 5.1.

Now it is straightforward to see that:

$$\begin{aligned}
Q_6 &= \{16tr(A^3) - 24J|A|^2 + 8J^3 + 8\langle \mathcal{B}, A \rangle\} \\
&= 16 \left[2 \left(\frac{\lambda-3}{5} \right)^3 + 4 \left(\frac{18-\lambda}{20} \right)^3 \right] - \frac{24(\lambda+12)}{5} \left[2 \left(\frac{\lambda-3}{5} \right)^2 + 4 \left(\frac{18-\lambda}{20} \right)^2 \right] \\
&\quad + 8 \frac{(\lambda+12)^3}{5^3} + \frac{3}{5} (\lambda+2)(\lambda-6)^2 \\
&= 0.48\lambda^3 - 6.72\lambda^2 + 63.36\lambda - 34.56
\end{aligned}$$

Next, we compute the P_6 operator. It is obvious that $\Delta^{(1)}$ and $\Delta^{(2)}$ commute.

$$\begin{aligned}
P_6 &= -\Delta^3 - 4\Delta\delta(J-2A)d - 4\delta(J-2A)d\Delta - 2\Delta(J\Delta) \\
&\quad - 8\delta(-J^2 + |A|^2 + 4JA - 6A^2 - \mathcal{B})d \\
&= -\Delta^3 - 2(\lambda-12)\Delta^{(1)}\Delta + 2\lambda\Delta^{(2)}\Delta \\
&\quad + 8 \left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25} \right) \Delta^{(1)} + 8 \left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25} \right) \Delta^{(2)}.
\end{aligned}$$

□

Example 5.1. We show that there exists $\lambda > 0$ such that $P_6\phi_1 = 6Q_6\phi_1$ where $\phi_1 = z_1\bar{z}_2 + z_2\bar{z}_1 + z_2\bar{z}_3 + z_3\bar{z}_2 + z_3\bar{z}_1 + z_1\bar{z}_3$ is the first eigenfunction of the Laplacian of $(\mathbb{C}P^2, g_{FS})$ with eigenvalue is 12. By using expressions computed above, the equation we need to solve is:

$$12^3 + 2\lambda \cdot 144 - 8 \left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25} \right) \cdot 12 = 60.48\lambda^3 - 6.72\lambda^2 + 63.36\lambda - 34.56.$$

Solving this equation numerically, we find a positive real root $\lambda_0 \approx 15.206$.

It is known that $\int_{\mathbb{C}P^2} \phi_1^3 \neq 0$ by [32]. Using Fubini's theorem, we get

$$\int_{S^2(\lambda_0) \times \mathbb{C}P^2} (1 \otimes \phi_1)^3 dV_g = Vol(S^2(\lambda_0)) \int_{\mathbb{C}P^2} \phi_1^3 \neq 0.$$

Thus, $(S^2(\lambda_0) \times \mathbb{C}P^2, g_{prod})$ is degenerate and the function F_3 is not everywhere zero on Λ_0 . This demonstrates that $(S^2(\lambda_0) \times \mathbb{C}P^2, g_{prod})$ satisfies AS_3 condition.

Next, we prove that for such λ_0 , $P_6 \geq 0$ with the kernel consisting of constant functions. We compute:

$$\begin{aligned}
P_6 &= -\Delta^3 - 2(\lambda - 12)\Delta^{(1)}\Delta + 2\lambda\Delta^{(2)}\Delta \\
&\quad + 8\left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25}\right)\Delta^{(1)} + 8\left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25}\right)\Delta^{(2)} \\
&= -\Delta^3 - 2(\lambda - 12)(\Delta^{(1)})^2 + 24\Delta^{(1)}\Delta^{(2)} + 2\lambda(\Delta^{(2)})^2 \\
&\quad + 8\left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25}\right)\Delta^{(1)} + 8\left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25}\right)\Delta^{(2)} \\
&\geq -(\Delta^{(1)})^3 - (\Delta^{(2)})^3 - 2(\lambda - 12)(\Delta^{(1)})^2 + 2\lambda(\Delta^{(2)})^2 \\
&\quad + 8\left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25}\right)\Delta^{(1)} + 8\left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25}\right)\Delta^{(2)} \\
&= -\Delta^{(1)}[(\Delta^{(1)})^2 + 2(\lambda - 12)\Delta^{(1)} - 8\left(-\frac{9}{50}\lambda^2 + \frac{42}{25}\lambda - \frac{198}{25}\right)] \\
&\quad - \Delta^{(2)}[(\Delta^{(2)})^2 - 2\lambda\Delta^{(2)} - 8\left(\frac{7}{100}\lambda^2 - \frac{33}{25}\lambda + \frac{27}{25}\right)].
\end{aligned}$$

We observe that $(2\lambda_0)^2 - 2(\lambda_0 - 12)2\lambda_0 - 8\left(-\frac{9}{50}\lambda_0^2 + \frac{42}{25}\lambda_0 - \frac{198}{25}\right) > 0$ and $12^2 - 2\lambda_0 * 12 - 8\left(\frac{7}{100}\lambda_0^2 - \frac{33}{25}\lambda_0 + \frac{27}{25}\right) > 0$. Hence $P_6 \geq 0$ and it is easy to see that the kernel consists of constant functions.

The total Q_6 -curvature is

$$\begin{aligned}
\int_{S^2(\lambda_0) \times CP^2} Q_6 dV_g &= (0.48\lambda_0^3 - 6.72\lambda_0^2 + 63.36\lambda_0 - 34.56) \cdot Vol(S^2(\lambda_0)) \cdot Vol(CP^2) \\
&= (0.48\lambda_0^3 - 6.72\lambda_0^2 + 63.36\lambda_0 - 34.56) \cdot \frac{4\pi}{\lambda_0} \cdot \frac{\pi^2}{2} \\
&< 144\pi^3
\end{aligned}$$

where $144\pi^3$ is the total Q_6 -curvature of the standard sphere S^6 . This confirms that the example satisfies the conditions of Theorem 1.3.

6. COMPARISON OF CASES FOR $n = 2, 4$ AND 6 .

In this section, we briefly discuss some differences in low-dimensional cases in terms of whether the operator \mathcal{L}_∞ has a non-trivial kernel.

When $n = 2$, i.e. (M, g) is a closed Riemannian manifold of dimension 2, the Q -curvature Q_g is simply the Gauss curvature K_g . Moreover, the Paneitz operator P_g is just but the negative Laplacian, ie. $P_g = -\Delta_g$. Therefore, the Q -curvature flow reduces to the two dimensional normalized Ricci flow:

$$(6.1) \quad \frac{\partial}{\partial t} g(t) = -(K_{g(t)} - \bar{K}_{g(t)})g(t).$$

The operator \mathcal{L}_∞ is given by

$$\mathcal{L}_\infty = -\Delta_{g_\infty} - 2K_{g_\infty}.$$

If $K_{g_\infty} \leq 0$, then $\ker \mathcal{L}_{g_\infty}$ is trivial. If $K_{g_\infty} \equiv \text{constant} > 0$, then (M, g) is either the standard sphere or the real projective plane. The standard sphere is integrable by Obata's theorem and for the real projective plane, \mathcal{L}_∞ is positive definite as the first eigenfunctions of Laplacian on the sphere are odd functions. Hence, we can easily observe that the 2-dimensional Q -curvature flow always converges exponentially.

For $n = 4$, we have the following explicit formulas for the fourth-order Paneitz operator and the Q_4 -curvature for (M, g_∞) :

$$(Q_4)_{g_\infty} = \frac{1}{6}(-\Delta_{g_\infty} R_{g_\infty} + R_{g_\infty}^2 - 3|Ric_{g_\infty}|^2)$$

$$(P_4)_{g_\infty} = \Delta_{g_\infty}^2 + \delta\left(\frac{2}{3}R_{g_\infty}g_\infty - 2Ric_{g_\infty}\right)d.$$

Whether the operator $\mathcal{L}_\infty = (P_4)_{g_\infty} - 4(Q_4)_\infty$ is integrable or non-negative is unknown. This is related to the uniqueness of solutions to the constant Q -curvature equation. If one proves that, for a Riemannian manifold (M, g) with $(Q_4)_g \equiv \text{constant} > 0$ and $R_g > 0$, the inequality

$$(6.2) \quad \int_M (P_4)_g \phi \cdot \phi dVol_g \geq 4(Q_4)_g \int_M \phi^2 dVol_g \text{ for } \int_M \phi = 0$$

is true with equality holding if and only if $\phi \equiv 0$ or (M, g) is the standard round sphere, then the solution for constant Q_4 -curvature equation with the constraint that the scalar curvature is positive is unique up to constants by the degree theory, as observed in [34, Section 5]. The uniqueness problem of the constant Q_4 -curvature equation is only solved when g_∞ is Einstein with a positive Einstein constant by [46], and the inequality (6.2) is true by the eigenvalue comparison theorem. Hence the Q_4 -curvature flow is converging exponentially to g_∞ when g_∞ is Einstein with a positive Einstein constant.

Example 5.1 indicates that we cannot expect such phenomena in dimension $n = 6$. Specifically, for $(S^2(\lambda_0) \times \mathbb{C}P^2, g_{prod})$, it is easy to see that

$$R_{g_{prod}} > 0$$

$$(Q_4)_{g_{prod}} = -\Delta_{g_{prod}} J_{g_{prod}} - 2|A_{g_{prod}}|^2 + 3J_{g_{prod}}^2 = -0.06\lambda_0^2 + 4.56\lambda_0 + 936 > 0$$

$$(Q_6)_{g_{prod}} > 0$$

while the operator $P_6 - 6Q_6$ has a non-constant function in the kernel.

7. ACKNOWLEDGEMENT

The first author was supported by the National Science and Technology Council (NSTC), Taiwan, with grant number: 112-2115-M-032-006-MY2. The second author was supported by KIAS Individual Grant, with grant number: MG096101.

REFERENCES

1. D. Adams and L. Simon, Rates of asymptotic convergence near isolated singularities of geometric extrema, *Indiana Univ. Math. J.* **37** (1988) 225-254
2. T. Aubin, Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire. *J. Math. Pures Appl. (9)* **55** (1976), 269-296.
3. S. Azami and A. Razavi, Yamabe flow on Berwald manifolds. *Int. J. Geom. Methods Mod. Phys.* **12** (2015), no. 2, 1550025, 27 pp.
4. J. Bergh and J. Löfström, Interpolation spaces. An introduction. *Grundlehren der Mathematischen Wissenschaften*, No. **223**. Springer-Verlag, Berlin-New York, 1976.
5. S. Brendle, Convergence of the Q -curvature flow on S^4 . *Adv. Math.* **205** (2006), no. 1, 1–32.
6. S. Brendle, Convergence of the Yamabe flow for arbitrary initial energy. *J. Differential Geom.* **69** (2005), 217-278.
7. S. Brendle, Convergence of the Yamabe flow in dimension 6 and higher. *Invent. Math.* **170** (2007), 541-576.
8. S. Brendle, Global existence and convergence for a higher order flow in conformal geometry. *Ann. of Math. (2)* **158** (2003), no. 1, 323–343.

9. C. Carlotto, O. Chodosh, and Y. Rubinstein, Slowly converging Yamabe flows. *Geom. Topol.* **19** (2015), 1523-1568.
10. S.-Y. A. Chang and P. C. Yang, Extremal metrics of zeta function determinants on 4-manifolds. *Ann. of Math. (2)* **142** (1995), no. 1, 171–212.
11. X. Chen and P. T. Ho, Conformal curvature flows on a compact manifold of negative Yamabe constant. *Indiana Univ. Math. J.* **67** (2018), no. 2, 537-581.
12. X. Chen and X. Xu, Q -curvature flow on the standard sphere of even dimension. *J. Funct. Anal.* **261** (2011), no. 4, 934–980.
13. L. Cheng and A. Zhu, Yamabe flow and ADM mass on asymptotically flat manifolds. *J. Math. Phys.* **56** (2015), no. 10, 101507, 21 pp.
14. R. Chill, On the Łojasiewicz-Simon gradient inequality. *J. Funct. Anal.* **201** (2003), 572–601.
15. B. Chow, The Yamabe flow on locally conformally flat manifolds with positive Ricci curvature. *Comm. Pure Appl. Math.* **45** (1992), 1003-1014.
16. F. Daneshvar and A. Razavi, A class of monotonic quantities along the Yamabe flow. *Bull. Belg. Math. Soc. Simon Stevin* **27** (2020), no. 1, 17–27.
17. Z. Djadli and A. Malchiodi, Existence of conformal metrics with constant Q -curvature. *Ann. of Math. (2)* **168** (2008), no. 3, 813–858.
18. H. Dong and H. Zhang, Schauder estimates for higher-order parabolic systems with time irregular coefficients. *Calc. Var. Partial Differential Equations* **54** (2015), no. 1, 47–74.
19. C. Fefferman and C. R. Graham, Conformal invariants, in *The Mathematical Heritage of Élie Cartan* (Lyon, 1984), *Astérisque* (1985), 95–116.
20. C. Fefferman, C. R. Graham, Q -curvature and Poincaré metrics. *Math. Res. Lett.* **9** (2002), 139–151.
21. P. T. Ho, Backwards uniqueness of the Yamabe flow. *Differential Geom. Appl.* **62** (2019), 184–189.
22. P. T. Ho, First eigenvalues of geometric operators under the Yamabe flow. *Ann. Global Anal. Geom.* **54** (2018), no. 4, 449–472.
23. P. T. Ho, Prescribed Q -curvature flow on S^n . *J. Geom. Phys.* **62** (2012), no. 5, 1233–1261.
24. P. T. Ho, Prescribing Q -curvature on S^n in the presence of symmetry. *Commun. Pure Appl. Anal.* **19** (2020), no. 2, 715–722.
25. P. T. Ho, Q -curvature flow on S^n . *Comm. Anal. Geom.* **18** (2010), no. 4, 791–820.
26. P. T. Ho, Results of prescribing Q -curvature on S^n . *Arch. Math. (Basel)* **100** (2013), no. 1, 85–93.
27. P. T. Ho, The Gauss-Bonnet-Chern mass under geometric flows. *J. Math. Phys.* **61** (2020), no. 11, 112501, 9 pp.
28. P. T. Ho and J. Shin, Slowly converging Yamabe-type flow on manifolds with boundary. *Commun. Contemp. Math.* **26** (2024), no. 3, Paper No. 2250072.
29. P. T. Ho, J. Shin, and Z. Yan, Convergence rate of the weighted Yamabe flow. *Differential Geom. Appl.* **93** (2024), Paper No. 102119.
30. A. Juhl, Families of Conformally Covariant Differential Operators, Q -Curvature and Holography, Progress in Mathematics, vol. 275, Birkhäuser Verlag, Basel (2009).
31. A. Juhl, Explicit formulas for GJMS-operators and Q -curvatures, *Geom. Funct. Anal.* **23** (2013), no. 4, 1278–1370.
32. K. Kröncke, Stability of Einstein metrics under Ricci flow, *Comm. Anal. Geom.*, **28** (2020), no.2, 351–394.
33. O. Lablee, Eigenvalues for a Schrödinger operator on a closed Riemannian manifold with holes. *Funct. Anal. Approx. Comput.* **5** (2013), no. 2, 43-58.
34. S. Lee, Some Uniformization Problems for a Fourth order Conformal Curvature, *arXiv:2305.08027*, preprint.
35. G. M. Lieberman, Second order parabolic differential equations, World Scientific, River Edge, NJ (1996)
36. S. Łojasiewicz, Une propriété topologique des sous-ensembles analytiques réels. from: “Les Équations aux Dérivées Partielles”, Éditions du Centre National de la Recherche Scientifique, Paris (1963), 87–89.
37. L. Ma and L. Cheng, Yamabe flow and Myers type theorem on complete manifolds. *J. Geom. Anal.* **24** (2014), no. 1, 246–270.
38. A. Malchiodi and M. Struwe, Q -curvature flow on S^4 . *J. Differential Geom.* **73** (2006), no. 1, 1–44.

39. C. B. Ndiaye, Constant Q -curvature metrics in arbitrary dimension. *J. Funct. Anal.* **251** (2007), no. 1, 1–58.
40. R. Schoen, Conformal deformation of a Riemannian metric to constant scalar curvature. *J. Differential Geom.* **20** (1984), 479–495.
41. M. B. Schulz, Instantaneously complete Yamabe flow on hyperbolic space. *Calc. Var. Partial Differential Equations* **58** (2019), no. 6, Paper No. 190, 30 pp.
42. H. Schwetlick and M. Struwe, Convergence of the Yamabe flow for “large” energies. *J. Reine Angew. Math.* **562** (2003), 59–100.
43. L. Simon, Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems. *Ann. of Math. (2)* **118** (1983), 525–571.
44. L. Simon, Theorems on regularity and singularity of energy minimizing maps, Birkhäuser, Basel (1996)
45. N. S. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds. *Ann. Scuola Norm. Sup. Pisa (3)* **22** (1968), 265–274.
46. J. Vétois, Uniqueness of Conformal Metrics with Constant Q -Curvature on Closed Einstein Manifolds, *Potential Anal.* (2023), <https://doi.org/10.1007/s11118-023-10117-1>.
47. J. Wei and X. Xu, On conformal deformations of metrics on S^n . *J. Funct. Anal.* **157** (1998), no. 1, 292–325.
48. R. Ye, Global existence and convergence of Yamabe flow. *J. Differential Geom.* **39** (1994), 35–50.
49. E. Zeidler, Nonlinear functional analysis and its applications, I : Fixed-point theorems, Springer, New York (1986)

DEPARTMENT OF MATHEMATICS, TAMKANG UNIVERSITY TAMSUI, NEW TAIPEI CITY 251301,
TAIWAN

Email address: paktungho@yahoo.com.hk

KOREA INSTITUTE FOR ADVANCED STUDY, HOEGIRO 85, SEOUL 02455, KOREA

Email address: s129@kias.re.kr