COLORING THE INTERSECTION OF TWO MATROIDS

ELI BERGER AND HE GUO

ABSTRACT. A result [1] from 2006 of Aharoni and the first author of this paper states that for any two natural numbers p, q, where p divides q, if a matroid \mathcal{M} is p-colorable and a matroid \mathcal{N} is q-colorable then $\mathcal{M} \cap \mathcal{N}$ is (p+q)-colorable. In this paper we show that the assumption that p divides q is in fact redundant, and we also prove that $\mathcal{M} \cap \mathcal{N}$ is even p + q list-colorable.

The result uses topology and relies on a new parameter yielding a lower bound for the topological connectivity of the intersection of two matroids.

1. INTRODUCTION

A hypergraph is a pair H = (V, E) where the vertex set V is a finite set and the edge set E is a set of subsets of V. For $U \subseteq V$, the induced subhypergraph of H on U is the hypergraph whose vertex set is U and whose edge set is $\{f \in E(H) \mid f \subseteq U\}$, and we denote the induced subhypergraph by H[U]. A set $X \subseteq V$ is called *independent* in H if there is no $e \in E$ such that $e \subseteq X$. We denote by $\mathcal{I}(H)$ the set of all independent sets in H.

A property of a set of the form $\mathcal{C} = \mathcal{I}(H)$ is that if $\sigma \in \mathcal{C}$ and $\tau \subseteq \sigma$ then $\tau \in \mathcal{C}$. A finite set of finite sets with such closed under taking subsets property is called an *(abstract simplicial) complex*. The reversed operator for \mathcal{I} can be defined as follows.

Definition 1.1. Let C be an abstract simplicial complex on the set V. We write

 $circ(\mathcal{C}) = \{ e \subseteq V : e \notin \mathcal{C}, \ (\forall x \in e)(e \setminus \{x\} \in \mathcal{C}) \}.$

Observation 1.2. If C is an abstract simplicial complex on V then $C = \mathcal{I}((V, circ(C)))$.

Definition 1.3. Let H = (V, E) be a hypergraph. For an edge $e \in E$, we write

$$H - e = (V, E \setminus \{e\}).$$

For a set $X \subseteq V$, we write

$$H/X = (V \setminus X, \{e \setminus X \mid e \in E, e \not\subseteq X\}),$$

$$H \setminus X = (V \setminus X, \{e \in E \mid e \cap X = \emptyset\}),$$

$$H \sim X = (V, \{e \in E \mid e \cap X = \emptyset\}).$$

If $v \in V$ then we write $H \sim v = H \sim \{v\}$.

Note that $H \setminus X$ and $H \sim X$ differ only by their vertex sets.

Definition 1.4. Let H = (V, E) be a hypergraph, and suppose the following properties hold:

- $\emptyset \notin E$,
- there are no two distinct edges $C_1, C_2 \in E$ such that $C_1 \subseteq C_2$, and
- (circuit elimination axiom) for every two edges $C_1, C_2 \in E$ and every $u \in C_1 \cap C_2$ and $v \in C_1 \setminus C_2$ there exists $C_3 \in E$, such that $C_3 \subseteq C_1 \cup C_2$ and $v \in C_3$ and $u \notin C_3$.

Date: June 2024.

Then we say that H is a circuit representation of a matroid, and that $\mathcal{I}(H)$ is the independent set representation of this matroid. We call V the ground set of the matroid. The elements of E are called the circuits of the matroid and the elements of $\mathcal{I}(H)$ are called the independent sets of the matroid.

In this paper, when saying that \mathcal{M} is a matroid, we mean that \mathcal{M} is the set of independent sets of some matroid. However, when applying the operators / and \sim , we refer to the application of these operators on the circuit representation of the matroid. Formally, if \mathcal{M} is a matroid on the ground set V, whose circuit representation is $H = (V, circ(\mathcal{M}))$, and if $X \subseteq V$, then we define

$$\mathcal{M}[X] = \mathcal{I}(H[X])$$
 and $\mathcal{M}/X = \mathcal{I}(H/X)$

and if $v \in V$ then we define

$$\mathcal{M} \sim v = \mathcal{I}(H \sim v)$$
 .

Note that the above definition of \mathcal{M}/X coincides with the usual definition of contraction in matroids. Equivalent to Definition 1.4, \mathcal{M} is a matroid if the following hold:

- $\emptyset \in \mathcal{M}$.
- If $T \in \mathcal{M}$ and $S \subseteq T$, then $S \in \mathcal{M}$.
- (Independence augmentation property) If $S, T \in \mathcal{M}$ and |S| < |T|, then there exists $v \in T \setminus S$ such that $S \cup \{v\} \in \mathcal{M}$.

Definition 1.5. Given a complex C, the chromatic number $\chi(C)$ of C is the minimum number of sets in C such that their union is $\bigcup C$. The list chromatic number $\chi_{\ell}(C)$ of C is the minimum number ksuch that for any lists of colors $(L_v)_{v \in \bigcup C}$ of size k, there exists a choice function $f : \bigcup C \to \bigcup_{v \in \bigcup C} L_v$ such that $f(v) \in L_v$ for every v and $f^{-1}(c) \in C$ for every color c.

Setting $L_v = \{1, \ldots, \chi_\ell(\mathcal{C})\}$ for each v proves

$$\chi(\mathcal{C}) \le \chi_{\ell}(\mathcal{C}).$$

In [1], it is proved that for two matroids \mathcal{M} and \mathcal{N} on the same ground set, then

 $\chi(\mathcal{M} \cap \mathcal{N}) \le 2 \max(\chi(\mathcal{M}), \chi(\mathcal{N})).$

And it is also proved that if $\chi(\mathcal{M})$ divides $\chi(\mathcal{N})$, then

$$\chi(\mathcal{M} \cap \mathcal{N}) \le \chi(\mathcal{M}) + \chi(\mathcal{N}).$$

In this paper, we extend these results.

Theorem 1.6. For two matroids \mathcal{M} and \mathcal{N} on the same ground set,

$$\chi_{\ell}(\mathcal{M} \cap \mathcal{N}) \le \chi(\mathcal{M}) + \chi(\mathcal{N})$$

2. The topological parameter η

Let \mathcal{C} be an abstract simplicial complex. Assuming we fix some ring R, we can apply homology theory on \mathcal{C} . We write $\eta(\mathcal{C})$ for the minimal value of k such that the reduced homology $\tilde{H}_{k-1}(\mathcal{C}, R)$ does not vanish. If \mathcal{C} is the empty complex then we write $\eta(\mathcal{C}) = 0$ and if all reduced homology groups of \mathcal{C} vanish then we write $\eta(\mathcal{C}) = \infty$.

For two complexes \mathcal{C}, \mathcal{D} on the disjoint sets, the *join* $\mathcal{C} * \mathcal{D}$ is $\{S \cup T \mid S \in \mathcal{C}, T \in \mathcal{D}\}$.

Theorem 2.1. [1] Let C and D be two abstract simplicial complexes on the disjoint sets. Then

$$\eta(\mathcal{C} * \mathcal{D}) \ge \eta(\mathcal{C}) + \eta(\mathcal{D}).$$

For two complexes on the same set, we can prove the following inequalities using the Mayer–Vietoris sequence:

Theorem 2.2. Let \mathcal{A} and \mathcal{B} be two abstract simplicial complexes on the same set V. Then

- (1) $\eta(\mathcal{A} \cup \mathcal{B}) \geq \min(\eta(\mathcal{A}), \eta(\mathcal{B}), \eta(\mathcal{A} \cap \mathcal{B}) + 1).$
- (2) $\eta(\mathcal{A} \cap \mathcal{B}) \ge \min(\eta(\mathcal{A}), \eta(\mathcal{B}), \eta(\mathcal{A} \cup \mathcal{B}) 1).$
- (3) $\eta(\mathcal{A}) \geq \min(\eta(\mathcal{A} \cup \mathcal{B}), \eta(\mathcal{A} \cap \mathcal{B})).$

Using inequality (1) for $\mathcal{A} = \mathcal{I}(H \setminus \{v\})$ and $\mathcal{B} = 2^{\{v\}} * \mathcal{I}(H \setminus (\{v\} \cup N_H(v)))$ of Theorem 2.2, where 2^S is the power set of the set S and $N_H(v)$ is the set of neighbors of v in H, we can deduce the following

Theorem 2.3. Let H = (V, E) be a hypergraph and let v be a vertex of H such that $\{v\} \notin E$. Then

$$\eta(\mathcal{I}(H)) \ge \min\left(\eta\left(\mathcal{I}(H \setminus \{v\})\right), \ \eta\left(\mathcal{I}(H/\{v\})\right) + 1\right)$$

Using inequality (3) for $\mathcal{A} = \mathcal{I}(H)$ and $\mathcal{B} = 2^e * \mathcal{I}(H/e)$ of Theorem 2.2 we can also deduce the following (see [2, Section 8.3] for the details of the proof.)

Theorem 2.4. Let H be a hypergraph and let e be an edge of H which does not contain any other edge. Then

$$\eta(\mathcal{I}(H)) \ge \min\left(\eta(\mathcal{I}(H-e)), \ \eta(\mathcal{I}(H/e)) + |e| - 1\right)$$

Theorem 2.4 was proved in [4] for the case that H is a graph, but the same proof holds for general hypergraphs as well. While Theorem 2.4 is extensively used for graphs, its use for hypergraphs is less common so far. It is used implicitly in [3].

Repeatedly applying Theorem 2.3 and Theorem 2.4 gives a very powerful tool for obtaining lower bounds for η .

Theorem 2.5. Let $\mathcal{M}_1, \ldots, \mathcal{M}_k$ be matroids on the ground set V, and let $v \in V$. Then either

$$\eta(\mathcal{M}_1 \cap \ldots \cap \mathcal{M}_k) \ge \eta((\mathcal{M}_1 \sim v) \cap \mathcal{M}_2 \cap \ldots \cap \mathcal{M}_k)$$

or there exists C such that

- (1) $v \in C$.
- (2) C is a circuit of \mathcal{M}_1 .
- (3) $C \in \bigcap_{i=2}^k \mathcal{M}_i$.
- (4) $\eta(\mathcal{M}_1 \cap \ldots \cap \mathcal{M}_k) \ge \eta((\mathcal{M}_1/C) \cap (\mathcal{M}_1/C) \cap \ldots \cap (\mathcal{M}_k/C)) + |C| 1.$

Proof. Let $H_i = (V, E_i) = (V, circ(\mathcal{M}_i))$ be the circuit representation of \mathcal{M}_i for $i = 1, \ldots, k$. For $I \subseteq \{1, \ldots, k\}$, let $\bigcup_{i \in I} H_i = (V, \bigcup_{i \in I} E_i)$ and let $H = \bigcup_{i=1}^k H_i$. So $\mathcal{I}(H) = \bigcap_{i=1}^k \mathcal{M}_i$.

Let C_1, \ldots, C_t be all the circuits of \mathcal{M}_1 satisfying that $v \in C_j$ and $C_j \in \bigcap_{i=2}^k \mathcal{M}_i$ for each $1 \leq j \leq t$.

Claim 2.6.

(1)
$$\mathcal{I}(H - C_1 - \dots - C_t) = (\mathcal{M}_1 \sim v) \cap \mathcal{M}_2 \cap \dots \cap \mathcal{M}_k,$$

(2)
$$\mathcal{I}((H-C_1-\cdots-C_{j-1})/C_j) = \mathcal{I}(H/C_j),$$

(3)
$$\mathcal{I}(H/C_j) = (\mathcal{M}_1/C_j) \cap \cdots \cap (\mathcal{M}_k/C_j).$$

Suppose the claim is true, then applying Theorem 2.4 repeatedly, where each C_j in turn takes the role of e in the theorem, yields either by (1) that

$$\eta(\mathcal{I}(H)) \ge \eta(\mathcal{I}(H - C_1 - \dots - C_t)) = \eta\Big((\mathcal{M}_1 \sim v) \cap \mathcal{M}_2 \cap \dots \cap \mathcal{M}_k\Big),$$

or for some $1 \le j \le t$ by (2) and (3) that

$$\eta(\mathcal{I}(H)) \ge \eta \Big(\mathcal{I}\Big((H - C_1 - \dots - C_{j-1})/C_j \Big) \Big) + |C_j| - 1$$
$$= \eta \Big(\mathcal{I}(H/C_j) \Big) + |C_j| - 1$$
$$= \eta \Big((\mathcal{M}_1/C_j) \cap \dots \cap (\mathcal{M}_k/C_j) \Big) + |C_j| - 1.$$

Setting $C = C_j$ completes the proof.

Proof of the claim. To verify (1), for S in the LHS, S does not contain edges of $\bigcup_{i=2}^{k} H_i$, therefore $S \in \bigcap_{i=2}^{k} \mathcal{M}_i$. And $S \setminus \{v\}$ does not contain any edge $f \in H_i$ satisfying $f \subseteq V \setminus \{v\}$, since such $f \neq C_j$ for every $1 \leq j \leq t$. Therefore $S \setminus \{v\} \in \mathcal{M}_1[V \setminus \{v\}]$ and $S \in \mathcal{M} \sim v$. Thus the LHS is contained in the RHS. For T in the RHS, $T \in \bigcap_{i=2}^{k} \mathcal{M}_i$ implies T is independent in $\bigcup_{i=2}^{k} H_i$. And T does not contain any edge of $H_1[V \setminus \{v\}]$. Furthermore, T does not contain any $C \in H_1$ such that $v \in C$ and $C \neq C_j$ for every $1 \leq j \leq t$. Suppose not, then such C must contain an edge of some H_i for $2 \leq i \leq k$ as C_1, \ldots, C_t are all the edges of H_1 including v and containing no edge of $\bigcup_{i=2}^{t} H_i$, which contradicts to the assumption $T \in \mathcal{M}_i$. Therefore the RHS is contained in the LHS.

To verify (2), the RHS is contained in the LHS, since $(H - C_1 - \cdots - C_{j-1})/C_j$ and H/C_j have the same vertex set and the edge set of the former is contained in that of the latter. And for any S in the LHS, we claim that S does not contain $C_{\ell} \setminus C_j$ for any $1 \le \ell \le j - 1$: suppose not, i.e., $C_{\ell} \setminus C_j \subseteq S$ for some $1 \le \ell \le j - 1$. Since $v \in C_j \cap C_{\ell}$, then by the circuit elimination axiom, there exists a circuit C' of \mathcal{M}_1 such that $C' \subseteq (C_j \cup C_\ell) \setminus \{v\}$. Then S contains $C' \setminus C_j$, contradicting the assumption $S \in \mathcal{I}(H/C_j)$. Therefore the LHS is contained in the RHS.

To verify (3), since $C_j \in \bigcap_{i=2}^k \mathcal{M}_i$, no edge of $\bigcup_{i=2}^k H_i$ is contained in C_j . And since $C_j \in circ(\mathcal{M}_1)$, no edge of H_1 other than C_j itself is contained in C_j . Therefore

$$H/C_j = (\bigcup_{i=1}^k H_i)/C_j = \bigcup_{i=1}^k (H_i/C_j).$$

By definition $\mathcal{I}(H_i/C_j) = \mathcal{M}_i/C_j$ for each i = 1, ..., k, therefore $\mathcal{I}(H/C_j) = \bigcap_{i=1}^k (\mathcal{M}_i/C_j)$.

3. The combinatorial parameter $\nu_{p,q}(\mathcal{M},\mathcal{N})$

For sets A_1, \ldots, A_k and an element v, we write $\#v(A_1, \ldots, A_k)$ for the number of sets among A_1, \ldots, A_k to which v belongs. For two natural numbers p, q and two matroids \mathcal{M}, \mathcal{N} on the same ground set V we define

$$\nu_{p,q}(\mathcal{M},\mathcal{N}) = \max_{\substack{A_1,\ldots,A_p \in \mathcal{M} \\ B_1,\ldots,B_q \in \mathcal{N}}} \sum_{v \in V} \min\left(\#v(A_1,\ldots,A_p), \ \#v(B_1,\ldots,B_q) \right).$$

Note that $\nu_{1,1}(\mathcal{M},\mathcal{N})$ is the size of the largest set which is independent in both \mathcal{M} and \mathcal{N} .

Observation 3.1. The parameter ν is monotone in p, q, \mathcal{M} and \mathcal{N} , i.e., if p, q, p', q' are natural numbers and $\mathcal{M}, \mathcal{N}, \mathcal{M}', \mathcal{N}'$ are matroids such that $p \leq p', q \leq q', \mathcal{M} \subseteq \mathcal{M}'$ and $\mathcal{N} \subseteq \mathcal{N}'$ then $\nu_{p,q}(\mathcal{M}, \mathcal{N}) \leq \nu_{p',q'}(\mathcal{M}', \mathcal{N}')$

Observation 3.2. For any two natural numbers p, q and two matroids \mathcal{M}, \mathcal{N} on the same ground set V, there exist sets $A_1, \ldots, A_p \in \mathcal{M}$ and $B_1, \ldots, B_q \in \mathcal{N}$ such that $\sum_{i=1}^p |A_i| = \sum_{j=1}^q |B_j| = \nu_{p,q}(\mathcal{M}, \mathcal{N})$ and every $v \in V$ satisfies $\#v(A_1, \ldots, A_p) = \#v(B_1, \ldots, B_q)$.

Lemma 3.3. If \mathcal{M}, \mathcal{N} are matroids on a common ground set then for every natural number q

$$\nu_{1,1}(\mathcal{M},\mathcal{N}) \ge \lceil \frac{\nu_{q,q}(\mathcal{M},\mathcal{N})}{q} \rceil.$$

Proof. Let V be the common ground set. By Edmonds' matroid intersection theorem there exist sets $I, V_1, V_2 \subseteq V$ such that $V = V_1 \cup V_2$ and $V_1 \cap V_2 = \emptyset$ and $I \in \mathcal{M} \cap \mathcal{N}$ and $I \cap V_1$ is the largest among subsets of V_1 which are independent in \mathcal{M} , and $I \cap V_2$ is the largest among subsets of V_2 which are independent in \mathcal{N} .

Now for every $A_1, \ldots, A_q \in \mathcal{M}$ and $B_1, \ldots, B_q \in \mathcal{N}$ we have

$$\sum_{v \in V} \min \left(\#v(A_1, \dots, A_q), \ \#v(B_1, \dots, B_q) \right)$$

= $\sum_{v \in V_1} \min \left(\#v(A_1, \dots, A_q), \ \#v(B_1, \dots, B_q) \right) + \sum_{v \in V_2} \min \left(\#v(A_1, \dots, A_q), \ \#v(B_1, \dots, B_q) \right)$
 $\leq \sum_{v \in V_1} \#v(A_1 \cap V_1, \dots, A_q \cap V_1) + \sum_{v \in V_2} \#v(B_1 \cap V_2, \dots, B_q \cap V_2)$
 $\leq q \cdot |I \cap V_1| + q \cdot |I \cap V_2| = q|I| \leq q\nu_{1,1}(\mathcal{M}, \mathcal{N}),$

which completes the proof.

Lemma 3.4. Let \mathcal{M}, \mathcal{N} be two matroids on the same ground set V such that $\nu_{1,1}(\mathcal{M}, \mathcal{N}) > 0$. Let p, q be natural numbers with $p \leq q$. Then there exist sets $X_1, \ldots, X_p \in \mathcal{M}$ and $Y_1, \ldots, Y_q \in \mathcal{N}$ and an element $z \in V$ with the following properties:

• Every $v \in V \setminus \{z\}$ satisfies $\#v(X_1, \ldots, X_p) = \#v(Y_1, \ldots, Y_q)$,

•
$$#z(X_1,\ldots,X_p)=p,$$

• $\sum_{i=1}^{q} |Y_i| = \nu_{p,q}(\mathcal{M}, \mathcal{N}).$

Proof. Write $r = \nu_{1,1}(\mathcal{M}, \mathcal{N})$. Let $Z \in \mathcal{M} \cap \mathcal{N}$ have size r, and let $A_1, \ldots, A_p \in \mathcal{M}$ and $B_1, \ldots, B_q \in \mathcal{N}$ be chosen such that

- (1) $\sum_{v \in V} \min \left(\# v(A_1, \dots, A_p), \ \# v(B_1, \dots, B_q) \right) = \nu_{p,q}(\mathcal{M}, \mathcal{N}) 1,$
- (2) Subject to the above condition $\sum_{v \in \mathbb{Z}} \min (\#v(A_1, \ldots, A_p), \#v(B_1, \ldots, B_q))$ is maximal,
- (3) Subject to the above two conditions $\sum_{i=1}^{p} |A_i| + \sum_{j=1}^{q} |B_j|$ is minimal.

Note that $\sum_{i=1}^{p} |A_i| + \sum_{j=1}^{q} |B_j|$ is minimal in condition (3) guarantees that for every $v \in V$,

(4)
$$\#v(A_1,\ldots,A_p) = \#v(B_1,\ldots,B_q),$$

since if $\#v(A_1, \ldots, A_p) > \#v(B_1, \ldots, B_q)$ for some $v \in V$, we can remove v from some of A_i including it, which does not violate conditions (1) and (2), but has a smaller total size, a contradiction. Similarly, we can get a contradiction if $\#v(A_1, \ldots, A_p) < \#v(B_1, \ldots, B_q)$ for some $v \in V$.

Therefore we have

$$\sum_{j=1}^{q} |B_j| = \sum_{v \in V} \#v(B_1, \dots, B_q) = \nu_{p,q}(\mathcal{M}, \mathcal{N}) - 1 < \nu_{p,q}(\mathcal{M}, \mathcal{N}) \le \nu_{q,q}(\mathcal{M}, \mathcal{N}) \le qr,$$

which implies at least one set among B_1, \ldots, B_q has size less than r. Without loss of generality $|B_q| < r$. By the independence augmentation property, this implies that for some $z \in Z \setminus B_q$ we have $B'_q = B_q \cup \{z\} \in \mathcal{N}$.

We now claim that for every $i \in \{1, ..., p\}$ we have $A_i \cup \{z\} \in \mathcal{M}$. Suppose not, then, say, $A_p \cup \{z\} \notin \mathcal{M}$, so it contains some circuit C of \mathcal{M} and $z \in C$. Take some $c \in C \setminus Z$. Then $c \in A_p$. By the independence augmentation property, $C \setminus \{c\} \in \mathcal{M}$ can be extended to a size $|A_p|$ set $A'_p = (A_p \cup \{z\}) \setminus \{c\} \in \mathcal{M}$. Since

$$#c(A_1, \ldots, A_{p-1}, A'_p) = 0 = #c(A_1, \ldots, A_{p-1}, A_p) - 1$$

and

$$\min\left(\#z(A_1,\ldots,A_{p-1},A'_p),\ \#z(B_1,\ldots,B_{q-1},B'_q)\right) = \min\left(\#z(A_1,\ldots,A_p),\ \#z(B_1,\ldots,B_q)\right) + 1,$$

by (4) we now have

$$\sum_{v \in V} \min \left(\# v(A_1, \dots, A_{p-1}, A'_p), \ \# v(B_1, \dots, B_{q-1}, B'_q) \right)$$

=
$$\sum_{v \in V} \min \left(\# v(A_1, \dots, A_{p-1}, A_p), \ \# v(B_1, \dots, B_{q-1}, B_q) \right) = \nu_{p,q}(\mathcal{M}, \mathcal{N}) - 1$$

and

$$\sum_{v \in Z} \min(\#v(A_1, \dots, A_{p-1}, A'_p), \#v(B_1, \dots, B_{q-1}, B'_q)) = \sum_{v \in Z} \min(\#v(A_1, \dots, A_p), \#v(B_1, \dots, B_q)) + 1$$

contradicting the way in which A_1, \ldots, A_q and B_1, \ldots, B_q were chosen. This finishes the proof of the claim.

Now the sets $X_i = A_i \cup \{z\}$ (for i = 1, ..., p) and $Y_j = B_j$ (for j = 1, ..., q-1) and $Y_q = B_q \cup \{z\}$ satisfy the required conditions of the lemma.

4. Relation between the parameters

Theorem 4.1. Let p, q be two natural numbers and let \mathcal{M}, \mathcal{N} be two matroids on the same ground set. Then

$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \frac{\nu_{p,q}(\mathcal{M}, \mathcal{N})}{p+q}$$

Proof. The proof is by induction on the size of the common ground set V. When $V = \emptyset$, $\eta(\mathcal{M} \cap \mathcal{N}) = \nu_{p,q}(\mathcal{M}, \mathcal{N}) = 0$. The statement is true.

Next we turn to $|V| \ge 1$. We assume without loss of generality $p \le q$ and by Lemma 3.4, there exist sets $X_1, \ldots, X_p \in \mathcal{M}$ and $Y_1, \ldots, Y_q \in \mathcal{N}$ and an element $z \in V$ with the following properties:

- Every $v \in V \setminus \{z\}$ satisfies $\#v(X_1, \ldots, X_p) = \#v(Y_1, \ldots, Y_q),$
- $#z(X_1,\ldots,X_p)=p,$
- $\sum_{i=1}^{q} |Y_i| = \nu_{p,q}(\mathcal{M}, \mathcal{N}).$

We apply Theorem 2.5 with \mathcal{M} taking the role of \mathcal{M}_1 . Then either

(5)
$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta((\mathcal{M} \sim z) \cap \mathcal{N})$$

or there is some circuit C in \mathcal{M} such that $z \in C, C \in \mathcal{N}$, and

(6)
$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta \big((\mathcal{M}/C) \cap (\mathcal{N}/C) \big) + |C| - 1$$

If (6) occurs, write $I = C \setminus \{z\}$ and s = |I| = |C| - 1. Let $A_1, \ldots, A_p \in \mathcal{M}$ and $B_1, \ldots, B_q \in \mathcal{N}$ satisfy $\nu_{p,q}(\mathcal{M}, \mathcal{N}) = \sum_{u \in V} \min \left(\#u(A_1, \ldots, A_p), \ \#u(B_1, \ldots, B_q) \right).$

We can find sets $S_1, \ldots, S_p, T_1, \ldots, T_q$, each of size s, such that $A_i \setminus S_i \in \mathcal{M}/C$ for all $i \in \{1, \ldots, p\}$ and $(B_j \setminus \{z\}) \setminus T_j \in \mathcal{N}/C$ for all $j \in \{1, \ldots, q\}$. In detail, to construct S_i , we take $A_i \cap C$ and add any $s - |A_i \cap C|$ other elements. Since C is a circuit in $\mathcal{M}, C \not\subseteq A_i$ and then $|A_i \cap C| < |C| = s + 1$. To construct T_i , we take $(B_j \setminus \{z\}) \cap C$ and add any $s - |(B_j \setminus \{z\}) \cap C|$ other elements.

Write $A'_i = A_i \setminus S_i$, which is in \mathcal{M}/C , for all $i \in \{1, \ldots, p\}$, and $B'_j = (B_j \setminus \{z\}) \setminus T_j$, which is in \mathcal{N}/C , for all $j \in \{1, \ldots, q\}$. We now claim that for each $u \in V$ we have

$$\min \left(\# u(A_1, \dots, A_p), \ \# u(B_1, \dots, B_q) \right)$$

$$\leq \min \left(\# u(A'_1, \dots, A'_p), \ \# u(B'_1, \dots, B'_q) \right) + \# u(S_1, \dots, S_p, T_1, \dots, T_q)$$

Indeed, when $u \neq z$ this is trivial, since $u \in A_i$ if and only if $u \in A'_i$ or S_i , and $u \in B_j$ if and only if $u \in B'_j$ or T_j ; and for u = z it follows from that fact that whenever $z \in A_i$ we must have also $z \in S_i$ so that

$$#z(A_1, \ldots, A_p) \le #z(S_1, \ldots, S_p, T_1, \ldots, T_q).$$

We thus have

$$\nu_{p,q}(\mathcal{M}, \mathcal{N}) = \sum_{u \in V} \min \left(\# u(A_1, \dots, A_p), \ \# u(B_1, \dots, B_q) \right)$$

$$\leq \sum_{u \in V} \left(\min \left(\# u(A'_1, \dots, A'_p), \ \# u(B'_1, \dots, B'_q) \right) + \# u(S_1, \dots, S_p, T_1, \dots, T_q) \right)$$

$$\leq \nu_{p,q}(\mathcal{M}/C, \mathcal{N}/C) + (p+q)s$$

and by the induction hypothesis

$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta\big((\mathcal{M}/C) \cap (\mathcal{N}/C)\big) + s \ge \frac{\nu_{p,q}(\mathcal{M}/C, \mathcal{N}/C)}{p+q} + s \ge \frac{\nu_{p,q}(\mathcal{M}, \mathcal{N})}{p+q}$$

If (5) occurs, applying Theorem 2.5 again to $\mathcal{M} \sim z$ and \mathcal{N} with \mathcal{N} taking the role of \mathcal{M}_1 , either

$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta((\mathcal{M} \sim z) \cap \mathcal{N}) \ge \eta((\mathcal{M} \sim z) \cap (\mathcal{N} \sim z)) \ge \eta(2^{\{z\}}) = \infty$$

Theorem 2.1

in which case we have $\eta(\mathcal{M} \cap \mathcal{N}) \geq \frac{\nu_{p+q}(\mathcal{M},\mathcal{N})}{p+q}$, or there exists a circuit D in \mathcal{N} such that $z \in D$ and $D \in \mathcal{M} \sim z$ and

$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta\Big((\mathcal{M} \sim z) \cap \mathcal{N}\Big) \ge \eta\Big(\big((\mathcal{M} \sim z)/D\big) \cap \big(\mathcal{N}/D\big)\Big) + |D| - 1.$$

In the last case, similar as above, we can find sets $S'_1, \ldots, S'_p, T'_1, \ldots, T'_q$, each of size t = |D| - 1, such that for each $1 \leq i \leq p, X'_i := (X_i \setminus \{z\}) \setminus S'_i \in (\mathcal{M} \sim z)/D$ using the fact that $z \in X_i$, and for each $1 \leq j \leq q, Y'_j := Y_j \setminus T'_j \in \mathcal{N}/D$. Thus

$$#u(Y_1,\ldots,Y_q) \le \min\left(#u(X'_1,\ldots,X'_p), \ #u(Y'_1,\ldots,Y'_q)\right) + #(S'_1,\ldots,S'_p,T'_1,\ldots,T'_q),$$

since for $u \neq z$, $\#u(Y_1, \ldots, Y_q) = \#u(X_1, \ldots, X_p)$, and for $u = z, z \in Y_j$ implies $z \in T'_j$. Therefore

$$\nu_{p,q}(\mathcal{M}, \mathcal{N}) = \sum_{i=1}^{q} |Y_i| = \sum_{u \in V} \#u(Y_1, \dots, Y_q)$$

$$\leq \sum_{u \in V} \left(\min\left(\#u(X'_1, \dots, X'_p), \ \#u(Y'_1, \dots, Y'_q) \right) + \#(S'_1, \dots, S'_p, T'_1, \dots, T'_q) \right)$$

$$\leq \nu_{p,q} \left((\mathcal{M} \sim z) / D, \mathcal{N} / D \right) + (p+q)t,$$

and by the induction hypothesis,

$$\eta(\mathcal{M} \cap \mathcal{N}) \ge \eta\Big(\big((\mathcal{M} \sim z)/D\big) \cap \big(\mathcal{N}/D\big)\Big) + t \ge \frac{\nu_{p,q}\big((\mathcal{M} \sim z)/D, \mathcal{N}/D\big)}{p+q} + t \ge \frac{\nu_{p,q}(\mathcal{M}, \mathcal{N})}{p+q},$$

ch completes the proof.

which completes the proof.

5. Proof of Theorem 1.6

Notation 5.1. For a complex C on the ground set V, let $\Delta_{\eta}(C) = \max_{\emptyset \neq S \subseteq V(C)} \frac{|S|}{\eta(C|S|)}$.

Applying Theorem 4.2 in [1], for a complex C, it is proved in Corollary 8.6 of [1] that $\chi(C) \leq \lceil \Delta_{\eta}(C) \rceil$. In [2], this bound is extended to the list chromatic number.

Theorem 5.2. For a complex C, $\chi_{\ell}(C) \leq \lceil \Delta_{\eta}(C) \rceil$.

Proof of Theorem 1.6. Let V be the common ground set of \mathcal{M} and \mathcal{N} and let $\mathcal{C} = \mathcal{M} \cap \mathcal{N}$.

Let $p = \chi(\mathcal{M})$ and $q = \chi(\mathcal{N})$. Let $A_1, \ldots, A_p \in \mathcal{M}$ satisfying that $\bigcup_{i=1}^p A_i = V$ and let $B_1, \ldots, B_q \in \mathcal{M}$ satisfying that $\bigcup_{i=1}^q B_j = V$. Then

$$\min(\#v(A_1,\ldots,A_p),\#v(B_1,\ldots,B_q)) \ge 1$$

for every $v \in V$, which implies that $\nu_{p,q}(\mathcal{M}, \mathcal{N}) \geq |V|$. Thus Theorem 4.1 implies that $\eta(\mathcal{C}) \geq \frac{|V|}{p+q}$, which is equivalent to

$$\frac{|V|}{\eta(\mathcal{C})} \le p + q.$$

Noting that $\chi(\mathcal{M}[S]) \leq \chi(\mathcal{M})$ and $\chi(\mathcal{N}[S]) \leq \chi(\mathcal{N})$ for every $S \subseteq V$, the above argument works for any non-empty subset S of V, therefore we have

$$\Delta_{\eta}(\mathcal{C}) \le \chi(\mathcal{M}) + \chi(\mathcal{N}),$$

which together with Theorem 5.2 completes the proof.

References

- R. Aharoni and E. Berger. The intersection of a matroid and a simplicial complex. Trans. Amer. Math. Soc. 358 (2006), 4895–4917.
- [2] R. Aharoni, E. Berger, H. Guo, and D. Kotlar. Intersection of matroids. In preparation.
- [3] R. Aharoni, E. Berger, and O. Kfir. Acyclic systems of representatives and acyclic colorings of digraphs. J. Graph Theory 59 (2008), 177–189.
- [4] R. Meshulam. Domination numbers and homology. J. Combin. Theory Ser. A 102 (2003), 321–330.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA 31905, ISRAEL *Email address:* berger.haifa@gmail.com

FACULTY OF MATHEMATICS, TECHNION, HAIFA 32000, ISRAEL *Email address:* hguo@campus.technion.ac.il