

SYMMETRY GROUPS AND DEFORMATIONS OF SUMS OF EXPONENTIALS

FLORIAN PAUSINGER AND DAVID PETRECCA

ABSTRACT. We study the symmetry groups and winding numbers of planar curves obtained by taking the image of the complex unit circle under a Laurent polynomial, a class that includes weighted sums of exponentials. We generalize various results on such sums of exponentials and relate them to other classes of curves present in the literature. Moreover, we consider the evolution under the wave equation of such curves for the case of binomials. Interestingly, our methods provide a unified and systematic way of constructing curves with prescribed properties, such as number of cusps, number of intersection points or winding number.

1. INTRODUCTION

In the late nineties, Farris [4] considered curves obtained through a *roulette*-like construction. They are the trajectories traced by a point on a wheel rolling at a certain speed that in turn rolls with another speed on another wheel of different radius. A weighted sum of exponentials

$$\sum_k w_k \exp(2\pi i a_k t)$$

can be used to describe such a construction in which the a_k are real exponents and the weights w_k are positive.

Farris noticed the symmetry of such curves, which motivated his later book [5], where he introduces the concept of a *rosette*. A rosette is a planar periodic curve admitting dihedral symmetry which is obtained from the image of the unit circle under a Laurent polynomial.¹

Several well-known curves can be included in this framework, such as roses or Rhodonea curves [3] or, closer to the original “wheels on wheels” constructions, the curves that can be drawn using the Spirograph™ device.

The former are obtained by polar plots of the form $\rho = \sin(n\theta)$ that can be seen as the image of the unit circle $t \mapsto \exp(2\pi i t)$ under the Laurent polynomial

$$f(z) = \frac{z^{1+n} - z^{1-n}}{2i}.$$

Such framework allows rephrasing neatly some open questions of Maurer [9]. The authors plan to address them in a future paper.

Date: July 15, 2024.

2020 Mathematics Subject Classification. 53A04, 33B10.

Key words and phrases. planar curves, symmetry groups, Laurent polynomials, wave equation.

¹This should not be confused with other definitions of rosette present in the literature that require convexity and/or positivity of the curvature as planar differentiable curves.

The latter examples are the trajectories of a point on a toothed wheel of radius r rolling along the interior or exterior of another of radius $R > r$. They can be seen as the image of the circle under the polynomial

$$p(z) = (R - r)z^r + rz^{R-r}$$

if the two radii are positive integers.

In two recent articles [10, 11] the first author together with Vartziotis studied the planar curves obtained by the sum of two complex exponentials. In particular, results about the symmetry groups of their graph, a description of the self-intersections as well as some differential geometric properties such as winding numbers and presence of cusps were obtained. Several questions were left open, such as proper generalizations to arbitrary sums of exponentials or the study of the case with irrational exponentials.

In this paper, we extend the work from [10, 11] and answer some open questions by exploring the connections with the more general rosette notion introduced by Farris.

We take a look at the more general level of Laurent series f defined on the complex plane, including weighted sums of exponentials and complex polynomials. We study the image $f(S^1)$ of the unit circle under such functions and, in particular, we study the symmetries of the image $f(S^1)$ and their effect on the zeros and poles of f . We use a notion of symmetry introduced by Farris [5] and relate our results to results of Quine [13] for the study of self-intersections.

In more detail, we answer in Theorem 1 and Prop. 1 the question stated in [10] about the symmetry group of the image of the unit circle through a Laurent series. Then we exhibit some examples of symmetries or lack thereof in case of polynomials, and finally we study the case of irrational exponents in Theorem 2 and Theorem 3 answering another question stated in [11]. In the context of the “wheels... on wheels” mechanical interpretation, this is an analog of the dense image of Lissajous curves in the case of incommensurable frequencies.

In addition, and following an idea of Farris [5], we investigate the evolution of our curves under the wave equation in Section 3. To simplify calculations, we restrict to the case of polynomials with two terms and study the variation of the winding number along such flow. We provide in Theorem 5 an analog of the winding number result proven in [11]. Moreover, the evolving curves can give, at special times that make the polynomial coefficients not integral, various examples of self-intersections and arising of cusps as investigated in Section 4.

2. SYMMETRY GROUPS

Farris [5] gives the following definition.

Definition 1. Let $k < m$ be a pair of coprime positive integers. A function $f: \mathbb{R} \rightarrow \mathbb{C}$ periodic of period 1 has symmetry of type (k, m) if

$$(1) \quad f\left(t + \frac{1}{m}\right) = \exp\left(2\pi i \frac{k}{m}\right) f(t)$$

for all $t \in \mathbb{R}$. Moreover, f has mirror symmetry along the axis generated by the line $\mathbb{R} \exp(i\sigma)$ if

$$(2) \quad f(-t) = \exp(2\pi i\sigma) \overline{f(t)}$$

for all $t \in \mathbb{R}$.

In what follows, we consider f to be a Laurent series on the complex plane, i.e.

$$f(z) = \sum_{n \in \mathbb{Z}} c_n z^n$$

and, if not otherwise stated, the sums will range for $n \in \mathbb{Z}$. Definition 1 can be generalized to such setting as follows.

Definition 2. Let $k < m$ be a pair of coprime positive integers. A Laurent series f defined on the complex plane \mathbb{C} has symmetry of type (k, m) if

$$(3) \quad f\left(\exp\left(2\pi i \frac{1}{m}\right) z\right) = \exp\left(2\pi i \frac{k}{m}\right) f(z)$$

for all $z \in \mathbb{C}$. Moreover, f has mirror symmetry along the axis generated by the line $\mathbb{R} \exp(i\sigma)$ if

$$(4) \quad f(\bar{z}) = \exp(2\pi i\sigma) \overline{f(z)}$$

for all $z \in \mathbb{C}$.

We consider the question of how symmetries relate to planar isometries that leave the image under f of the unit circle $S^1 = \{z \in \mathbb{C} : |z| = 1\}$ invariant.

Definition 3 ([10]). A subgroup $G \subseteq O(2)$ is a symmetry group for $f(S^1)$ if $G(f(S^1)) \subseteq f(S^1)$.

We immediately see that a symmetry group contains all the symmetries from Definition 1.

Proposition 1. *If f has symmetry of type (k, m) then the isometry group of $f(S^1)$ contains the cyclic group of order m . Moreover, if f also has a mirror symmetry along any axis, then G contains the dihedral group D_{2m} of order $2m$.*

Proof. Since $\text{GCD}(k, m) = 1$, we have that $g = \exp(2\pi i k/m)$ generates the cyclic group of order m , so the first inclusion follows from (1). If f is also mirror symmetric, then we see that G also must contain the map $\tau : z \mapsto \exp(2\pi i\sigma) \bar{z}$. The conclusion follows from the fact that g and τ generate D_{2m} . \square

Remark 1. If the Laurent polynomial f has symmetry of type (k, m) , the isometry group of $f(S^1)$ can strictly contain the cyclic group of order m . Indeed, consider the Laurent polynomial that describes a Rhodonea of four petals

$$(5) \quad f(z) = \frac{1}{2i}(z^3 - z^{-1}).$$

According to Farris's definition, f has symmetry of type $(1, 2)$ since all exponents are congruent to 1 modulo 2 but the isometry group of $f(S^1)$ is strictly bigger than the group of order two. In fact, f has also symmetry of type $(3, 4)$ for an analog reason.

It is a known fact that every subgroup of the planar isometry group $O(2)$ is either finite or dense. In the latter case, it is generated by a rotation of irrational angle $g : z \mapsto \exp(2\pi i\alpha z)$. The following lemma explores the relation to symmetries of f .

Lemma 1. *Let $f(z) = \sum_n c_n z^n$ be a Laurent series and let $\alpha \in \mathbb{R}$ and $k \in \mathbb{Z}$ be such that $\exp(2\pi i k \alpha) f(z) = f(\exp(2\pi i \alpha) z)$. Then α is rational. Moreover, a mirror symmetry $z \mapsto \exp(2\pi i\sigma) \bar{z}$ is a symmetry of $f(S^1)$ if and only if $\exp(2\pi i\sigma) \bar{c}_n = c_n$.*

Proof. For every n such that $c_n \neq 0$ we must have that the coefficient of z^n satisfies $\exp(2\pi i k \alpha) = \exp(2\pi i n \alpha)$. This means that $(k - n)\alpha \in \mathbb{Z}$ so α is rational.

If $g: z \mapsto \exp(2\pi i \sigma \bar{z})$ is a mirror symmetry that leaves f invariant, one has that

$$\sum_n c_n \bar{z}^n = f(\bar{z}) = \exp(2\pi i \sigma) \sum_n \overline{c_n z^n}$$

so that, by equating coefficients, for every n it must be either $c_n = 0$ or $c_n = \exp(2\pi i \sigma) \overline{c_n}$. \square

This allows us to obtain the following theorem.

Theorem 1. *Let f be a Laurent polynomial on the complex plane. Then the symmetry group G of $f(S^1)$ is either trivial, cyclic or dihedral.*

Proof. Let g be a rotational symmetry of f . By Lemma 1, it must be the rotation by a rational angle $\frac{p}{q} \in \mathbb{Q}$ with $\text{GCD}(p, q) = 1$. So g generates a cyclic group G of order q and hence we have that $G \subseteq \mathbb{Z}/q\mathbb{Z}$. If G also contains a mirror symmetry, this would together with g generate a dihedral group of order $2q$, so $G \subseteq D_{2q}$. \square

Remark 2. Let us step back to complex polynomials and use the notations of [10,11]. Consider $p(z) = \sum w_k z^{a_k}$, where $a_1 < a_2 < \dots < a_n$ and let $m = \text{GCD}(\{a_k - a_j : 1 \leq j < k \leq n\})$. Then p has always mirror symmetry with $\sigma = 0$ if $w_k \in \mathbb{R}$ by Lemma 1 and has symmetry of type (a_1, m) if $\text{GCD}(a_1, m) = 1$ since

$$\begin{aligned} p\left(\exp\left(2\pi i \frac{a_1}{m}\right) z\right) &= \exp\left(2\pi i \frac{a_1}{m}\right) z^{a_1} \sum_{k=1}^n w_k \exp\left(2\pi i \frac{a_k - a_1}{m}\right) z^{a_k - a_1} \\ &= \exp\left(2\pi i \frac{a_1}{m}\right) z^{a_1} \sum_{k=1}^n w_k z^{a_k - a_1} \end{aligned}$$

where the last equality holds since m divides $a_k - a_1$ for all k . We notice as well that this computation does not consider the polynomial coefficients w_k at all, meaning that they do not affect the type of symmetry (a_1, m) of p . This answers the question stated in the cited papers about the symmetry groups of an arbitrary weighted sum of exponentials.

Example 1. For all $a, b, c \in \mathbb{C}$, the polynomials $p(z) = az^2 + bz^7 + cz^{12}$ have symmetry of type $(2, 5)$ since the smallest exponent is 2 and is coprime with $\text{GCD}(5, 5, 10) = 5$. So the symmetry group of $p(S^1)$ contains D_{10} if $a, b, c \in \mathbb{R}$ and $\mathbb{Z}/5\mathbb{Z}$ otherwise. The plots can be seen in Figure 1.

Example 2. If the exponents of the polynomial p are such that a_1 is not coprime with $m = \text{GCD}(a_k - a_j)$ then the symmetry group of p need not contain the cyclic group of order m . Consider $p(z) = z^5 + z^{10} + z^{15}$ where we have that $a_1 = 5$ is not coprime with $m = 5$. We note that $p(\exp(2\pi i/5) \cdot z) = p(z)$ but $p(S^1)$ is not invariant under the rotation of angle $2\pi/5$.

2.1. Permutations on zeros and poles. Let us slightly generalize our setting and take f to be a meromorphic function defined on (a domain of) \mathbb{C} and let us study the relations between transformations of $f(S^1)$ and the zeros and the poles of f .

Proposition 2. *Let g be a symmetry of $f(S^1)$. Then g induces a permutation on the zeros of f and on the poles of f .*




FIGURE 1. Two polynomials of symmetry of type $(2, 5)$. On the left, $p(z) = z^2 + z^7 + z^{12}$ that has real coefficients. The symmetry group of $p(S^1)$ contains both a rotational symmetry of order 5 and a mirror symmetry along the x -axis. On the right, the polynomial $p(z) = 2z^2 - 2iz^7 + iz^{12}$ has no mirror symmetries hence the symmetry group of $p(S^1)$ contains no elements of order two

Proof. Let $z_0 \in \mathbb{C}$ be a zero (resp. a pole) of f and let g be a rotation of k/m . Then we have that $f(gz_0) = f(\exp(2\pi i/m) z_0) = \exp(2\pi i k/m) f(z_0) = 0$. If g is of the form $g(z) = \exp(2i\sigma) \bar{z}$ then $f(gz_0) = f(\exp(2\sigma i) \bar{z}_0) = \exp(2i\sigma) \overline{f(z_0)} = 0$. If z_0 is a pole of f , apply the same argument on $1/f$ in order to reach the same conclusion. \square

Proposition 2 allows us to construct curves that have trivial isometry group, inspired by the examples of [12].

Example 3. Consider the set $Z = \{0, \frac{1}{2}, i\} \subset \mathbb{C}$ that admits no planar nontrivial isometry carrying Z into Z . Consider the polynomial $p(z) = z(z - \frac{1}{2})(z - i)$ admitting such points as roots. Then the curve $p(S^1)$ has trivial isometry group and looks like in Figure 2.

2.2. Algebraic varieties. It has been proved by Quine [13] that the image of the circle under a complex polynomial p of degree n lies in a real algebraic variety $V \subset \mathbb{C}$ of degree $2n$. Generalizations of this to Laurent polynomials have been proved later [8]. Such variety has the rational parametrization

$$(6) \quad \begin{cases} w &= p(z) \\ \bar{w} &= \bar{p}(\frac{1}{z}). \end{cases}$$

An implicit equation can be found by elimination of z by computing the resultant $h(w, \bar{w})$ of the polynomials $A(z) = p(z) - w$ and $B(z) = z^n(\bar{p}(1/z) - \bar{w})$. If $\alpha_1, \dots, \alpha_n$ are the complex roots of A (depending on w), then $1/\alpha_1, \dots, 1/\alpha_n$ are the roots of B and by standard results (see e.g. the classical [6]) their resultant is

$$(7) \quad h(w, \bar{w}) = (-1)^{n^2} \prod_{k=1}^n A\left(\frac{1}{\alpha_k}\right)$$




FIGURE 2. Image of the unit circle under the polynomial $p(z) = z(z - \frac{1}{2})(z - i)$

where we consider conjugate coordinates (w, \bar{w}) of \mathbb{C} . If σ is a transformation of \mathbb{C} , then its action in such coordinates can be written as $\sigma \cdot (w, \bar{w}) = (\sigma w, \sigma^{-1} \bar{w})$.

If p admits a symmetry σ , we want to infer that the variety V is also invariant under σ or a power thereof.

Proposition 3. *Let p be a complex Laurent polynomial admitting a symmetry σ of type (k, m) and let $p(S^1)$ sit in the real variety $V = \{h = 0\} \subset \mathbb{C}$. Then V is σ^k -invariant. Moreover, V is invariant under mirror symmetries.*

Proof. We want to prove that the polynomial defining V is σ^k -invariant. Note that $\sigma \alpha_\ell$ is a root of $p(z) - \sigma^k w$. Let us then compute

$$\begin{aligned}
 h(\sigma^k w, \sigma^{-k} \bar{w}) &= (-1)^{n^2} \prod_{\ell=1}^n \left(p\left(\frac{1}{\sigma \alpha_\ell}\right) - \sigma^k w \right) \\
 &= (-1)^{n^2} \prod_{\ell=1}^n \left(\sigma^k p\left(\frac{1}{\alpha_\ell}\right) - \sigma^k w \right) \\
 &= (-1)^{n^2} \prod_{\ell=1}^n \sigma^k \left(p\left(\frac{1}{\alpha_\ell}\right) - w \right) \\
 &= (-1)^{n^2} \prod_{\ell=1}^n \sigma^k A\left(\frac{1}{\alpha_\ell}\right) \\
 &= \sigma^{kn} h(w, \bar{w})
 \end{aligned}$$

So if $w \in V$, then it satisfies $h(w, \bar{w}) = 0 = h(\sigma^k w, \sigma^{-k} \bar{w})$ and then $\sigma^k w \in V$.

If $\sigma: z \mapsto \exp(2\pi i \tau) \bar{z}$ is a mirror symmetry, we note that $\sigma \alpha_\ell$ is a root of $p(z) - \sigma w$ since

$$\begin{aligned}
 p(\sigma \alpha_\ell) - \sigma w &= \exp(2\pi i \tau) \overline{p(\alpha_\ell)} - \sigma w \\
 &= \exp(2\pi i \tau) \left(\overline{p(\alpha_\ell) - w} \right) \\
 &= \sigma \overline{A(\alpha_\ell)} \\
 &= 0.
 \end{aligned}$$

So $1/\sigma\alpha_\ell$ is a root of $p(1/z) - \sigma w$ and we can compute

$$\begin{aligned} h(\sigma w, \overline{\sigma w}) &= (-1)^{n^2} \prod_{\ell=1}^n \left(p(\sigma\alpha_\ell^{-1}) - \sigma w \right) \\ &= (-1)^{n^2} \prod_{\ell=1}^n \exp(2\pi i \tau) \left(\overline{p(\alpha_\ell^{-1})} - \sigma w \right) \\ &= (-1)^{n^2} \prod_{\ell=1}^n \sigma A \left(\frac{1}{\alpha_\ell} \right) \\ &= \sigma^n h(w, \overline{w}) \end{aligned}$$

that is the same as $h(w, \overline{w})$ if n is even or $\sigma h(w, \overline{w})$ if n is odd. \square

2.3. Irrational case. Let $a_1 < a_2 < \dots < a_n$ be a sequence of real numbers and do not set restrictions on $w_k \in \mathbb{C}$. Consider the curve

$$\gamma(t) = \sum_{j=1}^n w_j \exp(2\pi i a_j t)$$

to be defined on the whole real line and let us consider the symmetry group of $\gamma(\mathbb{R})$.

The necessity of considering the image of the whole real line is suggested from the following fact.

Proposition 4. *If the curve $\gamma: \mathbb{R} \rightarrow \mathbb{C}$ is periodic, then the a_k span a one-dimensional vector space over \mathbb{Q} .*

Proof. Let T be the period of γ . Then by definition it holds that

$$\gamma(t) = \sum_k w_k \exp(2\pi i a_k t) = \gamma(t + T) = \sum_k w_k \exp(2\pi i a_k (T + t))$$

from which we see that every exponential term must equal its counterpart by linear independence. Hence, there exists integers λ_k such that $a_k T = \lambda_k$. This means that every a_k is a rational multiple of $1/T$. \square

Concerning the symmetry group of $\gamma(\mathbb{R})$, we have the following.

Theorem 2. *Let the exponents a_k be real and rationally independent, i.e. linearly independent over \mathbb{Q} . Then the isometry group of $\gamma(\mathbb{R})$ is a group of order 2 consisting only of the conjugation map $z \mapsto \overline{z}$.*

Proof. The fact that the complex conjugation is always a transformation of the curve image of the whole line is due to the fact that $\overline{\gamma(t)} = \gamma(-t)$. Let σ be a planar rotation given by multiplication with the complex number we still call $\sigma = \exp(2\pi i \alpha)$. Then one needs to have

$$\sigma \gamma(0) = \exp(2\pi i \alpha) \sum_k w_k = \sum_k w_k \exp(2\pi i a_k t')$$

for some real t' . We must then have that $a_k t' - \alpha$ is an integer. Assume for a contradiction that $t' \neq 0$. Then by subtracting such relations for all k with the special one $a_1 t' - \alpha \in \mathbb{Z}$, we get that for all k

$$(a_k - a_1)t' = \lambda_k \in \mathbb{Z}$$

If $t' \neq 0$ we have that $\lambda_k \neq 0$ and can then write

$$\frac{a_i - a_j}{\lambda_{ij}} = \frac{a_k - a_\ell}{\lambda_{k\ell}}$$

that would mean there exist a rational linear combination of the a_k 's giving zero. This contradicts the assumption that the a_k are independent. Hence, $t' = 0$ which means that $\alpha \in \mathbb{Z}$ and $\sigma = 1$. \square

Theorem 3. *If the a_k and 1 are rationally independent, then the image $\gamma(\mathbb{R})$ is dense in an annulus of the complex plane.*

For this we need a lemma.

Lemma 2. *Let T^n be the n -dimensional torus and consider the map $\Pi: T^n \rightarrow A$ defined by $\Pi(x_1, \dots, x_n) = \sum_k w_k x_k$ and where A is the annulus*

$$A = \{z \in \mathbb{C} : \min_{\mathbb{R}} |\gamma(t)| \leq |z| \leq \max_{\mathbb{R}} |\gamma(t)|\}.$$

Then Π is continuous and surjective.

Proof. Let $z = \rho \exp(2\pi i \phi) \in A$. By construction, there exist a $t \in \mathbb{R}$ such that $|\gamma(t)| = \rho$ and such t is hence mapped by γ onto a point on the circle of radius ρ containing z . So we can write, for some angle σ , that $z = \exp(2\pi i \sigma) \gamma(t)$. If we take $x_k = \exp(2\pi i(a_k t + \sigma)) \in S^1$ we obtain that

$$\Pi(x_1, \dots, x_n) = \sum_k w_k \exp(2\pi i(a_k t + \sigma)) = \exp(2\pi i \sigma) \gamma(t) = z.$$

\square

Proof of Theorem 3. Consider the map $f: \mathbb{R} \rightarrow T^n$ given by

$$f(t) = (\exp(2\pi i a_1 t), \dots, \exp(2\pi i a_n t)).$$

It is well known to have the property that its image $f(\mathbb{R})$ is dense in the n -dimensional torus if the a_k are rationally independent, see e.g. [7, p. 32].

Compose f with the map Π of Lemma 2 to obtain $\Pi \circ f: \mathbb{R} \rightarrow A$. Since Π is continuous and surjective, the closure of $\Pi(f(\mathbb{R}))$ in A coincides with $\Pi(\overline{f(\mathbb{R})}) = \Pi(T^n) = A$, where $\overline{f(\mathbb{R})} = T^n$ denotes the closure in T^n of the dense subset $f(\mathbb{R}) \subset T^n$. This shows that $\gamma(\mathbb{R}) = \Pi(T^n)$ is dense in the annulus A . \square

Remark 3. The problem of estimating the thickness of such annulus boils down to the minimization of $\|\gamma\|$ over \mathbb{R} , since the maximum over \mathbb{R} of $\|\gamma\|$ is always attained at $t = 0$ and is equal to the sum of the coefficients w_k . In the binomial case, the square distance from the origin can be estimated as

$$\|\gamma(t)\|^2 = v^2 + w^2 - 2vw \cos 2\pi(b-a)t \geq v^2 + w^2 - 2vw = (v-w)^2 \geq 0,$$

so when the two weights are different the annulus is not a disk. Note that this is independent of the exponents $a, b \in \mathbb{R}$. The authors do not know any estimates for the case of more than two terms.

3. EVOLUTION UNDER THE WAVE EQUATION

Following the ideas of Farris [5] one can consider the initial curve

$$\gamma = \gamma(0): x \mapsto p(\exp(2\pi ix))$$

evolving under the wave equation, that is to consider a solution of the one-dimensional wave problem

$$(8) \quad \begin{aligned} u_{tt} &= c^2 u_{xx} \\ u(x, 0) &= \gamma(x) \\ u_t(x, 0) &= f(x) \end{aligned}$$

for $u: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ and some initial condition $f: [0, 1] \rightarrow \mathbb{R}$.

It is known that all solutions are given by the D'Alembert formula

$$(9) \quad u(x, t) = \frac{1}{2} \left(\gamma(x - ct) + \gamma(x + ct) + \frac{1}{c} \int_{x-ct}^{x+ct} f(s) ds \right).$$

One could write a solution explicitly in terms of sums and, for the simpler case $f = 0$ one ends up with

$$(10) \quad u(x, t) = \sum_n c_n \cos(2\pi nct) \exp(2\pi inx).$$

We can prove some properties of the function u and the family of curves it defines.

Theorem 4. *Let $\gamma(t) = u(\cdot, t)$ be the family of curves defined by u .*

- (i) *The family of curves $\gamma(t)$ satisfies the wave problem (8) with initial condition $f = 0$.*
- (ii) *The period with respect to t of $\gamma(t)$ is $T = \frac{1}{c \text{GCD}\{n: c_n \neq 0\}}$, that is $\gamma(t) = \gamma(t+T)$ as curves $[0, 1] \rightarrow \mathbb{C}$.*
- (iii) *The symmetry group of the curve $\gamma(t)$ contains the symmetry group of $\gamma = \gamma(0)$.*

Proof. (i) Solving the one-dimensional wave problem by series yields the expression (10) solving (8) with initial condition $u_t(x, 0) = 0$.

To prove (ii), consider the family of linear independent functions $x \mapsto \exp(2\pi inx)$ on the unit circle. By imposing the period condition $\gamma(t+T) = \gamma(t)$ and considering its components along such family, we obtain the system

$$c_n (\cos(2\pi cnt) - \cos(2\pi cn(t+T))) = 0$$

holding for all n and all $t \in \mathbb{R}$. This yields to either $c_n = 0$ or that cnT is an integer. So the smallest T that works for all n such that $c_n \neq 0$ is $T = \frac{1}{c \text{GCD}\{n: c_n \neq 0\}}$.

We have remarked after Proposition 1 that the symmetry group of γ does not depend on the coefficients c_n or $c_n \cos(2\pi nt)$ and, if c_n is real, then also $c_n \cos(2\pi nct)$. Hence, every element of such a group also defines a symmetry of $\gamma(t)$ and this proves (iii). \square

Remark 4. We note that the symmetry group of $\gamma(t)$ can “degenerate” for special values of t that make coefficients of $u(x, t)$ vanish, i.e. zeros of $\cos(2\pi cnt)$. We shall see some examples in our closer look at the sum of two exponentials.

3.1. Evolution of the sum of two exponentials. We consider the special case of the sum of two exponentials as it lets us carry out computations in details. Consider the complex Laurent polynomial

$$(11) \quad p(z) = c_a z^a + c_b z^b$$

for integers $a < b$. Then the one-dimensional solution of the wave problem (8) with $f = 0$ is

$$(12) \quad u(x, t) = c_a \cos(2\pi act) \exp(2\pi i ax) + c_b \cos(2\pi bct) \exp(2\pi ibx).$$

3.1.1. Differential geometric properties, winding number. Applying Theorem 4 we can consider $\gamma(t)$ for $t \in [0, T]$, where $T = 1/(c \text{GCD}(a, b))$. Moreover, every transformation in the symmetry group D_{b-a} is also a transformation of $\gamma(t)$. However, for t such that $\cos(2\pi act) = 0$ or $\cos(2\pi bct) = 0$, the curve $\gamma(t)$ “degenerates” to a circle, making the symmetry group “explode” to the non-discrete S^1 .

Analogous to [11, Theorem 1], it is possible to prove a theorem about the winding number of $\gamma(t)$. We start with two lemmas from calculus. First, recall that it is possible to express $\cos(dx)$ for integer d in terms of $y = \cos(x)$ through the d -th Chebyshev polynomial T_d , i.e. for all $x \in [-\pi, \pi]$ we have that $T_d(\cos(x)) = \cos(dx)$. Such polynomials satisfy the recurrence relation $T_0(y) = 1, T_1(y) = y$ and $T_{d+1}(x) = 2xT_d(x) - T_{d-1}(x)$. The Chebyshev polynomial of second kind $U_k(y)$ expresses $\sin(2\pi\theta) \cdot \sin(2\pi k\theta)$ in terms of $y = \cos(2\pi\theta)$. The range of U_n is $[-(n+1), n+1]$. For an introduction to the topic we refer to [1, Chap. 22].

Lemma 3. *For odd d , the Chebyshev polynomials satisfy*

$$(13) \quad |T_d(y)| \leq d|y|$$

for all $y \in [-1, 1]$

Proof. Note that $T_{2n+1}(0) = 0$ and that the statement can be rewritten as $|T_{2n+1}(y) - T_{2n+1}(0)| \leq |T'_{2n+1}(0)||y - 0|$ since T_{2n+1} is a Lipschitz continuous function. \square

For the polynomials of second kind we have the following.

Lemma 4. *Let $I := [0, 1] \setminus \{t : \sin 2\pi bt = 0\}$ and define $\psi: I \rightarrow \mathbb{R}$ be the function defined as $\psi(\theta) = \frac{\sin(2\pi a\theta)}{\sin(2\pi b\theta)}$. If $a|b$, extend ψ to the points θ_k of the form \mathbb{Z}/a where $\sin(2\pi a\theta_k) = \sin(2\pi b\theta_k) = 0$ to assume value a/b . Then such extension is continuous and*

$$(14) \quad |\psi(I)| = \begin{cases} [0, +\infty) & \text{if } a \nmid b \\ [\frac{a}{b}, +\infty) & \text{if } a|b. \end{cases}$$

In the latter case, the minimum $\frac{a}{b}$ is attained exactly at the points θ_k .

Proof. If $a \nmid b$ then there exists $\theta_0 \in [0, 1]$ such that $\sin(2\pi b\theta_0) = 0$ and $\sin(2\pi a\theta_0) \neq 0$, so we have that the range of $|\psi|$ is $[0, +\infty)$. If $b = da$ then we can write for $s = \cos(2\pi\theta)$ and $y = \cos(2\pi a\theta)$

$$\psi(\theta) = \frac{U_{a-1}(s)}{U_{da-1}(s)} = \frac{U_{a-1}(s)}{U_{d-1}(T_a(y))U_{a-1}(s)} = \frac{1}{U_{d-1}(y)}$$

Since it is known that the range of U_{d-1} is $[-d, d]$ we conclude: For θ_k as in the statement in the case $b = da$ let us compute

$$\lim_{\theta \rightarrow \theta_k} \frac{\sin(2\pi a\theta)}{\sin(2\pi b\theta)} = \lim_{\theta \rightarrow \theta_k} \frac{a \cos(2\pi a\theta)}{b \cos(2\pi b\theta)} = \frac{a}{b} \frac{1}{(-1)^d},$$

so we obtain the second statement. \square

An analog holds for the function defined by the ratio of cosines.

Lemma 5. *Let $I_T := [0, T] \setminus \{t : \cos 2\pi act = 0\}$. For fixed $c > 0$, consider the function $\phi: I_T \rightarrow \mathbb{R}$ defined by $\phi(t) = \frac{\cos(2\pi bct)}{\cos(2\pi act)}$. In case $a|b$ and $2a \nmid b$, extend it to the points of the form $t_k = \frac{1}{4ac} + \frac{k}{2ac}$ where $\cos(2\pi act) = \cos(2\pi bct) = 0$ to assume value b/a . Then the extension is continuous and*

$$(15) \quad |\phi(I_T)| = \begin{cases} [0, +\infty) & \text{if } a \nmid b \text{ or } 2a|b \\ [0, \frac{b}{a}] & \text{if } a|b \text{ and } 2a \nmid b. \end{cases}$$

In the latter case, the maximum $\frac{b}{a}$ is attained exactly at the points t_k .

Proof. If $a \nmid b$, then there exists $t_0 \in [0, T]$ such that $\cos(2\pi act_0) = 0$ and $\cos(2\pi bct_0) \neq 0$. Then

$$(16) \quad \lim_{t \rightarrow t_0} |\phi(t)| = +\infty$$

and since $[0, T]$ contains also points that make $\cos(2\pi bct)$ vanish, we get that the range of $|\phi|$ is $[0, \infty)$.

If $a|b$ we can write $b = da$ and express $\phi(t)$ as the ratio of the Chebyshev polynomials $T_d(y)/y$ for $y = \cos(2\pi act)$. We have that $T_d(0) = 0$ if d is odd. So if $2a|b$ we can pick d even and get that $\lim_{y \rightarrow 0} |T_d(y)/y| = +\infty$ to the range of $|\phi|$ is $[0, +\infty)$. If $2a \nmid b$, then d is odd, and we apply Lemma 3 to obtain that $|\phi| \leq d = \frac{b}{a}$.

For the last assertion, compute the limit

$$\lim_{t \rightarrow t_k} \phi(t) = \lim_{t \rightarrow t_k} \frac{b \sin(2\pi bct)}{a \sin(2\pi act)} = \frac{b}{a}. \quad \square$$

Remark 5. Note that although its period may be different from T , for the study of the evolution we are only interested in the behavior of ϕ for $t \leq T$.

We are now ready to prove our winding number theorem. Recall that by the classical argument principle, if 0 is not in its support, the winding number of the curve $f(S^1)$ with respect to 0 is the number of zeros of f inside the unit disk minus the number of poles of f , (see e.g. [2]).

Theorem 5. *Let $\gamma(t) = u(\cdot, t)$ be the family of curves defined above for $t \in [0, T] \setminus \{t : \cos 2\pi act = 0\}$ and consider the function $\phi(t) = \frac{\cos(2\pi bct)}{\cos(2\pi act)}$. Let $N(t, 0)$ be the winding number of $\gamma(t)$ with respect to zero. Then*

$$(17) \quad N(t, 0) = \begin{cases} a & \text{if } \cos(2\pi bct) = 0 \text{ or } |\phi(t)| > \frac{c_a}{c_b} \text{ and } a > 0 \\ -a & \text{if } \cos(2\pi bct) = 0 \text{ or } |\phi(t)| > \frac{c_a}{c_b} \text{ and } a < 0 \\ \text{undefined} & \text{if } |\phi(t)| = \frac{c_a}{c_b} \\ b & \text{if } \cos(2\pi act) = 0 \text{ or } |\phi(t)| < \frac{c_a}{c_b} \text{ and } a > 0 \\ b - 2a & \text{if } \cos(2\pi act) = 0 \text{ or } |\phi(t)| < \frac{c_a}{c_b} \text{ and } a < 0. \end{cases}$$

Proof. Write the family of curves as

$$u(x, t) = c_a \cos(2\pi act) z^a \left(1 + \frac{c_b}{c_a} \phi(t) z^{b-a} \right) \Big|_{z=\exp(2\pi ix)}.$$

We want to count the zeros and the poles of a complex polynomial of the form $z^a(1 + Kz^{b-a})$ as the real number $K = \frac{c_b}{c_a}\phi(t)$ varies. The possibilities can be summarized as follows:

- if $a > 0$ and $|K| < 1$ then the unit disk contains only the root 0 with multiplicity a ;
- if $a < 0$ and $|K| < 1$ then the unit disk contains only the pole 0 with multiplicity a ;
- if $a > 0$ and $|K| > 1$ then the unit disk contains the root 0 with multiplicity a and the root $-\frac{1}{K}$ with multiplicity $b - a$;
- if $a < 0$ and $|K| > 1$ then the unit disk contains the pole 0 with multiplicity a and the root $-\frac{1}{K}$ with multiplicity $b - a$.

This translates as

$$\begin{aligned} & \# \text{ of zeros} - \# \text{ of poles inside the unit disk} \\ &= \begin{cases} a & \text{if } |\phi(t)| > \frac{c_a}{c_b} \text{ and } a > 0 \\ -a & \text{if } |\phi(t)| > \frac{c_a}{c_b} \text{ and } a < 0 \\ b & \text{if } |\phi(t)| < \frac{c_a}{c_b} \text{ and } a > 0 \\ b - 2a & \text{if } |\phi(t)| < \frac{c_a}{c_b} \text{ and } a < 0 \end{cases} \end{aligned}$$

and that for $|\phi(t)| = \frac{c_a}{c_b}$ the curve crosses the origin, so the winding number is undefined. \square

3.1.2. *Singularities.* The singularities in the differential geometric sense of the curve $\gamma(t)$ are attained at points $x \in [0, 1]$ such that $u_x(x, t) = 0$. In general, we have

$$u_x(x, t) = 2\pi i \sum_n n \cos(2\pi nct) \exp(2\pi inx)$$

In our particular case it becomes

$$(18) \quad u_x(x, t) = 2\pi i(a \cos(2\pi act) \exp(2\pi iax) + b \cos(\pi bct) \exp(2\pi ibx))$$

$$(19) \quad = 2\pi ia \cos(2\pi act) \cdot z^a(1 + \frac{b}{a} \frac{c_b}{c_a} \phi(t) z^{b-a})|_{z=\exp(2\pi iax)}$$

The study of the existence of zeroes on the unit circle again boils down to the study of the function ϕ from Lemma 5, so we obtain the following facts.

Proposition 5. *The curve $\gamma(t)$ admits singular points for t_0 such that*

$$|\phi(t_0)| = \frac{|c_a|}{|c_b|} \frac{a}{b}$$

and x^k with $x = \frac{1}{2\pi(b-a)}$ for $k = 0, \dots, b - a - 1$, i.e., at the $b - a$ -th roots of unity. Such points are cusps.

Proof. The form of the solutions (x^k, t_0) follows directly from (18). Differentiate again and evaluate at such points to obtain

$$u_{xx}(x^k, t_0) = -4\pi^2 a \cos(2\pi act) x^{ka} \cdot 0 + 4\pi^2 a \cos(2\pi act) x^{ka} (b - a) \neq 0$$

\square

Example 4. Assume that γ has at least two points $x_k \in [0, 1]$ such that $\gamma'(x_k) = 0$. Consider two such points $x_1 < x_2$ and the intersection $(x_0 = \frac{x_1 + x_2}{2}, t_0)$ of the lines $x + ct = x_1$ and $x - ct = x_2$. Then at such point we have

$$u_x(x_0, t_0) = \gamma'(x_0 + ct_0) + \gamma'(x_0 - ct_0) = \gamma'(x_1) + \gamma'(x_2) = 0,$$

so we have constructed a singular point at a future time t_0 by “following the characteristic lines”.

Example 5. As in Example 4 for $n = 2$, we note that the presence of only one singularity at $t = 0$ and $x = \frac{1}{2}$ is not enough to infer the existence of more singularities for $u(\cdot, t)$ for future $t > 0$. Indeed, we have $u(x, t) = \cos(4\pi ct) \exp(2\pi ix) + \cos(2\pi ct) \exp(4\pi ix)$ and so

$$u_x(x, t) = 2\pi i \exp(2\pi ix) (\cos(4\pi ct) + 2 \cos(2\pi ct) \exp(2\pi ix))$$

that admits zeroes (x, t) only for $t = 0$ for which we recover the known $x = \frac{1}{2}$.

4. SELF-INTERSECTIONS

Quine [14] gives a characterization of the pairs $(z_1, z_2) \in S^1 \times S^1$ for which $p(z_1) = p(z_2)$ and estimates from above the number of such self-intersections of the image $p(S^1)$ by $2(n - 1)^2$ where n is the degree of p .

Furthermore, he shows that this estimate is sharp by proving that the polynomial $z^n + \varepsilon z$ verifies equality for small enough ε . Quine considers the open subsets of $S^1 \times S^1$ given by

$$(20) \quad A = (-1, 1) \times S^1$$

$$(21) \quad B = \{(z_1, z_2) \in S^1 \times S^1 : z_1 \neq z_2\}$$

and the diffeomorphism $\psi: A \rightarrow B$ given by $\psi(s, x) = (e^{i\theta}x, e^{-i\theta}x)$, where $\cos(2\pi\theta) = s$.

Quine’s method to find points in B that are mapped to the same value uses the properties of the *Dieudonné polynomial* that can be defined on B or on A as

$$\begin{aligned} G(z_1, z_2) &= \frac{p(z_1) - p(z_2)}{z_1 - z_2} \\ &= \frac{p(e^{2\pi i\theta}x) - p(e^{-2\pi i\theta}x)}{x(e^{2\pi i\theta} - e^{-2\pi i\theta})} \\ &= \sum_{k=0}^n c_{k+1} U_k(s) x^k \\ &=: g(s, x) \end{aligned}$$

where U_k is the k -th Chebishev polynomial of second kind, that expresses $\sin 2\pi(k + 1)\theta / \sin(2\pi\theta)$ in terms of $s = \cos(2\pi\theta)$.

A zero $(s, x) \in A$ for G is also a zero of $G(s, 1/\bar{x})$, hence of the polynomial $G^*(s, x) = x^n G(s, 1/\bar{x})$ and so of the resultant $R(t)$.

We state a lemma concerning a property of such intersection points. Analogously to g , let us define the polynomial h as

$$(22) \quad h(s, x) = \sum_{k=0}^n c_k T_k(s) x^k$$

Lemma 6. *For all $(s, x) \in A$ and its corresponding θ with $s = \cos(2\pi\theta)$ we have*

$$(23) \quad p(e^{2\pi i\theta}x) = h(s, x) + i \sin(\theta) \cdot xg(s, x).$$

In particular, if $(s, x) \in A$ is a zero of g , then the point $p(e^{i\theta}x) \in \mathbb{C}$ lies along the direction of $h(s, x)$.

Proof. Using the definition of Chebyshev polynomials we can expand

$$e^{2\pi ik\theta} = T_k(s) + i \sin(\theta) U_{k-1}(s)$$

for $k \geq 0$ with the convention $U_{-1}(s) = 0$. We then compute

$$(24) \quad p(e^{2\pi i\theta} x) = \sum_{k=0}^n c_k (T_k(s) + i \sin(\theta) U_{k-1}(s)) x^k$$

$$(25) \quad = h(s, x) + i \sin(\theta) \cdot xg(s, x).$$

To prove the last statement, multiply by $\overline{h(s, x)}$ and note that

$$p(e^{2\pi i\theta} x) \overline{h(s, x)} = |h(s, x)|^2 \in \mathbb{R}$$

has zero imaginary part, meaning that the directions of $p(e^{i\theta} x)$ and $h(s, x)$ are parallel in the plane. \square

Let us now consider the simple case $p(z) = vz^a + wz^b$ and apply Quine's method to find the self-intersections. Then we have the following result, interesting on its own.

Theorem 6. *Let γ be the image under p above of the circle. Then*

- (i) *The self-intersections of γ lie on the directions of $\pm\eta^k$ with $\eta = \exp\left(\frac{2\pi ia}{b-a}\right)$ for $k = 0, \dots, b-a-1$;*
- (ii) *On every such direction, the self-intersections are twice the number of solutions of the polynomial equation $U_{b-1}(s) = \mp v/(wU_{a-1}(s))$ and have modulus $|vT_a(s) \mp wT_b(s)|$.*

Proof. For our p we have

$$(26) \quad g(s, x) = vU_{a-1}(s)x^{a-1} + wU_{b-1}(s)x^{b-1}.$$

and

$$(27) \quad h(s, x) = vT_a(s)x^a + wT_b(s)x^b$$

If $U_{a-1}(s) = U_{b-1}(s) = 0$, then s solves $\sin(2\pi a\theta) = \sin(2\pi b\theta) = 0$ and hence

$$\theta \in \frac{\mathbb{Z}}{b} \cap \frac{\mathbb{Z}}{a} = \frac{\mathbb{Z}}{\text{GCD}(a, b)}$$

hence $2\theta = k/2g$ for some integer $0 \leq k < 2g$ and $x \in S^1$. This case leads to no solutions, since for algebraic reasons the number of solutions of $g = 0$ must be finite. If $U_{a-1}(s) \neq 0$, then we can rewrite our equation as

$$(28) \quad vx^{a-1}U_{a-1}(s) \left(1 + \frac{w}{v} \frac{U_{b-1}(s)}{U_{a-1}(s)} x^{b-a} \right) = 0$$

and its solutions are in the set

$$(29) \quad S_- \times R_+ \cup S_+ \times R_-$$

where

$$S_{\pm} = \left\{ s \in (-1, 1) : \frac{U_{b-1}(s)}{U_{a-1}(s)} = \pm \frac{v}{w} \right\}$$

and

$$R_{\pm} = \{x \in S^1 : x^{b-a} = \pm 1\}.$$

For such pairs, we have that $h(s, x) = x^a(vT_a(s) \mp wT_b(s))$ and hence the direction of $p(e^{i\theta}x)$ is parallel to the one of \bar{x}^a . The modulus of $p(e^{i\theta}x)$ coincides in this cases with

$$|h(s, x)| = |vT_a(s) \mp wT_b(s)|$$

proving the last statement. \square

Example 6. Consider $p(z) = 2z^3 + z^5$ and compute

$$g(s, x) = 2U_2(s)x^2 + U_4(s)x^4$$

that has as roots (s, x) where $x^2 = \pm 1$ and θ satisfies

$$\pm 2 \sin(6\pi\theta) = \sin(10\pi\theta).$$

Hence, the self-intersection points are to be expected on real and imaginary axes and their possible modulus is

$$|h(s, \pm 1)| = |h(s, \pm i)| = |2T_3(s) + T_5(s)|$$

Note that for $x = \pm i$ the possible s are solutions of

$$(4s^2 - 1)^2 / (16s^4 - 12s^2 - 1)^2 = 4.$$

We note that the solutions of g of the form $(1, x)$ correspond to the points where the parametrization is not regular. Indeed, at such points one has

$$\gamma'(t) = p'(e^{2\pi i\theta}) = 0$$

and p' coincides with $g(1, \cdot)$ on S^1 .

Example 7. Let us construct an example providing points of arbitrary multiplicity. Let n be a positive integer and consider $p(z) = 1 + z + \dots + z^n = \frac{z^{n+1} - z}{z - 1}$ and the points $0, -1 \in \mathbb{C}$ that have preimages

$$p^{-1}(0) \cap S^1 = \left\{ \exp\left(2\pi i \frac{k}{n}\right) : k = 1, \dots, n \right\}$$

and

$$p^{-1}(1) \cap S^1 = \left\{ \exp\left(2\pi i \frac{k}{n+1}\right) : k = 1, \dots, n+1 \right\}$$

of size n and $n+1$. This means that the curve crosses n times the origin and $n+1$ times the point $1 \in \mathbb{C}$. Such points are ordinary since at all such points the derivative p' assumes different values. The rosette is shown in Figure 3.

Example 8. The image of $p(z) = z^n + nz$ has a cusp. Differentiate to get $p'(z) = n(z^{n-1} + 1)$, that has the $(n-1)$ -th roots of -1 as roots, lying on the unit circle, and note that the derivative has all its $n-1$ zeros on the unit circle. Such points are cusps since they are not zeros of $p''(z) = n(n-1)z^{n-2}$. We note this is the only way to obtain singularities of the parametrization $t \mapsto p(\exp(2\pi it))$.




FIGURE 3. The rosette for $p(z) = 1 + z + z^2 + z^3 + z^4 + z^5$, that crosses 5 times the origin and 6 times the point $(1, 0)$

4.1. Self-intersections along the wave flow. Let us now apply Theorem 6 for $v = v(t) = c_a \cos(2\pi cat)$ and $w = w(t) = c_b \cos(2\pi cbt)$ to study the time evolution of the rosette associated to the polynomial (11) under the wave flow. We obtain then the following result.

Theorem 7. *Let $\gamma(t) = u(\cdot, t)$ be the wave evolution of a polynomial (11). Then*

- (i) *The self-intersections of $\gamma(t)$, if any, lie along the directions of the powers of $\exp\left(\frac{2\pi ia}{b-a}\right)$ for all $t > 0$.*
- (ii) *On every such line, the number of self-intersections is the number of roots of*

$$(30) \quad U_{a-1}(s) = \mp \frac{c_b}{c_a} \phi(t) U_{b-1}(s)$$

as rational function in $s \in [-1, 1]$, and have length

$$|c_a \cos(2\pi cat) T_a(s) \mp c_b \cos(2\pi cbt) T_b(s)|.$$

Proof. Apply Theorem 6 for $v = c_a \cos(2\pi cat)$ and $w = c_b \cos(2\pi cbt)$. □

Example 9. Consider the evolution $\gamma(t)$ of the curve defined by the polynomial $p(z) = vz^a + wz^b$. If $a \nmid b$, we see from Lemmas 4 and 5 that the range of ψ and the range of ϕ are both \mathbb{R} , so we can say that for all $t \in [0, T]$ there is $s \in [-1, 1]$ satisfying (30). In order to construct a family of curves that has no self-intersections at all times t one can pick $a|b$ and weights c_a, c_b such that $\frac{c_b}{c_a} < \frac{a^2}{b^2}$. In this case, the ranges

$$\frac{c_b}{c_a} \phi([0, T]) = \left[-\frac{bc_b}{ac_a}, \frac{bc_b}{ac_a} \right]$$

and

$$\psi([0, 1]) = \left(-\infty, \frac{a}{b} \right] \cup \left[\frac{a}{b}, +\infty \right)$$

will never intersect. A numerical example of such situation is displayed in Figure 4.




FIGURE 4. On the left, the evolution of $p(z) = 26z^2 + z^{10}$ where self-intersections are absent for all times. On the right, the plots of ψ and ϕ that never intersect




FIGURE 5. Evolution of $p(z) = 6z + z^6$ with $c = 1/6$ and period $T = 6$. For $t = 0$, the rosette has five cusps and self-intersections occur, and disappear, at later times $t \leq 1$.

Example 10. There can be cases where there are no self-intersections at $t = 0$, but for positive t . Take $p(z) = nz + z^n$ with n odd, that, as seen in Example 8 has $n - 1$ cusps, so no self-intersections. We notice that for $t_k = \frac{1}{4ct} + \frac{k}{2ct}$ the function ϕ reaches ± 1 . On the other hand, the ranges of $\frac{1}{n}\phi$ and ψ intersect exactly in

$$\frac{1}{n}\phi([0, T]) \cap \psi([0, 1]) = [-1, 1] \cap ((-\infty, -1] \cup [1, +\infty)) = \{-1, 1\}$$

where we have used that n is odd and Lemmas 5 and 4. A plot is shown in Figure 5

Example 11. Let us consider the evolution of the polynomial $p(z) = z^2 + 2z^5$ that is given by

$$u(x, t) = \cos(4\pi ct)e^{2\pi i 2x} + 2 \cos(10\pi ct)e^{2\pi i 5t}.$$




FIGURE 6. Plot of $|\phi| = |\cos(\pi t)|/|\cos(.4\pi t)|$ for $0 < t < 1$. The function vanishes at $t = .5$ and attains the value $c_b/c_a = .4$ at $t \approx .35$ and $t \approx .62$

If we pick $c = 1/10$, we obtain a period $T = 1$, and we see that the solutions of $|\phi| = c_b/c_a$ lie at $t \approx .35$ and $t \approx .62$ as we can see in Figure 6. Moreover, another remarkable time is $t = .5$ where $\cos 4\pi ct = 0$ making the curve degenerate do a circle traced twice. The shape of $u(\cdot, t)$ for several values of $t < 1$ are shown in Figure 7. According to Theorem 5, the winding number with respect to zero is then 2 for $t \in (.35, .62)$ and 5 outside this interval. Note also that, given the similarity of the equation for singularities $u_x(x, t) = 0$ with the one for self-intersection $g(x, s) = 0$, we also note that their solutions (x, t) and (x, s) must have their x -component along the same directions.

REFERENCES

1. M. Abramowitz and I.A. Stegun, *Handbook of mathematical functions: With formulas, graphs, and mathematical tables*, Applied mathematics series, Dover Publications, 1965.
2. L.V. Ahlfors and Karreman Mathematics Research Collection, *Complex analysis: An introduction to the theory of analytic functions of one complex variable*, International series in pure and applied mathematics, McGraw-Hill Education, 1979.
3. Hilário Alencar, Walcy Santos, and Gregório Silva Neto, *Differential geometry of plane curves*, Stud. Math. Libr., vol. 96, Providence, RI: American Mathematical Society (AMS), 2022.
4. Frank A Farris, *Wheels on wheels on wheels-surprising symmetry*, Mathematics Magazine **69** (1996), no. 3, 185–189.
5. ———, *Creating symmetry: The artful mathematics of wallpaper patterns*, Princeton University Press, 2015.
6. I.M. Gelfand, M. Kapranov, and A. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Modern Birkhäuser Classics, Birkhäuser Boston, 2009.
7. A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 1995.
8. Leonid V Kovalev and Xuerui Yang, *Algebraic structure of the range of a trigonometric polynomial*, Bulletin of the Australian Mathematical Society **102** (2020), no. 2, 251–260.
9. Peter M. Maurer, *A rose is a rose...*, The American Mathematical Monthly **94** (1987), no. 7, 631–702.
10. Florian Pausinger and Dimitris Vartziotis, *On the symmetry of finite sums of exponentials*, Elemente der Mathematik **76** (2021), no. 2, 62–73.




FIGURE 7. Time evolution of $z^2 + 2z^5$ for $0 < t < 1$. The curve degenerates to a circle traced twice for $t = 0.5$ and passes through zero (blue dot) at $t \approx .35$ and $t \approx .62$. The self-intersections as well as the cusps lie along the lines given by powers of $e^{\frac{4}{3}\pi i}$ at all times (dashed lines) and, for some value $t \in (.60, .62)$, the former are not simple (that is, they have the same tangent)

11. Florian Pausinger and Dimtris Vartziotis, *On weighted sums of two exponentials*, Mathematics Magazine (to appear).
12. Konstantin Poelke, Zoi Tokoutsis, and Konrad Polthier, *Complex polynomial mandalas and their symmetries*, Proceedings of Bridges 2014: Mathematics, Music, Art, Architecture, Culture, 2014, pp. 433–436.
13. John Quine, *The geometry of $p(s^1)$* , Pacific Journal of Mathematics **64** (1976), no. 2, 551–557.
14. JR Quine, *On the self-intersections of the image of the unit circle under a polynomial mapping*, Proceedings of the American Mathematical Society **39** (1973), no. 1, 135–140.

QUEEN'S UNIVERSITY BELFAST, BELFAST, UNITED KINGDOM.
 Email address: f.pausinger at qub.ac.uk

STUTTGART, GERMANY
 Email address: david.petrecca at gmail.com