

ULTRADIFFERENTIABLE FUNCTIONS VIA THE LAGUERRE OPERATOR

S. JAKŠIĆ, S. PILIPOVIĆ, N. TEOFANOV, AND ĐORĐE VUČKOVIĆ

ABSTRACT. We introduce spaces of test functions $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 0$, by using the iterates of the Laguerre operator, and show that these spaces represent a natural counterpart of global Pilipović spaces on \mathbb{R}^d . Moreover, we show that $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ coincide with nontrivial G -type spaces. We also consider the associated spaces of ultradistributions. The main tool in our study is the use of sequence spaces $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$ arising from the corresponding Laguerre expansions. In addition, by considering sequence spaces $\ell_{b,\sigma}(\mathbb{N}_0^d)$ and $\ell_{0,b,\sigma}(\mathbb{N}_0^d)$, we introduce and study flat Pilipović spaces on \mathbb{R}_+^d .

0. INTRODUCTION

Recently, Toft, Bhimani and Manna demonstrated that (global) Pilipović spaces on \mathbb{R}^d , $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, $\alpha > 0$, and their dual spaces of distributions, appear as a convenient framework in the study of the well-posedness of certain Schrödinger equations, which are ill-posed in the class of Gelfand-Shilov type distributions, see [24]. In this paper we address the following question:

Q: What are natural counterparts of global Pilipović spaces when considering the positive orthants \mathbb{R}_+^d instead of \mathbb{R}^d ?

Recall that $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, $\alpha > 0$, are defined in terms of the Hermite operator. We introduce Pilipović spaces (P -spaces) on the positive orthants, $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 0$, of Roumieu and Beurling type by utilizing the iterates of the Laguerre operator. We also consider their dual spaces of ultradistributions.

We show that elements of $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 0$, can be expressed through Laguerre expansions, and establish a topological isomorphism between these spaces and sequence spaces $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$ considered in [23]. As a consequence of Laguerre expansions, we deduce that $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ coincide with G -type spaces $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$, respectively, when these spaces are nontrivial. We refer to [5, 10, 11] for G -type spaces.

Furthermore, to answer the question Q, we establish a natural isomorphism between Pilipović spaces on \mathbb{R}^d consisting of even functions, and P -spaces on \mathbb{R}_+^d , extending main results from [12] where an isomorphism between subspaces of Gelfand-Shilov spaces and G -type spaces is shown. Finally, we conclude the paper by exploring the properties of

2010 *Mathematics Subject Classification.* 46F12, 46F05, 33C45, 42B10, 44A15, 47G10, 47G30, 35S05.

Key words and phrases. Spaces of ultradistributions of Beurling type over \mathbb{R}_+^d , Laguerre series expansions.

P -flat spaces on \mathbb{R}_+^d , which are associated with Pilipović flat spaces on \mathbb{R}^d considered in e.g. [23].

We may briefly summarize the context of our research as follows.

Spaces of smooth functions on \mathbb{R}_+^d , which might serve as counterparts for the well-known spaces of Schwartz functions $\mathcal{S}(\mathbb{R}^d)$, or Gelfand-Shilov functions $\Sigma_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 1/2$, and $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, $\alpha \geq 1/2$ (of Beurling and Roumieu type, respectively) have been studied for decades. In the 1990s, Duran studied Schwartz functions and tempered distributions on \mathbb{R}_+ , as well as Gelfand-Shilov type spaces and their distribution spaces on \mathbb{R}_+ in [1] and [5], respectively. Comparing to the standard toolbox used to study the spaces in the global (i.e. \mathbb{R}^d) setting, different techniques are employed when related spaces on orthants are considered. This includes, for example, Laguerre (eigen)expansions, and the Hankel-Clifford transform. Duran's results were extended in [10, 11, 12], where the test spaces of ultradifferentiable functions of Beurling and Roumieu type, denoted by $g_\alpha^\alpha(\mathbb{R}_+^d)$ and $G_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha \geq 1$, respectively, are introduced and characterized via decay properties of their Laguerre expansions coefficients. The spaces $g_\alpha^\alpha(\mathbb{R}_+^d)$ and $G_\alpha^\alpha(\mathbb{R}_+^d)$ are non-trivial when $\alpha > 1$ and $\alpha \geq 1$ respectively, and then we have dense and continuous inclusions:

$$g_\alpha^\alpha(\mathbb{R}_+^d) \hookrightarrow G_\alpha^\alpha(\mathbb{R}_+^d), \quad \alpha > 1.$$

In the global setting, the (isotropic) spaces of Gelfand - Shilov type $\Sigma_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 1/2$, and $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, $\alpha \geq 1/2$, are characterized by the decay of their coefficients in Hermite (eigen)expansions, cf. [15, 16, 19]. More generally, these spaces are characterized by the decay properties of the coefficients in expansions involving eigenfunctions of certain elliptic and positive Shubin-type operators $P(x, D)$, see [8, 26].

Pilipović spaces $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, $\alpha > 0$, were initially introduced in [19] by considering the iterates of linear Harmonic oscillator. These spaces are nontrivial for any $\alpha > 0$, and for $\alpha > \frac{1}{2}$ they coincide with Gelfand-Shilov spaces $\Sigma_\alpha^\alpha(\mathbb{R}^d)$. Similar relations hold for the Roumieu case (with $\alpha > \frac{1}{2}$) as well, cf. [23]. In this paper, we take a similar approach (see also [18, 29]), and introduce test function spaces on \mathbb{R}_+^d by using a suitable self-adjoint operator. Due to their close relation to Pilipović spaces on \mathbb{R}^d we call them Pilipović spaces on positive orthants, or P -spaces for short.

The paper is organized as follows. In Section 1 we set the stage for our results by collecting main notions which will be used in the sequel. P -spaces on \mathbb{R}_+^d , together with their representation in terms of Laguerre expansions are introduced in Section 2. Theorems 2.3 and 2.5 extend existing results [4, Theorem 3.6], [10, Theorem 5.7], and [11, Theorem 3.7]. In Section 3 we show that L^2 -norm in the definition of P -spaces can be replaced by any other L^p -norm, $p \in [1, \infty]$, Proposition 3.2. We establish in Section 4 an isomorphism between subspaces of global Pilipović spaces and P -spaces on \mathbb{R}_+^d , Theorems 4.5 and 4.6. These results extend Propositions 3.1 and 3.2 from [12]. Finally, in Section 5 we introduce and study flat Pilipović spaces on orthants.

We end this section with review of some basic notation.

0.1. **Notation.** We denote by \mathbb{N} , \mathbb{Z} , \mathbb{R} and \mathbb{C} the sets of positive integers, integers, real and complex numbers, respectively. Furthermore, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbb{R}_+ = (0, \infty)$, $\overline{\mathbb{R}_+} = [0, \infty)$. The upper index $d \in \mathbb{N}$ will denote the dimension, e.g. $\mathbb{R}_+^d = (0, \infty)^d$ and so on.

We use the standard multi-index notation: for $\alpha \in \mathbb{N}_0^d$ (or $\alpha \in \overline{\mathbb{R}_+^d}$) and $x \in \mathbb{R}^d$ (or $x \in \overline{\mathbb{R}_+^d}$), $x^\alpha = x_1^{\alpha_1} \dots x_d^{\alpha_d}$ and $\partial^\alpha = \partial^{\alpha_1} / \partial x_1^{\alpha_1} \dots \partial^{\alpha_d} / \partial x_d^{\alpha_d}$. We will use the convention $0^0 = 1$. The length of $\alpha \in \mathbb{N}_0^d$ is given by $|\alpha| = \sum_{i=1}^d \alpha_i$. For $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbb{R}^d$ such that $-\gamma_j \notin \mathbb{N}$, $j = 1, \dots, d$, and $m \in \mathbb{N}_0^d$, we use the abbreviation

$$\binom{\gamma}{m} = \prod_{j=1}^d \binom{\gamma_j}{m_j}.$$

We also use the standard notation for common spaces of functions, distributions, and sequences. For example, ℓ^p , $p \geq 1$, denotes the Banach space of Lebesgue sequences, endowed with the usual norm:

$$\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell^p \quad \Longleftrightarrow \quad \|a_n\|_{\ell^p} = \left(\sum_n |a_n|^p \right)^{1/p} < \infty,$$

and $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell^\infty$ if and only if $\|a_n\|_{\ell^\infty} = \sup_{n \in \mathbb{N}_0^d} |a_n| < \infty$.

A measurable function f belongs to the Hilbert space $L^2(\mathbb{R}_+^d)$ if

$$\|f\|_{L^2(\mathbb{R}_+^d)} := \left(\int_{\mathbb{R}_+^d} |f(x)|^2 dx \right)^{1/2} < \infty,$$

and the scalar product is given by $\langle f, g \rangle = \int_{\mathbb{R}_+^d} f(x) \overline{g(x)} dx$, $f, g \in L^2(\mathbb{R}_+^d)$.

By $\mathcal{S}(\mathbb{R}_+^d)$ we denote the space of all smooth functions on positive orthant, $f \in \mathcal{C}^\infty(\mathbb{R}_+^d)$ such that all of its derivatives $\partial^\alpha f$, $\alpha \in \mathbb{N}_0^d$, extend to continuous functions on $\overline{\mathbb{R}_+^d}$, and

$$\sup_{x \in \mathbb{R}_+^d} x^\beta |\partial^\alpha f(x)| < \infty, \quad \alpha, \beta \in \mathbb{N}_0^d.$$

If s denotes the space of rapidly decreasing sequences:

$$\{a_n\}_{n \in \mathbb{N}_0^d} \in s \quad \Longleftrightarrow \quad \sum_{n \in \mathbb{N}_0^d} |a_n| |n|^N < \infty, \quad \forall N \in \mathbb{N}, \quad (1)$$

then the mapping $\iota : \mathcal{S}(\mathbb{R}_+^d) \rightarrow s$, given by $\iota(f) = \{a_n(f)\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism, see [9, Theorem 3.1]). Here $\{a_n(f)\}_{n \in \mathbb{N}_0^d}$ are the Fourier-Laguerre coefficients of f , see (2) below.

Let I be a directed set and let $\{X_\alpha\}_{\alpha \in I}$ be a family of normed spaces. By $\bigcup_{\alpha \in I} X_\alpha$ we denote the union of $\{X_\alpha\}_{\alpha \in I}$, endowed with inductive limit topology, and by $\bigcap_{\alpha \in I} X_\alpha$ we denote the intersection of $\{X_\alpha\}_{\alpha \in I}$, endowed with projective limit topology.

If \mathcal{A} is a space of test functions, then \mathcal{A}' denotes its dual space of (ultra)distributions, and $\mathcal{A}'(\cdot, \cdot)_{\mathcal{A}} = (\cdot, \cdot)$ denotes the dual pairing.

1. PRELIMINARIES

In this section we collect some background notions and properties which will be used in the sequel.

1.1. Laguerre functions and Laguerre operator. For $n \in \mathbb{N}_0$ and $\gamma \geq 0$, the n -th Laguerre polynomial of order γ is defined by

$$L_n^\gamma(x) = \frac{x^{-\gamma} e^x}{n!} \frac{d^n}{dt^n} (e^{-x} x^{\gamma+n}), \quad x \geq 0,$$

and the n -th Laguerre function is defined by $l_n^\gamma(x) = L_n^\gamma(x) e^{-x/2}$, $x \geq 0$. The d -dimensional Laguerre polynomials and Laguerre functions are given by tensor products:

$$L_n^\gamma(x) = \prod_{k=1}^d L_{n_k}^\gamma(x_k) \quad \text{and} \quad l_n^\gamma(x) = \prod_{k=1}^d l_{n_k}^\gamma(x_k), \quad x \in \overline{\mathbb{R}_+^d}, \quad n \in \mathbb{N}_0^d.$$

We will be particularly interested in the case $\gamma = 0$, and write L_n and l_n instead of L_n^0 and l_n^0 , respectively. The Laguerre functions $\{l_n\}_{n \in \mathbb{N}_0^d}$ form an orthonormal basis in $L^2(\mathbb{R}_+^d)$, cf. e.g. [22, 27]. Thus, every $f \in L^2(\mathbb{R}_+^d)$ can be expressed via its Laguerre series expansion; by $a_n(f)$ we denote the corresponding Laguerre coefficients:

$$a_n(f) = \int_{\mathbb{R}_+^d} f(x) l_n(x) dx, \quad n \in \mathbb{N}_0^d. \quad (2)$$

Next, we introduce the Laguerre operator:

$$E = - \sum_{j=1}^d \left(\frac{\partial}{\partial x_j} (x_j \frac{\partial}{\partial x_j}) - \frac{x_j}{4} + \frac{1}{2} \right) = - \sum_{j=1}^d \left(x_j \frac{\partial^2}{\partial x_j^2} + \frac{\partial}{\partial x_j} - \frac{x_j}{4} + \frac{1}{2} \right).$$

Then the Laguerre functions are the eigenfunctions of E :

$$E(l_n(x)) = \left(\sum_{i=1}^d n_i \right) \cdot l_n(x) = |n| \cdot l_n(x), \quad x \in \overline{\mathbb{R}_+^d}, \quad n \in \mathbb{N}_0^d.$$

see [6, (11), p.188], wherefrom

$$E^N(l_n(x)) = |n|^N l_n(x), \quad x \in \overline{\mathbb{R}_+^d}, \quad N \in \mathbb{N}. \quad (3)$$

1.2. Hermite functions and Hermite operator. The Hermite polynomials H_n and the corresponding Hermite functions h_n are defined by

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2}), \quad h_n(x) = (2^n n! \sqrt{\pi})^{-1/2} e^{-x^2/2} H_n(x), \quad x \in \mathbb{R}, \quad n \in \mathbb{N}_0,$$

respectively. The d -dimensional Hermite polynomials H_n and Hermite functions h_n are then given by

$$H_n(x) = H_{n_1}(x_1) \dots H_{n_d}(x_d), \quad h_n(x) = h_{n_1}(x_1) \dots h_{n_d}(x_d), \quad x \in \mathbb{R}^d, \quad n \in \mathbb{N}_0^d,$$

respectively. Next, we introduce the Hermite operator:

$$H = |x|^2 - \Delta = \sum_{j=1}^d \left(x_j^2 - \frac{\partial^2}{\partial x_j^2} \right) \quad x \in \mathbb{R}^d, \quad n \in \mathbb{N}_0^d.$$

Then the Hermite functions are eigenfunctions of H , $H(h_n(x)) = (2|n| + d)h_n(x)$, $x \in \mathbb{R}^d$, $n \in \mathbb{N}_0^d$, see [23, pp. 8], and

$$H^N(h_n(x)) = (2|n| + d)^N h_n(x) \quad x \in \mathbb{R}^d, \quad N \in \mathbb{N}. \quad (4)$$

Hermite and Laguerre polynomials can be related as follows:

$$H_{2n}(x) = (-1)^n 2^{2n} n! L_n^{-1/2}(x^2), \quad H_{2n+1}(x) = (-1)^n 2^{2n} n! L_n^{1/2}(x^2) x, \quad x \in \mathbb{R}^d, n \in \mathbb{N}_0^d,$$

see (1.1.52) and (1.1.53) in [22].

1.3. The sequence spaces $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$. We will employ the notation from [23].

Definition 1.1. Let $\alpha > 0$, $h > 0$, and put $\vartheta_{h,\alpha}(n) := e^{h|n|^{1/(2\alpha)}}$, $n \in \mathbb{N}_0^d$. By $\ell_{[\vartheta_{h,\alpha}]}^p(\mathbb{N}_0^d)$, $p \geq 1$, we denote the space of all sequences $\{a_n\}_{n \in \mathbb{N}_0^d}$, $a_n \in \mathbb{C}$, $n \in \mathbb{N}_0^d$, which satisfy

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h,\alpha}]}^p} := \|\{|a_n| \cdot \vartheta_{h,\alpha}(n)\}_{n \in \mathbb{N}_0^d}\|_{l^p} < \infty. \quad (5)$$

The space $\ell_{[\vartheta_{h,\alpha}]}^p(\mathbb{N}_0^d)$ is a Banach space, with the norm $\|\cdot\|_{\ell_{[\vartheta_{h,\alpha}]}^p}$ given in (5), and with the usual modification when $p = \infty$. Next, we introduce locally convex spaces

$$\ell_\alpha(\mathbb{N}_0^d) = \bigcup_{h>0} \ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d) \quad \text{and} \quad \ell_{0,\alpha}(\mathbb{N}_0^d) = \bigcap_{h>0} \ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d),$$

in the sense of inductive and projective limit topology of $\ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d)$ with respect to $h > 0$, respectively. In addition, we consider their dual (locally convex) spaces

$$\ell'_\alpha(\mathbb{N}_0^d) = \bigcap_{h>0} \ell_{[1/\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d) \quad \text{and} \quad \ell'_{0,\alpha}(\mathbb{N}_0^d) = \bigcup_{h>0} \ell_{[1/\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d),$$

in the sense of projective and inductive limit topology of $\ell_{[1/\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d)$ with respect to $h > 0$, respectively. Note that a different notation is used in [10]. The reader may immediately see that for $\alpha \geq 1$, the sequence space $s^{\alpha,a}(\mathbb{N}_0^d)$ defined in [10] is the same as $\ell_{[\vartheta_{\ln a, \alpha/2}]}^\infty(\mathbb{N}_0^d)$, $a > 1$, and also $s^\alpha(\mathbb{N}_0^d) = \ell_{\alpha/2}^\infty(\mathbb{N}_0^d)$. When $\alpha = 0$, $\ell_0(\mathbb{N}_0^d)$ denotes the space of all sequences $\{a_n\}_{n \in \mathbb{N}_0^d}$ such that $a_n \neq 0$ for only finitely many $n \in \mathbb{N}_0^d$, and $\ell_{0,0}(\mathbb{N}_0^d) = \{0\}$.

Remark 1.2. Observe that different choices of $p \in [1, \infty]$ give rise to equivalent norms in (5). We are particularly interested in the case $p \in \{2, \infty\}$, i.e. $\ell_{[\vartheta]}^\infty(\mathbb{N}_0^d) = \ell_{[\vartheta]}^2(\mathbb{N}_0^d)$. Thus, if $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d)$ then for any given $h > 0$ there exist $h_1, h_2 > 0$ and $C_1, C_2 > 0$ such that

$$C_1 \|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d)} \leq \left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 \vartheta_{h,\alpha}(n) \right)^{1/2} \leq C_2 \|\{|a_n| \cdot \vartheta_{h_2,\alpha}(n)\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h,\alpha}]}^\infty}.$$

Regarding the topological structure of $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$, we have the following.

Proposition 1.3. *Let there be given $\alpha > 0$, and $h_1 > h_2 > 0$. Then the canonical inclusion*

$$\ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_2,\alpha}]}^\infty(\mathbb{N}_0^d) \quad (6)$$

is nuclear.

Proof. The proof is similar to the proof of [10, Proposition 2.1], with slight modifications, and we present it here for the sake of completeness. We will use the fact that a mapping is nuclear if it can be represented as a composition of two quasinuclear mappings, cf. [21]. Since the inclusion (6) can be considered as a composition of inclusions $\ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_3,\alpha}]}^\infty(\mathbb{N}_0^d)$ and $\ell_{[\vartheta_{h_3,\alpha}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_2,\alpha}]}^\infty(\mathbb{N}_0^d)$, for some/any $h_3 \in (h_2, h_1)$, it is enough to prove the quasinuclearity of the inclusion $I_{h_1}^{h_3} : \ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_3,\alpha}]}^\infty(\mathbb{N}_0^d)$.

Let there be given $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d)$ and let $e_m \in (\ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d))'$ be given by

$$e_m(\{a_n\}_{n \in \mathbb{N}_0^d}) = a_m e^{h_3|m|^{1/(2\alpha)}}, \quad m \in \mathbb{N}_0^d.$$

Then clearly $\|e_m\|_{(\ell_{[\vartheta_{h_1,\alpha}]}^\infty)'} \leq C e^{(h_3-h_1)|m|^{1/(2\alpha)}}$ for some $C > 0$, and therefore

$$\sum_{m \in \mathbb{N}_0^d} \|e_m\|_{(\ell_{[\vartheta_{h_1,\alpha}]}^\infty)'} < \infty,$$

since $h_3 < h_1$. Now we have

$$\begin{aligned} \|I_{h_1}^{h_3}(\{a_n\}_{n \in \mathbb{N}_0^d})\|_{\ell_{[\vartheta_{h_3,\alpha}]}^\infty} &= \|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_3,\alpha}]}^\infty} \\ &= \sup_{m \in \mathbb{N}_0^d} |a_m| e^{h_3|m|^{1/(2\alpha)}} = \sup_{m \in \mathbb{N}_0^d} |e_m(\{a_n\}_{n \in \mathbb{N}_0^d})| \leq \sum_{m \in \mathbb{N}_0^d} |e_m(\{a_n\}_{n \in \mathbb{N}_0^d})|, \end{aligned}$$

for every $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d)$. Then by [21, Definition 3.2.3] it follows that $I_{h_1}^{h_3}$ is quasinuclear. The same is true for $I_{h_3}^{h_2} : \ell_{[\vartheta_{h_3,\alpha}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_2,\alpha}]}^\infty(\mathbb{N}_0^d)$, so that (6) is nuclear. \square

By Proposition 1.3 and [25, Theorem 50.1, p.511] it follows that $\ell_\alpha(\mathbb{N}_0^d)$ is a (DFN) -space (dual of a Fréchet nuclear space), and that $\ell_{0,\alpha}(\mathbb{N}_0^d)$ is an (FN) -space (Fréchet nuclear space).

Thus $\ell'_\alpha(\mathbb{N}_0^d)$ is a (FN) -space and $\ell'_{0,\alpha}(\mathbb{N}_0^d)$ is an (DFN) -space (cf. [10, Proposition 2.2] and [11, Proposition 1.1]).

Remark 1.4. *When $\alpha = 0$, we have the following : $\ell_{0,0}(\mathbb{N}_0^d) = \{0\}$, and $\ell_0(\mathbb{N}_0^d) = \bigcup_{n \in \mathbb{N}} E_n$, where $E_n = \{\{a_k\}_{k \in \mathbb{N}_0}, k = 0 \text{ when } k > n\}$. Since the canonical inclusion $id : E_n \rightarrow E_{n+1}$ is nuclear, it follows that $\ell_0(\mathbb{N}_0^d)$ is a (DFN) -space.*

1.4. Test spaces of formal Laguerre and Hermite series. In what follows, we introduce spaces of formal Laguerre and Hermite series expansions.

We follow [23] and consider the spaces of formal Laguerre series expansions $f = \sum_{n \in \mathbb{N}_0^d} a_n l_n$ that correspond to the sequence $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$, or $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha/2}(\mathbb{N}_0^d)$, $\alpha \geq 0$.

These spaces will be denoted by $\mathcal{G}_\alpha(\mathbb{R}_+^d)$ and $\mathcal{G}_{0,\alpha}(\mathbb{R}_+^d)$, respectively, and the natural mappings

$$T : \ell_{\alpha/2}(\mathbb{N}_0^d) \rightarrow \mathcal{G}_\alpha(\mathbb{R}_+^d), \quad T(\{a_n\}_{n \in \mathbb{N}_0^d}) = \sum_{n \in \mathbb{N}_0^d} a_n l_n, \quad (7)$$

$$T : \ell_{0,\alpha/2}(\mathbb{N}_0^d) \rightarrow \mathcal{G}_{0,\alpha}(\mathbb{R}_+^d), \quad T(\{a_n\}_{n \in \mathbb{N}_0^d}) = \sum_{n \in \mathbb{N}_0^d} a_n l_n, \quad (8)$$

induce topologies on the target spaces.

The spaces of formal Hermite series expansions $f = \sum_{n \in \mathbb{N}_0^d} a_n h_n$ that correspond to the sequence $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_\alpha(\mathbb{N}_0^d)$, or $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha}(\mathbb{N}_0^d)$, $\alpha \geq 0$, are denoted by $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, respectively, and the natural mappings

$$T_h : \ell_\alpha(\mathbb{N}_0^d) \rightarrow \mathcal{H}_\alpha(\mathbb{R}^d), \quad T_h(\{a_n\}_{n \in \mathbb{N}_0^d}) = \sum_{n \in \mathbb{N}_0^d} a_n h_n, \quad (9)$$

$$T_h : \ell_{0,\alpha}(\mathbb{N}_0^d) \rightarrow \mathcal{H}_{0,\alpha}(\mathbb{R}^d), \quad T_h(\{a_n\}_{n \in \mathbb{N}_0^d}) = \sum_{n \in \mathbb{N}_0^d} a_n h_n, \quad (10)$$

induce topologies on the target spaces.

1.5. Pilipović and Gelfand-Shilov spaces on \mathbb{R}^d . Pilipović spaces are defined by using the iterates of Hermite operators as follows.

We say that $f \in C^\infty(\mathbb{R}^d)$ belongs to the (global) Pilipović spaces of Roumieu type $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ (of Beurling type $\Sigma_\alpha^\alpha(\mathbb{R}^d)$), $\alpha > 0$, if there exist $h > 0$ and $C > 0$ (for every $h > 0$ there exists $C = C_h > 0$) such that the following estimate holds:

$$\|H^N f\|_{L^\infty(\mathbb{R}^d)} \leq Ch^N N!^{2\alpha}.$$

From [23, Definition 3.1] and [23, Theorem 5.2] it follows that the spaces of formal Hermite expansions $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, coincide with Pilipović spaces $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ and $\Sigma_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 0$, respectively.

For that reason, from now on we will use the notation $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, instead of $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ and $\Sigma_\alpha^\alpha(\mathbb{R}^d)$, respectively.

Next we recall the definition of (isotropic) Gelfand-Shilov spaces.

Let there be given $\alpha, A > 0$. We denote by $\mathcal{S}_{\alpha,A}^{\alpha,A}(\mathbb{R}^d)$ the Banach space of all $\varphi \in C^\infty(\mathbb{R}^d)$ such that

$$\|\varphi\|_{\mathcal{S}_{\alpha,A}^{\alpha,A}} := \sup_{p,k \in \mathbb{N}_0^d} \frac{\|x^k \partial^p \varphi(x)\|_{L^2(\mathbb{R}^d)}}{A^{|p+k|} k!^\alpha p!^\alpha} < \infty,$$

and $\|\cdot\|_{\mathcal{S}_{\alpha,A}^{\alpha,A}}$ is the norm in $\mathcal{S}_{\alpha,A}^{\alpha,A}(\mathbb{R}^d)$.

The Gelfand-Shilov space (of Roumieu type) $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 0$ (cf. [7]) is defined by

$$\mathcal{S}_\alpha^\alpha(\mathbb{R}^d) = \bigcup_{A>0} \mathcal{S}_{\alpha,A}^{\alpha,A}(\mathbb{R}^d),$$

and it is endowed with inductive limit topology. $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ is nontrivial if and only if $\alpha \geq 1/2$, cf. [7]. Indeed, $h_n \in \mathcal{S}_{1/2}^{1/2}(\mathbb{R}^d)$, $n \in \mathbb{N}_0^d$, and Gelfand-Shilov spaces obviously increase with respect to α .

The Gelfand-Shilov distribution space of Roumieu type $(\mathcal{S}_\alpha^\alpha)'(\mathbb{R}^d)$ is the dual space of $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 0$, i.e. it is given by

$$(\mathcal{S}_\alpha^\alpha)'(\mathbb{R}^d) = \bigcap_{A>0} (\mathcal{S}_{\alpha,A}^{\alpha,A})'(\mathbb{R}^d),$$

and endowed with projective limit topology.

The spaces $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ and $(\mathcal{S}_\alpha^\alpha)'(\mathbb{R}^d)$ can be characterized through Hermite expansions as follows.

Proposition 1.5. *Let $\alpha \geq 1/2$. The mapping $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d) \rightarrow \ell_\alpha(\mathbb{N}_0^d)$, $f \mapsto \{\langle f, h_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle f, h_n \rangle h_n$ converges absolutely to f in $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$.*

The mapping $(\mathcal{S}_\alpha^\alpha)'(\mathbb{R}^d) \rightarrow \ell'_\alpha(\mathbb{N}_0^d)$, $u \mapsto \{\langle u, h_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle u, h_n \rangle h_n$ converges absolutely to u in $(\mathcal{S}_\alpha^\alpha)'(\mathbb{R}^d)$.

Proposition 1.5 is a special case of Theorem 3.4 and Corollary 3.5 in [15].

Together with $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ one can consider Gelfand-Shilov spaces of Beurling type

$$\Sigma_\alpha^\alpha(\mathbb{R}^d) = \bigcap_{A>0} \mathcal{S}_{\alpha,A}^{\alpha,A}(\mathbb{R}^d),$$

endowed with projective limit topology, and their dual spaces of Gelfand-Shilov distributions of Beurling type $(\Sigma_\alpha^\alpha)'(\mathbb{R}^d)$.

The space $\Sigma_\alpha^\alpha(\mathbb{R}^d)$ is nontrivial if and only if $\alpha > 1/2$, and

$$\Sigma_\alpha^\alpha(\mathbb{R}^d) \subset \mathcal{S}_\alpha^\alpha(\mathbb{R}^d) \subset \Sigma_\beta^\beta(\mathbb{R}^d) \subset \mathcal{S}_\beta^\beta(\mathbb{R}^d), \quad \frac{1}{2} < \alpha < \beta.$$

An analogue of Proposition 1.5 can be formulated for $\Sigma_\alpha^\alpha(\mathbb{R}^d)$ and $(\Sigma_\alpha^\alpha)'(\mathbb{R}^d)$, $\alpha > 1/2$. We leave details for the reader. In what follows, we use [23, Theorem 5.2], see also [19].

Proposition 1.6. *Let $\alpha > 1/2$. Then,*

$$\mathcal{H}_\alpha(\mathbb{R}^d) = \mathcal{S}_\alpha^\alpha(\mathbb{R}^d) \quad \text{and} \quad \mathcal{H}_{0,\alpha}(\mathbb{R}^d) = \Sigma_\alpha^\alpha(\mathbb{R}^d),$$

In addition, $\mathcal{H}_{1/2}(\mathbb{R}^d) = \mathcal{S}_{1/2}^{1/2}(\mathbb{R}^d)$ and $\mathcal{H}_{0,1/2}(\mathbb{R}^d) \neq \Sigma_{1/2}^{1/2}(\mathbb{R}^d) = \{0\}$.

Therefore, from now on we will use the notation $\mathcal{H}_\alpha(\mathbb{R}^d)$ and $\mathcal{H}_{0,\alpha}(\mathbb{R}^d)$, instead of $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$ and $\Sigma_\alpha^\alpha(\mathbb{R}^d)$, respectively.

Since Pilipović spaces are non-trivial even when $\alpha \in (0, 1/2)$, they can be considered as a natural extension of Gelfand-Shilov spaces when considering ultradifferentiable functions of ultra-rapid decay.

1.6. **G -type spaces on \mathbb{R}^d .** We end this section with the definition of G -type spaces and their distribution spaces, as it is given in [10, 11, 12]. In addition, we recall the Laguerre expansions in these spaces.

Let $A > 0$. By $g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d)$, we denote the space of all $f \in \mathcal{S}(\mathbb{R}_+^d)$ such that

$$\sup_{p,k \in \mathbb{N}_0^d} \frac{\|x^{(p+k)/2} \partial^p f(x)\|_{L^2(\mathbb{R}_+^d)}}{A^{|p+k|} k^{(\alpha/2)k} p^{(\alpha/2)p}} < \infty. \quad (11)$$

With the seminorms

$$\sigma_{A,j}^{\alpha,\alpha}(f) = \sup_{p,k \in \mathbb{N}_0^d} \frac{\|x^{(p+k)/2} \partial^p f(x)\|_{L^2(\mathbb{R}_+^d)}}{A^{|p+k|} k^{(\alpha/2)k} p^{(\alpha/2)p}} + \sup_{\substack{|p| \leq j \\ |k| \leq j}} \sup_{x \in \mathbb{R}_+^d} |x^k \partial^p f(x)|, \quad j \in \mathbb{N}_0, \quad (12)$$

it becomes a Fréchet space ((F) -space).

The spaces $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$ are then defined as union, respectively intersection of $g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d)$ with respect to A :

$$G_\alpha^\alpha(\mathbb{R}_+^d) = \bigcup_{A>0} g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d), \quad g_\alpha^\alpha(\mathbb{R}_+^d) = \bigcap_{A>0} g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d).$$

Then $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$ are endowed with inductive, respectively projective limit topology. We refer to $G_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha \geq 1$, and $g_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 1$ as G -type spaces of Roumieu and Beurling type, respectively.

The G -type distribution spaces $(G_\alpha^\alpha)'(\mathbb{R}_+^d)$ and $(g_\alpha^\alpha)'(\mathbb{R}_+^d)$ are then given by

$$(G_\alpha^\alpha)'(\mathbb{R}_+^d) = \bigcap_{A>0} (g_{\alpha,A}^{\alpha,A})'(\mathbb{R}_+^d), \quad \alpha \geq 1, \quad (g_\alpha^\alpha)'(\mathbb{R}_+^d) = \bigcup_{A>0} (g_{\alpha,A}^{\alpha,A})'(\mathbb{R}_+^d), \quad \alpha > 1,$$

and they are endowed with projective, respectively, inductive limit topology.

Next, we summarize results from [4, Theorem 3.6], [10, Theorem 5.7] and [11, Theorem 3.7] as follows:

Theorem 1.7. *Let $f \in \mathcal{S}(\mathbb{R}_+^d)$. Then the following is true.*

- (i) *Let $\alpha \geq 1$. The mapping $G_\alpha^\alpha(\mathbb{R}_+^d) \rightarrow \ell_{\alpha/2}(\mathbb{N}_0^d)$, $f \mapsto \{\langle f, l_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle f, l_n \rangle l_n$ converges absolutely to f in $G_\alpha^\alpha(\mathbb{R}_+^d)$.*
- (ii) *Let $\alpha > 1$. The mapping $g_\alpha^\alpha(\mathbb{R}_+^d) \rightarrow \ell_{0,\alpha/2}(\mathbb{N}_0^d)$, $f \mapsto \{\langle f, l_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle f, l_n \rangle l_n$ converges absolutely to f in $g_\alpha^\alpha(\mathbb{R}_+^d)$.*
- (iii) *Let $\alpha \geq 1$. The mapping $(G_\alpha^\alpha)'(\mathbb{R}_+^d) \rightarrow \ell'_{\alpha/2}(\mathbb{N}_0^d)$, $u \mapsto \{\langle u, l_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle u, l_n \rangle l_n$ converges absolutely to u in $(G_\alpha^\alpha)'(\mathbb{R}_+^d)$.*
- (iv) *Let $\alpha > 1$. The mapping $(g_\alpha^\alpha)'(\mathbb{R}_+^d) \rightarrow \ell'_{0,\alpha/2}(\mathbb{N}_0^d)$, $u \mapsto \{\langle u, l_n \rangle\}_{n \in \mathbb{N}_0^d}$, is a topological isomorphism. Moreover, $\sum_{n \in \mathbb{N}_0^d} \langle u, l_n \rangle l_n$ converges absolutely to u in $(g_\alpha^\alpha)'(\mathbb{R}_+^d)$.*

By Theorem 1.7 it follows that $\mathcal{G}_{0,\alpha}(\mathbb{R}_+^d) = g_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha > 1$ and $\mathcal{G}_\alpha(\mathbb{R}_+^d) = G_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha \geq 1$.

We refer to [5] for the triviality of $G_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha < 1$, and $g_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha \leq 1$.

2. PILIPOVIĆ SPACES ON \mathbb{R}_+^d

We start with a simple observation.

Lemma 2.1. *Let $f \in C^\infty(\mathbb{R}_+^d)$ such that $E^N f \in L^2(\mathbb{R}_+^d)$ for every $N \in \mathbb{N}_0$. Then $f \in \mathcal{S}(\mathbb{R}_+^d)$.*

Proof. Let us consider the Laguerre expansion of f , $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n$. By (3) it follows that $E^N f = \sum_{n \in \mathbb{N}_0^d} |n|^N a_n(f) l_n$, and since $E^N f \in L^2(\mathbb{R}_+^d)$ we get $\sum_{n \in \mathbb{N}_0^d} (a_n(f))^2 |n|^{2N} < \infty$ for every $N \in \mathbb{N}$, i.e. $a_n(f) \in s$ (cf. (1)) so that for every $N \in \mathbb{N}_0$ there exists $C_N > 0$ such that $|a_n| \leq \frac{C_N}{(1+|n|)^N}$.

Let $k, p \in \mathbb{N}_0^d$. We will use the estimate (3.1) from [9]:

$$\sup_{t \in \mathbb{R}_+^d} |t^k \partial^p l_n(t)| \leq C_{p,k} (1+|n|)^{|p+k|}. \quad (13)$$

Thus for every $N \in \mathbb{N}$ we have

$$|x^k \partial^p f(x)| \leq \sum_{n \in \mathbb{N}_0^d} |a_n(f)| \cdot |x^k \partial^p l_n(x)| \leq C_{p,k} C_N \sum_{n \in \mathbb{N}_0^d} \frac{(1+|n|)^{|p+k|}}{(1+|n|)^N}, x \in \mathbb{R}_+^d.$$

Since N can be chosen arbitrary large, it follows that $\sup_{x \in \mathbb{R}_+^d} |x^k \partial^p f(x)| < \infty$, so that $f \in \mathcal{S}(\mathbb{R}_+^d)$. \square

We proceed with the definition of P -spaces on positive orthants.

Definition 2.2. *Let $h > 0$, and $\alpha > 0$. Then a smooth functions f belongs to the space $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$, if*

$$\eta_h^\alpha(f) := \sup_{N \in \mathbb{N}} \frac{\|E^N f\|_{L^2(\mathbb{R}_+^d)}}{h^N N!^\alpha} < \infty. \quad (14)$$

The Pilipović spaces (P -spaces) over \mathbb{R}_+^d , $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ of Roumieu and Beurling type respectively, are defined by

$$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = \bigcup_{h>0} \mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d), \quad \text{and} \quad \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = \bigcap_{h>0} \mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d).$$

The spaces $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ are endowed with inductive, respectively projective limit topology.

By Lemma 2.1 it follows that $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d) \subset \mathcal{S}(\mathbb{R}_+^d)$, and

$$\eta_h^\alpha(f) < \infty \iff \eta_h^\alpha(f) + \sup_{\substack{|p| \leq j \\ |k| \leq j}} \sup_{x \in \mathbb{R}_+^d} |x^k \partial^p f(x)| < \infty, \quad j \in \mathbb{N}_0.$$

Clearly, if $0 < \alpha < \beta$, $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$ is continuously injected into $\mathbf{G}_{\beta,h}^{\beta,h}(\mathbb{R}_+^d)$.

The Banach space $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$, $\alpha > 0$, $h > 0$, is non-trivial. Indeed, Laguerre functions l_n , $n \in \mathbb{N}_0^d$, satisfy (14) for every $h > 0$ since

$$\sup_{N \in \mathbb{N}} \frac{\|E^N l_n\|_{L^2(\mathbb{R}_+^d)}}{h^N N!^\alpha} = \sup_{N \in \mathbb{N}} \frac{\left(\frac{|n|}{h}\right)^N}{N!^\alpha} < \infty,$$

so that $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$ contains finite linear combinations of Laguerre functions.

$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ are continuously injected into $\mathcal{S}(\mathbb{R}_+^d)$, and the following embeddings are obvious:

$$\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) \subset \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) \subset \mathbf{g}_\beta^\beta(\mathbb{R}_+^d) \subset \mathbf{G}_\beta^\beta(\mathbb{R}_+^d), \quad 0 < \alpha < \beta. \quad (15)$$

The corresponding dual spaces are denoted by $(\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^d)$ and $(\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^d)$.

The main result of this section is to characterize P -spaces and their duals on positive orthants in the spirit of Theorem 1.7.

We will frequently use the following estimates (cf. [7]): there exist positive constants C_1, C_2, A , and B such that

$$C_1 e^{As^{1/\alpha}} \leq \sup_{n \in \mathbb{N}_0} \frac{s^n}{n!^\alpha} \leq C_2 e^{Bs^{1/\alpha}}, \quad s \geq 0. \quad (16)$$

Theorem 2.3. *Let $\alpha > 0$.*

- i) *If $f \in \mathcal{S}(\mathbb{R}_+^d)$ and $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}$, then $f \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$.*
- ii) *If $f \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$, then $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}$.*
- iii) *If $a_n(f) \in \ell'_{\alpha/2}(\mathbb{N}_0^d)$ then the sequence*

$$\sum_{n \in \mathbb{N}_0^d} a_n(f) l_n$$

converges in the topology of $(\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^d)$, thus defining an element $f \in (\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^d)$.

- iv) *Let $f \in (\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^d)$. Then for the sequence of its Fourier-Laguerre coefficients we have $a_n(f) \in \ell'_{\alpha/2}(\mathbb{N}_0^d)$.*

Proof. We prove i) and ii), and iii) and iv) then follow by the usual duality arguments.

i) Let $f \in \mathcal{S}(\mathbb{R}_+^d)$ and $h > 0$ but such that $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h,\alpha/2}]}$. By the norm equivalence (cf. Remark 1.2) there exists $h' > 0$ such that $\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h',\alpha/2}]}} < \infty$.

We need to prove that

$$\eta_{h_1}^\alpha(f) = \sup_{N \in \mathbb{N}_0} \frac{\|E^N f\|_{L^2(\mathbb{R}_+^d)}}{h_1^N N!^\alpha} < \infty,$$

for some $h_1 > 0$. Choose $h_1 > 0$ such that $h' = B/h_1^{1/\alpha}$, where B is the constant in (16). Then, by (3) we have

$$\frac{\|E^N f\|_{L^2(\mathbb{R}_+^d)}}{h_1^N N!^\alpha} = \frac{\left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 \cdot |n|^{2N}\right)^{1/2}}{h_1^N N!^\alpha} = \left(\sum_{n \in \mathbb{N}_0^d} \frac{|a_n|^2 \cdot |n|^{2N}}{h_1^{2N} N!^{2\alpha}}\right)^{1/2} \leq C_2 \|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h', \alpha/2}]}^2},$$
(17)

for any $N \in \mathbb{N}_0$. Therefore,

$$\eta_{h_1}^\alpha(f) = \sup_{N \in \mathbb{N}} \frac{\left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 |n|^{2N}\right)^{1/2}}{h_1^N N!^\alpha} \leq C_2 \|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h', \alpha/2}]}^2} < \infty,$$

i.e. $f \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$.

ii) Now suppose that $f \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$, i.e. $\eta_h(f) < \infty$ for some $h > 0$. We need to show that there exists $h_1 > 0$ such that $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h_1, \alpha/2}]}^\infty$. By using (16) we get

$$e^{h_2|n|^{1/\alpha}} = e^{A \cdot \left(\frac{h_2}{A}\right)^\alpha |n|} \leq \frac{1}{C_1} \sup_{N \in \mathbb{N}} \frac{|n|^N}{h^N N!^{1/\alpha}},$$

where $h_2 := A/h^{1/\alpha}$. Now we have

$$\sup_{n \in \mathbb{N}_0^d} |a_n| e^{h_2|n|^{1/\alpha}} \leq \frac{1}{C_1} \sup_{n \in \mathbb{N}_0^d} \left(|a_n| \cdot \left(\sup_{N \in \mathbb{N}} \frac{|n|^N}{h^N N!^{1/\alpha}}\right)\right) \leq \frac{1}{C_1} \sup_{N \in \mathbb{N}} \frac{\left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 |n|^{2N}\right)^{1/2}}{N!^\alpha h^N} \leq C \frac{1}{C_1} \eta_h^\alpha(f),$$

wherefrom we get

$$\left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 e^{2h_1|n|^{1/\alpha}}\right)^{1/2} \leq \left(\sup_{n \in \mathbb{N}_0^d} |a_n| e^{3h_1|n|^{1/\alpha}}\right) \cdot \left(\sum_{n \in \mathbb{N}_0^d} e^{-h_1|n|^{1/2}}\right)^{1/2} \leq C \eta_h^\alpha(f) < \infty.$$

Hence, $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h_1, \alpha/2}]}^\infty$, and the proof is finished. \square

Remark 2.4. Notice that we just proved that for every sequence $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h, \alpha/2}]}^\infty$ there exists $h_1 > 0$ and $C_1 > 0$ such that

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h, \alpha/2}]}^\infty} \leq C_1 \cdot \eta_{h_1}^\alpha(T(\{a_n\}_{n \in \mathbb{N}_0^d})),$$

and, conversely, for every $f \in \mathbf{G}_{\alpha, h}^{\alpha, h}(\mathbb{R}^d)$ there exists $h_1 > 0$ and $C_1 > 0$ such that $\{a_n\}_{n \in \mathbb{N}_0^d} = T^{-1}(f) \in \ell_{[\vartheta_{h_1, \alpha/2}]}^\infty(\mathbb{N}_0^d)$, and

$$\eta_h^\alpha(f) \leq C_1 \cdot \|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_1, \alpha/2}]}^\infty}.$$

Therefore, the mapping $T : \ell_{\alpha/2}(\mathbb{N}_0^d) \rightarrow \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ is a well defined topological isomorphism.

Thus, Theorem 2.3 implies $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = \mathcal{G}_\alpha(\mathbb{R}_+^d)$, $\alpha > 0$.

When considering the Beurling space $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 0$, we have the following analogue of Theorem 2.3. The proof is omitted since it is similar to the proof of Theorem 2.3.

Theorem 2.5. *Let $\alpha > 0$.*

- i) *If $f \in \mathcal{S}(\mathbb{R}_+^d)$ and $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha/2}(\mathbb{N}_0^d)$, then $f \in \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$.*
- ii) *If $f \in \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, then $a_n(f) \in \ell_{0,\alpha/2}(\mathbb{N}_0^d)$.*
- iii) *If $a_n(f) \in \ell'_{0,\alpha/2}(\mathbb{N}_0^d)$ then the sequence*

$$\sum_{n \in \mathbb{N}_0^d} a_n(f) l_n$$

converges in the topology of dual space $(\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^d)$, thus defining an element $f \in (\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^d)$.

- iv) *Let $f \in (\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^d)$. Then for the sequence of its Fourier-Laguerre coefficients we have $a_n(f) \in \ell'_{0,\alpha/2}(\mathbb{N}_0^d)$.*

Thus, $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = \mathcal{G}_{0,\alpha}(\mathbb{R}_+^d)$, and the mapping $T : \ell_{0,\alpha/2}(\mathbb{N}_0^d) \rightarrow \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ is a well defined topological isomorphism. From Theorems 2.3 and 1.7, it also follows that

$$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = G_\alpha^\alpha(\mathbb{R}_+^d), \quad \alpha \geq 1, \quad \text{and} \quad \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = g_\alpha^\alpha(\mathbb{R}_+^d), \quad \alpha > 1. \quad (18)$$

Moreover,

$$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) \neq G_\alpha^\alpha(\mathbb{R}_+^d), \quad 0 < \alpha < 1, \quad \text{and} \quad \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) \neq g_\alpha^\alpha(\mathbb{R}_+^d), \quad 0 < \alpha \leq 1. \quad (19)$$

Concerning the dual spaces, by Theorems 2.3 and 2.5 it immediately follows that $(\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^d)$ and $(\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^d)$, are topologically isomorphic to sequence spaces $\ell'_{\alpha/2}(\mathbb{N}_0^d)$ and $\ell'_{0,\alpha/2}(\mathbb{N}_0^d)$, $\alpha > 0$, respectively.

Remark 2.6. *Let now $\alpha = 0$, and $h > 0$. We define $\mathbf{G}_{0,h}^{0,h}(\mathbb{R}_+^d)$, as the set of all smooth functions f such that $\|E^N f\|_{L^2(\mathbb{R}_+^d)} \leq Ch^{|N|}$, for some $C > 0$. Then*

$$\mathbf{G}_{0,h}^{0,h}(\mathbb{R}_+^d) = \left\{ \sum_{n \in \mathbb{N}_0^d} a_n l_n \mid a_n = 0 \text{ if } |n| > h \right\},$$

and it follows that $\mathbf{G}_0^0(\mathbb{R}_+^d) = \bigcup_{h>0} \mathbf{G}_{0,h}^{0,h}(\mathbb{R}_+^d)$ is the set of all (finite) linear combinations of Laguerre functions. Therefore $\mathbf{G}_0^0(\mathbb{R}_+^d) = \mathcal{G}_0(\mathbb{R}_+^d)$, see subsection 1.4. In addition, $\mathbf{g}_0^0(\mathbb{R}_+^d) = \bigcap_{h>0} \mathbf{G}_{0,h}^{0,h}(\mathbb{R}_+^d) = \{0\}$.

The kernel theorem for G -type spaces can be extended to P -spaces on positive orthants as follows.

Theorem 2.7. *Let $\alpha > 0$. We have the following:*

$$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^{d_1}) \hat{\otimes} \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^{d_2}) \cong \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^{d_1+d_2}) \quad \text{and} \quad (\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^{d_1}) \hat{\otimes} (\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^{d_2}) \cong (\mathbf{G}_\alpha^\alpha)'(\mathbb{R}_+^{d_1+d_2}),$$

$$\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^{d_1}) \hat{\otimes} \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^{d_2}) \cong \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^{d_1+d_2}) \quad \text{and} \quad (\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^{d_1}) \hat{\otimes} (\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^{d_2}) \cong (\mathbf{g}_\alpha^\alpha)'(\mathbb{R}_+^{d_1+d_2}).$$

Proof. Recall, the sequence spaces $\ell_{\alpha/2}(\mathbb{N}_0^d)$ and $\ell_{0,\alpha/2}(\mathbb{N}_0^d)$ are nuclear. The claim follows by calculations similar to the ones given in the proof of Theorem 6.4 in [10] (see also [11]). We leave details to the reader. \square

3. ALTERNATIVE NORMS FOR DEFINING P -SPACES

In Definition 2.2 we introduced P -spaces by considering L^2 -norms of the iterates of the Laguerre operator, and in Theorem 2.3 we relate them with spaces of sequences of the corresponding Laguerre coefficients. For such spaces of sequences we noted in Remark 1.2 that the weighted ℓ^2 -norm, $\|\cdot\|_{\ell^2_{[\vartheta_{h,\alpha}]}}$, which gives rise to $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$, can be replaced by a weighted ℓ_p norm, $p \in [1, \infty]$, $\|\cdot\|_{\ell^p_{[\vartheta_{h,\alpha}]}}$, without changing the spaces $\ell_\alpha(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d)$, and their topological structure.

In this subsection we demonstrate that the same is true for P -spaces. More precisely, the L^2 -norm, $\|\cdot\|_{L^2(\mathbb{R}_+^d)}$, in (14) can be replaced by $\|\cdot\|_{L^p(\mathbb{R}_+^d)}$, $p \in [1, \infty]$, yielding the same P -space, and preserving its topological structure.

Next we estimate L^p -norms of the Laguerre functions l_n , $n \in \mathbb{N}_0$. Although this result is probably known, we give its proof for the sake of completeness.

Lemma 3.1. *Let $p \in [1, \infty]$. For any Laguerre function l_n , $n \in \mathbb{N}_0^d$, the following estimate holds:*

$$\|l_n\|_{L^p(\mathbb{R}_+^d)} \leq C \prod_{j=1}^d n_j^2 \leq C |n|^{2d}, \quad (20)$$

where the constant $C > 0$ is independent on $n \in \mathbb{N}_0^d$ and $p \in [1, \infty]$.

Proof. Let $p \in [1, \infty)$ (the case $p = \infty$ is obvious), and let us consider the one dimensional case, i.e. $n \in \mathbb{N}_0$. By [2, Thm. 3] it follows that

$$|x^k l_n(x)| \leq 4^k (n+1)(n+2) \cdots (n+k), \quad x \in \mathbb{R}_+, \quad (21)$$

wherefrom

$$\int_0^\infty |l_n(x)|^p dx = \left(\int_0^1 + \int_1^\infty \right) |l_n(x)|^p dx \leq 1 + 16^p (n+1)^p (n+2)^p \int_1^\infty \frac{1}{x^{2p}} dx \leq C^p n^{2p},$$

for a suitable $C > 0$ that does not depend on $n \in \mathbb{N}_0$ and $p \in [1, \infty)$. The same calculation holds for $n \in \mathbb{N}_0^d$, and we get (20). \square

Proposition 3.2. *Let $p \in [1, \infty]$, $\alpha > 0$, and $f \in C^\infty(\mathbb{R}_+^d)$. There exist $h > 0$ and $C > 0$ such that*

$$\|E^N f\|_{L^p(\mathbb{R}^d)} \leq C h^N N!^\alpha, \quad (22)$$

for every $N \in \mathbb{N}_0$, if and only if there exists $h_1 > 0$ such that $f \in \mathbf{G}_{\alpha, h_1}^{\alpha, h_1}(\mathbb{R}_+^d)$.

Proof. Suppose $p \in [1, \infty)$, without loss of generality.

(\Rightarrow) By the Hölder inequality, (22) and Lemma 3.1, for any given $n, N \in \mathbb{N}_0$, we have that

$$|n|^N |a_n(f)| = \left| \int_{\mathbb{R}_+^d} E^N(f)(x) l_n(x) dx \right| \leq \|E^N(f)\|_{L^p(\mathbb{R}_+^d)} \cdot \|l_n\|_{L^{p'}(\mathbb{R}_+^d)} \leq C_1 h^N N!^\alpha |n|^{2d},$$

for a suitable constant $C_1 > 0$, where $\frac{1}{p} + \frac{1}{p'} = 1$. Hence, there exist $C_2, C_3, h_3 > 0$ such that

$$|a_n(f)| \leq \sup_{N \in \mathbb{N}_0} \frac{C_2 h^N N!^\alpha}{|n|^N} \cdot |n|^{2d} \leq C_3 e^{-h_3 |n|^{1/\alpha}} \cdot |n|^{2d},$$

and therefore $\|\{a_n(f)\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_3/2, \alpha/2}]}^\infty} < \infty$. By Remark 1.2 there exist $h_4 > 0$, so that $\|\{a_n(f)\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_4, \alpha/2}]}^2(\mathbb{N}_0^d)} < \infty$, and the claim follows from Theorem 2.3.

(\Leftarrow) Let $f \in \mathbf{G}_{\alpha, h_1}^{\alpha, h_1}(\mathbb{R}_+^d)$ for some $h_1 > 0$, and let $\{a_n(f)\}_{n \in \mathbb{N}_0^d}$ be the sequence of its Laguerre coefficients. By Theorem 2.3 we have that $|a_n(f)| \leq C_2 e^{-h_2 |n|^{1/\alpha}}$ for some $h_2, C_2 > 0$, which, together with Lemma 3.1 gives $\|\sum_{n \in \mathbb{N}_0^d} a_n(f) \cdot l_n\|_{L^p(\mathbb{R}_+^d)} < \infty$. Moreover,

$$\|E^N f\|_{L^p(\mathbb{R}^d)_+} \leq \sum_{n \in \mathbb{N}_0^d} |n|^N |a_n| \cdot \|l_n\|_{L^p(\mathbb{R}_+^d)} \leq C C_2 \sup_{n \in \mathbb{N}_0^d} (|n|^N e^{-\frac{h_2}{2} |n|^{1/\alpha}}) \cdot \sum_{n \in \mathbb{N}_0^d} |n|^{2d} e^{-\frac{h_2}{2} |n|^{1/\alpha}},$$

and (22) now follows from (16). \square

4. RELATION BETWEEN PILIPOVIĆ SPACE ON \mathbb{R}^d AND P -SPACES ON \mathbb{R}_+^d

The relation between G-type spaces and Gelfand-Shilov spaces is studied in [12]. In this section we extend these results to and relate P -spaces on orthants with their natural global counterparts. Throughout the section we will consider the Roumieu case, and note that analogous results hold for the Beurling case. Details are left for the reader.

By $\mathcal{H}_{\alpha, \text{even}}^\alpha(\mathbb{R}^d)$ we denote the space “even” functions in $\mathcal{H}_\alpha^\alpha(\mathbb{R}^d)$:

$$f \in \mathcal{H}_{\alpha, \text{even}}^\alpha(\mathbb{R}^d) \iff f(x) = f(x_1, x_2, \dots, x_d) = f(x_1, \dots, x_{j-1}, -x_j, x_{j+1}, \dots, x_d),$$

for every $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$.

In a similar fashion, one can introduce $\mathcal{S}_{\alpha, \text{even}}^\alpha(\mathbb{R}^d)$, $\Sigma_{\alpha, \text{even}}^\alpha(\mathbb{R}^d)$, and $\mathcal{H}_{0, \alpha, \text{even}}(\mathbb{R}^d)$, $\alpha > 0$. Then we have the following.

Proposition 4.1. [12, Proposition 2.2] *Let $\alpha > 0$. The space $\mathcal{H}_{\alpha, \text{even}}^\alpha(\mathbb{R}^d)$ is a closed subspace of $\mathcal{H}_\alpha^\alpha(\mathbb{R}^d)$, which consists of those $f \in \mathcal{H}_\alpha^\alpha(\mathbb{R}^d)$ which can be represented as $f = \sum_{n \in \mathbb{N}_0^d} a_{2n} h_{2n}$ where $\{a_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_\alpha(\mathbb{N}_0^d)$.*

Proof. The proof for $\alpha \geq 1/2$ is given in [12]. We notice that a more general case $\alpha > 0$ follows by using the same arguments as in the proof of Proposition 2.2 in [12]. \square

Remark 4.2. *The meaning of $\{a_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_\alpha(\mathbb{N}_0^d)$ should be understood in the following way: the sequence $\{b_k\}_{k \in \mathbb{N}_0^d}$ belongs to $\ell_\alpha(\mathbb{N}_0^d)$ if $b_k = a_k$ for all indices $k = 2n$, $n \in \mathbb{N}_0^d$, and if all other elements of $\{b_k\}_{k \in \mathbb{N}_0^d}$ are equal to 0. In the sequel, whenever we use this notation, it will have exactly this meaning.*

Next we establish the relationship between $\mathcal{H}_{\alpha, \text{even}}(\mathbb{R}^d)$ and $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha > 0$. We introduce the same notation as in [12]:

$$v : \mathbb{R}^d \rightarrow \mathbb{R}_+^d, v(x) = (x_1^2, \dots, x_d^2),$$

$$w : \mathbb{R}_+^d \rightarrow \mathbb{R}_+^d, w(x) = (\sqrt{x_1}, \dots, \sqrt{x_d}),$$

$\mathbf{1}/\mathbf{2} = (1/2, \dots, 1/2) \in \mathbb{R}_+^d$, and $\mathbf{3}/\mathbf{2} = (3/2, \dots, 3/2) \in \mathbb{R}_+^d$, and recall two important results given there.

Theorem 4.3. [12, Proposition 3.1] *If $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n \in G_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha \geq 1$. Then $f \circ v \in \mathcal{S}_{\alpha/2, \text{even}}^{\alpha/2}(\mathbb{R}^d)$ and we have*

$$f \circ v = \sum_{n \in \mathbb{N}_0^d} b_{2n} h_{2n},$$

where $\{b_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$, is given by

$$b_{2n}(f) = \frac{(-1)^n \pi^{d/4} \sqrt{(2n)!}}{2^{|n|} n!} \cdot \sum_{k \in \mathbb{N}_0^d} a_{k+n}(f) \binom{k - \mathbf{1}/\mathbf{2}}{k}, \quad n \in \mathbb{N}_0^d. \quad (23)$$

Theorem 4.4. [12, Proposition 3.2] *Let $\alpha \geq 1$, and $f = \sum_{n \in \mathbb{N}_0^d} a_{2n}(f) h_{2n} \in \mathcal{S}_{\alpha/2, \text{even}}^{\alpha/2}(\mathbb{R}^d)$.*

Then $f|_{\mathbb{R}_+^d} \circ w$, belongs to $G_\alpha^\alpha(\mathbb{R}_+^d)$, and $\{b_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$. Moreover

$$b_n = \frac{(-1)^n 2^{|n|}}{\pi^{d/4}} \sum_{k \in \mathbb{N}_0^d} \binom{k - \mathbf{3}/\mathbf{2}}{k} \frac{(-1)^{|k|} 2^{|k|} (k+n)! a_{2(k+n)}}{\sqrt{(2(k+n))!}}, \quad n \in \mathbb{N}_0^d. \quad (24)$$

From [23, Definition 3.1], it follows that image of the sequence space $\ell_\alpha(\mathbb{N}^d)$ under the Hermite expansion mapping $T_h : \{a_n\}_{n \in \mathbb{N}^d} \rightarrow \sum_{n \in \mathbb{N}^d} a_n h_n$ is in fact the Pilipović space $\mathcal{H}_\alpha(\mathbb{R}^d)$ ([23, Theorem 5.2, Lemma 5.7]), $\alpha > 0$.

Since Pilipović spaces extend the class of (Fourier transform invariant) Gelfand-Shilov spaces, it is then natural to ask oneself whether (23) and (24) have counterparts when considering P -spaces.

By the careful examination of the proof of [12, Proposition 3.1] we conclude that almost all estimates given there are valid also when $\alpha \in (0, 1)$, except the inequality

$$|n|^{1/\alpha} + |m|^{1/\alpha} \leq 2(|n| + |m|)^{1/\alpha}, \quad m, n \in \mathbb{N}_0^d \quad (25)$$

(see (3.4) in [12]). To prove extensions of Theorems 4.3 and 4.4 we use the following estimates instead. Let $s = (s_1, s_2, \dots, s_d) \in \mathbb{R}_+^d$ and $\|s\|_p = (\sum_{k=1}^d |s_k|^p)^{1/p}$, $s \in \mathbb{R}_+^d$, $p \in \mathbb{N}$. Then there exist $M_1, M_2 > 0$ (depending on $\alpha > 0$, and p) such that $M_1 \|s\|_p \leq (\sum_{j=1}^d s_j^{\frac{1}{\alpha}})^\alpha \leq M_2 \|s\|_p$, which implies

$$|m|^{1/\alpha} + |n|^{1/\alpha} \leq C|m + n|^{1/\alpha}, \quad m, n \in \mathbb{N}_0^d, \quad \alpha > 0. \quad (26)$$

Thus, if we replace (25) by (26), and otherwise follow the proofs of Propositions 3.1 and 3.2 in [12], we get the following theorem for P -spaces.

Theorem 4.5. *Let $\alpha > 0$. If $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$, then $f \circ v \in \mathcal{H}_{\alpha/2, \text{even}}(\mathbb{R}^d)$ and we have*

$$f \circ v = \sum_{n \in \mathbb{N}_0^d} b_{2n} h_{2n}$$

such that $\{b_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$ and (23) holds. If, on the other hand,

$$f = \sum_{n \in \mathbb{N}_0^d} b_{2n}(f) h_{2n} \in \mathcal{H}_{\alpha/2, \text{even}}(\mathbb{R}^d),$$

then $f|_{\mathbb{R}_+^d} \circ w \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$, $\{b_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$ and moreover, (24) holds.

An analogous result to Theorem 4.5 holds for the Beurling case.

Theorem 4.6. *Let $\alpha > 0$. If $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n \in \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, then $f \circ v \in \mathcal{H}_{0, \alpha/2, \text{even}}(\mathbb{R}^d)$ and we have $f \circ v = \sum_{n \in \mathbb{N}_0^d} b_{2n} h_{2n}$ such that $\{b_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_{0, \alpha/2}(\mathbb{N}_0^d)$ and (23) holds. If, on the other hand,*

$$f = \sum_{n \in \mathbb{N}_0^d} b_{2n}(f) h_{2n} \in \mathcal{H}_{0, \alpha/2, \text{even}}(\mathbb{R}^d),$$

then $f|_{\mathbb{R}_+^d} \circ w \in \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$, $\{b_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{0, \alpha/2}(\mathbb{N}_0^d)$ and (24) holds.

5. FLAT PILIPOVIĆ SPACES, EXTENSION TO \mathbb{R}_b

It is observed by Toft in [23] that together with Pilipović spaces $\mathcal{H}_{\alpha/2}(\mathbb{R}^d)$ and $\mathcal{H}_{0, \alpha/2}(\mathbb{R}^d)$ which are essentially linked to the weights of the form $\vartheta_{h, \alpha}(\cdot) = e^{h|\cdot|^{1/(2\alpha)}}$, $\alpha, h > 0$, one can observe another type of weights, which gives rise to "fine tuning" between Pilipović spaces when $\alpha \in (0, 1/2)$.

Such spaces are called flat Pilipović spaces in [23]. In this section we study their counterparts on positive orthants.

Following [23, Section 3.1], we consider the extension of \mathbb{R}_+ given by

$$\mathbb{R}_b := \mathbb{R}_+ \cup \{b_\sigma : \sigma > 0\}, \quad \text{and} \quad \overline{\mathbb{R}_b} = \mathbb{R}_b \cup \{0\},$$

where the usual ordering holds in \mathbb{R}_+ , and new elements are ordered in the following manner:

$$x_1 < b_{\sigma_1} < b_{\sigma_2} < x_2 \quad \text{if} \quad \sigma_1 < \sigma_2 \quad \text{and} \quad x_1 < 1/2 \leq x_2, \quad x_1, x_2 \in \mathbb{R}_+.$$

In addition, $b_\infty := \frac{1}{2}$.

Let $b_\sigma \in \mathbb{R}_b \setminus \mathbb{R}_+$, $\sigma > 0$, and $h > 0$. We define $\vartheta_{h,b_\sigma}(n) := h^{|n|} \cdot n!^{1/2\sigma}$, and consider the space $\ell_{[\vartheta_{h,b_\sigma}]}^\infty$ of sequences $\{a_n\}_{n \in \mathbb{N}_0^d}$ satisfying

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h,b_\sigma}]}^\infty(\mathbb{N}_0^d)} = \|\{a_n \cdot \vartheta_{h,b_\sigma}(n)\}_{n \in \mathbb{N}_0^d}\|_{l^\infty} = \sup_{n \in \mathbb{N}_0^d} |a_n| \cdot h^{|n|} \cdot n!^{\frac{1}{2\sigma}} < \infty. \quad (27)$$

Moreover,

$$\ell_{b_\sigma}(\mathbb{N}_0^d) := \bigcup_{h>0} \ell_{[\vartheta_{h,b_\sigma}]}^\infty(\mathbb{N}_0^d) \quad \text{and} \quad \ell_{0,b_\sigma}(\mathbb{N}_0^d) := \bigcap_{h>0} \ell_{[\vartheta_{h,b_\sigma}]}^\infty(\mathbb{N}_0^d),$$

in the sense of inductive and projective limit topology, respectively.

We first show inclusion relations between these spaces.

Lemma 5.1. *Let $\alpha, \beta \in \mathbb{R}_b$ and $\alpha < \beta$. Then $\ell_\alpha(\mathbb{N}_0^d) \subseteq \ell_\beta(\mathbb{N}_0^d)$ and $\ell_{0,\alpha}(\mathbb{N}_0^d) \subseteq \ell_{0,\beta}(\mathbb{N}_0^d)$.*

Proof. We will prove the Roumieu case $\ell_\alpha(\mathbb{N}_0^d) \subseteq \ell_\beta(\mathbb{N}_0^d)$, and leave the Beurling case for the reader. In addition, to simplify calculations, we will use $|n|!$ instead of $n!$ without losing generality.

Note that the claim is obvious when $\alpha, \beta \in \mathbb{R}_+$ or $\alpha = b_{\sigma_1}, \beta = b_{\sigma_2}, \sigma_1 < \sigma_2$.

Let $\alpha = b_\sigma, \sigma > 0$, $\beta = \frac{1}{2}$, and let $h > 0$ be chosen so that $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h,b_\sigma}]}^\infty(\mathbb{N}_0^d)$, i.e. $\sup_{n \in \mathbb{N}_0^d} |a_n| \cdot h^{|n|} \cdot |n|!^{\frac{1}{2\sigma}} < \infty$. Using the estimate $|n|^{|n|} \leq |n|!e^n$ we get

$$|a_n|e^{h_1|n|} \leq |a_n| \cdot |n|!^{\frac{1}{2\sigma}} \cdot e^{\frac{|n|}{2\sigma}} \cdot \frac{e^{h_1|n|}}{|n|^{\frac{|n|}{2\sigma}}} \leq C|a_n| \cdot |n|!^{\frac{1}{2\sigma}} \cdot h^{|n|} < \infty,$$

for a suitable $C > 0$, where $h_1 > 0$ is chosen so that $e^{\frac{1}{2\sigma}} e^{h_1} \leq h|n|^{-\frac{1}{2\sigma}}$, for $n \in \mathbb{N}_0^d$ large enough. It follows that $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{1/2}(\mathbb{N}_0^d)$.

Next we prove the claim when $\alpha \in (0, \frac{1}{2})$ and $\beta = b_\sigma, \sigma > 0$. Let $h > 0$ be chosen so that such that $|a_n| \cdot e^{h|n|^{1/(2\alpha)}} < \infty$. Then we have

$$\begin{aligned} |a_n| h_1^{|n|} \cdot |n|!^{\frac{1}{2\sigma}} &\leq |a_n| h_1^{|n|} e^{h|n|^{1/(2\alpha)}} \cdot \frac{|n|^{\frac{|n|}{2\sigma}}}{e^{h|n|^{1/(2\alpha)}}} \\ &\leq |a_n| e^{h|n|^{1/(2\alpha)}} \left(\frac{h_1 |n|^{\frac{1}{2\sigma}}}{e^{h|n|^{\frac{1}{2\alpha}-1}}} \right)^{|n|}, \quad h_1 > 0. \end{aligned}$$

Since $1 > 2\alpha$, we have that $\sup_{n \in \mathbb{N}_0^d} |n|^{\frac{1}{2\sigma}} e^{-h|n|^{\frac{1}{2\alpha}-1}} \leq C_h$, so we can choose $h_1 > 0$ such that $C_h h_1 < 1$, wherefrom it follows that

$$|a_n| h_1^{|n|} |n|!^{\frac{1}{2\sigma}} \leq |a_n| e^{h|n|^{1/(2\alpha)}} < \infty,$$

i.e. $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d)$, $\sigma > 0$.

The inclusion $\ell_{0,\alpha}(\mathbb{N}_0^d) \subseteq \ell_{0,\beta}(\mathbb{N}_0^d)$ can be proved in a similar fashion. \square

Remark 5.2. *The sequence $\{e^{-|n|}\}_{n \in \mathbb{N}_0^d}$ satisfies*

$$\{e^{-|n|}\}_{n \in \mathbb{N}_0^d} \in \ell_{1/2}(\mathbb{N}_0^d) \quad \text{and} \quad \{e^{-|n|}\}_{n \in \mathbb{N}_0^d} \notin \bigcup_{\sigma > 0} \ell_{b_\sigma}(\mathbb{N}_0^d),$$

hence the inclusion $\bigcup_{\sigma > 0} \ell_{b_\sigma}(\mathbb{N}_0^d) \subset \ell_{\frac{1}{2}}(\mathbb{N}_0^d)$ is strict.

The next result can be proved in a similar way as Proposition 1.3.

Proposition 5.3. *Fix an arbitrary $\alpha = b_\sigma$ and $h_1 > h_2$. Then the canonical inclusion $\ell_{[\vartheta_{h_1, b_\sigma}]}^\infty(\mathbb{N}_0^d) \rightarrow \ell_{[\vartheta_{h_2, b_\sigma}]}^\infty(\mathbb{N}_0^d)$ is nuclear.*

We recall that in [23], the spaces of formal expansions

$$\mathcal{H}_{b_\sigma}(\mathbb{R}^d) = \left\{ \sum_{n \in \mathbb{N}_0^d} a_n h_n : \{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d) \right\},$$

$$\mathcal{H}_{0, b_\sigma}(\mathbb{R}^d) = \left\{ \sum_{n \in \mathbb{N}_0^d} a_n h_n : \{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0, b_\sigma}(\mathbb{N}_0^d) \right\}$$

were considered. By $\mathcal{H}_{b_\sigma, \text{even}}(\mathbb{R}^d)$ and $\mathcal{H}_{0, b_\sigma, \text{even}}(\mathbb{R}^d)$ we denote the space of even functions from $\mathcal{H}_{b_\sigma}(\mathbb{R}^d)$ and $\mathcal{H}_{0, b_\sigma}(\mathbb{R}^d)$, respectively.

We introduce the space of formal expansions

$$\mathcal{G}_{b_\sigma}(\mathbb{R}^d) = \left\{ \sum_{n \in \mathbb{N}_0^d} a_n l_n : \{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{b_{\sigma/2}}(\mathbb{N}_0^d) \right\}$$

and its Beurling counterpart

$$\mathcal{G}_{0, b_\sigma}(\mathbb{R}^d) = \left\{ \sum_{n \in \mathbb{N}_0^d} a_n l_n : \{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0, b_{\sigma/2}}(\mathbb{N}_0^d) \right\}.$$

Let us note that, for the arbitrary $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{b_{\sigma/2}}(\mathbb{N}_0^d)$, the expansion $\sum_{n \in \mathbb{N}_0^d} a_n l_n$ is absolutely convergent, thus defining the function in $L^2(\mathbb{R}_+^d)$. The same holds for the Beurling case.

Now, Theorems 4.5 and 4.6 can be extended as follows.

Theorem 5.4. *Let $\sigma > 0$, and $v(x) = (x_1^2, \dots, x_d^2)$, $x \in \mathbb{R}^d$. If $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n \in \mathcal{G}_{b_\sigma}(\mathbb{R}_+^d)$,*

then $f \circ v \in \mathcal{H}_{b_\sigma, \text{even}}(\mathbb{R}^d)$ and we have

$$f \circ v = \sum_{n \in \mathbb{N}_0^d} b_{2n} h_{2n}$$

such that $\{b_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d)$ and (23) holds.

Proof. Suppose that $f = \sum_{n \in \mathbb{N}_0^d} a_n(f) l_n \in \mathcal{G}_{b_\sigma}(\mathbb{R}_+^d)$, i.e.

$$|a_n(f)| \leq C \frac{r^{|n|}}{|n|!^{1/\sigma}}, \quad n \in \mathbb{N}_0^d,$$

for some $C, r > 0$. Since $\sum_{n \in \mathbb{N}^d} |a_n| \leq C \sum_{n \in \mathbb{N}^d} \frac{r^{|n|}}{|n|!^{1/\sigma}} < \infty$, the absolute convergence argument allows us to apply formula (23). Then by using $\sqrt{(2n)!} \leq d^{|n|} n!$, $|k|! \cdot |n|! \leq |k+n|!$, and

$$\left| \binom{k - \mathbf{1}/2}{k} \right| \leq 1,$$

we obtain

$$|b_{2n}(f)| \leq \left| \frac{(-1)^n \pi^{\frac{d}{4}} \sqrt{(2n)!}}{2^{|n|} n!} \right| \cdot \sum_{k \in \mathbb{N}_0^d} |a_{k+n}(f)| \cdot \left| \binom{k - \mathbf{1}/2}{k} \right| \leq C_1 \frac{r^{|n|}}{|n|!^{1/\sigma}} \sum_{k \in \mathbb{N}_0^d} \frac{r^{|k|}}{|k|!^{1/\sigma}}, \quad n \in \mathbb{N}_0^d,$$

for a suitable $C_1 > 0$. Since $\sum_{k \in \mathbb{N}_0^d} \frac{r^{|k|}}{|k|!^{1/\sigma}} < \infty$, we get

$$|b_{2n}(f)| \leq C_2 \frac{r^{|n|}}{|n|!^{1/\sigma}} \leq C_3 \frac{\sqrt{r}^{2n}}{|2n|!^{1/(2\sigma)}},$$

for suitable constants $C_2, C_3 > 0$, which gives

$$\sup_{n \in \mathbb{N}_0^d} |b_{2n}(f)| h^{2n} |2n|!^{1/(2\sigma)} < \infty,$$

where $h := \frac{1}{\sqrt{r}} > 0$. Thus the sequence $\{b_{2n}\}_{n \in \mathbb{N}_0^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d)$ gives rise to the Hermite expansion, and defines a function in the space $\mathcal{H}_{b_\sigma, \text{even}}(\mathbb{R}^d)$. \square

Theorem 5.5. *Let $\sigma > 0$, $w(x) = (\sqrt{x_1}, \dots, \sqrt{x_d})$, $x \in \mathbb{R}_+^d$, and let $f = \sum_{n \in \mathbb{N}_0^d} a_{2n}(f) h_{2n} \in$*

$\mathcal{H}_{b_\sigma, \text{even}}(\mathbb{R}^d)$. *Then $f|_{\mathbb{R}_+^d} \circ w$ belongs to $\mathcal{G}_{b_\sigma}(\mathbb{R}_+^d)$,*

$$f \circ v = \sum_{n \in \mathbb{N}_0^d} b_{2n} h_{2n},$$

$\{b_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d)$, and (24) holds.

Proof. We sketch the main steps of the proof, and note that the absolute convergence can be justified by calculations similar to the ones given in [12]. Suppose that

$$|a_{2n}(f)| \leq C \frac{r^{|2n|}}{|2n|!^{1/(2\sigma)}}, \quad n \in \mathbb{N}_0^d$$

for some $C, r > 0$. From [12, (3.9)] (see also (24)) we get

$$\begin{aligned} |b_n(f)| &\leq \left| \frac{(-1)^n 2^{|n|}}{\pi^{d/4}} \right| \cdot \sum_{k \in \mathbb{N}_0^d} \left| \binom{k - \mathbf{3}/2}{k} \right| \cdot \left| \frac{(-1)^{|k|} 2^{|k|} (k+n)! a_{2k+2n}}{\sqrt{(2k+2n)!}} \right| \\ &\leq C_1 \frac{r^{2n}}{|2n|!^{1/(2\sigma)}} \sum_{k \in \mathbb{N}^d} \frac{r^{2k}}{|2k|!^{1/(2\sigma)}} \leq C_2 \frac{r^{2n}}{|n|!^{1/\sigma}} \end{aligned}$$

for suitable $C_i > 0$, $i = 1, 2$. Thus the sequence $\{b_{2n}\}_{n \in \mathbb{N}^d} \in \ell_{b_\sigma}(\mathbb{N}_0^d)$ gives rise to the Laguerre expansion, and defines a function $f|_{\mathbb{R}_+^d} \circ w \in \mathcal{G}_{b_\sigma}(\mathbb{R}_+^d)$, which concludes the proof. \square

Similar claims hold for $\mathcal{G}_{0,b_\sigma}(\mathbb{R}^d)$, we leave details for the reader.

6. ACKNOWLEDGEMENT

This research was supported by the Science Fund of the Republic of Serbia, #GRANT No. 2727, *Global and local analysis of operators and distributions - GOALS*. The first author is supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia Grant No. 451-03-65/2024-03/ 200169. The second author is supported by the project F10 financed by the Serbian Academy of Sciences and Arts. The third author is supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Grants No. 451-03-66/2024-03/200125 & 451-03-65/2024-03/200125).

REFERENCES

- [1] A. J. Duran, Laguerre expansions of Tempered Distributions and Generalized Functions, *J. Math. Anal. Appl.* 150 (1990), 166-180.
- [2] A. J. Duran, A bound on the Laguerre polynomials, *Studia Math.* 100 (1991), 169-181.
- [3] A. J. Duran, The analytic functionals in the lower half plane as a Gel'fand-Shilov space, *Math. Nachr.* 153 (1991), 145-167.
- [4] A. J. Duran, Laguerre expansions of Gel'fand-Shilov spaces, *J. Approx. Theory* 74 (1993), 280-300.
- [5] A. J. Duran, Gel'fand-Shilov spaces for the Hankel transform, *Indag. Math.* 3 (1992), 137-151.
- [6] A. Erdelyi, *Higher Transcendentals Function*, Vol. 2, McGraw-Hill, New York, 1953.
- [7] I. M. Gel'fand, G. E. Shilov, *Generalized Functions Volume 2 - Spaces of Fundamental and Generalized functions - Academic Press*, 1968.
- [8] T. Gramchev, S. Pilipović, L. Rodino, *Eigenfunction expansions in \mathbb{R}^n* , *Proc. Amer. Math. Soc.* **139** (2011), 4361-4368.
- [9] S. Jakšić, B. Prangoski, Extension theorem of Whitney type for $S(\mathbb{R}_+^d)$ by the use of the Kernel Theorem, *Publ. Inst. Math. Beograd*, 99(113)(2016), 59-65.
- [10] S. Jakšić, S. Pilipović, B. Prangoski, G-type spaces of ultradistributions over \mathbb{R}_+^d and the Weyl pseudo-differential operators with radial symbols, *RACSAM* 111 (2017), 613-640.
- [11] S. Jakšić, S. Pilipović, B. Prangoski, Spaces of ultradistributions of Beurling type over \mathbb{R}_+^d through Laguerre expansions, *Sarajevo J. Math.* 12 (25) (2016), no. 2, suppl., 385-399.
- [12] S. Jakšić, S. Maksimović, S. Pilipović, B. Prangoski, Relations between Hermite and Laguerre expansions of ultradistributions over \mathbb{R}^d and \mathbb{R}_+^d , *J. Pseudo-Differ. Oper. Appl.* 8 (2017), 275-296.
- [13] G. Köthe, *Topological vector spaces I*, Vol.I, Springer-Verlag, New York Inc., 1969.
- [14] G. Köthe, *Topological vector spaces II*, Vol.II, Springer-Verlag, New York Inc., 1979.

- [15] M. Langenbruch, Hermite functions and weighted spaces of generalized functions, *Manuscripta Mathematica* 119 (2006), 269-285.
- [16] Z. Lozanov–Crvenković, D. Perišić, M. Tasković, Hermite expansions of elements of Gelfand-Shilov spaces in quasianalytic and non quasianalytic case, *Novi Sad J. Math.*, 37 (2007), 129–147
- [17] F. Nicola, L. Rodino, *Global Pseudo-Differential Calculus on Euclidean Spaces*, Pseudo-Differential Operators, 4, Birkhäuser, Basel, 2010.
- [18] S. Pilipović, Generalization of Zemanian spaces of generalized functions which elements have series expansions, *SIAM J. Math. Anal.* 17 (1986), 477-484.
- [19] S. Pilipović, Tempered ultradistributions, *Bollettino della Unione Matematica Italiana*, 7(2-B) (1988), 235–251.
- [20] S. Pilipović, On the Laguerre expansions of generalized functions, *C. R. Math. Rep. Acad. Sci. Canada* 11 (1989), 23-27.
- [21] A. Pietsch, *Nuclear locally convex spaces*, Springer-Verlag, Berlin-Heidelberg- New York, 1972.
- [22] S. Thangavelu, *Lectures on Hermite and Laguerre Expansions*, Princeton University Press, Princeton New Jersey (1993)
- [23] J. Toft, Images of function and distribution spaces under the Bargmann transform, *J. Pseudo-Differ. Oper. Appl.* 8 (2017) 83–139
- [24] J. Toft, D. G. Bhimani, R. Manna, Fractional Fourier transforms, harmonic oscillator propagators and Strichartz estimates on Pilipović and modulation spaces, *Appl. Comput. Harmon. Anal.* 67 (2023) 101580
- [25] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*, Dover Publications, New York, 1995.
- [26] Đ. Vučković, J. Vindas, Eigenfunction expansions of ultradifferentiable functions and ultradistributions in \mathbb{R}^d , *J. Pseudo-Differ. Oper. Appl.* 7(4) (2016) 519- 531.
- [27] M. W. Wong, *Weyl Transform*, Springer-Verlag, New York, 1998.
- [28] A. I. Zayed, Laguerre series as boundary values. *SIAM J. Math. Anal.* 13 (1982), 263-279.
- [29] A. H. Zemanian, *Generalized integral transformations*, Interscience, New York, 1968.

FACULTY OF FORESTRY, UNIVERSITY OF BELGRADE, KNEZA VIŠESLAVA 1, BELGRADE, SERBIA
Email address: smiljana.jaksic@sfb.bg.ac.rs

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA OBRADOVIĆA 4, NOVI SAD, SERBIA
Email address: stevan.pilipovic@dmi.uns.ac.rs

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA OBRADOVIĆA 4, NOVI SAD, SERBIA
Email address: tnenad@dmi.uns.ac.rs@dmi.uns.ac.rs

ĐORĐE VUČKOVIĆ, TECHNICAL FACULTY “MIHAJLO PUPIN”, ĐURE ĐAKOVIĆA BB, 23000 ZRENJANIN, SERBIA
Email address: djordjeplusja@gmail.com