Integer programs with nearly totally unimodular matrices: the cographic case

Manuel Aprile Samuel Fiorini Gwenaël Joret Stefan Kober Michał T. Seweryn Stefan Weltge Yelena Yuditsky

July 15, 2024

Abstract

It is a notorious open question whether integer programs (IPs), with an integer coefficient matrix *M* whose subdeterminants are all bounded by a constant ∆ in absolute value, can be solved in polynomial time. We answer this question in the affirmative if we further require that, by removing a constant number of rows and columns from *M*, one obtains a submatrix *A* that is the transpose of a network matrix.

Our approach focuses on the case where *A* arises from *M* after removing *k* rows only, where *k* is a constant. We achieve our result in two main steps, the first related to the theory of IPs and the second related to graph minor theory.

First, we derive a strong proximity result for the case where *A* is a general totally unimodular matrix: Given an optimal solution of the linear programming relaxation, an optimal solution to the IP can be obtained by finding a constant number of augmentations by circuits of $[A \mid I]$.

Second, for the case where *A* is transpose of a network matrix, we reformulate the problem as a maximum constrained integer potential problem on a graph *G*. We observe that if *G* is 2-connected, then it has no rooted $K_{2,t}$ -minor for $t = \Omega(k\Delta)$. We leverage this to obtain a tree-decomposition of *G* into highly structured graphs for which we can solve the problem locally. This allows us to solve the global problem via dynamic programming.

1 Introduction

As for most computational problems that are NP-hard, the mere input size of an integer program (IP) does not seem to capture its difficulty. Instead, several works have identified additional parameters that significantly influence the complexity of solving IPs. These include the number of integer variables (Lenstra [\[LJ83\]](#page-57-0), see also [\[Kan87,](#page-57-1) [Dad12,](#page-56-0) [RR23\]](#page-58-0)), the number of inequalities for IPs in inequality form (Lenstra [\[LJ83\]](#page-57-0)), the number of equations for IPs in equality form (Papadimitriou [\[Pap81\]](#page-58-1), see also [\[EW19\]](#page-56-1)), and features capturing the block structure of coefficient matrices (see for instance [\[CEH](#page-55-0)+21, $CEP⁺21, EHK⁺22, BKK⁺24, CKL⁺24$ $CEP⁺21, EHK⁺22, BKK⁺24, CKL⁺24$]).

Another parameter that has received particular interest is the *largest subdeterminant* of the coefficient matrix, which already appears in several works concerning the complexity of linear programs (LPs) and the geometry of their underlying polyhedra [\[Tar86,](#page-59-0) [DF94,](#page-56-4) [BDSE](#page-55-3)⁺14, [EV17\]](#page-56-5) as well as proximity results relating optimal solutions of IPs and their LP relaxations [\[CGST86,](#page-56-6) [PWW20,](#page-58-2) [CKPW22\]](#page-56-7). Consider an IP of the form

$$
\max \{ p^{\mathsf{T}} x : Mx \le b, x \in \mathbb{Z}^n \},\tag{IP}
$$

where *M* is an integer matrix that is *totally* ∆*-modular*, i.e., the determinants of square submatrices of *M* are all in $\{-\Delta,\ldots,\Delta\}$. It is a basic fact that if *M* is totally unimodular ($\Delta = 1$), then the optimum value of [\(IP\)](#page-0-0) is equal to the optimum value of its LP relaxation, implying that [\(IP\)](#page-0-0) can be solved in polynomial time. In a seminal paper by Artmann, Weismantel & Zenklusen [\[AWZ17\]](#page-54-0), it is shown that [\(IP\)](#page-0-0) is still polynomial-time solvable if $\Delta = 2$, leading to the conjecture that this may hold for every constant Δ . Below, we refer to this conjecture as the *totally* ∆*-modular IP conjecture*. Recently, Fiorini, Joret, Yuditsky & Weltge [\[FJWY22\]](#page-56-8) answered this in the affirmative under the further restriction that *M* has only two nonzeros per row or column. In particular, they showed that in this setting, [\(IP\)](#page-0-0) can be reduced to the stable set problem in graphs with bounded odd cycle packing number [\[BFMRV14,](#page-55-4) [CFHW20,](#page-55-5) [CFH](#page-55-6)+20].

We remark that the algorithm of [\[AWZ17\]](#page-54-0) even applies to full column rank matrices $M \in \mathbb{Z}^{m \times n}$ for which only the ($n \times n$)-subdeterminants are required to be in { $-\Delta$,..., Δ } for $\Delta = 2$. Further results supporting the conjecture have been recently obtained by Nägele, Santiago & Zenklusen [\[NSZ22\]](#page-58-3) and Nägele, Nöbel, Santiago & Zenklusen [\[NNSZ23\]](#page-57-2) who considered the special case where all size-(*n* ×*n*) subdeterminants are in {−∆,0,∆}. Interestingly, the results of [\[AWZ17,](#page-54-0) [NSZ22,](#page-58-3) [NNSZ23\]](#page-57-2) are crucially centered around a reformulation of [\(IP\)](#page-0-0) where *M* becomes *totally unimodular up to removing a constant number of rows*, where the additional constraints capture a constant number of congruency constraints.

1.1 Main contribution

In an effort to provide more evidence for the totally ∆-modular IP conjecture, we initiate the study of IPs in which *M* is totally ∆-modular and *nearly totally unimodular*, i.e., *M* becomes totally unimodular after removing a constant number of rows and columns. Note that without requirements on the subdeterminants, IPs with nearly totally unimodular coefficient matrices are still NP-hard. A famous example is the densest *k*-subgraph problem [\[BCC](#page-54-1)⁺10, [Kho06,](#page-57-3) [Man17\]](#page-57-4), which can be seen as an IP defined by a totally unimodular matrix with two extra rows (modeling a single equality constraint). A closely related example is the partially ordered knapsack problem [\[KS02\]](#page-57-5), which is also strongly NP-hard. Another famous example is the exact matching (or red-blue matching) problem [\[Maa22,](#page-57-6) [MVV87\]](#page-57-7), for which no deterministic polynomial-time algorithm is known (yet).

As we will see, it still does not seem to be an easy task to prove the totally ∆-modular IP conjecture for the full class of nearly totally unimodular matrices. However, in our main result, we resolve the conjecture for a main building block of such matrices:

Theorem 1. *There is a strongly polynomial-time algorithm for solving the integer program* [\(IP\)](#page-0-0) *for the case where M is totally* ∆*-modular for some constant* ∆ *and becomes the transpose of a network matrix after removing a constant number of rows and columns.*

Recall that, to any given (weakly) connected directed graph *G* and spanning tree *T* of *G*, one associates the *network matrix* $A \in \{0, \pm 1\}^{E(T) \times E(G-T)}$ such that $A_{e,(v,w)}$ is equal to 1 if *e* is traversed in forward direction on the unique *v*-*w*-path in *T* , is equal to −1 if it is traversed in backward direction, and is equal to 0 otherwise.

By Seymour's celebrated decomposition theorem for regular matroids [\[Sey80a\]](#page-58-4), network matrices and their transposes are the main building blocks of totally unimodular matrices. While we have not been able to prove an analogous result for network matrices (instead of their transposes) yet, we believe that studying the totally ∆-modular IP conjecture for the above case yields several new insights that are relevant for resolving the general conjecture. It is known that ∆-modular matrices give rise to a proper minor-closed class of \mathbb{F}_p -representable matroids for $p > \Delta$, see for instance [\[GNW21\]](#page-57-8). A key result in the matroid minors project [\[GGW15\]](#page-56-9) states that, in each such family of matroids, every matroid whose vertical connectivity is sufficiently high is a low-rank perturbation^{[1](#page-1-0)} of a frame^{[2](#page-1-1)} or coframe matroid. The

¹ If *M* and *N* are matroids generated by matrices *A* and *B* respectively, where rank(*A*−*B*) ≤ *t*, then *M* is said to be a rank-(≤ *t*) perturbation of *N*.

 2 In this context, a matroid is said to be *frame* if it can be represented over \mathbb{F}_p by a matrix with at most two nonzeros per column, and *coframe* if it is the dual of a frame matroid.

structure of such matroids is highly governed by a graph, and so it is to be expected that questions about graphs become a central aspect of resolving the general totally ∆-modular IP conjecture. In fact, our approach for proving Theorem [1](#page-1-2) strongly builds upon results from the theory of graph minors that are complementary to those that have been used in the work in [\[FJWY22\]](#page-56-8). Thus, we believe that our work significantly enhances our understanding of which questions about graphs are relevant to resolve to general totally ∆-modular IP conjecture.

Moreover, note that nearly totally unimodular matrices *are* low-rank perturbations of well-behaved matrices, an aspect that has not been studied yet in the context of the conjecture. Finally, our insights also lead to a new proximity result (see Theorem [4\)](#page-6-0) in integer programming that generalizes a result by Eisenbrand & Weismantel [\[EW19\]](#page-56-1), yielding another motivation for studying the class of nearly totally unimodular matrices.

1.2 Approach

As indicated above, we achieve our result in two main steps, one related to the theory of integer programming and one related to graph minor theory. More specifically, the first step is concerned with bounds on distances between optimal solutions of IPs and their LP relaxations. A classic result of this type was established by Cook, Gerards, Schrijver, & Tardos [\[CGST86\]](#page-56-6) who showed that if *M* is totally ∆-modular, [\(IP\)](#page-0-0) is feasible, and *x* ∗ is an optimal solution of the LP relaxation, then there exists an optimal solution *z* [∗] of [\(IP\)](#page-0-0) with ∥*x* [∗] − *z* ∗ ∥[∞] ≤ *n*∆. It is still open whether this bound can be replaced with a function in ∆ only, see [\[CKPW22\]](#page-56-7).

A convenient consequence of this result is that, given *x* ∗ , one can efficiently enumerate the possible values of z^{*} for a constant number of variables. In particular, if we are given the integer program [\(IP\)](#page-0-0) with a totally ∆-modular coefficient matrix *M* that becomes totally unimodular after removing *k* rows and *ℓ* columns, we may simply guess the values of the variables corresponding to the *ℓ* columns and solve a smaller IP for each guess.

Thus, we may assume that *M* (is totally ∆-modular and) is of the form $M = \left[\begin{smallmatrix} A \ W \end{smallmatrix} \right]$, where *A* is totally unimodular and *W* is an integer matrix with only *k* rows. For this class of IPs, we derive a considerably strengthened proximity result: Given an optimal solution *x* [∗] of the corresponding LP relaxation, there is an optimal solution z^* of [\(IP\)](#page-0-0) where $||x^* - z^*||_{\infty} \le f(k,\Delta)$ for some function f that depends only on k and ∆ (see Theorem [4\)](#page-6-0), again provided that [\(IP\)](#page-0-0) is feasible. In fact, by bringing [\(IP\)](#page-0-0) into equality form, we show that x^* can be rounded to a closeby integer point from which z^* can be reached by adding a number of conformal *circuits* of $[A \mid I]$ that can be bounded in terms of *k* and ∆ only. Moreover, we observe that the fact that *M* is totally Δ -modular implies that every circuit *c* satisfies $||Wc||_{\infty} \leq \Delta$. (See the next section for definitions and a more precise statement.)

While these findings are valid for all totally unimodular matrices *A*, we will see that they can be crucially exploited for the case where *A* is the transpose of a network matrix, which we refer to as the *cographic case*. For these instances, it is convenient to reformulate the original problem [\(IP\)](#page-0-0) as a particular instance of a *maximum constrained integer potential problem*

$$
\max\{p^{\top}y: \ell(v, w) \le y(v) - y(w) \le u(v, w) \text{ for all } (v, w) \in E(G), Wy = d, y \in \mathbb{Z}^{V(G)}\},\tag{MCIPP}
$$

where *G* is a directed graph, $p \in \mathbb{Z}^{V(G)}$, $\ell, u \in \mathbb{Z}^{E(G)}$, $W \in \mathbb{Z}^{\lfloor k \rfloor \times V(G)}$ and $d \in \mathbb{Z}^k$, and moreover each row of *p*[⊺] or *W* sums up to zero. Notice that the first constraints are still given by a totally unimodular matrix, and hence we may regard $Wy = d$ as extra (or complicating) constraints. With this reformulation, the circuits of $[A \mid I]$ turn into vertex subsets $S \subseteq V(G)$ with the property that both induced subgraphs $G[S]$ and $G[\overline{S}]$ are (weakly) connected, where $\overline{S} := V(G) \setminus S$. We call such sets *doubly connected sets* or *docsets*. Using this notion, our previous findings translate to two strong properties of the instances of [\(MCIPP\)](#page-2-0) we have to solve: First, every feasible instance has an optimal solution that is the sum of at most *f* (*k*,∆) incidence vectors χ^S ∈ {0, 1} $^{V(G)}$, where *S* is a docset. Second, every docset *S* satisfies $\|W\chi^S\|_\infty$ ≤ ∆.

Referring to the vertices whose variables appear with a nonzero coefficient in at least one of the extra constraints as *roots*, the second property above implies that roots cannot be arbitrarily distributed in the input graph. Roughly speaking, by carefully exploiting the structure of the instance, we will be able to guess $y(v)$ for each root v . Note that once all of these variables are fixed, the resulting IP becomes easy since its constraint matrix is totally unimodular. In fact, the guessing cannot be done for the whole graph at once and we will have to do it locally, and then combine the local optimal solutions via dynamic programming.

Figure 1: Subgraph containing a rooted *K*2,3-minor. Roots are indicated with the red squares. Contracting all the edges in each of the five *branch sets* produces a properly rooted $K_{2,3}$.

Our structural insights are based on the observation that our input graphs do not contain a *rooted* $K_{2,t}$ *-minor*, where $t = 4k\Delta + 1$, provided that they are 2-connected. Here, the minors of a rooted graph (graph with a distinguished set of vertices called roots) are defined similarly as for usual graphs, with two differences: whenever some edge *e* is contracted we declare the resulting vertex as a root if and only if at least one of its ends is a root, and we have the possibility to remove a vertex from the set of roots. A rooted *K*2,*^t* is said to be *properly rooted* if each one of the *t* vertices in the "large" side is a root. For the sake of simplicity, we say that a rooted graph contains a *rooted K*_{2,t}*-minor* if it has a rooted minor that is a properly rooted *K*2,*^t* , see Figure [1.](#page-3-0)

Our main structural result is a decomposition theorem for rooted graphs without rooted $K_{2,t}$ -minor, see Theorem [2](#page-3-1) below. It relies partly on several works about the structure of graphs excluding a minor, extending the original results of Robertson & Seymour within the graph minors project, more specifically on works by Diestel, Kawarabayashi, Müller & Wollan [\[DiKMW12\]](#page-56-10), Kawarabayashi, Thomas & Wollan [\[KTW20\]](#page-57-9), and Thilikos & Wiederrecht [\[TW22a,](#page-59-1) [TW22b\]](#page-59-2). Furthermore, we use results of Böhme & Mohar [\[BM02\]](#page-55-7) and Böhme, Kawarabayashi, Maharry & Mohar [\[BKMM08\]](#page-55-8) (see Appendix [A\)](#page-60-0), to control the distribution of the roots in surface-embedded rooted graphs without rooted *K*2,*t*-minors.

Our decomposition theorem is formulated in terms of a tree-decomposition of graph *G*. Recall that a *tree-decomposition* is a pair (T, \mathcal{B}) where *T* is a rooted tree (tree with a unique *root node*) and $\mathcal{B} =$ ${B_u : u \in V(T)}$ is a collection of vertex subsets of *G*, called *bags*, such that for every vertex *v* of *G* the set of bags containing *v* induces a non-empty subtree of *T* , and for every edge *e* of *G* there is a bag that contains both ends of *e*. We define the *weak torso* of a bag *B^u* as the graph obtained from the induced subgraph $G[B_u]$ by adding a clique on $B_u \cap B_{u'}$ for each node $u' \in V(T)$ that is a child of *u*. Having stated these definitions, we are ready to state the decomposition theorem. See Figure [2](#page-4-0) for an illustration.

Theorem 2 (simplified version of Theorem [52\)](#page-33-0). *For every t* $\in \mathbb{Z}_{\geq 1}$ *there exists a constant* $\ell = \ell(t)$ *such that every* 3*-connected rooted graph G without a rooted K*2,*t-minor admits a tree-decomposition* (*T*,B)*, where* $\mathcal{B} = \{B_u : u \in V(T)\}\$, with the following properties:

- (*i*) the bags B_u and $B_{u'}$ of two adjacent nodes $u, u' \in V(T)$ have at most ℓ vertices in common, and
- *(ii)* for every node $u \in V(T)$, all but at most ℓ children $u' \in V(T)$ of u are leaves and the roots contained *in the corresponding bags* $B_{u'}$ *are all contained in bag* B_u *, and*
- *(iii) every node* $u \in V(T)$ *satisfies one of the following:*
	- (a) *bag* B_u *has at most* ℓ *vertices, or*
	- *(b) u is a leaf and B^u has at most ℓ roots, all contained in the bag of the parent of u, or*
	- *(c) after removing at most ℓ vertices, the weak torso of B^u becomes a* 3*-connected rooted graph that does not contain a rooted K*2,*t-minor and has an embedding in a surface of Euler genus at most ℓ such that every face is bounded by a cycle, and all its roots can be covered by at most ℓ facial cycles.*

Figure 2: Illustrating the decomposition of Theorem [2.](#page-3-1) The decomposition tree *T* is shown on the left. The weak torsos of the two bags B_u and $B_{u'}$ are shown on the right. The top one satisfies (iii).(c), and the bottom one (iii).(b).

As we show, there is a polynomial-time algorithm that finds the tree-decomposition of Theorem [2](#page-3-1) together with a polynomial-size collection \mathcal{X}_u for each node $u \in V(T)$, containing all the possible intersections of a docset of *G* with the roots contained in bag *Bu*. This yields an efficient dynamic programming algorithm to solve the instances of [\(MCIPP\)](#page-2-0) we are interested in, which proves Theorem [1.](#page-1-2)

In the next section, we give a more detailed overview of our approach together with important intermediate results, as well as an outline of the paper. The proofs of these results, as well as the missing definitions, can be found in the subsequent sections.

2 Overview

Suppose we are given an instance of [\(IP\)](#page-0-0), where *M* is totally ∆-modular and becomes totally unimodular (TU) after removing a bounded number of rows and columns. Using Tardos' algorithm [\[Tar86\]](#page-59-0), we can solve its LP relaxation in strongly polynomial time. We may assume that the latter has an optimal solu-tion.^{[3](#page-4-1)} Thus, by using the proximity result of [\[CGST86\]](#page-56-6), we may eliminate a bounded number of variables

 3 If the LP relaxation is infeasible, so is [\(IP\)](#page-0-0). If the LP is unbounded, we may first replace the objective by the all-zero vector and run our algorithm. If we find a feasible integer solution, we can conclude that the original instance is also unbounded. Otherwise, the original problem is also infeasible.

such that remaining problem is of the form $max\{p^{\top}x : Ax \leq b, Wx \leq d, x \in \mathbb{Z}^n\}$, where $A \in \mathbb{Z}^{m \times n}$ is TU, £ *A W* ¤ is totally ∆-modular, and *W* ∈ Z *^k*×*ⁿ* has a bounded number of rows. Now that our initial reduction is established, we proceed by explaining how we exploit the fact that $\left[\begin{smallmatrix}A\V\end{smallmatrix}\right]$ is totally ∆-modular. To this end, we consider the circuits of matrices, which are defined as follows.

2.1 Circuits

For a subset $U \subseteq \mathbb{R}^n$, we say that a vector $x \in U \setminus \{0\}$ is a *support-minimal* vector of U if there is no other vector in $U \setminus \{0\}$ whose support is strictly contained in the support of *x*. A *circuit* of a rational matrix $B \in \mathbb{Q}^{m \times n}$ is a support-minimal (nonzero) vector of the kernel of *B* that is integer and reduced (that is, the gcd of its components is 1). The collection of all the circuits of *B* is denoted by $\mathscr{C}(B)$. If $B \in \{0, \pm 1\}^{m \times n}$ is a TU matrix, then it is a basic fact that every circuit of *B* is in ${0, \pm 1}^n$, see for instance Onn [\[Onn10,](#page-58-5) §2.3.4]. We observe the following.[4](#page-5-0)

Lemma 3. Let $A \in \{0, \pm 1\}^{m \times n}$ be totally unimodular, $w \in \mathbb{Z}^n$, and $\Delta \ge 1$. Then $\begin{bmatrix} A \\ w^{\intercal} \end{bmatrix}$ is totally Δ -modular if *and only if every circuit* $\begin{bmatrix} x \\ y \end{bmatrix}$ *of* $\begin{bmatrix} A & I \end{bmatrix}$ *satisfies* $|w^{\top}x| \leq \Delta$ *.*

This implies that, for our instance, every circuit $\begin{bmatrix} x \\ y \end{bmatrix}$ of the matrix $\begin{bmatrix} A & I \end{bmatrix}$ satisfies $||Wx||_{\infty} \leq \Delta$.

2.2 Equality Form

For the next considerations, it is convenient to bring our integer program into equality form. By adding slack variables ν and z , we obtain the equivalent problem

$$
\max\left\{p^{\top}x: Ax+y=b, Wx+z=d, x\in\mathbb{Z}^n, y\in\mathbb{Z}_{\geq 0}^m, z\in\mathbb{Z}_{\geq 0}^k\right\}.
$$

By again using the proximity result of [\[CGST86\]](#page-56-6), we obtain finite lower and upper bounds for all variables that are satisfied by at least one optimal solution of this integer program. Thus, by setting \bar{p} = $[p^{\dagger} \quad \mathbf{0} \quad \mathbf{0}]^{\dagger}$, $\bar{A} = [A \quad \mathbf{I} \quad \mathbf{0}], \ \bar{W} = [W \quad \mathbf{0} \quad \mathbf{I}]$, and $\bar{n} = n + m + k$, our integer program turns into

$$
\max\{\bar{p}^{\top}\bar{x} : \bar{A}\bar{x} = b, \bar{W}\bar{x} = d, \bar{x} \in \mathbb{Z}^{\bar{n}}, \ell \le \bar{x} \le u\}
$$

for some $\ell, u \in \mathbb{Z}^{\bar{n}}$. Now, consider any circuit \bar{x} of \bar{A} . Note that it is of the form $\bar{x} = \begin{bmatrix} x^{\intercal} & y^{\intercal} & \mathbf{0} \end{bmatrix}^{\intercal}$, where $\begin{bmatrix} x \\ y \end{bmatrix}$ is a circuit of $[A \mid I]$. By Lemma [3,](#page-5-1) we obtain $\|\bar{W}\bar{x}\|_{∞} = \|Wx\|_{∞} ≤ ∆$. Thus, we end up with a problem of the following form.

Problem 1. Let $k, \Delta \in \mathbb{Z}_{\geq 1}$ be fixed. Given $p, \ell, u \in \mathbb{Z}^n$, $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Z}^m$, $W \in \mathbb{Z}^{k \times n}$, and $d \in \mathbb{Z}^k$ such that *A* is totally unimodular and $\max\{\|Wx\|_{\infty}: x \in \mathcal{C}(A)\} \leq \Delta$, find

$$
\max\{p^{\mathsf{T}}x: Ax = b, Wx = d, \ell \le x \le u, x \in \mathbb{Z}^n\}.
$$
 (IP₁)

Moreover, in the above transformation, note that if A is the transpose of a network matrix, so is \overline{A} .

Thus, in order to prove Theorem [1,](#page-1-2) we need to give a strongly polynomial time algorithm for instances of Problem [1](#page-5-2) where *A* is the transpose of a network matrix. Before we focus on such matrices, we derive a strengthened proximity result that holds for all totally unimodular matrices *A*.

⁴ Recall that all proofs are deferred to later sections.

2.3 Proximity

For Problem [1,](#page-5-2) we will prove a strengthened proximity result on distances between optimal solutions of (IP_1) (IP_1) and its LP relaxation

$$
\max \{ p^{T} x : Ax = b, Wx = d, \ell \le x \le u \}.
$$
 (LP₁)

Our first result is the following.

Theorem 4. If x^* is an optimal solution of [\(LP](#page-6-1)₁) and [\(IP](#page-5-3)₁) is feasible, then there is an optimal solution z^* *of* [\(IP](#page-5-3)₁) *with* $||x^* - z^*||_{\infty} \le f_4(k, \Delta)$ $||x^* - z^*||_{\infty} \le f_4(k, \Delta)$ $||x^* - z^*||_{\infty} \le f_4(k, \Delta)$ *. Here,* $f_4(k, \Delta)$ *is a function*^{[5](#page-6-2)} *of k and* ∆ *only.*

The proof of Theorem [4](#page-6-0) is based on the following result, which we prove following the strategy by Eisenbrand and Weismantel [\[EW19\]](#page-56-1). We remark that our proof of Proposition [5](#page-6-3) extends the proximity result in [\[EW19\]](#page-56-1) to general totally unimodular matrices with a constant number of additional equalities, and recovers the proximity bound for their setting, that is, an equality system with a constant number of constraints, and entries bounded by a constant in absolute value.

Here, we say that vectors c_1, \ldots, c_t $\in \mathbb{R}^n$ are *conformal* if, in every coordinate, they have the same sign, that is, c_1, \ldots, c_t are contained in the same orthant (see [\[Onn10,](#page-58-5) §3.1]).

Proposition 5. *Given an instance of Problem [1,](#page-5-2) in strongly polynomial time we can compute an integer point* $z \in \mathbb{Z}^n$ satisfying $Az = b$, $\ell \le z \le u$ such that there is an optimal solution to (IP_1) of the form $z +$ $\sum_{j=1}^t c_j$ where $c_1,...,c_t$ \in $\mathscr{C}(A)$ are conformal and $t \leq f_5(k,\Delta)$ $t \leq f_5(k,\Delta)$ $t \leq f_5(k,\Delta)$, or conclude that [\(IP](#page-5-3)₁) is infeasible.

Notice that after applying Proposition [5,](#page-6-3) we can perform a translation of the solution space mapping the integer point *z* to the origin. This allows us to reduce to the special case of Problem [1](#page-5-2) where $b = 0$, and look for an optimal solution $x \in \mathbb{Z}^n$ that is a conformal sum of at most $f_5(k, \Delta)$ $f_5(k, \Delta)$ $f_5(k, \Delta)$ circuits of *A*.

2.4 Increasing the connectivity of the TU matrix

In order to deal with Problem [1,](#page-5-2) it is convenient to ensure some connectivity properties of *A*. Suppose that, possibly after permuting the columns of *A*, we can write $A = \begin{bmatrix} A_1 & A_2 \end{bmatrix}$ in such a way that the column space of $A_1 \in \{0, \pm 1\}^{m \times n_1}$ and the column space of $A_2 \in \{0, \pm 1\}^{m \times n_2}$ intersect in a linear subspace of dimension at most *q* − 1, for some *q* ∈ $\mathbb{Z}_{\geq 1}$. Provided that min{*n*₁, *n*₂} ≥ *q*, the corresponding partition of the columns of *A* is a *separation of order q*, or shortly a *q-separation*. We define the *connectivity* of *A* as the minimum *q* such that *A* has a *q*-separation, and call *A q-connected* if its connectivity is at least *q*.

In case the connectivity of *A* is low, it is natural to split $(IP₁)$ $(IP₁)$ into two IPs with bounded interaction. We show how to do this for separations of order $q \le 2$, and give a (strongly) polynomial-time, black-box reduction of instances of Problem [1](#page-5-2) to instances such that *A* is *almost* 3*-connected*, in the sense that *A* becomes 3-connected when we delete from *A* every column that is a nonzero multiple of another column. We point out that special care should be taken because of the extra constraints $Wx = d$.

Connectivity is a fundamental concept for studying TU matrices and the corresponding matroids, which are known as *regular* matroids. By Seymour's decomposition theorem [\[Sey80a\]](#page-58-4), if *A* is 4-connected then there exists a network matrix $D \in \{0, \pm 1\}^{r \times (n-r)}$ such that $Ax = 0$ is equivalent either to $[D \quad I_r] x = 0$ or to $[I_{n-r}$ −*D*^T] $x = 0$. The two cases are dual of each other. In order to prove Theorem [1,](#page-1-2) we now focus on the latter case.

⁵All functions of *k* and ∆ defined in this paper are in fact computable.

2.5 Reducing to the maximum constrained integer potential problem

Assume that $Ax = 0$ is equivalent to $[I_{n-r} - D^{\dagger}]x = 0$ where $D \in \{0, \pm 1\}^{(n-r) \times r}$ is a network matrix, and $b = 0$. We perform a change of variables that transforms any such instance of Problem [1](#page-5-2) into an instance of the maximum constrained integer potential problem.

Let *G* denote a (weakly) connected directed graph such that $D = B^{-1}N$ where $M = \begin{bmatrix} N & B \end{bmatrix}$ is the vertex-edge incidence matrix of *G* with one row deleted, and *B* is a basis of *M*. Writing $x \in \mathbb{R}^n$ as $x = \left[\frac{x_N}{x_B}\right]$, we have

$$
Ax = \mathbf{0} \iff x_N - D^{\mathsf{T}} x_B = \mathbf{0} \iff x_N = D^{\mathsf{T}} x_B \iff x_N = N^{\mathsf{T}} B^{-\mathsf{T}} x_B.
$$

Now let $y := B^{-\mathsf{T}} x_B$. Then $x = \begin{bmatrix} N^{\mathsf{T}} y \\ B^{\mathsf{T}} y \end{bmatrix} = M^{\mathsf{T}} y$ and (IP₁) can be rewritten as

$$
\max \{ p^{\top} M^{\top} y : \ell \le M^{\top} y \le u, \ W M^{\top} y = d, \ y \in \mathbb{Z}^r \} .
$$
 (IP₁-cographic)

Note that $x = \begin{bmatrix} x_N \\ x_B \end{bmatrix} \in \mathbb{Z}^n$ holds if and only if $y \in \mathbb{Z}^r$, since *B* is unimodular and $x_N = N^{\top}B^{-\top}x_B$. We claim that (IP₁[-cographic\)](#page-7-0) is equivalent to [\(MCIPP\)](#page-2-0). In order to see this, let $v_0 \in V(G)$ denote the vertex whose row is missing from *M*. Notice that $r = |V(G - v_0)| = |V(G)| - 1$. Now append to *M* the missing row for v_0 , and similarly append a row to y for v_0 . After performing this modification and renaming the profit vector and weight matrix, $\text{(IP}_1\text{-cographic)}$ $\text{(IP}_1\text{-cographic)}$ $\text{(IP}_1\text{-cographic)}$ transforms precisely into [\(MCIPP\)](#page-2-0). (Now that *M* is the full incidence matrix of *G*, notice that we have M^{\dagger} **1** = **0**, hence each row of $p^{\dagger}M^{\dagger}$ or WM^{\dagger} in (IP₁[-cographic\)](#page-7-0) sums up to zero.)

Consider the resulting instance of the maximum constrained integer potential problem, see [\(MCIPP\)](#page-2-0). Let $R := \{v \in V(G) : \exists i \in [k] : W(i, v) \neq 0\}$ denote the set of roots of *G*. Below, we denote the resulting rooted graph as (*G*,*R*). (Also, we regard *G* as an undirected graph most of the time, since the edge directions are relevant only when we go back to solving [\(MCIPP\)](#page-2-0).)

By applying Lemma [3](#page-5-1) and Proposition [5,](#page-6-3) we infer the two properties of the resulting MCIPP instance mentioned in the previous section: First, provided that the instance is feasible, it has an optimal solution that is the sum of at most $f_5(k,\Delta)$ $f_5(k,\Delta)$ $f_5(k,\Delta)$ incidence vectors of docsets. Second, the weight vector $W\chi^S$ of every docset *S* has all its components in [−∆,∆] ∩ Z. We in fact obtain two further properties. Third, since *A* is (without loss of generality) almost 3-connected, *G* is a 3-connected graph some of whose edges are subdivided. Fourth, (G, R) does not contain a rooted $K_{2,t}$ -minor for $t = \Omega(k\Delta)$. This fourth property, which follows from the second, is the starting point of our structural analysis. The precise result we show is as follows.

Lemma 6. *Let k*,∆ ∈ Z≥¹ *be fixed, and let t* := 4*k*∆+1*. Consider an instance of the maximum constrained integer potential problem, see* [\(MCIPP\)](#page-2-0). If G is 2-connected and $W \in \mathbb{Z}^{\{k\} \times V(G)}$ satisfies $||W \chi^S||_{\infty} \leq \Delta$ for *all docsets S, then* (G, R) *does not contain a rooted* $K_{2,t}$ *-minor.*

2.6 Decomposing graphs forbidding a rooted *K*2,*t***-minor**

Lemma [6](#page-7-1) motivates the structural investigation of rooted graphs (G, R) with no rooted $K_{2,t}$ -minor. (We mention in passing that rooted *K*2,*t*-minors were studied before in the literature, for instance in connection to the problem of computing the genus of apex graphs, see Mohar [\[Moh01\]](#page-57-10).) On this front, the main result we prove is Theorem [2.](#page-3-1) We start here with a general discussion of the theorem and its context, follow this with an examination of its algorithmic consequences in Section [2.7,](#page-8-0) and conclude by outlining its proof in Section [2.8.](#page-10-0)

We point out that Theorem [2](#page-3-1) assumes *G* to be 3-connected. This is justified by our results on 1- and 2-separations. In case *G* is a 3-connected graph some of whose edges are subdivided, we may suppress the degree-2 vertices, and apply the theorem to the resulting 3-connected graph. For the rest of the discussion, we assume that *G* is 3-connected.

There are a number of basic cases for which Theorem [2](#page-3-1) holds with a trivial or almost trivial treedecomposition (T, \mathcal{B}) .

First, if the number of roots of (G, R) is bounded by some constant ℓ , then (i), (ii) and (iii).(b) always hold in Theorem [2](#page-3-1) for $T := (\{r, u\}, \{r u\}), B_r := R$ and $B_u := V(G)$. Hence, we may assume without loss of generality that (*G*,*R*) has many roots.

Second, it is easy to see that a 3-connected graph *G* with at least two roots always has a rooted *K*2,2 minor, hence the case *t* = 2 of Theorem [2](#page-3-1) is trivial. Robertson and Seymour [\[RS90\]](#page-58-6) characterized the 3-connected rooted graphs (*G*,*R*) with no rooted *K*2,3-minor: either *G* has at most 2 roots, or *G* is planar and has all its roots incident to a common face. This settles the case *t* = 3 of Theorem [2:](#page-3-1) we let *ℓ* := 2 and (T, \mathcal{B}) be the *trivial* tree-decomposition with $T := (\{u\}, \emptyset)$ and $B_u := V(G)$. Then (i) and (ii) trivially hold, and either (iii).(a) or (iii).(c) holds. Hence we may assume without loss of generality that $t \ge 4$.

Third, assume that *G* is a planar 3-connected graph without a rooted *K*2,*t*-minor, where *t* is any fixed integer, and consider again Theorem [2.](#page-3-1) We let once more (T,\mathscr{B}) be the trivial tree-decomposition, which in particular implies (i) and (ii). It turns out that (iii).(c) always holds, in virtue of the following result of Böhme & Mohar.

Theorem [7](#page-8-1). *[Theorem 1.2 of [\[BM02\]](#page-55-7)] There is a function* $f_7 : \mathbb{Z}_{\geq 1} \to \mathbb{Z}_{\geq 1}$ *such that the following holds. Let G be a 3-connected, planar graph and let R* ⊆ *V* (*G*) *be a set of roots. Assume that* (*G*,*R*) *does not have a rooted K*_{2,*t*}-minor. Then G has a collection of at most $f_7(t)$ $f_7(t)$ $f_7(t)$ facial cycles such that each root is contained in *at least one of these cycles.*

Fourth, assume that *G* is a 3-connected graph embedded in a surface of Euler genus $g \ge 1$, where *g* is bounded.^{[6](#page-8-2)} It turns out that Theorem [7](#page-8-1) extends to this case, assuming that the face-width of *G* is large enough:

Theorem [8](#page-8-3). *There is a function* f_8 : $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 2}$ *, such that the following holds. Let* (G, R) *be a* 3*connected rooted graph without a rooted K*2,*t-minor, embedded in a surface of Euler genus g with facewidth at least* $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ *. Then G has a collection of at most* $f_8(g, t)$ *facial cycles covering all the roots.*

This result was stated in Böhme, Kawarabayashi, Maharry & Mohar [\[BKMM08\]](#page-55-8) as a remark, without a proof. We provide a proof in Appendix [A.](#page-60-0) Theorem [8](#page-8-3) implies that Theorem [2](#page-3-1) holds with a trivial treedecomposition whenever *G* can be embedded in a surface of bounded genus, with large face-width. In case the face-width of the embedding is not large enough, then several things might happen. For instance, it might be that there exists a small vertex subset *Z* such that $G - Z$ is 3-connected and has large face-width. The vertices in *Z* are commonly known as *apices*. In this case, (iii).(c) of Theorem [2](#page-3-1) still holds. Beyond this case, we have to decompose the graph in a non-trivial way.

Roughly speaking, Theorem [2](#page-3-1) states that every 3-connected rooted graph forbidding a rooted *K*2,*t*minor can be obtained by gluing in a tree-like fashion the graphs that we examined so far: graphs with a bounded number of roots, and graphs with bounded genus whose roots can be covered with a bounded number of face boundaries, plus a bounded number of apices.

2.7 Algorithmic consequences of the decomposition theorem

We discuss here how Theorem [2](#page-3-1) leads to an efficient algorithm for solving the instances of the maximum constrained integer potential problem resulting from cographic instances of Problem [1.](#page-5-2)

Consider a node $u \in V(T)$ of the decomposition tree and the corresponding bag $B_u \subseteq V(G)$. We claim that every docset of *G* intersects the roots in B_u in a polynomial number of ways. If (iii).(a) or (iii).(b)

 6 See Section 6 for the missing definitions.

holds, there are at most ℓ roots in B_u . Hence, any docset of *G* can intersect the roots in B_u in at most 2^{ℓ} ways, which is a constant. (Recall that $t = 4k\Delta + 1$ and $\ell = \ell(t)$.)

From now on, assume that (iii).(c) holds. Let $G^{\#}[B_{u}]$ denote the weak torso of B_{u} and Z \subseteq $V(G^{\#}[B_{u}])$ denote the set of apices. That is, $|Z| \leq \ell$, and $G^{\#}[B_u] - Z$ is a 3-connected graph without a rooted $K_{2,t}$ minor that has an embedding in a surface of Euler genus at most *ℓ* such that every face is bounded by a cycle, and admits a collection *√* of at most *ℓ* facial cycles covering all its roots.

Consider any docset *S* of *G*, any facial cycle *C* in \mathcal{C} , and the roots covered by *C*. Observe that the roots in *C* are cyclically ordered. Our strategy to establish the claim is to prove that the intersection of *S* with the roots in *C* can be partitioned into a bounded number of blocks of consecutive elements in the cyclic ordering, say at most $\kappa = \kappa(t)$ intervals. Since every root in B_u is either in *Z* or in one of the cycles in $\mathscr C$, this directly yields a $n^{O(\kappa\ell)}$ bound on the number of possible intersections of S with the roots in $B_u.$ We elaborate briefly on the reasons why the above strategy is sound.

First, consider the case where there are no apices, that is $Z = \emptyset$. In $G^* [B_u]$, we add a clique on $B_u \cap B_{u'}$ where *u'* is the parent of *u*. The resulting graph is the (full) *torso* of B_u . Since $B_u \cap B_{u'}$ has at most ℓ vertices, the torso of B_u does not contain a rooted $K_{2,t'}$ -minor for $t' := 2^{\text{poly}(\ell)} t$. Notice that the intersection of docset *S* with the torso of B_u is in fact a docset of the latter. We show that if the intersection of *S* with the roots in some $C \in \mathcal{C}$ cannot be partitioned into a small number of blocks of consecutive elements, we find a $K_{3,s}$ -minor in $G^*[B_u]$ for some $s = s(\kappa,\ell)$ that is increasing in κ and decreasing in *ℓ*. If we choose *κ* large enough, then *s* can be made larger than any given linear function in ℓ . This contradicts the fact that the weak torso $G^{\#}[B_u]$ is embedded in a surface of Euler genus at most ℓ , since the Euler genus of *K*3,*^s* is linear in *s*.

Second, in the presence of apices, the argument gets slightly more involved and in one case we obtain a rooted $K_{2,t'}$ -minor in the torso of B_u , which is also a contradiction.

Finally, we state a result that formalizes the way in which we use our decomposition theorem algorithmically. It is also stated in terms of tree-decompositions. However, we slightly simplify the treedecomposition of Theorem [2](#page-3-1) by merging some leaf bags in their parents. This has some advantages, in particular for optimization purposes.

Let $\ell \in \mathbb{Z}_{\geq 1}$ be fixed. We say that a (rooted) tree-decomposition (T, \mathcal{B}) of *G* is ℓ -special if (i) for each *t* $t' \in E(T)$ we have $|B_t \cap B_{t'}| \leq \ell$; (ii) every node $t \in V(T)$ has at most ℓ children.

Let $\mathcal{S} = \mathcal{S}(G)$ denote the collection of all docsets in *G*. For $R' \subseteq R$, the *docset profile* of R' is the collection of sets $\mathcal{P}_G(R') := \{R' \cap S : S \in \mathcal{S}\}\$. A *docset superprofile* of R' is a collection of subsets of R' that is a superset of the docset profile $\mathscr{P}_G(R')$.

We prove the following result, which is the centerpiece of our algorithm.

Theorem [9](#page-9-0). *There exists a function* $f_9: \mathbb{Z}_{\geq 1} \to \mathbb{Z}_{\geq 1}$ *such that for every fixed* $t \in \mathbb{Z}_{\geq 1}$ *, there is a polynomialtime algorithm that given a rooted* 3*-connected graph* (*G*,*R*) *having no rooted K*2,*t-minor, outputs an f*_{[9](#page-9-0)}(*t*)*-special tree-decomposition* (*T*, \mathcal{B}) *of G and a collection* { \mathcal{X}_u : $u \in V(T)$ } *where each* \mathcal{X}_u *is a docset superprofile of* $R \cap B_u$ *in G of size polynomial in* $|V(G)|$ *.*

Given Theorem [9,](#page-9-0) it is easy to design a strongly polynomial time algorithm to solve the instances of the maximum constrained integer potential problem that originate from cographic instances of Problem [1,](#page-5-2) see [\(MCIPP\)](#page-2-0). Roughly speaking, the idea is to first compute polynomially many optimal local solutions for each bag B_u , namely, one for each guess on the variables $y(v)$ where $v \in B_u$ is a root or belongs to an adjacent bag *B^u* ′ . For each fixed bag and guess, an optimal local solution can be found in (strongly) polynomial time by solving a single IP on a TU constraint matrix. Next, we use a dynamic programming approach to find out optimal ways to combine the optimal local solutions. In virtue of Proposition [5](#page-6-3) and Theorem [9,](#page-9-0) the whole dynamic programming algorithm runs in (strongly) polynomial time.

2.8 Proving the decomposition theorem

The purpose of this section is to describe our proof strategy for Theorem [2.](#page-3-1) Before this, we briefly discuss how to deduce Theorem [9](#page-9-0) from Theorem [2.](#page-3-1) Let (*G*,*R*) be a rooted 3-connected graph without a rooted $K_{2,t}$ -minor, and let (T,\mathscr{B}) denote the tree-decomposition of *G* that we get as the output of Theorem [2.](#page-3-1) Consider any node $u \in V(T)$. We say that a child u' of u is *tame* if u' is a leaf and there is no root in $B_{u'} \setminus B_u$. We turn (T,\mathscr{B}) into an ℓ -special tree-decomposition (T',\mathscr{B}') satisfying the statement of Theorem [9,](#page-9-0) as follows: For each node $u \in V(T)$, we contract every edge connecting u to a tame child u' , adding all the vertices of $B_{u'}$ to B_u . After observing that the original tree-decomposition (*T, 3*) can be computed in polynomial time, we prove that the docset profile \mathscr{P}_G (*R* ∩ *B*^{*t*} u' is of polynomial size for each $u \in V(T')$, and that we can furthermore compute a docset superprofile $\mathscr{X}_u \supseteq \mathscr{P}_G(R \cap B'_i)$ u' for each $u \in V(T')$ in polynomial time.

Now, we turn to the proof of Theorem [2.](#page-3-1) We build the required tree-decomposition recursively. Assume we are sitting at some node $u \in V(T)$. The decomposition process defines for us a corresponding induced subgraph *H* of *G*. Let $R_H = R \cap V(H)$. We consider two cases for *H*. We say that *H* is *k*-interesting (for an integer $k = k(t)$ chosen large enough for the rest of our arguments) if *H* contains a set of vertices *X* such that $|X| = k$ and *X* is well-connected with respect to the vertices in it and is well-connected to *R^H* . If *H* is not *k*-interesting then we call it *k-boring*. It is easy to deal with the *k*-boring graphs. In each such graph, we find 3 subsets $B_0, B_1, B_2 \subseteq V(H)$ such that we can set B_0 to be a child bag of *u* in our decomposition and either further decompose *H*[*B*1] and *H*[*B*2] or just decompose one of them and the other becomes a leaf bag.

Figure 3: An elementary wall of height 6. Walls of height *h* are defined as subdivisions of elementary walls of the same height *h*.

In what follows, we focus on *k*-interesting subgraphs *H*. For an integer *h*, if *k* is large enough as a function of *h* then it is possible to find a *wall W* of height of *h* in *H* (see Figure [3\)](#page-10-1) such that there are *h* vertex-disjoint paths between the set *X* and any *transversal set* of *W* . A transversal set of a wall is a maximum collection of vertices of degree 3 in *W* such that no two vertices are chosen from the same column or the same row of *W*. Moreover, there are *h* vertex-disjoint paths between R_H and any transversal set of *W*. We show that because *H* does not contain a rooted $K_{2,t}$ -minor, *H* cannot contain a *K*^{*t*} - minor *grasped* by the wall *W* for $t' = t'(t)$ and $h \ge t'$.

Using the results in [\[KTW20\]](#page-57-9) and [\[DiKMW12\]](#page-56-10) we derive that graphs *H* have a restricted structure: There is a bounded-size set $Z \subseteq V(H)$ such that $H - Z$ has an embedding on a bounded genus surface Σ , with a bounded number of *large vortices* and a possibly unbounded number of *small vortices*. Moreover *H* − *Z* contains a large wall *W*^{*'*} which can be essentially found as a minor of *W*. The wall *W'* is drawn inside a disk on Σ and is far enough from any large vortex. We handle the large vortices using the results in [\[TW22b\]](#page-59-2). We again use that the fact that *H* does not contain a rooted $K_{2,t}$ -minor to show that after removal of a bounded number of vertices (which are added to *Z*), each large vortex can be either drawn on Σ or turned into a small vortex.

The graph we obtain after the removal of the new set *Z* is not necessarily 3-connected, but after additional steps, including considering the SPQR tree for $H - Z$, we can ensure 3-connectivity while keeping large face-width. This allows us to apply Theorem [8](#page-8-3) and find a small collection of facial cycles covering all the roots in *R^H* that are not in *Z*.

2.9 Outline

We begin the main part of our paper in Section [3](#page-11-0) with an in-depth treatment of circuits and proximity, including proofs for Lemma [3,](#page-5-1) Theorem [4](#page-6-0) and Proposition [5.](#page-6-3) We follow this in Section [4](#page-15-0) by designing black-box reductions in the presence of 1- or 2-separations in the TU matrix *A*. These first two sections consider general TU matrices. Starting from Section [5,](#page-24-0) we focus on the cographic case. We prove that none of our input graphs contains a rooted *K*2,*t*-minor, see Lemma [6,](#page-7-1) and give a polynomial-time dynamic programming algorithm solving the cographic instances of Problem [1,](#page-5-2) assuming Theorem [9.](#page-9-0) This proves Theorem [1.](#page-1-2) Section [6](#page-32-0) establishes our structural results on rooted graphs forbidding a rooted *K*2,*t*minor, including Theorem [2](#page-3-1) and Theorem [9.](#page-9-0) We close our paper in Section [7](#page-53-0) with remarks on possible approaches to generalize Theorem [1](#page-1-2) to all nearly TU matrices *M*, and a discussion of related research questions.

3 Circuits and Proximity

In this section, we collect some basic facts about circuits of totally unimodular matrices, including a proof of Lemma [3,](#page-5-1) and provide proofs for our main results on proximity, Theorem [4](#page-6-0) and Proposition [5.](#page-6-3)

3.1 Decompositions into Circuits

Let us start by proving Lemma [3,](#page-5-1) which states that $\left[\frac{A}{w^1}\right]$ is totally ∆-modular if and only if every circuit $\left[\frac{x}{y}\right]$ of $[A \mid I]$ satisfies $|w^{\top}x| \leq \Delta$. Our proof is based on the following observation. Here, for a matrix $A \in \mathbb{R}^{m \times n}$ and $I \subseteq [m], I \subseteq [n]$, the submatrix of *A* whose rows correspond to *I* and whose columns correspond to *J* is denoted by $A_{I,J}$. Similarly, the restriction of a vector $x \in \mathbb{R}^n$ to the entries indexed by *J* is denoted by *xJ* .

Remark 10. Let $A \in \mathbb{R}^{m \times n}$, $w \in \mathbb{Z}^n$, and $x \in \text{ker}(A)$ such that $w^\dagger x \neq 0$. Then x is support-minimal in $\text{ker}(A)$ if and only if $\left[\frac{A}{w^{\intercal}}\right]$ has an invertible (hence square) submatrix of the form $\left[\frac{A_{I,J}}{w^{\intercal}_I}\right]$ w_J^{\dagger} $}|$ *with* supp(*x*) ⊆ *J*.

Proof. Let *x* be support-minimal in ker(*A*) and set $J = \text{supp}(x)$. Then ker($A_{[m],J}$) is the 1-dimensional subspace spanned by x_j . Indeed, given any two linearly independent vectors y , y' in ker($A_{[m],j}$), there is a nonzero linear combination of *y*, *y'* whose support is strictly contained in *J*. Thus, rank($A_{[m],J}$) = |*J*|−1 and hence there is a subset *I* ⊆ [*m*] with rank(*A_{I,J}*) = |*J*| − 1. Consequently, $\begin{bmatrix} A_{I,J} \\ w^I_I \end{bmatrix}$ \vec{w}_J^{\dagger} i is invertible.

Let $I \subseteq [m], J \subseteq [n]$ such that $\begin{bmatrix} A_{I,j} \\ w^T \end{bmatrix}$ | is invertible and supp(*x*) ⊆ *J*. Since rank(A ^{*I*},*J*) = |*J*| − 1, we see \vec{w}_J^{\dagger} that ker($A_{I,J}$) is the 1-dimensional subspace spanned by x_J . Note that for every $y \in \text{ker}(A) \setminus \{0\}$ with $supp(y) \subseteq supp(x) \subseteq J$ we have $y_I \in ker(A_I,_I)$ and hence *y* is a scalar multiple of *x*, in particular, supp(*y*) = $supp(x)$. \Box

Proof of Lemma [3.](#page-5-1) Foremost, notice that *A* is TU if and only if $[A \mid I]$ is TU, and $\begin{bmatrix} A \\ w^T \end{bmatrix}$ is totally ∆-modular if and only if $\begin{bmatrix} A & \mathbf{I} \\ w^{\intercal} & \mathbf{0}^{\intercal} \end{bmatrix}$ is totally Δ -modular.

Suppose first that $\begin{bmatrix} A \\ w^T \end{bmatrix}$ is totally ∆-modular and consider any circuit $\begin{bmatrix} x \\ y \end{bmatrix} \in \{0, \pm 1\}^{n+m}$ of $\begin{bmatrix} A & I \end{bmatrix}$. If $w^{\top} x \neq 0$, then by Remark [10,](#page-11-1) there exist *I* ⊆ [*m*], *J* ⊆ [*n*], *K* ⊆ [*m*] with supp(*x*) ⊆ *J* and supp(*y*) ⊆ *K* such that $B := \begin{bmatrix} A_{I,J} & I_{I,K} \\ w^{\mathsf{T}} & \mathbf{0}^{\mathsf{T}} \end{bmatrix}$ $\left[\begin{smallmatrix} A_{I,J} & I_{I,K} \ w_I^\intercal & \mathbf{0}^\intercal \end{smallmatrix}\right]$ is invertible. Consider the adjugate matrix adj(*B*) of *B* and let $\left[\begin{smallmatrix} \bar{x}_I & \bar{y}_I \ \bar{y}_K & \bar{y}_I \end{smallmatrix}\right]$ $\left[\frac{\bar{x}_J}{\bar{y}_K}\right]$ ∈ $\{0, \pm 1\}^{J\cup K}$

denote the last column of adj(*B*). Since *B* adj(*B*) = det(*B*)**I**, we have $\begin{bmatrix} A_{I,J}\tilde{x}_J + \mathbf{I}_{I,K}\tilde{y}_K \\ \tilde{y}_I + \tilde{y}_I + \tilde{y}_I \end{bmatrix}$ $\begin{bmatrix} \bar{x}_J + \mathbf{I}_{I,K} \bar{y}_K \\ w_J^\intercal \bar{x}_J \end{bmatrix} = B \begin{bmatrix} \bar{x}_J \\ \bar{y}_K \end{bmatrix}$ $\begin{bmatrix} \bar{x}_J \\ \bar{y}_K \end{bmatrix} = \begin{bmatrix} 0 \\ det(B) \end{bmatrix},$ which implies $\begin{bmatrix} \bar{x}_j \\ \bar{y}_i \end{bmatrix}$ $\begin{bmatrix} \bar{x}_J \\ \bar{y}_K \end{bmatrix} = \pm \begin{bmatrix} x_J \\ y_K \end{bmatrix}$ and $|w^{\mathsf{T}}x| = |w^{\mathsf{T}}_J$ \overline{J} _{*x*}_{*J*} $| = |w$ _{*J*} $|$ $|\bar{x}_J|$ = $|\det(B)| \leq \Delta$.

Suppose now that every circuit $\begin{bmatrix} x \\ y \end{bmatrix}$ of $\begin{bmatrix} A & I \end{bmatrix}$ satisfies $|w^{\dagger}x| \leq \Delta$. Consider any invertible square sub- $\max_{\mathbf{U}} \mathbf{B}$ of $\left[\begin{array}{cc} A & \mathbf{I} \\ w^{\mathsf{T}} & \mathbf{0}^{\mathsf{T}} \end{array}\right]$. We have to show that $|\det(B)| \leq \Delta$. Notice that if $|\det(B)| \geq 2$, then $B = \left[\begin{array}{cc} A_{I,J} & \mathbf{I}_{I,K} \\ w^{\mathsf{T}}_{I,J} & \mathbf{0}^{\mathsf{T}} \end{array}\right]$ $\left[\begin{smallmatrix} A_{I,J} & \mathbf{I}_{I,K} \ w_J^\intercal & \mathbf{0}^\intercal \end{smallmatrix}\right]$ for some *I* ⊆ [*m*], *J* ⊆ [*n*], *K* ⊆ [*m*]. Notice that *K* ⊆ *I* since otherwise *B* has a zero column. By adding every index $i \in \overline{I} = [m] \setminus I$ both to *I* and *K*, we may even assume $B = \begin{bmatrix} A_{[m],J} & I_{[m],K} \\ w^{\text{T}} & 0 \end{bmatrix}$ $\left[\begin{smallmatrix} \bar{x}\ M_J^\top & \mathbf{0}^\top\ \bar{y}_K \end{smallmatrix}\right]$. Pick $\left[\begin{smallmatrix} \bar{x}_J\ \bar{y}_K \end{smallmatrix}\right]$ $\begin{bmatrix} \bar{x}_J \\ \bar{y}_K \end{bmatrix}$ ∈ {0, ±1}^{*J*∪*K*} in the kernel of [$A_{[m],J}$ $\mathbf{I}_{[m],K}$] such that w_J^{\dagger} *J*_{*x*} *z* f *b*. Extend *x* and *y* to *x* ∈ {0,±1}^{*n*} and *y* ∈ {0,±1}^{*m*} by adding zeros, respectively. By Remark [10,](#page-11-1) $\begin{bmatrix} x \\ y \end{bmatrix}$ is a circuit of $\begin{bmatrix} A & I \end{bmatrix}$. As above, we conclude $|\det(B)| = |w^{\top}x| \leq \Delta$. \Box

Next, we recall that circuits generalize edge directions of polyhedra. For instance, every edge direction of the polyhedron $\{x \in \mathbb{R}^n : Ax = b, \ell \le x \le u\}$ is a scalar multiple of a support-minimal vector of ker(*A*), see [\[Onn10,](#page-58-5) Lem. 2.18]. More importantly, we will crucially exploit the fact that nonzero vectors in kernels of totally unimodular matrices can be decomposed into circuits in a conformal way:

Lemma 11 (see [\[NSZ22,](#page-58-3) Lemma 7]). Let $P = \{x \in \mathbb{R}^n : Ax = b, \ell \le x \le u\}$ where $A \in \{0, \pm 1\}^{m \times n}$ is totally $unimodular, b \in \mathbb{R}^m$, and $d = \dim(P)$, $\ell \in \mathbb{Z}^n$, $u \in \mathbb{Z}^n$, and let $x, x' \in P$. In strongly polynomial time, we can find conformal $c_1,...,c_d\in \mathscr{C}(A)$ and $\lambda_1,...,\lambda_d\geq 0$ with $x'=x+\sum_{j=1}^d\lambda_jc_j.$ If x,x' are integer vectors, then *λ*1,...,*λ^d can be chosen to be nonnegative integers.*

Remark 12. If P, x, x', c₁,..., c_d, and λ_1 ,..., λ_d are as above, then for every μ_1 ,..., μ_d with $0 \le \mu_i \le \lambda_i$ for *all i* ∈ [*d*], the vector $x + \sum_{j=1}^d \mu_j c_j$ is also contained in P.

Proof. Let $x'' = x + \sum_{j=1}^{d} \mu_j c_j$. Since c_1, \ldots, c_d are circuits of A, we have $Ax'' = Ax = b$. To see that $\ell \le x \le u$ holds, consider any coordinate index $i \in [n]$. Suppose first that $x_i \leq x_i'$ C_i' . Since c_1, \ldots, c_d are conformal, this implies $(c_j)_i$ ≥ 0 for all $j \in [d]$, and hence

$$
\ell_i \le x_i \le x_i + \sum_{j=1}^d \mu_j(c_j)_i = x''_i \le x_i + \sum_{j=1}^d \lambda_j(c_j)_i = x'_i \le u_i.
$$

The case $x_i \geq x'_i$ $\binom{r}{i}$ follows analogously.

3.2 An Intermediate Result

In general, our proof of the proximity result follows the same framework as the proximity proof by Eisenbrand and Weismantel [\[EW19\]](#page-56-1), extending their results to totally unimodular inequality systems with a constant number of additional equality constraints. A similar proximity framework has recently been applied for the exact weight basis problem in matroids [\[ERW24\]](#page-56-11).

Our proofs of Theorem [4](#page-6-0) and Proposition [5](#page-6-3) are based on the following result.

Proposition 13. *Let* $k, \Delta \in \mathbb{Z}_{\geq 1}$ *be fixed, and let p, A, b, W, d,* ℓ *, u define an instance of Problem [1.](#page-5-2) Given an optimal vertex solution x*[∗] ∈ R *ⁿ of* [\(LP](#page-6-1)1)*, in strongly polynomial time we can compute an integer point* $z ∈ \mathbb{Z}^n$ *with*

- (i) $Az = b, l \leq z \leq u$,
- *(ii)* $||x^* - z||_{\infty} < k$ *, and*
- *(iii) if* [\(IP](#page-5-3)₁) *is feasible, then it has an optimal solution* $z^* = z + \sum_{j=1}^t c_j$ *where* $c_1, ..., c_t$ ∈ $C(A)$ *are conformal and t* $\leq k(2k\Delta+1)^k =: f_{13}(k,\Delta)$ $\leq k(2k\Delta+1)^k =: f_{13}(k,\Delta)$ $\leq k(2k\Delta+1)^k =: f_{13}(k,\Delta)$.

 \Box

Note that Proposition [5](#page-6-3) follows directly from Proposition [13](#page-12-0) by taking $f_5(k,\Delta) = f_{13}(k,\Delta)$. To see how Proposition [13](#page-12-0) implies Theorem [4,](#page-6-0) consider any instance of Problem [1.](#page-5-2) Let *x* [∗] be any optimal solution of (LP_1) (LP_1) , and assume that (IP_1) (IP_1) is feasible. Pick a vertex *x'* of $\{x \in \mathbb{R}^n : Ax = b, Wx \leq d, \lfloor x^* \rfloor \leq x \leq \lceil x^* \rceil \}$ with $p^{\top}x' = p^{\top}x^*$. Here, the vector $\lfloor x^* \rfloor$ arises from x^* by applying $\lfloor \cdot \rfloor$ to each entry of x^* , and $\lceil x^* \rceil$ is defined analogously. Let $I = \{i \in [n] : x'_i\}$ $i \in \mathbb{Z}$. Note that, for the sake of proving Theorem [4,](#page-6-0) we can assume that we know the optimal solution to [\(IP](#page-5-3)₁), hence we may partition *I* into I_{\le} and I_{\ge} such that the optimum value of

$$
\max\left\{p^{\top}x: Ax = b, Wx = d, \ell \le x \le u, x_i \le x_i' \text{ for all } i \in I_{\le}, x_i \ge x_i' \text{ for all } i \in I_{\ge}, x \in \mathbb{Z}^n\right\}
$$
(1)

is equal to (IP_1) . Since x' is an optimal vertex solution of the LP relaxation of (1) , we may apply Proposi-tion [13](#page-12-0) to obtain a point $z \in \mathbb{Z}^n$ with $||x'-z||_{\infty} < k$ such that [\(1\)](#page-13-0) has an optimal solution $z^* = z + \sum_{j=1}^{\tilde{t}} c_j$ where *c*₁,...,*c*_{*t*} ∈ $\mathcal{C}(A)$ and *t* ≤ *f*_{[13](#page-12-0)}(*k*,∆). Note that *z*^{*} is also an optimal solution of [\(IP](#page-5-3)₁) and observe that we have

$$
||x^* - z^*||_{\infty} \le ||x^* - x'||_{\infty} + ||x' - z||_{\infty} + ||z - z^*||_{\infty} \le 1 + k + ||z - z^*||_{\infty} \le 1 + k + \sum_{j=1}^t ||c_j||_{\infty} = 1 + k + t.
$$

The result follows by taking $f_4(k,\Delta) := 1 + k + f_{13}(k,\Delta)$ $f_4(k,\Delta) := 1 + k + f_{13}(k,\Delta)$.

The remainder of this section is devoted to the proof of Proposition [13.](#page-12-0) To this end, let *p*, *A*, *b*, *W* , *d*, ℓ , *u* define an instance of Problem [1](#page-5-2) and denote $P := \{x \in \mathbb{R}^n : Ax = b, \ell \leq x \leq u\}$. Moreover, let x^* be an optimal vertex solution of (LP_1) (LP_1) .

3.3 Proof of Proposition [13](#page-12-0)

Let us first describe how the integer point *z* can be computed, and then show that it satisfies [\(i\)](#page-12-1)[–\(iii\).](#page-12-2) To this end, consider denote the inclusionwise smallest face *F* of *P* that contains x^* . Recall that $Wx = d$ consists of only *k* equations, and hence $\dim(F) \le k$. Notice also that $F = \{x \in \mathbb{R}^n : Ax = b, \ell' \le x \le u'\}$ for some appropriately chosen $\ell', u' \in \mathbb{Z}^n$. Since A is totally unimodular, we can compute a vertex z' of F in strongly polynomial time. As *P* is an integer polytope, so is *F* and hence *z* ′ is integer. Moreover, we can invoke Lemma [11](#page-12-3) to find circuits *c* ′ $\frac{1}{1}, \ldots, c'_{\mathfrak{p}}$ χ'_{k} of *A* and $\lambda_1, ..., \lambda_k \in \mathbb{R}_{\geq 0}$ with $z' = x^* + \sum_{j=1}^{k} \lambda_j c'_j$ *j* . We define

$$
z = x^* + \sum_{j=1}^k (\lambda_j - \lfloor \lambda_j \rfloor) c_j'
$$

and note that, by Remark [12,](#page-12-4) it is contained in *F*. In particular, we see that *z* satisfies [\(i\).](#page-12-1) Note also that *z* is an integer vector since

$$
z = x^* + \sum_{j=1}^k (\lambda_j - \lfloor \lambda_j \rfloor) c'_j = \underbrace{z'}_{\in \mathbb{Z}^n} - \sum_{j=1}^k \underbrace{\lfloor \lambda_j \rfloor}_{\in \mathbb{Z}} \underbrace{c'_j}_{\in \mathbb{Z}^n}.
$$

Moreover, it satisfies [\(ii\)](#page-12-5) since

$$
\|x^*-z\|_{\infty} = \left\|\sum_{j=1}^k (\lambda_j - \lambda_j) c'_j\right\|_{\infty} \le \sum_{j=1}^k (\lambda_j - \lambda_j) \|c'_j\|_{\infty} < \sum_{j=1}^k \|c'_j\|_{\infty} \le k.
$$

Thus, it remains to show [\(iii\).](#page-12-2) To this end, we first observe that since *F* is the minimal face of *P* containing *x* ∗ , we have

$$
z + v \in P \implies x^* + \varepsilon v \in P \text{ for some } \varepsilon > 0 \tag{2}
$$

for every vector $v \in \mathbb{R}^n$.

Now, take any optimal solution *z* [∗] of the integer program, and decompose

$$
z^*-z=c_1+\cdots+c_t
$$

where $c_1, \ldots, c_t \in \mathcal{C}(A)$ are conformal (such a decomposition exists due to Lemma [11\)](#page-12-3), and choose z^* and this decomposition such that *t* is smallest possible. Recall that for every $I \subseteq [t]$, the vector $z + \sum_{i \in I} c_i$ is also contained in *P*.

3.3.1 Reordering Circuits

To obtain a bound on the number *t* of circuit vectors, let us first reorder c_1, \ldots, c_t such that all partial sums $\sum_{i=1}^{s} W c_i$ have small norm. For this purpose, we make use of the Steinitz lemma [\[Ste13\]](#page-59-3):

Lemma 14 (see [\[Sev78,](#page-58-7) [SB97\]](#page-58-8)). Let $\|\cdot\|$ be any norm on \mathbb{R}^k . If $y_1,\ldots,y_{t'}\in \mathbb{R}^k$ satisfy $y_1+\cdots+y_{t'}=\mathbf{0}$, then *there exists a permutation π on* [*t* ′] *such that*

$$
\left\|\sum_{i=1}^s y_{\pi(i)}\right\| \leq k \cdot \max\{\|y_1\|, \dots, \|y_t\|\}
$$

holds for all $s \in [t']$ *.*

To apply the above lemma, we first set $y_i = W c_i$ for $i = 1, ..., t$. Recall that we consider an instance of Problem [1,](#page-5-2) and hence

$$
||Wc_i||_{\infty} \leq \Delta
$$

holds for all $i \in [t]$. Moreover, we have

$$
||Wc_1 + \dots + Wc_t||_{\infty} = ||W(z^* - z)||_{\infty} \le ||W(z^* - x^*)||_{\infty} + ||W(x^* - z)||_{\infty} = ||W(x^* - z)||_{\infty} \le k\Delta,
$$

where the equality holds since we have $Wz^* = d$ and $Wx^* = d$. This means that we can find integer vectors $y_{t+1},...,y_{t+k} \in \mathbb{Z}^k$ such that $y_{t+1} + \cdots + y_{t+k} = -(Wc_1 + \cdots + Wc_t)$ and $||y_{t+j}||_{\infty} \leq \Delta$ holds for all *j* ∈ [*k*]. By construction, we have $y_1 + \cdots + y_{t+k} = 0$ and $||y_i||_{\infty} \leq \Delta$ for all *i* ∈ [*t* + *k*]. Thus, by Lemma [14](#page-14-0) there exists a permutation π on $[t + k]$ such that

$$
\left\| \sum_{i=1}^{s} y_{\pi(i)} \right\| \le k\Delta \tag{3}
$$

holds for all $s \in [t+k]$.

We now claim that

$$
t \le k(2k\Delta + 1)^k \tag{4}
$$

holds, which yields [\(iii\).](#page-12-2) For the sake of contradiction, suppose that the claim does *not* hold. Since there are (2*k*∆ + 1)*^k* integer points of infinity norm at most *k*∆ in Z *k* , we can apply the pigeonhole principle and [\(3\)](#page-14-1) to obtain $k+1$ indices $1 \le s_1 < s_2, \dots < s_{k+1} \le t+k$ such that $\sum_{i=1}^{s_1} y_{\pi(i)} = \sum_{i=1}^{s_j} y_{\pi(i)}$ for all $j \in [k+1]$. Thus, the (nonempty) sets

$$
S_i := \{ \pi(s_i + 1), \dots, \pi(s_{i+1}) \} \text{ for } i \in [k],
$$

and

$$
S_{k+1} := \{\pi(1), \ldots, \pi(s_1)\} \cup \{\pi(s_{k+1} + 1), \ldots, \pi(t + k)\}\
$$

yield $k + 1$ disjoint partial sums, summing up to zero each. By the pigenohole principle, one of the S_i does not contain any $j \in \{t+1,\ldots,t+k\}$, say *S*'. Setting $\bar{c} = \sum_{i \in S'} c_i$, we have $W\bar{c} = \mathbf{0}$.

3.3.2 An Optimal Solution with a Shorter Decomposition

Consider the vector $z' = z^* - \bar{c} = z + \sum_{i \in [t] \setminus S'} c_i \in P$. Since $Wz' = Wz^* - W\bar{c} = d$, we see that z' is feasible for the integer program. We claim that

$$
p^{\top} \bar{c} \le 0 \tag{5}
$$

holds. Note that the latter claim implies $p^{\top}z' = p^{\top}z^* - p^{\top}\bar{c} \geq p^{\top}z^*$. This shows that z' is also an optimal solution for the integer program with a shorter decomposition, a contradiction. Thus, our assumption (that *t* does not satisfy the bound in [\(4\)](#page-14-2)) was wrong and we are done.

To show [\(5\)](#page-15-1), note that $z + \bar{c} = z + \sum_{i \in I} c_i \in P$. Thus, by [\(2\)](#page-13-1) there exists some $\varepsilon > 0$ such that $x^* + \varepsilon \bar{c} \in P$. Moreover, since $W(x^* + \varepsilon \bar{c}) = Wx^* = d$ we see that $x^* + \varepsilon \bar{c}$ is a feasible solution for the LP relaxation. By the optimality of x^* , this implies $p^T \bar{c} \leq 0$.

4 Dealing with 1**-sums and** 2**-sums**

In Section [4.1,](#page-15-2) we define 1-sums and 2-sums for vector configurations. Applied to the vector configurations arising from the columns of a TU matrix, we recover the standard notions of 1-sum and 2-sum of TU matrices (see for instance [\[Sch98\]](#page-58-9)), and regular matroids (see for instance [\[Oxl06\]](#page-58-10)). In Section [4.2,](#page-17-0) we state the actual version of Problem [1](#page-5-2) that we solve. Next, in Section [4.3,](#page-18-0) we explain how to reduce Problem [1](#page-5-2) to the 2-connected case. Finally, in Section [4.4,](#page-19-0) we explain how to reduce to the almost 3-connected case. We point out that our reductions are all black-box.

4.1 Vector configurations

We regard matrices $A \in \mathbb{R}^{m \times n}$ as *(vector) configurations* in \mathbb{R}^m . In other words, we see $A \in \mathbb{R}^{m \times n}$ as an ordered multiset of *n* vectors $\{a_1, \ldots, a_n\}$ in \mathbb{R}^m , namely, the *n* columns of *A*. The *size* of a configuration is defined as its number of vectors, taking into account multiplicities. Slightly abusing notation, we write *v* ∈ *A* to mean that *v* is a column of *A*. For *v* ∈ *A*, we let *A* − *v* denote the configuration obtained from *A* by deleting the column for *v*, hence decreasing the multiplicity of *v* by 1. If $A_1 \in \mathbb{R}^{m \times n_1}$ and $A_2 \in \mathbb{R}^{m \times n_2}$ are configurations in \mathbb{R}^m , we let A_1 ⊎ A_2 := $\begin{bmatrix} A_1 & A_2 \end{bmatrix}$ denote the configuration obtained by concatenating the two configurations. Below, we let $\text{colsp}(A) := \text{span}(\{a_1, \ldots, a_n\})$ denote the column space of A.

We point out that configurations are, basically, representations of real matroids. Most of the notions we discuss here such as connectivity, 1-sum and 2-sum are the counterparts of standard matroid notions. The reader who has some knowledge on matroids should "feel at home".

Definition 15 (regular configuration)**.** *We call a configuration A* ∈ R *m*×*n* regular *whenever every circuit of A is in* {0,±1}*ⁿ . (This terminology is inspired by Tutte's notion of a regular subspace, see [\[Tut65\]](#page-59-4).)*

Every TU matrix *A* ∈ R *m*×*n* , seen as a vector configuration, is a regular configuration. Conversely, if $A \in \mathbb{R}^{m \times n}$ is a regular configuration then there exists a TU matrix $A' \in \{0, \pm 1\}^{m \times n}$ such that ker $A = \text{ker } A'$. Assume that *A* is a regular configuration. We say that a circuit $c \in \mathcal{C}(A)$ *uses* a vector $v \in A$ if $c(v) \neq 0$. Circuit *c* uses *v* positively if $c(v) = 1$, and *negatively* if $c(v) = -1$.

We recall the definition of connectivity given in the introduction.

Definition 16 (separation, connectivity). *A configuration A* $\in \mathbb{R}^{m \times n}$ *has a q*-separation *(for q* $\in \mathbb{Z}_{\geq 1}$ *) if it can be written, possibly after permuting its columns, as* $A = \begin{bmatrix} A_1 & A_2 \end{bmatrix}$ *, with* $dim(colsp(A_1) \cap colsp(A_2)) \le$ *q* −1 *where both A*¹ *and A*² *have at least q columns. The* connectivity *of A is defined as the smallest order of a separation of A. We say that A is q*-connected *if its connectivity is at least q. A configuration A is said to be* almost 3-connected *if it becomes* 3*-connected when one deletes from A every column that is a nonzero multiple of another column.*

Figure 4: An example of 2-sum of incidence matrices, represented as directed graphs. In order to fit Definition [18,](#page-16-0) *A*¹ should be the incidence matrix of *G*¹ with four extra zero rows for the vertices in $V(G_2) \setminus V(G_1)$, and A_2 should be defined similarly, so that A_1 and A_2 have as many rows as the number of vertices of *G* and share a column *v* corresponding to arc (i, j) . Then it is easy to check that $A_1 \oplus_2 A_2$ is the incidence matrix of *G*.

Next, for *q* = 1,2, we relate these definitions to the notion of a *q*-sum.

Definition 17 (1-sum). Let A_1 ∈ $\mathbb{R}^{m \times n_1}$ and A_2 ∈ $\mathbb{R}^{m \times n_2}$ be configurations such that $\text{colsp}(A_1) \cap \text{colsp}(A_2)$ = {**0***}. The* 1-sum *of* A_1 *and* A_2 *is the configuration* $A_1 \oplus_1 A_2 := A_1 \oplus A_2 \in \mathbb{R}^{m \times (n_1 + n_2)}$ *.*

The circuits of a 1-sum can be characterized as follows:

$$
\mathscr{C}(A_1\oplus_1 A_2)=\left\{\begin{bmatrix}c_1\\ \mathbf{0}\end{bmatrix}:c_1\in\mathscr{C}(A_1)\right\}\cup\left\{\begin{bmatrix}\mathbf{0}\\ c_2\end{bmatrix}:c_2\in\mathscr{C}(A_2)\right\}.
$$

A configuration *A* of size at least 2, that contains the zero vector is never 2-connected. We say that two vectors $v, v' \in \mathbb{R}^m$ are *parallel* if one is a nonzero multiple of the other. Hence the zero vector is only parallel to itself.

Definition 18 (2-sum)**.** *Let A*¹ ∈ R *^m*×(*n*1+1) *and A*² [∈] ^R *^m*×(*n*2+1) *be configurations with a common nonzero vector v, and such that* $\text{colsp}(A_1) \cap \text{colsp}(A_2) = \text{span}(\{v\})$ *. The* 2-sum *of* A_1 *and* A_2 *is the configuration A*₁ ⊕₂ *A*₂ := $(A_1 - v) \cup (A_2 - v) \in \mathbb{R}^{m \times (n_1 + n_2)}$.

Assuming that configurations A_1 and A_2 are regular, and that the common vector v is the last vector of *A*¹ and the first vector of *A*2, we have the following characterization of the circuits of a 2-sum:

$$
\mathcal{C}(A_1 \oplus_2 A_2) = \left\{ \begin{bmatrix} c_1 \\ \mathbf{0} \end{bmatrix} : \begin{bmatrix} c_1 \\ 0 \end{bmatrix} \in \mathcal{C}(A_1) \right\} \cup \left\{ \begin{bmatrix} \mathbf{0} \\ c_2 \end{bmatrix} : \begin{bmatrix} 0 \\ c_2 \end{bmatrix} \in \mathcal{C}(A_2) \right\}
$$

$$
\cup \left\{ \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} : \begin{bmatrix} c_1 \\ \pm 1 \end{bmatrix} \in \mathcal{C}(A_1), \begin{bmatrix} \mp 1 \\ c_2 \end{bmatrix} \in \mathcal{C}(A_2) \right\}.
$$

In other words, each circuit of $A_1 \oplus_2 A_2$ is either a circuit of A_i that does not use the common vector *v*, for some $i \in [2]$, or obtained by combining a circuit of A_i that uses *v* positively together with a circuit of *A*3−*ⁱ* that uses *v* negatively, for some *i* ∈ [2]. We say that the circuits of the second type *cross v*.

We call *circulation* of *A* any vector $x \in \mathbb{R}^n$ such that $Ax = 0$, that is, any vector in ker(*A*). The above characterization of the circuits of a 2-sum naturally extends to circulations: $\genfrac{[}{]}{0pt}{}{x_1}{x_2}$ is a circulation of A_1 \oplus_2 *A*₂ if and only if there exists a scalar φ such that $\begin{bmatrix} x_1 \\ -\varphi \end{bmatrix}$ is a circulation of A_1 and $\begin{bmatrix} \varphi \\ x_2 \end{bmatrix}$ is a circulation of *A*₂. We say that $\left[\frac{x_1}{-\varphi}\right]$ and $\left[\frac{\varphi}{x_2}\right]$ are obtained by *splitting* $\left[\frac{x_1}{x_2}\right]$. If $\left[\frac{x_1}{x_2}\right]$ is an integer circulation splitting into $\left[\begin{array}{c} x_1 \\ -\varphi \end{array}\right]$ and $\left[\begin{array}{c} \varphi \\ x_2 \end{array}\right]$, then it can be written as a conformal sum of circuits in $\mathcal{C}(A_1 \oplus_2 A_2)$ with exactly $|\varphi|$ circuits crossing *v*.

Notice that a configuration is 2-connected if and only if it cannot be written as the 1-sum of two smaller configurations, possibly changing the ordering of its vectors, and 3-connected if and only if it cannot be written as the 2-sum of two configurations (possibly changing the ordering of its vectors), each with at least 3 vectors.

Definition 19 (decomposition tree)**.** *A tree T is a* decomposition tree *of a* 2*-connected configuration A* ∈ R *^m*×*ⁿ of size at least* 3 *if it satisfies the following conditions:*

- *(i) each node t of T has a corresponding configuration* $A(t)$ *in* \mathbb{R}^m *of size at least 3,*
- (*ii*) each edge tt' of T has a corresponding nonzero vector^{[7](#page-17-1)} $\overline{v} = \overline{v}(t t') \in \mathbb{R}^m$ that belongs to both A(*t*) and $A(t')$ *, and such that* $\text{colsp}(A(t)) \cap \text{colsp}(A(t')) = \text{span}(\{\overline{v}\})$ *,*
- (*iii*) *if t and t' are distinct nodes of T then* dim(colsp($A(t)$)∩colsp($A(t')$)) ≤ 1*, and if t'' is a node of the t*−*t' path in T then* $\text{colsp}(A(t)) \cap \text{colsp}(A(t')) \subseteq \text{colsp}(A(t''))$ *,*
- *(iv) the configuration obtained by iteratively performing (in any order) the* 2*-sum corresponding to each edge of T , and then possibly reordering the vectors equals A.*

It follows directly from [\[CE80,](#page-55-9) [Sey81\]](#page-58-11) that every 2-connected configuration of at least 3 vectors admits a decomposition tree in which $A(t)$ is 3-connected for each node $t \in V(T)$. Moreover, if *A* is regular (in particular, when *A* is a TU matrix) then we may choose the configurations *A*(*t*) in such a way that *A*(*t*) is regular for every node $t \in V(T)$. We will always assume this.

4.2 The maximum constrained integer circulation problem

Consider any (feasible) instance of Problem [1.](#page-5-2) Recall that by Proposition [5,](#page-6-3) we may reduce to the case where $b = 0$ by performing a translation. We now define, for better reference, a new problem which is essentially the restriction of Problem [1](#page-5-2) to the instances that have $b = 0$. This is the version will need in our reduction.

Problem 2 (Maximum constrained integer circulation problem (MCICP)). *Let* $k, \Delta \in \mathbb{Z}_{\geq 1}$ *be constants. Given p* $\in \mathbb{Z}^n$, regular configuration $A \in \{0, \pm 1\}^{m \times n}$, $W \in \mathbb{Z}^{k \times n}$ such that $\|Wc\|_{\infty} \leq \Delta$ for all circuits $c \in \mathbb{Z}^n$ $\mathscr{C}(A)$, $d \in \mathbb{Z}^k$ and $\ell, u \in \mathbb{Z}^n$ such that $\ell \leq u$, find

$$
\max \{ p^{T} x : Ax = 0, Wx = d, \ell \le x \le u, x \in \mathbb{Z}^{n} \}.
$$
 (IP₂)

Letting $I := (p, A, W, d, \ell, u)$ *, we define* OPT(*I*) *as the optimum value of* [\(IP](#page-17-2)₂)*. If I is infeasible, then* $OPT(I) = -\infty$.

Here, we call a circulation $x \in \text{ker}(A)$ to be *feasible* if $\ell \leq x \leq u$. This terminology is meant to generalize to any regular configuration the usual notion of a circulation in a directed graph, which arises when *A* is an incidence matrix.

⁷The vector $\overline{v} = \overline{v}(tt')$ is considered as being *private* to $A(t)$ and $A(t')$. Formally, we would need to use labels for the vectors of the configurations under consideration in order to distinguish the different copies of the same vector. Then, \bar{v} could possibly appear as a vector in $A(t'')$ for some node t'' distinct from t and t' , but not with the same label. For the sake of simplicity, we will however not do this explicitly.

Figure 5: An example of a 2-sum decomposition of an incidence matrix into four matrices, represented as directed graphs as in Figure [4,](#page-16-1) with corresponding decomposition tree *T*. Arcs with the same color represent the common vector in the span of two incidence matrices.

Recall that Proposition [5](#page-6-3) implies the following: [\(IP](#page-17-2)₂) either is infeasible, or has an optimal solution *x* that is the conformal sum of at most $f_5(k,\Delta)$ $f_5(k,\Delta)$ $f_5(k,\Delta)$ circuits of *A*. Also, in the context of Proposition [5,](#page-6-3) we may assume that there is no *feasible c* $\in \mathcal{C}(A)$ such that $Wc = \mathbf{0}$ and $p^{\dagger}c > 0$. We will also always assume this.

Before discussing 1-sums and 2-sums further, we establish some useful notation and terminology relative to the MCICP. For $x \in \mathbb{R}^n$, it will be convenient to denote by $x(v)$ the coordinate of *x* corresponding to vector *v* ∈ *A*. For instance, using this notation, the equation $Ax = 0$ becomes $\sum_{v \in A} x(v) \cdot v = 0$. We will refer to $x(v)$ as the *flow* along vector v .

In the next section we show how to reduce any MCICP instance to a constant number MCICP instances that are 2-connected in the sense that their corresponding vector configuration is 2-connected. The reduction to the almost 3-connected case is more involved, and is discussed in the section after the next.

4.3 Reduction to the 2**-connected case**

Consider any MCICP instance $I = (p, A, W, d, \ell, u)$ that is not 2-connected. After permuting its *n* columns, *A* decomposes as a 1-sum $A = A_1 \oplus_1 \cdots \oplus_1 A_q$ of 2-connected configurations, where $2 \le q \le n$. We extend the decomposition of *A* to a corresponding decomposition of vectors *p*, *ℓ*, *u* and matrix *W* . For $j \in \{1, \ldots, q\}$, we write

$$
p_{\leq j} := \begin{bmatrix} p_1 \\ \vdots \\ p_j \end{bmatrix}, \quad W_{\leq j} := \begin{bmatrix} W_1 & \cdots & W_j \end{bmatrix}, \quad \ell_{\leq j} := \begin{bmatrix} \ell_1 \\ \vdots \\ \ell_j \end{bmatrix} \quad \text{and} \quad u_{\leq j} := \begin{bmatrix} u_1 \\ \vdots \\ u_j \end{bmatrix}.
$$

We use a straightforward dynamic program (DP) to solve [\(IP](#page-17-2)2). Let *B* := [−∆*f*[5](#page-6-3)(*k*,∆),∆*f*[5](#page-6-3)(*k*,∆)]*^k* ∩Z *k* be an integer box containing all target vectors that can be attained by solutions that are sums of at most *f*_{[5](#page-6-3)}(k , Δ) circuits. For each *j* ∈ [*q*] and $d_{\leq j}$ ∈ *B* we let

$$
F_j(d_{\leq j}) := \text{OPT}(p_{\leq j}, A_1 \oplus_1 \cdots \oplus_1 A_j, W_{\leq j}, d_{\leq j}, \ell_{\leq j}, u_{\leq j}).
$$

These optimum values can be easily computed through the following DP equation, whose proof is left to the reader.

Observation 20. *For all j* > 1 *and* $d ≤ j ∈ B$ *we have*

$$
F_j(d_{\leq j}) = \max\left\{F_{j-1}(d_{\leq j-1}) + \text{OPT}(p_j, A_j, W_j, d_j, \ell_j, u_j) : d_{\leq j-1} \in B, d_j \in B, d_{\leq j-1} + d_j = d_{\leq j}\right\}.
$$

Since OPT(*I*) = $F_q(d)$, and since it is straightforward to extend the DP approach to also recover corresponding optimal solutions, Observation [20](#page-19-1) implies the following result.

Theorem 21. *Let k*,∆ ∈ Z≥¹ *be constants. Any instance of the MCICP on a matrix with n columns can be solved after solving at most* (2∆*f*[5](#page-6-3)(*k*,∆)+1)*kn instances of the MCICP that are* 2*-connected and on matrices with at most n columns, and performing extra work in time* $(\Delta f_5(k,\Delta))^{O(k)}n$ $(\Delta f_5(k,\Delta))^{O(k)}n$ $(\Delta f_5(k,\Delta))^{O(k)}n$.

4.4 Reduction to the almost 3**-connected case**

In this section we consider a variant of Problem [2,](#page-17-3) which we call the *rooted* MCICP. This problem is defined exactly as Problem [2,](#page-17-3) with two additional input data: a *root vector* $\overline{v} \in A$ and a prescribed flow value $\varphi \in \mathbb{Z}$ for $\overline{\nu}$. We let $\ell(\overline{\nu}) = u(\overline{\nu}) := \varphi$, which forces $x(\overline{\nu}) = \varphi$ for all feasible circulations *x*.

It is easy to convert any MCICP instance into an equivalent rooted MCICP instance. For example, we may duplicate any column of *A*, name the new column $\overline{\nu}$ and let $\varphi := 0$. On the other hand, the rooted MCICP is a just special case of MCICP, hence the two problems are actually equivalent. For convenience, we focus on the rooted MCICP throughout this section.

Consider a rooted MCICP instance $I = (p, A, W, d, \ell, u, \overline{v}, \varphi)$, and consider a decomposition tree *T* for *A* such that $A(t)$ is regular and 3-connected for all nodes $t \in V(T)$. There exists a unique *root node r* ∈ *V*(*T*) such that \overline{v} is a column of *A*(*r*). This turns *T* into a rooted tree.

As in the previous section, let $B:=[-\Delta f_5(k,\Delta),\Delta f_5(k,\Delta)]^k\cap \mathbb{Z}^k$ $B:=[-\Delta f_5(k,\Delta),\Delta f_5(k,\Delta)]^k\cap \mathbb{Z}^k$ $B:=[-\Delta f_5(k,\Delta),\Delta f_5(k,\Delta)]^k\cap \mathbb{Z}^k$. Also, let $\Phi:=[-f_5(k,\Delta),f_5(k,\Delta)]\cap \mathbb{Z}$.

Definition 22 (rooted MCICP instances for subtrees). *For t* \in *V*(*T*)*, we let A_t denote the configuration obtained by performing all the* 2*-sums corresponding to the edges of the subtree of T rooted at t. Let* $\overline{v}_t \in A_t$ *denote the vector associated to the edge between t and its parent. If t is the root, then we let* $\overline{v}_t := \overline{v}$ *. For* $d_t \in B$ and $\varphi_t \in \Phi$ we let $I_t = I_t(d_t, \varphi_t)$ denote the rooted MCICP instance $I_t := (p_t, A_t, W_t, d_t, \ell_t, u_t, \overline{v}_t, \varphi_t)$ where p_t , W_t , ℓ_t and u_t are found by restricting the corresponding vectors in the original instance I to the columns of A_t and letting $p_t(\overline{v}_t) := 0$ and $\ell_t(\overline{v}_t) = u_t(\overline{v}_t) := \varphi_t$. We define $W_t(\cdot, \overline{v}_t)$ in such a way that $||W_t c_t||_\infty$ ≤ Δ *for all circuits c_t of A_t. One way to achieve this is to let* $W_t(i, \overline{v}_t)$ *be the "upper weight" of some circuit of A crossing* \overline{v}_t *with respect to* $W(i, \cdot)$ *, for each i* \in [*k*]*, see Lemma [23](#page-19-2) below. We call I_t a* rooted *instance at node t. We let*

$$
F_t(d_t, \varphi_t) := \text{OPT}(I_t) = \text{OPT}(p_t, A_t, W_t, d_t, \ell_t, u_t, \overline{v}_t, \varphi_t)
$$

denote the optimum value of the local MCICP instance corresponding to node $t \in V(T)$ *and to the choices of target vector* $d_t \in B$ *and prescribed flow value* $\varphi_t \in \Phi$ *for the root vector.*

Lemma 23. There exists integers $W_t(i,\overline{v}_t)$ for each $i \in [k]$ such that $||W_t c_t||_\infty \leq \Delta$ for all circuits c_t of A_t .

Proof. If *t* is the root, we simply let $W_t(i, \overline{v}_t) = W_t(i, \overline{v}) := 0$ for all $i \in [k]$. From now on, assume that *t* is not the root. Let t' denote the parent of t . Consider the configuration A^t obtained by performing each 2-sum corresponding to the edges of $T - tt'$ that are *not* in the subtree rooted at *t*. We get the decomposition $A = A^t \oplus_2 A_t$. Let $\overline{c} \in \mathscr{C}(A)$ be any circuit crossing $\overline{\nu}_t$. By negating \overline{c} if necessary, we may assume that $\overline{c} = \begin{bmatrix} \overline{c}_1 \\ \overline{c}_2 \end{bmatrix}$ *c*2 $\left[\begin{array}{c} \overline{c}_1 \\ -1 \end{array} \right] \in \mathscr{C}(A^t)$ and $\left[\frac{1}{c_2} \right] \in \mathscr{C}(A_t)$. For each *i* ∈ [*k*], we let

$$
W_t(i,\overline{\nu}_t):=\sum_{v\in A^t\atop v\neq \overline{\nu}_t}W(i,\nu)\overline{c}(v)
$$

denote the *upper weight* of *c* with respect to the *i*-th row of *W* .

Now consider any circuit c_t of A_t . If $c_t(\overline{v}_t) = 0$ then c_t yields a circuit of A , say $c \in \mathcal{C}(A)$, such that $||W_t c_t||_{\infty} = ||Wc||_{\infty} \leq \Delta$. Otherwise, we may assume that $c_t(\overline{v}_t) = 1$. Then c_t combines with $\begin{bmatrix} \overline{c}_1 \\ -1 \end{bmatrix}$ into a circuit of *A*, say $c' \in \mathcal{C}(A)$. By our choice of $W_t(\cdot, \overline{v}_t)$, we get $||W_t c_t||_{\infty} = ||Wc'||_{\infty} \leq \Delta$. \Box

If *t* is a leaf of *T* then we can compute $F_t(d_t, \varphi_t)$ for every fixed choices of $d_t \in B$ and $\varphi_t \in \Phi$ by solving a single 3-connected MCICP instance. Now consider a node *t* that is not a leaf, let t_1, \ldots, t_q denote the children of *t*. Hence, $\overline{v}_{t_1},...,\overline{v}_{t_q}$ are the vectors of $A(t)$ respectively associated to the edges $tt_1,...,tt_q.$

Consider a choice of $d_t^0, d_{t_1},..., d_{t_q} \in B$ and $\varphi_{t_1},..., \varphi_{t_q} \in \Phi$. Let $I_t^0 = I_t^0(d_t^0, \varphi_t, -\varphi_{t_1},..., -\varphi_{t_q})$ denote the rooted MCICP instance obtained from $I_t = I_t(d_t, \varphi_t)$ by restricting to the sole vectors of $A(t)$, redefining the target vector to d_t^0 and setting both bounds for each \overline{v}_{t_i} to − φ_{t_i} for $i \in [q]$. Moreover, we define the weight $W_t^0(i, \overline{\nu}_{t_j}) \in \mathbb{Z}$ for each $i \in [k]$ and $j \in [q]$ in such a way that $||W_t^0 c_t^0||_{\infty} \leq \Delta$ for all $c_t^0 \in \mathcal{C}(A(t))$. This is done by using the "lower weight" of any circuit crossing \overline{v}_{t_i} with respect to $W(i, \cdot)$, similarly as in the proof of Lemma [23.](#page-19-2) We call I_t^0 a *local* instance at node t . Next, let $F_t^0(d_t^0, \varphi_t, -\varphi_{t_1}, \ldots, -\varphi_{t_q})$ denote the optimum value of instance I^0_t . The following result generalizing Observation [20](#page-19-1) can be used to compute $F_t(d_t, \varphi_t)$ when the number of children of *t* is small (say, logarithmic in *n*).

Lemma 24. *With the above notation, we have*

$$
F_t(d_t, \varphi_t) = \max \left\{ F_t^0(d_t^0, \varphi_t, -\varphi_{t_1}, \dots, -\varphi_{t_q}) + \sum_{i=1}^q F_{t_i}(d_{t_i}, \varphi_{t_i}) : d_t^0, d_{t_1}, \dots, d_{t_q} \in B, \varphi_{t_1}, \dots, \varphi_{t_q} \in \Phi, \tag{6}
$$
\n
$$
d_t^0 + \sum_{i=1}^q \left(d_{t_i} - \left(W_t(\cdot, \overline{\nu}_{t_i}) - W_t^0(\cdot, \overline{\nu}_{t_i}) \right) \varphi_{t_i} \right) = d_t \right\}
$$

for every $d_t \in B$ *and* $\varphi_t \in \Phi$ *.*

Proof. Let x_t denote an optimum solution of $I_t = I_t(d_t, \varphi_t)$. Consider a conformal decomposition $x_t =$ $c_{t,1} + \cdots + c_{t,\ell}$ as a sum of $\ell \le f_5(k,\Delta)$ $\ell \le f_5(k,\Delta)$ $\ell \le f_5(k,\Delta)$ circuits of A_t . We split x_t into circulations x_t^0 in $A(t)$ and x_{t_1}, \ldots , x_{t_q} in A_{t_1}, \ldots, A_{t_q} respectively. For $i \in [q]$, let $\varphi_{t_i} := x_{t_i}(\overline{v}_{t_i}) = -x_t^0(\overline{v}_{t_i})$. Hence, $|\varphi_{t_i}| \leq \ell \leq f_5(k, \Delta)$ $|\varphi_{t_i}| \leq \ell \leq f_5(k, \Delta)$ $|\varphi_{t_i}| \leq \ell \leq f_5(k, \Delta)$ which implies $\varphi_{t_i}\in\Phi$. Next, let d_t^0 and d_{t_1},\ldots,d_{t_q} denote the weight vectors of x_t^0 and x_{t_1},\ldots,x_{t_q} respectively. That is, let $d_t^0 := W_t^0 x_t^0$ and $d_{t_i} := W_{t_i} x_{t_i}$ for $i \in [q]$ and $j \in [k]$. By our choice of $W_t^0(\cdot, \overline{v}_{t_i})$ and $W_{t_i}(\cdot, \overline{v}_{t_i})$ for $i \in [q]$, all these weight vectors are in *B*. The sum $d_t^0 + d_{t_1} + \cdots + d_{t_q}$ and the weight vector of x_t are related as follows:

$$
d_t^0 + d_{t_1} + \cdots + d_{t_q} = W_t x_t + \sum_{i=1}^q (W_t(\cdot, \overline{v}_{t_i}) - W_t^0(\cdot, \overline{v}_{t_i})) \varphi_{t_i}.
$$

This shows that, in [\(6\)](#page-16-2), the left hand side is at most the right hand side.

In order to show the converse inequality, fix some valid choice of $d_t^0, d_{t_1}, \ldots, d_{t_q} \in B$ and $\varphi_{t_1}, \ldots, \varphi_{t_q} \in B$ Φ . Let x_t^0 be an optimal solution for $I_t^0(d_t^0, \varphi_t, -\varphi_{t_1}, \dots, -\varphi_{t_q})$ and, for $i \in [q]$, let x_{t_i} be an optimal solution for $I_{t_i}(d_{t_i},\varphi_{t_i})$. By what precedes, these optimal solutions combine into a feasible solution x_t for $I_t(d_t, \varphi_t)$, proving that the left hand side of [\(6\)](#page-16-2) is at least the right hand side. \Box

In case *q* is large (super-logarithmic in *n*), Lemma [24](#page-20-0) cannot be used as is since the number of cases to consider might be super-polynomial. For this reason, we have to delve deeper in the properties of the rooted MCICP instances arising from node *t* and its children t_1, \ldots, t_d .

First, remark that if the weight matrix W_{t_i} is zero for some child t_i , then the corresponding rooted MCICP instances $I_{t_i} = I_{t_i}(d_{t_i}, \varphi_{t_i})$ are easy to solve since either $d_{t_i} = \mathbf{0}$ and the k complicating constraints $W_{t_i}x_{t_i} = d_{t_i}$ are trivially satisfied, or the instance is trivially infeasible. This holds more generally when t_i is "tame" in the following sense.

Definition 25 (tame and wild children)**.** *We say that a child tⁱ of t is* tame *if for all circuits c*,*c* ′ *of Atⁱ such* that $c(\overline{v}_{t_i}) = c'(\overline{v}_{t_i}) = 1$, we have $W_{t_i}c = W_{t_i}c'$. If t_i is not tame, we say that it is wild.

In case t_i is a tame child, as we show below, we can basically get rid of the whole subtree rooted at t_i and modify the configuration $A(t)$ by adding a constant number of copies of \overline{v}_{t_i} with carefully defined profits, see Lemma [31.](#page-22-0) This directly leads to an improvement in Lemma [24,](#page-20-0) since we no longer have to guess φ_{t_i} when t_i is a tame child of t . A further crucial result that we prove below is that every circuit $c_t \in \mathcal{C}(A_t)$ crosses a bounded number of vectors \overline{v}_{t_i} such that t_i is wild, see Lemma [32.](#page-23-0) Combining these two lemmas, we obtain an efficient algorithm for computing the optimal values $F_t(d_t, \varphi_t)$.

Tame children and gadgets

We start with a discussion of weight vectors of tame nodes, and follow this with a discussion of gadgets. In order to do this, we need further notions.

Definition 26 (linear equivalence, standardized configuration)**.** *Two configurations A* ∈ R *^m*×*ⁿ and A*′ ∈ R *^m*′×*ⁿ are said to be* linearly equivalent *if* ker(*A*) ⁼ ker(*^A* ′)*, or equivalently if there exists a bijective linear* $map \alpha : \text{colsp}(A) \rightarrow \text{colsp}(A')$ *that maps the j th column of A to the j th column of A' for each j* ∈ [*n*]*. Let* $A \in \mathbb{R}^{m \times n}$ *be a rank-r configuration. We say that A is* standardized *if* $AP = \begin{bmatrix} D & \mathbf{I}_r \end{bmatrix}$ *for some* $D \in \mathbb{R}^{r \times (n-r)}$ *, and some permutation matrix P* \in {0, 1}^{*n*×*n*}.

It is easy to see that every configuration is linearly equivalent to a standardized configuration. This can be achieved by pivoting and deleting zero rows (see for instance [\[Oxl06\]](#page-58-10)).

Definition 27 (dual configuration, cocircuit). Let $A = \begin{bmatrix} D & I_r \end{bmatrix} \in \mathbb{R}^{m \times n}$ denote a standardized rank-r con*figuration. The dual configuration of* $A = \begin{bmatrix} D & I_r \end{bmatrix}$ *is defined as* $\begin{bmatrix} I_{n-r} & -D^\top \end{bmatrix}$ *. It is known that every configuration has the same connectivity as its dual configuration. We call* cocircuit *of A every circuit of the dual* $\text{configuration} \begin{bmatrix} \mathbf{I}_{n-r} & -D^{\mathsf{T}} \end{bmatrix}$ and let

$$
\mathcal{C}^*(A) = \mathcal{C}^*\left(\begin{bmatrix} D & \mathbf{I}_r \end{bmatrix}\right) := \mathcal{C}\left(\begin{bmatrix} \mathbf{I}_{n-r} & -D^\top \end{bmatrix}\right)
$$

denote the set of cocircuits of A.

It is easy to check that every circuit of *A* is orthogonal to every cocircuit of *A*.

Now consider a regular configuration $A \in \mathbb{R}^{m \times n}$. If $A = [D \quad \mathbf{I}_r]$, then D is a TU matrix, which implies that the dual configuration $\begin{bmatrix} \mathbf{I}_{n-r} & -D^{\mathsf{T}} \end{bmatrix}$ is also regular.

Definition 28 (equivalence of weight vectors). *We say that two weight vectors* $w_1, w_2 \in \mathbb{Z}^n$ *are* equivalent *if* $w_1^{\mathsf{T}} c = w_2^{\mathsf{T}}$ $\frac{1}{2}$ c for all circuits c ∈ $\mathscr{C}(A)$. Notice that, in a MCICP instance, any row W(i,⋅) of the weight *matrix can be replaced by any equivalent weight vector (transposed).*

A *basis* of *A* is a subconfiguration of *A* forming a maximally linearly independent set.

Lemma 29. Let B denote a basis of A. For every weight vector $w_1 \in \mathbb{Z}^n$ there exists an equivalent weight *function* $w_2 \in \mathbb{Z}^n$ such that $w_2(v) = 0$ for all $v \in B$.

Proof. Notice that if $c^* \in \mathcal{C}^*(A)$ is any cocircuit of *A*, then w_1 and $w_1 + c^*$ are equivalent. Every basis element *v* ∈ *B* has a corresponding fundamental cocircuit c_v^* ∈ $\mathcal{C}^*(A)$ that uses *v* and no other element of *B*. By the remark above, w_1 and $w_2 := w_1 - \sum_{v \in B} w_1(v) c_v^*$ $_{\nu}^{\ast}$ are equivalent weight vectors. Moreover, $w_2(v) = 0$ for all $v \in B$, as required. \Box

Lemma 30. Let $A \in \mathbb{R}^{m \times n}$ be a 2-connected regular configuration, and let $\overline{v} \in A$. Let $w \in \mathbb{Z}^n$ be a weight *vector. If every circuit c* $\epsilon \mathcal{C}(A)$ *using* $\overline{\nu}$ has zero weight, then w is equivalent to the zero weight function.

Proof. We claim that every circuit of *A* has zero weight.

Before proving the claim, we verify that it implies the result. Assuming that the claim holds, pick a basis *B* and consider any weight vector *w* ′ equivalent to *w* that is zero on all elements of *B*. By Lemma [29,](#page-21-0) such a weight function exists. Now consider any $v \in A \setminus B$, and let $c \in \mathcal{C}(A)$ denote the unique circuit of *A* whose support is included in *B* ∪{*v*} and that uses *v* positively. We get

$$
w'(v) = \sum_{u \in \text{supp}(c)} w'(u) c(u) = (w')^{\text{T}} c = 0.
$$

We conclude that w' is identically zero.

In order to establish the claim, let $c_1 \in \mathcal{C}(A)$ be arbitrary. If c_1 uses $\overline{\nu}$, then $w^{\dagger}c_1 = 0$ by hypothesis.

From now on, assume that c_1 does not use $\overline{\nu}$, that is, $c_1(\overline{\nu}) = 0$. Since A is 2-connected, for each ν used by c_1 , there exists a circuit $c_2 \in \mathcal{C}(A)$ using both $\overline{\nu}$ and ν . Choose such a vector ν and circuit c_2 such that the union of the supports of c_1 and c_2 is inclusionwise minimal. By negating c_1 and/or c_2 if necessary, we may assume that $c_2(\overline{v}) = 1$, and $c_1(v) = -c_2(v)$.

Now consider the circulation $c_1 + c_2$, and any conformal decomposition $c_1 + c_2 = c_3 + c_4 + \cdots + c_\ell$ as a sum of circuits of *A*. After permuting the indices if necessary, we may assume that $c_3(\overline{v}) = 1$. Let C_i denote the support of c_i , for $1 \le i \le \ell$. (The C_i 's are circuits in the matroid $M(A)$ represented by A.)

Since the decomposition is conformal and $c_1(v) + c_2(v) = 0$, $C_3 \subseteq (C_1 \cup C_2) - v$. In particular, C_3 is distinct from C_2 . Since C_2 is a circuit, $C_3 \setminus C_2$ is nonempty. Hence, C_3 intersects C_1 . By choice of c_2 , we have $C_2 \setminus C_1 \subseteq C_3$.

Using again the fact that the decomposition of $c_1 + c_2$ is conformal, we have $c_3(u) = c_2(u)$ for all *u* ∈ $C_2 \setminus C_1$. This implies that C_i ⊆ C_1 − *v* for $4 \le i \le \ell$. We conclude that $\ell = 3$ and $c_1 + c_2 = c_3 \in \mathcal{C}(A)$. We get $w^{\top} c_1 = w^{\top} c_3 - w^{\top} c_2 = 0$. The claim is proved. \Box

We resume the discussion of the local MCICP instance $I = (p, A, W, d, \ell, u, \overline{v}, \varphi)$ and the corresponding decomposition tree *T* for *A*. Consider a node $t \in V(T)$. We can resort to the proof of Lemma [30](#page-22-1) to check efficiently if *t* is tame. Notice that being tame or wild does not depend on $W_t(\cdot, \overline{v}_t)$. We redefine $W_t(i, \overline{v}_t) \in \mathbb{Z}$ for each $i \in [k]$ in such a way that $W_t c_t = \mathbf{0}$, for all $c_t \in \mathcal{C}(A_t)$. By the proof of Lemma [30,](#page-22-1) t is tame if and only if the weight vector of every circuit of A_t is zero, that is, if and only if $A_t x = 0$ implies $W_t x = 0$ for all vectors x of the appropriate dimension. This can be easily checked by considering any basis B of A_t and checking that every fundamental circuit has a zero weight vector relatively to $W_t.$

Lemma 31. Let $t \in V(T)$, and assume that the weight vector of every circuit of A_t with respect to W_t is zero. Consider a new tree T' with a single node t', letting $A(t')$ denote the vector configuration that is formed by the root vector \overline{v}_t , $f_5(k,\Delta)$ $f_5(k,\Delta)$ $f_5(k,\Delta)$ extra copies of \overline{v}_t and $f_5(k,\Delta)$ further vectors that are all copies of $-\overline{v}_t$, $W_{t'}:=0$, $\ell_{t'}(v) := 0$ and $u_{t'}(v) := 1$ for every $v \in A_{t'} - \overline{v}_t$. Then for some choice of profits $p_{t'}(v)$ for $v \in A_{t'} - \overline{v}_t$, we *have* F_t ^{*'*} $(d_t, \varphi_t) = F_t(d_t, \varphi_t)$ for all $d_t \in B$ and $\varphi_t \in \Phi$.

Proof. First, we notice that if $d_t \neq \mathbf{0}$ then $F_{t'}(d_t, \varphi_t) = F_t(d_t, \varphi_t) = -\infty$ for all $\varphi_t \in \Phi$, hence we can fix $d_t = 0$ for the rest of the proof.

Consider the function of one variable $f(\varphi_t) := F_t(\mathbf{0}, \varphi_t)$. Since A_t is regular, we can omit the integrality condition in the local MCICP, and write

$$
f(\varphi_t) = \max \{ p_t^\top x : A_t x = \mathbf{0}, x(\overline{v}_t) = \varphi_t, \ell_t \le x \le u_t \}.
$$

It is easy to see that f is piecewise linear and concave. Indeed, given integers φ and φ' , corresponding feasible circulations *x* and *x'* respectively attaining *f*(*ϕ*) and *f*(*ϕ*[']), and *λ* ∈ [0, 1] such that *λφ* + (1 − *λ*)*ϕ*['], we see that the convex combination $\lambda x + (1-\lambda)x'$ is a feasible circulation giving a lower bound on $f(\lambda \varphi +$ $(1 - \lambda)\varphi'$).

Furthermore, $f(0) = 0$. Indeed, $f(0) \ge 0$ since $x = 0$ is a feasible solution for $\varphi_t = 0$. Moreover, $f(0) > 0$ would imply the existence of a feasible circuit of *A* with zero weight and positive profit, which we assume does not exist.

For the sake of simplicity, let $\Gamma = \Gamma(k,\ell) := f_5(k,\Delta)$ $\Gamma = \Gamma(k,\ell) := f_5(k,\Delta)$ $\Gamma = \Gamma(k,\ell) := f_5(k,\Delta)$. Label the copies of \overline{v}_t in $A(t')$ as $v_{-1}, \ldots, v_{-\Gamma}$ and the copies of $-\overline{v}_t$ as v_1, \ldots, v_{Γ} . We let $p_{t'}(v_i) := f(i) - f(i-1)$ and $p_{t'}(v_{-i}) := f(-i) - f(-i+1)$ for $i = 1,..., \Gamma$. Since f is concave, we have $p_{t'}(v_1) \ge p_{t'}(v_2) \ge \cdots \ge p_{t'}(v_{\Gamma})$ and $p_{t'}(v_{-1}) \ge p_{t'}(v_{-2}) \ge \cdots \ge \Gamma$ p_t ^{*'*}(v _{−Γ}</sub>). We leave it to the reader to check that F_t [']($\mathbf{0}, \varphi_t$) = $f(\varphi_t) = F_t(\mathbf{0}, \varphi_t)$ for all $\varphi_t \in \Phi$.

To show that F_t ['](0, φ_t) = $f(\varphi_t)$ for $\varphi_t \in \Phi$, consider an optimal solution x' of the rooted instance for node *t'*, and assume that $\varphi_t \ge 0$, the other case being similar. First, one checks that if $x'(v_i) = 1$ for any $i < 0$, there is some $j > 0$ with $x'(v_j) = 1$ and we can set both to zero without decreasing the objective value. Second, if there is some $0 < i < j$ with $x'(v_i) = 0$, $x'(v_j) = 1$, then we can swap the two values without decreasing the objective value. Hence, given that $x'(\bar{v}_t) = \varphi_t$ and A'_t $t'_t x' = 0$, we conclude that $x'(v_i) = 1$ for $i = 1, ..., \varphi_t$ and 0 for all other vectors. This implies that $F_{t'}(\mathbf{0}, \varphi_t)$, the objective value of x' , is equal to $\sum_{i=1}^{\varphi_t} f(i) - f(i-1) = f(\varphi_t)$. \Box

Thanks to Lemma [24](#page-20-0) and Lemma [31,](#page-22-0) we can replace the subtree rooted at any tame node of $V(T)$ by a single node whose corresponding configuration is a multiset of parallel vectors without changing any value function F_t for the other nodes. This allows us to assume that every tame node is a leaf. Next, for each tame leaf node $t \in V(T)$ with parent $u \in V(T)$ we replace $A(u)$ by $A(u) \oplus_2 A_t = A(u) \oplus_2 A(t)$ and delete node *t* from the decomposition tree. In this way, we obtain a new local MCICP instance $I' = (p', A', W', d', \ell', u', \overline{\nu}', \varphi')$ and decomposition tree *T*['] of *A*['] that is a subtree of *T* containing the root, in such a way that *T'* has no tame node, $A(t)$ is *almost* 3-connected for every $t \in V(T)$, and the value function F'_t with respect to *I'* equals the value function F_t with respect to the original instance *I*, for every node *t* ∈ $V(T')$.

Final DP

Consider again a local MCICP instance $I = (p, A, W, d, \ell, u, \overline{v}, \varphi)$ and decomposition tree *T* for *A*. Assume that no node of *T* is tame, and *A*(*t*) is almost 3-connected for all nodes $t \in V(T)$. By the previous section, we may reduce to this case. Fix some non-leaf node $t \in V(T)$ and denote its children by t_1, \ldots, t_q . The next result is the final piece in our reduction of MCICP to the almost 3-connected case.

Lemma 32. *Every circuit of c* ∈ $C(A_t)$ *crosses at most* 2k∆ *vectors* \overline{v}_{t_i} *such that node* t_i *is a wild child of t.*

Proof. After renumbering the children of *t*, we may assume that the indices $i \in [q]$ such that *c* crosses \overline{v}_{t_i} and t_i is wild are $i = 1, ..., q'$ for some $q' \leq q$. Then c splits into one circuit c_0 of $A(t) \oplus_2 A_{t_{q'+1}} \oplus_2 \cdots \oplus_2 A_{t_q}$ and *q'* circuits *c*₁, ..., *c*_{*q'*} of *A*_{*t*₁}, ..., *A*_{*t_q'*} respectively. For each *i* \in [*q'*], let *c'*_{*i*} c_i' and c_i'' i' denote two circuits of A_{t_i} such that c_i' $c_i^{\prime}(\overline{v}_{t_i}) = c_i^{\prime\prime}$ $W_i(\overline{v}_{t_i}) = 1$ and $W_{t_i}c_i'$ $\sum_{i}^{'} \neq W_{t_i} c''_i$ *i* .

Towards a contradiction, suppose that *q* ′ > 2*k*∆. By the pigeonhole principle, we can find a row index *j* ∈ [*k*] and a set of 2∆ + 1 indices *i* ∈ [*q*[']] such that $W_{t_i}(j,·)c_i^j$ $\sum_{i}^{J} \neq W_{t_i}(j,\cdot)c''_i$ i' . After permuting the indices

again, we may assume that these indices are $i = 1, ..., 2\Delta + 1$. By exchanging c_i' c_i ['] and c_i ^{''} $\binom{n}{i}$ if necessary, we may assume that $W_{t_i}(j, \cdot) c_i'$ $W'_{i} > W_{t_i}(j, \cdot) c''_{i}$ *^{''}* for all *i* ∈ [2∆ + 1].

Notice that replacing c_i for each $i \in [2\Delta + 1]$ with any other circuit $\tilde{c}_i \in \{\pm c_i\}$ c_i^{\prime} , $\pm c_i^{\prime\prime}$ $\binom{n}{i}$ of A_{t_i} such that $c_i(\overline{v}_{t_i}) = \tilde{c}_i(\overline{v}_{t_i})$ in the decomposition of *c* results in a new circuit of A_t . Consider the circuit c^+ obtained from *c* by replacing c_i with c'_i \int_i if $c_i(\overline{v}_{t_i}) = 1$, and with $-c_i''$ *^{''}*</sup> if $c_i(\overline{v}_{t_i}) = -1$, for $i \in [2\Delta+1]$. Consider the circuit c^- obtained similarly by exchanging the roles of c'_i c_i ['] and c_i ^{''} *i*^{*i*}. In this way, $W_t(j, ⋅) c^+$ and $W_t(j, ⋅) c^-$ differ by at least 2∆+1, contradicting the fact that both weights should be at most ∆ in absolute value.

This last result directly yields an improvement to Lemma [24.](#page-20-0) Let $(d_t^0,d_{t_1},\ldots,d_{t_q},\varphi_{t_1},\ldots,\varphi_{t_q})\in B^{q+1}\times$ Φ *^q* be a *guess* such that

$$
d_t^0 + \sum_{i=1}^q \left(d_{t_i} - \left(W_t(\cdot, \overline{\nu}_{t_i}) - W_t^0(\cdot, \overline{\nu}_{t_i}) \right) \varphi_{t_i} \right) = d_t,
$$

see [\(6\)](#page-16-2). Recall that every MCICP instance has an optimal solution that is a sum of at most *f*[5](#page-6-3)(*k*,∆) circuits. Together with Lemma [32,](#page-23-0) this implies that we can restrict to the guesses such that

$$
\sum_{i=1}^q |\varphi_{t_i}| \leq 2k\Delta f_5(k,\Delta).
$$

Moreover, we can further require that there are at most $f_5(k,\Delta)$ $f_5(k,\Delta)$ $f_5(k,\Delta)$ indices *i* such that $\varphi_{t_i} = 0$ and $d_{t_i} \neq 0$. Let Γ_t denote the set of all guesses $(d_t^0, d_{t_1},...,d_{t_q}, \varphi_{t_1},...,\varphi_{t_q}) \in B^{q+1} \times \Phi^q$ satisfying the three conditions above. Then we can change [\(6\)](#page-16-2) to

$$
F_t(d_t, \varphi_t) = \max \left\{ F_t^0(d_t^0, \varphi_t, -\varphi_{t_1}, \dots, -\varphi_{t_q}) + \sum_{i=1}^q F_{t_i}(d_{t_i}, \varphi_{t_i}) : (d_t^0, d_{t_1}, \dots, d_{t_q}, \varphi_{t_1}, \dots, \varphi_{t_q}) \in \Gamma_t \right\}
$$

where $d_t \in B$ and $\varphi_t \in \Phi$ are arbitrary. We point out that the DP can be once again extended to store an optimal solution for each table entry $F_t(d_t, \varphi_t)$. Putting everything together, we obtain the following result, which concludes this section.

Theorem [33](#page-24-1). *Let k*, $\Delta \in \mathbb{Z}_{\geq 1}$ *be constants, and let f*₃₃(*k*, Δ) := *k* Δf_5 Δf_5 (*k*, Δ)*. Any* 2*-connected instance of the MCICP on a matrix with n columns can be solved after solving at most* (*f*[33](#page-24-1)(*k*,∆))*kn ^O*(*f*[33](#page-24-1)(*k*,∆)) *instances of the MCICP that are almost* 3*-connected and on matrices with at most n columns, and performing extra work in time* $(f_{33}(k,\Delta)n)^{O(f_{33}(k,\Delta))}$ $(f_{33}(k,\Delta)n)^{O(f_{33}(k,\Delta))}$ $(f_{33}(k,\Delta)n)^{O(f_{33}(k,\Delta))}$.

5 Algorithm for the cographic case

This section provides a strongly polynomial time algorithm for solving cographic instances, relying on structural results that are established in Section [6.](#page-32-0) In Section [5.1,](#page-25-0) we discuss circuit vectors of (co) graphic instances in terms of the corresponding graphs. In Section [5.2,](#page-26-0) we recall how to reformulate any cographic instance in the vertex space through a change of variables that transforms the main part of the coefficient matrix into the transpose of an incidence matrix. By this change of variables, the *k* additional rows of the constraint matrix yield a *k*-dimensional weight vector for each vertex. Defining the set of roots as the set of vertices whose weight vector is nonzero, we turn the underlying graph into a rooted graph. In Section [5.3,](#page-27-0) we prove that these graphs do not have a rooted $K_{2,t}$ -minor for $t := 4k\Delta + 1$. This in turn implies the strong structural properties stated in Theorem [9,](#page-9-0) which are used in Section [5.4](#page-29-0) to prove Theorem [1.](#page-1-2)

5.1 Circuits and connectivity

Let *G* be any directed graph. We allow parallel and anti-parallel directed edges, but do not allow loops. Seen as a vector configuration, the incidence matrix of *G* associates to each directed edge (*v*, *w*) $\in E(G)$ the vector $\chi^{\nu} - \chi^{\nu \nu}$ in $\mathbb{R}^{V(G)}$, where { $\chi^{\nu} : \nu \in V(G)$ } denotes the canonical basis of $\mathbb{R}^{V(G)}$.

Definition 34 (graphic and cographic configurations)**.** *Let A* ∈ R *^m*×*ⁿ be a configuration. We call A* graphic *whenever A is linearly equivalent to the configuration defined by the incidence matrix of some directed graph G, and* cographic *whenever A is linearly equivalent to the dual of some graphic configuration.*

Up to a permutation of its columns, every rank-*r* graphic configuration is linearly equivalent to a standardized configuration of the form $[D \mid I_r]$, where $D \in \{0, \pm 1\}^{r \times (n-r)}$ is a network matrix. Conversely, every network matrix can be obtained in this way. Recall that the dual configuration of $\begin{bmatrix} D & \mathbf{I}_r \end{bmatrix}$ is $\begin{bmatrix} \mathbf{I}_{n-r} & -D^{\mathsf{T}} \end{bmatrix}$.

We remark that we may restrict to directed graphs *G* that are (weakly) connected. Indeed, if *G* has $\ell > 1$ connected components then we can pick any vertex v_i in the *i*th component for each $i \in [\ell]$, and merge *v*₁, ..., *v*_ℓ in a single vertex. The resulting graph is connected and the corresponding configuration is linearly equivalent to the original one.

In Lemma [35](#page-25-1) below, we give the well-known characterization of circuits of graphic and cographic configurations. Before doing this, we need more terminology regarding directed graphs (we generally follow [\[Sch03\]](#page-58-12)). Note that the lemma follows from [\[Oxl06,](#page-58-10) Proposition 2.3.1].

Let *G* be a connected directed graph. For $X \subseteq V(G)$, we let $\delta(X)$ denote the set of edges with one end in *X* and the other end in $\overline{X} := V(G) \setminus X$, $\delta^{out}(X)$ denote the set of edges leaving *X*, and $\delta^{in}(X)$ denote the set of edges entering *X*. Notice that $\delta(X) = \delta^{out}(X) \cup \delta^{in}(X)$. The set $\delta(X)$ is called an *undirected cut*, while $\delta^{\text{out}}(X)$ and $\delta^{\text{in}}(X)$ are *directed cuts*. If *X* is a proper and nonempty subset of $V(G)$, we say that $\delta(X)$, $\delta^{out}(X)$ and $\delta^{in}(X)$ are *nontrivial* cuts. It is easy to see that an inclusion-wise minimal nontrivial undirected cut, sometimes called *bond*, is a undirected cut $\delta(X)$ such that each of X and \overline{X} induces a nonempty connected subgraph of *G*. To every such cut there correspond two opposite *signed incidence vectors* in $\{0, \pm 1\}^{E(G)}$, namely, $\chi^{\delta^{\text{out}}(X)} - \chi^{\delta^{\text{in}}(X)}$ and $\chi^{\delta^{\text{in}}(X)} - \chi^{\delta^{\text{out}}(X)}$.

An *undirected circuit* of *G* is a cycle of its underlying undirected graph, that is, a sequence *C* = $(v_0, e_1, v_1, \ldots, e_k, v_k)$ where $k \ge 1$, v_0, v_1, \ldots, v_k are vertices, e_i is an edge incident to both v_{i-1} and v_i for each $i \in [k]$, $v_0 = v_k$, v_1, \ldots, v_k are pairwise distinct. To every undirected circuit there corresponds a *signed incidence vector* in $\{0, \pm 1\}^{E(G)}$, defined as $\chi^{C^+} - \chi^{C^-}$ where C^+ denotes set of edges of the circuit that are traversed forwards and *C* [−] the set of edges traversed backwards.

Lemma 35. *Let A* ∈ R *^m*×*ⁿ be a graphic configuration, and let G be a corresponding connected directed graph, with* $n = |E(G)|$ *. Then, the rank of A equals* $|V(G)| - 1$ *, the circuits of A are the signed incidence vectors of undirected circuits of G, and the cocircuits of A are the signed incidence vectors of minimal nontrivial undirected cuts of G.*

The next lemma relates the connectivity of a graphic or cographic configuration to the (undirected) vertex connectivity of the corresponding graph, see [\[Oxl06,](#page-58-10) Chapter 8]. Below, we say that directed graph *G* is *simple* if it contains no parallel or antiparallel directed edges.

Lemma 36. Let $A \in \mathbb{R}^{m \times n}$ be a rank-r configuration that is graphic or cographic, and let G be a corre*sponding (weakly) connected directed graph, with* $r = |V(G)| - 1$ *and* $n = |E(G)|$ *. Assuming* $r \ge 2$ *, A is* 2*connected if and only if G is 2-connected. Assuming* $n \geq 4$ *, A is 3-connected if and only if G is 3-connected and simple. Now assume that A is a cographic configuration. Then A is almost* 3*-connected if and only if G can be obtained from a* 3*-connected simple graph by subdividing some of its edges.*

5.2 Change of variables

Given a (feasible) cographic instance of Problem [1,](#page-5-2) we can assume without loss of generality that $A =$ $[I_{n-r}$ −*D*^T], where *D* ∈ {0, ±1}^{*r*×(*n*−*r*)} is a network matrix. Next, using Proposition [5](#page-6-3) and applying a translation mapping *z* to **0**, we reduce to an instance of Problem [2](#page-17-3) (MCICP), with the same matrix *A* (regarded as a vector configuration, A is a cographic configuration) and $b = 0$. Finally, by the results of Section [4,](#page-15-0) we further reduce this instance to a polynomial number of instances on almost 3-connected, cographic configurations. We point out that given the network matrix *D* (or its transpose *D* ⊺), one can efficiently find a corresponding directed graph *G*, see for instance [\[Tut60\]](#page-59-5).

Recall from Section [2.5](#page-7-2) that by letting $x = M^{\dagger}y$ where $M = \begin{bmatrix} N & B \end{bmatrix}$ is the incidence matrix of *G* with the row corresponding to some $v_0 \in V(G)$ removed and *B* is a basis of *M*, we can rewrite the cographic instance of Problem [2](#page-17-3) at hand almost as an instance of the maximal constrained integer potential problem. More precisely, we transform [\(IP](#page-17-2)₂) for $A = \begin{bmatrix} I_{n-r} & -D^T \end{bmatrix}$ into (IP₁[-cographic\)](#page-7-0). By combining Lemma [35](#page-25-1) and the following lemma, we see that circuits of *A* bijectively correspond to docsets of *G*.

Lemma 37. *A vector* $x \in \mathbb{R}^n \cong \mathbb{R}^{E(G)}$ *is a circuit of* $A = [\mathbf{I}_{n-r} \ -D^{\dagger}]$ *if and only if* $x = \pm M^{\dagger} \chi^S$ *where* $S \subseteq$ $V(G - v_0)$ *is a nontrivial docset of G, and the incidence vector is taken in* $\mathbb{R}^{V(G - v_0)} \cong \mathbb{R}^r$.

Proof. The map α : ker(*A*) $\rightarrow \mathbb{R}^r$ such that $\alpha(x) = B^{-\dagger}x_B$ for all $x = \begin{bmatrix} x_N \\ x_B \end{bmatrix} \in \text{ker}(A)$ is an isomorphism whose inverse is given by $\alpha^{-1}(y) = M^{\dagger}y$ for all $y \in \mathbb{R}^r$. If *S* is a nontrivial docset of *G* with $v_0 \notin S$, then $\pm M^{\dagger}\chi^{S} = \pm \chi^{\delta^{out}(S)} \mp \chi^{\delta^{in}(S)}$, which are both circuits of *A*. Moreover, every circuit of *A* can be obtained in this way. \Box

Recall also that the problem we will solve, namely the maximum constrained integer potential prob-lem, can be obtained from (IP₁[-cographic\)](#page-7-0) by adding one extra column to M^{\dagger} and one extra row to *y*, both corresponding to vertex v_0 , in such a way that M^{\dagger} becomes the transpose of the (full) incidence matrix of *G*. Notice that letting $y(v_0) := 0$ transforms back [\(MCIPP\)](#page-2-0) into (IP₁[-cographic\)](#page-7-0).

In order to ease the task of the reader, we state formally below the precise version of the MCIPP that we have to solve, including the condition on the weight of docsets.

Problem 3. *Let k*,∆ ∈ Z≥¹ *be constants. Given a connected directed graph G, profit vector p* ∈ Z *^V* (*G*) *with* $\sum_{v\in V(G)} p(v) = 0$, weight matrix $W\in \mathbb{Z}^{[k]\times V(G)}$ such that each row of W sums up to 0 and $\|W\chi^S\|_\infty \leq \Delta$ for all docsets S, target vector $d \in \mathbb{Z}^k$, bounding vectors $\ell, u \in \mathbb{Z}^{E(G)}$, find

$$
\max \{ p^{\top} y : \ell(v, w) \le y(v) - y(w) \le u(v, w) \,\forall (v, w) \in E(G), \, Wy = d, \, y \in \mathbb{Z}^{V(G)} \} \,.
$$
 (IP₃)

Using the results of Section [4](#page-15-0) and Lemma [36,](#page-25-2) we may assume without loss of generality that the input graph *G* is a 3-connected graph some of whose edges are subdivided. In fact, we can also assume that no vertex of degree 2 is a root. This follows from Lemma [38](#page-26-1) and the fact that the edges *e* incident to vertices of degree 2 have $W(\cdot, e) = 0$, see Lemma [31.](#page-22-0) Moreover, we can still argue that (IP_3) either is infeasible, or has an optimal solution *y* that is a sum of at most *f*[5](#page-6-3)(*k*,∆) incidence vectors of docsets of *G*. This follows from Proposition [5](#page-6-3) and Lemma [37.](#page-26-3) Notice that for any solution *y* to Problem [3,](#page-26-4) *y* ′ := *y* +**1** is a solution to Problem [3](#page-26-4) with $p^{\dagger} y' = p^{\dagger} y$.

The following lemma follows directly from the definition of the incidence matrix, and gives a concrete characterization of the profit vector *p* and weight matrix *W* in Problem [3](#page-26-4) after our change of variables.

Lemma 38. Let $M \in \{0, \pm 1\}^{V(G) \times E(G)}$ be the incidence matrix of a directed graph G and $w \in \mathbb{Z}^{E(G)}$ a weight *vector. Then for each* $v \in V(G)$ *,*

$$
(M^{\top}w)(v) = \sum_{e \in \delta^{\mathrm{out}}(v)} w(e) - \sum_{e \in \delta^{\mathrm{in}}(v)} w(e).
$$

5.3 Rooted $K_{2,t}$ -models

In this section we consider (vertex)-weighted undirected graphs (*G*,*a*) without a docset of large weight, and study related properties of rooted $K_{2,t}$ -minors in such graphs, where the roots are the vertices ν with $a(v) \neq 0$. We use the following notation.

 $\bf{Definition~39}$ (maximum weight of a docset $\beta(G,a)$). *Let G be a connected graph and a* $\in \mathbb{Z}^{V(G)}$ *be vertex weights such that* $\sum_{v \in V(G)} a(v) = 0$ *. For S* ⊆ *V*(*G*)*, we let* $a(S) := \sum_{v \in S} a(v)$ *. We define*

$$
\beta(G, a) := \max \{a(S) : S \subseteq V(G), S \text{ is a docset of } G\}.
$$

Notice that $a(\overline{S}) = -a(S)$, hence $\beta(G, a) \ge 0$. If *G* is 2-connected, then {*v*} is a docset for every vertex *v*, which implies $\beta(G, a) \ge \max\{|a(v)| : v \in V(G)\}\)$. Every instance of Problem [3](#page-26-4) has $\beta(G, a) \le \Delta$ for all weight vectors *a* which are (the transpose of) a row of *W* .

The main result we prove is that no weighted graph (G, a) with $\beta(G, a) \leq \Delta$ can contain a rooted $K_{2,t}$ minor for $t > 4\Delta$, taking as roots the vertices *v* with $a(v) \neq 0$. We need the following technical lemma. Below, we use $N(X)$ to denote the open neighborhood of $X \subseteq V(G)$ in *G*, that is, we let $N(X) := \{v \in V(G) | v \in V(G) \}$ *V*(*G* − *X*): ∃*w* ∈ *X* : *v w* ∈ *E*(*G*)}. For *X* ⊆ *V*(*G*) and *v* ∈ *V*(*G*) we denote by *X* + *v* the set *X* ∪ {*v*}.

Lemma 40. Let G be a 2-connected graph and let X, Y be two docsets of G. If $X \subseteq Y$, then there exists some *vertex* $v \in Y \setminus X$, such that $X + v$ is a docset.

Proof. Let $u \in \overline{Y}$ be arbitrary. Since *Y* is connected and *X* is a proper subset of *Y*, $N(X) \cap Y$ is nonempty. Let *v* denote any vertex of $N(X) ∩ Y$ whose distance from *u* in $G[\overline{X}] = G - X$ is maximum. Clearly, $G[X + v]$ is connected. We claim that $G[X + v]$ is also connected.

For the sake of contradiction, assume that $G[X + v]$ has several (connected) components. Since $G[Y]$ is connected, it is contained in a unique component of $G[X + v]$, say *K*. Let *w* be any vertex of $N(X) \cap Y$ distinct from *v*. By choice of *v*, no shortest path from *u* to *w* in $G[\overline{X}]$ contains *v*. This proves that $w \in K$.

Let *K'* be any component of $G[\overline{X+v}]$ distinct from *K*. Observe that $V(K') \subseteq Y$. Since *K'* is a component of $G[\overline{X+v}]$, we have $N(K') \subseteq X + v$. Since *G* is 2-connected, *K'* should have at least two neighbors, and at least one distinct from *v*. But this implies that *K'* contains at least one vertex $w \in N(X) \cap Y$ distinct from *v*. By the argument in the previous paragraph, $w \in K$, hence we get a contraditcion to the assumption that *K* and *K'* are different connected components in $G[X + v]$. Hence, the claim is proved and $X + v$ is a docset, as required. \Box

Definition 41 (model)**.** *Given a graph H, an H*-model *M in a graph G consists of one connected vertex subset* $M(v) ⊆ V(G)$ *for each vertex* $v ∈ V(H)$ *, called a* branch set*, and one edge* $M(uv) ∈ E(G)$ *with one endpoint in* $M(u)$ *and the other in* $M(v)$ *for each edge uv* $\in E(H)$ *such that all branch sets are pairwise vertex disjoint. Note that G contains H as a minor if and only if G has an H-model.*

Definition 42 (rooted *K*2,*t*-model)**.** *Given a rooted graph* (*G*,*R*)*, a* rooted *K*2,*t*-model *in* (*G*,*R*) *is a model of K*2,*^t in G such that each of the t branch sets corresponding to vertices on the "t side" of K*2,*^t contains a vertex from R. (Such branch sets will be called* central branch sets*.) Note that* (*G*,*R*) *contains a rooted* $K_{2,t}$ *-minor if and only if* (G, R) *contains a rooted* $K_{2,t}$ *-model.*

In the proof below, we say that two vertex subsets *X*,*Y* of a graph *G touch* if their distance in *G* is at most 1.

Lemma 43. Let (G, a) be a 2-connected weighted graph, where $a \in \mathbb{Z}^{V(G)}$ satisfies $\sum_{v \in V(G)} a(v) = 0$. Let $R = \{v \in V(G) : a(v) \neq 0\}$. If (G, R) *contains a rooted* $K_{2,t}$ *-model, then* $\beta(G, a) \geq t/4$.

Proof. Notice that we may assume that *t* ≥ 4 without loss of generality. Notice further that any rooted $K_{2,t}$ -model in (G, R) that avoids some vertex can be modified in order to cover the whole vertex set. This follows easily from the connectivity of *G*. We call such models *full*. Let *M* denote a full rooted $K_{2,t}$ model in (*G*,*R*) that maximizes the number of nonzero weight central branch sets. In other words, letting *V*($K_{2,t}$) := {−1,0}∪{1,...,*t*}, we assume that *M* maximizes

$$
f(M) := |\{i \in [t] : a(M(i)) \neq 0\}|\
$$

among all full rooted $K_{2,t}$ -models in (G, R) .

We claim that $f(M) \ge t/2$, that is, at least $t/2$ of the central branch sets have nonzero weight.

For notational convenience, we let $B_i := M(i)$ (where $i \in \{-1,0\} \cup [t]$) denote the branch sets of M. Assuming that the claim holds, consider the docsets

$$
X^+ := B_{-1} \cup \bigcup_{i \in [t]: a(B_i) > 0} B_i \quad \text{and} \quad X^- := B_{-1} \cup \bigcup_{i \in [t]: a(B_i) < 0} B_i \, .
$$

Notice that *a*(*X*⁺) − *a*(*X*[−]) ≥ *t*/2. Hence, *a*(*X*⁺) ≥ *t*/4 or *a*(*X*[−]) ≤ −*t*/4. This implies that *β*(*G*, *a*) ≥ *t*/4.

Consider any central branch set B_i of zero weight. Since M is full, B_i is a docset. We claim that B_i has a partition into docsets U_i , V_i , which both are of nonzero weight. Let $u_i \in B_i$ denote any vertex adjacent to some vertex of *B*−1. By applying iteratively Lemma [40](#page-27-1) starting from docsets {*ui*} and *Bⁱ* , we can find a docset $W_i \subsetneq B_i$ that contains u_i and a unique root. Thus, both $a(W_i) \neq 0$ and $a(B_i \setminus W_i) \neq 0$. Observe that $G[B_i \setminus W_i]$ decomposes into connected components, at least one of which has nonzero weight, which we denote by V_i . Let $U_i := B_i \setminus V_i$. By construction, U_i and V_i induce connected subgraphs and $a(V_i) \neq 0$ and thus $a(U_i)\neq 0.$ Observe that also $\overline{U_i}$ and $\overline{V_i}$ induce connected subgraphs. Indeed, the set $\overline{V_i}\cap B_i$ is *U*^{*i*} and therefore connected, the set *U*^{*i*} contains *W*^{*i*} and therefore is touching *B*^{−1}. The set *U*^{*i*} induces a connected subgraph because $G[\overline{W_i}]$ is connected and hence V_i is touching at least one B_j for $j\neq i$ (note that *j* can be also an index of a central branch set different from B_i). Thus, U_i and V_i are docsets.

By maximality of *M*, V_i touches neither B_{-1} nor B_0 and hence U_i touches both B_{-1} and B_0 . Otherwise for some *j* ∈ {−1,0} we could replace B_i by U_i and B_j by $B_j \cup V_i$ and strictly increase $f(M)$.

Recall that U_i is a docset, hence V_i touches B_j for some $j \in [t]$ different from *i*. If $a(V_i) + a(B_j) \neq 0$, we could merge *B^j* and *Vⁱ* and contradict the maximality of *M*. In particular, all the central branch sets *B*^{*j*} with $j \neq i$ touched by V_i have nonzero weight.

Assume that there is an index $i' \in [t]$ with $i' \neq i$, such that $a(B_i) = a(B_{i'}) = 0$ and V_i , $V_{i'}$ touch the same central branch set B_j with $a(B_j) \neq 0$. By the argument given above, we have $a(V_i) + a(B_j) = 0$. This implies that $a(V_i) + a(V_{i'}) + a(B_j) = a(V_{i'}) \neq 0$, in which case we can move all the vertices of V_i and $V_{i'}$ inside B_j to strictly increase $f(M)$, which contradicts again the maximality of M.

It follows that the number of central branch sets *B^j* with nonzero weight are at least as numerous as the central branch sets B_i with zero weight. That is, $f(M) \ge t/2$. This proves the claim.

Using the fact that $f(M) \ge t/2$ and that the model *M* is full, it is easily seen that $\beta(G, a) \ge t/4$. The lemma follows. \Box

Lemma [6](#page-7-1) from Section [2](#page-4-2) follows directly from the previous lemma.

Proof of Lemma [6.](#page-7-1) Assume to the contrary that (G, R) contains a rooted $K_{2,t}$ -minor, and consider a corresponding rooted $K_{2,t}$ -model. By the pigeonhole principle, there exists a row w_i , $i \in [k]$ of W , such that at least 4∆+1 central branch sets of the model contain a vertex with a non-zero entry in *wⁱ* . By Lemma [43,](#page-27-2) this implies that $\beta(G, w_i) \geq \Delta + 1$, a contradiction. \Box

We conclude with two simple results describing how the size of the largest rooted $K_{2,t}$ -model in a graph can change by adding or contracting an edge.

Lemma 44. Let (G, R) be a rooted graph, and let $e = uv$ where $u, v \in V(G)$. If $(G + e, R)$ contains a rooted *K*_{2,*t*}*-model* (*with* $t \geq 2$ *), then* (*G, R*) *contains a rooted K*_{2,} $\lceil t/2 \rceil$ *-model.*

Proof. Let *M* be a rooted $K_{2,t}$ -model in $(G+e, R)$. If the vertices *u* and *v* are contained in different branch sets of the model, then (G, R) contains a rooted $K_{2,t-1}$ -model, and hence a rooted $K_{2,\lceil t/2 \rceil}$ -model since *t* ≥ 2. If *e* is contained in one of the central branch sets $M(i)$ (with $i \in [t]$), then again (*G*,*R*) contains a rooted *K*2,*t*−1-model. Finally, suppose that *e* is contained in one of the branch sets *M*(−1) or *M*(0). Without loss of generality, assume that *e* is contained in *M*(−1). Then *G*[*M*(−1)] has at most two connected components and therefore (G, R) contains a rooted $K_{2, [t/2]}$ -model. \Box

The following lemma is straightforward. Below, we let (*G*,*R*)/*e* denote the rooted graph obtained from (G, R) by contracting edge $e = uv \in E(G)$: if none of *u* or *v* is a root then we keep the same roots, otherwise we remove u and/or v from the set of roots and replace them by the vertex arising from the contraction of *e*.

Lemma 45. *Let* (G, R) *be a rooted graph, and let* $e \in E(G)$ *. If* (G, R) */e contains a rooted* $K_{2,t}$ *-model, then* (G, R) *also contains a rooted* $K_{2,t}$ *-model.*

5.4 The algorithm

In this section, we describe how to solve instances of Problem [3,](#page-26-4) assuming the structural results from Section [6.](#page-32-0) We remark that the dynamic programming (DP) algorithm in this section is an adaptation of the DP algorithm in Section [4.4.](#page-19-0) Here, we consider local instances for each bag of the tree-decomposition and each possible way a superposition of $f_5(k,\Delta)$ $f_5(k,\Delta)$ $f_5(k,\Delta)$ docsets of *G* can intersect the roots within the bag, and then solving an IP with a TU constraint matrix for each fixed choice. The optimal solutions to the local instances are then composed recursively to an optimal solution of the whole instance.

Structural foundations

We start by recalling the relevant definitions and result from Section [2,](#page-4-2) which underlie our approach.

Definition 46 (tree-decomposition). A tree-decomposition of a graph G is a pair (T, \mathcal{B}) where T is a tree and $\mathcal{B} = \{B_t : t \in V(T)\}\$ *is a collection of subsets of* $V(G)$ *, called* bags. Moreover, we require the following *conditions:*

 (i) $\bigcup_{t \in V(T)} B_t = V(G)$ *,*

(ii) for every edge $e \in E(G)$ *, there is* $t \in V(T)$ *, such that* $e \subseteq B_t$ *,*

(*iii*) let t, t', t'' $\in V(T)$ be such that t' lies on the path between t and t'' in T, then $B_t \cap B_{t'} \subseteq B_{t'}$.

We always assume that the decomposition tree *T* is rooted at one of its nodes, called the *root node* of *T* (not to be confused with the root vertices of *G*).

Definition 47 (ℓ -special tree-decomposition). *Given a graph G and an integer* $\ell \geq 1$ *, we say that a treedecomposition* (*T*, \mathcal{B}) *is* ℓ *-special if*

(i) for each tt' $\in E(T)$ *we have* $|B_t \cap B_{t'}| \leq \ell$ *, and*

(ii) every node $t \in V(T)$ *has at most* ℓ *children.*

For any given node $t \in V(T)$, we define T_t as the subtree of T rooted at t, and denote by G_t the subgraph of *G* induced by $\bigcup_{u \in T_t} B_u$. If *t* is not the root node, let *t*^{*'*} denote its parent. We define the *adhesion* along the edge $e = t t'$ as the intersection of the corresponding bags and let $X_t := B_t \cap B_{t'}$. If *t* is the root node, we let $X_t := \emptyset$.

Our final definition concerns the possible intersections of docsets of *G* with a subset of the roots.

Definition 48 (docset profiles and superprofiles). Let $\mathcal{S}(G)$ denote the collection of all docsets in G. The docset profile *of a subset R'* \subseteq *R is the collection of sets* $\mathcal{P}(G, R') := \{R' \cap S \mid S \in \mathcal{S}(G)\}\)$. A docset superprofile of R' is a collection of subsets of R' that contains the docset profile $\mathcal{P}(G, R')$.

Consider an instance of Problem [3](#page-26-4) defined by a tuple (p, ℓ, u, G, W, d) . Recall that Theorem [9](#page-9-0) (see Section [2\)](#page-4-2) states that, for each fixed $s \in \mathbb{Z}_{\geq 1}$, we can efficiently obtain an $f_9(s)$ $f_9(s)$ $f_9(s)$ -special tree-decomposition of a 3-connected rooted graph (*G*,*R*) with no rooted *K*2,*s*-minor, together with a docset superprofile of R_t := $R \cap B_t$ for each $t \in V(T)$. We prove this in Section [6](#page-32-0) and extend the result to subdivisions of 3connected graphs without roots of degree 2, see Proposition [78.](#page-53-1)

The local instances

We proceed to define the *local docset completion problem* that our algorithm will solve within the DP, see Definition [49](#page-30-0) below.

Recall that since *G* is 2-connected, {*v*} is a docset for every $v \in V(G)$, implying that $||W||_{\infty} \leq \Delta$. Hence, if *y* ∈ $\mathbb{Z}^{V(G)}$ is the sum of (at most) *f*_{[5](#page-6-3)}(*k*, ∆) incidence vectors of docsets, and $y' \in \mathbb{Z}^{V(G)}$ is obtained from *y* by zeroing an arbitrary set of coordinates, we have $||Wy'||_{\infty} \le n\Delta f_5(k,\Delta)$ $||Wy'||_{\infty} \le n\Delta f_5(k,\Delta)$ $||Wy'||_{\infty} \le n\Delta f_5(k,\Delta)$. In view of this, we let $\mathcal{D} :=$ $[-n\Delta f_5(k,\Delta), n\Delta f_5(k,\Delta)]^k \cap \mathbb{Z}^k$ $[-n\Delta f_5(k,\Delta), n\Delta f_5(k,\Delta)]^k \cap \mathbb{Z}^k$ $[-n\Delta f_5(k,\Delta), n\Delta f_5(k,\Delta)]^k \cap \mathbb{Z}^k$. Thus \mathscr{D} is a polynomial-size set containing the weight vector of any partial solution we consider in the context of the DP. Finally, we let $\Phi := [0, f_5(k,\Delta)] \cap \mathbb{Z}$ $\Phi := [0, f_5(k,\Delta)] \cap \mathbb{Z}$ $\Phi := [0, f_5(k,\Delta)] \cap \mathbb{Z}$. Every time we need to guess the value of a variable *y*(*v*), we take the guess in *Φ*. Proposition [5](#page-6-3) and Lemma [37](#page-26-3) imply that this is safe. Observe that *Φ* is a constant-size set.

Definition 49 (local docset completion problem)**.** *Let k*,∆ ∈ Z≥¹ *be constants, let t* ∈ *V* (*T*) *be a fixed node of the decomposition tree and let* t_1, \ldots, t_q *denote the children of t (in case t is a leaf, q = 0). Let* ℓ_t *and* u_t *denote the respective restrictions of* ℓ *and u to* $E(G[B_t])$ *. Let* p_t *and* W_t *denote the restrictions of* p *and* W to B_t , redefining $p_t(v) = W_t(i, v) := 0$ for all $v \in X_t$ and all $i \in [k]$. Finally, let $Y_t := X_t \cup \bigcup_{i \in [q]} X_{t_i} \cup R_t$. The local docset completion problem *with respect to node t is as follows: given* $\varphi_t \in \Phi^{Y_t}$ *, find*

$$
\max \left\{ p_t^{\intercal} y : \ell_t(v, w) \le y(v) - y(w) \le u_t(v, w) \; \forall (v, w) \in E(G[B_t]), \; y(v) = \varphi_t(v) \; \forall \, v \in Y_t, \; y \in \mathbb{Z}^{B_t} \right\}.
$$
 (7)

We denote an instance of this problem by $I_t^0 = I_t^0(\varphi_t)$, and define $F_t^0(\varphi_t) := \text{OPT}(I_t^0)$ as the optimum value of [\(7\)](#page-26-5) (if I_t^0 is infeasible, then $\text{OPT}(I_t^0) = -\infty$). For each feasible instance $I_t^0 = I_t^0(\varphi_t)$, we let $y_t^0(\varphi_t) \in \mathbb{Z}^{B_t}$ *denote any optimal solution of* I^0_t *. We call this solution an* optimal local solution.

Notice that for every $t \in V(T)$ and $\varphi_t \in \Phi^{Y_t}$, we can compute $F_t^0(\varphi_t)$ as well as an optimal local solution $y_t^0(\varphi_t)$ (in case the instance $I_0^t(\varphi_t)$ is feasible) in strongly polynomial time, since the constraint matrix of [\(7\)](#page-26-5) is TU. Given a polynomial-size docset superprofile \mathscr{X}_{t} for node t , we can enumerate all relevant guesses $\varphi_t\in \varPhi^{Y_t}$ in polynomial time and consider them one after the other. In this way, we obtain a complete set of local optimal solutions for each $t \in V(T)$.

Moreover, for every optimal solution y^* of [\(IP](#page-26-2)₃) and every $t \in V(T)$, it should be clear that the restriction of y^* to B_t is an optimal solution of the local docset completion problem in which φ_t is the restriction of y^* to Y_t . In other words, we can think of any optimal solution y^* of [\(IP](#page-26-2)₃) as being obtained by composing 8 8 optimal local solutions $y_t^0(\varphi_t)$ for some choice of φ_t for $t\in V(T).$ Our DP finds an optimal way to compose the optimal local solutions in a recursive way, starting at the leaves of the decomposition tree and going up to the root. We formalize this below.

 8 Let *V* be any finite set and *V*₁,...,*V*_l be subsets of *V*. We say that a vector $y \in \mathbb{R}^V$ is obtained by *composing* vectors $y_1 \in$ $\mathbb{R}^{V_1},\ldots,y_\ell\in\mathbb{R}^{V_\ell}$ if the restriction of y to each V_i is exactly $y_i.$ In particular, this implies that the vectors $y_i,$ $i\in[\ell]$ agree on their common coordinates.

The dynamic program

Our DP is based on the concept of a *rooted solution,* which is a vector $y_t \in \mathbb{Z}^{V(G_t)}$ obtained by composing (optimal) local solutions $y_t^0(\varphi_t)$ where t' is a descendant of some fixed node t . Given a guess φ_t^{up} *t* ∈ *Φ Xt* on the variables corresponding to the vertices in the adhesion to the parent, and a target $d_t \in \mathcal{D}$ for the weight vector of the rooted solution, the DP seeks a maximum profit rooted solution *y^t* that *complies* with φ_t^{up} $\frac{dp}{dt}$ and d_t , in the sense that $y_t(v) = \varphi_t^{\text{up}}$ t_t ^{up}(*v*) for all *v* ∈ *X_t* and the weight vector of *y_t* with respect to the vertices of $V(G_t - X_t)$ is coordinate-wise equal to d_t .

Definition 50 (DP for composing optimal local solutions). Let $t \in V(T)$ be a node of the rooted decom*position tree, and let* t_1 , ..., t_q *denote the children of t (where* $q = 0$ *whenever t is a leaf). Let* \mathcal{X}_t *be a polynomial-size docset superprofile with respect to* $R_t = R \cap B_t$ *. For any tuple* $\mathfrak{S} = (S_i)_{i \in [f_5(k, \Delta)]} \in \mathscr{X}^{f_5(k, \Delta)}_t$ $\mathfrak{S} = (S_i)_{i \in [f_5(k, \Delta)]} \in \mathscr{X}^{f_5(k, \Delta)}_t$ $\mathfrak{S} = (S_i)_{i \in [f_5(k, \Delta)]} \in \mathscr{X}^{f_5(k, \Delta)}_t$ $t^{J5(k,\Delta)}$, *we let* $\sigma^{\mathfrak{S}} := \sum_{i=1}^{f_5(k,\Delta)} \chi^{S_i} \in \Phi^{B_i}$ $\sigma^{\mathfrak{S}} := \sum_{i=1}^{f_5(k,\Delta)} \chi^{S_i} \in \Phi^{B_i}$ $\sigma^{\mathfrak{S}} := \sum_{i=1}^{f_5(k,\Delta)} \chi^{S_i} \in \Phi^{B_i}$ denote the sum of the corresponding incidence vectors. For any given *ϕ up* $a_t^{up} \in \Phi^{X_t}$ and $d_t \in \mathcal{D}$, we define

$$
F_t(\varphi_t^{up}, d_t) := \max \Big\{ F_t^0(\varphi_t) + \sum_{i \in [q]} F_{t_i}(\varphi_{t_i}^{up}, d_{t_i}) : \varphi_t \in \Phi^{Y_t}, \varphi_{t_i}^{up} \in \Phi^{X_{t_i}} \ \forall i \in [q], d_{t_i} \in \mathcal{D} \ \forall i \in [q] : \\ \varphi_t(v) = \varphi_t^{up}(v) \ \forall v \in X_t, \ \varphi_{t_i}^{up}(v) = \varphi_t(v) \ \forall v \in X_{t_i} \ \forall i \in [q], \\ \exists \mathfrak{S} \in \mathcal{X}_t^{f_{\mathfrak{S}}(k,\Delta)} : \varphi_t(v) = \sigma^{\mathfrak{S}}(v) \ \forall v \in R_t, \\ W_t \sigma^{\mathfrak{S}} + \sum_{i \in [q]} d_{t_i} = d_t \Big\}.
$$

Furthermore, we define $y_t(\varphi_t^{up})$ *^{<i>up*}, *d*_{*t*}) ∈ $\mathbb{Z}^{V(G_t)}$ *as the rooted solution obtained by composing* $y_t^0(\varphi_t)$ *and* y_{t_i} ($\varphi_{t_i}^{up}$ $\theta_{t_i}^{up}, d_{t_i}$) for $i \in [q]$, where $\varphi_t \in \Phi^{Y_t}$, $\varphi_{t_i}^{up}$ $\mathcal{L}^{up}_{t_i} \in \Phi^{X_{t_i}}$ and $d_{t_i} \in \mathscr{D}$ for $i \in [q]$ achieve the maximum.

Notice that there is a constant number of choices for φ_t^{up} $t_t^{\text{up}} \in \Phi^{X_t}$ and a polynomial number of choices for $d_t \in \mathcal{D}$. All in all, there are a polynomial number of table entries $F_t(\varphi_t^{\text{up}})$ t_t^{up}, d_t) for each $t \in V(T)$. For every φ_t^{up} $\Phi_t^{\text{up}} \in \Phi^{X_t}$ and $d_t \in \mathcal{D}$, we can compute $F_t(\varphi_t^{\text{up}})$ t_t^{up} , d_t) in strongly polynomial time, from the table entries for the children t_1, \ldots, t_q (note that $q \le f_9(s)$ $q \le f_9(s)$ $q \le f_9(s)$) and the polynomially many optimal values $F_t^0(\varphi_t)$, which can all be computed in strongly polynomial time before running the DP.

Lemma 51. For every $t\in V(T)$, $\varphi_t^{up}\in \Phi^{X_t}$ and $d_t\in \mathscr{D}$: (i) the profit of $y_t(\varphi_t^{up},d_t)$ with respect to vertices in *t t t t t t d t t t d t t t t d t t t d t t t d t t t d t t d t t t d t t t d t t t t t t d t t d t t d t t d* θ_t^{up} , d_t), (ii) the weight vector of $y_t(\varphi_t^{up})$ \int_{t}^{up} , *d*_t) *with respect to vertices in* $V(G_t - X_t)$ *is equal to* d_t *, (iii)* $F_t(\varphi_t^{up})$ $\int_t^{\mu p} d_t$) is the maximum profit of a rooted solution complying with $\varphi_t^{\mu p}$ $\int_t^u \theta \, du \, d\theta \, d\theta$

Proof. All three results are easily proved by induction, starting from the leaves of *T*, using the fact that $p_t(v) = W_t(i, v) = 0$ for all $v \in X_t$ and $i \in [k]$. We leave the details to the reader. \Box

We are now in position to prove the main algorithmic result of this paper.

Proof of Theorem [1.](#page-1-2) We showed in Section [2.2](#page-5-4) that any integer program [\(IP\)](#page-0-0) on a totally ∆-modular constraint matrix which becomes the transpose of a network matrix after removing a constant number of rows and columns can be efficiently reduced to an instance of Problem [1](#page-5-2) in which *A* is the transpose of a network matrix. Using the results of Section [4](#page-15-0) we further reduce the problem to polynomially many instances of Problem [3,](#page-26-4) in which the graph is a subdivision of a 3-connected graph whose roots all have degree at least 3.

Consider any instance of Problem [3,](#page-26-4) defined by $I := (p, \ell, u, G, W, d)$, and let OPT denote the optimal value of instance *I*. Observe that, by Lemma [43,](#page-27-2) *G* does not contain any rooted *K*2,4*k*∆+1-model. Letting *s* := 4*k*∆ + 1, we apply Proposition [78](#page-53-1) in order to obtain a $f_9(s)$ $f_9(s)$ $f_9(s)$ -special tree-decomposition (*T*, \mathcal{B}) of *G* and a collection $\{\mathscr{X}_t : t \in V(T)\}$ of docset superprofiles.

As observed before, we can compute the table entry $F_t(\varphi^{\mathrm{up}}_t)$ $\mathcal{L}_t^{\text{up}}, d_t$) for all $t \in V(T)$, φ_t^{up} $t_t^{\text{up}} \in \Phi^{X_t}$ and $d_t \in \mathcal{D}$, in strongly polynomial time. Let t_0 be the root of *T* (remember that $X_{t_0} = \emptyset$), and let ε denotes the null vector in \mathbb{Z}^{\varnothing} . We claim that $y_{t_0}(\varepsilon, d)$ is an optimal solution of *I* whenever it is defined, and *I* is infeasible whenever $y_{t_0}(\varepsilon,d)$ is not defined. We split the proof in two parts: (i) $y_{t_0}(\varepsilon,d)$ is feasible for [\(IP](#page-26-2)₃) whenever it is defined, (ii) $F_{t_0}(\varepsilon, d) = \text{OPT}.$

For (i), assume that $y_{t_0}(\varepsilon, d)$ is defined, that is, $F_{t_0}(\varepsilon, d) > -\infty$. For simplicity, let $y := y_{t_0}(\varepsilon, d)$.

Observe that *y* is obtained by composing optimal local solutions. Hence, there exist $\varphi_t \in \Phi^{Y_t}$ for each *t* ∈ *V*(*T*) such that *y* is obtained by composing $y_t^0(\varphi_t)$ for *t* ∈ *V*(*T*). Each optimal local solution $y_t^0(\varphi_t)$ is the restriction of *y* to B_t . Moreover, each $y_t^0(\varphi_t)$ is in particular feasible for the corresponding instance of the local docset completion problem. Since each edge of G is contained in B_t for at least one *t* ∈ *V*(*T*) by Definition [46](#page-29-1) (ii), we see that $\ell(v, w) \leq \gamma(v) - \gamma(w) \leq u(v, w)$ for all $(v, w) \in E(G)$. By Lemma [51.](#page-31-0)(ii), $Wy = d$. We conclude that *y* is a feasible solution of [\(IP](#page-26-2)₃). Using Lemma 51.(i), this $implies$ OPT $\geq F_{t_0}(\varepsilon, d)$.

For (ii), it remains to prove that $F_{t_0}(\varepsilon, d) \geq \text{OPT}$. Without loss of generality, assume that [\(IP](#page-26-2)₃) is feasible and consider an optimal solution *y*¯ that is the sum of at most *f*[5](#page-6-3)(*k*,∆) incidence vectors of docsets of *G*. For each $t \in V(T)$, let $\bar{\varphi}_t \in \Phi^{Y_t}$ denote the restriction of \bar{y} to Y_t , let $\bar{\varphi}_t^{\text{up}}$ $t_t^{\text{up}} \in \Phi^{X_t}$ similarly denote the restriction of \bar{y} to X_t , and let \bar{d}_t denote the weight vector of \bar{y} restricted to the nodes of $V(G_t - X_t)$. Notice that, for every fixed node *t*, redefining $\bar{y}(v) := y_t^0(\bar{\varphi}_t)$ for $v \in B_t$ preserves both the feasibility and the optimality of \bar{y} . Now \bar{y} is obtained by composing optimal local solutions, hence \bar{y} is a rooted solution with respect to $t = t_0$ that complies with ε and d . By Lemma [51,](#page-31-0) we conclude that $F_{t_0}(\varepsilon, d)$ is at least the profit of \bar{y} , that is, at least OPT.

 \Box

6 Structure of graphs without a rooted *K*2,*^t***-minor**

This section has two main goals.

First, we prove our decomposition theorem, see Theorem [2](#page-3-1) in Section [1,](#page-0-1) or Theorem [52](#page-33-0) below for a precise statement. Roughly, our decomposition theorem states that every 3-connected rooted graph (G, R) without a rooted $K_{2,t}$ -minor has a tree-decomposition (T, \mathcal{B}) in which every bag has a "nice structure".

Second, we prove that given such a tree-decomposition, it is possible to find in polynomial time a collection $\mathcal{X} = \{\mathcal{X}_u \mid u \in V(T)\}\$ of subsets of vertices of *G* where each \mathcal{X}_u is a docset superprofile of $R \cap B_u$, of polynomial size. This is the content of Theorem [9.](#page-9-0)

Before stating Theorem [52,](#page-33-0) we need to introduce some terminology and notations about graphs embedded in surfaces. A *surface* Σ is a non-empty compact connected Hausdorff topological space in which every point has a neighborhood that is homeomorphic to the plane [\[MT01\]](#page-57-11). Let $\Sigma(h, c)$ denote the surface obtained by removing 2*h*+*c* open disks with disjoint closures from the sphere and gluing *h* cylinders and *c* Möbius strips onto the boundaries of these disks. As stated by the classification theorem of surfaces, every surface is topologically equivalent to Σ(*h*,*c*), for some *h* and *c*. The *Euler genus* eg(Σ) of a surface $\Sigma \cong \Sigma(h, c)$ is $2h + c$. A *curve* in Σ is the image of a continuous function $f : [0,1] \to \Sigma$. The curve is *simple* if *f* is injective. The curve $e = f([0,1])$ *connects* the endpoints $f(0)$ and $f(1)$. We say that a graph *H* is *embedded* in a surface Σ if the vertices of *H* are distinct points in Σ and every edge $uv \in E(H)$ is a simple curve in Σ which connects in Σ the points which are the images of *u* and *v*, such that its interior is disjoint from other edges and vertices. We also condider embeddings in a disk, and the definitions above extend from surfaces to a disk. For simplicity, we sometimes identify an embedded graph with the corresponding subset of the surface when no confusion can occur.

Consider a graph *H* embedded in Σ. The *faces* of the embedded graph *H* are the regions of Σ – H^9 H^9 . A curve *C* in Σ is said to be *H-normal* if it intersects *H* only at vertices. The *distance in* Σ between two points $x, y \in \Sigma$ is the minimum of $|C \cap V(H)|$ over all *H*-normal curves *C* linking *x* to *y*. The distance in Σ between two vertex subsets *A*,*B* of *H* is the minimum distance in Σ between a vertex of *A* and a vertex of *B*. A *noose* is a simple, closed, *H*-normal and non-contractible curve in Σ. For a surface Σ that is not a sphere, the *facewidth* (also known as *representativity*) of the embedding of *H* in Σ is the minimum of $|C \cap V(H)|$ over all nooses *C*. (The facewidth of any graph embedded on the sphere is set to be infinity.)

Let (T, \mathcal{B}) be a tree-decomposition of a graph *G*. Recall that the tree *T* is assumed to be rooted at some node of *T*. With respect to this root, the *weak torso* of a node $u \in V(T)$ is the graph $G[B_u]$ where we add edges between all (previously non-adjacent) pairs of vertices in each of the adhesion sets of $B_u \cap B_v$ where ν is a child of u . We denote the weak torso of u as $G^\#[B_u].$

Theorem 52. *For every integer* $t \ge 1$ *, there exists an integer* $\ell = \ell(t) \ge 1$ *such that the following holds. Let* (G, R) *be a* 3*-connected rooted graph without a rooted* $K_{2,t}$ *-minor. Then G has a tree-decomposition* (*T*, \mathcal{B}) *with the following properties.*

 (i) *the adhesion of* (T, \mathcal{B}) *is at most* ℓ *, and*

(ii) for every node $u \in V(T)$ *, all, but at most* ℓ *, children of* u *are leaves* v *with* $B_v \cap R \subseteq B_u$ *, and*

- *(iii) for every node* $u \in V(T)$ *, at least one of the following is true:*
	- (a) $|B_u| \leq \ell$,
	- *(b) u* is a leaf of T and $B_u \cap R \subseteq B_{u'}$, where u' is the parent of u in T, or
	- *(c) there is a set* $Z \subseteq B_u$ *such that* $|Z| \le \ell$, $G^#[B_u] Z$ *is* 3*-connected, does not contain a rooted* $K_{2,t}$ *minor with respect to the set of roots B^u* ∩ *R* − *Z , and has an embedding in a surface of Euler genus at most ℓ such that every face is bounded by a cycle of the graph, and all the vertices in* $B_u \cap R - Z$ *can be covered using at most* ℓ *facial cycles.*

Furthermore, for fixed t, the above tree-decomposition and the embeddings described in [\(c\)](#page-33-2) can be found in polynomial time given (*G*,*R*) *in input.*

We restate Theorem [9](#page-9-0) here, for the reader's convenience.

Theorem [9](#page-9-0). *There exists a function* $f_9: \mathbb{Z}_{\geq 1} \to \mathbb{Z}_{\geq 1}$ *such that for every fixed* $t \in \mathbb{Z}_{\geq 1}$ *, there is a polynomialtime algorithm that given a rooted* 3*-connected graph* (*G*,*R*) *having no rooted K*2,*t-minor, outputs an f*_{[9](#page-9-0)}(*t*)*-special tree-decomposition* (*T*, \mathcal{B}) *of G and a collection* { \mathcal{X}_u : $u \in V(T)$ } *where each* \mathcal{X}_u *is a docset superprofile of* $R \cap B_u$ *in G of size polynomial in* $|V(G)|$ *.*

Recall that a tree-decomposition (*T*, \mathscr{B}) of a graph *G* is ℓ -special for some integer $\ell \geq 1$ if $|B_u \cap B_v| \leq \ell$ for each edge $uv \in E(T)$, and every node $u \in V(T)$ has at most ℓ children.

6.1 Building the special tree-decomposition

A *star-decomposition* of a graph *H* is a tree-decomposition *H* where the tree is a star. Such a treedecomposition is denoted $(B_0; B_1, \ldots, B_m)$, where B_0 is the center bag, and B_1, \ldots, B_m are the leaf bags.

For a fixed rooted graph (*G*,*R*) without a rooted *K*2,*t*-minor, we construct the tree-decomposition of *G* as in Theorem [52](#page-33-0) recursively, one node at a time.

For any subgraph $H \subseteq G$, we denote by $\partial_G(H)$ the set of all vertices of *H* that are adjacent in *G* to a vertex not in *H*. When the graph *G* is clear from the context, we drop the subscript *G*.

Given a subgraph $H \subseteq G$, we construct a tree-decomposition of *H* recursively. Firstly, we find a stardecomposition $(B_0; B_1, \ldots, B_m)$ of *H* and secondly we find the subtrees attached to the root bag B_0 by

⁹ that is, the equivalence classes of the equivalence relation where two points of Σ−*H* are equivalent if there is a simple curve in Σ− *H* connecting them

applying recursion to the subgraphs H_1, \ldots, H_m , where $H_i = H[B_i]$. To do so, we need to introduce some definitions and technical lemmas.

A set of vertices *X* ⊆ *V*(*H*) is *well-linked* in a graph *H* if for every *X*₁, *X*₂ ⊆ *X* there exist min{ $|X_1|, |X_2|$ } vertex-disjoint *X*₁–*X*₂ paths in *H*. Given a rooted graph (*G*, *R*) and an integer $k \ge 1$, a subgraph $H \subseteq G$ is said to be *k*-interesting if there exists a set $X \subseteq V(H)$ such that

• $|X| = k$,

- *∂*(*H*) ⊆ *X*,
- *X* is well-linked in *H*, and

• there exist *k* vertex-disjoint R_H –*X* paths in *H* where $R_H = R \cap V(H)$.

Such a set *X* is called a *core* of *H*. In particular, for every *k*-interesting subgraph *H*, we have $|\partial(H)| \leq k$. A subgraph *H* ⊆ *G* is *k*-*boring* if $|\partial(H)|$ ≤ *k* and *H* is not *k*-interesting.

We will prove Theorem [52](#page-33-0) using the following two lemmas.

Lemma 53. *Let* $k ≥ 1$ *be an integer, let* (G, R) *be a rooted graph, and let* $H ⊆ G$ *be a k-boring subgraph. Then there exists a star-decomposition* $(B_0; B_1, B_2)$ *of H with* $\partial(H) \subseteq B_0$ *such that*

 (i) $|B_0| \leq 2k$,

(ii) $B_i \neq V(H)$ *for i* $\in \{1,2\}$ *.*

 (iii) | $B_0 ∩ B_1$ | ≤ *k*, and

(iv) either $|B_0 \cap B_2|$ ≤ *k, or* $B_2 \cap R$ ⊆ B_0 *.*

Furthermore, for fixed t, the star-decomposition can be found in polynomial time given (*G*,*R*) *in input.*

Lemma 54. For every integer $t \geq 1$, there exist integers $k = k(t)$, $k' = k'(t) \geq 1$ such that for every 3*connected rooted graph* (G, R) *without a rooted* $K_{2,t}$ *-minor, and every k-interesting subgraph* $H \subseteq G$ *with a* core X, there exists a star-decomposition $(B_0; B_1, \ldots, B_m)$ of H such that $X \subseteq B_0$, for every $i \in \{1, \ldots, m\}$, $I(we \text{ } h \text{ } a \text{ } we \text{ } |B_0 \cap B_i| \leq k, B_i \neq V(H)$, and if $i > k'$, then $B_i \cap R \subseteq B_0$, and there is a set $U \subseteq B_0$ such that

- (i) $|U| \leq k'$ *, and*
- (*ii*) $(H^* [B_0] - U, R \cap B_0 - U)$ *is* 3*-connected, has no rooted K*_{2,*t*}*-minor, and admits an embedding in a surface with Euler genus at most k such that every face is bounded by a cycle, and the set* $R \cap B_0 - U$ *can be covered using at most k*′ *facial cycles.*

Furthermore, for fixed t, the star-decomposition and the embedding described in [\(ii\)](#page-34-0) can be found in polynomial time given (*G*,*R*) *in input.*

Before proving these two lemmas, let us show that they easily imply Theorem [52.](#page-33-0)

Proof of Theorem [52.](#page-33-0) Fix an integer $t \ge 1$, and let $k = k(t)$ be as in Lemma [54.](#page-34-1) We prove the theorem for $l = 2k$. Fix a 3-connected rooted graph (*G*, *R*) that does not contain a rooted $K_{2,t}$ -minor.

We will prove the following claim: For every subgraph $H \subseteq G$ with $|\partial(H)| \leq k$, there exists a treedecomposition (*T*, \mathcal{B}) of *H* such that $\partial(H)$ is contained in the root bag, and

- (i) the adhesion of (T,\mathscr{B}) is at most ℓ , and
- (ii) for every node $u \in V(T)$, all but at most ℓ children of u are leaves v with $B_v \cap R \subseteq B_u$, and
- (iii) for every node $u \in V(T)$, at least one of the following is true:
	- $|B_u| \leq \ell$,
	- (b) *u* is a leaf of *T* and $B_u \cap R \subseteq B_{u'}$, where *u'* is the parent of *u* in *T*, or
	- (c) there is a set $Z \subseteq B_u$ such that $|Z| \le \ell$, $H^* [B_u] Z$ is 3-connected, does not contain a rooted *K*_{2,*t*}-minor with respect to the set of roots $B_u \cap R - Z$, and has an embedding in a surface of Euler genus at most *ℓ* such that every face is bounded by a cycle of the graph, and all the vertices in $B_u \cap R - Z$ can be covered using at most ℓ facial cycles.

Since $\partial_G(G) = \emptyset$, the theorem follows from the claim with *H* = *G*.

We prove the claim by induction on $|V(H)|$. If $|V(H)| = 0$, the theorem is satisfied by a tree-decomposition with only one, empty bag. For the induction step suppose that $|V(H)| \ge 1$, and let $(B_0; B_1, \ldots, B_m)$ be a

star-decomposition of *H* obtained by applying Lemma [53](#page-34-2) or Lemma [54,](#page-34-1) depending on whether *H* is *k*boring or *k*-interesting. Let $H_i \coloneqq H[B_i]$ for each $i \in \{0, 1, ..., m\}$. Note that we can remove from B_0 the set $∪_{i \in [m]}(B_0 ∩ B_i - ∂(H_i))$ while preserving the properties of the star-decomposition. This way we get that $\partial(H_i) = B_0 \cap B_i$ for each $i \in [m]$, since $\partial(H) \subseteq B_0$, and thus either $|\partial(H_i)| = |B_0 \cap B_i| \leq 2k$ and $B_i \cap R \subseteq B_0$, or $|\partial(H_i)| = |B_0 \cap B_i| \le k$. For each $i \in [m]$ with $|\partial(H_i)| = |B_0 \cap B_i| \le k$, apply the induction hypothesis to $H_i = H[B_i]$ to obtain a tree-decomposition (T_i, \mathcal{B}_i) , and let (T, \mathcal{B}) be the tree-decomposition of *H* obtained by the disjoint union of these tree decompositions by adding a new root node *u* whose children are the roots of T_1, \ldots, T_m , and whose bag is the center bag B_0 of the star-decomposition. It is easily checked that the resulting tree-decomposition of *H* satisfies the claim. \Box

A *separation* in a graph *G* is an ordered pair (*A*,*B*) of vertex subsets of *G* such that *A* ∪*B* = *V* (*G*) and there are no edges with one endpoint in $A \setminus B$ and the other in $B \setminus A$. The *order* of a separation is $|A \cap B|$. We conclude this subsection with the proof of Lemma [53.](#page-34-2)

Proof of Lemma [53.](#page-34-2) Let *H* be a *k*-boring subgraph of (G, R) . If $|V(H)| \leq 2k$, then the lemma is satisfied by the star-decomposition $(V(H); \phi, \phi)$. So we assume that $|V(H)| \geq 2k + 1$. Let *X* be any set with $\partial(H) \subseteq$ $X \subseteq V(H)$ and $|X| = k$.

Suppose that *X* is not well-linked in *H*, and let $X_1, X_2 \subseteq X$ be such that there are no min{ $|X_1|, |X_2|$ } vertex-disjoint X_1 – X_2 paths in *H*. By Menger's Theorem, there exists a separation (B_1 , B_2) of *H* with *X*^{*i*} ⊆ *B*^{*i*} for *i* ∈ {1, 2} and |*B*₁ ∩ *B*₂| < min{|*X*₁|, |*X*₂|}. Let *B*₀ = *X* ∪ (*B*₁ ∩ *B*₂), and observe that

• $|B_0| \leq |X| + |B_1 \cap B_2| < 2k$, and

• for *i* ∈ {1, 2}, $|B_0 \cap B_i|$ ≤ $|X \setminus X_{3-i}|$ + $|B_1 \cap B_2|$ < $|X \setminus X_{3-i}|$ + $|X_{3-i}|$ ≤ $|X|$ = k ,

so $(B_0; B_1, B_2)$ satisfies the lemma.

Now suppose that *X* is well-linked in *H*. Since *H* is *k*-boring, this implies that there are no *k* vertexdisjoint R_H –*X* paths in *H*, where $R_H = R \cap V(H)$. By Menger's theorem, there exists a separation (B_1, B_2) of *H* with $R_H \subseteq B_1$, $X \subseteq B_2$, and $|B_1 \cap B_2| < k$. Let $B_0 = X \cup (B_1 \cap B_2)$, and observe that

- $|B_0| \le |X| + |B_1 \cap B_2| < 2k$, and
- $|B_0 \cap B_1| = |B_1 \cap B_2| < k$, and
- $B_2 \cap R = B_2 \cap R_H \subseteq B_1 \cap B_2 \subseteq B_0$.

Thus, $(B_0; B_1, B_2)$ satisfies the lemma.

Finally, we remark that in both cases above, the resulting star-decomposition of *H* is easily computed in polynomial time. \Box

The following subsections are devoted to the proof of Lemma [54.](#page-34-1)

6.2 Finding a large wall

Let (G, R) be a rooted graph and let $k \geq 1$ be an integer. Suppose that *H* is a *k*-interesting subgraph of (G, R) and let $X \subseteq V(H)$ be a core of *H*. Thus, $|X| = k$, $\partial(H) \subseteq X$, *X* is well-linked in *H*, and there are *k* vertex-disjoint paths between $R_H = R \cap V(H)$ and *X* in *H*. The goal of this section is to show that *H* must contain a large *wall* as a subgraph and that there are many vertex-disjoint paths from *R^H* to any *transversal* set of vertices in the wall. We proceed with the necessary definitions.

Given an integer $h \ge 2$, an *elementary wall of height* h is the graph obtained from the $2h \times h$ grid with vertex set $[2h] \times [h]$ by removing all edges with endpoints $(2i - 1, 2j - 1)$ and $(2i - 1, 2j)$ for all $i \in [h]$ and $j \in [\lfloor h/2 \rfloor]$, all edges with endpoints $(2i,2j)$ and $(2i,2j+1)$ for all $i \in [h]$ and $j \in [\lfloor (h-1)/2 \rfloor]$, and removing the two vertices of degree at most 1 in the resulting graph.

In an elementary wall*W* of height *h*, there is a unique set of *h* vertex-disjoint paths linking the bottom row (vertices of the form $(i,1)$) to the top row (vertices of the form (i,h)). These paths are called the *vertical paths* of *W*. We enumerate the vertical paths as Q_1, \ldots, Q_h so that the first coordinates of their vertices are increasing. There is also a unique set of h vertex-disjoint paths linking Q_1 to Q_h , called the *horizontal paths* of *W* .

A subdivision of an elementary wall of height *h* is called a *wall of height h*. Vertical paths, and horizontal paths of a wall are defined as expected, as the subdivided version of their counterparts in the elementary wall.

Let *W* be a wall. Let*U* be the set of vertices of degree 3 in *W* , we call the vertices in*U branch vertices*. A *transversal set of vertices in W* is an inclusion-wise maximal subset of *U* such that no two vertices are in the same row or column of *W* . Observe that such a set necessarily has size *h*, where *h* is the height of *W* . One way to obtain a transversal set in *W* is to choose all the vertices on the "diagonal" of *W* .

Lemma [55](#page-36-0). *For every integer* $h \ge 1$ *there is an integer* $k = k_{55}(h) ≥ 1$ *so that the following holds. For every rooted graph* (*G*,*R*)*, every k-interesting subgraph H of G, and every core X of H, there exists a wall W of height h in H such that, for every transversal set Y of W , there are h vertex-disjoint X –Y paths and h vertex-disjoint* R_H -*Y* paths in *H*, where $R_H = R \cap V(H)$.

For the proof of the lemma we need a few definitions and a lemma from [\[KTW20\]](#page-57-9). Let *s* ∈ N. An *s*-tangle in a graph *G* is a collection $\mathcal T$ of (ordered) separations of *G* such that,

- for every separation (*A*, *B*) of *G* of order at most $s 1$, exactly one of (*A*, *B*), (*B*, *A*) is included in \mathcal{T} , and
- \bullet there are no three separations (A_1, B_1) , (A_2, B_2) , $(A_3, B_3) \in \mathcal{F}$ with $G[A_1] \cup G[A_2] \cup G[A_3] = G$.

Let *W* be a wall of height *h* in a graph *G*. The tangle \mathcal{T}_W induced by *W* is the collection of separations (A, B) of *G* of order at most $h - 1$ such that $B \setminus A$ contains the vertex set of both a horizontal and vertical path of W . A tangle \mathcal{T}' is a *restriction* of \mathcal{T} if $\mathcal{T}' \subseteq \mathcal{T}$.

Lemma 56 (Lemma 14.6 in [\[KTW20\]](#page-57-9))**.** *Let fwall be a function such that for every positive integer s, every graph with tree-width at least fwall*(*s*) *contains an s-wall as a subgraph. Let h* ≥ 1 *be an integer. Let G be a* graph and let $\mathcal T$ be a tangle of order at least 3 f_{wall} (6 h^2) + 1 *in G. Then there exists a wall W of height h in G* such that \mathcal{T}_W *is a restriction of* \mathcal{T} *.*

Note that such a function *f*wall exists by the Grid Minor Theorem of Robertson and Seymour [\[RS86\]](#page-58-13) (see [\[CT21\]](#page-56-12) for the current best known bound).

We may now turn to the proof of Lemma [55.](#page-36-0)

Proof of Lemma [55.](#page-36-0) Let $h \ge 1$ and let $k = 6 \cdot (3f_{wall}(6h^2) + 1)$. Let *H* be a *k*-interesting subgraph of a rooted graph (G, R) . Let $X \subseteq V(H)$ be a core of *H*. We define a $k/6$ -tangle \mathcal{T}_X in *H* with respect to *X*: For every separation (*A*, *B*) in *H* of order at most ($k/6-1$) we add either (*A*, *B*) or (*B*, *A*) to \mathcal{T}_X as follows. Since *X* is well-linked in *H*, either *A* \ *B* or *B* \ *A* contains at most $k/6 - 1$ elements from *X*. We add (*A*, *B*) to \mathcal{T}_X if $|B \cap X|$ ≥ $k/6$, otherwise we add (B, A) to \mathcal{T}_X . Note that for each separation $(A, B) \in \mathcal{T}_X$, $|A \cap X|$ ≤ $k/3-2$, and thus it is easily checked that \mathcal{T}_X is a tangle.

By Lemma [56,](#page-36-1) there is a wall *W* of height *h* in *H* such that the tangle \mathcal{T}_W induced by *W* is a restriction of \mathcal{T}_X .

Let *Y* be some transversal set of *W*. Thus, $|Y| = h$. First, suppose that there are no *h* vertex-disjoint *X*–*Y* paths in *H*. Then, by Menger's Theorem, there is a separation (*A*,*B*) in *H* such that $|A \cap B| \le h - 1$, *Y* ⊆ *A* and *X* ⊆ *B*. Note that $h \le k/6$ and hence $(A, B) \in \mathcal{T}_X$. Since $|A \cap B| \le h - 1$, there is an horizontal path of *W* and a vertical path of *W* that have an empty intersection with *A* ∩*B*. Each of these two paths contains some vertex in *Y* (otherwise we get a contradiction to the maximality of *Y* in the definition of a transversal in a wall), and thus they must be contained in $A \setminus B$. Hence, $(B, A) \in \mathcal{T}_W$. But this contradicts the fact that \mathcal{T}_W is a restriction of \mathcal{T}_X . Therefore, we conclude that there are *h* vertex-disjoint *X*–*Y* paths in *H*.

Assume now that there are no *h* vertex-disjoint R_H –*Y* paths in *H*. Then, similarly as before, let (A, B) be a separation in *H* such that $|A \cap B|$ ≤ h − 1, $Y \subseteq A$ and R ^{*H*} ≤ *B*. For the same reason as before, we must have $(B, A) \in \mathcal{T}_W$, and thus also $(B, A) \in \mathcal{T}_X$ since \mathcal{T}_W is a restriction of \mathcal{T}_X . By the definition of \mathcal{T}_X , this implies that $|(B \setminus A) \cap X| \leq k/6 - 1$. Since we also have $|A \cap B| \leq h - 1 \leq k/6 - 1$, it follows that $((B \setminus A) \cap X) \cup (A \cap B)$ has size at most $k/3 - 2 < k$ and meets all R_H –X paths in *H*, which contradicts the existence of k vertex-disjoint R ^{H} – X paths in H . Therefore, we conclude that there are h vertex-disjoint R_H –*Y* paths in *H*, as desired. \Box

6.3 Excluding a complete graph minor grasped by a wall

Given a wall W in G, and an H-model M in G for a graph H with $|V(H)| = s$, we say that the wall W grasps the model *M* if for every $v \in V(H)$ there are *s* distinct horizontal paths R_1^v, \ldots, R_s^v of *W* and *s* distinct vertical paths Q_1^v, \ldots, Q_s^v of *W* such that $V(R_i^v) \cap V(Q_i^v) \subseteq M(v)$ for each $i \in [s]$.

Lemma 57. *Let h*,*t* ∈ N *be such that h* ≥ 4*t* + 2*. Let* (*G*,*R*) *be a rooted graph with* |*R*| ≥ *h and having no rooted K*2,*t-minor. Assume that G contains an h-wall W such that, for every transversal set Y of W , there are h vertex-disjoint R–Y paths in G. Then G does not contain a* K_{4t+2} *-model grasped by the wall W.*

Proof. Assume to the contrary that *G* contains a *K*4*t*+2-model *M* grasped by the wall *W* . First, we show that one can find a set *Y* ′ of 2*t* branch vertices of *W* contained in distinct branch sets of *M* and such that each horizontal path of *W* and each vertical path of *W* contains at most one vertex in *Y* ′ . This can be seen as follows: Let $s = 4t + 2$, and consider 2*t* distinct vertices v_1, \ldots, v_{2t} of K_{4t+2} . For $j = 1, 2, \ldots, 2t$, consider *s* distinct horizontal paths $R_1^{\nu_j}$ $\sum_{1}^{v_j}$,...,*R*^v of *W* and *s* distinct vertical paths $Q_1^{\overline{v}_j}$ $\overline{Q}_1^{\overline{\nu}_j}$,..., $\overline{Q}_s^{\nu_j}$ of *W* such that $V(R_i^{v_j})$ $\binom{v_j}{i}$ ∩*V*($Q_i^{v_j}$ *^v*</sup>^{*j*} ⊆ *M*(*v*_{*j*}) for each *i* ∈ [*s*]. Since at most 2*j* − 2 of these paths were marked previously, we can find $i \in [s]$ such that neither R^{ν_j} nor Q^{ν_j} is marked. We mark these two paths and choose a branch vertex of *W* in $V(R_i^{v_j})$ $\binom{v_j}{i}$ ∩ $V(Q_i^{v_j})$ v_j). Finally, we let *Y*' be the set of chosen branch vertices.

Next, extend the set *Y'* to a transversal set *Y* in *W*. By our assumptions, there are *h* vertex-disjoint R –*Y* paths in *G*, and thus in particular there are 2*t* vertex-disjoint R –*Y'* paths in *G*.

In what follows, we call *feasible path system* a collection of 2*t* vertex-disjoint paths in *G* such that all the paths have one endpoint (called *source*) in *R*, and their other endpoint (called *destination*) in distinct branch sets of *M*. Note that there is at least one feasible path system in *G*, since there are 2*t* vertexdisjoint *R*–*Y* ′ paths in *G*. A path *P* in a feasible path system P is *good* if the branch set of *M* containing the destination of P avoids all the other paths in \mathcal{P} , and is *bad* otherwise. With respect to \mathcal{P} , we classify branch sets of *M* into four categories: A branch set *B* of *M* is

- *good* if *B* contains the destination of a *good* path of \mathcal{P} ;
- *bad* if *B* contains the destination of a *bad* path of \mathcal{P} ;
- *ugly* if *B* intersects some path in \mathcal{P} but *B* contains no destination of paths in \mathcal{P} ;
- *free* if *B* intersects no path in \mathcal{P} .

We let $g(\mathcal{P}), b(\mathcal{P}), u(\mathcal{P}), f(\mathcal{P})$ denote the number of branch sets in *M* that are respectively good, bad, ugly, and free w.r.t. \mathscr{P} . Finally, we let *i*(\mathscr{P}) denote the sum over every path $P \in \mathscr{P}$ of the number of branch sets of *M* intersected by *P*. (The letter *i* stands for 'intersection'.)

To each feasible path system P in *G* we associate a corresponding *vector* with two entries, defined as follows:

$$
(i(\mathscr{P}) + b(\mathscr{P}), b(\mathscr{P})). \tag{8}
$$

Now, choose a feasible path system $\mathcal P$ in *G* whose vector is lexicographically minimum. Let us make a few observations about $\mathcal P$. First, we claim that

$$
u(\mathscr{P}) \le g(\mathscr{P}).\tag{9}
$$

To see this, observe first that no bad path *P* of $\mathcal P$ intersects an ugly branch set *B* of *M*, since otherwise we could shorten *P* so that it ends in *B*, which would improve the vector of \mathcal{P} . Thus, ugly branch sets of *M* only intersects good paths of \mathcal{P} . If a good path *P* of \mathcal{P} intersects two ugly branch sets B_1 , B_2 of *M* in this order from its source, then one could shorten *P* so that it ends in B_1 . This possibly increases $b(\mathcal{P})$ by 1 (since *P* might become bad) but decreases $i(\mathcal{P})$ by at least 2 (since *P* no longer intersects B_2 nor the good branch set that contained its destination), thus resulting in a feasible path system with a better vector, a contradiction. Hence, every good path intersects at most one ugly branch set, and [\(9\)](#page-37-0) follows.

Observe that

$$
f(\mathscr{P}) \ge 2\tag{10}
$$

since $b(\mathcal{P}) + g(\mathcal{P}) + u(\mathcal{P}) = 2t + u(\mathcal{P}) \le 2t + g(\mathcal{P}) \le 4t$ by [\(9\)](#page-37-0).

Next, we claim that

$$
b(\mathcal{P}) \le g(\mathcal{P}).\tag{11}
$$

To show this, we are going to associate to each bad branch set *B* of *M* a corresponding good path of \mathcal{P} , in an injective manner. Since the number of good paths equals to the number of good branch sets, this will imply [\(11\)](#page-38-0). So, suppose *B* is a bad branch set of *M*. Let *F* be a free branch set of *M*, which exists by [\(10\)](#page-38-1), and let vw be an edge linking *B* and *F* in *G*, with v in *B*. Consider a spanning tree *T* of *G*[*B*] and let *P* be the path in $\mathscr P$ closest to *v* in the tree *T*. (Possibly *v* is in *P*.) If *P* is bad, then one could reroute *P* towards *v* in the tree *B* and use the edge *vw* so that it ends in *F*. Then *P* would become good, so $b(\mathcal{P})$ would decrease by 1, and $i(\mathcal{P})$ would increase by at most 1 (due to the intersection of *P* with *F*). Overall, this would result in a better vector for \mathcal{P} , which is a contradiction. Thus *P* must be good. Moreover, if *B* is not the last bad branch set intersected by *P* on its way to its destination, then performing the same rerouting gives again a contradiction, so *B* is the last such branch set. Hence, we may associate *B* to the good path *P*, and this way *P* will not be chosen by any other bad branch set, as desired. This shows [\(11\)](#page-38-0).

Since $b(\mathcal{P}) + g(\mathcal{P}) = 2t$, it follows from [\(11\)](#page-38-0) that one can find t good branch sets in M. Using the union of these branch sets with the corresponding good paths in $\mathcal P$ plus two additional free branch sets of *M* (which exist by [\(10\)](#page-38-1)), we find a rooted *K*2,*t*-model in (*G*,*R*). This final contradiction concludes the proof. \Box

6.4 The structure of graphs excluding a complete graph minor grasped by a wall

We combine results in [\[KTW20\]](#page-57-9) and [\[DiKMW12\]](#page-56-10) to derive a theorem that describes the structure of graphs without a complete graph minor grasped by a wall. Before we do so, we introduce the necessary definitions.

A *vortex* is a pair $V = (J, \Omega)$ where *J* is a graph and $\Omega = \Omega(V)$ is a linearly ordered set (u_1, u_2, \ldots, u_d) of a subset of vertices of *J*. With a slight abuse of notation, we will denote the unordered set of vertices $\{u_1, u_2, \ldots, u_d\}$ by Ω as well.

Given a vortex $V = (J, \Omega)$ and a vertex subset *Z*, we denote by $V - Z$ the vortex obtained by deleting the vertices in *Z* from *J* and Ω . If V is a set of vortices, we let $V - Z$ denote the set of vortices $V - Z$ with *V* ∈ $\mathcal V$ such that *V* − *Z* has at least one vertex.

Let Σ be a surface and let *D* be a closed disk in the surface. We denote by $bd(D)$ the boundary of *D*.

Let $V = (J, \Omega)$ be a vortex with $\Omega = (u_1, u_2, \dots, u_q)$. A *linear decomposition* of *V* is a collection of sets (X_1, X_2, \ldots, X_d) such that

• for each $i \in [q]$, $X_i \subseteq V(J)$ and $u_i \in X_i$, and moreover $\cup_{i=1}^q X_i = V(J)$;

- for every $uv \in E(J)$, there exists $i \in [q]$ such that $\{u, v\} \subseteq X_i$, and
- for every $x \in V(J)$, the set $\{i : x \in X_i\}$ is an interval in [q].

The *adhesion* of the linear decomposition is $max(|X_i \cap X_{i+1}| : 1 \le i \le q-1)$.

Let $\alpha_0, \alpha_1, \alpha_2 \in \mathbb{N}$, and let Σ be a surface. A graph *G* is $(\alpha_0, \alpha_1, \alpha_2)$ *-nearly embeddable in* Σ if there exist $Z \subseteq V(G)$ with $|Z| \le \alpha_0$ and an integer $\alpha' \le \alpha_1$ such that $G - Z$ can be written as the union of $p + 1$ edge-disjoint graphs G_0, G_1, \ldots, G_p with the following properties:

- (i) For each $i \in [p]$, $V_i := (G_i, \Omega_i)$ is a vortex where the set Ω_i is $V(G_i \cap G_0)$ (its linear ordering is speci-fied in [\(iv\)](#page-39-0) below). For all $1 \le i < j \le p$, $G_i \cap G_j \subseteq G_0$.
- (ii) The vortices $V_1, \ldots, V_{\alpha'}$ have a linear decomposition of adhesion at most α_2 . Let V be the collection of those vortices.
- (iii) The vortices $V_{\alpha'+1},...,V_p$ satisfy $|\Omega_i| \leq 3$ for each $\alpha'+1 \leq i \leq p$. Let W be the collection of those vortices.
- (iv) There are closed disks $D_1, ..., D_p$ in Σ with disjoint interiors and an embedding $\sigma : G_0 \hookrightarrow \Sigma$ – $\cup_{i=1}^p$ int(*Di*) such that $\sigma(G_0) \cap \text{bd}(D_i) = \sigma(\Omega_i)$ for each $i \in [p]$ and the linear ordering of Ω_i is compatible with the cyclic ordering of $\sigma(\Omega_i)$.

We call the tuple (σ, G_0, Z, V, W) an $(\alpha_0, \alpha_1, \alpha_2)$ *-near embedding* of *G*. We will denote by $D(V_i)$ the disk D_i corresponding to vortex V_i in the above definition. We call vortices in $\mathcal V$ *large vortices* and vortices in $\mathcal W$ *small vortices.* Given *α* ∈ ℕ, a tuple $(σ, G_0, Z, V, W)$ is said to be an *α-near embedding* if it is an $(α, α, α)$ near embedding. We note that if the constants are clear from the context, we sometimes omit them and simply write *near embedding*.

Given a near embedding (σ , G_0 , Z , V , W) of a graph *G* in a surface Σ, we define a corresponding graph G'_{0} C_0' , obtained from G_0 by adding an edge \it{vw} for every pair of non-adjacent vertices \it{v},w in G_0 that are in a common small vortex $V \in W$. These extra edges are drawn without crossings, in the disks accommodating the corresponding vortices. We will refer to these edges as the *virtual edges* of *G* ′ \int_0' . If H' and H are subgraphs of *G* ′ $\frac{1}{0}$ and *G* − *Z*, respectively, such that *H* is obtained from *H*^{\prime} by replacing each virtual edge uv of H' with an $u-v$ path contained in some vortex in W , in such a way that all the paths are internally vertex disjoint, then we call H a *lift* of H' (w.r.t. the near embedding (σ , G_0 , Z , V , W)), and say that H' can *be lifted to H*.

A cycle *C* in a graph *H* embedded in a surface Σ is *flat* if *C* bounds a disk in Σ. A wall *W* in *H* is *flat* if the boundary cycle of *W* (which is defined in the obvious way) bounds a closed disk *D*(*W*) with the wall *W* drawn inside it.

Vertex-disjoint cycles *C*1,...,*C^s* of *H* are *concentric* if they bound closed disks *D*1,...,*D^s* in Σ with $D_1 \supseteq \cdots \supseteq D_s$ in Σ . These cycles *enclose* a vertex subset Ω if $\Omega \subseteq D_s$. They *tightly enclose* Ω if moreover, for every *i* ∈ [*s*] and every point *v* ∈ bd(*Di*), there is a vertex *w* ∈ Ω at distance at most *s* −*i* +2 from *v* in Σ w.r.t. *H*. (See the beginning of Section [6](#page-32-0) for the definition of 'distance'.)

In the context of a near-embedding (*σ*, G_0 , Z , V , W) of a graph *G* in a surface Σ, concentric cycles C_1, \ldots, C_s in G'_0 \mathcal{O}_0' (*tightly) enclose* a vortex $V \in \mathcal{V}$ if they (tightly) enclose $\Omega(V)$.

For integers $3 \le s \le h$, an $(\alpha_0, \alpha_1, \alpha_2)$ -near embedding (σ, G_0, Z, V, W) of a graph *G* in a surface Σ is (*s*,*h*)*-good* if the following properties are satisfied.

- (i) ^{G'} $\frac{1}{0}$ contains a flat wall W_0' of height h .
- (ii) If Σ is not the sphere, then the facewidth of G' \int_0' in Σ is at least *s*.
- (iii) For every vortex $V \in V$ there are *s* concentric cycles $C_1(V),..., C_s(V)$ in G' 0 tightly enclosing *V* and bounding closed disks $D_1(V) \supseteq \cdots \supseteq D_s(V)$, such that $D_s(V)$ contains $\Omega(V)$ and $D(W'_0)$ does not meet $D_1(V)$. For distinct $V, V' \in V$, the disks $D_1(V)$ and $D_1(V')$ are disjoint.

We call the above wall W_0' a *good wall* with respect to the (*s*, *h*)-good ($\alpha_0, \alpha_1, \alpha_2$)-near embedding. Also, we let $G(D_i(V))$ denote be the subgraph of *G* contained in the disk $D_i(V)$ for each $i \in [s]$.

By combining the main result of [\[KTW20\]](#page-57-9) (Theorem 2.11) and results in [\[DiKMW12\]](#page-56-10), it is possible to deduce the following theorem. We remark that some extra properties of the embedding could be derived as well from these two papers, we only state here the properties that we will need.

Theorem [58](#page-39-1) ([\[KTW20,](#page-57-9) Theorem 2.11], [\[DiKMW12\]](#page-56-10)). Let t', s, $h \in \mathbb{N}$. Then there exist $h' = h'_{58}(t', s, h)$,

 $\alpha_0 = \alpha_0(t',s)$, and $\alpha = \alpha(t')$ such that the following holds. Let G be a graph and let W be a wall of height *h* ′ *in G. Then either G has a K^t* ′*-model grasped by W , or G has an* (*s*,*h*)*-good* (*α*0,*α*,*α*)*-near embedding* (σ,G_0,Z,V,W) in a surface Σ of Euler genus at most α , such that G'_0 contains a good wall W'_0 with respect *to the embedding that can be lifted to a subwall* W_0 *of* W .

Furthermore, for some function T , there is an algorithm with running time T (*t* ′ ,*s*,*h*)·*n ^O*(1) *that, given an n-vertex graph G and a wall W as above, finds one of the two structures guaranteed by the two outcomes of the theorem.*

Using this theorem, we prove the following lemma.

Lemma 59. Let t, s, h be positive integers. Let $t' = 4t + 2$, $h' = h'_{58}(t', s, h)$ $h' = h'_{58}(t', s, h)$ $h' = h'_{58}(t', s, h)$, $k = k_{55}(h')$ $k = k_{55}(h')$ $k = k_{55}(h')$, and let $\alpha_0 = \alpha_0(t', s)$ a *and* $\alpha = \alpha(t')$ *be as in Theorem [58.](#page-39-1) Then, for every rooted graph* (*G,R*) *without a rooted* $K_{2,t}$ *-minor and every k-interesting subgraph H of* (*G*,*R*) *with core X ,*

- *(i) there exists an* (*s*,*h*)*-good* (*α*0,*α*,*α*)*-near embedding* (*σ*,*H*0,*Z*,V ,W) *of H in a surface* Σ *of Euler genus at most α, and*
- (ii) H_0' σ_0' contains a good wall W_0' with respect to the embedding that can be lifted to a wall W_0 in H, and such that for every transversal Y' of W₀ there are h vertex-disjoint X–Y' paths in H, and there are h *vertex-disjoint* $R_H - Y'$ *paths in* H *, where* $R_H = R \cap V(H)$ *.*

Furthermore, for some function T', there is an algorithm with running time T'(t, s, h) \cdot n^{O(1)} that, given an *n-vertex rooted graph* (*G*,*R*) *and a k-interesting subgraph H of G with core X as above, finds this embed-* \ddot{a} *ding and the wall* W'_0 .

Proof. Let (G, R) be a rooted graph without a rooted $K_{2,t}$ -minor and let *H* be a *k*-interesting subgraph of (G, R) with core *X*. By Lemma [55,](#page-36-0) there is a wall *W* of height *h*^{\prime} in *H* such that for every transversal set *Y* of *W* there are *h*^{\prime} vertex-disjoint *X*–*Y* paths in *H* and *h*^{\prime} vertex-disjoint *R_H*–*Y* paths in *H*, where $R_H = R \cap V(H)$.

It follows from Lemma [57](#page-37-1) that H does not contain a $K_{t'}$ -model grasped by $W.$ Theorem [58](#page-39-1) implies then that *H* has an (*s*, *h*)-good ($α_0$, $α$, $α$)-near embedding ($σ$, H_0 , Z , V , W) in a surface Σ of Euler genus at most *α*. This shows property [\(i\).](#page-40-0)

Moreover, by Theorem [58,](#page-39-1) *H* ′ $\frac{1}{0}$ contains a good wall W_{0}' with respect to the embedding that can be lifted to a subwall W_0 of W . Finally, suppose Y' is a transversal set of W'_0 . Since W'_0 has height h , it follows that $|Y'| = h$. Since W'_0 can be lifted to the subwall W_0 of W , it follows that Y' can be extended to a transversal set *Y* of *W* . We know that are *h* ′ vertex-disjoint *X*–*Y* paths in *H* and *h* ′ vertex-disjoint *R^H* –*Y* paths in *H*. Keeping only the paths that have an endpoint in *Y* ′ give the desired paths in property [\(ii\).](#page-40-1)

6.5 Discarding large vortices

For an integer *q*, the *shallow vortex grid of order q* is the graph obtained from the Cartesian product of a cycle u_1, u_2, \ldots, u_{4q} with a path v_1, v_2, \ldots, v_q by adding the edges $\{(u_{4(i-1)+1}, v_1), (u_{4(i-1)+3}, v_1)\}$ and $\{(u_{4(i-1)+2}, v_1), (u_{4(i+1)+4}, v_1)\}\$ for every $i \in [q]$. For $i \in [q]$, the cycle $(u_1, v_i), (u_2, v_i), \ldots, (u_{4q}, v_i)$ is called the *i -th cycle* of the shallow vortex grid. A *sliced shallow vortex grid of order q* is defined as above with the difference that instead of taking the Cartesian product of a cycle and a path, we take the Cartesian product of a path with a path.

Let *G* be a graph and assume that there is an (s, h) -good near-embedding (σ, G_0, Z, V, W) of *G* in a surface Σ. Let $V = (J, \Omega) \in V$ be a large vortex of the near-embedding. We say that a model of a shallow vortex grid of order *q surrounds V* if for each *i* ∈ [*q* −1], the union of the vertex sets of the branch sets in the model that correspond to vertices of the *i*-th cycle separates that for the *q*-th cycle from Ω in *G*.

Thilikos and Wiederrecht [\[TW22b\]](#page-59-2) developed a useful tool that, given a near embedded graph, allows either to 'remove' a large vortex from the near embedding, or to find a shallow vortex grid minor of large order in the graph. This is Lemma 31 in [\[TW22b\]](#page-59-2). While this is not stated explicitly in [\[TW22b\]](#page-59-2), it can be checked that, in the second case, the proof in [\[TW22b\]](#page-59-2) produces a model of the shallow vortex grid that surrounds the vortex, which will be important for our purposes. (The authors are grateful to Sebastian Wiederrecht for helpful discussions regarding this matter.) One then can check that applying Lemma 31 from [\[TW22b\]](#page-59-2) with this extra "surrounding" property to the good near embeddings considered in this paper gives the following result.

Theorem 60 (Corollary from [\[TW22b,](#page-59-2) Lemma 31]). *For every positive integers q,* α_2 *with* $q \leq \alpha_2$ *, there exists a positive integer s'* = $s'_{60}(q)$ $s'_{60}(q)$ $s'_{60}(q)$ *such that for every positive integers s* \geq *s' and h the following holds. Let G* $be a graph that has an(s, h) -good(a_0, a_1, a_2) - near embedding(\sigma, G_0, Z, V, W)$ *in a surface* Σ *of Euler genus at most* α *for some integers* $\alpha_0, \alpha_1, \alpha$ *. Let* $V = (J, \Omega) \in V$ *and let* $G_V = G(D_1(V))$ *, where* $D_1(V)$ *is defined as in [\(iii\)](#page-39-2) in the definition of* (*s*,*h*)*-good embeddings. Then at least one of the following two outcomes holds.*

- *(a) There is a vertex subset S of G_{<i>V*} *with* $|S| \le 12\alpha_2 q$ *such that there is an* $(s |S|, h)$ -good $(\alpha_0 + |S|, \alpha_1 1,\alpha_2$)-near embedding ($\widetilde{\sigma},\widetilde{G}_0,Z\cup S,V\setminus\{V\},\widetilde{W})$ of G in Σ such that the embedding $\widetilde{\sigma}$ is an extension *of* σ *to the graph J* − *S and with* $W \subseteq \widetilde{W}$ *.*
- *(b) G^V contains a model of a shallow vortex grid of order q that surrounds V .*

Lemma 61. *Let S be a sliced shallow vortex grid of order t, that is, the Cartesian product of a path* u_1, u_2, \ldots, u_{4t} with a path v_1, v_2, \ldots, v_t plus crosses. Let $R = \{u_1, u_2, \ldots, u_{4t}\} \times \{v_t\}$. Then (S, R) contains *a rooted K*2,*t-model.*

Proof. Using the crosses on the boundary, we find two disjoint intertwining paths P_1 and P_2 with vertices in $\{u_1, u_2,..., u_{4t}\}\times \{v_1, v_2\}$ so that P_1 contains all vertices (u_i, v_2) with $i \mod 8 \in \{1, 2, 7, 8\}$, and P_2 contains all vertices (u_i, v_2) with i mod $8 \in \{3, 4, 5, 6\}$. For each $j \in [t]$, let $B_j = \{u_{4j-3}, u_{4j-2}, u_{4j-1}, u_{4j}\} \times \{v_3, \ldots, v_t\}$. Then the desired rooted $K_{2,t}$ -model has branch sets $V(P_i)$ for $i \in \{1,2\}$, and B_i for $i \in [t]$.

Lemma 62. *Let t, m, n be positive integers with* $min\{m, n\} \geq 3t$, and let *J be the n* × *m grid. Let X denote the set of all t² vertices with coordinates* (*i*, *j*) such that $1 \le i, j \le t$, and let Y be any set of 4t² vertices of J. *Then there exist* 2*t* −1 *vertex-disjoint X –Y paths in the grid.*

Proof. Suppose towards a contradiction, that the lemma does not hold. Then, by Menger's Theorem, there exists a set of vertices *S* with |*S*| < 2*t* − 1 such that after removing *S* from the grid, no component intersects both *X* and *Y*. For each $i \in [t]$, let P'_i a_i^\prime denote the path induced by the vertices of the grid with coordinates in the set $\{i\} \times \{t, \ldots, m\}$, and for each $j \in [t-1]$, let P''_i $_j^{\prime\prime}$ denote the path induced by the vertices of the grid with coordinates in the set $\{t,\ldots,n\}\times\{j\}$. Note that the paths P'_i P' _{*i*} and P'' _{*j*} *j* are all pairwise vertexdisjoint, and each of them intersects *X*. Since $|S|$ < 2*t*−1, there exists $P_X \in {P'_1}$ $P'_{1},...,P'_{t}$ $'_{t}$, P''_{1} $\binom{n}{1}, \ldots, P_{t-1}^{\prime\prime}$ disjoint from *S*.

The number of vertices in the set *Y* is at most the number of rows of the grid intersecting *Y* times the number of columns of the grid intersecting *Y*. As $|Y| = (2t)^2$, there exists a row or column P_Y that intersects *Y* and that is disjoint from *S*.

Since $\min\{m, n\} \geq 3t$, and the set *X* intersects *t* rows and *t* columns, there exist a column *Q'* and a row Q'' of the grid, that are disjoint from $X \cup S$. Thus, $P_X \cup P_Y \cup Q' \cup Q''$ is a connected subgraph of the grid which is disjoint from *S*, and intersects both *X* and *Y* , a contradiction. \Box

Note that the definition of a transversal set in a wall naturally extends to a transversal set in a grid. Let *M* be a model of a shallow vortex grid in some graph *G*, then a *transversal set of M* is a set *Y* of vertices chosen such that no two vertices are taken from the same branch set of *M*, and the branch sets which contain vertices from *Y* correspond to a transversal set of the grid we obtain by contracting the branch sets of *M*.

Lemma 63. Let t, s, h be positive integers with $h \ge 3t^2$, and let $t' = 4t + 2$. Then there exist positive integers $s' = s'(t, s)$ and $k = k(t, s, h)$ such that, letting $\alpha_0 = \alpha_0(t', s')$ and $\alpha = \alpha(t')$ be as in Theorem [58,](#page-39-1) the follow*ing holds. For every rooted graph* (*G*,*R*) *without a rooted K*2,*t-minor, and every k-interesting subgraph H of* (*G*,*R*) *with core X ,*

- *(i) there exist an* (*s*,*h*)*-good* (*α*0,0,*α*)*-near embedding* (*σ*,*H*0,*Z*,∅,^W) *of H in a surface* ^Σ *of Euler genus at most α, and*
- (ii) H_0' σ_0' contains a good wall W_0' with respect to the embedding that can be lifted to a wall W_0 in H and such that for every transversal Y' of W₀' there are h vertex-disjoint X-Y' paths in H.

Furthermore, for some function T'', there is an algorithm with running time $T''(t,s,h) \cdot n^{O(1)}$ *that, given an n-vertex rooted graph* (*G*,*R*) *and a k-interesting subgraph H of G with core X as above, finds this em*bedding and the wall W'_0 .

Proof. Let

$$
p = 4t2
$$

q = 2t(p + 1).

It can be checked that the constant $\alpha = \alpha(t')$ from Theorem [58](#page-39-1) is (much) larger than q ; in particular

 $\alpha \geq q/2$,

which will be used in the proof. Let also

$$
\beta = 12q\alpha^2 + 3p\alpha
$$

\n
$$
s' = s + s'_{60}(q) + \beta
$$

\n
$$
h'' = h + \beta
$$

\n
$$
h' = h'_{58}(t', s', h'')
$$

\n
$$
k = k_{55}(h').
$$

With these constants, Lemma [59](#page-40-2) implies that

- (i) there exists an (s', h') -good $(\alpha_0, \alpha, \alpha)$ -near embedding (σ, H_0, Z, V, W) of *H* in a surface Σ of Euler genus at most *α*, and
- (iii) H'_0 α_0' contains a good wall W_0' with respect to the embedding that can be lifted to a wall W_0 in H , and such that for every transversal *Y'* of W'_0 there are *h'* vertex-disjoint *X*–*Y'* paths in *H*, and there are *h*^{\prime} vertex-disjoint *R_H*-*Y*^{\prime} paths in *H*, where *R_H* = *R* \cap *V*(*H*).

If $V = \emptyset$ then the above near embedding is the required embedding. Indeed, any (s', h') -good embedding is also (s, h) -good since $s' \geq s$ and $h' \geq h$. Hence, we may assume that $V \neq \emptyset$.

The main idea of the proof below is to show that there is no shallow vortex grid model surrounding a vortex *V* \in *V* such that there are many vertex-disjoint paths between *R_H* and Ω. This is because, if such a structure existed, then we could find a rooted $K_{2,t}$ -minor in (H, R_H) , and thus also in (G, R) , which is a contradiction to our assumption about (*G*,*R*). Using the non-existence of such models, we use Theorem [60](#page-41-0) to find a small set of vertices, the removal of which "kills" the vortex *V* . We then iterate over all remaining vortices in $\mathcal V$.

First, assume that there is some vortex $V \in V$ such that

(i) there is a shallow vortex grid model *M* of order *q* in *H* that surrounds *V* , and

(ii) there are *p* vertex-disjoint paths in *H* between R_H and *p* distinct branch sets of *M*.

We will show that the existence of this model and these *p* paths together imply that (*G*,*R*) has a rooted *K*2,*t*-minor, a contradiction.

In what follows, we use a terminology similar to the one in the proof of Lemma [57.](#page-37-1) We call *feasible path system* a collection of *p* vertex-disjoint paths in *H* such that all the paths have one endpoint (called *source*) in *R^H* , and their other endpoint (called *destination*) in distinct branch sets of *M*. Note that there is at least one feasible path system in *H*, by our assumption above. A path *P* in a feasible path system $\mathcal P$ is *good* if the branch set of *M* containing the destination of *P* avoids all the other paths in \mathcal{P} , and is *bad* otherwise. With respect to \mathcal{P} , we classify branch sets of M into four categories: A branch set B of M is

- *good* if *B* contains the destination of a *good* path of \mathcal{P} ;
- *bad* if *B* contains the destination of a *bad* path of \mathcal{P} ;
- *ugly* if *B* intersects some path in \mathcal{P} but *B* contains no destination of paths in \mathcal{P} ;
- *free* if *B* intersects no path in \mathcal{P} .

We let $g(\mathcal{P}), b(\mathcal{P}), u(\mathcal{P}), f(\mathcal{P})$ denote the number of branch sets in *M* that are respectively good, bad, ugly, and free w.r.t. \mathcal{P} . Finally, we let *i*(\mathcal{P}) denote the sum over every path $P \in \mathcal{P}$ of the number of branch sets of *M* intersected by *P*.

Let $\mathscr P$ be a feasible path system that minimizes $i(\mathscr P)$. Observe that this implies

$$
u(\mathcal{P})=0,
$$

since otherwise some path P in $\mathcal P$ could be shortened so as to end in an ugly branch set, which would decrease $i(\mathcal{P})$. Thus,

$$
b(\mathcal{P})+g(\mathcal{P})=|\mathcal{P}|=p.
$$

The shallow vortex grid of order $q = 2t(p + 1)$ contains $p + 1$ disjoint copies of the sliced shallow vortex grid of order 2*t*, so one of these copies, which we denote by *S*, has the property that all branch sets *M*(*u*) with $u \in V(S)$ are free. In our shallow vortex grid of order q, we find a $4q \times q$ grid *J* such that the vertex set of *S* is in the corner of the grid; that is, for every $i, j \in [2t]$, the vertex of *S* corresponding to (u_i, v_j) coincides with the vertex of *J* with coordinates (i, j) (where u_i, v_j refer to the notations from Lemma [61\)](#page-41-1). Let *S* ′ be a sliced shallow vortex grid of order *t* contained in *S*, which contains the same corner of *J* as *S*, and thus, is induced by the vertices (u_i, v_j) of *S* with $i, j \le t$. We will build a rooted $K_{2,t}$ -model in (G, R) by rerouting some paths in ${\mathscr P}$ so that they hit S' in a way that allows applying Lemma $61.$

Let *U* denote the set of all $p = 4t^2$ vertices $u \in V(J)$ such that the branch set $M(u)$ contains the destination of a path of \mathcal{P} (i.e., such that $M(u)$ is good or bad).

We will now focus on paths in the grid *J*, and then we will use the model *M* to build a rooted $K_{2,t}$ minor in (*G*,*R*). By Lemma [62,](#page-41-2) there exists a set \mathcal{Q} of $2t - 1$ disjoint $U - V(S')$ paths in *J*. Recall that the sliced shallow vortex grid *S* ′ has order *t*, and that its vertex set resides in the corner of *J* and is disjoint from*U*, so each path in Q has its destination in one of the 2*t* −1 vertices of *S* ′ that are adjacent to vertices outside *S* in *J*. Since *J* is bipartite, we can find $Q_1, \ldots, Q_t \in \mathcal{Q}$ such that if w_c denotes the end of Q_c in *U*, then the vertices w_1, \ldots, w_t are pairwise nonadjacent in *J*.

Let s^* denote the vertex of S that corresponds to the corner (u_{2t}, v_{2t}) , and for each $c \in [t]$, let s'_c denote the end of Q_c in $V(S')$, and let s_c be the vertex of Q_c belonging to $V(S) \setminus \{s^*\}$ that is closest to w_c on Q_c (so that the s_c-w_c subpath of Q_c intersects $V(S) \setminus \{s^*\}$ only in s_c). Thus, each vertex s_c corresponds to some vertex (u_i, v_j) with exactly one of *i*, *j* equal to 2*t*. Let $R_{S'}$ denote the set of the vertices of *S'* corresponding to the vertices (u_i, v_j) with $j = t$. After reordering the paths Q_1, \ldots, Q_t and rerouting them using the buffer zone of *S*, we may assume that each *s* ′ c' is the vertex corresponding to the vertex $(u_c, v_t) \in R_{S'}$ of S' .

Let $c \in [t]$, and let w'_c denote the vertex of Q_c adjacent to its end w_c . Thus, the graph H contains an edge between $M(w_c)$ and $M(w_c')$ *c*^{\prime}. Since *w_c* ∈ *U*, the branch set *M*(*w_c*) intersects at least one path from $\mathscr{P}.$ Let us pick a path $P_c \in \mathscr{P}$ such that the subgraph of H induced by $M(w_c)$ contains a path P'_c c' such that one end of P'_c \int_{c}^{t} is in P_c , the other end of P'_c α'_{c} is a vertex adjacent to $M(w'_{c})$ c' _c), and no internal vertex of P'_c c' is in any of the paths in \mathcal{P} .

Since the vertices w_1, \ldots, w_t are pairwise non-adjacent in *J*, the paths P_1, \ldots, P_t are pairwise distinct: Indeed, if some P_{c_1} and P_{c_2} with $c_1 \neq c_2$ were equal, then we could reroute that path to either $M(w'_c)$ c_1) or $M(w'$ $'_{c_2}$) to make $i(\mathscr{P})$ smaller.

Let *Q* ∗ $x_1^*,...,Q_t^*$ t ^{*} be vertex-disjoint paths in *H* such that each Q_c^* \int_{c}^{*} is an $M(s)$ c')– R *H* path contained in $\bigcup_{w\in V(Q_c)} M(w) \cup V(P'_c)$ C'_c) ∪ *V*(P_c). By Lemma [61,](#page-41-1) there exists a rooted *K*_{2,*t*}-model *M'* in (*S'*, *R*_{*S'*}). We can then transform it into a rooted $K_{2,t}$ -model M'' in (G, R) , where for each $a \in V(K_{2,t})$, the branch set $M''(a)$ is defined as the union of all branch sets $M(u)$ of the shallow vortex grid model with $u \in M'(a)$, and the union over all $c \in [t]$ with $s'_c \in M'(a)$ of the vertex sets of the paths Q_c^* *c* . The existence of this rooted *K*2,*t*-model is a contradiction.

Therefore, we may assume that for each vortex $V \in V$, at least one of the two conditions [\(i\)](#page-42-0) and [\(ii\)](#page-42-1) above fails to hold.

Let $V \in V$. If [\(i\)](#page-42-0) does not hold for *V*, then we simply update the current near embedding using Theorem [60,](#page-41-0) allowing thus to get rid off *V* , at the price of a small increase in the size of the apex set, and a small decrease in the first parameter for the goodness of the near embedding, which is perfectly fine as *s* ′ was chosen large enough at the beginning to have room for up to *α* such reductions. This way, we may remove iteratively all vortices $V \in V$ for which [\(i\)](#page-42-0) does not hold, updating the near embedding each time.

Next, assume that there is a vortex $V = (J, \Omega) \in V$ for which condition [\(i\)](#page-42-0) holds but not condition [\(ii\).](#page-42-1) As we explain below, in this case there is a transversal set *Y'* of \widetilde{W}_0 , where \widetilde{W}_0 is the subwall of W_0 obtained after the previous steps of the near embedding updates, such that there is a separation (*A*,*B*) of *H* where $\Omega \subseteq A$, $Y' \subseteq B$ and $|A \cap B| < 3p$. In this case we update the near embedding as follows. We add $A \cap B$ to the apex set, remove *V* from *V* and add (A, \emptyset) to *W*. Again, this results in a small increase of the apex set, and a decrease of at most 3*p* in the first parameter for the goodness of the near embedding, which we have budgeted for thanks to the initial choice of *s'*. Proceeding iteratively this way with every remaining vortex in $\mathcal V$, we eventually get rid of all of them, as desired.

Hence, it only remains to show the existence of the transversal set *Y'* of \widetilde{W}_0 and the separation (A, B) of *H* mentioned above. Assume to the contrary that they do not exist, that is, for each transversal set *Y* ′ in \widetilde{W}_0 there is no such separation (*A*, *B*) of *H*. We will show that condition [\(ii\)](#page-42-1) then holds for *V*, which is a contradiction. Let *M* be a model of a shallow vortex grid of order *q* that surrounds *V* , as in [\(i\).](#page-42-0) Let *Y* ′ be a transversal set in \widetilde{W}_0 and let \mathscr{P}' be a collection of at least 3*p* vertex-disjoint paths in *H* between *Y'* and Ω (they exist by Menger's theorem and our assumption that there is no separation between them of a smaller size). Using the facts that $D(\widetilde{W}'_0)$ does not meet $D_1(V)$ and that *M* surrounds *V*, we deduce that each path $P \in \mathcal{P}'$ intersects the union of the vertex sets of the branch sets in the model that correspond to vertices of the *i*-th cycle for each $i \in [q]$, we refer to this collection as the *i*-th cycle of the model of M or simply *a cycle of M*. Indeed, if *P* avoids some cycle of *M*, then this cycle would separate *P* from Ω, in contradiction to the definition of *P*.

Let Y'' be some transversal set in M . Next, we show that there are 3 p vertex-disjoint paths between *Y* ′ and *Y* ′′. Arguing by contradiction, assume otherwise, and let (*A*,*B*) be a separation of *H* such that *Y*' ⊆ *B*, *Y*" ⊆ *A* and $|A \cap B|$ ≤ 3*p* − 1. As $|\mathcal{P}'|$ ≥ 3*p* and *M* is of order *q* > 3*p*, there is a path *P* ∈ \mathcal{P}' and a cycle *C* of *M* that are missed by *A*∩*B*. The path *P* contains a vertex from *Y* ′ , the cycle *C* contains a vertex from *Y*^{*''*}, and *P* intersects *C*. We deduce that $A \cap B$ does not separate *Y*^{*'*} from *Y*^{*''*} in *H*, a contradiction.

Finally, we show that there are p vertex-disjoint paths between R_H and $Y^{\prime\prime}$, so that [\(ii\)](#page-42-1) holds. Assume that this is not the case, and let (A, B) be a separation of *H* such that $Y'' \subseteq A, R_H \subseteq B$ and $|A \cap B| < p$. This implies that $|A \cap Y'| \geq 2p$ but this contradicts the fact that there are at least $|Y'| \geq 3p$ vertex-disjoint paths between R_H and Y' . \Box

6.6 Further refinement of the embedding

For two near embeddings (σ , G_0 , A , V , W) and ($\tilde\sigma$, $\tilde G_0$, $\tilde A$, $\tilde V$, $\tilde W$) of a graph G in a surface Σ, we say that $(\tilde{\sigma}, \tilde{G}_0, \tilde{A}, \tilde{V}, \tilde{W})$ is *finer* than (σ, G_0, A, V, W) if $G_0 \subseteq \tilde{G}_0$, $\tilde{\sigma}$ extends σ , $A = \tilde{A}$, $V = \tilde{V}$, and for every $\tilde{V} \in \tilde{W}$, there is $V \in W$ with $\tilde{V} \subseteq V$; moreover, it is *strictly finer* if the two near embeddings are not the same. A small vortex $(J, \Omega) \in W$ is *properly attached* if *J* is connected, every component of $J - \Omega$ is adjacent to all vertices in Ω , and either $|\Omega| \leq 2$, or $V(J) \setminus \Omega \neq \emptyset$ and for every three distinct vertices $x, y, z \in \Omega$ and $w \in V(J) \setminus \Omega$, there are an *x*–*y* path and a *z*–*w* path in *J* that are vertex disjoint. Note that our notion of proper attachment is stronger than the one in [\[DiKMW12\]](#page-56-10).

Lemma 64. *For every near embedding* (σ , G_0 , A , V , W) *of a graph G in a surface* $Σ$ *, there exists a finer near embedding* (*σ*˜ ,*G*˜ ⁰, *A*˜,V˜,W˜) *of G in* Σ *such that every small vortex V*˜ ∈ W˜ *is properly attached. Furthermore, the finer near embedding can be found in polynomial time given G and* (σ, G_0, A, V, W) *in input.*

To prove Lemma [64](#page-45-0) we need the following theorem.

Theorem 65 (Two Disjoint Paths Theorem [\[Jun70,](#page-57-12) [RS90,](#page-58-6) [Sey80b,](#page-58-14) [Shi80,](#page-58-15) [Tho80\]](#page-59-6))**.** *Let x*, *y*, *z*,*w be distinct vertices in a graph J. Then either there exist an x–y path and a z–w path in J that are vertex disjoint, or there is a near embedding* (σ^*, G_0^*) $\int_0^* A^*, V^*, W^*$ of *J* into a disk such that $A^* = \emptyset$, $V^* = \emptyset$, and the vertices *x*,*w*, *y*, *z are mapped by σ* ∗ *into the boundary of the disk in that cyclic order. Furthermore, there is a polynomial-time algorithm that finds either the two disjoint paths or the near embedding.*

We now turn to the proof of Lemma [64.](#page-45-0)

Proof of Lemma [64.](#page-45-0) Let $(\tilde{\sigma}, \tilde{G}_0, \tilde{A}, \tilde{V}, \tilde{W})$ be a near embedding of *G* in Σ that is finer than (σ, G_0, A, V, W) , and such that there is no near embedding of *G* in Σ that is strictly finer than this one. (Observe that there is such a near embedding, since there cannot be an infinite sequence of near embeddings of *G* in Σ such that each one is strictly finer than the previous one.)

We claim that ($\tilde{\sigma}$, \tilde{G}_0 , \tilde{A} , $\tilde{\mathcal{V}}$, $\tilde{\mathcal{W}}$) satisfies the lemma. Suppose towards a contradiction that some vortex $(J,\Omega) \in \tilde{V}$ is not properly attached.

If *J* is disconnected, say *J* is the disjoint union of nonempty graphs *J*¹ and *J*2, we can obtain a strictly finer near embedding by replacing (*J*,Ω) by two small vortices (*J*₁,Ω₁) and (*J*₂,Ω₂) where each Ω_{*i*} is the restriction of Ω to $\Omega \cap V(J_i)$, a contradiction. (Observe that we can find a disk in Σ accommodating (J_i, Ω_i) for $i = 1, 2$ since $|\Omega_i| \leq 2$.)

If $V(J) \setminus \Omega = \emptyset$, then we obtain a strictly finer near embedding by removing the small vortex (J,Ω) from \tilde{V} , replacing G_0 by $G_0 \cup I$, and extending $\tilde{\sigma}$ to an embedding of $G_0 \cup I$ by drawing the edges of *J* along the boundary of the disk Δ corresponding to (*J*, Ω), which is a contradiction. Thus, V (*J*) \ $\Omega \neq \emptyset$.

If some connected component *C* of *J* − Ω is only adjacent to a strict subset Ω_1 of the vertices in Ω , then letting *J*₁ = *J*[*V*(*C*) ∪ Ω₁], we can obtain a strictly finer near embedding by replacing (*J*, Ω) by two small vortices (J_1, Ω_1) and $(J - V(C), \Omega)$, a contradiction. Thus, every connected component of $J - \Omega$ is adjacent to all vertices in Ω .

Finally, suppose that there are distinct vertices $x, y, z \in \Omega$ and $w \in V(J) \setminus \Omega$ such that there do not exist an *x*–*y* path and a *z*–*w* path in *J* that are vertex disjoint. Then, by Theorem [65,](#page-45-1) there exists a near embedding of *J* in a disk with no apices and no large vortices such that the vertices x, w, y, z appear on the boundary of the disk in that cyclic order. In such case we can make the near embedding strictly finer by replacing the disk ∆*ⁱ* representing (*J*,Ω) with the near embedding of *J* in the disk.

We remark that this proof can be turned into a polynomial-time algorithm by iteratively applying each of the improvement step above, and observing that the total number of improvements that can be made is bounded by a polynomial in $|V(G)|$. \Box

6.7 Ensuring 3**-connectivity**

In this subsection, we describe how to transform a given near embedding (*σ*, G_0 , *A*, V, *W*) of a graph *G* in surface Σ with a large flat wall *W* so as to make *G* ′ $\frac{\pi}{0}$ 3-connected while keeping many properties of the original near embedding (in particular, we want that a large subwall of *W* is still flat). We use the notion of SPQR trees that has been introduced by Di Battista & Tamassia [\[DBT89\]](#page-56-13). We remark that there is a linear-time construction algorithm due to Gutwenger & Mutzel [\[GM00\]](#page-56-14).

In what follows, *multigraphs* may have parallel edges between pairs of vertices, but no loops. Let *G*¹ and G_2 be multigraphs such that $V(G_1) \cap V(G_2) = \{x, y\}$ for some distinct vertices *x* and *y*, and each of G_1 and G_2 has at least one edge between *x* and *y*. Then, the 2-sum of G_1 and G_2 is the graph *G* obtained by removing one edge between x and y from each of G_1 and G_2 , and taking the union of the resulting subgraphs. A *dipole* is a multigraph with 2 vertices (and any number of parallel edges between them). For instance, the graph *K*2,*^t* can be obtained by 2-summing a dipole with *t* edges with *t* triangles.

A multigraph is 2*-connected* if every edge lies on a cycle (possibly of length 2). For every 2-connected multigraph *G*, the *SPQR tree* of *G* is a canonical representation of *G* as the 2-sum of multigraphs of three types: dipoles, cycles (without parallel edges) and 3-connected graphs (without parallel edges). We do not provide a full definition of the SPQR tree, and only describe its properties required in our proof (see [\[DBT89,](#page-56-13) [GM00\]](#page-56-14) for more background). The SPQR tree of *G* is uniquely defined. It is a tree *T* in which every node *u* is associated with a graph *Gu*, and has one of four types

- if *u* is an *S node*, G_u is a cycle,
- if *u* is a *P node*, *G^u* is a dipole with at least three edges,
- if *u* is a *Q node*, *G^u* is a dipole with two edges, and
- if *u* is an *R node*, *G^u* is a 3-connected graph.

For each edge $uv \in E(T)$, the graphs G_u and G_v share two common vertices that are adjacent in both graphs, and thus their 2-sum is well defined. We can contract any edge *uv* of *T* into a new vertex *w* and associate with it the graph G_w which is the 2-sum of G_u and G_v . If we contract the edges of *T* in any order, then regardless the order in which we contract the edges, the graph *G^r* associated with the node *r* into which we contracted the tree is equal to *G*. Since a dipole with 2 edges is a neutral element for 2-sum, if the SPQR tree has a *Q* node, then it is the only node, and *G* is a dipole with 2 edges.

Theorem 66 (Mohar [\[Moh97\]](#page-57-13))**.** *Let* Σ *be a surface with strictly positive Euler genus, and let G be a* 2 *connected graph that can embedded in* Σ *with facewidth w* ≥ 3*. Then, there is exactly one node u in the SPQR tree of G such that* G_u *is not planar. Further,* G_u *can be embedded in* Σ *with facewidth w.*

Lemma 67. *Let r* ≥ 3*, let G be a graph, let* (*σ*,*G*0,*Z*,V ,W) *be an* (*α*,0,*α*)*-near embedding of G in a surface* Σ with all small vortices properly attached, let W_0' be an $(r+2)$ -wall in G_0' that can be lifted to an $(r+2)$ wall W in G−*Z* . Then, there is an (α ,0, α)-near embedding ($\tilde\sigma$, $\tilde G_0$, $\tilde Z$, $\tilde V$, $\tilde W$) of G in a surface $\tilde\Sigma$ with ${\rm eg}(\tilde\Sigma)\leq$ eg(Σ) *such that the facewidth of σ*˜ *is not smaller than the facewidth of σ, all small vortices are properly* attached, $\tilde{G_0}$ \tilde{N} contains an r-wall \tilde{W}'_0 that can be lifted to an r-subwall \tilde{W} of W, and $\tilde{G_0}$ ′ *is* 3*-connected. Furthermore, the near embedding can be found in polynomial time given G and* (*σ*,*G*0,*Z*,V ,W) *in input.*

Proof. First, suppose that *G* ′ J_0' is not connected, and let J_0' $_0^{\prime}$ be a connected component of G'_{0} $\frac{1}{0}$ distinct from the one containing W'_0 . Let W' denote the set of all small vortices $(J, \Omega) \in W$ with $\Omega \cap V(J'_0)$ $_{0}^{\prime}) \neq \emptyset$ (and thus $\Omega \subseteq V(J'_c)$ J_0)), and let J_1 denote the union of $J_0' \cap G_0$ and all small vortices in \mathcal{W}' . Then, after replacing *G*⁰ with $G_0 - V U_0'$ J_0') and $\mathscr W$ with $(\mathscr W\setminus \mathscr W')\cup \{(J_1,\varnothing)\}$, we obtain a near embedding, where J_0' $\frac{7}{0}$ is no longer a connected component of *G* ′ δ_0' . Hence, after removing similarly all connected components of G_0' which do not contain W'_0 , we may assume that G'_0 $\frac{1}{0}$ is connected.

Similarly, we argue that we may assume *G* ′ $\frac{1}{0}$ to be 2-connected. Suppose that G'_{0} $\frac{1}{0}$ is not 2-connected. Let G' $\frac{1}{1}$ be the block of G_0' $\frac{1}{0}$ containing W_0' , and let *x* be a cut-vertex of G_0' \int_0^{\prime} contained in G'_1 J_1' . Let J_0' $\frac{7}{0}$ denote the

subgraph of *G* ′ $\frac{1}{0}$ induced by *x* and all connected components of $G'_{0} - V(G'_{1})$ \mathcal{U}_1') that are adjacent to *x*. Let \mathcal{W}' denote the set of all small vortices $(J, \Omega) \in W$ with $\emptyset \neq \Omega \subseteq V(J'_{\Omega})$ J_0^{\prime}), and let J_1 denote the union of $J_0^{\prime} \cap G_0$ and all small vortices in W'. Then, after replacing G_0 with G_0 – ($\breve{V}(J'_0)$ $\binom{1}{0} \setminus \{x\}$ and W with $(W \setminus W') \cup \{(J_1, \{x\}\}),$ we obtain a near embedding, where *x* is no longer a cut-vertex of G'_{0} η_0' . Hence, we may assume that G_0' $_{0}^{\prime}$ is 2-connected.

Let *T* be the SPQR tree of *G* ′ \mathcal{C}_0 . With each edge $u_1u_2 \in E(T)$, we associate a separation (A_1, A_2) of G_0 $_{0}^{\prime}$, defined as follows. For each *i* ∈ {1,2}, let T_i denote the connected component of $T - u_1 u_2$ that contains u_i , and let $A_i = \bigcup_{u \in V(T_i)} V(G_u)$ (where G_u denotes the graph corresponding to node *u* in the SPQR tree of *G* ′ χ'_0). Since *W*₀ is a subdivision of a 3-connected graph, for some $i \in \{1,2\}$, the set of vertices of *W*₀ in A_i is either empty or induces a path, and A_{3-i} contains all branch vertices of W'_0 . We refer to A_i as the *small side* and to A_{3-i} as the *big side*. Let us orient every edge u_1u_2 towards the vertex u_i such that A_i is the big side of the separation, and let u_0 be a sink in this orientation. Hence, $V(G_{u_0})$ contains all branch vertices of W'_0 . Moreover, the graph G_{u_0} contains a graph obtained from W'_0 by suppressing some degree-2 vertices. In particular, G_{u_0} contains an *r* -wall (which can be lifted to a subwall of W), so u_0 must be an R node of the SPQR tree.

To complete the proof, we will construct a near embedding with $\tilde{G_0} = G_0 \cap G_{u_0}.$ This choice uniquely determines the small vortices, and we will have $G'_0 = G_{u_0}$, so G'_0 will be 3-connected. By Theorem [66,](#page-46-0) either G_u has an embedding in Σ with the same facewidth as σ , or G_u is planar (and thus has an embedding in the sphere with infinite facewidth). In both cases, there exists a surface $\tilde{\Sigma}$ with Euler genus not greater than the one of Σ and an embedding $\tilde{\sigma}$ of G_u in $\tilde{\Sigma}$ with facewidth not smaller than the one of σ . Let $\tilde{G}_0 = G_0 \cap G_{u_0}$, let $\tilde{Z} = Z$, let $\tilde{V} = V = \emptyset$, and let \tilde{W} be defined as follows: for each small vortex (*J*,Ω) ∈ W with $\Omega \subseteq V(G_{u_0})$, include it in $\tilde{\mathscr{W}}$. Next, turn each connected component T' of $T-u_0$ into a small vortex (J', Ω') defined as follows: Let J'_{0} \int_0^{\prime} denote the subgraph of G_0 induced by all vertices in $\bigcup_{u\in V(T')}V(G_u)$, and let *J'* denote the union of J'_{0} \mathcal{O}_0 and all small vortices $(J, \Omega) \in \mathcal{W}$ with $\Omega \subseteq V(J'_0)$ α_0'). Finally, let Ω' denote the only cyclic ordering of the two-element set $V(G_{u_0}) \cap V(J'_{c})$ $\tilde{\mathbf{v}}_0'$). This defines the set of small vortices \tilde{W} . Note that neither the Euler genus nor *Z* were increased during the construction. Therefore, we have $\tilde{G}_0 = G_{u_0}$, so $(\tilde{\sigma},\tilde{G}_0,\tilde{Z},\tilde{\mathcal{V}},\tilde{\mathcal{W}})$ is an $(\alpha,0,\alpha)$ -near embedding satisfying the lemma. П

6.8 Constraining the roots

In this section, we finally prove Lemma [54.](#page-34-1) By Lemma [63,](#page-41-3) for every rooted graph (*G*,*R*) without a rooted *K*2,*t*-minor and every *k*-interesting subgraph *H* of (*G*,*R*) with core *X*, there exists an (*s*,*h*)-good (*α*,0,*α*) near embedding (σ , G_0 , A , V , W) in a surface of Euler genus at most α with a flat wall W' of height h in G_0' 0 that can be lifted to a wall W'_0 in *G*, and there are *h* vertex-disjoint paths between *X* and any transversal set of W'_0 . It remains to show that

- (i) only a bounded number of small vortices in W contain a root from the set R that is not in G' $_0^{\prime}$, and
- (ii) the roots in G' $\frac{1}{0}$ have a small face cover.

First, we will prove that [\(i\)](#page-47-0) holds for the small vortices that have at most two vertices in G'_{0} v'_0 , and then we will use Theorem [8](#page-8-3) to simultanously handle the small vortices with three vertices in *G* ′ $\frac{1}{0}$ and prove [\(ii\).](#page-47-1)

Lemma 68. *Let* $t ≥ 2$ *be an integer, let G be a* 3*-connected graph, and let* $D ⊆ E(G)$ *be a set of edges with* $|D|$ ≥ 3*t* − 3. Then there exists a subset $D' \subseteq D$ with $|D'| = t$ such that $G - D'$ is connected.

Proof. We assume that $G - D$ is disconnected since otherwise the lemma is satisfied by any set $D' \subseteq$ *D* with $|D'| = t$. Let $c \ge 2$ denote the number of components of $G - D$. Since *G* is 3-connected, each component of *G* − *D* is linked to the remaining components with at least three edges from *D*, so $|D| \geq \frac{3}{2}c$. Combining this with the assumption $|D| \geq 3t - 3$, we deduce that $|D| \geq \frac{2}{3}(\frac{3}{2})$ $(\frac{3}{2}c) + \frac{1}{3}$ $\frac{1}{3}(3t-3) = c + t - 1$. Let *D*₀ ⊆ *D* be a set of *c* − 1 edges such that $(G - D) + D_0$ is connected. We have $|D \setminus D_0| = |D| - |D_0| \ge$ $(c+t-1)-(c-1) = t$. Let $D' \subseteq D \setminus D_0$ be any subset with $|D'| = t$. Then $G - D'$ is a supergraph of the graph $(G - D) + D_0$, so *D'* satisfies the lemma. \Box

Lemma 69. Let (G, R) be a 3-connected rooted graph without a rooted $K_{2,t}$ *-minor, let H be a k-interesting* s ubgraph, and let (σ, H_0, Z, V, W) be an $(\alpha, 0, \alpha)$ -near embedding of H with H'_0 3-connected and all small α *vortices properly attached. Then there are at most* $k + (t-1)\binom{\alpha}{2} + (3t-4)\alpha$ *small vortices* $(J,\Omega) \in W$ *with* $|\Omega| \leq 2$ *and* $(V(J) - \Omega) \cap R \neq \emptyset$.

Proof. Partition the set of all small vortices $(J, \Omega) \in W$ with $|\Omega| \leq 2$ and $(V(J) - \Omega) \cap R \neq \emptyset$ into three sets \mathcal{W}_∂ , $\mathcal{W}_{\leq 1}$, and \mathcal{W}_2 , by assigning (*J*, Ω) to \mathcal{W}_∂ if (*V*(*J*) − Ω) ∩ $\partial(H) \neq \emptyset$, to $\mathcal{W}_{\leq 1}$ if (*V*(*J*) − Ω) ∩ $\partial(H) = \emptyset$ and $|\Omega|$ ≤ 1, and to \mathcal{W}_2 if $(V(J) - \Omega) \cap \partial(H) = \emptyset$ and $|\Omega| = 2$. Since *H* is *k*-interesting, we have $|\partial(H)| \le k$, and since the graphs *J* − Ω are pairwise disjoint for $(J, \Omega) \in W$, we have $|\mathcal{W}_{\partial}| \leq k$.

Next, we claim that $|W_{\leq 1}| \leq (t-1)\binom{\alpha}{2}$. Suppose to the contrary that $|W_{\leq 1}| \geq t\binom{\alpha}{2}$. For each $(J,\Omega) \in W_{\leq 1}$, choose a component *C*(*J*, Ω) of *J* − Ω that intersects *R*. Since *G* is 3-connected, $|\Omega| \le 1$, and (*V*(*J*) − Ω) ∩ $\partial(H) = \emptyset$, the component $C(J,\Omega)$ must be adjacent to at least two vertices in *Z*. We have $|Z| \leq \alpha$, so by the pigeonhole principle, there exist *t* small vortices $(J_1, \Omega_1),..., (J_t, \Omega_t) \in \mathcal{W}_{\leq 1}$ and distinct vertices $z_1, z_2 \in Z$ such that $C(J_i, \Omega_i)$ is adjacent to both z_1 and z_2 for each $i \in [t]$. Therefore, $K_{2,t}$ is a rooted minor of (G, R) , as witnessed by the model with branch sets { z_i } for $i \in \{1,2\}$, and $V(C(J_j, \Omega_j))$ for $j \in [t]$, a contradiction.

Finally, we show that \mathcal{W}_2 has size at most $(3t-4)α$. For each $(J, \Omega) ∈ \mathcal{W}_2$, choose a component $C(J, \Omega)$ of *J* −Ω that intersects the root set *R*. Since (*J*,Ω) is properly attached, *C*(*J*,Ω) is adjacent to both vertices in Ω. Since *G* is 3-connected and $(V(J) - Ω) ∩ ∂(H) = ∅$, the component *C*(*J*, Ω) is adjacent to a vertex in *Z*. Suppose towards a contradiction that $|\mathcal{W}_2|$ > $(3t-4)\alpha$. By the pigeonhole principle, there exist a subset $\mathcal{W}'_2 \subseteq \mathcal{W}_2$ and a vertex $z \in Z$ such that $|\mathcal{W}'_2|$ \mathcal{Z}'_2 | ≥ 3*t* − 3 and for each (*J*, Ω) ∈ \mathcal{W}'_2 2 , the component *C*(*J*,Ω) is adjacent to *z*. Let *D* denote the set of all edges $xy \in E(H_0)$ \mathcal{U}_0) such that $\Omega = \{x, y\}$ for some $(J, \Omega) \in \mathcal{W}_2'$ 2^{\prime} . By Lemma [68](#page-47-2) (note that we can assume that $t \geq 2$ without loss of generality), there exists a subset $D^{\tilde{t}} \subseteq D$ with $|D'| = t$, such that $H'_0 - D'$ is connected. Let $(J_1, \Omega_1), \ldots, (J_t, \Omega_t)$ be distinct small vortices in W'_2 $\frac{2}{2}$ such that each Ω_i consists of the endpoints of a different edge in D' . Let H' denote the union of H_0 with all small vortices $(J, \Omega) \in W$ except those with $\Omega = \{x, y\}$ for some $xy \in D'$, or $\Omega = \emptyset$. Since the small vortices are properly attached, the graph H' is connected analogously to H'_0 ⁻ D' . In this case we obtained a rooted model of $K_{2,t}$ in (*G*,*R*), with branch sets {*z*}, $V(H')$, and $V(C(J_i, \Omega_i))$ for $i \in [t]$, contradiction. \Box

Lemma 70. Let (G, R) be a 3-connected rooted graph without a rooted $K_{2,t}$ *-minor, let H be a k-interesting subgraph, and let* (*σ*,*H*0,*Z*,V ,W) *be a near embedding of H in a surface* Σ *with Euler genus at most g such that H*′ 0 3*-connected and all small vortices are properly attached. Suppose further that the facewidth of* σ *is at least* $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ *. Then, there are at most* $f_8(g, t)$ *small vortices* $(J, \Omega) \in W$ *such that* $|\Omega| = 3$ *and* $(V(f) - \Omega) \cap R \neq \emptyset$, and there is a set of at most $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ facial cycles in σ that cover the set $R \cap V(H_0)$.

Furthermore, for fixed t, the set of facial cycles can be found in polynomial time given (*G*,*R*)*, H, and* (σ, H_0, Z, V, W) *in input.*

Proof. Let H_0'' $B_0^{\prime\prime}$ be obtained from H_0^\prime \mathcal{O}_0' by adding, for each $V = (J, \Omega) \in \mathcal{W}$ with $|\Omega| = 3$ and $(V(J) - \Omega) \cap R \neq \emptyset$, a new vertex x_V that is adjacent to all vertices in Ω , and extend the embedding σ of H_0' $\frac{7}{0}$ to an embedding *σ*['] of *H*₀["] $\frac{\partial}{\partial t}$ by embedding each x_V in the interior of the triangular face containing the three vertices in Ω. Note that H''_0 σ' is still 3-connected, and the facewidth of σ' is at least $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$. Let *R'* be the union of $R \cap V(H_0)$ and the set of all new vertices x_V . Since the small vortices are properly attached, we can see that there is no rooted $K_{2,t}$ -minor in (H''_0) C_0'' , *R*[']) (otherwise we could lift it to a rooted $K_{2,t}$ -minor in (G, R)). By Theorem [8,](#page-8-3) there is a set of at most $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ facial cycles in σ' that cover all vertices in R' . Since each face of σ' contains at most one vertex x_V , we conclude that there are at most $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ vertices x_V , and thus there are at most *f*_{[8](#page-8-3)}(*g*, *t*) small vortices (*J*, Ω) ∈ *W* with $|Ω| = 3$ and (V (*J*) − Ω) ∩ *R* ≠ \emptyset . We can also transform the collection of at most $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ facial cycles in σ' covering R' into a collection of at most $f_8(g, t)$ $f_8(g, t)$ $f_8(g, t)$ facial cycles in σ covering $R \cap V(H_0)$ by replacing any facial cycle containing a vertex x_V with $V = (J, \Omega)$ by the facial cycle H'_0 $\int_0^{\prime} [\Omega]$. \Box

An embedding of a graph *G* in a surface Σ is *cellular* if the closure of every face is homeomorphic to a disk. A graph *G* is *cellularly embedded* in Σ if the embedding is cellular.

Theorem 71 (Robertson and Vitray [\[RV90\]](#page-58-16), [\[MT01,](#page-57-11) Proposition 5.5.2]). *Let* Σ *be a surface with* eg(Σ) > 0 *and let G be a graph that is embedded in* Σ *with facewidth at least* 2*. Then there is precisely one block Q of G that contains a noncontractible cycle. Moreover, Q is cellularly embedded in* Σ *and all its faces are bounded by cycles. Each block Q*′ *of G distinct from Q is a planar subgraph of G contained in the closure of some face of Q. Finally, the facewidth of Q is equal to the facewidth of G.*

A corollary of this result is that if *G* is a 2-connected graph that is embedded in Σ with facewidth at least 2, then all the faces of the embedding are bounded by cycles.

Proof of Lemma [54.](#page-34-1) Let $\alpha = \alpha(t)$ be as in Lemma [63,](#page-41-3) let $h = 2\alpha + 8$ $h = 2\alpha + 8$. Let $w = f_8(\alpha, t)$. Let $k = k(t, w, 3t^2)$ (as in Lemma [63\)](#page-41-3), and let $k' = k + (t-1)\binom{\alpha}{2} + (3t-4)\alpha + w$. Let (G, R) be a rooted graph without a rooted *K*2,*t*-minor, and let *H* be a *k*-interesting subgraph. By Lemma [63,](#page-41-3) *H* admits a (*w*,*h* + 2)-good (*α*,0,*α*) near embedding (σ , H_0 , Z , V , W) (thus, $V = \emptyset$), and there is a good (h + 2)-wall W'_0 in H'_0 $\frac{7}{0}$ such that any transversal set of W'_0 is linked to *X* in *H* with *h* disjoint paths. By Lemma [64,](#page-45-0) we may assume that all vortices are properly attached, and by Lemma [67,](#page-46-1) we may alter the near embedding and the wall so that H_0' $\frac{1}{0}$ is 3-connected and W_0' is a good *h*-wall, but the facewidth is still at least *w*, and the small vortices are still properly attached. Note that by Theorem [71,](#page-49-0) the embedding σ is cellular, and thus each face is bounded by a cycle. By Lemma [69,](#page-48-0) there are at most k' − *w* small vortices (*J*, Ω) ∈ *W* with $|\Omega|$ ≤ 2 and $(V(J) - \Omega) \cap R \neq \emptyset$. By Lemma [70,](#page-48-1) there are at most *w* small vortices $(J, \Omega) \in W$ with $|\Omega| = 3$ and $(V(f) - \Omega) \cap R \neq \emptyset$, and there is a set of *w* facial cycles which cover all vertices in $V(H_0) \cap R$.

Let $(J_1, \Omega_1),..., (J_m, \Omega_m)$ denote the small vertices in W, listed in an order such that for any $i \in [m]$, if $i > 2k \ge k + (t+1)\binom{\alpha}{2} + 3(t+1)\alpha + w$, then $(V(J_i) - \Omega_i) \cap R = \emptyset$. We define a star-decomposition $(B_0; B_1, \ldots, B_m)$ by letting $B_0 = V(H_0) \cup X$, and $B_i = V(J_i) \cup Z$ for $i \in [m]$. Now, let $U = Z \cup (X \setminus V(H_0))$. Then, $|U| \leq |Z| + |X| \leq \alpha + k \leq k'$, and we have $H^* [B_0] - U = H_0'$ $\binom{1}{0}$, so it remains to verify that $|B_0 \cap B_i| \leq k$ for each $i \in [m]$. We have $B_0 \cap B_i \subseteq (V(I_i) \cap (V(H_0) \cup X)) \cup Z$ and $|V(I_i) \cap V(H_0)| \le 3$ and $|Z| \le \alpha$, so it suffices to bound $|V(J_i) \cap X|$.

Fix *i* ∈ $[m]$. Let $P_1, ..., P_h$ be disjoint paths in *G* such that each P_j is between a vertex x_j ∈ *X* and a v ertex y_j ∈ $V(W'_0)$ such that $\{y_1,..., y_h\}$ is transversal set in W'_0 . Since W'_0 is a good wall, $J_i - \Omega_i$ contains at most one vertex y_i . The set $Z \cup \Omega_i$ has at most $\alpha + 3$ elements, and separates $V(J_i)$ from $V(H) \setminus V(J_i)$. Since at most $\alpha + 3$ of the paths P_1, \ldots, P_h intersect $Z \cup \Omega_i$, we may assume that the paths $P_{\alpha+4}, \ldots, P_h$ are disjoint from $Z \cup \Omega_i$, and thus either contained in $J_i - \Omega_i$, or disjoint from $V(J_i) \cup \Omega_i$. Furthermore, since W'_0 is a good wall, $J_i - \Omega_i$ contains at most one vertex y_i , so at most one of the paths P_j is contained in $J_i - \Omega_i$. Thus, we may assume that the paths $P_{\alpha+5},...,P_h$ are disjoint from J_i . In particular, none of the $\alpha + 4$ vertices $x_{\alpha+5},...,x_h$ belongs to J_i . Once again, the set $Z \cup \Omega_i$ has at most $\alpha + 3$ elements and separates $V(J_i)$ from $\{x_{\alpha+5},...,x_h\}$, so by well-linkedness of *X*, we have $|V(J_i) \cap X| \le \alpha + 3$, and thus $|B_0 \cap B_i| \leq 2\alpha + 6 \leq k$, as required. \Box

6.9 Obtaining the docset superprofiles

The goal of this subsection is to prove Theorem [9,](#page-9-0) which is our main structural result. The proof consists of two main steps: First we modify the tree-decomposition from Theorem [52](#page-33-0) to be *ℓ*-special. Second we explicitly construct a docset superprofile for each bag of the resulting tree-decomposition.

For the proof of Theorem [9](#page-9-0) we need the following results.

Figure 6: A $K_{3,6}$ -minor as in the proof of Lemma [74.](#page-50-0) The auxiliary vertices and edges are drawn within the face *F* without crossing.

Theorem 72 ([\[Rin65,](#page-58-17) [Bou78\]](#page-55-10)). *The Euler genus of a surface that embeds* $K_{m,n}$ *is at least* $\left[\frac{(m-2)(n-2)}{2} \right]$ $\frac{2(n-2)}{2}$ *if* $m \geq 2$ *and* $n \geq 2$ *.*

The above theorem in particular gives a lower bound on the Euler genus of a surface where *K*3,*m*, $m \in \mathbb{N}$, can be embedded.

Corollary 73. *Let m* ∈ N *and let* Σ *be a surface of Euler genus g* := eg(Σ)*. Then K*3,*^m cannot be embedded* $in \Sigma$ *for* $m > 2g + 2$.

Lemma 74. *Let G be a graph embedded in a surface* Σ *of Euler genus g and let F be a face bounded by a cycle of G in the embedding. Let a*1,*b*1,*a*2,*b*2,...,*am*,*b^m be some subset of vertices of F ordered in a cyclic order. Then, if* $m > g + 1$ *, there are no two vertex-disjoint connected subgraphs A, B* \subseteq *G such that* {*a*1,*a*2,...,*am*} ⊆ *A and* {*b*1,*b*2,...,*bm*} ⊆ *B.*

Proof. Assume to the contrary that we have two vertex-disjoint connected subgraphs *A*,*B* satisfying the above condition. We claim that $K_{3,2m}$ can then be embedded in Σ , in contradiction to Corollary [73.](#page-50-1) This can be seen as follows. For any pair of consecutive vertices a_i , b_j in the cyclic order given by the face F , we add a vertex c_{i+j} (c_1 if $i = 1$ and $j = m$), and edges (a_i, c_{i+j}) and (c_{i+j}, b_j). We denote this set of 2m auxiliary vertices by *C*. Place one more auxiliary vertex *v* inside *F* and connect it to each of the vertices in *C*. This can be done without introducing edge intersections, since all a_i and b_i , $i \in [m]$ are incident to *F*. We obtain a model of $K_{3,2m}$ where the vertices in *C* are the vertices of degree 3 and $\{v\}$, $V(A)$, $V(B)$ are the branch sets of vertices of degree 2*m* (see Figure [6\)](#page-50-2), implying that *K*3,2*^m* can be embedded in Σ, as claimed.

Proof of Theorem [9.](#page-9-0) Let $\ell = \ell(t)$ and let (T', \mathcal{B}') , where $\mathcal{B}' = \{B'_t\}$ u' : $u \in V(T')$ }, be the tree-decomposition we get from Theorem [52,](#page-33-0) that is, a tree-decomposition satisfying the following properties.

- (i) the adhesion of (T', \mathcal{B}') is at most ℓ , and
- (ii) for every node $u \in V(T')$, all but at most ℓ children of u are leaves v with $B'_v \cap R \subseteq B'_v$ u' , and
- (iii) for every node $u \in V(T')$, at least one of the following is true
	- (a) $|B'_1\rangle$ $|u'_{u}| \leq \ell,$
	- (b) *u* is a leaf of *T'* and $B'_u \cap R \subseteq B'_v$ $'_{u'}$, where u' is the parent of u in T' , or

(c) there is a set $Z \subseteq B'_1$ ℓ ^{*u*} such that $|Z| \leq \ell$, G^* $[B'_i]$ u'_u] − *Z* is 3-connected, does not contain a rooted *K*_{2,*t*}-minor with respect to the set of roots $B'_u \cap R - Z$ and has an embedding in a surface of Euler genus at most *ℓ* such that every face is bounded by a cycle of the graph, and all the vertices in $B'_u \cap R - Z$ can be covered using at most ℓ facial cycles.

We can turn such a tree-decomposition into an ℓ -special tree-decomposition by the following simple procedure. We traverse the rooted tree bottom-up and for every node $u \in V(T')$, we add to B'_t μ ['] all the vertices in the bags B'_v where *v* is a leaf child of *u* in *T'* and $B'_v \cap R \subseteq B'_v$ μ [']. We denote the resulting treedecomposition by (T,\mathcal{B}) , where $\mathcal{B} = \{B_u : u \in V(T)\}$. Let $u \in V(T)$ be a node of the tree-decomposition (T, \mathcal{B}) and let $R_u := B_u \cap R$ be the set of roots contained in B_u .

If $|R_u| \leq \ell$ then we obtain a docset superprofile of R_u by taking all the possible subsets of R_u . More precisely, $\mathcal{X}_u := \{R' \mid R' \subseteq R_u\} \supseteq \mathcal{P}(G, R_u)$. In this case, $|\mathcal{X}_u| \leq 2^{\ell}$, which is an upper bound that depends only on *t* and is independent of the number of vertices in *G*.

Otherwise, $|R_u| > \ell$, and B_u was obtained from a bag B'_i $'_{u'}$ of type [\(iii\).](#page-50-3)[\(c\)](#page-51-0) by adding bags of children of *u*^{$′$} without additional roots. Let *Z* ⊆ *B*^{$′$} α [']_{*u'*} and let $\mathcal F$ be the set of faces that cover the elements in *R*_{*u*} − *Z*. We obtain a docset superprofile in this case by considering collections of roots in $R_u \cap Z$ together with collections of roots contained in a bounded number of intervals chosen from each $F \in \mathcal{F}$. Formally, let *F* ∈ $\mathcal F$ and let v_1, v_2, \ldots, v_f be a cyclic ordering of the vertices on the cycle C_F bounding *F*. For *i*, *j* ∈ [*f*], we define an interval $I_{i,j}$ on C_F as the set of consecutive vertices on C_F from v_i to v_j with respect to the cyclic ordering. That is, $I_{i,j} := \{v_i, v_{i+1}, \ldots, v_j\}$ (potentially containing \ldots, v_f, v_1, \ldots), and $I_{i,i-1} := \emptyset$ for all *i* ∈ [*f*]. Let *X*, *Y* ⊆ *V*(*G*) be disjoint. We call a tuple {*X*, *Y*} of vertex sets *k*-*interlacing* with respect to *F* if there is at least k pairs $i_1, i_2 \in [f]$, where $i_1 \neq i_2$ and $v_{i_1} \in X$, $v_{i_2} \in Y$, and $\{v_j : j \in I_{i_1+1, i_2-1}\} \cap (X \cup Y) = \emptyset$. We first show a statement on the properties of *k*-interlacing sets under partitions.

Claim 75. *Let* {*X*, *Y*} *be k*-interlacing with respect to some face F, and let $p, q \in \mathbb{N}$. If we partition X into X_1, X_2, \ldots, X_p and Y into Y_1, Y_2, \ldots, Y_q , then there is some $i \in [p]$ and $j \in [q]$ such that $\{X_i, Y_j\}$ is at least $\left(\frac{k}{pq}\right)$ -interlacing with respect to F.

Proof of the claim. Each of the *k* pairs of vertices from *X*,*Y* that certify that {*X*,*Y* } are *k*-interlacing corresponds to a tuple (i, j) of indices of subsets $X_i, Y_j.$ There are at most $p\bar{q}$ different tuples of this form. The claim follows by the pigeonhole principle. \Box

We proceed with the definition of the docset superprofile for the bag B_u . For each face $F \in \mathscr{F}$, let $R_F := R_u \cap V(F)$. Let $\ell' := (\ell + 2) \cdot (2^{\ell^3 + 1} \cdot t \cdot \ell^2)^2 \cdot \ell^4$, and

 $\mathscr{X}_u := \{ R' \mid R' \subseteq R_u, \, \{R' \cap F, (R_u - R') \cap F \} \text{ is not } \ell' \text{-interlacing w.r.t. } F \text{ for any } F \in \mathscr{F} \}$.

Note that we allow all combinations on the set of roots that are in *Z*. Since the intersection between the roots and each face corresponds to a bounded number of intervals, the size of \mathcal{X}_u is bounded. We claim that $\mathscr{X}_u \supseteq \mathscr{P}(G, R_u)$. Arguing by contradiction, suppose this is not the case. Then, there is a docset *S* in *G* and a face $F \in \mathcal{F}$ such that $\{S \cap R_F, \overline{S} \cap R_F\}$ is ℓ' -interlacing.

Observe that $(G[B_u], R_u)$ does not contain a rooted $K_{2,t}$ -minor, since (G, R) has no rooted $K_{2,t}$ -minor. We perform three modifications to the graph $G[B_u]$. Firstly we add each edge $e = xy$ where $x, y \in Z$ and *e* is not in the graph yet. In this way, the graph induced on *Z* is a clique. Secondly, we add each edge *e* = *xy* such that *x*, *y* ∈ *B*^{*u*} ∩ *B*^{*w*} where *w* is the parent of *u* in *T*, and *e* is not in the graph yet. In this way, the graph induced on $B_u \cap B_w$ is also a clique. We denote the second set of edges we add as *M*. Thirdly, for each child *v* of *u* in *T*, we add each edge $e = xy$ such that $x, y \in B_u \cap B_v$, and *e* is not in the graph yet. In this way, the graph induced on $B_u \cap B_v$ is a clique for each child v.

Let *H* be the resulting graph. Since *G*[*Bu*] does not contain a rooted *K*2,*t*-minor, by Lemma [44](#page-28-0) and the facts that $|Z| \leq \ell$, $|B_u \cap B_w| \leq \ell$, $|B_u \cap B_v| \leq \ell$, and there is at most ℓ children, (H, R_u) does not contain a rooted $K_{2,t'}$ -minor for $t':=2^{\ell^3}\cdot t$.

Note that the docset *S* restricted to *H* is a docset in *H*. Let *X* and *Y* be the corresponding partition of $V(H)$, such that $H[X]$ and $H[Y]$ are connected. After the removal of the edges in *M*, the subgraphs $H[X]$ and $H[Y]$ decompose into at most ℓ^2 connected components. By Claim [75](#page-51-1) there are sets X', Y' such that *H*[*X*[']] − *M* and *H*[*Y*[']] − *M* are connected and {*X*['] ∩ *R_F*, *Y*['] ∩ *R_F*} are ((ℓ + 2) · (2^{ℓ ³⁺¹ · *t* · ℓ ²)²) -interlacing. In} addition, observe that *H*[*B* ′ $\binom{n}{u'}$ − *M* − *Z* is a subgraph of G^* [B'_i *u* ′]− *Z*. The difference is that we do not add a clique in the adhesion to children of *u* ′ that were merged in order to obtain *u*. Hence we can efficiently find an embedding on the same surface as in [\(iii\)](#page-50-3)[.\(c\).](#page-51-0)

Consider the collection of connected components of $H[X'] - Z$. Let X_1, X_2, \ldots, X_p be the vertex sets those connected components that intersect $\overline{R_F}$. Define Y_1, Y_2, \ldots, Y_q similarly, replacing X' with $Y'.$

Let $t'' := 2^{\ell^3} \cdot t \cdot \ell^2$. Assume first that either $p \ge t''$ or $q \ge t''$, say without loss of generality $p \ge t''$. In this case we find a model of a rooted $K_{2,t'}$ in (H,R_u) as follows. The set $X' \cap Z$ is one of the branch sets of the vertices of degree *t* ′ . The *t* ′ branch sets of the vertices of degree 2 will be modeled by *t* ′ well-chosen sets among the \hat{X}_i s, as we explain below. Observe first that indeed, $H[X' \cap Z]$ is connected (since Z is a clique in *H*), each X_i contains a root (by definition) and each X_i sends an edge to $X' \cap Z$, since $H[X']$ is connected. Furthermore, each X_i intersects C_F . By the connectivity properties of this cycle, each X_i has at least one neighbor in $C_F \cap Y$. Choose one arbitrarily. Recall that $H[Y] - M$ decomposes into at most ℓ^2 connected components. Thus, by the pigeonhole principle there is one connected component of $H[Y]$ – *M* that is adjacent to at least $t''/\ell^2 = t'$ of the X_i s. Choosing those X_i s for the branch sets of the vertices of degree 2 and the vertex set of that connected component of *H*[*Y*]− *M* for the last branch set completes the description of our rooted model of $K_{2,t'}$ in (H,R_u) , This is a contradiction with the fact that (H, R_u) does not contain a rooted $K_{2,t'}$ -minor.

Therefore, $p + q < 2t''$. Note that $((\ell + 2) \cdot (2^{\ell^3 + 1} \cdot t \cdot \ell^2)^2)/(2t'')^2 = \ell + 2$. Hence by Claim [75,](#page-51-1) there are *i* ∈ [*p*] and *j* ∈ [*q*] such that {*X_i* ∩ *R_F*, *Y_j* ∩ *R_F*} are ℓ + 2-interlacing with respect to *F*. By Lemma [74,](#page-50-0) this implies a contradiction with the Euler genus of the surface embedding of $G^{\#}[B_{l}^{\prime}]$ u' _{*u*}</sub> $\overline{}$ *– Z* (if the model uses connectivity through some child, this can be captured by the virtual edges in $G^{\#}[B_{i}^{\dagger}]$ u'_{u}] – *Z*). \Box

6.10 Extending to subdivisions

We extend Theorem [9](#page-9-0) to subdivisions of rooted 3-connected graphs (*G*,*R*), where roots have degree at least 3. Let $t \in \mathbb{Z}_{\geq 0}$ be such that (G, R) does not contain a rooted $K_{2,t}$ -minor. We denote by (G', R) the rooted graph obtained from (*G*,*R*) by contracting each subdivided edge to a single edge. Note that by Lemma [45,](#page-29-2) (G', R) does not contain a rooted $K_{2,t}$ -minor. Furthermore, the sets of roots coincide since we never remove a vertex from the set of roots, and never contract an edge between two roots.

Lemma 76. Let $\ell \in \mathbb{Z}$. Given an ℓ -special tree-decomposition (T', \mathcal{B}') of G' , where $\mathcal{B}' = \{B'_{\ell}\}$ u' : $u \in V(T')$ }, *we can efficiently compute an* ℓ *-special tree-decomposition* (T, \mathcal{B}) of G, where $\mathcal{B} = \{B_u : u \in V(T)\}$, such *that* $T = T'$ *and* $R \cap B_u = R \cap B'_u$ u' for each $u \in V(T)$.

Proof. Let $F \subseteq E(G')$ denote the edges of *G*' that correspond to subdivided edges of *G*. Each $e \in F$ induces a path in *G*, which we denote by $P(e) := v_0, e_1, v_1, \ldots, v_p, e_{p+1}, v_{p+1}$, such that each v_i with $i \in [p]$ has degree 2 in *G*, and v_0 , v_{n+1} have degrees at least 3 in *G*.

Let u_0 be the root of the decomposition tree *T'*, and let $vv' = e \in F$. By property [\(ii\)](#page-29-3) of Definition [46,](#page-29-1) there is a node $u \in V(T')$, such that $v, v' \in B'_t$ \mathcal{U}_u . By property [\(iii\)](#page-29-4) of Definition [46,](#page-29-1) the bags of (T',\mathcal{B}') that contain both *v* and *v*['] correspond to a subtree of *T*['], which we denote by T_e' \overline{u}_e [']. Let \overline{u}' denote the least common ancestor of T'_e with respect to u_0 . Then we insert the inner nodes v_i for $i \in [p]$ of $P(e)$ into B'_i '_{u'},

in order to obtain a collection of subsets of $V(G)$, denoted by \mathcal{B} . It is easy to see that the resulting pair (*T*, \mathcal{B}) is an ℓ -special tree-decomposition of *G* having the desired properties. \Box

Lemma 77. *Let* $R' \subseteq R$ *be a subset of the roots. Then,* $\mathcal{P}(G, R') \subseteq \mathcal{P}(G', R')$ *.*

Proof. Consider an element $P \in \mathcal{P}(G, R')$, as well as $S \in \mathcal{S}(G)$ a docset of *G* with $S \cap R' = P$. Observe that the restriction of *S* to *G'* is a docset of *G'*. Thus $P = S \cap R' \in \mathcal{P}(G', R')$. \Box

Proposition 78. *Let* (*G*,*R*) *be a subdivision of a* 3*-connected rooted graph, such that each v* ∈ *R has degree at least* 3*, and* (*G*,*R*) *has no rooted K*2,*t-minor. Then the statement of Theorem [9](#page-9-0) applies, that is, there is a polynomial-time algorithm that outputs an f*[9](#page-9-0)(*t*)*-special tree-decomposition* (*T*,B) *of G and a collection* ${X_u : u ∈ V(T)}$ *where each* X_u *is a docset superprofile of* $R ∩ B_u$ *in G of size polynomial in* |*V*(*G*)|.

Proof. By Theorem [9,](#page-9-0) we obtain an $f_9(t)$ $f_9(t)$ $f_9(t)$ -special tree-decomposition (T, \mathcal{B}) of G' and a collection $\{\mathcal{X}_u:$ $u \in V(T)$ } where each \mathcal{X}_u is a docset superprofile of R_u in G' (of size polynomial in $|V(G)|$). By Lemma [76,](#page-52-0) and Lemma [77,](#page-53-2) both the tree-decomposition and the superprofiles extend to *G*. \Box

7 Discussion

In this paper, we initiated the study of integer programs (IPs) on constraint matrices that simultaneously have bounded subdeterminants and are nearly totally unimodular (TU), in the sense that they become TU after the deletion of a constant number of rows and columns. In view of the state of the art [\[AWZ17,](#page-54-0) [FJWY22,](#page-56-8) [NSZ22,](#page-58-3) [NNSZ23\]](#page-57-2), and also in view of the matroid minors project [\[GGW14,](#page-56-15) [GGW15\]](#page-56-9), this is a natural case of the conjecture that bounding all subdeterminants suffices to guarantee that an IP can be solved in polynomial time.

We reduced to the case in which the linear constraints are $Ax = b$, $Wx = d$, $\ell \le x \le u$, where *A* is a TU matrix and *W* is an integer matrix with a constant number of rows, see (IP_1) . The combinatorial structure of *A*, more specifically the collection of *circuits* of *A*, plays a crucial role in our analysis. We showed that bounding the subdeterminants of the original constraint matrix translates to bounding the weight of every circuit of *A* with respect to any row of *W*. Moreover, we proved that some optimal solution \bar{z} of the IP can be reached from some integer point *z*, that can be efficiently found through solving the LP relaxation, by adding (or augmenting on) a *bounded* number of circuits of *A*.

We proved that we can reduce in a black-box fashion to the case where *A* almost 3-connected. Furthermore, we proved that an optimal augmentation can be found efficiently in the cographic case, that is, when the circuits of *A* correspond to the minimal cuts of a graph. This leaves open the question of efficiently finding an optimal augmentation in case *A* is a *general* TU matrix.

By Seymour's decomposition of regular matroids [\[Sey80a\]](#page-58-4), if *A* is 4-connected then there exists a graph *G* such that the circuits of *A* correspond either to the circuits of *G* (graphic case), or to the minimal cuts of *G* (cographic case). Since we solved the cographic case in the present paper, the graphic case seems to be the second and last main case to consider. We have preliminary observations concerning this.

First, we expect that, in the graphic case, *G* admits a tree-decomposition similar to that of Theo-rem [52,](#page-33-0) replacing rooted $K_{2,t}$ minors with the dual class of obstructions. In view of this, the case where *G* can be embedded in a fixed surface Σ seems once again particularly interesting. If Σ is orientable, then we can reduce the problem to the cographic case, via taking the dual of the directed graph *G*. If Σ is non-orientable, then the dual of *G* is no longer a directed graph, but rather a bidirected graph. Thus, we are naturally led to consider the case where $A = \begin{bmatrix} B & I \end{bmatrix}$ and *B* is the transpose of a binet matrix. In fact, the "right" case to consider next toward the totally ∆-modular IP conjecture after the cographic case might be "nearly binet or transposed binet" case. We remark that for the corresponding instances, large vortices are unavoidable in near-embeddings and have to be dealt with in the optimization phase. To achieve this, the ideas of [\[FJWY22\]](#page-56-8) should be very useful.

Second, we leave open the task of handling 3-separations of *A*. Achieving this in a similar way as we dealt with 2-separation seems challenging. (The fact that 3-sums are much more difficult to handle than 1- and 2-sums is a recurring theme in recent work using Seymour's decomposition, see for instance [\[AF22,](#page-54-2) [AWZ17\]](#page-54-0).) Despite our efforts, we are not aware of any family of gadgets that could be used to reduce the problem in a black-box fashion to the case where the 3-separations of *A* are "under control". Instead, we suggest to view 3-separations of A as small order separations of $\left[\begin{smallmatrix} A \ W \end{smallmatrix} \right]$ and develop more systematic ways to handle small order separations in $[\begin{smallmatrix}A\V\end{smallmatrix}]$, if possible extending also the decomposition technique we developed for graphs forbidding a rooted $K_{2,t}$ -minor.

Third, we observe that the graphic case of our problem contains instances of (the optimization version of) the famous *red-blue matching problem* (also known as the *exact matching problem*): given a bipartite graph *G* whose edges are colored either red or blue, a positive integer *k*, and integer profits on the edges of *G*, find a maximum profit perfect matching of *G* containing exactly *k* red edges. The resulting IP reads max{ $p^{\top}x$: $Ax = 1$, $0 \le x \le 1$, $\sum_{e \text{ red}} x(e) = k$, $x \in \mathbb{Z}^{E(G)}$ }, where $A \in \{0,1\}^{V(G) \times E(G)}$ is the incidence matrix of *G*. Bounding the subdeterminants of the constraint matrix of this IP by ∆ (in absolute value) entails asking that every single augmentation changes the number of red edges in a (perfect) matching by at most ∆. (More precisely, the constraint matrix of the IP has all its subdeterminants bounded by ∆ if and only if the weight of every circuit in G augmented with an extra vertex ν_0 adjacent to all vertices through a zero-weight edge is at most ∆. Rewriting the IP in inequality form allows to get rid of *v*0, and consider a less restricted set of instances.) There is a randomized polynomial-time algorithm to check whether the IP is feasible, even without assuming that the subdeterminants are bounded, see [\[MVV87\]](#page-57-7). We believe that there is a deterministic polynomial time algorithm to solve the above IP to optimality, provided that the subdeterminants are bounded.

Acknowledgements

We thank Sebastian Wiederrecht for helpful discussions regarding [\[TW22a,](#page-59-1) [TW22b\]](#page-59-2).

Manuel Aprile and Samuel Fiorini acknowledge funding from *Fonds de la Recherche Scientifique* - FNRS through research project BD-DELTA (PDR 20222190, 2021–24). Samuel Fiorini was also funded by *King Baudouin Foundation* through project BD-DELTA-2 (convention 2023-F2150080-233051, 2023–26). Gwenaël Joret and Michał Seweryn acknowledge funding from FNRS (PDR "Product structure of planar graphs"). Stefan Kober and Stefan Weltge were supported by the Deutsche Forschungsgemeinschaft (German Research Foundation) under the projects 277991500/GRK2201 and 451026932, respectively. Yelena Yuditsky was supported by FNRS as a Postdoctoral Researcher.

References

van. Detecting high log-densities: An *O*(*n* 1/4) approximation for densest k-subgraph. In

Proceedings of the Forty-Second ACM Symposium on Theory of Computing, STOC '10, pages 201–210, New York, NY, USA, 2010. Association for Computing Machinery.

- [BDSE+14] Nicolas Bonifas, Marco Di Summa, Friedrich Eisenbrand, Nicolai Hähnle, and Martin Niemeier. On sub-determinants and the diameter of polyhedra. *Discrete & Computational Geometry*, 52(1):102–115, 2014.
- [BFMRV14] Adrian Bock, Yuri Faenza, Carsten Moldenhauer, and Andres Jacinto Ruiz-Vargas. Solving the stable set problem in terms of the odd cycle packing number. In *34th International Conference on Foundation of Software Technology and Theoretical Computer Science*, volume 29 of *LIPIcs. Leibniz Int. Proc. Inform.*, pages 187–198. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2014.
- [BKK⁺24] Marcin Briański, Martin Koutecký, Daniel Kráľ, Kristýna Pekárková, and Felix Schröder. Characterization of matrices with bounded graver bases and depth parameters and applications to integer programming. *Mathematical Programming*, 2024.
- [BKMM08] Thomas Böhme, Ken-ichi Kawarabayashi, John Maharry, and Bojan Mohar. *K*3,*^k* -minors in large 7-connected graphs, 2008.
- [BM02] Thomas Böhme and Bojan Mohar. Labeled *K*2,*^t* minors in plane graphs. *Journal of Combinatorial Theory, Series B*, 84(2):291–300, 2002.
- [Bou78] André Bouchet. Orientable and nonorientable genus of the complete bipartite graph. *Journal of Combinatorial Theory, Series B*, 24(1):24–33, 1978.
- [CE80] William H Cunningham and Jack Edmonds. A combinatorial decomposition theory. *Canadian Journal of Mathematics*, 32(3):734–765, 1980.
- [CEH+21] Jana Cslovjecsek, Friedrich Eisenbrand, Christoph Hunkenschröder, Lars Rohwedder, and Robert Weismantel. Block-structured integer and linear programming in strongly polynomial and near linear time. In *Proceedings of the Thirty-Second Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '21, pages 1666–1681, USA, 2021. Society for Industrial and Applied Mathematics.
- [CEP+21] Jana Cslovjecsek, Friedrich Eisenbrand, Michał Pilipczuk, Moritz Venzin, and Robert Weismantel. Efficient Sequential and Parallel Algorithms for Multistage Stochastic Integer Programming Using Proximity. In Petra Mutzel, Rasmus Pagh, and Grzegorz Herman, editors, *29th Annual European Symposium on Algorithms (ESA 2021)*, volume 204 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 33:1–33:14, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [CFH+20] Michele Conforti, Samuel Fiorini, Tony Huynh, Gwenaël Joret, and Stefan Weltge. The stable set problem in graphs with bounded genus and bounded odd cycle packing number. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 2896–2915. SIAM, 2020.
- [CFHW20] Michele Conforti, Samuel Fiorini, Tony Huynh, and Stefan Weltge. Extended formulations for stable set polytopes of graphs without two disjoint odd cycles. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 104–116. Springer, 2020.
- [CGST86] William Cook, Albertus MH Gerards, Alexander Schrijver, and Éva Tardos. Sensitivity theorems in integer linear programming. *Mathematical Programming*, 34(3):251–264, 1986.
- [CKL+24] Jana Cslovjecsek, Martin Koutecký, Alexandra Lassota, Michał Pilipczuk, and Adam Polak. *Parameterized algorithms for block-structured integer programs with large entries*, pages 740–751. SIAM, 2024.
- [CKPW22] Marcel Celaya, Stefan Kuhlmann, Joseph Paat, and Robert Weismantel. Improving the Cook et al. proximity bound given integral valued constraints. In Karen Aardal and Laura Sanità, editors, *Integer Programming and Combinatorial Optimization*, pages 84–97, Cham, 2022. Springer International Publishing.
- [CT21] Julia Chuzhoy and Zihan Tan. Towards tight(er) bounds for the excluded grid theorem. *J. Combin. Theory, Series B*, 146:219–265, 2021.
- [Dad12] Daniel Dadush. *Integer programming, lattice algorithms, and deterministic volume estimation*. PhD thesis, Georgia Institute of Technology, 2012.
- [DBT89] G. Di Battista and R. Tamassia. Incremental planarity testing. In *30th Annual Symposium on Foundations of Computer Science*, pages 436–441, 1989.
- [DF94] Martin Dyer and Alan Frieze. Random walks, totally unimodular matrices, and a randomised dual simplex algorithm. *Math. Program.*, 64(1-3):1–16, 1994.
- [DiKMW12] Reinhard Diestel, Ken ichi Kawarabayashi, Theodor Müller, and Paul Wollan. On the excluded minor structure theorem for graphs of large tree-width. *Journal of Combinatorial Theory, Series B*, 102(6):1189–1210, 2012.
- [EHK+22] Friedrich Eisenbrand, Christoph Hunkenschröder, Kim-Manuel Klein, Martin Koutecký, Asaf Levin, and Shmuel Onn. An algorithmic theory of integer programming, 2022.
- [ERW24] Friedrich Eisenbrand, Lars Rohwedder, and Karol W˛egrzycki. Sensitivity, proximity and FPT algorithms for exact matroid problems. *arXiv:2404.03747*, 2024.
- [EV17] Friedrich Eisenbrand and Santosh Vempala. Geometric random edge. *Math. Program.*, 164(1-2):325–339, 2017.
- [EW19] Friedrich Eisenbrand and Robert Weismantel. Proximity results and faster algorithms for integer programming using the Steinitz lemma. *ACM Transactions on Algorithms (TALG)*, 16(1):1–14, 2019.
- [FJWY22] Samuel Fiorini, Gwenaël Joret, Stefan Weltge, and Yelena Yuditsky. Integer programs with bounded subdeterminants and two nonzeros per row. In *2021 IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 13–24, 2022.
- [GGW14] Jim Geelen, Bert Gerards, and Geoff Whittle. Solving rota's conjecture. *Notices Amer. Math. Soc.*, 61:736–743, 2014.
- [GGW15] Jim Geelen, Bert Gerards, and Geoff Whittle. The highly connected matroids in minorclosed classes. *Annals of Combinatorics*, 19:107–123, 2015.
- [GM00] Carsten Gutwenger and Petra Mutzel. A linear time implementation of SPQR-trees. In *International Symposium on Graph Drawing*, pages 77–90. Springer, 2000.

A Proof of Theorem [8](#page-8-3)

We provide a proof of the following remark from [\[BKMM08\]](#page-55-8).

Theorem [8](#page-8-3). *There is a function* f_8 : $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 2}$ *, such that the following holds. Let* (G, R) *be a* 3*connected rooted graph without a rooted K*2,*t-minor, embedded in a surface of Euler genus g with facewidth at least f*[8](#page-8-3)(*g* ,*t*)*. Then G has a collection of at most f*[8](#page-8-3)(*g* ,*t*) *facial cycles covering all the roots.*

A.1 Finding a large Möbius wall

In non-orientable graphs, it will be helpful for us to find large twisted grids. Assuming large face-width, we follow the arguments by Reed [\[Ree99\]](#page-58-18) (for a shorter proof, see Kawarabayashi and Reed [\[KR10\]](#page-57-14)).

Given an integer $h \geq 2$, the *elementary wall of height h* is the graph obtained from the $2h \times h$ grid with vertex set $[2h] \times [h]$ by removing all edges with endpoints $(2i - 1, 2j - 1)$ and $(2i - 1, 2j)$ for all $i \in [h]$ and *j* ∈ [$[h/2]$], all edges with endpoints (2*i*,2*j*) and (2*i*,2*j* + 1) for all *i* ∈ [*h*] and *j* ∈ [[$(h-1)/2$]]. Note that unlike some definitions in the literature, we do not delete the two vertices of degree one in the resulting graph. For integers $k, h \ge 0$ with $1 \le k \le h/2$, the k -th *outermost cycle* in the elementary wall of height *h* is the only cycle contained in the subgraph induced by the set $\{2k-1, 2k, h-2k+1, h-2k+2\} \times \{k, h-k+1\}$. A subdivision of an elementary wall of height *h* is called a *wall of height h*. The notion of *k*-th outermost cycle naturally generalizes from elementary walls to general walls.

For an even integer $h \geq 2$, the *elementary Möbius wall of height* h is obtained from the elementary wall of height *h* by adding an edge with endpoints (2*i* − 1,1) and (2*h* − 2*i* + 1,*h*) for each *i* ∈ [*h*]. For an integer *k* with $1 \le k \le h/2$, the *k*-th outermost cycle in the elementary Möbius wall of height *h* is the only cycle contained in the subgraph induced by $\{2k-1, 2k, 2h-2k+1, 2h-2k+2\} \times [h]$. A cycle of length six in an elementary Möbius wall is called a *brick*. A subdivision of an elementary Möbius wall of height *h* is a *Möbius wall of height h*. The notion of *k*-th outermost cycle and a brick naturally generalizes from elementary Möbius walls to general Möbius walls. An *Escher wall* is a non-bipartitie Möbius wall in which every brick has even length.

Theorem [79](#page-60-1) (Reed [\[Ree99,](#page-58-18) Thm. 1]). *There is a computable function* t_{79} : $\mathbb{Z}^2_{\geq 0} \to \mathbb{Z}_{\geq 0}$ such that for all *integers k*,*h* ≥ 0*, if G is a graph with neither k vertex disjoint odd cycles nor an Escher wall of height h then there is an odd cycle cover X of G with* $|X| \le t_{79}(k, h)$ $|X| \le t_{79}(k, h)$ $|X| \le t_{79}(k, h)$ *.*

We call a Möbius wall *flat*, if every facial cycle of the wall corresponds to a contractible curve in the surface.

Corollary 80. Let $h \geq 1$. Let G be a graph embedded in the projective plane with face-width at least *t*[79](#page-60-1)(2,*h*)+1*. Then G contains a flat Möbius wall of height h.*

Proof. For a graph *G* embedded in the projective plane, we have a *signature* λ : $E(G) \rightarrow \{-1,1\}$, such that a cycle *C* is one-sided if and only if $\prod_{e \in C} \lambda(e) = -1$. We produce an auxiliary graph *G'* by replacing every edge $e \in E$ with $\lambda(e) = 1$ by a (once) subdivided edge e_1, e_2 with $\lambda(e_1) = \lambda(e_2) = 1$. Observe that $fw(G) = Tw(G')$, *G'* is embedded in the projective plane, and a cycle of *G'* is one-sided if and only if it is odd.

Since *G'* is embedded in the projective plane, it has no two vertex-disjoint one-sided (odd) cycles. Assume that we find $X \subseteq G'$ with $|X| \le t_{79}(2, h)$ $|X| \le t_{79}(2, h)$ $|X| \le t_{79}(2, h)$ such that $G'' := G' - X$ is bipartite. Since G'' does not contain a one-sided cycle, it is planar. This contradicts that $fw(G') \geq t_{79}(2, h) + 1$ $fw(G') \geq t_{79}(2, h) + 1$ $fw(G') \geq t_{79}(2, h) + 1$.

By Theorem [79,](#page-60-1) *G* ′ contains an Escher wall of height *h*. Clearly, this implies that *G* contains a Möbius wall of height *h*. In addition, this Möbius wall is flat, since the Möbius wall has a unique embedding in the projective plane if the facewidth is sufficiently large [\[Moh95\]](#page-57-15). \Box

A.2 Planarization

Let G' be a graph embedded in a disjoint union of surfaces Σ' . We define two operations which transform *G'* and Σ' into a graph *G* embedded in a disjoint union of surfaces Σ: adding a crosscap and adding a handle.

For an even-length facial cycle $C' = x_1'$ $x'_1 \cdots x'_k$ $k \cdot x_1''$ $x''_1 \cdots x''_k$ $\frac{u}{k}x_1'$ $\frac{1}{1}$ in *G'*, we *add a crosscap* at *C* by deleting in Σ' the face bounded by C' and identifying the two arcs between x_1 ['] x_1' and x_1'' $\frac{1}{1}$ in the loop $\bigcup E(C')$ so that for each $i \in \{1, ..., k\}$, the vertices x_i' \int_{i} an x''_i \int_{i}^{b} are identifed into a new vertex x_i , and the cycle C' turns into a cycle $C = x_1 \cdots x_k x_1$. The resulting union of disjoint surfaces Σ has the same components as Σ' except that the component of Σ ′ containing the cycle *C* ′ is replaced by a surface with Euler genus greater by one.

For two facial cycles of the same length $C' = x_1'$ $\frac{1}{1} \cdots x'_{k}$ $\int_k' x'_1$ x_1' and $C'' = x_1''$ $x''_1 \cdots x''_k$ $\int_k'' x_1''$ \int_1'' in *G*', we *add a handle* between *C'* and *C''* by deleting the faces bounded by C^{\prime} and C'' in Σ' and identifying the loops $\bigcup E(C')$ and $\bigcup E(C''$ so that for each $i \in \{1, ..., k\}$, the vertices x'_i \int_{i} and x''_i a''_i are identified into a new vertex x_i , and the cycles *C'* and *C''* are identified into a cycle $C = x_1 \cdots x_k x_1$. This operation is called *adding a handle*. The resulting sum of disjoint surfaces Σ has the same components as Σ' except that the one or two components of Σ' containing *C'* and *C''* are replaced by the component Σ_{*C*} of Σ containing *C*. If *C'* and *C''* are contained in one component $\Sigma'_{C',C''}$ of Σ' , then the Euler genus of Σ_C is the Euler genus of $\Sigma'_{C',C''}$ plus two, and if *C'* and *C''* are contained in distinct components $\Sigma'_{C'}$ and $\Sigma'_{C'}$, then the Euler genus of Σ_C is the sum of the Euler genera of $\Sigma_{C'}'$ and $\Sigma_{C'}'$.

The operation inverse to adding a crosscap or a handle is *cutting* along a cycle. Let *G* be a graph embedded in a disjoint union of surfaces Σ , and let $C = x_1 \cdots x_k x_1$ be a cycle in *G*. Then, cutting Σ and *G* along *C* yields a disjoint union of surfaces Σ' and a graph *G'* embedded in it, where each x_i will have two copies x_i' \int_{i} and x''_i \int_{i}^{n} in *G*_{*i*}. If *C* is one-sided, then there will exist a facial cycle *C*^{*'*} = *x*¹ $x'_1 \cdots x'_l$ k' _k x_1'' $x''_1 \cdots x''_k$ $\frac{u}{k}x_1'$ 1 in *G'* such that Σ and *G* can be restored by adding a crosscap at *C'* to *G'* and Σ'. If *C* is two-sided, then there will exist facial cycles $C' = x_1'$ $x'_1 \cdots x'_k$ $\int_k' x'_1$ \int_{1}^{7} and $C'' = x_1''$ $x''_1 \cdots x''_k$ $\int_k^l x''_1$ $\frac{1}{1}$ in *G*['] such that Σ and *G* can be restored by adding a handle between C' and C'' to G' and Σ'

A *planarizing set of cycles* is a set of disjoint cycles *€* in *G* such that cutting *G* and Σ along the cycles from *'€* yields a graph *G'* embedded in a disjoint union of spheres Σ'. Note that we will obtain the same *G*' and Σ' regardless of the order in which we choose the cycles from $\mathscr C$ to cut along. Observe that if $\mathscr C$ is an inclusion-minimal planarizing set of cycles, then Σ ′ has only one component, and thus is a sphere. If \mathscr{C}' denotes the set of cycles in *G*['] corresponding to the cycles from \mathscr{C} in *G*, then the tuple $(\mathscr{C}, G', \Sigma', \mathscr{C}')$ is a *planarization* of *G* and Σ.

For a graph *G* embedded in a disjoint union of surfaces Σ, a facial cycle *C* and an integer *d* ≥ 1, a *depth-d nest* around *C* is a sequence of disjoint cycles $\mathcal{D} = (D_0, \ldots, D_d)$ in *G* such that $D_0 = C$, and for each $i \in \{1, ..., d\}$, the loop $\bigcup E(D_i)$ is the boundary of a closed disk Δ_i in Σ which contains the cycles *D*₀,...,*D*_{*i*−1}. A cycle *C*['] in *G* is *outside* the nest \mathcal{D} if *C*['] is disjoint from the disk Δ_d . Two depth-*d* nests $\mathscr{D}' = (D'_0)$ $'_{0},\ldots,D'_{\ell}$ \mathcal{Q}'_d) and $\mathcal{D}'' = (D_0'')$ $\binom{n}{0}, \ldots, D_d''$ d ^{*d*}) around some facial cycles are *disjoint* if $V(D)$ ^{*i*} $\binom{l}{i} \cap D'$ j_j) = \emptyset for *i*, *j* ∈ $\{1,\ldots,d\}.$

Theorem 81 (Yu [\[Yu97,](#page-59-7) Thm. 4.3]). *Let* $g, d \ge 1$ *be integers. Let* Σ *be a surface with Euler genus g, and let G be a graph embedded in* Σ *with face-width at least* $w_{81}(g, d) = 8(2d + 1)(2^g - 1)$ $w_{81}(g, d) = 8(2d + 1)(2^g - 1)$ $w_{81}(g, d) = 8(2d + 1)(2^g - 1)$ *. Then there exists* a planarization ($\mathscr{C}, G', \Sigma', \mathscr{C}'$) of G and a depth-d nest $\mathscr{D}(C')$ in G' around each $C' \in \mathscr{C}'$ such that for any C' , $C'' \in \mathscr{C}'$, the nests $\mathscr{D}(C')$ and $\mathscr{D}(C'')$ are disjoint and C' is outside $\mathscr{D}(C'')$ or C'' is outside $\mathscr{D}(C')$.

A *holed surface* is a topological space Π which can be obtained from a surface Σ by deleting the interiors of disjoint open disks ∆1,...,∆*^c* ⊆ Σ. The boundaries of the disks ∆1,...,∆*^c* are the *cuffs* of Π. Note that the cuffs may intersect. The union of the cuffs is the *boundary* of Π.

For a cycle *C* in a graph *G* embedded in a surface Σ and an integer *r* ≥ 1, the *r -th neighborhood of C* is a subspace *N^r* (*C*) ⊆ Σ which is the topological closure of all points at distance at most *r* from *C* in Σ.

Let *G* be a graph embedded in a surface Σ with a sufficiently large face-width. Let *r* ≥ 1 be an integer, and let $(\mathscr{C}, G', \Sigma', \mathscr{C}')$ be a planarization of *G* obtained by applying Theorem [81](#page-61-0) with $d = 2r + 1$. Then, the *r*-th neighborhood of each cycle from \mathcal{C}' in Σ' is a holed sphere, and these neighborhoods are pairwise disjoint. The *r*-th neighborhood of a cycle $C \in \mathscr{C}$ in Σ is a holed projective plane if *C* is one-sided or a holed sphere if *C* is two-sided.

Lemma [82](#page-62-0). Let $g \ge 1$ *be an integer. Let* $r = t_{79}(2,6) + 1$ $r = t_{79}(2,6) + 1$ $r = t_{79}(2,6) + 1$ *, let* $d = 2r + 1$ *, and let* $w_{82}(g) = \max\{w_{81}(g,d), r\}$ $w_{82}(g) = \max\{w_{81}(g,d), r\}$ $w_{82}(g) = \max\{w_{81}(g,d), r\}$ *Let* Σ *be a surface with Euler genus g , and let G be a graph embedded in* Σ *with face-width at least w. Then there exist k* \leq 3*g and pairs of holed spheres* (Π') $'_{i}, \Pi''_{i}$ $\binom{n}{i}$ *in* Σ *for i* \in {1,...,*k*} *such that each* Π'_{i} *i is contained in the interior of* Π ′′ *i , every cuff of* Π ′ \int_i' or Π''_i *i coincides with a non-contractible cycle in G and every vertex of G is contained in the interior of some* Π ′ *i .*

Proof. Let $(\mathscr{C}, G', \Sigma', \mathscr{C}')$ be a planarization of *G* obtained by applying Theorem [81](#page-61-0) to *g*, *d*, Σ , and *G*, and let $\mathscr{D}(C')$ be the corresponding nests around the cycles $C' \in \mathscr{C}'$ in G' (note that all except the first cycle $D_0 = C'$ in $\mathscr{D}(C')$ are contained in *G* and disjoint from the cycles in \mathscr{C} . We assume that Σ' is a sphere, and thus, $|\mathscr{C}| \leq g$.

For an integer $s \ge 0$, and a cycle $C \in \mathcal{C}$, let $N_s(C)$ denote the set of all points of Σ at distance at most *s* from *C'*. Thus, for any *s* with $0 \le s \le r$, if *C* is one-sided, then $N_s(C)$ is a holed projective plane, and if *C* is two-sided, then $N_s(C)$ is a holed sphere. Furthermore, $N_s(C_1)$ and $N_s(C_2)$ are disjoint for distinct $C_1, C_2 \in \mathscr{C}$.

For each two-sided $C \in \mathscr{C}$, we define a pair of holed spheres (Π'(C), Π"(C)), where Π'(C) is the topo-*C* logical closure of the component of Σ \ $\bigcup_{C_1 \in \mathscr{C} \setminus \{C\}} N_r(C_1)$ containing $N_r(C)$ and $\Pi''(C)$ is the topological closure of the component of $\Sigma \setminus \bigcup_{C_1 \in \mathscr{C} \setminus \{C\}} N_{r-1}(C_1)$ containing $N_r(C)$.

For each one-sided $C \in \mathscr{C}$, we will define three pairs (Π' $\frac{i}{i}(C), \Pi''_i$ i' _{*i*}(*C*)) for *i* ∈ {1,...,3}. Together with the pairs $(\Pi'(C), \Pi''_i)$ $\binom{n}{i}(C)$) for two-sided cycles $C \in \mathscr{C}$, we will have up to 3*g* pairs satisfying the lemma.

First, for each one-sided $C \in \mathscr{C}$ we will find an Escher wall $W(C)$ of height 8 inside the holed projective plane *N^r* (*C*). We achieve this as follows. Let *H*(*C*) denote the subgraph of *G* consisting of the vertices and the edges contained in $N_r(C)$, and let $\Sigma(C)$ denote the projective plane obtained by capping the cuffs of *N*_{*r*}(*C*). Towards a contradiction, suppose that the face-width of *H*(*C*) in Σ(*C*) is less than *r*. Then, there is a noncontractible noose *o* in Σ(*C*) which intersects *H*(*C*) in fewer than *r* vertices. The noose *o* has to intersect *C*, and since the distance of *C* from the boundary of $N_r(C)$ is *r*, the noose *o* must be contained in the interior of $N_r(C)$, and thus is a non-contractible noose in Σ which intersects *G* in fewer than *r* vertices. This is a contradiction since the face-width of *C* is at least *w*, which is greater than or equal to *r* . Therefore, the face-width of *C* is at least *r* , and by Corollary [80,](#page-60-2) there exists a height-eight Escher wall $W(C)$ in $N_r(C)$.

For each one-sided $C \in \mathcal{C}$, we cover the height-eight Escher wall $W(C)$ with two (planar) walls $W_1(C)$ and $W_2(C)$ of height eight, so that their intersection is the disjoint union of two 2 × 8 walls. For $i \in \{1,2\}$ and *j* ∈ {1,2}, let *D*^(*j*) $\binom{f}{i}(C)$ denote respectively the *j*-th outermost cycle of *W*_{*i*}(*C*), and let *W*^(*j*) $\int_{i}^{(j)}(C)$ denote the height-(10−2*j*)subwall of $W_i(C)$ bounded by $D_i^{(j)}$ *i*^{*j*} (*C*). Similarly, for each *j* ∈ {1,...,4}, $D_3^{(j)}$ $3^{\prime\prime\prime}(C)$ denote the *j*-th outermost cycle of the Escher wall *W* (*C*).

For each $i \in \{1, 2\}$, we define a holed sphere Π_i' $\mathcal{L}_i^{\prime}(C)$ as the topological closure of the component of $\overline{C}_i^{(2)}(C)$, and a holed sphere Π_i'' $\Sigma \setminus (\bigcup E(D_i^{(2)}))$ *i*⁽²⁾(*C*))∪∪ $C_1 \in \mathscr{C} \setminus \{C\}$ *N_r*(*C*₁)) containing the interior of $W_i^{(2)}$ $i^{\prime\prime}(C)$ as the topological closure of the component of Σ \ ($\bigcup E(D_i^{(1)}\big)$ *i*⁽¹⁾(*C*))∪∪ $C_1 \in C_1$ (*C*₁)) containing the interior of $V_i^{(1)}(C) = W_i(C)$. Finally, let Π_i^{\prime} $W_i^{(1)}$ $\mathcal{S}'_3(C)$ denote the topological closure of the component of Σ \ ($\bigcup E(D_3^{(3)}\big)$ 3 (*C*))∪ $_3^{(1)}(C)$, and let Π_3'' $\bigcup_{C_1 \in \mathscr{C} \setminus \{C\}} N_r(C_1)$) which contains $D_3^{(1)}$ $C_3^{\prime\prime}(C)$ denote the topological closure of the component $^{(1)}_3$ (C). The resulting pairs (Π', Π'') satisfy the of Σ \ (∪ $E(D_3^{(4)}$ $B_3^{(4)}(C)$)∪ $\bigcup_{C_1 \in \mathscr{C} \setminus \{C\}} N_{r-1}(C_1)$) which contains $D_3^{(1)}$ lemma. \Box

A.3 Obtaining a planar minor

Lemma 83. *Let G be a 3-connected graph embedded in a surface* Σ *with face-width at least* 3*. Let* Π ′ *and* Π ′′ *be holed spheres in* Σ *such that* Π ′ *is contained in the interior of* Π ′′*, and every cuff of* Π ′ *or* Π ′′ *coincides with a non-contractible cycle in G. Then there exists a 3-connected plane minor G*′ *of G which contains the subgraph of G consisting of the edges and vertices in* Π ′ *.*

Proof. Let *G*["] denote the part of *G* embedded in Π["]. Let *G*' be the graph obtained from *G*["] by contracting the edge set of every component of $G'' - (V(G'') \cap \Pi')$ which intersects the boundary of Π'' . It remains to show that *G*["] is 3-connected.

Suppose that G' admits a 2-cut-set $\{v_1, v_2\}$. If all contracted vertices are in the same component of $G - \{v_1, v_2\}$, then $\{v_1, v_2\}$ is a 2-cut-set in *G*. Otherwise, there is a non-planarity on either side of the separation, implying that there is a non-contractible noose in *G* which intersects *G* in only two vertices, and is homotopic with a cuff of Π' , and thus non-contractible. This contradicts the assumption that the face-width is at least three. \Box

Theorem [7](#page-8-1). *[Theorem 1.2 of [\[BM02\]](#page-55-7)] There is a function* f_7 *:* $\mathbb{Z}_{\geq 1}$ \to $\mathbb{Z}_{\geq 1}$ *such that the following holds. Let G be a 3-connected, planar graph and let R* ⊆ *V* (*G*) *be a set of roots. Assume that* (*G*,*R*) *does not have a rooted K*_{2,*t*}-minor. Then G has a collection of at most $f_7(t)$ $f_7(t)$ $f_7(t)$ facial cycles such that each root is contained in *at least one of these cycles.*

Proof of Theorem [8.](#page-8-3) Let $f(g, t) = \max\{3g \cdot f_7(t), w_{82}(g)\}$ $f(g, t) = \max\{3g \cdot f_7(t), w_{82}(g)\}$. We apply Lemma 82 to *G* and obtain an integer $1 \leq k \leq 3g$, and a collection of pairs of holed spheres (Π') $'_{i}, \overline{\Pi}''_{i}$ $''_i$) in Σ. By Lemma [83](#page-63-0) applied to the pairs $(\Pi'_i$ $'_{i}, \Pi''_{i}$ *i*^{$'$}), there exists a collection of at most 3*g* 3-connected plane minors of *G* such that every *x* ∈ *V*(*G*) is incident with the same set of faces in *G* as in one of these faces. By Theorem [7,](#page-8-1) in each of these plane graphs we can find a set of at most $f_7(t)$ $f_7(t)$ $f_7(t)$ facial cycles which cover any terminals from *R* contained in the interior of the corresponding Π' . Hence, all terminals from *R* can be covered with at most 3g · $f_7(t)$ $f_7(t)$ $f_7(t)$ facial cycles, as claimed. \Box