

A POISONOUS EXAMPLE TO EXPLICIT RESOLUTIONS OF UNBOUNDED COMPLEXES

DOLORS HERBERA, WOLFGANG PITSCHE, MANUEL SAORÍN, AND SIMONE VIRILI

ABSTRACT. We show that various methods for explicitly building resolutions of unbounded complexes in fact fail when applied to a rather simple and explicit complex. We show that one way to rescue these methods is to assume Roos (Ab.4*)- k axiom, which we adapt to encompass also resolutions in the framework of relative homological algebra. In the end we discuss the existence of model structures for relative homological algebra for unbounded complex under the relative (Ab.4*)- k condition, and present a variety of examples where our results apply.

CONTENTS

Introduction	1
1. Preliminaries and notations	3
2. DG-injective resolutions of complexes via truncations	5
3. The “go-to” counter-example for unbounded resolutions	9
4. Cartan-Eilenberg resolutions and their totalization	11
5. Sanblidze’s construction through multicomplexes	14
6. Ding and Yang’s construction via repeated killing of coboundaries	15
7. Model structures for relative homological algebra	19
8. Examples and applications	23
References	32

INTRODUCTION

After the publication of Verdier’s seminal thesis in 1967 (see [Ver96] for the later complete publication), the study of the derived categories of Abelian categories has received considerable attention and found widespread applications across many areas of abstract mathematics. It is important to underline that the initial focus was mostly restricted to bounded or half-bounded derived categories, where it was already known that, when starting with an Abelian category with enough injectives (or projectives), each lower (upper) bounded complex is quasi-isomorphic to a lower (upper) bounded complex of injectives (projectives). Additionally, the morphisms in the derived category between two such bounded complexes of injectives (projectives) correspond precisely to the homotopy classes of chain maps between them. This suffices to ensure that the corresponding half-bounded derived categories are locally small, i.e., they have small Hom-sets. In contrast, the extension of this result to unbounded derived categories presented a considerable challenge.

One of the main difficulties that arises in the unbounded setting is the necessity to find, for each complex, a quasi-isomorphism to a complex of injectives that, furthermore, is K -injective

Date: July 16, 2024.

2020 *Mathematics Subject Classification.* Primary 18G25, Secondary 18E10, 18G10.

The first and the fourth authors were partially supported by the projects MIMECO PID2020-113047GB-I00 financed by the Spanish Government. The first, second, and fourth authors are members of the LIGAT (Interactions Lab between Geometry, Algebra, and Topology), Research Group of the Generalitat de Catalunya number 2021 SGR 01015. The second author was supported by the FEDER/MEC grant “Homotopy theory of combinatorial and algebraic structures” PID2020-116481GB-I00. The third author was supported by the grant PID2020-113206GB-I00, funded by MCIN/AEI/10.13039/501100011033, and the project 22004/PI/22 of the “Fundación Séneca” (Murcia).

(or, dually, from a K -projective complex of projectives). In other words, one needs to build DG-injective (DG-projective) resolutions of unbounded complexes. Even so, such constructions were not available at the time, even for commonly used Grothendieck categories, like categories of modules or of quasi-coherent sheaves. The first breakthrough in this direction was pioneered by Spaltenstein [Spa88], who was able to build a DG-projective resolution for each $X^\bullet \in \text{Ch}(\text{Mod}(R))$, with R a ring, and a DG-injective resolution for each $X^\bullet \in \text{Ch}(\mathcal{A})$, where $\mathcal{A} = \text{Sheaf}_\Theta(X)$ is the category of sheaves of modules, over a sheaf of rings Θ on a topological space X . While Spaltenstein’s approach does not extend to general Grothendieck categories, his constructions have the advantage of not being merely existence results, as they rely on very explicit constructions of resolutions.

To the best of our knowledge, the first complete proof of the existence of DG-injective resolutions of unbounded complexes over a completely general Grothendieck category was published by Hovey [Hov01], although a very similar argument had already been outlined by Joyal in a private correspondence with Grothendieck (see [Joy84]). Both authors use variations of the small object argument to establish the existence of DG-injective resolutions in complete generality. However, the resolutions produced by these methods are non-explicit in nature, rendering them impractical for the computation of DG-resolutions in concrete situations.

At this point, the following questions arise naturally:

Questions. *Let \mathcal{A} be a complete Abelian category with enough injectives, and take $X^\bullet \in \text{Ch}(\mathcal{A})$.*

(Q.1) *Is it always possible to find a DG-injective resolution for X^\bullet ?*

(Q.2) *If X^\bullet has a DG-injective resolution, can one build such a resolution explicitly (i.e., with a step-by-step procedure that allows for a complete description of the final result, thus excluding, in principle, all constructions based on the small object argument)?*

In the literature, one can find two separate papers – [San07] and [YD15] – each claiming to give positive answers to both questions. Unfortunately, both constructions fail when applied to the “poisonous example” mentioned in the title, which is an unbounded complex $X^\bullet \in \text{Ch}(\mathcal{G})$, where \mathcal{G} is a localization of $\text{Mod}(R)$, with R being Nagata’s classical example of a commutative Noetherian ring with infinite Krull dimension (see [Nag62]). Let us clarify that we are not claiming that the above questions have a negative answer, and so we do not discard the possibility that the derived category $\mathcal{D}(\mathcal{A})$ is locally small whenever \mathcal{A} is a complete Abelian category with enough injectives. For example, for our “poisonous example” X^\bullet , both (Q.1) and (Q.2) can be answered in the positive. Indeed, take a DG-injective resolution $X^\bullet \rightarrow E^\bullet$ in $\text{Ch}(\text{Mod}(R))$ (e.g., using Spaltenstein’s construction); the inclusion of the torsion part $T^\bullet \leq E^\bullet$ is degreewise split (and, arguably, easy to describe), so E^\bullet/T^\bullet is a complex of torsion-free injectives. In particular, this quotient lives in $\text{Ch}(\mathcal{G})$, and $X^\bullet \rightarrow E^\bullet/T^\bullet$ is a DG-injective resolution (see Section 3 for details).

On the positive side, the second named author together with Chacholski, Neeman and Scherer [CNPS18] gave affirmative answers to both questions, adapting Spaltenstein’s construction, under the extra hypothesis that the category \mathcal{A} is $(\text{Ab}4^*)$ - k (in the sense of Roos [Roo06]), for some $k \in \mathbb{N}$. More generally, given an injective class (i.e., a preenveloping class closed under summands) $\mathcal{I} \subseteq \mathcal{A}$, they proved that any complex $X^\bullet \in \text{Ch}(\mathcal{A})$ has an \mathcal{I} -fibrant replacement, provided \mathcal{A} satisfies the $(\text{Ab}4^*)$ - \mathcal{I} - k condition – a version of Roos’ $(\text{Ab}4^*)$ - k relative to \mathcal{I} . To recover the results about DG-injective resolutions, it is then enough to take $\mathcal{I} = \text{Inj}(\mathcal{A})$, the class of all injective objects in \mathcal{A} . The existence of \mathcal{I} -fibrant replacements, when combined with results of [CH02], allows us to build and study a suitable model structure in $\text{Ch}(\mathcal{A})$ whose homotopy category is the \mathcal{I} -derived category $\mathcal{D}(\mathcal{A}; \mathcal{I})$ from [CNPS18, CH02]; the usual derived category $\mathcal{D}(\mathcal{A})$ is equivalent to $\mathcal{D}(\mathcal{A}; \text{Inj}(\mathcal{A}))$.

The paper is organized as follows: In Section 1, after fixing the needed conventions and notations about cochain complexes, we review some basic definitions and results regarding weak factorization systems, model categories, cotorsion pairs and Hovey’s correspondence. In Section 2 we show that, whenever \mathcal{A} is a complete $(\text{Ab}4^*)$ - k Abelian category with enough injectives, any $X^\bullet \in \text{Ch}(\mathcal{A})$ has a DG-injective resolution. Although this may be deduced from more general results in [CNPS18], we include a short direct proof. The “poisonous example” mentioned in the title is introduced and studied in Section 3, where it is used to show that the $(\text{Ab}4^*)$ - k condition is really necessary for the constructions of Section 2. Section 4 is devoted to injective Cartan-Eilenberg (CE)-resolutions: after reviewing the definitions and some classical results, we prove that, to build a DG-injective resolution of an $X^\bullet \in \text{Ch}(\mathcal{A})$ (with \mathcal{A} as in Section 2) one can just totalize a CE-resolution of X^\bullet .

Furthermore, this strategy fails when applied to our “poisonous example”, showing that (Ab.4*)- k is necessary also in this approach. In Section 5 we briefly review the construction of [San07] and we show that it fails to produce a DG-injective resolution for the “poisonous example”. Similarly, in Section 6, we review the construction of [YD15] and we show that it does not work when applied to our “poisonous example”, also observing that the gap in the argument disappears if one assumes that the ambient category is (Ab.4*). In Section 7, we concentrate on the construction of the so-called \mathcal{I} -injective model structure on $\text{Ch}(\mathcal{A})$, relative to a given injective class $\mathcal{I} \subseteq \mathcal{A}$. After recalling some general results from [CH02], we describe the construction of Spaltenstein towers partial \mathcal{I} -injective resolutions given in [CNPS18]. In the last part of the section, we prove a number of results related to the (Ab.4*)- \mathcal{I} - k condition, showing that it implies the existence of the \mathcal{I} -injective model structure on $\text{Ch}(\mathcal{A})$. Finally, in Section 8, we include a number of examples and applications of the formalism introduced in Section 7. In particular, we first consider those injective classes with the additional property of being cogenerating, we then study the injective classes contained in $\text{Inj}(\mathcal{A})$, and the injective class of the pure-injective objects in a locally finitely presented Grothendieck category. Finally, we introduce a notion of global dimension for \mathcal{A} relative to an injective class \mathcal{I} ; if this invariant is finite, then \mathcal{A} is (Ab.4*)- \mathcal{I} - k . Using the relative global dimension, we give a new characterization of the n -tilting cotorsion pairs.

Acknowledgments. The initial results that persuaded us to write this paper date back to 2019, during an “Intensive Research Period in Small Groups” at the CRM (Barcelona), supported by the Barcelona Graduate School of Mathematics (BGSMath) as a María de Maeztu Excellence Centre 2015–2019.

We are glad to thank Sergio Estrada for several fruitful conversations on the topics of this paper and, in particular, for sharing with us his doubts regarding [YD15].

1. PRELIMINARIES AND NOTATIONS

Cochain complexes. Given an Abelian category \mathcal{A} , we denote by $\text{Ch}(\mathcal{A})$ the category of unbounded cochain complexes, and maps of complexes. In particular, given

$$X^\bullet := (\dots \longrightarrow X^{n-1} \xrightarrow{d^{n-1}} X^n \xrightarrow{d^n} X^{n+1} \longrightarrow \dots) \in \text{Ch}(\mathcal{A}),$$

we denote by $B^n(X^\bullet) := \text{Im}(d^{n-1})$, $Z^n(X^\bullet) := \text{Ker}(d^n)$, and $H^n(X^\bullet) := Z^n(X^\bullet)/B^n(X^\bullet)$, the n -th **coboundaries**, **cocycles**, and **cohomology** of X^\bullet , respectively. We let $\Sigma: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})$ be the “shift to the left” auto-equivalence of $\text{Ch}(\mathcal{A})$, that is, $(\Sigma X^\bullet)^n := X^{n+1}$.

A given morphism of complexes $\phi^\bullet: X^\bullet \rightarrow Y^\bullet$ is said to be a **quasi-isomorphism** if $H^n(\phi^\bullet)$ is an isomorphism for all $n \in \mathbb{Z}$. We also say that a complex X^\bullet is **exact** (or **acyclic**) if $H^n(X^\bullet) = 0$, for all $n \in \mathbb{Z}$ or, equivalently, $X^\bullet \rightarrow 0$ is a quasi-isomorphism.

For each $k \in \mathbb{Z}$, we denote by $\text{Ch}^{\geq k}(\mathcal{A})$ (resp., $\text{Ch}^{\leq k}(\mathcal{A})$) the full subcategory of the $X^\bullet \in \text{Ch}(\mathcal{A})$ such that $X^n = 0$, for all $n < k$ (resp., $n > k$). Moreover, $\text{Ch}^+(\mathcal{A}) := \bigcup_{n \in \mathbb{Z}} \text{Ch}^{\geq n}(\mathcal{A})$ denotes the full subcategory of bounded below complexes. Given $k \in \mathbb{Z}$, the inclusion $\text{Ch}^{\geq k}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})$ has a left adjoint, called the k -th **left truncation functor** $\tau^{\geq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}^{\geq k}(\mathcal{A})$ such that:

$$\tau^{\geq k}(X^\bullet) := (\dots \longrightarrow 0 \longrightarrow X^k/B^k(X^\bullet) \xrightarrow{\bar{d}^k} X^{k+1} \xrightarrow{d^{k+1}} \dots) \in \text{Ch}^{\geq k}(\mathcal{A}).$$

Similarly, the inclusion $\text{Ch}^{\leq k}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})$ has a right adjoint, called the k -th **right truncation functor** $\tau^{\leq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}^{\leq k}(\mathcal{A})$ such that:

$$\tau^{\leq k}(X^\bullet) := (\dots \longrightarrow X^{k-2} \xrightarrow{d^{k-2}} X^{k-1} \xrightarrow{d^{k-1}} Z^k(X^\bullet) \longrightarrow 0 \longrightarrow \dots) \in \text{Ch}^{\leq k}(\mathcal{A}).$$

Given an object $A \in \mathcal{A}$ and $k \in \mathbb{Z}$ we let $S^k(A)$ denote the following **stalk complex**:

$$S^k(A) := (\dots \longrightarrow 0 \longrightarrow A \longrightarrow 0 \longrightarrow \dots) \in \text{Ch}^{\geq k}(\mathcal{A}) \cap \text{Ch}^{\leq k}(\mathcal{A}),$$

while for $n = 0$ we sometimes denote $S^0(A)$ simply by A . Moreover, we denote by $D^k(A)$ the following **disk complex**:

$$D^k(A) := (\dots \longrightarrow 0 \longrightarrow A \xrightarrow{\text{id}_A} A \longrightarrow 0 \longrightarrow \dots) \in \text{Ch}^{\geq k}(\mathcal{A}) \cap \text{Ch}^{\leq k+1}(\mathcal{A}).$$

Observe that $D^k(A)$ is an exact complex, that is, $H^i(D^k(A)) = 0$ for all $i \in \mathbb{Z}$. Moreover, as co/products of isomorphisms are isomorphisms, any co/product of disc complexes is still exact. Finally, observe that, if Y^\bullet is such a co/product of disk complexes, and $X^\bullet \in \text{Ch}(\mathcal{A})$ is any complex, then $H^i(X^\bullet \oplus Y^\bullet) \cong H^i(X^\bullet) \oplus H^i(Y^\bullet) \cong H^i(X^\bullet)$, for all $i \in \mathbb{Z}$. This easy observation is often useful when we need to modify a given complex X^\bullet without changing its cohomologies.

Weak factorization systems. Given a category \mathcal{A} , denote by \mathcal{A}^2 its category of morphisms. Given ϕ and $\psi \in \mathcal{A}^2$, one says that ϕ is **left weakly orthogonal** to ψ (and, equivalently, that ψ is **right weakly orthogonal** to ϕ), in symbols $\phi \boxdot \psi$, if for any solid commutative square

$$\begin{array}{ccc} X_0 & \xrightarrow{a} & Y_0 \\ \phi \downarrow & \dashrightarrow^{\exists d} & \downarrow \psi \\ X_1 & \xrightarrow{b} & Y_1 \end{array}$$

there is a (not necessarily unique) diagonal $d: X_1 \rightarrow Y_0$ such that $a = d \circ \phi$ and $b = \psi \circ d$.

Given a class of objects $\mathcal{X} \subseteq \mathcal{A}^2$, we denote by

$$\boxdot \mathcal{X} := \{l \in \mathcal{A}^2 : l \boxdot x, \forall x \in \mathcal{X}\} \quad \text{and} \quad \mathcal{X} \boxdot := \{r \in \mathcal{A}^2 : x \boxdot r, \forall x \in \mathcal{X}\},$$

the classes of all morphisms in \mathcal{A} which are weakly left or right orthogonal, respectively, to \mathcal{X} . Using another common terminology, $l \in \boxdot \mathcal{X}$ (resp., $r \in \mathcal{X} \boxdot$), if it has the **left (right) lifting property** with respect to \mathcal{X} . The following general lemma is often useful in practice:

Lemma 1.1 ([Rie14, Lemma 11.1.4]). *Any class of arrows of the form $\boxdot \mathcal{X}$ is closed under coproducts, push-outs, transfinite composition, retracts, and it contains the isomorphisms. The class $\mathcal{X} \boxdot$ has the dual closure properties.*

In particular, we will need the following consequence of the above lemma: Suppose that $(A_n)_{n \in \mathbb{N}}$ is an inverse system in \mathcal{A} such that $\varprojlim_{\mathbb{N}} A_n$ exists in \mathcal{A} . If $(A_{n+1} \rightarrow A_n) \in \mathcal{X} \boxdot$ for all $n \in \mathbb{N}$, then the canonical map $\varprojlim_{\mathbb{N}} A_n \rightarrow A_0$ is also in $\mathcal{X} \boxdot$.

A pair of subclasses $(\mathcal{X}, \mathcal{Y})$ of \mathcal{A}^2 is a **weak factorization system** if the following hold:

(WFS.1) $\mathcal{X} \boxdot = \mathcal{Y}$ and $\mathcal{X} = \boxdot \mathcal{Y}$;

(WFS.2) each $\phi \in \mathcal{A}^2$ can be written as $\phi = y \circ x$, with $x \in \mathcal{X}$ and $y \in \mathcal{Y}$.

Model categories. A **model structure** on a bicomplete category \mathcal{M} is a triple $(\mathcal{W}, \mathcal{C}, \mathcal{F})$ of classes of maps, called respectively **weak equivalences**, **cofibrations** and **fibrations**, such that:

(MS.1) \mathcal{W} contains all the isomorphisms and has the 2-out-of-3 property;

(MS.2) both $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{F} \cap \mathcal{W})$ are weak factorization systems.

In this case, the quadruple $(\mathcal{M}, \mathcal{W}, \mathcal{C}, \mathcal{F})$ is called a **model category**. The morphisms in $\mathcal{C} \cap \mathcal{W}$ and $\mathcal{F} \cap \mathcal{W}$ are called **trivial cofibrations** and **trivial fibrations**, respectively. Furthermore, an object $F \in \mathcal{M}$ is said to be **fibrant** if the terminal map $F \rightarrow *$ is a fibration while, for \mathcal{M} pointed, we say that F is **trivial** if the trivial map $0 \rightarrow F$ is a weak equivalence. Finally, a **fibrant replacement** for an object $X \in \mathcal{M}$, is a weak equivalence $X \rightarrow F$, with F fibrant. Cofibrant objects and cofibrant replacements are defined dually.

Remark 1.2. *Suppose that we are in the following situation: we have a bicomplete category \mathcal{M} and two classes of maps, \mathcal{W} , whose members we call weak equivalences, and \mathcal{C} , whose members we call cofibrations, that we would like to use to construct a model structure on \mathcal{M} . In this setting, we are forced to take $\mathcal{F} := (\mathcal{C} \cap \mathcal{W}) \boxdot$ as a class of fibrations. Even if we do not know whether $(\mathcal{W}, \mathcal{C}, \mathcal{F})$ is a model structure, let us abuse terminology and call an object $X \in \mathcal{M}$ fibrant if $X \rightarrow *$ belongs to \mathcal{F} . Consider an inverse system $(F_n)_{\mathbb{N}}$ of fibrant objects and suppose that $(F_{n+1} \rightarrow F_n) \in \mathcal{F}$ for all $n \in \mathbb{N}$, then $F := \varprojlim_{\mathbb{N}} F_n$ is a fibrant object.*

Proof. Consider the inverse system $(F'_n)_{\mathbb{N}}$, where $F'_0 = *$ is the terminal object, and $F'_{n+1} := F_n$ for all $n \in \mathbb{N}$. Then, as $\mathcal{F} = (\mathcal{C} \cap \mathcal{W}) \boxdot$, the unique map $\varprojlim_{\mathbb{N}} F_n \cong \varprojlim_{\mathbb{N}} F'_n \rightarrow F'_0 = *$ does still belong to \mathcal{F} , as discussed briefly after Lemma 1.1. \square

Hovey correspondence. Let \mathcal{A} be a bicomplete Abelian category. A pair $(\mathcal{X}, \mathcal{Y})$ of classes in \mathcal{A} is said to be a **cotorsion pair** if:

- $\mathcal{X} = {}^{\perp 1}\mathcal{Y} := \{A \in \mathcal{A} : \text{Ext}_{\mathcal{A}}^1(A, Y) = 0, \text{ for all } Y \in \mathcal{Y}\}$, and
- $\mathcal{Y} = \mathcal{X}^{\perp 1} := \{A \in \mathcal{A} : \text{Ext}_{\mathcal{A}}^1(X, A) = 0, \text{ for all } X \in \mathcal{X}\}$.

A cotorsion pair $(\mathcal{X}, \mathcal{Y})$ is **left** (resp., **right**) **complete** if for all $A \in \mathcal{A}$ there is an exact sequence

$$0 \longrightarrow Y \longrightarrow X \longrightarrow A \longrightarrow 0 \quad (\text{resp.}, 0 \longrightarrow A \longrightarrow Y' \longrightarrow X' \longrightarrow 0),$$

with $X, X' \in \mathcal{X}$ and $Y, Y' \in \mathcal{Y}$, called a **left** (resp., **right**) **$(\mathcal{X}, \mathcal{Y})$ -approximation** of A . Moreover, we say that $(\mathcal{X}, \mathcal{Y})$ is **complete** if it is left and right complete, and **hereditary** if \mathcal{X} is closed under kernels of epimorphisms or, equivalently, \mathcal{Y} is closed under cokernels of monomorphisms.

Recall also that a **model structure** on \mathcal{A} is said to be **Abelian** if:

- the (trivial) cofibrations are the monomorphisms that have a (trivially) cofibrant cokernel;
- the (trivial) fibrations are the epimorphisms that have a (trivially) fibrant kernel.

By a famous result of Hovey (see [Hov02]), Abelian model structures on a bicomplete Abelian category \mathcal{A} are in bijection with the so-called **Hovey triples**, which are triples of classes of objects $(\mathcal{O}_{\mathcal{W}}, \mathcal{O}_{\mathcal{C}}, \mathcal{O}_{\mathcal{F}})$ such that $\mathcal{O}_{\mathcal{W}}$ is thick (i.e., it is closed under summands and, given any short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in \mathcal{A} , if two elements of the set $\{A, B, C\}$ belong in $\mathcal{O}_{\mathcal{W}}$, then so does the third) and both $(\mathcal{O}_{\mathcal{C}}, \mathcal{O}_{\mathcal{W}} \cap \mathcal{O}_{\mathcal{F}})$ and $(\mathcal{O}_{\mathcal{C}} \cap \mathcal{O}_{\mathcal{W}}, \mathcal{O}_{\mathcal{F}})$ are complete cotorsion pairs. In this case, the class $\mathcal{O}_{\mathcal{C}}$ is precisely the class of cofibrant objects, $\mathcal{O}_{\mathcal{F}}$ is the class of fibrant objects, and $\mathcal{O}_{\mathcal{W}}$ is the class of trivial objects in the corresponding Abelian model structure.

Lifting of cotorsion pairs. Given a cotorsion pair $(\mathcal{X}, \mathcal{Y})$ in \mathcal{A} , Gillespie [Gil06, Definition 3.3] introduced the following classes, with the goal of inducing a Hovey triple (and so also an Abelian model structure) in $\text{Ch}(\mathcal{A})$:

- $\tilde{\mathcal{X}}$ (resp., $\tilde{\mathcal{Y}}$) is the class of all acyclic $A^{\bullet} \in \text{Ch}(\mathcal{A})$ such that $Z^n(A^{\bullet}) \in \mathcal{X}$ ($Z^n(A^{\bullet}) \in \mathcal{Y}$);
- $\text{dg}\mathcal{X}$ is the class of those $A^{\bullet} \in \text{Ch}(\mathcal{A})$ such that $A^n \in \mathcal{X}$ for all $n \in \mathbb{Z}$ and for which the total hom-complex $\underline{\text{Hom}}(A^{\bullet}, Y^{\bullet})$ is exact for all $Y^{\bullet} \in \tilde{\mathcal{Y}}$;
- $\text{dg}\mathcal{Y}$ is the class of those $A^{\bullet} \in \text{Ch}(\mathcal{A})$ such that $A^n \in \mathcal{Y}$ for all $n \in \mathbb{Z}$ and for which the total hom-complex $\underline{\text{Hom}}(X^{\bullet}, A^{\bullet})$ is exact for all $X^{\bullet} \in \tilde{\mathcal{X}}$.

Proposition 1.3 ([Gil06, Lemma 3.4, Proposition 3.6, Lemma 3.14] and [Hov02, Lemma 6.2]). *Let \mathcal{A} be a bicomplete Abelian category, $(\mathcal{X}, \mathcal{Y})$ a cotorsion pair, and \mathcal{E} the class of exact complexes. Then, the following statements hold true*

- (1) *a complex $A^{\bullet} \in \text{Ch}^+(\mathcal{A})$ belongs to $\text{dg}\mathcal{Y}$ if and only if $A^n \in \mathcal{Y}$, for all $n \in \mathbb{Z}$;*
- (2) *if $(\mathcal{X}, \mathcal{Y})$ is complete, both $(\tilde{\mathcal{X}}, \text{dg}\mathcal{Y})$ and $(\text{dg}\mathcal{X}, \tilde{\mathcal{Y}})$ are cotorsion pairs in $\text{Ch}(\mathcal{A})$. Hence, if $(A_n)_{\mathbb{N}}$ is an inverse system such that $A_0 \in \text{dg}\mathcal{Y}$ and $A_{n+1} \rightarrow A_n$ is an epimorphism with $\ker(A_{n+1} \rightarrow A_n) \in \text{dg}\mathcal{Y}$ (for all $n \in \mathbb{N}$), then $\lim_{\mathbb{N}} A_n \in \text{dg}\mathcal{Y}$;*
- (3) *if $(\text{dg}\mathcal{X}, \tilde{\mathcal{Y}})$ is a complete cotorsion pair, and $\text{dg}\mathcal{X} \cap \mathcal{E} = \tilde{\mathcal{X}}$, then $\text{dg}\mathcal{Y} \cap \mathcal{E} = \tilde{\mathcal{Y}}$.*

As a consequence of the above proposition, one can deduce the following criterion for a cotorsion pair in \mathcal{A} to induce a model structure on complexes:

Corollary 1.4. *Let \mathcal{A} be a bicomplete Abelian category, let $(\mathcal{X}, \mathcal{Y})$ be a cotorsion pair in \mathcal{A} , let \mathcal{E} be the class of exact complexes and suppose that:*

- (1) *$(\mathcal{X}, \mathcal{Y})$, $(\text{dg}\mathcal{X}, \tilde{\mathcal{Y}})$ and $(\tilde{\mathcal{X}}, \text{dg}\mathcal{Y})$ are complete;*
- (2) *$\text{dg}\mathcal{X} \cap \mathcal{E} = \tilde{\mathcal{X}}$.*

Then $(\mathcal{E}, \text{dg}\mathcal{X}, \text{dg}\mathcal{Y})$ is a Hovey triple and, therefore, it induces an Abelian model structure on $\text{Ch}(\mathcal{A})$ such that the weak equivalences are the quasi-isomorphisms, the cofibrations are the monomorphisms with cokernel in $\text{dg}\mathcal{X}$, and the fibrations are the epimorphisms with kernel in $\text{dg}\mathcal{Y}$.

2. DG-INJECTIVE RESOLUTIONS OF COMPLEXES VIA TRUNCATIONS

Let \mathcal{A} be an Abelian category with enough injectives, that is, for all $A \in \mathcal{A}$ there is a monomorphism $A \rightarrow E$, with $E \in \text{Inj}(\mathcal{A})$, and consider the complete cotorsion pair $(\mathcal{A}, \text{Inj}(\mathcal{A}))$ in \mathcal{A} . Then:

- $\tilde{\mathcal{A}} = \mathcal{E}$ is the class of exact complexes;
- $\widetilde{\text{Inj}(\mathcal{A})}$ is the class of contractible complexes of injectives, that coincides with the class $\text{Inj}(\text{Ch}(\mathcal{A}))$ of injective objects in $\text{Ch}(\mathcal{A})$ (see [Wei94, Exercise 2.2.1]);
- $\text{dg}\mathcal{A} = \text{Ch}(\mathcal{A})$ and $\text{dg}(\text{Inj}(\mathcal{A}))$ is, by definition, the class of **DG-injective complexes**.

In particular, the cotorsion pairs induced in $\text{Ch}(\mathcal{A})$ are $(\text{Ch}(\mathcal{A}), \text{Inj}(\text{Ch}(\mathcal{A})))$, which is complete (see [Wei94, Exercise 2.2.2]), and $(\mathcal{E}, \text{dg}(\text{Inj}(\mathcal{A})))$. Furthermore, condition (2) in Corollary 1.4 is trivial in this case. Hence, when \mathcal{A} is bicomplete, we get a model structure in $\text{Ch}(\mathcal{A})$ provided each $X^\bullet \in \text{Ch}(\mathcal{A})$ has a **DG-injective resolution**, that is, a quasi-isomorphism $\lambda^\bullet: X^\bullet \rightarrow E^\bullet$, with $E^\bullet \in \text{dg}(\text{Inj}(\mathcal{A}))$. In fact, this is equivalent to the fact that, in the short exact sequence

$$(2.1) \quad 0 \longrightarrow \Sigma^{-1}E^\bullet \longrightarrow \Sigma^{-1}\text{cone}(\lambda^\bullet) \longrightarrow X^\bullet \longrightarrow 0,$$

$\text{cone}(\lambda^\bullet) \in \mathcal{E}$ and $E^\bullet \in \text{dg}(\text{Inj}(\mathcal{A}))$, so $(\mathcal{E}, \text{dg}(\text{Inj}(\mathcal{A})))$ is left complete. Note also that, up to adding a bunch of disc complexes to E^\bullet , we may assume that λ^\bullet is a monomorphism. Hence, $0 \rightarrow X^\bullet \rightarrow E^\bullet \rightarrow E^\bullet/X^\bullet \rightarrow 0$ is a right approximation sequence, and $(\mathcal{E}, \text{dg}(\text{Inj}(\mathcal{A})))$ is complete.

In this section we give a complete proof, based on ideas from [CNPS18], of the fact that Spaltenstein's construction of DG-injective resolutions in the category of complexes of modules (over a ring or, more generally, over a sheaf of rings) works in every complete Abelian category \mathcal{A} with enough injectives that satisfies Roos' condition (Ab.4^*) - k (for some $k \in \mathbb{N}$).

2.1. Resolution of bounded-below complexes. Consider a complex $X^\bullet \in \text{Ch}^{\geq k}(\mathcal{A})$ (for some $k \in \mathbb{Z}$), where \mathcal{A} is any Abelian category with enough injectives. In what follows we show how to construct a monomorphic quasi-isomorphism $\lambda^\bullet: X^\bullet \rightarrow E^\bullet$ with $E^\bullet \in \text{Ch}^{\geq k}(\mathcal{A})$ and E^i injective for all $i \in \mathbb{Z}$; then this is a DG-injective resolution, by Proposition 1.3(1). Suppose for simplicity that $k = 0$, that is, we start with $X^\bullet \in \text{Ch}^{\geq 0}(\mathcal{A})$, and we proceed inductively to construct

$$\lambda^\bullet: X^\bullet \longrightarrow E^\bullet := (\cdots \longrightarrow 0 \longrightarrow E^0 \xrightarrow{e^0} E^1 \xrightarrow{e^1} \cdots).$$

Indeed, let $E^n = 0$, $e^n = 0$ and $\lambda^n = 0$, for all $n < 0$ and, for $n = 0$, define $\lambda^0: X^0 \rightarrow E^0$ to be a chosen monomorphism with $E^0 \in \text{Inj}(\mathcal{A})$. Now, given $n \geq 0$, if we have already built $e^{i-1}: E^{i-1} \rightarrow E^i$ and $\lambda^i: X^i \rightarrow E^i$, for all $i \leq n$, such that:

- (1_n) $\lambda^i \circ d^{i-1} = e^{i-1} \circ \lambda^{i-1}$ for all $i \leq n$;
- (2_n) $e^{i-1} \circ e^{i-2} = 0$ for all $i \leq n$;
- (3_n) the map $H^i(X^\bullet) \rightarrow H^i(E^\bullet)$ induced by λ^i is an isomorphism for all $i \leq n-1$;
- (4_n) the map $\bar{\lambda}^n: X^n/B^n(X^\bullet) \rightarrow E^n/B^n(E^\bullet)$ induced by λ^n is a monomorphism;

we consider the following push-out diagram:

$$\begin{array}{ccc} X^n/B^n(X^\bullet) & \xrightarrow{\bar{d}^n} & X^{n+1} \\ \bar{\lambda}^n \downarrow & \text{P.O.} & \downarrow \mu^{n+1} \\ E^n/B^n(E^\bullet) & \xrightarrow{y^n} & P^{n+1}. \end{array}$$

In particular, the map $\bar{\mu}^{n+1}: (X^{n+1}/B^{n+1}(X^\bullet) \cong) \text{Coker}(\bar{d}^n) \rightarrow \text{Coker}(y^n)$ is an isomorphism. Moreover, as $\bar{\lambda}^n$ is a monomorphism by hypothesis, the above square is also a pullback, so that $\bar{\lambda}^n$ induces an isomorphism between $\text{Ker}(\bar{d}^n) = H^n(X^\bullet)$ and $\text{Ker}(y^n)$. Choose now a monomorphism $\iota^{n+1}: P^{n+1} \rightarrow E^{n+1}$, with $E^{n+1} \in \text{Inj}(\mathcal{A})$. Then, the following square remains a pullback:

$$\begin{array}{ccc} X^n/B^n(X^\bullet) & \xrightarrow{\bar{d}^n} & X^{n+1} \\ \bar{\lambda}^n \downarrow & \text{P.B.} & \downarrow \lambda^{n+1} := \iota^{n+1} \circ \mu^{n+1} \\ E^n/B^n(E^\bullet) & \xrightarrow{\iota^{n+1} \circ y^n} & E^{n+1}. \end{array}$$

Finally define $e^n := \iota^{n+1} \circ y^n \circ \pi^n: E^n \rightarrow E^{n+1}$, where $\pi^n: E^n \rightarrow E^n/B^n(E^\bullet)$ is the canonical projection. Conditions (1_{n+1})–(4_{n+1}) hold by construction, so we can continue with the induction.

2.2. Spaltenstein towers of partial resolutions. Let \mathcal{A} be an Abelian category with enough injectives and fix a complex $X^\bullet \in \text{Ch}(\mathcal{A})$. We start considering the following inverse system of successive truncations:

$$\cdots \xrightarrow{\rho_2^\bullet} \tau^{\geq -2}(X^\bullet) \xrightarrow{\rho_1^\bullet} \tau^{\geq -1}(X^\bullet) \xrightarrow{\rho_0^\bullet} \tau^{\geq 0}(X^\bullet),$$

where ρ_n^\bullet is the canonical projection. As $\tau^{\geq -n}(X^\bullet) \in \text{Ch}^{\geq -n}(\mathcal{A})$, the argument of Subsection 2.1 gives a monomorphic DG-injective resolution $\lambda_n^\bullet: \tau^{\geq -n}(X^\bullet) \rightarrow E_n^\bullet$, with $E_n^\bullet \in \text{Ch}^{\geq -n}(\text{Inj}(\mathcal{A}))$, for all $n \in \mathbb{N}$. We get the following solid diagram:

$$(2.2) \quad \begin{array}{ccccccc} \cdots & \xrightarrow{\rho_2^\bullet} & \tau^{\geq -2}(X^\bullet) & \xrightarrow{\rho_1^\bullet} & \tau^{\geq -1}(X^\bullet) & \xrightarrow{\rho_0^\bullet} & \tau^{\geq 0}(X^\bullet) \\ & & \downarrow \lambda_2^\bullet & & \downarrow \lambda_1^\bullet & & \downarrow \lambda_0^\bullet \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ & \cdots & \xrightarrow{t_2^\bullet} & E_2^\bullet & \xrightarrow{t_1^\bullet} & E_1^\bullet & \xrightarrow{t_0^\bullet} & E_0^\bullet \end{array};$$

and we claim that it can be completed in a commutative way by the dotted arrows. Indeed, consider the following short exact sequence for each $n > 0$

$$0 \rightarrow \tau^{\geq -n}(X^\bullet) \xrightarrow{\lambda_n^\bullet} E_n^\bullet \rightarrow \text{Coker}(\lambda_n^\bullet) \rightarrow 0,$$

and apply the functor $\text{Hom}_{\text{Ch}(\mathcal{A})}(-, E_{n-1}^\bullet)$ to get the following exact sequence in Ab:

$$\text{Hom}_{\text{Ch}(\mathcal{A})}(E_n^\bullet, E_{n-1}^\bullet) \xrightarrow{(\lambda_n^\bullet)^*} \text{Hom}_{\text{Ch}(\mathcal{A})}(\tau^{\geq -n}(X^\bullet), E_{n-1}^\bullet) \rightarrow \text{Ext}_{\text{Ch}(\mathcal{A})}^1(\text{Coker}(\lambda_n^\bullet), E_{n-1}^\bullet) = 0.$$

The last equality to zero is due to the fact $\text{Coker}(\lambda_n^\bullet) \in \mathcal{E}$, $E_{n-1}^\bullet \in \text{dg}(\text{Inj}(\mathcal{A}))$ and $(\mathcal{E}, \text{dg}(\text{Inj}(\mathcal{A})))$ is a cotorsion pair in $\text{Ch}(\mathcal{A})$. It then follows that $\lambda_{n-1}^\bullet \circ \rho_{n-1}^\bullet \in \text{Im}((\lambda_n^\bullet)^*)$, so we have a map $t_{n-1}^\bullet: E_n^\bullet \rightarrow E_{n-1}^\bullet$, such that $t_{n-1}^\bullet \circ \lambda_n^\bullet = \lambda_{n-1}^\bullet \circ \rho_{n-1}^\bullet$, as desired.

Observe also that, possibly adding a bunch of disk complexes to our partial resolutions if needed, one can always suppose that t_n^\bullet is a (degreewise split) epimorphism with $\ker(t_n^\bullet) \in \text{Ch}^+(\text{Inj}(\mathcal{A}))$, whence with DG-injective kernel, for all $n \in \mathbb{N}$.

Definition 2.1. A **Spaltenstein tower** of partial resolutions for $X^\bullet \in \text{Ch}(\mathcal{A})$ is a commutative diagram like the one in (2.2) where λ_n^\bullet is a DG-injective resolution for all $n \in \mathbb{N}$, and t_n^\bullet is a (degreewise split) epimorphism with DG-injective kernel, for each $n \in \mathbb{N}$.

In particular, in the above discussion we have just shown that:

Proposition 2.2. Let \mathcal{A} be an Abelian category with enough injectives. Then any complex in $\text{Ch}(\mathcal{A})$ has a Spaltenstein tower of partial resolutions.

2.3. Injective resolutions and the (Ab.4*)-k condition. The goal of this subsection is to prove that, under suitable conditions, the inverse limit of a Spaltenstein tower of partial resolutions actually produces a DG-injective resolution. We start with two easy observations:

Lemma 2.3. Let \mathcal{A} be a complete Abelian category and consider a sequence of complexes

$$\cdots \xrightarrow{t_2^\bullet} X_2^\bullet \xrightarrow{t_1^\bullet} X_1^\bullet \xrightarrow{t_0^\bullet} X_0^\bullet$$

such that, for each $n \in \mathbb{N}$, the chain map t_n^\bullet is degreewise a split-epimorphism. Then, there is the following degreewise split-exact sequence in $\text{Ch}(\mathcal{A})$:

$$0 \longrightarrow \varprojlim_{\mathbb{N}} X_n^\bullet \longrightarrow \prod_{\mathbb{N}} X_n^\bullet \xrightarrow{1-t} \prod_{\mathbb{N}} X_n^\bullet \longrightarrow 0,$$

where the map $1-t$ is described by a matrix with identities on the main diagonal, $-t_n^\bullet$ as the n -th entry of the first superdiagonal, and 0's everywhere else.

Proof. The universal property defining $\varprojlim_{\mathbb{N}} X_n^\bullet$ is clearly equivalent to the universal property defining the kernel of $1-t$. Moreover, to prove that $1-t$ is a degreewise split-epimorphism, it is enough to find, for each $k \in \mathbb{Z}$, a right-inverse for the following matrix:

$$\begin{pmatrix} \text{id}_{X_0^k} & -t_0^k & 0 & 0 & 0 & \cdots \\ 0 & \text{id}_{X_1^k} & -t_1^k & 0 & 0 & \cdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

Fix, for each $n \in \mathbb{N}$ and $k \in \mathbb{Z}$, a split-monomorphism $s_n^k: X_n^k \rightarrow X_{n+1}^k$ such that $t_n^k \circ s_n^k = \text{id}_{X_n^k}$. Then, the right-inverse we are looking for is the following row-finite matrix:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \dots \\ -s_0^k & 0 & 0 & 0 & 0 & \dots \\ -s_1^k \circ s_0^k & -s_1^k & 0 & 0 & 0 & \dots \\ -s_2^k \circ s_1^k \circ s_0^k & -s_2^k \circ s_1^k & -s_2^k & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots \end{pmatrix} \quad \square$$

Lemma 2.4. *Let \mathcal{A} be a complete Abelian category and fix some integer $k \in \mathbb{Z}$. Consider also two families of complexes $(X_n^\bullet)_{n \in \mathbb{N}}, (Y_n^\bullet)_{n \in \mathbb{N}} \subseteq \text{Ch}(\mathcal{A})$, and maps of complexes $\phi_n^\bullet: X_n^\bullet \rightarrow Y_n^\bullet$ (for each $n \in \mathbb{N}$), such that ϕ_n^i is an isomorphism for all $i \geq k$. Letting*

$$\phi^\bullet := \prod_{\mathbb{N}} \phi_n^\bullet: \prod_{\mathbb{N}} X_n^\bullet \longrightarrow \prod_{\mathbb{N}} Y_n^\bullet,$$

ϕ^i is an isomorphism for all $i \geq k$. In particular, $H^i(\prod_{\mathbb{N}} X_n^\bullet) \cong H^i(\prod_{\mathbb{N}} Y_n^\bullet)$, for all $i > k$.

Let us point out that the isomorphism for the i -th cohomologies of the products holds for all $i > k$, but it may actually fail for $i = k$, even if $H^k(X_n^\bullet) \cong H^k(Y_n^\bullet)$, for all $n \in \mathbb{N}$.

Proof. As products are built componentwise, the result follows by the observation that the class of isomorphisms in \mathcal{A} is closed under products. \square

The following definition was introduced by Roos [Roo06]:

Definition 2.5. *A complete Abelian category \mathcal{A} with enough injectives is (Ab.4^*) - k , for some $k \in \mathbb{N}$ if, for any family $(A_\lambda)_\Lambda \subseteq \mathcal{A}$, the n -th derived functor $\prod_{\Lambda}^{(n)} A_\lambda$ vanishes for all $n > k$.*

In particular, \mathcal{A} is (Ab.4^*) -0 if and only if it satisfies Grothendieck's axiom (Ab.4^*) , that is, products are exact in \mathcal{A} . In the following lemma we show a direct consequence of the (Ab.4^*) - k condition, which is precisely what is needed for the construction of DG-injective resolutions:

Lemma 2.6. *Fix $k \in \mathbb{N}$, $h \in \mathbb{Z}$, and let \mathcal{A} be a complete (Ab.4^*) - k Abelian category. Given a family $(X_n^\bullet)_{n \in \mathbb{N}} \subseteq \text{Ch}(\mathcal{A})$ such that $H^i(X_n^\bullet) = 0$ and with X_n^i an injective object, for all $i > h$, we have that $H^i(\prod_{\mathbb{N}} X_n^\bullet) = 0$, for all $i > h + k + 1$.*

Proof. Observe that $(\tau^{\geq h} X_n^\bullet)_{n \in \mathbb{N}}$ is a family of injective resolutions for $(S^h(X_n^h/B^h(X_n^\bullet)))_{n \in \mathbb{N}}$. Hence, the $h + i$ -th cohomology of $\prod_{\mathbb{N}} \tau^{\geq h}(X_n^\bullet)$ is precisely the i -th derived functor $\prod_{n \in \mathbb{N}}^{(i)} (X_n^h/B^h(X_n^\bullet))$. In particular, by Definition 2.5, we deduce that $H^i(\prod_{\mathbb{N}} \tau^{\geq h} X_n^\bullet) = 0$, for all $i > h + k$. Moreover, the canonical map $X_n^\bullet \rightarrow \tau^{\geq h} X_n^\bullet$ (for each $n \in \mathbb{N}$) is an isomorphism in all degrees $> h$. Conclude by the first part and Lemma 2.4. \square

Combining the above lemma with Lemma 2.3, we deduce the following:

Proposition 2.7. *Let \mathcal{A} be a complete (Ab.4^*) - k Abelian category (for some $k \in \mathbb{N}$) with enough injectives, and consider the following sequence in $\text{Ch}^+(\mathcal{A})$:*

$$\cdots \longrightarrow K_2^\bullet \xrightarrow{t_1^\bullet} K_1^\bullet \xrightarrow{t_0^\bullet} K_0^\bullet.$$

We suppose that this sequence satisfies the following conditions:

- (1) K_n^i is injective for all $i \in \mathbb{Z}$ and $n \in \mathbb{N}$ (so each K_n^\bullet is DG-injective);
- (2) t_n^\bullet is a degreewise split-epimorphism, for all $n \in \mathbb{N}$;
- (3) there is an integer $h \in \mathbb{Z}$ such that $H^i(K_n^\bullet) = 0$ for all $n \in \mathbb{N}$ and all $i \geq h$.

Then, $H^i(\varprojlim_{\mathbb{N}} K_n^\bullet) = 0$, for all $i > h + k + 2$.

Proof. By Lemma 2.3 and condition (2), there is a short exact sequence

$$(2.3) \quad 0 \longrightarrow \varprojlim_{\mathbb{N}} K_n^\bullet \longrightarrow \prod_{\mathbb{N}} K_n^\bullet \longrightarrow \prod_{\mathbb{N}} K_n^\bullet \longrightarrow 0.$$

Furthermore, by Lemma 2.6 and conditions (1) and (3), $H^i(\prod_{\mathbb{N}} K_n^\bullet) = 0$ for all $i \geq h + k + 1$. To conclude, just look at the long exact sequence in cohomologies induced by (2.3). \square

Let us also register the following useful consequence:

Corollary 2.8. *Let \mathcal{A} be a complete (Ab.4^*) - k Abelian category (for some $k \in \mathbb{N}$) with enough injectives, and let $E^\bullet := \varprojlim_{\mathbb{N}} E_n^\bullet$ by the limit of the following sequence of bounded below complexes:*

$$\cdots \longrightarrow E_2^\bullet \xrightarrow{t_1^\bullet} E_1^\bullet \xrightarrow{t_0^\bullet} E_0^\bullet.$$

We suppose that this sequence satisfies the following conditions:

- (1) E_n^\bullet is injective for all $i \in \mathbb{Z}$ and $n \in \mathbb{N}$ (so each E_n^\bullet is DG-injective);
- (2) t_n^\bullet is a degreewise split-epimorphism, for all $n \in \mathbb{N}$ (i.e., each t_n^\bullet is an $\text{Inj}(\mathcal{A})$ -fibration);
- (3) $H^{-n-1}(E_n^\bullet) = 0$, for all $n \in \mathbb{N}$;
- (4) $H^i(t_n^\bullet): H^i(E_{n+1}^\bullet) \rightarrow H^i(E_n^\bullet)$ is an isomorphism, for all $i \geq -n$ (with $i \in \mathbb{Z}$, $n \in \mathbb{N}$).

Then, for each $h \in \mathbb{N}$, $H^i(E^\bullet) \cong H^i(E_h^\bullet)$, for all $i \geq -h + k + 3$.

Proof. Fix an $h \in \mathbb{N}$ and, for each $n > h$, consider the composition $c_n^\bullet := t_h^\bullet \circ \cdots \circ t_{n-1}^\bullet: E_n^\bullet \rightarrow E_h^\bullet$. By (2), each c_n^\bullet is a degreewise split-epimorphism, so $K_n^\bullet := \text{Ker}(c_n^\bullet)$ is DG-injective and it fits into a degreewise split-exact sequence: $0 \rightarrow K_n^\bullet \rightarrow E_n^\bullet \rightarrow E_h^\bullet \rightarrow 0$. Consider the associated long exact sequence of cohomologies and deduce by (3) and (4) that $H^i(K_n^\bullet) = 0$, for all $i \geq -h$. Taking the limit for $n > h$, we get the following short exact sequence:

$$(2.4) \quad 0 \longrightarrow \varprojlim_{n>h} K_n^\bullet \longrightarrow E^\bullet \longrightarrow E_h^\bullet \longrightarrow 0,$$

where the central term is E^\bullet since $\mathbb{N}_{>h}$ is cofinal in \mathbb{N} . By Proposition 2.7, $H^i(\varprojlim_{\mathbb{N}} K_n^\bullet) = 0$, for all $i > -h + k + 2$. Conclude by looking at the long cohomology sequence associated to (2.4). \square

Finally, we are able to prove the main result of this section:

Theorem 2.9. *Let \mathcal{A} be a complete (Ab.4^*) - k (for some $k \in \mathbb{N}$) Abelian category with enough injectives and let $X^\bullet \in \text{Ch}(\mathcal{A})$. Then, there is a Spaltenstein tower of partial resolutions of X^\bullet :*

$$\begin{array}{ccccc} \cdots & \longrightarrow & \tau^{\geq -2}(X^\bullet) & \longrightarrow & \tau^{\geq -1}(X^\bullet) & \longrightarrow & \tau^{\geq 0}(X^\bullet) \\ & & \downarrow \lambda_2^\bullet & & \downarrow \lambda_1^\bullet & & \downarrow \lambda_0^\bullet \\ \cdots & \xrightarrow{t_2^\bullet} & E_2^\bullet & \xrightarrow{t_1^\bullet} & E_1^\bullet & \xrightarrow{t_0^\bullet} & E_0^\bullet. \end{array}$$

Define $E^\bullet := \varprojlim_{\mathbb{N}} E_n^\bullet$, and let $\lambda^\bullet := \varprojlim_{\mathbb{N}} \lambda_n^\bullet: X^\bullet \cong \varprojlim_{\mathbb{N}} \tau^{\geq -n}(X^\bullet) \rightarrow E^\bullet$ be the induced map. Then, $\lambda^\bullet: X^\bullet \rightarrow E^\bullet$ is a DG-injective resolution.

Proof. The existence of Spaltenstein towers follows by Proposition 2.2. Furthermore, by Proposition 1.3(2), E^\bullet is DG-injective, so it is enough to prove that λ^\bullet is a quasi-isomorphism. Observe that, for each $h \in \mathbb{N}$, Corollary 2.8 implies that $H^i(E^\bullet) \cong H^i(E_h^\bullet) \cong H^i(X^\bullet)$, for all $i \geq -h + k + 3$. As this holds for all $h \in \mathbb{N}$, we conclude that $H^i(E^\bullet) \cong H^i(X^\bullet)$ for all $i \in \mathbb{Z}$. \square

3. THE “GO-TO” COUNTER-EXAMPLE FOR UNBOUNDED RESOLUTIONS

The goal of this section is to construct a suitable bicomplete Abelian category with enough injectives \mathcal{G} (that, in fact, will be a Grothendieck category) which is not (Ab.4^*) - k for any $k \in \mathbb{N}$, and a particular unbounded complex $X^\bullet \in \text{Ch}(\mathcal{G})$ such that the construction discussed in Section 2 fails to produce a DG-injective resolution for X^\bullet , showing that the (Ab.4^*) - k condition is needed for that construction. As we will see in Sections 4, 5 and 6, the complex $X^\bullet \in \text{Ch}(\mathcal{G})$ can be used to show that several other constructions of DG-injective resolutions fail in general.

3.1. Construction of the category \mathcal{G} . We start recalling Nagata’s construction [Nag62, Example 1, Appendix A.1] of a commutative Noetherian ring of infinite Krull dimension. Let \mathbf{k} be a field, consider the polynomial ring on countably many variables

$$\mathbf{k}[\underline{x}] := \mathbf{k}[x_0, x_1, x_2, \dots],$$

and the following sequence of prime ideals in $\mathbf{k}[\underline{x}]$:

$$\mathfrak{p}_1 := (x_0, x_1), \quad \mathfrak{p}_2 := (x_2, x_3, x_4), \quad \mathfrak{p}_3 := (x_5, x_6, x_7, x_8), \quad \dots$$

where the depth of \mathfrak{p}_i is $i + 1$, for all $i \geq 1$. Let S be the multiplicative set of those elements of $\mathbf{k}[\underline{x}]$ which are not in any of the \mathfrak{p}_i 's, and define R as the ring of S -fractions of $\mathbf{k}[\underline{x}]$, that is,

$$S := \mathbf{k}[\underline{x}] \setminus \bigcup_{i=1}^{\infty} \mathfrak{p}_i \quad \text{and} \quad R := S^{-1}(\mathbf{k}[\underline{x}]).$$

Lemma 3.1 ([Nag62]). *In the above notation, R is a commutative Noetherian ring of infinite Krull dimension. In fact, the maximal ideals of R are of the form $\mathfrak{m}_i := S^{-1}\mathfrak{p}_i$, with $i \geq 1$, and this is a sequence of ideals of strictly increasing height.*

Consider the hereditary torsion class $\mathcal{T} := \text{Loc}(R/\mathfrak{m}_i : i \geq 1)$ generated by the simple R -modules (that is, the first layer of the Gabriel filtration of $\text{Mod}(R)$) and define

$$\mathcal{G} := \text{Mod}(R)/\mathcal{T}$$

as the Gabriel quotient of $\text{Mod}(R)$ over \mathcal{T} , identified with the Giraud subcategory of $\text{Mod}(R)$ of those R -modules M such that both M and $E(M)/M$ belong to

$$\mathcal{F} := \{R/\mathfrak{m}_i : i \geq 1\}^\perp = \{M \in \text{Mod}(R) : \text{Hom}_R(R/\mathfrak{m}_i, M) = 0, \text{ for all } i \geq 1\} \subseteq \text{Mod}(R).$$

It is well-known that \mathcal{G} is a Grothendieck category. Moreover, the inclusion $\iota: \mathcal{G} \rightarrow \text{Mod}(R)$ has an exact left adjoint $\mathbf{Q}: \text{Mod}(R) \rightarrow \mathcal{G}$ such that $\text{Ker}(\mathbf{Q}) := \{M \in \text{Mod}(R) : \mathbf{Q}(M) = 0\} = \mathcal{T}$. The product $\prod_I X_i$ in \mathcal{G} of a family $(X_i)_I \subseteq \mathcal{G}$ can be computed as $\mathbf{Q}(\prod_I X_i)$, where \prod_I denotes the product in the category $\text{Mod}(R)$.

The injective objects in \mathcal{G} are exactly the injective R -modules in \mathcal{F} . Furthermore, as R is commutative Noetherian, it is well-known (see [Ste75, Chap. VII, Proposition 4.5]) that \mathcal{T} is stable (i.e., closed under taking injective envelopes in $\text{Mod}(R)$). As a consequence, any injective module $E \in \text{Mod}(R)$ decomposes as

$$E \cong t(E) \oplus \mathbf{Q}(E),$$

where $t(E)$ is an injective module in \mathcal{T} and $\mathbf{Q}(E)$ is an injective object in \mathcal{G} . Note also that, if we denote by $\text{Sp}(R)$ the prime ideal spectrum of R , and by $\text{MSp}(R) = \{\mathfrak{m}_i : i \in \mathbb{N}\} \subseteq \text{Sp}(R)$ the maximal ideal spectrum, by the classification of injectives over commutative Noetherian rings given by Matlis [Mat58], we have that $E \cong \bigoplus_{\mathfrak{p} \in \text{Sp}(R)} E(R/\mathfrak{p})^{(I_{\mathfrak{p}})}$ (for suitable sets $I_{\mathfrak{p}}$), so that

$$t(E) \cong \bigoplus_{i=1}^{\infty} E(R/\mathfrak{m}_i)^{(I_{\mathfrak{m}_i})} \quad \text{and} \quad \mathbf{Q}(E) \cong \bigoplus_{\mathfrak{p} \in \text{Sp}(R) \setminus \text{MSp}(R)} E(R/\mathfrak{p})^{(I_{\mathfrak{p}})}.$$

3.2. The complex $X^\bullet \in \text{Ch}(\mathcal{G})$. With the usual abuse of notation, let $\mathbf{Q}: \text{Ch}(\text{Mod}(R)) \rightarrow \text{Ch}(\mathcal{G})$ be the functor on cochain complexes obtained by applying \mathbf{Q} componentwise. Observe that, as \mathbf{Q} is exact, it commutes with cohomologies, that is, given $X^\bullet \in \text{Ch}(\text{Mod}(R))$, we have:

$$H_{\mathcal{G}}^i(\mathbf{Q}(X^\bullet)) \cong \mathbf{Q}(H_R^i(X^\bullet)), \quad \text{for all } i \in \mathbb{Z},$$

where $H_{\mathcal{G}}^i(-)$ and $H_R^i(-)$ denote the i -th cohomology in $\text{Ch}(\mathcal{G})$ and in $\text{Ch}(\text{Mod}(R))$, respectively.

Let $\mu: R \rightarrow (\iota \circ \mathbf{Q})(R)$ be the component at R of the unit of the adjunction $\mathbf{Q} \dashv \iota: \text{Mod}(R) \rightleftarrows \mathcal{G}$ and fix an injective resolution $\lambda: R \rightarrow E^\bullet$ of R in $\text{Mod}(R)$. In the language of [CNPS18], the composition $\mathbf{Q}(\lambda) \circ \mu: R \rightarrow \mathbf{Q}(R) \rightarrow \mathbf{Q}(E^\bullet)$ is a relative $(\mathcal{F} \cap \text{Inj}(\text{Mod}(R))$ -injective resolution. By [CNPS18, Lemma 8.2], $H^0(\mathbf{Q}(E^\bullet)) \cong R$, that is, μ is an isomorphism. As a consequence:

Lemma 3.2 ([CNPS18, Lemma 8.2]). *Let $\lambda: R \rightarrow E^\bullet := (E^0 \xrightarrow{e^0} E^1 \xrightarrow{e^1} \dots \rightarrow E^n \xrightarrow{e^n} \dots)$ be an injective resolution of the right R -module R_R and consider $\mathbf{Q}(E^\bullet) \in \text{Ch}^{\geq 0}(\text{Mod}(R))$. Then,*

$$H_R^i(\mathbf{Q}(E^\bullet)) \cong \begin{cases} R & \text{if } i = 0; \\ E(R/\mathfrak{m}_i) & \text{otherwise;} \end{cases} \quad \text{and} \quad H_{\mathcal{G}}^i(\mathbf{Q}(E^\bullet)) \cong \begin{cases} \mathbf{Q}(R) \cong R & \text{if } i = 0; \\ \mathbf{Q}(E(R/\mathfrak{m}_i)) = 0 & \text{otherwise.} \end{cases}$$

In particular, $\mathbf{Q}(\lambda): R \rightarrow \mathbf{Q}(E^\bullet)$ is an injective resolution in \mathcal{G} .

We are now ready to introduce the central example for our paper. Indeed, consider:

$$\prod_{i \in \mathbb{Z}} S^i(R) := (\dots \xrightarrow{0} R \xrightarrow{0} R \xrightarrow{0} \dots \xrightarrow{0} R \xrightarrow{0} \dots) \in \text{Ch}(\text{Mod}(R)).$$

As in Lemma 3.2, let $E^\bullet \in \text{Ch}(\text{Mod}(R))$ be an injective resolution of R , viewed as a complex concentrated in non-negative degrees. Then, $\prod_{i \in \mathbb{Z}} \Sigma^i E^\bullet$ is a DG-injective resolution of $\prod_{i \in \mathbb{Z}} S^i(R)$. Our example is the complex

$$X^\bullet := \mathbf{Q}(\prod_{i \in \mathbb{Z}} S^i(R)) \cong \prod_{i \in \mathbb{Z}} S^i(R) \in \text{Ch}(\mathcal{G}).$$

Proposition 3.3. *In the above setting, let $\lambda: R \rightarrow E^\bullet$ be an injective resolution in $\text{Mod}(R)$. Then,*

- (1) $\prod_{i \in \mathbb{Z}} \Sigma^i \lambda: \prod_{i \in \mathbb{Z}} S^i(R) \rightarrow \prod_{i \in \mathbb{Z}} \Sigma^i E^\bullet$ is a quasi-isomorphism in $\text{Ch}(\text{Mod}(R))$;
- (2) $\prod_{i \in \mathbb{Z}} \Sigma^i \mathbf{Q}(\lambda): X^\bullet \rightarrow \prod_{i \in \mathbb{Z}} \Sigma^i \mathbf{Q}(E^\bullet)$ is not a quasi-isomorphism in $\text{Ch}(\mathcal{G})$.

Proof. (1) Since $\text{Mod}(R)$ is (Ab.4^*) , the class of quasi-isomorphisms is closed under products.

(2) Take the mapping cone $Z^\bullet := \text{cone}(\prod_{i \in \mathbb{Z}} \Sigma^i \mathbf{Q}(\lambda))$. By the proof of [CNPS18, Theorem 8.4], viewing Z^\bullet as a complex in $\text{Ch}(\text{Mod}(R))$, one has $H_R^n(Z^\bullet) \cong \prod_{j=1}^{\infty} E(R/\mathfrak{m}_j) \notin \mathcal{T}$, for all $n \in \mathbb{Z}$. Hence, $H_{\mathcal{G}}^n(Z^\bullet) \cong \mathbf{Q}(H_R^n(Z^\bullet)) \neq 0$, showing that Z^\bullet is not exact as a complex in $\text{Ch}(\mathcal{G})$. \square

Observe that, as \mathcal{G} is a Grothendieck category, our complex $X^\bullet \in \text{Ch}(\mathcal{G})$ is known to have a DG-injective resolution (see, e.g., [Hov01]). On the other hand, it is a consequence of the following corollary that the construction of Section 2 fails to produce a DG-injective resolution for X^\bullet :

Corollary 3.4. *In the above setting, the following statements hold for all $n \in \mathbb{N}$:*

- (1) $\tau^{\geq -n}(X^\bullet) = \prod_{i \geq -n} S^i(R)$;
- (2) the diagonal $\tau^{\geq -n}(X^\bullet) \rightarrow \prod_{i \leq n} \Sigma^i(\mathbf{Q}(E^\bullet))$ is a DG-injective resolution;
- (3) the kernel of the projection $\prod_{i \leq n+1} \Sigma^i(\mathbf{Q}(E^\bullet)) \rightarrow \prod_{i \leq n} \Sigma^i(\mathbf{Q}(E^\bullet))$ is DG-injective;
- (4) the diagonal map $X^\bullet \rightarrow \prod_{i \in \mathbb{Z}} \Sigma^i(\mathbf{Q}(E^\bullet))$ is the limit of a Spaltenstein tower of partial resolutions of X^\bullet , but it is not a DG-injective resolution of X^\bullet .

Proof. (1) is trivial while (2) follows since, even if we are considering infinite products, they are degreewise finite products, so they can be seen also as coproducts (which are exact in \mathcal{G}), showing that the map in the statement is a quasi-isomorphism. Moreover, $\prod_{i \leq n} \Sigma^i(\mathbf{Q}(E^\bullet))$ is DG-injective by Proposition 1.3(1). Part (3) is also trivial since the kernel of the projection in the statement is $\Sigma^{n+1}(\mathbf{Q}(E^\bullet))$. Finally, part (4) follows by (1–3) and Proposition 3.3(2). \square

4. CARTAN-EILENBERG RESOLUTIONS AND THEIR TOTALIZATION

In this section we recall the construction of the Cartan-Eilenberg resolutions of a complex and we show that its totalization, when applied to the example of Section 3 fails to produce a DG-injective resolution. On the other hand, as for Spaltenstein's approach, Cartan-Eilenberg resolutions produce DG-injective resolutions, provided the base category is $(\text{Ab.4}^*)\text{-}k$, for some $k \in \mathbb{N}$.

4.1. Bicomplexes and their totalizations. As a Cartan-Eilenberg resolution is, actually, a bicomplex, we start recalling that a **bicomplex** $C^{\bullet, \bullet}$ over an Abelian category \mathcal{A} is given by

- a family of objects $C^{\bullet, \bullet} := (C^{n, m})_{(n, m) \in \mathbb{Z} \times \mathbb{Z}}$;
- **differentials** $d_0^{\bullet, \bullet}: C^{\bullet, \bullet} \rightarrow C^{\bullet, \bullet+1}$ and $d_1^{\bullet, \bullet}: C^{\bullet, \bullet} \rightarrow C^{\bullet+1, \bullet}$, called **horizontal** and **vertical**, respectively, that satisfy the following formulas for all $(n, m) \in \mathbb{Z} \times \mathbb{Z}$:

$$d_0^{n, m+1} \circ d_0^{n, m} = 0, \quad d_1^{n+1, m} \circ d_1^{n, m} = 0, \quad \text{and} \quad d_1^{n, m+1} \circ d_0^{n, m} + d_0^{n+1, m} \circ d_1^{n, m} = 0.$$

In other words, a bicomplex $C^{\bullet, \bullet}$ is represented by the following diagram

$$(4.1) \quad \begin{array}{ccccccc} & & \vdots & & \vdots & & \vdots \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & C^{n, m-1} & \xrightarrow{d_0^{n, m-1}} & C^{n, m} & \xrightarrow{d_0^{n, m}} & C^{n, m+1} \longrightarrow \dots \\ & & \downarrow d_1^{n, m-1} & & \downarrow d_1^{n, m} & & \downarrow d_1^{n, m+1} \\ \dots & \longrightarrow & C^{n+1, m-1} & \xrightarrow{d_0^{n+1, m-1}} & C^{n+1, m} & \xrightarrow{d_0^{n+1, m}} & C^{n+1, m+1} \longrightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \vdots & & \vdots & & \vdots \end{array}$$

where all rows and columns are cochain complexes, and all the little squares anticommute. With reference to the representation (4.1), we say that $C^{\bullet, \bullet}$ is a **lower half-plane** bicomplex provided

$C^{n,m} = 0$ for all $n < 0$, $m \in \mathbb{Z}$. A **morphism of bicomplexes** is just a family of morphisms $\phi^{\bullet,\bullet}: X^{\bullet,\bullet} \rightarrow Y^{\bullet,\bullet}$ that commutes both with horizontal and vertical differentials. We denote by $\text{bCh}(\mathcal{A})$ the category of bicomplexes over \mathcal{A} .

In fact, $\text{bCh}(\mathcal{A})$ is equivalent to the category $\text{Ch}(\text{Ch}(\mathcal{A}))$ obtained as follows: start with \mathcal{A} and form the category of cochain complexes $\text{Ch}(\mathcal{A})$, which is itself an Abelian category; iterating the construction once more, one gets the category $\text{Ch}(\text{Ch}(\mathcal{A}))$ of **double complexes**, where a double complex can be represented by a diagram like (4.1), where all rows and columns are cochain complexes but the small squares actually commute. The canonical functor $\text{bCh}(\mathcal{A}) \rightarrow \text{Ch}(\text{Ch}(\mathcal{A}))$ is constructed via the following **sign trick**: given $C^{\bullet,\bullet} \in \text{bCh}(\mathcal{A})$ take, for each $m \in \mathbb{N}$, the complex

$$C_0^{\bullet,m} = (\dots \longrightarrow C_0^{n-1,m} \xrightarrow{(-1)^m d_1^{n-1,m}} C_0^{n,m} \xrightarrow{(-1)^m d_1^{n,m}} C_0^{n+1,m} \longrightarrow \dots)$$

Then, the horizontal differentials of $C^{\bullet,\bullet}$ induce chain maps $C_0^{\bullet,m} \rightarrow C_0^{\bullet,m+1}$ for all $m \in \mathbb{Z}$, so that

$$C_0^{\bullet,\bullet} := (\dots \longrightarrow C_0^{\bullet,m-1} \xrightarrow{d_0^{\bullet,m-1}} C_0^{\bullet,m} \xrightarrow{d_0^{\bullet,m}} C_0^{\bullet,m+1} \xrightarrow{d_0^{\bullet,m+1}} \dots) \in \text{Ch}(\text{Ch}(\mathcal{A})).$$

In particular, to $C^{\bullet,\bullet} \in \text{bCh}(\mathcal{A})$ we associate, for all $n \in \mathbb{Z}$, the complex of n -coboundaries $B^n(C_0^{\bullet,\bullet})$, that of n -cocycles $Z^n(C_0^{\bullet,\bullet})$ and that of n -cohomology $H^n(C_0^{\bullet,\bullet})$, which are all objects of $\text{Ch}(\mathcal{A})$.

Finally, if \mathcal{A} is complete, we can also consider the **totalization** of the bicomplex $C^{\bullet,\bullet}$. This is the cochain complex $\text{Tot}(C^{\bullet,\bullet}) \in \text{Ch}(\mathcal{A})$, defined by letting $\text{Tot}(C^{\bullet,\bullet})^n := \prod_{i \in \mathbb{Z}} C^{i,n-i}$, for all $n \in \mathbb{Z}$, and with the n -th differential $\partial^n := (\partial_i^n)_{i \in \mathbb{Z}}: \prod_{i \in \mathbb{Z}} C^{i,n-i} \rightarrow \prod_{i \in \mathbb{Z}} C^{i,n+1-i}$, where ∂_i^n is $(d_0^{i,n-i}, d_1^{i,n-i})^t: C^{i,n-i} \rightarrow C^{i,n+1-i} \times C^{i+1,n-i}$ composed with the obvious projection and inclusion.

4.2. Classical Cartan-Eilenberg resolutions. Let \mathcal{A} be an Abelian category with enough injectives, and consider a (possibly unbounded) complex $A^\bullet \in \text{Ch}(\mathcal{A})$. An **injective Cartan-Eilenberg resolution** or, more briefly, a **CE-resolution** of A^\bullet is given by a lower half-plane bicomplex $C^{\bullet,\bullet}$ of injectives, and a homomorphism of complexes $\lambda^\bullet: A^\bullet \rightarrow C_0^{\bullet,\bullet}$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & A^{n-1} & \xrightarrow{a^{n-1}} & A^n & \xrightarrow{a^n} & A^{n+1} & \xrightarrow{a^{n+1}} & \dots \\ & & \downarrow \lambda^{n-1} & & \downarrow \lambda^n & & \downarrow \lambda^{n+1} & & \\ \dots & \longrightarrow & C^{0,n-1} & \xrightarrow{d_0^{0,n-1}} & C^{0,n} & \xrightarrow{d_0^{0,n}} & C^{0,n+1} & \xrightarrow{d_0^{0,n+1}} & \dots \\ & & \downarrow d_1^{0,n-1} & & \downarrow d_1^{0,n} & & \downarrow d_1^{0,n+1} & & \\ \dots & \longrightarrow & C^{1,n-1} & \xrightarrow{d_0^{1,n-1}} & C^{1,n} & \xrightarrow{d_0^{1,n}} & C^{1,n+1} & \xrightarrow{d_0^{1,n+1}} & \dots \\ & & \downarrow d_1^{1,n-1} & & \downarrow d_1^{1,n} & & \downarrow d_1^{1,n+1} & & \\ & & \vdots & & \vdots & & \vdots & & \end{array}$$

that satisfy the following properties:

- (CE.1) for each $n \in \mathbb{Z}$, if $A^n = 0$, then the column $C^{\bullet,n}$ is constantly 0;
- (CE.2) $B^n(\lambda^\bullet): B^n(A^\bullet) \rightarrow B^n(C_0^{\bullet,\bullet})$ is an injective resolution of $B^n(A^\bullet)$, for all $n \in \mathbb{Z}$;
- (CE.3) $H^n(\lambda^\bullet): H^n(A^\bullet) \rightarrow H^n(C_0^{\bullet,\bullet})$ is an injective resolution of $H^n(A^\bullet)$, for all $n \in \mathbb{Z}$.

If these three properties hold, then one also gets the following two extra properties (which are sometimes included in the definition) for free (see [Wei94, Exercise 5.7.1]):

- (CE.4) $Z^n(\lambda^\bullet): Z^n(A^\bullet) \rightarrow Z^n(C_0^{\bullet,\bullet})$ is an injective resolution of $Z^n(A^\bullet)$, for all $n \in \mathbb{Z}$;
- (CE.5) $\lambda^n: A^n \rightarrow C_0^{\bullet,n}$ is an injective resolution of A^n , for all $n \in \mathbb{Z}$.

It is well-known (see, e.g., [Wei94, Lemma 5.7.2]) that any complex A^\bullet admits a CE-resolution. Let us briefly go through the (dual of the) construction given by Weibel. The whole process is based on the Horseshoe Lemma, so let us briefly recall it here:

Lemma 4.1 (Horseshoe Lemma). *Let \mathcal{A} be an Abelian category and consider the following solid diagram, whose first and third row are injective resolutions of X and Z , respectively, the columns*

are exact, with ι^n and π^n ($n \in \mathbb{N}$) being the obvious inclusions and projections:

$$\begin{array}{ccccccc}
 0 & & 0 & & 0 & & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \longrightarrow & E_X^0 & \longrightarrow & E_X^1 & \longrightarrow & E_X^2 \longrightarrow \dots \\
 \downarrow & & \downarrow \iota^0 & & \downarrow \iota^1 & & \downarrow \iota^2 \\
 Y & \cdots \longrightarrow & E_X^0 \oplus E_Z^0 & \cdots \longrightarrow & E_X^1 \oplus E_Z^1 & \cdots \longrightarrow & E_X^2 \oplus E_Z^2 \cdots \\
 \downarrow & & \downarrow \pi^0 & & \downarrow \pi^1 & & \downarrow \pi^2 \\
 Z & \longrightarrow & E_Z^0 & \longrightarrow & E_Z^1 & \longrightarrow & E_Z^2 \longrightarrow \dots \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 0 & & 0 & & 0 & & 0
 \end{array}$$

One can then complete the second row to an injective resolution, so that the diagram commutes.

To construct a CE-resolution of A^\bullet , one proceeds as follows:

- (step.1) for all $n \in \mathbb{Z}$, fix injective resolutions $\lambda_{B^n} : B^n(A^\bullet) \rightarrow E_{B^n}^\bullet$ and $\lambda_{H^n} : H^n(A^\bullet) \rightarrow E_{H^n}^\bullet$;
- (step.2) for all $n \in \mathbb{Z}$, use the Horseshoe Lemma to combine these resolutions along the sequence $0 \rightarrow B^n(A^\bullet) \rightarrow Z^n(A^\bullet) \rightarrow H^n(A^\bullet) \rightarrow 0$, to get a new resolution $\lambda_{Z^n} : Z^n(A^\bullet) \rightarrow E_{Z^n}^\bullet$;
- (step.3) for all $n \in \mathbb{Z}$, use the Horseshoe Lemma to combine the previous injective resolutions along $0 \rightarrow Z^n(A^\bullet) \rightarrow A^n \rightarrow B^{n+1}(A^\bullet) \rightarrow 0$, to get a new injective resolution $\lambda_{A^n} : A^n \rightarrow E_{A^n}^\bullet$;
- (step.4) for all $n \in \mathbb{Z}$ and $m \in \mathbb{N}$, let $C^{m,n} := E_{A^n}^m$, with $d_1^{\bullet,n}$ using the differentials of $E_{A^n}^\bullet$ multiplied by $(-1)^n$, build $d_0^{\bullet,n}$ as a composition $E_{A^n}^\bullet \rightarrow E_{B^{n+1}}^\bullet \hookrightarrow E_{Z^{n+1}}^\bullet \hookrightarrow E_{A^{n+1}}^\bullet$ of the projection and embeddings given by the Horseshoe Lemma, and let $\lambda^\bullet : A^\bullet \rightarrow C^{0,\bullet}$ be $\lambda^n := \lambda_{A^n}$.

By [Wei94, Lemma 5.7.2], these $C^{\bullet,\bullet}$ and $\lambda^\bullet : A^\bullet \rightarrow C^{0,\bullet}$ are a CE-resolution. Furthermore, the following lemma is a consequence of the dual of [Wei94, Exercise 5.7.1]:

Lemma 4.2. *Let \mathcal{A} be a complete Abelian category with enough injectives, $A^\bullet \in \text{Ch}^+(\mathcal{A})$, and take a CE-resolution $\lambda^\bullet : A^\bullet \rightarrow C^{\bullet,\bullet}$. Then, $\text{Tot}(\lambda^\bullet) : A^\bullet \rightarrow \text{Tot}(C^{\bullet,\bullet})$ is a quasi-isomorphism.*

Observe that, in the setting of the above lemma, $\text{Tot}(C^{\bullet,\bullet})$ is degreewise injective and, by (CE.1), this complex is bounded below. Hence, by Proposition 1.3, $\text{Tot}(\lambda^\bullet) : A^\bullet \rightarrow \text{Tot}(C^{\bullet,\bullet})$ is a DG-injective resolution. Therefore, the totalization of CE-resolutions provides a valid alternative to the construction in Subsection 2.1.

4.3. CE-resolutions and the (Ab.4*)- k condition. In [Wei94, Exercise 5.7.1], Weibel suggests that Lemma 4.2 works for unbounded complexes, provided \mathcal{A} is (Ab.4*). In fact, this assumption can be weakened to (Ab.4*)- k :

Theorem 4.3. *Let \mathcal{A} be a complete Abelian category with enough injectives, let $A^\bullet \in \text{Ch}(\mathcal{A})$ be a complex, and consider a CE-resolution $\lambda^\bullet : A^\bullet \rightarrow C^{\bullet,\bullet}$. If \mathcal{A} is (Ab.4*)- k for some $k \in \mathbb{N}$, the induced map $A^\bullet \rightarrow \text{Tot}(C^{\bullet,\bullet})$ is a DG-injective resolution of A^\bullet .*

Proof. By definition of CE-resolution, we get injective resolutions $\lambda_{B^n} : B^n(A^\bullet) \rightarrow E_{B^n}^\bullet := B^n(C_0^{\bullet,\bullet})$ and $\lambda_{H^n} : H^n(A^\bullet) \rightarrow E_{H^n}^\bullet := H^n(C_0^{\bullet,\bullet})$, for each $n \in \mathbb{Z}$, so that $C^{n,m} \cong E_{B^n}^m \oplus E_{H^n}^m \oplus E_{B^{n+1}}^m$, for all $m \in \mathbb{N}$ and $n \in \mathbb{Z}$. Observe that, for each $n \in \mathbb{N}$, there is a CE-resolution $\lambda_n^\bullet : \tau^{\geq -n}(A^\bullet) \rightarrow C_{\geq -n}^{\bullet,\bullet}$, where $C_{\geq -n}^{\bullet,i} = C^{\bullet,i}$ for all $i > -n$, $C_{\geq -n}^{\bullet,i} = 0$ for all $i < -n$, and $C_{\geq -n}^{\bullet,n} = E_{H^n}^n \oplus E_{B^{n+1}}^n$ (with differentials induced by those of $C^{\bullet,\bullet}$). Denoting by $\pi_n^\bullet : \tau^{\geq -n-1}A^\bullet \rightarrow \tau^{\geq -n}(A^\bullet)$ and $\gamma_n^{\bullet,\bullet} : C_{\geq -n-1}^{\bullet,\bullet} \rightarrow C_{\geq -n}^{\bullet,\bullet}$ the obvious projections and taking the totalizations, we obtain the following Spaltenstein tower of partial resolutions (see Lemma 4.2):

$$\begin{array}{ccccccc}
 \dots & \xrightarrow{\pi_{-2}^\bullet} & \tau^{\geq -2}(A^\bullet) & \xrightarrow{\pi_{-1}^\bullet} & \tau^{\geq -1}(A^\bullet) & \xrightarrow{\pi_0^\bullet} & \tau^{\geq 0}(A^\bullet) \\
 & & \downarrow \lambda_2^\bullet & & \downarrow \lambda_1^\bullet & & \downarrow \lambda_0^\bullet \\
 \dots & \xrightarrow{\text{Tot}(\gamma_{-2}^{\bullet,\bullet})} & \text{Tot}(C_{\geq -2}^{\bullet,\bullet}) & \xrightarrow{\text{Tot}(\gamma_{-1}^{\bullet,\bullet})} & \text{Tot}(C_{\geq -1}^{\bullet,\bullet}) & \xrightarrow{\text{Tot}(\gamma_0^{\bullet,\bullet})} & \text{Tot}(C_{\geq 0}^{\bullet,\bullet}).
 \end{array}$$

By Theorem 2.9, the complex $\varprojlim_{i \in \mathbb{N}} \text{Tot}(C_{\geq -i}^{\bullet,\bullet}) \cong \text{Tot}(C^{\bullet,\bullet})$ is a DG-injective resolution of A^\bullet . \square

To understand what can go wrong if we do not assume (Ab.4*)- k , let us construct a CE-resolution for the complex $X^\bullet := \bigsqcup_{i \in \mathbb{Z}} S^i(R) \in \text{Ch}(\mathcal{G})$ from Section 3. For this specific complex: $B^n(X^\bullet) = 0$

and $Z^n(X^\bullet) \cong X^n \cong H^n(X^\bullet) \cong R$, for all $n \in \mathbb{Z}$. As in Lemma 3.2, we choose an injective resolution $\lambda: R \rightarrow E^\bullet$ of $R_R \in \text{Mod}(R)$, so that $\mathbf{Q}(E^\bullet) \in \text{Ch}^{\geq 0}(\mathcal{G})$ is an injective resolution of $R \in \mathcal{G}$. Hence, the following is a CE-resolution of X^\bullet :

$$(4.2) \quad \begin{array}{ccccccc} \cdots & \longrightarrow & R & \xrightarrow{0} & R & \xrightarrow{0} & R & \xrightarrow{0} & \cdots \\ & & \downarrow \mathbf{Q}(\lambda) & & \downarrow \mathbf{Q}(\lambda) & & \downarrow \mathbf{Q}(\lambda) & & \\ \cdots & \longrightarrow & \mathbf{Q}(E^0) & \xrightarrow{0} & \mathbf{Q}(E^0) & \xrightarrow{0} & \mathbf{Q}(E^0) & \xrightarrow{0} & \cdots \\ & & \downarrow \mathbf{Q}(e^0) & & \downarrow \mathbf{Q}(e^0) & & \downarrow \mathbf{Q}(e^0) & & \\ \cdots & \longrightarrow & \mathbf{Q}(E^1) & \xrightarrow{0} & \mathbf{Q}(E^1) & \xrightarrow{0} & \mathbf{Q}(E^1) & \xrightarrow{0} & \cdots \\ & & \downarrow \mathbf{Q}(e^1) & & \downarrow \mathbf{Q}(e^1) & & \downarrow \mathbf{Q}(e^1) & & \\ & & \vdots & & \vdots & & \vdots & & \end{array}$$

Totalizing this CE-resolution one gets the diagonal map $\prod_{i \in \mathbb{Z}} \Sigma^i \mathbf{Q}(\lambda): X^\bullet \rightarrow \prod_{i \in \mathbb{Z}} \Sigma^i \mathbf{Q}(E^\bullet)$, which is not a quasi-isomorphism by Proposition 3.3.

5. SANEBLIDZE'S CONSTRUCTION THROUGH MULTICOMPLEXES

A different construction of general DG-injective resolutions for (possibly unbounded) complexes is proposed in [San07]. Given an Abelian category \mathcal{A} , which is just supposed to have countable products and enough injectives, and $X^\bullet \in \text{Ch}(\mathcal{A})$, Saneblidze constructs what he calls a homological injective multicomplex $E^{\bullet, \bullet}$ (see below for unexplained terminology) whose totalization provides, supposedly, a DG-injective resolution for X^\bullet . In this section, after recalling the necessary background and some details about the construction proposed in [San07], we show that the same example considered in Section 3 is also a counterexample to this construction.

5.1. Cohomological injective multicomplexes and resolutions. Let \mathcal{A} be an Abelian category. A **multicomplex** $C^{\bullet, \bullet}$ over \mathcal{A} is given by the following data:

- a family of objects $C^{\bullet, \bullet} = \{C^{n,m} : (n,m) \in \mathbb{Z} \times \mathbb{Z}\}$ in \mathcal{A} ;
- for each $r \in \mathbb{N}$ a degree r differential $d_r^{\bullet, \bullet}: C^{\bullet, \bullet} \rightarrow C^{\bullet+r, \bullet-r+1}$, such that the following condition is satisfied for all $i \in \mathbb{N}$ and all $(n,m) \in \mathbb{Z} \times \mathbb{Z}$

$$(5.1) \quad \sum_{r+s=i} d_s^{n+r, m-r+1} \circ d_r^{n,m} = 0.$$

As we did for bicomplexes in Subsection 4, we restrict our attention to **lower half-plan multicomplexes** $C^{\bullet, \bullet}$, which can be visualized as follows (where labels denote the degree of the arrows)

$$\begin{array}{cccccccc} \cdots & \xrightarrow{0} & C^{0,m-2} & \xrightarrow{0} & C^{0,m-1} & \xrightarrow{0} & C^{0,m} & \xrightarrow{0} & C^{0,m+1} & \xrightarrow{0} & \cdots \\ & & \downarrow 1 & & \downarrow 1 & & \downarrow 1 & & \downarrow 1 & & \\ \cdots & \xrightarrow{0} & C^{1,m-2} & \xrightarrow{0} & C^{1,m-1} & \xrightarrow{0} & C^{1,m} & \xrightarrow{0} & C^{1,m+1} & \xrightarrow{0} & \cdots \\ & & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & & \\ \cdots & \xrightarrow{0} & C^{2,m-2} & \xrightarrow{0} & C^{2,m-1} & \xrightarrow{0} & C^{2,m} & \xrightarrow{0} & C^{2,m+1} & \xrightarrow{0} & \cdots \\ & & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & & \\ \cdots & \xrightarrow{0} & C^{3,m-2} & \xrightarrow{0} & C^{3,m-1} & \xrightarrow{0} & C^{3,m} & \xrightarrow{0} & C^{3,m+1} & \xrightarrow{0} & \cdots \\ & & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & \swarrow 2 & \downarrow 1 & & \\ & & \vdots & & \vdots & & \vdots & & \vdots & & \end{array}$$

For $i = 0$, (5.1) means that each row is a cochain complex while, for $i = 1$, it says that the little squares anticommute. Moreover, if $d_0^{\bullet, \bullet} = 0$ or $d_2^{\bullet, \bullet} = 0$, then (5.1) for $i = 2$ says that each column is a cochain complex. Hence, a bicomplex is just a multicomplex such that $d_r^{\bullet, \bullet} = 0$, for all $r \geq 2$.

A **morphism of multicomplexes** $(\phi_r^{\bullet, \bullet})_{r \in \mathbb{N}}: X^{\bullet, \bullet} \rightarrow Y^{\bullet, \bullet}$ is specified by a family of morphisms $\phi_r^{(n,m)}: X^{(n,m)} \rightarrow Y^{(n+r, m-r)}$ which is compatible with differentials in the following sense:

$$\sum_{r+s=i} \phi_s^{(n+r, m-r+1)} \circ d_r^{(n,m)} = \sum_{r+s=i} d_s^{(n+r, m-r)} \circ \phi_r^{(i,j)} \quad \text{for all } (n,m) \in \mathbb{Z} \times \mathbb{Z} \text{ and } i \in \mathbb{N}.$$

Definition 5.1 ([San07]). *A lower half-plane multicomplex of injectives $E^{\bullet,\bullet}$ is said to be a **homological injective multicomplex** if:*

- $d_0^{\bullet,\bullet} = 0$ (so that each column $E^{\bullet,m} \in \text{Ch}^{\geq 0}(\mathcal{A})$ with the degree 1 differentials);
- for each $m \in \mathbb{Z}$, $H^i(E^{\bullet,m}) = 0$ for all $i \geq 1$, that is, the column $E^{\bullet,m}$ is an injective resolution of the object $H^0(E^{\bullet,m}) = \text{Ker}(d_1^{0,m}) \in \mathcal{A}$.

Finally, if \mathcal{A} is complete, given a lower half-plane multicomplex $C^{\bullet,\bullet}$, one can consider its **totalization**, which is a cochain complex $\text{Tot}(C^{\bullet,\bullet}) \in \text{Ch}(\mathcal{A})$ such that $\text{Tot}(C^{\bullet,\bullet})^i := \prod_{j=0}^{\infty} C^{j,i-j}$, for all $i \in \mathbb{Z}$, where the n -th differential is

$$\partial^i := (\partial_j^i)_{j \in \mathbb{N}}: \prod_{j \in \mathbb{N}} C^{j,i-j} \longrightarrow \prod_{j \in \mathbb{N}} C^{j,i+1-j},$$

where ∂_j^i is determined by its components $(d_0^{j,i-j}, d_1^{j,i-j}, d_2^{j,i-j}, \dots)^t: C^{j,i-j} \rightarrow \prod_{r \in \mathbb{N}} C^{j+r,i+1-j-r}$.

Definition 5.2 ([San07]). *A **homological injective resolution** of a complex $X^{\bullet} \in \text{Ch}(\mathcal{A})$ is a morphism of multicomplexes $\lambda = (\lambda_r^{\bullet})_{r \in \mathbb{N}}: X^{\bullet} \rightarrow E^{\bullet,\bullet}$ such that $E^{\bullet,\bullet}$ is a homological injective multicomplex and $\text{Tot}(\lambda): X^{\bullet} \rightarrow \text{Tot}(E^{\bullet,\bullet})$ is a quasi-isomorphism.*

5.2. A counterexample for the construction. In [San07, Proposition 3], Saneblidze states that, if \mathcal{A} is a cocomplete Abelian category with enough projectives, then every complex has a homological projective resolution. In the proof of [San07, Proposition 3] one can find a rather involved construction of a homological resolution of an arbitrary complex. Let us recall the first few steps (in the dual setting of homological injective resolutions). Indeed, fix a generic complex $A^{\bullet} \in \text{Ch}(\mathcal{A})$. To build a homological injective resolution $\lambda = (\lambda_r^{\bullet})_{r \in \mathbb{N}}: A^{\bullet} \rightarrow E^{\bullet,\bullet}$. Consider, for each $n \in \mathbb{Z}$, an injective resolution $\lambda^n: H^n(A^{\bullet}) \rightarrow E_{H^n}^{\bullet} = (E_{H^n}^0 \rightarrow E_{H^n}^1 \rightarrow \dots)$ and define:

- $E^{m,n} := E_{H^n}^m$ for all $m \in \mathbb{N}$ and $n \in \mathbb{Z}$;
- define the differentials of degree 0 of $E^{\bullet,\bullet}$ to be trivial;
- use the differentials of $E_{H^n}^{\bullet}$ to define the differentials of degree 1 in $E^{\bullet,\bullet}$;
- let $\lambda_0^n: A^n \rightarrow E_{H^n}^0$ be any lifting of the composition $Z^n(A^{\bullet}) \rightarrow H^n(A^{\bullet}) \rightarrow E_{H^n}^0$ along the inclusion $Z^n(A^{\bullet}) \rightarrow A^n$ (which exists by the injectivity of $E_{H^n}^0$).

Hence, after this first step of the construction, we have all the object components of $E^{\bullet,\bullet}$, its horizontal and vertical differentials, and the 0-degree component of λ . As it turns out, since our goal is to apply the construction to the complex $X^{\bullet} \in \text{Ch}(\mathcal{G})$ of Section 3, which is a complex with trivial differentials, we can just stop here with the construction:

Remark 5.3 ([San07, pag. 320]). *As observed by Saneblidze right after [San07, Proposition 3], and as it can be easily checked in his proof, if all the differentials of A^{\bullet} are trivial, that is, if $A^{\bullet} \cong \prod_{n \in \mathbb{Z}} S^{-n}(A^n)$, one can take $d_r^{\bullet,\bullet} = 0$ for all $r \geq 2$ and $\lambda_s^{\bullet} = 0$, for all $s \geq 1$. Hence, in this case, $E^{\bullet,\bullet}$ is just a bicomplex with trivial horizontal differentials, and $\lambda = \lambda^{\bullet}: A^{\bullet} \rightarrow E^{0,\bullet}$.*

By the above discussion and remark, it is easy to see that the result of applying Saneblidze's construction to the complex $X^{\bullet} \in \text{Ch}(\mathcal{G})$ from Section 3 is the bicomplex we have constructed in (4.2) when studying CE-resolutions. As we have already seen, the totalization of that bicomplex is not quasi-isomorphic to X^{\bullet} and, therefore, Saneblidze's construction does not produce a homological injective resolution in this case.

6. DING AND YANG'S CONSTRUCTION VIA REPEATED KILLING OF COBOUNDARIES

Let \mathcal{A} be a bicomplete Abelian category, $(\mathcal{X}, \mathcal{Y})$ a complete hereditary cotorsion pair in \mathcal{A} , and consider the induced cotorsion pairs $(\text{dg}\mathcal{X}, \tilde{\mathcal{Y}})$ and $(\tilde{\mathcal{X}}, \text{dg}\mathcal{Y})$ in $\text{Ch}(\mathcal{A})$, introduced by Gillespie [Gil08]. In many important cases, these cotorsion pairs are complete, hereditary and compatible, so they give rise to an Abelian model structure in $\text{Ch}(\mathcal{A})$. On the other hand, to the best of our knowledge, the following question remains open:

Question 6.1. *Let \mathcal{A} be a bicomplete Abelian category and let $(\mathcal{X}, \mathcal{Y})$ be a complete and hereditary cotorsion pair in \mathcal{A} . Are the induced cotorsion pairs $(\text{dg}\mathcal{X}, \tilde{\mathcal{Y}})$ and $(\tilde{\mathcal{X}}, \text{dg}\mathcal{Y})$ complete in $\text{Ch}(\mathcal{A})$?*

An attempt to solve this problem in the positive was made by N. Ding and X. Yang. The argument they used is based on [YD15, Lemma 2.1], whose proof contains a very concrete construction,

for each $A^\bullet \in \text{Ch}(\mathcal{A})$, of a $Y^\bullet \in \text{dg}\mathcal{Y}$ which is quasi-isomorphic to A^\bullet . In particular, if we start with the trivial cotorsion pair $(\mathcal{A}, \text{Inj}(\mathcal{A}))$ in \mathcal{A} (assuming that \mathcal{A} has enough injectives), this construction should produce a DG-injective resolution of A^\bullet .

In this section we test the proof of [YD15, Lemma 2.1] against the complex $X^\bullet \in \text{Ch}(\mathcal{G})$ from Section 3, showing that it actually fails to produce the desired resolution. Furthermore, we briefly analyze Ding and Yang's proof, pointing out a gap in the argument, and showing that the problem completely disappears if one assume that \mathcal{A} is $(\text{Ab}, 4^*)$.

6.1. The Ding-Yang construction in concrete situations. Let \mathcal{A} be a bicomplete Abelian category with enough injectives, so that the cotorsion pair $(\mathcal{A}, \text{Inj}(\mathcal{A}))$ is complete. Let us start recalling the idea of the proof of [YD15, Lemma 2.1] in this very special case. Indeed, the construction of Ding and Yang is based on a very simple operation, of “killing coboundaries” that we have condensed in the following

Construction 6.2 (Killing coboundaries). *Let \mathcal{A} be an Abelian category with enough injectives, fix a complex $X^\bullet \in \text{Ch}(\mathcal{A})$ and $n \in \mathbb{Z}$. Denote by $\rho^n: X^n \rightarrow X^n/B^n(X^\bullet)$ the obvious projection and by $\iota^n: X^n/B^n(X^\bullet) \rightarrow E$ an inclusion into an injective object $E \in \text{Inj}(\mathcal{A})$. Define a new complex:*

$$K(X^\bullet, n) : \quad \dots \longrightarrow X^{n-1} \xrightarrow{d^{n-1}} X^n \xrightarrow{\begin{bmatrix} d^n \\ \iota^n \rho^n \end{bmatrix}} X^{n+1} \oplus E \xrightarrow{\begin{bmatrix} d^{n+1} & 0 \end{bmatrix}} X^{n+2} \xrightarrow{d^{n+2}} \dots$$

and a morphism of complexes $\pi_n^\bullet: K(X^\bullet, n) \rightarrow X^\bullet$ such that

$$\pi_n^m := \begin{cases} \text{id}_{X^m} & \text{for all } m \neq n+1; \\ \text{the obvious projection } X^{n+1} \oplus E \rightarrow X^{n+1} & \text{otherwise.} \end{cases}$$

Then, $H^n(K(X^\bullet, n)) = 0$, π_n^\bullet is a degreewise split-epic, and $\text{Ker}(\pi_n^\bullet) = S^{n+1}(E)$ is DG-injective.

Observe that $K(X^\bullet, n)$ depends on the choice of the embedding $\iota^n: X^n/B^n(X^\bullet) \rightarrow E$. In some cases this choice can be made canonical, e.g., whenever $X^n/B^n(X^\bullet)$ is trivial or, more generally, when it is an injective object, it is natural to take $\iota^n := \text{id}$. To illustrate the construction in some basic case, take an injective object $E \in \mathcal{A}$ and consider the stalk complex $S^i(E)$ and the disk complex $D^i(E)$, for some $i \in \mathbb{Z}$. Then,

$$K(S^i(E), j) = \begin{cases} S^i(E) & \text{if } j \neq i; \\ D^i(E) & \text{if } j = i; \end{cases} \quad \text{and} \quad K(D^i(E), j) = \begin{cases} D^i(E) & \text{if } j \neq i; \\ D^i(E) \oplus S^{i+1}(E) & \text{if } j = i. \end{cases}$$

In fact, it is not completely trivial to verify that $K(D^i(E), i) \cong D^i(E) \oplus S^{i+1}(E)$ since the i -th differential of $K(D^i(E), i)$ is, a priori, just a triangular (but not diagonal) matrix. In fact, this kind of complications with the differentials can be solved with the following general lemma:

Lemma 6.3. *Given $X^\bullet = (X^n, d^n)_{n \in \mathbb{Z}} \in \text{Ch}(\mathcal{A})$ such that, for a given $n \in \mathbb{Z}$, $X^n \cong Y^{(k)}$ is a coproduct of k -many copies of a given object $Y \in \mathcal{A}$. If $d^{n-1} = (\varphi, \varphi, \dots, \varphi)^t: X^{n-1} \rightarrow Y^{(k)}$ is the diagonal map for some morphism $\varphi: X^{n-1} \rightarrow Y$. Then, there is an isomorphism of complexes $\phi^\bullet: X^\bullet \rightarrow (X')^\bullet = ((X')^n, (d')^n)_{n \in \mathbb{Z}}$, where $(X')^i := X^i$ for all $i \in \mathbb{Z}$, $(d')^i := d^i$ for all $i \neq n-1, n$, $(d')^{n-1} := (\varphi, 0, \dots, 0)^t$, and $(d')^n := (\psi_1 + \dots + \psi_k, \psi_2, \dots, \psi_k)$, where $d^n = (\psi_1, \psi_2, \dots, \psi_k): Y^{(k)} \rightarrow X^{n+1}$.*

Proof. One can just define $\phi^i := \text{id}_{X^i}$ for all $i \neq n$, and

$$\phi^n := \begin{bmatrix} +\text{id}_Y & 0 & \dots & 0 & 0 \\ -\text{id}_Y & \text{id}_Y & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -\text{id}_Y & 0 & \dots & \text{id}_Y & 0 \\ -\text{id}_Y & 0 & \dots & 0 & \text{id}_Y \end{bmatrix} \in \text{Mat}_{k \times k}(\text{End}_{\mathcal{A}}(Y)) \cong \text{End}_{\mathcal{A}}(Y^{(k)}).$$

□

Consider now a family of complexes $\{X_n^\bullet\}_{n \in \mathbb{N}} \in \text{Ch}(\mathcal{A})$ such that

$$|\{n \in \mathbb{N} : X_n^i \neq 0\}| < \infty, \quad \text{for each } i \in \mathbb{Z}.$$

Observe that, for such a family, $\prod_{\mathbb{N}} X_n^\bullet = \bigoplus_{\mathbb{N}} X_n^\bullet$. Hence, letting $Y^\bullet := \prod_{\mathbb{N}} X_n^\bullet = \bigoplus_{\mathbb{N}} X_n^\bullet$, we have that $B^i(Y^\bullet) = \bigoplus_{\mathbb{N}} B^i(X_n^\bullet) = \prod_{\mathbb{N}} B^i(X_n^\bullet)$ (where both equalities use that finite products coincide with finite coproducts in any Abelian category and, hence, they are exact). Similarly,

one can verify that $Z^i(Y^\bullet) = \prod_{\mathbb{N}} Z^i(X_n^\bullet) = \bigoplus_{\mathbb{N}} Z^i(X_n^\bullet)$ and, combining these two, one even gets $H^i(Y^\bullet) = \prod_{\mathbb{N}} H^i(X_n^\bullet) = \bigoplus_{\mathbb{N}} H^i(X_n^\bullet)$.

Lemma 6.4. *Let $\{X_n^\bullet\}_{n \in \mathbb{N}} \in \text{Ch}(\mathcal{A})$ be a family as above and let $Y^\bullet := \prod_{\mathbb{N}} X_n^\bullet$. Then, for each $i \in \mathbb{Z}$, one can choose suitable embeddings into injectives in Construction 6.2 so to obtain $K(Y^\bullet, i) = \prod_{\mathbb{N}} K(X_n^\bullet, i)$.*

Going back to Ding and Yang's proof of their Lemma 2.1, they fix an arbitrary surjection $\tau: \mathbb{N} \rightarrow \mathbb{Z}$ with $n_i := \tau(i)$, for all $i \in \mathbb{N}$, with the property of having infinite fibers. As we are trying to do some concrete computations, we need to make a choice. Indeed, we fix the following sequence of integers (in which every integer appears infinitely many times, as desired):

$$\begin{aligned} n_0 &= 0, \\ n_1 &= -1, \quad n_2 = 0, \quad n_3 = 1, \\ n_4 &= -2, \quad n_5 = -1, \quad n_6 = 0, \quad n_7 = 1, \quad n_8 = 2, \\ n_9 &= -3, \quad n_{10} = -2, \quad n_{11} = -1, \quad n_{12} = 0, \quad n_{13} = 1, \quad n_{14} = 2, \quad n_{15} = 3, \\ &\dots \end{aligned}$$

Now, starting with our complex $X^\bullet \in \text{Ch}(\mathcal{A})$, the idea of the Ding-Yang construction is to iterate Construction 6.2 in order to build the following sequential inverse system:

$$\dots \longrightarrow Y_i^\bullet \xrightarrow{\pi_i^\bullet} \dots \xrightarrow{\pi_2^\bullet} Y_1^\bullet \xrightarrow{\pi_1^\bullet} Y_0^\bullet \xrightarrow{\pi_0^\bullet} Y_{-1}^\bullet := X^\bullet \quad \text{in } \text{Ch}(\mathcal{A}),$$

where $Y_i^\bullet := K(Y_{i-1}^\bullet, n_i)$ and π_i^\bullet is the obvious degreewise split-epimorphism with DG-injective kernel, for all $i \in \mathbb{N}$. Defining $Y^\bullet := \varprojlim_{\mathbb{N}} Y_i^\bullet$, one gets an exact sequence $0 \rightarrow E^\bullet \rightarrow Y^\bullet \rightarrow X^\bullet \rightarrow 0$ in $\text{Ch}(\mathcal{A})$, with E^\bullet a DG-injective complex. The claim made in [YD15] is that the complex Y^\bullet is exact, so there is a quasi-isomorphism $X^\bullet \rightarrow \Sigma E^\bullet$, which is precisely the DG-injective resolution of X^\bullet we are looking for. In the rest of this section we will apply this strategy in some easy cases: first with X^\bullet a stalk complex, and then a product of stalk complexes. As we will see, if the product of stalk complexes X^\bullet is the complex from Section 3, then the resulting complex Y^\bullet is not exact, showing that the proof of [YD15, Lemma 2.1] may fail even for the trivial cotorsion pair $(\mathcal{G}, \text{Inj}(\mathcal{G}))$, with \mathcal{G} a Grothendieck category.

As a first example, let us apply the Ding-Yang construction to a stalk complex $X = S^0(A)$ concentrated in degree 0, for some $A \in \mathcal{A}$. Fix an injective resolution for A , and use it to build the following exact complex concentrated in non-negative degrees:

$$E_A^{[0, \infty)} := (\dots \longrightarrow 0 \longrightarrow A \xrightarrow{\varphi} E^1 \xrightarrow{\lambda^1} E^2 \xrightarrow{\lambda^2} E^3 \xrightarrow{\lambda^3} \dots).$$

For each $m \in \mathbb{N}$, let $E_A^{[0, m]}$ be the naive truncation of $E_A^{[0, \infty)}$ so that, for example,

$$E_A^{[0, 0]} = S^0(A), \quad E_A^{[0, 1]} = (\dots \longrightarrow 0 \longrightarrow A \longrightarrow E^1 \longrightarrow 0 \longrightarrow \dots), \quad \text{and so on.}$$

We can now start computing the complexes of the form $Y_i^\bullet := K(Y_{i-1}^\bullet, n_i)$, for $i \in \mathbb{N}$:

$$\begin{aligned} (i = 0): & Y_0^\bullet = K(S^0(A), 0) = E_A^{[0, 1]}; \\ (i = 1): & Y_1^\bullet = Y_0^\bullet; \\ (i = 2): & Y_2^\bullet = E_A^{[0, 1]} \oplus S^1(E^1) \text{ (by a suitable application of Lemma 6.3);} \\ (i = 3): & Y_3^\bullet = E_A^{[0, 2]} \oplus D^1(E^1) \text{ (use Lemma 6.4);} \\ (i = 4, 5): & Y_5^\bullet = Y_4^\bullet = Y_3^\bullet; \\ (i = 6): & Y_6^\bullet = E_A^{[0, 2]} \oplus S^1(E^1) \oplus D^1(E^1); \\ (i = 7): & Y_7^\bullet = E_A^{[0, 2]} \oplus D^1((E^1)^2) \oplus S^2(E^1 \oplus E^2); \\ (i = 8): & Y_8^\bullet = E_A^{[0, 3]} \oplus D^1((E^1)^2) \oplus D^2(E^1 \oplus E^2); \\ (i = 9, 10, 11): & Y_{11}^\bullet = Y_{10}^\bullet = Y_9^\bullet = Y_8^\bullet; \\ (i = 12): & Y_{12}^\bullet = E_A^{[0, 3]} \oplus S^1(E^1) \oplus D^1((E^1)^2) \oplus D^2(E^1 \oplus E^2); \\ & \dots \end{aligned}$$

After computing these first steps it seems clear that the limit $Y^\bullet := \varprojlim(\cdots \rightarrow Y_n^\bullet \rightarrow \cdots \rightarrow Y_0^\bullet)$ will be the following complex:

$$Y^\bullet \cong E_A^{[0,\infty)} \times \prod_{i=1}^{\infty} \left(\prod_{j=1}^i (D^i(E^j))^{\mathbb{N}} \right).$$

In other words, Y^\bullet is a product of our chosen resolution of A (including A , so this part is an exact complex) with a bunch of disk complexes, so Y^\bullet is an exact complex. This means that the Ding-Yang construction works just fine when applied to a single stalk complex.

Suppose now that, instead of starting with a single stalk complex, we start with the complex $X^\bullet = \bigoplus_{n \in \mathbb{Z}} S^n(A) = \prod_{n \in \mathbb{Z}} S^n(A)$, for some object $A \in \mathcal{A}$, that is:

$$X^\bullet = \cdots \xrightarrow{0} A \xrightarrow{0} A \xrightarrow{0} \cdots \xrightarrow{0} A \xrightarrow{0} \cdots$$

By Lemma 6.4, if we choose carefully the embeddings into injective objects, the resulting Y^\bullet is:

$$Y^\bullet = \prod_{n \in \mathbb{Z}} \Sigma^n \left(E_A^{[0,\infty)} \times D^0 \left(\prod_{j=0}^{\infty} (E^j)^{\mathbb{N}} \right) \right).$$

It is easy to deduce from this description that Y^\bullet is quasi-isomorphic to $\prod_{n \in \mathbb{Z}} \Sigma^n E_A^{[0,\infty)}$; we claim that such a product of exact complexes is not necessarily exact. Indeed, if $\mathcal{A} = \mathcal{G}$ is the Grothendieck category we have introduced in Section 3, and if $A := R \in \mathcal{G}$, then the object $X^\bullet := \prod_{i \in \mathbb{Z}} S^i(R) \in \text{Ch}(\mathcal{G})$ is the same complex we have considered in Section 3. Take the following short exact sequence of complexes:

$$0 \longrightarrow X^\bullet \longrightarrow \prod_{n \in \mathbb{Z}} \Sigma^n E_R^{[0,\infty)} \longrightarrow \prod_{n \in \mathbb{Z}} \Sigma^n (E^\bullet) \longrightarrow 0.$$

If, looking for a contradiction, $\prod_{n \in \mathbb{Z}} \Sigma^n E_R^{[0,\infty)}$ was acyclic, all of its cohomologies would be trivial and, analyzing the long exact sequence of cohomologies, this would force isomorphisms $H_G^i(\prod_{n \in \mathbb{Z}} \Sigma^n (E^\bullet)) \cong H_G^{i+1}(X^\bullet) = R$, for all $i \in \mathbb{Z}$. By Proposition 3.3, we know that this is not the case, so $\prod_{n \in \mathbb{Z}} \Sigma^n E_R^{[0,\infty)}$ cannot be exact in $\text{Ch}(\mathcal{G})$.

6.2. The problem in the proof of [YD15, Lemma 2.1]. As shown in the previous subsection, the proof of [YD15, Lemma 2.1] fails in general. For this reason, it may be interesting to identify the concrete problem in Ding and Yang's argument and, if possible, to find additional hypotheses under which the proposed construction can be made to work. The unique problem we could identify in [YD15] is the following: on page 3210 in [Op.Cit.], in the last part of the proof of Lemma 2.1, there is an inverse system (in the notation of [Op.Cit.]):

$$\cdots \xrightarrow{\mu^{i+1}} Y^i \xrightarrow{\mu^i} \cdots \xrightarrow{\mu^2} Y^1 \xrightarrow{\mu^1} Y^0 \quad \subseteq \text{Ch}(\mathcal{G})$$

and an $l \in \mathbb{Z}$ such that $H_l(Y^i) = 0$ for all $i \in \mathbb{N}$. The authors want to prove that the following map

$$1 - \nu: \prod_{i=0}^{\infty} Y^i \longrightarrow \prod_{i=0}^{\infty} Y^i$$

induces an isomorphism in homology at degree l . They call $\pi^j: \prod_{i=0}^{\infty} Y^i \rightarrow Y^j$ the canonical projection, for each $j \geq 0$, they consider $\pi^j \circ (1 - \nu) = \pi^j - \mu^{j+1} \circ \pi^{j+1}$ and correctly verify that

$$(6.1) \quad \mu_l^{j+1} \pi_l^{j+1} \left(Z_l^{\prod_{i=0}^{\infty} Y^i} \right) \subseteq B_l^{Y^j}, \quad \text{for all } j \in \mathbb{Z}.$$

Unfortunately, (6.1) does not imply that $Z_l^{\prod_{i=0}^{\infty} Y^i} \subseteq B_l^{\prod_{i=0}^{\infty} Y^i}$ but just the weaker inclusion: $Z_l^{\prod_{i=0}^{\infty} Y^i} \subseteq \prod_{i=0}^{\infty} B_l^{Y^i}$. In fact, it may happen that $\prod_{i=0}^{\infty} B_l^{Y^i} \neq B_l^{\prod_{i=0}^{\infty} Y^i}$ (e.g., in the setting of Section 3): the property “the boundaries of the product coincide with the product of boundaries” (if required for all possible products) is in fact equivalent to the (Ab.4*) condition on \mathcal{G} . In particular, adding the (Ab.4*) condition to the hypotheses, the proof of [YD15, Lemma 2.1(1)] works perfectly. We will also obtain the conclusion of [YD15, Lemma 2.1(1)] under a different set of hypotheses in Remark 8.6.

7. MODEL STRUCTURES FOR RELATIVE HOMOLOGICAL ALGEBRA

In this section we combine some of the main results of [CNPS18] about model approximations for relative homological algebra, and a general criterion for the existence of suitable model categories from [CH02] to verify that, if \mathcal{A} is a bicomplete Abelian category, and \mathcal{I} an injective class of objects such that \mathcal{A} is (Ab.4*)- \mathcal{I} - k (for some $k \in \mathbb{N}$), then there is an induced \mathcal{I} -injective model structure on $\text{Ch}(\mathcal{A})$; in particular, the \mathcal{I} -derived category $\mathcal{D}(\mathcal{A}; \mathcal{I})$ is locally small.

7.1. Injective classes. Given a class $\mathcal{I} \subseteq \mathcal{A}$, a morphism $\phi: X \rightarrow Y$ in \mathcal{A} is said to be an **\mathcal{I} -monomorphism** if

$$\text{Hom}_{\mathcal{A}}(\phi, I): \text{Hom}_{\mathcal{A}}(Y, I) \longrightarrow \text{Hom}_{\mathcal{A}}(X, I)$$

is surjective (in Ab), for all $I \in \mathcal{I}$. With this concept at hand, we can now introduce the following:

Definition 7.1. A class of objects \mathcal{I} in an Abelian category \mathcal{A} is said to be an **injective class** if:

(IC.1) \mathcal{I} is closed under products and summands;

(IC.2) for each $A \in \mathcal{A}$ there is an \mathcal{I} -monomorphism $\phi: A \rightarrow I$, with $I \in \mathcal{I}$.

Observe that a class \mathcal{I} that satisfies the property (IC.2) is usually called **pre-enveloping**. In the following lemma we show that the above definition is slightly redundant, in fact, there is no need to ask that \mathcal{I} is closed under products in (IC.1):

Lemma 7.2. Let \mathcal{A} be a complete Abelian category. Then, a pre-enveloping class $\mathcal{I} \subseteq \mathcal{A}$ which is closed under direct summands is also closed under products.

Proof. Consider a set $\{X_\lambda\}_{\lambda \in \Lambda}$ of objects in \mathcal{I} , let $X := \prod_{\lambda \in \Lambda} X_\lambda$ and for each $\lambda \in \Lambda$ denote by $\pi_\lambda: X \rightarrow X_\lambda$ the canonical projection. By hypothesis, there is an \mathcal{I} -monomorphism $\varphi: X \rightarrow I$, for some $I \in \mathcal{I}$. Since $X_\lambda \in \mathcal{I}$ for each $\lambda \in \Lambda$, the following map is surjective:

$$\text{Hom}_{\mathcal{A}}(\varphi, E_\lambda): \text{Hom}_{\mathcal{A}}(I, E_\lambda) \longrightarrow \text{Hom}_{\mathcal{A}}(X, E_\lambda),$$

so there is $\psi_\lambda \in \text{Hom}_{\mathcal{A}}(I, E_\lambda)$ such that $\psi_\lambda \circ \varphi = \pi_\lambda$. By the universal property of the product, there is a unique $\psi: I \rightarrow X$ such that $\pi_\lambda \circ \psi = \psi_\lambda$, for all $\lambda \in \Lambda$. Moreover, for each $\lambda \in \Lambda$:

$$\pi_\lambda \circ \psi \circ \varphi = \psi_\lambda \circ \varphi = \pi_\lambda = \pi_\lambda \circ \text{id}_X.$$

By the uniqueness in the universal property of the product, $\psi \circ \varphi = \text{id}_X$ and, therefore, X is a summand of $I \in \mathcal{I}$. Hence, $X \in \mathcal{I}$, as desired. \square

If \mathcal{A} is an Abelian category with enough injectives (e.g., if \mathcal{A} is Grothendieck) then $\mathcal{I} := \text{Inj}(\mathcal{A})$, the class of all the injective objects in \mathcal{A} , is an injective class for which the \mathcal{I} -monomorphisms are the usual monomorphisms. We refer to Section 8 for other concrete examples.

7.2. Relative \mathcal{I} -injective resolutions of objects. Let \mathcal{A} be an Abelian category. Given $X^\bullet \in \text{Ch}(\mathcal{A})$ and $A \in \mathcal{A}$, we define the cochain complex $\text{Hom}(X^\bullet, A) \in \text{Ch}(\text{Ab})$ as follows:

- for each $n \in \mathbb{Z}$, let $\text{Hom}(X^\bullet, A)^n := \text{Hom}_{\mathcal{A}}(X^{-n}, A)$;
- for each $n \in \mathbb{Z}$, the $(n-1)$ -th differential of $\text{Hom}(X^\bullet, A)$ is the following map:

$$(d^{-n})^*: \text{Hom}_{\mathcal{A}}(X^{-n+1}, A) \longrightarrow \text{Hom}_{\mathcal{A}}(X^{-n}, A),$$

such that $(d^{-n})^*(f) := f \circ d^{-n}$, for all $f \in \text{Hom}_{\mathcal{A}}(X^{-n+1}, A)$.

Moreover, any morphism $\phi^\bullet: X^\bullet \rightarrow Y^\bullet$ in $\text{Ch}(\mathcal{A})$, induces the following morphism in $\text{Ch}(\text{Ab})$:

$$\text{Hom}(\phi^\bullet, A): \text{Hom}(Y^\bullet, A) \longrightarrow \text{Hom}(X^\bullet, A) \quad \text{such that} \quad (\text{Hom}(\phi^\bullet, A))(g) := g \circ \phi^\bullet,$$

for all $g \in \text{Hom}(Y^\bullet, A)$. In particular, this gives a functor $\text{Hom}(-, A): (\text{Ch}(\mathcal{A}))^{\text{op}} \rightarrow \text{Ch}(\text{Ab})$.

Definition 7.3. Let \mathcal{A} be a complete Abelian category, let \mathcal{I} be an injective class, and let $A \in \mathcal{A}$. A **relative \mathcal{I} -injective resolution** of A is a pair $(I^\bullet, u: S^0(A) \rightarrow I^\bullet)$ such that:

- $I^\bullet \in \text{Ch}^{\geq 0}(\mathcal{I}) \subseteq \text{Ch}^{\geq 0}(\mathcal{A})$;
- $\text{Hom}(u, I): \text{Hom}(I^\bullet, I) \rightarrow \text{Hom}(S^0(A), I)$ is a quasi-isomorphism, for all $I \in \mathcal{I}$.

Let us give some equivalent reformulations of the above definition:

Lemma 7.4. *Let \mathcal{A} be a complete Abelian category, $\mathcal{I} \subseteq \mathcal{A}$ an injective class, and take a complex:*

$$\tilde{I}^\bullet: \quad \cdots \longrightarrow 0 \longrightarrow A \xrightarrow{d^{-1}} I^0 \xrightarrow{d^0} I^1 \xrightarrow{d^1} \cdots \longrightarrow I^{n-1} \xrightarrow{d^{n-1}} I^n \xrightarrow{d^n} \cdots \in \text{Ch}^{\geq -1}(\mathcal{A}),$$

with $I^j \in \mathcal{I}$, for all $j \geq 0$. Denote by $I^\bullet := (\cdots \rightarrow 0 \rightarrow I^0 \xrightarrow{d^0} \cdots \rightarrow I^{n-1} \xrightarrow{d^{n-1}} I^n \xrightarrow{d^n} \cdots)$ the naive truncation above 0, and by $d^{-1}: S^0(A) \rightarrow I^\bullet$ the obvious map. Then, the following are equivalent:

- (1) $(I^\bullet, d^{-1}: S^0(A) \rightarrow I^\bullet)$ is a relative \mathcal{I} -injective resolution of A ;
- (2) $\text{Hom}(\tilde{I}^\bullet, I)$ is exact, for all $I \in \mathcal{I}$;
- (3) $\tilde{I}^k/B^k(\tilde{I}^\bullet) \rightarrow I^{k+1}$ is an \mathcal{I} -monomorphism, for all $k \geq -1$.

Proof. The equivalence “(1) \Leftrightarrow (2)” is trivial; let us verify that (2) is also equivalent to (3). Indeed, for each $I \in \mathcal{I}$ and $k \geq -1$, consider the following sequence:

$$(7.1) \quad \text{Hom}_{\mathcal{A}}(\tilde{I}^{k+1}, I) \xrightarrow{(d^k)^*} \text{Hom}_{\mathcal{A}}(\tilde{I}^k, I) \xrightarrow{(d^{k-1})^*} \text{Hom}_{\mathcal{A}}(\tilde{I}^{k-1}, I),$$

Observe that, being $\text{Hom}_{\mathcal{A}}(-, I): \mathcal{A}^{\text{op}} \rightarrow \text{Ab}$ a left-exact functor, there is a canonical isomorphism $\text{Hom}_{\mathcal{A}}(\tilde{I}^k/B^k(\tilde{I}^\bullet), I) = \text{Hom}_{\mathcal{A}}(\text{Coker}(d^{k-1}), I) \cong \ker((d^{k-1})^*)$. In particular, the sequence (7.1) is exact precisely when the induced map $\tilde{I}^k/B^k(\tilde{I}^\bullet) \rightarrow I^{k+1}$ is an \mathcal{I} -monomorphism. \square

Given an Abelian category \mathcal{A} and an injective class $\mathcal{I} \subseteq \mathcal{A}$, since \mathcal{A} has enough \mathcal{I} -injectives, it is easy to construct a relative \mathcal{I} -injective resolution for any object $A \in \mathcal{A}$ by induction, just using again and again the equivalence “(1) \Leftrightarrow (3)” in the above lemma.

Lemma 7.5. *Let \mathcal{A} be an Abelian category, let \mathcal{I} be an injective class in \mathcal{A} , let $(C^\bullet, \partial^\bullet)$ and $(D^\bullet, d^\bullet) \in \text{Ch}(\mathcal{A})$ be two cochain complexes, and let $0 \leq r$ be a natural number such that,*

- (1) $D^i \in \mathcal{I}$, for all $i > r$;
- (2) $H^{-i}(\text{Hom}(C^\bullet, I)) = 0$, for all $I \in \mathcal{I}$ and all $i \geq r$.

Then, any family of morphisms $(f^k: C^k \rightarrow D^k)_{k \leq r}$ such that $f^k \circ \delta^{k-1} = d^{k-1} \circ f^{k-1}$, for all $k \leq r$, extends to a morphism of complexes $f^\bullet: C^\bullet \rightarrow D^\bullet$. Moreover, any two such extensions are homotopic via a homotopy $(h^n: C^n \rightarrow D^{n-1})_{n \in \mathbb{Z}}$ such that $h^k = 0$, for all $k \leq r$.

Proof. Our hypothesis (2) means that the following sequence is exact in Ab

$$(7.2) \quad \text{Hom}_{\mathcal{A}}(C^{j+1}, I) \xrightarrow{(\delta^j)^*} \text{Hom}_{\mathcal{A}}(C^j, I) \xrightarrow{(\delta^{j-1})^*} \text{Hom}_{\mathcal{A}}(C^{j-1}, I),$$

for all $I \in \mathcal{I}$ and all $j \geq r$. Then, by (1), the sequence (7.2) is then provided $I = D^{j+1}$ and, for $j > r$, also for $I = D^j$.

We can now proceed to construct $f^\bullet: C^\bullet \rightarrow D^\bullet$ by induction. Indeed, suppose that we have morphisms $(f^n: C^n \rightarrow D^n)_{n \leq j}$, for some $j \geq r$, such that $f^n \circ \delta^{n-1} = d^{n-1} \circ f^{n-1}$, for all $n \leq j$. In particular, taking $I = D^{j+1}$ in (7.2), we deduce that

$$(\delta^{j-1})^*(d^j \circ f^j) = d^j \circ f^j \circ \delta^{j-1} = d^j \circ d^{j-1} \circ f^{j-1} = 0,$$

that is, $d^j \circ f^j \in \ker((\delta^{j-1})^*) = \text{Im}((\delta^j)^*)$ and, therefore, there is some $f^{j+1} \in \text{Hom}_{\mathcal{A}}(C^{j+1}, D^{j+1})$ such that $(\delta^j)^*(f^{j+1}) = f^{j+1} \circ \delta^j = d^j \circ f^j$. The family $(f^n: C^n \rightarrow D^n)_{n \leq j+1}$ now satisfies $f^n \circ \delta^{n-1} = d^{n-1} \circ f^{n-1}$, for all $n \leq j+1$, and the induction can continue.

Assume now that $f^\bullet, g^\bullet \in \text{Hom}_{\text{Ch}(\mathcal{A})}(C^\bullet, D^\bullet)$ are both extensions of the family $(f^k)_{k \leq r}$. We proceed by induction to construct $(h^n: C^{n+1} \rightarrow D^n)_{n \in \mathbb{Z}}$ such that $f^n - g^n = d^{n-1} \circ h_{n-1} + h^n \circ \delta^n$, for all $n \in \mathbb{Z}$. Indeed, let $h^i = 0$, for all $i \leq r$ and suppose that, for some $j > r$ we have constructed a family $(h^n: C^{n+1} \rightarrow D^n)_{n \leq j}$ such that $f^n - g^n = d^{n-1} \circ h^{n-1} + h^n \circ \delta^n$, for all $n \leq j$. In particular, using that $(f^\bullet - g^\bullet)$ is a map of complexes, we deduce that

$$(f^{j+1} - g^{j+1}) \circ \delta^j = d^j \circ (f^j - g^j) = d^j \circ d^{j-1} \circ h^{j-1} + d^j \circ h^j \circ \delta^j = d^j \circ h^j \circ \delta^j.$$

Hence, taking $I = D^j$ in (7.2), and using the above computation, we get:

$$(\delta^j)^*(f^{j+1} - g^{j+1} - d^j \circ h^j) = (\delta^j)^*(f^{j+1} - g^{j+1}) - d^j \circ h^j \circ \delta^j = 0,$$

that is, $f^{j+1} - g^{j+1} - d^j \circ h^j \in \ker((\delta^j)^*) = \text{Im}((\delta^{j+1})^*)$ and so there is some $h^{j+1} \in \text{Hom}_{\mathcal{A}}(C^j, D^{j+1})$ such that $(\delta^{j+1})^*(h^{j+1}) = f^{j+1} - g^{j+1} - d^j \circ h^j$, that is, $f^n - g^n = d^{n-1} \circ h^{n-1} + h^n \circ \delta^n$ holds for all $n \leq j+1$, and the induction can continue. \square

As a consequence, we obtain that relative \mathcal{I} -injective resolutions are unique up to homotopy:

Corollary 7.6. *Let \mathcal{A} be an Abelian category and let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class. If $u: S^0(A) \rightarrow I^\bullet$ and $v: S^0(A) \rightarrow J^\bullet$ are both relative \mathcal{I} -injective resolutions of an object $A \in \mathcal{A}$, then there is a homotopy equivalence $f^\bullet: I^\bullet \rightarrow J^\bullet$ such that $f^\bullet \circ u = v$, and f^\bullet is unique up to homotopy.*

Proof. Use Lemma 7.5 to extend $\text{id}_A: A \rightarrow A$ to two morphisms $f^\bullet: I^\bullet \rightarrow J^\bullet$ and $g^\bullet: J^\bullet \rightarrow I^\bullet$, such that $f^\bullet \circ u = v$ and $g^\bullet \circ v = u$. The uniqueness of liftings up to homotopy forces $g^\bullet \circ f^\bullet$ and $f^\bullet \circ g^\bullet$ to be homotopic to id_{I^\bullet} and id_{J^\bullet} , respectively. \square

7.3. Generalities about the \mathcal{I} -injective model structure. Given an injective class \mathcal{I} in an Abelian category \mathcal{A} , one can introduce the following classes of morphisms in $\text{Ch}(\mathcal{A})$ that, under suitable hypotheses, make $\text{Ch}(\mathcal{A})$ into a model category:

Definition 7.7. *Let \mathcal{A} be an Abelian category and $\mathcal{I} \subseteq \mathcal{A}$ an injective class. We say that a morphism $\phi^\bullet: X^\bullet \rightarrow Y^\bullet$ is*

- an \mathcal{I} -**cofibration** if ϕ^n is an \mathcal{I} -monomorphism, for all $n \in \mathbb{Z}$;
- an \mathcal{I} -**weak equivalence** if $\text{Hom}(\phi^\bullet, I)$ is a quasi-isomorphism in $\text{Ch}(\text{Ab})$, for all $I \in \mathcal{I}$;
- an \mathcal{I} -**fibration** if it has the right lifting property with respect to the trivial \mathcal{I} -cofibrations.

We denote these classes of maps by $\mathcal{C}_{\mathcal{I}}$, $\mathcal{W}_{\mathcal{I}}$, and $\mathcal{F}_{\mathcal{I}}$, respectively. Furthermore, we say that a complex $X^\bullet \in \text{Ch}(\mathcal{A})$ is \mathcal{I} -**acyclic** provided $\text{Hom}(X^\bullet, I)$ is acyclic in $\text{Ch}(\text{Ab})$, for all $I \in \mathcal{I}$.

Remark 7.8. *We take advantage of this paper to correct an annoying misprint in the statement of [CNPS18, Theorem 2.3]: the \mathcal{I} -cofibrations in the relative model structure on $\text{Ch}_{\leq n}$ should be \mathcal{I} -monomorphisms in degrees $i < n$, and not $i \leq n$.*

In the following lemma, which is a consequence of [CH02, Proposition 2.5 and Lemma 2.7(b)], we collect some properties of \mathcal{I} -fibrations and of \mathcal{I} -fibrant objects:

Lemma 7.9. *Let \mathcal{A} be a bicomplete Abelian category and let \mathcal{I} be an injective class in \mathcal{A} . Then:*

- (1) *the \mathcal{I} -fibrations are precisely the degreewise split epimorphisms with \mathcal{I} -fibrant kernel;*
- (2) *a bounded below complex $X^\bullet \in \text{Ch}^+(\mathcal{A})$ is \mathcal{I} -fibrant, provided $X^i \in \mathcal{I}$, for all $i \in \mathbb{Z}$.*

More generally, let us recall the following important result by Christensen and Hovey (they actually state the dual result for projective classes, but it is an easy exercise to show that their statement is equivalent to the following one):

Theorem 7.10 ([CH02, Theorem 2.2]). *Let \mathcal{A} be a bicomplete Abelian category and $\mathcal{I} \subseteq \mathcal{A}$ an injective class such that, for each $X^\bullet \in \text{Ch}(\mathcal{A})$, there is an \mathcal{I} -fibrant replacement $\lambda^\bullet: X^\bullet \rightarrow F^\bullet$ (i.e., $\lambda^\bullet \in \mathcal{W}_{\mathcal{I}}$ and F^\bullet is \mathcal{I} -fibrant). Then, $(\text{Ch}(\mathcal{A}), \mathcal{W}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}})$ is a model category.*

In the following proposition (adapted from [CNPS18]) we establish the existence of ‘‘Spaltenstein towers of partial \mathcal{I} -fibrant replacements’’, in complete analogy to our discussion in Section 2 for the case $\mathcal{I} = \text{Inj}(\mathcal{A})$. This construction produces a standard candidate to \mathcal{I} -fibrant replacement for any given $X^\bullet \in \text{Ch}(\mathcal{A})$. In Subsection 7.4, we will discuss a sufficient condition to ensure that the construction below always produces an \mathcal{I} -fibrant replacement.

Proposition 7.11. *Let \mathcal{A} be a bicomplete Abelian category and let \mathcal{I} be an injective class in \mathcal{A} . Then, for each $X^\bullet \in \text{Ch}(\mathcal{A})$, there is an inverse system (a **tower**) of complexes*

$$(7.3) \quad \cdots \xrightarrow{t_2^\bullet} E_2^\bullet \xrightarrow{t_1^\bullet} E_1^\bullet \xrightarrow{t_0^\bullet} E_0^\bullet$$

that satisfies the following conditions, for all $n \in \mathbb{N}$:

- (1) *$E_n^\bullet \in \text{Ch}^{\geq -n}(\mathcal{A})$ and $E_n^\bullet \in \mathcal{I}$, for all $i \in \mathbb{Z}$ (so E_n^\bullet is \mathcal{I} -fibrant);*
- (2) *the morphism t_n^\bullet is an \mathcal{I} -fibration;*
- (3) *there is an \mathcal{I} -weak equivalence $\lambda_n^\bullet: \tau^{\geq -n}(X^\bullet) \rightarrow E_n^\bullet$ such that*

$$\lambda_n^\bullet \circ \pi_n^\bullet = t_n^\bullet \circ \lambda_{n+1}^\bullet, \quad \text{for all } n \in \mathbb{N},$$

where $\pi_n^\bullet: \tau^{\geq -n-1}(X^\bullet) \rightarrow \tau^{\geq -n}(X^\bullet)$ is the canonical projection.

Proof. By the results in Sections 4-5 in [CNPS18], there is model category $\text{Tow}(\mathcal{A}, \mathcal{I})$ of towers of complexes. Given $X^\bullet \in \text{Ch}(\mathcal{A})$, one can consider $\text{tow}(X^\bullet) \in \text{Tow}(\mathcal{A}, \mathcal{I})$, the tower of successive truncations of X^\bullet . A sequence (7.3), with the properties (1-3) in the statement, is precisely a fibrant replacement of $\text{tow}(X^\bullet)$ in the model category $\text{Tow}(\mathcal{A}, \mathcal{I})$. \square

7.4. The (Ab.4*)- k condition and the injective model structure relative to \mathcal{I} . Let \mathcal{A} be a bicomplete Abelian category, $X^\bullet \in \text{Ch}(\mathcal{A})$, $\mathcal{I} \subseteq \mathcal{A}$ an injective class, and define $E^\bullet := \varprojlim_{\mathbb{N}} E_n^\bullet$ as the inverse limit of the tower (7.3) described by Proposition 7.11. Then, $E^\bullet := \varprojlim_{\mathbb{N}} E_n^\bullet$ is \mathcal{I} -fibrant (by Remark 1.2) and, by condition (3) in the proposition, there is a canonical morphism $\lambda^\bullet: X^\bullet \rightarrow E^\bullet$. To determine whether λ^\bullet is an \mathcal{I} -weak equivalence (i.e., if E^\bullet is an \mathcal{I} -fibrant replacement for X^\bullet), it is useful to introduce a relative version of Roos' (Ab.4*)- k condition (see [CNPS18, Definition 6.1]). We do it after the following technical lemma:

Lemma 7.12. *Let \mathcal{A} be a complete Abelian category, let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class, and let $(u_\lambda: A_\lambda \rightarrow I_\lambda^\bullet)_\Lambda$ be a family of relative \mathcal{I} -injective resolutions of objects of \mathcal{A} . Then, the following assertions are equivalent for any $k \geq 0$:*

- (1) $H^{-n}(\text{Hom}(\prod_{\Lambda} I_\lambda^\bullet, I)) = 0$, for all $I \in \mathcal{I}$ and all $n > k$;
- (2) the induced map $\text{Coker}(\prod d_\lambda^{n-1}) \rightarrow \prod_{\Lambda} I_\lambda^{n+1}$ is an \mathcal{I} -monomorphism, for all $n > k$.

Proof. We set by convention $I_\lambda^{-1} := A_\lambda$, for all $\lambda \in \Lambda$. Condition (1) means that the following sequences are exact in Ab for all $I \in \mathcal{I}$ and all $n > k$:

$$(7.4) \quad \text{Hom}_{\mathcal{A}}(\prod_{\Lambda} I_\lambda^{n+1}, I) \xrightarrow{(\prod d_\lambda^n)^*} \text{Hom}_{\mathcal{A}}(\prod_{\Lambda} I_\lambda^n, I) \xrightarrow{(\prod d_\lambda^{n-1})^*} \text{Hom}_{\mathcal{A}}(\prod_{\Lambda} I_\lambda^{n-1}, I),$$

that is, we want the morphism $(\prod d_\lambda^n)^*: \text{Hom}_{\mathcal{A}}(\prod_{\Lambda} I_\lambda^{n+1}, I) \rightarrow \text{Ker}((\prod d_\lambda^{n-1})^*)$ to be surjective. As $\text{Hom}_{\mathcal{A}}(-, I)$ is left exact, there is an isomorphism $\text{Ker}((\prod d_\lambda^{n-1})^*) \cong \text{Hom}_{\mathcal{A}}(\text{Coker}(\prod d_\lambda^{n-1}), I)$. In particular, this shows that the sequences in (7.4) are exact for all $I \in \mathcal{I}$ if, and only if, $\text{Hom}_{\mathcal{A}}(\prod_{\Lambda} I_\lambda^{n+1}, I) \rightarrow \text{Hom}_{\mathcal{A}}(\text{Coker}(\prod d_\lambda^{n-1}), I)$ is surjective for all $I \in \mathcal{I}$, that is, if the morphism $\text{Coker}(\prod d_\lambda^{n-1}) \rightarrow \prod_{\Lambda} I_\lambda^{n+1}$ is an \mathcal{I} -monomorphism, which is condition (2). \square

Definition 7.13. *Let \mathcal{A} be a complete Abelian category, let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class, and let k be a natural number. Then, \mathcal{A} is said to be (Ab.4*)- \mathcal{I} - k when, for each family $(A_\lambda)_\Lambda$ of objects of \mathcal{A} , and for some (or, equivalently, any) choice of relative \mathcal{I} -injective resolutions $(I_\lambda^\bullet, u_\lambda: S^0(A_\lambda) \rightarrow I_\lambda^\bullet)_\Lambda$, the equivalent conditions (1) and (2) in Lemma 7.12 hold true.*

Observe that varying the chosen relative \mathcal{I} -injective resolutions does not affect the above definition. This is a consequence of Corollary 7.6, which implies that, if $(I_\lambda^\bullet, u_\lambda)_\Lambda$ and $(J_\lambda^\bullet, v_\lambda)_\Lambda$ are two families of relative \mathcal{I} -injective resolutions corresponding to $(A_\lambda)_\Lambda$, then there are homotopy equivalences $(f_\lambda: I_\lambda^\bullet \rightarrow J_\lambda^\bullet)_\Lambda$, whose product gives a homotopy equivalence $\prod_{\Lambda} f_\lambda: \prod_{\Lambda} I_\lambda^\bullet \rightarrow \prod_{\Lambda} J_\lambda^\bullet$, which then induces an homotopy equivalence between $\text{Hom}(\prod_{\Lambda} I_\lambda^\bullet, I)$ and $\text{Hom}(\prod_{\Lambda} J_\lambda^\bullet, I)$, for all $I \in \mathcal{I}$. In particular, these two complexes of Abelian groups have the same cohomology groups. Hence, condition (1) in Lemma 7.12 can be checked on either complex with the same result.

Corollary 7.14. *Let \mathcal{A} be a complete Abelian category, let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class, and consider the following assertions:*

- (1) each product of \mathcal{I} -preenvelopes is an \mathcal{I} -preenvelope;
- (2) each product of \mathcal{I} -monomorphisms is an \mathcal{I} -monomorphism;
- (3) \mathcal{A} is (Ab.4*)- $\mathcal{I} := (\text{Ab.4}^*)\text{-}\mathcal{I}\text{-0}$.

Then, the equivalence “(1) \Leftrightarrow (2)” holds in general, while “(1,2) \Rightarrow (3)” holds if \mathcal{A} is (Ab.4).*

Proof. As “(2) \Rightarrow (1)” is trivial, we just verify that “(1) \Rightarrow (2)”: let $(\iota_\lambda: X_\lambda \rightarrow Y_\lambda)_\Lambda$ be a family of \mathcal{I} -monomorphisms in \mathcal{A} . Choose, for each $\lambda \in \Lambda$, an \mathcal{I} -preenvelope $u_\lambda: X_\lambda \rightarrow I_\lambda$. As each ι_λ is an \mathcal{I} -monomorphism, there are morphisms $v_\lambda: Y_\lambda \rightarrow I_\lambda$ such that $v_\lambda \circ \iota_\lambda = u_\lambda$, for all $\lambda \in \Lambda$. By (1), the product map $\prod u_\lambda = (\prod v_\lambda) \circ (\prod \iota_\lambda)$ is an \mathcal{I} -preenvelope, and so also an \mathcal{I} -monomorphism. As a consequence, also $\prod \iota_\lambda$ is an \mathcal{I} -monomorphism, as desired.

For the implication “(2) \Rightarrow (3)” recall that, by Lemma 7.12, for \mathcal{A} to be (Ab.4*)- \mathcal{I} it is enough that, for any family $(X_\lambda)_\Lambda$ and a corresponding choice of relative \mathcal{I} -injective resolutions $(I_\lambda^\bullet, u_\lambda)_\Lambda$, the induced map $(\prod_{\Lambda} I_\lambda^n)/B^n(\prod_{\Lambda} I_\lambda^\bullet) \rightarrow \prod_{\Lambda} I_\lambda^{n+1}$ is an \mathcal{I} -monomorphism for all $n \geq -1$. Assuming that \mathcal{A} is (Ab.4*), there is an isomorphism $(\prod_{\Lambda} I_\lambda^n)/B^n(\prod_{\Lambda} I_\lambda^\bullet) \cong \prod_{\Lambda} (I_\lambda^n/B^n(I_\lambda^\bullet))$, so the condition becomes equivalent to the fact that $\prod_{\Lambda} (I_\lambda^n/B^n(I_\lambda^\bullet)) \rightarrow \prod_{\Lambda} I_\lambda^{n+1}$ is an \mathcal{I} -monomorphism, for all $n \geq -1$. This is then a consequence of (2), as the map in question is the product of the family of \mathcal{I} -monomorphisms $(I_\lambda^n/B^n(I_\lambda^\bullet) \rightarrow I_\lambda^{n+1})_\Lambda$. \square

The proof of the following theorem follows the same steps used in the proof of Theorem 2.9:

Theorem 7.15 ([CNPS18, Theorem 6.4]). *Let \mathcal{A} be a bicomplete Abelian category and $\mathcal{I} \subseteq \mathcal{A}$ an injective class such that \mathcal{A} is $(Ab.4^*)$ - \mathcal{I} - k for some $k \in \mathbb{N}$. Given $X^\bullet \in \text{Ch}(\mathcal{A})$ and a tower*

$$\cdots \xrightarrow{t_2^\bullet} E_2^\bullet \xrightarrow{t_1^\bullet} E_1^\bullet \xrightarrow{t_0^\bullet} E_0^\bullet$$

satisfying (1–3) in Proposition 7.11, the map $\lambda^\bullet: X^\bullet \rightarrow E^\bullet := \varprojlim_{\mathbb{N}} E_n^\bullet$ is an \mathcal{I} -weak equivalence.

In particular, in the setting of the above theorem (also using Proposition 7.11 and Remark 1.2), any complex $X^\bullet \in \text{Ch}(\mathcal{A})$ has an \mathcal{I} -fibrant replacement $\lambda^\bullet: X^\bullet \rightarrow E^\bullet$. Combining this result with Theorem 7.10, one immediately deduces that:

Corollary 7.16. *Let \mathcal{A} be a bicomplete Abelian category and let \mathcal{I} be an injective class in \mathcal{A} . If \mathcal{A} is $(Ab.4^*)$ - \mathcal{I} - k for some $k \in \mathbb{N}$, then $(\text{Ch}(\mathcal{A}), \mathcal{W}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}})$ is a model category.*

Observe that an advantage of Corollary 7.16 is that it applies to Abelian categories that are not necessarily Grothendieck, and where the small object argument may not be applicable.

8. EXAMPLES AND APPLICATIONS

8.1. On injective classes that are cogenerating. Let \mathcal{A} be an Abelian category. A full subcategory $\mathcal{I} \subseteq \mathcal{A}$ is called **cogenerating** if, for each $A \in \mathcal{A}$, there is a monomorphism $A \rightarrow I$, for some $I \in \mathcal{I}$.

Proposition 8.1. *The following are equivalent for an injective class \mathcal{I} in an Abelian category \mathcal{A} :*

- (1) \mathcal{I} is a cogenerating class in \mathcal{A} ;
- (2) \mathcal{I} -monomorphisms in \mathcal{A} are, in particular, also monomorphisms;
- (3) \mathcal{I} -acyclic complexes in $\text{Ch}(\mathcal{A})$ are, in particular, also acyclic;
- (4) \mathcal{I} -weak equivalences in $\text{Ch}(\mathcal{A})$ are, in particular, also quasi-isomorphisms.

Proof. (1) \Leftrightarrow (2). Given an \mathcal{I} -monomorphism $\phi: A \rightarrow B$, fix a monomorphism $u: A \rightarrow I \in \mathcal{I}$. The surjectivity of the map $\text{Hom}_{\mathcal{A}}(\phi, I): \text{Hom}_{\mathcal{A}}(B, I) \rightarrow \text{Hom}_{\mathcal{A}}(A, I)$ implies that $u = v \circ \phi$, for some $v \in \text{Hom}_{\mathcal{A}}(B, I)$ and, therefore, ϕ is a monomorphism. The converse is trivial.

(2) \Leftrightarrow (3). Given $X^\bullet = (\cdots \rightarrow X^n \xrightarrow{d^n} X^{n+1} \rightarrow \cdots) \in \text{Ch}(\mathcal{A})$, let $\bar{d}^n: X^n/B^n(X^\bullet) \rightarrow X^{n+1}$ be the morphism induced by the differential, so that $H^n(X^\bullet) = \text{Ker}(\bar{d}^n)$, for all $n \in \mathbb{Z}$. If X^\bullet is \mathcal{I} -acyclic then \bar{d}^n is an \mathcal{I} -monomorphism (by [CNPS18, Proposition 1.15(7)]) for all $n \in \mathbb{Z}$, and so, by (2), $H^n(X^\bullet) = \text{Ker}(\bar{d}^n) = 0$, for all $n \in \mathbb{Z}$. Conversely, a map $\psi: A \rightarrow B$ in \mathcal{A} is a(n) \mathcal{I} -monomorphism if and only if the complex $(\cdots \rightarrow 0 \rightarrow A \rightarrow B \rightarrow \text{Coker}(\psi) \rightarrow 0 \rightarrow \cdots)$ in $\text{Ch}^{\geq -1}(\mathcal{A}) \cap \text{Ch}^{\leq 1}(\mathcal{A})$ is \mathcal{I} -acyclic. Hence, also the implication “(3) \Rightarrow (2)” holds.

(3) \Leftrightarrow (4). It is well-known that a morphism $\phi^\bullet: A^\bullet \rightarrow B^\bullet$ in $\text{Ch}(\mathcal{A})$ is a quasi-isomorphism if and only if $\text{cone}(\phi^\bullet)$ is acyclic, while ϕ^\bullet is an \mathcal{I} -weak equivalence if and only if $\text{cone}(\phi^\bullet)$ is \mathcal{I} -acyclic (by [CNPS18, Proposition 1.15(2)]). These two characterizations imply the desired equivalence. \square

Remark 8.2. *Let \mathcal{A} be an $(Ab.4^*)$ Abelian category, and $(X_\lambda^\bullet)_\Lambda$ a family in $\text{Ch}(\mathcal{A})$, then there are isomorphisms $B^n(\prod_\Lambda X_\lambda^\bullet) \cong \prod_\Lambda B^n(X_\lambda^\bullet)$, for all $n \in \mathbb{Z}$. Take a cogenerating injective class $\mathcal{I} \subseteq \mathcal{A}$, and suppose that X_λ^\bullet is \mathcal{I} -acyclic, for all $\lambda \in \Lambda$. By Proposition 8.1, each X_λ^\bullet is also acyclic, and so $\text{Coker}(d_\lambda^{n-1}) = X_\lambda^n/B^n(X_\lambda^\bullet) = X_\lambda^n/Z^n(X_\lambda^\bullet) \cong B^{n+1}(X_\lambda^\bullet)$, for all $n \in \mathbb{Z}$. Combining these two observations we deduce that $\prod_\Lambda X_\lambda^\bullet$ is \mathcal{I} -acyclic if, and only if, the map $\prod_\Lambda B^n(X_\lambda^\bullet) \rightarrow \prod_\Lambda X_\lambda^n$ is an \mathcal{I} -monomorphism, for all $n \in \mathbb{Z}$.*

Using an analogous idea, one proves that an $(Ab.4^)$ Abelian category \mathcal{A} is $(Ab.4^*)$ - \mathcal{I} -d, with \mathcal{I} cogenerating, if and only if the following variation of the condition (2) in Lemma 7.12 holds:*

$$(2') \quad \prod_\Lambda B^{n+1}(I_\lambda^\bullet) \rightarrow \prod_\Lambda I_\lambda^{n+1} \text{ is an } \mathcal{I}\text{-monomorphism, for all } n > k;$$

that is, we want the product functor to preserve the \mathcal{I} -monomorphisms $(B^{n+1}(I_\lambda^\bullet) \rightarrow I_\lambda^{n+1})_\Lambda$.

Let $\mathcal{I} \subseteq \mathcal{A}$ be a cogenerating injective class. Then, a short exact sequence in \mathcal{A} :

$$0 \longrightarrow X \xrightarrow{\iota} Y \longrightarrow Z \longrightarrow 0$$

is called **\mathcal{I} -exact** if ι is an \mathcal{I} -monomorphism or, equivalently, if the following sequences are exact:

$$0 \longrightarrow \text{Hom}_{\mathcal{A}}(Z, I) \longrightarrow \text{Hom}_{\mathcal{A}}(Y, I) \xrightarrow{\iota^*} \text{Hom}_{\mathcal{A}}(X, I) \longrightarrow 0, \quad \text{for all } I \in \mathcal{I}.$$

We have the following variation of Corollary 7.14 for cogenerating injective classes:

Lemma 8.3. *Let \mathcal{A} be a complete Abelian category, and $\mathcal{I} \subseteq \mathcal{A}$ a cogenerating injective class. Suppose that any family of \mathcal{I} -exact sequences $(0 \rightarrow X_\lambda \rightarrow Y_\lambda \rightarrow Z_\lambda \rightarrow 0)_\Lambda$, gives a short exact sequence $0 \rightarrow \prod_\Lambda X_\lambda \rightarrow \prod_\Lambda Y_\lambda \rightarrow \prod_\Lambda Z_\lambda \rightarrow 0$, which is moreover \mathcal{I} -exact. Then, \mathcal{A} is (Ab.4*)- \mathcal{I} .*

Proof. Let $(X_\lambda)_\Lambda \subseteq \mathcal{A}$ and, for each $\lambda \in \Lambda$, consider the following relative \mathcal{I} -injective resolution:

$$(8.1) \quad 0 \longrightarrow X_\lambda \xrightarrow{u_\lambda} E_\lambda^0 \xrightarrow{d_\lambda^0} E_\lambda^1 \xrightarrow{d_\lambda^1} \cdots \xrightarrow{d_\lambda^{n-1}} E_\lambda^n \xrightarrow{d_\lambda^n} \cdots$$

with $E_\lambda^i \in \mathcal{I}$, for all $i \geq 0$. As \mathcal{I} is cogenerating, (8.1) is also an exact sequence. In particular, letting $B_\lambda^0 := X_\lambda$ and $B_\lambda^n := \text{Im}(d_\lambda^{n-1})$, for all $n \geq 1$, we have the following short exact sequences, for all $n \geq 0$ and $\lambda \in \Lambda$, that are also \mathcal{I} -exact:

$$0 \longrightarrow B_\lambda^n \longrightarrow E_\lambda^n \longrightarrow B_\lambda^{n+1} \longrightarrow 0.$$

By hypothesis, for each $n \geq 0$, the following short exact sequence is \mathcal{I} -exact:

$$0 \longrightarrow \prod_\Lambda B_\lambda^n \longrightarrow \prod_\Lambda E_\lambda^n \longrightarrow \prod_\Lambda B_\lambda^{n+1} \longrightarrow 0.$$

These conditions tell us that the following is a relative \mathcal{I} -injective resolution:

$$0 \longrightarrow \prod_\Lambda X_\lambda \xrightarrow{\prod u_\lambda} \prod_\Lambda E_\lambda^0 \xrightarrow{\prod d_\lambda^0} \prod_\Lambda E_\lambda^1 \xrightarrow{\prod d_\lambda^1} \cdots \xrightarrow{\prod d_\lambda^{n-1}} \prod_\Lambda E_\lambda^n \xrightarrow{\prod d_\lambda^n} \cdots,$$

showing that \mathcal{A} is (Ab.4*)- \mathcal{I} , as desired. \square

Let \mathcal{A} be an Abelian category with an injective class \mathcal{I} . As in [CNPS18], consider the localization of $\text{Ch}(\mathcal{A})$ with respect to $\mathcal{W}_\mathcal{I}$, the class of \mathcal{I} -weak equivalences, $\mathcal{D}(\mathcal{A}; \mathcal{I}) := \text{Ch}(\mathcal{A})[\mathcal{W}_\mathcal{I}^{-1}]$. In general this category could fail to be **locally small** (that is, the morphisms between two objects may form a proper class), while it is certainly locally small whenever $(\text{Ch}(\mathcal{A}), \mathcal{W}_\mathcal{I}, \mathcal{C}_\mathcal{I}, \mathcal{F}_\mathcal{I})$ is a model category (e.g., in the setting of Corollary 7.16) as, in that case, $\mathcal{D}(\mathcal{A}; \mathcal{I})$ is equivalent to the corresponding homotopy category. In what follows we denote by $\mathcal{K}(\mathcal{A})$ the homotopy category of \mathcal{A} , and by $\text{Ac}(\mathcal{A}) \subseteq \mathcal{K}(\mathcal{A})$ (resp., $\text{Ac}_\mathcal{I}(\mathcal{A}) \subseteq \mathcal{K}(\mathcal{A})$) its full subcategory of (\mathcal{I} -)acyclic complexes.

Corollary 8.4. *Let \mathcal{A} be an Abelian category and let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class. Then,*

- (1) *if $\mathcal{D}(\mathcal{A}; \mathcal{I})$ is locally small, it is a Verdier quotient $\mathcal{D}(\mathcal{A}; \mathcal{I}) \cong \mathcal{K}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})$;*
- (2) *if both $\mathcal{D}(\mathcal{A}; \mathcal{I})$ and $\mathcal{D}(\mathcal{A})$ are locally small and \mathcal{I} is cogenerating, then there is a canonical Verdier quotient functor $\mathcal{D}(\mathcal{A}; \mathcal{I}) \rightarrow \mathcal{D}(\mathcal{A})$.*

Proof. (1) follows by [CNPS18, Proposition 2.3 and Corollary 2.4].

(2) By Proposition 8.1, there are inclusions $\text{Ac}_\mathcal{I}(\mathcal{A}) \subseteq \text{Ac}(\mathcal{A}) \subseteq \mathcal{K}(\mathcal{A})$ of full triangulated subcategories. By [Ver96, Proposition 2.3.1 in Chap. II], we know that $\text{Ac}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})$ is a full triangulated subcategory of $\mathcal{K}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})$ and that there is a canonical triangulated functor

$$\pi_\mathcal{I}: \mathcal{D}(\mathcal{A}) \cong \mathcal{K}(\mathcal{A})/\text{Ac}(\mathcal{A}) \longrightarrow \frac{\mathcal{K}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})}{\text{Ac}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})} \cong \frac{\mathcal{D}(\mathcal{A}; \mathcal{I})}{\text{Ac}(\mathcal{A})/\text{Ac}_\mathcal{I}(\mathcal{A})}.$$

By Proposition 8.1, $\mathcal{W}_\mathcal{I} \subseteq \mathcal{W}$, so the canonical functor $p: \text{Ch}(\mathcal{A}) \rightarrow \mathcal{D}(\mathcal{A})$ sends \mathcal{I} -weak equivalences to isomorphisms. Therefore, p induces a unique functor $\mathcal{D}(\mathcal{A}; \mathcal{I}) = \text{Ch}(\mathcal{A})[\mathcal{W}_\mathcal{I}^{-1}] \rightarrow \mathcal{D}(\mathcal{A})$, which is a triangulated quasi-inverse to $\pi_\mathcal{I}$. \square

We now give an application to the derived category relative to a generator introduced in [Gil16]:

Example 8.5. *Given a complete Abelian category \mathcal{A} with a cogenerator E , we can consider the cogenerating injective class $\mathcal{I} = \text{Prod}(E)$ of all the summands of products of copies of E . In particular, take $\mathcal{A} = \mathcal{G}^{\text{op}}$ for a Grothendieck category \mathcal{G} , that is, \mathcal{A} is the category of strict complete topologically coherent left R -modules over some strict complete topologically left coherent and linearly compact ring R (see [Obe70]). Then, E is just a generator of \mathcal{G} , and the localization $\mathcal{D}(\mathcal{A}; \mathcal{I}) \cong \mathcal{D}_E(\mathcal{G})^{\text{op}}$ is the dual of the E -derived category defined in [Gil16]. Since $\mathcal{D}(\mathcal{G})^{\text{op}} \cong \mathcal{D}(\mathcal{A})$, Corollary 8.4 gives a Verdier quotient functor $\mathcal{D}_E(\mathcal{G}) \rightarrow \mathcal{D}(\mathcal{G})$, for any generator E of \mathcal{G} .*

Observe that, if $(\mathcal{X}, \mathcal{Y})$ is a right complete cotorsion pair in an Abelian category \mathcal{A} , then \mathcal{Y} is cogenerating (by the right completeness). Moreover, \mathcal{Y} is closed under summands in \mathcal{A} , and it is also pre-enveloping since any right $(\mathcal{X}, \mathcal{Y})$ approximation sequence is clearly \mathcal{Y} -exact. As we have seen in Subsection 6.2, the original proof of [YD15, Lemma 2.1] works if we assume that the ambient category is (Ab.4^*) . In fact, the conclusion of the lemma also holds for the injective classes \mathcal{Y} such that $({}^{\perp 1}\mathcal{Y}, \mathcal{Y})$ is a right complete cotorsion pair, provided \mathcal{A} is (Ab.4^*) - \mathcal{Y} - k , for some $k \in \mathbb{N}$:

Remark 8.6. *Let \mathcal{A} be a complete Abelian category and let $(\mathcal{X}, \mathcal{Y})$ be a right complete cotorsion pair in \mathcal{A} such that \mathcal{A} is (Ab.4^*) - \mathcal{Y} - k for some $k \in \mathbb{N}$. Given $X^\bullet \in \text{Ch}(\mathcal{A})$ one can find, as discussed in Subsection 7.4, a \mathcal{Y} -fibrant replacement $\lambda^\bullet: X^\bullet \rightarrow E^\bullet \in \text{dg}\mathcal{Y}$. As \mathcal{Y} is cogenerating, λ^\bullet is a quasi-isomorphism (by Proposition 8.1) and we get a short exact sequence:*

$$0 \longrightarrow \Sigma^{-1}E^\bullet \longrightarrow \Sigma^{-1}\text{cone}(\lambda^\bullet) \longrightarrow X^\bullet \longrightarrow 0,$$

with $\Sigma^{-1}E^\bullet \in \text{dg}\mathcal{Y}$ and $\Sigma^{-1}\text{cone}(\lambda^\bullet) \in \mathcal{E}$, proving [YD15, Lemma 2.1(1)] under these hypotheses.

8.2. Injective classes of injectives. Let \mathcal{A} be a complete Abelian category and $\mathcal{I} \subseteq \mathcal{A}$ an injective class. We have seen in Subsection 8.1 that, whenever \mathcal{I} is cogenerating, the classical derived category $\mathcal{D}(\mathcal{A})$ is a Verdier quotient of $\mathcal{D}(\mathcal{A}; \mathcal{I})$. In this subsection we concentrate on the injective classes \mathcal{I} of injectives in a Grothendieck category \mathcal{G} , that is, $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$. These injective classes have the property that $\mathcal{D}(\mathcal{G}; \mathcal{I})$ is always a Verdier quotient of $\mathcal{D}(\mathcal{G})$, so they are in some sense at the extreme opposite from the cogenerating ones; actually, the unique cogenerating injective class of injectives is $\text{Inj}(\mathcal{G})$, for which clearly $\mathcal{D}(\mathcal{G}; \text{Inj}(\mathcal{G})) \cong \mathcal{D}(\mathcal{G})$.

The key to understand the injective classes of injectives in a Grothendieck category \mathcal{G} is the observation that they correspond bijectively to the hereditary torsion pairs in \mathcal{G} (see [Vir17, Theorem 4.8]). More precisely, an injective class $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$ corresponds to the hereditary torsion pair $\tau_{\mathcal{I}} = (\mathcal{T} := {}^{\perp}\mathcal{I}, \mathcal{F} := ({}^{\perp}\mathcal{I})^{\perp})$, where \mathcal{F} can also be described as $\text{Cogen}(\mathcal{I})$, and $\mathcal{I} = \text{Inj}(\mathcal{G}) \cap \mathcal{F}$.

Lemma 8.7 ([Vir17, Lemmas 4.4 and 5.6]). *Let \mathcal{G} be a Grothendieck category, $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$ an injective class of injectives, and let $\tau_{\mathcal{I}} = (\mathcal{T}, \mathcal{F})$ be the associated hereditary torsion pair. Then*

- (1) *the following are equivalent for a morphism $\phi: X \rightarrow Y$ in \mathcal{G} :*
 - (1.1) *ϕ is an \mathcal{I} -monomorphism;*
 - (1.2) *the kernel of ϕ is $\tau_{\mathcal{I}}$ -torsion, that is, $\text{Ker}(\phi) \in \mathcal{T}$;*
- (2) *the following are equivalent for a morphism $\phi^\bullet: X^\bullet \rightarrow Y^\bullet$ in $\text{Ch}(\mathcal{G})$:*
 - (2.1) *ϕ^\bullet is an \mathcal{I} -weak equivalence;*
 - (2.2) *the cone $\text{cone}(\phi^\bullet)$ is \mathcal{I} -acyclic;*
 - (2.3) *the cone has $\tau_{\mathcal{I}}$ -torsion cohomologies, that is, $H^n(\text{cone}(\phi^\bullet)) \in \mathcal{T}$, for all $n \in \mathbb{Z}$.*

Consider the Gabriel quotient \mathcal{G}/\mathcal{T} , that comes with an adjunction $\mathbf{Q} \dashv \mathbf{S}: \mathcal{G} \rightleftarrows \mathcal{G}/\mathcal{T}$ such that:

- the right adjoint \mathbf{S} is fully faithful, so denoting by $\eta: \text{id}_{\mathcal{G}} \Rightarrow \mathbf{S} \circ \mathbf{Q}$ and $\varepsilon: \mathbf{Q} \circ \mathbf{S} \Rightarrow \text{id}_{\mathcal{G}/\mathcal{T}}$ the unit and the counit, both ε and $\mathbf{Q}(\eta)$ are natural isomorphisms. Abusing notation, we will often write $\mathbf{Q}(\mathbf{S}(Y)) = Y$ and $\mathbf{Q}(X) = \mathbf{Q}(\mathbf{S}(\mathbf{Q}(X)))$, if $Y \in \mathcal{G}/\mathcal{T}$ and $X \in \mathcal{G}$;
- the left adjoint \mathbf{Q} is an exact functor such that $\text{Ker}(\mathbf{Q}) = \mathcal{T}$, that is, $\mathbf{Q}(X) = 0$ if and only if $X \in \mathcal{T}$. In particular, given a morphism $\phi: X \rightarrow Y$ in \mathcal{G} , $\mathbf{Q}(\phi)$ is a monomorphism (resp., epimorphism) if, and only if, $\text{Ker}(\phi) \in \mathcal{T}$ (resp., $\text{Coker}(\phi) \in \mathcal{T}$). In particular, $\text{Ker}(\eta_X), \text{Coker}(\eta_X) \in \mathcal{T}$, for all $X \in \mathcal{G}$;
- as \mathbf{Q} is exact, \mathbf{S} sends injectives in \mathcal{G}/\mathcal{T} to injectives in \mathcal{G} . In fact, the adjunction $\mathbf{Q} \dashv \mathbf{S}$ restricts to an equivalence of categories $\text{Inj}(\mathcal{G}/\mathcal{T}) \cong \mathcal{F} \cap \text{Inj}(\mathcal{G}) = \mathcal{I}$.

With a slight abuse of notation, denote by $\mathbf{Q} \dashv \mathbf{S}: \text{Ch}(\mathcal{G}) \rightleftarrows \text{Ch}(\mathcal{G}/\mathcal{T})$ also the adjunction induced pointwise on complexes. It is easy to reformulate the above lemma as follows:

Corollary 8.8 ([Vir17, Lemmas 4.4 and 5.6]). *Let \mathcal{G} be a Grothendieck category, $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$ an injective class of injectives, and let $\tau_{\mathcal{I}} = (\mathcal{T}, \mathcal{F})$ be the associated hereditary torsion pair. Then*

- (1) *a morphism ϕ in \mathcal{G} is an \mathcal{I} -monomorphism if, and only, $\mathbf{Q}(\phi)$ is a monomorphism in \mathcal{G}/\mathcal{T} ;*
- (2) *a morphism ϕ^\bullet in $\text{Ch}(\mathcal{G})$ is an \mathcal{I} -weak equivalence if, and only if, $\mathbf{Q}(\phi^\bullet)$ is a quasi-isomorphism or, equivalently, if $\mathbf{Q}(\text{cone}(\phi^\bullet))$ is an acyclic complex.*

In particular, a morphism ϕ^\bullet in $\text{Ch}(\mathcal{G})$ is a (trivial) \mathcal{I} -cofibration if and only if $\mathbf{Q}(\phi^\bullet)$ is a (trivial) $\text{Inj}(\mathcal{G}/\mathcal{T})$ -cofibration in $\text{Ch}(\mathcal{G}/\mathcal{T})$.

The following is a more precise version of [Vir17, Theorem 5.7]:

Theorem 8.9. *Let \mathcal{G} be a Grothendieck category, $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$ an injective class of injectives, and let $\tau_{\mathcal{I}} = (\mathcal{T}, \mathcal{F})$ be the associated hereditary torsion pair. Then:*

- (1) ψ^\bullet is an $\text{Inj}(\mathcal{G}/\mathcal{T})$ -fibration in $\text{Ch}(\mathcal{G}/\mathcal{T})$, if and only if $\mathbf{S}(\psi^\bullet)$ is an \mathcal{I} -fibration;
- (2) $E^\bullet \in \text{Ch}(\mathcal{I}) \subseteq \text{Ch}(\mathcal{G})$ is \mathcal{I} -fibrant if, and only if, $\mathbf{Q}(E^\bullet)$ is DG-injective in $\text{Ch}(\mathcal{G}/\mathcal{T})$;
- (3) any complex $X^\bullet \in \text{Ch}(\mathcal{G})$ admits an \mathcal{I} -fibrant replacement $\lambda^\bullet: X^\bullet \rightarrow E^\bullet \in \text{Ch}(\mathcal{I})$.

In particular, $(\text{Ch}(\mathcal{G}), \mathcal{W}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}})$ is a model category, and the following is a Quillen equivalence

$$\mathbf{Q} \dashv \mathbf{S}: \text{Ch}(\mathcal{G}) \rightleftarrows \text{Ch}(\mathcal{G}/\mathcal{T}),$$

when $\text{Ch}(\mathcal{G}/\mathcal{T})$ is endowed with the usual injective model structure. As a consequence, we can describe the homotopy category $\mathcal{D}(\mathcal{G}; \mathcal{I}) := \text{Ch}(\mathcal{G})[\mathcal{W}_{\mathcal{I}}^{-1}]$ as a Verdier quotient of the usual derived category as follows: $\mathcal{D}(\mathcal{G}; \mathcal{I}) \cong \mathcal{D}(\mathcal{G}/\mathcal{T}) \cong \mathcal{D}(\mathcal{G})/\mathcal{D}_{\mathcal{T}}(\mathcal{G})$ (where $\mathcal{D}_{\mathcal{T}}(\mathcal{G})$ is the full, localizing, subcategory of those $X^\bullet \in \mathcal{D}(\mathcal{G})$ such that $H^n(X^\bullet) \in \mathcal{T}$, for all $n \in \mathbb{Z}$).

Proof. Fix a morphism $\psi^\bullet: Y_1^\bullet \rightarrow Y_2^\bullet$ in $\text{Ch}(\mathcal{G}/\mathcal{T})$ and let $\phi^\bullet: X_1^\bullet \rightarrow X_2^\bullet$ be a trivial \mathcal{I} -cofibration in $\text{Ch}(\mathcal{G})$. By Corollary 8.8 $\mathbf{Q}(\phi^\bullet)$ is a trivial $\text{Inj}(\mathcal{G}/\mathcal{T})$ -cofibration, and all the trivial $\text{Inj}(\mathcal{G}/\mathcal{T})$ -cofibrations in $\text{Ch}(\mathcal{G}/\mathcal{T})$ are of this form. Consider now the following commutative diagrams:

$$\begin{array}{ccc} X_1^\bullet & \xrightarrow{\alpha_1} & \mathbf{S}(Y_1^\bullet) \\ \phi^\bullet \downarrow & \nearrow & \downarrow \mathbf{S}(\psi^\bullet) \\ X_2^\bullet & \xrightarrow{\alpha_2} & \mathbf{S}(Y_2^\bullet) \end{array} \qquad \begin{array}{ccc} \mathbf{Q}(X_1)^\bullet & \xrightarrow{\beta_1} & Y_1^\bullet \\ \mathbf{Q}(\phi^\bullet) \downarrow & \nearrow & \downarrow \psi^\bullet \\ \mathbf{Q}(X_2)^\bullet & \xrightarrow{\beta_2} & Y_2^\bullet, \end{array}$$

where $\alpha_i^\bullet \in \text{Hom}_{\text{Ch}(\mathcal{G})}(X_i^\bullet, \mathbf{S}(Y_i^\bullet))$ and $\beta_i^\bullet \in \text{Hom}_{\text{Ch}(\mathcal{G}/\mathcal{T})}(\mathbf{Q}(X_i^\bullet), Y_i^\bullet)$ (for $i = 1, 2$) correspond to each other via the adjunction $\mathbf{Q} \dashv \mathbf{S}$. By adjunction, $\phi^\bullet \dashv \mathbf{S}(\psi^\bullet)$ if, and only if, $\mathbf{Q}(\phi^\bullet) \dashv \psi^\bullet$, proving (1). In particular, $E^\bullet \in \text{Ch}(\mathcal{I}) \subseteq \text{Ch}(\mathcal{G})$ is \mathcal{I} -fibrant if, and only if, $E^\bullet \rightarrow 0$ is an \mathcal{I} -fibration, if and only if $\mathbf{Q}(E^\bullet) \rightarrow 0$ is an $\text{Inj}(\mathcal{G}/\mathcal{T})$ -fibration, i.e., $\mathbf{Q}(E^\bullet) \in \text{Ch}(\text{Inj}(\mathcal{G}/\mathcal{T}))$ is a DG-injective complex, verifying (2). For (3), let $X^\bullet \in \text{Ch}(\mathcal{G})$ and consider a DG-injective resolution $\lambda^\bullet: \mathbf{Q}(X^\bullet) \rightarrow E^\bullet$ in $\text{Ch}(\mathcal{G}/\mathcal{T})$. We claim that the following composition is an \mathcal{I} -fibrant replacement

$$\mathbf{S}(\lambda^\bullet) \circ \eta_{X^\bullet}: X^\bullet \longrightarrow \mathbf{S}(\mathbf{Q}(X^\bullet)) \longrightarrow \mathbf{S}(E^\bullet).$$

Indeed, by Corollary 8.8, both the unit η_{X^\bullet} (which is sent to an isomorphism by \mathbf{Q}) and the map $\mathbf{S}(\lambda^\bullet)$ (which is sent by \mathbf{Q} to $\lambda^\bullet = \mathbf{Q}(\mathbf{S}(\lambda^\bullet))$, which is a quasi-isomorphism by construction) are \mathcal{I} -weak equivalences. Hence, the composition $\mathbf{S}(\lambda^\bullet) \circ \eta_{X^\bullet}$ is an \mathcal{I} -weak equivalence, while the complex $\mathbf{S}(E^\bullet) \in \text{Ch}(\mathcal{I})$ is \mathcal{I} -fibrant by part (2). By Theorem 7.10, this also shows that $(\text{Ch}(\mathcal{G}), \mathcal{W}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}})$ is a model category. By Corollary 8.8, \mathbf{Q} preserves both cofibrations and trivial cofibrations, so $\mathbf{Q} \dashv \mathbf{S}$ is a Quillen adjunction. Finally, to see that this is a Quillen equivalence, take $\phi^\bullet: X^\bullet \rightarrow \mathbf{S}(Y^\bullet)$ in $\text{Ch}(\mathcal{G})$, whose adjoint map is $\psi^\bullet := \mathbf{Q}(\phi^\bullet): \mathbf{Q}(X^\bullet) \rightarrow \mathbf{Q}(\mathbf{S}(Y^\bullet)) = Y^\bullet$; by Corollary 8.8, $\phi^\bullet \in \mathcal{W}_{\mathcal{I}}$ if and only if $\psi^\bullet \in \mathcal{W}_{\text{Inj}(\mathcal{G}/\mathcal{T})}$, concluding the proof. \square

Observe that, for any object $X \in \mathcal{G}$, a morphism $\lambda: S^0(X) \rightarrow I^\bullet \in \text{Ch}^{\geq 0}(\mathcal{I})$ is a relative \mathcal{I} -injective resolution if, and only if, $\mathbf{Q}(\lambda): S^0(\mathbf{Q}(X)) \rightarrow \mathbf{Q}(I^\bullet)$ is a DG-injective resolution of $\mathbf{Q}(X) \in \mathcal{G}/\mathcal{T}$. In particular, one can construct a resolution of a given $X \in \mathcal{G}$ as follows: first build a DG-injective resolution $S^0(\mathbf{Q}(X)) \rightarrow E^\bullet \in \text{Ch}^{\geq 0}(\text{Inj}(\mathcal{G}/\mathcal{T}))$ of $\mathbf{Q}(X)$ in \mathcal{G}/\mathcal{T} , and then note that the composition $S^0(X) \rightarrow S^0(\mathbf{S}(\mathbf{Q}(X))) \rightarrow \mathbf{S}(E^\bullet)$ is the desired \mathcal{I} -injective resolution. Recall also that, being \mathbf{S} a right adjoint, it commutes with all limits (and, in particular, with products). Thus, given $(Y_\lambda^\bullet)_\Lambda \subseteq \text{Ch}(\mathcal{G}/\mathcal{T})$, $\mathbf{S}(\prod_\Lambda Y_\lambda^\bullet) = \prod_\Lambda \mathbf{S}(Y_\lambda^\bullet)$. Exploiting these two ideas, we can show that the (Ab.4^*) - \mathcal{I} - k condition in \mathcal{G} is equivalent to the “absolute” (Ab.4^*) - k condition in \mathcal{G}/\mathcal{T} :

Proposition 8.10. *Let \mathcal{G} be a Grothendieck category, $\mathcal{I} \subseteq \text{Inj}(\mathcal{G})$ an injective class of injectives, and let $\tau_{\mathcal{I}} = (\mathcal{T}, \mathcal{F})$ be the associated hereditary torsion pair. Then, the following are equivalent:*

- (1) \mathcal{G} is (Ab.4^*) - \mathcal{I} - k ;
- (2) \mathcal{G}/\mathcal{T} is (Ab.4^*) - k (in the sense of Roos).

Proof. Let $(X_\lambda)_\Lambda \subseteq \mathcal{G}$ and, for each $\lambda \in \Lambda$, take a DG-injective resolution $\lambda_\lambda: S^0(\mathbf{Q}(X_\lambda)) \rightarrow E_\lambda^\bullet$ in $\text{Ch}^{\geq 0}(\text{Inj}(\mathcal{G}/\mathcal{T}))$, so that $\mu_\lambda: S^0(X_\lambda) \rightarrow \mathbf{S}(E_\lambda^\bullet) \in \text{Ch}^{\geq 0}(\mathcal{I})$ is the corresponding relative \mathcal{I} -injective

resolution, for all $\lambda \in \Lambda$, as discussed above. By Lemma 7.12(1), \mathcal{G} is (Ab.4*)- \mathcal{I} - k if, and only if:

$$(8.2) \quad H^{-n}(\mathrm{Hom}(\prod_{\Lambda} \mathbf{S}(E_{\lambda}^{\bullet}), I)) = 0, \quad \text{for all } n > k, \text{ and all } I \in \mathcal{I}.$$

Since $\mathcal{I} \subseteq \mathrm{Inj}(\mathcal{G})$ by hypothesis, the functor $\mathrm{Hom}_{\mathcal{G}}(-, I)$ is exact and, therefore, it commutes with cohomologies, so (8.2) becomes equivalent to:

$$\mathrm{Hom}_{\mathcal{G}}(H^n(\prod_{\Lambda} (\mathbf{S}(E_{\lambda}^{\bullet})), I) = 0, \quad \text{for all } n > k, \text{ and all } I \in \mathcal{I}.$$

As, by definition, $\mathcal{T} := {}^{\perp}\mathcal{I}$, the above condition means exactly that $H^n(\prod_{\Lambda} (\mathbf{S}(E_{\lambda}^{\bullet}))) \in \mathcal{T}$, for all $n > k$, that is, $\mathbf{Q}(H^n(\prod_{\Lambda} (\mathbf{S}(E_{\lambda}^{\bullet}))) = 0$, for all $n > k$. Using that \mathbf{Q} commutes with cohomologies, \mathbf{S} commutes with products, and $\mathbf{Q} \circ \mathbf{S} = \mathrm{id}_{\mathcal{G}/\mathcal{T}}$, condition (8.2) becomes equivalent to:

$$\prod_{\Lambda}^{(n)} \mathbf{Q}(X_{\lambda}) := H^n(\prod_{\Lambda} E_{\lambda}^{\bullet}) = H^n(\mathbf{Q} \circ \mathbf{S}(\prod_{\Lambda} E_{\lambda}^{\bullet})) = 0, \quad \text{for all } n > k.$$

This last equation is precisely the (Ab.4*)- k condition for the quotient category \mathcal{G}/\mathcal{T} . \square

8.3. The pure derived category of a locally finitely presented Grothendieck category.

Let \mathcal{G} be a Grothendieck category and recall that an object X in \mathcal{G} is said to be **finitely presented** if $\mathrm{Hom}_{\mathcal{G}}(X, -)$ commutes with directed colimits. We denote by $\mathrm{fp}(\mathcal{G})$ the full subcategory of finitely presented objects in \mathcal{G} , and we say that \mathcal{G} is **locally finitely presented** provided $\mathrm{fp}(\mathcal{G})$ is essentially small, and it generates \mathcal{G} . Observe that, whenever \mathcal{G} is locally finitely presented, the so-called **restricted Yoneda embedding** gives a fully faithful functor

$$\mathbf{y}: \mathcal{G} \rightarrow [\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}], \quad \text{such that } \mathbf{y}(X) = \mathbf{y}_X := \mathrm{Hom}_{\mathcal{G}}(-, X)|_{\mathrm{fp}(\mathcal{G})}.$$

Observe that \mathbf{y} is left exact, and that it commutes with products and directed colimits. The category $[\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}]$ is generated by $\mathbf{y}(\mathrm{fp}(\mathcal{G})) := \{\mathbf{y}_P : P \in \mathrm{fp}(\mathcal{G})\}$, the essentially small subcategory of the representable (=finitely presented projective) functors, and so $[\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}]$ is an (Ab.4*) Grothendieck category (it is equivalent to a category of modules over a ring with enough idempotents). Let us also recall that:

- $F: \mathrm{fp}(\mathcal{G})^{\mathrm{op}} \rightarrow \mathrm{Ab}$ is said to be **flat** if it belongs to the class $\mathcal{F} := \varinjlim \mathbf{y}(\mathrm{fp}(\mathcal{G}))$. The restricted Yoneda embedding induces an equivalence $\mathcal{G} \cong \mathbf{y}(\mathcal{G}) = \mathcal{F}$;
- the functors in the class $\mathcal{C} := \mathcal{F}^{\perp 1} = (\mathbf{y}(\mathrm{fp}(\mathcal{G})))^{\perp 1}$ are called **cotorsion**, and $(\mathcal{F}, \mathcal{C})$ is a complete cotorsion pair in $[\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}]$ (see [EEO04, Theorem 2.6]);
- a short exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ in \mathcal{G} is said to be **pure exact** if, and only if, $0 \rightarrow \mathbf{y}_X \rightarrow \mathbf{y}_Y \rightarrow \mathbf{y}_Z \rightarrow 0$ is exact in $[\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}]$. We say that $X \rightarrow Y$ is a **pure monomorphism** if it can be completed to a pure exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$;
- an object $E \in \mathcal{G}$ is said to be **pure injective** if it is injective with respect to all the pure exact sequences in \mathcal{G} , i.e., if $0 \rightarrow \mathrm{Hom}_{\mathcal{G}}(Z, E) \rightarrow \mathrm{Hom}_{\mathcal{G}}(Y, E) \rightarrow \mathrm{Hom}_{\mathcal{G}}(X, E) \rightarrow 0$ is exact in Ab , for any pure exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ in \mathcal{G} ;
- if we denote by $\mathrm{P.Inj}(\mathcal{G})$ the class of pure injectives in \mathcal{G} , the restricted Yoneda embedding induces an equivalence $\mathrm{P.Inj}(\mathcal{G}) \cong \mathbf{y}(\mathrm{P.Inj}(\mathcal{G})) = \mathcal{F} \cap \mathcal{C}$ (see [Her03, Lemma 3]).

Proposition 8.11. *Let \mathcal{G} be a locally finitely presented Grothendieck category, and $\mathcal{I} := \mathrm{P.Inj}(\mathcal{G})$:*

- (1) \mathcal{I} is an injective class in \mathcal{G} , and the \mathcal{I} -monomorphisms are the pure monomorphisms;
- (2) \mathcal{G} is (Ab.4*)- \mathcal{I} , and so $(\mathrm{Ch}(\mathcal{G}), \mathcal{W}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}})$ is a model category.

*In particular, the homotopy category $\mathcal{D}(\mathcal{G}; \mathcal{I}) := \mathrm{Ch}(\mathcal{G})[\mathcal{W}_{\mathcal{I}}^{-1}]$ is locally small, and it is equivalent to the so-called **pure derived category** $\mathcal{D}_{\mathrm{pure}}(\mathcal{G})$ introduced in [CH02, Section 5.3] when \mathcal{G} is a module category, and studied in [Kra12].*

Proof. Recall from [Her03, Theorem 6] that, for any $X \in \mathcal{G}$, there exists a pure exact sequence $0 \rightarrow X \rightarrow PE(X) \rightarrow PE(X)/X \rightarrow 0$ in \mathcal{G} , with $PE(X) \in \mathcal{I}$. In particular, \mathcal{I} is cogenerating.

(1) It is clear that, if $X \rightarrow Y$ is a pure monomorphism, then it is also an \mathcal{I} -monomorphism (by our definition of pure injective object). On the other hand, if $f: X \rightarrow Y$ is an \mathcal{I} -monomorphism, take a pure monomorphism $g: X \rightarrow PE(X)$, with $PE(X) \in \mathcal{I}$, and observe that (by definition of \mathcal{I} -monomorphism) $f^*: \mathrm{Hom}_{\mathcal{G}}(Y, PE(X)) \rightarrow \mathrm{Hom}_{\mathcal{G}}(X, PE(X))$ is surjective. In particular, there exists $\tilde{g}: Y \rightarrow PE(X)$, such that $\tilde{g} \circ f = f^*(\tilde{g}) = g$. In other words, $\mathbf{y}_g = \mathbf{y}_{\tilde{g}} \circ \mathbf{y}_f$ is a monomorphism in $[\mathrm{fp}(\mathcal{G})^{\mathrm{op}}, \mathrm{Ab}]$, and so \mathbf{y}_f is a monomorphism; i.e., f is a pure monomorphism.

(2) Consider a family of pure exact sequences $\{0 \rightarrow X_\lambda \rightarrow Y_\lambda \rightarrow Z_\lambda \rightarrow 0\}_\Lambda$ in \mathcal{G} , which gives a family of short exact sequences of functors $\{0 \rightarrow \mathbf{y}_{X_\lambda} \rightarrow \mathbf{y}_{Y_\lambda} \rightarrow \mathbf{y}_{Z_\lambda} \rightarrow 0\}_\Lambda$. Since $[\text{fp}(\mathcal{G})^{\text{op}}, \text{Ab}]$ is (Ab.4*), we obtain a short exact sequence $0 \rightarrow \prod_\Lambda \mathbf{y}_{X_\lambda} \rightarrow \prod_\Lambda \mathbf{y}_{Y_\lambda} \rightarrow \prod_\Lambda \mathbf{y}_{Z_\lambda} \rightarrow 0$. Moreover, since the restricted Yoneda embedding commutes with products, we obtain the following short exact sequence: $0 \rightarrow \mathbf{y}_{\prod_\Lambda X_\lambda} \rightarrow \mathbf{y}_{\prod_\Lambda Y_\lambda} \rightarrow \mathbf{y}_{\prod_\Lambda Z_\lambda} \rightarrow 0$. Hence, $0 \rightarrow \prod_\Lambda X_\lambda \rightarrow \prod_\Lambda Y_\lambda \rightarrow \prod_\Lambda Z_\lambda \rightarrow 0$ is pure exact (i.e., \mathcal{I} -exact) in \mathcal{G} . One can now conclude by Lemma 8.3 and Corollary 7.16.

For the last statements, it is enough to check that the \mathcal{I} -acyclic complexes are precisely the pure-acyclic complexes. As both the \mathcal{I} -acyclic and the pure-acyclic complexes are acyclic, this reduces to prove that a short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ in \mathcal{G} is pure if, and only if, it “is” an \mathcal{I} -acyclic complex which, in turn, is equivalent to prove that a monomorphism $u: A \rightarrow B$ in \mathcal{G} is pure if, and only if, it is an \mathcal{I} -monomorphism, that follows by (1). \square

8.4. Categories of finite \mathcal{I} -global dimension. A rich source of examples of (Ab.4*)- \mathcal{I} - k Abelian categories is derived from the following concepts (see Corollary 8.13 below):

Definition 8.12. *Let \mathcal{A} be a complete Abelian category and let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class. For each object A in \mathcal{A} , define the following set:*

$$\mathbb{N}(A, \mathcal{I}) := \{n \in \mathbb{N} : \text{there is a relative } \mathcal{I}\text{-injective resolution } A \rightarrow I^\bullet \in \text{Ch}^{\geq 0}(\mathcal{A}) \cap \text{Ch}^{\leq n}(\mathcal{A})\}.$$

Then, we define the \mathcal{I} -codimension of A as follows:

$$\mathcal{I}\text{-codim}(A) := \inf \mathbb{N}(A, \mathcal{I}),$$

with the convention that $\inf \emptyset := \infty$. Finally, we define the \mathcal{I} -global dimension of \mathcal{A} as:

$$\mathcal{I}\text{-gl.dim}(\mathcal{A}) := \sup\{\mathcal{I}\text{-codim}(A) : A \in \mathcal{A}\} \in \mathbb{N} \cup \{\infty\}.$$

We have the following immediate consequence.

Corollary 8.13. *Let \mathcal{A} be a complete Abelian category and let $\mathcal{I} \subseteq \mathcal{A}$ be an injective class such that $\mathcal{I}\text{-gl.dim}(\mathcal{A}) = d < \infty$. Then, \mathcal{A} is (Ab.4*)- \mathcal{I} - d .*

Proof. Let $(A_\lambda)_{\lambda \in \Lambda}$ be a family of objects in \mathcal{A} and choose, for each λ , a relative \mathcal{I} -injective resolution $u_\lambda: A_\lambda \rightarrow I_\lambda^\bullet \in \text{Ch}^{\geq 0}(\mathcal{A}) \cap \text{Ch}^{\leq d}(\mathcal{A})$. Observe then that $\prod_\Lambda I_\lambda^\bullet \in \text{Ch}^{\geq 0}(\mathcal{A}) \cap \text{Ch}^{\leq d}(\mathcal{A})$ and, therefore, $\text{Hom}(\prod_\Lambda I_\lambda^\bullet, I) \in \text{Ch}^{\leq 0}(\text{Ab}) \cap \text{Ch}^{\geq -d}(\text{Ab})$. In particular, $H^k(\text{Hom}(\prod_\Lambda I_\lambda^\bullet, I)) = 0$, for all $k < -d$, that is, \mathcal{A} is (Ab.4*)- \mathcal{I} - d . \square

We are now going to describe a few concrete examples of bicomplete Abelian categories \mathcal{A} that admit a suitable injective class $\mathcal{I} \subseteq \mathcal{A}$ for which $\mathcal{I}\text{-gl.dim}(\mathcal{A}) < \infty$. By the above corollary, these are also examples where \mathcal{A} is (Ab.4*)- \mathcal{I} - k , for some $k \in \mathbb{N}$. In particular, by Corollary 7.16, in each of these cases $(\text{Ch}(\mathcal{A}), \mathcal{W}_{\mathcal{I}}, \mathcal{C}_{\mathcal{I}}, \mathcal{F}_{\mathcal{I}})$ is a model category, and so the corresponding homotopy category $\mathcal{D}(\mathcal{A}; \mathcal{I}) := \text{Ch}(\mathcal{A})[\mathcal{W}_{\mathcal{I}}^{-1}]$ is locally small. The first two examples below have also appeared in [CNPS18, Section 9], with different terminology.

Example 8.14. *Let R be a d -Gorenstein ring, i.e., a (left and right) Noetherian ring that has injective dimension $= d$ on both sides, for a given $d \in \mathbb{N}$, and consider the following class:*

$$\mathcal{I} = \mathcal{GI}(R) := \{M \in \text{Mod-}R : M \text{ is Gorenstein injective}\} \subseteq \text{Mod-}R =: \mathcal{A}.$$

By [ELR02, Theorem 2.3], \mathcal{I} is an injective class in $\text{Mod-}R$, while it follows by [EJ11, Chapter 12] that $\mathcal{I}\text{-gl.dim}(\mathcal{A}) = d$.

Example 8.15. *Let R be a ring of pure global dimension $= d$, for some $d \in \mathbb{N}$ (e.g., if $d > 0$ and $|R| \leq \aleph_{d-1}$, then R has pure global dimension $\leq d$). Let $\mathcal{A} := \text{Mod-}R$, and consider the injective class $\mathcal{I} := \text{P.Inj}(\mathcal{A})$ of the pure-injective right R -modules. By definition (see [GJ81, Section 1]), the pure global dimension of R is the supremum of the pure-injective dimensions of its modules. Hence, $\mathcal{I}\text{-gl.dim}(\mathcal{A}) = d$ coincides with the pure global dimension of R .*

In a bicomplete Abelian category \mathcal{A} , a given $V \in \mathcal{A}$ is **1-tilting** ([PSV23, Definition 6.1]) if:

- $V^{\perp 1} = \text{Gen}(V) =: \mathcal{I}$;
- \mathcal{I} is cogenerating in \mathcal{A} .

We refer to [PSV23, Section 6] for several characterizations of 1-tilting objects and their connection with the traditional concept of a 1-tilting module. In the following example we show that, if V is 1-tilting, then $\text{Gen}(V)\text{-gl.dim}(\mathcal{A}) = 1$; for an extension to the n -tilting case, see Subsection 8.5.

Example 8.16. *Let \mathcal{A} be bicomplete Abelian category with a generator, $V \in \mathcal{A}$ a 1-tilting object, and let $\mathcal{I} := \text{Gen}(V) = V^{\perp 1}$. By [PSV23, Proposition 6.7], there is a generator G of \mathcal{A} together with a short exact sequence $0 \rightarrow G \xrightarrow{u} V_0 \rightarrow V_1 \rightarrow 0$ (with $V_0, V_1 \in \text{Add}(V)$), such that any coproduct of copies of the sequence remains exact. Now, given $A \in \mathcal{A}$, choose an epimorphism $\pi: G^{(S)} \rightarrow A$ and observe that $u^{(S)}: G^{(S)} \rightarrow V_0^{(S)}$ is an \mathcal{I} -preenvelope, since $\text{Ext}_{\mathcal{A}}^1(V_1^{(S)}, I) = 0$, for all $I \in \mathcal{I}$. Taking the push-out of $u^{(S)}$ along π , one gets a short exact sequence $0 \rightarrow A \xrightarrow{v} I_A \rightarrow V_1^{(S)} \rightarrow 0$, with $I_A \in \mathcal{I}$. By construction, v is clearly an \mathcal{I} -preenvelope, so $\mathcal{I}\text{-codim}(A) \leq 1$. Thus, $\mathcal{I}\text{-gl.dim}(\mathcal{A}) = 1$.*

Example 8.17. *Let V be a projective variety over a field K , i.e., a closed set for the Zariski topology of $\mathbf{P}^n(K)$, for some integer $n > 0$, let $\Gamma(V)$ be its (graded) ring of coordinates, and let $\mathcal{A} := \text{Gr-}\Gamma(V)$ be the category of graded $\Gamma(V)$ -modules. If V is regular of dimension d , let \mathcal{I} be the class of injective objects of $\text{Gr-}\Gamma(V)$ that have zero graded socle. Now, if $\mathcal{G} := \text{Qcoh}(V)$ is the Grothendieck category of quasi-coherent sheaves on V , when we view V as a scheme in the usual way, the regularity of V implies that its dimension coincides with the global dimension of \mathcal{G} , i.e., $\text{Inj}(\mathcal{G})\text{-gl.dim}(\mathcal{G})$ with our terminology. A well-known result of Serre (see Proposition 7.8 in Section 59, p. 252 in [Ser55]) says that there is an equivalence of categories $\mathcal{G} \cong (\text{Gr-}\Gamma(V))/\mathcal{T}$, where $\mathcal{T} \subseteq \text{Gr-}\Gamma(V)$ is the hereditary torsion class of the locally finite graded $\Gamma(V)$ -modules. This is exactly the torsion class generated by the class \mathcal{S} of graded-simple modules, and hence the corresponding torsion pair is cogenerated by $\mathcal{S}^{\perp} \cap \text{Inj}(\Gamma(V)) = \mathcal{I}$. By the discussion right before Proposition 8.10, it is now easy to see that $\mathcal{I}\text{-gl.dim}(\mathcal{A}) = d$.*

Proposition 8.18. *Let \mathcal{A} be an Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, and let $(\mathcal{X}, \mathcal{Y})$ be a right complete hereditary cotorsion pair. Then, the following are equivalent for any $n \geq 0$:*

- (1) $\text{Ext}_{\mathcal{A}}^k(X', X) = 0$, for all $X, X' \in \mathcal{X}$ and all $k > n$;
- (2) $\text{Ext}_{\mathcal{A}}^{n+1}(X', X) = 0$, for all $X, X' \in \mathcal{X}$;
- (3) for any $X \in \mathcal{X}$, there is an exact sequence of the form: $0 \rightarrow X \rightarrow W^0 \rightarrow \dots \rightarrow W^n \rightarrow 0$, with $W^i \in \mathcal{W} := \mathcal{X} \cap \mathcal{Y}$, for all $i = 0, \dots, n$.

Moreover, if $\mathcal{Y}\text{-gl.dim}(\mathcal{A}) \leq n$, then the above equivalent conditions are all verified.

Proof. (1) \Rightarrow (2) is clear.

(2) \Rightarrow (3). Let $X \in \mathcal{X}$ and consider a right $(\mathcal{X}, \mathcal{Y})$ -approximation $0 \rightarrow X \rightarrow W^0 \rightarrow X^1 \rightarrow 0$, where $W^0 \in \mathcal{Y}$, $X^1 \in \mathcal{X}$ and, since $\mathcal{X} = {}^{\perp 1}\mathcal{Y}$ is closed under extensions, also $W^0 \in \mathcal{X}$, so that $W^0 \in \mathcal{W}$. Continue by taking a right $(\mathcal{X}, \mathcal{Y})$ -approximation $0 \rightarrow X^1 \rightarrow W^1 \rightarrow X^2 \rightarrow 0$, with $W^1 \in \mathcal{W}$ and $X^2 \in \mathcal{X}$, and proceed inductively in this way to construct a \mathcal{W} -coresolution

$$(8.3) \quad 0 \longrightarrow X \xrightarrow{u} W^0 \xrightarrow{d^0} \dots \xrightarrow{d^{n-1}} W^n \xrightarrow{d^n} \dots,$$

with $\text{Im}(d^i) \in \mathcal{X}$, for all $i \geq 0$. Observe that, letting $X' := \text{Im}(d^n) \in \mathcal{X}$, any object $W \in \mathcal{W}$ is $\text{Hom}_{\mathcal{A}}(X', -)$ -acyclic (i.e., $\text{Ext}_{\mathcal{A}}^i(X', W) = 0$, for all $i > 0$), and so the \mathcal{W} -coresolution in (8.3) gives the following formula: $\text{Ext}_{\mathcal{A}}^i(X', X) \cong H^i(\text{Hom}_{\mathcal{A}}(X', W^{\bullet}))$, for all $i > 0$. By (2), we deduce that $H^{n+1}(\text{Hom}_{\mathcal{A}}(X', W^{\bullet})) = 0$ or, equivalently, that the map $\text{Hom}_{\mathcal{A}}(X', W^n) \rightarrow \text{Hom}_{\mathcal{A}}(X', X')$ is surjective, that is, $W^n \cong \text{Im}(d^{n-1}) \oplus X'$, showing that $\text{Im}(d^{n-1}) \in \mathcal{W}$, as this class is closed under summands. In particular, we obtain a \mathcal{W} -coresolution of the desired length:

$$0 \longrightarrow X \xrightarrow{u} W^0 \xrightarrow{d^0} \dots \xrightarrow{d^{n-2}} W^{n-1} \xrightarrow{d^{n-1}} \text{Im}(d^{n-1}) \longrightarrow 0 \longrightarrow \dots$$

(3) \Rightarrow (1). Given $X, X' \in \mathcal{X}$, take a \mathcal{W} -coresolution $0 \rightarrow X \rightarrow W^0 \rightarrow \dots \rightarrow W^n \rightarrow 0$ and observe that, as in the previous implication, $\text{Ext}_{\mathcal{A}}^i(X', X) \cong H^i(\text{Hom}_{\mathcal{A}}(X', W^{\bullet}))$, for all $i > 0$. Given $k > n$, $\text{Hom}_{\mathcal{A}}(X', W^k) = \text{Hom}_{\mathcal{A}}(X', 0) = 0$, and so $\text{Ext}_{\mathcal{A}}^k(X', X) \cong H^k(\text{Hom}_{\mathcal{A}}(X', W^{\bullet})) = 0$.

For the final statement, let $X, X' \in \mathcal{X}$. If $\mathcal{Y}\text{-gl.dim}(\mathcal{A}) \leq n$, take a relative \mathcal{Y} -injective resolution

$$(8.4) \quad 0 \longrightarrow X \xrightarrow{u} Y^0 \xrightarrow{d^0} \dots \xrightarrow{d^{n-1}} Y^n \longrightarrow 0 \longrightarrow \dots$$

Since \mathcal{Y} is cogenerating, the sequence (8.4) is exact while, as any $Y \in \mathcal{Y}$ is $\text{Hom}_{\mathcal{A}}(X', -)$ -acyclic, $\text{Ext}_{\mathcal{A}}^i(X', X) \cong H^i(\text{Hom}_{\mathcal{A}}(X', Y^{\bullet}))$, for all $i > 0$. This clearly implies (1). \square

As a direct consequence of Proposition 8.18, we get:

Corollary 8.19. *Let \mathcal{A} be an Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, let $(\mathcal{X}, \mathcal{Y})$ be a right complete hereditary cotorsion pair in \mathcal{A} , and fix $n \in \mathbb{N}$ such that the equivalent conditions of Proposition 8.18 are verified. Then, $\mathcal{Y}\text{-gl.dim}(\mathcal{A}) \leq n + 1$ and, therefore, \mathcal{A} is $(\text{Ab.4}^*)\text{-}\mathcal{Y}\text{-}(n + 1)$.*

Proof. Given $A \in \mathcal{A}$, take a right $(\mathcal{X}, \mathcal{Y})$ -approximation $0 \rightarrow A \rightarrow Y \rightarrow X \rightarrow 0$, with $Y \in \mathcal{Y}$ and $X \in \mathcal{X}$. Take now a \mathcal{W} -coresolution of X as in Proposition 8.18(3), that is, an exact sequence $0 \rightarrow X \rightarrow W^0 \rightarrow \dots \rightarrow W^n \rightarrow 0$, with $W^i \in \mathcal{W}$ for all $i = 0, \dots, n$. We obtain an exact sequence $0 \rightarrow A \rightarrow Y \rightarrow W^0 \rightarrow \dots \rightarrow W^n \rightarrow 0$, that is clearly a relative \mathcal{Y} -injective resolution of A . \square

8.5. n -Tilting cotorsion pairs. In this subsection we will work on an Abelian category \mathcal{A} that is (Ab.4) , and such that $\mathcal{D}(\mathcal{A})$ is locally small. In particular, $\mathcal{D}(\mathcal{A})$ has arbitrary coproducts, which are computed degreewise. These conditions are always verified in each of the following cases:

- if \mathcal{A} is a Grothendieck category;
- if \mathcal{A} is a bicomplete (Ab.4) Abelian category with enough projectives; this follows by [CNPS18, Theorem 6.4], or even Corollary 7.16, applied to \mathcal{A}^{op} with the injective class $\text{Inj}(\mathcal{A}^{\text{op}}) = (\text{Proj}(\mathcal{A}))^{\text{op}}$.

Recall that the **projective dimension** $\text{p.dim}_{\mathcal{A}}(X)$ of an object $X \in \mathcal{A}$ is defined as follows:

$$\text{p.dim}_{\mathcal{A}}(X) := \inf\{n \in \mathbb{N} : \text{Ext}_{\mathcal{A}}^k(T, A) = 0, \forall k > n, \forall A \in \mathcal{A}\},$$

with the convention that $\inf \emptyset = \infty$. The following definition comes from [NSZ19, Definition 6.8]:

Definition 8.20. *Let \mathcal{A} be an (Ab.4) Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, and let $n > 0$ be an integer. An object T of \mathcal{A} is called **n -tilting** when the following conditions hold:*

- (T.1) $\text{Ext}_{\mathcal{A}}^k(T, T^{(I)}) = 0$, for any set I and all $k > 0$;
- (T.2) $\text{p.dim}_{\mathcal{A}}(T) = n$;
- (T.3) there is a generating class $\mathcal{P} \subseteq \mathcal{A}$ such that, for each $P \in \mathcal{P}$, there is an exact sequence

$$0 \longrightarrow P \xrightarrow{u} T^0 \xrightarrow{d^0} \dots \xrightarrow{d^{n-1}} T^n \longrightarrow 0,$$

with $T^k \in \text{Add}(T)$, for all $k = 0, \dots, n$.

Any n -tilting object, has an associated cotorsion pair, that we describe in the following lemma:

Lemma 8.21. *Let \mathcal{A} be an (Ab.4) Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, let $T \in \mathcal{A}$ be n -tilting (for some $n \geq 1$), and let $(\mathcal{X}, \mathcal{Y}) := (\perp^1(T^{\perp > 0}), T^{\perp > 0})$ be the corresponding **n -tilting cotorsion pair**. Then, the following statements hold true:*

- (1) $(\mathcal{X}, \mathcal{Y})$ is a hereditary cotorsion pair in \mathcal{A} ;
- (2) $\mathcal{W} := \mathcal{X} \cap \mathcal{Y} = \text{Add}(T)$;
- (3) $(\mathcal{X}, \mathcal{Y})$ is right complete and, therefore, \mathcal{Y} is a cogenerating injective class;
- (4) $\mathcal{Y}\text{-gl.dim}(\mathcal{A}) \leq n$ and, therefore, \mathcal{A} satisfies the $(\text{Ab.4}^*)\text{-}\mathcal{Y}\text{-}n$ condition;
- (5) the class \mathcal{X} is generating in \mathcal{A} .

Proof. (1) follows immediately by the definition of n -tilting object.

(2). The inclusion “ $\text{Add}(T) \subseteq \mathcal{W}$ ” is clear. On the other hand, given $W \in \mathcal{W} \subseteq \mathcal{Y} \subseteq \text{Gen}(T)$ (see [NSZ19, Theorem 6.3 and Remark 6.2] and their proofs), if we denote by $\varepsilon_f : T \rightarrow T^{(\text{Hom}_{\mathcal{A}}(T, W))}$ the inclusion in the coproduct of the copy of T corresponding to the morphism $f \in \text{Hom}_{\mathcal{A}}(T, W)$, we get a canonical morphism $p_W : T^{(\text{Hom}_{\mathcal{A}}(T, W))} \rightarrow W$ such that $p_W \circ \varepsilon_f = f$, for all $f \in \text{Hom}_{\mathcal{A}}(T, W)$. The condition $W \in \text{Gen}(T)$ can be expressed equivalently by saying that p_W is an epimorphism; let $W' := \ker(p_W)$. The condition $f = p_W \circ \varepsilon_f =: p_W^*(\varepsilon_f)$ for all $f : T \rightarrow W$, shows the surjectivity of $p_W^* : \text{Hom}_{\mathcal{A}}(T, T^{(\text{Hom}_{\mathcal{A}}(T, W))}) \rightarrow \text{Hom}_{\mathcal{A}}(T, W)$, so $\text{Ext}_{\mathcal{A}}^1(T, W') = 0$ by the exact sequence below:

$$\text{Hom}_{\mathcal{A}}(T, T^{(\text{Hom}_{\mathcal{A}}(T, W))}) \xrightarrow{p_W^*} \text{Hom}_{\mathcal{A}}(T, W) \rightarrow \text{Ext}_{\mathcal{A}}^1(T, W') \rightarrow \text{Ext}_{\mathcal{A}}^1(T, T^{(\text{Hom}_{\mathcal{A}}(T, W))}) = 0,$$

where the last term on the right is $= 0$ by condition (T.1). Furthermore, $\text{Ext}_{\mathcal{A}}^k(T, W') = 0$, for all $k \geq 2$, as shown by the following exact sequence:

$$0 = \text{Ext}_{\mathcal{A}}^{k-1}(T, W) \rightarrow \text{Ext}_{\mathcal{A}}^k(T, W') \rightarrow \text{Ext}_{\mathcal{A}}^k(T, T^{(\text{Hom}_{\mathcal{A}}(T, W))}) = 0.$$

Thus, we have shown that $W' \in T^{\perp > 0} = \mathcal{Y}$. Moreover, $W' \in \mathcal{X}$, since \mathcal{X} is closed under kernels of epimorphisms, and so $W' \in \mathcal{W}$. As a consequence, $\text{Ext}_{\mathcal{A}}^1(W, W') = 0$, and so the exact sequence $0 \rightarrow W' \rightarrow T^{\text{Hom}_{\mathcal{A}}(T, W)} \rightarrow W \rightarrow 0$ has to split. In particular, $W \in \text{Add}(T)$ as desired.

(3)-(4). By [NSZ19, Lemma 6.13], given $A \in \mathcal{A}$, there is an exact sequence of the form:

$$(8.5) \quad 0 \longrightarrow A \xrightarrow{u} Y \xrightarrow{d^0} T^1 \xrightarrow{d^1} \dots \xrightarrow{d^{n-1}} T^n \longrightarrow 0,$$

such that $Y \in \mathcal{Y}$ and $T^k \in \text{Add}(T) = \mathcal{W}$, for all $k = 1, \dots, n$. Since \mathcal{X} is closed under kernels of epimorphisms, $X^{i-1} := \text{Ker}(d^{i-1}) \in \mathcal{X}$, for all $i = 2, \dots, n$. In particular, $\text{Coker}(u) \in \mathcal{X}$, showing that $0 \rightarrow A \rightarrow Y \rightarrow \text{Coker}(u) \rightarrow 0$ is a right $(\mathcal{X}, \mathcal{Y})$ -approximation of A , and concluding the proof of (3). Moreover, observe that $0 \rightarrow X^{i-1} \rightarrow T^{i-1} \rightarrow X^i \rightarrow 0$ is also a right $(\mathcal{X}, \mathcal{Y})$ -approximation (of X^{i-1}), for all $i = 2, \dots, n$ and, therefore, the exact sequence in (8.5) is a relative \mathcal{Y} -injective resolution of A , showing that $\mathcal{Y}\text{-codim}(A) \leq n$, as desired.

(5). Given $A \in \mathcal{A}$, we can take a right $(\mathcal{X}, \mathcal{Y})$ -approximation $0 \rightarrow A \rightarrow Y \rightarrow X \rightarrow 0$ of A , which is a short exact sequence with $Y \in \mathcal{Y}$, and $X \in \mathcal{X}$. Now, as at the beginning of the proof of part (2) above, we observe that, by results of [NSZ19], there is an inclusion $\mathcal{Y} \subseteq \text{Gen}(T)$. As a consequence, there is a canonical epimorphism $W \rightarrow Y \rightarrow 0$, where $W := T^{\text{Hom}_{\mathcal{A}}(T, Y)} \in \text{Add}(T) = \mathcal{W} \subseteq \mathcal{X}$. Consider now the following pullback diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & P & \longrightarrow & W & \longrightarrow & X & \longrightarrow & 0 \\ & & \downarrow & \text{P.B.} & \downarrow & & \parallel & & \\ 0 & \longrightarrow & A & \longrightarrow & Y & \longrightarrow & X & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & & & \\ & & 0 & & 0 & & & & \end{array}$$

We deduce that A is a quotient of P , which belongs to \mathcal{X} , since $(\mathcal{X}, \mathcal{Y})$ is hereditary. \square

In the following theorem we give a characterization of the tilting cotorsion pairs:

Theorem 8.22. *Let \mathcal{A} be an (Ab.4) Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, and let $(\mathcal{X}, \mathcal{Y})$ be a right complete, hereditary cotorsion pair, with \mathcal{X} generating in \mathcal{A} . If there is $W \in \mathcal{A}$ such that $\text{Add}(W) = \mathcal{X} \cap \mathcal{Y} =: \mathcal{W}$, and $\text{p.dim}_{\mathcal{A}}(W) = n \leq m \in \mathbb{N}$, then the following are equivalent:*

- (1) $\mathcal{Y}\text{-gl.dim}(\mathcal{A}) \leq m$;
- (2) $\text{Ext}_{\mathcal{A}}^{m+1}(X', X) = 0$, for all $X, X' \in \mathcal{X}$;
- (3) there is a subcategory $\mathcal{P} \subseteq \mathcal{X}$ which is generating in \mathcal{A} and such that, for each $P \in \mathcal{P}$, there is an exact sequence $0 \rightarrow P \rightarrow W^0 \rightarrow \dots \rightarrow W^m \rightarrow 0$, with $W^i \in \mathcal{W}$, for all $i = 0, \dots, m$.

If these conditions hold, then W is n -tilting, and $(\mathcal{X}, \mathcal{Y})$ is its associated n -tilting cotorsion pair.

Proof. The implications “(1) \Rightarrow (2)” and “(2) \Rightarrow (3)” follow by Proposition 8.18, while the implication “(3) \Rightarrow (1)” will follow, by Lemma 8.21(4), as a consequence of the last part of the statement. Thus, to conclude our proof, it is enough to show that (3) implies that W is n -tilting. In fact, W satisfies the condition (T.1) (see Definition 8.20), since $(\mathcal{X}, \mathcal{Y})$ is hereditary and $W \in \mathcal{X} \cap \mathcal{W}$, while (T.2) is given by hypothesis, so we just need to take care of the axiom (T.3). We proceed by induction on $m - n \geq 0$. Indeed, if $m = n$, then clearly (3) implies (T.3), so W is n -tilting. On the other hand, if $m > n$ and given $P \in \mathcal{P}$, fix the following exact sequence as in (3):

$$0 \longrightarrow P \xrightarrow{u} W^0 \xrightarrow{d^0} W^1 \xrightarrow{d^1} \dots \xrightarrow{d^{m-1}} W^m \longrightarrow 0,$$

Using again and again the fact that \mathcal{X} is closed under taking kernels of epimorphisms, one gets that $X^k := \text{Im}(d^k) \in \mathcal{X}$, for $k = -1, 0, \dots, m$, where $X^{-1} := P$. Hence, the induced exact sequence $0 \rightarrow X^{m-n-2} \rightarrow W^{m-n-1} \rightarrow \dots \rightarrow W^m \rightarrow 0$ yields an element of $\text{Ext}_{\mathcal{A}}^{n+1}(W^m, X^{m-n-2}) = 0$. Since \mathcal{W} consists of $\text{Hom}_{\mathcal{A}}(W^m, -)$ -acyclic objects, an argument similar to the one in the proof of implication “(2) \Rightarrow (3)” of Proposition 8.18 shows that $d^{m-1}: W^{m-1} \rightarrow W^m$ is a retraction, so that $X^{m-2} \in \mathcal{W}$. Letting $\overline{W}^k := W^k$ for $k = 0, 1, \dots, m-2$ and $\overline{W}^{m-1} := X^{m-2}$, we get an exact sequence $0 \rightarrow P \rightarrow \overline{W}^0 \rightarrow \dots \rightarrow \overline{W}^{m-1} \rightarrow 0$, so we can conclude by inductive hypothesis. \square

Example 8.23. *Let \mathcal{A} be a locally Noetherian Grothendieck category, and suppose that*

$$\sup\{\text{p.dim}_{\mathcal{A}}(I) : I \text{ is an indecomposable injective}\} = n > 0,$$

and that there is a set \mathcal{S} of generators, each of which has injective dimension $\leq n$. A coproduct W of all the indecomposable injectives, one for each isoclass, is an n -tilting object. Indeed, as $\text{Inj}(\mathcal{A})$ is closed under coproducts, one readily checks that conditions (T.1–3) of Definition 8.20 hold. In particular, if $\mathcal{Y} := W^{\perp > 0} = (\text{Inj}(\mathcal{A}))^{\perp > 0}$, Corollary 8.13 and Lemma 8.21 tell us that \mathcal{A} is $(\text{Ab.}\mathcal{A}^*)\text{-}\mathcal{Y}\text{-}n$.

Observe that any locally Noetherian Grothendieck category \mathcal{A} whose homological dimension is n (i.e., $\text{Ext}_{\mathcal{A}}^n(-, -) \neq 0 = \text{Ext}_{\mathcal{A}}^{n+1}(-, -)$) satisfies the conditions of the above example. In particular, we get the following case, where we can completely describe the classes \mathcal{X} and \mathcal{Y} :

Example 8.24. Let R be a n -Gorenstein ring, $\mathcal{A} := \text{Mod-}R$ and define $(\mathcal{X}, \mathcal{Y})$ as in Example 8.23. Then, the subcategory \mathcal{Y} is precisely that of the Gorenstein injectives, and $\mathcal{X} := {}^{\perp 1}\mathcal{Y}$ is the subcategory of modules of finite injective (=finite projective) dimension (see [GT12, Example 8.13]).

Remark 8.25. If \mathcal{A} is an $(\text{Ab.}\mathcal{A}^*)$ Abelian category such that $\mathcal{D}(\mathcal{A})$ is locally small, then \mathcal{A}^{op} is a category that satisfies the general hypotheses listed at the beginning of this subsection. This allows one to define an $(\text{Ab.}\mathcal{A})\text{-}\mathcal{X}\text{-}n$ condition and to relate it to n -cotilting objects and n -cotilting cotorsion pairs.

REFERENCES

- [CH02] J. Daniel Christensen and Mark Hovey. Quillen model structures for relative homological algebra. *Math. Proc. Camb. Philos. Soc.*, 133(2):261–293, 2002.
- [CNPS18] Wojciech Chachólski, Amnon Neeman, Wolfgang Pitsch, and Jérôme Scherer. Relative homological algebra via truncations. *Doc. Math.*, 23:895–937, 2018.
- [EEO04] J. R. García Rozas Edgar Enochs, S. Estrada and L. Oyonarte. Flat covers in the category of quasi-coherent sheaves over the projective line. *Communications in Algebra*, 32(4):1497–1508, 2004.
- [EJ11] E.E. Enochs and O.M.G. Jenda. *Relative Homological Algebra: Volume 1*. De Gruyter Expositions in Mathematics. De Gruyter, 2011.
- [ELR02] Edgar E. Enochs and J. A. López-Ramos. Kaplansky classes. *Rendiconti del Seminario Matematico della Università di Padova*, 107:67–79, 2002.
- [Gil06] James Gillespie. The flat model structure on complexes of sheaves. *Transactions of the American Mathematical Society*, 358(7):2855–2874, 2006.
- [Gil08] James Gillespie. Cotorsion pairs and degreewise homological model structures. *Homology, Homotopy and Applications*, 10(1):283–304, 2008.
- [Gil16] James Gillespie. The derived category with respect to a generator. *Annali di Matematica Pura ed Applicata (1923 -)*, 195(2):371–402, 2016.
- [GJ81] L. Gruson and C. U. Jensen. *Dimensions cohomologiques reliées aux foncteurs (i)*, volume 867, pages 234–294. Springer, Schweiz, 1981.
- [GT12] Rüdiger Göbel and Jan Trlifaj. *Approximations and endomorphism algebras of modules. Volume 2*, volume 41 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, extended edition, 2012. Predictions.
- [Her03] Ivo Herzog. Pure-injective envelopes. *Journal of Algebra and Its Applications*, 02(04):397–402, 2003.
- [Hov01] Mark Hovey. Model category structures on chain complexes of sheaves. *Transactions of the American Mathematical Society*, 353(6):2441–2457, 2001.
- [Hov02] Mark Hovey. Cotorsion pairs, model category structures, and representation theory. *Math. Z.*, 241(3):553–592, 2002.
- [Joy84] André Joyal. Lettre d’André Joyal à Alexandre Grothendieck. Éditée par Georges Maltsiniotis, 1984.
- [Kra12] Henning Krause. Approximations and adjoints in homotopy categories. *Mathematische Annalen*, 353(3):765–781, 2012.
- [Mat58] Eben Matlis. Injective modules over Noetherian rings. *Pacific J. Math.*, 8:511–528, 1958.
- [Nag62] Masayoshi Nagata. Local rings. New York and London: Interscience Publishers, a division of John Wiley and Sons. XIII, 234 pages, 1962.
- [NSZ19] Pedro Nicolás, Manuel Saorín, and Alexandra Zvonareva. Silting theory in triangulated categories with coproducts. *Journal of Pure and Applied Algebra*, 223(6):2273–2319, 2019.
- [Obe70] Ulrich Oberst. Duality theory for grothendieck categories and linearly compact rings. *Journal of Algebra*, 15(4):473–542, 1970.
- [PSV23] Carlos E. Parra, Manuel Saorín, and Simone Virili. Tilting preenvelopes and cotilting precovers in general abelian categories. *Algebras and Representation Theory*, 26(4):1087–1140, 2023.
- [Rie14] Emily Riehl. *Categorical homotopy theory*, volume 24. Cambridge University Press, 2014.
- [Roo06] Jan-Erik Roos. Derived functors of inverse limits revisited. *J. Lond. Math. Soc., II. Ser.*, 73(1):65–83, 2006.
- [San07] Samson Sanzblidze. On derived categories and derived functors. *arXiv preprint arXiv:0710.5065*, 2007.
- [Ser55] J. P. Serre. Faisceaux algébriques cohérents. *Ann. of Math.*, 61(2):197–278, 1955.
- [Spa88] N. Spaltenstein. Resolutions of unbounded complexes. *Compos. Math.*, 65(2):121–154, 1988.

- [Ste75] Bo Stenström. Rings of quotients. An introduction to methods of ring theory. Die Grundlehren der mathematischen Wissenschaften. Band 217. Berlin-Heidelberg-New York: Springer-Verlag. VIII, 1975.
- [Ver96] Jean-Louis Verdier. Des categories derivees des categories abeliennes. *Astérisque*, 239, 1996.
- [Vir17] Simone Virili. On the exactness of products in the localization of (ab. 4*) grothendieck categories. *Journal of Algebra*, 470:37–67, 2017.
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [YD15] Xiaoyan Yang and Nanqing Ding. On a question of Gillespie. *Forum Mathematicum*, 27(6):3205–3231, 2015.

DEPT. DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA (08193) BELLATERRA, BARCELONA, SPAIN,
CENTRE DE RECERCA MATEMÀTICA (08193) BELLATERRA, BARCELONA, SPAIN.
Email address: `Dolors.Herbera@uab.cat`

DEPT. DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA (08193) BELLATERRA, BARCELONA, SPAIN.
Email address: `Wolfgang.Pitsch@uab.cat`

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, AVDA. TENIENTE FLORESTA 5 (30003) MURCIA, SPAIN.
Email address: `MSaorinc@um.es`

DEPT. DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA (08193) BELLATERRA, BARCELONA, SPAIN.
Email address: `Simone.Virili@uab.cat` or `Virili.Simone@gmail.com`