

Vitalii Shpakivskyi

## Construction of an infinite-dimensional family of exact solutions of a three-dimensional biharmonic equation by the hypercomplex method

An infinite-dimensional family of exact solutions of a three-dimensional biharmonic equation was constructed by the hypercomplex method.

### Preface

In 2019, at the 12th ISAAC Congress (Aveiro, Portugal), I gave a report on the topic: Hypercomplex method for solving linear PDEs. After the report, Professor Yuri Grigoriev approached me and asked: can I write out the solutions of the three-dimensional biharmonic equation? I answered: yes, I can. Then he asked me to write a separate paper on the solutions of the three-dimensional biharmonic equation and give him a copy. I said: ok. But only now I am fulfilling my promise. It is a pity that Professor Yuri Grigoriev did not read it in his lifetime. Only now did I understand that he had been studying the spatial theory of elasticity throughout his life, in particular, the three-dimensional biharmonic equation...

### 1 Introduction

Consider the three-dimensional biharmonic equation

$$\Delta^2 U(x, y, z) := \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right)^2 U(x, y, z) = 0 \quad (1)$$

or

$$\frac{\partial^4 U}{\partial x^4} + \frac{\partial^4 U}{\partial y^4} + \frac{\partial^4 U}{\partial z^4} + 2 \frac{\partial^4 U}{\partial x^2 \partial y^2} + 2 \frac{\partial^4 U}{\partial x^2 \partial z^2} + 2 \frac{\partial^4 U}{\partial y^2 \partial z^2} = 0.$$

Not many authors have tried to solve the three-dimensional biharmonic equation. The reason, of necessity, is that three-dimensional problems require large computing power. Such computing power has only recently begun to become available for research.

However, in paper [1], Yu. Grigoriev considers equation (1) in star domains relative to the origin of coordinates. In such domains, he gives a representation of the general solution of equation (1) in spherical coordinates. We will also note

the paper [2] in which the theory of regular quaternion functions systematically were used for a solution of a three-dimensional problem of the elasticity theory.

In the paper [3], we are concerned with energy decay estimates for solutions of the biharmonic equation in a semi-infinite cylinder with nonzero boundary conditions on the finite end. In the work [4], it is considered several finite-difference approximations for the three-dimensional biharmonic equation.

In this paper we construct an infinite-dimensional family of exact solutions of equation (1) via holomorphic functions of a complex variable.

## 2 The method

This method originates from the works of P. W. Ketchum [5, 6, 7], and developed independently in the works of I. P. Mel'nichenko [8, 9], S. A. Plaksa [10] and ours. The final formation of the method was done in our papers [11, 12, 13, 14]. This method is systematically described in monograph [15, chapters 8, 11-13].

Let  $\mathbb{A}$  be an  $n$ -dimensional commutative associative algebra over the field of complex numbers  $\mathbb{C}$  and let  $e_1, e_2, e_3 \in \mathbb{A}$  be three linearly independent vectors over the real field  $\mathbb{R}$ . Let  $\zeta := xe_1 + ye_2 + ze_3$ ,  $x, y, z \in \mathbb{R}$ . Let  $E_3 := \{\zeta = xe_1 + ye_2 + ze_3 : x, y, z \in \mathbb{R}\}$  be the linear span of vectors  $e_1, e_2, e_3$  over the field  $\mathbb{R}$ . Let  $\Omega$  be a domain in  $E_3$ .

**Definition 1.** *We say that a continuous function  $\Phi : \Omega \rightarrow \mathbb{A}$  is monogenic in  $\Omega$  if  $\Phi$  is differentiable in the sense of Gateaux in every point of  $\Omega$ , i. e. if for every  $\zeta \in \Omega$  there exists an element  $\Phi'(\zeta) \in \mathbb{A}$  such that*

$$\lim_{\varepsilon \rightarrow 0+0} \frac{\Phi(\zeta + \varepsilon h) - \Phi(\zeta)}{\varepsilon} = h\Phi'(\zeta) \quad \forall h \in E_3. \quad (2)$$

$\Phi'(\zeta)$  is the Gateaux derivative of the function  $\Phi$  in the point  $\zeta$ .

In the paper [11] it is shown that under certain conditions, which are fulfilled in our case, the monogenic function  $\Phi$  has Gateaux derivatives of all orders  $\Phi^r$ ,  $r = 1, 2, 3, \dots$ . Therefore, if a function  $\Phi$  is monogenic in every point of  $\Omega$ , then

$$\frac{\partial^4 \Phi}{\partial x^4} = e_1^4 \Phi^{IV}(\zeta), \quad \frac{\partial^4 \Phi}{\partial x^2 \partial y^2} = e_1^2 e_2^2 \Phi^{IV}(\zeta),$$

etc. Consequently, due to the equality

$$\Delta^2 \Phi(\zeta) = \Phi^{IV}(\zeta)(e_1^2 + e_2^2 + e_3^2)^2 = 0$$

every monogenic in  $\Omega$  function  $\Phi$  satisfies the equation (1) everywhere in  $\Omega$  if and only if

$$(e_1^2 + e_2^2 + e_3^2)^2 = 0. \quad (3)$$

**Definition 2.** Equation (3) is called the characteristic equation.

Instead of the algebra  $\mathbb{A}$ , we will consider a sequence of algebras  $\{\mathbb{E}_\rho^n\}_{n=1}^\infty$ . For any natural  $n$ , let  $\mathbb{E}_\rho^n$  be a commutative associative algebra over the complex field with a basis  $\{1, \rho, \rho^2, \dots, \rho^{n-1}\}$ , where  $\rho^n = 0$ .

Let vectors  $e_1, e_2, e_3 \in \mathbb{E}_\rho^n$  have the following decompositions with respect to the basis of the algebra  $\mathbb{E}_\rho^n$ :

$$e_1 := \sum_{r=0}^{n-1} k_r \rho^r, \quad e_2 := \sum_{r=0}^{n-1} m_r \rho^r, \quad e_3 := \sum_{r=0}^{n-1} g_r \rho^r, \quad k_r, m_r, g_r \in \mathbb{C}. \quad (4)$$

Then

$$\zeta = xe_1 + ye_2 + ze_3 = \sum_{r=0}^{n-1} (k_r x + m_r y + g_r z) \rho^r =: \sum_{r=0}^{n-1} \xi_r \rho^r,$$

where  $\xi_0 := k_0 x + m_0 y + g_0 z$  and  $\xi_r := k_r x + m_r y + g_r z$ ,  $r = 1, 2, \dots, n-1$ . In this case, the decomposition of resolvent  $(t - \zeta)^{-1}$  is of the form

$$(t - \zeta)^{-1} = \sum_{k=0}^{n-1} A_k \rho^k \quad \forall \zeta \in E_3 \quad \forall t \in \mathbb{C} : t \neq \xi, \quad (5)$$

where the coefficients  $A_k$  are determined by the following recurrence relations:

$$A_0 = \frac{1}{t - \xi_0}, \quad A_s = \frac{1}{t - \xi_0} (\xi_s A_0 + \xi_{s-1} A_1 + \dots + \xi_1 A_{s-1}), \quad s = 1, 2, \dots \quad (6)$$

In particular, we have the following initial functions:

$$\begin{aligned} A_0 &= \frac{1}{t - \xi_0}, & A_1 &= \frac{\xi_1}{(t - \xi_0)^2}, & A_2 &= \frac{\xi_2}{(t - \xi_0)^2} + \frac{\xi_1^2}{(t - \xi_0)^3}, \\ A_3 &= \frac{\xi_3}{(t - \xi_0)^2} + \frac{2\xi_1\xi_2}{(t - \xi_0)^3} + \frac{\xi_1^3}{(t - \xi_0)^4}, \\ A_4 &= \frac{\xi_4}{(t - \xi_0)^2} + \frac{2\xi_1\xi_3 + \xi_2^2}{(t - \xi_0)^3} + \frac{3\xi_1^2\xi_2}{(t - \xi_0)^4} + \frac{\xi_1^4}{(t - \xi_0)^5}, \\ A_5 &= \frac{\xi_5}{(t - \xi_0)^2} + \frac{2\xi_1\xi_4 + 2\xi_2\xi_3}{(t - \xi_0)^3} + \frac{3\xi_1^2\xi_3 + 3\xi_1\xi_2^2}{(t - \xi_0)^4} + \frac{4\xi_1^3\xi_2}{(t - \xi_0)^5} + \frac{\xi_1^5}{(t - \xi_0)^6}, \\ A_6 &= \frac{\xi_6}{(t - \xi_0)^2} + \frac{\xi_3^2 + 2\xi_1\xi_5 + 2\xi_2\xi_4}{(t - \xi_0)^3} + \frac{\xi_2^3 + 6\xi_1\xi_2\xi_3 + 3\xi_1^2\xi_4}{(t - \xi_0)^4} \end{aligned}$$

$$+\frac{4\xi_1^3\xi_3+6\xi_1^2\xi_2^2}{(t-\xi_0)^5}+\frac{5\xi_1^4\xi_2}{(t-\xi_0)^6}+\frac{\xi_1^6}{(t-\xi_0)^7},$$

etc.

Let  $D$  be a domain in the complex plane and let  $\Pi := \{(x, y, z) \in \mathbb{R}^3 : \xi_0 = k_0x + m_0y + g_0z \in D\}$ .

As shown in [14] (see also subsections 13.2.1, 13.3.2 in [15]) that an infinite-dimensional family of solutions of equation (1) on the set  $\Pi$  is of the following form:

$$\left\{ \frac{1}{2\pi i} \int_{\Gamma} F(t) A_k dt \right\}_{k=0}^{\infty}, \quad (7)$$

where  $F : D \rightarrow \mathbb{C}$  may be arbitrary holomorphic functions in the domain  $D$ ,  $\Gamma$  is a closed Jordan rectifiable curve in  $D$  that surround the point  $\xi_0$ , and  $A_k$  are defined by the recurrent formulas (6), where  $n$  can increase to infinity.

Let us write some initial functions, which belong to this family and are also denoted as  $\{U_k(x, y, z)\}_{k=0}^{\infty}$ . After substituting the functions  $A_0, A_1, \dots, A_6$  into expressions (7), we get

$$U_0 = F_0(\xi_0), \quad U_1 = \xi_1 F_1'(\xi_0), \quad U_2 = \xi_2 F_2'(\xi_0) + \frac{\xi_1^2}{2!} F_2''(\xi_0),$$

$$U_3 = \xi_3 F_3'(\xi_0) + \xi_1 \xi_2 F_3''(\xi_0) + \frac{\xi_1^3}{3!} F_3'''(\xi_0),$$

$$U_4 = \xi_4 F_4'(\xi_0) + \frac{1}{2}(2\xi_1 \xi_3 + \xi_2^2) F_4''(\xi_0) + \frac{\xi_1^2 \xi_2}{2} F_4'''(\xi_0) + \frac{\xi_1^4}{4!} F_4^{(4)}(\xi_0),$$

$$U_5 = \xi_5 F_5'(\xi_0) + (\xi_1 \xi_4 + \xi_2 \xi_3) F_5''(\xi_0) + \frac{1}{2}(\xi_1 \xi_2^2 + \xi_1^2 \xi_3) F_5'''(\xi_0)$$

$$+\frac{1}{3!} \xi_1^3 \xi_2 F_5^{(4)}(\xi_0) + \frac{\xi_1^5}{5!} F_5^{(5)}(\xi_0),$$

$$U_6 = \xi_6 F_6'(\xi_0) + \frac{1}{2}(\xi_3^2 + 2\xi_1 \xi_5 + 2\xi_2 \xi_4) F_6''(\xi_0) + \frac{1}{6}(\xi_2^3 + 6\xi_1 \xi_2 \xi_3 + 3\xi_1^2 \xi_4) F_6'''(\xi_0)$$

$$+\frac{1}{4!}(4\xi_1^3 \xi_3 + 6\xi_1^2 \xi_2^2) F_6^{(4)}(\xi_0) + \frac{\xi_1^4 \xi_2}{4!} F_6^{(5)}(\xi_0) + \frac{\xi_1^6}{6!} F_6^{(6)}(\xi_0),$$

etc., where  $F_m$  for  $m = 0, 1, 2, 3, 4, 5, 6$  are arbitrary holomorphic functions of a complex variable.

Calculating the subsequent functions  $A_k$  by the recurrent formula (6), we can write the infinite-dimensional family of solutions of equation (1) that represent family of form (7).

### 3 An infinite-dimensional family of exact solutions of a three-dimensional biharmonic equation

We shall construct solutions of form (7) for the three-dimensional biharmonic equation (1). For this purpose, in the sequence of algebras  $\{\mathbb{E}_\rho^n\}_{n=1}^\infty$ , we shall find all triads of vectors  $e_1, e_2, e_3$  satisfying the characteristic equation (3).

Using decomposition (4) and multiplication table of the algebra  $\mathbb{E}_\rho^n$ , we have

$$e_1^2 = \sum_{r=0}^{n-1} B_r \rho^r, \quad (8)$$

where

$$B_0 = k_0^2, \quad B_1 = 2k_0k_1, \quad B_2 = k_1^2 + 2k_0k_2,$$

and, in the general case,

$$B_r \equiv B_r(k_0, k_1, \dots, k_r) := \begin{cases} k_{r/2}^2 + 2(k_0k_r + k_1k_{r-1} + \dots + k_{\frac{r}{2}-1}k_{\frac{r}{2}+1}) & \text{if } r \text{ is even,} \\ 2(k_0k_r + k_1k_{r-1} + \dots + k_{\frac{r-1}{2}}k_{\frac{r+1}{2}}) & \text{if } r \text{ is odd.} \end{cases}$$

In a similar way, we have

$$e_2^2 = \sum_{r=0}^{n-1} C_r \rho^r, \quad (9)$$

where the coefficients  $C_r$  satisfies the relations

$$C_r \equiv C_r(m_0, m_1, \dots, m_r) = B_r(m_0, m_1, \dots, m_r)$$

for  $r = 0, 1, \dots, n-1$ .

Similarly,

$$e_3^2 = \sum_{r=0}^{n-1} D_r \rho^r,$$

where the coefficients  $D_r$  satisfies the relations

$$D_r \equiv D_r(g_0, g_1, \dots, g_r) = B_r(g_0, g_1, \dots, g_r)$$

for  $r = 0, 1, \dots, n-1$ .

Now, using (8) and multiplication table of the algebra  $\mathbb{E}_\rho^n$ , we have

$$e_1^4 = \sum_{r=0}^{n-1} G_r \rho^r,$$

where the coefficients  $G_r$  satisfies the relations

$$G_r \equiv G_r(k_0, k_1, \dots, k_r) = B_r(B_0, B_1, \dots, B_r) \quad (10)$$

for  $r = 0, 1, \dots, n - 1$ .

Similarly,

$$e_2^4 = \sum_{r=0}^{n-1} H_r \rho^r,$$

where the coefficients  $H_r$  satisfies the relations

$$H_r \equiv H_r(m_0, m_1, \dots, m_r) = C_r(C_0, C_1, \dots, C_r) = G_r(m_0, m_1, \dots, m_r) \quad (11)$$

for  $r = 0, 1, \dots, n - 1$ , and

$$e_3^4 = \sum_{r=0}^{n-1} M_r \rho^r,$$

where the coefficients  $M_r$  satisfies the relations

$$M_r \equiv M_r(g_0, g_1, \dots, g_r) = D_r(D_0, D_1, \dots, D_r) = G_r(g_0, g_1, \dots, g_r) \quad (12)$$

for  $r = 0, 1, \dots, n - 1$ .

Equalities (8), (9) yield

$$e_1^2 e_2^2 = \sum_{r=0}^{n-1} P_r \rho^r,$$

where the coefficients  $P_r$  satisfies the relations

$$P_r = B_0 C_r + B_1 C_{r-1} + \dots + B_r C_0 \quad (13)$$

for  $r = 0, 1, \dots, n - 1$ .

Similarly,

$$e_1^2 e_3^2 = \sum_{r=0}^{n-1} R_r \rho^r,$$

where the coefficients  $R_r$  satisfies the relations

$$R_r = B_0 D_r + B_1 D_{r-1} + \dots + B_r D_0 \quad (14)$$

for  $r = 0, 1, \dots, n - 1$ , and

$$e_2^2 e_3^2 = \sum_{r=0}^{n-1} S_r \rho^r,$$

where the coefficients  $S_r$  satisfies the relations

$$S_r = C_0 D_r + C_1 D_{r-1} + \cdots + C_r D_0 \quad (15)$$

for  $r = 0, 1, \dots, n-1$ .

Thus, characteristic equation (3) is equivalent to the following system of equations

$$G_r + H_r + M_r + 2P_r + 2R_r + 2S_r = 0, \quad r = 0, 1, 2, \dots, \quad (16)$$

where  $G_r, H_r, M_r, P_r, R_r, S_r$  are defined by equalities (10), (11), (12), (13), (14), (15), respectively.

According to Remark 13.1 [15] in characteristic equation (3) we can set the vectors  $e_1, e_2$  arbitrarily, i.e.,  $k_r$  and  $m_r$  may be arbitrary complex numbers for  $r = 0, 1, \dots, n-1$  for any natural  $n$ . Then the vector  $e_3$  can be found, and the coefficients  $g_r$  can be found from system (16).

We note that for every fixed  $r$  equation (16) is linear with respect to  $g_r$ . Therefore, it is easy to find  $g_r$  for each fixed  $r = 0, 1, 2, \dots$ .

So, characteristic equation (3) is solved on the sequence of algebras  $\{\mathbb{E}_\rho^n\}_{n=1}^\infty$ .

Thus, we proved the following theorem

**Theorem.** *In a domain  $\Pi$  equation (1) has infinite-dimensional family of exact solutions of the form (7), where  $F : D \rightarrow \mathbb{C}$  may be arbitrary holomorphic functions in the domain  $D$ ,  $\Gamma$  is a closed Jordan rectifiable curve in  $D$  that surround the point  $\xi_0$ , and  $A_k$  are defined by the recurrent formulas (6), where  $n$  can increase to infinity. Moreover, in coefficients  $A_k$  the variables  $\xi_0, \xi_r$  are of the form  $\xi_0 = k_0 x + m_0 y + g_0 z$ ,  $\xi_r = k_r x + m_r y + g_r z$ ,  $r = 1, 2, \dots$ , where  $k_r, m_r$  for  $r = 0, 1, 2, \dots$ , are arbitrary complex numbers, and  $g_r$  for each fixed  $r = 0, 1, 2, \dots$  are determined by equality (16).*

#### 4 Several first solutions from family (7)

Now we will write some first solutions from family (7). For  $r = 0$  equation (16) has the form

$$(k_0^2 + m_0^2 + g_0^2)^2 = 0 \quad \text{or} \quad g_0 = \pm i \sqrt{k_0^2 + m_0^2}. \quad (17)$$

Therefore,  $\xi_0 = k_0 x + m_0 y \pm i \sqrt{k_0^2 + m_0^2} z$ , where  $k_0, m_0$  are arbitrary complex numbers.

Thus, the **first exact solution** of equation (1) from family (7) is of the form

$$U_0(x, y, z) = F_0(\xi_0) = F_0 \left( k_0 x + m_0 y \pm i \sqrt{k_0^2 + m_0^2} z \right),$$

where  $F_0$  is an arbitrary holomorphic function of the complex variable  $\xi_0 = k_0x + m_0y \pm i\sqrt{k_0^2 + m_0^2}z$  with arbitrary complex numbers  $k_0, m_0$ .

For  $r = 1$  equation (16) has the following form

$$\begin{aligned} & 2k_0^3k_1 + 2m_0^3m_1 + 2g_0^3g_1 + 4k_0^2m_0m_1 + 4m_0^2k_0k_1 \\ & + 4k_0^2g_0g_1 + 4k_0k_1g_0^2 + 4m_0^2g_0g_1 + 4m_0m_1g_0^2 = 0, \end{aligned} \quad (18)$$

i.e., we have the linear equation with respect to  $g_1$ . Substituting expression (17) into (18), we obtain

$$g_1 = \pm i \frac{k_0^3k_1 + m_0^3m_1}{2 \left( \sqrt{k_0^2 + m_0^2} \right)^3}. \quad (19)$$

Therefore,

$$\xi_1 = k_1x + m_1y \pm i \frac{k_0^3k_1 + m_0^3m_1}{2 \left( \sqrt{k_0^2 + m_0^2} \right)^3} z,$$

where  $k_0, m_0, k_1, m_1$  are arbitrary complex numbers.

Thus, the **second exact solution** of equation (1) from family (7) is of the form

$$\begin{aligned} & U_1(x, y, z) = \xi_1 F_1(\xi_0) \\ & = \left( k_1x + m_1y \pm i \frac{k_0^3k_1 + m_0^3m_1}{2 \left( \sqrt{k_0^2 + m_0^2} \right)^3} z \right) F_1 \left( k_0x + m_0y \pm i \sqrt{k_0^2 + m_0^2} z \right), \end{aligned}$$

where  $F_1$  is an arbitrary holomorphic function of the complex variable  $\xi_0 = k_0x + m_0y \pm i\sqrt{k_0^2 + m_0^2}z$ , and  $k_0, m_0, k_1, m_1$  are arbitrary complex numbers.

For  $r = 2$  equation (16) has the following form

$$\begin{aligned} & 4k_0^6k_1^2 + 4k_0^4k_2 + 2k_0^2k_1^2 + 4m_0^6m_1^2 + 4m_0^4m_2 + 2m_0^2m_1^2 + 4g_0^6g_1^2 + 4g_0^4g_2 + 2g_0^2g_1^2 \\ & + 2k_0^2m_1^2 + 4k_0^2m_0m_2 + 8k_0k_1m_0m_1 + 2m_0^2k_1^2 + 4m_0^2k_0k_2 + 2k_0^2g_1^2 + 4k_0^2g_0g_2 \\ & + 8k_0k_1g_0g_1 + 2k_1^2g_0^2 + 4k_0k_2g_0^2 + 2m_0^2g_1^2 + 4m_0^2g_0g_2 + 8m_0m_1g_0g_1 \\ & + 2m_1^2g_0^2 + 4m_0m_2g_0^2 = 0. \end{aligned} \quad (20)$$

i.e., we have the linear equation with respect to  $g_2$ . From (20) we obtain

$$\begin{aligned} g_2 &= \frac{1}{4(g_0^3 - g_0^4)} \left( 4k_0^6k_1^2 + 4k_0^4k_2 + 2k_0^2k_1^2 + 4m_0^6m_1^2 + 4m_0^4m_2 + 2m_0^2m_1^2 \right. \\ & + 4g_0^6g_1^2 + 2g_0^2g_1^2 + 2k_0^2m_1^2 + 4k_0^2m_0m_2 + 8k_0k_1m_0m_1 + 2m_0^2k_1^2 + 4m_0^2k_0k_2 \\ & \left. + 2k_0^2g_1^2 + 8k_0k_1g_0g_1 + 2k_1^2g_0^2 + 4k_0k_2g_0^2 + 2m_0^2g_1^2 + 8m_0m_1g_0g_1 \right) \end{aligned}$$



$$+2m_1^2g_0^2 + 4m_0m_2g_0^2), \quad (21)$$

where  $g_0, g_1$  are defined by (17), (19), and  $k_0, m_0, k_1, m_1, k_2, m_2$  are arbitrary complex numbers.

Therefore,

$$\begin{aligned} \xi_2 = & k_2x + m_2y + \frac{1}{4(g_0^3 - g_0^4)} \left( 4k_0^6k_1^2 + 4k_0^4k_2 + 2k_0^2k_1^2 + 4m_0^6m_1^2 + 4m_0^4m_2 \right. \\ & + 2m_0^2m_1^2 + 4g_0^6g_1^2 + 2g_0^2g_1^2 + 2k_0^2m_1^2 + 4k_0^2m_0m_2 + 8k_0k_1m_0m_1 + 2m_0^2k_1^2 \\ & + 4m_0^2k_0k_2 + 2k_0^2g_1^2 + 8k_0k_1g_0g_1 + 2k_1^2g_0^2 + 4k_0k_2g_0^2 + 2m_0^2g_1^2 \\ & \left. + 8m_0m_1g_0g_1 + 2m_1^2g_0^2 + 4m_0m_2g_0^2 \right) z, \end{aligned}$$

where  $k_0, m_0, k_1, m_1, k_2, m_2$  are arbitrary complex numbers, and  $g_0, g_1$  are defined by equalities (17), (19).

Thus, the **third exact solution** of equation (1) from family (7) is of the form

$$\begin{aligned} U_2 = & \xi_2 F_2(\xi_0) + \frac{\xi_1^2}{2!} F_2'(\xi_0) \\ = & \left[ k_2x + m_2y + \frac{1}{4(g_0^3 - g_0^4)} \left( 4k_0^6k_1^2 + 4k_0^4k_2 + 2k_0^2k_1^2 + 4m_0^6m_1^2 + 4m_0^4m_2 \right. \right. \\ & + 2m_0^2m_1^2 + 4g_0^6g_1^2 + 2g_0^2g_1^2 + 2k_0^2m_1^2 + 4k_0^2m_0m_2 + 8k_0k_1m_0m_1 + 2m_0^2k_1^2 \\ & + 4m_0^2k_0k_2 + 2k_0^2g_1^2 + 8k_0k_1g_0g_1 + 2k_1^2g_0^2 + 4k_0k_2g_0^2 + 2m_0^2g_1^2 \\ & \left. \left. + 8m_0m_1g_0g_1 + 2m_1^2g_0^2 + 4m_0m_2g_0^2 \right) z \right] F_2 \left( k_0x + m_0y \pm i\sqrt{k_0^2 + m_0^2} z \right) \\ & + \frac{1}{2} \left( k_1x + m_1y \pm i \frac{k_0^3k_1 + m_0^3m_1}{2(\sqrt{k_0^2 + m_0^2})^3} z \right)^2 F_2' \left( k_0x + m_0y \pm i\sqrt{k_0^2 + m_0^2} z \right), \end{aligned}$$

where  $F_2$  is an arbitrary holomorphic function of the complex variable  $\xi_0 = k_0x + m_0y \pm i\sqrt{k_0^2 + m_0^2} z$ , and  $k_0, m_0, k_1, m_1, k_2, m_2$  are arbitrary complex numbers, and  $g_0, g_1$  are defined by equalities (17), (19).

Continuing in this way we can write out an infinite number of exact solutions of equation (1).

### The algorithm.

1. • We put  $r = 0$ ;
- from the linear equation (16) we find  $g_0$  and we write out  $\xi_0 = k_0x + m_0y + g_0z$ . Here  $k_0, m_0$  are arbitrary complex numbers;

- $\xi_0$  we substitute in  $A_0$  from (6);
  - $A_0$  we substitute in (7) and we obtain the solution  $U_0$ .
- 2.
- We put  $r = 1$ ;
  - having  $g_0$  from the linear equation (16) we find  $g_1$  and we write out  $\xi_1 = k_1x + m_1y + g_1z$ . Here  $k_0, m_0, k_1, m_1$  are arbitrary complex numbers;
  - $\xi_0$  and  $\xi_1$  we substitute in  $A_1$  from (6);
  - $A_1$  we substitute in (7) and we obtain the solution  $U_1$ .
- 3.
- We put  $r = 2$ ;
  - having  $g_0, g_1$  from the linear equation (16) we find  $g_2$  and we write out  $\xi_2 = k_2x + m_2y + g_2z$ . Here  $k_0, m_0, k_1, m_1, k_2, m_2$  are arbitrary complex numbers;
  - $\xi_0, \xi_1$  and  $\xi_2$  we substitute in  $A_2$  from (6);
  - $A_2$  we substitute in (7) and we obtain the solution  $U_2$ .
4. etc.

### Acknowledgment

This work was supported by a grant from the Simons Foundation (1030291,V.S.Sh.).

### References

- [1] Grigor'ev, Yu. M.: Three-dimensional Quaternionic Analogue of the Kolosov–Muskhelishvili Formulae, in: New perspectives and applications, Trends in Mathematics, (Eds. S. Bernstein, U. Kaehler, I. Sabadini, F. Sommen), Basel, Birkhauser, pp. 145–166, 2014.
- [2] Grigor'ev, Yu. M.: Solution of spatial problems of an elasticity theory by means of a quaternionic functions theory. Candidat dissertation (PhD) [in Russian], Lavrent'ev Institute of Hydrodynamics, Novosibirsk, 1985.
- [3] Changhao, Lin: Energy estimates for the biharmonic equation in three dimensions. Quarterly of Applied Mathematics, **52** (4), 649–663 (1994).
- [4] Altas, I., Erhel, J., and Gupta, M. M.: High Accuracy Solution of Three-Dimensional Biharmonic Equations. Numerical Algorithms **29**, 1–19 (2002).

- [5] Ketchum, P. W.: Analytic functions of hypercomplex variables. *Trans. Amer. Math. Soc.*, **30**, 641–667 (1928).
- [6] Ketchum, P.W.: A complete solution of Laplace’s equation by an infinite hypervariable. *Amer. J. Math.* **51**, 179–188 (1929)
- [7] Ketchum, P.W.: Solution of partial differential equations by means of hypervariables. *American J. Math.* **54** (2), 253–264 (1932)
- [8] Mel’nichenko, I. P.: The representation of harmonic mappings by monogenic functions. *Ukr. Math. J.*, **27** (5), 499–505 (1975).
- [9] Mel’nichenko, I. P.: Algebras of functionally invariant solutions of the three-dimensional Laplace equation. *Ukr. Math. J.*, **55**, no. 9, 1551–1557 (2003)
- [10] Plaksa, S. A.: Commutative algebras associated with classic equations of mathematical physics. *Advances in Applied Analysis, Trends in Mathematics*. — Basel: Springer, 2012. — P. 177 – 223.
- [11] Shpakivskyi, V. S.: Constructive description of monogenic functions in a finite-dimensional commutative associative algebra. *Adv. Pure Appl. Math.*, **7**(1), 63–75, 2016.
- [12] Shpakivskyi, V. S.: On monogenic functions defined in different commutative algebras. *J. Math. Sci. (USA)*, **239**(1), 92–109 (2019)
- [13] Shpakivskyi, V. S.: On monogenic functions on extensions of commutative algebra (in Ukrainian). *Proceedings of the International Geometrical Center*, **11**(3), 1–18 (2018)
- [14] Shpakivskyi, V. S.: Hypercomplex method for solving linear partial differential equations (in Ukrainian). *Proc. of the Inst. Appl. Math. Mech. NAS Ukraine*, **32**, 147–168 (2018)
- [15] Plaksa, S. A., Shpakivskyi, V. S.: *Monogenic Functions in Spaces with Commutative Multiplication and Applications*. *Frontiers in Mathematics*, Birkhäuser Cham, 2023, 550 p.

Vitalii Shpakivskyi

Institute of Mathematics of the

National Academy of Sciences of Ukraine, Kyiv

e-mail: shpakivskyi86@gmail.com