

WEIGHTED CSCK METRIC (I): A PRIORI ESTIMATES

ELEONORA DI NEZZA, SIMON JUBERT AND ABDELLAH LAHDILI

ABSTRACT. Let X be a compact Kähler manifold. In this paper we study the existence of constant weighted scalar curvature Kähler (weighted cscK) metrics on X . More precisely, we establish a priori C^k -estimates ($k \geq 0$) for the Kähler potential associated with these metrics, thereby extending a result due to Chen and Cheng in the classical cscK setting.

CONTENTS

1. Introduction	1
1.1. Weighted cscK problem	2
1.2. Main results and strategy of the proof	3
1.3. Notations	4
1.4. Organisation of the paper	4
Acknowledgements	4
2. Preliminaries in pluripotential theory	4
3. Weighted cscK equation as a system of elliptic PDEs	5
4. C^0 -estimates	9
5. Integral C^2 -estimates	14
6. C^2 -estimates	22
Appendix A. Weighted Laplacian and weighted trace	30
A.1. The weighted Laplacian	30
A.2. The weighted trace	30
References	31

1. INTRODUCTION

A central theme in Kähler geometry is the search for canonical Kähler metrics. The concept of *constant weighted scalar curvature metrics* (weighted cscK for short), introduced by the third author in [25], provides a unification for various related notions of canonical Kähler metrics that satisfy special curvature conditions.

We present here the first of two papers, focused on investigating the existence of weighted cscK metrics in a given cohomology class. In this paper, we derive *a priori* C^k -estimates, $k \geq 0$, for the Kähler potential associated to the metric with constant weighted scalar curvature. This work extends the result of Chen and Cheng [12] concerning cscK metrics. In the subsequent second paper, we will demonstrate the existence of the weighted cscK metric provided that the weighted Mabuchi energy is relatively coercive, and we will explore several geometric applications.

1.1. Weighted cscK problem. We fix a compact Kähler manifold $(X, \omega_0, \mathbb{T})$ of complex dimension n . Let \mathbb{T} be an r -dimensional compact real torus in the reduced automorphism group $\text{Aut}_{\text{red}}(X)$ with Lie algebra \mathfrak{t} . Suppose ω_0 is a \mathbb{T} -invariant Kähler form and consider the set of smooth \mathbb{T} -invariant Kähler potentials $\mathcal{K}(X, \omega_0)^{\mathbb{T}}$ relative to ω_0 . For $\varphi \in \mathcal{K}(X, \omega_0)^{\mathbb{T}}$ we denote by $\omega_\varphi = \omega_0 + dd^c \varphi$ the corresponding Kähler metric. Here $d = \partial + \bar{\partial}$ and $d^c = i(\bar{\partial} - \partial)$ so that $dd^c = 2i\partial\bar{\partial}$ as in [17, page 29]. This choice of d and d^c is consistent with the fact that we choose the scalar curvature to be defined as two times the trace; note that this normalization has an impact also in the expression of the Ricci form, i.e. $\text{Ric}(\omega_\varphi) = \text{Ric}(\omega_0) + \frac{1}{2}dd^c \log \frac{\omega_\varphi^n}{\omega_0^n}$.

It is well-known that the \mathbb{T} -action on X is ω_φ -Hamiltonian (see e.g. [17, Section 2.5]) and that $P_\varphi := \mu_\varphi(X)$ is a convex polytope in \mathfrak{t}^* [5, 19, 25]. Here $\mu_\varphi : X \rightarrow \mathfrak{t}^*$ is the moment map associated to ω_φ and \mathfrak{t}^* stands for the dual vector space for the Lie algebra \mathfrak{t} of \mathbb{T} . We normalize μ_φ by

$$(1) \quad \mu_\varphi = \mu_0 + d^c \varphi,$$

in such a way that $P = P_\varphi$ is φ -independent, see [25, Lemma 1]. From now, we always suppose that the moment map satisfies (1). Let us stress that the identity in (1) means that for all $\xi \in \mathfrak{t}$,

$$\langle \mu_\varphi, \xi \rangle = \langle \mu_0, \xi \rangle + \langle d^c \varphi, \xi \rangle, \quad \text{with} \quad \langle d^c \varphi, \xi \rangle = \sum_{i=1}^r \langle d^c \varphi(\xi_i) \xi_i^*, \xi \rangle,$$

where (ξ_1, \dots, ξ_r) is a basis of the Lie algebra \mathfrak{t} of \mathbb{T} and $(\xi_1^*, \dots, \xi_r^*)$ is its dual basis.

For a given positive weight function $v \in C^\infty(P, \mathbb{R}_{>0})$, we define *the v-weighted Ricci form* by

$$(2) \quad \text{Ric}_v(\omega_\varphi) := \text{Ric}(\omega_\varphi) - \frac{1}{2}dd^c \log v(\mu_\varphi).$$

Then the v -weighted scalar curvature of a \mathbb{T} -invariant Kähler metric $\omega_\varphi \in [\omega_0]$ is defined by

$$(3) \quad \text{Scal}_v(\omega_\varphi) := 2v(\mu_\varphi)\Lambda_{\varphi, v}(\text{Ric}_v(\omega_\varphi)),$$

where $\Lambda_{\varphi, v}$ is the v -weighted trace with respect to ω_φ (see Appendix A.2 for more details about this operator). This definition is equivalent to the one introduced in [25] (see Lemma 3.2), but we do prefer the above more compact expression.

For a given weight $w \in C^\infty(P, \mathbb{R})$, a \mathbb{T} -invariant Kähler metric ω_φ is said *(v, w)-weighted cscK* if its v -weighted scalar curvature satisfies

$$(4) \quad \text{Scal}_v(\omega_\varphi) = w(\mu_\varphi).$$

The significance of (4) in relation to various geometric conditions is thoroughly examined in [25] (see also [4] for v -solitons). However, we shall mention a few specific cases below:

- (a) for $v = 1$ and $w = c$ is a constant, (4) corresponds to the classical cscK equation;
- (b) for $v = 1$ and $w = \ell_{\text{ext}}$, where ℓ_{ext} is the affine extremal function, the weighted cscK metrics are Calabi's extremal Kähler metrics [10];
- (c) for $v > 0$ is any smooth function and $w = 2v(x)(n + \langle d \log v(x), x \rangle)$, (4) corresponds to the weighted v -soliton examined in [4, 20], generalizing the well-studied Kähler-Ricci solitons (see e.g. [7, 8, 27, 28]);

- (d) for v and w polynomials, then (v, w) -cscK metrics on X correspond to Calabi's extremal Kähler metrics on the total space of an holomorphic fibration Y with fiber X , called semisimple principal fibration [2, 4, 22];
- (e) for $v = \ell^{-n-1}$, $w = a\ell^{-n-2}$ and $[\omega_0] = c_1(L)$, where ℓ is a positive affine-linear function on P , a is a constant, and L is a polarization of X , then a weighted cscK metric is a scalar flat cone Kähler metric on the affine cone $(L^{-1})^\times$ (see [1, 3]).

1.2. Main results and strategy of the proof. The starting point is to re-write the 4th order non-linear PDE in (4) as a system of two linear elliptic PDEs:

$$(5) \quad \begin{cases} F &= \log \left(v(\mu_\varphi) \frac{\omega_\varphi^n}{\omega_0^n} \right) \\ \Delta_{\varphi, v} F &= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi, v}(\text{Ric}(\omega_0)). \end{cases}$$

The equivalence between (4) and (5) is established in Proposition 3.3.

In the above equation $\Delta_{\varphi, v}$ is the v -weighted Laplacian with respect to ω_φ (we refer to Appendix A.1 for basic properties of this operator). Note that for $v = 1$ and w being constant equal to $\frac{nc_1(X) \cdot \{\omega_0\}^{n-1}}{\{\omega_0\}^n}$, we retrieve the system proposed by [12] in the cscK case.

In the original proof by Chen and Cheng [12] in the unweighted setting, the authors derive C^0 and C^2 a priori estimates by establishing an intermediate C^1 a priori estimate for the Kähler potential φ . With the C^0 and C^2 estimates established, higher order estimates are subsequently obtained through standard regularity results for complex Monge-Ampère equations and linear elliptic operators. It is important to emphasize that their proof of the C^0 estimate [12, Theorem 5.1] relies on the Alexandroff maximum principle for the real Monge-Ampère operator.

In this paper, following [16], we present a proof of the C^0 and C^2 estimates that bypass the need for an intermediate C^1 estimate. However, it is important to note that the “weighted” case introduces significantly more challenges and subtleties due to the presence of the weighted operators $\Delta_{\varphi, v}$ and $\Lambda_{\varphi, v}$ in (5). In practice, many additional terms require (uniform) bounds when establishing a priori estimates. We address these terms with detailed explanations to ensure clarity in the presentation.

Our first main result is the C^0 -estimate:

Theorem A. *The functions φ and F are uniformly bounded by a constant that depends on n, ω_0, v, w and the entropy $\text{Ent}(\varphi) := \int_X \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) \omega_\varphi^n$.*

The proof of the C^0 -estimate is an extension of the approach proposed in [16], which crucially employs pluripotential theory.

From this, we then get C^2 -estimate:

Theorem B. *There exists a positive constant $C > 0$ depending on $\omega_0, v, w, \|F\|_{C^0}$ and $\|\varphi\|_{C^0}$ such that*

$$\max_X (|dF|_\varphi^2 + \Lambda_0(\omega_\varphi)) \leq C$$

where Λ_0 is the trace with respect to ω_0 .

Two crucial intermediate steps consist in establishing a priori integral L^p -estimates ($p \geq 1$) for the Laplacian Δ_φ (Theorem 5.1) and a lower bound for the weighted Laplacian $\Delta_{\varphi, v} u$ for $u := e^{\frac{F}{2}} |dF|_\varphi^2 + K\Lambda_0(\omega_\varphi)$ (Proposition 6.3).

Noting that $\Delta_{\varphi, \nu}$ is elliptic (see Lemma A.2), the higher-order estimates can be derived using standard arguments of elliptic and complex Monge-Ampère theories [6, 24] similar to those used in the usual cscK problem (cf [12, page 3]).

1.3. Notations. In the text we will denote by $\Delta_{\varphi}, \Lambda_{\varphi}$ the Laplacian and the trace with respect to ω_{φ} . The scalar curvature Scal is defined as twice the trace of the Ricci form, i.e. $\text{Scal}(\omega_{\varphi}) = 2\Lambda_{\varphi}(\text{Ric}(\omega_{\varphi}))$ where the Ricci form is defined by $\text{Ric}(\omega_{\varphi}) := \text{ric}_{\varphi}(J \cdot, \cdot)$ with ric_{φ} the Ricci symmetric 2-tensor of the underlying riemannian metric g_{φ} . We denote by ∇^{φ} the gradient with respect to g_{φ} .

1.4. Organisation of the paper. In Section 2 we introduce notations and results which will be used throughout the paper: we provide definitions and references to pluripotential theory. Section 3 is dedicated to establish the equivalence between the fourth-order PDE (4) and the system of elliptic PDE's (5).

The proof of the C^0 -estimate, stated in Theorem A, is presented in Section 4. In Section 5, we prove the L^p -estimates needed for the proof of Theorem B which can be found finally in Section 6.

ACKNOWLEDGEMENTS

The first author is supported by the project SIGMA ANR-22-ERCS-0004-02. The authors are grateful to Vestislav Apostolov for valuable suggestions. We also thank Sebastien Boucksom and Mattias Jonsson for useful discussions.

We would like to acknowledge Jiyuan Han and Yaxiong Liu for bringing to our attention the overlaps between our project and their work [21]. Despite these similarities, our approach introduces some distinct methodologies. Notably, our C^0 -estimate leverages pluripotential theory, and we circumvent the need for a gradient estimate by directly deriving the C^2 -estimate from the C^0 one.

2. PRELIMINARIES IN PLURIPOTENTIAL THEORY

We recall below some ingredients from pluripotential theory that are going to be crucial in what follows in order to establish uniform C^0 -estimates. The first is a powerful integrability result which is known as a uniform version of Skoda's integrability theorem. We introduce

$$\nu_{\omega_0} := \sup_{u, x} \nu(u, x), \quad x \in X, u \in \text{PSH}(X, \omega_0),$$

where $\nu(u, x)$ denotes the Lelong number of u at x . We note that from the proof of [18, Lemma 8.10] one can deduce that $\nu_{\omega_0} \geq 1$.

Theorem 2.1. *Let $c < 2\nu_{\omega_0}^{-1}$. Then there exists a uniform constant $C > 0$ such that for all $u \in \text{PSH}(X, \omega_0)$ with $\sup_X u = 0$ we have*

$$\int_X e^{-cu} \omega_0^n \leq C.$$

We refer to [18, Theorem 8.11] for a proof. The following result is due to Kołodziej [23]:

Theorem 2.2. *Assume $\omega_u^n = f\omega_0^n$ with $f \in L^p$ for some $p > 1$. Then there exists $C > 0$ depending only on $\omega, n, \|f\|_{L^p}$ such that*

$$\text{Osc}_X u \leq C$$

Here $L^p := L^p(\omega_0^n)$. In the following we specify the reference volume form in the notation of the L^p norms only if it is different from the standard one.

At last, we recall [15, Theorem 3.3], that can be viewed as a generalization of Kołodziej's theorem:

Theorem 2.3. *Fix $a \in [0, 1)$, $A > 0$, $\chi \in \text{PSH}(X, \omega_0)$ and $0 \leq f \in L^p$ for some $p > 1$. Assume that $u \in \text{PSH}(X, \omega_0)$, normalized by $\sup_X u = 0$, satisfies*

$$\omega_u^n \leq f\omega_0^n + a\omega_\chi^n.$$

Assume also that

$$(6) \quad \int_E f\omega_0^n \leq A[\text{Cap}_\chi(E)]^2,$$

for every Borel subset $E \subset X$. If $P[u]$ is less singular than χ (i.e. $\chi \leq P[u] + C$, for some $C > 0$) then

$$\chi - \sup_X \chi - C \left(\|f\|_{L^p}, p, (1-a)^{-1}, A \right) \leq u.$$

It is worth it to mention that such a result is stated and proved in a much more general version in [15] to which we refer for a proof. Here $\text{Cap}_\chi(E)$ is the χ -relative capacity of E and it is defined as

$$\text{Cap}_\chi(E) := \sup \left\{ \int_E \omega_u^n : u \in \text{PSH}(X, \omega_0), \chi - 1 \leq u \leq \chi \right\}$$

and

$$P[u] = (\sup \{v \in \text{PSH}(X, \omega_0), v \leq 0 \text{ and } v \leq u + C, \text{ for some } C > 0\})^*,$$

where $*$ denotes the upper semi-continuous regularization. For later purposes we mention that $P[u] = 0$ if and only if u is such that $\int_X \omega_u^n = V$ ([13, Theorem 1.3]).

3. WEIGHTED CSCK EQUATION AS A SYSTEM OF ELLIPTIC PDES

The goal of this section is to write the weighted cscK equation (4) as a system of two elliptic PDEs. We define $F := \log \left(v(\mu_\varphi) \frac{\omega_\varphi^n}{\omega_0^n} \right)$.

We first show that the weighted scalar curvature introduced by [25] and that writes as

$$(7) \quad \text{Scal}_v(\omega_\varphi) = v(\mu_\varphi)\text{Scal}(\omega_\varphi) - 2\Delta_\varphi v(\mu_\varphi) + \sum_{i,j=1}^r v_{,ij}(\mu_\varphi)g_\varphi(\xi_i, \xi_j),$$

is equivalent to (3). We take (7) as a definition.

We develop the second term of the RHS of (7).

$$(8) \quad \begin{aligned} \Delta_\varphi v(\mu_\varphi) &= \Lambda_\varphi dd^c v(\mu_\varphi) \\ &= \Lambda_\varphi d \left(\sum_{i=1}^r v_{,i}(\mu_\varphi) d^c \mu_\varphi^i \right) \\ &= \Lambda_\varphi \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) dd^c \mu_\varphi^i + v_{,ij}(\mu_\varphi) d\mu_\varphi^i \wedge d^c \mu_\varphi^j \right) \\ &= \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i - v_{,ij}(\mu_\varphi) \Lambda_\varphi (\omega_\varphi(\xi_i, \cdot) \wedge \omega_\varphi(\xi_j, J \cdot)), \end{aligned}$$

where $\mu_\varphi^i := \langle \mu_\varphi, \xi_i \rangle$ and we use that $\omega_\varphi(\xi_i, \cdot) = -d\mu_\varphi^i$ and $\omega_\varphi(\xi_j, J\cdot) = d^c\mu_\varphi^j$ for the last equality. We continue the computation

$$\begin{aligned} \Delta_\varphi v(\mu_\varphi) &= \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i - v_{,ij}(\mu_\varphi) \Lambda_\varphi (g_\varphi(J\xi_i, \cdot) \wedge g_\varphi(\xi_j, \cdot)) \\ &= \sum_{i,j=1}^r \left(v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i - v_{,ij}(\mu_\varphi) \sum_{k=1}^{2n} g_\varphi(J\xi_i, e_k) g_\varphi(\xi_j, J e_k) \right) \\ &= \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i + v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j), \end{aligned}$$

where $(e_k)_{k=1}^{2n}$ is a local orthonormal frame w.r.t. g_φ . We showed

$$(9) \quad -\Delta_\varphi v(\mu_\varphi) = - \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i - v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j).$$

For later purposes, let us observe that the first 4th lines of the above computation give

$$\begin{aligned} \Delta_0 v(\mu_\varphi) &= \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_0 \mu_\varphi^i + v_{,ij}(\mu_\varphi) \Lambda_0 (d\mu_\varphi^i \wedge d^c \mu_\varphi^j) \\ (10) \quad &= \sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_0 \mu_\varphi^i + v_{,ij}(\mu_\varphi) g_0(d\mu_\varphi^i, d\mu_\varphi^j), \end{aligned}$$

where the last identity follows from a general fact, i.e. that for smooth functions f_1, f_2 ,

$$(11) \quad g_0(df_1, df_2) = \Lambda_0(df_1 \wedge d^c f_2).$$

Lemma 3.1. *The scalar curvature has the alternative expression*

$$\begin{aligned} \text{Scal}(\omega_\varphi) &= 2\Lambda_\varphi(\text{Ric}(\omega_0)) - \Delta_\varphi F + \frac{1}{v(\mu_\varphi)} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i + v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right) \\ (12) \quad &- \frac{1}{v(\mu_\varphi)^2} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right), \end{aligned}$$

where $F := \log \left(v(\mu_\varphi) \frac{\omega_\varphi^n}{\omega_0^n} \right)$.

Proof. By definition

$$(13) \quad \text{Scal}(\omega_\varphi) = 2\Lambda_\varphi(\text{Ric}(\omega_0)) - \Delta_\varphi \log \frac{\omega_\varphi^n}{\omega_0^n}.$$

Observe that there is a factor 2 since, with our notations the scalar curvature is twice the trace. Also, by definition of F

$$\Delta_\varphi \log \frac{\omega_\varphi^n}{\omega_0^n} = \Delta_\varphi F - \Delta_\varphi \log v(\mu_\varphi).$$

We develop

$$\begin{aligned}
\Delta_\varphi \log v(\mu_\varphi) &= \Lambda_\varphi dd^c \log v(\mu_\varphi) \\
&= \frac{1}{v(\mu_\varphi)} \Delta_\varphi v(\mu_\varphi) - \Lambda_\varphi \left(\frac{1}{v(\mu_\varphi)^2} dv(\mu_\varphi) \wedge d^c v(\mu_\varphi) \right) \\
&= \frac{1}{v(\mu_\varphi)} \Delta_\varphi v(\mu_\varphi) - \frac{1}{v(\mu_\varphi)^2} \Lambda_\varphi \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) d\mu_\varphi^i \wedge d^c \mu_\varphi^j \right) \\
&= \frac{1}{v(\mu_\varphi)} \Delta_\varphi v(\mu_\varphi) - \frac{1}{v(\mu_\varphi)^2} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right).
\end{aligned}$$

The last equality follows from the same computation as in (8). Then, using (9),

$$\begin{aligned}
\Delta_\varphi \log \frac{\omega_\varphi^n}{\omega_0^n} &= \Delta_\varphi F - \frac{1}{v(\mu_\varphi)} \Delta_\varphi v(\mu_\varphi) + \frac{1}{v(\mu_\varphi)^2} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right) \\
&= \Delta_\varphi F - \frac{1}{v(\mu_\varphi)} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i + v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right) \\
&\quad + \frac{1}{v(\mu_\varphi)^2} \left(\sum_{i,j} v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right).
\end{aligned}$$

Injecting in (13), we get the result. \square

Lemma 3.2. *The weighted scalar curvature has the alternative expression*

$$\text{Scal}_v(\omega_\varphi) = 2v(\mu_\varphi) \Lambda_{\varphi,v}(\text{Ric}_v(\omega_\varphi))$$

where $\Lambda_{\varphi,v}$ is defined in (54) and Ric_v is the v -weighted Ricci form, defined in (2).

Proof. We first re-write the term $\sum_{i=1}^r v_{,i}(\mu_\varphi) \Delta_\varphi \mu_\varphi^i$. Let $\xi_j \in \mathfrak{t}$. Then

$$\begin{aligned}
d^c F(\xi_j) &= \frac{d^c v(\mu_\varphi)(\xi_j)}{v(\mu_\varphi)} + d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) (\xi_j) \\
(14) \quad &= \frac{\sum_{i=1}^r v_{,i}(\mu_\varphi) d^c \mu_\varphi^i(\xi_j)}{v(\mu_\varphi)} + d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) (\xi_j) \\
&= \frac{\sum_{i=1}^r v_{,i}(\mu_\varphi) g_\varphi(\xi_i, \xi_j)}{v(\mu_\varphi)} + d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) (\xi_j).
\end{aligned}$$

Also, using Cartan's formula and the fact that $\xi_j \lrcorner \omega_\varphi = -d\mu_\varphi^j$ we deduce $\mathcal{L}_{J\xi_j} \omega_\varphi = -dd^c \mu_\varphi^j$, where $\mathcal{L}_{J\xi_j}$ denotes the Lie derivative in the direction $J\xi_j$. And the same holds for ω_0 . It follows from arguments in the proof of [25, Lemma 5]

$$(15) \quad d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) (\xi_j) = \Delta_\varphi \mu_\varphi^j - \Delta_0 \mu_0^j,$$

Multiplying (15) by $v_{,j}(\mu_\varphi)$, summing over $j = 1, \dots, r$ and using (14) we obtain

$$\begin{aligned}
(16) \quad \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_\varphi \mu_\varphi^j &= \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j + v_{,j}(\mu_\varphi) d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) (\xi_j) \\
&= \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j + v_{,j}(\mu_\varphi) d^c F(\xi_j) - \sum_{i,j=1}^r \frac{v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j)}{v(\mu_\varphi)}.
\end{aligned}$$

Injecting in (12), we find

$$\begin{aligned}
\text{Scal}(\omega_\varphi) &= 2\Lambda_\varphi(\text{Ric}(\omega_0)) - \Delta_\varphi F + \frac{1}{v(\mu_\varphi)} \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \\
&\quad - \frac{1}{v(\mu_\varphi)^2} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right) \\
&\quad + \frac{1}{v(\mu_\varphi)} \left(\sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j + v_{,j}(\mu_\varphi) d^c F(\xi_j) - \sum_{i,j=1}^r \frac{v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j)}{v(\mu_\varphi)} \right) \\
(17) \quad &= 2\Lambda_\varphi(\text{Ric}(\omega_0)) - \Delta_\varphi F - \frac{2}{v(\mu_\varphi)^2} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi) v_{,j}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right) \\
&\quad + \frac{1}{v(\mu_\varphi)} \left(\sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j + v_{,j}(\mu_\varphi) d^c F(\xi_j) + \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_\varphi(\xi_i, \xi_j) \right)
\end{aligned}$$

We conclude by injecting (9) and (17) in (7):

$$\begin{aligned}
\text{Scal}_v(\omega_\varphi) &= 2v(\mu_\varphi)\Lambda_\varphi(\text{Ric}(\omega_0)) - v(\mu_\varphi)\Delta_\varphi F + \sum_{j=1}^r v_{,j}(\mu_\varphi)\Delta_0\mu_0^j + v_{,j}(\mu_\varphi)d^c F(\xi_j) \\
&\quad + \sum_{i,j=1}^r v_{,ij}(\mu_\varphi)g_\varphi(\xi_i, \xi_j) - \frac{2}{v(\mu_\varphi)} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi)v_{,j}(\mu_\varphi)g_\varphi(\xi_i, \xi_j) \right) \\
&\quad - 2 \sum_{i,j=1}^r v_{,i}(\mu_\varphi)\Delta_\varphi\mu_\varphi^i - 2 \sum_{i,j=1}^r v_{,ij}(\mu_\varphi)g_\varphi(\xi_i, \xi_j) + \sum_{i,j=1}^r v_{,ij}(\mu_\varphi)g_\varphi(\xi_i, \xi_j) \\
&= 2v(\mu_\varphi)\Lambda_\varphi(\text{Ric}(\omega_0)) - v(\mu_\varphi)\Delta_\varphi F + \sum_{j=1}^r v_{,j}(\mu_\varphi)\Delta_0\mu_0^j + v_{,j}(\mu_\varphi)d^c F(\xi_j) \\
&\quad - 2 \sum_{i=1}^r v_{,i}(\mu_\varphi)\Delta_\varphi\mu_\varphi^i - \frac{2}{v(\mu_\varphi)} \left(\sum_{i,j=1}^r v_{,i}(\mu_\varphi)v_{,j}(\mu_\varphi)g_\varphi(\xi_i, \xi_j) \right) \\
&= v(\mu_\varphi)2\Lambda_\varphi(\text{Ric}(\omega_0)) - v(\mu_\varphi)\Delta_\varphi F - \sum_{j=1}^r v_{,j}(\mu_\varphi)d^c F(\xi_j) - v_{,j}(\mu_\varphi)\Delta_0\mu_0^j \\
&= v(\mu_\varphi)2\Lambda_\varphi(\text{Ric}(\omega_0)) + d^*(v(\mu_\varphi)dF) - \sum_{j=1}^r v_{,j}(\mu_\varphi)\Delta_0\mu_0^j \\
&= v(\mu_\varphi)(-\Delta_{\varphi,v}F + 2\Lambda_{\varphi,v}\text{Ric}(\omega_0)) \\
&= 2v(\mu_\varphi)\Lambda_{\varphi,v}(\text{Ric}_v(\omega_\varphi)),
\end{aligned}$$

where in the third equality we used (16). In the last line we use (56) for the expression of $\Lambda_{\varphi,v}\text{Ric}(\omega_0)$, the definition of the v -weighted Laplacian in (53) and that $\Delta_{\varphi,v}f = \Lambda_{\varphi,v}dd^c f$ (see Appendix A.1). \square

We then conclude this section:

Proposition 3.3. *A \mathbb{T} -invariant Kähler metric is a weighted cscK metric if and only if it is solution of the following system of elliptic PDEs*

$$(18) \quad \begin{cases} F &= \log\left(v(\mu_\varphi)\frac{\omega_\varphi^n}{\omega_0^n}\right) \\ \Delta_{\varphi,v}F &= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi,v}(\text{Ric}(\omega_0)), \end{cases}$$

where $\Delta_{\varphi,v}$ and $\Lambda_{\varphi,v}$ are introduced in Appendix A.

Proof. From the end of the proof of Lemma 3.2 we have that $\frac{1}{v(\mu_\varphi)}\text{Scal}_v(\omega_\varphi) = -\Delta_{\varphi,v}F + 2\Lambda_{\varphi,v}(\text{Ric}(\omega_0))$. Then $\text{Scal}_v(\omega_\varphi) = w(\mu_\varphi)$ if and only if the second equation in (18) is satisfied. \square

4. C^0 -ESTIMATES

The goal of this section is to prove a priori C^0 -estimates for φ and F , solutions of (18) where φ is normalized such that $\sup_X \varphi = 0$.

First, observe that since P is compact and v, w are smooth, there exist positive constants η, L, ν, M such that

$$(19) \quad \begin{aligned} 0 < \eta \leq v \leq L < +\infty, & \quad -\nu \leq w \leq M, \\ -\eta \leq v_{,j} \leq L < +\infty, & \quad -\nu \leq w_{,j} \leq M. \end{aligned}$$

Let ψ be the unique solution of the Monge-Ampère equation

$$\omega_\psi^{[n]} = b^{-1} \sqrt{F^2 + 1} \omega_\varphi^{[n]}, \quad \sup_X \psi = 0,$$

$b := \int_X \sqrt{F^2 + 1} \omega_\varphi^{[n]}$. We normalize ω_0 such that

$$(20) \quad \int_X \omega_0^{[n]} = 1,$$

where we recall that $\omega_0^{[k]} := \frac{\omega_0^k}{k!}$, for $1 \leq k \leq n$. Observe that, since $F^2 + 1 \leq 2F^2$ on $\{F \geq 1\}$,

$$\begin{aligned} 0 < b &= \int_{\{F < 1\}} v(\mu_\varphi)^{-1} e^F \sqrt{F^2 + 1} \omega_0^{[n]} + \int_{\{F \geq 1\}} v(\mu_\varphi)^{-1} e^F \sqrt{F^2 + 1} \omega_0^{[n]} \\ &\leq \sqrt{2} \eta^{-1} (e + \text{Ent}_v(\varphi)) \end{aligned}$$

where $\text{Ent}_v(\varphi) := \int_X \log \left(\frac{v(\mu_\varphi) \omega_\varphi^n}{\omega_0^n} \right) v(\mu_\varphi) \omega_\varphi^{[n]}$ denotes the *weighted entropy*.

We observe that the latter is comparable to the classical entropy $\text{Ent}(\varphi) := \text{Ent}_1(\varphi)$:

Lemma 4.1. *For any weight $v > 0$, the weighted entropy Ent_v is bounded if and only if the entropy Ent is:*

$$\text{Ent}_v(\varphi) < \infty \quad \Leftrightarrow \quad \text{Ent}(\varphi) < \infty,$$

for any $\varphi \in \mathcal{K}(X, \omega_0)^\mathbb{T}$.

Proof. Suppose first that the entropy $\text{Ent}(\varphi)$ is bounded. For simplicity, we write $g := e^F$ and $f := \frac{g}{v(\mu_\varphi)}$. Then $g = \frac{v(\mu_\varphi) \omega_\varphi^{[n]}}{\omega_0^{[n]}}$. We compute

$$\begin{aligned} \text{Ent}_v(\varphi) &= \int_X g \log g \omega_0^{[n]} = \int_X v(\mu_\varphi) f \log(v(\mu_\varphi) f) \omega_0^{[n]} \\ &\leq L \left(\int_X f \log f \omega_0^{[n]} + \log L \int_X f \omega_0^{[n]} \right) \\ &= L \text{Ent}(\varphi) + L \int_X \omega_\varphi^{[n]} \\ &= L \text{Ent}(\varphi) + L \log L, \end{aligned}$$

where we use (20) to pass to the last line and L is defined in (19). The converse is obtain via similar computations. \square

Therefore, if $\text{Ent}(\varphi)$ is uniformly bounded, so is b .

Let $A_0 > 0$ such that $-A_0 \omega_0 \leq \text{Ric}(\omega_0) \leq A_0 \omega_0$. We start with the following key observation:

$$\begin{aligned}
(21) \quad |\Lambda_{v,\varphi} \text{Ric}(\omega_0)| &= \left| \Lambda_\varphi(\text{Ric}(\omega_0)) - \frac{1}{2v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j \right| \\
&\leq A_0 \Lambda_\varphi(\omega_0) + \sup_X \left| \frac{1}{2v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j \right| \\
&\leq A_0 \Lambda_\varphi(\omega_0) + C.
\end{aligned}$$

It is important to note that the constant C above does not depend on φ by (19). Following [16] we show:

Theorem 4.2. *Given $\varepsilon \in (0, 1)$, there exists $C = C(\varepsilon, n, \omega_0, P, w, v, b)$ such that*

$$F + \varepsilon\psi - A\varphi \leq C,$$

where $A > 0$ is a uniform constant depending only on the lower bound of the Ricci curvature.

Proof. We let $H := F + \varepsilon\psi - A\varphi$ with $A = 2A_0 + 1$.

Then

$$\begin{aligned}
\Delta_{\varphi,v} H &= \Delta_{\varphi,v} F + \varepsilon \Delta_{\varphi,v} \psi - A \Delta_{\varphi,v} \varphi \\
&= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi,v} \text{Ric}(\omega_0) + \varepsilon \Delta_\varphi \psi - A \Delta_\varphi \varphi \\
&\quad + \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) (\varepsilon d^c \psi(\xi_j) - A d^c \varphi(\xi_j)) \\
&= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi,v} \text{Ric}(\omega_0) + \varepsilon \frac{dd^c \psi \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} - A \frac{dd^c \varphi \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} \\
&\quad + \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) (\varepsilon d^c \psi(\xi_j) - A d^c \varphi(\xi_j)) \\
&\geq -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} - 2A_0 \Lambda_\varphi(\omega_0) - C + \varepsilon \frac{dd^c \psi \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} - A \frac{dd^c \varphi \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} \\
&\quad + \sum_{j=1}^r \frac{1}{v(\mu_\varphi)} v_{,j}(\mu_\varphi) (\varepsilon d^c \psi(\xi_j) - A d^c \varphi(\xi_j)).
\end{aligned}$$

For the inequality, we use (21). We continue the computation

$$\begin{aligned}
\Delta_{\varphi, \nu} H &\geq -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} - 2A_0 \Lambda_\varphi(\omega_0) - C + \varepsilon \frac{(\omega_\psi - \omega_0) \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} \\
&\quad - A \frac{(\omega_\varphi - \omega_0) \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} + \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) (\varepsilon d^c \psi(\xi_j) - Ad^c \varphi(\xi_j)) \\
&\geq -\frac{M}{L} - An - C + (A - 2A_0 - \varepsilon) \Lambda_\varphi(\omega_0) + \varepsilon \frac{\omega_\psi \wedge \omega_\varphi^{[n-1]}}{\omega_\varphi^{[n]}} \\
&\quad + \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) (\varepsilon d^c \psi(\xi_j) - Ad^c \varphi(\xi_j)) \\
&\geq -C_1 + nb^{-1/n} \varepsilon (F^2 + 1)^{\frac{1}{2n}} + \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \left(\varepsilon (\mu_\psi^j - \mu_0^j) - A (\mu_\varphi^j - \mu_0^j) \right) \\
&\geq -C_1 + C_2 + nb^{-1/n} \varepsilon (F^2 + 1)^{\frac{1}{2n}}
\end{aligned}$$

where C_1 depends on (21), $C_2 := \inf_X \sum_{j=1}^r \frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \left(\varepsilon (\mu_\psi^j - \mu_0^j) - A (\mu_\varphi^j - \mu_0^j) \right)$. Observe that the constant C_2 is indeed independent of φ and ψ since the image of μ_φ is the moment polytope P for any \mathbb{T} -invariant Kähler potential in $[\omega_0]$ (see the introduction and [25, Lemma 1] for details).

From the second inequality to the third, we use the mixed Monge-Ampère inequality (see [9, Proposition 1.11]) ensuring that $\omega_\psi \wedge \omega_\varphi^{n-1} \geq b^{-1/n} (\sqrt{F^2 + 1})^{1/n} \omega_\varphi^n$. We also used that the moment map μ_φ of a Kähler potential φ satisfies $\mu_\varphi^j = \mu_0^j - d^c \varphi(\xi^j)$ for any $\xi^j \in \mathfrak{t}$.

By the maximum principle, applied to H , we can then infer that at a maximum point x_0 we have

$$nb^{-1/n} \varepsilon (F^2 + 1)^{1/2n}(x_0) \leq C.$$

Thus $F(x_0) \leq C_0$, $C_0 = C_0(\varepsilon, A_0, \omega_0, b, v, w)$.

We then claim that

$$\varepsilon \psi - A \varphi \leq C_3,$$

where $C_3 > 0$ depends on ε , A and b . Let us now prove the claim. First of all we observe that, for any $a, \delta \in (0, 1)$ we have either $\sqrt{F^2 + 1} \geq b/(a\delta^n)$ or $F \leq \sqrt{F^2 + 1} \leq b/(a\delta^n)$; thus

$$\begin{aligned}
\omega_\varphi^n &= v(\mu_\varphi) e^F \omega_0^n \leq La\delta^n b^{-1} e^F \sqrt{F^2 + 1} \omega_0^n + Le^{\frac{b}{a\delta^n}} \omega_0^n \\
&= La\delta^n \omega_\psi^n + Le^{\frac{b}{a\delta^n}} \omega_0^n \\
&\leq La\omega_{\delta\psi}^n + Le^{\frac{b}{a\delta^n}} \omega_0^n.
\end{aligned}$$

We are going to apply Theorem 2.3 with $u = \varphi$, $\chi = \delta\psi$ and $f = Le^{b/(a\delta^n)}$. In fact, we have that $e^{b/(a\delta^n)} \in L^p$, for any $p \geq 1$ and, since $\int_X \omega_\varphi^{[n]} = 1$, [13, Theorem 1.3] implies that $P[\varphi] = 0 \geq \delta\psi$. In particular $P[\varphi]$ is less singular than $\delta\psi$. Moreover, the assumption in (6) is satisfied. Indeed, for any Borel set $E \subset X$

$$\int_E f \omega_0^n = Le^{\frac{b}{a\delta^n}} \text{Vol}_{\omega_0}(E) \leq Le^{\frac{b}{a\delta^n}} \exp\left(\frac{-C_4}{\text{Cap}_{\omega_0}(E)^{1/n}}\right) \leq Le^{\frac{b}{a\delta^n}} C_4 \text{Cap}_{\omega_0}(E)^2.$$

The inequality

$$\text{Vol}_{\omega_0}(E) \leq \exp\left(\frac{-C_4}{\text{Cap}_{\omega_0}(E)^{1/n}}\right)$$

where $C_4 > 0$ depends on n and ω_0 , follows from [18, eq. (12.1.3) and Lemma 12.2] (see also [14, Proposition 2.10]). Using that $\text{Cap}_{\omega_0} \leq (1 - \delta)^{-n} \text{Cap}_{(1-\delta)\omega}$ and that $\text{Cap}_{(1-\delta)\omega} \leq \text{Cap}_{\delta\psi}$ (see [26, Lemma 2.7]) we get that

$$\int_E f \omega_0^n \leq C_4 (1 - \delta)^{-2n} \text{Cap}_{\delta\psi}(E)^2.$$

We can then infer that $\varphi \geq \delta\psi - C_4((1 - a)^{-1}, e^{b/a\delta^n}, (1 - \delta)^{-2n})$. Choosing δ small enough so that $\varepsilon - A\delta \geq 0$ we obtain the claim with $C_3 = AC_4$.

It then follows that for any $x \in X$

$$H(x) \leq H(x_0) \leq C_0 + C_3,$$

which concludes the proof. \square

Corollary 4.3. *The functions ψ, φ, F are uniformly bounded by a constant that only depends on n, ω_0, ν, w and $\text{Ent}(\varphi)$.*

Proof. From Theorem 4.2 we know that $F \leq C - \varepsilon\psi + A\varphi \leq C - \varepsilon\psi$, since $\sup_X \varphi = 0$. Therefore

$$\int_X e^{2F} \omega_0^n \leq \tilde{C} \int_X e^{-2\varepsilon\psi} \omega_0^n.$$

Choosing $\varepsilon < \nu_{\omega_0}^{-1}$, by Theorem 2.1 we get a uniform bound for $\|e^F\|_{L^2}$. It follows from Kołodziej uniform estimates (Theorem 2.2), applied to the equation $\omega_\varphi^n = \frac{1}{\nu(\mu_\varphi)} e^F \omega_0^n$, that $\varphi \geq -C(\nu, \|e^F\|_{L^2}, \omega_0)$. In particular, since $\sup_X \varphi = 0$ we do get a uniform control on $\|\varphi\|_{L^\infty}$. Also,

$$\int_X e^{2F} (F^2 + 1) \omega_0^n \leq \int_X e^{4F} \omega_0^n \leq C' \int_X e^{-4\varepsilon\psi} \omega_0^n.$$

Once again, thanks to Theorem 2.1, choosing $\varepsilon \leq (2\nu_{\omega_0})^{-1}$ we get a uniform bound for $\|e^F \sqrt{F^2 + 1}\|_{L^2}$. Theorem 2.2 then gives a uniform control for $\|\psi\|_{L^\infty}$.

We can then conclude from Theorem 4.2 together with the arguments above that

$$F \leq C - \varepsilon\psi + A\varphi \leq -\varepsilon \inf_X \psi \leq C_4$$

for some uniform positive constant C_4 .

It remains to prove a uniform lower bound for F . For this purpose we apply the minimum principle to $F + t\varphi$, with $t = 2A_0 + 1$ where we recall that $A_0 > 0$ is such that $\text{Ric}(\omega_0) \leq A_0\omega_0$. Using Definition (56), Lemma A.2 and (21), we find

$$\begin{aligned}
\Delta_{\varphi, \nu}(F + t\varphi) &= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi, \nu}(\text{Ric}(\omega_0)) + t\Delta_\varphi\varphi + \frac{t}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) d^c\varphi(\xi_j) \\
&\leq -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_\varphi(\text{Ric}(\omega_0)) + \tilde{C} + t(n - \Lambda_\varphi(\omega_0)) + \frac{t}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi)(\mu_\varphi^j - \mu_0^j) \\
&\leq \frac{\nu}{\eta} + tn + (2A_0 - t)\Lambda_\varphi(\omega_0) + \tilde{C} + \frac{t}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi)(\mu_\varphi^j - \mu_0^j) \\
&\leq C - \Lambda_\varphi(\omega_0) \\
&\leq C - \eta^{\frac{1}{n}} n e^{-F/n},
\end{aligned}$$

where $C = \sup_X \frac{1}{v(\mu_\varphi)} \left(t \sum_{j=1}^r v_{,j}(\mu_\varphi)(\mu_\varphi^j - \mu_0^j) \right) + \nu + tn + \tilde{C}$. It is worth to note again that the constant C is indeed independent of φ since the image of μ_φ is the moment polytope P for any \mathbb{T} -invariant Kähler potential in $\varphi \in \mathcal{K}(X, \omega_0)^\mathbb{T}$.

The last inequality follows from the fact that, given two positive $(1,1)$ -forms α, β we have $\Lambda_\beta(\alpha) \geq n \left(\frac{\alpha^n}{\beta^n} \right)^{1/n}$. In particular

$$(22) \quad \Lambda_\varphi(\omega_0) \geq n v(\mu_\varphi)^{1/n} e^{-F/n}.$$

Now, let x_0 be a minimum point of the function $F + t\varphi$, then $0 \leq C - \eta^{\frac{1}{n}} n e^{-F(x_0)/n}$, or equivalently $F(x_0) \geq -n \log(C \eta^{-\frac{1}{n}}/n)$. For any $x \in X$, $F(x) + t\varphi(x) \geq F(x_0) + t\varphi(x_0)$, hence $F \geq -n \log(C \eta^{-\frac{1}{n}}/n) - t \|\varphi\|_{L^\infty}$. \square

5. INTEGRAL C^2 -ESTIMATES

The goal in this section is to show the theorem below.

Theorem 5.1. *Let φ be a solution of (18). Then for any $p \geq 1$; there exists a constant $C > 0$, depending on $p, n, \nu, \|\varphi\|_{C^0}, \|F\|_{C^0}$, an upper bound on the Ricci form of ω_0 and a lower bound of the holomorphic bisectional curvature of ω_0 so that*

$$\|\Lambda_0(\omega_\varphi)\|_{L^p} \leq C.$$

We start with some lemmas which will be useful in the following.

Lemma 5.2. *Let φ be solution of (18). Then*

$$(23) \quad \Lambda_0(\omega_\varphi) \leq C \Lambda_\varphi(\omega_0)^{n-1}, \quad \Lambda_\varphi(\omega_0) \leq C \Lambda_0(\omega_\varphi)^{n-1},$$

where C is a positive constant depending on $\|F\|_{C^0}, \nu$ and n . Moreover,

$$(24) \quad \Lambda_0(\omega_\varphi) C_1 \geq 1 \quad \Lambda_\varphi(\omega_0) C_2 \geq 1,$$

for positive constants C_1, C_2 depending on $\|F\|_{C^0}, \nu$, and n .

Proof. Recall that for any smooth $(1,1)$ -form α, β

$$n \left(\frac{\alpha^n}{\beta^n} \right)^{1/n} \leq \Lambda_\beta(\alpha) \leq n \frac{\alpha^n}{\beta^n} \Lambda_\alpha(\beta)^{n-1}.$$

By (18) and the second inequality above, we infer

$$\Lambda_0(\omega_\varphi) \leq n \frac{e^F}{v(\mu_\varphi)} \Lambda_\varphi(\omega_0)^{n-1}.$$

Similarly $\Lambda_\varphi(\omega_0) \leq nv(\mu_\varphi)e^{-F} \Lambda_0(\omega_\varphi)^{n-1}$.

On the other side, $\Lambda_0(\omega_\varphi) \geq n \left(\frac{e^F}{v(\mu_\varphi)} \right)^{1/n}$ and $\Lambda_\varphi(\omega_0) \geq n (v(\mu_\varphi)e^{-F})^{1/n}$. \square

The following lemma is well-known; however, as it will be used very often, we give a statement below without proof.

Lemma 5.3. *Let S be a covariant 2-tensor S and f_1, f_2 two smooth functions on X . Then*

$$g_\varphi(df_1, g_\varphi(S, df_2)) = S(\nabla^\varphi f_1, \nabla^\varphi f_2)$$

and

$$(25) \quad g_\varphi(S, df_1 \otimes df_2) = S(\nabla^\varphi f_1, \nabla^\varphi f_2).$$

In particular, when $S = \nabla^\varphi dh$, for some smooth function $h : X \rightarrow \mathbb{R}$ we have

$$g_\varphi(df_1, g_\varphi(\nabla^\varphi dh, df_2)) = \nabla^\varphi dh(\nabla^\varphi f_1, \nabla^\varphi f_2)$$

Let us stress that $g_\varphi(S, df_2)$ has to be understood as the 1-form: $V \rightarrow g_\varphi(S(V, \cdot), df_2)$.

Lemma 5.4. *For any smooth (1, 1)-form β we have the following identities between 1-forms*

$$(26) \quad d\Lambda_\varphi(\beta) = \Lambda_\varphi(\mathcal{L}\beta) - g_\varphi(\beta, \mathcal{L}\omega_\varphi), \quad d^c\Lambda_\varphi(\beta) = -\Lambda_\varphi(\mathcal{L}_J\beta) + g_\varphi(\beta, \mathcal{L}_J\omega_\varphi),$$

where $g_\varphi(\beta, \mathcal{L}\omega_\varphi)$ is the 1-form defined as $V \rightarrow g_\varphi(\beta, \mathcal{L}_V\omega_\varphi)$ and $\Lambda_\varphi(\mathcal{L}\beta)$ is the 1-form defined as $V \rightarrow \Lambda_\varphi(\mathcal{L}_V\beta)$.

Proof. By definition of the trace

$$(27) \quad \beta \wedge \omega_\varphi^{[n-1]} = \Lambda_\varphi(\beta)\omega_\varphi^{[n]}.$$

Let V be a vector field. Differentiating the LHS of the above inequality in the direction V we obtain

$$\begin{aligned} \mathcal{L}_V(\beta \wedge \omega_\varphi^{[n-1]}) &= \mathcal{L}_V(\beta) \wedge \omega_\varphi^{[n-1]} + \beta \wedge \mathcal{L}_V(\omega_\varphi) \wedge \omega_\varphi^{[n-2]} \\ &= \Lambda_\varphi(\mathcal{L}_V(\beta))\omega_\varphi^{[n]} + \beta \wedge \mathcal{L}_V(\omega_\varphi) \wedge \omega_\varphi^{[n-2]}. \end{aligned}$$

Differentiating (27) and rearranging the terms, we get

$$\begin{aligned} \mathcal{L}_V(\Lambda_\varphi(\beta))\omega_\varphi^{[n]} &= \Lambda_\varphi(\mathcal{L}_V(\beta))\omega_\varphi^{[n]} + \beta \wedge \mathcal{L}_V(\omega_\varphi) \wedge \omega_\varphi^{[n-2]} - \Lambda_\varphi(\beta)\mathcal{L}_V(\omega_\varphi) \wedge \omega_\varphi^{[n-1]} \\ &= \Lambda_\varphi(\mathcal{L}_V(\beta))\omega_\varphi^{[n]} + \beta \wedge \mathcal{L}_V(\omega_\varphi) \wedge \omega_\varphi^{[n-2]} - \Lambda_\varphi(\beta)\Lambda_\varphi(\mathcal{L}_V(\omega_\varphi))\omega_\varphi^{[n]} \\ &= \Lambda_\varphi(\mathcal{L}_V(\beta))\omega_\varphi^{[n]} - g_\varphi(\beta, \mathcal{L}_V\omega_\varphi)\omega_\varphi^{[n]}, \end{aligned}$$

where to pass to the last line we use the well know identity $\Lambda_\varphi(\beta)\Lambda_\varphi(\alpha)\omega_\varphi^{[n]} = g_\varphi(\beta, \alpha)\omega_\varphi^{[n]} + \beta \wedge \alpha \wedge \omega_\varphi^{[n-2]}$ (see e.g. [17, eq. (1.12.5)]), for any (1, 1)-form α, β .

This conclude the proof of the first identity. The second one simply follows from the first one and the that $d^c f = -df(J\cdot)$ for any smooth function f . \square

The following lemma is a generalization of Yau inequality [29, (2.10)] for F solution of the weighted Monge-Ampère equation (18) and for the weighted Laplacian.

Lemma 5.5. *The following holds true:*

$$(28) \quad \Delta_{\varphi, \mathfrak{v}} \Lambda_0(\omega_\varphi) \geq \Delta_0 F - \Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi)) - C_1 \Lambda_0(\omega_\varphi) \Lambda_\varphi(\omega_0) - C_2,$$

where $C_1 > 0$ depends on ω_0 , $\|F\|_{C^0, \mathfrak{v}, n}$ and $C_2 > 0$ on \mathfrak{v} and a lower bound for the holomorphic bisectional curvature of ω_0 .

In particular,

$$(29) \quad \Delta_{\varphi, \mathfrak{v}} \Lambda_0(\omega_\varphi) \geq \Delta_0 F - \Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi)) - C_1 \Lambda_0(\omega_\varphi)^n - C_2.$$

In the above statement, $g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi)$ is understood as the 2-form $(V, W) \rightarrow g_\varphi(\mathcal{L}_V\omega_\varphi, \mathcal{L}_W\omega_\varphi)$. Recall that in an orthonormal frame $(e_k)_{k=1}^{2n}$ with respect to ω_0 we have that $\Lambda_0(\beta) = \frac{1}{2} \sum_{k=1}^{2n} \beta(e_k, J e_k)$, for any 2-form β . Hence

$$-\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi)) = \frac{1}{2} \sum_{k=1}^{2n} |\mathcal{L}_{e_k}\omega_\varphi|_\varphi^2 \geq 0.$$

Proof. We follow Yau's computations [29]. First, we take the Laplacian with respect to ω_0 of Monge-Ampère equation in (18). We have that

$$\begin{aligned} \Delta_0 \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) &= \Delta_0 \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) = \Lambda_0 dd^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) \\ &= \Lambda_0 d \left(\Lambda_0(\mathcal{L}_J\omega_0) - \Lambda_\varphi(\mathcal{L}_J\omega_\varphi) \right) \\ &= \Lambda_0 \left(\Lambda_0(\mathcal{L}\mathcal{L}_J\omega_0) - g_0(\mathcal{L}\omega_0, \mathcal{L}_J\omega_0) - \Lambda_\varphi(\mathcal{L}\mathcal{L}_J\omega_\varphi) + g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi) \right). \end{aligned}$$

Let us observe that in orthonormal frame $(e_k)_{k=1}^{2n}$ w.r.t. ω_0 we write $-\Lambda_0(\mathcal{L}\mathcal{L}_J\omega_0) = \frac{1}{2} \sum_{k=1}^{2n} \mathcal{L}_{e_k} \mathcal{L}_{e_k} \omega_0$.

The second and the third equalities are due to the fact that by the computation in (15) we have

$$d \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) = \Lambda_\varphi(\mathcal{L}\omega_\varphi) - \Lambda_0(\mathcal{L}\omega_0), \quad d^c \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) = -\Lambda_\varphi(\mathcal{L}_J\omega_\varphi) + \Lambda_0(\mathcal{L}_J\omega_0).$$

The last equality follows from Lemma 5.4.

Recall that by (18), $\Delta_0 \log \left(\frac{\omega_\varphi^n}{\omega_0^n} \right) = \Delta_0(F - \log \mathfrak{v}(\mu_\varphi))$. Then, taking normal coordinate with respect to ω_0 and doing local computations we obtain

$$(30) \quad \begin{aligned} \Delta_0(F - \log \mathfrak{v}(\mu_\varphi)) &= \Lambda_0(\Lambda_0(\mathcal{L}\mathcal{L}_J\omega_0)) - \Lambda_0(\Lambda_\varphi(\mathcal{L}\mathcal{L}_J\omega_\varphi)) \\ &\quad + \Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}_J\omega_\varphi)) - \Lambda_0 g_0(\mathcal{L}\omega_0, \mathcal{L}_J\omega_0). \end{aligned}$$

On the other hand, using (several times) (26) and the fact that ∇^0 preserves ω_0 , we obtain

$$\begin{aligned}
\Delta_{\varphi, v} \Lambda_0(\omega_\varphi) &= \Delta_\varphi \Lambda_0(\omega_\varphi) + g_\varphi(d \log v(\mu_\varphi), d\Lambda_0(\omega_\varphi)) \\
&= \Lambda_\varphi dd^c(\Lambda_0(\omega_\varphi)) + g_\varphi(d \log v(\mu_\varphi), \Lambda_0(\mathcal{L}.\omega_\varphi)) - g_\varphi(d \log v(\mu_\varphi), g_0(\mathcal{L}.\omega_0, \omega_\varphi)) \\
&= \Lambda_\varphi d \left(g_0(\mathcal{L}.\omega_0, \omega_\varphi) - \Lambda_0(\mathcal{L}.\omega_\varphi) \right) + g_\varphi(d \log v(\mu_\varphi), \Lambda_0(\mathcal{L}.\omega_\varphi)) \\
&\quad - g_\varphi(d \log v(\mu_\varphi), g_0(\mathcal{L}.\omega_0, \omega_\varphi)) \\
&= \Lambda_\varphi \left(g_0(\nabla^0 \mathcal{L}.\omega_0, \omega_\varphi) + g_0(\mathcal{L}.\omega_0, \nabla^0 \omega_\varphi) - \Lambda_0(\mathcal{L}.\mathcal{L}.\omega_\varphi) + g_0(\mathcal{L}.\omega_0, \mathcal{L}.\omega_\varphi) \right) \\
&\quad + g_\varphi(d \log v(\mu_\varphi), \Lambda_0(\mathcal{L}.\omega_\varphi)) - g_\varphi(d \log v(\mu_\varphi), g_0(\mathcal{L}.\omega_0, \omega_\varphi)).
\end{aligned}$$

Then, in normal coordinates with respect to ω_0 (where $\nabla^0 = \mathcal{L}$) we have that

$$(31) \quad \Delta_{\varphi, v} \Lambda_0(\omega_\varphi) = \Lambda_\varphi g_0(\mathcal{L}.\mathcal{L}.\omega_0, \omega_\varphi) - \Lambda_\varphi(\Lambda_0(\mathcal{L}.\mathcal{L}.\omega_\varphi)) + g_\varphi(d \log v(\mu_\varphi), \Lambda_0(\mathcal{L}.\omega_\varphi)).$$

Observe that in a orthonormal frame $(e_k)_{k=1}^{2n}$ with respect to ω_0 we have

$$\begin{aligned}
\Lambda_\varphi(\Lambda_0(\mathcal{L}.\mathcal{L}.\omega_\varphi)) &= \frac{1}{2} \Lambda_\varphi \left(\sum_{k=1}^{2n} (\mathcal{L}.\mathcal{L}.\omega_\varphi)(e_k, J e_k) \right) \\
&= \frac{1}{4} \sum_{k,j=1}^{2n} \frac{\mathcal{L}_{e_j} \mathcal{L}_{e_j} \omega_\varphi(e_k, J e_k)}{\omega_\varphi(e_j, J e_j)} \\
&= \frac{1}{4} \sum_{k,j=1}^{2n} \frac{\mathcal{L}_{e_k} \mathcal{L}_{e_k} \omega_\varphi(e_j, J e_j)}{\omega_\varphi(e_k, J e_k)} \\
&= \Lambda_0(\Lambda_\varphi(\mathcal{L}.\mathcal{L}.\omega_\varphi)),
\end{aligned}$$

i.e. the two traces commute.

Combining (30) with (31) and using (18) we get

$$\begin{aligned}
\Delta_{\varphi, v} \Lambda_0(\omega_\varphi) &= \Delta_0(F - \log(v(\mu_\varphi))) + \Lambda_\varphi g_0(\mathcal{L}.\mathcal{L}.\omega_0, \omega_\varphi) + g_\varphi(d \log(v(\mu_\varphi)), \Lambda_0(\mathcal{L}.\omega_\varphi)) \\
&\quad - \Lambda_0(\Lambda_0(\mathcal{L}.\mathcal{L}.\omega_0)) - \Lambda_0(g_\varphi(\mathcal{L}.\omega_\varphi, \mathcal{L}.\omega_\varphi)) \\
(32) \quad &= \Delta_0 F - \Delta_0 \log v(\mu_\varphi) + \Lambda_\varphi g_0(\mathcal{L}.\mathcal{L}.\omega_0, \omega_\varphi) + g_\varphi(d \log(v(\mu_\varphi)), \Lambda_0(\mathcal{L}.\omega_\varphi)) \\
&\quad - \Lambda_0(\Lambda_0(\mathcal{L}.\mathcal{L}.\omega_0)) - \Lambda_0(g_\varphi(\mathcal{L}.\omega_\varphi, \mathcal{L}.\omega_\varphi)).
\end{aligned}$$

The second equality is due to the definition of $\Delta_{0, v}$.

Also, since $-J\xi_j$ is the gradient flow of μ_φ^j w.r.t. g_φ we infer that $g_\varphi(d\mu_\varphi^j, \Lambda_0(\mathcal{L}.\omega_\varphi)) = -\Lambda_0(\mathcal{L}.\xi_j \omega_\varphi)$. Using once again that $\mathcal{L}.\xi_j \omega_\varphi = -dd^c \mu_\varphi^j$ and (10) we get

(33)

$$\begin{aligned}
g_\varphi(d \log v(\mu_\varphi), \Lambda_0(\mathcal{L}\cdot\omega_\varphi)) &= \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) g_\varphi(d\mu_\varphi^j, \Lambda_0(\mathcal{L}\cdot\omega_\varphi)) \\
&= -\frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \Lambda_0(\mathcal{L}J\xi_j\omega_\varphi) \\
&= \frac{1}{v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_\varphi^j \\
&= \frac{1}{v(\mu_\varphi)} \Delta_0 v(\mu_\varphi) - \frac{1}{v(\mu_\varphi)} \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_0(d\mu_\varphi^i, d\mu_\varphi^j) \\
&= \Delta_0 \log v(\mu_\varphi) + \frac{1}{v(\mu_\varphi)^2} |dv(\mu_\varphi)|_0^2 - \frac{1}{v(\mu_\varphi)} \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_0(d\mu_\varphi^i, d\mu_\varphi^j) \\
&= \Delta_0 \log v(\mu_\varphi) + |d \log v(\mu_\varphi)|_0^2 - \frac{1}{v(\mu_\varphi)} \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_0(d\mu_\varphi^i, d\mu_\varphi^j).
\end{aligned}$$

Injecting (33) in (32), we obtain that, in normal coordinates, we have

$$\begin{aligned}
\Delta_{\varphi,v} \Lambda_0(\omega_\varphi) &= \Delta_0 F + |d \log v(\mu_\varphi)|_0^2 \\
&\quad + \Lambda_\varphi g_0(\mathcal{L}\cdot\mathcal{L}J\omega_0, \omega_\varphi) - \frac{1}{v(\mu_\varphi)} \sum_{i,j=1}^r v_{,ij}(\mu_\varphi) g_0(d\mu_\varphi^i, d\mu_\varphi^j) \\
&\quad - \Lambda_0(\Lambda_0(\mathcal{L}\cdot\mathcal{L}J\omega_0)) - \Lambda_0(g_\varphi(\mathcal{L}\cdot\omega_\varphi, \mathcal{L}J\omega_\varphi))
\end{aligned}$$

We bound $g_0(d\mu_\varphi^i, d\mu_\varphi^j)$:

$$\begin{aligned}
g_0(d\mu_\varphi^i, d\mu_\varphi^j) &= \sum_{k=1}^{2n} d\mu_\varphi^i(e_k) d\mu_\varphi^j(e_k) = \sum_{k=1}^{2n} \omega_\varphi(e_k, \xi_i) \omega_\varphi(e_k, \xi_j) \\
&= \sum_{k=1}^{2n} g_\varphi(e_k, J\xi_i) g_\varphi(e_k, J\xi_j) = g_\varphi(J\xi_j, J\xi_i) = g_\varphi(\xi_j, \xi_i) \\
&\leq C_1 \Lambda_0(\omega_\varphi) \leq C_2 \Lambda_\varphi(\omega_0) \Lambda_0(\omega_\varphi).
\end{aligned}$$

where the last inequality follows from (24) and $(e_k)_{k=1}^{2n}$ is an orthonormal frame for ω_0 .

Also, locally, we have $-\Lambda_0(\Lambda_0(\mathcal{L}\cdot\mathcal{L}J\omega_0)) = \frac{1}{4} \sum_{k=1}^{2n} \mathcal{L}_{e_k} \mathcal{L}_{e_k} \omega_0(e_k, J e_k)$ and is uniformly bounded by a constant B depending only on the holomorphic bisectional curvature of ω_0 . For the last term, we use Cauchy-Schwarz inequality and (23)

$$\begin{aligned}
|\Lambda_\varphi g_0(\mathcal{L}\cdot\mathcal{L}J\omega_0, \omega_\varphi)| &\leq |\Lambda_\varphi(\mathcal{L}\cdot\mathcal{L}J\omega_0)|_0 |\omega_\varphi|_0 \\
&\leq C \Lambda_\varphi(\omega_0) \Lambda_0(\omega_\varphi).
\end{aligned}$$

We then get

$$\Delta_{\varphi, \nu} \Lambda_0(\omega_\varphi) \geq \Delta_0 F - \Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi)) - C\Lambda_0(\omega_\varphi)\Lambda_\varphi(\omega_0) + |d \log v(\mu_\varphi)|_0^2 - B,$$

which give (28) since $|d \log v(\mu_\varphi)|_0^2$ is positive. The inequality (29) follows from Lemma 5.2. \square

We now present a generalization of [11, Lemma 2.2] to the weighted setting:

Lemma 5.6. *The following inequality holds true*

$$\Delta_{\varphi, \nu} \log \Lambda_0(\omega_\varphi) \geq \frac{1}{\Lambda_0(\omega_\varphi)} \Delta_0 F - B\Lambda_\varphi(\omega_0),$$

where $B > 0$ depends on $\omega_0, \nu, n, \|F\|_{C^0}$ and a lower bound for the holomorphic bisectional curvature of ω_0 .

Proof. A direct computation (using the explicit expression of the weighted Laplacian given in Lemma A.2) shows that

$$\Delta_{\varphi, \nu} \log \Lambda_0(\omega_\varphi) = \frac{1}{\Lambda_0(\omega_\varphi)} \Delta_{\varphi, \nu} \Lambda_0(\omega_\varphi) - \frac{1}{\Lambda_0(\omega_\varphi)^2} |d\Lambda_0(\omega_\varphi)|_\varphi^2.$$

We focus on the second term of the RHS of the above equality. We let $(e_k)_{k=1}^{2n}$ be a orthonormal frame with respect to ω_0 . By Lemma 5.4 it then follows that in these coordinates

$$|d\Lambda_0(\omega_\varphi)|_\varphi^2 = |\Lambda_0(\mathcal{L}\omega_\varphi)|_\varphi^2,$$

Moreover,

$$\begin{aligned} |\Lambda_0(\mathcal{L}\omega_\varphi)|_\varphi^2 &= \left| \frac{1}{2} \sum_{k=1}^n \mathcal{L}\omega_\varphi(e_k, Je_k) \right|_\varphi^2 \\ &= \frac{1}{4} g_\varphi \left(\sum_{k=1}^n \mathcal{L}\omega_\varphi(e_k, Je_k), \sum_{\ell=1}^n \mathcal{L}\omega_\varphi(e_\ell, Je_\ell) \right) \\ &= \frac{1}{4} \sum_{k,j=1}^n \frac{(\mathcal{L}_{e_j} \omega_\varphi(e_k, Je_k))^2}{\omega_\varphi(e_j, Je_j)} \\ &= \frac{1}{4} \sum_{i,j,k=1}^n \frac{\omega_\varphi(e_i, Je_i) (\mathcal{L}_{e_j} \omega_\varphi(e_k, Je_k))^2}{\omega_\varphi(e_i, Je_i) \omega_\varphi(e_j, Je_j)} \\ &\leq \frac{1}{2} \sum_{p=1}^n \omega_\varphi(e_p, Je_p) \frac{1}{2} \sum_{i,j,k=1}^n \frac{(\mathcal{L}_{e_j} \omega_\varphi(e_k, Je_k))^2}{\omega_\varphi(e_j, Je_j) \omega_\varphi(e_i, Je_i)} \\ &= \Lambda_0(\omega_\varphi) \frac{1}{2} \left(\sum_{k=1}^n |\mathcal{L}_{e_k} \omega_\varphi|_\varphi^2 \right). \end{aligned}$$

By (28) we have

$$\Delta_{\varphi, \nu} \Lambda_0(\omega_\varphi) \geq \Delta_0 F + \frac{1}{2} \sum_{k=1}^{2n} |\mathcal{L}_{e_k} \omega_\varphi|_\varphi^2 - C_1 \Lambda_0(\omega_\varphi) \Lambda_\varphi(\omega_0) - C_2.$$

We then get

$$\begin{aligned}
\Delta_{\varphi, \nu} \log \Lambda_0(\omega_\varphi) &\geq \frac{1}{\Lambda_0(\omega_\varphi)} \left(\Delta_0 F + \frac{1}{2} \sum_{k=1}^{2n} |\mathcal{L}_{e_k} \omega_\varphi|_\varphi^2 - C_1 \Lambda_0(\omega_\varphi) \Lambda_\varphi(\omega_0) - C_2 \right) \\
&\quad - \frac{1}{2\Lambda_0(\omega_\varphi)} \left(\sum_{k=1}^{2n} |\mathcal{L}_{e_k} \omega_\varphi|_\varphi^2 \right) \\
&\geq \frac{1}{\Lambda_0(\omega_\varphi)} \Delta_0 F - C_1 \Lambda_\varphi(\omega_0) - \frac{C_2}{\Lambda_0(\omega_\varphi)} \\
&\geq \frac{1}{\Lambda_0(\omega_\varphi)} \Delta_0 F - (C_1 + C_2) \Lambda_\varphi(\omega_0),
\end{aligned}$$

where in the last inequality we used that $\Lambda_\varphi(\omega_0) \Lambda_0(\omega_\varphi) \geq 1$. □

We are now ready to give the proof of Theorem 5.1:

proof of Theorem 5.1. Consider

$$u := e^{-\gamma(F+\lambda\varphi)} \Lambda_0(\omega_\varphi) > 0,$$

where $\gamma, \lambda > 1$ are uniform constants to be chosen in a suitable way in the following. The ν -Laplacian of u is

$$\begin{aligned}
\Delta_{\varphi, \nu} u &= \Delta_{\varphi, \nu} e^{\log u} \\
&= \Delta_\varphi e^{\log u} + g_\varphi \left(d \log \nu(\mu_\varphi), d \left(e^{\log u} \right) \right) \\
&= \Lambda_\varphi \left(\frac{e^{\log u}}{u^2} du \wedge d^c u \right) + e^{\log u} \Delta_{\varphi, \nu} \log u \\
&= \frac{e^{\log(u)}}{u^2} |du|_\varphi^2 + e^{\log u} \Delta_{\varphi, \nu} \log u \\
&\geq e^{\log u} \Delta_{\varphi, \nu} \log u \\
&= -\gamma u \Delta_{\varphi, \nu} (F + \lambda\varphi) + u \Delta_{\varphi, \nu} (\log \Lambda_0(\omega_\varphi)).
\end{aligned}$$

From (18), the fact that $\text{Ric}(\omega_0) \leq A_0 \omega_0$ and that by (54) $\Lambda_{\varphi, \nu}(\omega_\varphi) \leq n + C$ we deduce that

$$\begin{aligned}
\Delta_{\varphi, \nu} (F + \lambda\varphi) &= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi, \nu}(\text{Ric}(\omega_0)) + \lambda \Lambda_{\varphi, \nu} dd^c \varphi \\
&= -\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi, \nu}(\text{Ric}(\omega_0)) + \lambda (\Lambda_{\varphi, \nu}(\omega_\varphi) - \Lambda_{\varphi, \nu}(\omega_0)) \\
&\leq C_1 + (2A_0 - \lambda) \Lambda_\varphi(\omega_0) + \lambda n + C_2 \\
&= C_3 + (2A_0 - \lambda) \Lambda_\varphi(\omega_0)
\end{aligned}$$

Thus, combining the above inequalities together with Lemma 5.6 leads us to:

$$\Delta_{\varphi, \nu} u \geq e^{-\gamma(F+\lambda\varphi)} \left(-\gamma C_3 \Lambda_0(\omega_\varphi) + \Delta_0 F + (\lambda\gamma - 2A_0\gamma - B) \Lambda_0(\omega_\varphi) \Lambda_\varphi(\omega_0) \right).$$

Using (18), (the proof of) Lemma 5.2 and the fact that $n \geq 2$, we have

$$\Lambda_0(\omega_\varphi) \Lambda_\varphi(\omega_0) \geq n^{-\frac{1}{n-1}} e^{-\frac{F}{n-1}} \eta^{\frac{1}{n-1}} \Lambda_0(\omega_\varphi)^{1+\frac{1}{n-1}} := a(n, \eta) e^{-\frac{F}{n-1}} \Lambda_0(\omega_\varphi)^{1+\frac{1}{n-1}},$$

where η is defined in (19).

We now choose $\lambda \geq 4 \max(2A_0, B)$ (in order to have $\lambda\gamma - 2A_0\gamma - B \geq \frac{\lambda\gamma}{2}$) so that

$$(34) \quad \Delta_{\varphi, \nu} u \geq -\gamma C_3 u + \frac{\lambda\gamma}{2} a(n, \eta) e^{-\frac{F}{n-1}} \Lambda_0(\omega_\varphi)^{\frac{1}{n-1}} u + e^{-\gamma(F+\lambda\varphi)} \Delta_0 F.$$

Now, since $|du|_\varphi^2 \Lambda_0 \omega_\varphi \geq |du|_0^2$ holds pointwise, we write

$$\frac{1}{2p+1} \Delta_{\varphi, \nu} u^{2p+1} = u^{2p} \Delta_{\varphi, \nu} u + 2pu^{2p-1} |du|_\varphi^2 \geq u^{2p} \Delta_{\varphi, \nu} u + 2pu^{2p-2} e^{-\gamma(F+\lambda\varphi)} |du|_0^2.$$

Thus, combining the above inequality with (34), and since $\Delta_{\varphi, \nu}$ is self-adjoint w.r.t. to $\nu(\mu_\varphi)\omega_\varphi^n$ (Lemma A.1), we get

$$(35) \quad \begin{aligned} 0 &= \frac{1}{2p+1} \int_X \Delta_{\varphi, \nu} u^{2p+1} \nu(\mu_\varphi)\omega_\varphi^{[n]} \geq 2p \int_X u^{2p-2} |du|_0^2 e^{-\gamma(F+\lambda\varphi)+F} \omega_0^{[n]} - \gamma C_3 \int_X u^{2p+1} e^F \omega_0^{[n]} \\ &\quad + \frac{\gamma\lambda}{2} a(n, \eta) \int_X u^{2p+1} e^{\left(\frac{n-2}{n-1}\right)F} \Lambda_0(\omega_\varphi)^{\frac{1}{n-1}} \omega_0^{[n]} \\ &\quad + \int_X u^{2p} e^{-\gamma(F+\lambda\varphi)+F} \Delta_0 F \omega_0^{[n]}. \end{aligned}$$

Next, we focus on finding a suitable lower bound for the last term involving the Laplacian of F . This estimate goes as in [16] but we show it for completeness and for reader's convenience.

Set $G := (1-\gamma)F - \gamma\lambda\varphi$. A formal trick gives that

$$\begin{aligned} I &:= - \int_X u^{2p} \Delta_0 F e^G \omega_0^{[n]} \\ &= \frac{1}{\gamma-1} \int_X u^{2p} \Delta_0 G e^G \omega_0^{[n]} + \frac{\gamma\lambda}{\gamma-1} \int_X u^{2p} \Delta_0 \varphi e^G \omega_0^{[n]} \\ &:= I_1 + I_2. \end{aligned}$$

Integration by part gives

$$(36) \quad \begin{aligned} I_1 &= -\frac{1}{\gamma-1} \int_X u^{2p} |dG|_0^2 e^G \omega_0^{[n]} - \frac{2p}{\gamma-1} \int_X u^{2p-1} e^G du \wedge d^c G \wedge \omega_0^{[n-1]} \\ &\leq -\frac{1}{2(\gamma-1)} \int_X u^{2p} |dG|_0^2 e^G \omega_0^{[n]} + \frac{2p^2}{\gamma-1} \int_X u^{2p-2} |du|_0^2 e^G \omega_0^{[n]} \\ &\leq \frac{2p^2}{\gamma-1} \int_X u^{2p-2} |du|_0^2 e^G \omega_0^{[n]}, \end{aligned}$$

where in the first inequality we used the fact that

$$\left| 2pu^{2p-1} \frac{du \wedge d^c G \wedge \omega_0^{[n-1]}}{\omega_0^{[n]}} \right| \leq \frac{(2p)^2}{2} u^{2p-2} |du|_0^2 + \frac{1}{2} u^{2p} |dG|_0^2,$$

by Young's inequality. Also,

$$(37) \quad \begin{aligned} I_2 &= \frac{\gamma\lambda}{\gamma-1} \int_X u^{2p} (\Lambda_0(\omega_\varphi) - n) e^G \omega_0^{[n]} \\ &\leq \frac{\gamma\lambda}{\gamma-1} \int_X u^{2p} \Lambda_0(\omega_\varphi) e^G \omega_0^{[n]} = \frac{\gamma\lambda}{\gamma-1} \int_X u^{2p+1} e^F \omega_0^{[n]}. \end{aligned}$$

Suppose $p > 1$. Then Combining (35), (36), (37) and choosing γ big enough (say $\gamma = ap$, with $a \gg 1$) we obtain

$$\begin{aligned}
(38) \quad 0 &\geq 2 \left(p - \frac{p^2}{\gamma - 1} \right) \int_X u^{2p-2} |du|_0^2 e^G \omega_0^{[n]} - \gamma \left(C_3 + \frac{\lambda}{\gamma - 1} \right) \int_X u^{2p+1} e^F \omega_0^{[n]} \\
&\quad + \frac{\gamma \lambda}{2} a(n, \eta) \int_X u^{2p+1} e^{\left(\frac{n-2}{n-1}\right)F} \Lambda_0(\omega_\varphi)^{\frac{1}{n-1}} \omega_0^{[n]} \\
&\geq -C_4 \int_X \Lambda_0(\omega_\varphi)^{2p+1} \omega_0^{[n]} + C_5 \int_X \Lambda_0(\omega_\varphi)^{2p+1+\frac{1}{n-1}} \omega_0^{[n]},
\end{aligned}$$

where the constant $C_4, C_5 > 0$ depends on $\|F\|_{C^0}$ and $\|\varphi\|_{C^0}$. Observe that in (38), the choice of γ ensures that $p - \frac{p^2}{\gamma-1} > 0$. Using Hölder inequality (with $r = (2p+n)/(n-1)$ and s its conjugate) we can conclude that

$$\|\Lambda_0(\omega_\varphi)\|_{L^{2p+1+\frac{1}{n-1}}}^{2p+1+\frac{1}{n-1}} \leq C \|\Lambda_0(\omega_\varphi)\|_{L^{2p+1}}^{2p+1} \leq C' \|\Lambda_0(\omega_\varphi)\|_{L^{2p+1+\frac{1}{n-1}}}^{2p+1}.$$

This gives the statement for $p > 3$, hence for $p \geq 1$ thanks to Hölder inequality. \square

6. C^2 -ESTIMATES

The goal to this section is to prove the theorem below.

Theorem 6.1. *Let φ be a solution of (18). Then there exists a positive constant C depending on $\omega_0, v, w, \|F\|_{C^0}$ and $\|\varphi\|_{C^0}$ such that*

$$\max_X (|dF|_\varphi^2 + \Lambda_0(\omega_\varphi)) \leq C.$$

We begin by a technical lemma:

Lemma 6.2. *For any smooth function f on X we have*

$$dd^c f(\cdot, J\cdot) = 2\nabla^{\varphi,+} df(\cdot, \cdot),$$

where $\nabla^{\varphi,+} \alpha$ for a 1-form α is defined by

$$\nabla_V^{\varphi,\pm} \alpha(W) := \frac{1}{2} (\nabla_V^\varphi \alpha(W) \pm \nabla_{JV}^\varphi \alpha(JW)).$$

Let us stress that the above identity holds for the Levi-Civita connection ∇^φ associated to any Kähler metric ω_φ . Observe as well that $\nabla^{\varphi,+}$ and $\nabla^{\varphi,-}$ are orthogonal w.r.t. g_φ .

Proof. Let V, W be two vector fields, we compute

$$\begin{aligned}
dd^c f(V, W) &= \nabla_V^\varphi d^c f(W) - \nabla_W^\varphi d^c f(V) \\
&= (\nabla_V^\varphi Jdf)(W) - (\nabla_W^\varphi Jdf)(V) \\
&= (J\nabla_V^\varphi df)(W) - (J\nabla_W^\varphi df)(V) \\
&= -(\nabla_V^\varphi df)(JW) + (\nabla_W^\varphi df)(JV),
\end{aligned}$$

where we use that $\nabla^\varphi J = 0$. It follows that

$$dd^c f(V, JW) = -\nabla_V^\varphi df(J^2W) + \nabla_{JV}^\varphi df(JW) = 2\nabla_V^{\varphi,+} df(W).$$

\square

The first main step to prove Theorem 6.1 is this key Proposition:

Proposition 6.3. *Let $u := e^{\frac{F}{2}}|dF|_{\varphi}^2 + K\Lambda_0(\omega_{\varphi})$. Then there exists positive constants K and C depending on ω_0 , n , v , w , $\|F\|_{C^0}$ and $\|\varphi\|_{C^0}$ and an upper bound for the Ricci curvature, such that the function u satisfies the following differential inequality*

$$\Delta_{\varphi,v}u \geq -C\Lambda_0(\omega_{\varphi})^{3n-3}u.$$

As a remark, we note that u is uniformly bounded. Indeed, by the proof of Lemma 5.2

$$u \geq K\Lambda_0(\omega_{\varphi}) \geq Kn \left(\frac{e^F}{v(\mu_{\varphi})} \right)^{1/n} \geq \frac{nK}{L^{1/n}} e^{\frac{-\|F\|_{C^0}}{n}},$$

where we recall that $v \leq L$.

Our first step to show Proposition 6.3 is to bound from below uniformly the weighted Laplacian of $e^{F/2}|dF|_{\varphi}^2$.

Lemma 6.4. *The following inequality holds*

(39)

$$\Delta_{\varphi,v} \left(e^{F/2}|dF|_{\varphi}^2 \right) \geq -C_1 \left(-\Lambda_0(g_{\varphi}(\mathcal{L}\omega_{\varphi}, \mathcal{L}J\omega_{\varphi})) + \Lambda_0(\omega_{\varphi})^{3n-3}|dF|_{\varphi}^2 + 1 \right) + C_2|\nabla^{\varphi,+}dF|_{\varphi}^2,$$

where C_1 is a positive constant depending on $\omega_0, n, \|F\|_{C^0}$, an upper bound for the Ricci curvature, the bounds for v, w and their derivatives, while $C_2 > 0$ depends only on $\|F\|_{C^0}$.

We recall that $g_{\varphi}(\mathcal{L}\omega_{\varphi}, \mathcal{L}J\omega_{\varphi})$ is understood as the 2-form $(V, W) \rightarrow g_{\varphi}(\mathcal{L}_V\omega_{\varphi}, \mathcal{L}_{JW}\omega_{\varphi})$. In particular, in a orthonormal frame $(e_k)_{k=1}^{2n}$ with respect to ω_0

$$-\Lambda_0(g_{\varphi}(\mathcal{L}\omega_{\varphi}, \mathcal{L}J\omega_{\varphi})) = \frac{1}{2} \sum_{k=1}^{2n} |\mathcal{L}_{e_k}\omega_{\varphi}|_{\varphi}^2 \geq 0.$$

Proof. We compute

$$\begin{aligned} e^{-F/2}\Delta_{\varphi,v} \left(e^{F/2}|dF|_{\varphi}^2 \right) &= \Delta_{\varphi,v} (|dF|_{\varphi}^2) + 2g_{\varphi}(d(F/2), d|dF|_{\varphi}^2) + |dF|_{\varphi}^2 e^{-F/2}\Delta_{\varphi,v} \left(e^{F/2} \right) \\ (40) \qquad \qquad \qquad &= \Delta_{\varphi} (|dF|_{\varphi}^2) + g_{\varphi}(d \log v(\mu_{\varphi}), d|dF|_{\varphi}^2) + g_{\varphi}(dF, d|dF|_{\varphi}^2) \\ &\quad + \frac{1}{2}|dF|_{\varphi}^2 \Delta_{\varphi,v} F + \frac{1}{4}|dF|_{\varphi}^4, \end{aligned}$$

where $|dF|_{\varphi}^2 := g_{\varphi}(dF, dF)$. In the above we used (11) for the first identity and Lemma A.2 to pass to the second line. We now expand each term.

Using the fact that the metric g_{φ} is parallel w.r.t. ∇_{φ} and the expression for the adjoint operator given in [17, eq. 1.10.13], we arrive at

$$\begin{aligned} \Delta_{\varphi} (|dF|_{\varphi}^2) &= -d_{\varphi}^* d (|dF|_{\varphi}^2) \\ &= -2d_{\varphi}^* (g_{\varphi}(\nabla^{\varphi}dF, dF)) \\ &= 2 \sum_{k=1}^{2n} \nabla_{e_k}^{\varphi} (g_{\varphi}(\nabla_{e_k}^{\varphi}dF, dF)) \\ &= -2g_{\varphi}(d_{\varphi}^* (\nabla^{\varphi}dF), dF) + 2|\nabla^{\varphi}dF|_{\varphi}^2 \\ &= 2g_{\varphi}(d\Delta_{\varphi}F, dF) + 2\text{ric}_{\varphi}(\nabla^{\varphi}F, \nabla^{\varphi}F) + 2|\nabla^{\varphi}dF|_{\varphi}^2, \end{aligned}$$

where in the fifth line we used the Bochner formula $-d_\varphi^*(\nabla^\varphi dF) = \Delta_\varphi(dF) + \text{ric}_\varphi(\nabla^\varphi F, \cdot)$ (cf [17, eq. 1.22.1]). Here $\text{ric}_\varphi(\cdot, \cdot)$ denotes the Ricci symmetric 2-tensor of g_φ .

For the second term in (40), by Lemma 5.3 we have

$$\begin{aligned} g_\varphi(d \log v(\mu_\varphi), d|dF|_\varphi^2) &= 2g_\varphi(d \log v(\mu_\varphi), g_\varphi(\nabla^\varphi dF, dF)) \\ &= 2g_\varphi(dF, g_\varphi(d \log v(\mu_\varphi), \nabla^\varphi dF)) \\ &= 2g_\varphi(dF, d(g_\varphi(d \log v(\mu_\varphi), dF))) - 2g_\varphi(dF, g_\varphi(\nabla^\varphi d \log v(\mu_\varphi), dF)) \\ &= 2g_\varphi(dF, d(g_\varphi(d \log v(\mu_\varphi), dF))) - 2\nabla^\varphi d \log v(\mu_\varphi)(\nabla^\varphi F, \nabla^\varphi F). \end{aligned}$$

By the fact that $d|dF|_\varphi^2 = d(g_\varphi(dF, dF)) = 2g_\varphi(\nabla^\varphi dF, dF)$ and again by Lemma 5.3, the next term in (40) is

$$g_\varphi(dF, d|dF|_\varphi^2) = 2(\nabla^\varphi dF)(\nabla^\varphi F, \nabla^\varphi F).$$

Moreover, by definition of $\Delta_{\varphi, v}$ we have

$$g_\varphi(d\Delta_\varphi F, dF) + g_\varphi(d(g_\varphi(d \log v(\mu_\varphi), dF)), dF) = g_\varphi(d\Delta_{\varphi, v} F, dF).$$

Substituting back and decomposing $\nabla^\varphi dF$ as a sum of J -invariant part $\nabla^{\varphi, +} dF$ and J -anti-invariant part $\nabla^{\varphi, -} dF$ we obtain

$$\begin{aligned} e^{-F/2} \Delta_{\varphi, v} \left(e^{F/2} |dF|_\varphi^2 \right) &= 2g_\varphi(d\Delta_{\varphi, v} F, dF) + 2(\text{ric}_\varphi - \nabla^\varphi d \log v(\mu_\varphi))(\nabla^\varphi F, \nabla^\varphi F) \\ &\quad + 2|\nabla^{\varphi, +} dF|_\varphi^2 + 2|\nabla^{\varphi, -} dF|_\varphi^2 \\ &\quad + 2\nabla^{\varphi, +} dF(\nabla^\varphi F, \nabla^\varphi F) + 2\nabla^{\varphi, -} dF(\nabla^\varphi F, \nabla^\varphi F) \\ &\quad + \left(\frac{1}{2} \Delta_{\varphi, v} F + \frac{1}{4} |dF|_\varphi^2 \right) |dF|_\varphi^2. \end{aligned}$$

Notice that by (25) we have

$$\begin{aligned} 0 &\leq 2|\nabla^{\varphi, -} dF + \frac{1}{2}(dF \otimes dF)^-|_\varphi^2 \\ &= 2|\nabla^{\varphi, -} dF|_\varphi^2 + \frac{1}{8}(|dF|_\varphi^4 + |d^c F|_\varphi^4) \\ &\quad + (\nabla^{\varphi, -} dF(\nabla^\varphi F, \nabla^\varphi F) - \nabla^{\varphi, -} dF(J\nabla^\varphi F, J\nabla^\varphi F)) \\ &= 2|\nabla^{\varphi, -} dF|_\varphi^2 + \frac{1}{4}|dF|_\varphi^4 + 2\nabla^{\varphi, -} dF(\nabla^\varphi F, \nabla^\varphi F), \end{aligned}$$

where $(\alpha \otimes \beta)^-(X, Y) := \frac{1}{2}(\alpha(X)\beta(Y) - \alpha(JX)\beta(JY))$ and we use that $J\nabla^\varphi F$ is the dual of $d^c F$ with respect to g_φ . We then get

$$\begin{aligned} (41) \quad e^{-F/2} \Delta_{\varphi, v} \left(e^{F/2} |dF|_\varphi^2 \right) &\geq 2g_\varphi(d\Delta_{\varphi, v} F, dF) + 2|\nabla^{\varphi, +} dF|_\varphi^2 \\ &\quad + 2(\text{ric}_\varphi + \nabla^{\varphi, +} dF - \nabla^\varphi d \log v(\mu_\varphi))(\nabla^\varphi F, \nabla^\varphi F) \\ &\quad + \frac{1}{2} \Delta_{\varphi, v} F |dF|_\varphi^2. \end{aligned}$$

Letting $\text{Ric}(\omega_\varphi)$ be the corresponding Ricci form of ω_φ , from (18) we get

$$\text{Ric}(\omega_\varphi) - \frac{1}{2} dd^c \log v(\mu_\varphi) = \text{Ric}(\omega_0) - \frac{1}{2} dd^c F.$$

Composing with the complex structure J and applying Lemma 6.2 we deduce that

$$\text{ric}_0 = \text{ric}_\varphi + \nabla^{\varphi,+} dF - \nabla^{\varphi,+} d \log v(\mu_\varphi)$$

Substituting back in (41) we obtain

$$(42) \quad e^{-F/2} \Delta_{\varphi,v} \left(e^{F/2} |dF|_\varphi^2 \right) \geq 2g_\varphi(d\Delta_{\varphi,v} F, dF) + 2\text{ric}_0(\nabla^\varphi F, \nabla^\varphi F) + \frac{1}{2} \Delta_{\varphi,v} F |dF|_\varphi^2 \\ + 2|\nabla^{\varphi,+} dF|_\varphi^2 - 2\nabla^{\varphi,-} d \log v(\mu_\varphi)(\nabla^\varphi F, \nabla^\varphi F)$$

We give a bound for the last term of (42). A direct computation shows that

$$\begin{aligned} \nabla^{\varphi,-} d \log v(\mu_\varphi) &= \sum_{j=1}^r \nabla^{\varphi,-} \left(\frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} d\mu_\varphi^j \right) \\ &= \sum_{j=1}^r \left(d \left(\frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \right) \otimes d\mu_\varphi^j + \frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \nabla^\varphi d\mu_\varphi^j \right)^- \\ &= \sum_{i,j=1}^r \left(\frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \right)_{,i} (d\mu_\varphi^i \otimes d\mu_\varphi^j)^- + \sum_{j=1}^r \frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \nabla^{\varphi,-} d\mu_\varphi^j \\ &= \sum_{i,j=1}^r \left(\frac{v_{,j}(\mu_\varphi)}{v(\mu_\varphi)} \right)_{,i} (d\mu_\varphi^i \otimes d\mu_\varphi^j)^- \end{aligned}$$

To pass to the last line, we use [17, Lemma 1.23.2] which insures that $\nabla^{\varphi,-} d\mu_\varphi^j = 0$ since μ_φ^j is Killing potential. On the other hand we have

$$\begin{aligned} |(d\mu_\varphi^i \otimes d\mu_\varphi^j)^- (\nabla^\varphi F, \nabla^\varphi F)| &= \frac{1}{2} |(d^c F(\xi_i) d^c F(\xi_j) - dF(\xi_i) dF(\xi_j))| \\ &= \frac{1}{2} |(g_0(J\nabla^0 F, \xi_i) g_0(J\nabla^0 F, \xi_j) - g_0(\nabla^0 F, \xi_i) g_0(\nabla^0 F, \xi_j))| \\ &\leq C |dF|_0^2 \\ &\leq C_1 |dF|_\varphi^2 \Lambda_0(\omega_\varphi). \end{aligned}$$

In the above we used that $d\mu_\varphi^i(\nabla^\varphi F) = -g_\varphi(J\xi_i, \nabla^\varphi F) = -dF(J\xi_i) = d^c F(\xi_i)$. It follows that

$$|\nabla^{\varphi,-} d \log v(\mu_\varphi)(\nabla^\varphi F, \nabla^\varphi F)| \leq C_2 |dF|_\varphi^2 \Lambda_0(\omega_\varphi).$$

Notice that

$$(43) \quad \begin{aligned} |\text{ric}_0(\nabla^\varphi F, \nabla^\varphi F)| &= |g_\varphi(\text{ric}_0, dF \otimes dF)| \\ &\leq |\text{Ric}(\omega_0)|_\varphi |dF|_\varphi^2 \\ &\leq A_0 |dF|_\varphi^2 \Lambda_\varphi(\omega_0) \\ &\leq A_1 |dF|_\varphi^2 \Lambda_0(\omega_\varphi)^{3n-3}. \end{aligned}$$

Let us stress that in the second line we apply Cauchy-Schwarz inequality for 2-tensors and we end up the norm of the (1,1)-form $\text{Ric}(\omega_0)$ since

$$|\text{ric}_0|_\varphi = |\text{ric}_0(\cdot, J\cdot)|_\varphi = |\text{Ric}(\omega_0)|_\varphi,$$

where we use that J is preserving the metric g_φ .

In the third inequality we used that $\text{Ric}(\omega_0) \leq A_0\omega_0$ and the well known inequality (see e.g. [17, (1.12.5)])

$$|\omega_0|_\varphi^2 = \Lambda_\varphi(\omega_0)^2 - \frac{\omega_0^2 \wedge \omega_\varphi^{[n-2]}}{\omega_0^{[n]}} \leq \Lambda_\varphi(\omega_0)^2.$$

The last inequality follows from (23) and (22). Also,

$$\begin{aligned} \Delta_{\varphi, \mathbf{v}} F |dF|^2 &= \left(-\frac{\mathbf{w}(\mu_\varphi)}{\mathbf{v}(\mu_\varphi)} + 2\Lambda_{\varphi, \mathbf{v}}(\text{Ric}(\omega_0)) \right) |dF|_\varphi^2 \\ (44) \quad &\geq -C \left(\left\| \frac{\mathbf{w}}{\mathbf{v}} \right\|_{C^0} + \Lambda_\varphi(\omega_0) + \tilde{C} \right) |dF|_\varphi^2 \\ &\geq -C' \left(\left\| \frac{\mathbf{w}}{\mathbf{v}} \right\|_{C^0} + \Lambda_0(\omega_\varphi)^{3n-3} + \tilde{C}' \right) |dF|_\varphi^2 \end{aligned}$$

where C, C' depend only on $\omega, \text{Ric}(\omega_0), \mathbf{v}$, and \tilde{C} is the constant which appears in (21). The last line, follows again from (22) and (23).

Using the second equation of (18) and (56) we obtain

$$\begin{aligned} g_\varphi(d\Delta_{\varphi, \mathbf{v}} F, dF) &= -g_\varphi \left(d \left(\frac{\mathbf{w}(\mu_\varphi)}{\mathbf{v}(\mu_\varphi)} \right), dF \right) + 2g_\varphi(d\Lambda_{\varphi, \mathbf{v}}(\text{Ric}(\omega_0)), dF) \\ (45) \quad &= -g_\varphi \left(d \left(\frac{\mathbf{w}(\mu_\varphi)}{\mathbf{v}(\mu_\varphi)} + \frac{1}{2\mathbf{v}(\mu_\varphi)} \sum_{j=1}^r \mathbf{v}_{,j}(\mu_\varphi) \Delta_0 \mu_0^j \right), dF \right) + \\ &\quad + 2g_\varphi(d\Lambda_\varphi(\text{Ric}(\omega_0)), dF). \end{aligned}$$

We consider the last term of the above equality. From (26) we get that

$$(46) \quad g_\varphi(d\Lambda_\varphi(\text{Ric}(\omega_0)), dF) = g_\varphi(\Lambda_\varphi(\mathcal{L}.\text{Ric}(\omega_0)), dF) - g_\varphi(g_\varphi(\text{Ric}(\omega_0), \mathcal{L}.\omega_\varphi), dF)$$

We bound the first term of (46). By Cauchy-Swcharz inequality we obtain that

$$\begin{aligned} g_\varphi(\Lambda_\varphi(\mathcal{L}.\text{Ric}(\omega_0)), dF) &\leq |\Lambda_\varphi(\mathcal{L}.\text{Ric}(\omega_0))|_\varphi |dF|_\varphi \\ &\leq C_1 \sqrt{\Lambda_\varphi(\omega_0)^3} |dF|_\varphi \\ (47) \quad &\leq \frac{1}{2} \Lambda_\varphi(\omega_0)^3 |dF|_\varphi^2 + \frac{C_1}{2} \\ &\leq \frac{C}{2} \Lambda_0(\omega_\varphi)^{3n-3} |dF|_\varphi^2 + \frac{C_1}{2}. \end{aligned}$$

The second inequality follows from the following local computations

$$\begin{aligned}
|\Lambda_\varphi(\mathcal{L}.\text{Ric}(\omega_0))|_\varphi^2 &= \left| \frac{1}{2} \sum_{j=1}^{2n} \frac{\mathcal{L}.\text{Ric}(\omega_0)(e_j, e_j)}{\omega_\varphi(e_j, J e_j)} \right|_\varphi^2 = \frac{1}{4} \sum_{j,k=1}^{2n} \frac{(\mathcal{L}_k \text{Ric}(\omega_0)(e_j, e_j))^2}{\omega_\varphi(e_j, J e_j)^2 \omega_\varphi(e_k, J e_k)} \\
&\leq \frac{1}{4} A \sum_{j,k=1}^{2n} \frac{1}{\omega_\varphi(e_j, J e_j)^2 \omega_\varphi(e_k, J e_k)} \leq \frac{1}{4} A \sum_{j=1}^{2n} \frac{1}{\omega_\varphi(e_j, J e_j)^2} \sum_{k=1}^{2n} \frac{1}{\omega_\varphi(e_k, J e_k)} \\
&\leq \frac{1}{4} A \sum_{j=1}^{2n} \frac{1}{\omega_\varphi(e_j, J e_j)} \sum_{i=1}^{2n} \frac{1}{\omega_\varphi(e_i, J e_i)} \sum_{k=1}^{2n} \frac{1}{\omega_\varphi(e_k, J e_k)} \\
&\leq 2A \Lambda_\varphi(\omega_0)^3,
\end{aligned}$$

where in the above we used that for a 1-form α we have $|\alpha|_\varphi^2 = \sum_{k=1}^{2n} \frac{\alpha_k^2}{\omega_\varphi(e_k, J e_k)}$. Also, the third line we make use of Young's inequality and in the last line we use (23).

Since $\text{Ric}(\omega_0) \leq A_0 \omega_0$ and applying once again Cauchy-Schwarz inequality we have

$$\begin{aligned}
(48) \quad g_\varphi(g_\varphi(\text{Ric}(\omega_0), \mathcal{L}.\omega_\varphi), dF) &\leq A_0 g_\varphi(g_\varphi(\omega_0, \mathcal{L}.\omega_\varphi), dF) \\
&\leq A_0 |g_\varphi(\omega_0, \mathcal{L}.\omega_\varphi)|_\varphi |dF|_\varphi.
\end{aligned}$$

We already observed (cf. the lines after the statement of Lemma 6.3) that

$$|g_\varphi(\omega_0, \mathcal{L}.\omega_\varphi)|_\varphi = \left(\sum_{k=1}^{2n} |\mathcal{L}_{e_k} \omega_\varphi|_\varphi^2 \right)^{\frac{1}{2}} = \left(-2\Lambda_0 g_\varphi(\mathcal{L}.\omega_\varphi, \mathcal{L}_J \omega_\varphi) \right)^{\frac{1}{2}}.$$

Injecting the above in (48) and using Young inequality, we find

$$(49) \quad g_\varphi(g_\varphi(\text{Ric}(\omega_0), \mathcal{L}.\omega_\varphi), dF) \leq C(|dF|_\varphi^2 - \Lambda_0 g_\varphi(\mathcal{L}.\omega_\varphi, \mathcal{L}_J \omega_\varphi)).$$

Combining (47) and (49), we get an upper bound for (46)

$$(50) \quad g_\varphi(d\Lambda_\varphi(\text{Ric}(\omega_0)), dF) \leq C \left(-\Lambda_0 (g_\varphi(\mathcal{L}.\omega_\varphi, \mathcal{L}_J \omega_\varphi)) + (\Lambda_0(\omega_\varphi)^{3n-3} + 1) |dF|_\varphi^2 + 1 \right).$$

We now bound the first term of the last line of (45).

Let $f_\varphi := \frac{w(\mu_\varphi)}{v(\mu_\varphi)} + \frac{1}{2v(\mu_\varphi)} \sum_{j=1}^r v_{,j}(\mu_\varphi) \Delta_0 \mu_0^j$. The differential of f_φ is a linear combination of terms of the form $\tilde{f}_\varphi^i d\mu_\varphi^i$, where \tilde{f}_φ^i is bounded independently of φ . Thus, in order to bound $g_\varphi(df_\varphi, dF)$, it is sufficient to bound $g(d\mu_\varphi^i, dF)$ for any $i = 1, \dots, r$. Applying Cauchy-Schwarz and Young inequalities once again we deduce

$$\begin{aligned}
(51) \quad g_\varphi(d\mu_\varphi^i, dF) &\leq |d\mu_\varphi^i|_\varphi |dF|_\varphi \\
&= |\xi^i|_\varphi |dF|_\varphi \\
&\leq \frac{1}{2} |\xi^i|_\varphi^2 |dF|_\varphi^2 + \frac{1}{2} \\
&\leq C \Lambda_0(\omega_\varphi) |dF|_\varphi^2 + \frac{1}{2} \\
&\leq C_1 \Lambda_0(\omega_\varphi)^{3n-3} |dF|_\varphi^2 + \frac{1}{2}
\end{aligned}$$

where C, C_1 are constant independent of φ . Let us observe that in the second line we have $|d\mu_\varphi^i|_\varphi = |\xi^i|_\varphi$ since the metric is invariant by duality, while in the last line we use (22).

Combining (50) and (51), we deduce an upper bound for (45):

$$(52) \quad g_\varphi(d\Delta_{\varphi,v}F, dF) \leq C \left(-\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi)) + (\Lambda_0(\omega_\varphi)^{3n-3} + 1) |dF|_\varphi^2 + 1 \right),$$

for a positive constant C independent of φ .

Finally, using (43), (44), (52) we obtain a lower bound for (42):

$$\begin{aligned} \Delta_{\varphi,v} \left(e^{F/2} |dF|_\varphi^2 \right) &\geq -e^{F/2} C \left(-\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi)) + (\Lambda_0(\omega_\varphi)^{3n-3} + 1) |dF|_\varphi^2 + 1 \right) \\ &\quad + 2e^{F/2} |\nabla^{\varphi,+} dF|_\varphi^2. \end{aligned}$$

This concludes the proof since $\|F\|_{C^0}$ is uniformly under control by Corollary 4.3 and since $1 \leq C\Lambda_0(\omega_\varphi)^{3n-3}$ by (24). \square

We are now in position to prove Proposition 6.3:

proof of Proposition 6.3. Recall that $u := e^{\frac{F}{2}} |dF|_\varphi^2 + K\Lambda_0(\omega_\varphi)$. From (29) together with (39), we obtain a lower bound of the weighted Laplacian of u :

$$\begin{aligned} \Delta_{\varphi,v} u &\geq -C_1 \left(-\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi)) + \Lambda_0(\omega_\varphi)^{3n-3} |dF|_\varphi^2 + 1 \right) + C_2 |\nabla^{\varphi,+} dF|_\varphi^2 \\ &\quad - K\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi)) - K\tilde{C}_1 \Lambda_0(\omega_\varphi)^n + K(\Delta_0 F - \tilde{C}_2). \end{aligned}$$

We now fix K large enough so that one can drop the term $-\Lambda_0(g_\varphi(\mathcal{L}\omega_\varphi, \mathcal{L}J\omega_\varphi))$. Observe that, since $n \geq 2$, $C\Lambda_0(\omega_\varphi)^{2n-2} \geq \Lambda_0(\omega_\varphi)^n$ thanks to (22). We then deduce

$$\Delta_{\varphi,v} u \geq C_2 |\nabla^{\varphi,+} dF|_\varphi^2 - C_3 \Lambda_0(\omega_\varphi)^{3n-3} u + K(\Delta_0 F - \tilde{C}_2) - C_1,$$

where $C_3 = \max(C_1 \sup_X e^{-F/2}, C\tilde{C}_1)$. Observe that $\sup_X e^{-F/2}$ is uniformly under control by Corollary 4.3.

Now, in orthonormal frame we find

$$|\Delta_0 F| = \left| \sum_{j=1}^{2n} \frac{F_{,jj} \omega_\varphi(e_j, J e_j)}{\omega_\varphi(e_j, J e_j)} \right|,$$

where $F_{,jj}$ denote the second derivative with respect to the vector e_j . Also by Lemma 6.2

$$|\nabla^{\varphi,+} dF|_\varphi^2 = \sum_{j=1}^{2n} \frac{(\nabla^{\varphi,+} dF(e_j, e_j))^2}{\omega_\varphi(e_j, J e_j)^2} = \frac{1}{4} \sum_{j=1}^{2n} \frac{(dd^c F(e_j, J e_j))^2}{\omega_\varphi(e_j, J e_j)^2} = \sum_{j=1}^{2n} \frac{F_{,jj}}{\omega_\varphi(e_j, J e_j)^2}$$

Hence, by Young's inequality

$$\begin{aligned} K|\Delta_0 F| &\leq \varepsilon |\nabla^{\varphi,+} dF|_\varphi^2 + \varepsilon^{-1} K^2 \Lambda_0(\omega_\varphi)^2 \\ &\leq \varepsilon |\nabla^{\varphi,+} dF|_\varphi^2 + C\varepsilon^{-1} K^2 \Lambda_0(\omega_\varphi)^{3n-2} u, \end{aligned}$$

where in the above we used Lemma 5.2 (since $n \geq 2$) and the fact that u is uniformly bounded. Choosing ε smaller than C_2 so that one can drop the term $|\nabla^{\varphi,+} dF|_\varphi^2$, we arrive at

$$\Delta_{\varphi,v} u \geq -(C_3 + CK\varepsilon^{-1}) \Lambda_0(\omega_\varphi)^{3n-3} u - K\tilde{C}_2 - C_1 \geq -C_4 \Lambda_0(\omega_\varphi)^{3n-3} u,$$

where C_4 depends on $\omega_0, K, \varepsilon, n, v$ and $\|F\|_{C^0}$. \square

Another key lemma is the following:

Lemma 6.5. *The following L_1 -estimate holds:*

$$\|u\|_{L_1} \leq C(\omega_0, v, \|F\|_{C^0}, \text{Ent}(\varphi)).$$

Proof. Using the second equation in (18):

$$\begin{aligned} \Delta_{v,\varphi} F^2 &= 2F \Delta_{v,\varphi} F + 2|dF|_\varphi^2 \\ &= 2F \left(-\frac{w(\mu_\varphi)}{v(\mu_\varphi)} + 2\Lambda_{\varphi,v}(\text{Ric}(\omega_0)) \right) + 2|dF|_\varphi^2. \end{aligned}$$

Since $\Delta_{v,\varphi}$ is self-adjoint with respect to $v(\mu_\varphi)\omega_\varphi^{[n]}$ (Lemma A.1), we deduce

$$\begin{aligned} \int_X |dF|_\varphi^2 v(\mu_\varphi)\omega_\varphi^{[n]} &= \int_X F \left(\frac{w(\mu_\varphi)}{v(\mu_\varphi)} - 2\Lambda_{\varphi,v}(\text{Ric}(\omega_0)) \right) v(\mu_\varphi)\omega_\varphi^{[n]} \\ &\leq \frac{M}{\eta} \int_X |F| e^F \omega_0^{[n]} + \left(2A_0 L \int_X \Lambda_\varphi(\omega_0)\omega_\varphi^{[n]} + 2C' L \right) \|F\|_{C^0}. \end{aligned}$$

In the above we use that $\Lambda_{\varphi,v}(\text{Ric}(\omega_0)) \leq A_0 \Lambda_\varphi(\omega_0) + C'$ (thanks to (56)) where C' depends on v and its derivatives, and that $\int_X \Lambda_\varphi(\omega_0)\omega_\varphi^{[n]} = 1$. \square

We can now conclude:

Proof of Theorem 6.1. We assume that φ is a (v, w) -cscK metric. The only adjustment needed for the proof is to consider the Leibniz rule for the v -weighted Laplacian:

$$\begin{aligned} \Delta_{\varphi,v} u^{2p+1} &= u^{2p} \Delta_{\varphi,v} u + 2p u^{2p-1} |du|_\varphi^2 \\ &= u^{2p} \Delta_{\varphi,v} u + \frac{8p}{(2p+1)^2} \left| d \left(u^{p+\frac{1}{2}} \right) \right|_\varphi^2, \end{aligned}$$

With the above identity and Proposition 6.3 in hands we can deduce the following bound for $p \geq 1$:

$$\begin{aligned} \int_X \left| d \left(u^{p+\frac{1}{2}} \right) \right|_\varphi^2 \omega_\varphi^{[n]} &\leq \frac{1}{\eta} \int_X \left| d \left(u^{p+\frac{1}{2}} \right) \right|_\varphi^2 v(\mu_\varphi)\omega_\varphi^{[n]} \\ &= \frac{(2p+1)^2}{8p\eta} \int_X (-u^{2p} \Delta_{\varphi,v} u + \Delta_{\varphi,v} u^{2p+1}) v(\mu_\varphi)\omega_\varphi^{[n]} \\ &= -\frac{(2p+1)^2}{8p\eta} \int_X u^{2p} \Delta_{\varphi,v} u \\ &\leq C \int_X \Lambda_0(\omega_\varphi)^{3n-3} u^{2p+1} \omega_\varphi^{[n]}, \end{aligned}$$

Observe also that at the third line we also make use of the fact that $\Delta_{\varphi,v}$ is self-adjoint with respect to $v(\mu_\varphi)\omega_\varphi^{[n]}$.

Once the latter inequality in hand, the exact same arguments in [16, Proof of Theorem 5.1] allow to conclude. \square

APPENDIX A. WEIGHTED LAPLACIAN AND WEIGHTED TRACE

A.1. The weighted Laplacian. For any smooth function f on X , the v -weighted Laplacian is defined as

$$(53) \quad \Delta_{\varphi, v} f := -\frac{1}{v(\mu_{\varphi})} d_{\varphi}^* (v(\mu_{\varphi}) df).$$

It follows straightforward from the above expression that:

Lemma A.1. *The weighted Laplacian $\Delta_{\varphi, v}$ is self-adjoint and Leibniz rule holds.*

Lemma A.2. *For any smooth function f on X , the weighted Laplacian $\Delta_{\varphi, v}$ can be expressed as*

$$\begin{aligned} \Delta_{\varphi, v} f &= \Delta_{\varphi} f + \frac{1}{v(\mu_{\varphi})} \sum_{j=1}^r v_{,j}(\mu_{\varphi}) d^c f(\xi_j) \\ &= \Delta_{\varphi} f + g_{\varphi}(d \log v(\mu_{\varphi}), df). \end{aligned}$$

In particular $\Delta_{\varphi, v}$ is elliptic.

Proof. We apply the well-know identity (see e.g. [17, (1.10.13)])

$$d_{\varphi}^*(h\alpha) = h d_{\varphi}^* \alpha - \alpha(\nabla^{\varphi} h),$$

where h is a smooth function and α a 1-form, for $h = v(\mu_{\varphi})$ and $\alpha = df$. We then use the fact that $d^c f(\cdot) = Jdf(\cdot) = -df(J\cdot)$ and that $-J\xi_j$ is the gradient of μ_{φ}^j with respect to g_{φ} since $d\mu_{\varphi}^j = -\omega_{\varphi}(\xi_j, \cdot) = -g_{\varphi}(J\xi_j, \cdot)$. \square

A.2. The weighted trace. Let f be a \mathbb{T} -invariant smooth function $f : X \rightarrow \mathbb{R}$. By Cartan's formula and the \mathbb{T} -invariance of $d^c f$ we get that

$$dd^c f(\xi, \cdot) = \xi \lrcorner dd^c f(\cdot, \cdot) = \mathcal{L}_{\xi} d^c f - d(d^c f(\xi)) = -d(d^c f(\xi)),$$

for any $\xi \in \mathfrak{t}$. Hence we choose the normalization of the moment map by defining $\mu_{dd^c f}^j := d^c f(\xi_j)$.

We consider a smooth path $(\varphi_t)_t$ in $\mathcal{K}(X, \omega_0)^{\mathbb{T}}$ with variation $\frac{d}{dt} \varphi_t|_{t=0} = \dot{\varphi}$. Taking the variation of the weighted volume form along φ_t and then computing it at $t = 0$, we obtain (see proof of [25, Lemma 4])

$$\frac{d}{dt} \left(v(\mu_{\varphi_t}) \omega_{\varphi_t}^{[n]} \right) \Big|_{t=0} = \left(\Lambda_{\varphi_0}(dd^c \dot{\varphi}) + \langle d \log v(\mu_{\varphi_0}), \mu_{dd^c \dot{\varphi}} \rangle \right) v(\mu_{\varphi_0}) \omega_{\varphi_0}^{[n]}.$$

The above identity motivates the following definition: for any $(1, 1)$ -form θ with moment map μ_{θ} , we define the v -weighted trace of θ by

$$(54) \quad \Lambda_{\varphi, v}(\theta) := \Lambda_{\varphi}(\theta) + \langle d \log v(\mu_{\varphi}), \mu_{\theta} \rangle$$

for any $\varphi \in \mathcal{K}(X, \omega_0)^{\mathbb{T}}$. Observe that the above definition depends on the choice of the normalization of the momentum map μ_{θ} . Equivalently, in a fixed basis (ξ_1, \dots, ξ_r) of the Lie algebra \mathfrak{t} of \mathbb{T}

$$(55) \quad \Lambda_{\varphi, \nu}(\theta) = \Lambda_{\varphi}(\theta) + \frac{1}{\nu(\mu_{\varphi})} \sum_{j=1}^r \nu_{,j}(\mu_{\varphi}) \mu_{\theta}^j.$$

In particular, since $\mu_{\text{Ric}(\omega_0)} = -\frac{1}{2}\Delta_0\mu_0$ with μ_0 being the moment map associated to ω_0 (see [25, Lemma 5 (i)]), we have

$$(56) \quad \Lambda_{\varphi, \nu}(\text{Ric}(\omega_0)) = \Lambda_{\varphi}\text{Ric}(\omega_0) - \frac{1}{2\nu(\mu_{\varphi})} \sum_{j=1}^r \nu_{,j}(\mu_{\varphi}) \Delta_0\mu_0^j.$$

Lemma A.3. *For any \mathbb{T} -invariant smooth function $f : X \rightarrow \mathbb{R}$, the following identity holds true*

$$\Delta_{\varphi, \nu}f = \Lambda_{\varphi, \nu}(dd^c f).$$

Proof. It is a direct consequence of (55) and Lemma A.2. □

REFERENCES

- [1] V. Apostolov and D. Calderbank. The CR geometry of weighted extremal Kähler and Sasaki metrics. *Math. Ann.*, 379(3-4):1047–1088, 2021.
- [2] V. Apostolov, D. Calderbank, P. Gauduchon, and C. Tønnesen-Friedman. Extremal Kähler metrics on projective bundles over a curve. *Adv. Math.*, 227:2385–2424, 2011.
- [3] V. Apostolov, D. Calderbank, and E. Legendre. Weighted K-stability of polarized varieties and extremality of Sasaki manifolds. *to appear in Adv. Math.*, *arXiv:2012.08628*, 2020.
- [4] V. Apostolov, S. Jubert, and A. Lahdili. Weighted K-stability and coercivity with applications to extremal Kähler and Sasaki metrics. *to appear in Geometry and Topology*, *arXiv:2104.09709*, 2021.
- [5] M. F. Atiyah. Convexity and commuting hamiltonians. *Bull. London Math. Soc.*, 14:1–15, 1982.
- [6] T. Aubin. Equations de type Monge-Ampère sur les variétés kählériennes compactes. *C. R. Acad. Sci. Paris*, 283:119–121, 1976.
- [7] R. J. Berman and D. Witt-Nyström. Complex optimal transport and the pluripotential theory of Kähler-Ricci solitons. *arXiv:1401.8264*, 2014.
- [8] Z. Blocki. On geodesics in the space of Kähler metrics, proceedings of the “Conference in Geometry” dedicated to Shing-Tung Yau. *Advances in Geometric Analysis, Advanced Lectures in Mathematics*, 20:3–20, 2009.
- [9] S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi. Monge-Ampère equations in big cohomology classes. *Acta Math.*, 205(2):199–262, 2010.
- [10] E. Calabi. Extremal Kähler metrics. *Seminar on Differential Geometry*, *Ann. of Math. Stud.*, Princeton Univ. Press, Princeton., 102:259–290, 1982.
- [11] F. Campana, H. Guenancia, and M. Paun. Metrics with cone singularities along normal crossing, divisors and holomorphic tensor fields. *Ann. Scient. Ec. Norm. Sup.*, page 879–916, 2013.
- [12] X. Chen and J. Cheng. On the constant scalar curvature Kähler metrics: apriori estimates. *J. Amer. Math. Soc.*, 2021.
- [13] T. Darvas, E. Di Nezza, and C. H. Lu. Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity. *Anal. PDE*, 11(8):2049–2087, 2018.
- [14] T. Darvas, E. Di Nezza, and C. H. Lu. On the singularity type of full mass currents in big cohomology classes. *Compos. Math.*, 154(2):380–409, 2018.
- [15] T. Darvas, E. Di Nezza, and C.H. Lu. Log-concavity of volume and complex Monge-Ampère equations with prescribed singularity. *Mathematische Annalen*, pages 1–38, 2019.
- [16] A. Deruelle and E. Di Nezza. Uniform estimates for cscK metrics. *to appear in Ann. Fac. Sci. Toulouse Math.*, 2021.
- [17] P. Gauduchon. Calabi’s extremal Kähler metrics: an elementary introduction. *Lecture Notes*, 2022.
- [18] V. Guedj and A. Zeriahi. *Degenerate complex Monge-Ampère equations*, volume 26 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2017.
- [19] V. Guillemin and S. Sternberg. Convexity properties of the moment mapping. *Invent. Math.*, 67:491–513, 1982.

- [20] J. Han and C.i Li. On the Yau-Tian-Donaldson conjecture for generalized Kähler-Ricci soliton equations. *Communications on Pure and Applied Mathematics*, 76(9):1793–1867, September 2023. Publisher Copyright: © 2022 Wiley Periodicals LLC.
- [21] J. Han and Y. Liu. On the existence of weighted-csck metrics. *arXiv:2406.10939*, 2024.
- [22] S. Jubert. A Yau–Tian–Donaldson correspondence on semisimple principale toric fibrations. *Annales de l’Institut Fourier*, 73, 2023.
- [23] Sławomir Kołodziej. The complex Monge-Ampère operator. *Acta Math.*, 180(1):69–117, 1998.
- [24] O. Ladyzenskaya and N. Uralsteva. Linear and quasilinear elliptic partial differential equations. *Academic Press*, 1968.
- [25] A. Lahdili. Kähler metrics with weighted constant scalar curvature and weighted K-stability. *Proc. London Math. Soc.*, 119:1065 – 114, 2019.
- [26] E. Di Nezza and C. Lu. Complex Monge-Ampère equations on quasi-projective varieties. *J. Reine Angew. Math.*, 727:145–167, 2017.
- [27] G. Tian and X. Zhu. Uniqueness of Kähler-Ricci solitons. *Acta Mathematica*, 184:271–305, 01 2000.
- [28] G. Tian and X. Zhu. A new holomorphic invariant and uniqueness of Kähler-Ricci solitons. *Commentarii Mathematici Helvetici*, 77(2):297–325, 2002.
- [29] S. Yau. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. *I. Commun. Pure Appl. Math.*, 21, 1978.