

An Optimal Pricing Formula for Smart Grid based on Stackelberg Game

Jiangjiang Cheng, Ge Chen, Zhouming Wu, and Yifen Mu

Abstract—The dynamic pricing of electricity is one of the most crucial demand response (DR) strategies in smart grid, where the utility company typically adjust electricity prices to influence user electricity demand. This paper models the relationship between the utility company and flexible electricity users as a Stackelberg game. Based on this model, we present a series of analytical results under certain conditions. First, we give an analytical Stackelberg equilibrium, namely the optimal pricing formula for utility company, as well as the unique and strict Nash equilibrium for users' electricity demand under this pricing scheme. To our best knowledge, it is the first optimal pricing formula in the research of price-based DR strategies. Also, if there exist prediction errors for the supply and demand of electricity, we provide an analytical expression for the energy supply cost of utility company. Moreover, a sufficient condition has been proposed that all electricity demands can be supplied by renewable energy. When the conditions for analytical results are not met, we provide a numerical solution algorithm for the Stackelberg equilibrium and verify its efficiency by simulation.

Index Terms— Stackelberg game, Stackelberg equilibrium, optimal pricing formula, smart grid, demand response

I. INTRODUCTION

Smart grid refers to an advanced electricity distribution system that incorporates smart metering infrastructure capable of monitoring and measuring power consumption of users with sophisticated communication network technology [1], [2], to achieve an efficient, reliable, and sustainable electric grid operation. [3], [4]. One of the key features of smart grid is the ability to regulate users' electricity demand with the aim of balancing supply and demand as well as reducing power generation costs, known as demand response (DR) [5]–[7]. DR commonly refers to the energy consumption changes of users in response to factors such as electricity prices or incentive payments [8], [9], which plays a crucial role in optimizing energy allocation, alleviating peak load, and ensuring the stability of the power grid [10]–[12]. On the other hand, the development of renewable energy technology has a significant effect on addressing the challenges of greenhouse gas emissions and environmental pollution, gradually occupying a dominant position in the energy supply of power grid [3], [13]. Therefore, it is natural to design DR mechanisms that take into account renewable energy to achieve the goal of sustainable development [14], [15].

Among various DR strategies, price-based DR strategy is a prevalent research topic, including day-ahead pricing, real-time pricing (RTP) and peak load pricing, etc [16]–[18]. In terms of improving the performance of energy market, RTP is the hottest area of DR mechanism, in which the electricity prices announced by the utility company change typically to regulate user electricity demand [19], [20]. The study of RTP mechanism can help the smart grid achieve different objectives, including maximizing the social welfare [21],

minimizing the energy expenses [22], mitigating peak load [23], and maintaining the stability of energy networks [24]. Through the dynamic change in electricity prices, RTP incentivizes users to adjust their electricity consumption patterns to match real-time electricity supply, in order to realize more flexible and sustainable power system operation.

Because of the coupling relationship between multiple interactive entities in smart grid, such as the energy service provider and various users, network game theory provides a natural framework for the interaction among various participators with different objectives [25]–[28]. Most network game models typically exhibit complicated dynamic behaviors, making the theoretical analysis of these models very difficult [29], [30]. Current mainstream methods include mean field dynamics [31], [32], semi-tensor product method [33], [34], and so on. Zhu et al. [35] formulated and solved a demand-side management problem of networked smart grid based on evolutionary game theory. By using semi-tensor product, they proposed a new binary optimal control algorithm to minimize the overall cost of smart grid. Based on the randomness and dynamicity of wind and solar power generation, it is natural to employ a richer class of network games, namely stochastic game, to capture such dynamic uncertain environments. Etesami et al. [36] introduced a novel model for energy trading in smart grid employing a stochastic game framework on the demand side and an optimization problem on the utility side. They innovatively proposed a distributed algorithm to obtain equilibrium points of the system and adopted online convex optimization to solve the utility side problem. Besides, there are numerous network game models used to assist utility company in estimating energy demand and supply costs [37], achieving load balancing [38], and managing energy storage and distributed microgrids [39]. However, due to the large-scale, distributed, multi-agent nature of smart grid, and the uncertainty of renewable energy, the equilibrium analysis and solution of network game models in smart grid are still highly challenging, which will become an important direction for future endeavors.

Due to the typical hierarchical structure inherent in the electricity market, numerous price-based DR problems with two decision levels can be effectively modeled as a special type of network game in smart grid – Stackelberg game. The Stackelberg game model can be used to analyze the interactions between various entities in different scenarios. A large amount of existing research focuses on the game relationship between utility company and users, and/or the interaction among users. Concretely, the utility company set dynamic electricity prices as a leader, and users adapt their electricity consumption patterns as followers, thereby achieving coordination of energy consumption and load distribution [40], [41]. Additionally, the competition between multiple power utilities has gradually attracted widespread attention, with the market regulator setting price schemes to improve social welfare, and other power utilities adjusting pricing strategies accordingly to maximize their own profits while considering market competition [42], [43].

Although there have been many studies on price-based DR strategies in smart grid, a fundamental issue for utility companies is still open: what is the optimal pricing formula? There are some numerical solutions to this problem [16], [20], [21], but its analytical expression

This research is supported by the National Key Research and Development Program of China (2022YFA1004600), and the National Natural Science Foundation of China (12288201, 12071465).

Jiangjiang Cheng and Zhouming Wu are with the School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China (e-mail: chengjiangjiang@amss.ac.cn; wuzhouming@amss.ac.cn).

Ge Chen and Yifen Mu are with the Key Laboratory of Systems and Control, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China (e-mail: chenge@amss.ac.cn; mu@amss.ac.cn).

has not yet been provided. This paper addresses this problem and its main contributions are as follows:

- This paper proposes a novel price-based DR model of Stackelberg game between the electricity company and users, in which the goal of the electricity company is to minimize the power supply cost, while the goals of users are to minimize electricity consumption costs. Different from previous works [10], [40], [41], our proposed model comprehensively considers the uncontrollability of wind and solar energy, the diversity of electricity users, and some constraints on users' electricity consumption, which makes it more realistic in certain scenarios.
- Based on our proposed model, we present a series of analytical results under certain inequality conditions. First, if the supply and demand of electricity can be predicted exactly in a certain time window, we give an analytical expression for the Stackelberg equilibrium, in which the unique strict Nash equilibrium among users is also the optimal solution for the utility company, therefore the price expression in the Stackelberg equilibrium can be treated as an optimal pricing formula. To the best of our knowledge, it is the first optimal pricing formula in the research of price-based DR strategies.

Also, if there exist prediction errors for the supply and demand of electricity, we provide an analytical expression for the Nash equilibrium among users and the corresponding energy supply cost of utility company. Moreover, we find that under the same prediction accuracy, by shortening the sampling length of the time slot, the utility company's cost function under the Nash equilibrium can be greatly reduced.

Based on above results, we further provide a sufficient condition for an ideal scenario where all electricity demand can be supplied by renewable energy sources such as wind and photovoltaic power.

- When the inequality conditions for the analytical solution are not satisfied, a search algorithm is developed to obtain the numerical solution of Stackelberg equilibrium, and a simulation example is presented to demonstrate the effectiveness of our proposed algorithm.

The rest of this paper is organized as follows. Section II introduces the constructed Stackelberg game model and some assumptions. Section III analyzes the proposed game model and provides analytical solutions for the corresponding Stackelberg equilibrium under certain conditions, and the proofs of the corresponding theoretical results are presented in Section IV. When the conditions are not met, Section V provides a numerical solution algorithm for Stackelberg equilibrium. The simulation results are presented and discussed subsequently. Finally, the conclusion and outlook are made in Section VI.

II. PROBLEM FORMULATION AND ASSUMPTIONS

This section provides the formulation of the problem to be solved in this paper. Without loss of generality, our system includes one supply side, one service side and one demand side. The supply side consists of various types of power supplies, which are divided into two categories: one is controllable power supply (e.g. thermal power generation, hydro power generation, storage); the other one is uncontrollable power supply (e.g. wind power generation, photovoltaic power generation). On the service side, there is a utility company engaged in electricity trading between the supply side and demand side. By setting an appropriate price, the utility company can regulate users' electricity demands, and then feedback energy consumption information to supply side to achieve supply-demand balance. The demand side is composed of a large number of users, which can be divided into two categories: flexible electricity users (e.g. electric

vehicles, hydrogen production plant, some hot water supply systems, some discrete electricity enterprises) and regular electricity users (e.g., household electricity, continuous electricity enterprises). The difference between these two categories of users is: the flexible users determine their electricity consumption according to the spot price, while the electricity demands of regular users do not depend on the spot price. Fig. 1 illustrates the process of energy trading between power supplies, utility company, and users. For ease of understanding, Table I summarizes the main notations used in this paper.




Fig. 1. The process of energy trading between power supplies, utility company, and users.

A. Stackelberg Game Model

We model the interaction between the utility company and flexible electricity users as a Stackelberg game with one leader and n followers. The utility company is the leader who regulates users' energy demands by setting proper electricity prices, thereby reducing the cost of controllable power generation; the flexible users are the followers who determine their electricity demand patterns after obtaining corresponding price schemes, thereby minimizing their own electricity costs. The mathematical representations of our proposed Stackelberg game are introduced as follows.

Demand side: Let $\mathcal{N} := \{1, 2, \dots, n\}$ be the set of n followers, where each follower denotes a flexible electricity user. Considering some practical scenarios, such as electric vehicle charging or electricity consumption of factories produced according to orders, we assume that each user $i \in \mathcal{N}$ has a total electricity demand $g_i > 0$ that must be satisfied within T time periods, where T represents the scheduling time window. To finish the energy consumption goals while minimizing their own electricity costs, each user $i \in \mathcal{N}$ needs to determine the electricity demand $\nu_i(t)$, $0 \leq \nu_i(t) \leq \nu_i^{\max}$ at each time slot $t, t = 1, \dots, T$. Here ν_i^{\max} represents the maximum electricity consumption of user i at each time slot, which satisfies $\nu_i^{\max} \geq g_i/T$. The strategy of user i is set to be

$$\boldsymbol{\nu}_i := (\nu_i(1), \nu_i(2), \dots, \nu_i(T))^T.$$

Therefore, the total electricity demand of all flexible electricity users at time slot t is $\nu_{\mathcal{N}}(t) := \sum_{i \in \mathcal{N}} \nu_i(t)$, $t = 1, \dots, T$, and the total electricity demand of all flexible electricity users is $g_{\mathcal{N}} := \sum_{i \in \mathcal{N}} g_i$.

TABLE I
LIST OF MAIN NOTATIONS.

Notation	Description
\mathcal{N}	Set of all followers, i.e., $\mathcal{N} = \{1, 2, \dots, n\}$
T	Number of scheduling time window
$\mathbf{1}$	A column vector whose elements are all 1
$\nu_i(\cdot)$	Electricity demand of flexible user i at each time slot
$\boldsymbol{\nu}_i$	The strategy of flexible user i , i.e., $\boldsymbol{\nu}_i = (\nu_i(1), \nu_i(2), \dots, \nu_i(T))^\top$
$\boldsymbol{\nu}$	The strategy combination of all flexible users, i.e., $\boldsymbol{\nu} = (\boldsymbol{\nu}_1, \boldsymbol{\nu}_2, \dots, \boldsymbol{\nu}_n)^\top$
$\boldsymbol{\nu}_{-i}$	The strategies of all flexible users except the i th one, i.e., $\boldsymbol{\nu}_{-i} = (\boldsymbol{\nu}_1, \dots, \boldsymbol{\nu}_{i-1}, \boldsymbol{\nu}_{i+1}, \dots, \boldsymbol{\nu}_n)^\top$
ν_i^{\max}	Maximum electricity consumption of flexible user i at each time slot
$\nu_{\mathcal{N}}(\cdot)$	Total electricity demand of all flexible users at each time slot, i.e., $\nu_{\mathcal{N}}(\cdot) = \sum_{i \in \mathcal{N}} \nu_i(\cdot)$
g_i	Total electricity demand of flexible user i
$g_{\mathcal{N}}$	Total electricity demand of all flexible users, i.e., $g_{\mathcal{N}} = \sum_{i \in \mathcal{N}} g_i$
$r(\cdot)$	Total electricity demand of all regular users at each time slot
$w(\cdot)$	Uncontrollable power generation at each time slot
$c(\cdot)$	Controllable energy supply at each time slot
$\pi(\cdot)$	Electricity price function at each time slot, i.e., $\pi(\cdot) = \pi(\nu_{\mathcal{N}}(\cdot), r(\cdot), w(\cdot))$
$\boldsymbol{\pi}$	Leader's strategy, i.e., $\boldsymbol{\pi} = (\pi(1), \pi(2), \dots, \pi(T))^\top$
$u^l(\cdot)$	The cost function of leader
$u_i^f(\cdot)$	The electricity cost of flexible user i

Let $\boldsymbol{\nu}$ be the strategy combination of all flexible users, i.e., $\boldsymbol{\nu} = (\boldsymbol{\nu}_1, \boldsymbol{\nu}_2, \dots, \boldsymbol{\nu}_n)^\top$.

On the other hand, the regular electricity users have their own electricity demands at each time slot and are not affected by real-time electricity prices. We assume that the total electricity demand of all regular users at time slot t is $r(t)$, $t = 1, \dots, T$, which is assumed to be a known sequence.

Supply side: We denote the energy supply of controllable and uncontrollable energy at time slot t as $c(t)$ and $w(t)$, respectively. It is worth mentioning that the uncontrollable power generation $w(t)$ significantly depends on climate conditions and should be a stochastic sequence. However, to obtain some deterministic results we assume $\{w(t)\}_{t=1, \dots, T}$ is a deterministic sequence.

Service side: As the leader, the utility company regulates the electricity demands of users by setting an appropriate electricity price $\pi(t) = \pi(\nu_{\mathcal{N}}(t), r(t), w(t))$, $t = 1, \dots, T$, which is a function with respect to $\nu_{\mathcal{N}}(t)$, $r(t)$, and $w(t)$. We denote the strategy of utility company as

$$\boldsymbol{\pi} := (\pi(1), \pi(2), \dots, \pi(T))^\top.$$

The utility company need purchase $c(t)$, $t = 1, \dots, T$ electricity quantity from controllable power generation to guarantee the dynamic

balance between electricity demand and supply, i.e., $\nu_{\mathcal{N}}(t) + r(t) = w(t) + c(t)$. Since

$$\begin{aligned} \sum_{t=1}^T c(t) &= \sum_{t=1}^T [\nu_{\mathcal{N}}(t) + r(t) - w(t)] \\ &= \sum_{i=1}^n g_i + \sum_{t=1}^T [r(t) - w(t)] \\ &= g_{\mathcal{N}} + \sum_{t=1}^T [r(t) - w(t)] \end{aligned}$$

is assumed to be a constant, we set the cost function $u^l(\boldsymbol{\pi}, \boldsymbol{\nu})$ of utility company to be the variance of controllable power $c(t)$. Let $\bar{c} := \frac{1}{T} \sum_{t=1}^T c(t)$ be the average value of $c(t)$ over the entire time periods. Therefore, the leader-level optimization problem is formulated in (1).

For each flexible electricity user $i \in \mathcal{N}$, let $\boldsymbol{\nu}_{-i}$ denote the strategies of all users except the i th one. The utility of flexible user i is set to be its electricity cost $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$, which is a function with respect to the strategies of the utility company and all flexible users. Then, the follower-level game among all flexible users can be described by (2).

Leader-Level Optimization:

$$\min_{\boldsymbol{\pi}} u^l(\boldsymbol{\pi}, \boldsymbol{\nu}) = \frac{1}{T} \sum_{t=1}^T [c(t) - \bar{c}]^2, \quad (1)$$

$$\text{s.t. } \nu_{\mathcal{N}}(t) + r(t) = w(t) + c(t), \quad t = 1, \dots, T.$$

Follower-Level Game: For each flexible user $i = 1, \dots, n$,

$$\begin{aligned} \min_{\boldsymbol{\nu}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}) &= \sum_{t=1}^T \pi(\nu_{\mathcal{N}}(t), r(t), w(t)) \cdot \nu_i(t), \\ \text{s.t. } \sum_{t=1}^T \nu_i(t) &= g_i, \\ 0 \leq \nu_i(t) &\leq \nu_i^{\max}, \quad t = 1, \dots, T. \end{aligned} \quad (2)$$

Overall process of the Stackelberg Game: The utility company first announces the price formula to all flexible users. Then, each user selects the corresponding optimal demand strategy as the “best response” to the leader’s strategy, and feeds its strategy back to the utility company. Therefore, the solution of the Stackelberg game problem can be ultimately transformed into the choice of the optimal pricing formula of the utility company, based on the rational assumption that each flexible user selects its best-response strategy.

III. ANALYSIS OF STACKELBERG GAME MODEL

In this section, we analyze the proposed Stackelberg game (1)-(2). First, we provide a strict definition of Stackelberg equilibrium as the solution of the Stackelberg game. Then, we present a series of analytical results for equilibria in Stackelberg game (1)-(2) under certain conditions. Concretely, we give an analytical Stackelberg equilibrium, namely the optimal pricing formula for utility company, as well as the unique and strict Nash equilibrium for users’ electricity demand under this pricing scheme. Also, if there exist prediction errors for the supply and demand of electricity, we provide an analytical expression for the energy supply cost of utility company. Moreover, a sufficient condition has been proposed that all electricity demands can be supplied by new energy sources.

A. Stackelberg Equilibrium

For the proposed Stackelberg game (1)-(2), we introduce the classic Stackelberg equilibrium (SE) [44] formulated as follows.

Given a leader's strategy π , the best response strategy $\nu(\pi) = (\nu_1(\pi), \nu_2(\pi), \dots, \nu_n(\pi))^\top$ of followers is the Nash equilibrium of game model (2), that is, for any user $i \in \mathcal{N}$ and any strategy ν'_i , we have

$$u_i^f(\pi, \nu'_i, \nu_{-i}(\pi)) \geq u_i^f(\pi, \nu_i(\pi), \nu_{-i}(\pi)). \quad (3)$$

Further, if the inequality in (3) strictly holds for any $\nu'_i \neq \nu_i(\pi)$, we say that $\nu(\pi)$ is the *strict* Nash equilibrium. Meanwhile, based on the assumption that the followers reach a Nash equilibrium, the leader should try to minimize its cost function $u^l(\pi, \nu(\pi))$ by optimizing its price function π . Let $\hat{\pi}$ be the solution of the problem $\min_{\pi} u^l(\pi, \nu(\pi))$, and $\hat{\nu} = \nu(\hat{\pi})$ be the best responses of all followers to the leader. Then $(\hat{\pi}, \hat{\nu})$ is the SE of the Stackelberg game (1)-(2).

In addition, we say $(\hat{\pi}, \hat{\nu})$ is a *perfect SE* of the Stackelberg game (1)-(2) if $u^l(\hat{\pi}, \hat{\nu}) = 0$.

B. Main Results

A natural idea is to set the price function as the ratio of total electricity demand to power generation capacity. In order to fully utilize the power generation capacity of wind and solar power generation, we define a price function as

$$\pi(t) = \frac{\nu_{\mathcal{N}}(t) + r(t) + a_2(t)}{w(t) + a_1(t)}, \quad \forall 1 \leq t \leq T, \quad (4)$$

where $a_1(t)$ and $a_2(t)$ are two adjustable factors. We aim to find the SE of the Stackelberg game (1)-(2) by adjusting the values of $a_1(t)$ and $a_2(t)$. Set

$$\tilde{w}(t) := w(t) + a_1(t), \quad \tilde{r}(t) := r(t) + a_2(t), \quad \forall 1 \leq t \leq T. \quad (5)$$

Under the leader's strategy (4), we first analyze the Nash equilibrium of the following game model:

$$\begin{cases} \min_{\nu_i} u_i^f(\pi, \nu_i, \nu_{-i}), \\ \text{s.t. } \sum_{i=1}^T \nu_i(t) = g_i, \end{cases} \quad i = 1, \dots, n. \quad (6)$$

Proposition 3.1: Assume that the leader's strategy adopts (4), and $\tilde{w}(t) > 0$ for any $1 \leq t \leq T$, then the strategy combination ν of all followers with

$$\begin{aligned} \nu_i(t) &= \frac{\tilde{w}(t)}{\sum_s \tilde{w}(s)} g_i + \frac{1}{(n+1) \sum_s \tilde{w}(s)} \\ &\quad \times \sum_{s=1}^T \tilde{w}(s) \tilde{w}(t) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(t)}{\tilde{w}(t)} \right], \end{aligned} \quad (7)$$

$$i = 1, \dots, n, \quad t = 1, \dots, T$$

is a strict and unique Nash equilibrium of the game model (6).

For the model (2), we can prove that the Nash equilibrium is strict and unique, however its explicit expression is unknown:

Proposition 3.2: Assume that the leader's strategy adopts (4) with $\tilde{w}(t) > 0$ for any $1 \leq t \leq T$, then the model (2) admits a strict and unique Nash equilibrium.

For any $1 \leq t \leq T$, if we choose

$$\begin{aligned} a_1^*(t) &:= \frac{g_{\mathcal{N}}}{T} - r(t) + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)], \\ a_2^*(t) &:= \frac{n+1}{n} [w(t) + a_1^*(t)] - r(t), \end{aligned} \quad (8)$$

then the price function (4) becomes

$$\begin{aligned} \pi^*(t) &:= \frac{\nu_{\mathcal{N}}(t) + r(t) + a_2^*(t)}{w(t) + a_1^*(t)} \\ &= 1 + \frac{1}{n} + \frac{\nu_{\mathcal{N}}(t)}{w(t) + \frac{g_{\mathcal{N}}}{T} - r(t) + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)]}. \end{aligned} \quad (9)$$

We set

$$\begin{aligned} \tilde{w}^*(t) &:= w(t) + a_1^*(t) \\ &= \frac{g_{\mathcal{N}}}{T} + w(t) - r(t) + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)], \\ \tilde{r}^*(t) &:= r(t) + a_2^*(t) = \frac{n+1}{n} \tilde{w}^*(t), \quad \forall 1 \leq t \leq T. \end{aligned} \quad (10)$$

Then, with (10), the strict and unique Nash equilibrium (7) of model (6) becomes

$$\begin{aligned} \nu_i^*(t) &:= \frac{\tilde{w}^*(t)}{\sum_s \tilde{w}^*(s)} g_i + \frac{1}{(n+1) \sum_s \tilde{w}^*(s)} \\ &\quad \times \sum_{s=1}^T \tilde{w}^*(s) \tilde{w}^*(t) \left[\frac{\tilde{r}^*(s)}{\tilde{w}^*(s)} - \frac{\tilde{r}^*(t)}{\tilde{w}^*(t)} \right] \\ &= \frac{g_i}{\sum_s \tilde{w}^*(s)} \tilde{w}^*(t) = \frac{g_i}{g_{\mathcal{N}}} \tilde{w}^*(t) \\ &= \frac{g_i}{T} + \frac{g_i}{g_{\mathcal{N}}} [w(t) - r(t)] + \frac{g_i}{g_{\mathcal{N}} T} \sum_{s=1}^T [r(s) - w(s)], \\ &\quad i = 1, \dots, n, \quad t = 1, \dots, T. \end{aligned} \quad (11)$$

Let $\pi^* := (\pi^*(1), \pi^*(2), \dots, \pi^*(T))^\top$ denote the leader's strategy, $\nu_i^* := (\nu_i^*(1), \nu_i^*(2), \dots, \nu_i^*(T))^\top$ denote the strategy of follower i , and $\nu^* := (\nu_1^*, \nu_2^*, \dots, \nu_n^*)^\top$ denote the strategy combination of all followers. We have

Proposition 3.3: When the leader's strategy is π^* and the followers' strategy is ν^* , the leader's cost function $u^l(\pi^*, \nu^*) = 0$.

The proofs of Proposition 3.1-3.3 are put in Subsection 4-A.

Based on Proposition 3.1-3.3, we now demonstrate that the Stackelberg game (1)-(2) can achieve a perfect SE if the following inequalities hold:

$$\begin{aligned} 0 &< \frac{g_{\mathcal{N}}}{T} + w(t) - r(t) + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] = \tilde{w}^*(t) \\ &\leq \min_{i \in \mathcal{N}} \left\{ \frac{\nu_i^{\max}}{g_i} \right\} g_{\mathcal{N}}, \quad \forall 1 \leq t \leq T. \end{aligned} \quad (12)$$

We show that, under the condition (12), ν^* is the best-response strategy to the leader's strategy π^* , and they contribute a perfect SE of the Stackelberg game (1)-(2). Thus, $\pi^*(t)$ can be treated as an optimal pricing formula for the utility company.

Theorem 3.1 (Perfect SE with optimal pricing formula $\pi^*(t)$): If the inequalities (12) hold then

- i) ν^* is a strict and unique Nash equilibrium of model (2) when the leader's strategy is π^* ;
- ii) (π^*, ν^*) is a perfect SE of the Stackelberg game (1)-(2).

Fig. 2 shows an example of the curves of $\{w(t), r(t), c(t)\}_{t=1, \dots, T}$ and $\{\nu_i^*(t)\}_{i=1, \dots, n; t=1, \dots, T}$, where we make $n = 10$, $T = 24$ and they satisfy the inequality (12), thus presenting a perfect SE profile.

Remark 1: From Theorem 3.1, if the utility company adopts the price formula $\pi^*(t)$, all flexible users have a stable electricity strategy combination ν^* , in which each flexible user will not proactively change its strategy to avoid an increase in electricity cost. Meanwhile, all electricity needs are met, all new energy generation is utilized, and




Fig. 2. An example of achieving a perfect SE, in which the uncontrollable energy accounts for 65.84% of energy supply, and flexible users account for 25.29% of the total energy demand.

the controllable power supply $c(t) \equiv \text{const}$, thereby social benefits are maximized.

An interesting question is, if we ignore the stability factors of the grid and only consider the supply-demand balance of electricity, under what conditions can all electricity be provided by new energy sources? The following corollary provides a solution for this ideal situation.

Corollary 3.1: Assume that the conditions

$$0 < w(t) - r(t) \leq \min_{i \in \mathcal{N}} \left\{ \frac{\nu_i^{\max}}{g_i} \right\} g_{\mathcal{N}}, \quad \forall 1 \leq t \leq T, \quad (13)$$

and

$$\sum_{t=1}^T w(t) = g_{\mathcal{N}} + \sum_{t=1}^T r(t) \quad (14)$$

hold. Let

$$\begin{aligned} \pi_0^* &:= 1 + \frac{1}{n} + \left(\frac{\nu_{\mathcal{N}}(1)}{w(1) - r(1)}, \dots, \frac{\nu_{\mathcal{N}}(T)}{w(T) - r(T)} \right)^\top, \\ \chi_i^* &:= \frac{g_i}{g_{\mathcal{N}}} (w(1) - r(1), \dots, w(T) - r(T))^\top, \quad i \in \mathcal{N}, \\ \chi^* &:= (\chi_1^*, \chi_2^*, \dots, \chi_n^*)^\top. \end{aligned}$$

Then, the Stackelberg game (1)-(2) has a perfect SE (π_0^*, χ^*) , in which all electricity is provided by uncontrollable energy sources such as wind and photovoltaic power, without energy storage or waste.

Fig. 3 shows an example of the curves of $\{w(t), r(t)\}_{t=1, \dots, T}$ and $\{\nu_i^*(t)\}_{i=1, \dots, n; t=1, \dots, T}$, which satisfies the conditions in Corollary 3.1. Therefore, it also presents a perfect SE profile, where all electricity is provided by uncontrollable energy sources.

Remark 2: One possible application of Corollary 3.1 is to plan the electricity consumption and power of flexible electricity users. For example, to absorb excess new energy generation we need build a hydrogen production plant. Using Corollary 3.1, we can get the required amount of hydrogen production power and capacity to achieve complete consumption of new energy.

The price formula (9) has a defect that the item $\frac{1}{T} \sum_{s=1}^T [r(s) - w(s)]$ requires the future information of $r(t)$ and $w(t)$. To avoid this




Fig. 3. An ideal situation of perfect SE, in which the uncontrollable energy accounts for 100% of energy supply, and flexible users account for 33.64% of the total energy demand.

defect we use the prediction value b instead of $\frac{1}{T} \sum_{s=1}^T [r(s) - w(s)]$, and let

$$\pi(b, t) := 1 + \frac{1}{n} + \frac{\nu_{\mathcal{N}}(t)}{w(t) + \frac{g_{\mathcal{N}}}{T} - r(t) + b}, \quad \forall 1 \leq t \leq T. \quad (15)$$

Denote

$$\delta = \delta(b) := b - \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \quad (16)$$

as the prediction error. If the utility company adopts the price formula $\pi(b, t)$, a natural problem is that how much does the power generation cost of utility company increase compared to the perfect SE in Theorem 3.1? The following theorem provides a precise result.

Theorem 3.2: Let

$$\begin{aligned} \pi_0 &:= (\pi(b, 1), \pi(b, 2), \dots, \pi(b, T))^\top, \\ \hat{w}(t) &:= w(t) - r(t) + \frac{g_{\mathcal{N}}}{T} + b, \quad t = 1, \dots, T, \\ \sigma_i &:= \frac{g_i}{g_{\mathcal{N}} + T\delta} (\hat{w}(1), \dots, \hat{w}(T))^\top, \quad i = 1, \dots, n, \\ \sigma &:= (\sigma_1, \sigma_2, \dots, \sigma_n)^\top, \end{aligned}$$

and assume that

$$0 < \hat{w}(t) \leq (g_{\mathcal{N}} + T\delta) \min_{i \in \mathcal{N}} \left\{ \frac{\nu_i^{\max}}{g_i} \right\}, \quad \forall 1 \leq t \leq T. \quad (17)$$

Then, σ is a strict and unique Nash equilibrium of model (2) when the leader's strategy is π_0 , while the cost function of the leader is

$$u^l(\pi_0, \sigma) = \left(\frac{T\delta}{g_{\mathcal{N}} + T\delta} \right)^2 \text{Var} \left(\{w(t) - r(t)\}_{t=1}^T \right), \quad (18)$$

where

$$\begin{aligned} \text{Var} \left(\{w(t) - r(t)\}_{t=1}^T \right) &:= \\ &\frac{1}{T} \sum_{t=1}^T \left(w(t) - r(t) - \frac{1}{T} \sum_{s=1}^T [w(s) - r(s)] \right)^2 \end{aligned}$$

denotes the variance of $\{w(t) - r(t)\}_{t=1}^T$.

The proofs of Theorem 3.1, Corollary 3.1 and Theorem 3.2 are put in Subsection 4-B.

From Theorem 3.2, an interesting conclusion can be deduced that under the same prediction accuracy of new energy generation and regular user electricity consumption, the utility company's cost




Fig. 4. A numerical case of the new energy generation (top) and the regular users' electricity consumption (bottom) in 24 hours. The blue curves denote the power curves of new energy generation (top) and regular users' electricity consumption (bottom), while each scatter plot ($\bullet + \blacktriangle \star$) denotes the new energy generation (top) or regular users' electricity consumption (bottom) at each sampling slot under different sampling numbers T .

function $u^l(\boldsymbol{\pi}_0, \boldsymbol{\sigma})$ under the Nash equilibrium can be reduced greatly by shortening the length of the time slot, or increasing the sampling number T . In fact, when the prediction value of $\sum_s [r(s) - w(s)]$ in a certain time window (such as 24 hours) remains unchanged, if we increase the sampling number from T to $2T$, then the prediction error δ will decrease to half by (16). On the other hand, we recall that $w(t)$ and $r(t)$ denote the new energy generation and regular user electricity consumption at time slot t respectively, so if we decrease the length of each time slot to half of the original, the values of $w(\cdot)$ and $r(\cdot)$ will also be down by nearly half at corresponding times. By (18), the leader's cost function $u^l(\boldsymbol{\pi}_0, \boldsymbol{\sigma})$ will reduce to nearly 1/4 of the original.

To verify this conclusion we make some simulations. We predict that the 24-hour new energy generation and regular users' electricity consumption are 125GWh and 120GWh respectively in a certain area. However, their real data are 110.1GWh and 121.1GWh respectively, whose power distributions are shown in Fig. 4. Then, by (16) the difference between the prediction value and the actual value of $\sum_s [r(s) - w(s)]$ is

$$\begin{aligned} T\delta &= Tb - \sum_s [r(s) - w(s)] \\ &= (120 - 125) - (121.1 - 110.1) = -16\text{GWh}. \end{aligned}$$

Besides, the total demand of flexible electricity users is $g_N = 41.6\text{GWh}$ in 24 hours. With above data, we calculate the prediction error δ , the variance of $\{w(t) - r(t)\}_{t=1}^T$, and the cost function $u^l(\boldsymbol{\pi}_0, \boldsymbol{\sigma})$ of the utility company for different sampling numbers T , as shown in Table II. According to Table II, as the sampling number T increases, the cost function of utility company gradually decreases.

IV. PROOFS OF MAIN RESULTS

In this section, we provide the proofs of the corresponding theoretical results in Section III.

To simplify the exposition, for any $t = 1, \dots, T$ and $i = 1, \dots, n$ we set

$$\begin{aligned} \mathbf{d}_i &:= (\nu_i(1), \nu_i(2), \dots, \nu_i(T-1))^\top, \\ h_i(t) &:= \frac{1}{\tilde{w}(t)} \left[\sum_{j \neq i} \nu_j(t) + \tilde{r}(t) \right], \end{aligned} \quad (19)$$

TABLE II
THE COST FUNCTION $u^l(\boldsymbol{\pi}_0, \boldsymbol{\sigma})$ FOR DIFFERENT T .

T	24	36	48	60
$\delta(\text{GWh})$	$-2/3$	$-4/9$	$-1/3$	$-4/15$
$\text{Var}(\{w(t) - r(t)\}_{t=1}^T)(\text{GWh}^2)$	1.74	0.84	0.48	0.31
$u^l(\boldsymbol{\pi}_0, \boldsymbol{\sigma})(\text{GWh}^2)$	0.68	0.33	0.19	0.12

and

$$\begin{aligned} \boldsymbol{\mu}_i &:= \begin{pmatrix} h_i(1) - h_i(T) - \frac{2g_i}{\tilde{w}(T)} \\ h_i(2) - h_i(T) - \frac{2g_i}{\tilde{w}(T)} \\ \vdots \\ h_i(T-1) - h_i(T) - \frac{2g_i}{\tilde{w}(T)} \end{pmatrix}, \\ \mathbf{C} &:= \begin{pmatrix} \frac{1}{\tilde{w}(1)} & 0 & \cdots & 0 \\ 0 & \frac{1}{\tilde{w}(2)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{1}{\tilde{w}(T-1)} \end{pmatrix} + \frac{1}{\tilde{w}(T)} \mathbf{1}\mathbf{1}^\top. \end{aligned} \quad (20)$$

In the following discussion, the definitions of variables $\bar{\mathbf{d}}_i$ and $\bar{\mathbf{d}}'_i$ are the same as \mathbf{d}_i in (19) after replacing $\nu_i(t)$ with $\bar{\nu}_i(t)$ and $\nu'_i(t)$.

A. Proofs of Propositions 3.1-3.3

One important factor why we chose (4) as the leader's strategy is that under this strategy, the objective function of each follower is a strictly convex quadratic function, which can be formulated as the following lemma:

Lemma 4.1: Assume that the leader's strategy adopts (4), and each follower i 's strategy ν_i satisfies $\mathbf{1}^\top \nu_i = g_i$. Then, the objective function $u_i^f(\boldsymbol{\pi}, \nu_i, \nu_{-i})$ of each follower i in model (2) can be written as

$$u_i^f(\boldsymbol{\pi}, \nu_i, \nu_{-i}) = \mathbf{d}_i^\top \mathbf{C} \mathbf{d}_i + \boldsymbol{\mu}_i^\top \mathbf{d}_i + h_i(T)g_i + \frac{g_i^2}{\tilde{w}(T)}. \quad (21)$$

In addition, if $\tilde{w}(t) > 0$ for any $1 \leq t \leq T$, then $u_i^f(\boldsymbol{\pi}, \nu_i, \nu_{-i})$ is a strictly convex quadratic function.

The proof of Lemma 4.1 is postponed to Appendix A-A.

According to Lemma 4.1 and the convex analysis method [45], the objective function of each follower i has a unique global minimum point $\bar{\mathbf{d}}_i$ satisfying

$$2\mathbf{C}\bar{\mathbf{d}}_i + \boldsymbol{\mu}_i = 0. \quad (22)$$

We get that (7) is the unique solution of linear equation system formed by uniting (22) of all followers.

Lemma 4.2: If $\tilde{w}(t) > 0$ for any $1 \leq t \leq T$, then the strategy combination (7) is the unique solution of linear equation system

$$2\mathbf{C}\bar{\mathbf{d}}_i + \boldsymbol{\mu}_i = 0, \quad i = 1, 2, \dots, n.$$

The proof of Lemma 4.2 is postponed to Appendix A-B.

To prove the uniqueness of the Nash equilibrium in game model (6) we need the well-known Banach's fixed point theorem:

Lemma 4.3 (Banach's Fixed Point Theorem [46]): Let (X, dist) be a complete metric space and let $\tilde{F} : X \rightarrow X$ be a contraction mapping on X , that is, there exists a positive constant $\kappa < 1$ such that

$$\text{dist}(\tilde{F}(x), \tilde{F}(y)) \leq \kappa \text{dist}(x, y), \quad \forall x, y \in X.$$

Then \tilde{F} has a unique fixed point $x \in X$ (such that $\tilde{F}(x) = x$).

Now we construct a contraction mapping for the game model (6). For each user i , we define its strategy hyperplane by $\mathbb{H}_i := \{\mathbf{x} \in \mathbb{R}^T : \mathbf{1}^\top \mathbf{x} = g_i\}$. Set $\mathbb{H}_{-i} := \mathbb{H}_1 \times \cdots \times \mathbb{H}_{i-1} \times \mathbb{H}_{i+1} \times \cdots \times \mathbb{H}_n$ and $\mathbb{H} := \mathbb{H}_1 \times \cdots \times \mathbb{H}_n$. By Lemma 4.1 and (22), we know that for each user i and any strategy combination $\boldsymbol{\nu}_{-i} \in \mathbb{H}_{-i}$ of other users, the best response strategy of user i in \mathbb{H}_i is unique. Thus, for any $\boldsymbol{\nu} \in \mathbb{H}$, we set the ‘‘sequential’’ best response mapping $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)^\top : \mathbb{H} \rightarrow \mathbb{H}$ with

$$\begin{aligned} \mathbf{f}_i(\boldsymbol{\nu}) := & \arg \min_{\boldsymbol{\nu}_i \in \mathbb{H}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, (\mathbf{f}_1(\boldsymbol{\nu}), \dots, \mathbf{f}_{i-1}(\boldsymbol{\nu}), \boldsymbol{\nu}_{i+1}, \dots, \boldsymbol{\nu}_n)^\top) \end{aligned} \quad (23)$$

for $i = 1, 2, \dots, n$. We show that:

Lemma 4.4: \mathbf{F} is a contraction mapping on \mathbb{H} .

The proof of Lemma 4.4 is postponed to Appendix A-C.

Proof of Proposition 3.1: For any user $i \in \mathcal{N}$ and any strategy $\boldsymbol{\nu}'_i = (\nu'_i(1), \dots, \nu'_i(T))^\top \neq \boldsymbol{\nu}_i$ with $\mathbf{1}^\top \boldsymbol{\nu}'_i = g_i$, according to (21) we have

$$\begin{aligned} & u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}'_i, \boldsymbol{\nu}_{-i}) - u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}) \\ &= \mathbf{d}'_i{}^\top \mathbf{C} \mathbf{d}'_i + \boldsymbol{\mu}_i{}^\top \mathbf{d}'_i + h_i(T)g_i + \frac{g_i^2}{\tilde{w}(T)} \\ & \quad - \left(\bar{\mathbf{d}}_i{}^\top \mathbf{C} \bar{\mathbf{d}}_i + \boldsymbol{\mu}_i{}^\top \bar{\mathbf{d}}_i + h_i(T)g_i + \frac{g_i^2}{\tilde{w}(T)} \right) \\ &= (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \mathbf{C} (\mathbf{d}'_i - \bar{\mathbf{d}}_i) + \mathbf{d}'_i{}^\top \mathbf{C} \bar{\mathbf{d}}_i + \bar{\mathbf{d}}_i{}^\top \mathbf{C} \mathbf{d}'_i \\ & \quad - 2\bar{\mathbf{d}}_i{}^\top \mathbf{C} \bar{\mathbf{d}}_i + (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \boldsymbol{\mu}_i \\ &= (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \mathbf{C} (\mathbf{d}'_i - \bar{\mathbf{d}}_i) + 2(\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \mathbf{C} \bar{\mathbf{d}}_i \\ & \quad + (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \boldsymbol{\mu}_i \\ &= (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \mathbf{C} (\mathbf{d}'_i - \bar{\mathbf{d}}_i) + (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top (2\mathbf{C} \bar{\mathbf{d}}_i + \boldsymbol{\mu}_i) \\ &= (\mathbf{d}'_i - \bar{\mathbf{d}}_i)^\top \mathbf{C} (\mathbf{d}'_i - \bar{\mathbf{d}}_i) > 0, \end{aligned}$$

where the third equality and the last inequality use the fact that \mathbf{C} is symmetric positive definite, and the last equality uses (22). So $\boldsymbol{\nu}$ is a strict Nash equilibrium of model (6) when the leader’s strategy is $\boldsymbol{\pi}$.

It remains to prove the uniqueness of the Nash equilibrium of model (6). Assume that $\tilde{\boldsymbol{\nu}} \in \mathbb{H}$ is an arbitrary Nash equilibria, then by (3) we have

$$\tilde{\boldsymbol{\nu}}_i = \arg \min_{\boldsymbol{\nu}_i \in \mathbb{H}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \tilde{\boldsymbol{\nu}}_{-i}), \quad \forall i \in \mathcal{N}. \quad (24)$$

Here we recall that the best response strategy of user i in \mathbb{H}_i is unique. By (23) and (24) we can sequentially get

$$\mathbf{f}_1(\tilde{\boldsymbol{\nu}}) = \arg \min_{\boldsymbol{\nu}_1 \in \mathbb{H}_1} u_1^f(\boldsymbol{\pi}, \boldsymbol{\nu}_1, (\tilde{\boldsymbol{\nu}}_2, \dots, \tilde{\boldsymbol{\nu}}_n)^\top) = \tilde{\boldsymbol{\nu}}_1$$

and

$$\begin{aligned} \mathbf{f}_i(\tilde{\boldsymbol{\nu}}) &= \arg \min_{\boldsymbol{\nu}_i \in \mathbb{H}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, (\mathbf{f}_1(\tilde{\boldsymbol{\nu}}), \dots, \mathbf{f}_{i-1}(\tilde{\boldsymbol{\nu}}), \tilde{\boldsymbol{\nu}}_{i+1}, \dots, \tilde{\boldsymbol{\nu}}_n)^\top) \\ &= \arg \min_{\boldsymbol{\nu}_i \in \mathbb{H}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, (\tilde{\boldsymbol{\nu}}_1, \dots, \tilde{\boldsymbol{\nu}}_{i-1}, \tilde{\boldsymbol{\nu}}_{i+1}, \dots, \tilde{\boldsymbol{\nu}}_n)^\top) \\ &= \tilde{\boldsymbol{\nu}}_i, \quad i = 2, \dots, n. \end{aligned}$$

So we have $\mathbf{F}(\tilde{\boldsymbol{\nu}}) = \tilde{\boldsymbol{\nu}}$, that is, $\tilde{\boldsymbol{\nu}}$ is a fixed point of \mathbf{F} .

Therefore, all Nash equilibria of model (6) are fixed points of the mapping \mathbf{F} . Based on Lemmas 4.3-4.4, \mathbf{F} has a unique fixed point, thus model (6) has a unique Nash equilibrium. Since $\boldsymbol{\nu} \in \mathbb{H}$ is a strict Nash equilibrium for model (6), $\boldsymbol{\nu}$ must be the unique Nash equilibrium of model (6). \square

Set $\mathbf{d} := (\mathbf{d}_1^\top, \dots, \mathbf{d}_n^\top)^\top$ to be the strategy vector of all followers, and

$$\mathbf{g}(\mathbf{d}) := \left(\frac{\partial u_1^f(\boldsymbol{\pi}, \boldsymbol{\nu}_1, \boldsymbol{\nu}_{-1})}{\partial \mathbf{d}_1^\top}, \dots, \frac{\partial u_n^f(\boldsymbol{\pi}, \boldsymbol{\nu}_n, \boldsymbol{\nu}_{-n})}{\partial \mathbf{d}_n^\top} \right)^\top.$$

Further, we make $\mathbf{J}(\mathbf{d})$ the Jacobian with respect to \mathbf{d} of $\mathbf{g}(\mathbf{d})$, i.e.,

$$\mathbf{J}(\mathbf{d}) := \begin{pmatrix} \frac{\partial^2 u_1^f(\boldsymbol{\pi}, \boldsymbol{\nu}_1, \boldsymbol{\nu}_{-1})}{\partial \mathbf{d}_1^2} & \cdots & \frac{\partial^2 u_1^f(\boldsymbol{\pi}, \boldsymbol{\nu}_1, \boldsymbol{\nu}_{-1})}{\partial \mathbf{d}_1 \partial \mathbf{d}_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 u_n^f(\boldsymbol{\pi}, \boldsymbol{\nu}_n, \boldsymbol{\nu}_{-n})}{\partial \mathbf{d}_n \partial \mathbf{d}_1} & \cdots & \frac{\partial^2 u_n^f(\boldsymbol{\pi}, \boldsymbol{\nu}_n, \boldsymbol{\nu}_{-n})}{\partial \mathbf{d}_n^2} \end{pmatrix}.$$

In the model (2), for each user $i \in \mathcal{N}$, we define the strategy space $\Omega_i := \mathbb{H}_i \cap [0, \nu_i^{\max}]^T$ as the set of its feasible strategies, which is a bounded closed convex set. According to Theorem 2 in [47], we can get that the Nash equilibrium of model (2) is unique, which can be formulated as the following lemma:

Lemma 4.5 (Theorem 2 in [47]): For each user $i \in \mathcal{N}$, if $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$ is a convex function on the bounded closed convex region Ω_i , and $\mathbf{J}(\mathbf{d})$ is symmetric positive definite for all \mathbf{d} , then model (2) admits a unique Nash equilibrium.

Proof of Proposition 3.2: For any $1 \leq i \leq n$, according to Lemma 4.1, $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$ is a strictly convex quadratic function. By (32) and (34), we have

$$\mathbf{g}(\mathbf{d}) = \begin{pmatrix} \mathbf{C} [2\mathbf{d}_1 + \sum_{j \neq 1} \mathbf{d}_j] - \tilde{\mathbf{g}}_1 \\ \vdots \\ \mathbf{C} [2\mathbf{d}_n + \sum_{j \neq n} \mathbf{d}_j] - \tilde{\mathbf{g}}_n \end{pmatrix}$$

and

$$\mathbf{J}(\mathbf{d}) = \begin{pmatrix} 2\mathbf{C} & \cdots & \mathbf{C} \\ \vdots & \ddots & \vdots \\ \mathbf{C} & \cdots & 2\mathbf{C} \end{pmatrix} = \begin{pmatrix} 2 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 2 \end{pmatrix} \otimes \mathbf{C},$$

where ‘‘ \otimes ’’ denotes the Kronecker product. Since the Kronecker product of two positive definite matrices is still positive definite [48], we can get that $\mathbf{J}(\mathbf{d})$ is symmetric positive definite for all \mathbf{d} . According to Lemma 4.5, we can conclude that model (2) admits a unique Nash equilibrium denoted as $\boldsymbol{\nu}$.

Now we show that $\boldsymbol{\nu}$ is a strict Nash equilibrium of model (2). For any user i and any strategy $\boldsymbol{\nu}'_i = (\nu'_i(1), \dots, \nu'_i(T))^\top \in \Omega_i$ with $\boldsymbol{\nu}'_i \neq \boldsymbol{\nu}_i$, by (3) we have

$$u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}'_i, \boldsymbol{\nu}_{-i}) \geq u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}).$$

It is clear that $\frac{\boldsymbol{\nu}'_i + \boldsymbol{\nu}_i}{2} \in \Omega_i$, so

$$u_i^f(\boldsymbol{\pi}, \frac{\boldsymbol{\nu}'_i + \boldsymbol{\nu}_i}{2}, \boldsymbol{\nu}_{-i}) \geq u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}). \quad (25)$$

If $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}'_i, \boldsymbol{\nu}_{-i}) = u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$, then according to strict convexity of $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$, we can get

$$\begin{aligned} u_i^f(\boldsymbol{\pi}, \frac{\boldsymbol{\nu}'_i + \boldsymbol{\nu}_i}{2}, \boldsymbol{\nu}_{-i}) &< \frac{u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}'_i, \boldsymbol{\nu}_{-i}) + u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})}{2} \\ &= u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}), \end{aligned}$$

which contradicts (25). So

$$u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}'_i, \boldsymbol{\nu}_{-i}) > u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}),$$

which implies $\boldsymbol{\nu}$ is a strict Nash equilibrium of model (2). \square

Proof of Proposition 3.3: Substituting (11) into the problem (1), we have the controllable energy generation $c^*(t)$ satisfies:

$$\begin{aligned} & c^*(t) - \bar{c} \\ &= \sum_{i \in \mathcal{N}} \nu_i^*(t) + r(t) - w(t) - \frac{g_{\mathcal{N}}}{T} - \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \\ &= \frac{g_{\mathcal{N}}}{T} + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] - \frac{g_{\mathcal{N}}}{T} - \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \\ &= 0, \quad \forall 1 \leq t \leq T, \end{aligned}$$

which indicates the leader's cost function $u^l(\pi^*, \nu^*) = 0$. \square

B. Proofs of Theorem 3.1, Corollary 3.1 and Theorem 3.2

Proof of Theorem 3.1: i) Assume the leader's strategy is π^* . First, by Proposition 3.1 and the discussions from (8) to (11) we have ν^* is a strict Nash equilibrium of model (6). Also, by (11) and (12) we have

$$0 < \nu_i^*(t) \leq g_i \min_{j \in \mathcal{N}} \left\{ \frac{\nu_j^{\max}}{g_j} \right\} \leq g_i \frac{\nu_i^{\max}}{g_i} = \nu_i^{\max},$$

which indicates that $\nu_i^*(t)$ satisfies the bound constraint condition in model (2). Therefore, ν^* is a strict Nash equilibrium of model (2). By Proposition 3.2, ν^* is the unique Nash equilibrium of model (2).

ii) By i) and Proposition 3.3, we have (π^*, ν^*) is a perfect SE of the Stackelberg game (1)-(2). \square

Proof of Corollary 3.1: Under the condition (14), the price function (9) becomes

$$\begin{aligned} \pi^*(t) &= 1 + \frac{1}{n} + \frac{\nu_{\mathcal{N}}(t)}{w(t) + \frac{g_{\mathcal{N}}}{T} - r(t) + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)]} \\ &= 1 + \frac{1}{n} + \frac{\nu_{\mathcal{N}}(t)}{w(t) - r(t)}, \quad t = 1, \dots, T, \end{aligned}$$

and the Nash equilibrium (11) is

$$\begin{aligned} \nu_i^*(t) &= \frac{g_i}{T} + \frac{g_i}{g_{\mathcal{N}}} [w(t) - r(t)] + \frac{g_i}{g_{\mathcal{N}} T} \sum_{s=1}^T [r(s) - w(s)] \\ &= \frac{g_i}{g_{\mathcal{N}}} [w(t) - r(t)], \quad i = 1, \dots, n, t = 1, \dots, T. \end{aligned}$$

Therefore, according to the inequality (13) and Theorem 3.1, χ^* is a strict and unique Nash equilibrium of model (2) when the leader's strategy is π_0^* , and (π_0^*, χ^*) is a perfect SE of the Stackelberg game (1)-(2). \square

Proof of Theorem 3.2: When (15) is chosen as the leader's strategy, by (4) and (5), we have $\tilde{w}(t) = w(t) - r(t) + \frac{g_{\mathcal{N}}}{T} + b$ and $\tilde{r}(t) = \frac{n+1}{n} \tilde{w}(t)$. According to Proposition 3.1, the strict and unique Nash equilibrium in (7) becomes

$$\begin{aligned} \nu_i(t) &= \frac{\tilde{w}(t)}{\sum_s \tilde{w}(s)} g_i + \frac{1}{(n+1) \sum_s \tilde{w}(s)} \\ &\quad \times \sum_{s=1}^T \tilde{w}(s) \tilde{w}(t) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(t)}{\tilde{w}(t)} \right] \\ &= \frac{g_i}{g_{\mathcal{N}} + Tb + \sum_{s=1}^T [w(s) - r(s)]} \\ &\quad \times [w(t) - r(t) + \frac{g_{\mathcal{N}}}{T} + b] \\ &= \frac{g_i}{g_{\mathcal{N}} + T\delta} [w(t) - r(t) + \frac{g_{\mathcal{N}}}{T} + b], \\ &\quad i = 1, \dots, n, t = 1, \dots, T, \end{aligned} \tag{26}$$

where the last equality uses (16). With the condition (17) and the similar process of the proof of Theorem 3.1, we can conclude that

σ is a strict and unique Nash equilibrium of model (2) when the leader's strategy is π_0 .

Further, for any $1 \leq t \leq T$, substituting (26) into the problem (1), we have the controllable energy generation $c(t)$ satisfies:

$$\begin{aligned} & c(t) - \bar{c} \\ &= \sum_{i \in \mathcal{N}} \nu_i(t) + r(t) - w(t) - \frac{g_{\mathcal{N}}}{T} - \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \\ &= \frac{g_{\mathcal{N}}}{g_{\mathcal{N}} + T\delta} \left(w(t) - r(t) + \frac{g_{\mathcal{N}}}{T} + \delta + \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \right) \\ &\quad + r(t) - w(t) - \frac{g_{\mathcal{N}}}{T} - \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] \\ &= \left(\frac{g_{\mathcal{N}}}{g_{\mathcal{N}} + T\delta} - 1 \right) \frac{1}{T} \sum_{s=1}^T [r(s) - w(s)] + \left(\frac{g_{\mathcal{N}}}{g_{\mathcal{N}} + T\delta} - 1 \right) \\ &\quad \times [w(t) - r(t)] \\ &= \frac{\delta}{g_{\mathcal{N}} + T\delta} \sum_{s=1}^T [w(s) - r(s)] - \frac{T\delta}{g_{\mathcal{N}} + T\delta} [w(t) - r(t)], \end{aligned}$$

where the second equality uses (16). From this, the cost function of the leader is

$$\begin{aligned} u^l(\pi_0, \sigma) &= \frac{1}{T} \sum_{t=1}^T [c(t) - \bar{c}]^2 \\ &= \frac{1}{T} \sum_{t=1}^T \left(\frac{\delta}{g_{\mathcal{N}} + T\delta} \sum_{s=1}^T [w(s) - r(s)] - \frac{T\delta}{g_{\mathcal{N}} + T\delta} \right. \\ &\quad \left. \times [w(t) - r(t)] \right)^2 \\ &= \frac{(T\delta)^2}{(g_{\mathcal{N}} + T\delta)^2} \frac{1}{T} \sum_{t=1}^T \left(w(t) - r(t) - \frac{1}{T} \sum_{s=1}^T [w(s) - r(s)] \right)^2 \\ &= \left(\frac{T\delta}{g_{\mathcal{N}} + T\delta} \right)^2 \text{Var} \left(\{w(t) - r(t)\}_{t=1}^T \right). \end{aligned} \quad \square$$

V. NUMERICAL SOLUTION OF SE FOR MODEL (1)-(2) WITHOUT CONDITION (12)

In Subsection 3-B we give the analytic expression of SE for the Stackelberg game (1)-(2) based on the inequality condition (12). A natural problem is whether we can still obtain the SE if the condition (12) is not satisfied. The analytic expression of SE should be hard for this case, however we can provide a numerical solution.

First, let us review the optimization problem (1) of the leader side. In fact, the leader optimizes its objective by adjusting the electricity demand of followers through price function. If we omit the price function, the optimization problem (1) can be rewritten as

$$\begin{aligned} & \min_{\{\nu_{\mathcal{N}}(t)\}_{t=1}^T} \frac{1}{T} \sum_{t=1}^T [c(t) - \bar{c}]^2, \\ & \text{s.t.} \quad \sum_{t=1}^T \nu_{\mathcal{N}}(t) = g_{\mathcal{N}}, \\ & \quad \nu_{\mathcal{N}}(t) + r(t) = w(t) + c(t), \\ & \quad 0 \leq \nu_{\mathcal{N}}(t) \leq \nu_{\mathcal{N}}^{\max}, \quad t = 1, \dots, T, \end{aligned} \tag{27}$$

where $\nu_{\mathcal{N}}^{\max} := \sum_{i \in \mathcal{N}} \nu_i^{\max}$.

The optimization problem (27) is a quadratic program with a strictly convex quadratic function. According to the convex analysis method [45], the problem (27) has a unique global optimal solution, which we denote as $\nu_{\mathcal{N}}^* = (\nu_{\mathcal{N}}^*(1), \nu_{\mathcal{N}}^*(2), \dots, \nu_{\mathcal{N}}^*(T))^T$, and it

can be numerically solved by several methods, such as the active-set method [49].

Algorithm 1: Best response algorithm for solving the game model (2)

Input: The number of flexible users n , the number of time periods T , the value of g_i and ν_i^{\max} for each user $i \in \mathcal{N}$, the pricing sequence $(\tilde{r}(t), \tilde{w}(t))_{1 \leq t \leq T}$, the tolerance ε_0 .

Initialization: Set $k = 0$. Set the initial strategy for each flexible user $i \in \mathcal{N}$ as the average strategy $\nu_i^{(0)} = (\frac{g_i}{T}, \dots, \frac{g_i}{T})^\top$.

While not converged do

1. **for** $i = 1, 2, \dots, n$ **do**
2. Calculate the best response strategy $\nu_i^{(k+1)}$ for game model (2) when fix $\nu_{-i} = (\nu_1^{(k+1)}, \dots, \nu_{i-1}^{(k+1)}, \nu_{i+1}^{(k)}, \dots, \nu_n^{(k)})^\top$.
3. **end for**
4. Calculate $er = \|\nu^{(k+1)} - \nu^{(k)}\|_1$.
5. Update $k = k + 1$.

Until $er < \varepsilon_0$

Output: The approximative Nash equilibrium $\nu^{(k)}$.

Algorithm 2: Search algorithm for the optimal pricing sequence

Input: The number of flexible users n , the number of time periods T , the value of g_i and ν_i^{\max} for each user $i \in \mathcal{N}$, the regular user demand and new energy supply sequence $(r(t), w(t))_{1 \leq t \leq T}$, the tolerance ε_0 .

Initialization:

1. Set $k = 0$. Randomly initialize a positive sequence $(\tilde{r}^{(0)}(t), \tilde{w}(t))_{1 \leq t \leq T}$.
2. Solve optimization problem (27) and obtain the unique global optimal solution $\nu_{\mathcal{N}}^*$.
3. Calculate Nash equilibrium $\nu^{(0)} = \text{NE}(\tilde{r}^{(0)}(t), \tilde{w}(t))$ (Alg. 1) for the game model (2), and denote the total electricity demand of flexible users in each time slot as $\nu_{\mathcal{N}}^{(0)} = (\nu_{\mathcal{N}}^{(0)}(1), \nu_{\mathcal{N}}^{(0)}(2), \dots, \nu_{\mathcal{N}}^{(0)}(T))^\top$.
4. Calculate $er = \|\nu_{\mathcal{N}}^{(0)} - \nu_{\mathcal{N}}^*\|_\infty$.

While not converged do

5. **for** $t = 1, 2, \dots, T$ **do**
6. **if** $\nu_{\mathcal{N}}^{(k)}(t) > \nu_{\mathcal{N}}^*(t)$, **then let**
7. $\tilde{r}^{(k+1)}(t) = \tilde{r}^{(k)}(t) + \varepsilon_0$;
8. **if** $\nu_{\mathcal{N}}^{(k)}(t) < \nu_{\mathcal{N}}^*(t)$, **then let**
9. $\tilde{r}^{(k+1)}(t) = \tilde{r}^{(k)}(t) - \varepsilon_0$.
10. **end for**
11. Calculate Nash equilibrium $\nu^{(k+1)} = \text{NE}(\tilde{r}^{(k+1)}(t), \tilde{w}(t))$ (Alg. 1) and total electricity demand $\nu_{\mathcal{N}}^{(k+1)}$ for game model (2).
12. Calculate $er = \|\nu_{\mathcal{N}}^{(k+1)} - \nu_{\mathcal{N}}^*\|_\infty$.
13. Update $k = k + 1$.

Until $er < \varepsilon_0$

Output: The optimal pricing sequence $\{\tilde{r}^{(k)}(t)\}_{1 \leq t \leq T}$.

Next we consider the game model (2) of followers. By (4) and (5) the price formula can be written as

$$\pi(\nu_{\mathcal{N}}(t), \tilde{r}(t), \tilde{w}(t)) = \frac{\nu_{\mathcal{N}}(t) + \tilde{r}(t)}{\tilde{w}(t)}, t = 1, \dots, T. \quad (28)$$

By Proposition 3.2, the model (2) with the leader's strategy (28) admits a strict and unique Nash equilibrium. Since the analytic expression of the Nash equilibrium is unknown, we try to find its numerical solution. Algorithm 1 provides a numerical solution of model (2): when we input a sequence $(\tilde{r}(t), \tilde{w}(t))_{1 \leq t \leq T}$ for (28), a Nash equilibrium of the strategy combination of all flexible users can be approximatively obtained through Algorithm 1.

Finally, we need to find an optimal pricing formula $\pi^*(t)$ to obtain the numerical solution of SE for Stackelberg game (1)-(2). That is to say, we manage to find the optimal sequence $(\tilde{r}^*(t), \tilde{w}^*(t))_{1 \leq t \leq T}$

so that the Nash equilibrium ν^* of game model (2) with the leader's strategy (28) satisfies

$$\sum_{i \in \mathcal{N}} \nu_i^*(t) = \nu_{\mathcal{N}}^*(t), \quad \forall 1 \leq t \leq T. \quad (29)$$

The detailed search algorithm is shown in Algorithm 2. In this algorithm, inspired by (7), we can randomly initialize a positive sequence $(\tilde{r}(t), \tilde{w}(t))_{1 \leq t \leq T}$, and then regulate only the sequence $\{\tilde{r}(t)\}_{t=1, \dots, T}$ to adjust the Nash equilibrium of game model (2), until (29) approximatively holds.

We provide a simulation example as follows. Set the number of flexible electricity users as $n = 20$, the number of scheduling time window $T = 24$, with each time slot being 1 hour. Moreover, the total electricity demands of flexible users are 1.55, 1.49, 1.04, 4.13, 3.7, 2.03, 2.29, 0.92, 4.47, 0.81, 8.98, 0.02, 0.29, 0.37, 0.13, 4.64, 4.65, 4.83, 4.49, and 0.26, respectively. For any $1 \leq i \leq n$, we set $\nu_i^{\max} = 2g_i/T$. The regular user electricity consumption and new energy generation sequence $(r(t), w(t))_{1 \leq t \leq T}$ are shown in Fig. 5 (a). It can be verified that the inequalities (12) do not hold in this situation. Through Algorithms 1-2, we can achieve the numerical solution of SE for model (1)-(2), in which the controllable energy generation $\{c(t)\}_{t=1, \dots, T}$ and the Nash equilibrium $\{\nu_i^*(t)\}_{i=1, \dots, n; t=1, \dots, T}$ of game model (2) are shown in Fig. 5 (a). Correspondingly, Fig. 5 (b) shows the variation curve of the utility company's cost $u^l(\pi^{(k)}, \nu^{(k)})$ with iteration number k , from which we can see that the cost function quickly converges to the optimal cost of the utility company, which can be obtained by solving problem (27).

From above example, for the case when the condition (12) is not satisfied, the optimal pricing formula for the utility company may be achieved by the Algorithms 1-2.

VI. CONCLUSION

With the rapid development of smart grid, RTP has become one of the most important DR strategies. In this paper, we propose a Stackelberg game model to describe the interaction between utility company and users, and provide analytical expressions and numerical solutions for the optimal pricing formula under different supply and demand relationships. Under our optimal pricing scheme, flexible electricity users have an unique and strict Nash equilibrium for electricity demand, in which new energy can be sufficiently utilized and the cost of traditional power generation can be minimized, thereby the social benefit can be maximized.

This work can be extended from different aspects. For example, we can incorporate energy storage companies into the game, consider the equilibrium solving problem when new energy generation is a random variable, and consider the game behavior between different power suppliers. These questions are meaningful and provide valuable directions for future research.

APPENDIX A

A. Proof of Lemma 4.1

With (5), the price formula (4) can be rewritten as

$$\pi(t) = \frac{\nu_{\mathcal{N}}(t) + \tilde{r}(t)}{\tilde{w}(t)}, \quad \forall 1 \leq t \leq T.$$




Fig. 5. An example of achieving numerical solution for SE through Algorithm 1-2, in which the uncontrollable numerical accounts for 62.22% of energy supply, and flexible users account for 29.41% of the total energy demand. The curves of $\{w(t), r(t), c(t), \nu_i^*(t)\}$ with respect to time slot t are shown in (a), while the cost function $u^l(\pi^{(k)}, \nu^{(k)})$ with respect to iteration number k is shown in (b).

Substitute this into (2), the cost function of each user i can be rewritten as

$$\begin{aligned}
 u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}) &= \sum_{t=1}^T \frac{\nu_{\mathcal{N}}(t) + \tilde{r}(t)}{\tilde{w}(t)} \nu_i(t) \\
 &= \sum_{t=1}^T \frac{1}{\tilde{w}(t)} \left[\nu_i(t) \sum_{j \in \mathcal{N}} \nu_j(t) + \tilde{r}(t) \nu_i(t) \right] \\
 &= \sum_{t=1}^T \frac{1}{\tilde{w}(t)} \left\{ \nu_i^2(t) + \left[\sum_{j \neq i} \nu_j(t) + \tilde{r}(t) \right] \nu_i(t) \right\}.
 \end{aligned} \tag{30}$$

According to the constraint condition $\mathbf{1}^\top \boldsymbol{\nu}_i = g_i$, we can obtain $\nu_i(T) = g_i - \sum_{t=1}^{T-1} \nu_i(t)$. From this and (19) we have

$$\begin{aligned}
 &\frac{1}{\tilde{w}(T)} \left\{ \nu_i^2(T) + \left[\sum_{j \neq i} \nu_j(T) + \tilde{r}(T) \right] \nu_i(T) \right\} \\
 &= \frac{\nu_i^2(T)}{\tilde{w}(T)} + h_i(T) \nu_i(T) \\
 &= \frac{1}{\tilde{w}(T)} \left[g_i - \sum_{t=1}^{T-1} \nu_i(t) \right]^2 + h_i(T) \left[g_i - \sum_{t=1}^{T-1} \nu_i(t) \right].
 \end{aligned}$$

Substitute this into (30) we get

$$\begin{aligned}
 &u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i}) \\
 &= \sum_{t=1}^{T-1} \left[\frac{1}{\tilde{w}(t)} \nu_i^2(t) + h_i(t) \nu_i(t) \right] + \frac{1}{\tilde{w}(T)} \\
 &\quad \times \left[g_i - \sum_{t=1}^{T-1} \nu_i(t) \right]^2 + h_i(T) \left[g_i - \sum_{t=1}^{T-1} \nu_i(t) \right] \\
 &= \sum_{t=1}^{T-1} \frac{1}{\tilde{w}(t)} \nu_i^2(t) + \frac{1}{\tilde{w}(T)} \left[\sum_{t=1}^{T-1} \nu_i(t) \right]^2 \\
 &\quad + \sum_{t=1}^{T-1} \left[h_i(t) - h_i(T) - \frac{2g_i}{\tilde{w}(T)} \right] \nu_i(t) + h_i(T) g_i + \frac{g_i^2}{\tilde{w}(T)} \\
 &= \mathbf{d}_i^\top \mathbf{C} \mathbf{d}_i + \boldsymbol{\mu}_i^\top \mathbf{d}_i + h_i(T) g_i + \frac{g_i^2}{\tilde{w}(T)},
 \end{aligned}$$

where the last line uses (20).

Now we show the convex property of $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$. By (19)-(20), the vector $\boldsymbol{\mu}_i$ does not depend on \mathbf{d}_i , which means

$$\frac{\partial \boldsymbol{\mu}_i^\top \mathbf{d}_i}{\partial \mathbf{d}_i} = \boldsymbol{\mu}_i, \quad \frac{\partial^2 \boldsymbol{\mu}_i^\top \mathbf{d}_i}{\partial \mathbf{d}_i^2} = \mathbf{0}, \quad \forall i \in \mathcal{N}. \tag{31}$$

Based on (21) and (31), we can get

$$\begin{cases} \frac{\partial u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})}{\partial \mathbf{d}_i} = 2\mathbf{C} \mathbf{d}_i + \boldsymbol{\mu}_i, \\ \frac{\partial^2 u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})}{\partial \mathbf{d}_i^2} = 2\mathbf{C}, \end{cases} \quad \forall i \in \mathcal{N}. \tag{32}$$

Also, by (20) and the condition that $\tilde{w}(t) > 0$ for any $t = 1, \dots, T$, we have

$$\mathbf{x}^\top \mathbf{C} \mathbf{x} = \sum_{t=1}^{T-1} \frac{1}{\tilde{w}(t)} x_t^2 + \frac{1}{\tilde{w}(T)} \left[\sum_{t=1}^{T-1} x_t \right]^2 > 0$$

for any $\mathbf{x} = (x_1, x_2, \dots, x_{T-1})^\top \in \mathbb{R}^{T-1}$ with $\mathbf{x} \neq \mathbf{0}$, which means \mathbf{C} is symmetric positive definite. From this and (32), the objective function $u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$ is a strictly convex quadratic function.

B. Proof of Lemma 4.2

For each follower i , by (19) and $\mathbf{1}^\top \boldsymbol{\nu}_j = g_j, j = 1, \dots, n$, we have

$$\begin{aligned}
 h_i(T) &= \frac{1}{\tilde{w}(T)} \left[\sum_{j \neq i} \nu_j(T) + \tilde{r}(T) \right] \\
 &= \frac{1}{\tilde{w}(T)} \sum_{j \neq i} \left[g_j - \sum_{t=1}^{T-1} \nu_j(t) \right] + \frac{\tilde{r}(T)}{\tilde{w}(T)} \\
 &= \frac{g_{\mathcal{N}} - g_i + \tilde{r}(T)}{\tilde{w}(T)} - \frac{1}{\tilde{w}(T)} \left[\sum_{j \neq i} \sum_{t=1}^{T-1} \nu_j(t) \right].
 \end{aligned} \tag{33}$$

Combining (20) and (33), $\boldsymbol{\mu}_i$ can be written as

$$\begin{aligned}
\boldsymbol{\mu}_i &= \begin{pmatrix} h_i(1) \\ \vdots \\ h_i(T-1) \end{pmatrix} - [h_i(T) + \frac{2g_i}{\tilde{w}(T)}] \mathbf{1} \\
&= \begin{pmatrix} \frac{1}{\tilde{w}(1)} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \frac{1}{\tilde{w}(T-1)} \end{pmatrix} \begin{pmatrix} \sum_{j \neq i} \nu_j(1) \\ \vdots \\ \sum_{j \neq i} \nu_j(T-1) \end{pmatrix} \\
&\quad + \begin{pmatrix} \frac{\tilde{r}(1)}{\tilde{w}(1)} \\ \vdots \\ \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \end{pmatrix} + \frac{1}{\tilde{w}(T)} \left[\sum_{j \neq i} \sum_{t=1}^{T-1} \nu_j(t) \right] \mathbf{1} \\
&\quad - \frac{g_{\mathcal{N}} + g_i + \tilde{r}(T)}{\tilde{w}(T)} \mathbf{1} \\
&= \left\{ \begin{pmatrix} \frac{1}{\tilde{w}(1)} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \frac{1}{\tilde{w}(T-1)} \end{pmatrix} + \frac{1}{\tilde{w}(T)} \mathbf{1} \mathbf{1}^\top \right\} \\
&\quad \times \begin{pmatrix} \sum_{j \neq i} \nu_j(1) \\ \vdots \\ \sum_{j \neq i} \nu_j(T-1) \end{pmatrix} + \begin{pmatrix} \frac{\tilde{r}(1)}{\tilde{w}(1)} \\ \vdots \\ \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \end{pmatrix} \\
&\quad - \frac{g_{\mathcal{N}} + g_i + \tilde{r}(T)}{\tilde{w}(T)} \mathbf{1} \\
&= \mathbf{C} \sum_{j \neq i} \mathbf{d}_j - \tilde{\mathbf{g}}_i,
\end{aligned} \tag{34}$$

where $\tilde{\mathbf{g}}_i = \frac{g_{\mathcal{N}} + g_i + \tilde{r}(T)}{\tilde{w}(T)} \mathbf{1} - \left(\frac{\tilde{r}(1)}{\tilde{w}(1)}, \dots, \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \right)^\top$, and the last equality uses the fact

$$\begin{aligned}
&\left(\sum_{j \neq i} \nu_j(1), \dots, \sum_{j \neq i} \nu_j(T-1) \right)^\top \\
&= \sum_{j \neq i} \left(\nu_j(1), \dots, \nu_j(T-1) \right)^\top = \sum_{j \neq i} \mathbf{d}_j.
\end{aligned}$$

Using (34), Eq. (22) can be rewritten as

$$\mathbf{C} \left[2\bar{\mathbf{d}}_i + \sum_{j \neq i} \mathbf{d}_j \right] = \tilde{\mathbf{g}}_i. \tag{35}$$

By uniting all followers through (35), we can obtain the following linear equation system

$$\begin{aligned}
\mathbf{C}\bar{\mathbf{d}}_1 + \mathbf{C}(\bar{\mathbf{d}}_1 + \cdots + \bar{\mathbf{d}}_n) &= \tilde{\mathbf{g}}_1 \\
\vdots & \\
\mathbf{C}\bar{\mathbf{d}}_n + \mathbf{C}(\bar{\mathbf{d}}_1 + \cdots + \bar{\mathbf{d}}_n) &= \tilde{\mathbf{g}}_n
\end{aligned} \tag{36}$$

Sum up all rows of (36) yields

$$\mathbf{C}(\bar{\mathbf{d}}_1 + \cdots + \bar{\mathbf{d}}_n) = \frac{1}{n+1} (\tilde{\mathbf{g}}_1 + \cdots + \tilde{\mathbf{g}}_n).$$

Substituting this into (36) we have

$$\bar{\mathbf{d}}_i = \mathbf{C}^{-1} \left[\tilde{\mathbf{g}}_i - \frac{1}{n+1} (\tilde{\mathbf{g}}_1 + \cdots + \tilde{\mathbf{g}}_n) \right], \quad \forall i \in \mathcal{N}. \tag{37}$$

According to the Sherman-Morrison formula¹, we have

$$\begin{aligned}
\mathbf{C}^{-1} &= \text{diag}(\tilde{w}(1), \dots, \tilde{w}(T-1)) - \frac{1}{1 + \frac{\sum_{t=1}^{T-1} \tilde{w}(t)}{\tilde{w}(T)}} \\
&\quad \times \frac{1}{\tilde{w}(T)} (\tilde{w}(1), \dots, \tilde{w}(T-1))^\top (\tilde{w}(1), \dots, \tilde{w}(T-1)) \\
&= \text{diag}(\tilde{w}(1), \dots, \tilde{w}(T-1)) - \frac{1}{\sum_{t=1}^T \tilde{w}(t)} \\
&\quad \times (\tilde{w}(1), \dots, \tilde{w}(T-1))^\top (\tilde{w}(1), \dots, \tilde{w}(T-1)).
\end{aligned} \tag{38}$$

On the other hand, we have

$$\begin{aligned}
\tilde{\mathbf{g}}_i - \frac{1}{n+1} \sum_{i=1}^n \tilde{\mathbf{g}}_i &= \frac{g_{\mathcal{N}} + g_i + \tilde{r}(T)}{\tilde{w}(T)} \mathbf{1} - \begin{pmatrix} \frac{\tilde{r}(1)}{\tilde{w}(1)} \\ \vdots \\ \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \end{pmatrix} \\
&\quad - \frac{(n+1)g_{\mathcal{N}} + n\tilde{r}(T)}{(n+1)\tilde{w}(T)} \mathbf{1} + \frac{n}{n+1} \begin{pmatrix} \frac{\tilde{r}(1)}{\tilde{w}(1)} \\ \vdots \\ \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \end{pmatrix} \\
&= \frac{(n+1)g_i + \tilde{r}(T)}{(n+1)\tilde{w}(T)} \mathbf{1} - \frac{1}{n+1} \begin{pmatrix} \frac{\tilde{r}(1)}{\tilde{w}(1)} \\ \vdots \\ \frac{\tilde{r}(T-1)}{\tilde{w}(T-1)} \end{pmatrix}, \quad \forall i \in \mathcal{N}.
\end{aligned} \tag{39}$$

Substituting (38) and (39) into (37), we can get

$$\begin{aligned}
\bar{\mathbf{d}}_i &= \frac{(n+1)g_i + \tilde{r}(T)}{(n+1)\tilde{w}(T)} \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} - \frac{1}{n+1} \\
&\quad \times \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T-1) \end{pmatrix} - \frac{\sum_{t=1}^{T-1} \tilde{w}(t)}{\sum_{t=1}^T \tilde{w}(t)} \frac{(n+1)g_i + \tilde{r}(T)}{(n+1)\tilde{w}(T)} \\
&\quad \times \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} + \frac{\sum_{t=1}^{T-1} \tilde{r}(t)}{\sum_{t=1}^T \tilde{w}(t)} \frac{1}{n+1} \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} \\
&= \left[\frac{\tilde{w}(T)}{\sum_{t=1}^T \tilde{w}(t)} \frac{(n+1)g_i + \tilde{r}(T)}{(n+1)\tilde{w}(T)} + \frac{\sum_{t=1}^{T-1} \tilde{r}(t)}{(n+1)\sum_{t=1}^T \tilde{w}(t)} \right] \\
&\quad \times \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} - \frac{1}{n+1} \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T-1) \end{pmatrix} \\
&= \left[\frac{g_i}{\sum_{t=1}^T \tilde{w}(t)} + \frac{\sum_{t=1}^T \tilde{r}(t)}{(n+1)\sum_{t=1}^T \tilde{w}(t)} \right] \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} \\
&\quad - \frac{\sum_{t=1}^T \tilde{w}(t)}{(n+1)\sum_{t=1}^T \tilde{w}(t)} \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T-1) \end{pmatrix}
\end{aligned} \tag{40}$$

¹Sherman-Morrison formula [50]: Given $\mathbf{A} \in \mathbb{R}^{n \times n}$, $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$, such that $\det(\mathbf{A}) \neq 0$, $1 + \mathbf{v}^\top \mathbf{A}^{-1} \mathbf{u} \neq 0$, then $\mathbf{A} + \mathbf{u}\mathbf{v}^\top$ is invertible, and we have

$$(\mathbf{A} + \mathbf{u}\mathbf{v}^\top)^{-1} = \mathbf{A}^{-1} - \frac{\mathbf{A}^{-1} \mathbf{u}\mathbf{v}^\top \mathbf{A}^{-1}}{1 + \mathbf{v}^\top \mathbf{A}^{-1} \mathbf{u}}.$$

$$\begin{aligned}
&= \frac{g_i}{\sum_{t=1}^T \tilde{w}(t)} \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} + \frac{1}{(n+1) \sum_{t=1}^T \tilde{w}(t)} \\
&\times \begin{pmatrix} \sum_{t=1}^T [\tilde{r}(t) \tilde{w}(1) - \tilde{w}(t) \tilde{r}(1)] \\ \vdots \\ \sum_{t=1}^T [\tilde{r}(t) \tilde{w}(T-1) - \tilde{w}(t) \tilde{r}(T-1)] \end{pmatrix} \\
&= (\nu_i(1), \nu_i(2), \dots, \nu_i(T-1))^\top, \quad \forall i \in \mathcal{N}.
\end{aligned}$$

Besides, we have

$$\begin{aligned}
&\sum_{t=1}^T \sum_{s=1}^T \tilde{w}(s) \tilde{w}(t) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(t)}{\tilde{w}(t)} \right] \\
&= \sum_{t=1}^T \tilde{w}(t) \sum_{s=1}^T \tilde{r}(s) - \sum_{t=1}^T \tilde{r}(t) \sum_{s=1}^T \tilde{w}(s) = 0,
\end{aligned}$$

so

$$\begin{aligned}
&\sum_{t=1}^{T-1} \sum_{s=1}^T \tilde{w}(s) \tilde{w}(t) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(t)}{\tilde{w}(t)} \right] \\
&= - \sum_{s=1}^T \tilde{w}(s) \tilde{w}(T) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(T)}{\tilde{w}(T)} \right]. \quad (41)
\end{aligned}$$

According to (40) and the constraint condition $\sum_{t=1}^T \nu_i(t) = g_i$, we can obtain

$$\begin{aligned}
\nu_i(T) &= g_i - \sum_{t=1}^{T-1} \nu_i(t) \\
&= g_i - \frac{\sum_{t=1}^{T-1} \tilde{w}(t)}{\sum_s \tilde{w}(s)} g_i - \frac{1}{(n+1) \sum_s \tilde{w}(s)} \\
&\quad \times \sum_{t=1}^{T-1} \sum_{s=1}^T \tilde{w}(s) \tilde{w}(t) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(t)}{\tilde{w}(t)} \right] \\
&= \frac{\tilde{w}(T)}{\sum_s \tilde{w}(s)} g_i + \frac{1}{(n+1) \sum_s \tilde{w}(s)} \\
&\quad \times \sum_{s=1}^T \tilde{w}(s) \tilde{w}(T) \left[\frac{\tilde{r}(s)}{\tilde{w}(s)} - \frac{\tilde{r}(T)}{\tilde{w}(T)} \right], \quad \forall i \in \mathcal{N},
\end{aligned}$$

where the last equality uses (41).

C. Proof of Lemma 4.4

Before the proof of Lemma 4.4, we need to introduce some lemmas as follows.

Lemma A.1: For each user $i \in \mathcal{N}$ and any strategy combination $\boldsymbol{\nu}_{-i} \in \mathbb{H}_{-i}$, the best response strategy $\bar{\boldsymbol{\nu}}_i = (\bar{\nu}_i(1), \dots, \bar{\nu}_i(T))^\top := \arg \min_{\boldsymbol{\nu}_i \in \mathbb{H}_i} u_i^f(\boldsymbol{\pi}, \boldsymbol{\nu}_i, \boldsymbol{\nu}_{-i})$ is

$$\begin{aligned}
\bar{\boldsymbol{\nu}}_i &= -\frac{1}{2} \sum_{j \neq i} \boldsymbol{\nu}_j + \frac{g_N + g_i + \sum_s \tilde{r}(s)}{2 \sum_s \tilde{w}(s)} \\
&\quad \times \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T) \end{pmatrix} - \frac{1}{2} \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T) \end{pmatrix}. \quad (42)
\end{aligned}$$

Proof. For each user i , by (22) the unique global minimum point $\bar{\boldsymbol{\nu}}_i$ satisfies

$$\bar{\boldsymbol{d}}_i = -\frac{1}{2} \mathbf{C}^{-1} \boldsymbol{\mu}_i. \quad (43)$$

Substituting (34) and (38) into (43), we have

$$\begin{aligned}
\bar{\boldsymbol{d}}_i &= -\frac{1}{2} \sum_{j \neq i} \boldsymbol{d}_j + \frac{1}{2} \mathbf{C}^{-1} \tilde{\boldsymbol{g}}_i \\
&= -\frac{1}{2} \sum_{j \neq i} \boldsymbol{d}_j + \frac{g_N + g_i + \tilde{r}(T)}{2 \tilde{w}(T)} \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} \\
&\quad - \frac{1}{2} \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T-1) \end{pmatrix} - \frac{g_N + g_i + \tilde{r}(T)}{2 \tilde{w}(T)} \frac{\sum_{s=1}^{T-1} \tilde{w}(s)}{\sum_s \tilde{w}(s)} \\
&\quad \times \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} + \frac{\sum_{s=1}^{T-1} \tilde{r}(s)}{2 \sum_s \tilde{w}(s)} \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} \\
&= -\frac{1}{2} \sum_{j \neq i} \boldsymbol{d}_j + \frac{g_N + g_i + \sum_s \tilde{r}(s)}{2 \sum_s \tilde{w}(s)} \\
&\quad \times \begin{pmatrix} \tilde{w}(1) \\ \vdots \\ \tilde{w}(T-1) \end{pmatrix} - \frac{1}{2} \begin{pmatrix} \tilde{r}(1) \\ \vdots \\ \tilde{r}(T-1) \end{pmatrix}.
\end{aligned}$$

Besides, due to $\mathbf{1}^\top \bar{\boldsymbol{\nu}}_i = g_i$, we have

$$\begin{aligned}
\bar{\nu}_i(T) &= g_i - \mathbf{1}^\top \bar{\boldsymbol{d}}_i \\
&= g_i + \frac{1}{2} \sum_{j \neq i} \sum_{t=1}^{T-1} \nu_j(t) - \frac{(g_N + g_i) \sum_{s=1}^{T-1} \tilde{w}(s)}{2 \sum_s \tilde{w}(s)} \\
&\quad - \left[\frac{\sum_s \tilde{r}(s) \sum_{s=1}^{T-1} \tilde{w}(s)}{2 \sum_s \tilde{w}(s)} - \frac{\sum_s \tilde{w}(s) \sum_{s=1}^{T-1} \tilde{r}(s)}{2 \sum_s \tilde{w}(s)} \right] \\
&= g_i + \frac{1}{2} \sum_{j \neq i} [g_j - \nu_j(T)] - \frac{(g_N + g_i) \sum_{s=1}^{T-1} \tilde{w}(s)}{2 \sum_s \tilde{w}(s)} \\
&\quad + \frac{\sum_s \tilde{r}(s) \tilde{w}(T) - \sum_s \tilde{w}(s) \tilde{r}(T)}{2 \sum_s \tilde{w}(s)} \\
&= -\frac{1}{2} \sum_{j \neq i} \nu_j(T) + g_i + \frac{1}{2} (g_N - g_i) \\
&\quad - \frac{(g_N + g_i) \sum_{s=1}^{T-1} \tilde{w}(s)}{2 \sum_s \tilde{w}(s)} + \frac{\sum_s \tilde{r}(s) \tilde{w}(T)}{2 \sum_s \tilde{w}(s)} - \frac{1}{2} \tilde{r}(T) \\
&= -\frac{1}{2} \sum_{j \neq i} \nu_j(T) + \frac{g_N + g_i + \sum_s \tilde{r}(s)}{2 \sum_s \tilde{w}(s)} \tilde{w}(T) - \frac{1}{2} \tilde{r}(T),
\end{aligned}$$

where the third equality uses (41). So the best response strategy is as (42). \square

For any matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$, let $\|\mathbf{A}\|_1 := \max_j \sum_{i=1}^n |\mathbf{A}_{ij}|$ denote the maximum absolute column sum norm.

Lemma A.2: For any $\mathbf{A} \in \mathbb{C}^{n \times n}$ and any constant $\varepsilon > 0$, we can find an invertible matrix $\mathbf{S}_{\mathbf{A}, \varepsilon} \in \mathbb{C}^{n \times n}$ such that

$$\|\mathbf{S}_{\mathbf{A}, \varepsilon}^{-1} \mathbf{A} \mathbf{S}_{\mathbf{A}, \varepsilon}\|_1 \leq \rho(\mathbf{A}) + \varepsilon,$$

where $\rho(\mathbf{A})$ is the spectral radius of \mathbf{A} .

Proof. For any matrix \mathbf{A} , there exists an invertible matrix \mathbf{P} such that $\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \mathbf{J}$, and \mathbf{J} is the Jordan canonical form of \mathbf{A} . Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the n eigenvalues of \mathbf{A} , so \mathbf{J} can be written as

$$\mathbf{J} = \boldsymbol{\Lambda} + \tilde{\mathbf{I}},$$

where $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ and

$$\tilde{\mathbf{I}} = \begin{pmatrix} 0 & \delta_1 & & & \\ & 0 & \delta_2 & & \\ & & \ddots & \ddots & \\ & & & 0 & \delta_{n-1} \\ & & & & 0 \end{pmatrix} (\delta_i = 0 \text{ or } 1).$$

Let $\mathbf{H} := \text{diag}(1, \varepsilon, \dots, \varepsilon^{n-1})$, and we have

$$(\mathbf{P}\mathbf{H})^{-1}\mathbf{A}(\mathbf{P}\mathbf{H}) = \mathbf{H}^{-1}\mathbf{J}\mathbf{H} = \mathbf{H}^{-1}\Lambda\mathbf{H} + \mathbf{H}^{-1}\tilde{\mathbf{I}}\mathbf{H} = \Lambda + \varepsilon\tilde{\mathbf{I}}.$$

Finally, we make $\mathbf{S}_{\mathbf{A},\varepsilon} = \mathbf{P}\mathbf{H}$, then $\mathbf{S}_{\mathbf{A},\varepsilon}$ is invertible, and

$$\|\mathbf{S}_{\mathbf{A},\varepsilon}^{-1}\mathbf{A}\mathbf{S}_{\mathbf{A},\varepsilon}\|_1 = \|\Lambda + \varepsilon\tilde{\mathbf{I}}\|_1 \leq \max_i |\lambda_i| + \varepsilon = \rho(\mathbf{A}) + \varepsilon.$$

□

Proof of Lemma 4.4: First, we provide an explicit expression for the mapping \mathbf{F} . For any $i \in \mathcal{N}$, denote

$$\alpha_i := \frac{g_{\mathcal{N}} + g_i + \sum_s \tilde{r}(s)}{2 \sum_s \tilde{w}(s)}$$

and

$$\mathbf{e}_i := \underbrace{(0, \dots, 0)}_i, 1, \dots, 1)^\top.$$

According to (42) and (23), we have

$$\begin{aligned} \mathbf{f}_1(\boldsymbol{\nu}) &= -\frac{1}{2}\boldsymbol{\nu}^\top \mathbf{e}_1 + \alpha_1(\tilde{w}(1), \dots, \tilde{w}(T))^\top \\ &\quad - \frac{1}{2}(\tilde{r}(1), \dots, \tilde{r}(T))^\top \\ \mathbf{f}_2(\boldsymbol{\nu}) &= -\frac{1}{2}(\mathbf{f}_1(\boldsymbol{\nu}) + \boldsymbol{\nu}^\top \mathbf{e}_2) + \alpha_2(\tilde{w}(1), \dots, \tilde{w}(T))^\top \\ &\quad - \frac{1}{2}(\tilde{r}(1), \dots, \tilde{r}(T))^\top \\ &= -\frac{1}{2}\boldsymbol{\nu}^\top \left(\mathbf{e}_2 - \frac{1}{2}\mathbf{e}_1 \right) + \left(\alpha_2 - \frac{\alpha_1}{2} \right) \\ &\quad \times (\tilde{w}(1), \dots, \tilde{w}(T))^\top - \frac{1}{2^2}(\tilde{r}(1), \dots, \tilde{r}(T))^\top, \\ &\quad \vdots \\ \mathbf{f}_n(\boldsymbol{\nu}) &= -\frac{1}{2}\boldsymbol{\nu}^\top \left(\mathbf{e}_n - \sum_{j=1}^{n-1} \frac{1}{2^{n-j}} \mathbf{e}_j \right) + \left(\alpha_n - \sum_{j=1}^{n-1} \frac{\alpha_j}{2^{n-j}} \right) \\ &\quad \times (\tilde{w}(1), \dots, \tilde{w}(T))^\top - \frac{1}{2^n}(\tilde{r}(1), \dots, \tilde{r}(T))^\top. \end{aligned} \quad (44)$$

Finally, by (44) we can obtain

$$\mathbf{F}(\boldsymbol{\nu}) = (\mathbf{f}_1(\boldsymbol{\nu}), \mathbf{f}_2(\boldsymbol{\nu}), \dots, \mathbf{f}_n(\boldsymbol{\nu}))^\top = \mathbf{L}_n \boldsymbol{\nu} + \mathbf{M}_n, \quad (45)$$

where

$$\begin{aligned} \mathbf{L}_n &:= -\frac{1}{2} \begin{pmatrix} \mathbf{e}_1^\top \\ \mathbf{e}_2^\top - \frac{1}{2}\mathbf{e}_1^\top \\ \vdots \\ \mathbf{e}_n^\top - \sum_{j=1}^{n-1} \frac{1}{2^{n-j}} \mathbf{e}_j^\top \end{pmatrix} \\ &= \begin{pmatrix} 0 & -\frac{1}{2} & -\frac{1}{2} & \cdots & -\frac{1}{2} & -\frac{1}{2} \\ 0 & \frac{1}{2^2} & -\frac{1}{2^2} & \cdots & -\frac{1}{2^2} & -\frac{1}{2^2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \frac{1}{2^{n-1}} & \frac{2^2-1}{2^{n-1}} & \cdots & \frac{2^{n-2}-1}{2^{n-1}} & -\frac{1}{2^{n-1}} \\ 0 & \frac{1}{2^n} & \frac{2^2-1}{2^n} & \cdots & \frac{2^{n-2}-1}{2^n} & \frac{2^{n-1}-1}{2^n} \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} \mathbf{M}_n &:= \left(\alpha_1, \dots, \alpha_n - \sum_{j=1}^{n-1} \frac{\alpha_j}{2^{n-j}} \right)^\top (\tilde{w}(1), \dots, \tilde{w}(T)) \\ &\quad - \left(\frac{1}{2}, \dots, \frac{1}{2^n} \right)^\top (\tilde{r}(1), \dots, \tilde{r}(T)). \end{aligned}$$

For above matrix $\mathbf{L}_n \in \mathbb{R}^{n \times n}$, it can be calculated that $\rho(\mathbf{L}_n) < 1$, so we can find a constant $\varepsilon_0 > 0$, such that $\rho(\mathbf{L}_n) + \varepsilon_0 < 1$. According to Lemma A.2, we can find the invertible matrix $\mathbf{S} := \mathbf{S}_{\mathbf{L}_n, \varepsilon_0}$ such that

$$\|\mathbf{S}^{-1}\mathbf{L}_n\mathbf{S}\|_1 \leq \rho(\mathbf{L}_n) + \varepsilon_0. \quad (46)$$

For any matrix $\mathbf{B} \in \mathbb{R}^{n \times T}$, we set

$$\|\mathbf{B}\|_{\mathbf{S}} := \|\mathbf{S}^{-1}\mathbf{B}\|_1. \quad (47)$$

We can verify that i) $\|\mathbf{B}_1 - \mathbf{B}_2\|_{\mathbf{S}} = 0$ if and only if $\mathbf{B}_1 = \mathbf{B}_2$; ii) $\|\mathbf{B}_1 - \mathbf{B}_2\|_{\mathbf{S}} = \|\mathbf{B}_2 - \mathbf{B}_1\|_{\mathbf{S}}$; iii) $\|\mathbf{B}_1 - \mathbf{B}_3\|_{\mathbf{S}} \leq \|\mathbf{B}_1 - \mathbf{B}_2\|_{\mathbf{S}} + \|\mathbf{B}_2 - \mathbf{B}_3\|_{\mathbf{S}}$. Therefore, $\|\cdot\|_{\mathbf{S}}$ is a metric on $\mathbb{R}^{n \times T}$, and $(\mathbb{R}^{n \times T}, \|\cdot\|_{\mathbf{S}})$ is a complete metric space. Further, $\mathbb{H} \subset \mathbb{R}^{n \times T}$ is a closed set, therefore $(\mathbb{H}, \|\cdot\|_{\mathbf{S}})$ is also a complete metric space.

For any $\boldsymbol{\nu}_1, \boldsymbol{\nu}_2 \in \mathbb{H}$, based on (45)-(47), we have

$$\begin{aligned} \|\mathbf{F}(\boldsymbol{\nu}_1) - \mathbf{F}(\boldsymbol{\nu}_2)\|_{\mathbf{S}} &= \|\mathbf{L}_n \boldsymbol{\nu}_1 - \mathbf{L}_n \boldsymbol{\nu}_2\|_{\mathbf{S}} \\ &= \|\mathbf{S}^{-1}\mathbf{L}_n(\boldsymbol{\nu}_1 - \boldsymbol{\nu}_2)\|_1 \\ &= \|\mathbf{S}^{-1}\mathbf{L}_n\mathbf{S}\mathbf{S}^{-1}(\boldsymbol{\nu}_1 - \boldsymbol{\nu}_2)\|_1 \\ &\leq \|\mathbf{S}^{-1}\mathbf{L}_n\mathbf{S}\|_1 \|\mathbf{S}^{-1}(\boldsymbol{\nu}_1 - \boldsymbol{\nu}_2)\|_1 \\ &\leq (\rho(\mathbf{L}_n) + \varepsilon_0) \|\boldsymbol{\nu}_1 - \boldsymbol{\nu}_2\|_{\mathbf{S}}, \end{aligned}$$

where the first inequality uses the norm compatibility of $\|\cdot\|_1$. Since $\rho(\mathbf{L}_n) + \varepsilon_0 < 1$, the mapping \mathbf{F} is a contraction mapping on \mathbb{H} . □

REFERENCES

- [1] S. Gupta, "From gadgets to the smart grid," *Nature*, vol. 526, no. 7575, pp. S90–S91, 2015.
- [2] J. Beyea, "The smart electricity grid and scientific research," *Science*, vol. 328, no. 5981, pp. 979–980, 2010.
- [3] O. Smith, O. Cattell, E. Farcot, R. D. O’Dea, and K. I. Hopcraft, "The effect of renewable energy incorporation on power grid stability and resilience," *Science Advances*, vol. 8, no. 9, p. eabj6734, 2022.
- [4] K. Moslehi and R. Kumar, "A reliability perspective of the smart grid," *IEEE Transactions on Smart Grid*, vol. 1, no. 1, pp. 57–64, 2010.
- [5] P. Siano, "Demand response and smart grids—a survey," *Renewable and Sustainable Energy Reviews*, vol. 30, pp. 461–478, 2014.
- [6] L. Liu, G. He, M. Wu, G. Liu, H. Zhang, Y. Chen, J. Shen, and S. Li, "Climate change impacts on planned supply–demand match in global wind and solar energy systems," *Nature Energy*, vol. 8, no. 8, pp. 870–880, 2023.
- [7] Y. Li, M. Han, Z. Yang, and G. Li, "Coordinating flexible demand response and renewable uncertainties for scheduling of community integrated energy systems with an electric vehicle charging station: A bi-level approach," *IEEE Transactions on Sustainable Energy*, vol. 12, no. 4, pp. 2321–2331, 2021.
- [8] Z. Wang, B. Lu, B. Wang, Y. Qiu, H. Shi, B. Zhang, J. Li, H. Li, and W. Zhao, "Incentive based emergency demand response effectively reduces peak load during heatwave without harm to vulnerable groups," *Nature Communications*, vol. 14, no. 1, p. 6202, 2023.
- [9] B. Xu, J. Wang, M. Guo, J. Lu, G. Li, and L. Han, "A hybrid demand response mechanism based on real-time incentive and real-time pricing," *Energy*, vol. 231, p. 120940, 2021.
- [10] T. Jiang, C. Chung, P. Ju, and Y. Gong, "A multi-timescale allocation algorithm of energy and power for demand response in smart grids: A stackelberg game approach," *IEEE Transactions on Sustainable Energy*, vol. 13, no. 3, pp. 1580–1593, 2022.
- [11] E. S. Parizy, H. R. Bahrami, and S. Choi, "A low complexity and secure demand response technique for peak load reduction," *IEEE Transactions on Smart Grid*, vol. 10, no. 3, pp. 3259–3268, 2018.

- [12] W. Zhong, R. Yu, S. Xie, Y. Zhang, and D. K. Yau, "On stability and robustness of demand response in v2g mobile energy networks," *IEEE Transactions on Smart Grid*, vol. 9, no. 4, pp. 3203–3212, 2016.
- [13] S. Sterl, I. Vanderkelen, C. J. Chawanda, D. Russo, R. J. Brecha, A. Van Griensven, N. P. van Lipzig, and W. Thiery, "Smart renewable electricity portfolios in west africa," *Nature Sustainability*, vol. 3, no. 9, pp. 710–719, 2020.
- [14] C. Cecati, C. Citro, and P. Siano, "Combined operations of renewable energy systems and responsive demand in a smart grid," *IEEE Transactions on Sustainable Energy*, vol. 2, no. 4, pp. 468–476, 2011.
- [15] Y. Dai, X. Sun, Y. Qi, and M. Leng, "A real-time, personalized consumption-based pricing scheme for the consumptions of traditional and renewable energies," *Renewable Energy*, vol. 180, pp. 452–466, 2021.
- [16] L. P. Qian, Y. J. A. Zhang, J. Huang, and Y. Wu, "Demand response management via real-time electricity price control in smart grids," *IEEE Journal on Selected Areas in Communications*, vol. 31, no. 7, pp. 1268–1280, 2013.
- [17] H. Li, Z. Wan, and H. He, "Real-time residential demand response," *IEEE Transactions on Smart Grid*, vol. 11, no. 5, pp. 4144–4154, 2020.
- [18] N. Mahmoudi, T. K. Saha, and M. Eghbal, "Wind power offering strategy in day-ahead markets: Employing demand response in a two-stage plan," *IEEE Transactions on Power Systems*, vol. 30, no. 4, pp. 1888–1896, 2014.
- [19] T. Ding, M. Qu, N. Amjadi, F. Wang, R. Bo, and M. Shahidepour, "Tracking equilibrium point under real-time price-based residential demand response," *IEEE Transactions on Smart Grid*, vol. 12, no. 3, pp. 2736–2740, 2020.
- [20] P. Samadi, H. Mohsenian-Rad, V. W. Wong, and R. Schober, "Real-time pricing for demand response based on stochastic approximation," *IEEE Transactions on Smart Grid*, vol. 5, no. 2, pp. 789–798, 2014.
- [21] T. Namerikawa, N. Okubo, R. Sato, Y. Okawa, and M. Ono, "Real-time pricing mechanism for electricity market with built-in incentive for participation," *IEEE Transactions on Smart Grid*, vol. 6, no. 6, pp. 2714–2724, 2015.
- [22] I. Atzeni, L. G. Ordóñez, G. Scutari, D. P. Palomar, and J. R. Fonollosa, "Noncooperative day-ahead bidding strategies for demand-side expected cost minimization with real-time adjustments: A gnep approach," *IEEE Transactions on Signal Processing*, vol. 62, no. 9, pp. 2397–2412, 2014.
- [23] A. Baniasadi, D. Habibi, O. Bass, and M. A. Masoum, "Optimal real-time residential thermal energy management for peak-load shifting with experimental verification," *IEEE Transactions on Smart Grid*, vol. 10, no. 5, pp. 5587–5599, 2018.
- [24] S. Choi, "Practical coordination between day-ahead and real-time optimization for economic and stable operation of distribution systems," *IEEE Transactions on Power Systems*, vol. 33, no. 4, pp. 4475–4487, 2017.
- [25] A. Galeotti, S. Goyal, M. O. Jackson, F. Vega-Redondo, and L. Yariv, "Network games," *The Review of Economic Studies*, vol. 77, no. 1, pp. 218–244, 2010.
- [26] G. Chen and Y. Yu, "Convergence analysis and strategy control of evolutionary games with imitation rule on toroidal grid," *IEEE Transactions on Automatic Control*, vol. 68, no. 12, pp. 8185–8192, 2023.
- [27] J. Cheng, W. Mei, W. Su, and G. Chen, "Evolutionary games on networks: Phase transition, quasi-equilibrium, and mathematical principles," *Physica A: Statistical Mechanics and its Applications*, vol. 611, p. 128447, 2023.
- [28] J. R. Riehl and M. Cao, "Towards optimal control of evolutionary games on networks," *IEEE Transactions on Automatic Control*, vol. 62, no. 1, pp. 458–462, 2016.
- [29] D. Madeo and C. Mocenni, "Game interactions and dynamics on networked populations," *IEEE Transactions on Automatic Control*, vol. 60, no. 7, pp. 1801–1810, 2014.
- [30] S. Tan, Y. Wang, and J. Lü, "Analysis and control of networked game dynamics via a microscopic deterministic approach," *IEEE Transactions on Automatic Control*, vol. 61, no. 12, pp. 4118–4124, 2016.
- [31] J. Moon and T. Başar, "Linear quadratic risk-sensitive and robust mean field games," *IEEE Transactions on Automatic Control*, vol. 62, no. 3, pp. 1062–1077, 2016.
- [32] T. Morimoto, T. Kanazawa, and T. Ushio, "Subsidy-based control of heterogeneous multiagent systems modeled by replicator dynamics," *IEEE Transactions on Automatic Control*, vol. 61, no. 10, pp. 3158–3163, 2015.
- [33] P. Guo, Y. Wang, and H. Li, "Algebraic formulation and strategy optimization for a class of evolutionary networked games via semi-tensor product method," *Automatica*, vol. 49, no. 11, pp. 3384–3389, 2013.
- [34] D. Cheng, T. Liu, K. Zhang, and H. Qi, "On decomposed subspaces of finite games," *IEEE Transactions on Automatic Control*, vol. 61, no. 11, pp. 3651–3656, 2016.
- [35] B. Zhu, X. Xia, and Z. Wu, "Evolutionary game theoretic demand-side management and control for a class of networked smart grid," *Automatica*, vol. 70, pp. 94–100, 2016.
- [36] S. R. Etesami, W. Saad, N. B. Mandayam, and H. V. Poor, "Stochastic games for the smart grid energy management with prospect prosumers," *IEEE Transactions on Automatic Control*, vol. 63, no. 8, pp. 2327–2342, 2018.
- [37] A.-H. Mohsenian-Rad, V. W. Wong, J. Jatskevich, R. Schober, and A. Leon-Garcia, "Autonomous demand-side management based on game-theoretic energy consumption scheduling for the future smart grid," *IEEE Transactions on Smart Grid*, vol. 1, no. 3, pp. 320–331, 2010.
- [38] I. S. Bayram, G. Michailidis, and M. Devetsikiotis, "Unsplittable load balancing in a network of charging stations under qos guarantees," *IEEE Transactions on Smart Grid*, vol. 6, no. 3, pp. 1292–1302, 2014.
- [39] H. Li, Z. Ren, A. Trivedi, P. P. Verma, D. Srinivasan, and W. Li, "A noncooperative game-based approach for microgrid planning considering existing interconnected and clustered microgrids on an island," *IEEE Transactions on Sustainable Energy*, vol. 13, no. 4, pp. 2064–2078, 2022.
- [40] S. Maharjan, Q. Zhu, Y. Zhang, S. Gjessing, and T. Basar, "Dependable demand response management in the smart grid: A stackelberg game approach," *IEEE Transactions on Smart Grid*, vol. 4, no. 1, pp. 120–132, 2013.
- [41] M. Yu and S. H. Hong, "A real-time demand-response algorithm for smart grids: A stackelberg game approach," *IEEE Transactions on Smart Grid*, vol. 7, no. 2, pp. 879–888, 2015.
- [42] M. Yan, M. Shahidepour, A. Paaso, L. Zhang, A. Alabdulwahab, and A. Abusorrah, "Distribution network-constrained optimization of peer-to-peer transactive energy trading among multi-microgrids," *IEEE Transactions on Smart Grid*, vol. 12, no. 2, pp. 1033–1047, 2020.
- [43] J. Li, G. Ma, T. Li, W. Chen, and Y. Gu, "A stackelberg game approach for demand response management of multi-microgrids with overlapping sales areas," *Science China Information Sciences*, vol. 62, pp. 1–13, 2019.
- [44] T. Başar and G. J. Olsder, *Dynamic noncooperative game theory*. SIAM, 1998.
- [45] J. Borwein and A. Lewis, *Convex Analysis*. Springer, 2006.
- [46] L. V. Kantorovich and G. P. Akilov, *Functional analysis*. Elsevier, 2016.
- [47] J. B. Rosen, "Existence and uniqueness of equilibrium points for concave n-person games," *Econometrica: Journal of the Econometric Society*, pp. 520–534, 1965.
- [48] A. J. Laub, *Matrix analysis for scientists and engineers*. SIAM, 2004.
- [49] N. Gould, D. Orban, and P. Toint, "Numerical methods for large-scale nonlinear optimization," *Acta Numerica*, vol. 14, pp. 299–361, 2005.
- [50] W. W. Hager, "Updating the inverse of a matrix," *SIAM Review*, vol. 31, no. 2, pp. 221–239, 1989.