

# F-PURE THRESHOLD FOR THE SYMMETRIC DETERMINANTAL RING

JUSTIN FONG

**ABSTRACT.** We give a value for the  $F$ -pure threshold at the maximal homogeneous ideal  $\mathfrak{m}$  of the symmetric determinantal ring over a field of prime characteristic. The answer is characteristic independent, so we immediately get the log canonical threshold in characteristic zero as well.

## 1. INTRODUCTION

The  $F$ -pure threshold, introduced by Takagi and Watanabe in [TW04], is a numerical invariant of a ring in positive characteristic associated to the  $F$ -singularity class of  $F$ -purity. Suppose  $R$  is a reduced  $F$ -finite  $F$ -pure ring of characteristic  $p > 0$ , and let  $\mathfrak{a} \subseteq R$  be a nonzero ideal. For a real number  $t \in \mathbb{R}_{\geq 0}$ , the pair  $(R, \mathfrak{a}^t)$  is called  $F$ -pure if for every  $q = p^e \gg 0$ , there exists an element  $f \in \mathfrak{a}^{\lfloor (q-1)t \rfloor}$  (where  $\lfloor (q-1)t \rfloor$  is the greatest integer less than or equal to the real number  $(q-1)t$ ) such that the inclusion  $f^{1/q}R \hookrightarrow R^{1/q}$  splits as an  $R$ -module homomorphism. The  $F$ -pure threshold of  $\mathfrak{a}$  is defined to be

$$\text{fpt}(\mathfrak{a}) := \sup\{t \in \mathbb{R}_{\geq 0} \mid (R, \mathfrak{a}^t) \text{ is } F\text{-pure}\}.$$

When  $\mathfrak{a} = \mathfrak{m}_R$  is the maximal homogeneous ideal of  $R$ , we will simply refer to  $\text{fpt}(\mathfrak{m}_R)$  as the  $F$ -pure threshold of  $R$ .

The  $F$ -pure threshold has been calculated for the determinantal rings of generic matrices [STV17, Proposition 4.3], Hankel matrices [CMSV18, Theorem 4.5], and is trivial for skew-symmetric matrices (see section 2.1 for an explanation). However, to the author's knowledge,  $\text{fpt}(\mathfrak{m}_R)$  has been unknown for the symmetric matrix case. We provide an answer in Theorem 2.5, as well as a proof that the symmetric determinantal ring is  $F$ -pure, a fact not clearly stated in the literature.

**Note.** At the time of writing this article, the author was notified of the recent paper [CSV24], which includes a proof of the more general property of  $F$ -regularity for symmetric determinantal rings, and mentions the same conclusion of Proposition 2.4 in this article. Our work is independent of that paper.

**Acknowledgements.** I would like to thank Irena Swanson, Vaibhav Pandey, and my advisor Uli Walther for providing helpful discussions.

## 2. SYMMETRIC DETERMINANTIAL RINGS

Suppose that  $X = (x_{ij})$  is a symmetric  $n \times n$  generic matrix (i.e.,  $x_{ij} = x_{ji}$ ), let  $\mathbb{k}[X] = \mathbb{k}[x_{ij} \mid 1 \leq i \leq j \leq n]$  be the polynomial ring over a field  $\mathbb{k}$  with the entries of  $X$  as its variables, and  $I_t(X)$  be the ideal generated by the  $t$ -minors of  $X$ . For an integer

$1 \leq t \leq n$  let  $R = \mathbb{k}[X]/I_t(X)$  be the *symmetric determinantal ring*. A formula for the  $a$ -invariant of  $R$  has been given by Barile in [Bar94]. When  $R$  is given the standard grading this is given as follows:

$$a(R) = \begin{cases} -(t-1)\frac{n}{2}, & \text{if } n \equiv t \pmod{2} \\ -(t-1)\frac{n+1}{2}, & \text{if } n \not\equiv t \pmod{2}. \end{cases} \quad (2.1)$$

In [Con94] it is shown that the graded canonical module of  $R$  is

$$\omega_R = \begin{cases} R \left( -(t-1)\frac{n}{2} \right), & \text{if } n \equiv t \pmod{2} \\ \mathfrak{p} \left( -(t-1)\frac{n-1}{2} \right), & \text{if } n \not\equiv t \pmod{2}, \end{cases} \quad (2.2)$$

where  $\mathfrak{p}$  is the prime ideal generated by the  $t-1$  minors of the first  $t-1$  rows of  $X$ . In [Got79] and [Got77] Goto shows that  $R$  is Gorenstein if and only if  $n \equiv t \pmod{2}$ , and the divisor class group of  $R$  is  $\mathbb{Z}/2\mathbb{Z}$  generated by  $[\mathfrak{p}]$ .

**Lemma 2.1.** [DSMnNb23, (4.2)] *Suppose  $S$  is a polynomial ring over a field of characteristic  $p > 0$ , and let  $I \subset S$  be a prime ideal of  $h = \text{height}(I)$ . Then  $I^{(h(p-1))} \subseteq (I^{[p]} :_S I)$ , i.e.,  $I^{(h(p-1))+1} \subseteq I^{[p]}$ .*

*Proof.* To show the containment  $I^{(h(p-1))+1} \subseteq I^{[p]}$ , it is enough to show it locally at all the associated primes of  $S/I^{[p]}$ . However, since the Frobenius map is flat, the set of associated primes of  $S/I^{[p]}$  is the same as for  $S/I$ , which consists only of  $I$  since it is prime. In the regular local ring  $(S_I, IS_I)$  the symbolic power of  $IS_I$  coincides with its ordinary power since  $IS_I$  is a maximal ideal. Hence it follows that

$$(IS_I)^{(h(p-1))} I = (IS_I)^{h(p-1)} I = (IS_I)^{h(p-1)+1}.$$

Since  $(S_I, IS_I)$  has dimension  $h = \text{height}(I)$ , its unique maximal ideal  $IS_I$  is generated by  $h$  elements  $g_1, \dots, g_h$ . The power  $(IS_I)^{h(p-1)+1}$  is generated by  $g_1^{a_1} \cdots g_h^{a_h}$  such that  $a_1 + \cdots + a_h = h(p-1) + 1$ . By the pigeonhole principle, at least one  $a_i \geq p$ , hence it follows that  $(IS_I)^{h(p-1)+1} \subseteq I^{[p]} S_I$ .  $\square$

**Proposition 2.2.** *Over a field of characteristic  $p > 0$  the symmetric determinantal ring  $R$  is  $F$ -pure.*

*Proof.* Let  $I_t = I_t(X)$ , and  $S = \mathbb{k}[X]$  be the polynomial ring in  $n(n+1)/2$  many variables. We will apply Fedder's criterion:  $R$  is  $F$ -pure if and only if  $\mathfrak{m}_S^{[p]} \not\subseteq (I_t^{[p]} :_S I_t)$ . Thus, in order to show that  $R$  is  $F$ -pure, we need to find an element  $f \in (I_t^{[p]} :_S I_t) \setminus \mathfrak{m}_S^{[p]}$ .

Let  $\mathfrak{D} = d_1 \cdots d_n$  be a product of  $i$ -minors  $d_i$  of the  $i \times i$  submatrices in the bottom-left corner of  $X$  (for  $i = 1, \dots, n$ ), whose diagonals are the  $n$  distinct diagonals of  $X$  (see Example 2.3 below). Our proposed element will be  $f = \mathfrak{D}^{p-1}$ . The leading term of  $\mathfrak{D}$  with respect to the diagonal term ordering

$$x_{11} > x_{12} > \cdots > x_{1n} > x_{21} > \cdots > x_{n-1n} > x_{nn}$$

is  $\text{in}(\mathfrak{D}) = \prod_{1 \leq i \leq j \leq n} x_{ij}$ , which is the product of all the distinct variables of  $X$ , hence it is a squarefree monomial. This implies  $\mathfrak{D}^{p-1} \notin \mathfrak{m}_S^{[p]}$ .

Next, we need to show  $\mathfrak{D}^{p-1}I_t \subseteq I_t^{[p]}$ , i.e.,  $\mathfrak{D}^{p-1} \in (I_t^{[p]} :_S I_t)$ . It suffices to prove this containment after localizing at the prime ideal  $I_t$ . If one can show that  $\mathfrak{D}$  belongs to the symbolic power  $I_t^{(h)}$ , where  $h = \text{height}(I_t)$ , then it would follow

$$\mathfrak{D}^{p-1}I_t S_{I_t} \subseteq ((I_t S_{I_t})^{(h)})^{p-1} I_t S_{I_t} = (I_t S_{I_t})^{h(p-1)+1} \subseteq I_t^{[p]} S_{I_t},$$

where  $(I_t S_{I_t})^{(h)} = (I_t S_{I_t})^h$ , because  $I_t S_{I_t}$  is the (unique) maximal ideal of  $S_{I_t}$ , and the last containment follows from Lemma 2.1. We will now show  $\mathfrak{D} \in I_t^{(h)}$ . In [Kut74, Theorem 1], Kutz has shown that  $\dim R = n(t-1) - \frac{1}{2}(t-1)(t-2)$ , hence it follows

$$h = \dim S - \dim R = \frac{n(n+1)}{2} - n(t-1) + \frac{(t-1)(t-2)}{2}.$$

In general, suppose  $\Delta$  is a product  $\delta_1 \cdots \delta_u$  of  $a_i$ -minors  $\delta_i$  of  $X$ , where the vector  $(a_1, \dots, a_u) \in \mathbb{N}^u$  is called the *shape* of  $\Delta$ . With the previous notation, the symbolic power  $I_t^{(k)}$  is described by

$$\left[ \Delta \in I_t^{(k)} \right] \iff \left[ a_i \leq n \ \forall i \text{ and } \gamma_t(\Delta) := \sum_{i=1}^u \max\{0, a_i - t + 1\} \geq k \right]$$

(see [JMnV15, Proposition 4.3].) We must show  $\gamma_t(\mathfrak{D}) \geq h$ . Observe that the product  $\mathfrak{D} = d_1 d_2 \cdots d_n$  has shape  $(1, 2, \dots, n)$ , and note that  $i - t + 1 > 0 \iff i > t - 1$ , hence  $\max\{0, i - t + 1\} = i - t + 1$  for  $t \leq i \leq n$ . Thus, one has

$$\begin{aligned} \gamma_t(\mathfrak{D}) &= \sum_{i=t}^n (i - t + 1) \\ &= \sum_{i=t}^n i + \sum_{i=t}^n (1 - t) \\ &= \frac{n(n+1)}{2} - \frac{t(t-1)}{2} + (n-t+1)(1-t) \\ &= \frac{n(n+1)}{2} - n(t-1) + \frac{(t-1)(t-2)}{2}, \end{aligned}$$

i.e.,  $\gamma_t(\mathfrak{D}) = \text{height}(I_t) = h$ , so it follows that  $\mathfrak{D} \in I_t^{(h)}$ . □

**Example 2.3.** For the  $3 \times 3$  symmetric matrix

$$X = \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{12} & x_{22} & x_{23} \\ x_{13} & x_{23} & x_{33} \end{pmatrix}$$

the  $i$ -minors  $d_i$  ( $i = 1, 2, 3$ ) of the  $i \times i$  botom-left corner of  $X$  are

$$d_1 = x_{13}, \quad d_2 = \begin{vmatrix} x_{12} & x_{22} \\ x_{13} & x_{23} \end{vmatrix}, \quad d_3 = |X|,$$

which have leading terms  $\text{in}(d_1) = x_{13}$ ,  $\text{in}(d_2) = x_{12}x_{23}$ , and  $\text{in}(d_3) = x_{11}x_{22}x_{33}$ , hence  $\mathfrak{D} = d_1d_2d_3$  has squarefree leading term  $x_{11}x_{12}x_{13}x_{22}x_{23}x_{33}$ .

**Proposition 2.4.** *Over a field  $\mathbb{k}$  of arbitrary characteristic, let  $\mathfrak{p}$  be the prime ideal of  $R = \mathbb{k}[X]/I_t(X)$  generated by the  $t - 1$  minors of the first  $t - 1$  rows of  $X$ . Then the second symbolic power  $\mathfrak{p}^{(2)}$  is principally generated by the  $(t - 1)$ -minor in the top left-hand corner of  $X$ .*

*Proof.* Since the class group  $\mathbb{Z}/2\mathbb{Z}$  of  $R$  is generated by the class of  $\mathfrak{p}$ , it follows  $\mathfrak{p}^{(2)}$  is isomorphic to a principal ideal of  $R$ . Consider the  $n \times n$  symmetric matrix

$$X = \begin{pmatrix} A & \cdots & B \\ \vdots & & \vdots \\ B^T & \cdots & C \end{pmatrix}$$

where  $A, B, C$ , and  $B^T$  (transpose of  $B$ ) are  $(t - 1) \times (t - 1)$  submatrices of  $X$  (it is possible that these four submatrices have overlapping entries). Their determinants will be denoted  $|A|, |B|$ , and  $|C|$ , which are  $(t - 1)$ -minors of  $X$ . We have  $\mathfrak{p} = I_{t-1} \left( \begin{matrix} A & \cdots & B \end{matrix} \right)$  is an ideal generated by the top  $(t - 1)$ -minors. It is enough to show  $|A| \in \mathfrak{p}^{(2)}$ , since the principal ideal generated by  $|A|$  is  $\mathfrak{p}$ -primary [Got77, Proposition (1)] and contains  $\mathfrak{p}^2$ , and  $\mathfrak{p}^{(2)}$  is the smallest  $\mathfrak{p}$ -primary ideal containing  $\mathfrak{p}^2$ . By definition,

$$\mathfrak{p}^{(2)} := \{r \in R \mid sr \in \mathfrak{p}^2 \text{ for some } s \notin \mathfrak{p}\},$$

so it suffices to exhibit an  $s \notin \mathfrak{p}$  such that  $s|A| \in \mathfrak{p}^2$ . We claim  $s = |C|$  is such an element. Since  $C$  is the bottom-right corner of  $X$ , it is not completely in the first  $(t - 1)$ -rows  $\left( \begin{matrix} A & \cdots & B \end{matrix} \right)$  of  $X$ , so  $|C| \notin \mathfrak{p}$ . On the other hand,  $|B|^2$  is one of the generators of  $\mathfrak{p}^2$ . We will show

$$|A||C| - |B|^2 \in I_t(X), \tag{2.3}$$

for then it will follow in the quotient  $R = \mathbb{k}[X]/I_t(X)$  that  $|A||C| = |B|^2 \in \mathfrak{p}^2$ , and hence  $\mathfrak{p}^{(2)}$  is principally generated by  $|A|$ .

By the nullstellensatz, after taking the algebraic closure  $\overline{\mathbb{k}}$ , (2.3) will follow if it holds for all matrices with entries in  $\overline{\mathbb{k}}$ . Assume  $A, B, C, X$  have  $\overline{\mathbb{k}}$ -entries, and  $X$  belongs to the variety  $\text{Var}(I_t)$  of  $t$ -minors, so  $\text{rank}(X) \leq t - 1$ . Omit all rows and columns of  $X$  that do not involve  $A, B, B^T, C$ . The resulting matrix, denoted by  $X'$ , is now a  $2t - 2$  square. If the top  $t - 1$  rows of  $X'$  have rank less than  $t - 1$ , then  $\text{rank}(A), \text{rank}(B) < t - 1$  and hence  $|A| = |B| = 0$ , so there is nothing to show. The same is true for the bottom  $t - 1$  rows. Thus, the top half  $(A, B)$  and bottom half  $(B^T, C)$  of  $X$  can be assumed to have rank  $t - 1$ . However, since  $\text{rank}(X) < t$ , it must be that  $\text{rank}(X) = t - 1$ , so the rows of the top  $(A, B)$  and bottom  $(B^T, C)$  are bases for the same row space  $W$ . This same argument applies to the left half  $\begin{pmatrix} A \\ B^T \end{pmatrix}$  and right half  $\begin{pmatrix} B \\ C \end{pmatrix}$  of  $X$  by symmetry, so the column space is also  $W$ . If  $A$  is full rank, then so is  $B$ , otherwise the column

space of  $A$  would not be contained in the column space of  $B$ . Similarly,  $C$  is full rank as well. Now, since  $\text{rank}(X) = \text{rank}(A) = t - 1$ , there exist invertible matrices  $D$  and  $E$  over  $\bar{\mathbb{k}}$  such that  $D(A, B) = (B^T, C)$ , and  $\begin{pmatrix} A \\ B^T \end{pmatrix} E = \begin{pmatrix} B \\ C \end{pmatrix}$ , which gives

$$\begin{aligned} DA &= B^T, & AE &= B, \\ DB &= C, & B^T E &= C. \end{aligned}$$

Using these, one has  $AC = A(DB) = ADAE$  and  $B^T B = (DA)(AE) = DAAE$ , which shows that  $|A||C| - |B|^2 = 0$  in  $\bar{\mathbb{k}}$ .

In the case where  $A, B, B^T, C$  do overlap, consider  $Y = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix}$  of size  $2t - 2$ .

Both  $Y$  and  $X'$ , from the non-overlap case above, share the same variables in the non-overlap regions of  $Y$ . As for the variables in the overlap region of  $Y$ , these are expressed in terms of the corresponding variables of  $X'$ , hence the overlap variables of  $Y$  are identified with the variables of the non-overlap case. Thus, the identity proven in the non-overlap case remains the same in the quotient  $R$ , hence (2.3) holds in general.  $\square$

**Theorem 2.5.** *The  $F$ -pure threshold of the symmetric determinantal ring  $R$  is*

$$\text{fpt}(\mathfrak{m}_R) = \frac{n(t-1)}{2}.$$

*In characteristic zero, this is equal to the log canonical threshold as well.*

*Proof.* The ring  $R = \mathbb{k}[X]/I_t$  is Gorenstein if and only if  $n \equiv t \pmod{2}$ , hence it follows from [DSNnB18, Theorem 5.2] and equation (2.1) that  $\text{fpt}(\mathfrak{m}_R) = -a(R) = (t-1)\frac{n}{2}$ , when  $n \equiv t \pmod{2}$ . Now, suppose  $R$  is non-Gorenstein, i.e.,  $n \not\equiv t \pmod{2}$ . Its  $a$ -invariant and graded canonical module are respectively  $a(R) = -(t-1)\frac{n+1}{2}$  and  $\omega_R = \mathfrak{p} \left( -(t-1)\frac{n-1}{2} \right)$  (see (2.2)). Since the class group  $\text{Cl}(R)$  is cyclic of order 2, the ring  $R$  is  $\mathbb{Q}$ -Gorenstein, so we will apply [STV17, Proposition 4.5]. We determine the generating degree  $D$  of  $\omega_R^{(2)}$ . First, note that  $\mathfrak{p}^{(2)}$  is principally generated by a  $(t-1)$ -minor, by Theorem 2.4, hence as graded modules  $\mathfrak{p}^{(2)} \cong R(-(t-1))$ . The second symbolic power of  $\omega_R$  is then

$$\begin{aligned} \omega_R^{(2)} &= \left( \mathfrak{p} \left( \frac{-(t-1)(n-1)}{2} \right) \right)^{(2)} = \mathfrak{p}^{(2)} \left( 2 \cdot \frac{-(t-1)(n-1)}{2} \right) \\ &= R(-(t-1))(-(t-1)(n-1)) \\ &= R(-(t-1) - (t-1)(n-1)) \\ &= R(-(t-1)n), \end{aligned}$$

which is generated in degree  $D = (t-1)n$ . It follows that  $\text{fpt}(\mathfrak{m}_R) = D/c = n(t-1)/2$ .

In characteristic zero, one has by [MTW05, Theorem 3.4] that the log canonical threshold of  $R$  is  $\text{lct}(\mathfrak{m}_R) = \lim_{p \rightarrow \infty} \text{fpt}(\mathfrak{m}_{R_p}) = n(t-1)/2$ , where  $R_p$  is the prime reduction modulo  $p$  of  $R$ .  $\square$

**2.1. Ring of Pfaffians.** Let  $X = (x_{ij})$  be an  $n \times n$  skew-symmetric matrix (i.e.,  $x_{ii} = 0$ ,  $x_{ij} = -x_{ji}$  for  $i < j$ ),  $\mathbb{k}[X]$  be the polynomial ring in the entries of  $X$ , and  $\text{Pf}_{2t}(X)$  be the ideal generated by the  $2t$ -Pfaffians of  $X$ . The ring of Pfaffians  $R = \mathbb{k}[X]/\text{Pf}_{2t}(X)$  is Gorenstein [Avr79] and strongly  $F$ -regular [Bae01]. Its  $a$ -invariant has been computed in [BH92, Corollary 1.7], and assuming it is standard graded, is given by  $a(R) = -n(t - 1)$ , if  $2t \leq n$ . By [DSNnB18, Theorem 5.2] we immediately get the following.

**Proposition 2.6.** *For  $2t \leq n$ , the  $F$ -pure threshold of the ring of Pfaffians  $R = \mathbb{k}[X]/\text{Pf}_{2t}(X)$  is  $\text{fpt}(\mathfrak{m}_R) = n(t - 1)$ .*

## REFERENCES

- [Avr79] Luchezar L. Avramov. A class of factorial domains. *Serdica*, 5(4):378–379, 1979.
- [Bae01] C. Baetica.  $F$ -rationality of algebras defined by Pfaffians. volume 3(53), pages 139–144. 2001. Memorial issue dedicated to Nicolae Radu.
- [Bar94] Margherita Barile. The Cohen-Macaulayness and the  $a$ -invariant of an algebra with straightening laws on a doset. *Comm. Algebra*, 22(2):413–430, 1994.
- [BH92] Winfried Bruns and Jürgen Herzog. On the computation of  $a$ -invariants. *Manuscripta Math.*, 77(2-3):201–213, 1992.
- [CMSV18] Aldo Conca, Maral Mostafazadehfard, Anurag K. Singh, and Matteo Varbaro. Hankel determinantal rings have rational singularities. *Adv. Math.*, 335:111–129, 2018.
- [Con94] Aldo Conca. Symmetric ladders. *Nagoya Math. J.*, 136:35–56, 1994.
- [CSV24] Aldo Conca, Anurag K. Singh, and Matteo Varbaro. Invariant rings of the special orthogonal group have nonunimodal  $h$ -vectors. arXiv:2406.14439 [math.AC], 2024.
- [DSMnNnB23] Alessandro De Stefani, Jonathan Montaña, and Luis Núñez Betancourt. Frobenius methods in combinatorics. *São Paulo J. Math. Sci.*, 17(1):387–429, 2023.
- [DSNnB18] Alessandro De Stefani and Luis Núñez Betancourt.  $F$ -thresholds of graded rings. *Nagoya Math. J.*, 229:141–168, 2018.
- [Got77] Shiro Goto. The divisor class group of a certain Krull domain. *J. Math. Kyoto Univ.*, 17(1):47–50, 1977.
- [Got79] Shiro Goto. On the Gorensteinness of determinantal loci. *J. Math. Kyoto Univ.*, 19(2):371–374, 1979.
- [JMnV15] Jack Jeffries, Jonathan Montaña, and Matteo Varbaro. Multiplicities of classical varieties. *Proc. Lond. Math. Soc. (3)*, 110(4):1033–1055, 2015.
- [Kut74] Ronald E. Kutz. Cohen-Macaulay rings and ideal theory in rings of invariants of algebraic groups. *Trans. Amer. Math. Soc.*, 194:115–129, 1974.
- [MTW05] Mircea Mustață, Shunsuke Takagi, and Kei-ichi Watanabe.  $F$ -thresholds and Bernstein-Sato polynomials. In *European Congress of Mathematics*, pages 341–364. Eur. Math. Soc., Zürich, 2005.
- [STV17] Anurag K. Singh, Shunsuke Takagi, and Matteo Varbaro. A Gorenstein criterion for strongly  $F$ -regular and log terminal singularities. *Int. Math. Res. Not. IMRN*, (21):6484–6522, 2017.
- [TW04] Shunsuke Takagi and Kei-ichi Watanabe. On  $F$ -pure thresholds. *J. Algebra*, 282(1):278–297, 2004.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA  
*Email address:* fong11@purdue.edu