ON ADICALLY COMPLETE D-MODULES IN CHARACTERISTIC ZERO

AMNON YEKUTIELI

ABSTRACT. Let (X, \mathcal{O}_X) be an algebraic manifold in characteristic 0, or an analytic manifold over \mathbb{C} . A standard theorem says that a left \mathcal{D}_X -module \mathcal{M} , which is coherent as an \mathcal{O}_X -module, is locally free. This theorem has a generalization to the adically complete algebraic setting, in a paper by Ogus from 1973.

In the present paper we take a new look at the work of Ogus. We provide a detailed proof of the theorem on \mathcal{D} -modules, and extend it to the non-noetherian setting. We also give another proof of an interesting result of Ogus about adically complete modules (slightly extended).

In the Appendix we discuss a related error in a book by Björk.

Contents

0.	Introduction	1
1.	Adically Complete Modules	3
2.	Rings of Differential Operators	6
3.	Complete \mathcal{D} -Modules – the One Dimensional Case	9
4.	Complete \mathcal{D} -Modules – the High Dimensional Case	13
Ap	ppendix A. An Error in a Book by Björk	15
References		16

0. INTRODUCTION

Let (X, \mathcal{O}_X) be an algebraic manifold over a field \mathbb{K} of characteristic 0, or an analytic manifold over \mathbb{C} . A standard theorem says that a left \mathcal{D}_X -module \mathcal{M} , which is coherent as an \mathcal{O}_X -module, is a locally free \mathcal{O}_X -module. See [Be], [Bj] or [HTT].

This geometric theorem has a vast generalization to the adically complete algebraic setting, namely Theorem 1.3 in the paper [Og] by A. Ogus from 1973. In Remark 4.7 we explain how the algebraic theorem of Ogus implies the geometric theorem.

In our present paper we give a detailed proof of the theorem of Ogus, and extend it to the non-noetherian setting. This is our Theorem 0.6. We also provide another proof of an interesting result of Ogus about adically complete modules (see [Og, Proposition A1]), and extend it slightly; this is our Theorem 0.1 below. Thus our paper is not original research, but rather an attempt to throw new light on these remarkable results.

Before stating our first theorem, we need a bit of background. Consider a commutative ring A and a finitely generated ideal $\mathfrak{a} \subseteq A$. Let M be an A-module. We do not assume that A is noetherian, nor that M is finitely generated. The module M is equipped with its \mathfrak{a} -adic filtration $\{\mathfrak{a}^i \cdot M\}_{i \in \mathbb{N}}$. The \mathfrak{a} -adic completion of M is the A-module

Date: 13 July 2024.

 $\Lambda_{\mathfrak{a}}(M) := \lim_{i \to i} M/\mathfrak{a}^i \cdot M$, and M is called \mathfrak{a} -adically complete (and separated) if the canonical homomorphism $\tau_{M,\mathfrak{a}}: M \to \Lambda_{\mathfrak{a}}(M)$ is an isomorphism. If $\tau_{M,\mathfrak{a}}$ is only injective, then M is called \mathfrak{a} -adically separated.

Suppose the *A*-module *M* has some other descending filtration $\{F^i(M)\}_{i \in I}$ by *A*-submodules, indexed by a directed set (I, \leq) . Namely, $\{F^i(M)\}_{i \in I}$ is an inverse system of submodules of *M*. The *F*-adic completion of *M* is the *A*-module $\Lambda_F(M) := \lim_{\leftarrow i} M/F^i(M)$, and *M* is called *F*-adically complete (and separated) if the canonical homomorphism $\tau_{M,F} :$ $M \to \Lambda_F(M)$ is an isomorphism.

Both the \mathfrak{a} -adic filtration and the *F*-adic filtration determine *A*-linear topologies on the *A*-module *M*. Let's denote the corresponding topological spaces by (M, \mathfrak{a}) and (M, F). We say that the *F*-adic topology on *M* is weaker than the \mathfrak{a} -adic topology on it if the identity of *M* is a continuous function $(M, \mathfrak{a}) \to (M, F)$. Concretely, this means that for every $i \in I$ there exists $j \in \mathbb{N}$ such that $\mathfrak{a}^j \cdot M \subseteq F^i(M)$.

The first theorem is a slight improvement of [Og, Proposition A1].

Theorem 0.1. Let A be a commutative ring and $\mathfrak{a} \subseteq A$ a finitely generated ideal. Let M be an A-module, with a filtration $\{F^i(M)\}_{i \in I}$ indexed by a directed set (I, \leq) . Assume that M is F-adically complete, and the F-adic topology on M is weaker than the \mathfrak{a} -adic topology. Then M is \mathfrak{a} -adically complete.

This is repeated as Theorem 1.5 in Section 1 and proved there. Note that in [Og] it is assumed that the indexing set *I* is countable; but our proof does not require this. Here are a few useful corollaries.

Corollary 0.2. With $\mathfrak{a} \subseteq A$ as in the theorem, suppose $\mathfrak{b} \subseteq A$ is an ideal such that $\mathfrak{a} \subseteq \mathfrak{b}$. If M is \mathfrak{b} -adically complete, then it is \mathfrak{a} -adically complete.

The next is [Og, Corollary A3].

Corollary 0.3. With $a \subseteq A$ as in the theorem, assume M is a-adically complete, and $N \subseteq M$ is an A-submodule that is closed for the a-adic topology. Then N is a-adically complete.

The next is a slight improvement of [Og, Corollary A2]. Again we do not require the set *I* to be countable.

Corollary 0.4. With $\mathfrak{a} \subseteq A$ as in the theorem, let $\{M_i\}_{i \in I}$ be an inverse system of A-modules, indexed by a directed set (I, \leq) . Assume every M_i is an \mathfrak{a} -adically complete module. Define the A-module $M := \lim_{t \to i} M_i$. Then M is \mathfrak{a} -adically complete.

The corollaries are repeated and proved in Section 1 of the paper.

We now move to \mathcal{D} -modules. Let A be a commutative ring of characteristic 0, i.e. A contains the field of rational numbers \mathbb{Q} . We do not assume A is noetherian. Let $B := A[[t_1, \ldots, t_n]]$, the ring of power series over A in n variables, and let $\mathfrak{b} \subseteq B$ be the ideal generated be the variables. We know that B is \mathfrak{b} -adically complete.

Consider the ring $\mathcal{D}_{B/A}$ of *A*-linear differential operators on *B* (in the sense of Grothendieck, see [EGA IV, Section 16]). For every *j* let $\partial_j := \partial/\partial t_j$, the partial derivative along t_j . In Proposition 2.8 we show that $\mathcal{D}_{B/A}$ is a Weyl algebra, namely as a left *B*module it is free with basis the collection of monomial operators $\{\partial_1^{k_1} \cdots \partial_n^{k_n}\}$, indexed by $(k_1, \ldots, k_n) \in \mathbb{N}^n$, and the usual relations hold.

Let *M* be a *B*-module. A left $\mathcal{D}_{B/A}$ -module structure on *M* is the same as an *inte*grable continuous connection $\nabla : M \to \widehat{\Omega}^1_{B/A} \otimes_B M$. Here $\widehat{\Omega}^1_{B/A}$ is the complete module of differential 1-forms, which is a free *B*-module with basis (dt_1, \ldots, dt_n) . The passage between connections and \mathcal{D} -modules is this: for an element $m \in M$ there is equality $\nabla(m) = \sum_j dt_j \otimes \partial_j(m)$ in $\widehat{\Omega}^1_{B/A} \otimes_B M$. Unfortunately, we could not find a reference in the literature for this interplay between \mathcal{D} -modules and connections. Anyhow, this is not required for our work, and it is mentioned only for the sake of comparison with the work of Ogus.

Given a left $\mathcal{D}_{B/A}$ -module *M*, its *horizontal submodule* is the *A*-module

$$(0.5) Mhor := \{ m \in M \mid \partial_j(m) = 0 \text{ for all } j \}.$$

In terms of the corresponding connection ∇ we have $M^{\text{hor}} = \text{Ker}(\nabla)$.

The following theorem is a generalization of [Og, Theorem 1.3], stated in terms of \mathcal{D} -modules. Ogus works with integrable connections, and assumes that the ring A is noetherian. We do not put any finiteness condition on A.

Theorem 0.6. Let A be a commutative ring containing \mathbb{Q} , let $B := A[[t_1, ..., t_n]]$, and let $\mathfrak{b} \subseteq B$ be the ideal generated by the variables. Suppose M is a left $\mathcal{D}_{B/A}$ -module, which is \mathfrak{b} -adically complete as a B-module. Then the canonical $\mathcal{D}_{B/A}$ -module homomorphism

$$B \widehat{\otimes}_A M^{\mathrm{hor}} \to M$$

is bijective.

In the theorem, $B \otimes_A M^{\text{hor}}$ is the b-adic completion of the *B*-module $B \otimes_A M^{\text{hor}}$. It has an induced left $\mathcal{D}_{B/A}$ -module structure. Ogus calls the corresponding connection a *completely constant connection*.

The 1-dimensional case of Theorem 0.6 is Theorem 3.13. For $n \ge 2$ it is Theorem 4.1, which is proved using Theorem 3.13 and induction on *n*. The induction requires Corollaries 0.2 and 0.3.

Corollary 0.7. Under the assumptions of the theorem, the canonical A-module homomorphism $M^{\text{hor}} \rightarrow M/\mathfrak{b} \cdot M$ is bijective.

A *B*-module *M* is called *b*-adically free if *M* is the *b*-adic completion of a free *B*-module.

Corollary 0.8. Under the assumptions of the theorem, if $M/\mathfrak{b} \cdot M$ is a free A-module, then *M* is a \mathfrak{b} -adically free *B* module.

The following corollary is a special case of the previous one. It is [Ka, Proposition 8.9] (which is phrased in terms of connections).

Corollary 0.9. Assume A is a field and M is a finitely generated B-module. Then $M \cong B^{\oplus r}$ as $\mathcal{D}_{B/A}$ -modules, for some natural number r.

These corollaries are proved in Section 4.

Finally, in Appendix A, we discuss an error we discovered in the book [Bj], in the proof of the analytic variant of Theorem 0.6.

Acknowledgments. I wish to thank Reinhold Hübl, Rolf Källström and Pierre Schapira for their useful advice in preparing this article.

1. Adically Complete Modules

In this section we study adic completions of modules. Theorem 1.5 is a slight improvement of [Og, Proposition A1].

Let A be a nonzero commutative ring, and let M be an A-module. We do not assume that A is noetherian, nor that M is finitely generated..

AMNON YEKUTIELI

Suppose $\{F^i(M)\}_{i \in I}$ is an inverse system of A-submodules of M, indexed by a directed set (I, \leq) . We call such a system a *filtration of* M. The collection of quotients $\{M/F^i(M)\}_{i \in I}$ is also an inverse system, and its inverse limit

(1.1)
$$\Lambda_F(M) := \lim_{t \to \infty} M/F^{\iota}(M)$$

is called the *F*-adic completion of *M*. There is a canonical homomorphism $\tau_{M,F} : M \to \Lambda_F(M)$, and we say that *M* is *F*-adically complete if $\tau_{M,F}$ is an isomorphism.

The next proposition is probably known, but we are not aware of a published reference.


Proposition 1.2. Let $\{M_i\}_{i \in I}$ be an inverse system of A-modules, indexed by a directed set (I, \leq) , and let $M := \lim_{i \to i} M_i$. Define the filtration $\{F^i(M)\}_{i \in I}$ of M by $F^i(M) := \text{Ker}(M \to M_i)$. Then the A-module M is F-adically complete.

We do not assume that the indexing set *I* is countable.

Proof. For every *i* let $N_i := M/F^i(M)$. There are injections $\psi_i : N_i \rightarrow M_i$, and in the limit we obtain an injection

$$\psi := \lim_{\leftarrow i} \psi_i := \Lambda_F(M) = \lim_{\leftarrow i} N_i \rightarrowtail \lim_{\leftarrow i} M_i = M.$$

It fits in the commutative diagram of A-modules



It follows that ψ is surjective, and thus it is bijective. Therefore $\tau_{M,F}$ is bijective.

Corollary 1.3. Let M be an A-module, with a filtration $\{F^i(M)\}_{i \in I}$ indexed by a directed set (I, \leq) . Define $\widehat{M} := \Lambda_F(M)$, and put on \widehat{M} the filtration $\{F^i(\widehat{M})\}_{i \in I}$, where $F^i(\widehat{M}) := \operatorname{Ker}(\widehat{M} \to M/F^i(M))$. Then \widehat{M} is F-adically complete.

Proof. For every *i* let $M_i := M/F^i(M)$. Then $\widehat{M} = \lim_{i \in I} M_i$, and its filtration *F* is the same as in Proposition 1.2. The proposition says that \widehat{M} is *F*-adically complete.

An important source of filtrations comes from ideals $\mathfrak{a} \subseteq A$. Namely, given an ideal \mathfrak{a} , we define $F^i(M) := \mathfrak{a}^i \cdot M$. This is the \mathfrak{a} -adic filtration of M, indexed by $I = \mathbb{N}$, and the completion $\Lambda_\mathfrak{a}(M) := \Lambda_F(M)$ is called the \mathfrak{a} -adic completion of M. Observe that $\Lambda_\mathfrak{a} : \operatorname{\mathsf{Mod}}(A) \to \operatorname{\mathsf{Mod}}(A)$ is an A-linear functor, and $\tau_\mathfrak{a} : \operatorname{Id} \to \Lambda_\mathfrak{a}$ is a morphism of functors.

We know (see [Ye3, Example 1.8]) there there are A-modules M whose \mathfrak{a} -adic completion \widehat{M} is *not* \mathfrak{a} -adically complete. How can this be reconciled with Corollary 1.3 ? The answer is that there are two, possibly distinct, filtrations on \widehat{M} , namely $\operatorname{Ker}(\widehat{M} \to M / \mathfrak{a}^i \cdot M)$ and $\mathfrak{a}^i \cdot \widehat{M}$. When the ideal \mathfrak{a} is finitely generated these two filtrations agree. Indeed:

Proposition 1.4. Let \mathfrak{a} be a finitely generated ideal in the ring A, and let M be an arbitrary A-module. Consider the \mathfrak{a} -adic filtration $\{\mathfrak{a}^i \cdot M\}_{i \in \mathbb{N}}$ on M, and the completion $\widehat{M} := \Lambda_{\mathfrak{a}}(M)$. Then:

- (1) The A-module \widehat{M} is a-adically complete.
- (2) For every *i* there is equality $a^i \cdot M = \text{Ker}(\widehat{M} \to M / a^i \cdot M)$.


Proof. According to [Ye3, Corollary 3.6], the *A*-module \widehat{M} is \mathfrak{a} -adically complete. By [Ye3, Theorem 1.2] the equality in item (2) holds.

Item (1) of the proposition says that the endofunctor $\Lambda_{\mathfrak{a}}$ on Mod(A) is idempotent.

The next theorem is an improvement of [Og, Proposition A1], where the indexing set *I* was assumed to be countable. We give a new proof, relying on Proposition 1.4.

Theorem 1.5 (Ogus). Let a be a finitely generated ideal in the ring A, and let M be an arbitrary A-module, with a filtration $\{F^i(M)\}_{i \in I}$ indexed by some directed set (I, \leq) . Assume that M is F-adically complete, and that the F-adic topology on M is weaker than the a-adic topology on it. Then M is a-adically complete.

Proof. Since the *F*-adic topology is weaker than the \mathfrak{a} -adic topology, for any $i \in I$ there is some $j \in \mathbb{N}$ such that $\mathfrak{a}^j \cdot M \subseteq F^i(M)$. Composing the canonical surjections $\Lambda_\mathfrak{a}(M) \to M/\mathfrak{a}^j \cdot M$ and $M/\mathfrak{a}^j \cdot M \to M/F^i(M)$ we get a homomorphism $\Lambda_\mathfrak{a}(M) \to M/F^i(M)$. These homomorphisms assemble into an inverse system indexed by *I*, and they induce a homomorphism $\psi : \Lambda_\mathfrak{a}(M) \to \Lambda_F(M)$. This homomorphism fits into the commutative diagram



We see that $\tau_{M,\mathfrak{a}}$ is a split injection, so there is an *A*-module isomorphism $\Lambda_{\mathfrak{a}}(M) \cong M \oplus K$, where $K := \text{Ker}(\psi)$.

Since $\Lambda_{\mathfrak{a}}(M)$ is a-adically complete (by Proposition 1.4), and since $\Lambda_{\mathfrak{a}}$ is an *A*-linear functor, the direct summand *M* of $\Lambda_{\mathfrak{a}}(M)$ is also a-adically complete. (So in fact ψ and $\tau_{M,\mathfrak{a}}$ are bijective, and K = 0.)

Corollary 1.6. With $\mathfrak{a} \subseteq A$ as in the theorem, suppose $\mathfrak{b} \subseteq A$ is an ideal such that $\mathfrak{a} \subseteq \mathfrak{b}$. If M is \mathfrak{b} -adically complete, then it is \mathfrak{a} -adically complete.

Proof. This is because the b-adic topology on *M* is weaker than the a-adic topology. \Box

The next is [Og, Corollary A3].

Corollary 1.7. With $a \subseteq A$ as in the theorem, assume M is a-adically complete, and $N \subseteq M$ is an A-submodule that is closed for the a-adic topology. Then N is a-adically complete.

Proof. Take any submodule *N* of *M*. Consider the filtration $\{F^i(N)\}_{i\in\mathbb{N}}$ of *N*, $F^i(N) := N \cap \mathfrak{a}^i \cdot M$. An element $m \in M$ is in the closure of *N*, for the \mathfrak{a} -adic topology, if and only if for every *i* the image of *m* in $M/\mathfrak{a}^i \cdot M$ belongs to the submodule $N/F^i(N)$. This means that the closure of *N* inside $M = \lim_{i \to i} M/\mathfrak{a}^i \cdot M$ is $\lim_{i \to i} N/F^i(N)$.

We are given that the submodule N is closed. Hence $N = \lim_{i \to i} N/F^i(N)$, i.e. it is *F*-adically complete. Since the *F*-adic topology on N is weaker than its *a*-adic topology, we conclude that N is *a*-adically complete.

The next is a slight improvement of [Og, Corollary A2]; again we do not require the set *I* to be countable.

Corollary 1.8. With $\mathfrak{a} \subseteq A$ as in the theorem, let $\{M_i\}_{i \in I}$ be an inverse system of A-modules, indexed by a directed set (I, \leq) . Assume every M_i is an \mathfrak{a} -adically complete module. Define the A-module $M := \lim_{t \in I} M_i$. Then M is \mathfrak{a} -adically complete.

Proof. Consider the set $K := I \times \mathbb{N}$, with this quasi-order: $k_1 = (i_1, j_1) \leq k_2 = (i_2, j_2)$ if $i_1 \leq i_2$ and $j_1 \leq j_2$. So (K, \leq) is a directed set. For $k = (i, j) \in K$ define the Amodule $M_k := M_i / \mathfrak{a}^j \cdot M_i$. We know that $M_i \cong \lim_{k \to j \in \mathbb{N}} M_{i,j}$ and $M = \lim_{k \to i \in I} M_i$. By

AMNON YEKUTIELI

the associativity property of inverse limits, we get $M \cong \lim_{k \in K} M_k$. Letting $F^k(M) := \text{Ker}(M \to M_k)$, Proposition 1.2 tells us that M is F-adically complete.

The *F*-adic topology is weaker that the \mathfrak{a} -adic topology on *M*, because $\mathfrak{a}^j \cdot M \subseteq F^{(i,j)}(M)$ for all $i \in i$ and $j \in \mathbb{N}$. According to Theorem 1.5, *M* is \mathfrak{a} -adically complete. \Box

2. Rings of Differential Operators

In this section we discuss differential operators between adically complete rings and modules.

We begin by recalling material from [EGA IV, Section 16]. Let $A \to B$ be a homomorphism between nonzero commutative rings. Given *B*-modules *M* and *N*, consider the *A*-central *B*-bimodule Hom_{*A*}(*M*, *N*), where the action by *B* is $(b_1 \cdot \theta \cdot b_2)(m) := b_1 \cdot \theta(b_2 \cdot m)$ for $b_i \in B$, $m \in M$ and $\theta \in \text{Hom}_A(M, N)$. We refer to the homomorphism θ as an *operator*. The operator θ is said to have *differential order* ≤ 0 if it is *B*-linear. For $k \geq 1$ the definition is inductive: an operator $\theta \in \text{Hom}_A(M, N)$ has differential order $\leq k$ if for every $b \in B$ the operator $[b, \theta] := b \cdot \theta - \theta \cdot b$ has differential order $\leq k - 1$. The set of all differential operators of order $\leq k$ is the *A*-submodule $\text{Diff}_{B/A}^k(M, N)$ of $\text{Hom}_A(M, N)$, and the set of all differential operators is

(2.1)
$$\operatorname{Diff}_{B/A}(M,N) := \bigcup_{k \ge 0} \operatorname{Diff}_{B/A}^k(M,N) \subseteq \operatorname{Hom}_A(M,N).$$

It is not hard to see that for *B*-modules M_0 , M_1 and M_2 , composition is a function

$$\operatorname{Diff}_{B/A}^{k_1}(M_0, M_1) \times \operatorname{Diff}_{B/A}^{k_2}(M_1, M_2) \to \operatorname{Diff}_{B/A}^{k_1+k_2}(M_0, M_2).$$

Thus for a single module M, $\text{Diff}_{B/A}(M, M)$ is a filtered B-ring, central over A.

For M = B we obtain the ring of differential operators $\mathcal{D}_{B/A} := \text{Diff}_{B/A}(B, B)$, which comes with the filtration $\{\mathcal{D}_{B/A}^k\}_{k \in \mathbb{N}}$. In this case $\mathcal{D}_{B/A}^0 = B$, but *B* is not in the center of $\mathcal{D}_{B/A}$.

Proposition 2.2. Let *M* be a left $\mathcal{D}_{B/A}$ -module. For any operator $\theta \in \mathcal{D}_{B/A}^k$, the operator $\theta_M \in \text{End}_A(M), \theta_M(m) := \theta \cdot m$, is a differential operator of order $\leq k$.

Proof. By induction on k. For k = 0 we have $\mathcal{D}_{B/A}^0 = B$, so θ_M has order ≤ 0 . Now take $k \geq 1$. For every $b \in B$ and $m \in M$ we have

$$[b, \theta_M](m) = b \cdot \theta \cdot m - \theta \cdot b \cdot m = [b, \theta]_M(m)$$

in *M*. Now $[b, \theta] \in \mathcal{D}_{B/A}^{k-1}$, so by the induction hypothesis $[b, \theta]_M \in \text{Diff}_{B/A}^{k-1}(M, M)$. We conclude that $[b, \theta_M]$ is a differential operator of order $\leq k - 1$ for all *b*. Therefore θ_M is a differential operator of order $\leq k$.

Let *I* be the kernel of the multiplication homomorphism $B \otimes_A B \to B$. The ring $\mathcal{P}_{B/A}^k := (B \otimes_A B)/I^{k+1}$ is called the *module of principal parts of order k*. See [EGA IV, Subsection 16.3]. We view $\mathcal{P}_{B/A}^k$ as a left *B*-module by the ring homomorphism $B \to \mathcal{P}_{B/A}^k$, $b \mapsto b \otimes 1$. Then the ring homomorphism $d^k : B \to \mathcal{P}_{B/A}^k$, $b \mapsto 1 \otimes b$, is a differential operator of order *k*. It is universal in the following sense: for *B* modules *M* and *N*, the *A*-module homomorphism

(2.3)
$$\operatorname{Hom}_{B}(\mathcal{P}^{k}_{B/A} \otimes_{B} M, N) \to \operatorname{Diff}^{k}_{B/A}(M, N), \ \phi \mapsto \phi \circ d^{k}_{M},$$

is bijective. Here $d_M^k(m) := d^k(1_B) \otimes m \in \mathcal{P}_{B/A}^k \otimes_B M$.

Lemma 2.4. Let $\mathfrak{b} \subseteq B$ be an ideal, and let $\theta \in \text{Diff}_{B/A}^k(M, N)$. Then $\theta(\mathfrak{b}^{i+k} \cdot M) \subseteq \mathfrak{b}^i \cdot N$ for all $i \in \mathbb{N}$.

Proof. This is [Ye1, Proposition 1.4.6].

Suppose *M* and *N* are *B*-modules, $\mathfrak{b} \subseteq B$ is a finitely generated ideal, and $\theta : M \to N$ is an A-linear homomorphism that's continuous for the b-adic topologies on these modules. Then θ extends uniquely to a continuous *A*-linear homomorphism $\hat{\theta} : \hat{M} \to \hat{N}$ between the b-adic completions.

Proposition 2.5. Let $\mathfrak{b} \subseteq B$ be a finitely generated ideal, and let \widehat{B} be the \mathfrak{b} -adic completion of B. Let M and N be B-modules, and let $\theta \in \text{Diff}_{B/A}^k(M, N)$.

- (1) The A-linear homomorphism $\theta: M \to N$ is continuous for the b-adic topologies.
- (2) The unique continuous extension $\widehat{\theta}: \widehat{M} \to \widehat{N}$ of θ belongs to $\text{Diff}_{\widehat{B}/A}^k(\widehat{M}, \widehat{N})$.

Proof. (1) Clear from Lemma 2.4.

(2) The operator $\hat{\theta}$ is a differential operator of order $\leq k$ iff $[\cdots [\hat{\theta}, b_0], \dots, b_k] = 0$ for all $b_0, \ldots, b_k \in \widehat{B}$. This commutator expression is a continuous multilinear function $\widehat{B}^{k+1} \to \widehat{B}$ \widehat{B} , and it vanishes on the image of B^{k+1} . Therefore it is zero on \widehat{B}^{k+1} .

Corollary 2.6. Let $\mathfrak{b} \subseteq B$ be a finitely generated ideal, and let \widehat{B} be the \mathfrak{b} -adic completion of B.

- (1) Any differential operator $\theta \in \mathcal{D}_{B/A}^k$ extends uniquely to a differential operator $\widehat{\theta} \in$ $\mathcal{D}_{\widehat{B}/A}^{k}.$ (2) The function $\theta \mapsto \widehat{\theta}$ is a filtered A-ring homomorphism $\mathcal{D}_{B/A} \to \mathcal{D}_{\widehat{B}/A}.$

Proof. (1) This is clear from Proposition 2.5.

(2) This follows from the uniqueness in (1).

Now assume $B = A[t_1, ..., t_n]$, the polynomial ring in *n* variables. *A* is still an arbitrary nonzero commutative ring. For a multi-index $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{N}^n$ we write $t^i := t_1^{i_1} \cdots t_n^{i_n}$ and $|\mathbf{i}| := i_1 + \cdots + i_n$. Then (see [Ye1, Section 1.4]) $\mathcal{P}_{B/A}^k$ is a free left *B*-module with basis the collection $\{d^k(t^i)\}_{|i| \le k}$. From this we get the dual basis $\{\theta_i\}_{|i| \le k}$ of $\mathcal{D}_{B/A}^k$. Namely, the differential operator θ_i is the unique differential operator such that $\theta_i(t^i) = 1$ and $\theta_i(t^j) = 0$ for $j \neq i$. For k = 1 we have the partial derivatives $\partial_i = \partial/\partial_{t_i} \in \mathcal{D}^1_{B/A}$, which satisfy $\partial_i(t_i) = 1$, and $\partial_i(t_j) = 0$ if $j \neq i$.

If A contains \mathbb{Q} , then, due to the uniqueness above, for every $\mathbf{i} = (i_1, \dots, i_n)$ there is equality

(2.7)
$$\theta_i = (i_1! \cdots i_n!)^{-1} \cdot \partial_1^{i_1} \cdots \partial_n^{i_n} \in \mathcal{D}_{B/A}.$$

It follows that as a left *B*-module, $\mathcal{D}_{B/A}$ is free with basis the collection of monomial operators $\{\partial_1^{i_1}\cdots\partial_n^{i_n}\}_{i\in\mathbb{N}^n}$. The relations satisfied by the multiplication in $\mathcal{D}_{B/A}$ are these: $\partial_i \cdot \partial_i = \partial_i \cdot \partial_i$, and $\partial_i \cdot b = b \cdot \partial_i + \partial_i(b)$ for $b \in B$.

Consider the ideal $\mathfrak{b} \subseteq B$ generated by the variables. The \mathfrak{b} -adic completion of B is the ring $B = A[[t_1, ..., t_n]]$ of power series. The next proposition describes the structure of the ring $\mathcal{D}_{\widehat{B}/A}$.

Proposition 2.8. Let A be a commutative ring containing \mathbb{Q} , let $B := A[t_1, \ldots, t_n]$, and let $\overline{B} := A[[t_1, \ldots, t_n]].$

AMNON YEKUTIELI

- (1) Any differential operator $\theta \in \mathcal{D}_{B/A}$ of order $\leq k$ extends uniquely to a differential operator $\hat{\theta} \in \mathcal{D}_{\widehat{B}/A}$ of order $\leq k$.
- (2) The function $\theta \mapsto \hat{\theta}$ is a filtered ring homomorphism $\mathcal{D}_{B/A} \to \mathcal{D}_{\hat{B}/A}$.
- (3) The left \widehat{B} -module homomorphism $\widehat{B} \otimes_B \mathcal{D}_{B/A} \to \mathcal{D}_{\widehat{B}/A}$ is bijective.
- (4) $\mathcal{D}_{\widehat{B}/A}$ is a free left \widehat{B} -module with basis $\{\partial_1^{i_1}\cdots\partial_n^{i_n}\}_{i\in\mathbb{N}^n}$.
- (5) The relations satisfied by the multiplication in $\mathcal{D}_{\widehat{B}/A}$ are these: $\partial_i \cdot \partial_j = \partial_j \cdot \partial_i$, and $\partial_i \cdot b = b \cdot \partial_i + \partial_i(b)$ for $b \in \widehat{B}$.

Proof. (1-2) These are special cases of items (1-2) of Corollary 2.6.

(3) By item (1) there is a canonical A-module isomorphism

(2.9)
$$\mathcal{D}_{\widehat{B}/A} = \mathrm{Diff}_{\widehat{B}/A}^k(\widehat{B},\widehat{B}) \cong \mathrm{Diff}_{B/A}^k(B,\widehat{B})$$

For every k, the left *B*-module $\mathcal{P}_{B/A}^k$ is free of finite rank. Therefore, using (2.9), we have isomorphisms of *A*-modules

$$\mathcal{D}_{\widehat{B}/A}^{k} \cong \operatorname{Hom}_{B}(\mathcal{P}_{B/A}^{k}, \widehat{B}) \cong \widehat{B} \otimes_{B} \operatorname{Hom}_{B}(\mathcal{P}_{B/A}^{k}, B) \cong \widehat{B} \otimes_{B} \mathcal{D}_{B/A}^{k}.$$

The composed isomorphism $\widehat{B} \otimes_B \mathcal{D}_{B/A}^k \xrightarrow{\simeq} \mathcal{D}_{\widehat{B}/A}^k$ is compatible with the ring homomorphism $\mathcal{D}_{B/A} \to \mathcal{D}_{\widehat{B}/A}$ from item (1).

(4) This is clear from item (3) and formula (2.7).

(5) These relations hold in $\mathcal{D}_{B/A}$, so by continuity they also hold in $\mathcal{D}_{\widehat{B}/A}$.

Corollary 2.10. In the setting of Proposition 2.8, let M be a left $\mathcal{D}_{B/A}$ -module, and let M be the \mathfrak{b} -adic completion of M. Then \widehat{M} has a unique left $\mathcal{D}_{\widehat{B}/A}$ -module structure, such that $\tau_M : M \to \widehat{M}$ is $\mathcal{D}_{B/A}$ -linear.

Proof. Of course \widehat{M} is a \widehat{B} -module. By Propositions 2.2 and 2.5, the partial derivatives ∂_i act continuously on M, and hence they extend uniquely to continuous A-linear endomorphisms on \widehat{M} . By continuity, the relations in Proposition 2.8(5) are satisfied in $\operatorname{End}_A^{\operatorname{cont}}(M)$. So we get a left $\mathcal{D}_{\widehat{R}/A}$ -module structure on M.

Remark 2.11. Let $A \to B$ be a ring homomorphism, $\mathfrak{b} \subseteq B$ a finitely generated ideal, and \widehat{B} the \mathfrak{b} -adic completion of B. We can give a description of the ring $\mathcal{D}_{\widehat{B}/A}$ as the *dir-inv completion* of the ring $\mathcal{D}_{B/A}$.


Dir-inv structures where introduced in [Ye2, Section 1]. In our setting we put on $\mathcal{D}_{B/A}$ the dir structure (an ascending filtration) $\{\mathcal{D}_{B/A}^k\}_{k\in\mathbb{N}}$, and on each $\mathcal{D}_{B/A}^k$ we put the inv structure (a descending filtration) $\{F^i(\mathcal{D}_{B/A}^k)\}_{i\in\mathbb{Z}}$, where

$$F^{i}(\mathcal{D}^{k}_{B/A}) \coloneqq \left\{ \theta \in \mathcal{D}^{k}_{B/A} \mid \theta(\mathfrak{b}^{j}) \subseteq \mathfrak{b}^{\max(j+i,0)} \text{ for all } j \in \mathbb{N} \right\}.$$

The dir-inv completion of $\mathcal{D}_{B/A}$ is

$$\Lambda_{\mathrm{di}}(\mathcal{D}_{B/A}) := \lim_{k \to i} \lim_{\leftarrow i} \mathcal{D}^{k}_{B/A} / F^{i}(\mathcal{D}^{k}_{B/A}).$$

This completion has a unique A-ring structure such that the diagram of A-rings



is commutative, and the horizontal arrow is an isomorphism.

This description of the ring $\mathcal{D}_{\widehat{B}/A}$ can be used to prove a generalization of Corollary 2.10 to the setting of this remark.

3. Complete D-Modules – the One Dimensional Case

This section is devoted to the proof of Theorem 3.13, which is the 1-dimensional case of Theorem 0.6 from the introduction. We give a rather detailed proof; but the essential idea already appears in the prior papers of Ogus and others, see Remark 3.15.

Throughout this section we assume the following:

Setup 3.1. We are given a commutative ring *A* containing \mathbb{Q} . Let B := A[[t]], the ring of power series in the variable *t*, and let $\mathfrak{b} \subseteq B$ be the ideal generated by *t*. We are also given a left $\mathcal{D}_{B/A}$ -module *M*, which is \mathfrak{b} -adically complete as a *B*-module.

Observe that the ring A is not assumed to be noetherian, and the B-module M is not assumed to be finitely generated.

By the definition of the b-adic topology on M, the submodules $b^i \cdot M$ are open in M, and thus – as always in linearly topologized modules – also closed.

As explained in Section 2, the ring of differential operators $\mathcal{D}_{B/A}$ is the subring of End_{*A*}(*B*) generated by *B* and the derivation $\partial := \frac{\partial}{\partial t}$. An operator $\phi \in \mathcal{D}_{B/A}$ of differential order $\leq i$ has a unique sum expansion

(3.2)
$$\phi = \sum_{0 \le j \le i} b_j \cdot \partial^j$$

in $\mathcal{D}_{B/A}$, with coefficients $b_j \in B$. According to Proposition 2.8(5), the relation satisfied by the element ∂ in the noncommutative ring $\mathcal{D}_{B/A}$ is

$$(3.3) \qquad \qquad \partial \cdot b = b \cdot \partial + \partial(b),$$

where $\partial(b) \in B$ is the result of action of the operator ∂ on an element $b \in B$. For every $m \in M$ and $b \in B$ we have the equality

(3.4)
$$\partial(b \cdot m) = (\partial \cdot b) \cdot m = (b \cdot \partial + \partial(b)) \cdot m = b \cdot \partial(m) + \partial(b) \cdot m.$$

This formula implies that $\partial(\mathfrak{b}^i \cdot M) \subseteq \mathfrak{b}^{i-1} \cdot M$, so ∂ acts continuously on M for its \mathfrak{b} -adic topology. By formula (3.2) we see that every operator $\phi \in \mathcal{D}_{B/A}$ acts continuously on M.

Definition 3.5.

(1) For every $i \in \mathbb{N}$ define the order *i* differential operator

$$\psi_i := (i!)^{-1} \cdot (-t)^i \cdot \partial^i \in \mathcal{D}_{B/A}.$$

(2) For every element $m \in M$ define the element

$$\psi(m) := \sum_{i \in \mathbb{N}} \psi_i(m) = \sum_{i \in \mathbb{N}} (i!)^{-1} \cdot (-t)^i \cdot \partial^i(m) \in M.$$

Since $\psi_i(m) \in \mathfrak{b}^i \cdot M$, the sum in item (2) converges in the \mathfrak{b} -adically complete module M. See Remark 3.16 regarding the possible convergence of the operator sum, namely making sense of the formula $\psi = \sum_{i \in \mathbb{N}} \psi_i$.

Recall that the module of horizontal elements of *M* is $M^{\text{hor}} = \{m \in M \mid \partial(m) = 0\}.$

Lemma 3.6. Under Setup 3.1, the following hold for the function $\psi : M \to M$ from Definition 3.5(2).

- (1) The function ψ is A-linear.
- (2) $\psi(m) \in M^{\text{hor}}$ for every $m \in M$.
- (3) If $m \in M^{\text{hor}}$ then $\psi(m) = m$.
- (4) $m \psi(m) \in \mathfrak{b} \cdot M$ for every $m \in M$.

Proof. (1) This is because each ψ_i is A-linear, and by the pointwise convergence of the sum $\psi(m) = \sum_{i \in \mathbb{N}} \psi_i(m)$ in M.

(2) Due to the continuity of ∂ , and using formula (3.4), for every $m \in M$ we have

$$\begin{split} \partial(\psi(m)) &= \sum_{i \in \mathbb{N}} \partial\big((i!)^{-1} \cdot (-t)^i \cdot \partial^i(m)\big) \\ &= \sum_{i \in \mathbb{N}} \left((i!)^{-1} \cdot (-t)^i \cdot \partial^{i+1}(m) - (i!)^{-1} \cdot i \cdot (-t)^{i-1} \cdot \partial^i(m)\right) \\ &= \left(\sum_{i \in \mathbb{N}} (i!)^{-1} \cdot (-t)^i \cdot \partial^{i+1}(m)\right) - \left(\sum_{j \in \mathbb{N}} (j!)^{-1} \cdot (-t)^j \cdot \partial^{j+1}(m)\right) = 0. \end{split}$$

(3) Take some $m \in M^{\text{hor}}$. Then $\psi_0(m) = m$, and $\psi_i(m) = 0$ for $i \ge 1$. Thus $\psi(m) = m$.

(4) Take some element $m \in M$. By definition we have $\psi_0(m) = m$, and $\psi_i(m) \in \mathfrak{b}^i \cdot M \subseteq \mathfrak{b} \cdot M$ for $i \ge 1$. Since $\mathfrak{b} \cdot M$ is closed, we have $\psi(m) - m = \sum_{i \ge 1} \psi_i(m) \in \mathfrak{b} \cdot M$. \Box

It is not immediately clear that $\psi : M \to M$ is continuous. One can show it by a messy calculation; but it will be an easy consequence later, see Corollary 3.14 below.

Define the *B*-module $\overline{M} := M / \mathfrak{b} \cdot M$, with canonical projection $\pi : M \twoheadrightarrow \overline{M}$.

Proposition 3.7 (Existence of a Solution). Under Setup 3.1, given an element $\overline{m} \in \overline{M}$, there exists an element $m \in M$ such that $\partial(m) = 0$ and $\pi(m) = \overline{m}$.

Later, in Corollary 4.3, we shall see that the element *m* is unique.

Proof. Let m' be some lifting of \overline{m} to M; namely $m' \in M$ satisfies $\pi(m') = \overline{m}$. Define $m := \psi(m') \in M$, where ψ is the operator from Definition 3.5(2). According to Lemma 3.6(2) the element m satisfies $\partial(m) = 0$, and by Lemma 3.6(4) we have $\pi(m) = \pi(m') = \overline{m}$.

Here is a result on induced $\mathcal{D}_{B/A}$ -modules. Let L be an A-module. The B-module $B \otimes_A L$ has an induced left $\mathcal{D}_{B/A}$ -module structure, with action $\partial(b \otimes l) = \partial(b) \otimes l$ for $b \in B$ and $l \in L$. Now define the B-module $N := B \widehat{\otimes}_A L$, the b-adic completion of $B \otimes_A L$ (where L is given the discrete topology). By continuity, the B-module N has a left $\mathcal{D}_{B/A}$ -module structure.

Lemma 3.8. For the B-module $N = B \widehat{\otimes}_A L$, with its induced $\mathcal{D}_{B/A}$ -module structure, the following assertions hold.

- (1) Every element $n \in N$ has a unique convergent power series expansion $n = \sum_{i\geq 0} t^i \otimes l_i$, with $l_i \in L$.
- (2) For $n \in N$, with its power series expansion from item (1), there is equality $\partial(n) = \sum_{i \ge 1} i \cdot t^{i-1} \otimes l_i$ in N.

(3) The A-module homomorphism $\alpha : L \to N^{\text{hor}}, \alpha(l) := 1_B \otimes l$, is bijective.

Proof. (1) For $i \in \mathbb{N}$ let $B_i := B/\mathfrak{b}^{i+1}$ and $N_i := N/\mathfrak{b}^{i+1} \cdot N$. Then $B_i \cong \bigoplus_{0 \le j \le i} A \cdot t^j$ and

$$(3.9) N_i \cong B_i \otimes_B N \cong B_i \otimes_A L \cong \bigoplus_{0 \le i \le i} t^j \otimes L$$

as *A*-modules, and these isomorphisms are compatible with the change of *i*. The image of *n* in N_i is uniquely a sum $\sum_{0 \le j \le i} t^j \otimes l_j$ with $l_j \in L$. In the inverse limit $N \cong \lim_{i \to i} N_i$ we obtain $n = \sum_{j \ge 0} t^j \otimes l_j$.

(2) For each *i* the derivation ∂ acts on $t^i \otimes l_i$ as follows: $\partial(t^i \otimes l_i) = \partial(t^i) \otimes l_i$. Also $\partial(t^i) = i \cdot t^{i-1}$. By *A*-linearity and continuity we get the stated expression for $\partial(n)$.

(3) Items (1)-(2) say that $n \in N^{\text{hor}}$ iff $l_i = 0$ for all $i \ge 1$, namely iff $n = t^0 \otimes l_0 = \alpha(l_0)$. \Box

Now we take the A-module $L := M^{hor}$. This gives us the complete B-module $N := B \widehat{\otimes}_A M^{hor}$ and the B-module homomorphism

(3.10)
$$\theta: N = B \widehat{\otimes}_A M^{\text{hor}} \to M.$$

Lemma 3.11. Consider $N = B \widehat{\otimes}_A M^{\text{hor}}$ as above, with the induced left $\mathcal{D}_{B/A}$ -module structure.

- (1) The B-module homomorphism $\theta : N \to M$ is a homomorphism of left $\mathcal{D}_{B/A}$ -modules.
- (2) The A-module homomorphism $\theta^{\text{hor}} := \theta|_{N^{\text{hor}}} : N^{\text{hor}} \to M^{\text{hor}}$ is bijective.

Proof. (1) It suffices to show that θ respects the action of the derivation ∂ . Let's denote by ∂_M and ∂_N the action of ∂ on the two modules. Take $n \in N$, with power series expansion $n = \sum_{i \ge 0} t^i \otimes l_i, l_i \in L = M^{\text{hor}}$, see Lemma 3.8(1). Since θ , ∂_M and ∂_N are continuous, and $\partial(l_i) = 0$, we get

$$\theta(\partial_N(n)) = \theta\left(\sum_{i\geq 1} i \cdot t^{i-1} \otimes l_i\right) = \sum_{l\geq 1} \theta(i \cdot t^{i-1} \otimes l_i) = \sum_{i\geq 1} i \cdot t^{i-1} \cdot l_i$$


and

$$\partial_N(\theta(n)) = \partial_N\left(\sum_{i\geq 0} t^i \cdot l_i\right) = \sum_{i\geq 1} i \cdot t^{i-1} \cdot l_i .$$

We see that $\theta \circ \partial_N = \partial_M \circ \theta$.

(2) According to Lemma 3.8(3) the homomorphism $\alpha : L \to N^{\text{hor}}$ is bijective. And $\theta^{\text{hor}} \circ \alpha = \text{id}_L$. Hence θ^{hor} is bijective.

We have the inclusion $\epsilon_M : M^{\text{hor}} \to M$, the projection $\pi_M : M \to \overline{M} = M / \mathfrak{b} \cdot M$, and their composition $\gamma_M := \pi_M \circ \epsilon_M$. There are similar homomorphisms for *N*. These sit in the following commutative diagram of *A*-modules.



Here is the main result of this section.

Theorem 3.13. Under Setup 3.1, the $\mathcal{D}_{B/A}$ -module homomorphism

$$\theta: B \widehat{\otimes}_A M^{\mathrm{hor}} \to M$$

is bijective.

Proof. The proof is divided into two steps. As before we write $N := B \widehat{\otimes}_A M^{\text{hor}}$.

Step 1. In this step we prove that $\theta : N \to M$ is surjective.

According to Proposition 3.7, the homomorphism $\gamma_M : M^{\text{hor}} \to \overline{M}$ is surjective. By Lemma 3.11(2) we know that $\theta^{\text{hor}} : N^{\text{hor}} \to M^{\text{hor}}$ is bijective. It follows that the homomorphism $\pi_M \circ \theta : N \to \overline{M}$ is surjective. Since both *B*-modules *M* and *N* are b-adically complete, the Complete Nakayama Theorem [Ye4, Theorem 1.10] says that $\theta : N \to M$ is surjective.

Step 2. In this step we prove that $\theta : N \to M$ is injective, and hence it is bijective.

Assume, for the sake of arriving at a contradiction, that $\theta : N \to M$ is not injective. Then there exists some nonzero element $n \in N$ such that $\theta(n) = 0$. The element n has a unique convergent power series expansion $n = \sum_{i\geq 0} t^i \otimes l_i$, with $l_i \in L = M^{\text{hor}}$; see Lemma 3.8(1).

Since *n* is nonzero, its b-adic order is finite, say $i_0 \in \mathbb{N}$. So in the power series expansion of *n* we have $l_i = 0$ for all $i < i_0$, and $l_{i_0} \neq 0$. Define $n' = \partial^{i_0}(n) \in N$. Let $n' = \sum_{i\geq 0} t^i \cdot l'_i$ be the power series expansion of *n'*, with $l'_i \in L$. According to Lemma 3.8(2) we have $l'_0 = i_0! \cdot l_{i_0}$, which is a nonzero element of $L = M^{\text{hor}}$. This means that $n' \notin \mathfrak{b} \cdot N$.

Let ψ_N be the idempotent operator from Definition 3.5(2), but for the complete \mathcal{D}_{B/A^-} module *N*. Define the element $n'' := \psi_N(n') \in N^{\text{hor}}$. By Lemma 3.6(4) we have $n' - n'' \in \mathfrak{b} \cdot N$, so $n'' \notin \mathfrak{b} \cdot N$, and hence n'' is a nonzero element of N^{hor} . By Lemma 3.11(2) the element $\theta(n'') = \theta^{\text{hor}}(n'') \in M^{\text{hor}}$ is nonzero.

On the other hand, we are assuming that $\theta(n) \in M$ is zero. Since the homomorphism θ is $\mathcal{D}_{B/A}$ -linear, see Lemma 3.11(1), we get $\theta(n') = \theta(\partial^{i_0}(n)) = \partial^{i_0}(\theta(n)) = 0$. For every j the operator ψ_j belongs to $\mathcal{D}_{B/A}$, and thus $\theta(\psi_j(n')) = \psi_j(\theta(n')) = 0$. But θ is continuous, so

$$\theta(n^{\prime\prime}) = \theta(\psi_N(n^\prime)) = \theta\left(\sum_{j\geq 0} \psi_j(n^\prime)\right) = \sum_{j\geq 0} \theta(\psi_j(n^\prime)) = 0.$$

This contradicts the calculation in the previous paragraph.

Corollary 3.14. Under Setup 3.1, the following hold:

- (1) There is an A-linear direct sum decomposition $M = M^{hor} \oplus \mathfrak{b} \cdot M$.
- (2) The homomorphism $\psi : M \to M$ is the A-linear projection onto M^{hor} in the decomposition in item (1). In particular, $\text{Ker}(\psi) = \mathfrak{b} \cdot M$.
- (3) The homomorphism $\psi: M \to M$ is continuous for the b-adic topology of M.

Proof. (1) The *A*-module *B* decomposes into $B = A \oplus \mathfrak{b}$. This gives a decomposition of $N := B \widehat{\otimes}_A M^{\text{hor}}$ into $N = M^{\text{hor}} \oplus \mathfrak{b} \widehat{\otimes}_A M^{\text{hor}}$. The isomorphism θ in the theorem carries this direct sum decomposition to M.

(2) By Lemma 3.6(2-3) we know that ψ is a projection (an idempotent operator) with image M^{hor} , and by Lemma 3.6(4) we know that $\text{Ker}(\psi) \subseteq \mathfrak{b} \cdot M$. This forces the equality $\text{Ker}(\psi) = \mathfrak{b} \cdot M$.

(3) The kernel of ψ is the open *A*-submodule $\mathfrak{b} \cdot M$.

Remark 3.15. The projection operator ψ that we use here is the same as the operator called *P* in formula (8.9.5) of [Ka], and in formula (3) of [Og]. We thank R. Källstrom for clarifying this to us.

Remark 3.16. By definition, for every $m \in M$ we have $\psi(m) = \sum_{i \geq \psi} \psi_i(m)$. Each ψ_i is a differential operator; but the operator ψ is merely continuous.

It seems plausible that after endowing the ring $\operatorname{End}_{A}^{\operatorname{cont}}(M)$ of continuous A-linear endomorphisms of M with a suitable topology, there will be equality $\psi = \sum_{i\geq 0} \psi_i$ inside $\operatorname{End}_{A}^{\operatorname{cont}}(M)$. Moreover, if such a topology can be produced, then the topological closure $\widehat{\mathcal{D}}_{B/A}$ of $\mathcal{D}_{B/A}$ inside $\operatorname{End}_{A}^{\operatorname{cont}}(M)$ will be some sort of ring of "pro-differential operators", containing the projection operator ψ . It is conceivable that the suitable topology on $\operatorname{End}_{A}^{\operatorname{cont}}(M)$ will make it into a *semi-topological ring*, in the sense of [Ye1, Section 1.2].

4. Complete \mathcal{D} -Modules – the High Dimensional Case

In this section we prove Theorem 4.1, which is Theorem 0.6 in the Introduction. As already mentioned, it is a rephrasing, and an improvement to the non-noetherian setting, of [Og, Theorem 1.3]; which itself is an improvement of [Ka, Proposition 8.9].

Let us recall a few facts from Sections 1 and 2. Consider a commutative ring A containing \mathbb{Q} . There are no finiteness assumptions on A. Let $B := A[[t_1, \ldots, t_n]]$, the ring of power series over A in n variables. The ring $\mathcal{D}_{B/A}$ of A-linear differential operators of B is generated by B and the partial derivatives $\partial_i = \partial/\partial_{t_i}$. Given a left $\mathcal{D}_{B/A}$ -module M, its horizontal submodule is the A-module

$$M^{\text{hor}} = \{m \in M \mid \partial_i(m) = 0 \text{ for all } i\} \subseteq M.$$

If *L* is an *A*-module, then the *B*-module $M := B \widehat{\otimes}_A L$ has an induced left $\mathcal{D}_{B/A}$ -module structure.

Theorem 4.1. Let A be a commutative ring containing \mathbb{Q} , let $B := A[[t_1, ..., t_n]]$, the ring of power series over A in n variables, and let $\mathfrak{b} \subseteq B$ be the ideal generated by the variables. Let M be a left $\mathcal{D}_{B/A}$ -module, which is \mathfrak{b} -adically complete as a B-module. Then the canonical $\mathcal{D}_{B/A}$ -module homomorphism

$$\theta: B \widehat{\otimes}_A M^{\mathrm{hor}} \to M$$

is bijective.

Proof. As done in [Be] and [Bj], our proof is by induction on *n*. For n = 1 this is Theorem 3.13.

Now assume that $n \ge 2$, and that the theorem is true for the ring $B' := A[[t_1, \ldots, t_{n-1}]]$. Writing $t := t_n$, we have an *A*-ring isomorphism $B \cong B'[[t]]$. Also, writing $\partial := \partial_n \in \mathcal{D}_{B/A}$, the operator ∂ commutes with the operators t_i and ∂_i , for all $1 \le i \le n-1$, inside the noncommutative ring $\mathcal{D}_{B/A}$. The ring $\mathcal{D}_{B/B'}$ is generated by *B* and ∂ .

For a subsequence (i_1, \ldots, i_k) of $(1, \ldots, n)$ let

$$M^{\partial_{i_1},\ldots,\partial_{i_k}} := \{m \in M \mid \partial_{i_1}(m) = \cdots = \partial_{i_k}(m) = 0\} \subseteq M.$$

Let us define the B'-module $M' := M^{\partial} = M^{\partial_n} \subseteq M$. The commutation relations mentioned above imply that M' is actually a $\mathcal{D}_{B'/A}$ -submodule of M. By definition we have $M^{\text{hor}} = M^{\partial_1,...,\partial_n}$, and hence $M^{\text{hor}} = (M')^{\partial_1,...,\partial_{n-1}}$.

The ideal $(t) \subseteq B$ is contained in the ideal \mathfrak{b} , and M is \mathfrak{b} -adically complete. Corollary 1.6 tells us that M is (t)-adically complete. Applying Theorem 3.13 to the ring homomorphism $B' \to B = B'[[t]]$, and the (t)-adically complete $\mathcal{D}_{B/B'}$ -module M, we conclude that the $\mathcal{D}_{B/B'}$ -module homomorphism $B \widehat{\otimes}_{B'} M' \to M$ is bijective. Let $\mathfrak{b}' \subseteq B'$ be the ideal (t_1, \ldots, t_{n-1}) . Using Corollary 1.6 once more, we know that M is \mathfrak{b}' -adically complete. Now $M' = \operatorname{Ker}(\partial)$, so it is a closed submodule of M for the \mathfrak{b}' -adic topology. According to Corollary 1.7, M' is a \mathfrak{b}' -adically complete B'-module. Since Theorem 4.1 is assumed to be true for B', and since $(M')^{\partial_1,\ldots,\partial_{n-1}} = M^{\operatorname{hor}}$, it follows that the $\mathcal{D}_{B'/A}$ -module homomorphism $B' \otimes_A M^{\operatorname{hor}} \to M'$ is bijective.

The associativity of complete tensor products implies that there are *B*-module isomorphisms

$$B \widehat{\otimes}_A M^{\text{hor}} \cong B \widehat{\otimes}_{B'} (B' \widehat{\otimes}_A M^{\text{hor}}) \cong B \widehat{\otimes}_{B'} M' \cong M.$$

The composed isomorphism is precisely $\theta : B \otimes_A M^{\text{nor}} \to M$.

In the corollaries below we assume the setting of the theorem.

Define the A-module $\overline{M} := M/\mathfrak{b} \cdot M$. Consider the inclusion $\epsilon : M^{\text{hor}} \to M$ and the projection $\pi : M \to \overline{M}$. Let $\gamma : M^{\text{hor}} \to \overline{M}$ be $\gamma := \pi \circ \epsilon$.

Corollary 4.2. The A-linear homomorphism $\gamma: M^{hor} \to \overline{M}$ is bijective.

Proof. This is obvious for the induced module $N := B \widehat{\otimes}_A M^{\text{hor}}$. According to Theorem 3.13 there is an isomorphism of left $\mathcal{D}_{B/A}$ -modules $N \cong M$, so the same holds for M. \Box

In view of Corollary 4.2, we can define the *A*-linear bijection $\sigma : \overline{M} \xrightarrow{\simeq} M^{\text{hor}}$ to be the inverse of *y*. We then have:

Corollary 4.3. Given an element $\overline{m} \in \overline{M}$, the element $m := \sigma(\overline{m}) \in M$ is the unique solution in M of the differential equation $\partial_1(m) = \cdots = \partial_n(m) = 0$ with initial condition $\pi(m) = \overline{m}$.

Corollary 4.4. There is an A-linear direct sum decomposition $M = M^{hor} \oplus \mathfrak{b} \cdot M$.

Proof. The same as the proof of Corollary 3.14(1).

A *B*-module *M* is called \mathfrak{b} -adically free if *M* is the \mathfrak{b} -adic completion of a free *B*-module. It is known that a \mathfrak{b} -adically free *B*-module *M* is isomorphic to the module of \mathfrak{b} -adically decaying functions $F_{dec}(Z, B)$ for some set *Z*; see [Ye3, Section 2].

Corollary 4.5. If $M/\mathfrak{b} \cdot M$ is a free A-module, then M is a \mathfrak{b} -adically free B-module.

Proof. The *A*-module $M^{\text{hor}} \cong M/\mathfrak{b} \cdot M$, so it is free. Then the *B*-module $B \otimes_A M^{\text{hor}}$ is free, and its \mathfrak{b} -adic completion $B \otimes_A M^{\text{hor}}$ is \mathfrak{b} -adically free.

The following corollary is [Ka, Proposition 8.9], which is phrased in terms of connections.

Corollary 4.6. Assume A is a field and M is a finitely generated B-module. Then $M \cong B^{\oplus r}$ as $\mathcal{D}_{B/A}$ -modules, for some natural number r.

Proof. Here $M/\mathfrak{b} \cdot M \cong A^{\oplus r}$ for some r, and $B^{\oplus r}$ is already complete. \Box

Remark 4.7. Suppose \mathbb{K} is a field of characteristic 0, X is a smooth \mathbb{K} -scheme, and \mathcal{M} is a sheaf of left $\mathcal{D}_{X/\mathbb{K}}$ -modules, which is coherent as an \mathcal{O}_X -module. According to [Be, Proposition 1.a] or [HTT, Theorem 1.4.10], \mathcal{M} is a locally free \mathcal{O}_X -module. The direct proof in [HTT] is very easy. They consider a closed point $x \in X$, the local ring $A := \mathcal{O}_{X,x}$, and the A-module $\mathcal{M} := \mathcal{M}_x$. It suffices to prove that \mathcal{M} is a free A-module. This is done by a calculation almost identical to the the one in the proof of Theorem 3.13.

It is possible to deduce the algebraic result above from the corresponding complete result, namely Theorem 4.1. Let \widehat{A} be the completion of the local ring A at its maximal ideal, and let \widehat{M} be the completion of the module M. Since $A \to \widehat{A}$ is faithfully flat,

it suffices to prove that \widehat{M} is a free \widehat{A} -module. The completed module \widehat{M} has a $\mathcal{D}_{\widehat{A}/\mathbb{K}}$ structure. The residue field \mathbb{K}' of A is a finite separable extension of \mathbb{K} , so it can be lifted uniquely into \widehat{A} , and then $\widehat{A} \cong \mathbb{K}'[[t_1, \ldots, t_n]]$. Now Theorem 4.1 can be applied.

The reverse direction – namely using [HTT, Theorem 1.4.10] to prove Theorem 4.1 – does not work, because we do not know a priori that \widehat{M} descends to some A-module M.

Furthermore, there are plenty of nonisomorphic $\mathcal{D}_{A/\mathbb{K}}$ -modules that trivialize when passing to the completion \widehat{A} . To see this explicitly, let's assume that $\mathbb{K} = \mathbb{C}$, n = 1 and the rank of M is 1. For every $\lambda \in \mathbb{C}$ there is a $\mathcal{D}_{A/\mathbb{C}}$ -module M_{λ} , which is a free A-module of rank 1 with basis e_{λ} , and such that $\partial(e_{\lambda}) = \lambda \cdot e_{\lambda}$. When passing to the completion, the \widehat{A} -module \widehat{M}_{λ} is still free with basis e_{λ} , but now the ring \widehat{A} contains the invertible element $u_{\lambda} := \exp(-\lambda \cdot t)$, and the element $u_{\lambda} \cdot e_{\lambda} \in \widehat{M}_{\lambda}$ is a horizontal basis, so that \widehat{M}_{λ} is a trivial $\mathcal{D}_{\widehat{A}/\mathbb{C}}$ -module.

Appendix A. An Error in a Book by Björk

While studying the literature on \mathcal{D} -modules, we looked at the proof of Theorem 1.1.25 in the book [Bj] by J.-E. Björk. This theorem is an analytic version of Corollary 0.9, which is a special case of Theorem 0.6. While reading the proof, we discovered an error in it. In this appendix we shall explain the error, and how it can be resolved.

Before doing so, we should address a question of etiquette. As customary, we wanted to communicate our discovery to the author of the book, giving him the possibility to address it. Unfortunately, Björk passed away several years ago. Therefore we communicated his student Rolf Källström, who is working in this same area. He confirmed that there is indeed an error, that this error had not been observed previously, and that we ought to write about it.

Now to the mathematics. We prefer to use our notation, and to simplify the situation as much as possible, so that the salient points will be clear. We also want to treat two settings at once:

- (*) The setting of our paper, specialized: $A = \mathbb{K}$ is a field of characteristic 0, and $B = \mathbb{K}[[t]]$. Convergence is for the b-adic topology, where $\mathfrak{b} = (t) \subseteq B$.
- (**) The complex analytic setting that is treated in the book [Bj]. Here $A = \mathbb{C}$, the field of complex numbers with its standard norm, and *B* is the subring of $\mathbb{C}[[t]]$ consisting of convergent power series, namely series $b = \sum_{k\geq 0} \lambda_k \cdot t^k$ with coefficients $\lambda_k \in \mathbb{C}$ satisfying the asymptotic condition $\limsup_{k\to\infty} (|\lambda_k|)^{1/k} < \infty$. In other words, *B* is the ring of germs of analytic functions in one variable at the origin. It is a noetherian local ring, with maximal ideal b generated by *t*.

The ring of differential operators $\mathcal{D}_{B/A}$ has the same structure in both settings. As before we write $\partial := \frac{\partial}{\partial t}$.

Suppose M is a left $\mathcal{D}_{B/A}$ -module that's finitely generated as a B-module. We can assume that M is a free B-module of rank r, with basis $\mathbf{m}' = (m'_1, \ldots, m'_r)$; for this we can either rely on [HTT, Theorem 1.4.10], or on the first half of the proof of [Bj, Theorem 1.1.25], which is correct (and is similar to the proof in [HTT]). The issue is how to replace the basis \mathbf{m}' with another basis \mathbf{m} , which is horizontal, and is congruent to \mathbf{m}' modulo the maximal ideal $\mathfrak{b} \subseteq B$. In other words, we want to find a matrix $\mathbf{g} = \mathbf{g}(t) \in \operatorname{GL}_r(B)$, $\mathbf{g} \equiv \mathbf{1} \mod t$, such that

(A.1)
$$\partial(\boldsymbol{g} \cdot \boldsymbol{m}') = 0$$

Then the basis $m := g \cdot m'$ of *M* will be horizontal.

The difficulty only arises when $r \ge 2$, so let us assume this. Björk considers the matrix $\mathbf{b} \in \operatorname{Mat}_r(B)$ such that $\partial(\mathbf{m}') = \mathbf{b} \cdot \mathbf{m}'$. The matrix \mathbf{b} has a power series expansion $\mathbf{b} = \sum_{k\ge 0} t^k \cdot \mathbf{a}_k$ with $\mathbf{a}_k \in \operatorname{Mat}_r(A)$. Convergence of this series in the setting (*) is easy; we did not verify convergence in the setting (**), but presumably this is automatic for standard reasons.

Next Björk defines the matrix

$$\boldsymbol{c} := \sum_{k\geq 0} (k+1)^{-1} \cdot \boldsymbol{t}^{k+1} \cdot \boldsymbol{a}_k \in \operatorname{Mat}_r(B),$$

which satisfies $\partial(c) = b$. Again, convergence is obvious in the adic setting (*), and requires justification in the analytic setting (**). Then he defines the matrix

$$\boldsymbol{g} := \exp(-\boldsymbol{c}) = \sum_{k \ge 0} (k!)^{-1} \cdot (-\boldsymbol{c})^k \in \operatorname{Mat}_r(B).$$

Observe that this sum converges in the adic setting (*) because the matrix c is divisible by t. In the complex analytic setting (**) convergence is because of the rapid decay of $(k!)^{-1}$.

The erroneous claim of Björk (implicit in the text, yet present in formula (2) there) is that the matrix $g \in Mat_r(B)$ satisfies the differential equation

(A.2)
$$\partial(\boldsymbol{g}) = \operatorname{talse} -\boldsymbol{g} \cdot \boldsymbol{b}$$

with initial condition $g(0) = 1 \in Mat_r(A)$. For the differential equation (A.2) to hold it is sufficient, and very likely also necessary, that the matrices a_k commute with each other.

Anyhow, here is a counterexample: take the ring $A := \mathbb{Q}$, so $B = \mathbb{Q}[[t]]$, and the rank r := 2. The matrix $\mathbf{c} = \mathbf{c}(t) \in \operatorname{Mat}_2(B)$ is $\mathbf{c} := t \cdot \mathbf{a}_1 + t^2 \cdot \mathbf{a}_2$, with $\mathbf{a}_1 := \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $\mathbf{a}_2 := \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. Recall that $\mathbf{b} = \partial(\mathbf{c})$ and $\mathbf{g} = \exp(-\mathbf{c})$. A calculation shows that $\partial(\mathbf{g}) \neq -\mathbf{g} \cdot \mathbf{b}$.

Björk proceeds as follows. He takes the sequence $m := g \cdot m'$ of elements of M. Here is our interpretation of Björk's formula (2):

(A.3)
$$\partial(\boldsymbol{m}) = \partial(\boldsymbol{g} \cdot \boldsymbol{m}') = \partial(\boldsymbol{g}) \cdot \boldsymbol{m}' + \boldsymbol{g} \cdot \partial(\boldsymbol{m}') = {}^{\text{false}} - \boldsymbol{g} \cdot \boldsymbol{b} \cdot \boldsymbol{m}' + \boldsymbol{g} \cdot \boldsymbol{b} \cdot \boldsymbol{m}' = 0.$$

The false equality $=^{\text{false}}$ relies on the false formula (A.2).

In the setting (**) of complex analytic functions, the correct way to solve the ODE (A.1) is by appealing to the standard existence results. Presumably this is what Deligne had in mind when he wrote "Il est bien connu que..." regarding [De, Theorem 2.17]. In [HTT, section 4.2], when discussing analytic \mathcal{D} -modules, they refer to the Frobenius Theorem.

In our *t*-adic setting (*), Lemma 3.6 and its proof give us an explicit formula for the matrix g solving the ODE (A.1). Given the basis $m' = (m'_1, \ldots, m'_r)$ of M, let $m_i := \psi(m'_i) \in M$ and $m := (m_1, \ldots, m_r)$, which is a horizontal basis of M. Then define $g \in GL_r(B)$ to be the matrix such that $g \cdot m' = m$. This matrix g will solve the ODE (A.1).

An alternative approach for obtaining the matrix g is by 1-dimensional nonabelian multiplicative integration, also known as path ordered exponential. This method would present g as the limit of *Riemann products*, cf. [Ye5, Chapter 3].

References

- [Be] J. Bernstein, Algebraic Theory of D-modules, preprint (1982-83), online http://www.math.tau.ac. il/~bernstei/Unpublished_texts/unpublished_texts/Bernstein82-dmod-from-chicago.pdf
- [Bj] J.-E. Björk, "Analytic D-Modules and Applications", Springer, 1993. Online https://doi.org/10. 1007/978-94-017-0717-6_1.
- [De] P. Deligne, "Equations Différentielles à Points Singuliers Reguliers", Lecture Notes in Mathematics 163, Springer, 1970. https://doi.org/10.1007/BFb0061194.

- [EGA IV] A. Grothendieck and J. Dieudonné, "Éléments de géometrie algébrique", Chapitre IV, Quatrième Partie, Publ. math. I.H.É.S. 32, 1967, online http://www.numdam.org/item/?id=PMIHES_1967__32__5_0.
- [HTT] R. Hotta, K. Takeuchi and T. Tanisaki, "D-modules, perverse sheaves, and representation theory", Progress in Mathematics 236, Birkhäuser, 2008, online https://link.springer.com/book/10.1007/ 978-0-8176-4523-6.
- [Ka] N.M. Katz, Nilpotent connections and the monodromy theorem: applications of a result of Turrittin, Publications mathématiques de l'I.H.É.S. 39 (1970), 175-232. Online http://www.numdam.org/item? id=PMIHES_1970_39_175_0.
- [Og] A. Ogus, Local Cohomological Dimension of Algebraic Varieties, Annals of Mathematics 98 (1973), 327-365. Online https://doi.org/10.2307/1970785.
- [Ye1] A. Yekutieli, "An Explicit Construction of the Grothendieck Residue Complex", Astérisque 208, 1992. Online http://www.numdam.org/item/AST_1992_208_3_0.
- [Ye2] A. Yekutieli, Continuous and Twisted L-infinity Morphisms, J. Pure Appl. Algebra 207 (2006), 575-606. DOI https://doi.org/10.1016/j.jpaa.2005.10.006.
- [Ye3] A. Yekutieli, On Flatness and Completion for Infinitely Generated Modules over Noetherian Rings, Comm. Algebra 39 (2011), DOI https://doi.org/10.1080/00927872.2010.522159.
- [Ye4] A. Yekutieli, Derived Complete Complexes at Weakly Proregular Ideals, eprint https://arxiv.org/ abs/2309.01687v3.
- [Ye5]A. Yekutieli, "Nonabelian Multiplicative Integration on Surfaces", World Scientific, 2015. DOI https:
//doi.org/10.1142/9537.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY, BE'ER SHEVA 84105, ISRAEL. Email address: amyekut@gmail.com URL: https://sites.google.com/view/amyekut-math/home