ON THE BLOW-UP FORMULA OF THE CHOW WEIGHTS FOR POLARIZED TORIC MANIFOLDS

KING LEUNG LEE AND NAOTO YOTSUTANI

Abstract. Let X be a smooth projective toric variety and let \tilde{X} be the blow-up manifold of X at finitely many distinct tours invariants points of X. In this paper, we give an explicit combinatorial formula of the Chow weight of \tilde{X} in terms of the base toric manifold X and the symplectic cuts of the Delzant polytope. We then apply this blow-up formula to the projective plane and see the difference of Chow stability between the toric blow-up manifolds and the manifolds of blow-ups at general points. Finally, we detect the blow-up formula of the Futaki-Ono invariant which is an obstruction for asymptotic Chow semistability of a polarized toric manifold.

1. INTRODUCTION

Through the paper, we use the same notation and terminology in [LY24]. Let (X, L) be an *n*-dimensional polarized manifold, where $L \to X$ is an ample line bundle over an *n*-dimensional compact Kähler manifold. Let $\operatorname{Aut}(X, L)$ be the group of holomorphic vector fields of (X, L) modulo the trivial automorphism $\mathbb{C}^{\times} := \mathbb{C} \setminus \{0\}$. Historically, it was proved by Donaldson that the existence of a constant scalar curvature Kähler (cscK) metric in the first Chern class $c_1(L)$ implies asymptotic Chow stability of (X, L) when $\operatorname{Aut}(X, L)$ is trivial [Don01]. As shown in [OSY12, DVZ12, Nak15, LLSW19], Donaldson's result can not be extended to the case where $\operatorname{Aut}(X, L)$ is non-trivial. In particular, Della Vedova and Zuddas used the blow-up argument effectively for constructing examples of asymptotically Chow unstable cscK classes. Remark that Odaka also constructed asymptotically Chow unstable cscK orbifolds with trivial $\operatorname{Aut}(X, L)$ by using a blow-up formula of the Donaldson-Futaki (DF) invariant along the flag ideals [Oda12].

Now we briefly recall the main argument in [DVZ12]. Della Vedova and Zuddas firstly observed that the obstruction for asymptotic Chow semistability of (X, L) does not vanish after taking the blow-ups at distinct points. 'based on this observation, they detected that

- (1) the blow-up formula of the obstruction for asymptotic Chow semistability, and
- (2) if we an appropriate polarization of the blow-up manifold (X, L), then it admits a cscK class by the result of Arezzo-Pacard [AP09].

Consequently, they found many new examples of asymptotically Chow unstable cscK classes in two dimension. Hence, these previous works show that the importance of the effect of blow-ups of a polarized manifold when we consider the existence problem of cscK metrics.

Remark 1.1. Concerning (1), we call the *Futaki-Ono invariant* an obstruction for a polarized toric variety (X, L) to be asymptotically Chow semistable (see, Section 5). On the other hand, the DF invariant of the blow-up manifold had been formulated and worked out by Stoppa in [Sto10].

The current paper aims to provide an explicit formula of the Chow weight of a toric blow-up manifold using symplectic cutting [Ler95, LT97]. To state our main result more precisely, for an

Date: July 16, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary 51M20; Secondary 53C55, 14M25.

Key words and phrases. Toric varieties, Chow weight, Blow-ups, Lattice polytope.

n-dimensional lattice Delzant moment polytope Δ , let $E_{\Delta}(t)$ and $s_{\Delta}(t)$ be the Ehrhart polynomial and the lattice sum points polynomial respectively. We refer the reader Section 2 for more detailed description. Let $(X_{\Delta}, \mathcal{O}_{X_{\Delta}}(i))$ be the associated polarized toric manifold, where *i* is any positive integer. Then we consider the *Chow weight* of $(X_{\Delta}, \mathcal{O}_{X_{\Delta}}(i))$ defined by

$$\operatorname{Chow}_{\Delta}(\boldsymbol{x}; i) = \operatorname{vol}(\Delta)\boldsymbol{s}_{\Delta}(i) - E_{\Delta}(i) \int_{\Delta} \boldsymbol{x} \, dv.$$

For an *n*-dimensional Delzant lattice polytope Δ , let $(X_{\Delta}, \omega_{\Delta}, \mathbb{T}^n, \mu_{\Delta})$ denote the associated symplectic toric manifold, where $\mathbb{T}^n = (S^1)^n$ and $\mu_{\Delta} : X_{\Delta} \to \operatorname{Lie}(\mathbb{T}^n) = \mathbb{R}^n$ is the corresponding moment map. Then, the symplectic blow-up of $(X_{\Delta}, \omega_{\Delta})$ at a fixed point of \mathbb{T}^n -action on $(X_{\Delta}, \omega_{\Delta})$ gives a new symplectic toric manifold. More precisely, let p be a fixed point of \mathbb{T}^n -action on $(X_{\Delta}, \omega_{\Delta})$ and let $p = \mu_{\Delta}(p)$ the corresponding vertex of Δ . For the symplectic blow-up

$$\varpi: X := \mathrm{Bl}_p(X_\Delta) \dashrightarrow X_\Delta,$$

it is known that the moment polytope $\widetilde{\Delta}$ for the blow-up $\widetilde{X}, \widetilde{\omega}$ of $(X_{\Delta}, \omega_{\Delta})$ at p is obtained from Δ by chopping off the corner $p \in \mathcal{V}(\Delta)$ (cf.[CdS08, Chapter 29]). Here $\mathcal{V}(\Delta)$ denotes the set of vertices of Δ . Regarding this operation, we have the decomposition of the polytope Δ such as

(1.1)
$$\Delta = \Delta' \cup \bigcup_{a=1}^{\ell} D_a,$$

if we consider the blow-up of $(X_{\Delta}, \omega_{\Delta})$ at \mathbb{T}^n -invariant points $p_1, p_2, \ldots, p_{\ell}$:

$$Y_{\Delta'} = \operatorname{Bl}_{(p_1, \dots, p_\ell)} \dashrightarrow X_{\Delta}.$$

Here each D_a in (1.1) denotes the *n*-dimensional simplex containing a vertex of Δ uniquely. See Section 3.1 for more detailed terminologies. The main theorem in the current paper is the following.

Theorem 1.2 (See, Theorem 3.3). Let $\Delta \subset \mathbb{R}^2$ be a two dimensional lattice Delzant polytope with the decomposition (1.1) which corresponds to the toric symplectic blow-up of X_{Δ} at the torus fixed points $p_1, p_2, \ldots, p_{\ell}$:

$$\varpi: Y_{\Delta'} = \operatorname{Bl}_{(p_1, \dots, p_\ell)} \dashrightarrow X_{\Delta}.$$

Let $k \in \mathbb{N}$ be the minimum integer such that $k\Delta'$ is a lattice Delzant polytope, and let $\widetilde{Y} := Y_{k\Delta'}$ denote the associated toric manifold. For any positive integer *i*, the Chow weight of $(\widetilde{Y}, \mathcal{O}_{\widetilde{Y}}(i))$ is

$$\operatorname{Chow}_{k\Delta'}(\boldsymbol{x};i) = \operatorname{Chow}_{k\Delta}(\boldsymbol{x};i) + i\mathcal{DF}_{k\Delta,1} + \mathcal{DF}_{k\Delta,2}$$

where $\mathcal{DF}_{k\Delta,1}$ and $\mathcal{DF}_{k\Delta,2}$ are combinatorial invariants defined in (3.9) and (3.10) respectively.

See Proposition 3.1 for arbitrary dimensional case. We then apply Theorem 1.2 to the projective plane $\mathbb{C}P^2$ and detect that a toric blow-up manifold $\mathbb{C}P^2\#6\overline{\mathbb{C}P^2}$ (a blow-up of $\mathbb{C}P^2$ at six torus fixed points of the \mathbb{T}^n -action) is Chow unstable. Meanwhile, let dP_3 be the del Pezzo surface of degree three which is obtained by the blow-up of $\mathbb{C}P^2$ at six distinct points in general position. Here six points in general position means that no three points are collinear and no six points lie on a conic. Then, it is known that $(dP_3, -K_{dP_3})$ is (asymptotically) Chow stable. Hence, the examples show that the significant difference of geometric properties of the blow-up manifolds when we consider (a) the toric blow-ups and (b) the blow-ups at points in general positions. See, Section 4.3 for these observations. Finally, we consider the blow-up formula of the FO invariant which was defined in our previous paper [LY24]. We also remark that the asymptotic behavior of such a combinatorial invariant (which is called the k-quantized barycenter in [JR24]) is well investigated by Jin and Rubinstein. In [JR24], they also established asymptotic expansion with respect to the tensor power k of the line bundle for algebro-geometric invariants used in Kähler geometry.

This paper is organized as follows. In Section 2, we briefly explain our notation and terminology used in this article. We also describe the Chow weight of a toric manifold in terms of the associated Delzant polytope Δ , and show the fundamental properties of the Chow weight under affine transformations of Δ (Proposition 2.3). In Section 3, we give an explicit characterization of the blow-up formula of the Chow weight for a polarized toric manifold. We prove Theorem 1.2 in Section 3.3. Section 4 is devoted to applying Theorem 1.2 and producing explicit examples which show the difference of geometric properties between toric blow-ups and blow-ups in general setting. In Section 4.4, we derive an explicit formula of the Chow weights of *all* toric surfaces whose moment polytopes have five vertices. In the final section, we shall see the blow-up for the FO invariant and an example of toric surface whose FO-invariant vanishes identically.

Acknowledgements. The second author (NY) would like to thank Professor Hajime Ono for inspiring discussions on this subject when NY was a postdoc researcher at USTC in China. The main idea on the construction of the blow-up formula in Section 3 is based on his unpublished work. This work was supported by the first author's ANR-21-CE40-0011 JCJC project MARGE, and the second author's JSPS KAKENHI 22K03316 (Grant-in-Aid for Scientific Research (C)).

2. TRANSFORMATION AND MULTIPLICATION RULES OF COMBINATORIAL INVARIANTS OF DELZANT POLYTOPES

Let M be a free abelian group of rank n such that $M \cong \mathbb{Z}^n$. Let Δ be an n-dimensional lattice Delzant polytope in $M_{\mathbb{R}} := M \otimes \mathbb{R} \cong \mathbb{R}^n$. Let $E_{\Delta}(t)$ be the Ehrhart polynomial of Δ which is written in the form of

(2.1)
$$E_{\Delta}(t) = \operatorname{Vol}(\Delta)t^{n} + \frac{\operatorname{Vol}(\partial\Delta)}{2}t^{n-1} + O(t^{n-2}) \quad \text{and} \\ E_{\Delta}(i) = \#(i\Delta \cap \mathbb{Z}^{n}) = \#(\Delta \cap (\mathbb{Z}/i)^{n})$$

for any positive integer *i*, where $\partial \Delta$ denotes the boundary of Δ . In particular, $E_{\Delta}(i)$ counts the number of lattice points in the *i*-th dilation of a polytope Δ .

Similar to the Ehrhart polynomial, it is known that we have the \mathbb{R}^n -valued polynomial satisfying

(2.2)
$$s_{\Delta}(i) := \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} \boldsymbol{a} = \frac{1}{i} \sum_{\boldsymbol{a} \in i \Delta \cap \mathbb{Z}^n} \boldsymbol{a}$$
$$= i^n \int_{\Delta} \boldsymbol{x} \, dv + \frac{i^{n-1}}{2} \int_{\partial \Delta} \boldsymbol{x} \, d\sigma + \dots + \boldsymbol{s}_{\Delta}$$

for any positive integer *i*. Here $d\sigma$ is the (n-1)-dimensional Lebesgue measure of $\partial\Delta$ defined as follows: let $h_i(\mathbf{x}) = \langle \mathbf{x}, \mathbf{v}_i \rangle + a_i$ be the defining function of the facet $F_i = \{ \mathbf{x} \in \Delta \mid h(\mathbf{x}) = 0 \}$ of Δ . Let $dv = dx_1 \wedge \cdots \wedge dx_n$ be the standard volume form of \mathbb{R}^n . On each facet $F_i \subset \partial\Delta$, we define the (n-1)-dimensional Lebesgue measure $d\sigma_i = d\sigma|_{F_i}$ by

$$dv = \pm d\sigma_i \wedge dh_i.$$

We call $s_{\Delta}(t)$ the *lattice sum points polynomial*.

Now we recall that the general theory of these phenomena, namely the *Euler-Maclaurin sum*mation formula for lattice simple polytopes. This provides a powerful connection between the integral over a lattice Delzant polytope Δ and the summation of lattice points in Δ as in (2.2). More specifically, let $f(\mathbf{x}) = A\mathbf{x}$ with $A \in GL_n(\mathbb{R})$ be an affine linear function on \mathbb{R}^n . For a positive integer *i*, we consider the summation

(2.3)
$$P_{\Delta}(f;i) := \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} f(\boldsymbol{a})$$

and see the polynomial behaviour in *i* of (2.3). By (2.2) and the linearity of f(x), we see that $P_{\Delta}(f;i)$ can be written in the form of

$$P_{\Delta}(f;i) = i^n \int_{\Delta} f(\boldsymbol{x}) dv + \frac{i^{n-1}}{2} \int_{\partial \Delta} f(\boldsymbol{x}) d\sigma + O(i^{n-2})$$

Definition 2.1. Let Δ be an *n*-dimensional Delzant lattice polytope in $M_{\mathbb{R}} \cong \mathbb{R}^n$. Let $f(\boldsymbol{x})$ be an affine linear function on \mathbb{R}^n . We define the *Chow weight* of $i\Delta$ with respect to f by

(2.4)
$$\operatorname{Chow}_{\Delta}(f;i) := \operatorname{Vol}(\Delta) P_{\Delta}(f;i) - E_{\Delta}(i) \int_{\Delta} f(\boldsymbol{x}) dv.$$

As a special case, if we take f(x) to be the coordinate function (i.e., f(x) = x), then (2.4) becomes

(2.5)
$$\operatorname{Chow}_{\Delta}(\boldsymbol{x};i) = \operatorname{Vol}(\Delta)\boldsymbol{s}_{\Delta}(i) - E_{\Delta}(i) \int_{\Delta} \boldsymbol{x} \, dv,$$

which agrees with Ono's obstruction for Chow semistability of a toric manifold $(X_{\Delta}, \mathcal{O}_{X_{\Delta}}(i))$ given in [Ono11, Theorem 1.2]. In particular, we remark the followings.

Remark 2.2. (1) $\operatorname{Chow}_{\Delta}(f; i)$ is an \mathbb{R}^n -valued polynomial of degree n-1 in i.

- (2) If the corresponding polarized toric manifold (X_{Δ}, L_{Δ}) with $L_{\Delta} = \mathcal{O}_{X_{\Delta}}(1)$ is asymptotically Chow semistable, then $\text{Chow}_{\Delta}(f; i) \equiv \mathbf{0}$ for any positive integer i and $f(\mathbf{x})$.
- (3) As it was mentioned in [LY24, (1.2)], the coefficients of $\operatorname{Chow}_{\Delta}(f; i)$ in (2.5) coincides with the linear span of $\{ \mathcal{F}_{\mathrm{Td}^p} |_{\mathbb{C}^n} : p = 1, \ldots, n \}$, where $\mathcal{F}_{\mathrm{Td}^p}$ is a family of integral invariants defined in [Fut04].

Chow weights defined in (2.4) enjoy the following properties under affine transformations of Δ .

Proposition 2.3. Let Δ be an *n*-dimensional lattice Delzant polytope in \mathbb{R}^n . Let *i* be any positive integer, and let $f(\mathbf{x})$ be an affine linear function on \mathbb{R}^n .

(1) Under any parallel transformation $\Delta + c$ in \mathbb{R}^n with $c \in \mathbb{Z}^n$, we have

$$\operatorname{Chow}_{\Delta+\boldsymbol{c}}(f;i) = \operatorname{Chow}_{\Delta}(f;i).$$

(2) Under any $SL_n(\mathbb{Z})$ -transformation $A\Delta$ in \mathbb{R}^n with $A \in SL_n(\mathbb{Z})$, we have

$$\operatorname{Chow}_{A\Delta}(f;i) = A\operatorname{Chow}_{\Delta}(f;i).$$

(3) For any constant multiplication $k\Delta$ in \mathbb{R}^n with $k \in \mathbb{N}$, we have

$$\operatorname{Chow}_{k\Delta}(f;i) = k^{n+1} \operatorname{Chow}_{\Delta}(f;ki).$$

Proof. (1). For $c \in \mathbb{Z}^n$, we observe that

$$E_{\Delta+\boldsymbol{c}} = E_{\Delta}(i), \quad \operatorname{Vol}(\Delta+\boldsymbol{c}) = \operatorname{Vol}(\Delta),$$

$$P_{\Delta+\boldsymbol{c}}(f;i) = \sum_{\boldsymbol{a}-\boldsymbol{c}\in\Delta\cap(\mathbb{Z}/i)^n} \boldsymbol{f}(a) = \sum_{\boldsymbol{b}\in\Delta\cap(\mathbb{Z}/i)^n} \boldsymbol{f}(\boldsymbol{b}+\boldsymbol{c})$$

$$= \sum_{\boldsymbol{b}\in\Delta\cap(\mathbb{Z}/i)^n} \boldsymbol{f}(\boldsymbol{b}) + \sum_{\boldsymbol{c}\in\Delta\cap(\mathbb{Z}/i)^n} \boldsymbol{c} = P_{\Delta}(f;i) + E_{\Delta}(i)\boldsymbol{c}, \quad \text{and}$$

$$\int_{\Delta+\boldsymbol{c}} \boldsymbol{f}(\boldsymbol{x})dv = \int_{\Delta} \boldsymbol{f}(\boldsymbol{x})dv + \boldsymbol{c} \int_{\Delta} 1 \, dv = \int_{\Delta} \boldsymbol{f}(\boldsymbol{x})dv + \operatorname{Vol}(\Delta)\boldsymbol{c}.$$

These yield that

$$\operatorname{Chow}_{\Delta+\boldsymbol{c}}(f;i) = \operatorname{Vol}(\Delta+\boldsymbol{c}) \left(P_{\Delta}(f;i) + E_{\Delta}(i)\boldsymbol{c} \right) - E_{\Delta+\boldsymbol{c}}(i) \left\{ \int_{\Delta} f(\boldsymbol{x}) dv + \operatorname{Vol}(\Delta)\boldsymbol{c} \right\}$$
$$= \operatorname{Vol}(\Delta) P_{\Delta}(f;i) - E_{\Delta}(i) \int_{\Delta} f(\boldsymbol{x}) dv = \operatorname{Chow}_{\Delta}(f;i).$$

(2). For $A \in SL_n(\mathbb{Z})$, we see that

$$E_{A\Delta}(i) = \#(i(A\Delta) \cap \mathbb{Z}^n) = \#(A(i\Delta \cap \mathbb{Z}^n)) = E_{\Delta}(i),$$

$$\operatorname{Vol}(A\Delta) = |\det A| \cdot \operatorname{Vol}(\Delta) = \operatorname{Vol}(\Delta),$$

$$P_{A\Delta}(f;i) = \sum_{\boldsymbol{a} \in A\Delta \cap (\mathbb{Z}/i)^n} f(\boldsymbol{a}) = \sum_{\boldsymbol{a} \in A(\Delta \cap (\mathbb{Z}/i)^n)} f(\boldsymbol{a}) = A \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} f(\boldsymbol{a}) = A \cdot P_{\Delta}(f;i),$$

$$\int_{A\Delta} f(\boldsymbol{x}) dv = |\det A| \cdot A \int_{\Delta} f(\boldsymbol{x}) dv = A \int_{\Delta} f(\boldsymbol{x}) dv,$$

where $|\det A| = 1$. Therefore, we find that

because $|\det A| = 1$. Therefore, we find that

$$\operatorname{Chow}_{A\Delta}(f;i) = \operatorname{Vol}(A\Delta)A \cdot P_{\Delta}(f;i) - E_{A\Delta}(i) \cdot A \int_{\Delta} f(\boldsymbol{x}) d\boldsymbol{v} = A \operatorname{Chow}_{\Delta}(f;i).$$

(3). For $k \in \mathbb{N}$, we have

$$E_{k\Delta}(i) = \#(i(k\Delta) \cap \mathbb{Z}^n) = E_{\Delta}(ki), \quad \operatorname{Vol}(k\Delta) = k^n \operatorname{Vol}(\Delta),$$

$$P_{k\Delta}(f;i) = \sum_{\boldsymbol{a} \in k\Delta \cap (\mathbb{Z}/i)^n} f(\boldsymbol{a}) = \sum_{\boldsymbol{a} \in k(\Delta \cap (\mathbb{Z}/ki)^n)} f(\boldsymbol{a}) = k \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/ki)^n} f(\boldsymbol{a}) = k P_{\Delta}(f;ki),$$

$$\int_{k\Delta} f(\boldsymbol{x}) dv = \int_{\Delta} f(k\boldsymbol{x}) k^n dv = k^{n+1} \int_{\Delta} f(\boldsymbol{x}) dv$$

by linearity of $f(\boldsymbol{x})$. Hence, we conclude that

$$\operatorname{Chow}_{k\Delta}(f;i) = k^{n} \operatorname{Vol}(\Delta) \cdot k P_{\Delta}(f;ki) - E_{\Delta}(ki) \cdot k^{n+1} \int_{\Delta} f(\boldsymbol{x}) d\boldsymbol{x}$$
$$= k^{n+1} \operatorname{Chow}_{\Delta}(f;ki).$$

Thus, the assertions are verified.

3. BLOW-UP FORMULA OF CHOW WEIGHTS OF PROJECTIVE TORIC MANIFOLDS

In this section, we shall give an explicit form of the blow-up formula of Chow weights of a polarized toric manifold.

3.1. Setting up. Let $\Delta \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ be an *n*-dimensional lattice Delzant polytope. Suppose that Δ is decomposed into

$$\Delta = \Delta' \cup \bigcup_{a=1}^{\ell} D_a$$

with the following properties:

- there exists $k \in \mathbb{N}$ such that $k\Delta'$ is an *n*-dimensional lattice Delzant polytope. We further assume that k is the minimum integer such that $k\Delta'$ is an *n*-dimensional lattice Delzant polytope.
- for each a = 1,..., l, D_a is an n-dimensional simplex which contains a vertex of Δ uniquely. We denote this unique vertex of Δ by p_a.
- for each distinct $a, b = 1, \ldots, \ell$, we set

$$D_a \cap \Delta' = \partial D_a \cap \Delta' =: L_a, \quad \text{and} \quad D_a \cap D_b = \emptyset.$$

In the geometry of symplectic toric manifolds, the blow-up of a toric symplectic manifold X_{Δ} at fixed points under the torus action corresponds to the operation such that chopping off the corners of the Delzant polytope Δ . Hence, the decomposition (3.1) induces a toric morphism

$$(3.2) X_{\Delta'} := \operatorname{Bl}_{(p_1,\dots,p_\ell)}(X_{\Delta}) \dashrightarrow X_{\Delta},$$

where we denote by p_1, \ldots, p_ℓ which are torus invariant points in X_Δ corresponding to the vertices p_1, \ldots, p_ℓ of Δ .

3.2. Arbitrary dimensional case. Firstly, we establish a combinatorial description of the blow-up formula of Chow weights in (3.2) in terms of the decomposition (3.1). Let us define

$$P_{kL_a}(f;i) := \sum_{\boldsymbol{a} \in kL_a \cap (\mathbb{Z}/i)^n} f(\boldsymbol{a}) \quad \text{and} \quad E_{kL_a}(i) := \#(kL_a \cap (\mathbb{Z}/i)^n)$$

respectively. Through the decomposition (3.1), we observe that each invariant in Section 2 satisfies the following property:

(3.3)

$$\operatorname{Vol}(k\Delta) = \operatorname{Vol}(k\Delta') + \sum_{a=1}^{\ell} \operatorname{Vol}(kD_a), \qquad \int_{k\Delta} f(\boldsymbol{x}) d\boldsymbol{v} = \int_{k\Delta'} f(\boldsymbol{x}) d\boldsymbol{v} + \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) d\boldsymbol{v},$$
$$P_{k\Delta}(f;i) = P_{k\Delta'}(f;i) + \sum_{a=1}^{\ell} (P_{kD_a}(f;i) - P_{kL_a}(f;i)), \qquad \text{and}$$
$$E_{k\Delta}(i) = E_{k\Delta'}(i) + \sum_{a=1}^{\ell} (E_{kD_a}(i) - E_{kL_a}(i)).$$

The following proposition is the blow-up formula of Chow weights with respect to $X_{\Delta'} \dashrightarrow X_{\Delta}$ in arbitrary dimensional case.

Proposition 3.1. We use the same notation as in Section 3.1. Then, we have

$$\begin{aligned} \operatorname{Chow}_{k\Delta'}(f;i) &= \operatorname{Chow}_{k\Delta}(f;i) \\ &- \left(\sum_{a=1}^{\ell} \operatorname{Vol}(kD_a)\right) P_{k\Delta}(f;i) - \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \operatorname{Vol}(kD_b)\right) \sum_{a=1}^{\ell} (P_{kD_a}(f;i) - P_{kL_a}(f;i)) \\ &+ E_{k\Delta}(i) \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) dv + \sum_{b=1}^{\ell} (E_{kD_b}(i) - E_{kL_b}(i)) \left(\int_{k\Delta} f(\boldsymbol{x}) dv - \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) dv\right). \end{aligned}$$

Proof. Plugging each value of (3.3) into (2.4), we find that

$$\begin{aligned} \operatorname{Chow}_{k\Delta'}(f;i) &= \operatorname{Vol}(k\Delta') P_{k\Delta'}(f;i) - E_{k\Delta'}(i) \int_{k\Delta'} f(\boldsymbol{x}) dv \\ &= \left(\operatorname{Vol}(k\Delta) - \sum_{a=1}^{\ell} \operatorname{Vol}(kD_a) \right) \left\{ P_{k\Delta}(f;i) - \sum_{a=1}^{\ell} (P_{kD_a}(f;i) - P_{kL_a}(f;i)) \right\} \\ &- \left\{ E_{k\Delta}(i) - \sum_{a=1}^{\ell} (E_{kD_a}(i) - E_{kL_a}(i)) \right\} \left(\int_{k\Delta} f(\boldsymbol{x}) dv - \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) dv \right) \\ &= \operatorname{Chow}_{k\Delta}(i) \\ &- \left(\sum_{a=1}^{\ell} \operatorname{Vol}(kD_a) \right) P_{k\Delta}(f;i) - \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \operatorname{Vol}(kD_b) \right) \sum_{a=1}^{\ell} (P_{kD_a}(f;i) - P_{kL_a}(f;i)) \\ &+ E_{k\Delta}(i) \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) dv + \sum_{b=1}^{\ell} (E_{kD_b}(i) - E_{kL_b}(i)) \left(\int_{k\Delta} f(\boldsymbol{x}) dv - \sum_{a=1}^{\ell} \int_{kD_a} f(\boldsymbol{x}) dv \right). \end{aligned}$$

This completes the proof.

3.3. Two dimensional case. Now we consider the case where a full-dimensional lattice polytope $\Delta \subset \mathbb{R}^n$ is a two dimensional Delzant polygon. In the case of n = 2, the Ehrhart polynomial in (2.1) is expressed as

$$E_{\Delta}(i) = \operatorname{Vol}(\Delta)i^2 + \frac{\#(\partial \Delta \cap \mathbb{Z}^2)}{2}i + 1$$

by Pick's formula. Here and hereafter, we only consider the case of f(x) = x for simplicity.

We note that each 2-dimensional simplex D_a in (3.1) can be written as follows:

let $\begin{pmatrix} \alpha_a \\ \beta_a \end{pmatrix}$, $\begin{pmatrix} \gamma_a \\ \delta_a \end{pmatrix} \in \mathbb{Z}^2$ be primitive generators of each edge emanating from the vertex \boldsymbol{p}_a . The orientation of each $\begin{pmatrix} \alpha_a \\ \beta_a \end{pmatrix}$ and $\begin{pmatrix} \gamma_a \\ \delta_a \end{pmatrix}$ can be uniquely determined by the condition

(3.4)
$$\det A_a = 1, \qquad A_a := \begin{pmatrix} \alpha_a & \gamma_a \\ \beta_a & \delta_a \end{pmatrix}.$$

Since $k\Delta'$ is a lattice Delzant polygon by our assumption (see, Section 3.1), there is some positive integer $m_a \in \mathbb{N}$ such that

$$D_a = \Delta(\boldsymbol{p}_a, \boldsymbol{q}_a, \boldsymbol{r}_a) := \operatorname{conv} \left\{ \boldsymbol{p}_a, \boldsymbol{q}_a, \boldsymbol{r}_a \right\}, \text{ where}$$
$$\boldsymbol{q}_a = \boldsymbol{p}_a + \frac{m_a}{k} \begin{pmatrix} \alpha_a \\ \beta_a \end{pmatrix} \quad \text{and} \quad \boldsymbol{r}_a = \boldsymbol{p}_a + \frac{m_a}{k} \begin{pmatrix} \gamma_a \\ \delta_a \end{pmatrix}.$$

Let Δ_{m_a} be the rectangular triangle

$$\operatorname{conv}\left\{ \begin{pmatrix} 0\\0 \end{pmatrix}, \begin{pmatrix} m_a\\0 \end{pmatrix}, \begin{pmatrix} 0\\m_a \end{pmatrix} \right\}.$$

Then, we see that the equality

$$kD_a = A_a \Delta_{m_a} + k \boldsymbol{p}_a$$

holds. Let l_{m_a} be the line connecting $\binom{m_a}{0}$ and $\binom{0}{m_a}$. Then, straightforward computations show that we have the following equalities.

Claim 3.2.

(3.6)
$$\operatorname{Vol}(\Delta_{m_a}) = \frac{m_a^2}{2},$$
$$s_{\Delta_{m_a}}(i) - s_{l_{m_a}}(i) = \frac{m_a}{6}(im_a + 1)(im_a - 1)\begin{pmatrix}1\\1\end{pmatrix},$$

(3.7)
$$E_{\Delta m_a}(i) - E_{l_{m_a}}(i) = \frac{im_a}{2}(im_a + 1) \quad and$$
$$\int_{\Delta m_a} \boldsymbol{x} \, dv = \frac{m_a^3}{6} \begin{pmatrix} 1\\ 1 \end{pmatrix}.$$

Proof. It suffices to compute (3.6) and (3.7) for the proof. Since our polygon Δ_{m_a} is symmetric with respect to the x-axes and the y-axes, we shall only compute the x-axes. For each $k = 0, 1, \ldots, im_a$, there are $im_a - k$ points in $(i\Delta_{m_a} \setminus il_{m_a}) \cap \mathbb{Z}^2$ whose the x component is k. Therefore, the x component of the left hand side of (3.6) is

$$\frac{1}{i}\sum_{k=0}^{im_a} k(im_a - k) = \frac{im_a^2}{2}(im_a + 1) - \frac{m_a}{6}(im_a + 1)(2im_a + 1)$$
$$= \frac{m_a}{6}(im_a + 1)(3im_a - 2im_a - 1) = \frac{m_a}{6}(im_a + 1)(im_a - 1).$$

Similarly, (3.7) is given by

$$E_{\Delta_{m_a}}(i) - E_{l_{m_a}}(i) = \sum_{k=0}^{im_a} (im_a - k) = \sum_{j=1}^{im_a} j = \frac{1}{2}im_a(im_a + 1).$$

This completes the proof of Claim 3.2.

-		
г		
L		

Using Claim 3.2, (3.5) and the transformation rules discussed in Section 2, we find that

(3.8)

$$\operatorname{Vol}(kD_a) = \frac{m_a^2}{2},$$

$$s_{kD_a}(i) - s_{kL_a}(i) = \frac{m_a}{6}(im_a + 1)(im_a - 1) \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + \frac{im_a}{2}(im_a + 1)kp_a,$$

$$E_{kD_a}(i) - E_{kL_a}(i) = \frac{im_a}{2}(im_a + 1),$$

$$\int_{kD_a} \boldsymbol{x} \, dv = \frac{m_a^3}{6} \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + \frac{m_a^2}{2}kp_a.$$

Setting

$$M := \sum_{a=1}^{\ell} m_a, \quad \widetilde{M} := \sum_{a=1}^{\ell} m_a^2, \quad A := \#(\partial(k\Delta) \cap \mathbb{Z}^2) - M \quad \text{and} \quad B := 2\text{Vol}(k\Delta) - \widetilde{M},$$

respectively, we define $\mathcal{DF}_{k\Delta,1}$ and $\mathcal{DF}_{k\Delta,2}$ by

$$\mathcal{DF}_{k\Delta,1} = \frac{A}{12} \sum_{a=1}^{\ell} m_a^3 \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + \frac{k}{4} \left(A \sum_{a=1}^{\ell} m_a^2 \boldsymbol{p}_a - B \sum_{a=1}^{\ell} m_a \boldsymbol{p}_a \right) \\ + \frac{M}{2} \int_{k\Delta} \boldsymbol{x} \, dv - \frac{\widetilde{M}}{4} \int_{\partial(k\Delta)} \boldsymbol{x} \, d\sigma, \quad \text{and}$$

$$(3.10) \qquad \mathcal{DF}_{k\Delta,2} = \frac{B}{12} \sum_{a=1}^{\ell} m_a \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + \frac{1}{6} \sum_{a=1}^{\ell} m_a^3 \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + \frac{k}{2} \sum_{a=1}^{\ell} m_a^2 \boldsymbol{p}_a - \frac{\widetilde{M}}{2} \boldsymbol{s}_{k\Delta} \boldsymbol{x} \, d\sigma.$$

Now we state the main theorem of this paper.

Theorem 3.3. Let $\Delta \subset \mathbb{R}^2$ be a lattice Delzant polygon with the decomposition

$$\Delta = \Delta' \cup \bigcup_{a=1}^{\ell} D_a$$

satisfying the properties described in Section 3.1. We take $k \in \mathbb{N}$ as the minimum integer such that $k\Delta'$ is a lattice Delzant polygon. Let $\varpi : Y'_{\Delta} \dashrightarrow X_{\Delta}$ be the corresponding symplectic blow-up of X_{Δ} at the torus fixed points p_1, \ldots, p_ℓ . For any positive integer *i*, the Chow weight of the toric manifold $\widetilde{Y} := Y_{k\Delta'}$ with respect to $\mathcal{O}_{\widetilde{Y}}(i)$ is given by

(3.11)
$$\operatorname{Chow}_{k\Delta'}(\boldsymbol{x};i) = \operatorname{Chow}_{k\Delta}(\boldsymbol{x};i) + i\mathcal{DF}_{k\Delta,1} + \mathcal{DF}_{k\Delta,2},$$

where $\mathcal{DF}_{k\Delta,1}$ and $\mathcal{DF}_{k\Delta,2}$ are invariants defined in (3.9) and (3.10) respectively.

Proof. Plugging each value in (3.8) into Proposition 3.1, we find that (3.12)

$$\begin{aligned} \operatorname{Chow}_{k\Delta'}(\boldsymbol{x};i) &= \operatorname{Chow}_{k\Delta}(\boldsymbol{x};i) - \left(\sum_{a=1}^{\ell} \frac{m_a^2}{2}\right) \boldsymbol{s}_{k\Delta}(i) \\ &- \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \frac{m_b^2}{2}\right) \sum_{a=1}^{\ell} \left(\frac{m_a}{6}(im_a+1)(im_a-1)\left(\frac{\alpha_a+\gamma_a}{\beta_a+\delta_a}\right) + \frac{im_a}{2}(im_a+1)k\boldsymbol{p}_a\right) \\ &+ E_{k\Delta}(i) \sum_{a=1}^{\ell} \left\{\frac{m_a^3}{6}\left(\frac{\alpha_a+\gamma_a}{\beta_a+\delta_a}\right) + \frac{m_a^2}{2}k\boldsymbol{p}_a\right\} \\ &+ \sum_{b=1}^{\ell} \left(\frac{im_b}{2}(im_b+1)\right) \left(\int_{k\Delta} \boldsymbol{x} \, dv - \sum_{a=1}^{\ell} \left\{\frac{m_a^3}{6}\left(\frac{\alpha_a+\gamma_a}{\beta_a+\delta_a}\right) + \frac{m_a^2}{2}k\boldsymbol{p}_a\right\}\right). \end{aligned}$$

In (3.12), since the last four terms can be simplified as $c_1i + c_0$ with appropriate coefficients c_0 and c_1 with respect to *i* (see, Remark 2.2 (1)). Hence, we further proceed our computations.

• The coefficient of
$$\begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix}$$
:

$$\sum_{a=1}^{\ell} \left[-\frac{1}{6} \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \frac{m_b^2}{2} \right) m_a(im_a + 1)(im_a - 1) + \frac{E_{k\Delta}(i)m_a^3}{6} - \frac{m_a^3}{12} \sum_{b=1}^{\ell} im_b(im_b + 1) \right] \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix}$$

$$= \sum_{a=1}^{\ell} \left[-\frac{1}{6} \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \frac{m_b^2}{2} \right) (m_a^3 i^2 - m_a) + \left(\operatorname{Vol}(k\Delta) i^2 + \frac{\#(\partial(k\Delta) \cap \mathbb{Z}^2)}{2} i + 1 \right) \frac{m_a^3}{6} - \frac{m_a^3}{12} \left(i^2 \sum_{b=1}^{\ell} m_b^2 + i \sum_{b=1}^{\ell} m_b \right) \right] \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix}$$

$$= \sum_{a=1}^{\ell} \left[\frac{m_a^3}{12} \left(\#(\partial(k\Delta) \cap \mathbb{Z}^2) - \sum_{b=1}^{\ell} m_b \right) i + \frac{m_a}{12} \left(2\operatorname{Vol}(k\Delta) + 2m_a^2 - \sum_{b=1}^{\ell} m_b^2 \right) \right] \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix}.$$

• The coefficient of $k p_a$:

$$\sum_{a=1}^{\ell} \left[-\frac{1}{2} \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \frac{m_b^2}{2} \right) im_a(im_a + 1) + E_{k\Delta}(i) \frac{m_a^2}{2} - \frac{1}{4} \sum_{b=1}^{\ell} im_b(im_b + 1)m_a^2 \right] k \boldsymbol{p}_a$$
$$= \sum_{a=1}^{\ell} \left[-\frac{1}{2} \left(\operatorname{Vol}(k\Delta) - \sum_{b=1}^{\ell} \frac{m_b^2}{2} \right) (m_a^2 i^2 + m_a i) + \frac{1}{2} \left(\operatorname{Vol}(k\Delta) i^2 + \frac{\#(\partial(k\Delta) \cap \mathbb{Z}^2)}{2} i + 1 \right) m_a^2 - \frac{m_a^2}{4} \left(i^2 \sum_{b=1}^{\ell} m_b^2 + i \sum_{b=1}^{\ell} m_b \right) \right] k \boldsymbol{p}_a$$

$$=\sum_{a=1}^{\ell} \left[\frac{m_a}{4} \left(-2\operatorname{Vol}(k\Delta) + \sum_{b=1}^{\ell} m_b^2 + \#(\partial(k\Delta) \cap \mathbb{Z}^2)m_a - m_a \sum_{b=1}^{\ell} m_b \right) i + \frac{m_a^2}{2} \right] k \boldsymbol{p}_a.$$

• The remaining terms:

$$-\frac{1}{2}\left(\sum_{a=1}^{\ell}m_{a}^{2}\right)\boldsymbol{s}_{k\Delta}(i)+\frac{1}{2}\sum_{a=1}^{\ell}(m_{a}^{2}i^{2}+m_{a}i)\int_{k\Delta}\boldsymbol{x}\,dv$$
$$=\frac{1}{2}\sum_{a=1}^{\ell}\left\{-m_{a}^{2}\left(i^{2}\int_{k\Delta}\boldsymbol{x}\,dv+\frac{i}{2}\int_{\partial(k\Delta)}\boldsymbol{x}\,d\sigma+\boldsymbol{s}_{k\Delta}\right)+\left(m_{a}^{2}i^{2}+m_{a}i\right)\int_{k\Delta}\boldsymbol{x}\,dv\right\}$$
$$=\frac{1}{4}\sum_{a=1}^{\ell}\left\{m_{a}\left(-m_{a}\int_{\partial(k\Delta)}\boldsymbol{x}\,d\sigma+2\int_{k\Delta}\boldsymbol{x}\,dv\right)i-2m_{a}^{2}\boldsymbol{s}_{k\Delta}\right\}.$$

Plugging these values into (3.12), we conclude that

$$\begin{aligned} \operatorname{Chow}_{k\Delta'}(i) &= \operatorname{Chow}_{k\Delta}(i) \\ &+ i \sum_{a=1}^{\ell} \frac{m_a}{4} \left[\frac{m_a^2}{3} \left(\#(\partial(k\Delta) \cap \mathbb{Z}^2) - \sum_{b=1}^{\ell} m_b \right) \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} \right. \\ &+ \left(-2\operatorname{Vol}(k\Delta) + \sum_{b=1}^{\ell} m_b^2 + \#(\partial(k\Delta) \cap \mathbb{Z}^2)m_a - m_a \sum_{b=1}^{\ell} m_b \right) k \boldsymbol{p}_a \\ &- m_a \int_{\partial(k\Delta)} \boldsymbol{x} \, d\sigma + 2 \int_{k\Delta} \boldsymbol{x} \, dv \right] \\ &+ \sum_{a=1}^{\ell} \frac{m_a}{12} \left[\left(2\operatorname{Vol}(k\Delta) + 2m_a^2 - \sum_{b=1}^{\ell} m_b^2 \right) \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} + 6m_a (k \boldsymbol{p}_a - \boldsymbol{s}_{k\Delta}) \right] \\ &= \operatorname{Chow}_{k\Delta}(i) + i \, \mathcal{DF}_{k\Delta,1} + \mathcal{DF}_{k\Delta,2}. \end{aligned}$$

Hence, the assertion is verified.

4. APPLICATION TO TORIC SURFACES

For any lattice Delzant polygon $\Delta \subset \mathbb{R}^2$, we observe that

(4.1)
$$E_{\Delta}(i) = \operatorname{Vol}(\Delta)i^{2} + (E_{\Delta}(1) - \operatorname{Vol}(\Delta) - 1)i + 1, \text{ and}$$
$$\mathbf{s}_{\Delta}(i) = i^{2} \int_{\Delta} \mathbf{x} \, dv + i \left(\mathbf{s}_{\Delta}(2) - \mathbf{s}_{\Delta}(1) - 3 \int_{\Delta} \mathbf{x} \, dv \right) + 2 \int_{\Delta} \mathbf{x} \, dv - \mathbf{s}_{\Delta}(2) + 2\mathbf{s}_{\Delta}(1).$$

In this section, we apply our blow-up formula (Theorem 3.3) into the projective plane.

4.1. The blow-up of $\mathbb{C}P^2$ at three distinct points. Let X be the blow-up of $\mathbb{C}P^2$ at three torus fixed points. Without loss of generality, we may assume that these three points are $p_1 = [1 : 0 : 0]$, $p_2 = [0 : 1 : 0]$ and $p_3 = [0 : 0 : 1]$. It is well-known that $(X, \mathcal{O}_X(-K_X))$ is a smooth symmetric Kähler-Einstein toric del Pezzo surface with the vanishing Futaki invariant. In particular, $(X, \mathcal{O}_X(-K_X))$ is asymptotically Chow polystable by our result in [LY24, Corollary 4.3]. See also, [Lee22, Example 9.2] for this example. We can also confirm Chow polystability of this example by direct computation with the blow-up formula as follows.

Let

(4.2)

$$\Delta = \operatorname{Conv}\left\{ \begin{pmatrix} 0\\0 \end{pmatrix}, \begin{pmatrix} 3\\0 \end{pmatrix}, \begin{pmatrix} 0\\3 \end{pmatrix} \right\}, \quad \Delta' = \operatorname{Conv}\left\{ \begin{pmatrix} 1\\0 \end{pmatrix}, \begin{pmatrix} 2\\0 \end{pmatrix}, \begin{pmatrix} 2\\1 \end{pmatrix}, \begin{pmatrix} 1\\2 \end{pmatrix}, \begin{pmatrix} 0\\2 \end{pmatrix}, \begin{pmatrix} 0\\1 \end{pmatrix} \right\},$$
$$D_1 = \operatorname{Conv}\left\{ \begin{pmatrix} 0\\0 \end{pmatrix}, \begin{pmatrix} 1\\0 \end{pmatrix}, \begin{pmatrix} 0\\1 \end{pmatrix} \right\}, \quad D_2 = \operatorname{Conv}\left\{ \begin{pmatrix} 3\\0 \end{pmatrix}, \begin{pmatrix} 2\\1 \end{pmatrix}, \begin{pmatrix} 2\\0 \end{pmatrix} \right\}, \quad \text{and}$$
$$D_3 = \operatorname{Conv}\left\{ \begin{pmatrix} 0\\3 \end{pmatrix}, \begin{pmatrix} 0\\2 \end{pmatrix}, \begin{pmatrix} 1\\2 \end{pmatrix} \right\}.$$

Then, Δ can be decomposed into

$$\Delta = \Delta' \cup \bigcup_{a=1}^3 D_a$$

with $p_1 = (0,0)$, $p_2 = (3,0)$ and $p_3 = (0,3)$. Since Δ' is a lattice polygon, we take k = 1. We readily see that $m_a = 1$ for each a = 1, 2, 3, and the matrices defined in (3.4) are given by

$$A_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} \text{ and } A_3 = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}.$$

Therefore,

$$M = \sum_{a=1}^{3} m_a = 3, \quad \widetilde{M} = \sum_{a=1}^{3} m_a^2 = 3, \text{ and}$$
$$A = \#(\partial \Delta \cap \mathbb{Z}^2) - M = 6, \qquad B = 2\text{vol}(\Delta) - \widetilde{M} = 6.$$

Since

$$\int_{\Delta} \boldsymbol{x} \, dv = \left(\frac{9}{2}, \frac{9}{2}\right)$$

by direct computation, this yields that

$$\boldsymbol{s}_{\Delta}(i) = \frac{9}{2} \begin{pmatrix} 1\\1 \end{pmatrix} i^2 + \frac{9}{2} \begin{pmatrix} 1\\1 \end{pmatrix} i + \begin{pmatrix} 1\\1 \end{pmatrix}$$

from (4.1). Plugging these values into (3.9) and (3.10), we conclude that

$$\mathcal{DF}_{\Delta,1} = \frac{27}{4} \left\{ \begin{pmatrix} 1\\1 \end{pmatrix} - \begin{pmatrix} 1\\1 \end{pmatrix} \right\} = \begin{pmatrix} 0\\0 \end{pmatrix}, \quad \text{and} \\ \mathcal{DF}_{\Delta,2} = \frac{1}{2} \sum_{a=1}^{3} p_a - \frac{3}{2} \boldsymbol{s}_{\Delta} = \frac{1}{2} \left\{ \begin{pmatrix} 0\\0 \end{pmatrix} + \begin{pmatrix} 3\\0 \end{pmatrix} + \begin{pmatrix} 0\\3 \end{pmatrix} \right\} - \frac{3}{2} \begin{pmatrix} 1\\1 \end{pmatrix} = \begin{pmatrix} 0\\0 \end{pmatrix}.$$

Hence,

$$\operatorname{Chow}_{\Delta'}(\boldsymbol{x};i) = \operatorname{Chow}_{\Delta}(\boldsymbol{x};i) + i \begin{pmatrix} 0\\ 0 \end{pmatrix} + \begin{pmatrix} 0\\ 0 \end{pmatrix} = \begin{pmatrix} 0\\ 0 \end{pmatrix}$$

by (3.11) which is consistent with our result in [LY24, Corollary 4.3].

4.2. The blow-up of $\mathbb{C}P^2$ at four distinct points. Now we consider the blow-up of $\mathbb{C}P^2$ at four points. In order to verify Chow stability of points in $\mathbb{C}P^2$ (or more generally, points in $\mathbb{C}P^n$), there is a nice geometric criterion as in [Muk03, Proposition 7.27].

Proposition 4.1 (Mukai). Let $\Lambda := \{p_1, \ldots, p_\ell\}$ be a set of points in $\mathbb{C}P^2$. Then the blow-up manifold $\widetilde{X} := Bl_{\Lambda}(\mathbb{C}P^2) \dashrightarrow \mathbb{C}P^2$ is:

(1) Chow stable if and only if

$$\frac{\#(\Lambda \cap V)}{\ell} < \frac{\dim V + 1}{3}$$

holds for any projective subspace $V \subsetneq \mathbb{C}P^n$. (2) Chow unstable if and only if there is some projective subspace $V \subsetneq \mathbb{C}P^n$ such that

$$\frac{\#(\Lambda \cap V)}{\ell} > \frac{\dim V + 1}{3}.$$

Proof. See, p. 232 in [Muk03].

Therefore, if we take distinct four points $\Lambda = \{p_1, p_2, p_3, p_4\} \subset \mathbb{C}P^2$ in general position which means that no three of which are collinear, we see that $Y := Bl_{\Lambda}(\mathbb{C}P^2)$ (this is the del Pezzo surface of degree five) is Chow stable as follows: for any projective line $V \cong \mathbb{C}P^1$ in $\mathbb{C}P^2$, we observe that V contains exactly two distinct points p_i and p_j in Λ . Thus, we find that

$$\frac{\#(\Lambda \cap V)}{\ell} = \frac{2}{4} < \frac{2}{3} = \frac{\dim V + 1}{3},$$

which yields that $(Y, \mathcal{O}_Y(1))$ is Chow stable by Proposition 4.1 (1).

Meanwhile, if we consider the *toric blow-up* for obtaining the blow-up manifold $Bl_{(4pts)}(\mathbb{C}P^2)$, we have to take the set of points $\Lambda = \{p_1, p_2, p_3, p_4\}$ so that p_1, p_3 and p_4 lie in the same projective line $\mathbb{C}P^1 \subset \mathbb{C}P^2$ as illustrated in Figure 1. More specifically, let $\Delta' \subset M_{\mathbb{R}}$ be the moment polytope in (4.2) associated with the toric surface $X := Bl_{(3pts)}(\mathbb{C}P^2)$ constructed in Section 4.1. We take the point p_4 in X corresponding to the vertex $p_4 = (0, 2)$ of Δ' . Let Z be the blow-up of X at p_4 which is given by a sequence of blow-ups of $\mathbb{C}P^2$ such as

$$Z := \operatorname{Bl}_{(p_4)}(X) \dashrightarrow X \dashrightarrow \mathbb{C}P^2.$$

As drawn in Figure 1, the corresponding polytope Δ'' of the toric manifold Z is obtained by chopping off the vertex p_4 of Δ' and we shall consider the decomposition $\Delta' = \Delta'' \cup D_1$ with

$$\Delta'' = \operatorname{Conv}\left\{ \begin{pmatrix} 1\\0 \end{pmatrix}, \begin{pmatrix} 2\\0 \end{pmatrix}, \begin{pmatrix} 2\\1 \end{pmatrix}, \begin{pmatrix} 1\\2 \end{pmatrix}, \begin{pmatrix} 1/2\\2 \end{pmatrix}, \begin{pmatrix} 0\\3/2 \end{pmatrix}, \begin{pmatrix} 0\\1 \end{pmatrix} \right\} \quad \text{and} \\ D_1 = \operatorname{Conv}\left\{ \begin{pmatrix} 0\\2 \end{pmatrix}, \begin{pmatrix} 0\\3/2 \end{pmatrix}, \begin{pmatrix} 1/2\\2 \end{pmatrix} \right\}.$$



FIGURE 1. The decomposition of the polygon.

In order to make $k\Delta''$ a lattice polygon, we need to take k = 2. Then, the matrix defined in (3.4) is $A_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Moreover, we see that $M = m_1 = 1, \quad \widetilde{M} = m_1^2 = 1, \quad E_{\Delta'}(i) = 3i^2 + 3i + 1,$ $A = \#(\partial(2\Delta') \cap \mathbb{Z}^2) - M = 11, \qquad B = 2\text{vol}(2\Delta') - \widetilde{M} = 23,$ (4.3) $\int_{\Delta'} \mathbf{x} \, dv = (3, 3), \qquad \mathbf{s}_{\Delta'}(i) = \begin{pmatrix} 3 \\ 3 \end{pmatrix} i^2 + \begin{pmatrix} 3 \\ 3 \end{pmatrix} i + \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \qquad \text{and}$ $\mathbf{s}_{2\Delta'}(i) = 2\mathbf{s}_{\Delta'}(2i) = \begin{pmatrix} 24 \\ 24 \end{pmatrix} i^2 + \begin{pmatrix} 12 \\ 12 \end{pmatrix} i + \begin{pmatrix} 2 \\ 2 \end{pmatrix}.$

Plugging these values into (3.9) and (3.10), we find that

$$\begin{aligned} \mathcal{DF}_{2\Delta',1} &= \frac{1}{12} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \frac{2}{4} \left\{ 11 \begin{pmatrix} 0 \\ 2 \end{pmatrix} - 23 \begin{pmatrix} 0 \\ 2 \end{pmatrix} \right\} + \frac{1}{2} \int_{2\Delta'} \boldsymbol{x} \, d\boldsymbol{v} - \frac{1}{4} \int_{\partial(2\Delta')} \boldsymbol{x} \, d\sigma \\ &= \frac{83}{12} \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \text{and} \\ \mathcal{DF}_{2\Delta',2} &= \frac{23}{12} \begin{pmatrix} \alpha_1 + \gamma_1 \\ \beta_1 + \delta_1 \end{pmatrix} + \frac{1}{6} \begin{pmatrix} \alpha_1 + \gamma_1 \\ \beta_1 + \delta_1 \end{pmatrix} + \frac{2}{2} \boldsymbol{p}_4 - \frac{1}{2} \boldsymbol{s}_{2\Delta'} \\ &= \frac{23}{12} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \frac{1}{6} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \begin{pmatrix} 0 \\ 2 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \frac{13}{12} \begin{pmatrix} 1 \\ -1 \end{pmatrix}. \end{aligned}$$

This yields that

(4.4)
$$\operatorname{Chow}_{2\Delta''}(\boldsymbol{x};i) = \operatorname{Chow}_{2\Delta'}(\boldsymbol{x};i) + \frac{83}{12} \begin{pmatrix} 1\\-1 \end{pmatrix} i + \frac{13}{12} \begin{pmatrix} 1\\-1 \end{pmatrix}$$

be (3.11). Consequently, $(Z, \mathcal{O}_Z(1))$ is Chow *unstable*. Note that the coefficient of i in (4.4) and the constant term are proportional to each other as $\operatorname{Chow}_{2\Delta'}(\boldsymbol{x}; i) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$. In other word, we find that

$$\dim \operatorname{Span}_{\mathbb{R}} \left\{ \operatorname{Coeff} \left(\operatorname{Chow}_{2\Delta''}(\boldsymbol{x}; i), i \right), \operatorname{Coeff} \left(\operatorname{Chow}_{2\Delta''}(\boldsymbol{x}; i), \operatorname{const} \right) \right\} = 1$$

This is explained as follows: let X_{Σ} be an *n*-dimensional smooth projective toric variety with the associated complete fan Σ in N. Let $\mathcal{W}(X_{\Sigma})$ be the Weyl group of X_{Σ} . By [BS99, Proposition 3.1], we have

$$\mathcal{W}(X_{\Sigma}) \cong G = \{ \gamma \in \mathrm{GL}(n, \mathbb{Z}) \mid \gamma(\Sigma) = \Sigma \}.$$

In our case, we see that Δ'' is invariant under unimodular transformation $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, which gives the coordinate exchange $x_1 \mapsto -x_1$ and $x_2 \mapsto -x_2$ in $M_{\mathbb{R}} \cong \mathbb{R}^2$. By this symmetry, we conclude that

(4.5)
$$\gamma = \begin{pmatrix} -1 & 0\\ 0 & -1 \end{pmatrix} \in \mathcal{W}(Z)$$

for the associated toric surface $Z = X_{\Sigma''}$, where Σ'' is the normal fan of Δ'' . Let $N_{\mathbb{R}}^{\mathcal{W}(Z)}$ be the Weyl group invariant subspace of $N_{\mathbb{R}}$. Then, (4.5) implies $\dim N_{\mathbb{R}}^{\mathcal{W}(Z)} = 2 - 1 = 1$. Since the coefficients of $\operatorname{Chow}_{2\Delta''}(\boldsymbol{x}; i)$ are invariant under $\mathcal{W}(Z)$ -action, we observe that

$$\dim \operatorname{Span}_{\mathbb{R}} \left\{ \operatorname{Coeff} \left(\operatorname{Chow}_{2\Delta''}(\boldsymbol{x}; i) \right) \right\} = \dim N_{\mathbb{R}}^{\mathcal{W}(Z)} = 1$$

4.3. The blow-up of $\mathbb{C}P^2$ at five and six distinct points. Let Z be the toric surface constructed in Section 4.2. We further take the point p_5 in Z corresponding to the vertex $p_5 = (2,0)$ of Δ'' . Let Z_1 be the blow-up of Z at p_5 given by

(4.6)
$$Z_1 := \operatorname{Bl}_{(p_5)}(Z) \dashrightarrow Z \dashrightarrow X \dashrightarrow \mathbb{C}P^2.$$

In this case, if we consider $\Lambda = \{p_1, \ldots, p_5\}$ and take the blow-up $\widetilde{Z} := Bl_{\Lambda}(\mathbb{C}P^2) \dashrightarrow \mathbb{C}P^2$ simultaneously, then we see that there are three projective lines V_i (i = 1, 2, 3) in $\mathbb{C}P^2$ satisfying

$$\{p_1, p_2, p_5\} \subset V_1, \{p_2, p_3\} \subset V_2 \text{ and } \{p_1, p_3, p_4\} \subset V_3.$$

Thus, we have

•
$$\frac{\#(\Lambda \cap V_i)}{\#\Lambda} = \frac{3}{5} < \frac{2}{3}$$
 for $i = 1, 3$, and
• $\frac{\#(\Lambda \cap V_2)}{\#\Lambda} = \frac{2}{5} < \frac{2}{3}$,

which implies that $(\widetilde{Z}, \mathcal{O}_{\widetilde{Z}}(1))$ is Chow polystable by Proposition 4.1.

Let us observe this by using our blow-up formula. The corresponding decompositions of the polygon with respect to (4.6) are

(4.7)
$$\Delta' = \Delta_1 \cup \bigcup_{a=1}^2 D_a = \Delta'' \cup D_1,$$

where Δ_1 and D_2 are convex polygons defined by

$$\Delta_{1} = \operatorname{Conv}\left\{ \begin{pmatrix} 1\\0 \end{pmatrix}, \begin{pmatrix} 3/2\\0 \end{pmatrix}, \begin{pmatrix} 2\\1/2 \end{pmatrix}, \begin{pmatrix} 2\\1 \end{pmatrix}, \begin{pmatrix} 1\\2 \end{pmatrix}, \begin{pmatrix} 1/2\\2 \end{pmatrix}, \begin{pmatrix} 0\\3/2 \end{pmatrix}, \begin{pmatrix} 0\\1 \end{pmatrix} \right\} \quad \text{and} \\ D_{2} = \operatorname{Conv}\left\{ \begin{pmatrix} 2\\0 \end{pmatrix}, \begin{pmatrix} 2\\1/2 \end{pmatrix}, \begin{pmatrix} 3/2\\0 \end{pmatrix} \right\}.$$

Then k = 2 and the matrix defined in (3.4) is $A_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Similar to the computations in (4.3), we find that

$$M = \sum_{j=1}^{2} m_j = 2, \qquad \widetilde{M} = \sum_{j=1}^{2} m_j^2 = 2,$$
$$A = \# \left(\partial (2\Delta') \cap \mathbb{Z}^2 \right) - M = 10, \qquad B = 2 \operatorname{Vol}(2\Delta') - \widetilde{M} = 22$$

through the decomposition (4.7). Thus, the Chow weight of $(Z_1, \mathcal{O}_{Z_1}(1))$ equals

$$\operatorname{Chow}_{\Delta_1}(\boldsymbol{x};i) = \operatorname{Chow}_{\Delta'}(\boldsymbol{x};i) + i\mathcal{DF}_{2\Delta',1} + \mathcal{DF}_{2\Delta',2}$$

by (3.11), where each $\mathcal{DF}_{2\Delta',j}$ is given by

$$\mathcal{DF}_{2\Delta',1} = \frac{10}{12} \sum_{j=1}^{2} \binom{\alpha_{j} + \gamma_{j}}{\beta_{j} + \delta_{j}} + \frac{2}{4} \left(10 \sum_{\ell=4}^{5} \boldsymbol{p}_{\ell} - 22 \sum_{\ell=4}^{5} \boldsymbol{p}_{\ell} \right) + \frac{2}{2} \binom{24}{24} - \frac{2}{4} \binom{24}{24} = \binom{0}{0} \quad \text{and} \\ \mathcal{DF}_{2\Delta',2} = \frac{22}{12} \sum_{j=1}^{2} \binom{\alpha_{j} + \gamma_{j}}{\beta_{j} + \delta_{j}} + \frac{1}{6} \sum_{j=1}^{2} \binom{\alpha_{j} + \gamma_{j}}{\beta_{j} + \delta_{j}} + \frac{2}{2} \sum_{\ell=4}^{5} \boldsymbol{p}_{\ell} - \frac{2}{2} \boldsymbol{s}_{2\Delta'} \\ = \sum_{\ell=4}^{5} \boldsymbol{p}_{\ell} - \boldsymbol{s}_{2\Delta'} = \left\{ \binom{0}{2} + \binom{2}{0} \right\} - \binom{2}{2} = \binom{0}{0}$$

respectively. Since $\operatorname{Chow}_{\Delta_1}(\boldsymbol{x}; i)$ vanishes identically, we further conclude that $(Z_1, \mathcal{O}_{Z_1}(1))$ is asymptotically Chow polystable (see, [LY24, Remark 4.4]).

Finally, we take p_6 in Z_1 , where p_6 corresponds to the vertex $p_6 = (1, 2)$ of Δ_1 . The induced toric blow-up of Z_1 at p_6 is

(4.8)
$$Z_2 := \operatorname{Bl}_{(p_6)}(Z_1) \dashrightarrow Z_1 \dashrightarrow \mathbb{C}P^2.$$

Let $\Delta_1 = \Delta_2 \cup D_3$ be the decomposition of the polygon corresponds to (4.8), where Δ_1 and Δ_2 are convex polygons defined by

$$\Delta_{2} = \operatorname{Conv}\left\{ \begin{pmatrix} 1\\0 \end{pmatrix}, \begin{pmatrix} 3/2\\0 \end{pmatrix}, \begin{pmatrix} 2\\1/2 \end{pmatrix}, \begin{pmatrix} 2\\1 \end{pmatrix}, \begin{pmatrix} 5/4\\7/4 \end{pmatrix}, \begin{pmatrix} 3/4\\2 \end{pmatrix}, \begin{pmatrix} 1/2\\2 \end{pmatrix}, \begin{pmatrix} 0\\3/2 \end{pmatrix}, \begin{pmatrix} 0\\1 \end{pmatrix} \right\} \quad \text{and} \\ D_{3} = \operatorname{Conv}\left\{ \begin{pmatrix} 1\\2 \end{pmatrix}, \begin{pmatrix} 3/4\\2 \end{pmatrix}, \begin{pmatrix} 5/4\\2 \end{pmatrix}, \begin{pmatrix} 5/4\\7/4 \end{pmatrix} \right\}$$

respectively (see Figure 2). Thus the minimum integer k such that $k\Delta_2$ will be a lattice polygon is four. Furthermore, the matrix in (3.4) is $A_3 = \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}$ and

$$M = m_3 = 1, \qquad \widetilde{M} = m_3^2 = 1, \qquad A = \# (\partial (4\Delta_1) \cap \mathbb{Z}^2) - M = 20 - 1 = 19 \qquad \text{and} \\ B = 2 \text{Vol}(4\Delta_1) - \widetilde{M} = 2 \cdot 44 - 1 = 87.$$



FIGURE 2. The blow-up of $\mathbb{C}P^2$ at the torus invariant six points.

The straightforward computation shows that

$$E_{4\Delta_2}(i) = 44i^2 + 10i + 1, \quad \int_{4\Delta_2} \boldsymbol{x} \, d\boldsymbol{v} = \begin{pmatrix} 176\\176 \end{pmatrix}, \quad \boldsymbol{s}_{4\Delta_2}(1) = \begin{pmatrix} 220\\220 \end{pmatrix}, \quad \boldsymbol{s}_{4\Delta_2}(2) = \begin{pmatrix} 1576\\1576 \end{pmatrix}$$

and we find

$$\mathbf{s}_{4\Delta_2}(i) = 176 \begin{pmatrix} 1\\1 \end{pmatrix} i^2 + 828 \begin{pmatrix} 1\\1 \end{pmatrix} i - 784176 \begin{pmatrix} 1\\1 \end{pmatrix}$$

by (4.1). Thus we can calculate $\mathcal{DF}_{4\Delta_2,j}$ as

$$\mathcal{DF}_{4\Delta_{2},1} = \frac{19}{12} \begin{pmatrix} \alpha_{3} + \gamma_{3} \\ \beta_{3} + \delta_{3} \end{pmatrix} + \frac{4}{4} (19\boldsymbol{p}_{6} - 87\boldsymbol{p}_{6}) + \frac{1}{2} \begin{pmatrix} 176 \\ 176 \end{pmatrix} - \frac{1}{4} \begin{pmatrix} 1656 \\ 1656 \end{pmatrix}$$
$$= \frac{19}{12} \begin{pmatrix} 0 \\ -1 \end{pmatrix} - 68 \begin{pmatrix} 1 \\ 2 \end{pmatrix} + (88 - 414) \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -394 \\ -\frac{4747}{12} \end{pmatrix} \quad \text{and}$$
$$\mathcal{DF}_{4\Delta_{2},2} = \frac{87}{12} \begin{pmatrix} 0 \\ -1 \end{pmatrix} + \frac{1}{6} \begin{pmatrix} 0 \\ -1 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 2 \end{pmatrix} - 392 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -390 \\ -\frac{4745}{12} \end{pmatrix}.$$

As a consequence, we have dim $\operatorname{Span}_{\mathbb{R}} \{ \operatorname{Coeff}(\operatorname{Chow}_{4\Delta_2}(\boldsymbol{x}; i)) \} = 2$ and

which implies that $(X_{\Delta_2}, \mathcal{O}_{X_{\Delta_2}}(4))$ is Chow unstable.

However, if we take $\Lambda = \{p_1, p_2, \dots, p_6\} \subset \mathbb{C}P^2$ to be a set of points in general position, then

 $\widetilde{X} := \mathrm{Bl}_{\Lambda}(\mathbb{C}P^2) \dashrightarrow \mathbb{C}P^2$

is the del Pezzo surface of degree three. In particular, \widetilde{X} admits a Kähler-Einstein metric in its first Chern class due to a result of Tian (see, [Tia90], [Tos12, Theorem 1.4]). On the other hand, it is known that the automorphism groups $\mathfrak{h}(\widetilde{X})$ of the blow-up of $\mathbb{C}P^2$ ad four or more general points satisfy $\mathfrak{h}(\widetilde{X}) = \{0\}$. Then, Donaldson's theorem in [Don01] implies that $(\widetilde{X}, -K_{\widetilde{X}})$ is asymptotically Chow stable. Thus, our computation in (4.9) shows the significant difference of Chow weight under the toric blow-ups and the blow-ups at general points.

4.4. The blow-up at a point in the Hirzebruch surfaces. We finish this section with the following observation. If the polygon $\Delta \subset \mathbb{R}^2$ is a rectangle, then we have $\operatorname{Chow}_{\Delta}(\boldsymbol{x}; i) = 0$. Thus, without loss of generality we may assume that $\Delta \subset \mathbb{R}^2$ with $\#\mathcal{V}(\Delta) = 4$ has vertices

$$\boldsymbol{p}_1 = \begin{pmatrix} 0\\ 0 \end{pmatrix}, \quad \boldsymbol{p}_2 = \begin{pmatrix} 0\\ a \end{pmatrix}, \quad \boldsymbol{p}_3 = \begin{pmatrix} b\\ a \end{pmatrix}, \quad \boldsymbol{p}_4 = \begin{pmatrix} b+an\\ 0 \end{pmatrix},$$

where a, b and n are positive integers. Let $\Delta_n(a, b) = \operatorname{conv} \{ \mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4 \}$. We remark that every 2-dimensional Delzant polytope with four vertices except for rectangles can be written in the form of $\Delta_n(a, b)$ up to parallel transformations and $SL_2(\mathbb{Z})$ -transformations. Firstly, we calculate the following quantity for later use.

Proposition 4.2.

$$\begin{aligned} \operatorname{Vol}(\Delta_n(a,b)) &= ab + \frac{1}{2}a^2n, \\ E_{\Delta_n(a,b)}(i) &= \left(ab + \frac{1}{2}a^2n\right)i^2 + \left(a + b + \frac{1}{2}an\right)i + 1, \\ \int_{\Delta_n(a,b)} \boldsymbol{x} \, dv &= \frac{a}{6} \begin{pmatrix} a^2n^2 + 3abn + 3b^2 \\ a(an + 3b) \end{pmatrix}, \\ \boldsymbol{s}_{\Delta_n(a,b)}(i) &= \frac{a}{6} \begin{pmatrix} a^2n^2 + 3abn + 3b^2 \\ a(an + 3b) \end{pmatrix}i^2 + \frac{1}{4} \begin{pmatrix} a^2n^2 + a^n + 2abn + 2ab + 2b^2 \\ 2a(a + b) \end{pmatrix}i \\ &+ \frac{1}{12} \begin{pmatrix} an^2 + 3an + 6b \\ 2a(3 - n) \end{pmatrix}. \end{aligned}$$

Proof. All calculations are straightforward and we leave them to the reader.

Plugging each value in Proposition 4.2 into (2.4), we find

Chow_{$$\Delta_n(a,b)$$} $(i) = \frac{1}{24}a^2n(ai+1)(an-a+2b)\binom{n}{-2}.$

Next, we shall compute the Chow weight of the manifold after taking the blow-ups at the points p_1, p_2, p_3 and p_4 . For these, we see that

$$\begin{pmatrix} \alpha_1 + \gamma_1 \\ \beta_1 + \delta_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} \alpha_2 + \gamma_2 \\ \beta_2 + \delta_2 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \begin{pmatrix} \alpha_3 + \gamma_3 \\ \beta_3 + \delta_3 \end{pmatrix} = \begin{pmatrix} n-1 \\ -1 \end{pmatrix}, \quad \begin{pmatrix} \alpha_4 + \gamma_4 \\ \beta_4 + \delta_4 \end{pmatrix} = \begin{pmatrix} -1-n \\ 1 \end{pmatrix},$$

respectively. Thus, we conclude that

$$\sum_{a=1}^{4} m_a^3 \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} = \begin{pmatrix} m_1^3 + m_2^3 + (n-1)m_3^3 - (n+1)m_4^3 \\ m_1^3 - m_2^3 - m_3^3 + m_4^3 \end{pmatrix},$$

$$(4.10) \quad \sum_{a=1}^{4} m_a \begin{pmatrix} \alpha_a + \gamma_a \\ \beta_a + \delta_a \end{pmatrix} = \begin{pmatrix} m_1 + m_2 + (n-1)m_3 - (n+1)m_4 \\ m_1 - m_2 - m_3 + m_4 \end{pmatrix},$$

$$\sum_{a=1}^{4} m_a^2 \boldsymbol{p}_a = \begin{pmatrix} bm_3^2 + (b+an)m_4^2 \\ a(m_2^2 + m_3^2) \end{pmatrix}, \qquad \sum_{a=1}^{4} m_a \boldsymbol{p}_a = \begin{pmatrix} bm_3 + (b+an)m_4 \\ a(m_2 + m_3) \end{pmatrix}.$$

Since $k\Delta_n(a,b) = \Delta_n(ka,kb)$, we conclude that

(4.11)

$$Chow_{k\Delta_n(a,b)}(i) = \frac{k^3}{24}a^2n(aki+1)(an-a+2b)\binom{n}{-2},$$

$$A = k(2a+an+2b) - M, \quad B = ak^2(an+2b) - \widetilde{M},$$

$$\int_{k\Delta_n(a,b)} \boldsymbol{x} \, dv = \frac{ak^3}{6} \binom{a^2n^2+3abn+3b^2}{a(an+3b)},$$

$$\int_{\partial(k\Delta_n(a,b))} \boldsymbol{x} \, d\sigma = \frac{k^2}{2} \binom{a^2n^2+a^2n+2abn+2ab+2b^2}{2a(a+b)},$$

$$\boldsymbol{s}_{k\Delta_n(a,b)} = \frac{k}{12} \binom{an^2+3an+6b}{2a(3-n)}.$$

Plugging (4.10) and (4.11) into (3.9) and (3.10), we can find the explicit value of $\operatorname{Chow}_{k\Delta'}(i)$ by (3.11).

5. Relationship with the vanishing condition of the Futaki-Ono invariant

In this section, we consider the Futaki-Ono (FO) invariant which is an obstruction for a polarized toric variety (X, L) to be asymptotically Chow semistable (see, (5.2)). We firstly see that the FO invariant vanishes identically for some special Kähler class of the toric del Pezzo surface with degree six (Example 5.5). Secondary, we give the formula for computing the FO invariant of the blow-up manifold explicitly (Proposition 5.6).

5.1. An example of toric surface with the vanishing FO invariant.

Definition 5.1. Let Δ be an *n*-dimensional lattice (not necessarily Delzant) polytope in $M_{\mathbb{R}}$. Δ is called *centrally symmetric* if it satisfies $\Delta = -\Delta$. We say that Δ is *weakly symmetric* if there exists a finite group $G < SL(n, \mathbb{Z})$ acting on Δ such that for any $\mathbf{x} \in (x_1, \ldots, x_n) \in \Delta$, we have

(5.1)
$$\sum_{g \in G} g \cdot \boldsymbol{x} = \boldsymbol{0}$$

Example 5.2. Let $\Delta = \operatorname{Conv} \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ -1 \end{pmatrix} \right\}$ and let $G < \operatorname{SL}(2, \mathbb{Z})$ be the finite cyclic group of order three generated by

$$G = \left\langle \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix} \right\rangle \cong \mathbb{Z}_3.$$

Then, we readily see that Δ and G satisfy (5.1). Thus Δ is weakly symmetric.

For a positive integer *i* and an affine function $\ell(\boldsymbol{x}) = \sum_j c_j x_j + c_0$ on $i\Delta$, we recall that the *Futaki-Ono invariant* $FO(\ell; i)$ is defined by

(5.2)
$$FO(\ell;i) := \frac{\sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} \ell(\boldsymbol{a})}{\# \left(\Delta \cap (\mathbb{Z}/i)^n\right)} - \frac{\int_{\Delta} \ell(\boldsymbol{x}) dv}{\operatorname{Vol}(\Delta)}$$

It is known that the associated toric variety (X_{Δ}, L_{Δ}) is asymptotically Chow semistable, then $FO(\ell; i)$ vanishes for any affine function.

Lemma 5.3. Suppose Δ is weakly symmetric, then the FO invariant vanishes.

Proof. Let G be the finite subgroup in $SL(n, \mathbb{Z})$ satisfying (5.1). For any $g \in G$, we observe that

$$\sum_{oldsymbol{a}\in\Delta\cap(\mathbb{Z}/i)^n}g\cdotoldsymbol{a}=\sum_{oldsymbol{a}\in\Delta\cap(\mathbb{Z}/i)^n}oldsymbol{a},$$

and

$$\int_{\Delta} g \cdot \boldsymbol{x} \, dv = \int_{g^* \Delta} \boldsymbol{x} \, dv = \int_{\Delta} \boldsymbol{x} \, dv.$$

Let |G| denote the order of G. Then we have

(5.3)
$$|G| \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} \boldsymbol{a} = \sum_{g \in G} \sum_{\boldsymbol{a} \in \Delta \cap (\mathbb{Z}/i)^n} g \cdot \boldsymbol{a} = \boldsymbol{0}.$$

because Δ is weakly symmetric. Moreover, we also find that

(5.4)
$$|G| \int_{\Delta} \boldsymbol{x} \, dv = \sum_{g \in G} \int_{\Delta} g \cdot \boldsymbol{x} \, dv = \boldsymbol{0}$$

Then we conclude that $FO(\ell; i)$ vanishes by (5.3) and (5.4).

Lemma 5.4. Let G be the finite cyclic group of order two generated by $x \mapsto -x$:

$$G = \left\langle \begin{pmatrix} -1 & & \\ & \ddots & \\ & & -1 \end{pmatrix} \right\rangle \cong \mathbb{Z}_2.$$

Suppose Δ is centrally symmetric, then Δ is weakly symmetric with the finite group $G < SL(n, \mathbb{Z})$. *Proof.* For any $x \in \Delta$, we have $-x \in \Delta$ by assumption. Thus, we see that

$$\sum_{g \in G} g \cdot \boldsymbol{x} = \boldsymbol{x} + g \cdot \boldsymbol{x} = \boldsymbol{x} - \boldsymbol{x} = \boldsymbol{0}.$$

Combining Lemmas 5.3 and 5.4, we can construct an example of toric surface with the vanishing FO invariant.

Example 5.5. We now consider the polarized toric surface (X_{Δ}, L_{Δ}) associated with the following moment polytope (Figure 3).

We readily see that the corresponding toric surface X_{Δ} is $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ (the del Pezzo surface with six), but it is not anticanonically polarized. Δ is centrally symmetric as every vertex $p \in \mathcal{V}(\Delta)$ satisfies $-p \in \mathcal{V}(\Delta)$. Thus $FO(\ell; i) \equiv 0$ by Lemmas 5.3 and 5.4.



FIGURE 3. A centrally symmetric polygon with six vertices.

5.2. The blow-up formula of the FO invariant. Finally, we compute the FO invariant of the blow-up manifold $\varpi : X_{\Delta'} \dashrightarrow X_{\Delta}$ assuming that all of the FO invariants of the base toric manifold X_{Δ} vanish.

In this section, we use the following terminology. Let S_n be the *n*-dimensional standard simplex and let $\ell(\mathbf{x}) = \sum_{i=1}^{n} c_i x_i + c_0$ be any affine function on S_n . Then we have

(5.5)
$$\sum_{\boldsymbol{a}\in S_n\cap\mathbb{Z}^n}\ell(\boldsymbol{a})=c_{S_n}\int_{S_n}\ell(\boldsymbol{x})dv,$$

where c_{S_n} is the constant defined by

(5.6)
$$c_{S_n} = \frac{\#(S_n \cap \mathbb{Z}^n)}{\operatorname{Vol}(S_n)} = \frac{n+1}{1/n!} = (n+1)!.$$

Let Δ be a lattice Delzant polytope in $M_{\mathbb{R}}$. We now consider a simultaneous blow-up of Δ at torus fixed k points p_1, \ldots, p_k . As we already saw in Section 3.1, this corresponds to remove the simplices D_1, \ldots, D_k , where Δ can be decomposed into

$$\Delta = \Delta' \cup \bigcup_{a=1}^{k} D_a.$$

Let p_1, \ldots, p_k with $p_a \in \mathcal{V}(D_a)$ be the vertices of Δ being blowing up. For any affine function $\ell(bsx)$ on Δ , we see that

(5.8)
$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}\ell(\boldsymbol{a})=\sum_{\boldsymbol{a}\in\Delta\cap\mathbb{Z}^n}\ell(\boldsymbol{a})-\sum_{a=1}^{\kappa}\ell(\boldsymbol{p}_a)$$

by (5.7). Furthermore, let F_a denote the only face of D_a so that $p_a \notin F_a$. Then we have

(5.9)
$$\ell(\boldsymbol{p}_a) = \sum_{\boldsymbol{a} \in D_a \cap \mathbb{Z}^n} \ell(\boldsymbol{a}) - \sum_{\boldsymbol{a} \in F_a \cap \mathbb{Z}^n} \ell(\boldsymbol{a})$$

By (5.8) and (5.9), we find the equality

(5.10)
$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}\ell(\boldsymbol{a})=\sum_{\boldsymbol{a}\in\Delta\cap\mathbb{Z}^n}\ell(\boldsymbol{a})-\sum_{a=1}^k\sum_{\boldsymbol{a}\in D_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a})+\sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a}).$$

Combining (5.10) with (5.5), we have the following.

Proposition 5.6. Suppose Δ has the decomposition (5.7). Let c_{Δ} and $c_{\Delta'}$ be the constants defined in (5.6) with respect to Δ and Δ' . If all the FO invariants of Δ vanish, then we have

(5.11)
$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) = c_{\Delta'}\int_{\Delta'}\ell(\boldsymbol{x})dv + (c_{\Delta} - c_{\Delta'})\int_{\Delta}\ell(\boldsymbol{x})dv + \left(c_{\Delta'} - (n+1)!\right)\sum_{a=1}^k\int_{D_a}\ell(\boldsymbol{x})dv + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_i\cap\mathbb{Z}^n}\ell(\boldsymbol{a}).$$

Proof. By (5.10) and (5.5), we see that

$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) = \sum_{\boldsymbol{a}\in\Delta\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) - \sum_{a=1}^k\sum_{\boldsymbol{a}\in D_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a})$$
$$= \sum_{\boldsymbol{a}\in\Delta\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) - \sum_{a=1}^k(n+1)!\int_{D_a}\ell(\boldsymbol{x})dv + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a})$$
$$= c_{\Delta}\int_{\Delta}\ell(\boldsymbol{x})dv - \sum_{a=1}^k(n+1)!\int_{D_a}\ell(\boldsymbol{x})dv + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a}),$$

,

where in the last equality, we used the assumption that the FO invariants vanish. Arranging these coefficients, we conclude that

$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) = (c_{\Delta} - c_{\Delta'})\int_{\Delta}\ell(\boldsymbol{x})dv + c_{\Delta'}\int_{\Delta}\ell(\boldsymbol{x})dv - c_{\Delta'}\sum_{a=1}^k\int_{D_a}\ell(\boldsymbol{x})dv + \left(c_{\Delta'} - (n+1)!\right)\sum_{a=1}^k\int_{D_a}\ell(\boldsymbol{x})dv + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a}) = c_{\Delta'}\int_{\Delta'}\ell(\boldsymbol{x})dv + (c_{\Delta} - c_{\Delta'})\int_{\Delta}\ell(\boldsymbol{x})dv + \left(c_{\Delta'} - (n+1)!\right)\left(\sum_{a=1}^k\int_{D_a}\ell(\boldsymbol{x})dv + \sum_{a=1}^k\sum_{\boldsymbol{a}\in F_a\cap\mathbb{Z}^n}\ell(\boldsymbol{a})\right)$$

through $\Delta' = \Delta \setminus \bigcup_{a=1}^k D_a$. The assertions is verified.

As a special case, we consider the case of $\ell(x) \equiv 1$ in (5.5). By (5.6), we have

$$\sum_{\boldsymbol{a}\in\Delta'\cap\mathbb{Z}^n}1=c_{\Delta'}\int_{\Delta'}1\,dv.$$

Plugging this into (5.11), we obtain the condition

$$(c_{\Delta} - c_{\Delta'}) \operatorname{Vol}(\Delta) + (c_{\Delta'} - (n+1)!) \sum_{a=1}^{k} \operatorname{Vol}(D_a) + \sum_{a=1}^{k} \#(F_a \cap \mathbb{Z}^n) = \mathbf{0}.$$

REFERENCES

[AP09] Claudio Arezzo and Frank Pacard, *Blowing up Kähler manifolds with constant scalar curvature, II.*, Ann. of Math. **170** (2009), no. 2, 685-738.

- [BS99] Victor. V. Batyrev and Elena. N. Selivanova, *Einstein-Kähler metrics on symmetric toric Fano manifolds.*, J. Reine Angew. Math. **512** (1999), 225-236.
- [CdS08] Ana Cannas da Silva, Lectures on Symplectic Geometry., Second, Lecture Notes in Mathematics 1764, Springer Berlin, Heidelberg, 2008.
- [DVZ12] Alberto Della Vedova and Fabio Zuddas, *Scalar curvature and asymptotic Chow stability of projective bundles and blowups.*, Trans. Am. Math. Soc. **364** (2012), no. 12, 6495-6511.
- [Don01] Simon. K. Donaldson, Scalar curvature and projective embeddings, I, J. Differential Geom. 59 (2001), no. 3, 479-522.
- [Fut04] Akito Futaki, *Asymptotic Chow semi-stability and integral invariants*, Internat. J. Math. **15** (2004), no. 9, 967-979.
- [JR24] Chenzi Jin and Yanir A. Rubinstein, *Asymptotics of quantized barycenters of lattice polytopes with applications to algebraic geometry*, arXiv:2406.18969 (2024).
- [Lee22] King Leung Lee, Asymptotic Chow stability of symmetric reflexive toric varieties, arXiv:2212.05000v4 (2022).
- [LLSW19] King Leung Lee, Zhiyuan Li, Jacob Sturm, and Xiaowei Wang, Asymptotic Chow stability of toric Del Pezzo surfaces, Math. Res. Lett. 26 (2019), no. 6, 1759-1787.
 - [LY24] King Leung Lee and Naoto Yotsutani, *Chow stability of* λ -stable toric varieties, arXiv:2405.06883 (2024).
 - [Ler95] Eugene Lerman, Symplectic cuts., Math. Res. Lett. 2 (1995), 247-258.
 - [LT97] Eugene Lerman and Susan Tolman, *Hamiltonian torus actions on symplectic orbifolds and toric varieties.*, Trans. A.M.S. **349** (1997), no. 10, 4201-4230.
 - [Muk03] Shigeru Mukai, *An Introduction to Invariants and Moduli.*, First, Cambridge Studies in Advanced Mathematics **81**, Cambridge Univ. Press, Cambridge, 2003.
 - [Nak15] Yasuhiro Nakagawa, On the examples of Nill and Paffenholz., Int. J. Math. 26 (2015), no. 4, 1377-1389.
 - [Oda12] Yuji Odaka, The Calabi Conjecture and K-stability., Int. Math. Res. Not. IMRN. 2012 (2012), no. 10, 2272-2288.
 - [Ono11] Hajime Ono, A necessary condition for Chow semistability of polarized toric manifolds., J. Math. Soc. Japan. **63** (2011), no. 4, 1377-1389.
 - [OSY12] Hajime Ono, Yuji Sano, and Naoto Yotsutani, An example of an asymptotically Chow unstable manifold with constant scalar curvature., Ann. Inst. Fourier (Grenoble) **62** (2012), no. 4, 1265-1287.
 - [Sto10] Jacopo Stoppa, Unstable blowups, J. Algebraic Geom 19 (2010), no. 1, 1-17.
 - [Tia90] Gang Tian, On Calabi's conjecture for complex surfaces with positive first Chern class, Invent. Math. 101 (1990), no. 1, 101-172.
 - [Tos12] Valentino Tosatti, Kähler-Einstein metrics on Fano surfaces, Expo. Math. 30 (2012), 11-31.

IMAG, UNIVERSITÉ DE MONTPELLIER, 499-554 RUE DU TRUEL 9, 34090 MONTPELLIER, FRANCE *Email address*: king-leung.lee@umontpellier.fr

KAGAWA UNIVERSITY, FACULTY OF EDUCATION, MATHEMATICS, SAIWAICHO 1-1, TAKAMATSU, KAGAWA, 760-8522, JAPAN

Email address: yotsutani.naoto@kagawa-u.ac.jp