

LOCALLY MEASURE PRESERVING PROPERTY OF BI-LIPSCHITZ MAPS BETWEEN MORAN SETS

LIANG-YI HUANG AND SHISHUANG LIU*

ABSTRACT. In literature it is shown that bi-Lipschitz maps between self-similar sets or self-affine sets enjoy a locally measure preserving property, namely, if $f : (E, \mu) \rightarrow (F, \nu)$ is a bi-Lipschitz map, then the Radon-Nykodym derivative $df^*\nu/d\mu$ is a constant function on a subset $E' \subset E$ with $\mu(E') > 0$, where $f^*\nu(\cdot) = \nu(f(\cdot))$. Indeed, this measure preserving property plays an important role in Lipschitz classification of fractal sets. In this paper, we show that such measure preserving property also holds for bi-Lipschitz maps between two Moran sets in a certain class.

1. Introduction

Let (E, d_E, μ) and (F, d_F, ν) be two metric spaces with finite Borel measures μ and ν , respectively. We say these two spaces enjoy a *universal locally measure preserving property*, if for any bi-Lipschitz map $f : E \rightarrow F$, the Radon-Nykodym derivative

$$\frac{df^*\nu}{d\mu} = \text{constant}$$

on a subset $E' \subset E$ with $\mu(E') > 0$, where

$$f^*\nu(\cdot) = \nu(f(\cdot)).$$

Cooper and Pignataro [1] proved that for two self-similar sets $E, F \subset \mathbb{R}$ satisfy the strong separation condition equipped with the Hausdorff measure (positive in its Hausdorff dimension), all monotone bi-Lipschitz maps enjoy the locally measure preserving bi-Lipschitz. Later, Falconer and Marsh [5] and Xi and Ruan [23] showed that the same result holds for $E, F \subset \mathbb{R}^d$ without the orientation preserving assumption. Rao, Yang and Zhang [18] showed the universal locally measure preserving property for Bedford-McMullen carpets equipped with the uniform Bernoulli measures.

This locally measure preserving property turns out to play a very important role in Lipschitz classification of self-similar sets and self-affine sets, see [5, 8, 11–14, 16, 21–23].

Recall that two metric spaces (X, d_X) and (Y, d_Y) are said to be *Lipschitz equivalent*, denoted by $(X, d_X) \sim (Y, d_Y)$, if there exists a map $f : X \rightarrow Y$ which is

Date: July 16, 2024.

2010 Mathematics Subject Classification: 28A80, 26A16

Key words and phrases: Moran set, Measure preserving property .

* Corresponding author.

bi-Lipschitz, that is, there exists a constant $C > 0$ such that for all $x, y \in X$,

$$C^{-1}d_X(x, y) \leq d_Y(f(x), f(y)) \leq Cd_X(x, y).$$

The main goal of the present paper is to show that the universal locally measure preserving property holds for a class of homogeneous Moran sets equipped with the uniform Bernoulli measures.

Let us recall the definition of Moran set. Let $\{n_k\}_{k \geq 1}$ be a sequence of positive integers with $n_k \geq 2$. Set $\Sigma^0 = \emptyset$, and $\Sigma^k = \prod_{j=1}^k \{0, \dots, n_j - 1\}$ for $k \geq 1$. Denote $\Sigma^* = \bigcup_{k \geq 0} \Sigma^k$. For $\sigma = (\sigma_1, \dots, \sigma_k)$ and $\tau = (\tau_1, \dots, \tau_s)$, let $\sigma * \tau = (\sigma_1, \dots, \sigma_k, \tau_1, \dots, \tau_s)$ be their concatenation. Let J be a closed interval. A collection of closed subintervals $\{J_\sigma : \sigma \in \Sigma^*\}$ of J is said to have a *Moran structure*, if

- (i) $J_\emptyset = J$;
- (ii) for any $k \geq 1$, $\sigma \in \Sigma^{k-1}$ and $i \in \{0, \dots, n_k - 1\}$, $J_{\sigma * i}$ are subintervals of J_σ with disjoint interiors.

Let

$$(1.1) \quad \mathbf{F}_k = \bigcup_{\sigma \in \Sigma^k} J_\sigma, \quad F = \bigcap_{k \geq 0} \mathbf{F}_k.$$

Then F is a compact set and we call F a *Moran set* with parameters $(J, \{n_k\}_{k \geq 1})$. Let $\sigma \in \Sigma^k$. We call J_σ a *basic interval* of rank k , and $F_\sigma = J_\sigma \cap F$ a *cylinder* of rank k .

We call F a *homogeneous Moran set* with parameters $(J, \{n_k\}_{k \geq 1}, \{r_k\}_{k \geq 1})$, if in addition, F satisfies that

- (iii) for any $k \geq 1$, $\sigma \in \Sigma^{k-1}$ and $0 \leq i \leq n_k - 1$, $\frac{|J_{\sigma * i}|}{|J_\sigma|} = r_k$, where $|A|$ denotes the diameter of set A .

We denote the collection of such Moran sets by

$$(1.2) \quad \mathcal{F}(J, \{n_k\}_{k \geq 1}, \{r_k\}_{k \geq 1}),$$

see [4, 6, 20]. Let F be a homogeneous Moran set in (1.2), we define $\Delta_1 = |J|(1 - n_1 r_1)$,

$$\Delta_k = |J|r_1 \dots r_{k-1}(1 - n_k r_k), \quad \forall k \geq 2$$

and we call Δ_k the *total gap of rank k* of F .

For $E, F \subset \mathbb{R}$, we denote $\text{dist}(E, F) = \inf\{|x - y|; x \in E, y \in F\}$.

Definition 1.1. We say a Moran set F satisfies the *weak separation condition* (WSC), if there is a constant $\eta_0 > 0$ such that for any $k \geq 0$, $\sigma \in \Sigma^k$ and $i \neq j$,

$$(1.3) \quad \text{either } J_{\sigma * i} \cap J_{\sigma * j} \neq \emptyset \text{ or } \text{dist}(J_{\sigma * i}, J_{\sigma * j}) > \eta_0 \left(|J_\sigma| - \sum_{i=0}^{n_{k+1}-1} |J_{\sigma * i}| \right).$$

Our main result is the following.

Theorem 1.1. *Let $E, F \in \mathcal{F}([0, 1], \{n_k\}_{k \geq 1}, \{r_k\}_{k \geq 1})$. Let $f : E \rightarrow F$ be a bi-Lipschitz map. Suppose that*

- (i) $\sup_{k \geq 1} n_k < \infty$;
- (ii) $\gamma = \inf_{k \geq 1} \frac{\Delta_k}{\Delta_{k+1}} > 1$;

(iii) both E and F satisfy the WSC.

Then for any cylinder E_τ of E , there exists a sub-cylinder $E_{\tau\sigma}$ of E_τ such that $df^*\nu/d\mu$ is a constant function of $E_{\tau\sigma}$, equivalently

$$\frac{\nu(f(\mathbf{A}))}{\mu(\mathbf{A})} = \text{constant}$$

for any Borel set $\mathbf{A} \subset E_{\tau\sigma}$.

2. Proof of Theorem 1.1

Measure preserving properties of bi-Lipschitz maps of self-similar sets and self-affine sets have been studied in [5, 7, 18, 23]. In this section, we investigate such property related to homogenous Moran sets.

In this section, unless otherwise specified, we always assume that E, F are two homogeneous Moran sets in $\mathcal{F}([0, 1], \{n_k\}_{k \geq 1}, \{r_k\}_{k \geq 1})$. Let μ be the Borel probability measure supported on E such that for each k -th cylinder E_τ ,

$$\mu(E_\tau) = \frac{1}{n_1 \dots n_k}.$$

We call μ the *uniform Benoulli measure* of E . Let ν be the uniform Bernoulli measure of F .

Theorem 2.1. *Let $f : E \rightarrow F$ be a bi-Lipschitz map, then $f^*\nu$ is equivalent to μ , that is, there exists $\alpha > 0$ such that*

$$(2.1) \quad \alpha^{-1}\mu(\mathbf{A}) \leq \nu(f(\mathbf{A})) \leq \alpha\mu(\mathbf{A}),$$

for any Borel set $\mathbf{A} \subset E$.

Proof. Let C be a Lipschitz constant of f .

First, we claim that there exists $\alpha > 0$ such that for each cylinder E_τ , it holds that

$$\nu(f(E_\tau)) \leq \alpha\mu(E_\tau).$$

Denote $k = |\tau|$, then $\mu(E_\tau) = \frac{1}{n_1 \dots n_k}$ and

$$\text{diam } f(E_\tau) \leq C(\text{diam } E_\tau) = Cr_1 \dots r_k.$$

Therefore, there are at most $C + 2$ cylinders of rank k of F intersecting $f(E_\tau)$, so we have

$$\nu(f(E_\tau)) \leq \frac{C + 2}{n_1 \dots n_k} = (C + 2)\mu(E_\tau).$$

The claim is proved.

Since the Borel σ -algebra of E can be generated by the collection of cylinders, it follows that $\nu(f(\mathbf{A})) \leq \alpha\mu(\mathbf{A})$ holds for any Borel set $\mathbf{A} \subset E$. By changing the role of E and F , we can get the other side inequality. This completes the proof of the theorem. \square

Definition 2.1 (δ -connected). A compact set $X \subset \mathbb{R}$ is said to be δ -connected, if for any $x, y \in X$, there exists a sequence $\{x = z_1, \dots, z_n = y\} \subset X$ such that $|z_i - z_{i+1}| \leq \delta$ holds for $1 \leq i \leq n - 1$. The sequence above is called a δ -chain connecting x and y .

Recall that \mathbf{E}_k is the union of basic intervals of rank k of E ; we call \mathbf{E}_k the k -th approximation of E . We use \mathcal{E}_k to denote the collection of connected component of \mathbf{E}_k . Similarly, we define \mathbf{F}_k and \mathcal{F}_k . Recall that Δ_k is the total gap of rank k .

The following lemma is a variation of Lemma 3.2 in Falconer and Marsh [5].

Lemma 2.1. *Let $f : E \rightarrow F$ be a bi-Lipschitz map with Lipschitz constant C . Suppose that both E and F satisfy the WSC and*

$$(2.2) \quad \gamma = \inf_{k \geq 1} \frac{\Delta_k}{\Delta_{k+1}} > 1.$$

Then there exists a positive integer p_0 such that, for any $k \geq p_0$ and $I \in \mathcal{E}_k$, there exist $J^ \in \mathcal{F}_{k-p_0}$ and $\{J_j\}_{j=1}^h \subset \mathcal{F}_{k+p_0}$ such that $\{J_j\}_{j=1}^h \subset J^*$ and*

$$f(I \cap E) = \bigcup_{j=1}^h (J_j \cap F).$$

Proof. Let η_0 be a common constant in the weak separation condition of E and F . Let p_0 be the smallest integer satisfying $(\gamma - 1)\gamma^{p_0} > \frac{2C}{\eta_0}$.

Let U, U' be two distinct elements of \mathcal{E}_k . By the weak separation condition, we have

$$\text{dist}(U, U') \geq \min\{\eta_0 \Delta_j; 1 \leq j \leq k\} = \eta_0 \Delta_k.$$

On the other hand, for any $U \in \mathcal{E}_k$, $U \cap E$ is $(\frac{2}{\gamma-1}\Delta_k)$ -connected since the maximal gap of $U \cap E$ is no larger than $2 \sum_{j=1}^{\infty} \Delta_{k+j} \leq 2 \sum_{j=1}^{\infty} \frac{\Delta_k}{\gamma^j} = \frac{2}{\gamma-1}\Delta_k$. Moreover, these two facts are also true for \mathcal{F}_k .

Pick any $J \in \mathcal{F}_{k+p_0}$. We claim that

$$\text{either } (J \cap F) \subset f(I \cap E), \text{ or } (J \cap F) \cap f(I \cap E) = \emptyset.$$

First, $J \cap F$ is $(\frac{2}{\gamma-1}\Delta_{k+p_0})$ -connected implies $f^{-1}(J \cap F)$ is $(\frac{2C}{\gamma-1}\Delta_{k+p_0})$ -connected. By our choice of p_0 , we have $(\frac{2C}{\gamma-1}\Delta_{k+p_0}) \leq (\frac{2C}{\gamma-1}\gamma^{-p_0}\Delta_k) < \eta_0 \Delta_k$, which implies that either $f^{-1}(J \cap F) \subset I$ or $f^{-1}(J \cap F) \cap I = \emptyset$. The claim is proved.

Applying the above claim to the map $f^{-1} : F \rightarrow E$, there exists $J \in \mathcal{F}_{k-p_0}$ such that $f(I \cap E) \subset J \cap F$. This finishes the proof of the lemma. \square

We denote by $\mathcal{C}_E = \bigcup_{k \geq 0} \mathcal{E}_k$, where $\mathcal{E}_0 = \{[0, 1]\}$ by convention. Let $U, U' \in \mathcal{C}_E$, we say U' is an offspring of U if $U' \subset U$, and it is called a direct offspring of U if the rank of U equals the rank of U' plus 1.

Proof of Theorem 1.1. Let E_τ be a cylinder of E . Let U_0 be a component in \mathcal{C}_E such that $U_0 \cap E \subset E_\tau$. Set

$$\mathcal{C}_{E, U_0} = \{V \in \mathcal{C}_E; V \subset U_0\}.$$

Define

$$\varphi(U) = \frac{\nu(f(U \cap E))}{\mu(U)}, \text{ where } U \in \mathcal{C}_{E, U_0}.$$

By Theorem 2.1, we have

$$\chi = \sup_{U \in \mathcal{C}_{E,U_0}} \varphi(U) < \infty.$$

Since each $U \in \mathcal{C}_{E,U_0}$ contains a basic interval and vice versa, the theorem holds if and only if there exists $U \in \mathcal{C}_{E,U_0}$ such that f is measure preserving on $U \cap E$.

Suppose on the contrary that $f|_{U \cap E}$ is not measure preserving for any $U \in \mathcal{C}_{E,U_0}$. Then we can get that $\varphi(U) < \chi$ for all $U \in \mathcal{C}_{E,U_0}$, for otherwise, $f|_{U \cap E}$ is measure preserving for these U such that $\varphi(U)$ attains χ .

Denote $\beta = \sup_{k \geq 1} n_k$. Set $\varepsilon = \frac{1}{2(1+8\beta^{2p_0+5})}$, where p_0 is the constant in Lemma 2.1. By the definition of χ , there is $U \in \mathcal{C}_{E,U_0}$ such that

$$\chi(1 - \varepsilon) < \varphi(U) < \chi.$$

Let k be the rank of U and let U_1, \dots, U_ℓ be the direct offsprings of U , we have either $\varphi(U) = \varphi(U_j)$ for all $1 \leq j \leq \ell$, or there exists $j_0 \in \{1, \dots, \ell\}$ such that $\varphi(U_{j_0}) > \varphi(U)$. Since $f|_{U \cap E}$ is not measure preserving, there exists a sequence $U = V_0, V_1, \dots, V_{t-1}, V_t$ in \mathcal{C}_{E,U_0} such that V_j is directed offspring of V_{j-1} for $j = 1, \dots, t$ and $\varphi(U) = \varphi(V_0) = \varphi(V_1) = \dots = \varphi(V_{t-1}) \neq \varphi(V_t)$. Without loss of generality, we may assume that $\varphi(V_t) > \varphi(V_{t-1})$ and we can replace U by V_{t-1} to start our discussion.

Next we estimate $\varphi(U_{j_0}) = \frac{\nu(f(U_{j_0} \cap E))}{\mu(U_{j_0})}$. Notice that $\varphi(U) = \frac{\nu(f(U \cap E))}{\mu(U)}$, we have

$$(2.3) \quad \varphi(U_{j_0}) = \frac{\nu(f(U_{j_0} \cap E))}{\nu(f(U \cap E))} \cdot \frac{\mu(U)}{\mu(U_{j_0})} \cdot \varphi(U).$$

By Lemma 2.1, $f(U \cap E)$ can be decomposed into

$$f(U \cap E) = \bigcup_{j=1}^h (J_j \cap F),$$

where $\{J_j\}_{j=1}^h \subset \mathcal{F}_{k+p_0}$, and are offsprings of $J^* \in \mathcal{F}_{k-p_0}$. Since each $J \in \mathcal{F}_{k-p_0}$ contains at most $2n_{k-p_0}$ basic intervals of rank $k - p_0$, we have

$$h \leq (2n_{k-p_0}) \prod_{j=1}^{2p_0} n_{k-p_0+j} \leq 2\beta^{2p_0+1}.$$

Applying Lemma 2.1 to U_{j_0} , there exists $\{I_j\}_{j=1}^s \subset \mathcal{F}_{k+1+p_0}$ such that $f(U_{j_0} \cap E) = \bigcup_{j=1}^s (I_j \cap F)$.

For any connected component Q of rank k , denote by $\#_k Q$ the number of k -th basic intervals contained in Q . We have

$$\nu(f(U \cap E)) = \sum_{j=1}^h \nu(J_j \cap F) = \sum_{j=1}^h \frac{\#_{(k+p_0)} J_j}{n_1 \dots n_{k+p_0}},$$

and

$$\nu(f(U_{j_0} \cap E)) = \sum_{j=1}^s \nu(I_j \cap F) = \sum_{j=1}^s \frac{\#_{(k+p_0+1)} I_j}{n_1 \dots n_{k+p_0} n_{k+p_0+1}}.$$

Substituting these two relations to (2.3), we obtain

$$\frac{\varphi(U_{j_0})}{\varphi(U)} = \frac{\sum_{j=1}^s \#_{(k+p_0+1)} I_j}{\sum_{j=1}^h \#_{(k+p_0)} J_j} \cdot \frac{1}{n_{k+p_0+1}} \cdot \frac{\#_k U}{\#_{(k+1)} U_{j_0}} \cdot \frac{1}{n_{k+1}}.$$

Notice that the denominator of the right hand side is no larger than $8\beta^{2p_0+5}$, by $\varphi(U_{j_0}) > \varphi(U)$, we obtain

$$(2.4) \quad \varphi(U_{j_0}) > \left(1 + \frac{1}{8\beta^{2p_0+5}}\right)\varphi(U) > \chi,$$

which contradicts the definition of χ . This completes the proof of the Theorem. \square

REFERENCES

- [1] D. Cooper and T. Pignataro, *On the shape of Cantor sets*, J. Differential Geom., **28** (1988), 203-221.
- [2] G. David and S. Semmes, *Fractured fractals and broken dreams: self-similar geometry through metric and measures*, Oxford Univ. Press, (1997).
- [3] J. Deng, Z. Y. Wen, Y. Xiong and L. F. Xi, *Bilipschitz embedding of self-similar sets*, J. Anal. Math., **114** (2011), 63-97.
- [4] D. J. Feng, Z. Y. Wen and J. Wu, *Some dimensional results for homogeneous Moran sets*, Sci. China Ser. A, **40** (1999), 475-482.
- [5] K. J. Falconer and D. T. Marsh, *On the Lipschitz equivalence of Cantor sets*, Mathematika, **39** (1992), 223-233.
- [6] S. Hua, H. Rao, Z. Y. Wen and J. Wu, *On the structures and dimensions of Moran sets*, Sci. China Ser. A, **43** (2000), 836-852.
- [7] L.Y. Huang, H. Rao, Z.Y. Wen and Y.L. Xu, *Box-counting measure of metric spaces*, Preprint (2024) (arXiv:2111.00752).
- [8] M. Llorente and P. Mattila, *Lipschitz equivalence of subsets of self-conformal sets*, Nonlinearity, **23** (2010), 875-882.
- [9] B. M. Li, W. X. Li and J. J. Miao, *Lipschitz equivalence of McMullen sets*, Fractals, **21** (2013), 1350022, 11 pp.
- [10] Z. Liang, J. J. Miao and H. J. Ruan, *Gap sequences and topological properties of Bedford-McMullen sets*, Nonlinearity, **35** (2022), 4043-4063.
- [11] J. J. Luo and K. S. Lau, *Lipschitz equivalence of self-similar sets and hyperbolic boundaries*, Adv. Math., **235** (2013), 555-579.
- [12] H. Rao, H. J. Ruan and L. F. Xi, *Lipschitz equivalence of self-similar sets*, C. R. Math. Acad. Sci. Paris, **342** (2006), 191-196.
- [13] H. Rao, H. J. Ruan and Y. Wang, *Lipschitz equivalence of Cantor sets and algebraic properties of contraction ratios*, Trans. Amer. Math. Soc., **364** (2012), 1109-1126.
- [14] H. Rao, H. J. Ruan and Y. Wang, *Lipschitz equivalence of self-similar sets: algebraic and geometric properties*, Contemp. Math., **600** (2013) 349-364.
- [15] H. Rao and Y. Zhang, *Higher dimensional Frobenius problem and Lipschitz equivalence of Cantor sets*, J. Math. Pures Appl. **104** (2015), 868-881.
- [16] H. J. Ruan, Y. Wang and L. F. Xi, *Lipschitz equivalence of self-similar sets with touching structures*, Nonlinearity, **27** (2014), 1299-1321.
- [17] H. J. Ruan and Y. Wang, *Topological invariants and Lipschitz equivalence of fractal squares*, J. Math. Anal. Appl., **451** (2017), 327-344.
- [18] H. Rao, Y. M. Yang and Y. Zhang, *Invariance of multifractal spectrum of uniform self-affine measures and its applications*, Preprint (2021) (arXiv:2005.07451).
- [19] Y. M. Yang and Y. Zhang, *Lipschitz classification of Bedford-McMullen carpets with uniform horizontal fibers*, J. Math. Anal. Appl., **495** (2021), 124742.

- [20] Z. Y. Wen, *Moran sets and Moran classes*, Chinese Sci. Bull., **46** (2001), 1849-1856.
- [21] L. F. Xi, *Lipschitz equivalence of dust-like self-similar sets*, Math. Z., **266** (2010), 683-691.
- [22] L. F. Xi and Y. Xiong, *Algebraic criteria for Lipschitz equivalence of dust-like self-similar sets*, J. Lond. Math. Soc., **103** (2021), 760-780.
- [23] L. F. Xi and H. J. Ruan, *Lipschitz equivalence of self-similar sets satisfying the strong separation condition*, Acta Math. Sinica (Chinese Ser.), **51** (2008), 493-500.

COLLEGE OF SCIENCE, WUHAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, 430070,
CHINA

Email address: liangyihuang@wust.edu.cn

DEPARTMENT OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY,
WUHAN, 430079, CHINA

Email address: shishuangliu@mails.ccnu.edu.cn