# ON THE SET OF EXTREME POINTS OF THE UNIT BALL OF A HARDY-LORENTZ SPACE

SERGEY V. ASTASHKIN

ABSTRACT. We prove that every measurable function  $f : [0, a] \to \mathbb{C}$  such that |f| = 1a.e. on [0, a] is an extreme point of the unit ball of the Lorentz space  $\Lambda(\varphi)$  on [0, a]whenever  $\varphi$  is a not linear, strictly increasing, concave, continuous function on [0, a]with  $\varphi(0) = 0$ . As a consequence, we complement the classical de Leeuw-Rudin theorem on a description of extreme points of the unit ball of  $H^1$  showing that  $H^1$ is a unique Hardy-Lorentz space  $H(\Lambda(\varphi))$ , for which every extreme point of the unit ball is a normed outer function. Moreover, assuming that  $\varphi$  is strictly increasing and strictly concave, we prove that every function  $f \in H(\Lambda(\varphi))$ ,  $||f||_{H(\Lambda(\varphi))} = 1$ , such that the absolute value of its nontangential limit  $f(e^{it})$  is a constant on some set of positive measure of  $[0, 2\pi]$ , is an extreme point of the unit ball of  $H(\Lambda(\varphi))$ .

Primary classification: 46E30

).

Secondary classification(s): 30H10, 30J05, 46A55, 46B22

Lorentz space, Hardy-Lorentz space,  $H^p$  space, symmetric space, rearrangement, extreme point, inner function, outer function

## 1. INTRODUCTION

In 1958, de Leeuw and Rudin gave the following remarkable characterization of extreme points of the unit ball of the Hardy space  $H^1 := H^1(\mathbb{D})$  (see [6, Theorem 1] or [9, Chapter 9]).

**Theorem 1** (de Leeuw-Rudin). A function in  $H^1$  is an extreme point of the unit ball of  $H^1$  if and only if it has norm one and is an outer function.

In 1974, this result has been partially extended by Bryskin and Sedaev to general Hardy-type spaces H(X) such that the norm  $\|\cdot\|_X$  of a symmetric space X on  $[0, 2\pi]$  is strictly monotone [2, Theorem 1].

The norm  $\|\cdot\|_X$  of a symmetric space X = X[0, a] is strictly monotone whenever from the conditions  $x, y \in X$ ,  $|x| \leq |y|$  a.e. on [0, a] and

$$m\{t \in [0, a] : |x(t)| < |y(t)|\} > 0$$

it follows that  $||x||_X < ||y||_X$ .

<sup>&</sup>lt;sup>†</sup> This work was performed at the Saint Petersburg Leonhard Euler International Mathematical Institute and supported by the Ministry of Science and Higher Education of the Russian Federation (agreement no. 075-15-2022-287.

**Theorem 2** (Bryskin-Sedaev). If the norm of a symmetric space X is strictly monotone, then every normed outer function in H(X) is an extreme point of the unit ball of H(X).

In this paper, we will be interested mainly in Lorentz and Hardy-Lorentz spaces. Let  $\varphi$  be an increasing, concave, continuous, real-valued function on [0, a] such that  $\varphi(0) = 0$  and  $\varphi(a) = 1$ . The Lorentz space  $\Lambda(\varphi) := \Lambda(\varphi)[0, a]$  consists of all complexvalued functions x(t) measurable on [0, a] and satisfying the condition:

(1) 
$$\|x\|_{\Lambda(\varphi)} := \int_0^a x^*(t) \, d\varphi(t) < \infty,$$

where  $x^*(t)$  is the decreasing rearrangement of |x|. Since  $\varphi$  being strictly monotone implies that the norm in  $\Lambda(\varphi)$  is strictly monotone (see Lemma 4 below), the result of Theorem 2 holds, in particular, under the above condition imposed on  $\varphi$ , for the space  $H(\Lambda(\varphi))$  (see [2, Corollary]).

In 1978, E. Semenov stated the problem of describing the set of extreme points of the unit ball of a Hardy-Lorentz space  $H(\Lambda(\varphi))$ , [16] as a part of a collection of 99 problems in linear and complex analysis from the Leningrad branch of the Steklov Mathematical Institute, [10, 5.1 pp.23-24]. Later on, the same problem appeared in the subsequent list of problems in 1984, [14, 1.6 pp.22-23], and in 1994, [15, 1.5 p.12].

Observe that the mapping

(2) 
$$f \mapsto \tilde{f} := f(e^{it}),$$

where we denote by  $f(e^{it})$  the nontangential limit of f(z) as  $z \to e^{it}$ , is an isometric inclusion of a Hardy-type space H(X) into the underlying symmetric space  $X = X[0, 2\pi]$ . Therefore, in connection with the above problem, it is important to find out which functions are being extreme points of the unit ball of a Lorentz space. The following result (see [4, Proposition 2.2], [5, Theorem 1] and [16]) contains a description of extreme points for the unit ball of  $\Lambda(\varphi)$  whenever  $\varphi$  is strictly concave.

**Theorem 3.** If f is an extreme point of the unit ball of the space  $\Lambda(\varphi) := \Lambda(\varphi)[0, a]$ , then

(3) 
$$f(t) = \frac{\varepsilon(t)}{\varphi(m(E))} \chi_E(t), \quad 0 \le t \le a,$$

where  $E \subset [0, a]$  is a measurable set and  $\varepsilon$  is a complex-valued measurable function such that  $|\varepsilon| = 1$  a.e.

Moreover, if the function  $\varphi$  is strictly concave, f is an extreme point of the unit ball of  $\Lambda(\varphi)$  if and only if f can be represented in the form (3).

Our first main result refines Theorem 3 in the case when the function  $\varphi$ , in general, fails to be strictly concave, showing that every measurable function  $f : [0, 2\pi] \to \mathbb{C}$ such that |f| = 1 a.e. on  $[0, 2\pi]$  is an extreme point of the unit ball of the space  $\Lambda(\varphi)$  if  $\varphi$  is not linear and strictly increasing. This result can be compared with the easy fact

 $\mathbf{2}$ 

that the set of extreme points of the unit ball of the space  $L^1$  (which coincides with  $\Lambda(\varphi_0)$ ), where  $\varphi_0(t) = t/(2\pi)$ ), is empty. As a consequence, we get that  $H^1 := H^1(\mathbb{D})$  is a unique space in the class of Hardy-Lorentz spaces  $H(\Lambda(\varphi))$ , where  $\varphi$  is a strictly increasing concave function on  $[0, 2\pi]$ , for which every extreme point of the unit ball is a normed outer function.

In the recent note [3], it was proved that every inner function is an extreme point of the unit ball of  $H(\Lambda(\varphi))$  whenever  $\varphi$  is strictly increasing and strictly concave on  $[0, 2\pi]$ . However, one can easily see that this result is an immediate consequence of Theorem 3. Indeed, for each inner function f we certainly have  $|\tilde{f}| = 1$  a.e. on  $[0, 2\pi]$ . Therefore,  $\tilde{f}$  is an extreme point of the unit ball of the space  $\Lambda(\varphi)$ . Since mapping (2) includes  $H(\Lambda(\varphi))$  into  $\Lambda(\varphi)$  as an isometric subspace, it follows then that f is an extreme point of the unit ball of the space  $H(\Lambda(\varphi))$ .

The second result of the paper resolves the above Semenov's problem for the class of functions  $f \in H(\Lambda(\varphi))$  such that  $|\tilde{f}|$  is constant on some subset of positive measure of  $[0, 2\pi]$ . More precisely, we prove that such a normed function is an extreme point of the unit ball of  $H(\Lambda(\varphi))$  if  $\varphi$  is strictly increasing and strictly concave on  $[0, 2\pi]$ . Note that this class contains, in particular, functions which are being neither inner, nor outer ones.

In a subsequent paper<sup>1</sup>, we investigate a different case when f is a normed function in  $H(\Lambda(\varphi))$  such that  $m\{t : |f(e^{it})| = c\} = 0$  for each  $c \ge 0$  and f(a) = 0 for some point  $a \in \mathbb{D}$ .

It is worth also to say briefly about resent investigations of the structure of extreme points of the unit ball for other types of function spaces which can be treated as a generalization of the classical  $H^1$ . We mention here only the paper [8], where a similar study was carried out for a punctured Hardy space  $H^1_K$  consisting of the integrable functions f on the unit circle whose Fourier coefficients  $c_k(f)$  vanish either k < 0 or  $k \in K$  (K is a fixed finite set of positive integers).

The author thanks Professors A. Baranov, G. Curbera and I. Kayumov for useful discussions related to matters considered in the paper.

#### 2. Preliminaries.

2.1. Symmetric spaces. A detailed exposition of the theory of symmetric spaces see in [1, 11, 13].

Let a > 0. A Banach space X := X[0, a] of complex-valued Lebesgue-measurable functions on the measure space ([0, a], m), where m is the Lebesgue measure on the interval [0, a], is called *symmetric* (or *rearrangement invariant*) if from the conditions  $y \in X$  and  $x^*(t) \leq y^*(t)$  almost everywhere (a.e.) on [0, a] it follows that  $x \in X$  and  $\|x\|_X \leq \|y\|_X$ . Here,  $x^*(t)$  is the right-continuous nonincreasing *rearrangement* of |x|,

<sup>&</sup>lt;sup>1</sup>S.V. Astashkin, On the set of extreme points of the unit ball of a Hardy-Lorentz space, 2 (manuscript).

i.e.,

4

$$x^*(t) := \inf\{\tau \ge 0 : m\{s \in [0, a] : |x(s)| > \tau\} \le t\}, \ 0 < t \le a.$$

Measurable functions x and y on [0, a] are said to be *equimeasurable* whenever

$$m\{s \in [0,a]: |x(s)| > \tau\} = m\{s \in [0,a]: |y(s)| > \tau\}, \ \tau > 0.$$

In particular, a complex-valued measurable function x(t) and its rearrangement  $x^*(t)$  are equimeasurable. Clearly, if X is a symmetric space,  $y \in X$  and x and y are equimeasurable on [0, a], then  $x \in X$  and  $||x||_X = ||y||_X$ .

Every symmetric space X on [0, a] satisfies the embeddings  $L^{\infty}[0, a] \subset X \subset L^{1}[0, a]$ .

The family of symmetric spaces includes many classical spaces appearing in analysis, in particular,  $L^p$ -spaces with  $1 \le p \le \infty$ , which are defined in the usual way, Orlicz, Lorentz, Marcinkiewicz spaces and many others. Next, we will be interested primarily in Lorentz spaces.

In what follows,  $\varphi$  is an increasing, concave, continuous function on [0, a] such that  $\varphi(0) = 0$  and  $\varphi(a) = 1$ . Then (see Section 1), the Lorentz space  $\Lambda(\varphi) := \Lambda(\varphi)[0, a]$  is equipped with norm (1). In particular, if  $\varphi(t) = a^{-1/p}t^{1/p}$ ,  $1 , this space is denoted, as usual, by <math>L^{p,1}$  (see, e.g.,  $[1, \S4.4]$  or  $[11, \SII.6.8]$ ).

For every  $\varphi$ ,  $\Lambda(\varphi)$  is a separable symmetric space. Moreover, such a space has the Fatou property, that is, the conditions  $x_n \in \Lambda(\varphi)$ ,  $n = 1, 2, \ldots$ ,  $\sup_{n=1,2,\ldots} \|x_n\|_{\Lambda(\varphi)} < \infty$ , and  $x_n \to x$  a.e. on [0, a] imply that  $x \in \Lambda(\varphi)$  and  $\|x\|_{\Lambda(\varphi)} \leq \liminf_{n\to\infty} \|x_n\|_{\Lambda(\varphi)}$ .

2.2. Hardy-type spaces of analytic functions. Let X be a symmetric space with the Fatou property on the unit circle

$$\mathbb{T} := \{ e^{it} : 0 \le t \le 2\pi \},\$$

which will be identified further with the interval  $[0, 2\pi]$ . The Hardy-type space H(X) consists of all  $f: \mathbb{D} \to \mathbb{C}$ , where  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ , such that

$$||f||_{H(X)} := \sup_{0 \le r < 1} ||f_r||_X < \infty,$$

where  $f_r(e^{it}) := f(re^{it})$  if  $0 \le r < 1$  and  $t \in [0, 2\pi]$  (see [2]). In particular, if  $X = L^p$ ,  $1 \le p \le \infty$ , we get the classical  $H^p$ -spaces,  $H^p = H(L^p)$  [7, 9, 12].

Since  $X \subset L^1$  for every symmetric space X, we have  $H(X) \subseteq H^1$ . Consequently, for each function  $f \in H(X)$  and almost all  $t \in [0, 2\pi]$  there exists the nontangential limit  $f(e^{it}) := \lim_{z \to e^{it}} f(z)$ . Moreover, precisely as in the classical  $H^p$  case, the mapping (2) is an isometry between H(X) and the subspace of X, consisting of all functions  $g \in X$ such that the Fourier coefficients

$$c_k(g) := \frac{1}{2\pi} \int_0^{2\pi} g(t) e^{-ikt} dt$$

vanish if k < 0.

Recall the canonical (inner-outer) factorization theorem for  $H^1$  functions. A function  $I \in H^{\infty}$  is called *inner* if  $|\tilde{I}| = 1$  a.e. on  $[0, 2\pi]$ . Also, a non-null function  $F \in H^1$  is termed *outer* if

$$\ln|F(0)| = \frac{1}{2\pi} \int_0^{2\pi} \ln|\tilde{F}(s)| \, ds.$$

As is well known, every function  $f \in H^1$ ,  $f \not\equiv 0$ , can be represented as f = IF, where I is inner and F is outer (see, for instance, [12, §IV.D.4<sup>0</sup>]).

Let  $g \in H(X)$ , where X is a symmetric space on  $[0, 2\pi]$ . Since  $g \in H^1$ , we have

$$\int_0^{2\pi} |\ln|\tilde{g}(t)|| \, dt < \infty$$

(see, for instance,  $[12, \S IV.D.4^0]$ ). Conversely, if  $\mu \in X$ ,  $\mu \ge 0$ , and  $\ln \mu \in L^1$ , then there are functions  $g \in H(X)$  such that

$$(4) \qquad \qquad |\tilde{g}| = \mu.$$

Indeed, for example, the outer function

$$F(z) := \exp\left\{\frac{1}{2\pi} \int_0^{2\pi} \frac{e^{it} + z}{e^{it} - z} \ln \mu(t) \, dt + i\lambda\right\}, \quad z \in \mathbb{D},$$

where  $\lambda \in \mathbb{R}$ , satisfies condition (4) [12, § *IV*.*D*.4<sup>0</sup>] and hence belongs to H(X).

2.3. Extreme points and some related definitions. Given Banach space  $X = (X, \|\cdot\|)$ , denote by ball (X) the closed unit ball of X, that is,

$$ball(X) := \{ x \in X : ||x|| \le 1 \}.$$

A point  $x_0 \in \text{ball}(X)$  is said to be *extreme* for the set ball(X) if  $x_0$  is not the midpoint of any nondegenerate segment contained in ball(X).

Clearly, every extreme point  $x_0$  of ball (X) is a normed element, i.e.,  $||x_0|| = 1$ . Moreover,  $x_0$  is an extreme point of ball (X) if and only if from  $y \in X$  and  $||x \pm y||_X \le 1$  it follows that y = 0.

We say that a function  $\varphi : [0, a] \to \mathbb{R}$  is *increasing* (resp. *strictly increasing*) if from  $0 \le t_1 < t_2 \le a$  it follows  $\varphi(t_1) \le \varphi(t_2)$  (resp.  $\varphi(t_1) < \varphi(t_2)$ ). A (strictly) decreasing function is defined similarly.

A real-valued function  $\varphi$  defined on [0, a] is called *strictly concave* if

$$\varphi((1-\alpha)t_1 + \alpha t_2) < (1-\alpha)\varphi(t_1) + \alpha\varphi(t_2)$$

for any  $\alpha \in (0, 1)$  and  $0 \le t_1, t_2 \le a, t_1 \ne t_2$ .

### 3. AUXILIARY RESULTS

The proof of the following assertion is immediate and hence we skip it.

**Lemma 4.** If  $\varphi$  is a strictly increasing, concave function on [0, a],  $\varphi(0) = 0$ ,  $\varphi(a) = 1$ , then the norm of the Lorentz space  $\Lambda(\varphi)$  is strictly monotone.

An inspection of the proof of de Leeuw-Rudin theorem in [6, Theorem 1] or [9, Chapter 9] implies the following useful result (see also Lemma 1 in [2]).

**Lemma 5.** Suppose the norm of a symmetric space X on [0, a] is strictly monotone. If  $||f||_X = 1$ ,  $g \in X$ ,  $m\{t : f(t) = 0\} = 0$ , and  $||f \pm g||_X \le 1$ , then g = hf, where h is a real-valued function,  $|h| \le 1$  a.e. on [0, a].

A proof of the following result can be found in [3, Lemma 6] (for the special case  $\varphi(t) = a^{-1/p} t^{1/p}$ , 1 , see also [5, Lemma 1]).

**Lemma 6.** Let  $\varphi$  be a strictly increasing and strictly concave function on [0, a] such that  $\varphi(0) = 0$ ,  $\varphi(a) = 1$ . Assuming that  $f, g \in \Lambda(\varphi)$  satisfy

$$||f||_{\Lambda(\varphi)} + ||g||_{\Lambda(\varphi)} = ||f + g||_{\Lambda(\varphi)},$$

we have

$$f^* + g^* = (f + g)^*$$
 a.e. on  $[0, a]$ 

As is known (see, for instance, [11, § II.2, the proof of Property 7<sup>0</sup>], for every function  $x \in L_1[0, a]$  there exists a family of measurable subsets  $\{E_t(x)\}_{0 \le t \le a}$  of the interval [0, a] such that  $m(E_t(x)) = t$ ,  $0 < t \le a$ ,  $E_{t_1}(x) \subset E_{t_2}(x)$  if  $0 < t_1 < t_2 \le a$ , the embeddings

(5)  $\{s \in [0, a] : |x(s)| > x^*(t)\} \subset E_t(x) \subset \{s \in [0, a] : |x(s)| \ge x^*(t)\}, 0 < t \le a,$  hold, and

(6) 
$$\int_{E_t(x)} |x(s)| \, ds = \int_0^t x^*(s) \, ds, \quad 0 < t \le a.$$

**Proposition 7.** Let  $\varphi$  be a strictly increasing and strictly concave function on [0, a],  $\varphi(0) = 0$ ,  $\varphi(a) = 1$ . Suppose  $u, v \in \Lambda(\varphi)[0, a]$ , u is nonnegative, v is real-valued,  $\|u\|_{\Lambda(\varphi)} = 1$  and  $\|u \pm v\|_{\Lambda(\varphi)} \leq 1$ .

Then both functions  $u(s) \pm v(s)$  are nonnegative a.e. on the interval [0, a] and there exists a family of measurable subsets  $\{E_t\}_{0 \le t \le a}$  of [0, a] which satisfies the following properties:  $m(E_t) = t$ ,  $0 \le t \le a$ ,  $E_{t_1} \subset E_{t_2}$ ,  $0 \le t_1 \le t_2 \le a$ ,

(7) 
$$\int_{E_t} u(s) \, ds = \int_0^t u^*(s) \, ds, \quad 0 < t \le a,$$

and

(8) 
$$\int_{E_t} (u(s) \pm v(s)) \, ds = \int_0^t (u \pm v)^*(s) \, ds, \quad 0 < t \le a.$$

 $\mathbf{6}$ 

*Proof.* First, from the hypothesis of the proposition it follows that  $||u \pm v||_{\Lambda(\varphi)} = 1$ . Hence,

$$||u+v||_{\Lambda(\varphi)} + ||u-v||_{\Lambda(\varphi)} = 2 = ||2u||_{\Lambda(\varphi)}.$$

Thus, the functions u+v and u-v satisfy the conditions of Lemma 6, and consequently

(9) 
$$(u+v)^* + (u-v)^* = 2u^*$$
 a.e. on  $[0,a]$ .

In view of [11, §II.2, Property 9<sup>0</sup>], the last inequality is equivalent to the fact that the functions  $u \pm v$  are of the same sign a.e. on [0, a] and as families of subsets  $\{E_t(u+v)\}_{0 \le t \le a}$  and  $\{E_t(u-v)\}_{0 \le t \le a}$  (see the discussion before the proposition), can be chosen the same family  $\{E_t\}_{0 \le t \le a}$ . Since  $u \ge 0$ , this implies, in particular, that the both functions  $u \pm v$  are a.e. nonnegative. Therefore, by (6), we obtain (8), that is, we have

$$\int_{E_t} (u(s) + v(s)) \, ds = \int_0^t (u+v)^*(s) \, ds, \ 0 < t \le a,$$

and

$$\int_{E_t} (u(s) - v(s)) \, ds = \int_0^t (u - v)^*(s) \, ds, \ 0 < t \le a.$$

Summing up these equations and applying (9), we get

$$2\int_{E_t} u(s) \, ds = \int_0^t ((u+v)^*(s) + (u-v)^*(s)) \, ds$$
$$= 2\int_0^t u^*(s) \, ds, \quad 0 < t \le a,$$

which implies (7). This completes the proof.

**Lemma 8.** Let a > 0 and let  $\varphi$  be a nonlinear, strictly increasing, concave, continuous function on [0, a] such that  $\varphi(0) = 0$  and  $\varphi(a) = 1$ . Assuming that a function  $\rho$ :  $[0, a] \to \mathbb{R}$  satisfies the condition:  $0 < m(\operatorname{supp} \rho) < a$ , we have

$$\int_0^a \rho^*(t) \, d\varphi(t) > -\int_0^a \rho^*(t) \, d\varphi(a-t).$$

*Proof.* First, we put  $b := m(\operatorname{supp} \rho)$ . Since the function  $\varphi$  is concave, the derivative  $\varphi'(t)$  exists a.e. on [0, a] and decreases. Moreover, by the assumption,  $\varphi$  is not linear, and hence  $\varphi'(t)$  is not constant on [0, a]. Therefore, since b < a, there exists  $0 < \delta < a - b$  such that for all  $0 < t < \delta$  we have

$$\varphi'(a-t) < \varphi'(t).$$

Because  $\rho \neq 0$ , this implies that

(10) 
$$\int_0^\delta \rho^*(t)\varphi'(t)\,dt > \int_0^\delta \rho^*(t)\varphi'(a-t)\,dt.$$

Next, from the inequality  $a > b + \delta$  it follows

$$\varphi'(a+t-b-\delta) \le \varphi'(t), \ \delta < t < b.$$

Moreover, the increasing function  $t \mapsto \varphi'(a-t)$  and decreasing function  $t \mapsto \varphi'(a+t-b-\delta)$  are equimeasurable, when restricted to the interval  $[\delta, b]$ . Thus, in view of the equality  $\rho'(t) = 0, t \in (b, a]$ , and a well-known property of the decreasing rearrangement (see, for instance, [11, §II.2, Property 16<sup>0</sup>]), we have

$$\begin{split} \int_{\delta}^{a} \rho^{*}(t)\varphi'(t) \, dt &= \int_{\delta}^{b} \rho^{*}(t)\varphi'(t) \, dt \\ &\geq \int_{\delta}^{b} \rho^{*}(t)\varphi'(a+t-b-\delta) \, dt \\ &\geq \int_{\delta}^{b} \rho^{*}(t)\varphi'(a-t) \, dt \\ &= \int_{\delta}^{a} \rho^{*}(t)\varphi'(a-t) \, dt. \end{split}$$

Combining this together with (10), we obtain

$$\int_0^a \rho^*(t)\varphi'(t) dt = \int_0^\delta \rho^*(t)\varphi'(t) dt + \int_\delta^a \rho^*(t)\varphi'(t) dt$$
  
> 
$$\int_0^\delta \rho^*(t)\varphi'(a-t) dt + \int_\delta^a \rho^*(t)\varphi'(a-t) dt$$
  
= 
$$\int_0^a \rho^*(t)\varphi'(a-t) dt = -\int_0^a \rho^*(t) d\varphi(a-t),$$

and thus everything is done.

Remark 9. The simple example  $\rho(t) = 1, \ 0 \le t \le a$  shows that the assumption  $m(\operatorname{supp} \rho) < a$  in the last lemma cannot be skipped.

# 4. Main results

Our first result refines Theorem 3 in the case when a concave function  $\varphi$ , in general, fails to be strictly concave. As a consequence, we get that the classical  $H^1$  is a unique space in the class of Hardy-Lorentz spaces  $H(\Lambda(\varphi))$ , with strictly increasing, concave  $\varphi$ , such that each extreme point of the unit ball of  $H(\Lambda(\varphi))$  is a normed outer function.

**Theorem 10.** Suppose  $\varphi$  is a nonlinear, strictly increasing, concave, continuous function on [0, a] such that  $\varphi(0) = 0$ ,  $\varphi(a) = 1$ . Then, every measurable function  $f : [0, a] \to \mathbb{C}$  such that |f| = 1 a.e. on [0, a] is an extreme point of the unit ball of the space  $\Lambda(\varphi)$ .

*Proof.* Assuming that f is not an extreme point of the unit ball of the space  $\Lambda(\varphi)$ , we find a function  $g \in \Lambda(\varphi)$ ,  $g \neq 0$ , such that

(11) 
$$\|f \pm g\|_{\Lambda(\varphi)} = 1.$$

Since  $\varphi$  is strictly increasing,  $m\{t: f(t) = 0\} = 0$  and  $||f||_{\Lambda(\varphi)} = 1$ , by Lemmas 4 and 5, we conclude that  $g = h \cdot f$ , where h is a real-valued function,  $|h| \leq 1$  a.e. on [0, a]. Thus, from the assumption |f| = 1 a.e. on [0, a] and (11) it follows that

(12) 
$$\|1 \pm h\|_{\Lambda(\varphi)} = 1.$$

Setting

$$E_{+} := \{t \in [0, a] : h(t) > 0\}, \quad E_{-} := \{t \in [0, a] : h(t) < 0\},$$

by the definition of the norm in a Lorentz space, we have

(13) 
$$\|1+h\|_{\Lambda(\varphi)} = \int_0^{m(E_+)} (1+h)^*(t) \, d\varphi(t) + \int_{a-m(E_-)}^a (1+h)^*(t) \, d\varphi(t) + \varphi(a-m(E_-)) - \varphi(m(E_+)).$$

A direct calculation shows that

$$(1+h)^*(t) = (h\chi_{E_+})^*(t) + 1, \ 0 \le t \le m(E_+).$$

and

$$(1+h)^*(t) = 1 - (h\chi_{E_-})^*(a-t), \ a - m(E_-) \le t \le a.$$

Therefore,

$$\|1+h\|_{\Lambda(\varphi)} = \varphi(m(E_{+})) + \int_{0}^{m(E_{+})} (h\chi_{E_{+}})^{*}(t) \, d\varphi(t) \\ + \int_{a-m(E_{-})}^{a} (1 - (h\chi_{E_{-}})^{*}(a-t)) \, d\varphi(t) \\ + \varphi(a - m(E_{-})) - \varphi(m(E_{+})) \\ = \int_{0}^{m(E_{+})} (h\chi_{E_{+}})^{*}(t) \, d\varphi(t) + 1 - \varphi(a - m(E_{-})) \\ - \int_{a-m(E_{-})}^{a} (h\chi_{E_{-}})^{*}(a-t) \, d\varphi(t) + \varphi(a - m(E_{-})) \\ = 1 + \int_{0}^{m(E_{+})} (h\chi_{E_{+}})^{*}(t) \, d\varphi(t) + \int_{0}^{m(E_{-})} (h\chi_{E_{-}})^{*}(t) \, d\varphi(a-t).$$

Similarly, we have

(15) 
$$\|1-h\|_{\Lambda(\varphi)} = 1 + \int_0^{m(E_-)} (h\chi_{E_-})^*(t) \, d\varphi(t) + \int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(a-t).$$
  
Note that

Note that

(16) 
$$0 < m(E_+) < a.$$

Indeed, assume that  $m(E_+) = 0$  or  $m(E_+) = a$ . Then, one of the following conditions holds: (a)  $m(E_+) = 0$  and  $m(E_-) > 0$ , (b)  $m(E_-) = 0$  and  $m(E_+) > 0$ , (c)  $m(E_+) = m(E_-) = 0$ .

In the case (a) we have

$$\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(t) = 0$$

and

$$\int_0^{m(E_-)} (h\chi_{E_-})^*(t) \, d\varphi(a-t) < 0.$$

since  $\varphi$  is strictly increasing and h(t) < 0 if  $t \in E_-$ . Consequently, by equation (14), we see that  $\|1+h\|_{\Lambda(\varphi)} < 1$ . But this impossible, because it contradicts condition (12).

In the case (b), arguing similarly, we obtain that  $||1+h||_{\Lambda(\varphi)} > 1$ , which is impossible also in view of (12).

Consider the case (c). The condition  $m(E_+) = m(E_-) = 0$  is equivalent to the fact that h = 0 a.e. on [0, a]. Then, g = 0 a.e. on [0, a], which contradicts the assumption. As a result, (16) is proved.

Next, in view of (16) and the definition of the sets  $E_+$  and  $E_-$ , the functions  $\rho_1 := h\chi_{E_+}$  and  $\rho_2 := h\chi_{E_-}$  satisfy the conditions of Lemma 8. Therefore, we have

(17) 
$$\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(t) > -\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(a-t).$$

and

(18) 
$$\int_{0}^{m(E_{-})} (h\chi_{E_{-}})^{*}(t) \, d\varphi(t) > -\int_{0}^{m(E_{-})} (h\chi_{E_{-}})^{*}(t) \, d\varphi(a-t).$$

Assume that

$$\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(t) \ge \int_0^{m(E_-)} (h\chi_{E_-})^*(t) \, d\varphi(t).$$

Then from (18) it follows

$$\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(t) > -\int_0^{m(E_-)} (h\chi_{E_-})^*(t) \, d\varphi(a-t),$$

or

$$\int_{0}^{m(E_{+})} (h\chi_{E_{+}})^{*}(t) \, d\varphi(t) + \int_{0}^{m(E_{-})} (h\chi_{E_{-}})^{*}(t) \, d\varphi(a-t) > 0.$$

As a result, appealing to (14), we conclude that  $||1 + h||_{\Lambda(\varphi)} > 1$ , which contradicts (12).

In the case when

$$\int_0^{m(E_+)} (h\chi_{E_+})^*(t) \, d\varphi(t) \le \int_0^{m(E_-)} (h\chi_{E_-})^*(t) \, d\varphi(t),$$

applying analogous arguments and (15), we conclude that  $||1-h||_{\Lambda(\varphi)} > 1$ , which again contradicts (12). So, the proof is completed.

# **Corollary 11.** $L^1$ is a unique Lorentz space, whose unit ball has no extreme points.

Since  $H(\Lambda(\varphi))$  is isometric to a subspace of the space  $\Lambda(\varphi)$ , we get the following extension of the above-mentioned result of the paper [3] (see Section 1) to the case when, in general,  $\varphi$  fails to be strictly concave.

**Corollary 12.** Suppose  $\varphi$  is a nonlinear, strictly increasing, concave, continuous function on  $[0, 2\pi]$  with  $\varphi(0) = 0$  and  $\varphi(2\pi) = 1$ . Then, each inner function is an extreme point of the unit ball of  $H(\Lambda(\varphi))$ .

Combining Corollary 12 with de Leeuw-Rudin theorem (see Section 1) gives the following compliment to the latter theorem.

**Theorem 13.** Suppose  $\varphi$  is a strictly increasing, concave, continuous function on  $[0, 2\pi]$  such that  $\varphi(0) = 0$ ,  $\varphi(2\pi) = 1$ . The set of extreme points of the unit ball of the space  $H(\Lambda(\varphi))$  coincides with the set of normed outer functions if and only if  $\varphi(t) = t/(2\pi)$ , i.e.,  $H(\Lambda(\varphi)) = H^1$ .

Let us now move on to the second main result of the paper.

Suppose  $\varphi$  is a strictly increasing and strictly concave function on  $[0, 2\pi]$  with  $\varphi(0) = 0$  and  $\varphi(2\pi) = 1$ . Then, according to the main result of the note [3], every inner function is an extreme point of the unit ball of  $H(\Lambda(\varphi))$ . However, as was explained already in Section 1, this is really a very easy consequence of Theorem 3 and the facts that the function  $\tilde{f}(t) := f(e^{it}), 0 \le t \le 2\pi$ , is an extreme point of the unit ball of the space  $\Lambda(\varphi)$  and mapping (2) includes  $H(\Lambda(\varphi))$  into  $\Lambda(\varphi)$  as an isometric subspace.

In the following theorem we show that the same assertion holds (under the same conditions imposed on  $\varphi$ ) for each function  $f \in H(\Lambda(\varphi))$ ,  $||f||_{H(\Lambda(\varphi))} = 1$ , satisfying the much weaker condition that the function  $|\tilde{f}|$  is a constant on some set of positive measure. Note that such a function needs to be neither inner, nor outer.

**Theorem 14.** Let  $\varphi$  be a strictly increasing and strictly concave function on  $[0, 2\pi]$ such that  $\varphi(0) = 0$  and  $\varphi(2\pi) = 1$ . Assume that  $f \in H(\Lambda(\varphi))$ ,  $||f||_{H(\Lambda(\varphi))} = 1$ . Moreover, let the function  $\mu := |\tilde{f}|$  be a constant on some subset of positive measure of the interval  $[0, 2\pi]$ .

Then, f is an extreme point of the unit ball of the space  $H(\Lambda(\varphi))$ .

*Proof.* By the assumption, we have that  $m\{s \in [0, 2\pi] : \mu(s) = c\} > 0$  for some  $c \ge 0$ . To the contrary, assume that f fails to be an extreme point of the unit ball of  $H(\Lambda(\varphi))$ . Thus, there exists a function  $g \in H(\Lambda(\varphi))$ ,  $g \ne 0$ , for which it holds

(19) 
$$\|\tilde{f} \pm \tilde{g}\|_{\Lambda(\varphi)} = \|f \pm g\|_{H(\Lambda(\varphi))} = 1.$$

Since  $\varphi$  is being strictly increasing, the norm of the space  $\Lambda(\varphi)$  is strictly monotone (see Lemma 4). Moreover, Lusin-Privalov uniqueness theorem (see, e.g., [12, §III.3<sup>0</sup>]) combined with the condition  $||f||_{H(\Lambda(\varphi))} = 1$  implies that  $m\{t : \tilde{f}(t) = 0\} = 0$ . Therefore, by Lemma 5, we have  $\tilde{g} = \tilde{f} \cdot h$ , where h is real-valued and  $|h(t)| \leq 1$  a.e. on  $[0, 2\pi]$ . Hence, both functions  $\mu \pm \mu h = \mu(1 \pm h)$  are nonnegative, and from (19) it follows that

(20) 
$$\|\mu(1\pm h)\|_{\Lambda(\varphi)} = 1.$$

Next, according to Proposition 7, there exists a family of subsets  $\{E_t\}_{0 < t \leq 2\pi}$  of the interval  $[0, 2\pi]$  such that

(21) 
$$\int_{E_t} \mu(s) \, ds = \int_0^t \mu^*(s) \, ds, \quad 0 < t \le 2\pi,$$

and

(22) 
$$\int_{E_t} \mu(s)(1\pm h(s)) \, ds = \int_0^t (\mu(1\pm h))^*(s) \, ds, \quad 0 < t \le 2\pi.$$

Combining (22) with  $[11, \S II.2, Property 8^0]$ ), for both plus and minus signs we get:

(23) 
$$\int_{E_t} \mu(s)(1\pm h(s))\,ds = \sup\Big\{\int_A \mu(s)(1\pm h(s))\,ds:\,A\subset[0,2\pi], m(A)=t\Big\}.$$

Let us denote

$$C := \{ s \in [0, 2\pi] : \ \mu(s) = c \}, \ C_{>} := \{ s \in [0, 2\pi] : \ \mu(s) > c \}.$$

Since  $\mu^*$  and  $\mu$  are equimeasurable and the rearrangement  $\mu^*$  decreases, it follows (up to a set of zero measure) that

$$\{s \in [0, 2\pi] : \mu^*(s) = c\} = (a, b),$$

where

$$a = m(C_{>}) = m\{s \in [0, 2\pi] : \mu^*(s) > c\}, \ b = m(C) + a.$$

Let  $t \in (a, b)$  be arbitrary. Since  $\mu^*(t) = c$ , by the definition of the sets C and  $C_>$ , we have

$$\{s \in [0, 2\pi] : \mu(s) = \mu^*(t)\} = C, \ \{s \in [0, 2\pi] : \mu(s) > \mu^*(t)\} = C_>.$$

Consequently, from the fact that  $m(E_t) = t$  and the embeddings

$$\{s \in [0, 2\pi] : \, \mu(s) > \mu^*(t)\} \subset E_t \subset \{s \in [0, 2\pi] : \, \mu(s) \ge \mu^*(t)\}$$

(see (5)) it follows that

(24) 
$$E_t = C_> \cup F_t,$$

where  $F_t \subset C$  and  $m(F_t) = t - a$  (in the case when a = 0 we have  $C_> = \emptyset$ ). Thus, since  $\mu(s) = c$  if  $s \in C$ , for all  $t \in (a, b)$  it holds

(25) 
$$\int_{E_t} \mu(s)(1\pm h(s)) \, ds = \int_{C_>} \mu(s)(1\pm h(s)) \, ds + \int_{F_t} \mu(s)(1\pm h(s)) \, ds \\ = \int_{C_>} \mu(s)(1\pm h(s)) \, ds + c\Big(t-a\pm \int_{F_t} h(s) \, ds\Big).$$

We claim that the function h is constant a.e. on the set C, i.e.,

(26) 
$$h(t) = c_0$$
 a.e. on *C* for some  $c_0 \in [-1, 1]$ .

Assuming that it is not the case, by continuity of the Lebesgue measure, we can find  $t_0 \in (a, b)$  and two subsets F' and F'' of C such that  $m(F') = m(F'') = t_0 - a$  and

(27) 
$$\alpha := \sup_{s \in F''} h(s) < \beta := \inf_{s \in F'} h(s)$$

Let us fix such a  $t_0$ .

Setting  $E' := C_{>} \cup F'$  and  $E'' := C_{>} \cup F''$ , as above, we obtain

(28) 
$$\int_{E'} \mu(s)(1\pm h(s))\,ds = \int_{C_{>}} \mu(s)(1\pm h(s))\,ds + c\Big(t_0 - a \pm \int_{F'} h(s)\,ds\Big)$$

and

(29) 
$$\int_{E''} \mu(s)(1\pm h(s)) \, ds = \int_{C_{>}} \mu(s)(1\pm h(s)) \, ds + c \Big(t_0 - a \pm \int_{F''} h(s) \, ds\Big).$$

Since  $m(E') = t_0$ , from (23) it follows

$$\int_{E_{t_0}} \mu(s)(1+h(s)) \, ds \ge \int_{E'} \mu(s)(1+h(s)) \, ds$$

Hence, comparing the right-hand sides of equations (25) (with  $t = t_0$ ) and (28) for the plus sign and applying (27), we obtain

$$\int_{F_{t_0}} h(s) \, ds \ge \int_{F'} h(s) \, ds \ge \beta(t_0 - a) > \alpha(t_0 - a) \ge \int_{F''} h(s) \, ds.$$

Consequently, by (25) (with  $t = t_0$ ) and (29) for the minus sign,

$$\int_{E_{t_0}} \mu(s)(1-h(s)) \, ds < \int_{C_{>}} \mu(s)(1-h(s)) \, ds + c \Big(t_0 - a - \int_{F''} h(s) \, ds\Big)$$
$$= \int_{E''} \mu(s)(1-h(s)) \, ds.$$

Since  $m(E'') = t_0$ , the last inequality contradicts (23) for  $t = t_0$  and the minus sign. As a result, claim (26) is proved.

In view of (26) and the definition of h, we have that  $\tilde{g} = c_0 \cdot \tilde{f}$  a.e. on C. This implies that the nontangential limit of the analytic function  $g(z) - c_0 f(z)$  as  $|z| \to 1$  vanishes

on some subset of positive measure of the circle T. Hence, applying Lusin-Privalov uniqueness theorem (see e.g. [12, §III.3<sup>0</sup>]) once more, we conclude that  $g(z) = c_0 f(z)$ for all  $z \in \mathbb{D}$ . Hence,  $h = c_0$  a.e. on  $[0, 2\pi]$ . In view of (20), this implies that

$$(1+c_0)||f||_{H(\Lambda(\varphi))} = (1-c_0)||f||_{H(\Lambda(\varphi))} = 1,$$

whence  $c_0 = 0$  or equivalently q = 0, which contradicts the assumption. This completes the proof.

## References

[1] Bennett C., Sharpley R., Interpolation of Operators, Academic Press, Boston, 1988.

- [2] Bryskin I. B., Sedaev A. A., Geometric properties of the unit ball in Hardy-type spaces, J. Soviet Math., 8 (1977), 1–9. https://doi.org/10.1007/BF01455319
- [3] Carrillo-Alanís J., Curbera G. P., A note on extreme points of the unit ball of Hardy-Lorentz spaces, Proc. Amer. Math. Soc., 152 (2024), 2551-2554.
- [4] Carothers N. L., Dilworth S. J. and Trautman D. A., On the geometry of the unit sphere of the Lorentz space  $L_{w,1}$ , Glasgow Math. J., **34** (1992), 21–25.
- Carothers N. L., Turett B., Isometries on  $L_{p,1}$ , Trans. Amer. Math. Soc., **297** (1986), 95–103.
- [6] de Leeuw K., Rudin W., Extreme points and extreme problems in  $H_1$ , Pacific J. Math., 8 (1958), 467 - 485.
- [7] Duren P. L., Theory of H<sup>p</sup> Spaces, Academic Press, New York-London, 1970.
- [8] Dyakonov K. M., Nearly outer functions as extreme points in punctured Hardy spaces, Adv. Math. **401** (2022), 108330, 22 pp.
- [9] Hoffman K., Banach Spaces of Analytic Functions, Prentice-Hall, Englewood Cliffs, NJ, 1962.
- [10] Investigations on linear operators and function theory, 99 unsolved problems in linear and complex analysis, V. P. Khavin, S. V. Khrushchëv, N. K. Nikol'skii, eds., Zap. Nauchn. Sem. LOMI, 81, "Nauka", Leningrad. Otdel., Leningrad, (1978). http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=znsl&paperid=2989
- [11] Krein S. G., Petunin Ju. I., Semenov E. M., Interpolation of Linear Operators, Amer. Math. Soc., Providence, R. I., 1982.
- [12] Koosis P., Introduction to  $H_p$  Spaces. With an Appendix on Wolff's Proof of the Corona Theorem, Cambridge University Press, London-New York, 1980.
- [13] Lindenstrauss L. and Tzafriri L., Classical Banach Spaces, II. Function Spaces. Springer-Verlag, Berlin-New York, 1979.
- [14] Linear and Complex Analysis. Problem Book. 199 Research Problems, V. P. Khavin, S. V. Khrushchëv, N. K. Nikol'skii, eds., Lecture Notes in Math. 1043, Springer-Verlag, Berlin, 1984.
- [15] Linear and Complex Analysis. Problem Book 3. Part I, V. P. Khavin, N. K. Nikol'skii, eds., Lecture Notes in Math. 1573, Springer-Verlag, Berlin, 1994.
- [16] Semenov E. M., 5.1. Hardy-type spaces, J. Soviet Math., 26:5 (1984), 2100 https://doi.org/10.1007/BF01221488

ASTASHKIN: DEPARTMENT OF MATHEMATICS, SAMARA NATIONAL RESEARCH UNIVERSITY, Moskovskoye shosse 34, Samara, 443086, Russian Federation; Saint Petersburg Uni-VERSITY, 7/9 UNIVERSITETSKAYA NAB., SAINT PETERSBURG, 199034, RUSSIAN FEDERATION; BAHCESEHIR UNIVERSITY, 34353, ISTANBUL, TURKEY

Email address: astash56@mail.ru

URL: www.mathnet.ru/rus/person/8713