

On vertex peripherians and Wiener index of graphs with fixed number of cut vertices

Dinesh Pandey

Lazaridis School of Business and Economics, Wilfrid Laurier University, 75 University Avenue West, Waterloo, Ontario, N2L3C5, Canada.

Email: dpandey@wlu.ca

Abstract

The distance of a vertex in a graph is the sum of distances from that vertex to all other vertices of the graph. The Wiener index of a graph is the sum of distances between all its unordered pairs of vertices. A graph has been obtained that contains a vertex achieving the maximum distance among all graphs on n vertices with fixed number of cut vertices. Further the graphs having maximum Wiener index among all graphs on n vertices with at most 3 cut vertices have been characterised.

Key words: cut vertex, distance, Wiener index, extremal problems

AMS subject classification. 05C09, 05C12, 05C35

1 Introduction

Throughout the paper the graphs are simple, finite, connected and undirected. Let G be a graph with vertex set $V(G)$ and the edge set $E(G)$. The edge joining the vertices u and v of G is denoted by uv . A *cut vertex* in G is a vertex whose removal makes the graph disconnected. A vertex of degree one in G is called a *pendant vertex*. An edge containing a pendant vertex of G is called a *pendant edge*. For two isomorphic graphs G_1 and G_2 , we use the notation $G_1 \cong G_2$. The path and the cycle on n vertices are denoted by P_n and C_n , respectively. The complete bipartite graph $K_{1,n-1}$ is called a *star* on n vertices. A *block* in G is a maximal 2 – connected subgraph of G . A *pendant block* of G is a block containing exactly one cut vertex of G . Two blocks of G are *adjacent* if they share a cut vertex. For $u, v \in V(G)$, the distance $d_G(u, v)$ or $d(u, v)$ is the number of edges in a shortest path joining u and v . The *distance* $D_G(v)$ of the vertex v in G is defined as $D_G(v) = \sum_{u \in V(G)} d(v, u)$. A vertex having maximum distance in G is called a *peripherian vertex* of G (see [12], page 93). The *Wiener index* of G is defined as the sum of distances between all its unordered pairs of vertices and denoted by $W(G)$. From this definition it follows that $W(G) = \frac{1}{2} \sum_{v \in V(G)} D_G(v)$. Other terminologies which are used in this article and not defined here can be found in [9].

The Wiener index is the most studied topological index in graph theory. It was introduced by the Chemist H. Wiener in [10] in 1947. In mathematical literature, [5] seems to be the first paper studying Wiener index. The graphs having extremal (maximal or minimal) Wiener index among various classes of graph have been studied extensively in last two decades. Two such recent studies can be seen in [2] and [3]. Over certain classes of graphs, the problem of maximizing the Wiener index seems comparatively difficult than the corresponding minimization problem. We denote the set of all connected graph on n vertices with k cut vertices by $\mathfrak{C}_{n,k}$. In

[7], the graphs having minimum Wiener index in $\mathfrak{C}_{n,k}$ have been characterised. In this paper we find a graph containing a vertex peripherian which attains maximum distance among all vertex peripherians in $\mathfrak{C}_{n,k}$. Further in the obtained graph we identify the vertex peripherians. Using this, we characterise the graphs having maximum Wiener index in $\mathfrak{C}_{n,k}$, $0 \leq k \leq 3$. The problem remains open for graphs having more than 3 cut vertices.

The paper is organised in the following way. In section 2, some results from the literature and some expressions for distance of a vertex and Wiener index of some specific graphs have been presented. In Section 3, special kind of pendant vertices and pendant blocks are introduced which are useful for the study. Section 4 presents the vertex peripherians which achieve maximum distances among all vertex peripherians in $\mathfrak{C}_{n,k}$. In Section 5, the graphs having maximum Wiener index in $\mathfrak{C}_{n,k}$ for $0 \leq k \leq 3$ have been characterised.

2 Preliminaries

The following lemma shows the effect of deleting an edge on the distance of a vertex and on the Wiener index of a graph, and they follow from the definitions.

Lemma 2.1. *Let G be a graph and $e \in E(G)$ such that $G - e$ is connected. Then*

- (i) *for any $v \in V(G)$, $D_{G-e}(v) \geq D_G(v)$.*
- (ii) *$W(G - e) > W(G)$.*

If $|V(G)| = n$, then G has at most $n - 2$ cut vertices and P_n , $n > 2$ is the only graph with $n - 2$ cut vertices. So we consider $\mathfrak{C}_{n,k}$ where $0 \leq k \leq n - 3$. Let G be a graph and w be a cut vertex of G . Then there always exist two subgraphs G_1 and G_2 (both on at least 2 vertices) such that $G \cong G_1 \cup G_2$ and $V(G_1) \cap V(G_2) = \{w\}$. The next lemma is frequently used in counting the distance of a vertex and the Wiener index of a graph, in $\mathfrak{C}_{n,k}$, $k \geq 1$.

Lemma 2.2 ([1], Lemma 1.1). *Let G be a graph and w be a cut vertex in G . Let G_1 and G_2 be two subgraphs of G such that $G \cong G_1 \cup G_2$ and $V(G_1) \cap V(G_2) = \{w\}$. Then*

- (i) *for any $v \in V(G_1)$, $D_G(v) = D_{G_1}(v) + (|V(G_2)| - 1)d(v, w) + D_{G_2}(w)$ and*
- (ii) *$W(G) = W(G_1) + W(G_2) + (|V(G_1)| - 1)D_{G_2}(w) + (|V(G_2)| - 1)D_{G_1}(w)$.*

A graph is called minimally 2-connected if it is 2-connected and deleting any edge gives a graph which is not 2-connected.

Lemma 2.3 ([4], Theorem 2). *A minimally 2-connected graph with more than 3 vertices is triangle free.*

Let G has maximum Wiener index over $\mathfrak{C}_{n,k}$, $n \geq 4$. Then by Lemma 2.1, blocks of G are minimally 2-connected and hence by Lemma 2.3 the blocks of size more than 3 are triangle free. Let B be a block of size 3 in G containing the cut vertex w of G . Then B is isomorphic to a triangle wxy . If B contains exactly one cut vertex w , then $G - xy \in \mathfrak{C}_{n,k}$ and by Lemma 2.1 (ii), $W(G - xy) > W(G)$, a contradiction. If B contains two cut vertices say w and x , then $G - wy \in \mathfrak{C}_{n,k}$ and $W(G - wy) > W(G)$, a contradiction. If all three vertices w, x and y are cut vertices in G , then for any $e \in \{wx, xy, yw\}$, $G - e \in \mathfrak{C}_{n,k}$ and $W(G - e) > W(G)$, a contradiction.

Similarly it can be shown that if $G \in \mathfrak{C}_{n,k}$, $n \geq 4$ and $v_0 \in V(G)$ such that $D_G(v_0) = \max\{D_G(v) : v \in V(G)\}$, then there exists a triangle free graph $G' \in \mathfrak{C}_{n,k}$ obtained by removing some edges (if necessary) from G such that $D_{G'}(v_0) \geq D_G(v_0)$. So, we conclude the following.

Remark 2.4. If G has maximum Wiener index over $\mathfrak{C}_{n,k}$, $n \geq 4$, then G is triangle free.

Remark 2.5. Among all G' satisfying $D_{G'}(v_0) = \max\{D_G(v) : G \in \mathfrak{C}_{n,k}, n \geq 4, v \in V(G)\}$, there exists one which is triangle free.

We now recall the distance of vertices and the Wiener indices of some known graphs which will be used in later sections. First, consider the path $P_n : v_1 v_2 \dots v_n$. Then for the vertex $v_i, 1 \leq i \leq n$,

$$D_{P_n}(v_i) = \frac{i(i-1)}{2} + \frac{(n-i)(n-i-1)}{2} \leq \frac{n(n-1)}{2} = D_{P_n}(v_1) = D_{P_n}(v_n) \quad (1)$$

and

$$W(P_n) = \binom{n+1}{3}.$$

By $L_{n,g}$, we denote the graph in $\mathfrak{C}_{n,n-g}$ obtained by identifying a pendant vertex of P_{n-g+1} with a vertex of C_g (see Figure 1). Note that $L_{n,n-g}$ has g cut vertices. The graph C_{m_1, m_2}^n is defined for $n \geq m_1 + m_2 - 1$ and $m_1, m_2 \geq 3$ in [7]. For $n \geq m_1 + m_2$, C_{m_1, m_2}^n is the graph obtained by identifying one pendant vertex of the path $P_{n+2-(m_1+m_2)}$ with a vertex of C_{m_1} and the other pendant vertex of $P_{n+2-(m_1+m_2)}$ with a vertex of C_{m_2} . For $n = m_1 + m_2 - 1$, C_{m_1, m_2}^n is the graph obtained by identifying a vertex of C_{m_1} with a vertex of C_{m_2} (see Figure 2). Note that C_{m_1, m_2}^n has $n + 2 - (m_1 + m_2)$ cut vertices.




Figure 1: The graph $L_{n,g}$




Figure 2: The graphs C_{m_1, m_2}^n

Lemma 2.6 ([7], Lemma 3.5). Let $m_1, m_2 \geq 3$ be two integers and let $n = m_1 + m_2 - 1$. Then $W(C_n) > W(C_{m_1, m_2}^n)$.

The graphs $L_{n,g}$ and C_{m_1, m_2}^n are important for us. So we recall the known expressions for the distance of a vertex in these graphs and for their Wiener indices. Wiener index of $L_{n,g}$ can be found in [11] (see Theorem 1.1) as following.

$$W(L_{n,g}) = \begin{cases} \frac{g^3}{8} + (n-g)\left(\frac{n^2+ng+3g-1}{6} - \frac{g^2}{12}\right) & \text{if } g \text{ is even,} \\ \frac{g(g^2-1)}{8} + (n-g)\left(\frac{n^2+ng+3g-1}{6} - \frac{g^2}{12} - \frac{1}{4}\right) & \text{if } g \text{ is odd.} \end{cases} \quad (2)$$

Also the distance of the pendant vertex v in $L_{n,g}$ can be computed using Lemma 2.2 (i) as

$$D_{L_{n,g}}(v) = \begin{cases} \frac{g^2}{4} + \frac{(n-g)(n+g-1)}{2} & \text{if } g \text{ is even,} \\ \frac{g^2-1}{4} + \frac{(n-g)(n+g-1)}{2} & \text{if } g \text{ is odd.} \end{cases} \quad (3)$$

The Wiener index of C_{m_1, m_2}^n when $n = m_1 + m_2 - 1$ is computed in [7] (see proof of Lemma 3.5 in [7]). We generalise that by counting the Wiener index of $C_{m_1, m_2}^n \in \mathfrak{C}_{n, k}$ i.e. when $n = m_1 + m_2 + k - 2$. Let w be the cut vertex of C_{m_1, m_2}^n lying in C_{m_1} . Using Lemma 2.2 (ii)

$$W(C_{m_1, m_2}^n) = W(C_{m_1}) + W(L_{m_2+k-1, m_2}) + (m_2 + k - 2)D_{C_{m_1}}(w) + (m_1 - 1)D_{L_{m_2+k-1, m_2}}(w).$$

From (2), we get

$$W(L_{m_2+k-1, m_2}) = \begin{cases} \frac{1}{24}(3m_2^3 + 6m_2^2k + 4k^3 + 12m_2k^2 - 12k^2 - 6m^2 - 12m_2k + 8k) & m_2 \text{ even,} \\ \frac{1}{24}(3m_2^3 + 6m_2^2k + 4k^3 + 12m_2k^2 - 12k^2 - 6m^2 - 12m_2k - 3m_2 + 2k + 6) & m_2 \text{ odd.} \end{cases}$$

Further, from (3), we get

$$D_{L_{m_2+k-1, m_2}}(w) = \begin{cases} \frac{1}{4}(m_2^2 + 4m_2k + 2k^2 - 6k - 4m_2 + 4) & m_2 \text{ even,} \\ \frac{1}{4}(m_2^2 + 4m_2k + 2k^2 - 6k - 4m_2 + 3) & m_2 \text{ odd.} \end{cases}$$

Performing some intricate calculations, the Wiener index of C_{m_1, m_2}^n can be obtained as follows.

$$\begin{aligned} & W(C_{m_1, m_2}^n) \\ = & \begin{cases} \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2k + 2m_2^2k + 4m_1k^2 + 4m_2k^2 + 8m_1m_2k \\ - 4m_1^2 - 4m_2^2 - 8m_1m_2 - 12m_1k - 12m_2k - 8k^2 + 8m_1 + 8m_2 - 8) + \frac{1}{6}(k^3 + 11k) & \text{both } m_1, m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2k + 2m_2^2k + 4m_1k^2 + 4m_2k^2 + 8m_1m_2k \\ - 4m_1^2 - 4m_2^2 - 8m_1m_2 - 12m_1k - 12m_2k - 8k^2 + 6m_1 + 7m_2 - 4) + \frac{1}{12}(2k^3 + 19k) & m_1 \text{ even, } m_2 \text{ odd,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2k + 2m_2^2k + 4m_1k^2 + 4m_2k^2 + 8m_1m_2k \\ - 4m_1^2 - 4m_2^2 - 8m_1m_2 - 12m_1k - 12m_2k - 8k^2 + 7m_1 + 6m_2 - 4) + \frac{1}{12}(2k^3 + 19k) & m_1 \text{ odd, } m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2k + 2m_2^2k + 4m_1k^2 + 4m_2k^2 + 8m_1m_2k \\ - 4m_1^2 - 4m_2^2 - 8m_1m_2 - 12m_1k - 12m_2k - 8k^2 + 5m_1 + 5m_2) + \frac{1}{8}(k^3 + 8k) & \text{both } m_1, m_2 \text{ odd.} \end{cases} \end{aligned} \quad (4)$$

3 s-pendant vertices and s-pendant blocks

We introduce a special kind of pendant blocks and pendant vertices in $\mathfrak{C}_{n, k}$, $k \geq 2$, which are used later in Section 5.

Definition 3.1. Let G be a graph with at least two cut vertices. An **s-pendant block** of G is a pendant block which shares its cut vertex with exactly one non-pendant block of G . If size of an s-pendant block is 2, then it is called an **s-pendant edge**. A pendant vertex lying on an s-pendant edge is called an **s-pendant vertex**.

In Figure 3, B_2 and e_4 are non pendant blocks. B_1 is an s-pendant block, v_3 is an s-pendant vertex and e_3 is an s-pendant edge. B_3 is a pendant block but not s-pendant, v_1, v_2 are pendant vertices but not s-pendant and e_1, e_2 are pendant edges but not s-pendant.

Let G be a graph with $k \geq 2$ cut vertices. Let w be a cut vertex and B_1, B_2, \dots, B_r be all the s-pendant blocks sharing the cut vertex w . Then the graph obtained from G by detaching all the s-pendant blocks from w i.e. $(G \setminus \cup_{i=1}^r B_i) \cup w$ has $k - 1$ cut vertices. The distance $d_G(B, B')$ or $d(B, B')$ between two blocks B and B' of G is defined as $d_G(B, B') = \min\{d(u, v) : u \in V(B), v \in V(B')\}$.

Lemma 3.2 ([8], Lemma 2.5). *If $d(B_1, B_2) = \max\{d(B, B') : B, B' \text{ are blocks of } G\}$, then both B_1 and B_2 are pendant blocks in G .*




Figure 3: s-pendant blocks and s-pendant vertices in a graph

Proposition 3.3. *Let $G \in \mathfrak{C}_{n,k}$, $k \geq 2$. Then G has at least two s-pendant blocks.*

Proof. Let B_1, B_2 be two blocks in G such that $d(B_1, B_2) = \max\{d(B, B') : B, B' \text{ are blocks of } G\}$. By Lemma 3.2, B_1 and B_2 are pendant blocks. We claim that both B_1 and B_2 are s-pendant blocks. Suppose B_1 is not s-pendant. Let $d(B_1, B_2) = d(w_1, w_2)$ where w_1 and w_2 are cut vertices lying in B_1 and B_2 , respectively. Let $P_{w_1 w_2}$ be a corresponding path joining w_1 and w_2 . As B_1 is not s-pendant there exist a non-pendant block B sharing the cut vertex w_1 with B_1 such that no vertex of B other than w_1 lies on $P_{w_1 w_2}$. Also as B is non-pendant, B is adjacent to a block B' disjoint from $P_{w_1 w_2}$. Let B and B' are adjacent via the cut vertex w' . Then $d(B_2, B') = d(w_2, w_1) + d(w_1, w') > d(B_1, B_2)$, a contradiction. Hence B_1 is s-pendant. Similarly B_2 is s-pendant. \square

4 Vertex peripherians attaining maximum distance in $\mathfrak{C}_{n,k}$

This section focuses on finding a graph that exhibit the vertex peripherians having maximum distance within the class $\mathfrak{C}_{n,k}$. For graphs devoid of cut vertices, the following is known.

Lemma 4.1 ([6], Lemma 2). *Let $G \in \mathfrak{C}_{n,0}$, $n \geq 3$ and $u \in V(G)$. Then for any $v \in V(C_n)$, $D_{C_n}(v) \geq D_G(u)$. Moreover*

$$D_{C_n}(v) = \begin{cases} \frac{n^2}{4} & \text{if } n \text{ is even,} \\ \frac{n^2-1}{4} & \text{if } n \text{ is odd.} \end{cases}$$

We generalise Lemma 4.1 for graphs with k cut vertices i.e. find a $G_0 \in \mathfrak{C}_{n,k}$ and $v_0 \in V(G_0)$ such that $D_{G_0}(v_0) = \max\{D_G(v) : G \in \mathfrak{C}_{n,k}, v \in V(G)\}$.

Lemma 4.2 ([7], Lemma 3.6). *Let u be the pendant vertex and v be a non-pendant vertex of $L_{n,n-k}$, $k \geq 1$. Then $D_{L_{n,n-k}}(u) > D_{L_{n,n-k}}(v)$.*

Lemma 4.3. *Let $G \in \mathfrak{C}_{n,k}$, $n \geq 5$ and $k \geq 1$. Let B be a pendant block in G with $|V(B)| = m \geq 4$ containing the cut vertex w . Let G' be the graph obtained from G by replacing B by C_m such that w remains a cut vertex of G' in C_m . Then for any $v \in V(G)$, there exists a $v' \in V(G')$ such that $D_{G'}(v') \geq D_G(v)$. In particular, if $v \in V(G) \setminus V(B) \cup \{w\}$, then $D_{G'}(v) \geq D_G(v)$.*

Proof. Let H be the subgraph of G induced by $V(G) \setminus V(B) \cup \{w\}$. Then $G \cong H \cup B$ with $V(H) \cap V(B) = w$ and $G' \cong H \cup C_m$ with $V(H) \cap V(C_m) = w$.

If $v \in V(H)$, then by Lemma 2.2 (i) we have

$$\begin{aligned} D_G(v) &= D_H(v) + (m-1)d(v, w) + D_B(w) \\ &\leq D_H(v) + (m-1)d(v, w) + D_{C_m}(w) && \text{[using Lemma 4.1]} \\ &= D_{G'}(v). \end{aligned}$$

If $v \in V(B)$, take v' to be the vertex in the cycle C_m of G' such that $d(v', w) = \lfloor \frac{m}{2} \rfloor$. Then

$$\begin{aligned}
D_G(v) &= D_B(v) + (|V(H)| - 1)d(v, w) + D_H(w) \\
&\leq D_{C_m}(v') + (|V(H)| - 1)\left\lfloor \frac{m}{2} \right\rfloor + D_H(w) \\
&= D_{C_m}(v') + (|V(H)| - 1)d(v', w) + D_H(w) \\
&= D_{G'}(v').
\end{aligned}$$

This completes the proof. \square

Proposition 4.4. *For $n \geq 4$ and $k \geq 1$ there exists a triangle free graph $G_0 \in \mathfrak{C}_{n,k}$ with all its pendant blocks are either K_2 or a cycle on at least 4 vertices such that $D_{G_0}(v_0) = \max\{D_G(v) : G \in \mathfrak{C}_{n,k}, k \geq 1, v \in V(G)\}$.*

Proof. Let $D_{G'}(v') = \max\{D_G(v) : G \in \mathfrak{C}_{n,k}, k \geq 1, v \in V(G)\}$. By Remark 2.5, we may assume that G' is triangle free. Replace each pendant blocks B of G' which is not cyclic by a cycle of size $|B|$. Let the resulting graph be G_0 . By Lemma 4.3, there exists a $v_0 \in V(G_0)$ such that $D_{G_0}(v_0) \geq D_{G'}(v')$. Since $G_0 \in \mathfrak{C}_{n,k}$ and $D_{G'}(v') = \max\{D_G(v) : G \in \mathfrak{C}_{n,k}, k \geq 1, v \in V(G)\}$, $D_{G_0}(v_0) = D_{G'}(v')$. \square

Lemma 4.5. *Let H be a graph containing a vertex w and $m_1, m_2 \geq 3$ be two integers such that $m_1 + m_2 - 1 = m$. Let G_1 be the graph obtained by identifying the cut vertex of C_{m_1, m_2}^m with w , G_2 be the graph obtained by identifying the cut vertex of $L_{m, m-1}$ with w and G be the graph obtained by identifying a vertex of C_m with w . Then for any $v \in V(H)$, $D_G(v) > D_{G_1}(v)$ and $D_G(v) > D_{G_2}(v)$.*

Proof.

$$\begin{aligned}
D_{G_1}(v) &= D_H(v) + (m-1)d(v, w) + D_{C_{m_1, m_2}^m}(w) \\
&\leq D_H(v) + (m-1)d(v, w) + \frac{m_1^2 + m_2^2}{4} \quad [\text{equality holds if both } m_1, m_2 \text{ are even}] \\
&< D_H(v) + (m-1)d(v, w) + \frac{(m_1 + m_2 - 1)^2 - 1}{4} \\
&\leq D_H(v) + (m-1)d(v, w) + D_{C_m}(w) \\
&= D_G(v)
\end{aligned}$$

and

$$\begin{aligned}
D_{G_2}(v) &= D_H(v) + (m-1)d(v, w) + D_{L_{m, m-1}^m}(w) \\
&\leq D_H(v) + (m-1)d(v, w) + \frac{(m-1)^2}{4} + 1 \quad [\text{equality holds if } m \text{ is odd}] \\
&< D_H(v) + (m-1)d(v, w) + \frac{m^2 - 1}{4} \quad [\text{for } m \geq 4] \\
&\leq D_G(v).
\end{aligned}$$

\square

Lemma 4.6. *Let v_0 be the pendant vertex of $L_{n, n-k}$ and let $n = m_1 + m_2 + k - 2$, $m_1, m_2 \geq 3$ and $k \geq 1$. Then $D_{L_{n, n-k}}(v_0) > D_{C_{m_1, m_2}^n}(v)$ for any $v \in V(C_{m_1, m_2}^n)$.*

Proof. First suppose $k \geq 2$. Let w and w' be the cut vertices of C_{m_1, m_2}^n lying in C_{m_1} and C_{m_2} , respectively. Let $P_k : w = v_1 v_2 \cdots v_k = w'$ be the path in C_{m_1, m_2}^n joining w and w' . Then $V(C_{m_1}), V(C_{m_2})$ and $V(P) \setminus \{w, w'\}$ partition $V(C_{m_1, m_2}^n)$. First suppose v belongs to the part $V(C_{m_1})$. Then

$$\begin{aligned} D_{C_{m_1, m_2}^n}(v) &= D_{C_{m_1}}(v) + (m_2 + k - 2)d(v, w) + D_{L_{m_2+k-1, m_2}}(w) \\ &\leq \frac{m_1^2}{4} + (m_2 + k - 2)\frac{m_1}{2} + \frac{m_2^2}{4} + \frac{(k-1)(2m_2 + k - 2)}{2} \\ &= \frac{1}{4}(m_1^2 + m_2^2 + 2m_1m_2 + 2m_1k + 4m_2k + 2k^2 - 4m_1 - 4m_2 - 6k + 4). \end{aligned}$$

Similarly if v belongs to the part $V(C_{m_2})$, we get

$$D_{C_{m_1, m_2}^n}(v) \leq \frac{1}{4}(m_1^2 + m_2^2 + 2m_1m_2 + 4m_1k + 2m_2k + 2k^2 - 4m_1 - 4m_2 - 6k + 4).$$

Finally suppose $v \in V(P) \setminus \{w, w'\}$. Let $v = v_i, 2 \leq i \leq k-1$. W.L.O.G. assume that $m_1 \leq m_2$. Then we have

$$\begin{aligned} D_{C_{m_1, m_2}^n}(v) &= D_{L_{m_2+k-1, m_2}}(v_i) + (i-1)(m_1-1) + D_{C_{m_1}}(w) \\ &= D_{P_k}(v_i) + (k-i)(m_2-1) + D_{C_{m_2}}(w') + (i-1)(m_1-1) + D_{C_{m_1}}(w) \\ &\leq \frac{k(k-1)}{2} + (m_2-1)(k-1) + \frac{m_2^2}{4} + \frac{m_1^2}{4} && \text{[using 1]} \\ &= \frac{1}{4}(m_1^2 + m_2^2 + 4m_2k + 2k^2 - 4m_2 - 6k + 4). \end{aligned}$$

Further, we have

$$\begin{aligned} D_{L_{n, n-k}}(v_0) &\geq \frac{(n-k)^2 - 1}{4} + \frac{k(2n-k-1)}{2} && \text{[from (3)]} \\ &= \frac{1}{4}(m_1^2 + m_2^2 + 2m_1m_2 + 4m_1k + 4m_2k + 2k^2 - 4m_1 - 4m_2 - 10k + 3). \end{aligned}$$

By comparing, we get $D_{L_{n, n-k}}(v_0) > D_{C_{m_1, m_2}^n}(v)$ for any $v \in V(C_{m_1, m_2}^n)$.

Now suppose $k = 1$. Then $w = w'$ and $V(P) \setminus \{w, w'\} = \emptyset$. By similar calculations as above it follows that $D_{L_{n, n-k}}(v_0) > D_{C_{m_1, m_2}^n}(v)$ for any $v \in V(C_{m_1, m_2}^n)$. \square

Lemma 4.7. *If $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n, k}, k \geq 1\}$, then v_0 must lie in some pendant block of G_0 . Furthermore, if $n \geq 5$, then the pendant block containing v_0 shares its cut vertex with exactly one other block.*

Proof. Let $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n, k}, k \geq 1\}$. We consider the following two cases.

Case I: $k = 1$

In this case every block of G_0 is a pendant block and hence v_0 lies in a pendant block. Suppose $n \geq 5$ and G_0 has more than two blocks. Let B_1, B_2, \dots, B_s be the blocks where $s \geq 3$ and $|V(B_i)| = m_i$ for $1 \leq i \leq s$. Assume that $v_0 \in V(B_1)$ and w is the cut vertex of G_0 . If $m_i = 2$ for $2 \leq i \leq s$, then $n = m_1 + s - 1$ and

$$\begin{aligned} D_{G_0}(v_0) &= D_{B_1}(v_0) + (s-1)d(v_0, w) + s - 1 \\ &\leq \frac{m_1^2}{4} + (s-1)\left(\frac{m_1}{2} + 1\right) && \text{[by Lemma 4.1]} \\ &= \frac{1}{4}(m_1^2 + 2m_1s + 4s - 2m_1 - 4) \\ &\leq D_{C_{m_1, s}^n}(v_0) && \text{[equality holds for } s = 3\text{]} \\ &< D_{L_{n, n-1}}(v'_0) && \text{[by Lemma 4.6]} \end{aligned}$$

where v'_0 is the pendant vertex of $L_{n,n-1}$. This contradicts that $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n,k}, k \geq 1\}$.

Other possibility is $m_i \geq 3$ for some $2 \leq i \leq s$. For $2 \leq i \leq s$, if $m_i \geq 3$ and $B_i \not\cong C_{m_i}$, then replace B_i by C_{m_i} such that w remains a cut vertex in the new graph. Let the resulting graph be G'_0 . Then by Lemma 4.3, it follows that $D_{G'_0}(v_0) = D_{G_0}(v_0)$. W. L. O. G. assume that $m_2 \geq 3$. Let H' be the subgraph of G'_0 corresponding to $B_2 \cup B_3$ in G_0 . Then $H' \cong C_{m_2+m_3-1}^{m_2, m_3}$ or $H' \cong L_{m_2+m_3-1, m_2+m_3-2}$ and $G'_0 \cong H \cup H'$ such that $V(H) \cap V(H') = \{w\}$. The subgraph H contains B_1 . Construct a new graph G''_0 from G'_0 by replacing H' by $C_{m_2+m_3-1}$. Then by Lemma 4.5, $D_{G''_0}(v_0) > D_{G'_0}(v_0) = D_{G_0}(v_0)$, which is a contradiction. Hence B_1 shares its cut vertex with exactly one other block.

Case II: $k \geq 2$

Suppose v_0 lies in a non-pendant block B of G_0 . Then there are at least two cut vertices w and w' of G_0 in B . Assume $d(v_0, w) \geq d(v_0, w')$. G_0 has two subgraphs G_1 and G_2 such that $G_0 \cong G_1 \cup G_2$ with $V(G_1) \cap V(G_2) = \{w'\}$. Assume that B lies in G_1 . Let $z \in V(G_1)$ such that $d(v_0, z) = \max\{d(v_0, v) : v \in V(G_1)\}$. Note that z can not be a cut vertex of G_0 and $d(v_0, w) < d(v_0, z)$. Construct a new graph $G'_0 \in \mathfrak{C}_{n,k}$ from G_1 and G_2 by identifying z and w' (of G_2). i.e. $G'_0 \cong G_1 \cup G_2$ and $V(G_1) \cap V(G_2) = \{z\}$. By Lemma 2.2 (i),

$$\begin{aligned} D_{G_0}(v_0) &= D_{G_1}(v_0) + (|V(G_2)| - 1)d(v_0, w') + D_{G_2}(w') \\ &< D_{G_1}(v_0) + (|V(G_2)| - 1)d(v_0, z) + D_{G_2}(w') \\ &= D_{G'_0}(v_0). \end{aligned}$$

which is a contradiction. So B must be pendant.

Let w be the cut vertex of G_0 in B . Suppose the cut vertex w is shared by at least two other blocks B_1 and B_2 . Then there exist two subgraphs G_1 and G_2 of G_0 such that $G_0 \cong G_1 \cup G_2$ and $V(G_1) \cap V(G_2) = \{w\}$ where G_1 contains B and has at least 2 cut vertices. Let w_1 be a cut vertex in $G_1 \setminus B$ at maximum distance from v_0 . Construct $G'_0 \in \mathfrak{C}_{n,k}$ from G_1 and G_2 by identifying w_1 and w (of G_2) i.e. $G'_0 \cong G_1 \cup G_2$ and $V(G_1) \cap V(G_2) = \{w_1\}$. Then by Lemma 2.2 (i), it follows that $D_{G'_0}(v_0) > D_{G_0}(v_0)$, which is a contradiction. Hence B shares its cut vertex with exactly one other block. This completes the proof. \square

Theorem 4.8. *Let z be the pendant vertex of $L_{n,n-k}$ and $G \in \mathfrak{C}_{n,k}$, $1 \leq k \leq n - 3$. Then $D_G(v) \leq D_{L_{n,n-k}}(z)$ for any $v \in V(G)$.*

Proof. We use induction on the number of cut vertices k .

Base step: $\mathfrak{C}_{4,1} = \{L_{4,3}, K_{1,3}\}$ and the result can easily be verified for $n = 4$. Now suppose $n \geq 5$. By Proposition 4.4, there exists a triangle free graph $G_0 \in \mathfrak{C}_{n,1}$ having its blocks either K_2 or cycles on at least 4 vertices such that $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n,1}\}$. By Lemma 4.7, either $G_0 \cong C_{m_1, m_2}^n$ for some m_1, m_2 satisfying $m_1 + m_2 - 1 = n$ or $G_0 \cong L_{n, n-1}$. But by Lemma 4.6, $D_{L_{n, n-1}}(z) > D_{C_{m_1, m_2}^n}(v)$ for any $v \in V(C_{m_1, m_2}^n)$ where z is the pendant vertex of $L_{n, n-1}$. Hence $G_0 \cong L_{n, n-1}$. Now by Lemma 4.2, v_0 is the pendant vertex of $G_0 (\cong L_{n, n-1})$. So the result is true for $k = 1$.

Induction hypothesis: Let the result be true for $G \in \mathfrak{C}_{n, k-1}$, $k \geq 2$ and $n \geq k + 2$.

Induction step: By Proposition 4.4, there exists a triangle free graph $G_0 \in \mathfrak{C}_{n,k}$ having each of its pendant blocks is either K_2 or a cycle on at least 4 vertices such that $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n,k}\}$. It is sufficient to show that $D_{L_{n, n-k}}(z) = D_{G_0}(v_0)$. As $k \geq 2$, $n \geq 5$. By Lemma 4.7, v_0 belongs to some pendant block B of G_0 and B shares its cut vertex with exactly

one other block. Let w be the cut vertex of G_0 in B . Then $G_0 \cong B \cup H$ with $V(B) \cap V(H) = \{w\}$. Let $|V(B)| = n_1$ and $|V(H)| = n_2 = n - n_1 + 1$. Then $H \in \mathfrak{C}_{n_2, k-1}$. So $n_2 \geq k + 1$.

Suppose $n_2 = k + 1$, then H is a path and $B \cong C_{n_1, n_1} \geq 4$. So $G_0 \cong L_{n, n_1}$ and we get $D_{L_{n, n_1}}(v_0) > D_{L_{n, n_1}}(z)$, where v_0 is a non pendant vertex of L_{n, n_1} . This is a contradiction to Lemma 4.2. So $n_2 \geq k + 2$. Suppose w as a vertex of H is not the pendant vertex of L_{n_2, n_2-k+1} (H may be isomorphic to L_{n_2, n_2-k+1}). Construct a new graph $G'_0 \in \mathfrak{C}_{n, k}$ from G_0 by replacing H by L_{n_2, n_2-k+1} such that $G'_0 \cong B \cup L_{n_2, n_2-k+1}$ and $V(B) \cap V(L_{n_2, n_2-k+1}) = \{w\}$ where w is the pendant vertex of L_{n_2, n_2-k+1} . Then by induction hypothesis $D_{L_{n_2, n_2-k+1}}(w) \geq D_H(w)$. By Lemma 2.2 we have

$$\begin{aligned} D_{G'_0}(v_0) &= D_B(v_0) + (n_2 - 1)d(v_0, w) + D_{L_{n_2, n_2-k+1}}(w) \\ &\geq D_B(v_0) + (n_2 - 1)d(v_0, w) + D_H(w) \\ &= D_{G_0}(v_0). \end{aligned}$$

As $D_{G_0}(v_0) = \max\{D_G(v) : v \in V(G), G \in \mathfrak{C}_{n, k}\}$, $D_{G'_0}(v_0) = D_{G_0}(v_0)$.

We now claim that $B \cong K_2$. Suppose $B \not\cong K_2$. Then $B \cong C_{n_1, n_1} \geq 4$ and $G'_0 \cong C_{n_1, n_3}^n$ where $n_3 = n_2 - k + 1$. By Lemma 4.6, $D_{G'_0}(v_0) < D_{L_{n, n-k}}(u_0)$ where u_0 is the pendant vertex of $L_{n, n-k}$, which is a contradiction. Therefore $B \cong K_2$ and hence $G'_0 \cong L_{n, n-k}$. Now by Lemma 4.2, v_0 is the pendant vertex z of $L_{n, n-k}$. Thus $D_{L_{n, n-k}}(z) = D_{G'_0}(v_0) = D_{G_0}(v_0)$. This completes the induction step and hence the proof. \square

5 Maximum Wiener index over $\mathfrak{C}_{n, k}$

Let $G \in \mathfrak{C}_{n, 0}$. Then G is a 2-connected graph and we have the following characterisation of graphs having maximum Wiener index over 2-connected graphs.

Lemma 5.1 ([6], Theorem 5). *Let $G \in \mathfrak{C}_{n, 0}$, $n \geq 3$. Then $W(G) \leq W(C_n)$ and equality holds if and only if $G \cong C_n$. Furthermore,*

$$W(C_n) = \begin{cases} \frac{n^3}{8} & \text{if } n \text{ is even,} \\ \frac{n(n^2-1)}{8} & \text{if } n \text{ is odd.} \end{cases}$$

Lemma 5.2. *Let G be a graph in $\mathfrak{C}_{n, k}$, $k \geq 1$ having maximum Wiener index. Then every pendant block of G is either K_2 or a cycle on at least 4 vertices.*

Proof. Let B be a pendant block in G and let w be the cut vertex of G in B . Then there exists a subgraph H (on at least 2 vertices) of G such that $G = H \cup B$ with $V(H) \cap V(B) = \{w\}$. Suppose $B \not\cong K_2$. Then by Remark 2.4, $|V(B)| \geq 4$. Let $|V(B)| = m$. Suppose $B \not\cong C_m$. Construct a graph G' from G by replacing B by C_m . Then using Lemma 2.2 (ii), we get

$$\begin{aligned} W(G) &= W(H) + W(B) + (|V(H)| - 1)D_B(w) + (m - 1)D_H(w) \\ &\text{using Lemma 4.1 and Lemma 5.1} \\ &< W(H) + W(C_m) + (|V(H)| - 1)D_{C_m}(w) + (m - 1)D_H(w) \\ &= W(G'), \end{aligned}$$

which is a contradiction. This completes the proof. \square

Lemma 5.3. *Let $m_1, m_2 \geq 4$ be two integers such that $n = m_1 + m_2 - 1$. Then*

$$W(C_n) \geq W(L_{n, n-1}) > W(C_{m_1, m_2}^n).$$

Proof. By (4),

$$W(C_{m_1, m_2}^n) = \begin{cases} \frac{1}{8}(m_1^3 + m_2^3 + 2m_2m_1^2 + 2m_1m_2^2 - 2m_1^2 - 2m_2^2) & \text{both } m_1, m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 - 2m_1^2 - 2m_2^2 - 2m_1 - m_2 + 2) & m_1 \text{ even, } m_2 \text{ odd,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 - 2m_1^2 - 2m_2^2 - m_1 - 2m_2 + 2) & m_1 \text{ odd, } m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_2m_1^2 + 2m_1m_2^2 - 2m_1^2 - 2m_2^2 - 3m_1 - 3m_2 + 4) & \text{both } m_1, m_2 \text{ odd.} \end{cases}$$

$$\leq \frac{1}{8}(m_1^3 + m_2^3 + 2m_2m_1^2 + 2m_1m_2^2 - 2m_1^2 - 2m_2^2).$$

From (2) we find

$$W(L_{n, n-1}) = \begin{cases} \frac{1}{8}(n^3 - n^2 + 6n - 8) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^3 - n^2 + 7n - 7) & \text{if } n \text{ is odd.} \end{cases}$$

$$\geq \frac{1}{8}(n^3 - n^2 + 6n - 8)$$

$$= \frac{1}{8}(m_1^3 + m_2^3 + 3m_1^2m_2 + 3m_1m_2^2 - 4m_1^2 - 4m_2^2 - 8m_1m_2 + 11m_1 + 11m_2 - 16).$$

So we get

$$\begin{aligned} W(L_{n, n-1}) - W(C_{m_1, m_2}^n) &\geq m_1^2m_2 + m_1m_2^2 + 11m_1 + 11m_2 - 2(m_1^2 + m_2^2 + 4m_1m_2 + 8) \\ &= m_1m_2(m_1 + m_2) + 11(m_1 + m_2) - 2(m_1 + m_2)^2 - 4m_1m_2 - 16 \\ &= m_1m_2(m_1 + m_2 - 4) + 11(m_1 + m_2) - 2(m_1 + m_2)^2 - 16 \\ &\geq 2(m_1 + m_2)(m_1 + m_2 - 4) + 11(m_1 + m_2) - 2(m_1 + m_2)^2 - 16 \end{aligned}$$

(This inequality holds because $m_1 + m_2 - 4 > 0$ and $m_1m_2 \geq 2(m_1 + m_2)$ for $m_1, m_2 \geq 4$)

$$\begin{aligned} &= 3(m_1 + m_2) - 16 \\ &> 0 \end{aligned}$$

Further we have

$$W(C_n) - W(L_{n, n-1}) = \begin{cases} \frac{1}{8}(n^2 - 6n + 8) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^2 - 8n + 7) & \text{if } n \text{ is odd.} \end{cases}$$

$$\geq 0 \quad (> 0 \text{ for } n \geq 8).$$

This completes the proof. \square

Proposition 5.4. *Let G has maximum Wiener index over $\mathfrak{C}_{n, k}, n \geq 7$. Then a cut vertex of G is shared by at most two pendant blocks. Furthermore, for $k \geq 2$, if a cut vertex is shared by two pendant blocks then both of them are K_2 .*

Proof. Let B_1 and B_2 be two pendant blocks of G sharing the cut vertex w such that $|V(B_1)| = m_1$ and $|V(B_2)| = m_2$. We consider the following two cases.

Case-I: $n = m_1 + m_2 - 1$

By Lemma 5.2, G is isomorphic to either $L_{n, n-1}$ or to C_{m_1, m_2}^n . By Lemma 5.3, $W(L_{n, n-1}) > W(C_{m_1, m_2}^n)$ and the result follows.

Case II: $n > m_1 + m_2 - 1$

In this cases there exists a subgraph H (on at least two vertices) of G such that $G \cong H \cup (B_1 \cup B_2)$ and $V(H) \cap V(B_1 \cup B_2) = \{w\}$. First suppose both $m_1, m_2 > 2$. Then by Lemma 5.2, $m_1, m_2 \geq 4$

and $B_1 \cup B_2 \cong C_{m_1, m_2}^{m_1+m_2-1}$. Construct a graph G' from G by replacing $B_1 \cup B_2$ by $C_{m_1+m_2-1}$ such that w remains a cut vertex shared by H and $C_{m_1+m_2-1}$. Then

$$\begin{aligned} W(G) &= W(H) + W(C_{m_1, m_2}^{m_1+m_2-1}) + (m_1 + m_2 - 2)D_H(w) + (|V(H)| - 1)D_{C_{m_1, m_2}^{m_1+m_2-1}}(w) \\ &\text{using Lemma 5.3} \\ &< W(H) + W(C_{m_1+m_2-1}) + (m_1 + m_2 - 2)D_H(w) + \frac{1}{4}(|V(H)| - 1)(m_1^2 + m_2^2) \\ &< W(H) + W(C_{m_1+m_2-1}) + (m_1 + m_2 - 2)D_H(w) + \frac{1}{4}(|V(H)| - 1)D_{C_{m_1+m_2-1}}(w) \\ &= W(G'), \end{aligned}$$

which is a contradiction.

Now suppose exactly one of m_1 or m_2 is greater than 2. Let $m_1 > 2$ and $m_2 = 2$. Then by Lemma 5.2, $B_1 \cup B_2 \cong L_{m_1+1, m_1}$ and $G \cong H \cup L_{m_1+1, m_1}$ such that $V(H) \cup V(L_{m_1+1, m_1}) = \{w\}$, where w is the cut vertex of L_{m_1+1, m_1} . Construct G' from G by replacing $B_1 \cup B_2$ by C_{m_1+1} such that $V(H) \cap V(C_{m_1+1}) = \{w\}$. Then

$$\begin{aligned} W(G) &= W(H) + W(L_{m_1+1, m_1}) + m_1 D_H(w) + (|V(H)| - 1)D_{L_{m_1+1, m_1}}(w) \\ &\text{using Lemma 5.3} \\ &\leq W(H) + W(C_{m_1+1}) + m_1 D_H(w) + \frac{1}{4}(|V(H)| - 1)(m_1^2 + 4) \\ &< W(H) + W(C_{m_1+1}) + m_1 D_H(w) + \frac{1}{4}(|V(H)| - 1)((m_1 + 1)^2 - 1) \\ &\leq W(H) + W(C_{m_1+1}) + m_1 D_H(w) + (|V(H)| - 1)D_{C_{m_1+1}}(w) \\ &= W(G'), \end{aligned}$$

which is a contradiction. This shows that if 2 or more pendant blocks share a cut vertex, then all of them are K_2 . So, as $n \geq 7$, H can not be a block and hence $k \geq 2$. Suppose there are more than 2 pendant blocks in G isomorphic to K_2 sharing w . Then there exists a subgraph H' (on at least 4 vertices) of G such that $G \cong H' \cup K_{1,3}$ such that $V(H') \cap V(K_{1,3}) = \{w\}$. Construct G' from G by replacing $K_{1,3}$ by C_4 such that $V(H') \cap V(C_4) = \{w\}$. Then

$$\begin{aligned} W(G) &= W(H') + W(K_{1,3}) + 3D_{H'}(w) + (|V(H')| - 1)D_{K_{1,3}}(w) \\ &= W(H') + 9 + 3D_{H'}(w) + 3(|V(H')| - 1) \\ &< W(H') + 8 + 3D_{H'}(w) + 4(|V(H')| - 1) \quad [\text{as } |V(H')| \geq 4] \\ &= W(G'), \end{aligned}$$

which is a contradiction. This completes the proof. \square

For $4 \leq n \leq 6$, the graphs having maximum Wiener index in $\mathfrak{C}_{n,1}$ have been shown in Table 1. The following theorem characterises the graphs having maximum Wiener index in $\mathfrak{C}_{n,1}$ for $n \geq 7$.

Theorem 5.5. *Let $G \in \mathfrak{C}_{n,1}$, $n \geq 7$. Then $W(G) \leq W(L_{n,n-1})$ and equality holds if and only if $G \cong L_{n,n-1}$.*

Proof. Let G' be a graph having maximum Wiener index in $\mathfrak{C}_{n,1}$. Then all the blocks of G' are pendant. So by Lemma 5.2 and Proposition 5.4, $G' \cong L_{n,n-1}$ or $G' \cong C_{m_1, m_2}^{m_1+m_2-1}$. By Lemma 5.3, $W(L_{n,n-1}) > W(C_{m_1, m_2}^{m_1+m_2-1})$. So $G' \cong L_{n,n-1}$ and the result follows. \square

Lemma 5.6. *Let $m_1, m_2 \geq 4$ and $n = m_1 + m_2$. Then $W(C_{m_1, m_2}^n) \leq W(L_{n,n-2})$ and equality holds iff $m_1 = m_2 = 4$.*




n	Graphs G having maximum W.I. over $\mathfrak{C}_{n,1}$	$W(G)$
4		9
5	$L_{5,4}$ and 	16
6	$L_{6,5}$ and 	26

Table 1: The graphs having maximum Wiener index over $\mathfrak{C}_{n,1}$, $4 \leq n \leq 6$

Proof. Using $k = n - 2$ in (2) we get

$$W(L_{n,n-2}) = \begin{cases} \frac{1}{8}(n^3 - 2n^2 + 20n - 32) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^3 - 2n^2 + 19n - 34) & \text{if } n \text{ is odd.} \end{cases}$$

Using $n = m_1 + m_2$ and $k = 2$ in (4) the Wiener index of C_{m_1, m_2}^n can be computed as

$$W(C_{m_1, m_2}^n) = \begin{cases} \frac{1}{8}(n^3 - m_1 m_2 (n - 8)) & \text{if both } m_1 \text{ and } m_2 \text{ are even,} \\ \frac{1}{8}(n^3 - m_1 m_2 (n - 8) - n - m_1) & \text{if } m_1 \text{ is even and } m_2 \text{ odd,} \\ \frac{1}{8}(n^3 - m_1 m_2 (n - 8) - n - m_2) & \text{if } m_1 \text{ is odd } m_2 \text{ is even,} \\ \frac{1}{8}(n^3 - m_1 m_2 (n - 8) - 3n) & \text{if both } m_1 \text{ and } m_2 \text{ are odd.} \end{cases}$$

First suppose n is even. In this case

$$W(C_{m_1, m_2}^n) \leq \frac{1}{8}(n^3 - m_1 m_2 (n - 8))$$

and equality holds if and only if both m_1 and m_2 are even. As $n \geq 8$, $m_1 m_2 \geq 2(m_1 + m_2) = 2n$. Now

$$\begin{aligned} W(L_{n,n-2}) - W(C_{m_1, m_2}^n) &\geq \frac{1}{8}(m_1 m_2 (n - 8) - 2n^2 + 20n - 32) \quad [\text{equality iff both } m_1, m_2 \text{ are even}] \\ &\geq \frac{1}{8}(2n(n - 8) - 2n^2 + 20n - 32) \\ &\geq \frac{1}{8}(4n - 32) \\ &\geq 0 \end{aligned}$$

and equality holds if and only if $m_1 = m_2 = 4$. Now suppose n is odd and hence $n \geq 9$. Without loss of generality assume that m_1 is even and m_2 is odd. Then

$$\begin{aligned} W(L_{n,n-2}) - W(C_{m_1, m_2}^n) &= \frac{1}{8}(m_1 m_2 (n - 8) - 2n^2 + 20n - 34 + m_1) \\ &\geq \frac{1}{8}(2n(n - 8) - 2n^2 + 20n - 34 + m_1) \\ &= \frac{1}{8}(4n - 34 + m_1) \\ &> 0 \end{aligned}$$

This completes the proof. □

There are not many non-isomorphic graphs in $\mathfrak{C}_{5,2}$, $\mathfrak{C}_{6,2}$, $\mathfrak{C}_{7,2}$, $\mathfrak{C}_{8,2}$ and $\mathfrak{C}_{9,2}$. The graphs having maximum Wiener index over $\mathfrak{C}_{n,2}$, $5 \leq n \leq 9$ have been shown in Table 2. The following theorem characterises the graph having maximum Wiener index in $\mathfrak{C}_{n,2}$ for $n \geq 10$.





n	Graphs G having maximum W.I. over $\mathfrak{C}_{n,2}$	$W(G)$
5		18
6	$L_{6,4}$ and 	29
7		44
8	$C_{4,4}^8$ and $L_{8,6}$	64
9	$L_{9,7}$ and 	88

Table 2: The graphs having maximum Wiener index over $\mathfrak{C}_{n,2}$ for $5 \leq n \leq 9$

Theorem 5.7. *Let $G \in \mathfrak{C}_{n,2}$, $n \geq 10$. Then $W(G) \leq W(L_{n,n-2})$ and the equality holds if and only if $G \cong L_{n,n-2}$.*

Proof. Let $G \in \mathfrak{C}_{n,2}$ and w, w' be the two cut vertices in G . As we are maximizing the Wiener index, by Lemma 5.2 we can assume that each pendant block of G is either K_2 or a cycle on at least 4 vertices. Since $G \in \mathfrak{C}_{n,2}$, it has exactly one non-pendant block, say B . As $n \geq 10$, either the maximal connected subgraph containing B and no other blocks at w or the maximal connected subgraph containing B and no other blocks at w' has 6 or more vertices. Let H_1 be the maximal connected subgraph containing B and no other blocks at w and assume that $|V(H_1)| = n_1 \geq 6$. Take the union of pendant blocks at w as H_2 and let $|V(H_2)| = n_2 \geq 2$. Then $G \cong H_1 \cup H_2$ and $V(H_1) \cap V(H_2) = \{w\}$. Also $H_1 \in \mathfrak{C}_{n_1,1}$. Construct G' from G by replacing H_1 by L_{n_1, n_1-1} such that w is the pendant vertex and w' is the cut vertex of L_{n_2, n_2-1} . Then we have

$$\begin{aligned}
W(G) &= W(H_1) + W(H_2) + (n_2 - 1)D_{H_1}(w) + (n_1 - 1)D_{H_2}(w) \\
&\text{observing Theorem 4.8, Theorem 5.5 and Table 1} \\
&\leq W(L_{n_1, n_1-1}) + W(H_2) + (n_2 - 1)D_{L_{n_1, n_1-1}}(w) + (n_1 - 1)D_{H_2}(w) \\
&= W(G')
\end{aligned}$$

and equality holds iff $H_1 \cong L_{n_1, n_1-1}$ and w is the pendant vertex of H_1 . $G' \cong H_2 \cup L_{n_1, n_1-1}$ and $V(H_2) \cap V(L_{n_1, n_1-1}) = \{w\}$. As we are maximizing the Wiener index, by Proposition 5.4, we may assume that either $H_2 \cong K_2$, $H_2 \cong K_{1,2}$ with w as its non pendant vertex or $H_2 \cong C_{n_2}$ for $n_2 \geq 4$. If $H_2 \cong K_2$, then $G' \cong L_{n, n-2}$ and the result follows. For the remaining two possibilities of H_2 , $G \not\cong L_{n, n-2}$.

If $H_2 \cong K_{1,2}$ with w as its non pendant vertex, then $n = n_1 + 2$ and $G' \cong K_{1,3} \cup C_{n_1-1}$ with $V(K_{1,3}) \cap V(C_{n_1-1}) = \{w'\}$ where w' is a pendant vertex of $K_{1,3}$. By counting we get

$$W(G') \leq \frac{1}{8}(n_1^3 + 3n_1^2 + 31n_1 - 3)$$

and

$$W(L_{n, n-2}) = W(L_{n_1+2, n_1}) \geq \frac{1}{8}(n_1^3 + 4n_1^2 + 23n_1 + 4).$$

This gives

$$W(L_{n, n-2}) - W(G') \geq \frac{1}{8}(n_1^2 - 8n_1 + 7) > 0. \quad [\text{as } n_1 \geq 8]$$

Thus $W(G') < W(L_{n, n-2})$ and hence $W(G) < W(L_{n, n-2})$.

Finally, if $H_2 \cong C_{n_2}$, $n_2 \geq 4$, then $G' \cong C_{n_1-1, n_2}^n$. By Lemma 5.6, $W(G') = W(C_{n_1-1, n_2}^n) < W(L_{n, n-2})$ and hence $W(G) < W(L_{n, n-2})$. This completes the proof. \square

Lemma 5.8. *Let $m_1, m_2 \geq 4$ and $n = m_1 + m_2 + 1$. If $n \geq 14$, then $W(C_{m_1, m_2}^n) < W(L_{n, n-3})$.*

Proof. As $n = m_1 + m_2 + 1$, $C_{m_1, m_2}^n \in \mathfrak{C}_{n, 3}$. Using $k = 3$ in (4), we get

$$W(C_{m_1, m_2}^n) = \begin{cases} \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2 + 2m_2^2 + 16m_1m_2 + 8m_1 + 8m_2) & m_1 \text{ even, } m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2 + 2m_2^2 + 16m_1m_2 + 6m_1 + 7m_2 - 2) & m_1 \text{ even, } m_2 \text{ odd,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2 + 2m_2^2 + 16m_1m_2 + 7m_1 + 6m_2 - 2) & m_1 \text{ odd, } m_2 \text{ even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2 + 2m_2^2 + 16m_1m_2 + 5m_1 + 5m_2 - 4) & m_1 \text{ odd, } m_2 \text{ odd,} \end{cases}$$

and for $n = m_1 + m_2 + 1$,

$$W(L_{n, n-3}) = \begin{cases} \frac{1}{8}(m_1^3 + m_2^3 + 3m_1^2m_2 + 3m_1m_2^2 + 35m_1 + 35m_2 - 52) & \text{if } n \text{ is even,} \\ \frac{1}{8}(m_1^3 + m_2^3 + 3m_1^2m_2 + 3m_1m_2^2 + 36m_1 + 36m_2 - 48) & \text{if } n \text{ is odd,} \end{cases}$$

Suppose n is even. Note that in this case either m_1 is even and m_2 is odd, or m_1 is odd and m_2 is even. The resulting C_{m_1, m_2}^n are isomorphic. So we may assume m_1 is even and m_2 is odd. This gives

$$\begin{aligned} W(L_{n, n-3}) - W(C_{m_1, m_2}^n) &= \frac{1}{8}(m_1^2m_2 + m_1m_2^2 - 2m_1^2 - 2m_2^2 - 16m_1m_2 + 29m_1 + 28m_2 - 50) \\ &= \frac{1}{8}(m_1m_2(m_1 + m_2 - 12) - 2(m_1 + m_2)^2 + 28(m_1 + m_2) + m_1 - 52) \\ &> \frac{1}{8}(2(m_1 + m_2)(m_1 + m_2 - 12) - 2(m_1 + m_2)^2 + 28(m_1 + m_2) + m_1 - 52) \\ &\quad [\text{Inequality holds because } m_1 + m_2 - 12 > 0 \text{ and here } m_1m_2 > 2(m_1 + m_2)] \\ &= \frac{1}{8}(4(m_1 + m_2) + m_1 - 50) \\ &> 0. \end{aligned} \quad [\text{as } m_1 + m_2 \geq 13]$$

Now suppose n is odd. In this case m_1, m_2 are either both even or both odd. $W(C_{m_1, m_2}^n)$ is maximum when both m_1, m_2 are even. So

$$W(C_{m_1, m_2}^n) \leq \frac{1}{8}(m_1^3 + m_2^3 + 2m_1^2m_2 + 2m_1m_2^2 + 2m_1^2 + 2m_2^2 + 16m_1m_2 + 8m_1 + 8m_2).$$

Similar counting as in the n even case gives

$$W(L_{n, n-3}) - W(C_{m_1, m_2}^n) > \frac{1}{8}(4(m_1 + m_2) - 48) > 0. \quad [\text{as } m_1 + m_2 \geq 14]$$

This completes the proof. \square

Theorem 5.9. *Let $G \in \mathfrak{C}_{n, 3}$, $n \geq 14$. Then $W(G) \leq W(L_{n, n-3})$ and equality holds if and only if $G \cong L_{n, n-3}$.*

Proof. Let $G \in \mathfrak{C}_{n, 3}$. As $n \geq 14$, there exists an s-pendant block B in G such that $n - |V(B)| + 1 \geq 9$. By Lemma 5.2, we may assume that B is either K_2 or a cycle on at least 4 vertices. Let $|V(B)| = n_1$ and w be the cut vertex of G in B .

If B does not share its cut vertex with any other pendant block, then there exists $H \in \mathfrak{C}_{n-n_1+1,2}$ such that $G \cong B \cup H$ and $V(B) \cap V(H) = \{w\}$. Construct a graph G' from G by replacing H by $L_{n-n_1+1,n-n_1-1}$ such that w is the pendant vertex of $L_{n-n_1+1,n-n_1-1}$. Then

$$\begin{aligned} W(G) &= W(B) + W(H) + (|V(H)| - 1)D_B(w) + (|V(B)| - 1)D_H(w) \\ &\leq W(B) + W(L_{n-n_1+1,n-n_1-1}) + (|V(H) - 1|)D_B(w) + (|V(B)| - 1)D_{L_{n-n_1+1,n-n_1-1}}(w) \\ &\quad \text{[inequality follows from Theorem 4.8 and Theorem 5.7]} \\ &= W(G') \end{aligned}$$

and equality holds if and only if $H \cong L_{n-n_1+1,n-n_1-1}$ and w is the pendant vertex of H . If $B \cong K_2$, then $G' \cong L_{n,n-3}$ and the result follows. If $B \cong C_{n_1}$, $n_1 \geq 4$, then $G' \cong C_{n_1,n_2}^n$ where $n_1 + n_2 + 1 = n$. By Lemma 5.8, $W(G') < W(L_{n,n-3})$ and hence $W(G) < W(L_{n,n-3})$.

If B shares its cut vertex with another pendant block, then by Lemma 5.4, we may assume that $B \cong K_2$ and it shares its cut vertex with another pendant block K_2 . By similar argument as above we get $W(G) \leq W(G'')$ where $G'' \cong K_{1,2} \cup L_{n-2,n-4}$ with $V(K_{1,2}) \cap V(L_{n-2,n-4}) = \{w\}$ where w is the non pendant vertex of $K_{1,2}$. The Wiener indices of G'' and $L_{n,n-3}$ can be computed as

$$W(G'') = \begin{cases} \frac{1}{8}(n^3 - 4n^2 + 56n - 152) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^3 - 4n^2 + 55n - 156) & \text{if } n \text{ is odd,} \end{cases}$$

and

$$W(L_{n,n-3}) = \begin{cases} \frac{1}{8}(n^3 - 3n^2 + 38n - 88) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^3 - 3n^2 + 39n - 85) & \text{if } n \text{ is odd.} \end{cases}$$

So

$$\begin{aligned} W(L_{n,n-3}) - W(G'') &= \begin{cases} \frac{1}{8}(n^2 - 18n + 64) & \text{if } n \text{ is even,} \\ \frac{1}{8}(n^2 - 16n + 71) & \text{if } n \text{ is odd,} \end{cases} \\ &> 0. \quad \text{[as } n \geq 14] \end{aligned}$$

Thus $W(G'') < W(L_{n,n-3})$ and hence $W(G) < W(L_{n,n-3})$. This completes the proof. \square

In Theorem 5.9, the graphs maximizing the Wiener index over $\mathfrak{C}_{n,3}$ for $n \geq 14$ have been characterised. The graphs having maximum Wiener index over $\mathfrak{C}_{n,3}$, $6 \leq n \leq 13$ have been listed in Table 3.

In conclusion, the investigation into maximal graphs in $\mathfrak{C}_{n,k}$ for $k \geq 4$ presents intriguing avenues for further exploration. It is believed that there exists an integer n_0^k such that the graph $L_{n,n-k}$ maximizes the Wiener index over $C_{n,k}$ for $n \geq n_0^k$. The comparison between $W(L_{n,n-k})$ and $W(C_{m_1,m_2}^n)$ for arbitrary k may involve some intricate calculations. Additionally, exploring the determination of m_1, m_2, k for which $W(C_{m_1,m_2}^n)$ is maximum (or minimum) over $\mathfrak{C}_{n,k}$ opens a new avenue for study, which could be helpful in complementing the current study.






n	Graphs G having maximum W.I. over $\mathfrak{C}_{n,3}$	$W(G)$
6		32
7	$L_{7,4}$ and 	48
8		69
9	$C_{4,4}^9$	96
10	$C_{4,5}^{10}$ and 	126
11	$C_{4,6}^{11}$	166
12	$C_{4,7}^{12}$ and 	209
13	$C_{6,6}^{13}$ and $L_{13,10}$	264

Table 3: The graphs having maximum Wiener index over $\mathfrak{C}_{n,3}$ for $6 \leq n \leq 13$

References

- [1] R. Balakrishnan, N. Sridharan, K. Viswanathan Iyer, Wiener index of graphs with more than one cut-vertex. *Appl. Math. Lett.* 21 (2008), no. 9, 922-927.
- [2] É. Czabarka, T. V. Olsen, S. J. Smith, L. Székly, Minimum Wiener index of triangulations and quadrangulations. *Discrete Appl. Math.* 322 (2022), 295-310.
- [3] P. Dankelmann, On the Wiener index of orientations of graphs. *Discrete Appl. Math.* 336 (2023), 125-131.
- [4] G. A. Dirac, Minimally 2-connected graphs. *J. Reine Angew. Math.* 228 (1967), 204-216.
- [5] R. C. Entringer, D. E. Jackson, D. A. Snyder, Distance in graphs. *Czechoslovak Math. J.* 26(101) (1976), no. 2, 283-296.
- [6] J. Plesník, On the sum of all distances in a graph or digraph. *J. Graph Theory.* 8 (1984), no. 1, 1-21.
- [7] D. Pandey, K. L. Patra, Wiener index of graphs with fixed number of pendant or cut vertices. *Czechoslovak Math. J.* 72(147) (2022), no. 2, 411-431.
- [8] D. Pandey, K. L. Patra, Total eccentricity index of graphs with fixed number of pendant or cut vertices. *Ric. Mat.*(2022), <https://doi.org/10.1007/s11587-022-00756-8>
- [9] D. B. West, *Introduction to Graph Theory*. Prentice Hall, Inc. Upper Saddle River, NJ. (1996) xvi+512pp.
- [10] H. Wiener, Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* 69 (1947), 17-20.
- [11] G. Yu, L. Feng, On the Wiener index of unicyclic graphs with given girth. *Ars Combin.* 94 (2010), 361-369.
- [12] B. Zelinka, Medians and peripherians of trees. *Arch. Math. (Brno)* 4 (1968), 87-95.