

# ON THE HIGHER-RANK ASKEY-WILSON ALGEBRAS

Wanxia Wang, Shilin Yang<sup>†</sup>

Department of Mathematics, Beijing University of Technology  
Beijing 100124, P. R. China

**Abstract:** In the paper, a new algebra  $\mathcal{A}(n)$ , which is generated by an upper triangular generating matrix with triple relations, is introduced. It is shown that there exists an isomorphism between the algebra  $\mathcal{A}(n)$  and the higher Askey-Wilson algebra  $\mathfrak{aw}(n)$  introduced by Crampé, Frappat et al. Furthermore, we establish a series of automorphisms of  $\mathcal{A}(n)$ , which satisfy braid group relations and coincide with those in  $\mathfrak{aw}(n)$ .

**Keywords:** Askey-Wilson algebra, braid group

**Mathematics subject classification:** 16T10; 33D45; 81R12

## Introduction

The study of algebraic structures has been a cornerstone of mathematical research, providing profound insights and applications in various fields. The Askey-Wilson algebras, introduced by Zhedanov in [1], have significant implications in mathematical physics, quantum groups and orthogonal polynomials. For instance, Koelink and Stokman [2] explored the relationships between the Askey-Wilson algebras and  $q$ -hypergeometric functions, and further applied these relationships into the study of  $q$ -hypergeometric functions. Crampé and Gaboriaud et al. [3] investigated the connections between Askey-Wilson algebras and the universal  $R$ -matrix of  $U_q(\mathfrak{sl}_2)$ , providing a new perspective for understanding these algebras. Terwilliger and Vidunas [4] studied the relationships between the Askey-Wilson algebras and the corresponding Leonard pairs, to highlight the importance of these algebras in representation theory. Lavrenov [5] demonstrated their applications in integrable system, to show strongly that their significance in theoretical physics. More relevant references can be found in [6–8].

As a natural extension, it is interesting to address the definition of the higher-rank Askey-Wilson algebras and explore their structures and properties, which can provide more complex descriptions with richer symmetries and representation theories. These descriptions can demonstrate advantages in solving complex physical systems and high-dimensional spaces. Baseilhac and Koizumi [9] have explored applications of a deformed analogue of Onsager's symmetry, which can be viewed as an analog of higher-rank Askey-Wilson algebra, in physical systems such as the XXZ open chain. De Bie and van de Vijver [10] investigated the discrete realization of the higher rank Racah algebras. The readers can be referred to [11–14] for more applications.

---

<sup>†</sup> Corresponding author: slyang@bjut.edu.cn

Supported by National Natural Science Foundation of China (Grant No. 11671024)

To the best of our knowledge, Crampé and Frappat et al. [15] recently introduced definition of higher-rank Askey-Wilson algebras explicitly. This definition is strongly general and it provides a broader framework to understand higher-rank Askey-Wilson algebras. In the same paper, the authors explored Casimir elements and a series of automorphisms, which enjoy braid group relations. We hope to find more intuitive definition of the higher-rank Askey-Wilson algebras by the generators and relations, and furthermore to recover the above braided group automorphisms. Fortunately, these higher-rank Askey-Wilson algebras can be defined equivalently by the generators located in upper-triangular matrix with triple relations. We hope that this viewpoint is to provide easy methods to study the higher-rank Askey-Wilson algebras.

Now we prepare to give the outline of the paper. In Section 1, the crucial notations and fundamental concepts are introduced, and we define the algebra  $\mathcal{A}(n)$  by an upper triangular generating matrix  $\mathcal{A}$  with relations. Several basic properties for these algebras are described. In Section 2, it is shown that  $\mathcal{A}(n)$  is isomorphic to the higher-rank Askey-Wilson algebra  $\mathfrak{aw}(n)$  given by Crampé et al. in [15]. In Section 3, we establish a series of automorphisms of  $\mathcal{A}(n)$ , which coincide with those automorphisms of  $\mathfrak{aw}(n)$  given in [15]. In Section 4, we prove that those automorphisms established in the above section satisfy the braid group relations independently in our approach. We provide the detailed proofs of these results. The exploration may make us to understand  $\mathfrak{aw}(n)$  in a more intuitive way, to construct the PBW basis of the algebra  $\mathcal{A}(n)$  or equivalently  $\mathfrak{aw}(n)$ .

## 1 The algebra $\mathcal{A}(n)$

In this section, we list some notations to use in the sequel, then we give the definition of the algebra  $\mathcal{A}(n)$ .

Always assume that  $\mathbb{K}$  is an algebraically closed field with  $\text{char } \mathbb{K} = 0$ , and  $q \in \mathbb{K}$  with  $q^2 \neq 1$ . Let  $\mathbb{Z}$  be the ring of integers,  $\mathbb{N}$  the set of the nonnegative integers and  $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$ . For  $i, j \in \mathbb{Z}$ , we denote by  $\llbracket i, j \rrbracket$  the set  $\{i, i+1, \dots, j\}$ , where  $i < j$ .

Suppose that  $A, B$  are the operators and set

$$[A, B] \triangleq AB - BA, \quad [A, B]_q \triangleq \frac{qAB - q^{-1}BA}{q - q^{-1}}.$$

Then, for given operators  $A, B, C$ , one easily sees that

$$[A, B] + [B, A] = 0, \tag{1.1}$$

$$[A, B] = \frac{q - q^{-1}}{q + q^{-1}} \left( [A, B]_q - [B, A]_q \right), \tag{1.2}$$

$$\left[ [A, B]_q, C \right]_q - \left[ A, [B, C]_q \right]_q = \frac{1}{(q - q^{-1})^2} [B, [C, A]], \tag{1.3}$$

$$\left[ A, [B, C]_q \right]_q - \left[ B, [A, C]_q \right]_q = \frac{q + q^{-1}}{q - q^{-1}} [[A, B], C]_{q^2}, \tag{1.4}$$

$$\left[ A, [B, C]_q \right] + \left[ C, [A, B]_q \right] + \left[ B, [C, A]_q \right] = 0. \tag{1.5}$$

Also, if  $[A, B] = 0$ , then

$$[A, B]_q = AB, \quad [A, CB]_q = [A, C]_q B, \quad [A, BC]_q = B[A, C]_q, \quad (1.6)$$

$$[A, [C, B]_q]_q = [[A, C]_q, B]_q, \quad [A, [B, C]_q]_q = [B, [A, C]_q]_q. \quad (1.7)$$

In particular, if  $[A, B] = [A, C] = 0$ , then

$$[A, [B, C]_q]_q = A[B, C]_q. \quad (1.8)$$

We denote by  $\mathcal{A} \stackrel{\triangle}{=} (a_{i,j})_{n \times n}$  an upper triangular generating matrix:  $a_{i,j} = 0$  whenever  $i > j$ . In general, the matrix  $\mathcal{A}$  can be partitioned as follows:

$$\left( \begin{array}{c|cc|cc|c} \mathcal{A}_{11}(i-1, i-1) & \mathcal{A}_{12}(i-1, j-i) & & a_{1,j} & \mathcal{A}_{14}(i-1, n-j) \\ \hline & & & \vdots & \\ & & a_{i-1,j} & & \\ \hline a_{i,i} & \cdots & a_{i,j-1} & a_{i,j} & a_{i,j+1} & \cdots & a_{i,n} \\ \hline & \mathcal{A}_{32}(j-i, j-i) & & a_{i+1,j} & & \mathcal{A}_{34}(j-i, n-j) \\ \hline & & & \vdots & & \\ & & a_{j,j} & & & \\ \hline & & & & \mathcal{A}_{44}(n-j, n-j) & \end{array} \right), \quad (1.9)$$

where

$$\begin{aligned} \mathcal{A}_{11}(i-1, i-1) &= \begin{pmatrix} a_{1,1} & \cdots & a_{1,i-1} \\ & \ddots & \vdots \\ & & a_{i-1,i-1} \end{pmatrix}, & \mathcal{A}_{12}(i-1, j-i) &= \begin{pmatrix} a_{1,i} & \cdots & a_{1,j-1} \\ \vdots & \ddots & \vdots \\ a_{i-1,i} & \cdots & a_{i-1,j-1} \end{pmatrix}, \\ \mathcal{A}_{14}(i-1, n-j) &= \begin{pmatrix} a_{1,j} & \cdots & a_{1,n} \\ \vdots & \ddots & \vdots \\ a_{i-1,j} & \cdots & a_{i-1,n} \end{pmatrix}, & \mathcal{A}_{32}(j-i, j-i) &= \begin{pmatrix} a_{i+1,i+1} & \cdots & a_{i+1,j-1} \\ & \ddots & \vdots \\ & & a_{j-1,j-1} \end{pmatrix}, \\ \mathcal{A}_{34}(j-i, n-j) &= \begin{pmatrix} a_{i+1,j+1} & \cdots & a_{i+1,n} \\ \vdots & \ddots & \vdots \\ a_{j,j+1} & \cdots & a_{j,n} \end{pmatrix}, & \mathcal{A}_{44}(n-j, n-j) &= \begin{pmatrix} a_{j+1,j+1} & \cdots & a_{j+1,n} \\ & \ddots & \vdots \\ & & a_{n,n} \end{pmatrix}. \end{aligned}$$

The following notations are assigned.

- For generators  $a_{i,j} \in \mathcal{A}$  and  $a_{k,l} \in \mathcal{A}_{34}(j-i, n-j)$ , where  $i < k \leq j < l$ , we denote

$$\mathcal{A}_{i,k;j+1}^{k-1,j,l} = \begin{pmatrix} a_{i,k-1} & a_{i,j} & a_{i,l} \\ & a_{k,j} & a_{k,l} \\ & & a_{j+1,l} \end{pmatrix}, \quad (1.10)$$

and let

$$f(\mathcal{A}_{i,k;j+1}^{k-1,j,l}) \stackrel{\triangle}{=} a_{i,j} + a_{k,l} (a_{i,k-1} a_{j+1,l} + a_{k,j} a_{i,l}) - (a_{i,k-1} a_{k,j} + a_{i,l} a_{j+1,l}), \quad (1.11)$$

$$g\left(\mathcal{A}_{i,k,j+1}^{k-1,j,l}\right) \triangleq a_{k,l} + a_{i,j} (a_{i,k-1}a_{j+1,l} + a_{k,j}a_{i,l}) - (a_{i,k-1}a_{i,l} + a_{k,j}a_{j+1,l}). \quad (1.12)$$

For example, if we choose

$$\mathcal{A}_{1,2,4}^{1,3,4} = \begin{pmatrix} a_{1,1} & a_{1,3} & a_{1,4} \\ & a_{2,3} & a_{2,4} \\ & & a_{4,4} \end{pmatrix},$$

then

$$\begin{aligned} f\left(\mathcal{A}_{1,2,4}^{1,3,4}\right) &= a_{1,3} + a_{2,4} (a_{1,1}a_{4,4} + a_{2,3}a_{1,4}) - (a_{1,1}a_{2,3} + a_{1,4}a_{4,4}), \\ g\left(\mathcal{A}_{1,2,4}^{1,3,4}\right) &= a_{2,4} + a_{1,3} (a_{1,1}a_{4,4} + a_{2,3}a_{1,4}) - (a_{1,1}a_{1,4} + a_{2,3}a_{4,4}). \end{aligned}$$

- For generators  $a_{i,j} \in \mathcal{A}$  and  $a_{k,l} \in \mathcal{A}_{34}(j-i, n-j)$ , where  $i < k \leq j < l < m \leq n$ , we denote

$$\mathcal{A}_{i,k,j+1}^{j,l,m} = \begin{pmatrix} a_{i,j} & a_{i,l} & a_{i,m} \\ a_{k,j} & a_{k,l} & a_{k,m} \\ & a_{j+1,l} & a_{j+1,m} \end{pmatrix}, \quad (1.13)$$

and let

$$\begin{aligned} \det_q\left(\mathcal{A}_{i,k,j+1}^{j,l,m}\right) &\triangleq \left[ [a_{i,j}, a_{k,l}]_q, a_{j+1,m} \right]_q + \left[ [a_{i,l}, a_{k,m}]_q, a_{j+1,l} \right]_q \\ &\quad - \left[ [a_{i,l}, a_{k,j}]_q, a_{j+1,m} \right]_q - \left[ [a_{i,j}, a_{k,m}]_q, a_{j+1,l} \right]_q - [a_{i,m}, a_{k,l}]_q, \end{aligned} \quad (1.14)$$

$$\begin{aligned} \det^q\left(\mathcal{A}_{i,k,j+1}^{j,l,m}\right) &\triangleq \left[ [a_{j+1,m}, a_{k,l}]_q, a_{i,j} \right]_q + \left[ [a_{k,m}, a_{i,l}]_q, a_{i,m} \right]_q \\ &\quad - \left[ [a_{j+1,m}, a_{k,j}]_q, a_{i,l} \right]_q - \left[ [a_{j+1,l}, a_{k,m}]_q, a_{i,j} \right]_q - [a_{k,l}, a_{i,m}]_q. \end{aligned} \quad (1.15)$$

For example, if we choose

$$\mathcal{A}_{1,2,4}^{3,4,5} = \begin{pmatrix} a_{1,3} & a_{1,4} & a_{1,5} \\ a_{2,3} & a_{2,4} & a_{2,5} \\ & a_{4,4} & a_{4,5} \end{pmatrix},$$

then

$$\begin{aligned} \det_q(\mathcal{A}_{1,2,4}^{3,4,5}) &= \left[ [a_{1,3}, a_{2,4}]_q, a_{4,5} \right]_q + \left[ [a_{1,4}, a_{2,5}]_q, a_{4,4} \right]_q \\ &\quad - \left[ [a_{1,4}, a_{2,3}]_q, a_{4,5} \right]_q - \left[ [a_{1,3}, a_{2,5}]_q, a_{4,4} \right]_q - [a_{1,5}, a_{2,4}]_q, \\ \det^q(\mathcal{A}_{1,2,4}^{3,4,5}) &= \left[ [a_{4,5}, a_{2,4}]_q, a_{1,3} \right]_q + \left[ [a_{2,5}, a_{1,4}]_q, a_{1,5} \right]_q \\ &\quad - \left[ [a_{4,5}, a_{2,3}]_q, a_{1,4} \right]_q - \left[ [a_{4,4}, a_{2,5}]_q, a_{1,3} \right]_q - [a_{2,4}, a_{1,5}]_q. \end{aligned}$$

Now we can introduce the definition of the algebra  $\mathcal{A}(n)$  explicitly.

**Definition 1.1.** *The  $\mathbb{K}$ -algebra  $\mathcal{A}(n)$  is an associative algebra generated by the upper generating matrix  $\mathcal{A}$  in (1.9), subjecting to the following defining relations:*

R1: *The entries  $a_{i,j}$  commutes with all entries in  $\mathcal{A}$  except in  $\mathcal{A}_{12}(i-1, j-i)$  and  $\mathcal{A}_{34}(j-i, n-j)$ .*

R2: *Any sub-matrix  $\mathcal{A}_{i,k,j+1}^{k-1,j,l}$  as (1.10) enjoys the relations*

$$\left[ a_{k,l}, [a_{i,j}, a_{k,l}]_q \right]_q = f(\mathcal{A}_{i,k,j+1}^{k-1,j,l}), \quad \left[ a_{i,j}, [a_{k,l}, a_{i,j}]_q \right]_q = g(\mathcal{A}_{i,k,j+1}^{k-1,j,l}). \quad (1.16)$$

R3: *Any sub-matrix  $\mathcal{A}_{i,k,j+1}^{j,l,m}$  as (1.13) enjoys the relations*

$$\det_q(\mathcal{A}_{i,k,j+1}^{j,l,m}) = \det^q(\mathcal{A}_{i,k,j+1}^{j,l,m}) = 0. \quad (1.17)$$

In this situation, we say that the algebra  $\mathcal{A}(n)$  is generated by the matrix  $\mathcal{A}$  with the relations (R1)-(R3).

**Example 1.2.** *The algebra  $\mathcal{A}(4)$  is generated by  $a_{i,j}$  ( $1 \leq i \leq j \leq 4$ ) with the relations as follows:*

R1:  $a_{1,1}, a_{2,2}, a_{3,3}, a_{4,4}, a_{1,4}$  are in the center of  $\mathcal{A}(4)$  and

$$[a_{1,2}, a_{1,3}] = [a_{1,2}, a_{3,4}] = [a_{1,3}, a_{2,3}] = [a_{2,3}, a_{2,4}] = [a_{2,4}, a_{3,4}] = 0;$$

R2:

$$\begin{aligned} \left[ a_{2,3}, [a_{1,2}, a_{2,3}]_q \right]_q &= f(\mathcal{A}_{1,2,3}^{1,2,3}), & \left[ a_{1,2}, [a_{2,3}, a_{1,2}]_q \right]_q &= g(\mathcal{A}_{1,2,3}^{1,2,3}), \\ \left[ a_{3,4}, [a_{2,3}, a_{3,4}]_q \right]_q &= f(\mathcal{A}_{2,3,4}^{2,3,4}), & \left[ a_{2,3}, [a_{3,4}, a_{2,3}]_q \right]_q &= g(\mathcal{A}_{2,3,4}^{2,3,4}), \\ \left[ a_{2,4}, [a_{1,3}, a_{2,4}]_q \right]_q &= f(\mathcal{A}_{1,2,4}^{1,3,4}), & \left[ a_{1,3}, [a_{2,4}, a_{1,3}]_q \right]_q &= g(\mathcal{A}_{1,2,4}^{1,3,4}), \\ \left[ a_{2,4}, [a_{1,2}, a_{2,4}]_q \right]_q &= f(\mathcal{A}_{1,2,3}^{1,2,4}), & \left[ a_{1,2}, [a_{2,4}, a_{1,2}]_q \right]_q &= g(\mathcal{A}_{1,2,3}^{1,2,4}), \\ \left[ a_{3,4}, [a_{1,3}, a_{3,4}]_q \right]_q &= f(\mathcal{A}_{1,3,4}^{2,3,4}), & \left[ a_{1,3}, [a_{3,4}, a_{1,3}]_q \right]_q &= g(\mathcal{A}_{1,3,4}^{2,3,4}); \end{aligned}$$

R3:

$$\det_q(\mathcal{A}_{1,2,3}^{2,3,4}) = \det^q(\mathcal{A}_{1,2,3}^{2,3,4}) = 0.$$

The following relations are used to prove the main results.

**Lemma 1.3.** *For the sub-matrix  $\mathcal{A}_{j,k,i+1}^{i,l,m}$  as (1.13), the following relations hold in  $\mathcal{A}(n)$ :*

$$\begin{aligned} &\left[ [a_{j,l}, a_{k,m}]_q, a_{j,i} \right]_q + a_{j,k-1} a_{i+1,l} a_{j,m} + [a_{k,l}, a_{i+1,m}]_q = a_{j,m} [a_{k,l}, a_{j,i}]_q \\ &\quad + a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q + a_{i+1,l} a_{k,m}, \\ &\left[ [a_{j,i}, a_{k,m}]_q, a_{j,l} \right]_q + a_{j,k-1} a_{i+1,l} a_{j,m} + [a_{i+1,m}, a_{k,l}]_q = a_{j,k-1} [a_{i+1,m}, a_{j,l}]_q \end{aligned} \quad (1.a)$$

$$+ a_{j,m} [a_{j,i}, a_{k,l}]_q + a_{i+1,l} a_{k,m}; \quad (1.b)$$

$$\left[ [a_{k,m}, a_{j,l}]_q, a_{i+1,m} \right]_q + a_{k,i} a_{l+1,m} a_{j,m} + [a_{k,l}, a_{j,i}]_q = a_{j,m} [a_{k,l}, a_{i+1,m}]_q$$

$$+ a_{l+1,m} [a_{k,m}, a_{j,i}]_q + a_{k,i} a_{j,l}, \quad (2.a)$$

$$\left[ [a_{i+1,m}, a_{j,l}]_q, a_{k,m} \right]_q + a_{k,i} a_{l+1,m} a_{j,m} + [a_{j,i}, a_{k,l}]_q = [a_{i+1,m}, a_{k,l} a_{j,m}]_q$$

$$+ a_{l+1,m} [a_{j,i}, a_{k,m}]_q + a_{k,i} a_{j,l}; \quad (2.b)$$

$$\left[ [a_{k,l}, a_{j,i}]_q, a_{k,m} \right]_q + a_{j,k-1} a_{k,i} a_{l+1,m} + [a_{j,l}, a_{i+1,m}]_q = a_{k,i} [a_{j,l}, a_{k,m}]_q$$

$$+ a_{j,k-1} [a_{k,l}, a_{i+1,m}]_q + a_{l+1,m} a_{j,i}, \quad (3.a)$$

$$\left[ [a_{k,m}, a_{j,i}]_q, a_{k,l} \right]_q + a_{j,k-1} a_{k,i} a_{l+1,m} + [a_{i+1,m}, a_{j,l}]_q = a_{k,i} [a_{k,m}, a_{j,l}]_q$$

$$+ a_{j,k-1} [a_{i+1,m}, a_{k,l}]_q + a_{l+1,m} a_{j,i}; \quad (3.b)$$

$$\left[ [a_{j,l}, a_{i+1,m}]_q, a_{k,l} \right]_q + a_{j,k-1} a_{i+1,l} a_{l+1,m} + [a_{j,i}, a_{k,m}]_q = a_{i+1,l} [a_{j,l}, a_{k,m}]_q$$

$$+ a_{l+1,m} [a_{j,i}, a_{k,l}]_q + a_{j,k-1} a_{i+1,m}, \quad (4.a)$$

$$\left[ [a_{k,l}, a_{i+1,m}]_q, a_{j,l} \right]_q + a_{j,k-1} a_{i+1,l} a_{l+1,m} + [a_{k,m}, a_{j,i}]_q = a_{i+1,l} [a_{k,m}, a_{j,l}]_q$$

$$+ a_{l+1,m} [a_{k,l}, a_{j,i}]_q + a_{j,k-1} a_{i+1,m}; \quad (4.b)$$

$$\left[ [a_{i,l}, a_{j,i}]_q, a_{k,l} \right]_q + a_{k,i-1} a_{i+1,l} a_{j,l} + [a_{j,i-1}, a_{k,i}]_q = a_{i+1,l} [a_{j,i-1}, a_{k,l}]_q$$

$$+ a_{j,l} [a_{i,l}, a_{k,i}]_q + a_{k,i-1} a_{j,i}, \quad (5.a)$$

$$\left[ [a_{k,l}, a_{j,i}]_q, a_{i,l} \right]_q + a_{k,i-1} a_{i+1,l} a_{j,l} + [a_{k,i}, a_{j,i-1}]_q = a_{i+1,l} [a_{k,l}, a_{j,i-1}]_q$$

$$+ a_{j,l} [a_{k,i}, a_{j,k}]_q + a_{k,i-1} a_{j,i}; \quad (5.b)$$

$$\left[ [a_{i,l}, a_{j,i}]_q, a_{i,m} \right]_q + a_{j,i-1} a_{i,i} a_{l+1,m} + [a_{j,l}, a_{i+1,m}]_q = a_{i,i} [a_{j,l}, a_{i,m}]_q$$

$$+ a_{j,i-1} [a_{i,l}, a_{i+1,m}]_q + a_{j,i} a_{l+1,m}, \quad (6.a)$$

$$\left[ [a_{i,m}, a_{j,i}]_q, a_{i,l} \right]_q + a_{j,i-1} a_{i,i} a_{l+1,m} + [a_{i+1,m}, a_{j,l}]_q = a_{i,i} [a_{i,m}, a_{j,l}]_q$$

$$+ a_{j,i-1} [a_{i+1,m}, a_{i,l}]_q + a_{j,i} a_{l+1,m}. \quad (6.b)$$

*Proof.* We only verify the relation (1.a) here. The proofs of the other relations follows in a similar way. We have

$$\begin{aligned} & \left[ [a_{j,l}, a_{k,m}]_q, a_{j,i} \right]_q + a_{j,k-1} a_{i+1,l} a_{j,m} + [a_{k,l}, a_{i+1,m}]_q \\ & - a_{j,m} [a_{k,l}, a_{j,i}]_q - a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q - a_{i+1,l} a_{k,m} \\ \equiv & \left[ [a_{j,l}, a_{k,m}]_q - a_{j,m} a_{k,l}, a_{j,i} \right]_q + a_{j,k-1} a_{i+1,l} a_{j,m} - a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q \\ & + [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l} a_{k,m} \\ \stackrel{(1.17)}{=} & - \left[ [a_{j,i}, [a_{k,l}, a_{i+1,m}]_q]_q - a_{i+1,l} [a_{j,i}, a_{k,m}]_q - a_{k,i} [a_{j,l}, a_{i+1,m}]_q + a_{k,i} a_{i+1,l} a_{j,m}, \right. \end{aligned}$$

$$\begin{aligned}
& [a_{j,i}]_q + a_{j,k-1}a_{i+1,l}a_{j,m} - a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q + [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l}a_{k,m} \\
&= - \left[ \left[ [a_{j,i}, a_{k,l}]_q, a_{i+1,m} \right]_q, a_{j,i} \right]_q + a_{i+1,l} \left[ [a_{j,i}, a_{k,m}]_q, a_{j,i} \right]_q \\
&\quad + a_{k,i} \left[ [a_{j,l}, a_{i+1,m}]_q, a_{j,i} \right]_q - a_{j,i}a_{k,i}a_{i+1,l}a_{j,m} + a_{j,k-1}a_{i+1,l}a_{j,m} \\
&\quad - a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q + [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l}a_{k,m} \\
&\stackrel{(1.16)}{=} - \left[ g \left( \mathcal{A}_{j,k,i+1}^{k-1,i,l} \right), a_{i+1,m} \right]_q + a_{i+1,l}g \left( \mathcal{A}_{j,k,i+1}^{k-1,i,m} \right) + a_{k,i} \left[ [a_{j,l}, a_{i+1,m}]_q, a_{j,i} \right]_q \\
&\quad - a_{j,i}a_{k,i}a_{i+1,l}a_{j,m} + a_{j,k-1}a_{i+1,l}a_{j,m} - a_{j,k-1} [a_{j,l}, a_{i+1,m}]_q \\
&\quad + [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l}a_{k,m} \\
&\stackrel{(1.7)}{=} - \left[ g \left( \mathcal{A}_{j,k,i+1}^{k-1,i,l} \right) + a_{j,k-1}a_{j,l} - a_{k,l}, a_{i+1,m} \right]_q + a_{k,i} \left[ [a_{j,l}, a_{i+1,m}]_q, a_{j,i} \right]_q \\
&\quad + a_{i+1,l} \left( g \left( \mathcal{A}_{j,k,i+1}^{k-1,i,m} \right) + a_{j,k-1}a_{j,m} - a_{k,m} \right) - a_{j,i}a_{k,i}a_{i+1,l}a_{j,m} \\
&\stackrel{(1.12)}{=} (a_{k,i}a_{i+1,l} - a_{j,k-1}a_{j,i}a_{i+1,l})a_{i+1,m} - a_{i+1,l}(a_{k,i}a_{i+1,m} - a_{j,i}a_{j,k-1}a_{i+1,m}) \\
&= 0.
\end{aligned}$$

This proof is finished.  $\square$

**Lemma 1.4.** For the sub-matrix  $\mathcal{A}_{j,k,i+1}^{i,l,m}$  as (1.13), the following relations hold in  $\mathcal{A}(n)$ :

$$[a_{k,i}a_{j,l} - [a_{j,i}, a_{k,l}]_q, a_{i+1,l}a_{k,m} - [a_{i+1,m}, a_{k,l}]_q] = 0, \quad (1.19)$$

$$[a_{k,i}a_{j,l} - [a_{j,i}, a_{k,l}]_q, a_{j,k-1}a_{i+1,m} - [a_{j,i}, a_{k,m}]_q] = 0, \quad (1.20)$$

$$[a_{j,k-1}a_{i+1,m} - [a_{j,i}, a_{k,m}]_q, a_{k,l}a_{j,m} - [a_{j,l}, a_{k,m}]_q] = 0, \quad (1.21)$$

$$[a_{j,i}a_{l+1,m} - [a_{j,l}, a_{i+1,m}]_q, a_{i+1,l}a_{k,m} - [a_{k,l}, a_{i+1,m}]_q] = 0, \quad (1.22)$$

$$[a_{j,i}a_{l+1,m} - [a_{j,l}, a_{i+1,m}]_q, a_{k,l}a_{j,m} - [a_{j,l}, a_{k,m}]_q] = 0. \quad (1.23)$$

*Proof.* Here we only verify the relation (1.19). The proofs of the other relations follows in a similar way. We have

$$\begin{aligned}
& \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, -[a_{i+1,m}, a_{k,l}]_q + a_{i+1,l}a_{k,m} \right] \\
&= - \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, [a_{i+1,m}, a_{k,l}]_q \right] + a_{i+1,l} \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, a_{k,m} \right] \\
&\stackrel{(1.5)}{=} \left[ a_{i+1,m}, \left[ a_{k,l}, -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l} \right]_q \right] + \left[ a_{k,l}, \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, a_{i+1,m} \right]_q \right] \\
&\quad + a_{i+1,l} \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, a_{k,m} \right] \\
&\stackrel{(1.16)}{=} \left[ a_{i+1,m}, -a_{j,i} + a_{j,k-1}a_{k,i} + a_{i+1,l}a_{j,l} \right] + \left[ a_{k,l}, \left[ a_{j,l}, a_{k,m} \right]_q - a_{j,k-1}a_{l+1,m} \right. \\
&\quad \left. - a_{k,l}a_{j,m} + a_{j,k-1}a_{l+1,m} + a_{i+1,l} \left( -[a_{j,i}, a_{k,m}]_q + a_{j,k-1}a_{i+1,m} + a_{k,i}a_{j,m} \right) \right] \\
&\quad + a_{i+1,l} \left[ -[a_{j,i}, a_{k,l}]_q + a_{k,i}a_{j,l}, a_{k,m} \right]
\end{aligned}$$

$$\begin{aligned}
& \stackrel{(R1)}{=} a_{i+1,l} [a_{i+1,m}, a_{j,l}] + a_{i+1,l} [a_{k,l}, -[a_{j,i}, a_{k,m}]_q + a_{j,k-1} a_{i+1,m} + a_{k,i} a_{j,m}] \\
& \quad + a_{i+1,l} [-[a_{j,i}, a_{k,l}]_q + a_{k,i} a_{j,l}, a_{k,m}] \\
& = a_{i+1,l} \left( [a_{i+1,m}, a_{j,l}] - [a_{k,l}, [a_{j,i}, a_{k,m}]_q - a_{j,k-1} a_{i+1,m} - a_{k,i} a_{j,m}] \right. \\
& \quad \left. - [[a_{j,i}, a_{k,l}]_q - a_{k,i} a_{j,l}, a_{k,m}] \right) \\
& \stackrel{(1.2)}{=} \frac{q-q^{-1}}{q+q^{-1}} a_{i+1,l} \left( [a_{i+1,m}, a_{j,l}]_q - [a_{j,l}, a_{i+1,m}]_q \right. \\
& \quad + [a_{k,l}, -[a_{j,i}, a_{k,m}]_q + a_{j,k-1} a_{i+1,m} + a_{k,i} a_{j,m}]_q \\
& \quad - [[- [a_{j,i}, a_{k,l}]_q + a_{k,i} a_{j,l}, a_{k,m}]_q - [a_{k,m}, (-[a_{j,i}, a_{k,l}]_q + a_{k,i} a_{j,l})]_q] \\
& = \frac{q-q^{-1}}{q+q^{-1}} a_{i+1,l} \left( ([a_{i+1,m}, a_{j,l}]_q + [[a_{k,m}, a_{j,i}]_q, a_{k,l}]_q - a_{k,i} [a_{k,m}, a_{j,l}]_q \right. \\
& \quad - a_{j,k-1} [a_{i+1,m}, a_{k,l}]_q) - ([a_{j,l}, a_{i+1,m}]_q + [[a_{k,l}, a_{j,i}]_q, a_{k,m}]_q \\
& \quad \left. - a_{k,i} [a_{j,l}, a_{k,m}]_q - a_{j,k-1} [a_{k,l}, a_{i+1,m}]_q) \right) \\
& \stackrel{(3.a)}{=} 0.
\end{aligned}$$

This proof is finished.  $\square$

**Proposition 1.5.** *The sub-algebra  $\mathcal{W}$  of  $\mathcal{A}(n)$  generated by  $\mathcal{A}_{i,k,j+1}^{k-1,j,l}$  as (1.10) is isomorphic to  $\mathcal{A}(3)$ .*

*Proof.* By definition, the algebra  $\mathcal{A}(3)$  is generated by the entries in the matrix

$$\begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ & a_{2,2} & a_{2,3} \\ & & a_{3,3} \end{pmatrix}$$

with the relations (R1)-(R3) in Definition 1.1.

Set a map  $\psi : \mathcal{W}$  to  $\mathcal{A}(3)$  such that

$$\psi(\mathcal{A}_{i,k,j+1}^{k-1,j,l}) = \psi \left( \begin{pmatrix} a_{i,k-1} & a_{i,j} & a_{i,l} \\ & a_{k,j} & a_{k,l} \\ & & a_{j+1,l} \end{pmatrix} \right) = \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ & a_{2,2} & a_{2,3} \\ & & a_{3,3} \end{pmatrix}.$$

It is obvious that  $\psi$  is an algebraic isomorphism. Finally, we note that  $a_{i,k-1}, a_{k,j}, a_{j+1,l}, a_{i,j}$  are in the center of the subalgebra  $\mathcal{W}$ .  $\square$

**Corollary 1.6.** *Given two generators  $a_{i,j}, a_{k,l} \in \mathcal{A}$ , then  $[a_{i,j}, a_{k,l}] = 0$  if one of the following holds.*

1.  $i = k$ ;
2.  $j = l$ ;
3.  $a_{i,j}, a_{k,l}$  locate on the secondary diagonal of a  $3 \times 3$  sub-matrix of the matrix  $\mathcal{A}$ .

*Proof.* It is obvious by Definition 1.1 and Proposition 1.5.  $\square$

We also have the following obvious results.

**Corollary 1.7.** *The generators  $a_{i,i}$  ( $i \in [\![1, n]\!]$ ) and  $a_{1,n}$  are in the center of  $\mathcal{A}(n)$ .*

**Proposition 1.8.** *We have the following filtration of algebras:*

$$\mathcal{A}(1) \subseteq \mathcal{A}(2) \subseteq \cdots \subseteq \mathcal{A}(n-1) \subseteq \mathcal{A}(n).$$

## 2 The relations between algebra $\mathcal{A}(n)$ and $\text{aw}(n)$

In this section, we first recall the concept of the higher-rank Askey-Wilson algebras  $\text{aw}(n)$  introduced in [15]. Then we explore the relationship between algebras  $\mathcal{A}(n)$  and  $\text{aw}(n)$ . In fact, we shall see that  $\mathcal{A}(n)$  is just isomorphic to the algebra  $\text{aw}(n)$ . In other word we get a equivalent definition of the higher-rank Askey-Wilson algebras  $\text{aw}(n)$ .


To see the fact, firstly, let us review the definition of  $\text{aw}(n)$  in [15].

The following relevant notations and terminology on sets and subsets are used.

- For two subsets  $I, J \subseteq \{1, \dots, n\}$ , we say that  $I < J$  if and only if

$$i < j, \quad \forall i \in I, \quad j \in J.$$

- A non-empty subset  $I \subseteq \{1, \dots, n\}$  is said to be connected if it consists in a subset of consecutive integers.
- Two disjoint connected subsets  $I, J \subseteq \{1, \dots, n\}$  are said to be adjacent if their union is connected.
- A hole  $H$  between two disjoint connected subsets  $I_1$  and  $I_2$  means that  $H$  consists in the connected subset between  $I_1$  and  $I_2$ , as the picture:



In this picture and in all the similar pictures below, the integers dotted by •'s are ordered either from left to right or from right to left (depending on the respective positions of  $I_1$  and  $I_2$  in the natural order). So such a picture does not mean that  $I_1 < I_2$ , but  $I_1, H, I_2$  are adjacent connected subsets.

- A sequence  $(I_1, \dots, I_k)$  of non-empty connected subsets of  $\{1, \dots, n\}$  is said monotonic if either  $I_1 < I_2 < \dots < I_k$  or  $I_1 > I_2 > \dots > I_k$ .

Let us consider  $I_1, I_2$  two non-empty and disjoint connected subsets of  $\{1, \dots, n\}$  with a non-empty hole  $H$  between them:

$$\cdots \bullet, \underbrace{\bullet, \dots, \bullet}_{I_1}, \underbrace{\bullet, \dots, \bullet}_{H}, \underbrace{\bullet, \dots, \bullet}_{I_2}, \bullet, \dots$$

The element  $C_{I_1 I_2}$  is defined by

$$C_{I_1 I_2} := -[C_{I_1 H}, C_{H I_2}]_q + C_{I_1} C_{I_2} + C_H C_{I_1 H I_2}. \quad (2.1)$$

Now we can describe the definition of higher-rank Askey-Wilson algebra  $\text{aw}(n)$ .

**Definition 2.1.** [15] *The algebra  $\text{aw}(n)$  is the unital associative algebra generated by the elements  $C_I$ , where  $I$  is any non-empty connected subset of  $\{1, \dots, n\}$ , satisfying the following relations :*

- for any two connected subsets  $I$  and  $J$ ,

$$[C_I, C_J] = 0 \quad \text{if } I \cap J = \emptyset \text{ or } I \subset J;$$

- for any monotonic sequence of three adjacent non-empty connected subsets  $(I_1, I_2, I_3)$ ,

$$C_{I_1 I_2} = -[C_{I_2 I_3}, C_{I_1 I_3}]_q + C_{I_1} C_{I_2} + C_{I_3} C_{I_1 I_2 I_3},$$

where  $C_{I_1 I_3}$  is defined by (2.1);

- for any monotonic sequence of four adjacent non-empty connected subsets  $(I_1, I_2, I_3, I_4)$ ,

$$C_{I_1 I_4} = -[C_{I_1 I_3}, C_{I_3 I_4}]_q + C_{I_1} C_{I_4} + C_{I_3} C_{I_1 I_3 I_4},$$

where  $C_{I_1 I_3}, C_{I_1 I_4}$  and  $C_{I_1 I_3 I_4}$  are defined by (2.1).

**Theorem 2.2.** *The algebra  $\mathcal{A}(n)$  is isomorphic to  $\text{aw}(n)$ .*

*Proof.* To see this, we define a map  $\phi : \mathcal{A}(n) \rightarrow \text{aw}(n)$ , which assigns  $a_{ij}$  to  $C_{\{i, i+1, \dots, j\}}$ , where  $\{i, i+1, \dots, j\} \subseteq \{1, 2, \dots, n\}$ .

It is evident that the mapping  $\phi$  is well-defined. In the following, we show that  $\phi$  is indeed an algebra homomorphism.

- (a).  $\phi$  keeps the relations (R1) in Definition 1.1:

For a given generator  $a_{i,j} \in \mathcal{A}$ , if there exists  $a_{k,l} \in \mathcal{A}$  such that  $[a_{i,j}, a_{k,l}] = 0$ , then  $a_{k,l} \notin \mathcal{A}_{12}(i-1, j-i) \cup \mathcal{A}_{34}(j-i, n-j)$ . Now, we set

$$I_1 = \{i, i+1, \dots, j\}, I_2 = \{k, k+1, \dots, l\}.$$

If  $a_{k,l}$  belongs to  $\mathcal{A}_{11}(n-j, n-j)$  or  $\mathcal{A}_{44}(i-1, i-1)$ , then

$$I_1 \cap I_2 = \emptyset.$$

If  $a_{k,l} \in \mathcal{A}_{14}(i-1, n-j)$ , we have

$$I_1 \subseteq I_2.$$

If  $a_{k,l} \in \mathcal{A}_{32}(j-i, j-i)$ , we have

$$I_2 \subseteq I_1.$$

In addition, if  $a_{k,l}$  is located on the same row or column as  $a_{i,j}$ , then

$$I_1 \subseteq I_2 \text{ or } I_2 \subseteq I_1.$$

Therefore, in all the aforementioned cases, we have

$$\begin{aligned} & \phi(a_{i,j})\phi(a_{k,l}) - \phi(a_{k,l})\phi(a_{i,j}) \\ &= C_{I_1}C_{I_2} - C_{I_2}C_{I_1} \\ &= 0. \end{aligned}$$

(b).  $\phi$  keeps the relations (R2) in Definition 1.1:

Recall that  $\mathcal{A}_{i,k,j+1}^{k-1,j,l}$  as in (1.10), we have  $i < k \leq j < l$  and define

$$I_1 = \{i, i+1, \dots, k-1\}, I_2 = \{k, k+1, \dots, j\}, I_3 = \{j+1, \dots, l\}.$$

The sequences  $(I_1, I_2, I_3)$  and  $(I_3, I_2, I_1)$  are two monotonic sequences.

Accordingly, we have

$$\begin{aligned} & \left[ \phi(a_{k,l}), [\phi(a_{i,j}), \phi(a_{k,l})]_q \right]_q \\ &= \left[ C_{I_2 I_3}, [C_{I_1 I_2}, C_{I_2 I_3}]_q \right]_q \\ &= [C_{I_2 I_3}, -C_{I_1 I_3} + C_{I_1} C_{I_3} + C_{I_2} C_{I_1 I_2 I_3}]_q \\ &= -[C_{I_2 I_3}, C_{I_1 I_3}]_q + C_{I_2 I_3} C_{I_1} C_{I_3} + C_{I_2 I_3} C_{I_2} C_{I_1 I_2 I_3} \\ &= C_{I_1 I_2} - C_{I_1} C_{I_2} - C_{I_3} C_{I_1 I_2 I_3} + C_{I_2 I_3} C_{I_1} C_{I_3} + C_{I_2 I_3} C_{I_2} C_{I_1 I_2 I_3} \\ &= \phi(a_{i,j}) - \phi(a_{i,k-1}) \phi(a_{k,j}) - \phi(a_{i,l}) \phi(a_{j+1,l}) \\ &\quad + \phi(a_{k,l}) (\phi(a_{i,k-1}) \phi(a_{j+1,l}) + \phi(a_{k,j}) \phi(a_{i,l})), \end{aligned}$$

and

$$\begin{aligned} & \left[ \phi(a_{i,j}), [\phi(a_{k,l}), \phi(a_{i,j})]_q \right]_q \\ &= \left[ C_{I_2 I_1}, [C_{I_3 I_2}, C_{I_2 I_1}]_q \right]_q \\ &= [C_{I_2 I_1}, -C_{I_3 I_1} + C_{I_3} C_{I_1} + C_{I_2} C_{I_3 I_2 I_1}]_q \\ &= -[C_{I_2 I_1}, C_{I_3 I_1}]_q + C_{I_2 I_1} C_{I_3} C_{I_1} + C_{I_2 I_1} C_{I_2} C_{I_3 I_2 I_1} \end{aligned}$$

$$\begin{aligned}
&= C_{I_3 I_2} - C_{I_3} C_{I_2} - C_{I_1} C_{I_3 I_2 I_1} + C_{I_2 I_1} C_{I_3} C_{I_1} + C_{I_2 I_1} C_{I_2} C_{I_3 I_2 I_1} \\
&= \phi(a_{k,l}) - \phi(a_{i,k-1}) \phi(a_{i,l}) - \phi(a_{k,j}) \phi(a_{j+1,l}) \\
&\quad + \phi(a_{i,j}) (\phi(a_{i,k-1}) \phi(a_{j+1,l}) + \phi(a_{k,j}) \phi(a_{i,l})).
\end{aligned}$$

Hence,

$$\begin{aligned}
[\phi(a_{k,l}), [\phi(a_{i,j}), \phi(a_{k,l})]]_q &= f(\phi(\mathcal{A}_{i,k,j+1}^{k-1,j,l})), \\
[\phi(a_{i,j}), [\phi(a_{k,l}), \phi(a_{i,j})]]_q &= g(\phi(\mathcal{A}_{i,k,j+1}^{k-1,j,l})). 
\end{aligned}$$

(c).  $\phi$  keeps the relations (R3) in Definition 1.1:

Let us consider  $\mathcal{A}_{i,k,j+1}^{j,l,m}$  as in (1.13), we must have  $i < k \leq j < l < m$ . Set

$$I_1 = \{i, i+1, \dots, k-1\}, I_2 = \{k, k+1, \dots, j\}, I_3 = \{j+1, \dots, l\}, I_4 = \{l+1, \dots, m\}.$$

Then,  $(I_1, I_2, I_3, I_4)$  and  $(I_4, I_3, I_2, I_1)$  are two monotonic sequences and we have

$$\begin{aligned}
&[[\phi(a_{i,j}), \phi(a_{k,l})], \phi(a_{j+1,m})]_q + [\phi(a_{i,l}), \phi(a_{k,m})]_q + [[\phi(a_{i,m}), \phi(a_{k,j})], \phi(a_{j+1,l})]_q \\
&\quad - [[\phi(a_{i,l}), \phi(a_{k,j})], \phi(a_{j+1,m})]_q - [[\phi(a_{i,j}), \phi(a_{k,m})], \phi(a_{j+1,l})]_q - [\phi(a_{i,m}), \phi(a_{k,l})]_q \\
&= [[C_{I_1 I_2}, C_{I_2 I_3}], C_{I_3 I_4}]_q + [C_{I_1 I_2 I_3}, C_{I_2 I_3 I_4}]_q + [[C_{I_1 I_2 I_3 I_4}, C_{I_2}], C_{I_3}]_q \\
&\quad - [[C_{I_1 I_2 I_3}, C_{I_2}], C_{I_3 I_4}]_q - [[C_{I_1 I_2}, C_{I_2 I_3 I_4}], C_{I_3}]_q - [C_{I_1 I_2 I_3 I_4}, C_{I_2 I_3}]_q \\
&= [[C_{I_1 I_2}, C_{I_2 I_3}], C_{I_2} C_{I_1 I_2 I_3}, C_{I_3 I_4}]_q + [C_{I_1 I_2 I_3}, C_{I_2 I_3 I_4}]_q \\
&\quad + C_{I_3} (C_{I_2} C_{I_1 I_2 I_3 I_4} - [C_{I_1 I_2}, C_{I_2 I_3 I_4}]) - C_{I_1 I_2 I_3 I_4} C_{I_2 I_3} \\
&= [-C_{I_1 I_3} + C_{I_1} C_{I_3}, C_{I_3 I_4}]_q + [C_{I_1 I_2 I_3}, C_{I_2 I_3 I_4}]_q + C_{I_3} (C_{I_1 I_3 I_4} - C_{I_1} C_{I_3 I_4}) - C_{I_1 I_2 I_3 I_4} C_{I_2 I_3} \\
&= -[C_{I_1 I_3}, C_{I_3 I_4}]_q + [C_{I_1 I_2 I_3}, C_{I_2 I_3 I_4}]_q - C_{I_1 I_2 I_3 I_4} C_{I_2 I_3} + C_{I_3} C_{I_1 I_3 I_4} \\
&= -[C_{I_1 I_3}, C_{I_3 I_4}]_q - C_{I_1 I_4} + C_{I_1} C_{I_4} + C_{I_3} C_{I_1 I_3 I_4} \\
&= 0.
\end{aligned}$$

We also have

$$\begin{aligned}
&[[\phi(a_{j+1,m}), \phi(a_{k,l})], \phi(a_{i,j})]_q + [\phi(a_{k,m}), \phi(a_{i,l})]_q + [[\phi(a_{j+1,l}), \phi(a_{k,j})], \phi(a_{i,m})]_q \\
&\quad - [[\phi(a_{j+1,m}), \phi(a_{k,j})], \phi(a_{i,l})]_q - [[\phi(a_{j+1,l}), \phi(a_{k,m})], \phi(a_{i,j})]_q - [\phi(a_{k,l}), \phi(a_{i,m})]_q \\
&= [[C_{I_3 I_4}, C_{I_2 I_3}], C_{I_1 I_2}]_q + [C_{I_2 I_3 I_4}, C_{I_1 I_2 I_3}]_q + [[C_{I_3}, C_{I_2}], C_{I_1 I_2 I_3 I_4}]_q \\
&\quad - [[C_{I_3 I_4}, C_{I_2}], C_{I_1 I_2 I_3}]_q - [[C_{I_3}, C_{I_2 I_3 I_4}], C_{I_1 I_2}]_q - [C_{I_2 I_3}, C_{I_1 I_2 I_3 I_4}]_q \\
&= [[C_{I_3 I_4}, C_{I_2 I_3}], C_{I_3} C_{I_2 I_3 I_4}, C_{I_1 I_2}]_q + [C_{I_2 I_3 I_4}, C_{I_1 I_2 I_3}]_q \\
&\quad + C_{I_2} (C_{I_3} C_{I_1 I_2 I_3 I_4} - [C_{I_3 I_4}, C_{I_1 I_2 I_3}]) - C_{I_2 I_3} C_{I_1 I_2 I_3 I_4} \\
&= [-C_{I_4 I_2} + C_{I_2} C_{I_4}, C_{I_1 I_2}]_q + [C_{I_2 I_3 I_4}, C_{I_1 I_2 I_3}]_q + C_{I_2} (C_{I_4 I_2 I_1} - C_{I_4} C_{I_1 I_2}) - C_{I_2 I_3} C_{I_1 I_2 I_3 I_4} \\
&= -[C_{I_4 I_2}, C_{I_1 I_2}]_q + C_{I_2} C_{I_4 I_2 I_1} + ([C_{I_2 I_3 I_4}, C_{I_1 I_2 I_3}]_q - C_{I_2 I_3} C_{I_1 I_2 I_3 I_4})
\end{aligned}$$

$$\begin{aligned}
&= -[C_{l_4 l_2}, C_{l_1 l_2}]_q + C_{l_2} C_{l_4 l_2 l_1} - C_{l_4 l_1} + C_{l_1} C_{l_4} \\
&= 0.
\end{aligned}$$

Hence,

$$\det_q \left( \phi \left( \mathcal{A}_{i,k,j+1}^{j,l,m} \right) \right) = \det^q \left( \phi \left( \mathcal{A}_{i,k,j+1}^{j,l,m} \right) \right) = 0.$$

The above statements mean that  $\phi$  is indeed an algebra homomorphism.

Furthermore, we define a map  $\varphi : \mathfrak{aw}(n) \rightarrow \mathcal{A}(n)$ , which sends generator  $C_I$  of  $\mathfrak{aw}(n)$  to  $a_{i_I, j_I}$ , where  $i_I = \min I$  and  $j_I = \max I$ .

Now, we have  $i \leq j$  and  $I = \{i, i+1, \dots, j\}$  be a non-empty connected subset of  $\{1, 2, \dots, n\}$ . Similarly, it is straightforward to see that  $\varphi$  is an algebra homomorphism. Also,  $\varphi\phi(a_{i,j}) = \varphi(C_I) = a_{i,j}$  and  $\varphi\varphi(C_I) = \phi(a_{i,j}) = C_I$ . It follows that  $\phi$  is an isomorphism of algebras.

Thus

$$\mathcal{A}(n) \cong \mathfrak{aw}(n)$$

and the proof is finished.  $\square$

### 3 Automorphisms of $\mathcal{A}(n)$

In this section, we construct a series of automorphisms of  $\mathcal{A}(n)$ , which coincide with those of  $\mathfrak{aw}(n)$  in [15]. The proof of the results is sketch in [15] and slightly difficult to handle. Here we give the proof in detail.

For the upper triangular generating matrix  $\mathcal{A} = (a_{ij})_{n \times n}$  of the algebra  $\mathcal{A}(n)$  and the map  $f : \mathcal{A}(n) \rightarrow \mathcal{A}(n)$ , the notation  $f(\mathcal{A})$  is defined by the upper triangular matrix  $(f(\mathcal{A})_{i,j})_{n \times n}$ , where  $f(\mathcal{A})_{i,j} := f(a_{i,j})$ . For convenience, we denote

$$a_{1,j} \delta_0 \triangleq -[a_{2,n}, a_{1,j}]_q + a_{j+1,n} a_{1,1} + a_{2,j} a_{1,n}, \quad (3.1)$$

$$a_{1,j} \delta'_0 \triangleq -[a_{1,j}, a_{2,n}]_q + a_{j+1,n} a_{1,1} + a_{2,j} a_{1,n}, \quad (3.2)$$

$$a_{k,i} \delta_i \triangleq -[a_{i,i+1}, a_{k,i}]_q + a_{i+1,i+1} a_{k,i-1} + a_{i,i} a_{k,i+1}, \quad (3.3)$$

$$a_{i+1,l} \delta_i \triangleq -[a_{i,i+1}, a_{i+1,l}]_q + a_{i,i} a_{i+2,l} + a_{i+1,i+1} a_{i,l}, \quad (3.4)$$

$$a_{k,i} \delta'_i \triangleq -[a_{k,i}, a_{i,i+1}]_q + a_{i+1,i+1} a_{k,i-1} + a_{i,i} a_{k,i+1}, \quad (3.5)$$

$$a_{i+1,l} \delta'_i \triangleq -[a_{i,i+1}, a_{i+1,l}]_q + a_{i,i} a_{i+2,l} + a_{i+1,i+1} a_{i,l}, \quad (3.6)$$

where  $1 \leq i \leq n-1$ ,  $2 \leq j \leq n-1$ ,  $1 \leq k \leq i-1$  and  $i+1 < l \leq n-1$ .

Now, let  $\delta_0, \delta'_0 : \mathcal{A}(n) \rightarrow \mathcal{A}(n)$  the maps be given by

$$\delta_0(\mathcal{A}) = \begin{pmatrix} a_{1,n} & a_{1,2}^{\delta_0} & \cdots & a_{1,j}^{\delta_0} & \cdots & a_{1,n-1}^{\delta_0} & a_{1,1} \\ & a_{2,2} & \cdots & a_{2,j} & \cdots & a_{2,n-1} & a_{2,n} \\ & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ & & a_{j,j} & \cdots & a_{j,n-1} & a_{j,n} & \\ & & & \ddots & \vdots & \vdots & \\ & & & & a_{n-1,n-1} & a_{n-1,n} & \\ & & & & & a_{n,n} & \end{pmatrix},$$

$$\delta'_0(\mathcal{A}) = \begin{pmatrix} a_{1,n} & a_{1,2}^{\delta'_0} & \cdots & a_{1,j}^{\delta'_0} & \cdots & a_{1,n-1}^{\delta'_0} & a_{1,1} \\ & a_{2,2} & \cdots & a_{2,j} & \cdots & a_{2,n-1} & a_{2,n} \\ & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ & & a_{j,j} & \cdots & a_{j,n-1} & a_{j,n} & \\ & & & \ddots & \vdots & \vdots & \\ & & & & a_{n-1,n-1} & a_{n-1,n} & \\ & & & & & a_{n,n} & \end{pmatrix},$$

the map  $\delta_i, \delta'_i : \mathcal{A}(n) \rightarrow \mathcal{A}(n)$  ( $i \in [\![1, n-1]\!]$ ) be given by

$$\delta_i(\mathcal{A}) = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,i}^{\delta_i} & a_{1,i+1} & a_{1,i+2} & \cdots & a_{1,n} \\ & \ddots & & \vdots & \vdots & \vdots & \vdots & \vdots \\ & & a_{i-1,i}^{\delta_i} & a_{i-1,i+1} & a_{i-1,i+2} & \cdots & a_{i-1,n} & \\ & & a_{i+1,i+1} & a_{i,i+1} & a_{i,i+2} & \cdots & a_{i,n} \\ & & a_{i,i} & a_{i+1,i+2}^{\delta_i} & \cdots & a_{i+1,n}^{\delta_i} & & \\ & & & a_{i+2,i+2} & \cdots & a_{i+2,n} & & \\ & & & \ddots & & \vdots & & \\ & & & & & a_{n,n} & & \end{pmatrix},$$

$$\delta'_i(\mathcal{A}) = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,i}^{\delta'_i} & a_{1,i+1} & a_{1,i+2} & \cdots & a_{1,n} \\ & \ddots & & \vdots & \vdots & \vdots & \vdots & \vdots \\ & & a_{i-1,i}^{\delta'_i} & a_{i-1,i+1} & a_{i-1,i+2} & \cdots & a_{i-1,n} & \\ & & a_{i+1,i+1} & a_{i,i+1} & a_{i,i+2} & \cdots & a_{i,n} \\ & & a_{i,i} & a_{i+1,i+2}^{\delta'_i} & \cdots & a_{i+1,n}^{\delta'_i} & & \\ & & & a_{i+2,i+2} & \cdots & a_{i+2,n} & & \\ & & & \ddots & & \vdots & & \\ & & & & & a_{n,n} & & \end{pmatrix}.$$

For the maps  $\delta_0, \delta'_0$ , the first row  $a_{1,j}$  ( $2 \leq j \leq n-1$ ) of  $\mathcal{A}$  are mapping to  $a_{1,j}^{\delta_0}$  (resp.  $a_{1,j}^{\delta'_0}$ ),  $\delta_0(a_{1,1}) = a_{1,n}$ ,  $\delta_0(a_{1,n}) = a_{1,1}$ , and the other generators are fixed. Similarly, For the maps  $\delta_i, \delta'_i$  ( $i \in [\![1, n-1]\!]$ ), the generators of  $\mathcal{A}$  are fixed except those of the  $(i+1)$ -th row and the  $i$ -th column.

Subsequently, we aim to demonstrate that the maps  $\delta_i, \delta'_i (i \in \llbracket 0, n-1 \rrbracket)$  defined in this way are automorphisms and satisfy the braid group relations. We begin by introducing two lemmas.

**Lemma 3.1.** (1) For  $\mathcal{A}_{1,k,i+1}^{i,j,m}$  as (1.13), we have

$$\begin{aligned} \left[ a_{1,i}^{\delta_0}, a_{1,k-1}^{\delta_0} \right]_q &= a_{1,i} a_{1,k-1} - a_{1,1} [a_{2,i}, a_{1,k-1}]_q - a_{i+1,n} a_{1,n} a_{1,k-1} + a_{1,1} \left[ a_{1,i}^{\delta_0}, a_{k,n} \right]_q \\ &\quad + a_{2,k-1} a_{1,i}^{\delta_0} a_{1,n} + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ a_{1,k-1}, a_{1,i}^{\delta_0} \right] \right], \end{aligned} \quad (3.7)$$

$$\begin{aligned} \left[ a_{1,i}^{\delta_0}, a_{1,j}^{\delta_0} \right]_q &= a_{1,i} a_{1,j} - a_{1,1} a_{2,i} a_{1,j} - a_{1,n} [a_{i+1,n}, a_{1,j}]_q + a_{1,1} a_{1,i}^{\delta_0} a_{j+1,n} \\ &\quad + a_{1,n} \left[ a_{1,i}^{\delta_0}, a_{2,j} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ a_{1,j}, a_{1,i}^{\delta_0} \right] \right], \end{aligned} \quad (3.8)$$

$$\begin{aligned} \left[ a_{1,k-1}^{\delta_0}, a_{1,j}^{\delta_0} \right]_q &= a_{1,k-1} a_{1,j} - a_{1,1} a_{2,k-1} a_{1,j} - a_{1,n} [a_{k,n}, a_{1,j}]_q + a_{1,1} a_{1,k-1}^{\delta_0} a_{j+1,n} \\ &\quad + a_{1,n} \left[ a_{1,k-1}^{\delta_0}, a_{2,j} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ a_{1,j}, a_{1,k-1}^{\delta_0} \right] \right], \end{aligned} \quad (3.9)$$

$$\begin{aligned} \left[ a_{1,m}^{\delta_0}, a_{1,j}^{\delta_0} \right]_q &= a_{1,m} a_{1,j} - a_{1,1} [a_{2,m}, a_{1,j}]_q - a_{1,n} a_{m+1,n} a_{1,j} + a_{1,1} \left[ a_{1,m}^{\delta_0}, a_{j+1,n} \right]_q \\ &\quad + a_{1,n} a_{1,m}^{\delta_0} a_{2,j} + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ a_{1,j}, a_{1,m}^{\delta_0} \right] \right], \end{aligned} \quad (3.10)$$

$$\begin{aligned} \left[ a_{1,m}^{\delta_0}, a_{1,i}^{\delta_0} \right]_q &= a_{1,m} a_{1,i} - a_{1,1} [a_{2,m}, a_{1,i}]_q - a_{1,n} a_{m+1,n} a_{1,i} + a_{1,1} \left[ a_{1,m}^{\delta_0}, a_{i+1,n} \right]_q \\ &\quad + a_{1,n} a_{1,m}^{\delta_0} a_{2,i} + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ a_{1,i}, a_{1,m}^{\delta_0} \right] \right]. \end{aligned} \quad (3.11)$$

(2) For  $\mathcal{A}_{j,k,i+1}^{k-1,i,l}$  as (1.10), we have

$$\begin{aligned} \left[ a_{k,i}^{\delta_i}, a_{l+1,l}^{\delta_i} \right]_q &= a_{k,i} a_{l+1,l} - a_{k,i-1} a_{i,i} a_{l+1,l} + a_{k,i}^{\delta_i} a_{i,i} a_{l+2,l} - a_{l+1,i+1} [a_{k,l+1}, a_{l+1,l}]_q \\ &\quad + a_{i+1,i+1} \left[ a_{k,i}^{\delta_i}, a_{i,l} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{i+1,l}, a_{k,i}^{\delta_i} \right] \right], \end{aligned} \quad (3.12)$$

$$\begin{aligned} \left[ a_{j,i}^{\delta_i}, a_{k,i}^{\delta_i} \right]_q &= a_{j,i} a_{k,i} - a_{i+1,i+1} a_{j,i+1} a_{k,i} - a_{i,i} [a_{j,i-1}, a_{k,i}]_q + a_{j,i}^{\delta_i} a_{k,i-1} a_{i+1,i+1} \\ &\quad + a_{i,i} \left[ a_{j,i}^{\delta_i}, a_{k,i+1} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{k,i}, a_{j,i}^{\delta_i} \right] \right], \end{aligned} \quad (3.13)$$

$$\begin{aligned} \left[ a_{j,i}^{\delta_i}, a_{i+1,l}^{\delta_i} \right]_q &= a_{j,i} a_{i+1,l} - a_{j,i-1} a_{i,i} a_{i+1,l} + a_{j,i}^{\delta_i} a_{i,i} a_{i+2,l} - a_{i+1,i+1} [a_{j,i+1}, a_{i+1,l}]_q \\ &\quad + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, a_{i,l} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{i+1,l}, a_{j,i}^{\delta_i} \right] \right]. \end{aligned} \quad (3.14)$$

(2) For  $\mathcal{A}_{j,k,i+1}^{i,l,m}$  as (1.13), we have

$$\begin{aligned} \left[ a_{k,i}^{\delta_i}, a_{i+1,m}^{\delta_i} \right]_q &= a_{k,i} a_{i+1,m} - a_{k,i-1} a_{i,i} a_{i+1,m} + a_{k,i}^{\delta_i} a_{i,i} a_{i+2,m} - a_{i+1,i+1} [a_{k,i+1}, a_{i+1,m}]_q \\ &\quad + a_{i+1,i+1} \left[ a_{k,i}^{\delta_i}, a_{i,m} \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{i+1,m}, a_{k,i}^{\delta_i} \right] \right], \end{aligned} \quad (3.15)$$

$$\left[ a_{j,i}^{\delta_i}, a_{i+1,m}^{\delta_i} \right]_q = a_{j,i} a_{i+1,m} - a_{j,i-1} a_{i,i} a_{i+1,m} + a_{j,i}^{\delta_i} a_{i,i} a_{i+2,m} - a_{i+1,i+1} [a_{j,i+1}, a_{i+1,m}]_q$$

$$+a_{i+1,i+1}\left[a_{j,i}^{\delta_i}, a_{i,m}\right]_q + \frac{1}{(q-q^{-1})^2} \left[a_{i,i+1}, \left[a_{i+1,m}, a_{j,i}^{\delta_i}\right]\right]. \quad (3.16)$$

*Proof.* We only focus on proving only one of them, the others are similar and straightforward. For example

$$\begin{aligned} & \left[a_{1,i}^{\delta_0}, a_{1,k-1}^{\delta_0}\right]_q \\ \stackrel{(3.1)}{=} & \left[a_{1,i}^{\delta_0}, -[a_{2,n}, a_{1,k-1}]_q + a_{1,1}a_{k,n} + a_{2,k-1}a_{1,n}\right]_q \\ = & -\left[a_{1,i}^{\delta_0}, [a_{2,n}, a_{1,k-1}]_q\right]_q + a_{1,1}\left[a_{1,i}^{\delta_0}, a_{k,n}\right]_q + a_{2,k-1}a_{1,i}^{\delta_0}a_{1,n} \\ \stackrel{(1.3)}{=} & -\left[\left[a_{1,i}^{\delta_0}, a_{2,n}\right]_q, a_{1,k-1}\right]_q + a_{1,1}\left[a_{1,i}^{\delta_0}, a_{k,n}\right]_q + a_{2,k-1}a_{1,i}^{\delta_0}a_{1,n} \\ & + \frac{1}{(q-q^{-1})^2} \left[a_{2,n}, \left[a_{1,k-1}, a_{1,i}^{\delta_0}\right]\right] \\ \stackrel{(3.1)}{=} & [a_{1,i} - a_{1,1}a_{2,i} - a_{i+1,n}a_{1,n}, a_{1,k-1}]_q + a_{1,1}\left[a_{1,i}^{\delta_0}, a_{k,n}\right]_q + a_{2,k-1}a_{1,i}^{\delta_0}a_{1,n} \\ & + \frac{1}{(q-q^{-1})^2} \left[a_{2,n}, \left[a_{1,k-1}, a_{1,i}^{\delta_0}\right]\right] \\ = & a_{1,i}a_{1,k-1} - a_{1,1}[a_{2,i}, a_{1,k-1}]_q - a_{i+1,n}a_{1,n}a_{1,k-1} + a_{1,1}\left[a_{1,i}^{\delta_0}, a_{k,n}\right]_q \\ & + a_{2,k-1}a_{1,i}^{\delta_0}a_{1,n} + \frac{1}{(q-q^{-1})^2} \left[a_{2,n}, \left[a_{1,k-1}, a_{1,i}^{\delta_0}\right]\right]. \end{aligned}$$

This proof is finished.  $\square$

**Lemma 3.2.** 1. For  $\mathcal{A}_{1,k,i+1}^{k-1,i,j}$  as (1.10), we have

$$\begin{aligned} & \left[a_{2,i}, [a_{k,j}, a_{1,i}]_q - a_{i+1,j}a_{1,k-1} - a_{k,i}a_{1,j}\right]_q + a_{2,k-1}a_{1,j} \\ = & \left[a_{1,i}^{\delta_0}, [a_{k,j}, a_{i+1,n}]_q - a_{i+1,j}a_{k,n} - a_{k,i}a_{j+1,n}\right]_q + \left[a_{1,k-1}^{\delta_0}, a_{j+1,n}\right]_q, \end{aligned} \quad (3.17)$$

$$\begin{aligned} & \left[a_{i+1,n}, [a_{k,j}, a_{1,i}]_q - a_{i+1,j}a_{1,k-1} - a_{k,i}a_{1,j}\right]_q + [a_{k,n}, a_{1,j}]_q \\ = & \left[a_{1,i}^{\delta_0}, [a_{k,j}, a_{2,i}]_q - a_{k,i}a_{2,j} - a_{2,k-1}a_{i+1,j}\right]_q + \left[a_{1,k-1}^{\delta_0}, a_{2,j}\right]_q. \end{aligned} \quad (3.18)$$

2. For  $\mathcal{A}_{j,k,i+1}^{k-1,i,l}$  as (1.10), we have

$$\begin{aligned} & \left[a_{i+2,l}, [a_{k,l}, a_{j,i-1}]_q - a_{j,k-1}a_{i,l} - a_{k,i-1}a_{j,l}\right]_q + a_{k,i-1}a_{j,i+1} \\ = & \left[[a_{k,l}, a_{j,i}]_q - a_{j,k-1}a_{i+1,l} - a_{k,i}a_{j,l}, a_{i+1,l}^{\delta_i}\right]_q + \left[a_{k,i}, a_{j,i}^{\delta_i}\right]_q, \end{aligned} \quad (3.19)$$

$$\begin{aligned} & \left[[a_{k,l}, a_{j,i-1}]_q - a_{j,k-1}a_{i,l} - a_{k,i-1}a_{j,l}, a_{k,i+1}\right]_q + [a_{i,l}, a_{j,i+1}]_q \\ = & \left[[a_{k,l}, a_{j,i}]_q - a_{j,k-1}a_{i+1,l} - a_{k,i}a_{j,l}, a_{k,i}^{\delta_i}\right]_q + \left[a_{i+1,l}, a_{j,i}^{\delta_i}\right]_q. \end{aligned} \quad (3.20)$$

3. For  $\mathcal{A}_{j,k,i+1}^{i,l,m}$  as (1.13), we have

$$\begin{aligned} & a_{k,i-1} \left( [a_{j,l}, a_{i+1,m}]_q - a_{k,i-1} a_{i+1,l} a_{j,m} \right) + \left[ a_{j,i-1}, [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l} a_{k,m} \right]_q \\ = & a_{k,i} \delta_i \left( [a_{j,l}, a_{i+2,m}]_q - a_{i+2,l} a_{j,m} \right) + \left[ a_{j,i} \delta_i, a_{i+2,l} a_{k,m} - [a_{k,l}, a_{i+2,m}]_q \right]_q, \end{aligned} \quad (3.21)$$

$$\begin{aligned} & \left[ a_{j,i+1}, a_{i+1,l} a_{k,m} - [a_{k,l}, a_{i+1,m}]_q \right]_q + \left[ a_{k,i+1}, [a_{j,l}, a_{i+1,m}]_q - a_{i+1,l} a_{j,m} \right]_q \\ = & \left[ a_{j,i} \delta_i, a_{i,l} a_{k,m} - [a_{k,l}, a_{i,m}]_q \right]_q + \left[ a_{k,i} \delta_i, [a_{j,l}, a_{i,m}]_q - a_{i,l} a_{j,m} \right]_q, \end{aligned} \quad (3.22)$$

$$\begin{aligned} & a_{i+1,l} \delta_i \left( [a_{k,m}, a_{j,i-1}]_q - a_{k,i-1} a_{j,m} \right) + \left[ a_{i+1,m} \delta_i, a_{k,i-1} a_{j,l} - [a_{k,l}, a_{j,i-1}]_q \right]_q \\ = & a_{i+2,l} \left( [a_{k,m}, a_{j,i}]_q - a_{i+2,l} a_{k,i} a_{j,m} \right) + \left[ a_{i+2,m}, [a_{k,l}, a_{j,i}]_q - a_{k,i} a_{j,l} \right]_q, \end{aligned} \quad (3.23)$$

$$\begin{aligned} & \left[ a_{i+1,m} \delta_i, a_{k,i+1} a_{j,l} - [a_{k,l}, a_{j,i+1}]_q \right]_q + \left[ a_{i+1,l} \delta_i, [a_{k,m}, a_{j,i+1}]_q - a_{k,i+1} a_{j,m} \right]_q \\ = & \left[ a_{i,m}, a_{k,i} a_{j,l} - [a_{k,l}, a_{j,i}]_q \right]_q + \left[ a_{i,l}, [a_{k,m}, a_{j,i}]_q - a_{k,i} a_{j,m} \right]_q. \end{aligned} \quad (3.24)$$

*Proof.* We only focus on proving only one of them, the others are similar and straightforward. For example

$$\begin{aligned} & \left[ a_{2,i}, [a_{k,j}, a_{1,i}]_q - a_{i+1,j} a_{1,k-1} \right]_q - a_{k,i} a_{2,i} a_{1,j} + a_{2,k-1} a_{1,j} \\ \stackrel{(4.b)}{=} & \left( a_{k,i} [a_{2,j}, a_{1,i}]_q - a_{1,1} a_{k,i} a_{i+1,j} - [a_{2,j}, a_{1,k-1}]_q + a_{1,1} a_{k,j} \right) + a_{k,i} a_{2,i} a_{1,j} + a_{2,k-1} a_{1,j} \\ = & \left( - [a_{2,j}, a_{1,k-1}]_q + a_{2,k-1} a_{1,j} \right) - a_{k,i} \left( - [a_{2,j}, a_{1,i}]_q + a_{2,i} a_{1,j} \right) + a_{1,1} a_{k,j} - a_{1,1} a_{k,i} a_{i+1,j} \\ \stackrel{(2.a)}{=} & \left( \left[ a_{2,n}, [a_{1,j}, a_{k,n}]_q \right]_q - a_{1,n} [a_{2,j}, a_{k,n}]_q + a_{2,k-1} a_{j+1,n} a_{1,n} - a_{j+1,n} [a_{2,n}, a_{1,k-1}]_q \right) \\ & - a_{k,i} \left( \left[ a_{2,n}, [a_{1,j}, a_{i+1,n}]_q - a_{j+1,n} [a_{2,n}, a_{1,i}]_q + a_{j+1,n} a_{2,i} a_{1,n} - a_{1,n} [a_{2,j}, a_{i+1,n}]_q \right]_q \right. \\ & \left. + a_{1,1} a_{k,j} - a_{1,1} a_{k,i} a_{i+1,j} \right) \\ = & \left[ a_{2,n}, [a_{1,j}, a_{k,n}]_q - a_{k,i} \left( [a_{1,j}, a_{i+1,n}]_q + a_{j+1,n} a_{1,i} \right) \right]_q + a_{1,1} (a_{k,j} - a_{k,i} a_{i+1,j}) \\ & - a_{1,n} \left( [a_{2,j}, a_{k,n}]_q - a_{2,k-1} a_{j+1,n} + a_{k,i} \left( [a_{2,j}, a_{i+1,n}]_q - a_{j+1,n} a_{2,i} \right) \right) \\ & - a_{j+1,n} [a_{2,n}, a_{1,k-1}]_q \\ \stackrel{(1.16)}{=} & \left[ [a_{2,n}, a_{1,i}]_q, - [a_{k,j}, a_{i+1,n}]_q + a_{k,i} a_{j+1,n} + a_{i+1,j} a_{k,n} \right]_q - a_{j+1,n} [a_{2,n}, a_{1,k-1}]_q \\ & - a_{1,1} \left( \left[ a_{i+1,n}, - [a_{k,j}, a_{i+1,n}]_q + a_{k,i} a_{j+1,n} + a_{i+1,j} a_{k,n} \right]_q - a_{j+1,n} a_{k,n} \right) \\ & - a_{1,n} \left( \left[ a_{2,i}, - [a_{k,j}, a_{i+1,n}]_q + a_{k,i} a_{j+1,n} + a_{i+1,j} a_{k,n} \right]_q + a_{2,k-1} a_{j+1,n} \right) \\ = & \left[ a_{1,i} \delta_0, [a_{k,j}, a_{i+1,n}]_q - a_{i+1,j} a_{k,n} - a_{k,i} a_{j+1,n} \right]_q + \left[ a_{1,k-1} \delta_0, a_{j+1,n} \right]_q. \end{aligned}$$

This proof is finished.  $\square$

**Proposition 3.3.**  $\delta_0, \delta'_0$  are the automorphisms of algebra  $\mathcal{A}(n)$  and  $\delta_0 \delta'_0 = \delta'_0 \delta_0 = \text{id}$ .

*Proof.* To prove that  $\delta_0$  is an algebra homomorphism, it is sufficient to demonstrate that  $\delta_0$  keeps the relations of  $\mathcal{A}(n)$  associating with  $a_{1,i}, i \in [\![1, n]\!]$ .

For the relations (R1). If  $i = 1$  or  $i = n$ ,  $a_{1,1}$  and  $a_{1,n}$  are in the centers of  $\mathcal{A}(n)$ . By the definition of  $\delta_0$ ,  $\delta_0(a_{1,1}) = a_{1,n}$ ,  $\delta_0(a_{1,n}) = a_{1,1}$ , which are also in the center of  $\mathcal{A}(n)$ . We suppose that  $i \in \llbracket 2, n-1 \rrbracket$ . If there exists  $a_{j,k}$  such that  $[a_{1,i}, a_{k,l}] = 0$ , we consider two cases:

When  $j = 1$  and  $k \neq n$ , let's assume that  $i < k$ , we have

$$\begin{aligned}
& [\delta_0(a_{1,i}), \delta_0(a_{1,k})] \\
\stackrel{(3.1)}{=} & \left[ a_{1,i}^{\delta_0}, -[a_{2,n}, a_{1,k}]_q + a_{k+1,n}a_{1,1} + a_{2,k}a_{1,n} \right] \\
\stackrel{(R1)}{=} & \left[ a_{1,i}^{\delta_0}, -[a_{2,n}, a_{1,k}]_q + a_{2,k}a_{1,n} \right] \\
\stackrel{(1.17)}{=} & \left[ a_{1,i}^{\delta_0}, \left[ [a_{i+1,n}, a_{2,k}]_q, a_{1,i} \right]_q - a_{2,i}[a_{i+1,n}, a_{1,k}]_q + a_{i+1,k} \left( a_{2,i}a_{1,n} - [a_{2,n}, a_{1,i}]_q \right) \right] \\
\stackrel{(3.1)}{=} & \left[ a_{1,i}^{\delta_0}, \left[ [a_{i+1,n}, a_{2,k}]_q, a_{1,i} \right]_q - a_{2,i}[a_{i+1,n}, a_{1,k}]_q + a_{i+1,k} \left( a_{1,i}^{\delta_0} - a_{i+1,n}a_{1,1} \right) \right] \\
\stackrel{(R1)}{=} & \left[ a_{1,i}^{\delta_0}, \left[ [a_{i+1,n}, a_{2,k}]_q, a_{1,i} \right]_q - a_{2,i}[a_{i+1,n}, a_{1,k}]_q \right] \\
\stackrel{(1.7)}{=} & \left[ a_{1,i}^{\delta_0}, \left[ a_{i+1,n}, [a_{2,k}, a_{1,i}]_q \right]_q - a_{2,i}[a_{i+1,n}, a_{1,k}]_q \right] \\
= & \left[ a_{1,i}^{\delta_0}, \left[ a_{i+1,n}, [a_{2,k}, a_{1,i}]_q - a_{2,i}a_{1,k} \right]_q \right] \\
\stackrel{(1.5)}{=} & - \left[ a_{i+1,n}, \left[ [a_{2,k}, a_{1,i}]_q - a_{2,i}a_{1,k}, a_{1,i}^{\delta_0} \right]_q \right] - \left[ [a_{2,k}, a_{1,i}]_q - a_{2,i}a_{1,k}, \right. \\
& \quad \left. \left[ a_{1,i}^{\delta_0}, a_{i+1,n} \right]_q \right] \\
\stackrel{(1.8)}{=} & -a_{1,i}^{\delta_0} \left[ a_{i+1,n}, [a_{2,k}, a_{1,i}]_q - a_{2,i}a_{1,k} \right] - a_{1,i}^{\delta_0} \left[ [a_{2,k}, a_{1,i}]_q - a_{2,i}a_{1,k}, a_{i+1,n} \right] \\
\stackrel{(1.1)}{=} & 0.
\end{aligned}$$

When  $\delta_0(a_{k,l}) = a_{k,l}$ , we have

$$\begin{aligned}
& [\delta_0(a_{1,i}), \delta_0(a_{j,k})] \\
\stackrel{(\delta_0)}{=} & \left[ a_{j,i}^{\delta_0}, a_{j,k} \right] \\
\stackrel{(3.1)}{=} & \left[ a_{j,k}, -[a_{2,n}, a_{1,i}]_q + a_{i+1,n}a_{1,1} + a_{2,i}a_{1,n} \right] \\
= & - \left[ [a_{2,n}, a_{1,i}]_q, a_{j,k} \right] + \left[ a_{i+1,n}a_{1,1}, a_{j,k} \right] + \left[ a_{2,i}a_{1,n}, a_{j,k} \right] \\
\stackrel{(R1)}{=} & 0.
\end{aligned}$$

Consequently, if  $[a_{i,j}, a_{k,l}] = 0$ , then  $\left[ a_{i,j}^{\delta_0}, a_{k,l}^{\delta_0} \right] = 0$ , which implies that

$$\left[ a_{i,j}^{\delta_0}, a_{k,l}^{\delta_0} \right]_q \stackrel{(1.6)}{=} a_{i,j}^{\delta_0}a_{k,l}^{\delta_0},$$

In the subsequent, we directly use this fact without further explanation..

For the relations (R2). If we have chosen the sub-matrix  $\mathcal{A}_{1,k,i+1}^{k-1,i,j}$  as (1.10), the proof deduce to the simpler case when  $k = 2$  or  $j = n$  and straightforward. In more tedious calculations, we concentrate on exploring the general cases where  $k \neq 2$  and  $j \neq n$  and have

$$\begin{aligned}
& \left[ \left[ \delta_0(a_{k,j}), \delta_0(a_{1,i}) \right]_q, \delta_0(a_{k,j}) \right]_q \\
\stackrel{(3.1)}{\stackrel{(1.7)}{=}} & - \left[ a_{2,n}, \left[ [a_{k,j}, a_{1,i}]_q, a_{k,j} \right]_q + a_{1,1} \left[ [a_{k,j}, a_{i+1,n}]_q, a_{k,j} \right]_q \right. \\
& \left. + a_{1,n} \left[ [a_{k,j}, a_{2,i}]_q, a_{k,j} \right]_q \right. \\
\stackrel{(1.16)}{=} & - \left[ a_{2,n}, f(\mathcal{A}_{1,k,i+1}^{k-1,i,j}) \right]_q + a_{1,1}g(\mathcal{A}_{k,i+1,n}^{i,j,n}) + a_{1,n}f(\mathcal{A}_{2,k,i+1}^{k-1,i,j}) \\
\stackrel{(1.11)}{\stackrel{(1.12)}{=}} & - [a_{2,n}, a_{1,i}]_q + a_{1,1}a_{i+1,n} + a_{2,i}a_{1,n} \\
& + a_{i+1,j}a_{k,j} \left( - [a_{2,n}, a_{1,k-1}]_q + a_{1,1}a_{k,n} + a_{2,k-1}a_{1,n} \right) \\
& + a_{k,i}a_{k,j} \left( - [a_{2,n}, a_{1,j}]_q + a_{1,1}a_{n,n} + a_{2,j}a_{1,n} \right) \\
& - a_{k,i} \left( - [a_{2,n}, a_{1,k-1}]_q + a_{1,1}a_{k,n} + a_{2,k-1}a_{1,n} \right) \\
& - a_{i+1,j} \left( - [a_{2,n}, a_{1,j}]_q + a_{1,1}a_{n,n} + a_{2,j}a_{1,n} \right) \\
\stackrel{(3.1)}{=} & a_{1,i}^{\delta_0} + a_{i+1,j}a_{k,j}a_{1,k-1}^{\delta_0} + a_{k,i}a_{k,j}a_{1,j}^{\delta_0} - a_{k,i}a_{1,k-1}^{\delta_0} - a_{i+1,j}a_{1,j}^{\delta_0} \\
\stackrel{(\delta_0)}{=} & \delta_0(a_{1,i}) + \delta_0(a_{1,k-1}) \delta_0(a_{i+1,j}) \delta_0(a_{k,j}) + \delta_0(a_{k,i}) \delta_0(a_{1,j}) \delta_0(a_{k,j}) \\
& - \delta_0(a_{1,k-1}) \delta_0(a_{k,i}) - \delta_0(a_{i+1,j}) \delta_0(a_{1,j})
\end{aligned}$$

and

$$\begin{aligned}
& \left[ \delta_0(a_{1,i}), \left[ \delta_0(a_{k,j}), \delta_0(a_{1,i}) \right]_q \right]_q \\
\stackrel{(3.1)}{\stackrel{(\delta_0)}{=}} & - \left[ a_{1,i}^{\delta_0}, \left[ a_{2,n}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q + a_{1,1} \left[ a_{k,j}, a_{i+1,n} \right]_q + a_{1,n} \left[ a_{k,j}, a_{2,i} \right]_q \right]_q \\
\stackrel{(1.3)}{=} & \left( - \left[ \left[ a_{1,i}^{\delta_0}, a_{2,n} \right]_q, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ \left[ a_{k,j}, a_{1,i} \right]_q, a_{1,i}^{\delta_0} \right] \right] \right) \\
& + a_{1,1} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{i+1,n} \right]_q \right]_q + a_{1,n} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{2,i} \right]_q \right]_q \\
\stackrel{(3.1)}{\stackrel{(1.16)}{=}} & \left[ a_{1,i}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q - a_{1,1} \left[ a_{2,i}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q - a_{1,n} \left[ a_{i+1,n}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q \\
& + a_{1,1} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{i+1,n} \right]_q \right]_q + a_{1,n} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{2,i} \right]_q \right]_q \\
& + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ \left[ a_{k,j}, a_{1,i} \right]_q, a_{1,i}^{\delta_0} \right] \right] \\
\stackrel{(1.16)}{=} & g(\mathcal{A}_{1,k,i+1}^{k-1,i,j}) - a_{1,1} \left[ a_{2,i}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q - a_{1,n} \left[ a_{i+1,n}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q \\
& + a_{1,1} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{i+1,n} \right]_q \right]_q + a_{1,n} \left[ a_{1,i}^{\delta_0}, \left[ a_{k,j}, a_{2,i} \right]_q \right]_q \\
& + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ \left[ a_{k,j}, a_{1,i} \right]_q, a_{1,i}^{\delta_0} \right] \right] \\
\stackrel{(1.12)}{=} & a_{k,j} - a_{2,i}a_{i+1,j} - a_{1,1} \left[ a_{2,i}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q - a_{1,n} \left[ a_{i+1,n}, \left[ a_{k,j}, a_{1,i} \right]_q \right]_q
\end{aligned}$$

$$\begin{aligned}
& + a_{1,1} \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{i+1,n}]_q \right]_q + a_{1,n} \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{2,i}]_q \right]_q + a_{1,i} a_{1,k-1} a_{i+1,j} \\
& + a_{1,i} a_{k,i} a_{1,j} - a_{1,k-1} a_{1,j} + \frac{1}{(q-q^{-1})^2} \left[ a_{2,n}, \left[ [a_{k,j}, a_{1,i}]_q, a_{1,i}^{\delta_0} \right] \right] \\
\stackrel{(3.7)(3.8)}{\stackrel{(3.9)}{=}} & a_{k,j} - a_{2,i} a_{i+1,j} - a_{1,1} \left[ a_{2,i}, [a_{k,j}, a_{1,i}]_q \right]_q - a_{1,n} \left[ a_{i+1,n}, [a_{k,j}, a_{1,i}]_q \right]_q \\
& + a_{1,1} \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{i+1,n}]_q \right]_q + a_{1,n} \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{2,i}]_q \right]_q \\
& + a_{i+1,j} \left( a_{1,i}^{\delta_0} a_{1,k-1}^{\delta_0} + a_{1,1} [a_{2,i}, a_{1,k-1}]_q + a_{i+1,n} a_{1,n} a_{1,k-1} \right. \\
& \quad \left. - a_{1,1} \left[ a_{1,i}^{\delta_0}, a_{k,n} \right]_q - a_{2,k-1} a_{1,i}^{\delta_0} a_{1,n} \right) + a_{k,i} \left( a_{1,i}^{\delta_0} a_{1,j}^{\delta_0} + a_{1,1} a_{2,j} a_{1,j} \right. \\
& \quad \left. + a_{1,n} [a_{i+1,n}, a_{1,j}]_q - a_{1,1} a_{1,i}^{\delta_0} a_{j+1,n} - a_{1,n} \left[ a_{1,i}^{\delta_0}, a_{2,j} \right]_q \right) - \left( a_{1,k-1}^{\delta_0} a_{1,j}^{\delta_0} \right. \\
& \quad \left. + a_{1,1} a_{2,k-1} a_{1,j} + a_{1,n} [a_{k,n}, a_{1,j}]_q - a_{1,1} a_{1,k-1}^{\delta_0} a_{j+1,n} - a_{1,n} \left[ a_{1,k-1}^{\delta_0}, a_{2,j} \right]_q \right) \\
& + \frac{1}{(q-q^{-1})^2} \left( \left[ a_{2,n}, \left[ [a_{k,j}, a_{1,i}]_q - a_{i+1,j} a_{1,k-1} - a_{k,i} a_{1,j}, a_{1,i}^{\delta_0} \right] + \left[ a_{1,j}, a_{1,k-1}^{\delta_0} \right] \right] \right) \\
\stackrel{(1.2)}{=} & a_{k,j} - a_{2,i} a_{i+1,j} + a_{i+1,j} a_{1,i}^{\delta_0} a_{1,k-1}^{\delta_0} + a_{k,i} a_{1,i}^{\delta_0} a_{1,j}^{\delta_0} - a_{1,k-1}^{\delta_0} a_{1,j}^{\delta_0} \\
& - a_{1,1} \left( \left[ a_{2,i}, [a_{k,j}, a_{1,i}]_q - a_{i+1,j} a_{1,k-1} - a_{k,i} a_{1,j} \right]_q + a_{1,k-1}^{\delta_0} a_{j+1,n} \right) \\
& + a_{1,1} \left( \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{i+1,n}]_q - a_{i+1,j} a_{k,n} - a_{k,i} a_{j+1,n} \right]_q + a_{2,k-1} a_{1,j} \right) \\
& - a_{1,n} \left( \left[ a_{i+1,n}, [a_{k,j}, a_{1,i}]_q - a_{i+1,j} a_{1,k-1} - a_{k,i} a_{1,j} \right]_q + [a_{k,n}, a_{1,j}]_q \right) \\
& + a_{1,n} \left( \left[ a_{1,i}^{\delta_0}, [a_{k,j}, a_{2,i}]_q - a_{k,i} a_{2,j} - a_{2,k-1} a_{i+1,j} \right]_q + \left[ a_{1,k-1}^{\delta_0}, a_{2,j} \right]_q \right) \\
\stackrel{(3.17)}{\stackrel{(3.18)}{=}} & a_{k,j} - a_{2,i} a_{i+1,j} + a_{i+1,j} a_{1,i}^{\delta_0} a_{1,k-1}^{\delta_0} + a_{k,i} a_{1,i}^{\delta_0} a_{1,j}^{\delta_0} - a_{1,k-1}^{\delta_0} a_{1,j}^{\delta_0} \\
\equiv & \delta_0(a_{k,j}) - \delta_0(a_{2,i}) \delta_0(a_{i+1,j}) + \delta_0(a_{i+1,j}) \delta_0(a_{1,i}) \delta_0(a_{1,k-1}) \\
& + \delta_0(a_{k,i}) \delta_0(a_{1,i}) \delta_0(a_{1,j}) - \delta_0(a_{1,k-1}) \delta_0(a_{1,j}).
\end{aligned}$$

Hence,

$$\begin{aligned}
f \left( \delta_0 \left( \mathcal{A}_{1,k,i+1}^{k-1,i,j} \right) \right) & = \left[ \delta_0(a_{k,j}), [\delta_0(a_{1,i}), \delta_0(a_{k,j})]_q \right]_q, \\
g \left( \delta_0 \left( \mathcal{A}_{1,k,i+1}^{k-1,i,j} \right) \right) & = \left[ \delta_0(a_{1,i}), [\delta_0(a_{k,j}), \delta_0(a_{1,i})]_q \right]_q.
\end{aligned}$$

For the relations (R3). Choosing the sub-matrix  $\mathcal{A}_{1,k,i+1}^{i,j,l}$  ( $l \neq n$ ), we have

$$\begin{aligned}
& \left[ [\delta_0(a_{i+1,l}), \delta_0(a_{k,j})]_q, \delta_0(a_{1,i}) \right]_q + [\delta_0(a_{k,l}), \delta_0(a_{1,j})]_q + \delta_0(a_{i+1,j}) \delta_0(a_{k,i}) \delta_0(a_{1,l}) \\
& - \delta_0(a_{k,i}) [\delta_0(a_{i+1,l}), \delta_0(a_{1,j})]_q - \delta_0(a_{i+1,j}) [\delta_0(a_{k,l}), \delta_0(a_{1,i})]_q - \delta_0(a_{k,j}) \delta_0(a_{1,l}) \\
\stackrel{(3.1)}{\stackrel{(\delta_0)}{=}} & \left[ [a_{i+1,l}, a_{k,j}]_q, -[a_{2,n}, a_{1,i}]_q + a_{1,1} a_{i+1,n} + a_{2,i} a_{1,n} \right]_q \\
& + \left[ a_{k,l}, -[a_{2,n}, a_{1,j}]_q + a_{1,1} a_{j+1,n} + a_{2,j} a_{1,n} \right]_q
\end{aligned}$$

$$\begin{aligned}
& + a_{i+1,j} a_{k,i} \left( -[a_{2,n}, a_{1,l}]_q + a_{1,1} a_{l+1,n} + a_{2,l} a_{1,n} \right) \\
& - a_{k,i} \left[ a_{i+1,l}, -[a_{2,n}, a_{1,j}]_q + a_{1,1} a_{j+1,n} + a_{2,j} a_{1,n} \right]_q \\
& - a_{i+1,j} \left[ a_{k,l}, -[a_{2,n}, a_{1,i}]_q + a_{1,1} a_{i+1,n} + a_{2,i} a_{1,n} \right]_q \\
& - a_{k,j} \left( -[a_{2,n}, a_{1,l}]_q + a_{1,1} a_{l+1,n} + a_{2,l} a_{1,n} \right) \\
\underline{\underline{(1.7)}} \quad & - \left[ a_{2,n}, \left[ [a_{i+1,l}, a_{k,j}]_q, a_{1,i} \right]_q \right]_q + a_{1,1} \left[ [a_{i+1,l}, a_{k,j}]_q, a_{i+1,n} \right]_q \\
& + a_{1,n} \left[ [a_{i+1,l}, a_{k,j}]_q, a_{2,i} \right]_q - \left[ a_{2,n}, [a_{k,l}, a_{1,j}]_q \right]_q + a_{1,1} [a_{k,l}, a_{j+1,n}]_q \\
& + a_{1,n} [a_{k,l}, a_{2,j}]_q - [a_{2,n}, a_{i+1,j} a_{k,i} a_{1,l}]_q + a_{1,1} a_{i+1,j} a_{k,i} a_{l+1,n} + a_{i+1,j} a_{k,i} a_{2,l} a_{1,n} \\
& + \left[ a_{2,n}, a_{k,i} [a_{i+1,l}, a_{1,j}]_q \right]_q - a_{1,1} a_{k,i} [a_{i+1,l}, a_{j+1,n}]_q - a_{k,i} a_{1,n} [a_{i+1,l}, a_{2,j}]_q \\
& + \left[ a_{2,n}, a_{i+1,j} [a_{k,l}, a_{1,i}]_q \right]_q - a_{1,1} a_{i+1,j} [a_{k,l}, a_{i+1,n}]_q - a_{i+1,j} a_{1,n} [a_{k,l}, a_{2,i}]_q \\
& + \left[ a_{2,n}, a_{k,j} a_{1,l} \right]_q - a_{1,1} a_{l+1,n} a_{k,j} - a_{k,j} a_{2,l} a_{1,n} \\
\underline{\underline{(1.15)}} \quad & - \left[ a_{2,n}, \det^q \left( \mathcal{A}_{1,k,i+1}^{i,j,l} \right) \right]_q + a_{1,1} \left( \left[ a_{i+1,l}, [a_{k,j}, a_{i+1,n}]_q \right]_q + a_{i+1,j} a_{k,i} a_{l+1,n} \right. \\
& \left. + [a_{k,l}, a_{j+1,n}]_q - a_{k,i} [a_{i+1,l}, a_{j+1,n}]_q - a_{i+1,j} [a_{k,l}, a_{i+1,n}]_q - a_{l+1,n} a_{k,j} \right) \\
& + a_{1,n} \det^q \left( \mathcal{A}_{2,k,i+1}^{i,j,l} \right) \\
\underline{\underline{(1.17)}} \quad & 0 \\
\underline{\underline{(6.a)}} \quad &
\end{aligned}$$

and

$$\begin{aligned}
& \left[ [\delta_0(a_{1,i}), \delta_0(a_{k,j})]_q, \delta_0(a_{i+1,l}) \right]_q + [\delta_0(a_{1,j}), \delta_0(a_{k,l})]_q + \delta_0(a_{k,i}) [\delta_0(a_{i+1,j}), \delta_0(a_{1,l})]_q \\
& - \delta_0(a_{i+1,j}) [\delta_0(a_{1,i}), \delta_0(a_{k,l})]_q - \delta_0(a_{k,i}) [\delta_0(a_{1,j}), \delta_0(a_{i+1,l})]_q - \delta_0(a_{k,j}) \delta_0(a_{1,l}) \\
\underline{\underline{(3.1)}} \quad & \left[ -[a_{2,n}, a_{1,i}]_q + a_{1,1} a_{i+1,n} + a_{2,i} a_{1,n}, [a_{k,j}, a_{i+1,l}]_q \right]_q \\
& + \left[ -[a_{2,n}, a_{1,j}]_q + a_{1,1} a_{j+1,n} + a_{2,j} a_{1,n}, a_{k,l} \right]_q \\
& + a_{k,i} a_{i+1,j} \left( -[a_{2,n}, a_{1,l}]_q + a_{1,1} a_{l+1,n} + a_{2,l} a_{1,n} \right) \\
& - a_{i+1,j} \left[ -[a_{2,n}, a_{1,i}]_q + a_{1,1} a_{i+1,n} + a_{2,i} a_{1,n}, a_{k,l} \right]_q \\
& - a_{k,i} \left[ -[a_{2,n}, a_{1,j}]_q + a_{1,1} a_{j+1,n} + a_{2,j} a_{1,n}, a_{i+1,l} \right]_q \\
& - a_{k,j} \left( -[a_{2,n}, a_{1,l}]_q + a_{1,1} a_{l+1,n} + a_{2,l} a_{1,n} \right) \\
\underline{\underline{(1.7)}} \quad & - \left[ a_{2,n}, \left[ a_{1,i}, [a_{k,j}, a_{i+1,l}]_q \right]_q \right]_q + a_{1,1} \left[ a_{i+1,n}, [a_{k,j}, a_{i+1,l}]_q \right]_q \\
& + a_{1,n} \left[ a_{2,i}, [a_{k,j}, a_{i+1,l}]_q \right]_q - \left[ a_{2,n}, [a_{1,j}, a_{k,l}]_q \right]_q + a_{1,1} [a_{j+1,n}, a_{k,l}]_q \\
& + a_{1,n} [a_{2,j}, a_{k,l}]_q - [a_{2,n}, a_{k,i} a_{i+1,j} a_{1,l}]_q + a_{1,1} a_{k,i} a_{i+1,j} a_{l+1,n} + a_{k,i} a_{i+1,j} a_{2,l} a_{1,n}
\end{aligned}$$

$$\begin{aligned}
& + \left[ a_{2,n}, a_{i+1,j} [a_{1,i}, a_{k,l}]_q \right]_q - a_{i+1,j} a_{1,1} [a_{i+1,n}, a_{k,l}]_q - a_{i+1,j} a_{1,n} [a_{2,i}, a_{k,l}]_q \\
& + \left[ a_{2,n}, a_{k,i} [a_{1,j}, a_{i+1,l}]_q \right]_q - a_{k,i} a_{1,1} [a_{j+1,n}, a_{i+1,l}]_q - a_{k,i} a_{1,n} [a_{2,j}, a_{i+1,l}]_q \\
& + \left[ a_{2,n}, a_{k,j} a_{1,l} \right]_q - a_{1,1} a_{l+1,n} a_{k,j} - a_{k,j} a_{2,l} a_{1,n} \\
\stackrel{(1.14)}{=} & - \left[ a_{2,n}, \det_q (\mathcal{A}_{1,k,i+1}^{i,j,l}) \right]_q + a_{1,1} \left( \left[ a_{i+1,n}, [a_{k,j}, a_{i+1,l}]_q \right]_q + [a_{j+1,n}, a_{k,l}]_q \right. \\
& \left. + a_{k,i} a_{i+1,j} a_{l+1,n} - a_{i+1,j} [a_{i+1,n}, a_{k,l}]_q - a_{k,i} [a_{j+1,n}, a_{i+1,l}]_q - a_{1,1} a_{l+1,n} a_{k,j} \right) \\
& + a_{1,n} \det_q (\mathcal{A}_{2,k,i+1}^{i,j,l}) \\
\stackrel{(1.17)}{=} & 0. \\
\end{aligned}$$

Hence,

$$\det^q (\delta_0 (\mathcal{A}_{1,k,i+1}^{i,j,l})) = \det_q (\delta_0 (\mathcal{A}_{1,k,i+1}^{i,j,l})) = 0.$$

In particular, when  $l = n$ , we have

$$\det_q (\delta_0 (\mathcal{A}_{1,k,i+1}^{i,j,n})) = \det_q (\delta_0 (\mathcal{A}_{1,k,i+1}^{i,j,n})) = 0.$$

Therefore,  $\delta_0$  is an algebra homomorphism of  $\mathcal{A}(n)$ . Similarly,  $\delta'_0$  is also an algebra homomorphism. In fact, we also have

$$\begin{aligned}
\delta_0 \delta'_0 (a_{1,1}) &= \delta_0 (a_{1,n}) = a_{1,1}, \\
\delta_0 \delta'_0 (a_{1,n}) &= \delta_0 (a_{1,1}) = a_{1,n}, \\
\delta_0 \delta'_0 (a_{1,i}), &= \delta_0 \left( a_{1,i} \delta'_0 \right) = a_{1,i} + [[a_{2,n}, a_{1,i}]_q, a_{2,n}]_q - f (\mathcal{A}_{1,2,i+1}^{1,i,n}) = a_{1,i}, \\
\delta_0 \delta'_0 (a_{i,l}) &= \delta_0 (a_{i,l}) = a_{i,l},
\end{aligned}$$

where  $i \neq 1$  and  $l \in \llbracket 1, n \rrbracket$ . This means that  $\delta_0 \delta'_0 = \text{id}$ . Similarly,  $\delta'_0 \delta_0 = \text{id}$ .

The proof is finished.  $\square$

**Proposition 3.4.**  $\delta_i, \delta'_i, i \in \llbracket 1, n-1 \rrbracket$  are the automorphisms of algebra  $\mathcal{A}(n)$  and  $\delta_i \delta'_i = \delta'_i \delta_i = \text{id}$ .

*Proof.* To prove that  $\delta_i$  is an algebra homomorphism, it is sufficient to demonstrate that  $\delta_i$  keeps the relations of  $\mathcal{A}(n)$ . It is similar for  $\delta'_i$ .

By definition of  $\delta_i$ , we only need to show that  $\delta_i$  keeps those relations that concern with the  $i+1$ -th row's generators  $a_{i+1,k}$  and  $i$ -column's generators  $a_{j,i}$ .

For the relations (R1). Noting that  $a_{i,i}$  and  $a_{i+1,i+1}$  belong to the center of  $\mathcal{A}(n)$ , we easily see that  $\delta_i (a_{i,i}) = a_{i+1,i+1}$ ,  $\delta_i (a_{i+1,i+1}) = a_{i,i}$ . Consequently,  $\delta_i$  must keep the relations associated with these elements.

If  $[a_{j,i}, a_{k,l}] = 0$  and  $k > j$ , we have for  $l = i$  that

$$[\delta_i (a_{j,i}), \delta_i (a_{k,i})]$$

$$\begin{aligned}
& \xrightarrow{(3.3)} && \left[ -[a_{i,i+1}, a_{j,i}]_q + a_{i+1,i+1} a_{j,i-1}, a_{k,i}^{\delta_i} \right] \\
& \xrightarrow{(3.b)} && \left[ \left[ a_{k,i+1}, [a_{j,i-1}, a_{k,i}]_q \right]_q - a_{k,i-1} [a_{k,i+1}, a_{j,i}]_q \right. \\
& && \left. - a_{j,k-1} ([a_{i,i+1}, a_{k,i}]_q - a_{i+1,i+1} a_{k,i-1}), a_{k,i}^{\delta_i} \right] \\
& \xrightarrow{(3.3)} && \left[ \left[ a_{k,i+1}, [a_{j,i-1}, a_{k,i}]_q \right]_q - a_{k,i-1} [a_{k,i+1}, a_{j,i}]_q \right. \\
& && \left. + a_{j,k-1} (a_{k,i}^{\delta_i} - a_{i,i} a_{k,i+1}), a_{k,i}^{\delta_i} \right] \\
& \xrightarrow{(R1)} && \left[ \left[ a_{k,i+1}, [a_{j,i-1}, a_{k,i}]_q \right]_q - a_{k,i-1} [a_{k,i+1}, a_{j,i}]_q, a_{k,i}^{\delta_i} \right] \\
& = && \left[ \left[ a_{k,i+1}, [a_{j,i-1}, a_{k,i}]_q - a_{k,i-1} a_{j,i} \right]_q, a_{k,i}^{\delta_i} \right] \\
& \xrightarrow{(1.5)} && - \left[ \left[ [a_{j,i-1}, a_{k,i}]_q - a_{k,i-1} a_{j,i}, a_{k,i}^{\delta_i} \right]_q, a_{k,i+1} \right] \\
& && - \left[ \left[ a_{k,i}^{\delta_i}, a_{k,i+1} \right]_q, [a_{j,i-1}, a_{k,i}]_q - a_{k,i-1} a_{j,i} \right] \\
& \xrightarrow{(1.8)} && - a_{k,i}^{\delta_i} \left[ [a_{j,i-1}, a_{k,i}]_q - a_{k,i-1} a_{j,i}, a_{k,i+1} \right] \\
& \xrightarrow{(1.19)} && - a_{k,i}^{\delta_i} \left[ a_{k,i+1}, [a_{j,i-1}, a_{k,i}]_q - a_{k,i-1} a_{j,i} \right] \\
& \xrightarrow{(1.1)} && 0.
\end{aligned}$$

Similarly, we can prove that

$$[\delta_i(a_{i+1,j}), \delta_i(a_{i+1,k})] = 0.$$

If  $k = i + 1$ , we have

$$\begin{aligned}
& [\delta_i(a_{j,i}), \delta_i(a_{i+1,l})] \\
& \xrightarrow{(3.3)} && \left[ -[a_{i,i+1}, a_{j,i}]_q + a_{i,i} a_{j,i+1}, a_{i+1,l}^{\delta_i} \right] \\
& \xrightarrow{(2.a)} && \left[ \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q \right]_q - a_{i+2,l} [a_{i,l}, a_{j,i}]_q - a_{j,l} ([a_{i,i+1}, a_{i+1,l}]_q + a_{i,i} a_{i+2,l}), a_{i+1,l}^{\delta_i} \right] \\
& \xrightarrow{(3.3)} && \left[ \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q \right]_q - a_{i+2,l} [a_{i,l}, a_{j,i}]_q + a_{j,l} (a_{i+1,l}^{\delta_i} - a_{i+1,i+1} a_{i,l}), a_{i+1,l}^{\delta_i} \right] \\
& \xrightarrow{(R1)} && \left[ \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q \right]_q - a_{i+2,l} [a_{i,l}, a_{j,i}]_q, a_{i+1,l}^{\delta_i} \right] \\
& = && \left[ \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i} \right]_q, a_{i+1,l}^{\delta_i} \right] \\
& \xrightarrow{(1.5)} && - \left[ \left[ [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i}, a_{i+1,l}^{\delta_i} \right]_q, a_{i,l} \right] - \left[ \left[ a_{i+1,l}^{\delta_i}, a_{i,l} \right]_q, [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i} \right] \\
& \xrightarrow{(1.8)} && - a_{i+1,l}^{\delta_i} \left[ [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i}, a_{i,l} \right] - a_{i+1,l}^{\delta_i} \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i} \right] \\
& \xrightarrow{(1.21)} && - a_{i+1,l}^{\delta_i} \left[ [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i}, a_{i,l} \right] - a_{i+1,l}^{\delta_i} \left[ a_{i,l}, [a_{j,i+1}, a_{i+1,l}]_q - a_{i+2,l} a_{j,i} \right] \\
& \xrightarrow{(1.1)} && 0.
\end{aligned}$$

If  $\delta_i(a_{k,l}) = a_{k,l}$ , we have

$$[\delta_i(a_{j,i}), \delta_i(a_{k,l})] = - \left[ [a_{i,i+1}, a_{j,i}]_q, a_{k,l} \right] + [a_{i+1,i+1} a_{j,i-1}, a_{k,l}] + [a_{i,i} a_{j,i+1}, a_{k,l}] = 0.$$

Similarly, we can prove that

$$[\delta_i(a_{i+1,j}), \delta_i(a_{k,l})] = 0.$$

For the relations (R2). If we have chosen the sub-matrix  $\mathcal{A}_{j,k,i+1}^{k-1,i,l}$  as (1.10), the proof deduce to the simpler case when  $k = i$  or  $l = i + 1$  and straightforward. In more tedious calculations, we concentrate on exploring the general cases where  $k \neq i$  and  $l \neq i + 1$ .

$$\begin{aligned} & \left[ \delta_i(a_{k,l}), [\delta_i(a_{j,i}), \delta_i(a_{k,l})]_q \right]_q \\ \stackrel{(3.3)}{=} & - \left[ a_{k,l}, \left[ [a_{i,i+1}, a_{j,i}]_q, a_{k,l} \right]_q \right]_q + a_{i+1,i+1} \left[ a_{k,l}, [a_{j,i-1}, a_{k,l}]_q \right]_q \\ & + a_{i,i} \left[ a_{k,l}, [a_{j,i+1}, a_{k,l}]_q \right]_q \\ \stackrel{(1.16)}{=} & - \left[ a_{k,l}, \left[ [a_{i,i+1}, a_{j,i}]_q, a_{k,l} \right]_q \right]_q + a_{i+1,i+1} f(\mathcal{A}_{j,k,i}^{k-1,i-1,l}) + a_{i,i} f(\mathcal{A}_{j,k,i+2}^{k-1,i+1,l}) \\ \stackrel{(1.7)}{=} & - \left[ a_{i,i+1}, \left[ a_{k,l}, [a_{j,i}, a_{k,l}]_q \right]_q \right]_q + a_{i+1,i+1} f(\mathcal{A}_{j,k,i}^{k-1,i-1,l}) + a_{i,i} f(\mathcal{A}_{j,k,i+2}^{k-1,i+1,l}) \\ \stackrel{(1.16)}{=} & - \left[ a_{i,i+1}, f(\mathcal{A}_{j,k,i+1}^{k-1,i,l}) \right]_q + a_{i+1,i+1} f(\mathcal{A}_{j,k,i}^{k-1,i-1,l}) + a_{i,i} f(\mathcal{A}_{j,k,i+2}^{k-1,i+1,l}) \\ \stackrel{(1.11)}{=} & - [a_{i,i+1}, a_{j,i}]_q + a_{j,i-1} a_{i+1,i+1} + a_{i,i} a_{j,i+1} \\ & + a_{j,k-1} a_{k,l} \left( - [a_{i,i+1}, a_{i+1,l}]_q + a_{i,i} a_{i+2,l} + a_{i+1,i+1} a_{i,l} \right) \\ & + a_{j,l} a_{k,l} \left( - [a_{i,i+1}, a_{k,i}]_q + a_{k,i-1} a_{i+1,i+1} + a_{i,i} a_{k,i+1} \right) \\ & - a_{j,k-1} \left( - [a_{i,i+1}, a_{k,i}]_q + a_{k,i-1} a_{i+1,i+1} + a_{i,i} a_{k,i+1} \right) \\ & - a_{j,l} \left( - [a_{i,i+1}, a_{i+1,l}]_q + a_{i,i} a_{i+2,l} + a_{i+1,i+1} a_{i,l} \right) \\ \stackrel{(3.3)}{=} & a_{j,i} \delta_i + a_{j,k-1} a_{k,l} a_{i+1,l} \delta_i + a_{j,l} a_{k,l} a_{k,i} \delta_i - a_{j,k-1} a_{k,i} \delta_i - a_{j,l} a_{i+1,l} \delta_i \\ = & \delta_i(a_{j,i}) + \delta_i(a_{j,k-1}) \delta_i(a_{i+1,l}) \delta_i(a_{k,l}) + \delta_i(a_{k,i}) \delta_i(a_{j,l}) \delta_i(a_{k,l}) \\ & - \delta_i(a_{j,k-1}) \delta_i(a_{k,i}) - \delta_i(a_{j,l}) \delta_i(a_{i+1,l}) \end{aligned}$$

and

$$\begin{aligned} & \left[ \delta_i(a_{j,i}), [\delta_i(a_{k,l}), \delta_i(a_{j,i})]_q \right]_q \\ \stackrel{(3.3)}{=} & - \left[ a_{j,i} \delta_i, \left[ a_{i,i+1}, [a_{k,l}, a_{j,i}]_q \right]_q \right]_q + a_{i+1,i+1} \left[ a_{j,i} \delta_i, [a_{k,l}, a_{j,i-1}]_q \right]_q \\ & + a_{i,i} \left[ a_{j,i} \delta_i, [a_{k,l}, a_{j,i+1}]_q \right]_q \\ \stackrel{(1.3)}{=} & - \left[ \left[ a_{j,i} \delta_i, a_{i,i+1} \right]_q, [a_{k,l}, a_{j,i}]_q \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{j,i} \delta_i \right] \right]_q \\ & + a_{i+1,i+1} \left[ a_{j,i} \delta_i, [a_{k,l}, a_{j,i-1}]_q \right]_q + a_{i,i} \left[ a_{j,i} \delta_i, [a_{k,l}, a_{j,i+1}]_q \right]_q \\ \stackrel{(3.3)}{=} & \left[ a_{j,i} - a_{j,i-1} a_{i,i} - a_{i+1,i+1} a_{j,i+1}, [a_{k,l}, a_{j,i}]_q \right]_q + a_{i+1,i+1} \left[ a_{j,i} \delta_i, [a_{k,l}, a_{j,i-1}]_q \right]_q \end{aligned}$$

$$\begin{aligned}
& + a_{i,i} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{j,i}^{\delta_i} \right] \right] \\
& = \left[ a_{j,i}, [a_{k,l}, a_{j,i}]_q \right]_q - a_{i,i} \left[ a_{j,i-1}, [a_{k,l}, a_{j,i}]_q \right]_q - a_{i+1,i+1} \left[ a_{j,i+1}, [a_{k,l}, a_{j,i}]_q \right]_q \\
& \quad + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q \right]_q + a_{i,i} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q \\
& \quad + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{j,i}^{\delta_i} \right] \right] \\
& \stackrel{(1.12)}{=} \stackrel{(1.16)}{=} a_{k,l} - a_{j,k-1} a_{j,l} - a_{i,i} \left[ a_{j,i-1}, [a_{k,l}, a_{j,i}]_q \right]_q + a_{i,i} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q \\
& \quad - a_{i+1,i+1} \left[ a_{j,i+1}, [a_{k,l}, a_{j,i}]_q \right]_q + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q \right]_q - a_{k,i} a_{i+1,l} \\
& \quad + a_{j,i} a_{k,i} a_{j,l} + a_{j,i} a_{j,k-1} a_{i+1,l} + \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{j,i}^{\delta_i} \right] \right] \\
& \stackrel{(3.12)(3.13)}{=} \stackrel{(3.14)}{=} a_{k,l} - a_{j,k-1} a_{j,l} - a_{i,i} \left[ a_{j,i-1}, [a_{k,l}, a_{j,i}]_q \right]_q + a_{i,i} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q \\
& \quad - a_{i+1,i+1} \left[ a_{j,i+1}, [a_{k,l}, a_{j,i}]_q \right]_q + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q \right]_q \\
& \quad - \left( a_{k,i}^{\delta_i} a_{i+1,l}^{\delta_i} + a_{k,i-1} a_{i,i} a_{i+1,l} - a_{k,i}^{\delta_i} a_{i,i} a_{i+2,l} + a_{i+1,i+1} \left[ a_{k,i+1}, a_{i+1,l} \right]_q \right. \\
& \quad \left. - a_{i+1,i+1} \left[ a_{k,i}^{\delta_i}, a_{i,l} \right]_q \right) + \left( a_{j,i}^{\delta_i} a_{k,i}^{\delta_i} + a_{i+1,i+1} a_{j,i+1} a_{k,i} + a_{i,i} \left[ a_{j,i-1}, a_{k,i} \right]_q \right. \\
& \quad \left. - a_{j,i}^{\delta_i} a_{k,i-1} a_{i+1,i+1} - a_{i,i} \left[ a_{j,i}^{\delta_i}, a_{k,i+1} \right]_q \right) a_{j,l} + a_{j,k-1} \left( a_{j,i}^{\delta_i} a_{i+1,l}^{\delta_i} \right. \\
& \quad \left. + a_{j,i-1} a_{i,i} a_{i+1,l} + a_{j,i}^{\delta_i} a_{i,i} a_{i+2,l} + a_{i+1,i+1} \left[ a_{j,i+1}, a_{i+1,l} \right]_q - a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, a_{i,l} \right]_q \right) \\
& \quad + \frac{1}{(q-q^{-1})^2} \left( \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q - a_{j,l} a_{k,i} - a_{j,k-1} a_{i+1,l} + \left[ a_{i+1,l}, a_{k,i}^{\delta_i} \right] \right] \right] \right) \\
& \stackrel{(1.2)}{=} a_{k,l} - a_{j,k-1} a_{j,l} - a_{k,i}^{\delta_i} a_{i+1,l}^{\delta_i} + a_{j,i}^{\delta_i} a_{k,i}^{\delta_i} a_{j,l} + a_{j,i}^{\delta_i} a_{j,k-1} a_{i+1,l}^{\delta_i} \\
& \quad + a_{i,i} \left( \left[ a_{j,i}^{\delta_i}, \left[ [a_{k,l}, a_{j,i+1}]_q - a_{j,l} a_{k,i+1} - a_{j,k-1} a_{i+2,l} \right]_q + a_{k,i}^{\delta_i} a_{i+2,l} \right] \right. \\
& \quad \left. - a_{i,i} \left( \left[ a_{j,i-1}, \left[ [a_{k,l}, a_{j,i}]_q - a_{j,k-1} a_{i+1,l} - a_{j,l} a_{k,i} \right]_q + a_{k,i-1} a_{i+1,l} \right] \right) \right. \\
& \quad \left. + a_{i+1,i+1} \left( \left[ a_{j,i}^{\delta_i}, \left[ [a_{k,l}, a_{j,i-1}]_q - a_{k,i-1} a_{j,l} - a_{j,k-1} a_{i,l} \right]_q + \left[ a_{k,i}^{\delta_i}, a_{i,l} \right]_q \right] \right) \right. \\
& \quad \left. - a_{i+1,i+1} \left( \left[ a_{j,i+1}, \left[ [a_{k,l}, a_{j,i}]_q - a_{j,k-1} a_{i+1,l} - a_{k,i} a_{j,l} \right]_q + \left[ a_{k,i+1}, a_{i+1,l} \right]_q \right] \right) \right) \\
& \stackrel{(3.19)}{=} \stackrel{(3.20)}{=} a_{k,l} - a_{j,k-1} a_{j,l} - a_{k,i}^{\delta_i} a_{i+1,l}^{\delta_i} + a_{j,i}^{\delta_i} a_{k,i}^{\delta_i} a_{j,l} + a_{j,i}^{\delta_i} a_{j,k-1} a_{i+1,l}^{\delta_i} \\
& \quad = \delta_i(a_{k,l}) - \delta_i(a_{j,k-1}) \delta_i(a_{j,l}) - \delta_i(a_{k,i}) \delta_i(a_{i+1,l}) + \delta_i(a_{j,i}) \delta_i(a_{k,i}) \delta_i(a_{j,l}) \\
& \quad + \delta_i(a_{j,i}) \delta_i(a_{j,k-1}) \delta_i(a_{i+1,l}),
\end{aligned}$$

we have

$$\left[ \delta_i(a_{k,l}), [\delta_i(a_{j,i}), \delta_i(a_{k,l})]_q \right]_q = f \left( \delta_i \left( \mathcal{A}_{j,k,i+1}^{k-1,i,l} \right) \right),$$

$$\left[ \delta_i(a_{j,i}), [\delta_i(a_{k,l}), \delta_i(a_{j,i})]_q \right]_q = g\left(\delta_i\left(\mathcal{A}_{j,k,i+1}^{k-1,i,l}\right)\right).$$

Similarly, we can get that

$$\begin{aligned} \left[ \delta_i(a_{i+1,l}), [\delta_i(a_{j,k}), \delta_i(a_{i+1,l})]_q \right]_q &= f\left(\delta_i\left(\mathcal{A}_{j,i+1,k+1}^{i,k,l}\right)\right), \\ \left[ \delta_i(a_{j,k}), [\delta_i(a_{i+1,l}), \delta_i(a_{j,k})]_q \right]_q &= g\left(\delta_i\left(\mathcal{A}_{j,i+1,k+1}^{i,k,l}\right)\right); \\ \left[ \delta_i(a_{k,i}), [\delta_i(a_{j,l}), \delta_i(a_{k,i})]_q \right]_q &= f\left(\delta_i\left(\mathcal{A}_{j,k,l+1}^{k-1,l,i}\right)\right), \\ \left[ \delta_i(a_{j,l}), [\delta_i(a_{k,i}), \delta_i(a_{j,l})]_q \right]_q &= g\left(\delta_i\left(\mathcal{A}_{j,k,l+1}^{k-1,l,i}\right)\right); \\ \left[ \delta_i(a_{k,l}), [\delta_i(a_{i+1,j}), \delta_i(a_{k,l})]_q \right]_q &= f\left(\delta_i\left(\mathcal{A}_{i+1,k,j+1}^{k-1,j,l}\right)\right), \\ \left[ \delta_i(a_{i+1,j}), [\delta_i(a_{k,l}), \delta_i(a_{i+1,j})]_q \right]_q &= g\left(\delta_i\left(\mathcal{A}_{i+1,k,j+1}^{k-1,j,l}\right)\right). \end{aligned}$$

For the relations (R3). Choosing the sub-matrix  $\mathcal{A}_{j,k,i+1}^{i,l,m}$  as (1.13), we explore the cases when  $k \neq i$  and  $l \neq i+1$  and have that

$$\begin{aligned} &\left[ \delta_i(a_{j,l}), \delta_i(a_{k,m}) \right]_q + \left[ [\delta_i(a_{j,i}), \delta_i(a_{k,l})]_q, \delta_i(a_{i+1,m}) \right]_q + \delta_i(a_{k,i}) \delta_i(a_{i+1,l}) \delta_i(a_{j,m}) \\ &- \delta_i(a_{k,i}) [\delta_i(a_{j,l}), \delta_i(a_{i+1,m})]_q - \delta_i(a_{i+1,l}) [\delta_i(a_{j,i}), \delta_i(a_{k,m})]_q - \delta_i(a_{k,l}) \delta_i(a_{j,m}) \\ \xrightarrow{(1.7)} \quad &\left[ a_{j,l}, a_{k,m} \right]_q - \left[ a_{j,i}^{\delta_i}, \left[ a_{i,i+1}, [a_{k,l}, a_{i+1,m}]_q \right]_q \right]_q + \left[ \left[ a_{j,i}^{\delta_i}, a_{k,l} \right]_q, a_{i,i} a_{i+1,m} \right]_q \\ &+ \left[ \left[ a_{j,i}^{\delta_i}, a_{k,l} \right]_q, a_{i+2,i+1} a_{i,m} \right]_q + a_{j,m} a_{k,i}^{\delta_i} a_{i+1,l}^{\delta_i} - a_{k,i}^{\delta_i} \left[ a_{j,l}, a_{i+1,m} \right]_q \\ &- a_{i+1,l}^{\delta_i} \left[ a_{j,i}^{\delta_i}, a_{k,m} \right]_q - a_{k,l} a_{j,m} \\ \xrightarrow{(1.3)} \quad &\left[ a_{j,l}, a_{k,m} \right]_q - a_{k,l} a_{j,m} - \left[ \left[ a_{j,i}^{\delta_i}, a_{i,i+1} \right]_q, \left[ a_{k,l}, a_{i+1,m} \right]_q \right]_q \\ &- \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ \left[ a_{k,l}, a_{i+1,m} \right]_q, a_{j,i}^{\delta_i} \right] \right]_q + a_{i,i} \left[ a_{j,i}^{\delta_i}, \left[ a_{k,l}, a_{i+2,m} \right]_q \right]_q \\ &+ a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, \left[ a_{k,l}, a_{i,m} \right]_q \right]_q - \left[ a_{j,l}, a_{k,i}^{\delta_i} a_{i+1,m}^{\delta_i} \right]_q \\ &- \left[ a_{i+1,l}^{\delta_i} a_{j,i}^{\delta_i}, a_{k,m} \right]_q + a_{j,m} a_{k,i}^{\delta_i} a_{i+1,l}^{\delta_i} \\ \xrightarrow{(1.16)(3.12)} \quad &\left[ a_{j,l}, a_{k,m} \right]_q - a_{k,l} a_{j,m} + \left[ a_{j,i}, \left[ a_{k,l}, a_{i+1,m} \right]_q \right]_q - a_{i,i} \left[ a_{j,i-1}, \left[ a_{k,l}, a_{i+1,m} \right]_q \right]_q \\ &- a_{i+1,i+1} \left[ a_{j,i+1}, \left[ a_{k,l}, a_{i+1,m} \right]_q \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ \left[ a_{k,l}, a_{i+1,m} \right]_q, a_{j,i}^{\delta_i} \right] \right]_q \\ &+ a_{i,i} \left[ a_{j,i}^{\delta_i}, \left[ a_{k,l}, a_{i+2,m} \right]_q \right]_q + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, \left[ a_{k,l}, a_{i,m} \right]_q \right]_q \\ &- \left[ a_{j,l}, a_{k,i} a_{i+1,m} - a_{k,i-1} a_{i,i} a_{i+1,m} + a_{k,i}^{\delta_i} a_{i,i} a_{i+2,m} - a_{i+1,i+1} \left[ a_{k,i+1}, a_{i+1,m} \right]_q \right]_q \\ &+ a_{i+1,i+1} \left[ a_{k,i}^{\delta_i}, a_{i,m} \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{i+1,m}, a_{k,i}^{\delta_i} \right] \right]_q \\ &- \left[ a_{j,i} a_{i+1,l} - a_{j,i-1} a_{i,i} a_{i+1,l} + a_{j,i}^{\delta_i} a_{i,i} a_{i+2,l} - a_{i+1,i+1} \left[ a_{j,i+1}, a_{i+1,l} \right]_q \right]_q \end{aligned}$$

$$\begin{aligned}
& + a_{i+1,i+1} \left[ a_{j,i}^{\delta_i}, a_{i,l} \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{i+1,l}, a_{j,i}^{\delta_i} \right] \right], a_{k,m} \Big]_q \\
& + a_{j,m} \left( a_{k,i} a_{i+1,l} - a_{k,i-1} a_{i,i} a_{i+1,l} + a_{k,i}^{\delta_i} a_{i,i} a_{i+2,l} - a_{i+1,i+1} [a_{k,i+1}, a_{i+1,l}]_q \right. \\
& \quad \left. + a_{i+1,i+1} \left[ a_{k,i}^{\delta_i}, a_{i,l} \right]_q + \left[ a_{i,i+1}, \left[ a_{i+1,l}, a_{k,i}^{\delta_i} \right] \right] \right) \\
\stackrel{(1.17)}{=} & \det_q (\mathcal{A}_{j,k,i+1}^{i,l,m}) - a_{i,i} \left( \left[ a_{j,i-1}, [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l} a_{k,m} \right]_q - a_{k,i-1} ([a_{j,l}, a_{i+1,m}]_q - \right. \\
& \quad \left. a_{i+1,l} a_{j,m}) - \left[ a_{j,i}^{\delta_i} [a_{k,l}, a_{i+2,m}]_q + a_{i+2,l} a_{k,m} \right]_q + a_{k,i}^{\delta_i} ([a_{j,l}, a_{i+2,m}]_q + a_{i+2,l} a_{j,m}) \right) \\
& - a_{i+1,i+1} \left( \left[ a_{j,i+1}, [a_{k,l}, a_{i+1,m}]_q - [a_{i+1,l}, a_{k,m}]_q \right]_q - \left[ a_{j,i}^{\delta_i} [a_{k,l}, a_{i,m}]_q - [a_{i,l}, a_{k,m}]_q \right]_q \right. \\
& \quad \left. - \left[ a_{k,i+1}, [a_{j,l}, a_{i+1,m}]_q - a_{j,m} a_{i+1,l} \right]_q + \left[ a_{k,i}^{\delta_i} a_{j,m} a_{i,l} - [a_{j,l}, a_{i,m}]_q \right]_q \right) \\
& + a_{i,i} a_{l+1,m} \left( - \left[ a_{j,i}^{\delta_i} a_{k,i+1} \right]_q + a_{k,i}^{\delta_i} a_{j,i+1} + [a_{j,i-1}, a_{k,i}]_q - a_{j,i} a_{k,i-1} \right) \\
& - a_{i+1,i+1} a_{l+1,m} \left( a_{k,i} a_{j,i+1} - a_{k,i-1} a_{j,i}^{\delta_i} - [a_{k,i+1}, a_{j,i}]_q + \left[ a_{k,i}^{\delta_i} a_{j,i-1} \right]_q \right) \\
\stackrel{(1.17)}{=} & a_{i,i} \left( \left[ a_{j,i}^{\delta_i} [a_{k,l}, a_{i+2,m}]_q - a_{k,i+1} a_{l+1,m} - a_{i+2,l} a_{k,m} \right]_q \right. \\
& \quad \left. - a_{k,i}^{\delta_i} ([a_{j,l}, a_{i+2,m}]_q - a_{j,i+1} a_{l+1,m} - a_{i+2,l} a_{j,m}) \right) \\
& - a_{i,i} \left( \left[ a_{j,i-1}, [a_{k,l}, a_{i+1,m}]_q - a_{k,i} a_{l+1,m} - a_{i+1,l} a_{k,m} \right]_q \right. \\
& \quad \left. - a_{k,i-1} ([a_{j,l}, a_{i+1,m}]_q - a_{j,i} a_{l+1,m} - a_{i+1,l} a_{j,m}) \right) \\
& - a_{i+1,i+1} \left( \left[ a_{j,i+1}, [a_{k,l}, a_{i+1,m}]_q - a_{k,i} a_{l+1,m} - a_{i+1,l} a_{k,m} \right]_q \right. \\
& \quad \left. - \left[ a_{k,i+1}, [a_{j,l}, a_{i+1,m}]_q - a_{j,i} a_{l+1,m} - a_{i+1,l} a_{j,m} \right]_q \right) \\
& + a_{i+1,i+1} \left( \left[ a_{k,i}^{\delta_i} [a_{j,l}, a_{i,m}]_q - a_{j,i-1} a_{l+1,m} - a_{i,l} a_{j,m} \right]_q \right. \\
& \quad \left. - \left[ a_{j,i}^{\delta_i} [a_{k,l}, a_{i,m}]_q - a_{k,i-1} a_{l+1,m} - a_{i,l} a_{k,m} \right]_q \right) \\
\stackrel{(2.b)}{=} & a_{i,i} \left( a_{k,i-1} ([a_{j,l}, a_{i+1,m}]_q - a_{k,i-1} a_{i+1,l} a_{j,m}) + [a_{j,i-1}, [a_{k,l}, a_{i+1,m}]_q - a_{i+1,l} a_{k,m}]_q \right. \\
& \quad \left. - a_{k,i}^{\delta_i} ([a_{j,l}, a_{i+2,m}]_q - a_{i+2,l} a_{j,m}) - \left[ a_{j,i}^{\delta_i}, a_{i+2,l} a_{k,m} - [a_{k,l}, a_{i+2,m}]_q \right]_q \right) \\
& + a_{i+1,i+1} \left( \left[ a_{j,i+1}, a_{i+1,l} a_{k,m} - [a_{k,l}, a_{i+1,m}]_q \right]_q + \left[ a_{k,i+1}, [a_{j,l}, a_{i+1,m}]_q - a_{i+1,l} a_{j,m} \right]_q \right. \\
& \quad \left. - \left[ a_{j,i}^{\delta_i}, a_{i,l} a_{k,m} - [a_{k,l}, a_{i,m}]_q \right]_q - \left[ a_{k,i}^{\delta_i}, [a_{j,l}, a_{i,m}]_q - a_{i,l} a_{j,m} \right]_q \right) \\
\stackrel{(3.21)}{=} & 0
\end{aligned}$$

and

$$[\delta_i(a_{k,m}), \delta_i(a_{j,l})]_q + \left[ [\delta_i(a_{i+1,m}), \delta_i(a_{k,l})]_q, \delta_i(a_{j,i}) \right]_q - \delta_i(a_{i+1,l}) [\delta_i(a_{k,m}), \delta_i(a_{j,i})]_q$$

$$\begin{aligned}
& -\delta_i(a_{k,i}) [\delta_i(a_{i+1,m}), \delta_i(a_{j,l})]_q + \delta_i(a_{i+1,l}) \delta_i(a_{k,i}) \delta_i(a_{j,m}) - \delta_i(a_{k,l}) \delta_i(a_{j,m}) \\
\stackrel{(1.7)}{=} & [a_{k,m}, a_{j,l}]_q - \left[ \left[ \left[ a_{i+1,m}^{\delta_i}, a_{i,i+1} \right]_q, a_{k,l} \right]_q, a_{j,i} \right]_q + \left[ \left[ a_{i+1,m}^{\delta_i}, a_{k,l} \right]_q, a_{i+1,i+1} a_{j,i-1} \right]_q \\
& + \left[ \left[ a_{i+1,m}^{\delta_i}, a_{k,l} \right]_q, a_{i,i} a_{j,i+1} \right]_q - a_{i+1,l}^{\delta_i} [a_{k,m}, a_{j,i}^{\delta_i}]_q \\
& - a_{k,i}^{\delta_i} [a_{i+1,m}^{\delta_i}, a_{j,l}]_q + a_{i+1,l}^{\delta_i} a_{k,i}^{\delta_i} a_{j,m} - a_{k,l} a_{j,m} \\
\stackrel{(1.3)}{=} & [a_{k,m}, a_{j,l}]_q - a_{k,l} a_{j,m} - \left[ \left[ a_{i+1,m}^{\delta_i}, a_{i,i+1} \right]_q, [a_{k,l}, a_{j,i}]_q \right]_q \\
& - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{i+1,m}^{\delta_i} \right] \right] + a_{i+1,i+1} \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q \right]_q \\
& + a_{i,i} \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q - \left[ a_{k,m}, a_{i+1,l}^{\delta_i} a_{j,i}^{\delta_i} \right]_q \\
& - \left[ a_{k,i}^{\delta_i} a_{i+1,m}^{\delta_i}, a_{j,l} \right]_q + a_{i+1,l}^{\delta_i} a_{k,i}^{\delta_i} a_{j,m} \\
\stackrel{(1.16)(3.12)}{=} & [a_{k,m}, a_{j,l}]_q - a_{k,l} a_{j,m} + \left[ a_{i+1,m}, [a_{k,l}, a_{j,i}]_q \right]_q - a_{i+1,i+1} \left[ a_{i+2,m}, [a_{k,l}, a_{j,i}]_q \right]_q \\
& - a_{i,i} \left[ a_{i,m}, [a_{k,l}, a_{j,i}]_q \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ [a_{k,l}, a_{j,i}]_q, a_{i+1,m}^{\delta_i} \right] \right] \\
& + a_{i+1,i+1} \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q \right]_q + a_{i,i} \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q \right]_q \\
& - \left[ a_{k,m}, a_{i+1,l} a_{j,i} - a_{i+2,l} a_{i+1,i+1} a_{j,i} + a_{i+1,l}^{\delta_i} a_{i+1,i+1} a_{j,i-1} - a_{i,i} [a_{i,l}, a_{j,i}]_q \right. \\
& \quad \left. + a_{i,i} \left[ a_{i+1,l}^{\delta_i}, a_{j,i+1} \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{j,i}, a_{i+1,l}^{\delta_i} \right] \right] \right]_q \\
& - \left[ a_{i+1,m} a_{k,i} - a_{i+2,m} a_{i+1,i+1} a_{k,i} + a_{i+1,m}^{\delta_i} a_{i+1,i+1} a_{k,i-1} - a_{i,i} [a_{i,m}, a_{k,i}]_q \right. \\
& \quad \left. + a_{i,i} \left[ a_{i+1,m}^{\delta_i}, a_{k,i+1} \right]_q - \frac{1}{(q-q^{-1})^2} \left[ a_{i,i+1}, \left[ a_{k,i}, a_{i+1,m}^{\delta_i} \right] \right], a_{j,l} \right]_q \\
& + \left( a_{i+1,l} a_{k,i} - a_{i+2,l} a_{i+1,i+1} a_{k,i} + a_{i+1,l}^{\delta_i} a_{i+1,i+1} a_{k,i-1} - a_{i,i} [a_{i,l}, a_{k,i}]_q \right. \\
& \quad \left. + a_{i,i} \left[ a_{i+1,l}^{\delta_i}, a_{k,i+1} \right]_q + \left[ a_{i,i+1}, \left[ a_{k,i}, a_{i+1,l}^{\delta_i} \right] \right] \right) a_{j,m} \\
\stackrel{(1.17)}{=} & \det^q \left( \mathcal{A}_{j,k,i+1}^{i,l,m} \right) - a_{i+1,i+1} \left( \left[ a_{i+2,m}, [a_{k,l}, a_{j,i}]_q - a_{k,i} a_{j,l} \right]_q - a_{i+2,l} \left( [a_{k,m}, a_{j,i}]_q - a_{k,i} a_{j,m} \right) \right. \\
& \quad \left. - \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q - a_{k,i-1} a_{j,l} \right]_q + a_{i+1,l}^{\delta_i} \left( [a_{k,m}, a_{j,i-1}]_q - a_{k,i-1} a_{j,m} \right) \right) \\
& - a_{i,i} \left( \left[ a_{i,m}, [a_{k,l}, a_{j,i}]_q - [a_{k,i}, a_{j,l}]_q \right]_q - \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q - [a_{k,i+1}, a_{j,l}]_q \right]_q \right. \\
& \quad \left. - \left[ a_{i,l}, [a_{k,m}, a_{j,i}]_q - a_{j,m} a_{k,i} \right]_q + \left[ a_{i+1,l}^{\delta_i}, a_{j,m} a_{k,i+1} - [a_{k,m}, a_{j,i+1}]_q \right]_q \right) \\
& - a_{i+1,i+1} a_{j,k-1} \left( \left[ a_{i+1,m}^{\delta_i}, a_{i,l} \right]_q - a_{i+1,l}^{\delta_i} a_{i,m} - \left[ a_{i+2,m}, a_{i+1,l} \right]_q + a_{i+1,m} a_{i+2,l} \right) \\
& + a_{i,i} a_{j,k-1} \left( -a_{i+1,l} a_{i,m} + a_{i+2,l} a_{i+1,m}^{\delta_i} + \left[ a_{i,l}, a_{i+1,m} \right]_q - \left[ a_{i+1,l}^{\delta_i}, a_{i+2,m} \right]_q \right)
\end{aligned}$$

$$\begin{aligned}
& \stackrel{(1.17)}{=} a_{i+1,i+1} \left( \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i-1}]_q - a_{i,l} a_{j,k-1} - a_{k,i-1} a_{j,l} \right]_q \right. \\
& \quad \left. - a_{i+1,l}^{\delta_i} \left( [a_{k,m}, a_{j,i-1}]_q - a_{i,m} a_{j,k-1} - a_{k,i-1} a_{j,m} \right) \right) \\
& \quad - a_{i+1,i+1} \left( \left[ a_{i+2,m}, [a_{k,l}, a_{j,i}]_q - a_{i+1,l} a_{j,k-1} - a_{k,i} a_{j,l} \right]_q \right. \\
& \quad \left. - a_{i+2,l} \left( [a_{k,m}, a_{j,i}]_q - a_{i+1,m} a_{j,k-1} - a_{k,i} a_{j,m} \right) \right) \\
& \quad - a_{i,i} \left( \left[ a_{i,m}, [a_{k,l}, a_{j,i}]_q - a_{i+1,l} a_{j,k-1} - a_{k,i} a_{j,l} \right]_q \right. \\
& \quad \left. - \left[ a_{i,l}, [a_{k,m}, a_{j,i}]_q - a_{i+1,m} a_{j,k-1} - a_{k,i} a_{j,m} \right]_q \right) \\
& \quad + a_{i,i} \left( \left[ a_{i+1,l}^{\delta_i}, [a_{k,m}, a_{j,i+1}]_q - a_{i+2,m} a_{j,k-1} - a_{k,i+1} a_{j,m} \right]_q \right. \\
& \quad \left. - \left[ a_{i+1,m}^{\delta_i}, [a_{k,l}, a_{j,i+1}]_q - a_{i+2,l} a_{j,k-1} - a_{k,i+1} a_{j,l} \right]_q \right) \\
& \stackrel{(2.a)}{=} \stackrel{(1.17)}{=} a_{i+1,i+1} \left( a_{i+1,l}^{\delta_i} \left( [a_{k,m}, a_{j,i-1}]_q - a_{k,i-1} a_{j,m} \right) + \left[ a_{i+1,m}^{\delta_i}, a_{k,i-1} a_{j,l} - [a_{k,l}, a_{j,i-1}]_q \right]_q \right. \\
& \quad \left. - a_{i+2,l} \left( [a_{k,m}, a_{j,i}]_q - a_{i+2,l} a_{k,i} a_{j,m} \right) - \left[ a_{i+2,m}, [a_{k,l}, a_{j,i}]_q - a_{k,i} a_{j,l} \right]_q \right) \\
& \quad + a_{i,i} \left( \left[ a_{i+1,m}^{\delta_i}, a_{k,i+1} a_{j,l} - [a_{k,l}, a_{j,i+1}]_q \right]_q + \left[ a_{i+1,l}^{\delta_i}, [a_{k,m}, a_{j,i+1}]_q - a_{k,i+1} a_{j,m} \right]_q \right. \\
& \quad \left. - \left[ a_{i,m}, a_{k,i} a_{j,l} - [a_{k,l}, a_{j,i}]_q \right]_q - \left[ a_{i,l}, [a_{k,m}, a_{j,i}]_q - a_{k,i} a_{j,m} \right]_q \right) \\
& \stackrel{(3.23)}{=} \stackrel{(3.24)}{=} 0,
\end{aligned}$$

Hence,

$$\det_q \left( \delta_i \left( \mathcal{A}_{j,k,i+1}^{i,l,m} \right) \right) = \det^q \left( \delta_i \left( \mathcal{A}_{j,k,i+1}^{i,l,m} \right) \right) = 0.$$

Similarly, we can get that

$$\begin{aligned}
\det_q \left( \delta_i \left( \mathcal{A}_{j,k,l+1}^{l,i,m} \right) \right) &= \det^q \left( \delta_i \left( \mathcal{A}_{j,k,l+1}^{l,i,m} \right) \right) = 0; \\
\det_q \left( \delta_i \left( \mathcal{A}_{j,k,l+1}^{l,m,i} \right) \right) &= \det^q \left( \delta_i \left( \mathcal{A}_{j,k,l+1}^{l,m,i} \right) \right) = 0; \\
\det_q \left( \delta_i \left( \mathcal{A}_{i+1,j,k+1}^{k,l,m} \right) \right) &= \det^q \left( \delta_i \left( \mathcal{A}_{i+1,j,k+1}^{k,l,m} \right) \right) = 0; \\
\det_q \left( \delta_i \left( \mathcal{A}_{j,i+1,k+1}^{k,l,m} \right) \right) &= \det^q \left( \delta_i \left( \mathcal{A}_{j,i+1,k+1}^{k,l,m} \right) \right) = 0.
\end{aligned}$$

Hence, the maps  $\delta_i$  are algebra homomorphism of  $\mathcal{A}(n)$ . Similarly,  $\delta'_i$  is algebra homomorphism.

In fact,  $\delta_i$  (resp.  $\delta'_i$ ) are automorphisms of  $\mathcal{A}(n)$ . Indeed,

$$\begin{aligned}
\delta_i \delta'_i (a_{i,i}) &= \delta_i (a_{i+1,i+1}) = a_{i,i}, \\
\delta_i \delta'_i (a_{i+1,i+1}) &= \delta_i (a_{i,i}) = a_{i+1,i+1}, \\
\delta_i \delta'_i (a_{j,i}) &= \delta_i (a_{j,i} \delta'_i)
\end{aligned}$$

$$\begin{aligned}
&= - \left[ a_{j,i} \delta_i, a_{i,i+1} \right]_q + a_{j,i-1} a_{i,i} + a_{i+1,i+1} a_{j,i+1} \\
&= \left[ [a_{i,i+1}, a_{j,i}]_q, a_{i,i+1} \right]_q - f(\mathcal{A}_{j,i,i+1}^{i-1,i,i+1}) + a_{j,i} \\
&= a_{j,i}, \\
\delta_i \delta'_i(a_{i+1,l}) &= \delta_i \left( a_{i+1,l} \delta'_i \right) \\
&= - \left[ a_{i+1,l} \delta_i, a_{i,i+1} \right]_q + a_{i+1,i+1} a_{i+2,l} + a_{i,l} a_{i,l} \\
&= \left[ [a_{i,i+1}, a_{i+1,l}]_q, a_{i,i+1} \right]_q - g(\mathcal{A}_{i,i+1,i+2}^{i,i+1,l}) + a_{i+1,l} \\
&= a_{i+1,l}, \\
\delta_i \delta'_i(a_{j,l}) &= \delta_i(a_{j,l}) = a_{j,l},
\end{aligned}$$

where  $j \neq i+1$  and  $l \neq i$ . Hence  $\delta_i \delta'_i = \text{id}$  and similarly, we have  $\delta'_i \delta_i = \text{id}$ .

This proof is finished.  $\square$

**Remark 3.5.** The automorphisms  $\delta_i$  of  $\mathcal{A}(n)$  coincide with those given in [15]. Indeed, one can easily see that

$$\begin{array}{ccc}
\mathcal{A}(n) & \xrightarrow{\delta_i} & \mathcal{A}(n) \\
\downarrow \phi & & \downarrow \phi \\
\mathfrak{aw}(n) & \xrightarrow{r_i} & \mathfrak{aw}(n).
\end{array}$$

## 4 The braid relations

In this section, we prove that the automorphisms established in Section 3 satisfy the braid group relations.

A group  $B_n$  is called a braid group if  $B_n$  is generated by  $\sigma_1, \dots, \sigma_n$  with the following relations:

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad i \in \llbracket 1, n-1 \rrbracket, \quad (4.1)$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, |i-j| \geq 2, \quad i, j \in \llbracket 1, n \rrbracket. \quad (4.2)$$

Accordingly, we have

**Theorem 4.1.** The algebra automorphisms  $\delta_i$  where  $i \in \llbracket 0, n-1 \rrbracket$  of  $\mathcal{A}(n)$ , satisfy the braid relations

$$\delta_i \delta_{i+1} \delta_i = \delta_{i+1} \delta_i \delta_{i+1}, \quad i \in \llbracket 0, n-2 \rrbracket;$$

$$\delta_i \delta_j = \delta_j \delta_i, |i-j| \geq 2, \quad i, j \in \llbracket 0, n-1 \rrbracket.$$

*Proof.* The proof includes more tedious calculations, but to understand them in a better way, we write them down in detail.

Firstly, we prove that  $\delta_i \delta_{i+1} \delta_i = \delta_{i+1} \delta_i \delta_{i+1}, i \in \llbracket 0, n-2 \rrbracket$ .

(a). When  $i = 0$ .

If  $k \in \llbracket 3, n \rrbracket$  and  $j \in \llbracket 2, n \rrbracket$ , obviously we have

$$\delta_0 \delta_1 \delta_0 (a_{k,j}) = a_{k,j} = \delta_1 \delta_0 \delta_1 (a_{k,j}).$$

If  $k = 1$  or  $k = 2$  or  $j = 3$ , then we have

$$\begin{aligned}
& \delta_0 \delta_1 \delta_0 (a_{1,1}) = a_{1,1} = \delta_1 \delta_0 \delta_1 (a_{1,1}), \\
& \delta_0 \delta_1 \delta_0 (a_{1,n}) = a_{2,2} = \delta_1 \delta_0 \delta_1 (a_{1,n}), \\
& \delta_0 \delta_1 \delta_0 (a_{2,2}) = a_{1,n} = \delta_1 \delta_0 \delta_1 (a_{2,2}), \\
& \delta_0 \delta_1 \delta_0 (a_{1,j}) \\
\stackrel{(3.1)}{=} & \left[ a_{1,2}, [a_{2,n}, a_{1,j}]_q \right]_q + a_{2,2} a_{j+1,n} - a_{1,n} [a_{1,2}, a_{2,j}]_q + a_{1,1} a_{3,j} a_{1,n} \\
& - a_{1,1} \left( a_{j+1,n} a_{1,2} + \left[ a_{1,2}^{\delta_0}, a_{2,j} \right]_q - a_{2,2} a_{1,j}^{\delta_0} \right) \\
\stackrel{(3.b)}{=} & \left[ a_{1,2}, [a_{2,n}, a_{1,j}]_q \right]_q + a_{2,2} a_{j+1,n} - a_{1,n} [a_{1,2}, a_{2,j}]_q \\
& + a_{1,1} a_{3,j} a_{1,n} - a_{1,1} [a_{3,n}, a_{1,j}]_q \\
\stackrel{(1.7)}{=} & \left[ [a_{1,2}, a_{2,n}]_q, a_{1,j} \right]_q - a_{1,1} [a_{3,n}, a_{1,j}]_q + a_{2,2} a_{j+1,n} \\
& - a_{1,n} \left( [a_{1,2}, a_{2,j}]_q - a_{1,1} a_{3,j} \right) \\
\stackrel{(3.3)}{=} & \left[ [a_{1,2}, a_{2,n}]_q - a_{1,1} a_{3,n}, a_{1,j} \right]_q + a_{2,2} a_{j+1,n} + a_{1,n} \left( a_{2,j}^{\delta_1} - a_{2,2} a_{1,j} \right) \\
\stackrel{(3.3)}{=} & - \left[ a_{2,n}^{\delta_1}, a_{1,j} \right]_q + a_{2,2} a_{j+1,n} + a_{2,j}^{\delta_1} a_{1,n} \\
\stackrel{(3.1)}{=} & \delta_1 \delta_0 \delta_1 (a_{1,j}), \\
& \delta_0 \delta_1 \delta_0 (a_{2,j}) \\
\stackrel{(3.1)}{=} & \left[ [a_{2,n}, a_{1,2}]_q, a_{2,j} \right]_q - a_{1,1} [a_{3,n}, a_{2,j}]_q + a_{3,j} a_{1,n} \\
& - a_{2,2} [a_{2,n}, a_{1,j}]_q + a_{1,1} a_{2,2} a_{j+1,n} \\
\stackrel{(1.7)}{=} & \left[ a_{2,n}, [a_{1,2}, a_{2,j}]_q \right]_q - a_{1,1} \left( a_{3,j} a_{2,n} - a_{2,j}^{\delta_1} a_{1,n} + \left[ a_{2,n}^{\delta_1}, a_{1,j} \right]_q \right) \\
& + a_{3,j} a_{1,n} - a_{2,2} [a_{2,n}, a_{1,j}]_q + a_{1,1} a_{2,2} a_{j+1,n} \\
= & \left[ a_{2,n}, [a_{1,2}, a_{2,j}]_q - a_{1,1} a_{3,j} - a_{2,2} a_{1,j} \right]_q + a_{1,n} a_{3,j} \\
& + a_{1,1} \left( a_{2,j}^{\delta_1} a_{1,n} - \left[ a_{2,n}^{\delta_1}, a_{1,j} \right]_q \right) + a_{1,1} a_{2,2} a_{j+1,n} \\
\stackrel{(1.b)}{=} & - \left[ a_{2,n}, a_{2,j}^{\delta_1} \right]_q + a_{1,n} a_{3,j} + a_{1,1} \left( a_{2,j}^{\delta_1} a_{1,n} - \left[ a_{2,n}^{\delta_1}, a_{1,j} \right]_q \right) + a_{1,1} a_{2,2} a_{j+1,n} \\
\stackrel{(3.1)}{=} & \delta_1 \delta_0 \delta_1 (a_{2,j}).
\end{aligned}$$

(b). When  $i \in \llbracket 1, n-2 \rrbracket$ .

If  $k \neq i+1, i+2$  and  $j \neq i, i+1$ , obviously we have

$$\delta_i \delta_{i+1} \delta_i (a_{k,j}) = a_{k,j} = \delta_{i+1} \delta_i \delta_{i+1} (a_{k,j}).$$

If  $k = i+1$  or  $k = i+2$  or  $j = i$  or  $j = i+1$ , we have

$$\begin{aligned} & \delta_i \delta_{i+1} \delta_i (a_{i,i}) = a_{i+2,i+2} = \delta_{i+1} \delta_i \delta_{i+1} (a_{i,i}), \\ & \delta_i \delta_{i+1} \delta_i (a_{i+1,i+1}) = a_{i+1,i+1} = \delta_{i+1} \delta_i \delta_{i+1} (a_{i+1,i+1}), \\ & \delta_i \delta_{i+1} \delta_i (a_{i+2,i+2}) = a_{i,i} = \delta_{i+1} \delta_i \delta_{i+1} (a_{i+2,i+2}), \\ & \delta_i \delta_{i+1} \delta_i (a_{k,i}) \\ \xrightarrow[(3.4)]{(3.3)} & - \left[ a_{i+1,i+2}, a_{k,i} \right]_q^{\delta_i} + a_{i+1,i+1} a_{i+2,i+2} a_{k,i}^{\delta_i} + a_{k,i-1} a_{i+2,i+2} \\ & - a_{i+1,i+1} \left[ a_{i+1,i+2}^{\delta_i}, a_{1,i+1} \right]_q + a_{i,i} a_{i+1,i+1} a_{1,i+2} \\ \xrightarrow[(3.4)]{(3.3)} & \left[ a_{i+1,i+2}, [a_{i,i+1}, a_{k,i}]_q \right]_q - a_{i+1,i+2} a_{k,i-1} a_{i+1,i+1} \\ & - a_{i,i} [a_{i+1,i+2}, a_{k,i+1}]_q + a_{i+1,i+1} a_{i+2,i+2} a_{k,i}^{\delta_i} + a_{k,i-1} a_{i+2,i+2} \\ & - a_{i+1,i+1} \left[ a_{i+1,i+2}^{\delta_i}, a_{k,i+1} \right]_q + a_{i,i} a_{i+1,i+1} a_{k,i+2} \\ \xrightarrow[(3.4)]{(1.7)} & \left[ [a_{i+1,i+2}, a_{i,i+1}]_q - a_{i+1,i+1} a_{i,i+2}, a_{k,i} \right]_q + a_{k,i-1} a_{i+2,i+2} \\ & - a_{i,i} \left( [a_{i+1,i+2}, a_{k,i+1}]_q - a_{i+1,i+1} a_{k,i+2} \right) \\ \xrightarrow[(3.4)]{(3.3)} & - \left[ a_{i,i+1}^{\delta_{i+1}}, a_{k,i} \right]_q + a_{k,i-1} a_{i+2,i+2} + a_{i,i} a_{k,i+1}^{\delta_{i+1}} \\ \xrightarrow[(3.4)]{(3.3)} & \delta_{i+1} \delta_i \delta_{i+1} (a_{k,i}), \\ & \delta_i \delta_{i+1} \delta_i (a_{k,i+1}) \\ \xrightarrow[(3.4)]{(3.3)} & \left[ [a_{i,i+1}, a_{i+1,i+2}]_q, a_{k,i+1} \right]_q - a_{i+1,i+1} [a_{i,i+2}, a_{k,i+1}]_q \\ & - a_{i+2,i+2} [a_{i,i+1}, a_{k,i}]_q + a_{k,i-1} a_{i+1,i+1} a_{i+2,i+2} + a_{i,i} a_{k,i+2} \\ \xrightarrow[(3.4)]{(1.17)} & \left[ a_{i,i+1}, [a_{i+1,i+2}, a_{k,i+1}]_q \right]_q - a_{i+1,i+1} \left( \left[ a_{i,i+1}^{\delta_{i+1}}, a_{k,i} \right]_q - a_{i,i} a_{k,i+1}^{\delta_{i+1}} \right. \\ & \left. + a_{i,i+1} a_{k,i+2} \right) - a_{i+2,i+2} \left( [a_{i,i+1}, a_{k,i}]_q + a_{k,i-1} a_{i+1,i+1} a_{i+2,i+2} \right) + a_{i,i} a_{k,i+2} \\ \xrightarrow[(3.4)]{(1.16)} & \left[ \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,i+2} \right]_q - a_{i,i} a_{i+1,i+1} - a_{i+2,i+2} a_{i,i+2}, a_{k,i+1}^{\delta_{i+1}} \right]_q \\ & - a_{i+1,i+1} \left[ a_{i,i+1}^{\delta_{i+1}}, a_{k,i} \right]_q + a_{k,i-1} a_{i+1,i+1} a_{i+2,i+2} \\ & + a_{i,i} a_{k,i+1}^{\delta_{i+1}} a_{i+1,i+1} + a_{i,i} a_{k,i+2}^{\delta_{i+1}} \\ \equiv & \left[ \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,i+2} \right]_q, a_{k,i+1}^{\delta_{i+1}} \right]_q - a_{i,i} a_{i+1,i+1} a_{k,i+1}^{\delta_{i+1}} \end{aligned}$$

$$\begin{aligned}
& -a_{i+2,i+2} \left[ a_{i,i+2}, a_{k,i+1}^{\delta_{i+1}} \right]_q - a_{i+1,i+1} \left[ a_{i,i+1}^{\delta_{i+1}}, a_{k,i} \right]_q \\
& + a_{k,i-1} a_{i+1,i+1} a_{i+2,i+2} + a_{i,i} a_{k,i+1}^{\delta_{i+1}} a_{i+1,i+1} + a_{i,i} a_{k,i+2}^{\delta_{i+1}} \\
\stackrel{(3.3)}{=} & \delta_{i+1} \left( - \left[ a_{i+1,i+2}^{\delta_i}, a_{k,i+1} \right]_q + a_{k,i} \delta_i a_{i+2,i+2} + a_{i,i} a_{k,i+2} \right) \\
\stackrel{(3.3)}{=} & \stackrel{(3.4)}{\delta_{i+1} \delta_i \delta_{i+1} (a_{k,i+1}),} \\
& \delta_i \delta_{i+1} \delta_i (a_{i+1,j}) \\
\stackrel{(3.3)}{=} & \stackrel{(3.4)}{\left[ \left[ a_{i+1,i+2}^{\delta_i}, a_{i,i+1} \right]_q, a_{i+2,j}^{\delta_{i+1}} \right]_q - a_{i,i} \left[ a_{i,i+2}, a_{i+2,j}^{\delta_{i+1}} \right]_q} \\
& - a_{i+1,i+1} \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q + a_{i,i} a_{i+1,i+1} a_{i+3,j} + a_{i+2,i+2} a_{i,j} \\
= & \left[ \left[ a_{i+1,i+2}^{\delta_i}, a_{i,i+1} \right]_q - a_{i,i} a_{i,i+2}, a_{i+2,j}^{\delta_{i+1}} \right]_q \\
& - a_{i+1,i+1} \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q + a_{i,i} a_{i+1,i+1} a_{i+3,j} + a_{i+2,i+2} a_{i,j} \\
\stackrel{(1.16)}{=} & - \left[ a_{i+1,i+2}, a_{i+2,j}^{\delta_{i+1}} \right]_q + a_{i+1,i+1} a_{i+2,i+2} a_{i+2,j}^{\delta_{i+1}} \\
& - a_{i+1,i+1} \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q + a_{i,i} a_{i+1,i+1} a_{i+3,j} + a_{i+2,i+2} a_{i,j} \\
\stackrel{(3.4)}{=} & \left[ a_{i+1,i+2}, [a_{i,i+1}, a_{i+1,j}]_q \right]_q - a_{i,i} [a_{i+1,i+2}, a_{i+2,j}]_q + a_{i+1,i+1} \left( a_{i+2,i+2} a_{i+2,j}^{\delta_{i+1}} \right. \\
& \left. - a_{i+1,i+2} a_{i,j} - \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q \right) + a_{i,i} a_{i+1,i+1} a_{i+3,j} + a_{i+2,i+2} a_{i,j} \\
\stackrel{(1.17)}{=} & \left[ [a_{i+1,i+2}, a_{i,i+1}]_q, a_{i+1,j} \right]_q - a_{i,i} [a_{i+1,i+2}, a_{i+2,j}]_q - a_{i+1,i+1} [a_{i,i+2}, a_{i+1,j}]_q \\
& + a_{i,i} a_{i+1,i+1} a_{i+3,j} + a_{i+2,i+2} a_{i,j} \\
\stackrel{(3.3)}{=} & \stackrel{(3.4)}{\delta_{i+1} \delta_i \delta_{i+1} (a_{i+1,j}),} \\
& \delta_i \delta_{i+1} \delta_i (a_{i+2,j}) \\
\stackrel{(3.3)}{=} & \stackrel{(3.4)}{- \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q + a_{i,i} a_{i+3,j} + a_{i+2,i+2} a_{i+2,j}^{\delta_{i+1}}} \\
\stackrel{(1.17)}{=} & \left[ [a_{i,i+1}, a_{i+1,i+2}]_q, a_{i+2,j} \right]_q - a_{i+1,i+1} [a_{i,i+2}, a_{i+2,j}]_q + a_{i,i} a_{i+3,j} \\
& - a_{i+2,i+2} [a_{i,i+1}, a_{i+1,j}]_q + a_{i+1,i+1} a_{i+2,i+2} a_{i,j} \\
\stackrel{(1.17)}{=} & \left[ a_{i,i+1}, [a_{i+1,i+2}, a_{i+2,j}]_q \right]_q - a_{i+1,i+1} \left( a_{i,i+1} a_{i+3,j} - a_{i,i} a_{i+2,j}^{\delta_{i+1}} \right. \\
& \left. + \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,j} \right]_q \right) - a_{i+2,i+2} [a_{i,i+1}, a_{i+1,j}]_q + a_{i,i} a_{i+3,j} \\
& + a_{i+1,i+1} a_{i+2,i+2} a_{i,j} \\
\stackrel{(3.4)}{=} & - \left[ a_{i,i+1}, a_{i+2,j}^{\delta_{i+1}} \right]_q + a_{i,i} a_{i+1,i+1} a_{i+2,j}^{\delta_{i+1}} + a_{i,i} a_{i+3,j}
\end{aligned}$$

$$\begin{aligned}
& -a_{i+1,i+1} \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,j} \right]_q + a_{i+1,i+1} a_{i+2,i+2} a_{i,j} \\
\stackrel{(1.16)}{=} & \left[ \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,i+2} \right]_q - a_{i+2,i+2} a_{i,i+2}, a_{i+2,j}^{\delta_{i+1}} \right]_q + a_{i,i} a_{i+3,j} \\
& -a_{i+1,i+1} \left[ a_{i,i+1}^{\delta_{i+1}}, a_{i+1,j} \right]_q + a_{i+1,i+1} a_{i+2,i+2} a_{i,j} \\
\stackrel{(3.3)}{=} & \delta_{i+1} \left( - \left[ a_{i+1,i+2}^{\delta_i}, a_{i+2,j} \right]_q + a_{i,i} a_{i+3,j} + a_{i+2,i+2} a_{i+2,j}^{\delta_{i+1}} \right) \\
\stackrel{(3.3)}{=} & \delta_{i+1} \delta_i \delta_{i+1} (a_{i+2,j}).
\end{aligned}$$

Hence,  $\delta_i$  satisfy the relations (4.1). Now, let us prove that  $\delta_i$  satisfy the relation (4.2).

(a). If one of  $i$  and  $j$  is equal to 0, let us say  $i = 0$ , then  $j \geq 2$ . If  $k \neq 1, j+1$ , and  $l \neq j$ , we have

$$\delta_0 \delta_j (a_{k,l}) = a_{k,l} = \delta_j \delta_0 (a_{k,l}).$$

If  $k = 1$  or  $k = j+1$  or  $l \neq j$ , we have

$$\begin{aligned}
\delta_0 \delta_j (a_{1,1}) &= \delta_0 (a_{1,1}) = a_{1,n} = \delta_j (a_{1,n}) = \delta_j \delta_0 (a_{1,1}), \\
\delta_0 \delta_j (a_{1,n}) &= \delta_0 (a_{1,n}) = a_{1,1} = \delta_j (a_{1,1}) = \delta_j \delta_0 (a_{1,n}), \\
\delta_0 \delta_j (a_{1,m}) &= \delta_0 (a_{1,m}) = a_{1,m}^{\delta_0} = \delta_j \left( a_{1,m}^{\delta_0} \right) = \delta_j \delta_0 (a_{1,m}), \\
\delta_0 \delta_j (a_{j+1,j+1}) &= \delta_0 (a_{j,j}) = a_{j,j} = \delta_j (a_{j+1,j+1}) = \delta_j \delta_0 (a_{j+1,j+1}), \\
\delta_0 \delta_j (a_{j+1,l}) &= \delta_0 \left( a_{j+1,l}^{\delta_j} \right) = a_{j+1,l}^{\delta_j} = \delta_j (a_{j+1,l}) = \delta_j \delta_0 (a_{j+1,l}), \\
\delta_0 \delta_j (a_{j,j}) &= \delta_0 (a_{j+1,j+1}) = a_{j+1,j+1} = \delta_j (a_{j,j}) = \delta_j \delta_0 (a_{j,j}), \\
\delta_0 \delta_j (a_{k,j}) &= \delta_0 \left( a_{k,j}^{\delta_j} \right) = \delta_j (a_{k,j}) = \delta_j \delta_0 (a_{k,j}).
\end{aligned}$$

(b). If  $0 < i < j$ , then if  $k \neq i+1, j+1$ , and  $l \neq i, j$ , we have

$$\delta_i \delta_j (a_{k,l}) = a_{k,l} = \delta_j \delta_i (a_{k,l}).$$

If  $k = i+1$  or  $k = j+1$  or  $l = i$  or  $l = j$ , we have

$$\begin{aligned}
\delta_i \delta_j (a_{i,i}) &= \delta_i (a_{i,i}) = a_{i+1,i+1} = \delta_j (a_{i+1,i+1}) = \delta_j \delta_i (a_{i,i}), \\
\delta_i \delta_j (a_{i+1,i+1}) &= \delta_i (a_{i+1,i+1}) = a_{i,i} = \delta_j (a_{i,i}) = \delta_j \delta_i (a_{i+1,i+1}), \\
\delta_i \delta_j (a_{j,j}) &= \delta_i (a_{j+1,j+1}) = a_{j+1,j+1} = \delta_j (a_{j,j}) = \delta_j \delta_i (a_{j,j}), \\
\delta_i \delta_j (a_{j+1,j+1}) &= \delta_i (a_{j,j}) = a_{j,j} = \delta_j (a_{j+1,j+1}) = \delta_j \delta_i (a_{j+1,j+1}), \\
\delta_i \delta_j (a_{k,i}) &= \delta_i (a_{k,i}) = a_{k,i}^{\delta_i} = \delta_j \left( a_{k,i}^{\delta_i} \right) = \delta_j \delta_i (a_{k,i}), \\
\delta_i \delta_j (a_{k,j}) &= \delta_i \left( a_{k,j}^{\delta_j} \right) = a_{k,j}^{\delta_j} = \delta_j (a_{k,j}) = \delta_j \delta_i (a_{k,j}), \\
\delta_i \delta_j (a_{i+1,l}) &= \delta_i (a_{i+1,l}) = a_{i+1,l}^{\delta_i} = \delta_j \left( a_{i+1,l}^{\delta_i} \right) = \delta_j \delta_i (a_{i+1,l}), \\
\delta_i \delta_j (a_{j+1,l}) &= \delta_i \left( a_{j+1,l}^{\delta_j} \right) = a_{j+1,l}^{\delta_j} = \delta_j (a_{j+1,l}) = \delta_j \delta_i (a_{j+1,l}).
\end{aligned}$$

Up to now,  $\delta_i$  also satisfy the relation (4.2).

In summary, we complete the proof.  $\square$

## Conclusion

In the paper, we offer an equivalent perspective on defining the higher-rank Askey-Wilson algebras  $\mathcal{A}(n)$  provided by the authors in [15]. We also write down a series of automorphisms in our settings, which coincide with those in [15]. The detailed proofs are done here that satisfy the braid group relations in  $\mathcal{A}(n)$ .

In the future work, we will explore to construct the PBW basis of  $\mathcal{A}(n)$  and to investigate the theory of their representations. It is interesting to find a more intuitive geometric interpretation of  $\mathcal{A}(n)$ .

## References

- [1] A. S. Zhedanov, "Hidden symmetry" of Askey-Wilson polynomial, *Theor. Math. Phys.*, 89(1991): 1146-1157.
- [2] E. Koelink, J.V. Stokman, The Askey-Wilson function transform, *Internat. Math. Res. Notices*, 22(2001): 1203-1227.
- [3] N. Crampé, J. Gaboriaud, L. Vinet, M. Zaimi, Revisiting the Askey-Wilson algebra with the universal  $R$ -matrix of  $U_q(\mathfrak{sl}_2)$ , *J. Phys. A*, 53(2020), 05LT01: 10 pp.
- [4] P. Terwilliger, R. Vidunas, Leonard pairs and the Askey-Wilson relations, *J. Algebra Appl.*, 3(2004): 411-426.
- [5] A. N. Lavrenov, On Askey-Wilson algebra, in Quantum Groups and intergrable system, II(Prague 1997), *Czechoslovak J. Phys.*, 47(1997): 1213-1219.
- [6] N. Crampé, L. Frappat, J. Gaboriaud, L. Poulain d'Andecy, E. Ragoucy, L. Vinet, The Askey-Wilson algebra and its avatars, *J. Phys. A*, 54(2021), 063001: 32 pp.
- [7] N. Crampé, L. Vinet L., M. Zaimi, Temperley-Lieb, Birman-Murakami-Wenzl and Askey-Wilson algebras and other centralizers of  $U_q(\mathfrak{sl}_2)$ , *Ann. Henri Poincaré*, 22(2021): 3499-3528.
- [8] W. Groenevelt, C. Wagenaar, An Askey-Wilson algebra of rank 2, *SIGMA*, 19(2023), 008: 35 pp.
- [9] P. Baseilhac, K. Koizumi, A deformed analogue of Onsager's symmetry in the  $XXZ$  open spin chain, *J. Stat. Mech. Theory Exp.*, 10(2005), P10005: 15 pp.
- [10] H. De Bie, W. van de Vijver, A discrete realization of the higher rank Racah algebra, *Constructive Approximation*, 52(2020): 1-29.
- [11] H. De Bie, H. De Clercq, W. van de Vijver, The higher rank  $q$ -deformed Bannai-Ito and Askey-Wilson algebra, *Comm. Math. Phys.*, 374(2020): 277-316.

- [12] S. Post, A. Walter, A higher rank extension of the Askey-Wilson algebra, arXiv:1705.01860.
- [13] J. Cooke, A. Lacabanne, Higher rank Askey-Wilson algebras as Skein algebras, arxiv:2205.04414.
- [14] H. De Clercq, Higher rank relations for the Askey-Wilson and  $q$ -Bannai-Ito algebra, SIGMA, 15(2019), 099: 32 pp.
- [15] N. Crampé, L. Frappat, L. Poulain d'Andecy, E. Ragoucy, The higher-rank Askey-Wilson algebra and its braid group automorphisms, SIGMA, 19(2023), 077: 36 pp.