

# A NEW DIFFERENTIAL SUBORDINATION TECHNIQUE FOR A SUBCLASS OF STARLIKE FUNCTIONS

S. SIVAPRASAD KUMAR AND POOJA YADAV

ABSTRACT. In the present investigation, we employ a new technique to find several first and second order differential subordination implications involving the following starlike class associated with a bean shaped domain:

$$\mathcal{S}_{\mathfrak{B}}^* := \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} \prec \sqrt{1 + \tanh z} =: \mathfrak{B}(z) \right\}.$$

Also, we give several applications stemming from our derived results.

## 1. INTRODUCTION

Let  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  and  $\mathcal{H}$  be the class of functions  $f$  analytic in  $\mathbb{D}$  with  $f(z) = 1 + a_1z + a_2z^2 + \dots$ . Also, let  $\mathcal{A}$  be the class of analytic functions  $f$  normalized by  $f(0) = f'(0) - 1 = 0$  in  $\mathbb{D}$  and  $\mathcal{S}$  be the subclass of  $\mathcal{A}$  consisting of all univalent functions. Further  $\mathcal{S}^*$  is a subclass of  $\mathcal{S}$ , which consists of all functions  $f$  such that  $f(\mathbb{D})$  is starshaped with respect to origin and these functions are known as starlike functions with respect to origin. Analytically,  $f \in \mathcal{S}^*$  if we have  $\operatorname{Re} zf'(z)/f(z) > 0$ . Furthermore, to compare the inclusion properties of two analytic functions  $f$  and  $g$ , having same origin, i.e.  $f(0) = g(0)$ , we say that  $f$  is subordinate to  $g$ , written as  $f \prec g$ , if and only if  $f(z) = g(w(z))$  where  $w(z)$  is a Schwarz function. In terms of subordination, Ma and Minda [9] unified all the subclasses of  $\mathcal{S}^*$  by defining the following class:

$$\mathcal{S}^*(\phi) = \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} \prec \phi(z) \right\}, \quad (1.1)$$

where  $\phi$  is a univalent and analytic function with positive real part, starlike with respect to  $\phi(0) = 1$ ,  $\phi'(0) > 0$  and  $\phi(\mathbb{D})$  is symmetric with respect to the real axis. Several well-known classes can be obtained by specializing  $\phi$ , such as the Janowski strongly starlike class of order  $\alpha$   $\mathcal{SS}^*(A, B, \alpha) := \mathcal{S}^*((1 + Az)/(1 + Bz))^\alpha$  where  $\alpha \in (0, 1]$  and  $A, B \in \mathbb{C}$  with  $|B| \leq 1$  and  $A \neq B$ , studied by Kumar and Yadav in [7]. Here  $\mathcal{SS}^*(1, -1, \alpha) =: \mathcal{SS}^*(\alpha)$ ; the class of strongly starlike functions of order  $\alpha$ . And for  $A, B \in [-1, 1]$  and  $\alpha = 1$  we obtain the class of Janowski starlike functions, denoted as  $\mathcal{S}^*(A, B)$ , which specifically reduces to many well-known classes, for instance see [5]. Sokól and Stankiewicz [15] considered the class  $\mathcal{S}_L^*$  by opting  $\phi(z) = \sqrt{1 + z}$ . Geometrically  $\mathcal{S}_L^* = \{f \in \mathcal{S}; |(zf'(z)/f(z))^2 - 1| < 1\}$ , that is,  $zf'(z)/f(z)$  lies in the domain bounded by the right lemniscate of Bernoulli. Moreover, Goel and Kumar [4] studied the starlike class associated with sigmoid,  $\mathcal{S}^*(2/(1 + e^{-z})) =: \mathcal{S}_{SG}^*$  and Mendiratta et al. [11] introduced and studied the class associated with exponential,  $\mathcal{S}^*(e^z) =: \mathcal{S}_e^*$ . Similarly, in [8] the following class was introduced by Kumar and Yadav:

$$\mathcal{S}_{\mathfrak{B}}^* := \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} \prec \sqrt{1 + \tanh z} =: \mathfrak{B}(z) \right\}.$$

---

2010 *Mathematics Subject Classification.* 30C45, 30C80.

*Key words and phrases.* Analytic functions, Starlike functions, Subordination, Admissibility condition.

Observe that the function  $\mathfrak{B}(z) := \sqrt{1 + \tanh z} = \sqrt{2/(1 + e^{-2z})}$  conformally maps  $\mathbb{D}$  onto the following bean shaped region

$$\Omega_{\mathfrak{B}} := \left\{ w \in \mathbb{C} : \left| \log \left( \frac{w^2}{2 - w^2} \right) \right| < 2 \right\}. \quad (1.2)$$

In [8], authors obtained various geometric characteristics of  $\mathfrak{B}(z)$  and derived several inclusion and radii results associated with  $\mathcal{S}_{\mathfrak{B}}^*$ . The present article is an extension of that work in terms of differential subordination. Here, we are establishing certain first and second order differential subordination implications related to the function  $\mathfrak{B}(z)$ , utilizing a lemma that significantly simplifies the calculations. Differential subordination, analogous to differential inequalities in real analysis, has numerous applications. A pivotal contribution to this field was made by Miller and Mocanu's influential monograph [12], which sparked a major revolution in the geometric function theory research community. Many researchers have addressed problems related to differential subordination, such as those in [3, 6, 13, 14, 16]. Chapter 2 of the aforementioned monograph presents a set of results that establish admissibility conditions for second-order differential subordination. Significant work has been built upon these results. For instance, Madaan et al. [10] investigated the class of admissible functions associated with the lemniscate of Bernoulli and proved several first- and second-order differential subordination implications. In [16], the author established various second- and third-order differential subordination implications by defining admissibility conditions for the exponential function. In [1], the authors modified the concept of second-order differential subordination by introducing  $\beta$ -admissible functions, and in [2], the authors extended the concept of differential subordination and admissibility conditions to third order.

We now present essential definitions, lemmas, and notations necessary for further discussion.

**Lemma 1.1.** [8] *Let  $w \in \mathbb{C}$  such that  $|w| > \sqrt{2}$  and  $R_0 = e\sqrt{2/(e^2 - 1)} \simeq 1.5208$ . Then*

$$\left| \log \left( \frac{w^2}{2 - w^2} \right) \right| \geq 2, \quad \text{whenever } |w| \geq R_0.$$

**Definition 1.2.** [12] Let  $\mathcal{Q}$  denotes the set of functions  $q$  that are analytic and univalent on  $\overline{\mathbb{D}} \setminus \mathcal{E}(q)$ , where

$$\mathcal{E}(q) = \left\{ \zeta \in \partial\mathbb{D} : \lim_{z \rightarrow \zeta} q(z) = \infty \right\}$$

such that  $q'(\zeta) \neq 0$  for  $\zeta \in \partial\mathbb{D} \setminus \mathcal{E}(q)$ .

**Definition 1.3.** [12] Let  $\Omega$  be a set in  $\mathbb{C}$ ,  $q \in \mathcal{Q}$  and  $n$  be a positive integer, the class of admissible functions  $\Psi_n(\Omega, q)$ , consists of those functions  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  that satisfy the admissibility condition  $\psi(r, s, t; z) \notin \Omega$ , whenever  $r = q(\zeta)$ ,  $s = m\zeta q'(\zeta)$ ,

$$\operatorname{Re} \left( \frac{t}{s} + 1 \right) \geq m \operatorname{Re} \left( \frac{\zeta q''(\zeta)}{q'(\zeta)} + 1 \right),$$

$z \in \mathbb{D}$ ,  $\zeta \in \partial\mathbb{D} \setminus \mathcal{E}(q)$  and  $m \geq n$ . In particular, we denote  $\Psi_1(\Omega, q)$  as  $\Psi(\Omega, q)$ .

**Lemma 1.4.** [12] *Let  $\psi \in \Psi(\Omega, q)$  with  $q(0) = 1$ . If  $p \in \mathcal{H}$  satisfies*

$$\psi(p(z), zp'(z), z^2 p''(z); z) \in \Omega,$$

*then  $p \prec q$ .*

Throughout this article, we consider the function  $q(z) = \mathfrak{B}(z)$  and define the admissibility class  $\Psi(\Omega, q)$ , where  $\Omega \subset \mathbb{C}$ . Since  $\mathcal{E}(\mathfrak{B}) = \emptyset$ , therefore  $\mathfrak{B} \in \mathcal{Q}$ . For  $\zeta = e^{i\theta}$  ( $0 \leq \theta < 2\pi$ ) we have,

$$q(\zeta) = \sqrt{\frac{2}{1 + e^{-2e^{i\theta}}}}, \quad \zeta q'(\zeta) = \frac{\sqrt{2}e^{i\theta}e^{-2e^{i\theta}}}{(1 + e^{-2e^{i\theta}})^{3/2}}$$

and

$$\operatorname{Re} \left( \frac{\zeta q''(\zeta)}{q'(\zeta)} \right) = \frac{\cos \theta - 2e^{4 \cos \theta} \cos \theta + e^{2 \cos \theta} (\cos(\theta - 2 \sin \theta) - 2 \cos(\theta + 2 \sin \theta))}{1 + e^{4 \cos \theta} + 2e^{2 \cos \theta} \cos(2 \sin \theta)} =: g(\theta). \quad (1.3)$$

Here,

$$|q(\zeta)| = \frac{\sqrt{2}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{1}{4}}} =: \omega(\theta). \quad (1.4)$$

From [8], we obtain that the minimum value of  $\omega(\theta)$  is  $\omega(\pi) = \sqrt{2/(1+e^2)}$  and maximum is  $\omega(\theta_0) =: \omega_{\theta_0} \simeq 1.438$ , where  $\theta_0 \simeq 1.31364$ . By simple analysis we also obtain that  $\min g(\theta) = g(0)$ , and

$$|\zeta q'(\zeta)| = \frac{\sqrt{2}e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{3/4}} := d(\theta). \quad (1.5)$$

Further, a computation shows that  $d'(\theta) = 0$  if and only if  $\theta \in \{0, \theta_1, \pi\}$  where  $\theta_1 \simeq 1.639$ . We observe that  $d(\theta)$  is increasing in  $[0, \theta_1]$  and decreasing in  $[\theta_1, \pi]$ . Therefore,  $\max d(\theta) = d(1.639) =: d_{\theta_1} \simeq 0.905$  and  $\min d(\theta) = \min\{d(0), d(\pi)\} = \min\{0.158, 0.304\} = d(0)$ . Moreover,  $\Psi(\Omega, \mathfrak{B})$  is defined to be the class of functions  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$ , which satisfies the following conditions:

$$\psi(r, s, t; z) \notin \Omega,$$

whenever

$$r = \mathfrak{B}(\zeta) = \sqrt{\frac{2}{1 + e^{-2e^{i\theta}}}}; \quad s = m\zeta \mathfrak{B}'(\zeta) = \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1 + e^{-2e^{i\theta}})^{3/4}}; \quad \operatorname{Re} \left( 1 + \frac{s}{t} \right) = m(1 + g(\theta)),$$

where  $z \in \mathbb{D}$ ,  $0 \leq \theta < 2\pi$  and  $m \geq 1$ . Here, we start from first order differential subordination.

## 2. FIRST ORDER DIFFERENTIAL SUBORDINATION

We first establish the following lemmas to prove our main results:

**Lemma 2.1.** *Let  $-1 < B \leq 0 < A \leq 1$  and  $w \in \mathbb{C}$  such that  $|w| > 1$ , then*

$$\left| \frac{w-1}{A-Bw} \right| \geq 1, \quad \text{whenever } |w| \geq \frac{1+A}{1+B}.$$

*Proof.* Let  $w = Re^{i\theta}$  where  $R > 1$  and  $\theta \in (0, 2\pi]$ . Then

$$\left| \frac{w-1}{A-Bw} \right| \geq 1$$

is equivalent to

$$g(R, \theta) := \frac{1 + R^2 - 2R \cos \theta}{A^2 + B^2 R^2 - 2ABR \cos \theta} \geq 1.$$

A simple computation shows that  $g(R, \theta)$  attains its minimum at  $\theta = 0$ . Thus the above inequality holds if  $g(R, 0) \geq 1$ , i.e.

$$\frac{(R-1)^2}{(A-BR)^2} \geq 1. \quad (2.1)$$

Since  $|w| = R > 1$  and  $B \leq 0$  then  $BR \leq B < A$ , thus (2.1) reduces to

$$\frac{R-1}{A-BR} \geq 1 \implies R \geq \frac{1+A}{1+B}.$$

Thus the result holds. ■

**Lemma 2.2.** *Let  $w \in \mathbb{C}$  such that  $|w| > 1$ , then*

$$|w^2 - 1| \geq 1, \quad \text{whenever } |w| \geq \sqrt{2}.$$

*Proof.* Let  $w = Re^{i\theta}$  where  $R > 1$  and  $\theta \in (0, 2\pi]$ . Then

$$|w^2 - 1| \geq 1$$

is equivalent to

$$L(R, \theta) := 1 + R^4 - 2R^2 \cos 2\theta \geq 1.$$

Clearly, the above inequality holds if  $\min L(R, \theta) \geq 1$ . Since,  $\min L(R, \theta) = L(R, 0)$  and  $R > 1$ , thus the result holds.  $\blacksquare$

In this section, we find conditions on  $\psi(p(z), zp'(z); z)$ , so that the following implication holds:

$$\psi(p(z), zp'(z); z) \prec h(z) \implies p(z) \prec \mathfrak{B}(z),$$

where  $h(z)$  is either  $\mathfrak{B}(z)$ ,  $(1 + Az)/(1 + Bz)$  ( $-1 < B \leq 0 < A \leq 1$ ) or  $\sqrt{1+z}$ . Here,  $\psi(p(z), zp'(z); z)$  is taken to be as follows:  $(p(z))^\delta + \beta(zp'(z))^n$ ,  $(p(z))^\delta + \beta zp'(z)/(p(z))^n$  and  $p(z) + zp'(z)/(\beta p(z) + \gamma)$ .

**Theorem 2.3.** *Let  $n$  be a positive integer,  $\delta \in \{0, 1\}$ ,  $R_0 = e\sqrt{2/(e^2 - 1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta \in \mathbb{C}$  such that  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{3n/2}/(\sqrt{2}e)^n$ . Suppose  $p(z) \in \mathcal{H}$  and satisfies the subordination*

$$(p(z))^\delta + \beta(zp'(z))^n \prec \mathfrak{B}(z),$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1 + e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = q(\mathbb{D}) = \Omega_{\mathfrak{B}}$ . Now consider  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  defined by  $\psi(r, s; z) = r^\delta + \beta s^n$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega_{\mathfrak{B}}, \mathfrak{B})$ , it must satisfy

$$\psi \left( \sqrt{\frac{2}{1 + e^{-2e^{i\theta}}}}, \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1 + e^{-2e^{i\theta}})^{3/2}}; z \right) \notin \Omega_{\mathfrak{B}} \quad (z \in \mathbb{D}, 0 \leq \theta < 2\pi, m \geq 1),$$

which is equivalent to

$$\left| \log \left( \frac{(r^\delta + \beta s^n)^2}{2 - (r^\delta + \beta s^n)^2} \right) \right| \geq 2.$$

We now use the Lemma 1.1 to prove the result, for which, let us assume  $w = r^\delta + \beta s^n$ . From (1.4) and (1.5) respectively, we have  $|r| = |\mathfrak{B}(\zeta)| = |q(\zeta)| = \omega(\theta) \leq \max \omega(\theta) = \omega_{\theta_0}$  and  $|s|/m = |\zeta \mathfrak{B}'(\zeta)| = |\zeta q'(\zeta)| = d(\theta) \geq \min d(\theta) = d(0)$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \beta s^n| \\ &\geq |\beta| |s|^n - |r|^\delta \\ &\geq |\beta| m^n (d(0))^n - \omega_{\theta_0}^\delta. \end{aligned}$$

Since  $m \geq 1$  and  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{3n/2}/(\sqrt{2}e)^n$ , we have

$$|w| \geq |\beta| \left( \frac{\sqrt{2}e}{(1 + e^2)^{3/2}} \right)^n - \omega_{\theta_0}^\delta \geq R_0 > \sqrt{2},$$

therefore the result follows at once by Lemma 1.1.  $\blacksquare$

**Corollary 2.4.** *Let  $n$  be a positive integer,  $\delta \in \{0, 1\}$ ,  $R_0 = e\sqrt{2/(e^2 - 1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta \in \mathbb{C}$  such that  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{3n/2}/(\sqrt{2}e)^n$ . Suppose  $f \in \mathcal{A}$  such that*

$$\left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right)^n \prec \mathfrak{B}(z),$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta(zp'(z))^n = \left(\frac{zf'(z)}{f(z)}\right)^\delta + \beta\left(\frac{zf'(z)}{f(z)}\left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)}\right)\right)^n.$$

Therefore the result holds using Theorem 2.3.  $\blacksquare$

**Theorem 2.5.** *Let  $n$  be a positive integer,  $\delta \in \{0, 1\}$ ,  $R_0 = e\sqrt{2/(e^2 - 1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta \in \mathbb{C}$  such that  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{(3-n)/2}/(\sqrt{2}e)^{1-n}$ . Suppose  $p(z) \in \mathcal{H}$  and satisfies the subordination*

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} \prec \mathfrak{B}(z),$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1 + e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = q(\mathbb{D}) = \Omega_{\mathfrak{B}}$ . Now consider  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be as  $\psi(r, s; z) = r^\delta + \beta s/r^n$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega_{\mathfrak{B}}, \mathfrak{B})$ , we must have

$$\psi\left(\sqrt{\frac{2}{1 + e^{-2e^{i\theta}}}}, \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1 + e^{-2e^{i\theta}})^{3/2}}; z\right) \notin \Omega_{\mathfrak{B}} \quad (z \in \mathbb{D}, 0 \leq \theta < 2\pi, m \geq 1),$$

which is equivalent to

$$\left|\log\left(\frac{(r^\delta + \beta s/r^n)^2}{2 - (r^\delta + \beta s/r^n)^2}\right)\right| \geq 2.$$

We now use the Lemma 1.1 to prove the result, for which, let us assume  $w = r^\delta + \beta s/r^n$ . From (1.4), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= \left|r^\delta + \beta \frac{s}{r^n}\right| \\ &\geq |\beta| \left(\frac{2^{\frac{1-n}{2}} m e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}}\right) - (\omega(\theta))^\delta \\ &\geq |\beta| m \chi_n(\theta) - (\omega(\theta))^\delta, \end{aligned}$$

where

$$\chi_n(\theta) := \frac{2^{\frac{1-n}{2}} e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}}.$$

After a simple computation we obtain that  $\min \chi_n(\theta) = \chi_n(0)$  for all  $n \geq 1$ . Since  $m \geq 1$  and  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{(3-n)/2}/(\sqrt{2}e)^{1-n}$ , we have

$$|w| \geq |\beta| \left(\frac{(\sqrt{2}e)^{1-n}}{(1 + e^2)^{\frac{3-n}{2}}}\right) - \omega_{\theta_0}^\delta \geq R_0 > \sqrt{2}.$$

Therefore, the result follows immediately from Lemma 1.1.  $\blacksquare$

**Corollary 2.6.** *Let  $n$  be a positive integer,  $\delta \in \{0, 1\}$ ,  $R_0 = e\sqrt{2/(e^2 - 1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta \in \mathbb{C}$  such that  $|\beta| \geq (R_0 + \omega_{\theta_0}^\delta)(1 + e^2)^{(3-n)/2}/(\sqrt{2}e)^{1-n}$ . Suppose  $f$  be a function in  $\mathcal{A}$  such that*

$$\left(\frac{zf'(z)}{f(z)}\right)^\delta + \beta \left(\frac{zf'(z)}{f(z)}\right)^{1-n} \left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)}\right) \prec \mathfrak{B}(z),$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} = \left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \right)^{1-n} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right).$$

Now the result follows by an application of Theorem 2.5.  $\blacksquare$

**Theorem 2.7.** Let  $R_0 = e\sqrt{2/(e^2-1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$  with  $|\beta|\omega_{\theta_0} + |\gamma| \leq \sqrt{2}e/((1+e^2)^{3/2}(R_0 + \omega_{\theta_0}))$ . Suppose  $p(z) \in \mathcal{H}$  and satisfies the subordination

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} \prec \mathfrak{B}(z),$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = q(\mathbb{D}) = \Omega_{\mathfrak{B}}$ . Now consider  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be as  $\psi(r, s; z) = r + s/(\beta r + \gamma)$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega_{\mathfrak{B}}, \mathfrak{B})$ , we must have

$$\psi \left( \sqrt{\frac{2}{1+e^{-2e^{i\theta}}}}, \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1+e^{-2e^{i\theta}})^{3/2}}; z \right) \notin \Omega_{\mathfrak{B}} \quad (z \in \mathbb{D}, 0 \leq \theta < 2\pi, m \geq 1),$$

which is equivalent to

$$\left| \log \left( \frac{(r + s/(\beta r + \gamma))^2}{2 - (r + s/(\beta r + \gamma))^2} \right) \right| \geq 2.$$

We now use the Lemma 1.1 to prove the result, for which, let us assume  $w = r + s/(\beta r + \gamma)$ . From (1.4) and (1.5), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$  and  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$ . Thus

$$\begin{aligned} |w| &= \left| r + \frac{s}{\beta r + \gamma} \right| \\ &\geq \frac{|s|}{|\beta||r| + |\gamma|} - |r| \\ &\geq \frac{md(0)}{|\beta|\omega_{\theta_0} + |\gamma|} - \omega_{\theta_0}. \end{aligned}$$

Since  $m \geq 1$  and  $|\beta|\omega_{\theta_0} + |\gamma| \leq \sqrt{2}e/((1+e^2)^{3/2}(R_0 + \omega_{\theta_0}))$ , we have

$$|w| \geq \frac{\sqrt{2}e}{(1+e^2)^{3/2}(|\beta|\omega_{\theta_0} + |\gamma|)} - \omega_{\theta_0} \geq R_0 > \sqrt{2},$$

therefore the result follows at once by Lemma 1.1.  $\blacksquare$

**Corollary 2.8.** Let  $R_0 = e\sqrt{2/(e^2-1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$  with  $\omega_{\theta_0}|\beta| + |\gamma| \leq \sqrt{2}e/((1+e^2)^{3/2}(R_0 + \omega_{\theta_0}))$ . Suppose  $f$  be a function in  $\mathcal{A}$  such that

$$\frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right) \prec \mathfrak{B}(z),$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} = \frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right).$$

Therefore the result holds using Theorem 2.7.  $\blacksquare$

**Theorem 2.9.** Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\beta \in \mathbb{C} \setminus \{0\}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $-1 < B \leq 0 < A \leq 1$ . Suppose  $p \in \mathcal{H}$  such that it satisfies

$$(p(z))^\delta + \beta(zp'(z))^n \prec \frac{1 + Az}{1 + Bz}$$

where

$$|\beta| \geq \left( \frac{(1 + e^2)^{3/2}}{\sqrt{2}e} \right)^n \left( \frac{1 + A}{1 + B} + \omega_{\theta_0}^\delta \right) \quad (2.2)$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1 + e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |(w-1)/(A-Bw)| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r^\delta + \beta s^n$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi\mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1 + e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \mathbb{D}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , which is equivalent to

$$\left| \frac{r^\delta + \beta s^n - 1}{A - B(r^\delta + \beta s^n)} \right| \geq 1.$$

We now use the Lemma 2.1 to prove the result, for which, let us assume  $w = r^\delta + \beta s^n$ . From (1.4) and (1.5), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$  and  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \beta s^n| \\ &\geq |\beta||s|^n - |r|^\delta \\ &\geq |\beta|m^n(d(0))^n - \omega_{\theta_0}^\delta \end{aligned}$$

Since  $m \geq 1$  and using (2.2), we have

$$|w| \geq |\beta| \left( \frac{\sqrt{2}e}{(1 + e^2)^{3/2}} \right)^n - \omega_{\theta_0}^\delta \geq \frac{1 + A}{1 + B} > 1,$$

therefore the result follows at once from Lemma 2.1.  $\blacksquare$

**Corollary 2.10.** Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$  and  $\beta \in \mathbb{C} \setminus \{0\}$  such that  $\beta$  satisfies inequality (2.2). Moreover, let  $\omega_{\theta_0} \simeq 1.438$  and  $-1 < B \leq 0 < A \leq 1$ . Suppose  $f \in \mathcal{A}$  such that

$$\left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right)^n \prec \frac{1 + Az}{1 + Bz},$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta(zp'(z))^n = \left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right)^n.$$

Now the result follows directly from Theorem 2.9.  $\blacksquare$

**Theorem 2.11.** Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\beta \in \mathbb{C} \setminus \{0\}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $-1 \leq B < 0 \leq A \leq 1$ . Suppose  $p \in \mathcal{H}$  such that it satisfies

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} \prec \frac{1 + Az}{1 + Bz},$$

where

$$|\beta| \geq \left( \frac{(1 + e^2)^{\frac{3-n}{2}}}{(\sqrt{2}e)^{1-n}} \right) \left( \frac{1 + A}{1 + B} + \omega_{\theta_0}^\delta \right). \quad (2.3)$$

Then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |(w-1)/(A-Bw)| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r^\delta + \beta s/r^n$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi\mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1+e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \overline{\mathbb{D}}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , which is equivalent to

$$\left| \frac{r^\delta + \beta \frac{s}{r^n} - 1}{A - B(r^\delta + \beta \frac{s}{r^n})} \right| \geq 1.$$

We now use the Lemma 2.1 to prove the result, for which, let us assume  $w = r^\delta + \beta s/r^n$ . From (1.4), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= \left| r^\delta + \beta \frac{s}{r^n} \right| \\ &\geq |\beta| \left( \frac{2^{\frac{1-n}{2}} m e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}} \right) - (\omega(\theta))^\delta \\ &\geq |\beta| m \chi_n(\theta) - (\omega(\theta))^\delta, \end{aligned}$$

where

$$\chi_n(\theta) := \frac{2^{\frac{1-n}{2}} e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}}.$$

After a simple computation we obtain that  $\min \chi_n(\theta) = \chi_n(0)$  for all  $n \geq 1$ . Since  $m \geq 1$  and using (2.3), we have

$$|w| \geq |\beta| \left( \frac{(\sqrt{2}e)^{1-n}}{(1+e^2)^{\frac{3-n}{2}}} \right) - \omega_{\theta_0}^\delta \geq \frac{1+A}{1+B} > 1,$$

therefore the result follows from Lemma 2.1. ■

**Corollary 2.12.** *Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$  and  $\beta \in \mathbb{C} \setminus \{0\}$  such that  $\beta$  satisfies inequality (2.3). Moreover, let  $\omega_{\theta_0} \simeq 1.438$  and  $-1 < B \leq 0 < A \leq 1$ . Suppose  $f$  be a function in  $\mathcal{A}$  such that*

$$\left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \right)^{1-n} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \frac{1+Az}{1+Bz},$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} = \left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \right)^{1-n} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right).$$

Now the result follows by an application of Theorem 2.11. ■

**Theorem 2.13.** *Let  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $-1 < B \leq 0 < A \leq 1$ . Suppose  $p \in \mathcal{H}$  such that it satisfies*

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} \prec \frac{1+Az}{1+Bz}$$

provided

$$|\beta| \omega_{\theta_0} + |\gamma| \leq \left( \frac{(1+e^2)^{3/2}}{\sqrt{2}e} \left( \frac{1+A}{1+B} + \omega_{\theta_0} \right) \right)^{-1} \quad (2.4)$$

where Then  $p(z) \prec \mathfrak{B}(z)$ .



*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |(w-1)/(A-Bw)| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r + s/(\beta r + \gamma)$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi\mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1+e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \overline{\mathbb{D}}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , which is equivalent to

$$\left| \frac{r + \frac{s}{\beta r + \gamma}}{A - B\left(r + \frac{s}{\beta r + \gamma}\right)} \right| \geq 1.$$

We now use the Lemma 2.1 to prove the result, for which, let us assume  $w = r + s/(\beta r + \gamma)$ . From (1.4) and (1.5), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$  and  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$ . Thus

$$\begin{aligned} |w| &= \left| r + \frac{s}{\beta r + \gamma} \right| \\ &\geq \frac{|s|}{|\beta||r| + |\gamma|} - |r| \\ &\geq \frac{md(0)}{|\beta|\omega_{\theta_0} + |\gamma|} - \omega_{\theta_0}. \end{aligned}$$

Since  $m \geq 1$  and using (2.4), we have

$$|w| \geq \frac{\sqrt{2}e}{(1+e^2)^{3/2}(|\beta|\omega_{\theta_0} + |\gamma|)} - \omega_{\theta_0} \geq \frac{1+A}{1+B} > 1,$$

therefore the result follows at once from Lemma 2.1.  $\blacksquare$

**Corollary 2.14.** *Let  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$  and satisfies inequality (2.4). Moreover, let  $\omega_{\theta_0} \simeq 1.438$  and  $-1 < B \leq 0 < A \leq 1$ . Suppose  $f$  be a function in  $\mathcal{A}$  such that*

$$\frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right) \prec \frac{1+Az}{1+Bz},$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} = \frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right).$$

Thus the result follows by an application of Theorem 2.13.  $\blacksquare$

**Theorem 2.15.** *Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\beta \in \mathbb{C} \setminus \{0\}$  and  $\omega_{\theta_0} \simeq 1.438$ . Suppose  $p \in \mathcal{H}$  such that it satisfies*

$$(p(z))^\delta + \beta(zp'(z))^n \prec \sqrt{1+z}$$

where

$$|\beta| \geq \left( \frac{(1+e^2)^{3/2}}{\sqrt{2}e} \right)^n \left( \sqrt{2} + \omega_{\theta_0}^\delta \right), \quad (2.5)$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |w^2 - 1| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r^\delta + \beta s^n$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi\mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1+e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \overline{\mathbb{D}}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , which is equivalent to

$$\left| (r^\delta + \beta s^n)^2 - 1 \right| \geq 1.$$

We now use the Lemma 2.2 to prove the result, for which, let us assume  $w = r^\delta + \beta s^n$ . From (1.4) and (1.5), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$  and  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \beta s^n| \\ &\geq |\beta| |s|^n - |r|^\delta \\ &\geq |\beta| m^n (d(0))^n - \omega_{\theta_0}^\delta. \end{aligned}$$

Since  $m \geq 1$  and using (2.5), we have

$$|w| \geq |\beta| \left( \frac{\sqrt{2}e}{(1+e^2)^{3/2}} \right)^n - \omega_{\theta_0}^\delta \geq \sqrt{2} > 1,$$

therefore the result follows from Lemma 2.2.  $\blacksquare$

**Corollary 2.16.** *Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\beta \in \mathbb{C} \setminus \{0\}$  such that it satisfies (2.5) and  $\omega_{\theta_0} \simeq 1.438$ . Further if  $f \in \mathcal{A}$  and satisfies*

$$\left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right)^n \prec \sqrt{1+z}$$

Then  $p(z) \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta (zp'(z))^n = \left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right)^n.$$

Now the result follows from Theorem 2.15.  $\blacksquare$

**Theorem 2.17.** *Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\beta \in \mathbb{C} \setminus \{0\}$  and  $\omega_{\theta_0} \simeq 1.438$ . Suppose  $p \in \mathcal{H}$  such that it satisfies*

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} \prec \sqrt{1+z}$$

where

$$|\beta| \geq \left( \frac{(1+e^2)^{\frac{3-n}{2}}}{(\sqrt{2}e)^{1-n}} \right) (\sqrt{2} + \omega_{\theta_0}^\delta) \quad (2.6)$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |w^2 - 1| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r^\delta + \beta s/r^n$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi \mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1+e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \overline{\mathbb{D}}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , which is equivalent to

$$\left| \left( r^\delta + \beta \frac{s}{r^n} \right)^2 - 1 \right| \geq 1.$$

We now use the Lemma 2.2 to prove the result, for which, let us assume  $w = r^\delta + \beta s/r^n$ . From (1.4), we have  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= \left| r^\delta + \beta \frac{s}{r^n} \right| \\ &\geq |\beta| \left( \frac{2^{\frac{1-n}{2}} m e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}} \right) - (\omega(\theta))^\delta \\ &\geq |\beta| m \chi_n(\theta) - (\omega(\theta))^\delta, \end{aligned}$$

where

$$\chi_n(\theta) := \frac{2^{\frac{1-n}{2}} e^{-2 \cos \theta}}{(1 + e^{-4 \cos \theta} + 2e^{-2 \cos \theta} \cos(2 \sin \theta))^{\frac{3-n}{4}}}.$$

After a simple computation we obtain that  $\min \chi_n(\theta) = \chi_n(0)$  for all  $n \geq 1$ . Since  $m \geq 1$  and using (2.6), we have

$$|w| \geq |\beta| \left( \frac{(\sqrt{2}e)^{1-n}}{(1+e^2)^{\frac{3-n}{2}}} \right) - \omega_{\theta_0}^\delta \geq \sqrt{2} > 1,$$

therefore the result follows from Lemma 2.2.  $\blacksquare$

**Corollary 2.18.** *Let  $n \in \mathbb{N}$ ,  $\delta \in \{0, 1\}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta \in \mathbb{C} \setminus \{0\}$  such that  $\beta$  satisfies inequality (2.6). Suppose  $f$  be a function in  $\mathcal{A}$  such that*

$$\left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \right)^{1-n} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \sqrt{1+z},$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$(p(z))^\delta + \beta \frac{zp'(z)}{(p(z))^n} = \left( \frac{zf'(z)}{f(z)} \right)^\delta + \beta \left( \frac{zf'(z)}{f(z)} \right)^{1-n} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right).$$

Now the result follows as an application of Theorem 2.17.  $\blacksquare$

**Theorem 2.19.** *Let  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$  and  $\omega_{\theta_0} \simeq 1.438$ . Suppose  $p \in \mathcal{H}$  such that it satisfies*

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} \prec \sqrt{1+z}$$

provided

$$|\beta|\omega_{\theta_0} + |\gamma| \leq \left( \frac{(1+e^2)^{3/2}}{\sqrt{2}e} (\sqrt{2} + \omega_{\theta_0}) \right)^{-1} \quad (2.7)$$

where Then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \sqrt{2/(1+e^{-2z})} =: \mathfrak{B}(z)$  and  $\Omega = \{w \in \mathbb{C} : |w^2 - 1| < 1\}$ . Let  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  be defined as  $\psi(r, s; z) = r + s/(\beta r + \gamma)$  where  $r = \mathfrak{B}(\xi)$ ,  $s = m\xi\mathfrak{B}'(\xi) = m\sqrt{2}\xi e^{-2\xi}/(1+e^{-2\xi})^{3/2}$  for  $z \in \mathbb{D}$ ,  $\xi \in \mathbb{D}$  and  $m \geq 1$ . For  $\psi$  to be an admissible function and for  $\psi \in \Psi[\Omega, \mathfrak{B}]$  we need to show  $\psi(\mathbb{D}) \not\subseteq \Omega$ , i.e.

$$\left| \left( r + \frac{s}{\beta r + \gamma} \right)^2 - 1 \right| \geq 1.$$

We now use the Lemma 2.2 to prove the result, for which, let us assume  $w = r + s/(\beta r + \gamma)$ . From (1.4) and (1.5), we have  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$  and  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= \left| r + \frac{s}{\beta r + \gamma} \right| \\ &\geq \frac{|s|}{|\beta||r| + |\gamma|} - |r| \\ &\geq \frac{md(0)}{|\beta|\omega_{\theta_0} + |\gamma|} - \omega_{\theta_0}. \end{aligned}$$

Since  $m \geq 1$  and using (2.7), we have

$$|w| \geq \frac{\sqrt{2}e}{(1+e^2)^{3/2}(|\beta|\omega_{\theta_0} + |\gamma|)} - \omega_{\theta_0} \geq \sqrt{2} > 1,$$

therefore, by Lemma 2.2, the result holds.  $\blacksquare$

**Corollary 2.20.** *Let  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C}$  such that  $\beta \neq 0$  and satisfies inequality (2.7). Suppose  $f$  be a function in  $\mathcal{A}$  such that*

$$\frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right) \prec \sqrt{1+z},$$

then  $f \in \mathcal{S}_{\mathfrak{B}}^*$ .

*Proof.* Let  $p(z) = zf'(z)/f(z)$ , then

$$p(z) + \frac{zp'(z)}{\beta p(z) + \gamma} = \frac{zf'(z)}{f(z)} \left( 1 + \left( \beta \frac{zf'(z)}{f(z)} + \gamma \right)^{-1} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right).$$

Therefore the result holds using Theorem 2.19.  $\blacksquare$

### 3. SECOND ORDER DIFFERENTIAL SUBORDINATION

In this section, we consider the following second order differential subordination implication

$$(p(z))^\delta + \gamma zp'(z) + \beta z^2 p''(z) \prec h(z) \implies p(z) \prec \mathfrak{B}(z).$$

Taking  $h(z)$  as  $\mathfrak{B}(z)$ ,  $(1+Az)/(1+Bz)$  ( $-1 < B \leq 0 < A \leq 1$ ) and  $\sqrt{1+z}$ .

**Theorem 3.1.** *Let  $\delta \in \{0, 1\}$ ,  $R_0 = e\sqrt{2/(e^2-1)}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C} \setminus \{0\}$ . If  $p(z) \in \mathcal{H}$  such that*

$$(p(z))^\delta + \gamma zp'(z) + \beta z^2 p''(z) \prec \mathfrak{B}(z),$$

provided

$$\sqrt{2}e(\gamma(1+e^2)^2 + \beta(1-2e^4-e^2)) \geq (R_0 + \omega_{\theta_0}^\delta)(e^2+1)^{7/2} \quad (3.1)$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Let  $q(z) = \mathfrak{B}(z)$  and  $\Omega = q(\mathbb{D}) = \Omega_{\mathfrak{B}}$ . Now consider  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be as  $\psi(r, s, t; z) = r^\delta + \gamma s + \beta t$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega_{\mathfrak{B}}, \mathfrak{B})$ , we must have

$$\psi(r, s, t; z) \notin \Omega_{\mathfrak{B}},$$

whenever

$$r = \mathfrak{B}(\zeta) = \sqrt{\frac{2}{1+e^{-2e^{i\theta}}}}; \quad s = m\zeta \mathfrak{B}'(\zeta) = \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1+e^{-2e^{i\theta}})^{3/4}}; \quad \operatorname{Re}\left(1 + \frac{s}{t}\right) = m(1+g(\theta)),$$

where  $z \in \mathbb{D}$ ,  $0 \leq \theta < 2\pi$  and  $m \geq 1$ . Thus, it is sufficient to show that

$$\left| \log \left( \frac{(r^\delta + \gamma s + \beta t)^2}{2 - (r^\delta + \gamma s + \beta t)^2} \right) \right| \geq 2.$$

We now use the Lemma 1.1 to prove the result, for which, let us assume  $w = r^\delta + \gamma s + \beta t$ . From (1.3), (1.4) and (1.5), we have  $\min g(\theta) = g(0)$ ,  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$  and

$|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \gamma s + \beta t| \\ &\geq \gamma |s| \operatorname{Re} \left( 1 + \frac{\beta t}{\gamma s} \right) - |r|^\delta \\ &\geq \gamma m d(\theta) \left( 1 + \frac{\beta}{\gamma} (m g(\theta) + m - 1) \right) - (\omega(\theta))^\delta \\ &\geq m \frac{\sqrt{2}e}{(e^2 + 1)^{3/2}} \left( \gamma + \beta \left( m \frac{1 - 2e^4 - e^2}{(e^2 + 1)^2} + m - 1 \right) \right) - \omega_{\theta_0}^\delta. \end{aligned}$$

Since  $m \geq 1$  and using (3.1), we have

$$|w| \geq \frac{\sqrt{2}e}{(e^2 + 1)^{3/2}} \left( \gamma + \beta \left( \frac{1 - 2e^4 - e^2}{(e^2 + 1)^2} \right) \right) - \omega_{\theta_0}^\delta \geq R_0 > \sqrt{2},$$

therefore the result follows at once by Lemma 1.1.  $\blacksquare$

**Theorem 3.2.** *Let  $\delta \in \{0, 1\}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C} \setminus \{0\}$ . If  $p(z) \in \mathcal{H}$  such that*

$$(p(z))^\delta + \gamma z p'(z) + \beta z^2 p''(z) \prec \frac{1 + Az}{1 + Bz}, \quad -1 < B \leq 0 < A \leq 1,$$

provided

$$\sqrt{2}e(\gamma(1 + e^2)^2 + \beta(1 - 2e^4 - e^2)) \geq \left( \frac{1 + A}{1 + B} + \omega_{\theta_0}^\delta \right) (e^2 + 1)^{7/2} \quad (3.2)$$

then  $p(z) \prec \mathfrak{B}(z)$ .

*Proof.* Here  $q(z) = \mathfrak{B}(z)$  and  $h(z) = (1 + Az)/(1 + Bz)$ , thus

$$\Omega = h(\mathbb{D}) = \left\{ w \in \mathbb{C} : \left| \frac{w - 1}{A - Bw} \right| < 1 \right\},$$

Now consider  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be as  $\psi(r, s, t; z) = r^\delta + \gamma s + \beta t$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega, \mathfrak{B})$ , we must have

$$\psi(r, s, t; z) \notin \Omega,$$

whenever

$$r = \mathfrak{B}(\zeta) = \sqrt{\frac{2}{1 + e^{-2e^{i\theta}}}}; \quad s = m\zeta \mathfrak{B}'(\zeta) = \frac{\sqrt{2}m e^{i\theta} e^{-2e^{i\theta}}}{(1 + e^{-2e^{i\theta}})^{3/4}}; \quad \operatorname{Re} \left( 1 + \frac{s}{t} \right) = m(1 + g(\theta)),$$

where  $z \in \mathbb{D}$ ,  $0 \leq \theta < 2\pi$  and  $m \geq 1$ . i.e. it is sufficient to show that

$$\left| \frac{r^\delta + \gamma s + \beta t - 1}{A - B(r^\delta + \gamma s + \beta t)} \right| \geq 1.$$

We now use the Lemma 2.1 to prove the result, for which, let us assume  $w = r^\delta + \gamma s + \beta t$ . From (1.3), (1.4) and (1.5), we have  $\min g(\theta) = g(0)$ ,  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$  and  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \gamma s + \beta t| \\ &\geq \gamma |s| \operatorname{Re} \left( 1 + \frac{\beta t}{\gamma s} \right) - |r|^\delta \\ &\geq \gamma m d(\theta) \left( 1 + \frac{\beta}{\gamma} (m g(\theta) + m - 1) \right) - (\omega(\theta))^\delta \\ &\geq m \frac{\sqrt{2}e}{(e^2 + 1)^{3/2}} \left( \gamma + \beta \left( m \frac{1 - 2e^4 - e^2}{(e^2 + 1)^2} + m - 1 \right) \right) - \omega_{\theta_0}^\delta. \end{aligned}$$

Since  $m \geq 1$  and by (3.2), we have

$$|w| \geq \frac{\sqrt{2}e}{(e^2+1)^{3/2}} \left( \gamma + \beta \left( \frac{1-2e^4-e^2}{(e^2+1)^2} \right) \right) - \omega_{\theta_0}^\delta \geq \frac{1+A}{1+B} > 1,$$

therefore, by Lemma 2.1, the result holds. ■

**Theorem 3.3.** *Let  $\delta \in \{0, 1\}$ ,  $\omega_{\theta_0} \simeq 1.438$  and  $\beta, \gamma \in \mathbb{C} \setminus \{0\}$ . If  $p(z) \in \mathcal{H}$  such that*

$$(p(z))^\delta + \gamma z p'(z) + \beta z^2 p''(z) \prec \sqrt{1+z},$$

*provided*

$$\sqrt{2}e(\gamma(1+e^2)^2 + \beta(1-2e^4-e^2)) \geq \left( \sqrt{2} + \omega_{\theta_0}^\delta \right) (e^2+1)^{7/2} \quad (3.3)$$

*then  $p(z) \prec \mathfrak{B}(z)$ .*

*Proof.* Here  $q(z) = \mathfrak{B}(z)$  and  $h(z) = \sqrt{1+z}$ , thus

$$\Omega = h(\mathbb{D}) = \{w \in \mathbb{C} : |w^2 - 1| < 1\},$$

Now consider  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  be as  $\psi(r, s, t; z) = r^\delta + \gamma s + \beta t$ . For  $\psi$  to lie in the admissible class  $\Psi(\Omega, \mathfrak{B})$ , we must have

$$\psi(r, s, t; z) \notin \Omega,$$

whenever

$$r = \mathfrak{B}(\zeta) = \sqrt{\frac{2}{1+e^{-2e^{i\theta}}}}; \quad s = m\zeta \mathfrak{B}'(\zeta) = \frac{\sqrt{2}me^{i\theta}e^{-2e^{i\theta}}}{(1+e^{-2e^{i\theta}})^{3/4}}; \quad \operatorname{Re}\left(1 + \frac{s}{t}\right) = m(1+g(\theta)),$$

where  $z \in \mathbb{D}$ ,  $0 \leq \theta < 2\pi$  and  $m \geq 1$ . i.e. it is sufficient to show that

$$\left| (r^\delta + \gamma s + \beta t)^2 - 1 \right| \geq 1.$$

We now use the Lemma 2.2 to prove the result, for which, let us assume  $w = r^\delta + \gamma s + \beta t$ . From (1.3), (1.4) and (1.5), we have  $\min g(\theta) = g(0)$ ,  $|s|/m = d(\theta) \geq \min d(\theta) = d(0)$  and  $|r| = \omega(\theta) \leq \omega_{\theta_0}$ . Thus

$$\begin{aligned} |w| &= |r^\delta + \gamma s + \beta t| \\ &\geq \gamma |s| \operatorname{Re}\left(1 + \frac{\beta t}{\gamma s}\right) - |r|^\delta \\ &\geq \gamma m d(\theta) \left(1 + \frac{\beta}{\gamma}(mg(\theta) + m - 1)\right) - (\omega(\theta))^\delta \\ &\geq m \frac{\sqrt{2}e}{(e^2+1)^{3/2}} \left( \gamma + \beta \left( m \frac{1-2e^4-e^2}{(e^2+1)^2} + m - 1 \right) \right) - \omega_{\theta_0}^\delta. \end{aligned}$$

Since  $m \geq 1$  and by (3.3), we have

$$|w| \geq \frac{\sqrt{2}e}{(e^2+1)^{3/2}} \left( \gamma + \beta \left( \frac{1-2e^4-e^2}{(e^2+1)^2} \right) \right) - \omega_{\theta_0}^\delta \geq \sqrt{2} > 1,$$

therefore, by Lemma 2.2, the result holds. ■

#### DECLARATIONS

**Ethical Approval** Not Applicable

**Conflicts of interest** The authors declare that they have no competing interests.

**Authors' contributions** All authors contributed equally.

**Funding** Not Applicable

**Availability of data and materials** Not Applicable

## REFERENCES

- [1] R.M. Ali, S. Nagpal, V. Ravichandran, Second-order differential subordination for analytic functions with fixed initial coefficient. *Bull. Malays. Math. Sci. Soc.* **34** (2011), 611–629.
- [2] J. Antonino and S. S. Miller, Third-order differential inequalities and subordinations in the complex plane, *Complex Var. Elliptic Equ.* **56** (2011), no. 5, 439–454.
- [3] N.E. Cho, S. Kumar, V. Kumar and V. Ravichandran, Differential subordination and radius estimates for starlike functions associated with the Booth lemniscate. *Turk. J. Math.* **42** (2018), no. 3, 1380–1399.
- [4] P. Goel and S. S. Kumar, Certain class of starlike functions associated with modified sigmoid function, *Bull. Malays. Math. Sci. Soc.* (2019). <https://doi.org/10.1007/s40840-019-00784-y>.
- [5] W. Janowski, Extremal problems for a family of functions with positive real part and for some related families, *Ann. Polon. Math.* **23** (1970/71), 159–177.
- [6] S. S. Kumar and P. Goel, Starlike functions and higher order differential subordinations, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **114** (2020), no. 4, Paper No. 192, 23 pp.
- [7] S. S. Kumar and P. Yadav, On Oblique Domains of Janowski Function, *Math. Slovaca* **73** (2023), no. 2, 407–424.
- [8] S. S. Kumar and P. Yadav, On starlike functions associated with the bean shaped domain, arXiv preprint arXiv:2403.14162, (2024).
- [9] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, Conf. Proc. Lecture Notes Anal., I, Int. Press, Cambridge, MA.
- [10] V. Madaan, A. Kumar and V. Ravichandran, Starlikeness associated with lemniscate of Bernoulli. *Filomat* **33** (2019), no. 7, 1937–1955.
- [11] R. Mendiratta, S. Nagpal and V. Ravichandran, On a subclass of strongly starlike functions associated with exponential function, *Bull. Malays. Math. Sci. Soc.* **38** (2015), no. 1, 365–386.
- [12] S. S. Miller and P. T. Mocanu, *Differential subordinations*, Monographs and Textbooks in Pure and Applied Mathematics, **225** (2000), Marcel Dekker, Inc., New York.
- [13] S.S. Miller and P.T. Mocanu, Subordinants of differential superordinations. *Complex Var. Theory Appl.* **48** (2003), no. 10, 815–826.
- [14] A. Saliu, K. Jabeen and V. Ravichandran, Differential subordination for certain strongly starlike functions, *Rendiconti del Circolo Matematico di Palermo Series*, **2** (2023), 1–18.
- [15] J. Sokół and J. Stankiewicz, Radius of convexity of some subclasses of strongly starlike functions, *Zeszyty Nauk. Politech. Rzeszowskiej Mat. No. 19* (1996), 101–105.
- [16] N. Verma and S. S. Kumar, Differential subordination of certain class of starlike functions, *Complex Var. Elliptic Equ.* (2024), 1–19. doi: 10.1080/17476933.2024.2326833.

DEPARTMENT OF APPLIED MATHEMATICS, DELHI TECHNOLOGICAL UNIVERSITY, DELHI-110042, INDIA

*Email address:* `spkumar@dce.ac.in`

DEPARTMENT OF APPLIED MATHEMATICS, DELHI TECHNOLOGICAL UNIVERSITY, DELHI-110042, INDIA

*Email address:* `poojayv100@gmail.com`