

FOURTH-ORDER OPERATORS WITH UNBOUNDED COEFFICIENTS IN L^1 SPACES

FEDERICA GREGORIO¹, CHIARA SPINA², CRISTIAN TACELLI¹

¹Dipartimento di Matematica, Università degli Studi di Salerno, Fisciano, Italy

²Dipartimento di Matematica e Fisica "Ennio De Giorgi", Università del Salento, Lecce, Italy

ABSTRACT. We prove that operators of the form $A = -a(x)^2\Delta^2$, with suitable growth conditions on the coefficient $a(x)$, generate analytic semigroups in $L^1(\mathbb{R}^N)$. In particular, we deduce generation results for the operator $A := -(1 + |x|^2)^\alpha\Delta^2$, $0 \leq \alpha \leq 2$. Moreover, we characterise the maximal domain of A in $L^1(\mathbb{R}^N)$.

1. INTRODUCTION

In this paper we are interested in elliptic and parabolic solvability in $L^1(\mathbb{R}^N)$ of the problems associated with fourth order operators of the form

$$A := -a(x)^2\Delta^2$$

with suitable growth conditions on the coefficient $a(x)$. In particular, the results proved in the paper apply to the model operator $A = -(1 + |x|^2)^\alpha\Delta^2$ in the case $0 \leq \alpha \leq 2$. In the language of semigroups, we prove that A , endowed with the domain

$$D(A_1) = \{u \in L^1(\mathbb{R}^N) : a^{\frac{1}{2}}Du, aD^2u, a^{\frac{3}{2}}D^3u, a^2\Delta^2u \in L^1(\mathbb{R}^N)\},$$

generates an analytic semigroup.

Although second order elliptic operators with unbounded coefficients and, in particular, polynomial growing coefficients, have been widely studied, see for example [6, 11, 10, 2], where unbounded coefficients similar to ours have been introduced, it seems that the case of higher order operators has been less exploited.

The semigroup generated by a class of higher order elliptic operators with bounded measurable coefficients in $L^p(\mathbb{R}^N)$ has been systematically studied by Davies, cf. [4]. Unbounded coefficients for higher order operators have been considered in recent times. In [1] the authors obtained generation results for the square of the Kolmogorov operator $L = \Delta + \frac{\nabla\mu}{\mu} \cdot \nabla$ in the weighted space $L^2(\mathbb{R}^N, d\mu)$. In [8] polynomial growing coefficients are considered.

2020 *Mathematics Subject Classification.* 35K35, 35B45, 47D06.

Key words and phrases. Higher order elliptic and parabolic equations, analytic semigroups, a priori estimates, unbounded coefficients, domain characterization.

The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). This article is based upon work from COST Action CA18232 MAT-DYN-NET, supported by COST (European Cooperation in Science and Technology), www.cost.eu.

More precisely, the operator $A = (1 + |x|^\alpha)^2 \Delta^2 + |x|^{2\beta}$ with $\alpha > 0$ and $\beta > (\alpha - 2)^+$ has been studied in the space $L^2(\mathbb{R}^N)$ for $N > 4$.

The present paper extends the results proved in [7] where we studied the operator A in $C_b(\mathbb{R}^N)$ and $L^p(\mathbb{R}^N)$ spaces for $1 < p \leq \infty$. By proving the sectoriality of the operator A , we obtained generation of an analytic semigroup and characterised the maximal domain of such operators in $L^p(\mathbb{R}^N)$ for $1 < p < \infty$.

Here we prove generation results also in $L^1(\mathbb{R}^N)$ by proving that the operator $(A, D(A_1))$ is sectorial. To get the main result, we first prove the resolvent estimate in L^1 in a suitable sector. In order to do this we use a duality argument which is based upon the generation result in continuous function spaces for the adjoint operator

$$-\hat{A}u = a^2 \Delta^2 u + 4Da^2 D\Delta u + 2\Delta a^2 \Delta u + 4trD^2 a^2 D^2 u + 4D\Delta a^2 Du + \Delta^2 a^2 u.$$

Since the last operator contains also explicit lower order terms with unbounded coefficients, we need a preliminary generalization of the above mentioned results in $L^p(\mathbb{R}^N)$, $1 < p \leq \infty$, and then in $C_b(\mathbb{R}^N)$ for operators of this form. For this reason Section 2 is devoted to the study of more general operators with lower order terms via perturbation methods. Once the resolvent estimate is known, an approximation argument with operators with smooth and bounded coefficients allows us to prove that a suitable sector is contained in the resolvent set.

We have a precise description of the domain, indeed we prove that the maximal domain of A in $L^1(\mathbb{R}^N)$ coincides with $D(A_1)$.

Notations. We use standard notations for function spaces. We denote by $L^p(\mathbb{R}^N)$ and $W^{k,p}(\mathbb{R}^N)$ the standard Lebesgue and Sobolev spaces, respectively. $C_c^\infty(\mathbb{R}^N)$ is the space of test functions and $C_b(\mathbb{R}^N)$ is the space of bounded continuous functions. We denote by $W_{\text{comp}}^{3,p}(\mathbb{R}^N)$ the space of functions in $W^{3,p}(\mathbb{R}^N)$ with compact support. B_R denotes the open ball of radius R centred at 0, whereas $B_R(x_0)$ is the ball centred at x_0 for some $x_0 \in \mathbb{R}^N$. We denote by B_R^c the complementary set of the open ball, $B_R^c = \mathbb{R}^N \setminus B_R$. Moreover, we use the symbol $\|\cdot\|_{p,R}$ to shorten the notation $\|\cdot\|_{L^p(B_R(x_0))}$. In the proofs, C is a positive constant that may vary from line to line. Finally, for every function u smooth enough we set

$$\begin{aligned} |D^4 u| &= \left(\sum_{i,j,k,l=1}^N |D_{ijkl} u|^2 \right)^{\frac{1}{2}}, & |D^3 u| &= \left(\sum_{i,j,k=1}^N |D_{ijk} u|^2 \right)^{\frac{1}{2}} \\ |D^2 u| &= \left(\sum_{i,j=1}^N |D_{ij} u|^2 \right)^{\frac{1}{2}}, & |Du| &= \left(\sum_{i=1}^N |D_i u|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

2. GENERATION RESULTS FOR MORE GENERAL OPERATORS

In this section we study generation results in C_b and in L^p spaces, $1 < p \leq \infty$, for operators with lower order terms of the form

$$-Lu = a^2 \Delta^2 u + \sum_{i,j,k=1}^N b_{ijk} D_{ijk} u + \sum_{i,j=1}^N c_{ij} D_{ij} u + \sum_{i=1}^N d_i D_i u + eu$$

where the lower order coefficients satisfy the following growth assumptions

$$(H_p) \quad \begin{aligned} a &\in C^1(\mathbb{R}^N), \quad a(x) > 0, \quad |Da(x)| \leq \kappa a(x)^{\frac{1}{2}}, \\ b_{ijk} &\in C(\mathbb{R}^N), \quad |b_{ijk}(x)| \leq \kappa a^{\frac{3}{2}}(x) \text{ for all } i, j, k = 1, \dots, N, \\ c_{ij} &\in C(\mathbb{R}^N), \quad |c_{ij}(x)| \leq \kappa^2 a(x) \text{ for all } i, j = 1, \dots, N, \\ d_i &\in C(\mathbb{R}^N), \quad |d_i(x)| \leq \kappa^3 a^{\frac{1}{2}}(x) \text{ for all } i = 1, \dots, N, \\ e &\in C(\mathbb{R}^N), \quad |e(x)| \leq \kappa^4, \end{aligned}$$

for all $x \in \mathbb{R}^N$ and some positive constant κ . We endow L with the following domain

$$D(A_p) = \{u \in W_{loc}^{4,p}(\mathbb{R}^N) \cap L^p(\mathbb{R}^N) : a^{\frac{1}{2}} Du, aD^2u, a^{\frac{3}{2}} D^3u, a^2 D^4u \in L^p(\mathbb{R}^N)\}.$$

We write the operator L as a sum operator

$$L = A + B$$

where $A = -a^2 \Delta^2$ is the principal part of L . The operator A has been studied in [7]. In particular, the following generation result holds for $(A, D(A_p))$ in $L^p(\mathbb{R}^N)$, $1 < p < \infty$.

Theorem 1. *Let $1 < p < \infty$ and assume that a satisfies (H_p) . Then, $(A, D(A_p))$ generates an analytic C_0 -semigroup in $L^p(\mathbb{R}^N)$.*

The generation of analytic semigroups was showed by proving the sectoriality of $(A, D(A_p))$ in a sector depending only on N, p and κ .

Having in mind to extend the result to the operator L , we recall the following density result (see [7, Lemma 2.2] for the proof).

Lemma 1. *Let $1 < p < \infty$ and assume that a satisfies (H_p) . Then $C_c^\infty(\mathbb{R}^N)$ is dense in $D(A_p)$ with respect to the norm*

$$\|u\|_{D(A_p)} = \|u\|_p + \|a^{\frac{1}{2}} Du\|_p + \|aD^2u\|_p + \|a^{\frac{3}{2}} D^3u\|_p + \|a^2 D^4u\|_p.$$

By arguing as in [7, Lemma 2.4] we prove the following interpolation estimates. The main novelty is in the last inequality (2.4) where the right hand side contains the whole operator L .

Lemma 2. *Let $1 < p < \infty$ and assume that a, b, c, d, e satisfy (H_p) . Then, there exists a positive constant $C = C(N, p, \kappa)$ such that for every $\varepsilon > 0$ the following inequalities hold*

$$(2.1) \quad \|a^{\frac{1}{2}} Du\|_p \leq \varepsilon^3 \|a^2 D^4u\|_p + \frac{C}{\varepsilon} \|u\|_p$$

$$(2.2) \quad \|aD^2u\|_p \leq \varepsilon^2 \|a^2 D^4u\|_p + \frac{C}{\varepsilon^2} \|u\|_p$$

$$(2.3) \quad \|a^{\frac{3}{2}}D^3u\|_p \leq \varepsilon \|a^2D^4u\|_p + \frac{C}{\varepsilon^3} \|u\|_p$$

$$(2.4) \quad \|a^2D^4u\|_p \leq C (\|Lu\|_p + \|u\|_p)$$

for every $u \in C_c^\infty(\mathbb{R}^N)$.

Proof. By [7, Lemma 2.4] the inequalities (2.1), (2.2), (2.3) and

$$(2.5) \quad \|a^2D^4u\|_p \leq C (\|Au\|_p + \|u\|_p)$$

hold with $C = C(N, p, \kappa)$. Then, we have

$$\|a^2D^4u\|_p \leq C (\|Lu\|_p + \|Bu\|_p + \|u\|_p).$$

On the other hand, since

$$-Bu = \sum_{i,j,k=1}^N b_{ijk}D_{ijk}u + \sum_{i,j=1}^N c_{ij}D_{ij}u + \sum_{i=1}^N d_iD_iu + eu$$

we have

$$\begin{aligned} \|Bu\|_p &\leq C \left(\sum_{i,j,k=1}^N \|b_{ijk}D_{ijk}u\|_p + \sum_{i,j=1}^N \|c_{ij}D_{ij}u\|_p + \sum_{i=1}^N \|d_iD_iu\|_p + \|eu\|_p \right) \\ &\leq C \left(\kappa \|a^{\frac{3}{2}}D^3u\|_p + \kappa^2 \|aD^2u\|_p + \kappa^3 \|a^{\frac{1}{2}}Du\|_p + \kappa^4 \|u\|_p \right). \end{aligned}$$

Then for every $\varepsilon > 0$ we have

$$\|Bu\|_p \leq \varepsilon \|a^2D^4u\|_p + C\kappa^4 \|u\|_p.$$

By (2.5) and the last estimate,

$$\|a^2D^4u\|_p \leq C (\|Au\|_p + \|u\|_p) \leq C (\|Lu\|_p + \varepsilon \|a^2D^4u\|_p + \|u\|_p)$$

and the claimed inequality (2.4) follows by choosing ε small enough. \square

By the previous Lemma we deduce that B is a small perturbation of A and then $L = A+B$ generate an analytic semigroup.

Theorem 2. *Let $1 < p < \infty$ and assume that a, b, c, d, e satisfy (H_p) . Then $(L, D(A_p))$ generates an analytic C_0 -semigroup in $L^p(\mathbb{R}^N)$. Moreover, the analyticity sector depends on N, p and κ . In particular, there exists $r_p > 0$ such that the equation $\lambda u - Lu = f$ is uniquely solvable in $D(A_p)$ for $f \in L^p(\mathbb{R}^N)$ and $\operatorname{Re} \lambda > r_p$.*

Proof. Let us endow B with its maximal domain

$$D(B) = \{u \in W_{\text{loc}}^{3,p}(\mathbb{R}^N) \cap L^p(\mathbb{R}^N) \mid Bu \in L^p(\mathbb{R}^N)\}.$$

We prove that B is relatively A -bounded with A -bound 0. First we observe that $D(A_p) \subset D(B)$. Indeed, let $u \in D(A_p)$, we have

$$\|Bu\|_p \leq C \left(\|a^{\frac{3}{2}}D^3u\|_p + \|aD^2u\|_p + \|a^{\frac{1}{2}}Du\|_p + \|u\|_p \right) = C \|u\|_{D(A_p)}.$$

Now, let $u \in D(A_p)$ and, by Lemma 3, let $u_n \in C_c^\infty(\mathbb{R}^N)$ be a sequence converging to u with respect to the norm $\|\cdot\|_{D(A_p)}$. By Lemma 2, for every $\varepsilon > 0$ there exists $C = C(N, p, \kappa)$ such that

$$\begin{aligned} \|Bu_n\|_p &\leq C \left(\|a^{\frac{3}{2}}D^3u_n\|_p + \|aD^2u_n\|_p + \|a^{\frac{1}{2}}Du_n\|_p + \|u_n\|_p \right) \\ &\leq \varepsilon \|a^2D^4u_n\| + C\|u_n\|_p \leq \varepsilon C_1 \|Au_n\|_p + C_2 \|u_n\|_p. \end{aligned}$$

Taking the limit as $n \rightarrow \infty$ and taking into account that $Bu_n \rightarrow Bu$ and $Au_n \rightarrow Au$ we have that for every $\varepsilon > 0$ there exists $C > 0$ such that

$$\|Bu\|_p \leq \varepsilon \|Au\|_p + C\|u\|_p.$$

By [5, Theorem III.2.10] and since $C = C(N, p, \kappa)$ we have the desired result. \square

As a consequence we obtain the following result.

Corollary 1. *Let $1 < p < \infty$ and assume that a, b, c, d, e satisfy (H_p) . Then, there exist $r_p = r_p(N, \kappa) > 0$ and $C = C(N, p, \kappa)$ such that for every $\operatorname{Re} \lambda > r_p$ and $u \in D(A_p)$*

$$|\lambda| \|u\|_p + |\lambda|^{\frac{3}{4}} \|a^{\frac{1}{2}}Du\|_p + |\lambda|^{\frac{1}{2}} \|aD^2u\|_p + |\lambda|^{\frac{1}{4}} \|a^{\frac{3}{2}}D^3u\|_p + \|a^2D^4u\|_p \leq C \|\lambda u - Lu\|_p.$$

Proof. By Theorem 2, L generates an analytic semigroup and there exists a sector Σ depending on N, p and κ such that

$$|\lambda| \|u\|_p \leq C \|\lambda u - Lu\|_p$$

for some positive constant C and for every $\lambda \in \Sigma$. The thesis now follows by Lemma 2. Indeed, for $h = 1, 2, 3$

$$|\lambda|^{\frac{4-h}{4}} \|a^{\frac{h}{2}}D^h u\|_p \leq |\lambda|^{\frac{4-h}{4}} \varepsilon^{4-h} \|a^2D^4u\|_p + |\lambda|^{\frac{4-h}{4}} \frac{C}{\varepsilon^h} \|u\|_p.$$

Setting $\frac{|\lambda|^{\frac{4-h}{4}}}{\varepsilon^h} = \frac{|\lambda|}{(\varepsilon')^h}$ we have

$$\begin{aligned} |\lambda|^{\frac{4-h}{4}} \|a^{\frac{h}{2}}D^h u\|_p &\leq (\varepsilon')^{4-h} \|a^2D^4u\|_p + \frac{C}{(\varepsilon')^h} |\lambda| \|u\|_p \\ &\leq (\varepsilon')^{4-h} (\|Lu\|_p + \|u\|_p) + C |\lambda| \|u\|_p \\ &\leq C \|\lambda u - Lu\|_p \end{aligned}$$

for $|\lambda|$ large enough. \square

At this point, using Corollary 1 and arguing as in [7, Section 4] we can prove the generation results in $C_b(\mathbb{R}^N)$ and $L^\infty(\mathbb{R}^N)$ for L . As for the operator A , we should assume more restrictive assumptions on a .

Let us define

$$\begin{aligned} &D(L_0) \\ &= \left\{ u \in \bigcap_{1 \leq p < \infty} W_{\text{loc}}^{4,p}(\mathbb{R}^N) \cap C_b(\mathbb{R}^N) \mid a^{\frac{h}{2}}D^h u, a^2\Delta^2 u \in C_b(\mathbb{R}^N), h = 0, 1, 2, 3 \right\}. \end{aligned}$$

Theorem 3. *Assume that a, b, c, d, e satisfy (H_p) and, in addition, $a(x) \geq \nu > 0$. Then $(L, D(L_0))$ generates an analytic C_0 -semigroup in $C_b(\mathbb{R}^N)$. Moreover, the analyticity sector depends on N, p and κ .*

3. GENERATION RESULTS IN L^1

For simplicity, here we consider operators containing only the higher order derivatives, that is $A = -a(x)^2 \Delta^2$. We require the assumptions:

$$(H_1) \quad \begin{aligned} a &\in C^4(\mathbb{R}^N), \quad a(x) \geq \nu > 0 \\ |D_{ijkl}a(x)| &\leq \kappa^4 a^{-1}(x) \text{ for all } i, j, k, l = 1, \dots, N, \\ |D_{ijk}a(x)| &\leq \kappa^3 a^{-\frac{1}{2}}(x) \text{ for all } i, j, k = 1, \dots, N, \\ |D_{ij}a(x)| &\leq \kappa^2 \text{ for all } i, j = 1, \dots, N, \\ |D_i a(x)| &\leq \kappa a^{\frac{1}{2}}(x) \text{ for all } i = 1, \dots, N, \end{aligned}$$

for all $x \in \mathbb{R}^N$ and some positive constants κ, ν . We will deal with the formal adjoint operator

$$-\hat{A}u = a^2 \Delta^2 u + 4Da^2 D\Delta u + 2\Delta a^2 \Delta u + 4tr D^2 a^2 D^2 u + 4D\Delta a^2 Du + \Delta^2 a^2 u.$$

Observe that conditions (H_1) on A insure that Theorem 3 applies and \hat{A} , endowed with the domain described before, generates an analytic semigroup in $C_b(\mathbb{R}^N)$.

Remark 1. *Observe that, if $0 \leq \alpha \leq 2$, the function $a(x) = (1 + |x|^2)^{\frac{\alpha}{2}}$ satisfies (H_1) .*

We consider the operator $A = -a(x)^2 \Delta^2$ endowed with the domain

$$D(A_1) = \{u \in L^1(\mathbb{R}^N) : a^{\frac{1}{2}} Du, aD^2 u, a^{\frac{3}{2}} D^3 u, a^2 \Delta^2 u \in L^1(\mathbb{R}^N)\}.$$

By [12, Theorem 5.8 (i)], $D(A_1)$ continuously embeds into $W^{3,p}(\mathbb{R}^N)$ for every $p \in \left[1, \frac{N}{N-1}\right]$ and, for every $u \in D(A_1)$,

$$\|u\|_{W^{3,p}} \leq C_p (\|u\|_1 + \|\Delta^2 u\|_1) \leq C_p \max\{1, \nu^{-2}\} (\|u\|_1 + \|a^2 \Delta^2 u\|_1).$$

3.1. A-priori estimates and domain characterization. We need a density result, useful to work with smooth functions. The proof follows as in [7, Lemma 2.2].

Lemma 3. *Assume that a satisfies (H_1) . Then $C_c^\infty(\mathbb{R}^N)$ is dense in $D(A_1)$ with respect to the norm*

$$\|u\|_{D(A_1)} = \|u\|_1 + \|a^{\frac{1}{2}} Du\|_1 + \|aD^2 u\|_1 + \|a^{\frac{3}{2}} D^3 u\|_1 + \|a^2 \Delta^2 u\|_1.$$

In order to prove some a-priori estimates, we need the following covering result with variable radius balls (see [3] for its proof).

Proposition 1. [3, Proposition 6.1] *Let $\mathcal{F} = \{B_{\rho(x)}(x)\}_{x \in \mathbb{R}^N}$, where $\rho : \mathbb{R}^N \rightarrow \mathbb{R}_+$ is a Lipschitz continuous function with Lipschitz constant κ strictly less than $\frac{1}{2}$. Then, there exist a countable subcovering $\{B_{\rho(x_n)}(x_n)\}_{n \in \mathbb{N}}$ and a natural number $\xi = \xi(N, \kappa)$ such that at most ξ among the doubled balls $\{B_{2\rho(x_n)}(x_n)\}_{n \in \mathbb{N}}$ overlap.*

Observe that by the assumption $|Da(x)| \leq \kappa a(x)^{\frac{1}{2}}$, the function $\rho(x) = \frac{\eta}{2\kappa} a(x)^{\frac{1}{2}}$, $0 < \eta \leq 1$, satisfies the assumptions of the previous covering Proposition with Lipschitz constant smaller than $\frac{\eta}{4}$.

We show some preliminary interpolative inequalities for the bi-harmonic operator.

Lemma 4. *There exists $\bar{\varepsilon}$ such that for every $0 < \varepsilon < \bar{\varepsilon}$, $u \in C_c^\infty(\mathbb{R}^N)$,*

$$(3.1) \quad \|Du\|_1 \leq \varepsilon^3 \|\Delta u\|_1 + \frac{C}{\varepsilon} \|u\|_1$$

and

$$(3.2) \quad \|D^3u\|_1 \leq \varepsilon \|\Delta^2 u\|_1 + \frac{C}{\varepsilon^3} \|u\|_1.$$

Proof. In the case $N = 1$ the inequalities (3.1) and (3.2) follow by straightforward computations using Taylor's formula with integral remainder.

Let now $N > 1$ and $\lambda > \bar{\lambda} > 0$. By setting $m = 4$ in [12, Theorem 5.7], the fundamental solution $K_\lambda(x, y)$ of the operator $\lambda + \Delta^2$ satisfies

$$|D_x^3 K_\lambda(x, y)| \leq C e^{-|\lambda|^{\frac{1}{4}}|x-y|} |x-y|^{1-N}.$$

It follows that for every $u \in C_c^\infty(\mathbb{R}^N)$

$$\|D^3u\|_1 \leq C |\lambda|^{-\frac{1}{4}} \|\lambda u + \Delta^2 u\|_1 \leq C \left(|\lambda|^{\frac{3}{4}} \|u\|_1 + |\lambda|^{-\frac{1}{4}} \|\Delta^2 u\|_1 \right).$$

Then, setting $|\lambda| = \varepsilon^{-4}$, we get (3.2).

The inequality (3.1) in the case $N > 1$ follows arguing as before setting $m = 2$ in [12, Theorem 5.7]. \square

The next lemma constitutes a first step toward the a-priori estimates.

Lemma 5. *Assume a satisfies (H_1) . Then, there exists a positive constant $C = C(N, \kappa)$ such that for every $\varepsilon > 0$ small enough the following inequalities hold*

$$(3.3) \quad \|a^{\frac{h}{2}} D^h u\|_1 \leq \varepsilon^{4-h} \|Au\|_1 + \frac{C}{\varepsilon^h} \|u\|_1$$

for $h = 1, 2, 3$ and for every $u \in C_c^\infty(\mathbb{R}^N)$.

Proof. We show that for $v \in C_c^\infty(\mathbb{R}^N)$

$$(3.4) \quad \|a^{\frac{h}{2}} D_i v\|_1 \leq \varepsilon \|a^{\frac{h+1}{2}} \Delta v\|_1 + \frac{C}{\varepsilon} \|a^{\frac{h-1}{2}} v\|_1$$

holds for $h \in \mathbb{N}$.

Let us set $\rho(x) = \frac{1}{2\kappa} a(x)^{\frac{1}{2}}$ and note that, as stated before, by (H_1) it follows that $\rho(x)$ satisfies the assumptions of Proposition 1 with Lipschitz constant smaller than $\frac{1}{4}$. We fix $x_0 \in \mathbb{R}^N$ and set $R = \rho(x_0) = \frac{1}{2\kappa} a^{\frac{1}{2}}(x_0)$. Then, if $x \in B_{2R}(x_0)$ the following inequalities hold

$$\frac{1}{2} a(x_0)^{\frac{1}{2}} \leq a(x)^{\frac{1}{2}} \leq \frac{3}{2} a(x_0)^{\frac{1}{2}}$$

and consequently

$$\left(\frac{1}{2}\right)^k a(x_0)^{\frac{k}{2}} \leq a(x)^{\frac{k}{2}} \leq \left(\frac{3}{2}\right)^k a(x_0)^{\frac{k}{2}}$$

for $k = h - 1, h, h + 1$.

Let now $\vartheta \in C_c^\infty(\mathbb{R}^N)$ be such that $0 \leq \vartheta \leq 1$, $\vartheta(x) = 1$ for $x \in B_R(x_0)$, $\vartheta(x) = 0$ for $x \in B_{2R}^c(x_0)$ and $|D^h \vartheta| \leq CR^{-h}$ for $h = 1, 2, 3, 4$.

By (3.1) we have

$$\begin{aligned} \|a^{\frac{h}{2}} Dv\|_{1,R} &\leq C \|a(x_0)^{\frac{h}{2}} Dv\|_{1,R} \leq C \|a(x_0)^{\frac{h}{2}} D(\vartheta v)\|_1 \\ &\leq \varepsilon \|a(x_0)^{\frac{h+1}{2}} \Delta(\vartheta v)\|_1 + \frac{C}{\varepsilon} \|a(x_0)^{\frac{h-1}{2}} \vartheta v\|_1 \\ &\leq \varepsilon \|a(x_0)^{\frac{h+1}{2}} \Delta v\|_{1,2R} + \frac{C}{R} \varepsilon \|a(x_0)^{\frac{h+1}{2}} Dv\|_{1,2R} + \frac{C}{R^2} \varepsilon \|a(x_0)^{\frac{h+1}{2}} v\|_{1,2R} \\ &\quad + \frac{C}{\varepsilon} \|a(x_0)^{\frac{h-1}{2}} v\|_{1,2R}. \end{aligned}$$

Then taking into account that $\frac{1}{R} a(x_0)^{\frac{h+1}{2}} = 2\kappa a(x_0)^{\frac{h}{2}} \leq C a(x)^{\frac{h}{2}}$ and $\frac{1}{R^2} a(x_0)^{\frac{h+1}{2}} = 4\kappa^2 a(x_0)^{\frac{h-1}{2}} \leq C a(x)^{\frac{h-1}{2}}$ in $B_{2R}(x_0)$ one has

$$(3.5) \quad \|a^{\frac{h}{2}} Dv\|_{1,R} \leq C \left(\varepsilon \|a^{\frac{h+1}{2}} \Delta v\|_{1,2R} + \varepsilon \|a^{\frac{h}{2}} Dv\|_{1,2R} + \left(\varepsilon + \frac{1}{\varepsilon} \right) \|a^{\frac{h-1}{2}} v\|_{1,2R} \right).$$

Let $\{B_{\rho(x_n)}(x_n)\}$ be a countable covering of \mathbb{R}^N as in Proposition 1 such that at most ζ among the double balls $\{B_{2\rho(x_n)}(x_n)\}$ overlap.

We write (3.5) with x_0 replaced by x_n and sum over n . Taking into account the above covering result, we get

$$\|a^{\frac{h}{2}} Dv\|_1 \leq C \left(\varepsilon \|a^{\frac{h+1}{2}} \Delta v\|_1 + \varepsilon \|a^{\frac{h}{2}} Dv\|_1 + \left(1 + \frac{1}{\varepsilon} \right) \|a^{\frac{h-1}{2}} v\|_1 \right).$$

Choosing $\varepsilon' = \frac{C\varepsilon}{1 - C\varepsilon}$ one can find a suitable positive C such that

$$\|a^{\frac{h}{2}} Dv\|_1 \leq \varepsilon' \|a^{\frac{h+1}{2}} \Delta v\|_1 + \frac{C}{\varepsilon'} \|a^{\frac{h-1}{2}} v\|_1$$

from which (3.4) follows.

Now for every $i, j = 1, \dots, N$ we have

$$\begin{aligned} \|aD_{ij}u\|_1 &\leq \varepsilon_1 \|a^{\frac{3}{2}} \Delta D_j u\|_1 + \frac{C}{\varepsilon_1} \|a^{\frac{1}{2}} D_j u\|_1 \\ &\leq \varepsilon_1 \varepsilon_2 \|a^2 \Delta^2 u\|_1 + C \frac{\varepsilon_1}{\varepsilon_2} \|a \Delta u\|_1 + C \frac{\varepsilon_3}{\varepsilon_1} \|a \Delta u\|_1 + C^2 \frac{1}{\varepsilon_1 \varepsilon_3} \|u\|_1 \\ &\leq \varepsilon_1 \varepsilon_2 \|a^2 \Delta^2 u\|_1 + C \left(\frac{\varepsilon_1}{\varepsilon_2} + \frac{\varepsilon_3}{\varepsilon_1} \right) \|a D^2 u\|_1 + C \frac{1}{\varepsilon_1 \varepsilon_3} \|u\|_1 \end{aligned}$$

and then

$$\|a D^2 u\|_1 \leq C \left(\varepsilon_1 \varepsilon_2 \|a^2 \Delta^2 u\|_1 + \left(\frac{\varepsilon_1}{\varepsilon_2} + \frac{\varepsilon_3}{\varepsilon_1} \right) \|a D^2 u\|_1 + \frac{1}{\varepsilon_1 \varepsilon_3} \|u\|_1 \right).$$

Setting $\varepsilon_2 = 4C\varepsilon_1$, $\varepsilon_3 = \frac{\varepsilon_1}{4C}$ and $\varepsilon^2 = 8C^2\varepsilon_1^2$ we obtain (3.3) when $h = 2$.

Then, we deduce

$$\|a^{\frac{1}{2}}Du\|_1 \leq \varepsilon_1\|a\Delta u\|_1 + \frac{C}{\varepsilon_1}\|u\|_1 \leq \varepsilon_1\varepsilon_2^2\|Au\|_1 + C\frac{\varepsilon_1}{\varepsilon_2}\|u\|_1 + C\frac{1}{\varepsilon_1}\|u\|_1.$$

Setting $\varepsilon_1 = \varepsilon_2$ we obtain (3.3) when $h = 1$.

Arguing as done before in order to prove (3.4), applying inequality (3.2), one gets

$$\|a^{\frac{3}{2}}D^3u\|_1 \leq \varepsilon\|Au\|_1 + \frac{C}{\varepsilon_3}\|u\|_1.$$

□

The following proposition is an immediate consequence of the previous lemma and gives the announced a priori estimates.

Proposition 2. *Assume a satisfies (H_1) . Then, for $u \in D(A_1)$ the following inequality holds*

$$\|a^{\frac{1}{2}}Du\|_1 + \|aD^2u\|_1 + \|a^{\frac{3}{2}}D^3u\|_1 \leq C(\|Au\|_1 + \|u\|_1)$$

where the constant C depends on N and κ .

We are now able to characterise the maximal domain of A in $L^1(\mathbb{R}^N)$ arguing as in [7, Proposition 2.8].

Proposition 3. *Assume a satisfies (H_1) . Then*

$$D(A_1) = D(A_{1,\max}) = \{u \in L^1(\mathbb{R}^N) : Au \in L^1(\mathbb{R}^N)\}.$$

3.2. Sectoriality and solvability. To investigate the generation of an analytic semigroup in $L^1(\mathbb{R}^N)$, we start by proving an a-priori sectoriality estimate for the resolvent in a suitable sector of the complex plane. Then we prove the surjectivity.

Theorem 4. *Assume that a satisfies (H_1) . Then, there exist $\Sigma \subseteq \mathbb{C}$ such that, if $\lambda \in \Sigma$, the following inequality holds for every $u \in D(A_1)$*

$$|\lambda|\|u\|_1 \leq C\|\lambda u - Au\|_1$$

where $C = C(N, \kappa)$ and the sector Σ depends on κ .

Proof. We use a duality argument to get the sectoriality estimate in L^1 . Consider the formal adjoint

$$\hat{A}v = -\hat{A}u = a^2\Delta^2u + 4Da^2D\Delta u + 2\Delta a^2\Delta u + 4\text{tr}D^2a^2D^2u + 4D\Delta a^2Du + \Delta^2a^2u.$$

By the conditions (H_1) on A , the operator \hat{A} generates an analytic semigroup in $C_b(\mathbb{R}^N)$ (Theorem 3). Therefore there exists a sector Σ contained in the resolvent set of \hat{A} and

$$|\lambda|\|u\|_\infty \leq C\|\lambda u - Au\|_\infty$$

for every $\lambda \in \Sigma$, $u \in C_b(\mathbb{R}^N)$ and for some constant C independent of λ and u . In view of Lemma 3, we prove the sectoriality estimate for functions in $C_c^\infty(\mathbb{R}^N)$. Let $\lambda \in \Sigma$, $u \in C_c^\infty(\mathbb{R}^N)$. Then, since $(\lambda I + \hat{A})(D(\hat{A})) = C_b(\mathbb{R}^N)$, we have

$$\|u\|_1 \leq \sup \left\{ \int_{\mathbb{R}^N} u(\lambda I - \hat{A})v \, dx : v \in D(\hat{A}), \|\lambda v - \hat{A}v\|_\infty \leq 1 \right\}$$

$$\leq \sup \left\{ \int_{\mathbb{R}^N} u(\lambda I - \hat{A})v \, dx : v \in D(\hat{A}), \|v\|_\infty \leq \frac{C}{|\lambda|} \right\}.$$

Integrating by parts, we get

$$\left| \int_{\mathbb{R}^N} u(\lambda I - \hat{A})v \right| = \left| \int_{\mathbb{R}^N} v(\lambda I - A)u \right| \leq \frac{C}{|\lambda|} \|\lambda u - Au\|_1.$$

□

The previous estimate leads, through an approximation procedure, to the solvability of the elliptic problem.

Theorem 5. *Assume that a satisfies (H_1) . Then there exists $\Sigma \subseteq \mathbb{C}$ such that the equation $\lambda u - Au = f$ is uniquely solvable in $D(A_1)$ for $f \in L^1(\mathbb{R}^N)$ and $\lambda \in \Sigma$ and the resolvent estimate*

$$|\lambda| \|u\|_1 \leq C \|\lambda u - Au\|_1$$

holds.

Proof. Let $\sigma > 0$ and define the function

$$a_\sigma(x) = \frac{a(x)}{1 + \sigma a(x)}, \quad x \in \mathbb{R}^N.$$

Observe that $a_\sigma(x)$ satisfies (H_1) with κ independent of σ . Consider the operators with bounded coefficients $A_\sigma = -(a_\sigma(x))^2 \Delta^2$. By [12, Section 5.4], there exist $\omega_\sigma \in \mathbb{R}$, $M_\sigma > 0$ such that $\{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \geq \omega_\sigma\} \subset \rho(A_\sigma)$ and

$$|\lambda| \|u\|_1 \leq M_\sigma \|\lambda u - A_\sigma u\|_1$$

for every $u \in D(A_{\sigma,1,\max})$ and $\lambda \in \mathbb{C}$ with $\operatorname{Re} \lambda \geq \omega_\sigma$. Moreover, by applying Theorem 4 to A_σ , since the condition in (H_1) is satisfied with a constant independent of σ , we have that there exists $r_1 > 0$ (independent of σ) such that for every $\lambda \in \mathbb{C}$ with $\operatorname{Re} \lambda > r_1$ and for every $u \in D(A_{\sigma,1,\max})$

$$(3.6) \quad |\lambda| \|u\|_1 \leq C_1 \|\lambda u - A_\sigma u\|_1$$

for some constant C_1 independent of σ . Let $\bar{\lambda} \in \rho(A_\sigma)$ and $\operatorname{Re} \bar{\lambda} > r_1$, then the inequality (3.6) gives that

$$(3.7) \quad \|R(\bar{\lambda}, A_\sigma)\|_1 \leq \frac{C_1}{|\bar{\lambda}|} \leq \frac{C_1}{r_1}.$$

By classical result, if $|\lambda - \bar{\lambda}| < \frac{1}{\|R(\bar{\lambda}, A_\sigma)\|_1}$, then $\lambda \in \rho(A_\sigma)$. By (3.7), if $|\lambda - \bar{\lambda}| < \frac{r_1}{C_1}$, then $\lambda \in \rho(A_\sigma)$. By covering with balls of radius $\frac{r_1}{C_1}$, it follows that, if $\operatorname{Re} \lambda > r_1$, then $\operatorname{Re} \lambda \in \rho(A_\sigma)$. In particular, $\lambda = \bar{\lambda} - \frac{\bar{\lambda}}{2C_1}$ satisfies $|\lambda - \bar{\lambda}| < \frac{|\bar{\lambda}|}{C_1}$ and then belongs to $\rho(A_\sigma)$.

Let $\operatorname{Re} \lambda > r_1$, $f \in L^1(\mathbb{R}^N)$. Denote by $u_\sigma \in D(A_{\sigma,1,\max})$ the unique solution of $\lambda u_\sigma - A_\sigma u_\sigma = f$. Then by [12, Theorem 5.8 (i)], $D(A_{\sigma,1,\max})$ continuously embeds into $W^{3,p}(\mathbb{R}^N)$ for every $p \in \left[1, \frac{N}{N-1}\right]$ and, observing that $a_\sigma \geq \frac{\nu}{\nu+1}$, for every $u \in D(A_{\sigma,1,\max})$,

$$(3.8) \quad \|u_\sigma\|_{W^{3,p}} \leq C_p (\|u_\sigma\|_1 + \|\Delta^2 u_\sigma\|_1) \leq C_p(\nu) (\|u_\sigma\|_1 + \|a_\sigma^2 \Delta^2 u_\sigma\|_1) \leq C_p(\nu, r_1) \|f\|_1.$$

From (3.8) we deduce that, up to considering a subsequence, we can assume that u_σ converges weakly in $W^{3,p}(\mathbb{R}^N)$ to some function $u \in W^{3,p}(\mathbb{R}^N)$ as σ goes to 0 and strongly in $L^p_{\text{loc}}(\mathbb{R}^N)$. In particular, u_σ converges to u in $L^1_{\text{loc}}(\mathbb{R}^N)$. From (3.6), we deduce that $u \in L^1(\mathbb{R}^N)$ and

$$|\lambda||u|_1 \leq C_1 \|f\|_1.$$

It remains to show that u belongs to $D(A_1)$ and solves the equation $\lambda u - Au = f$. Fix $v \in C_c^4(\mathbb{R}^N)$ and observe that

$$(3.9) \quad \int_{\mathbb{R}^N} a_\sigma^2 \Delta^2 u_\sigma v \, dx = \int_{\mathbb{R}^N} u_\sigma \Delta^2 (a_\sigma^2 v) \, dx.$$

By easy computations, $\Delta^2(a_\sigma^2 v)$ converges uniformly in \mathbb{R}^N to $\Delta^2(a^2 v)$. Letting σ to 0 in $\lambda u_\sigma - A_\sigma u_\sigma = f$, we get $A_\sigma u_\sigma \rightarrow \lambda u - f$ and in the equation (3.9) we obtain

$$\int_{\mathbb{R}^N} \frac{\lambda u - f}{a^2} a^2 v \, dx = \int_{\mathbb{R}^N} u \Delta^2 (a^2 v) \, dx.$$

It follows that

$$\int_{\mathbb{R}^N} \frac{\lambda u - f}{a^2} w \, dx = \int_{\mathbb{R}^N} u \Delta^2 w \, dx$$

for every $w \in C_c^4(\mathbb{R}^N)$. This means that $a^2 \Delta^2 u \in L^1(\mathbb{R}^N)$ and u satisfies $\Delta^2 u = \frac{\lambda u - f}{a^2}$ in the distributional sense. \square

In view of [9, Proposition 3.2.8] the operator A is sectorial. Therefore, by standard generation results, cf. [5, Ch. II, Theorem 4.6], we get the following generation theorem.

Theorem 6. *Assume that a satisfies (H_1) . Then, $(A, D(A_1))$ generates an analytic C_0 -semigroup in $L^1(\mathbb{R}^N)$.*

REFERENCES

1. D. Addona, F. Gregorio, A. Rhandi, and C. Tacelli, *Bi-Kolmogorov type operators and weighted Rellich's inequalities*, Nonlinear Differ. Equ. Appl. **29** (2022), no. 2, Paper No. 13, 37 pp.
2. S.E. Boutiah, L. Caso, F. Gregorio, and C. Tacelli, *Some results on second-order elliptic operators with polynomially growing coefficients in L^p -spaces*, J. Math. Anal. Appl. **501** (2021), 21 pp.
3. G. Cupini and S. Fornaro, *Maximal regularity in $L^p(\mathbb{R}^N)$ for a class of elliptic operators with unbounded coefficients*, Diff. Int. Eqs. **17** (2004), 259–296.
4. E. B. Davies, *Uniformly elliptic operators with measurable coefficients*, J. Funct. Anal. **132** (1995), 141–169.
5. K.J. Engel and R. Nagel, *One parameter semigroups for linear evolution equations*, Springer-Verlag, Berlin, 2000.
6. S. Fornaro and L. Lorenzi, *Generation results for elliptic operators with unbounded diffusion coefficients in L^p and C_b -spaces*, Discr. Cont. Dyn. Syst. A **18** (2007), 747–772.
7. F. Gregorio, C. Spina, and C. Tacelli, *Fourth-order operators with unbounded coefficients*, Communications on Pure & Applied Analysis **23** (2024), 463–479.
8. F. Gregorio and C. Tacelli, *Fourth-order Schrödinger type operator with unbounded coefficients in $L^2(\mathbb{R}^N)$* , Math Meth Appl Sci. **45** (2022), 11246–11261.
9. L. Lorenzi and A. Rhandi, *Semigroups of bounded operators and Second-order elliptic and parabolic partial differential equations*, Monogr. Res. Notes Math. CRC Press, Boca Raton, FL, 2021.
10. G. Metafume, N. Okazawa, M. Sobajima, and C. Spina, *Scale invariant elliptic operators with singular coefficients*, J. Evol. Equ. **16** (2016), 391–439.

11. G. Metafune and C. Spina, *Elliptic operators with unbounded diffusion coefficients in L^p spaces*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **11** (2012), no. 2, 303–340.
12. H. Tanabe, *Functional analytic methods for partial differential equations*, M. Dekker, 1997.
Email address: fgregorio@unisa.it
Email address: chiara.spina@unisalento.it
Email address: ctacelli@unisa.it