

# Coupling Fluid Plasma and Kinetic Neutral Models using Correlated Monte Carlo Methods

G.J. Parker, M.V. Umansky, B.D. Dudson

*Stanford University, Department of Mathematics, Stanford CA 94305, USA*

*Lawrence Livermore National Lab, Livermore, CA 94550, USA*

---

## Abstract

While boundary plasmas in present day tokamaks generally fall in a fluid regime, neutral species near the boundary often require kinetic models due to long mean-free-paths compared to characteristic spatial scales in the region. Monte-Carlo (MC) methods provide a complete, high-fidelity approach to solving kinetic models, and must be coupled to fluid plasma models to simulate the full plasma-neutrals system. The statistical nature of MC methods, however, prevents convergence of coupled fluid-kinetic simulations to an exact self-consistent steady-state. Moreover, this forces the use of explicit methods that can suffer from numerical errors and require huge computational resources.

Correlated Monte-Carlo (CMC) methods are expected to alleviate these issues, but have historically enjoyed only mixed success. Here, a fully implicit method for coupled plasma-neutral systems is demonstrated in 1D using the UEDGE plasma code and a homemade CMC code. In particular, it is shown that ensuring the CMC method is a differentiable function of the background plasma is sufficient to employ a Jacobian-Free Newton-Krylov solver for implicit time steps. The convergence of the implicit coupling method is explored and compared with explicit coupling and uncorrelated methods. It is shown that ensuring differentiability by controlling random seeds in the MC is sufficient to achieve convergence, and that the use of implicit time-stepping methods has the potential for improved stability and runtimes over explicit coupling methods.

*Keywords:* fusion, boundary plasma, numerical algorithms, Monte-Carlo methods

---

## 1. Introduction

High fidelity modeling for coupled systems of plasma and neutral species is essential in the design and study of divertor regions in fusion devices, where strong interactions occur [1, 2]. Fluid models provide accurate approximations for the plasma behavior, and robust approaches using finite-volume methods have now been employed with great success for several decades [3]. In cases where the mean-free paths of neutral particles in edge plasmas are long, neutral species exhibit non fluid-like behavior and are more accurately modeled using the kinetic Boltzman equation [4, 5]. Monte-Carlo (MC) methods [1, 6, 7, 8] are by far the most developed among the several known approaches [9, 10] to solving the kinetic equation.

The stochastic nature of Monte-Carlo methods however, results in unfavorable convergence properties when coupling to fluid plasma models. Simulations typically converge to stochastic equilibria, for which robust convergence criteria may be difficult to determine, and which are known to exhibit bias compared to true solutions [1, 11, 12]. More importantly, the stochastic nature of Monte-Carlo methods typically forces the use of explicit coupling methods. By standard numerical stability theory [13, 14], such explicit methods can result in large numerical errors

and require excessively small timesteps to achieve desired accuracy. As a result, resolving long timescale (1s) effects to necessary accuracies using explicit methods can require prohibitively long runtimes or heavy computational resources. Implicit methods enjoy much more desirable stability properties and, in theory, have the potential to be more accurate by an exponential factor for the same size timestep, but are impossible when coupling to standard Monte-Carlo methods.

It has been known to experts for many years that correlated Monte-Carlo (CMC) methods have the potential to address these problems, but attempts to realize this have had only mixed success. To the authors' knowledge, correlated methods have only been successfully demonstrated in one spatial dimension, and only using explicit coupling for which the advantages of correlation were shown to be negligible [11]. The purpose of this study is to demonstrate a fully implicit coupling scheme in one spatial dimension with robust stability and convergence properties employing correlated methods.

Section 2 explains the coupled model studied in detail, and reviews the difference between explicit and implicit coupling methods. Section 3 offers a new perspective on Correlated Monte-Carlo methods that is useful in the development of implicit coupling algorithms. Section 4 in-

50 introduces the UEDGE plasma code, a homemade Monte-Carlo code using the JULIA programming language and the methods of coupling. Section 5 displays the results of coupled simulations, and Sections 6 and 7 provide the discussion and conclusion respectively.

## 2. Numerical Methods

The model investigated couples a set of fluid-plasma evolution equations to a chosen equation modeling neutral species with two basic interaction: (i) ionization, and (ii) charge exchange. Let  $p(x)$  and  $n(x)$  denote the plasma and neutral state vectors, so that e.g.  $p = (n_i, n_e, T_i, T_e, u_i)$  is a vector-valued function combining the ion and electron densities, temperatures, and velocities. The coupled model leads to a system of equations with the schematic form

$$\partial_t p + P(p) = S_1(p, n) \quad (1)$$

$$N(n) = S_2(p) \quad (2)$$

wherein  $P, N$  are the plasma and neutral evolution equations respectively, and  $S_i$  comprise the interaction terms. We make the additional assumption that the neutral species' relaxation time is on a significantly faster timescale than the plasma timescale – in this situation it is only necessary to consider the time-independent version of  $N(n)$ . It is additionally assumed that recombination is negligible, and that the source of neutrals from the wall is independent of the neutral distribution, i.e.  $S_2$  is independent of  $n$ .

While we focus here on two specific plasma models encapsulated by a simple non-linear diffusion equation and the UEDGE model [15, 16], we emphasize that the techniques described are equally applicable to any chosen pairs of plasma and neutral models, or more generally to any chosen pair of continuum (e.g. finite volume) and particle-based (e.g. Monte Carlo) simulations. In particular, including additional physics effects such as recombination of ions or additional neutral species should constitute no fundamental change.

### 2.1. Explicit vs. Implicit Coupling

Suppose that known techniques may be applied to time evolve the plasma for a given fixed neutral distribution, and vice versa. That is, assume that one has known numerical algorithms implementing the time evolution operator  $U_{\Delta t}(p, S)$  for plasma with timestep  $\Delta t$  and fixed source term  $S$ , and  $N^{-1}$  the inverse operator of the time-independent neutral equation. The system (1–2) may then be coupled using either explicit or implicit methods. Note that this terminology refers to the method used to couple the system, while the individual solvers for  $U_{\Delta t}, N^{-1}$  may themselves include an internal choice of an explicit or implicit method.

#### 2.1.1. Explicit Coupling

Given a coupled state  $(p_j, n_j)$  at time  $t_j = j\Delta t$ , an explicit method advances the state by setting

$$p_{j+1} = U_{\Delta t}(p_j, S_1(p_j, n_j)) \quad (3)$$

$$n_{j+1} = N^{-1}(S_2(p_{j+1})). \quad (4)$$

The method is explicit in the sense that the states at timestep  $j + 1$  are defined entirely using quantities calculated in the previous steps. The above demonstrates a standard first-order splitting method, but higher order or Strang splitting methods would be equally applicable. The above time-step is iterated until a sufficient equilibrium is reached.

Explicit methods have several key advantages and disadvantages. Advantages include that they are relatively easy to implement and parallelize, and can be implemented treating the existing individual solvers or codebases for  $U_{\Delta t}, N^{-1}$  as “black boxes”. The one main disadvantage is that explicit coupling methods display the same quite severe numerical instability as explicit time evolution for a single equation: small errors may compound exponentially as they propagate forward in time. To guarantee a desired accuracy on a time interval  $[0, T]$ , the time-step must be taken to be of size  $O(T^{-1}e^{-\lambda T})$ , where  $\lambda$  is the largest eigenvalue of the (linearization of) the time-independent equation [13, Ch. 7]. The second disadvantage is that explicit coupling schemes rely on the assumption that the interaction terms are relatively weak, and so for strongly coupled systems convergence is not guaranteed. Despite these significant shortcomings, *explicit methods are currently the only method available for coupling kinetic models using Monte-Carlo methods*. Contrasting the above with implicit methods makes the reason for this clear.

#### 2.1.2. Implicit Coupling

A first-order implicit method advances the state by  $\Delta t$  by taking  $(p_{j+1}, n_{j+1})$  as the unique solution of the non-linear system:

$$(I + \Delta t P)p_{j+1} + \Delta t S_1(p_{j+1}, n_{j+1}) = p_j \quad (5)$$

$$n_{j+1} = N^{-1}(S_2(p_{j+1})), \quad (6)$$

where  $P$  now denotes the time-independent plasma equation. Substituting (6) into (5) results in an equation purely for  $p_{j+1}$  defining the evolution of the plasma. For  $\Delta t$  sufficiently small, the equation is well-conditioned and may be solved using a suitable implementation of Newton's method.

Implicit methods enjoy far better stability properties than explicit ones, and in practice the time-step may usually be taken as large as the Newton solver will allow without impacting accuracy. The seemingly innocuous disadvantage of implicit methods is that they require the functions  $N, P, S_i$  to be differentiable functions of the plasma and neutral states. This is the requirement that precludes the use of Monte-Carlo methods (which are *a priori* not

even constant for repeated evaluation at the same input). From a mathematical perspective, *the key result of this study is to demonstrate that correlated Monte Carlo methods can be made sufficiently differentiable as a function of parameters to apply Newton solvers*, and thus implicit methods.

### 3. A New View of Correlated MC

When the model  $N(n)$  for the neutral species is the kinetic Boltzman equation, Monte-Carlo methods are the most widely-used technique for finding the equilibrium distribution. The standard viewpoint of a Monte-Carlo method (see e.g. [6, 7, 8, 1]) is the following: particle testflights are run using (pseudo) random numbers to generate particle trajectories, and each testflight gives a sample from the equilibrium particle distribution. Thus with sufficiently<sup>210</sup> many flights, the tallied output converges to the solution of  $N(n) = S_2(p)$ .

A correlated Monte-Carlo method does the same, but fixes the random numbers generated for each individual testflight to make the output reproducible. The result is a sample from the same equilibrium distribution, and the distribution is now obtained by averaging over increasingly large numbers of possibilities for the fixed random seeds.

#### 3.1. Correlated MC Constructs the Inverse Operator

When considering a Monte-Carlo method as a particu-<sup>220</sup> lar step in a larger algorithm as in (5–6), it is advantageous adopt a more abstract viewpoint. Rather than working at the level of equations as explained above, let us consider the situation at the level of operators. Suppose that the neutral state is described by a vector  $n(x) \in \mathbb{R}^m$  on a discretized spatial grid of size  $m$ . Thus the Boltzman equation  $N_p(n) = N(n) - S_2(p)$  may be considered as a matrix  $N_p \in M(\mathbb{R}^m, \mathbb{R}^m)$  in the space of operators on the state space. In our case, this operator is linear, but the same framework applies equally well to a space of nonlinear op-<sup>230</sup> erators including neutral-neutral interactions.

The existence of a physical equilibrium distribution dictates that the equation should be always be solvable for reasonable plasma backgrounds  $p$ , i.e. that  $N_p^{-1} \in M(\mathbb{R}^m, \mathbb{R}^m)$  exists. One may combine all the random choices required for Monte-Carlo flights into a joint probability distribution  $\mathbb{X}$ . Then, at an operator level, a correlated Monte-Carlo method is a (parameterized) operator-valued probability distribution

$$P \times \mathbb{X} \longrightarrow M(\mathbb{R}^m, \mathbb{R}^m) \quad (7)$$

$$(p, \xi) \mapsto N_{p,\xi}^{-1}. \quad (8)$$

That associates to any pair of a background plasma state  $p$  and joint draw of pseudorandom numbers  $\xi$  a particular approximate inverse to  $N_p$ . The true inverse is then obtained as  $N_p^{-1} = \int N_{p,\xi}^{-1} d\mathbb{X}(\xi)$ , i.e. averaging over increasingly large samples of the joint probability distribution  $\mathbb{X}$  gives better approximations to the true inverse.

For algorithms that require multiple applications of the operator  $N_p^{-1}$  (e.g. explicit or implicit coupling), it can be advantageous to fix a particular choice and carry out the algorithm to completion, rather than generate a new inverse each time  $N_p^{-1}$  is called within the algorithm.

#### 3.1.1. Smooth Dependence of MC

The space  $M(\mathbb{R}^m, \mathbb{R}^m)$  where the approximate inverse operators are valued is a vector space (more generally a manifold for non-linear operators), as is the space of plasma parameters  $p$ . It therefore makes sense to ask whether, for fixed  $\xi \in \mathbb{X}$ , the operator inverse  $N_{p,\xi}^{-1}$  is a continuous, differentiable, or smooth function of the plasma state  $p$ . Note that without correlation, continuity certainly fails — even for fixed  $p$  each call to an uncorrelated MC method produces a different result. If differentiability is correctly imposed for fixed  $\xi$ , this function may be called repeatedly as a step in larger algorithms relying on the linearized approximation of a function, in particular various version of Newton’s method.

In practice, there are at least two challenges in making a MC output (i.e.  $N_{p,\xi}^{-1}$ ) depend differentiably on  $p$ . These include:

- (i) De-synchronization of trajectories
- (ii) Discreteness of estimators

If one naively fixes the random seeds, particle flights can de-synchronize for infinitesimal changes in background parameters. To elaborate, the cut-off determining whether a particular collision is a charge-exchange or ionization is a discrete transition; thus a tiny variation in background parameters causing a fixed random seed to land on the other side of this cut-off may cause the particle flight to proceed for more steps, thereby shifting the seeds for subsequent flights. Even if this shift is remedied, the estimator used to tally the output will change by a discrete transition when the particle flight elongates, which creates a point of discontinuity with respect to the parameters.

To fix the first of these issues, it is necessary to identify the joint probability distribution  $\mathbb{X}$  and have a reliable method of generating joint draws so that these shifts do not occur, as well as imposing continuity of the tallies. In practice, the former means pre-generating random numbers for all possible branchings of the particle trajectory before the flight begins. To fix the second, one must use more advanced estimators with better continuity properties. In this study, the attenuated absorption method [6] was used with a track length estimator [17, Eq. 2.35]. If one prefers a collisional estimator, then finite-size particle methods provide continuity properties. *In general, it is advantageous to identify any source of discreteness within the MC simulation and replace this by a continuous quantity.*

When smoothness is imposed correctly, the derivative

$$\frac{\partial N_{p,\xi}^{-1}}{\partial p} \approx \frac{N_{p+\Delta p,\xi}^{-1} - N_{p,\xi}^{-1}}{\Delta p}$$




Figure 1: The differentiability of the correlated Monte-Carlo method (blue) compared to the discontinuous uncorrelated method (orange). The derivative displayed is the partial derivative at a particular volume cell with respect to varying the plasma background  $p$  at equilibrium in the direction of a gaussian perturbation  $dp$ .

exists, and can be calculated through finite-difference approximation schemes as we use here, or potentially by auto-differentiation methods.

#### 4. Simulation Setup

The system of equations (1–2) is discretized on a 1-dimensional grid, assuming both rotational and poloidal symmetry. The grid spacing decreases toward the domain wall to better resolve edge effects, and upstream boundary conditions are imposed on the inner boundary.

##### 4.1. Simulation Codes

Two plasma models are implemented: first, a simplified model denoted  $P_0$  for troubleshooting. This model assumes constant velocity, and temperatures, and that the ion densities satisfies a second-order non-linear diffusion model. The second model, denoted  $P_1$  is given by the UEDGE code, described in detail in [15, 16]. The code implements finite-volume methods for a non-linear system of coupled fluid and transport equations in the variables  $p = (n_i, T_i, T_e, u_i)$  of ion density, temperature, and velocity, and electron temperature  $t_e$ .

The Correlated Monte-Carlo methods are implemented using a simplified, homemade code written in JULIA. This code includes 1-dimensional capabilities following large codebases such as DEGAS2 [6, 17]. Testflights are run using the attenuated absorption method, and tallied with a track length estimator.

##### 4.2. Coupling Methods

Although the UEDGE code includes a sophisticated and optimized JFNK solver for time-stepping, we here opt for a more accessible setup using an external solver. Thus the above codes are used simply to call their respective

functions  $P_i$  and  $N_{p,\xi}^{-1}$ . The coupling schemes (3–4) and (5–6) are implemented in a Python overhead using SciPy’s JFNK and finite-difference methods. The workflow of the implicit solver is depicted in Figure 2 below.




Figure 2: The code workflow for fully implicit coupling. The derivative of the Monte-Carlo method is called repeatedly to generate the Krylov subspace for the Jacobian during Newton iteration.

#### 4.3. Simulation Parameters

Simulations are run beginning from a uniform ion density of  $10^{19}$  particles/m<sup>2</sup>, with uniform starting temperatures 10 eV, and velocity  $10^4$  m/s toward the divertor. The a fixed value  $R$  is chosen for the wall recycling coefficient (generally  $R = 1$  is taken). Dirichlet boundary conditions are imposed on the upstream plasma boundary, and Robbin boundary conditions are imposed at the plate. Neutrals are assumed to have perfect reflection at walls.

### 5. Simulation results

#### 5.1. Correlated vs. Uncorrelated Explicit Methods

The convergence properties of the coupled system (1–2) was investigated using the explicit coupling scheme (3–4) for the UEDGE plasma model with both correlated and uncorrelated Monte-Carlo Methods. The rate of convergence is displayed in Figure 3 on a logarithmic scale as a function of time.

With uncorrelated MC, the coupled system reaches a stochastic equilibrium after only a few orders of magnitude of convergence (Figure 3a). The magnitude of the stochastic oscillations and the amount of convergence displayed is proportional to  $\sqrt{M}$  for  $M$  the number of flights. In contrast, the correlated method produces convergence to the machine floating-point precision (or set tolerance of the plasma step), regardless of the number of flights. For the uncorrelated method, true equilibria are obtained by averaging the stochastic oscillations, whereas for correlated methods by averaging over different seeds (Figure 3b). These averages are known to each also include a bias arising from the non-linearity of the plasma model, but this is a second-order (i.e.  $O(M)$ ) effect and becomes negligible for large particle numbers [11].




Figure 3: (Top) Comparison of residual difference from converged solution for correlated vs uncorrelated explicit coupling on a log scale as a function of time (Bottom) The average of the stochastic equilibrium at two selected volume elements compared to the converged correlated solution for a variety of fixed random seeds.

### 5.2. Explicit vs. Implicit Correlated Methods

To begin, implicit methods were tested using a simplified plasma model given described by the non-linear diffusion equation

$$\frac{\partial p}{\partial t} - \nabla(D(p)\nabla p) = 0$$

where  $D(p) = cp^n$  is the diffusion constant. The converged equilibria for the explicit and implicit coupling schemes (3–4) and (5–6) are compared below in Figure 4.

The implicit coupling method is found to obtain an equilibrium which is *independent* of the time-step used up to machine accuracy, while the explicit coupling method produces an equilibrium which depends on the time-step, and approaches that of the implicit method as  $dt \rightarrow 0$  (Figure 4a).

Figure 4b compares the runtimes of the two methods to attain a certain accuracy. The explicit coupling method is found to require exponentially more computation to reach the same levels of accuracy as the implicit method.

### 5.3. Implicit Coupling to UEDGE

The fully implicit coupling scheme was also tested for the UEDGE plasma model in one spatial dimension. Con-




Figure 4: (Top) Time evolution of plasma density at a single cell  $x = 3.8$  for implicit and explicit coupling with various timesteps  $dt$ . (Bottom) residual difference from converged solution as a function of machine runtime for both explicit and implicit methods plotted on log-log scale.

vergence to machine floating-point accuracy was achieved for a variety of time steps, up to approximately  $\sim 0.1$  ms. The results displayed in Figure 4 are reproducible using UEDGE, though the trends persist only for several orders of magnitude smaller than the starting residual, after which long time-scale effects in the UEDGE model dominate. The exploration of the capabilities of this fully implicit model using UEDGE will be the subject of future work.

## 6. Discussion

The use of correlated Monte-Carlo methods in coupled Plasma-neutral systems was investigated. Previous results [11] demonstrating convergence to machine precision were reproduced using the UEDGE plasma code and a homemade Monte-Carlo code.

Implicit methods are demonstrated to have significant advantage over explicit coupling schemes. From a naive perspective, it is not so clear that the implicit method should be advantageous; indeed, to do a single implicit time-step requires calling the Monte-Carlo code  $O(10^3)$  times to differentiate, generate the Krylov subspace, and

perform Newton iteration. In contrast each time-step of the explicit scheme requires only one invocation of the Monte-Carlo solver. The advantage of the implicit scheme appears when considering the number of calls to achieve a particular accuracy or a particular simulation time. Standard numerical theory shows explicit methods can result in numerical error proportional to  $dt \cdot Te^{\lambda T}$  where  $dt$  is the timestep,  $T$  is the time to convergence, and  $\lambda$  is the (maximum in  $t$  of the) largest eigenvalue of the Jacobian, requiring the number of timesteps to increase exponentially as a function of desired accuracy. In contrast, implicit methods have accuracy independent of stepsize provided stability conditions are met, allowing timesteps to be taken several orders of magnitude larger to model effects on longer timescales without significantly increasing runtime. Testing both coupling schemes confirms that the fully implicit coupling scheme indeed realizes these improvements, despite the apparent difficulty in performing a single time-step.

The use of fully implicit methods presents several avenues for further study. The robust stability and convergence properties of these methods make them ideal candidates for generalizing to study the convergence of two and three-dimensional coupled systems using correlated methods, which have eluded previous efforts. It is also an interesting prospect to explore whether the same differentiability properties of the Monte-Carlo output may be obtained using existing functions in established codebases such as DEGAS2 [8] and EIRENE [7]. Finally, there are many avenues for optimizing and improving the performance of the method. In particular, one interesting prospect is to investigate the use of auto-differentiation methods to generate the Jacobian of the Monte-Carlo output.

## 7. Conclusions

The methods investigated here display significant potential advantages over the current coupling schemes of fluid plasma and kinetic neutral models. First, a shift in viewpoint of Monte-Carlo methods is proposed, to one in which the code is considered as a “black box” producing an approximate inverse operator, which allows the use and analysis of more sophisticated (in particular implicit) coupling methods. Then, by imposing smoothness of the correlated Monte-Carlo output with respect to background plasma parameters, we show that the Monte-Carlo code may be differentiated using finite-difference schemes making it viable for using in solving algorithms based on Newton’s method, in particular implicit time steps using Krylov based methods. Converged equilibria using fully implicit coupled simulations are obtained for a variety of parameters. Comparisons of runtimes demonstrate that implicit methods enjoy an exponential speedup in machine time over explicit methods to obtain the same accuracy. Moreover, the equilibria obtained are shown to be independent of the timestep to machine accuracy, allowing

the use of long timesteps to investigate effects on different timescales without significantly increasing runtime.

## 8. Acknowledgments

G.P. is supported by an NSF Mathematical Sciences Postdoctoral Research Fellowship (Award No. 2303102). Performed in part by LLNL under Contract DE-AC52-07NA27344. G.P. wishes to acknowledge Andreas Holm and George Wilkie for helpful discussions.

## References

- [1] S. Krasheninnikov, A. Smolyakov, and A. Kukushkin, *On the edge of magnetic fusion devices* (Springer, 2020).
- [2] F. Militello, *Boundary Plasma Physics: An Accessible Guide to Transport, Detachment, and Divertor Design*, Vol. 123 (Springer Nature, 2022).
- [3] S. Wiesen, D. Reiter, V. Kotov, M. Baelmans, W. Dekeyser, A. Kukushkin, S. Lisgo, R. Pitts, V. Rozhansky, G. Saibene, *et al.*, *Journal of nuclear materials* **463**, 480 (2015).
- [4] W. Van Uytven, W. Dekeyser, M. Blommaert, S. Carli, and M. Baelmans, *Nuclear Fusion* **62**, 086023 (2022).
- [5] M. Blommaert, N. Horsten, P. Börner, and W. Dekeyser, *Nuclear Materials and Energy* **19**, 28 (2019).
- [6] D. Heifetz, in *Physics of Plasma-Wall Interactions in Controlled Fusion* (Springer, 1986) pp. 695–771.
- [7] D. Reiter, M. Baelmans, and P. Boerner, *Fusion science and technology* **47**, 172 (2005).
- [8] D. Stotler and C. Karney, *Contributions to Plasma Physics* **34**, 392 (1994).
- [9] A. Hakim, M. Francisquez, J. Juno, and G. W. Hammett, *Journal of Plasma Physics* **86**, 905860403 (2020).
- [10] C. Wersal, (2017).
- [11] K. Ghooos, W. Dekeyser, G. Samaey, P. Börner, and M. Baelmans, *Journal of computational physics* **322**, 162 (2016).
- [12] D. Boeyaert, S. Carli, K. Ghooos, W. Dekeyser, S. Wiesen, and M. Baelmans, *Nuclear Fusion* **63**, 016005 (2022).
- [13] R. J. LeVeque, *Finite difference methods for ordinary and partial differential equations: steady-state and time-dependent problems* (SIAM, 2007).
- [14] P. J. Roache, *Computational Fluid Dynamics* (1976).
- [15] T. Rognlien, J. Milovich, M. Rensink, and G. Porter, *J. Nuc. Mat.* **196**, 347 (1992).
- [16] T. D. Rognlien, D. D. Ryutov, N. Mattor, and G. D. Porter, *Physics of Plasmas* **6**, 1851 (1999).
- [17] D. Stotler, C. Karney, G. Wilkie, R. Kanzleiter, and S. Jaisankar, (2022).